

# On parametric resonance in the laser action <sup>1</sup>

A. I. Komech

Faculty of Mathematics, University of Vienna  
alexander.komech@univie.ac.at

## Abstract

We consider the selfconsistent semiclassical Maxwell–Schrödinger system for the solid state laser. The system consists of the Maxwell equations coupled to  $N \sim 10^{20}$  Schrödinger equations for active molecules. The system includes time-periodic pumping and a weak dissipation. We introduce the corresponding Poincaré map  $\mathcal{P}$  and consider the differential  $D\mathcal{P}(Y^0)$  at suitable ground state  $Y^0$ .

We conjecture that the *stable laser action* is due to the *parametric resonance* (PR) which means that the maximal absolute value of the corresponding multipliers is sufficiently large. The multipliers are defined as eigenvalues of  $D\mathcal{P}(Y^0)$ . The PR makes the stationary state  $Y^0$  highly unstable, and we suppose that this instability maintains the *coherent laser radiation*. We prove that the spectrum  $\text{Spec } D\mathcal{P}(Y^0)$  is approximately symmetric with respect to the unit circle  $|\mu| = 1$  if the dissipation is sufficiently small.

More detailed results are obtained for the Maxwell–Bloch system which is a finite-dimensional approximation of the Maxwell–Schrödinger system. The approximation consists of one-mode Maxwell field coupled to  $N$  two-level molecules. We calculate the corresponding Poincaré map  $P$  by successive approximations. The key role in calculation of the multipliers is played by the sum of  $N$  positive terms arising in the second-order approximation for the total current. This fact can be interpreted as the *synchronization of molecular currents* in all active molecules, which is provisionally in line with the *superradiance* and with the role of *stimulated emission* in the laser action. The calculation of the sum relies on probabilistic arguments which is one of main novelties of our approach. Other main novelties are i) the calculation of the differential  $DP$  in the “Hopf representation” which corresponds to the factorization of the dynamics by the action of the symmetry gauge group, ii) the justification of this representation, iii) the block structure of the differential, and iv) the justification of the “rotating wave approximation” by a new estimate for the averaging of slow rotations.

The main peculiarity of the eigenvalue problem is that  $DP$  is a matrix of size  $\sim N \times N$  which depends on random distribution of active molecules in the resonance cavity. The block structure of  $DP$  allows us to reduce the eigenvalue problem to polynomial equation of degree four. This reduction relies on a novel probabilistic arguments.

*MSC classification:* 78A60, 37N20, 81V80, 70K28, 35P30.

*Keywords:* laser; Maxwell–Schrödinger equations; Maxwell–Bloch equations; parametric resonance; Poincaré map; multiplier; spectrum; theorem of Lyapunov–Poincaré; perturbation theory; synchronization.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Laser equations</b>	<b>4</b>
<b>3</b>	<b>Parametric resonance</b>	<b>5</b>
<b>4</b>	<b>Symmetry of spectrum</b>	<b>6</b>
<b>5</b>	<b>Dipole approximation</b>	<b>6</b>
<b>6</b>	<b>Gauge transform</b>	<b>7</b>
<b>7</b>	<b>Decoupled system and eigenfunction expansions</b>	<b>7</b>
<b>8</b>	<b>Resonance condition</b>	<b>9</b>
<b>9</b>	<b>The harmonic pumping</b>	<b>9</b>

---

<sup>1</sup>Preprint [arXiv:2208.10179](https://arxiv.org/abs/2208.10179) [quant-ph]

<b>10</b>	<b>The Maxwell–Bloch equations</b>	<b>10</b>
<b>11</b>	<b>Molecular currents</b>	<b>10</b>
<b>12</b>	<b>Time scaling</b>	<b>11</b>
<b>13</b>	<b>Parametric resonance for the Maxwell–Bloch equations</b>	<b>12</b>
<b>14</b>	<b>Successive approximations</b>	<b>13</b>
14.1	The first-order approximation . . . . .	14
14.2	The second-order approximation . . . . .	15
<b>15</b>	<b>The Poincaré map</b>	<b>17</b>
<b>16</b>	<b>Synchronisation of molecular currents</b>	<b>18</b>
<b>17</b>	<b>Factordynamics and the Hopf fibration</b>	<b>19</b>
<b>18</b>	<b>Reduction of spectral problem by the symmetry group</b>	<b>20</b>
<b>19</b>	<b>Differential of the Poincaré map on the factorspace</b>	<b>21</b>
<b>20</b>	<b>Block-matrix approximation</b>	<b>22</b>
20.0.1	Reduction to polynomial equation . . . . .	23
<b>A</b>	<b>On averaging of slow rotations</b>	<b>25</b>
<b>B</b>	<b>Some integrals</b>	<b>27</b>
2.1	Integrals (14.26) . . . . .	27
2.2	Integrals (14.32) . . . . .	28
2.3	Integrals (15.5) . . . . .	28

## 1 Introduction

The article addresses the problems of laser and maser radiation. In the existing approaches, time-periodic pumping is not included in the dynamical equations, so the corresponding resonance features are absent. This fact was one of our main motivations.

We consider the selfconsistent semiclassical Maxwell–Schrödinger system for the solid state laser. The system consists of the Maxwell equations coupled to  $N \sim 10^{20}$  Schrödinger equations for active molecules with one active electron. The system includes time-periodic pumping and a weak dissipation. We introduce the corresponding Poincaré map  $\mathcal{P}$  and consider its differential  $D\mathcal{P}(Y^0)$  at suitable ground state  $Y^0$ . Let  $\mu$  denote the multipliers which are eigenvalues of the differential  $D\mathcal{P}(Y^0)$ . We conjecture that the laser action is due to the parametric resonance, when

$$\max_{\mu \in \text{Spec } D\mathcal{P}(Y^0)} |\mu| > 1. \quad (1.1)$$

For solutions to the linearized dynamical equation with the spectrum satisfying (1.1), the components with  $|\mu| \leq 1$  are bounded, while the ones with  $|\mu| > 1$  grow exponentially for large times. As a result, *almost all* solutions grow exponentially. This is why we expect that under the condition (1.1), similar growth takes place in the case of sufficiently small nonlinear coupling (the growth only occurs for limited time because of nonlinear effects and dissipation). Thus, our point of view is in line with the conventional interpretation of the laser action as the appearance of exponentially growing solutions of the corresponding linearized dynamics, see ([28, (6.5.12) and (6.6.23)]). The conjecture is confirmed by the fact that laser action is possible only for the pumping above a threshold that is quite similar to the parametric resonance [2].

**Remark 1.1.** We suppose that the overwhelming predominance of a single mode, corresponding to the maximal value of  $|\mu|$ , can be responsible for the *monochromatic laser radiation*.

The main goal of the present article is an analysis of properties of the multipliers and development of methods for their calculation. Our main results are as follows.

I. We develop the Hamiltonian formalism for the ideal case of the Maxwell–Schrödinger system without dissipation. Using this formalism and the Lyapunov–Poincaré theorem [41, (3.28)], we establish the approximate symmetry of the spectrum  $\text{Spec } D\mathcal{P}(Y^0)$  with respect to the unit circle  $|\mu| = 1$  when the dissipation is sufficiently small (Lemma 4.1 and (4.7)).

More detailed results (II and III below) are obtained for the Maxwell–Bloch system which is a finite-dimensional approximation of the Maxwell–Schrödinger system. The approximation consists of one-mode Maxwell field coupled to  $N$  two-level molecules. The system is time-periodic in the case of periodic pumping.

II. We calculate the corresponding Poincaré map  $P$  by successive approximations. The key role in calculation of the multipliers is played by the sum of  $N$  positive terms (16.2) arising in the second-order approximation for the total current. This fact can be interpreted as the *synchronization of molecular currents* in all active molecules, which is provisionally in line with the role of *stimulated emission* in the laser action.

III. We calculate suitable approximation for the matrix  $DP(Y^0)$  corresponding to material parameters of the ruby laser. The main peculiarity of the corresponding eigenvalue problem is that  $DP(Y^0)$  is a matrix of size  $\sim N \times N$  which depends on a random distribution of active molecules in the resonance cavity. Using the block structure of this matrix, we reduce the eigenvalue problem to polynomial equation of degree four.

Our main novelties are as follows: i) calculation of the sum (16.6) and the reduction to polynomial equation of degree six relying on Shnirelman’s Ergodic Theorem [39, 40] and probabilistic arguments (20.17)–(20.23), ii) the block structure of the differential (20.8), and iii) the justification of the “rotating wave approximation” by a new estimate for averaging of slow rotations (Lemma A.1).

**Remark 1.2.** The condition (1.1) alone is not sufficient for the stable laser/maser action: to be sure that the energy of the Maxwell field increases indeed, we check that the projections of the corresponding eigenvectors onto the Maxwell field do not vanish, see (13.8) and Remark 20.4.

Let us comment on previous theories of laser radiation. The today laser/maser theory exists on the following three levels.

*Rate equations.* This approach is based on balance equations for the numbers of emitted and absorbed photons [1, 14, 28, 33, 42].

*Semiclassical Maxwell–Schrödinger theory.* This theory is based on finite-dimensional approximations of the Schrödinger equations for each active molecule coupled to the classical Maxwell equations [1, 3, 14, 28, 33, 36, 42].

*Quantum Electrodynamics.* This approach is based on quantized Maxwell equations in the Dicke model [9, 14, 28, 33, 42] and [15]–[19].

The theory of the rate equation comes back to famous Einstein’s article [11] developing the theory of *stimulated and spontaneous emission and absorption*.

The semiclassical approach uses method of slowly varying amplitudes [28, Sections 6.2 and 6.3]. The formation of the coherent monochromatic radiation is explained by an instability in the equation for amplitude of the Maxwell mode under various hypothesis on the density of population of active molecules [28, Sections 6.5 and 6.6]. The criterion of the instability is formulated in terms of presence of an unstable (nonnegative) root of the corresponding characteristic equation ([28, (6.5.12) and (6.6.23)]).

The quantum theory of laser action relies on quantized Maxwell equations in the Dicke–Haken–Lax model [19]. This theory resulted in successful explanation of many laser phenomena: superradiance, photon echoes, self-induced transparency and others [12, 14, 28, 33, 36, 42]. The results by K. Hepp and E. Lieb [19] establish the phase transitions which occur in the thermodynamical limit as  $N \rightarrow \infty$  if the coupling constant is sufficiently large.

Note that the time-dependent pumping with a resonant frequency plays the crucial role sustaining the population inversion. At the same time, the existing theories [1, 3, 14, 27, 28, 33, 36, 42] ignore the time-dependence of the pumping, modelling the maintenance of the population inversion by *phenomenological constant terms* in the corresponding dynamical equations; see [28, (6.4.11), (6.4.12)] for the semiclassical model and the beginning of Section 7.3 in [28] for the quantum model. So, these models do not include an external periodic pumping, and the resonance features are not considered.

In conclusion, the case of quantum Maxwell field will be considered elsewhere.

## 2 Laser equations

We use the Heaviside–Lorentz units (*unrationalized Gaussian units*), in which the main physical constants (electron charge and mass, Planck’s constant, and the speed of light in vacuum) read [43, p. 221]

$$e = -4.8 \times 10^{-10} \text{esu}, \quad m = 9.1 \times 10^{-28} \text{g}, \quad \hbar = 1.055 \times 10^{-27} \text{erg} \cdot \text{s}, \quad c = 3.0 \times 10^{10} \text{cm/s}. \quad (2.1)$$

**Laser cavity and active molecules.** The laser resonator is a (usually cylindrical) cavity  $V \subset \mathbb{R}^3$  with metallic walls (ideal conductor) connected with the output waveguide. Active molecules occupy a subregion  $V_a \subset V$ . Typical industrial laser consists of two cylindrical lamps of identical size with glass walls placed at the foci of a metallic cavity  $V$  with an elliptic cross-section. The *active lamp*  $V_a$  is filled with a dielectric medium doped with identical active molecules. The other *pumping lamp* is the discharge light source which produces the pumping field. In particular, for the typical cylindrical lamp of length  $12 \text{cm}$  and diameter  $0.6 \text{cm}$  we have

$$|V_a| = \frac{12\pi \cdot 0.36}{4} \approx 3.4 \text{cm}^3. \quad (2.2)$$

The active region  $V_a$  is filled with a dielectric medium of electrical conductivity  $\sigma$ . For example, the ruby laser is filled with corundum of electrical conductivity

$$\sigma = 10^{-14} \text{Om}^{-1} \cdot \text{cm}^{-1} = 10^{-2} \text{s}^{-1}. \quad (2.3)$$

The active region  $V_a$  is filled with active molecules located at the points  $x_n$  with numbers  $n = 1, \dots, N \sim 10^{20}$ . Suppose, for simplicity of notation, that in each active molecule only one electron is involved in this interaction, and it is subject to an effective static molecular potential

$$\Phi_n(x) = \Phi(R_n(x - x_n)), \quad x \in \mathbb{R}^3, \quad R_n \in SO(3). \quad (2.4)$$

Here  $\Phi$  is the potential of the ion (or nucleus) with the total charge  $|e| > 0$ . Further we consider two different hypothesis on the distribution of  $x_n$  and  $R_n$  with  $n \in \bar{N} := (1, \dots, N)$ .

**H1. Polycrystalline medium (or ”glass medium” [37]).** The values  $x_n \in V$  and  $R_n \in SO(3)$  are uniformly distributed and independent, and almost independent from  $x_{n'}, R_{n'}$  with large  $|x_n - x_{n'}|$ .

**H2. Crystalline medium.** The values  $x_n \in V$  are uniformly distributed, while  $R_n$  do not depend on  $n \in \bar{N}$ .

**Maxwell–Schrödinger equations.** Each active molecule is described by the corresponding wave function  $\psi_n(x, t)$ , and in Heaviside–Lorentz units, the corresponding Maxwell–Schrödinger system reads (see [4, 5, 7, 13, 22, 23, 24, 25, 29, 32, 38] for the equations without dissipation)

$$\begin{cases} \frac{1}{c^2} \ddot{\mathbf{A}}(x, t) = \Delta \mathbf{A}(x, t) - \frac{\sigma}{c^2} \dot{\mathbf{A}}(x, t) + \frac{1}{c} P_s \mathbf{j}(\cdot, t), & x \in V \\ i\hbar \dot{\psi}_n(t) = H_n(t) \psi_n(t), & n = 1, \dots, N \end{cases}, \quad (2.5)$$

where  $\sigma > 0$  is electrical conductivity of the cavity medium. We assume the Coulomb gauge [20]

$$\text{div} \mathbf{A}(x, t) \equiv 0, \quad (2.6)$$

and we denote by  $P_s$  the orthogonal projection onto divergent-free vector fields in the Hilbert space  $L^2(V) \otimes \mathbb{R}^3$ . Further,

$$H_n(t) := \frac{1}{2m} D^2(t) + e\Phi_n(x). \quad (2.7)$$

Here

$$D(t) = -i\hbar \nabla - \frac{e}{c} [\mathbf{A}(x, t) + \mathbf{A}_p(x, t)], \quad (2.8)$$

where  $\mathbf{A}_p(x, t)$  is an external pumping potential, also in the Coulomb gauge and time-periodic:

$$\text{div} \mathbf{A}_p(x, t) \equiv 0; \quad \mathbf{A}_p(x, t + T) = \mathbf{A}_p(x, t), \quad t > 0, \quad (2.9)$$

where  $T > 0$ . The current density is defined by

$$\mathbf{j}(x, t) = \frac{e}{m} \sum_n \text{Re} [\bar{\psi}_n(x, t) D(t) \psi_n(x, t)]. \quad (2.10)$$

This formula neglects overlapping of the supports of the wave functions since the distance between active molecules is sufficiently large: it is of order 10 molecular diameters when the density of active molecules is of order  $10^{20}$ .

Note that the pumping field is a solution to the Maxwell equations in the cavity  $V$  excited by the discharge currents in the pumping lamp.

**Boundary conditions.** We choose the boundary conditions modelling ideally conducting diamagnetic materials (like cooper, silver, gold, etc). More precisely, we assume that

$$\mathbf{n}(x) \times \mathbf{A}(x, t) = 0, \quad \mathbf{n}(x) \cdot \text{rot} \mathbf{A}(x, t) = 0; \quad \psi_n(x, t) = 0, \quad n \in \bar{N}; \quad x \in \Gamma, \quad (2.11)$$

where  $\mathbf{n}(x)$  is the outward normal to  $\Gamma$  at a point  $x \in \Gamma$ .

**Remark 2.1.** The boundary conditions (2.11) must to be valid for the total Maxwell field  $\mathbf{A}(x, t) + \mathbf{A}_p(x, t)$ . We require the same boundary conditions for  $\mathbf{A}(x, t)$  to separate the *own field*  $\mathbf{A}(x, t)$  from the pumping field  $\mathbf{A}_p(x, t)$ . In this case the pumping field  $\mathbf{A}_p(x, t)$  satisfies the same boundary conditions as  $\mathbf{A}(x, t)$ .

Under the Dirichlet boundary conditions for the wave functions, the total charge of each active molecule is constant, i.e.,

$$\int_V |\psi_n(x, t)| dx = 1, \quad t > 0; \quad n \in \bar{N}. \quad (2.12)$$

**Hamiltonian structure.** In the *ideal case*, when  $\sigma = 0$ , the system (2.5), under the boundary conditions (2.11), is formally Hamiltonian, with the Hamiltonian functional (which is the energy up to a factor)

$$\mathcal{H}(\mathbf{A}, \mathbf{\Pi}, \boldsymbol{\psi}, t) = \frac{1}{2} [\|\frac{1}{c} \mathbf{\Pi}\|^2 + \|\text{rot} \mathbf{A}\|^2] + \frac{1}{2} \sum_n \langle \psi_n, H_n(\mathbf{A}, t) \psi_n \rangle, \quad (2.13)$$

where  $\boldsymbol{\psi} := (\psi_1, \dots, \psi_N)$ ,  $\|\cdot\|$  stands for the norm in the real Hilbert space  $L^2(\mathbb{R}^3) \otimes \mathbb{R}^3$ , and the brackets  $\langle \cdot, \cdot \rangle$  stand for the inner product in  $L^2(\mathbb{R}^3) \otimes \mathbb{C}$ . The Schrödinger operators are defined by (2.7), (2.8):

$$H_n(\mathbf{A}, t) := \frac{1}{2m} [-i\hbar \nabla - \frac{e}{c} (\mathbf{A}(x) + \mathbf{A}_p(x, t))]^2 + e\Phi_n(x). \quad (2.14)$$

The system (2.5) with  $\sigma \geq 0$  can be written in the Hamiltonian form with variational derivatives,

$$\left\{ \begin{array}{l} \frac{1}{c^2} \dot{\mathbf{A}}(t) = D_{\mathbf{\Pi}} \mathcal{H}(\mathbf{A}(t), \mathbf{\Pi}(t), \boldsymbol{\psi}(t), t), \quad \frac{1}{c^2} \dot{\mathbf{\Pi}}(t) = -D_{\mathbf{A}} \mathcal{H}(\mathbf{A}(t), \mathbf{\Pi}(t), \boldsymbol{\psi}(t), t) - \frac{\sigma}{c^2} \dot{\mathbf{A}}(x, t) \\ i\hbar \dot{\psi}_n(t) = D_{\psi_n} \mathcal{H}(\mathbf{A}(t), \mathbf{\Pi}(t), \boldsymbol{\psi}(t), t) = H_n(\mathbf{A}(t), \boldsymbol{\psi}(t), t) \psi_n(t), \end{array} \right. \quad (2.15)$$

**Symmetry group.** For any  $\theta_n \in \mathbb{R}$ , the functions  $\mathbf{A}(x, t)$ ,  $e^{i\theta_n} \psi_n(x, t)$  are solutions of the system (2.5) if  $\mathbf{A}(x, t)$ ,  $\psi_n(x, t)$  is a solution. In other words, the dynamics (2.5) under the boundary conditions (2.11) commutes with the action of the *symmetry group*  $G := [U(1)]^N$  with the action

$$(e^{i\theta_1}, \dots, e^{i\theta_N})(\mathbf{A}, \psi_1, \dots, \psi_N) = (\mathbf{A}, e^{i\theta_1} \psi_1, \dots, e^{i\theta_N} \psi_N). \quad (2.16)$$

### 3 Parametric resonance

Let us denote  $L^2 = L^2(\mathbb{R}^3)$ . The system (2.15) can be written as the dynamical equation on the Hilbert phase space  $\mathbb{Y} = [L^2(\mathbb{R}) \otimes \mathbb{R}^3]^2 \oplus [L^2]^{2N}$ :

$$\dot{Y}(t) = \mathcal{F}(Y(t), t), \quad Y(t) := (\mathbf{A}(t), \mathbf{\Pi}(t), \boldsymbol{\psi}(t)). \quad (3.1)$$

This system is  $T$ -periodic by (2.9), i.e.,

$$\mathcal{F}(Y, t+T) \equiv \mathcal{F}(Y, t). \quad (3.2)$$

The corresponding Poincaré map is defined by

$$\mathcal{P} : Y(0) \mapsto Y(T). \quad (3.3)$$

Let  $Y^0(t)$  be a “stationary state” of the system (3.1) with the zero pumping  $\mathbf{A}_p(x, t) = 0$ , i.e., a solution of type

$$Y^0(t) = (\mathbf{A}, \mathbf{\Pi}, \boldsymbol{\psi}(0) e^{-i\omega t}), \quad (3.4)$$

where  $\omega \in \mathbb{R}$ . The following definition is relevant for the system (3.1) with sufficiently small pumping.

**Definition 3.1.** The parametric resonance occurs for  $Y^0 := Y^0(0)$  when the condition (1.1) holds.

**Remark 3.2.** The parametric resonance means an exponential divergence of trajectories close to  $Y^0$ .

## 4 Symmetry of spectrum

In the *ideal case*, when  $\sigma = 0$ , the dissipation is absent, and the system (3.1) is Hamiltonian by (2.15), so it can be written as

$$\dot{Y}(t) = JD_Y \mathcal{H}(Y(t), t), \quad (4.1)$$

where  $J$  is a skewadjoint operator, and  $J^2$  is proportional to the identity operator.

**Lemma 4.1.** (Lyapunov–Poincaré theorem) *In the case  $\sigma = 0$ , the symmetry holds*

$$\text{Spec } D\mathcal{P}(Y^0) = R(\text{Spec } D\mathcal{P}(Y^0)), \quad (4.2)$$

where  $R\mu = \bar{\mu}^{-1}$  is the inversion, which is the reflection in the unit circle  $|\mu| = 1$ .

*Proof.* The differential  $D\mathcal{P}(Y^0)$  admits the representation

$$D\mathcal{P}(Y^0)\mathbf{Y}(0) = \mathbf{Y}(T), \quad (4.3)$$

where  $\mathbf{Y}(t)$  is the solution to the linearized equation

$$\dot{\mathbf{Y}}(t) = K(t)\mathbf{Y}(t), \quad K(t) = D_Y \mathcal{F}(Y^0(t), t). \quad (4.4)$$

However, (4.1) implies that

$$\mathcal{F}(Y, t) = JD_Y \mathcal{H}(Y, t),$$

and therefore,

$$K(t) = JD_Y^2 \mathcal{H}(Y^0(t), t) = JA(t), \quad (4.5)$$

where  $A(t)$  is the selfadjoint operator. Denote the map  $U(t) : \mathbf{Y}(0) \rightarrow \mathbf{Y}(t)$ , so  $D\mathcal{P}(X^0) = U(2\pi)$  by (4.3). Finally, the Lyapunov–Poincaré arguments [41, (3.28)] imply that  $\text{Spec } U^* = \text{Spec } U^{-1}$ , where  $U = U(2\pi)$ . Namely, we have  $\dot{U}(t) = JA(t)U(t)$ . Hence, differentiating  $W(t) := U^*(t)JU(t)$ , we obtain

$$\dot{W} = \dot{U}^*JU + U^*J\dot{U} = U^*A(-J^2)U + U^*J^2AU = 0 \quad (4.6)$$

since  $J^2$  commutes with  $A$ . Therefore,  $U^*(t)JU(t) = J$ , and hence  $U^*(t) = JU^{-1}(t)J^{-1}$ . This similarity implies that the spectra of  $U^*(t)$  and  $U^{-1}(t)$  coincide. Now the symmetry (4.2) is proved.  $\square$

**Remark 4.2.** The symmetry (4.2) can be considered as a nonlinear version of the Lyapunov–Poincaré theorem, which plays the key role in M.G. Krein’s theory of parametric resonance [8, 41]. The ground for this extension is the Hamiltonian structure of the dynamics (4.1).

Thus, in the ideal case, the set of multipliers is symmetric with respect to the unit circle. In general case, when the dissipation is included, the symmetry (4.2) is broken, though it holds approximately for sufficiently small  $\sigma > 0$ :

$$\text{Spec } D\mathcal{P}(Y^0) \approx R(\text{Spec } D\mathcal{P}(Y^0)). \quad (4.7)$$

**Remark 4.3.** Similar symmetry holds for the case of the Maxwell–Bloch equations which are finite-dimensional approximation of the Maxwell–Schrödinger equations, see (12.9) below. This symmetry is important by the following reason. Our ultimate goal is the reduction of the corresponding finite-dimensional eigenvalue problem to a polynomial of a lower degree (see Section 20.0.1 below). In the ideal case, when  $\sigma = 0$ , the symmetry of type (4.2) allows us to reduce the degree twice. We expect that even for the non-ideal case, the approximate symmetry of type (4.7) also can help in similar reduction.

## 5 Dipole approximation

The theory admits a significant simplification in the case when the wavelength of the pumping is large with respect to the size of active molecules  $D$ .

**H3. The dipole radiation condition.** We will assume that

$$\lambda = cT = 2\pi c/\Omega_p \gg D, \quad (5.1)$$

where  $\Omega_p = 2\pi/T$  is the frequency of the pumping,

**Example 5.1.** This condition holds for the ruby laser since the wavelength  $\lambda \approx 7 \times 10^{-5} \text{ cm}$  while the size of the chromium molecule is

$$D \approx 2.5 \text{ pm} = 2.5 \times 10^{-10} \text{ cm}. \quad (5.2)$$

Taking into account the condition (5.1), we approximate the system (2.5) by

$$\left\{ \begin{array}{l} \frac{1}{c^2} \ddot{\mathbf{A}}(x, t) = \Delta \mathbf{A}(x, t) - \frac{\sigma}{c} \dot{\mathbf{A}}(x, t) + \frac{1}{c} P_3 \hat{\mathbf{j}}(x, t) \\ i\hbar \dot{\psi}_n(x, t) = \hat{H}_n(t) \psi_n(t), \quad n = 1, \dots, N \end{array} \right\}, \quad x \in V. \quad (5.3)$$

Here

$$\hat{H}_n(t) := \frac{1}{2m} D_n^2(t) + e\Phi_n(x), \quad D_n(t) = -i\hbar \nabla - \frac{e}{c} [\mathbf{A}(x, t) + \mathbf{A}_p(x_n, t)], \quad (5.4)$$

The current (2.10) in the dipole approximation becomes

$$\mathbf{j}(x, t) \approx \frac{e}{m} \text{Re} \sum_n \bar{\psi}_n(x, t) D_n(t) \psi_n(x, t). \quad (5.5)$$

The system (5.3) with  $\sigma = 0$  is also Hamiltonian with the Hamilton functional

$$\hat{\mathcal{H}}(\mathbf{A}, \mathbf{\Pi}, \psi, t) = \frac{1}{2} \left[ \|\frac{1}{c} \mathbf{\Pi}\|^2 + \|\text{rot } \mathbf{A}\|^2 \right] + \frac{1}{2} \sum_n \langle \psi_n, \hat{H}_n(\mathbf{A}, t) \psi_n \rangle, \quad (5.6)$$

where

$$\hat{H}_n(\mathbf{A}, t) := \frac{1}{2m} D_n^2(\mathbf{A}, t) + e\Phi_n(x), \quad D_n(\mathbf{A}, t) = -i\hbar \nabla - \frac{e}{c} [\mathbf{A}(x) + \mathbf{A}_p(x_n, t)], \quad (5.7)$$

## 6 Gauge transform

The dipole approximation (5.3) allows us to apply the gauge transform [36, (5.1.15)]:

$$\psi_n(x, t) = e^{i\chi_n(x, t)} \tilde{\psi}_n(x, t), \quad \chi_n(x, t) := \frac{e}{\hbar c} (x - x_n) [\mathbf{A}(x_n, t) + \mathbf{A}_p(x_n, t)]. \quad (6.1)$$

Now the dipole approximation (5.3) becomes

$$\left\{ \begin{array}{l} \frac{1}{c^2} \ddot{\mathbf{A}}(x, t) = \Delta \mathbf{A}(x, t) - \frac{\sigma}{c} \dot{\mathbf{A}}(x, t) + \frac{1}{c} P \mathbf{j}(x, t) \\ i\hbar \dot{\tilde{\psi}}_n(x, t) = \tilde{H}_n(t) \tilde{\psi}_n(t), \quad n = 1, \dots, N \end{array} \right\}, \quad x \in V. \quad (6.2)$$

The Schrödinger operator now reads

$$\tilde{H}_n(t) = -\frac{1}{2m} (-i\hbar \nabla - \frac{e}{c} [\mathbf{A}(x, t) - \mathbf{A}(x_n, t)])^2 + e\Phi_n(x) + \hbar \dot{\chi}_n(x, t) \approx -\frac{\hbar^2}{2m} \Delta + e\Phi_n(x), \quad (6.3)$$

where we used the condition (5.1). The current (5.5) now reads as

$$\mathbf{j}(x, t) \approx \frac{e}{m} \text{Re} \sum_n \bar{\tilde{\psi}}_n(x, t) (-i\hbar \nabla - \frac{e}{c} [\mathbf{A}(x, t) - \mathbf{A}(x_n, t)]) \tilde{\psi}_n(x, t) \approx \frac{e}{m} \text{Re} \sum_n \bar{\tilde{\psi}}_n(x, t) (-i\hbar \nabla) \tilde{\psi}_n(x, t). \quad (6.4)$$

## 7 Decoupled system and eigenfunction expansions

**Decoupled system.** The system (3.1) is a weak perturbation of the corresponding “decoupled” system obtained by neglecting the interaction terms with the coupling constant  $e/c$ :

$$\left\{ \begin{array}{l} \dot{\mathbf{A}}(x, t) = \mathbf{\Pi}(x, t) \\ \frac{1}{c^2} \dot{\mathbf{A}}(x, t) = \Delta \mathbf{A}(x, t) - \frac{\sigma}{c^2} \dot{\mathbf{A}}(x, t) \\ i\hbar \dot{\psi}_n(t) = H_n^0 \psi_n(t), \quad n \in \bar{N} \end{array} \right\}, \quad (7.1)$$

where

$$H_n^0 := -\frac{\hbar^2}{2m}\Delta + e\Phi_n(x). \quad (7.2)$$

**Schrödinger modes.** The decoupled system (7.1) can be solved explicitly using the orthonormal eigenfunctions of the Schrödinger operators  $H_n^0$  under the Dirichlet boundary conditions (2.11):

$$H_n^0 \varphi_{n,l}(x) = \omega_{n,l} \varphi_{n,l}(x), \quad l = 1, 2, \dots \quad (7.3)$$

Hence, the system (3.1) can be analyzed by perturbation methods. The eigenfunctions  $\varphi_{nl}$  differ by rotations (2.4), while the eigenvalues  $\omega_{n,l} = \omega_l$  do not depend on  $n$  with a great accuracy. The decoupled system (3.1) admits the “stationary” solutions

$$Z^0(t) \equiv (0, 0, (e^{-i\omega_{l(n)}t} \varphi_{n,l(n)} : n \in \bar{N})), \quad (7.4)$$

where  $l(n) = 1, 2, \dots$

**Maxwell modes.** The first equation of the decoupled system (7.1) can be solved using the orthonormal eigenmodes

$$\left\{ \begin{array}{l} \Delta \mathbf{X}_v(x) = -\frac{\Omega_v^2}{c^2} \mathbf{X}_v(x), \quad \text{div } \mathbf{X}_v(x) = 0, \quad x \in V \\ \mathbf{n}(x) \times \mathbf{X}_v(x) = 0, \quad \mathbf{n}(x) \cdot \text{rot } \mathbf{X}_v(x) = 0, \quad x \in \partial V \end{array} \right\}, \quad (7.5)$$

and  $\Omega_v > 0$ .

**Example 7.1.** The eigenmodes can be calculated explicitly for the rectangular cuboid  $V = [0, l_1] \times [0, l_2] \times [0, l_3]$ :

$$\mathbf{X}_k = C \begin{pmatrix} a_k^1 \cos \frac{k_1 \pi x_1}{l_1} \sin \frac{k_2 \pi x_2}{l_2} \sin \frac{k_3 \pi x_3}{l_3} \\ a_k^2 \sin \frac{k_1 \pi x_1}{l_1} \cos \frac{k_2 \pi x_2}{l_2} \sin \frac{k_3 \pi x_3}{l_3} \\ a_k^3 \sin \frac{k_1 \pi x_1}{l_1} \sin \frac{k_2 \pi x_2}{l_2} \cos \frac{k_3 \pi x_3}{l_3} \end{pmatrix}, \quad C = \sqrt{\frac{8}{l_1 l_2 l_3}}, \quad k = (k_1, k_2, k_3), \quad k_j = 1, 2, \dots, \quad (7.6)$$

where the amplitudes  $a_k = (a_k^1, a_k^2, a_k^3)$  are unit vectors, orthogonal to the wave vectors  $k$ . In particular, consider the cuboid  $V$  with dimensions  $l_1 = 12\text{cm}$ ,  $l_2 = l_3 = 2\text{cm}$ . Then “the dispersion” of the “random value”  $\mathbf{X}_k(x_n)$  on the probability space  $\{n \in \bar{N}\}$  is

$$D(\mathbf{X}(x_n)) = \frac{1}{N} \sum \mathbf{X}_k^2(x_n) \approx \frac{1}{|V_a|} \int_{V_a} \mathbf{X}_k^2(x) dx \approx \frac{1}{|V_a|} \frac{|V_a|}{|V|} = \frac{1}{|V|} \approx 0.02. \quad (7.7)$$

Accordingly, the mean value is

$$|\mathbf{X}(x_n)| \approx 0.14. \quad (7.8)$$

**Eigenfunction expansions.** We will expand  $\tilde{\psi}_n(\cdot, t)$  in the orthonormal eigenfunctions (7.3):

$$\tilde{\psi}_n(x, t) = \sum_{l=1}^{\infty} c_{n,l}(t) e^{-i\omega_l t} \varphi_{n,l}(x) = e^{-\frac{i}{\hbar} H_n^0 t} \phi_n(t), \quad \phi_n(t) := \sum_{l=1}^{\infty} c_{n,l}(t) \varphi_{n,l}(x). \quad (7.9)$$

Now each Schrödinger equation in (6.2) with  $n \in \bar{N}$  is equivalent to the system

$$\dot{c}_{n,l}(t) = -i \sum_{l'} e^{-i\omega_{l'} t} \langle \dot{\chi}_n(x, t) \varphi_{n,l'}(x), \varphi_{n,l}(x) \rangle c_{n,l'}(t), \quad l \geq 1; \quad \omega_{l,l'} = \omega_{l'} - \omega_l. \quad (7.10)$$

Similarly, the Maxwell field can be expanded in the orthonormal eigenmodes (7.5):

$$\mathbf{A}(x, t) = \sum_v a_v(t) \mathbf{X}_v(x). \quad (7.11)$$

Now the Maxwell equation in (6.2) is equivalent to the system

$$\ddot{a}_v(t) + c\sigma \dot{a}_v(t) + \Omega_v^2 a_v(t) = c \langle \mathbf{j}(t), \mathbf{X}_v \rangle, \quad v = 1, \dots \quad (7.12)$$



## 8 Resonance condition

The resonance condition means that

$$\Delta = \Omega_p = \Omega_{\bar{\nu}}, \quad \Delta := \omega_a - \omega_b, \quad (8.1)$$

where  $\omega_b < \omega_a$  are some fixed eigenvalues (7.3),  $\Omega_{\bar{\nu}}$  is one of the Maxwell eigenfrequencies (7.5), and  $\Omega_p = 2\pi/T$  is the frequency of the pumping (2.9).

We can renumerate the eigenfunctions  $\varphi_{n,l}$  in such a way that  $b = 1$  and  $a = 2$ . Then the decoupled system (7.1) admits solutions (7.4) with  $l(n) \equiv 1$ :

$$Y^0(t) \equiv (0, 0, (e^{-i\omega_l t} \varphi_{n,1} : n \in \bar{N})), \quad (8.2)$$

Let us call as ‘‘ground states’’ all values of these solutions:

$$Y^0 = (0, 0, (\varphi_{n,1} : n \in \bar{N})). \quad (8.3)$$

The perturbation theory allows us to calculate solutions to the coupled system (2.5) with initial states close to (8.3). We expect that these solutions are rapidly growing due to the resonance condition (8.1) and the parametric resonance (1.1).

## 9 The harmonic pumping

We consider the Maxwell–Schrödinger system (2.5) with the pumping  $\mathbf{A}_p(x, t)$ , which is switched on at  $t = 0$  and is monochromatic, i.e.,

$$\mathbf{A}_p(x, t) = \mathbf{a}_p(x) \sin \Omega_p t, \quad (9.1)$$

where  $\Omega_p$  is the pumping frequency. For example, for the ruby laser the wavelength and the resonant pumping frequency are

$$\lambda = 694.3 \text{ nm} = 694.3 \times 10^{-9} \text{ m} \approx 7 \times 10^{-5} \text{ cm}, \quad \Omega_p = \frac{2\pi c}{\lambda} = \frac{19 \times 10^{10}}{7 \times 10^{-5}} \approx 3 \times 10^{15} \text{ s}^{-1}. \quad (9.2)$$

**The magnitude of pumping.** The pumping field in a neighborhood of any active molecule is approximately a plane wave

$$\mathbf{A}_p(x, t) \approx \mathbf{a}_p \sin(\Omega_p t - kx), \quad |k| = |\Omega_p|/c, \quad (9.3)$$

or a mixture of such waves. Let us estimate the pumping field in a typical cylindrical cavity of length 12 cm and diameter 0.6 cm illuminated by the pumping lamp of the typical power  $W = 1 \text{ Kw}t = 10^{10} \text{ erg/s}$ . The side surface  $S$  of the cavity is about  $12\pi \cdot 0.6 \text{ cm}^2 \approx 23 \text{ cm}^2$ . The intensity  $I$  of the plane wave (9.3) is  $\frac{ck^2 |\mathbf{a}_p|^2}{2} \frac{\text{erg/s}}{\text{cm}^2}$ , so

$$W = IS = \frac{ck^2 |\mathbf{a}_p|^2}{2} 12\pi \cdot 0.6 = 10^{10}. \quad (9.4)$$

Substituting  $|k| = |\Omega_p|/c$ , we find

$$\frac{\Omega_p^2 |\mathbf{a}_p|^2}{2c} 23 = 10^{10}. \quad (9.5)$$

For example, for the ruby laser with frequency (9.2), we have

$$|\mathbf{a}_p|^2 \approx \frac{c}{\Omega_p^2} \times 10^9 = 0.3 \times 10^{-11}. \quad (9.6)$$

Finally,

$$|\mathbf{a}_p| \approx 1.7 \times 10^{-6} \text{ esu/cm}. \quad (9.7)$$

**H4. Distribution of pumping.** We assume that i)  $\mathbf{a}_p(x_n)$  is the random vector distributed uniformly on the sphere  $|\mathbf{a}_p(x)| = a_p = 1.7 \times 10^{-6}$ , and ii)  $\mathbf{a}_p(x_n)$  and  $\mathbf{a}_p(x_{n'})$  are almost independent at large distances  $|x_n - x_{n'}|$ .

## 10 The Maxwell–Bloch equations

Now we consider the finite-dimensional Faedo–Galerkin approximation of the Maxwell–Schrödinger system (3.1). Namely, we will consider solutions

$$\mathbf{A}(x, t) = a(t)\mathbf{X}_{\bar{v}}(x), \quad \tilde{\Psi}_n(x, t) = c_{n,1}(t)\varphi_{n,1}(x) + c_{n,2}(t)\varphi_{n,2}(x) \quad (10.1)$$

which satisfy the system consisting of two equations (7.10) with  $n = 1$  and  $n = 2$  for each  $n \in \bar{N}$ , and one equation of (7.12) with  $v = \bar{v}$ . Below, we will write  $a(t)$ ,  $\Omega(\tau)$  and  $\mathbf{X}(x)$  instead of  $a_{\bar{v}}(t)$ ,  $\Omega_{\bar{v}}(\tau)$  and  $\mathbf{X}_{\bar{v}}(x)$  respectively. Thus, the approximation of the Maxwell–Schrödinger equations reads as the Maxwell–Bloch equations

$$\ddot{a}(t) + c\sigma\dot{a}(t) + \Omega^2 a(t) = c\langle \mathbf{j}(t), \mathbf{X} \rangle, \quad (10.2)$$

$$\left\{ \begin{array}{l} \dot{c}_{n,1}(t) = -i\sum_{l'=1}^2 e^{-i\omega_{1,l'}t} \langle \dot{\chi}_n(x, t)\varphi_{n,l'}(x), \varphi_{n,1}(x) \rangle c_{n,l'}(t) \\ \dot{c}_{n,2}(t) = -i\sum_{l'=1}^2 e^{-i\omega_{2,l'}t} \langle \dot{\chi}_n(x, t)\varphi_{n,l'}(x), \varphi_{n,2}(x) \rangle c_{n,l'}(t) \end{array} \right\}, \quad n \in \bar{N}. \quad (10.3)$$

## 11 Molecular currents

By (5.1), the matrix entries in (10.3) read as

$$\begin{aligned} b_n^{ll'}(t) &:= \langle \dot{\chi}_n(x, t)\varphi_{n,l'}(x), \varphi_{n,l}(x) \rangle \approx \frac{e}{\hbar c} [\dot{\mathbf{A}}(x_n, t) + \dot{\mathbf{A}}_p(x_n, t)] \langle (x - x_n)\varphi_{n,l'}(x), \varphi_{n,l}(x) \rangle \\ &= \frac{e}{\hbar c} [\dot{a}_1(t)\mathbf{X}(x_n) + \mathbf{a}_p(x_n)\Omega_p \cos \Omega_p t] \mathbf{P}_n^{ll'} \\ &= \frac{\mathbf{P}_n^{ll'}}{\hbar c} [\dot{a}_1(t)\mathbf{X}(x_n) + \mathbf{a}_p(x_n)\Omega_p \cos \Omega_p t], \quad l, l' = 1, 2, \end{aligned} \quad (11.1)$$

where  $\mathbf{P}_n^{ll'}$  are the dipole momenta

$$\mathbf{P}_n^{ll'} := e \langle \varphi_{n,l'}(x), (x - x_n)\varphi_{n,l}(x) \rangle. \quad (11.2)$$

These momenta are real vectors since we have chosen real eigenfunctions  $\varphi_{n,l}(x)$ .

**Remark 11.1.** In the case of the ruby laser, the matrix of type (11.2) corresponds to the dipole momenta of chromium ions  $\text{Cr}^{3+}$ . For definiteness of calculations, we accept

$$|\mathbf{P}| \approx 4 \text{ Debye} = 4 \times 10^{-18} \text{ esu} \cdot \text{cm}, \quad (11.3)$$

which is the *permanent dipole moment* of chromium oxide molecules  $\text{CrO}$ ; see [10, p. 10]. However, actual magnitude of the matrix elements (11.2) in the case of the ruby laser must be revised.

From now on, we will assume that the potential  $\Phi(x)$  is an even function, i.e.,

$$\Phi(-x) = \Phi(x), \quad x \in \mathbb{R}^3. \quad (11.4)$$

In this case, all eigenfunctions  $\varphi_{n,l}$  can be chosen odd or even, and hence,

$$\mathbf{P}_n^{11} = \mathbf{P}_n^{22} = 0. \quad (11.5)$$

Therefore, the system (10.3) reads as

$$\left\{ \begin{array}{l} \dot{c}_{n,1}(t) = -ib_n(t)e^{-i\Delta t} c_{n,2}(t) \\ \dot{c}_{n,2}(t) = -ib_n(t)e^{i\Delta t} c_{n,1}(t) \end{array} \right\}, \quad n \in \bar{N}, \quad (11.6)$$

where we denote

$$b_n(t) := \frac{1}{\hbar c} [\dot{a}_1(t)\mathbf{P}_n\mathbf{X}_1(x_n) + \mathbf{P}_n\mathbf{a}_p(x_n)\Omega_p \cos \Omega_p t], \quad \mathbf{P}_n := \mathbf{P}_n^{12}. \quad (11.7)$$

The matrix of this system is skew-symmetric, which corresponds to the electron charge conservation in each active molecule i.e.,

$$|c_{n,1}(t)|^2 + |c_{n,2}(t)|^2 = 1, \quad t > 0, \quad n \in \bar{N}. \quad (11.8)$$

We now calculate the current (6.4) corresponding to the wave functions (10.1), and the right hand side of (7.12) with  $\mathbf{X}_v = \mathbf{X}$ : taking into account the dipole approximation (5.1), we obtain that

$$\begin{aligned} \langle \mathbf{j}(t), \mathbf{X} \rangle &\approx \frac{e\hbar}{m} \sum_n \mathbf{X}(x_n) \text{Im} \left\{ \bar{c}_{n,1}(t) c_{n,2}(t) e^{-i\Delta t} \int \bar{\varphi}_{n,1}(x) \nabla \varphi_{n,2}(x) dx + \bar{c}_{n,2}(t) c_{n,1}(t) e^{i\Delta t} \int \bar{\varphi}_{n,2}(x) \nabla \varphi_{n,1}(x) dx \right\} \\ &= 2\Delta \sum_n \mathbf{P}_n \mathbf{X}(x_n) \text{Im} \left\{ \bar{c}_{n,1}(t) c_{n,2}(t) e^{-i\Delta t} \right\}, \end{aligned} \quad (11.9)$$

since

$$\int \bar{\varphi}_{n,1} \nabla \varphi_{n,2} dx = \frac{\Delta m}{\hbar} \int \bar{\varphi}_{n,1} x \varphi_{n,2} dx = \frac{\Delta m \mathbf{P}_n}{\hbar e} \quad (11.10)$$

by [35, (44.20)].

## 12 Time scaling

Now the Maxwell–Bloch equations (10.2)–(10.3) become

$$\ddot{a}(t) + \Omega^2 a(t) + c\sigma \dot{a}(t) = j(t) := 2c\Delta \sum_n \mathbf{P}_n \mathbf{X}(x_n) \text{Im} \left\{ \bar{c}_{n,1}(t) c_{n,2}(t) e^{-i\Delta t} \right\}, \quad (12.1)$$

$$\left\{ \begin{array}{l} \dot{c}_{n,1}(t) = -ib_n(t) e^{-i\Delta t} c_{n,2}(t) \\ \dot{c}_{n,2}(t) = -ib_n(t) e^{i\Delta t} c_{n,1}(t) \end{array} \right\}, \quad n \in \bar{N}. \quad (12.2)$$

Let us transform the system (12.1) to the natural time scale

$$\tau := \Omega_p t \approx 3 \times 10^{15} t, \quad (12.3)$$

in which the period of pumping is  $2\pi$ . Below, we denote any function  $f(t)$  in the units of time (12.3) also as  $f(\tau)$ . In particular, (11.7) becomes

$$b_n(\tau) = \frac{\Omega_p}{\hbar c} [\partial_\tau a(\tau) \mathbf{P}_n \mathbf{X}(x_n) + \mathbf{P}_n \mathbf{a}_p(x_n) \cos \tau]. \quad (12.4)$$

Now the Maxwell–Bloch equations (12.1), (12.2) read as

$$\partial_\tau^2 a(\tau) + \sigma_1 \partial_\tau a(\tau) + \frac{\Omega^2}{\Omega_p^2} a(\tau) = j(\tau), \quad \left\{ \begin{array}{l} \partial_\tau c_{n,1}(\tau) = -i \frac{b_n(\tau)}{\Omega_p} e^{-i \frac{\Delta}{\Omega_p} \tau} c_{n,2}(\tau) \\ \partial_\tau c_{n,2}(\tau) = -i \frac{b_n(\tau)}{\Omega_p} e^{i \frac{\Delta}{\Omega_p} \tau} c_{n,1}(\tau) \end{array} \right\}, \quad n \in \bar{N} \quad (12.5)$$

where

$$j(\tau) := \frac{2c\Delta}{\Omega_p^2} \sum_n \mathbf{P}_n \mathbf{X}(x_n) \text{Im} \left\{ \bar{c}_{n,1}(\tau) c_{n,2}(\tau) e^{-i \frac{\Delta}{\Omega_p} \tau} \right\} \quad (12.6)$$

and

$$\sigma_1 := \frac{c\sigma}{\Omega_p} \approx \sigma 10^{-5} \approx 10^{-7}. \quad (12.7)$$

Let us denote the real constants

$$\alpha_n = \frac{2c}{\Omega_p} \mathbf{P}_n \mathbf{X}(x_n), \quad \beta_n = \frac{\mathbf{P}_n \mathbf{X}(x_n)}{\hbar c}, \quad \gamma_n = \frac{\mathbf{P}_n \mathbf{a}_p(x_n)}{\hbar c}. \quad (12.8)$$

Then the system (12.5) becomes

$$\ddot{a}(\tau) + \sigma_1 \dot{a}(\tau) + a(\tau) = j(\tau), \quad \left\{ \begin{array}{l} \dot{c}_{n,1}(\tau) = -i\omega_n(\tau) c_{n,2}(\tau) \\ \dot{c}_{n,2}(\tau) = -i\bar{\omega}_n(\tau) c_{n,1}(\tau) \end{array} \right\}, \quad n \in \bar{N}, \quad (12.9)$$

where we denote  $\dot{a} := \partial_\tau a$ , etc, and

$$j(\tau) = \sum_n \alpha_n \operatorname{Im} \left\{ \bar{c}_{n,1}(\tau) c_{n,2}(\tau) e^{-i\tau} \right\}, \quad \omega_n(\tau) = [\beta_n \dot{a}(\tau) + \gamma_n \cos \tau] e^{-i\tau}, \quad (12.10)$$

where we used that  $\Delta = \Omega_p$  according to (8.1). The matrix of system (12.9) is skew-adjoint, and the charge conservation holds: similarly to (11.8),

$$|c_{n,1}(\tau)|^2 + |c_{n,2}(\tau)|^2 = 1, \quad \tau \in \mathbb{R}. \quad (12.11)$$

All coefficients  $\alpha_n$ ,  $\beta_n$  and  $\gamma_n$  are small. In particular, for the ruby laser with the chromium dipole moment (11.3), and the pumping frequency (9.2), we have

$$\frac{c|\mathbf{P}_n|}{\Omega_p} \sim \frac{12 \times 10^{10} 10^{-18}}{3 \times 10^{15}} \sim 4 \times 10^{-23}, \quad \frac{|\mathbf{P}_n|}{\hbar c} \sim \frac{4 \times 10^{-18}}{10^{-27} 3 \times 10^{10}} \sim 1.3 \times 10^{-1}. \quad (12.12)$$

Now (7.8) and (9.7) imply that

$$|\alpha_n| \sim 10^{-23}, \quad |\beta_n| \sim 0.2 \times 10^{-2}, \quad |\gamma_n| \sim 0.7 \times 10^{-7}. \quad (12.13)$$

In all calculations below we will consider these values.

**Remark 12.1.** By our hypotheses **H1–H4**, the parameters  $\alpha_n, \beta_n, \gamma_n$  and  $\alpha_{n'}, \beta_{n'}, \gamma_{n'}$  are almost independent for sufficiently large distances  $|x_{n'} - x_n|$ .

### 13 Parametric resonance for the Maxwell–Bloch equations

Let us reformulate the concept of parametric resonance (1.1) for the system (12.9). In the case of zero pumping  $\mathbf{a}_p = 0$ , this system admits “ground states”, which are solutions corresponding to (8.2):

$$X^0(t) \equiv X^0 := (a^0, b^0, (c_n^0 : n \in \bar{N})) \quad (13.1)$$

with

$$a^0 = b^0 = 0; \quad c_n^0 = (e^{i\theta_n}, 0), \quad n \in \bar{N}. \quad (13.2)$$

The solutions describe the laser dynamics before the pumping is switched on. Let us write the system (12.9) as

$$\dot{a}(\tau) = b(\tau), \quad \dot{b}(\tau) + \sigma_1 b(\tau) + a(\tau) = j(\tau), \quad \dot{c}_n(\tau) = -i\Omega_n(\tau) c_n(\tau), \quad n \in \bar{N}, \quad (13.3)$$

where we denote

$$c_n(\tau) = (c_{n,1}(\tau), c_{n,2}(\tau)), \quad \Omega_n(\tau) = \begin{pmatrix} 0 & \omega_n(\tau) \\ \bar{\omega}_n(\tau) & 0 \end{pmatrix}, \quad \omega_n(\tau) = [\beta_n b(\tau) + \gamma_n \cos \tau] e^{-i\tau}. \quad (13.4)$$

The system (13.3) can be written as

$$\dot{X}(\tau) = F(X(\tau), \tau), \quad X(\tau) := (a(\tau), b(\tau), (c_n(\tau) : n \in \bar{N})). \quad (13.5)$$

This system (13.5) is  $2\pi$ -periodic, i.e.,

$$F(X, \tau + 2\pi) \equiv F(X, \tau). \quad (13.6)$$

The corresponding Poincaré map is defined by

$$P : X(0) \mapsto X(2\pi). \quad (13.7)$$

The following definition is relevant for the system (13.5) in the case of sufficiently small pumping.

**Definition 13.1.** The parametric resonance occurs for the ground state (13.1) when the condition (1.1) (with  $DP(X^0)$  instead of  $D\mathcal{P}(Y^0)$ ) holds with a sufficiently large  $\rho > 0$ .

We also need to check the additional condition for all the corresponding eigenvectors  $\mathbf{X} := (\mathbf{a}, \mathbf{b}, (\mathbf{c}_n : n \in \bar{N}))$ :

$$(\mathbf{a}, \mathbf{b}) \neq 0, \quad (13.8)$$

which guarantees an exponential increment of the Maxwell field. We have verified this condition in the case of ruby laser; see Remark 20.4.

## 14 Successive approximations

We will construct the Poincaré map (13.7) calculating successive approximations for the equations (13.3) on the interval  $[0, 2\pi]$  with initial data

$$a(0), \quad b(0); \quad c_n(0), \quad n \in \overline{N}, \quad (14.1)$$

which are close to the set of stationary states (13.2), i.e.,

$$|a(0)| + |b(0)| + \sum_n ||c_{n,1}(0)| - 1| + |c_{n,2}(0)| \leq \varepsilon, \quad (14.2)$$

where  $\varepsilon > 0$  is sufficiently small. As the *zero-order approximation*, we choose the stationary solutions (13.1):

$$a^{(0)}(\tau) \equiv b^{(0)}(\tau) \equiv 0; \quad c_{n,1}^{(0)}(\tau) \equiv e^{-i\theta_n}, \quad c_{n,2}^{(0)}(\tau) \equiv 0 \quad (14.3)$$

since we plan to study instability of solutions which are close to these stationary solutions. The higher-order approximations are defined as follows.

**The Schrödinger amplitudes.** We define the approximations  $c_n^{(k)}$  with  $k = 1, \dots$  as solutions of equations

$$\dot{c}_n^{(k)}(\tau) = -i\Omega_n^{(k-1)}(\tau)c_n^{(k)}(\tau), \quad n \in \overline{N}, \quad (14.4)$$

with the initial data (14.1),

$$c_n^{(k)}(0) = c_n(0), \quad n \in \overline{N}. \quad (14.5)$$

Here we denote

$$\Omega_n^{(k-1)}(\tau) = \begin{pmatrix} 0 & \omega_n^{(k-1)}(\tau) \\ \overline{\omega}_n^{(k-1)}(\tau) & 0 \end{pmatrix}, \quad \omega_n^{(k-1)}(\tau) = [\beta_n b^{(k-1)}(\tau) + \gamma_n \cos \tau] e^{-i\tau}. \quad (14.6)$$

To solve the Schrödinger equations (14.4), we will replace them by the corresponding averaged equations

$$\dot{c}_n^{(k)}(\tau) = -i\tilde{\Omega}_n^{(k-1)} c_n^{(k)}(\tau), \quad n \in \overline{N}, \quad (14.7)$$

where

$$\tilde{\Omega}_n^{(k-1)} := \frac{1}{2\pi} \int_0^{2\pi} \Omega_n^{(k-1)}(\tau) d\tau. \quad (14.8)$$

The matrix functions  $\Omega_n^{(k-1)}(\tau)$  are sufficiently small by (12.13), so the replacement is justified by Lemma A.1.

All the matrices (14.6) and (14.8) are skewsymmetric, and hence,

$$c_n^{(k)}(\tau) = U_n^{(k)}(\tau) c_n(0), \quad (14.9)$$

where  $U_n^{(k)}(\tau)$  is a unitary matrix.

**Remark 14.1.** The last formula does not mean that the map  $c_n(0) \mapsto c_n^{(k)}(\tau)$  is linear since the matrix  $U_n^{(k)}(\tau)$  generally depends on  $c_n(0)$  as well as on all initial data (14.1).

**The Maxwell amplitude.** We define the approximations  $a^{(k)}(\tau)$  with  $k = 1, \dots$  as solutions of the equations

$$\ddot{a}^{(k)}(\tau) + \sigma_1 \dot{a}^{(k)}(\tau) + a^{(k)}(\tau) = j^{(k)}(\tau), \quad j^{(k)}(\tau) := \sum_n \alpha_n \operatorname{Im} \left\{ \overline{c}_{n,1}^{(k)}(\tau) c_{n,2}^{(k)}(\tau) e^{-i\tau} \right\}, \quad (14.10)$$

with the initial data (14.1):

$$a^{(k)}(0) = a(0), \quad \dot{a}^{(k)}(0) = b(0). \quad (14.11)$$

Let us denote  $\varkappa := \sigma_1/2 \approx 10^{-7}$ . The solution of the Maxwell equation (14.10) is given by the convolution

$$a^{(k)}(\tau) = a_0(\tau) + \int_0^\tau j^{(k)}(\tau') E(\tau - \tau') d\tau', \quad a_0(\tau) := a(0) \dot{E}(\tau) + [\dot{a}(0) + 2\varkappa a(0)] E(\tau), \quad (14.12)$$

where  $E(\tau)$  is the retarded fundamental solution,

$$\ddot{E}(\tau) + 2\kappa\dot{E}(\tau) + E(\tau) = \delta(\tau), \quad \tau \in \mathbb{R}; \quad E(\tau) = 0, \quad \tau < 0. \quad (14.13)$$

This fundamental solution is expressed as follows. Denote by  $\lambda_{\pm}$  the roots of characteristic equation

$$\lambda^2 + 2\kappa\lambda + 1 = 0 \quad (14.14)$$

which are

$$\lambda_{\pm} = -\kappa \pm \sqrt{\kappa^2 - 1} = -\kappa \pm i + O(\kappa^2). \quad (14.15)$$

Thus,  $\text{Re } \lambda_{\pm} = -\kappa < 0$ , and neglecting terms which are  $O(\kappa^2)$ , we obtain that the functions

$$a_1(\tau) = \frac{\lambda_+ e^{\lambda_- \tau} - \lambda_- e^{\lambda_+ \tau}}{\lambda_+ - \lambda_-} = e^{-\kappa\tau} \cos \tau + O(\kappa^2), \quad a_2(\tau) = \frac{e^{\lambda_+ \tau} - e^{\lambda_- \tau}}{\lambda_+ - \lambda_-} = e^{-\kappa\tau} \sin \tau + O(\kappa^2), \quad (14.16)$$

are solutions of the homogeneous Maxwell equation (14.10) with initial data (1,0) and (0,1) respectively. Finally, the fundamental solution is the real-valued function

$$E(\tau) = \theta(\tau) a_2(\tau) = \theta(\tau) e^{-\kappa\tau} \sin \tau + O(\kappa^2), \quad \tau \in \mathbb{R}. \quad (14.17)$$

## 14.1 The first-order approximation

**The Schrödinger amplitudes.** Formulas (14.6)–(14.8) with  $k = 1$  give the averaged matrices

$$\tilde{\Omega}_n^{(0)}(\tau) = \begin{pmatrix} 0 & -i\frac{\gamma_n}{2} \\ -i\frac{\gamma_n}{2} & 0 \end{pmatrix}. \quad (14.18)$$

Hence, equation (14.4) with  $k = 1$  reads as the system in (12.9) with  $\frac{\gamma_n}{2}$  instead of  $\omega_n(\tau)$ :

$$\begin{cases} \dot{c}_{n,1}^{(1)}(\tau) &= -i\frac{\gamma_n}{2} c_{n,2}^{(1)}(\tau) \\ \dot{c}_{n,2}^{(1)}(\tau) &= -i\frac{\gamma_n}{2} c_{n,1}^{(1)}(\tau) \end{cases}, \quad n \in \bar{N}. \quad (14.19)$$

Its solutions are

$$c_{n,1}^{(1)}(\tau) = d_{n,1}^{(1)} \cos \frac{\gamma_n \tau}{2} + d_{n,2}^{(1)} \sin \frac{\gamma_n \tau}{2}, \quad c_{n,2}^{(1)}(\tau) = -id_{n,1}^{(1)} \sin \frac{\gamma_n \tau}{2} + id_{n,2}^{(1)} \cos \frac{\gamma_n \tau}{2}, \quad (14.20)$$

where

$$d_{n,1}^{(1)} = c_{n,1}(0), \quad d_{n,2}^{(1)} = -ic_{n,2}(0). \quad (14.21)$$

Hence, the matrix in the formula (14.9) with  $n = 1$  reads

$$U_n^{(1)}(\tau) = \begin{pmatrix} \cos \frac{\gamma_n \tau}{2} & -i \sin \frac{\gamma_n \tau}{2} \\ -i \sin \frac{\gamma_n \tau}{2} & \cos \frac{\gamma_n \tau}{2} \end{pmatrix} + O(\gamma_n^2) \quad (14.22)$$

by Lemma A.1.

**The current.** Substituting (14.20) and (14.21) in the second formula of (14.10) with  $k = 1$ , we obtain

$$\begin{aligned} j^{(1)}(\tau) &= \sum_n \alpha_n \text{Im} \left\{ \bar{c}_{n,1}^{(1)}(\tau) c_{n,2}^{(1)}(\tau) e^{-i\tau} \right\} \\ &= \sum_n \alpha_n \text{Im} \left\{ \left[ \bar{d}_{n,1}^{(1)} \cos \frac{\gamma_n \tau}{2} + \bar{d}_{n,2}^{(1)} \sin \frac{\gamma_n \tau}{2} \right] \left[ -id_{n,1}^{(1)} \sin \frac{\gamma_n \tau}{2} + id_{n,2}^{(1)} \cos \frac{\gamma_n \tau}{2} \right] e^{-i\tau} \right\} \\ &= \sum_n \alpha_n \text{Im} \left\{ \left[ i\bar{d}_{n,1}^{(1)} d_{n,2}^{(1)} - |d_{n,1}^{(1)}|^2 \sin \frac{\gamma_n \tau}{2} + |d_{n,2}^{(1)}|^2 \sin \frac{\gamma_n \tau}{2} \right] e^{-i\tau} \right\} + O\left(\sum_n \alpha_n \gamma_n^2\right) \\ &= \sum_n \alpha_n \text{Re} \left\{ \left[ \bar{d}_{n,1}^{(1)} d_{n,2}^{(1)} - |d_{n,1}^{(1)}|^2 \sin \frac{\gamma_n \tau}{2} + |d_{n,2}^{(1)}|^2 \sin \frac{\gamma_n \tau}{2} \right] e^{-i\tau} \right\} + O\left(\sum_n \alpha_n \gamma_n^2\right) \\ &= \sum_n \alpha_n \left[ \text{Im} \left\{ \bar{c}_{n,1}^{(1)}(0) c_{n,2}^{(1)}(0) e^{-i\tau} \right\} + \left[ |c_{n,2}^{(1)}(0)|^2 - |c_{n,1}^{(1)}(0)|^2 \right] \frac{\gamma_n \tau}{2} \cos \tau \right] + O(10^{-17}), \quad \tau \in [0, 2\pi], \quad (14.23) \end{aligned}$$

since  $\sum_n \alpha_n \gamma_n^2 \sim 10^{-17}$  by (12.13).

**Notation.** For each number  $\varepsilon > 0$ , the equality  $r = O(\varepsilon)$  means that  $|r| \leq 10\varepsilon$ .

**The Maxwell amplitude.** Now (14.12) with  $k = 1$  becomes

$$a^{(1)}(\tau) = a_0(\tau) + \int_0^\tau j^{(1)}(\tau') E(\tau - \tau') d\tau'. \quad (14.24)$$

Substituting here the current (14.23), we obtain

$$a^{(1)}(\tau) = a_0(\tau) + \sum_n \alpha_n \left[ \text{Im} \left\{ \bar{c}_{n,1}(0) c_{n,2}(0) I_1(\tau) \right\} + [|c_{n,2}(0)|^2 - |c_{n,1}(0)|^2] \gamma_n I_2(\tau) \right] + O(10^{-17}), \quad (14.25)$$

where we denote

$$I_1(\tau) := \int_0^\tau e^{-i\tau'} E(\tau - \tau') d\tau' = -\frac{i}{2} \sin \tau - \frac{\tau}{2i} e^{-i\tau} + \varkappa \left[ \frac{-\sin \tau + \tau e^{i\tau}}{4} + \frac{\tau^2 e^{-i\tau}}{4i} \right] + O_2,$$

$$I_2(\tau) := \int_0^\tau \frac{\tau'}{2} \cos \tau' E(\tau - \tau') d\tau' = \frac{1}{4} \left[ \frac{\tau^2}{2} \sin \tau + \frac{\tau}{2} \cos \tau - \frac{1}{2} \sin \tau \right] + O_2,$$

where  $O_2 := O(\varkappa^2) = O(10^{-14})$ . The details of calculation can be found in Appendix B.

## 14.2 The second-order approximation

**The Schrödinger amplitudes.** Formulas (14.6) with  $k = 2$  give

$$\Omega_n^{(1)}(\tau) = \begin{pmatrix} 0 & \omega_n^{(1)}(\tau) \\ \bar{\omega}_n^{(1)}(\tau) & 0 \end{pmatrix}, \quad \omega_n^{(1)}(\tau) = [\beta_n \dot{a}^{(1)}(\tau) + \gamma_n \cos \tau] e^{-i\tau}. \quad (14.26)$$

Let us calculate the averaged matrices  $\tilde{\Omega}_n^{(1)}$ . First, (14.25) and (14.12) imply that

$$\mathbf{v} := \frac{1}{2\pi} \int_0^{2\pi} \dot{a}^{(1)}(\tau) e^{-i\tau} d\tau = \mathbf{v}_1 + \mathbf{v}_2, \quad (14.27)$$

where

$$\mathbf{v}_1 := \frac{1}{2\pi} \int_0^{2\pi} \dot{a}_0(\tau) e^{-i\tau} d\tau = \frac{1}{2\pi} \int_0^{2\pi} \{a(0) \dot{E}(\tau) + [\dot{a}(0) + 2\varkappa a(0)] \dot{E}(\tau)\} e^{-i\tau} d\tau,$$

$$\mathbf{v}_2 := \frac{1}{2\pi} \int_0^{2\pi} \sum_n \alpha_n \left[ \text{Im} \left\{ \bar{c}_{n,1}(0) c_{n,2}(0) I_1'(\tau) e^{-i\tau} \right\} + [|c_{n,2}(0)|^2 - |c_{n,1}(0)|^2] \gamma_n I_2'(\tau) e^{-i\tau} \right] d\tau$$

First, let us calculate  $\mathbf{v}_1$ . Neglecting errors  $O(\varkappa^2) = O(10^{-14})$ , we obtain:

$$\begin{aligned} \int_0^{2\pi} \dot{E}(\tau) e^{-i\tau} d\tau &\approx i \int_0^{2\pi} E(\tau) e^{-i\tau} d\tau \approx i \int_0^{2\pi} e^{-\varkappa\tau} \sin \tau e^{-i\tau} d\tau = \frac{1}{2} \int_0^{2\pi} e^{-\varkappa\tau} (e^{i\tau} - e^{-i\tau}) e^{-i\tau} d\tau \\ &= -\left[ \frac{1}{2\varkappa} + \frac{1}{2\varkappa + 4i} \right] (e^{-2\varkappa\pi} - 1) = -\frac{2\varkappa + 2i}{\varkappa + 2i} \frac{-\varkappa\pi + (\varkappa\pi)^2 + O(\varkappa^3)}{\varkappa} \\ &\approx -\left[ 1 + \frac{\varkappa}{\varkappa + 2i} \right] (-\pi + \varkappa\pi^2) \approx -\left[ 1 + \frac{\varkappa}{2i} \right] (-\pi + \varkappa\pi^2) \approx \pi + \varkappa \left( \frac{\pi}{2i} - \pi^2 \right). \end{aligned} \quad (14.28)$$

Similarly, using (14.13), (14.17), and (14.28), we obtain

$$\begin{aligned} \int_0^{2\pi} \dot{E}(\tau) e^{-i\tau} d\tau &= \int_0^{2\pi} [-2\varkappa \dot{E}(\tau) - E(\tau)] e^{-i\tau} d\tau \approx -(i2\varkappa + 1) \int_0^{2\pi} E(\tau) e^{-i\tau} d\tau \\ &\approx i(i2\varkappa + 1) \left[ \pi + \varkappa \left( \frac{\pi}{2i} - \pi^2 \right) \right] = (-2\varkappa + i) \left[ \pi + \varkappa \left( \frac{\pi}{2i} - \pi^2 \right) \right] \\ &\approx i\pi + \varkappa \left[ -2\pi + \frac{\pi}{2} - i\pi^2 \right] = i\pi - \varkappa \left[ \frac{3\pi}{2} + i\pi^2 \right]. \end{aligned} \quad (14.29)$$

As a result,

$$\left. \begin{aligned} v_1 &\approx a(0) \left[ \frac{i}{2} - \varkappa \left[ \frac{3}{4} + i \frac{\pi}{2} \right] \right] + [\dot{a}(0) + 2\varkappa a(0)] \left[ \frac{1}{2} + \varkappa \left( \frac{1}{4i} - \frac{\pi}{2} \right) \right] \\ &\approx a(0) \left[ \frac{i}{2} + \varkappa \left( \frac{1}{4} - i \frac{\pi}{2} \right) \right] + \dot{a}(0) \left[ \frac{1}{2} + \varkappa \left( \frac{1}{4i} - \frac{\pi}{2} \right) \right] = v_{11} + i v_{12}, \\ v_{11} &= \frac{\varkappa}{4} a(0) + \frac{1}{2} (1 - \varkappa \pi) \dot{a}(0), \quad v_{12} = \frac{1}{2} (1 - \varkappa \pi) a(0) - \frac{\varkappa}{4} \dot{a}(0) \end{aligned} \right\}. \quad (14.30)$$

It remains to note that

$$v_2 := \sum_n \alpha_n \left[ \operatorname{Im} \left\{ \bar{c}_{n,1}(0) c_{n,2}(0) J_1 \right\} + [|c_{n,2}(0)|^2 - |c_{n,1}(0)|^2] \gamma_n J_2 \right], \quad (14.31)$$

where  $J_1$  and  $J_2$  are the integrals

$$J_1 := \frac{1}{2\pi} \int_0^{2\pi} I'_1(\tau) e^{-i\tau} d\tau \approx 0, \quad J_2 := \frac{1}{2\pi} \int_0^{2\pi} I'_2(\tau) e^{-i\tau} d\tau \approx \frac{\pi^2}{12}. \quad (14.32)$$

We calculate these integrals in Appendix B. As a result, the averaged matrix reads

$$\tilde{\Omega}_n^{(1)}(\mathbf{v}) = \begin{pmatrix} 0 & \tilde{\omega}_n^{(1)} \\ \bar{\tilde{\omega}}_n^{(1)} & 0 \end{pmatrix}, \quad \tilde{\omega}_n^{(1)} = \beta_n \mathbf{v} + \frac{\gamma_n}{2}. \quad (14.33)$$

Now equation (14.7) with  $k = 2$  reads as the system

$$\left\{ \begin{aligned} \dot{c}_{n,1}^{(2)}(\tau) &= -i \tilde{\omega}_n^{(1)} c_{n,2}^{(2)}(\tau) \\ \dot{c}_{n,2}^{(2)}(\tau) &= -i \bar{\tilde{\omega}}_n^{(1)} c_{n,1}^{(2)}(\tau) \end{aligned} \right\}, \quad n \in \bar{N}. \quad (14.34)$$

Its solution reads similarly to (14.20):

$$c_{n,1}^{(2)}(\tau) = d_{n,1}^{(2)} \cos |\tilde{\omega}_n^{(1)}| \tau + d_{n,2}^{(2)} \sin |\tilde{\omega}_n^{(1)}| \tau, \quad c_{n,2}^{(2)}(\tau) = -i d_{n,1}^{(2)} \frac{|\tilde{\omega}_n^{(1)}|}{\tilde{\omega}_n^{(1)}} \sin |\tilde{\omega}_n^{(1)}| \tau + i d_{n,2}^{(2)} \frac{|\tilde{\omega}_n^{(1)}|}{\bar{\tilde{\omega}}_n^{(1)}} \cos |\tilde{\omega}_n^{(1)}| \tau, \quad (14.35)$$

where

$$d_{n,1}^{(2)} = c_{n,1}(0), \quad d_{n,2}^{(2)} = -i \frac{\tilde{\omega}_n^{(1)}}{|\tilde{\omega}_n^{(1)}|} c_{n,2}(0). \quad (14.36)$$

Thus, (14.35) can be rewritten as

$$\left\{ \begin{aligned} c_{n,1}^{(2)}(\tau) &= c_{n,1}(0) \cos |\tilde{\omega}_n^{(1)}| \tau - i s_n c_{n,2}(0) \sin |\tilde{\omega}_n^{(1)}| \tau \\ c_{n,2}^{(2)}(\tau) &= -i \bar{s}_n c_{n,1}(0) \sin |\tilde{\omega}_n^{(1)}| \tau + c_{n,2}(0) \cos |\tilde{\omega}_n^{(1)}| \tau, \end{aligned} \right\}, \quad (14.37)$$

where  $s_n = \frac{\tilde{\omega}_n^{(1)}}{|\tilde{\omega}_n^{(1)}|}$ . Hence, the matrix in the formula (14.9) with  $k = 2$  reads

$$U_n^{(2)}(\tau) = \begin{pmatrix} \cos |\tilde{\omega}_n^{(1)}| \tau & -i s_n \sin |\tilde{\omega}_n^{(1)}| \tau \\ -i \bar{s}_n \sin |\tilde{\omega}_n^{(1)}| \tau & \cos |\tilde{\omega}_n^{(1)}| \tau \end{pmatrix}. \quad (14.38)$$

**The current.** Expanding (14.37) into the Taylor series, we obtain

$$\left\{ \begin{aligned} c_{n,1}^{(2)}(\tau) &= c_{n,1}(0) - i c_{n,2}(0) \tilde{\omega}_n^{(1)} \tau + O(|\tilde{\omega}_n^{(1)}|^2) \\ c_{n,2}^{(2)}(\tau) &= -i c_{n,1}(0) \bar{\tilde{\omega}}_n^{(1)} \tau + c_{n,2}(0) + O(|\tilde{\omega}_n^{(1)}|^2) \end{aligned} \right\}, \quad (14.39)$$



Here the error  $O(|\tilde{\omega}_n^{(1)}|^2) \sim 10^{-14}$  uniformly in  $\tau \in [0, 2\pi]$ . Indeed,  $\tilde{\omega}_n^{(1)} = \beta_n \mathbf{v} + \frac{\gamma_n}{2}$ , where  $\mathbf{v}$  is expressed by (14.27) and (14.30)–(14.32):

$$\left\{ \begin{array}{l} \mathbf{v} = v_{11} + i v_{12} + v_2 \\ v_{11} \approx \frac{b(0)}{2} + \frac{\varkappa}{4}[a(0) - 2\pi b(0)], \quad v_{12} \approx \frac{a(0)}{2} - \frac{\varkappa}{4}[2\pi a(0) + b(0)] \\ v_2 \approx J_2 \sum_n \alpha_n \gamma_n [|c_{n,2}(0)|^2 - |c_{n,1}(0)|^2] \end{array} \right., \quad (14.40)$$

Hence at the points (13.2), we have  $v_{11} = v_{12} = 0$ , so

$$\beta_n \mathbf{v} = \beta_n v_2 = -\beta_n J_2 \sum_n \alpha_n \gamma_n \sim 10^{-12} \quad (14.41)$$

by (12.13) and (14.32). Therefore,

$$\tilde{\omega}_n^{(1)} = \beta_n \mathbf{v} + \frac{\gamma_n}{2} \approx \frac{\gamma_n}{2} \sim 10^{-7}. \quad (14.42)$$

Finally, substituting (14.39) into the second formula of (14.10) with  $k = 2$ , we obtain similarly to (14.23),

$$\begin{aligned} j^{(2)}(\tau) &= \sum_n \alpha_n \operatorname{Im} \left\{ \bar{c}_{n,1}^{(2)}(\tau) c_{n,2}^{(2)}(\tau) e^{-i\tau} \right\} \\ &= \sum_n \alpha_n \operatorname{Im} \left\{ \bar{c}_{n,1}(0) c_{n,2}(0) e^{-i\tau} + i \bar{\omega}_n^{(1)} [|c_{n,2}(0)|^2 - |c_{n,1}(0)|^2] \tau e^{-i\tau} \right\} + O(|\tilde{\omega}_n^{(1)}|^2) \end{aligned} \quad (14.43)$$

**The Maxwell amplitude.** The last formula of (14.12) with  $k = 2$  gives

$$\left\{ \begin{array}{l} a^{(2)}(\tau) = a_0(\tau) + \int_0^\tau j^{(2)}(\tau') E(2\pi - \tau') d\tau' \\ b^{(2)}(\tau) = \dot{a}_0(\tau) + \int_0^\tau j^{(2)}(\tau') \dot{E}(2\pi - \tau') d\tau' \end{array} \right. \quad (14.44)$$

## 15 The Poincaré map

Let us denote

$$a^0 := a(0), \quad b^0 := \dot{a}(0), \quad c_n^0 := c_n(0); \quad a := a^{(2)}(2\pi), \quad b := \dot{a}^{(2)}(2\pi), \quad c_n := c_n^{(2)}(2\pi). \quad (15.1)$$

Then (14.44) gives

$$a = a_0(2\pi) + \int_0^{2\pi} j^{(2)}(\tau') E(2\pi - \tau') d\tau', \quad b = \dot{a}_0(2\pi) + \int_0^{2\pi} j^{(2)}(\tau') \dot{E}(2\pi - \tau') d\tau'. \quad (15.2)$$

Let us calculate these expressions. First, (14.12) and (14.13), (14.17) imply that

$$\begin{aligned} a_0(2\pi) &= a^0 \dot{E}(2\pi), \quad \dot{a}_0(2\pi) = a^0 \ddot{E}(2\pi) + [b^0 + \sigma_1 a^0] \dot{E}(2\pi) \\ &= -a^0 E(2\pi) + b^0 \dot{E}(2\pi) \approx b^0 \dot{E}(2\pi) + O_2, \end{aligned} \quad (15.3)$$

where  $O_2 := O(\varkappa^2) \sim 10^{-14}$ . Second, substituting  $\tilde{\omega}_n^{(1)} = \beta_n \mathbf{v} + \frac{\gamma_n}{2}$  into (14.43), we obtain

$$\begin{aligned} j^{(2)}(\tau) &\approx \sum_n \alpha_n \operatorname{Im} \left\{ \bar{c}_{n,1}^0 c_{n,2}^0 e^{-i\tau} \right\} \\ &+ \sum_n \alpha_n \beta_n [|c_{n,2}^0|^2 - |c_{n,1}^0|^2] \operatorname{Re} \left\{ \bar{\mathbf{v}} \tau e^{-i\tau} \right\} + \frac{1}{2} \sum_n \alpha_n \gamma_n [|c_{n,2}^0|^2 - |c_{n,1}^0|^2] \tau \cos \tau. \end{aligned} \quad (15.4)$$

Introducing this current into (15.2), we obtain the formulas for  $a$  and  $b$  expressed via the following integrals which are calculated in Appendix B:

$$\left\{ \begin{array}{l} A_1 := \int_0^{2\pi} e^{-i\tau'} E(2\pi - \tau') d\tau', A_2 := \int_0^{2\pi} \tau' e^{-i\tau'} E(2\pi - \tau') d\tau', A_3 := \int_0^{2\pi} \tau' \cos \tau' E(2\pi - \tau') d\tau' = \text{Re} A_2 \\ B_1 = \int_0^{2\pi} e^{-i\tau'} \dot{E}(2\pi - \tau') d\tau', B_2 := \int_0^{2\pi} \tau' e^{-i\tau'} \dot{E}(2\pi - \tau') d\tau', B_3 := \int_0^{2\pi} \tau' \cos \tau' \dot{E}(2\pi - \tau') d\tau' = -\text{Im} B_2 \end{array} \right. \quad (15.5)$$

With this notation, (15.2) becomes

$$\left\{ \begin{array}{l} a \approx a_0(2\pi) + \sum_n \alpha_n \text{Im} \left\{ \bar{c}_{n,1}^0 c_{n,2}^0 A_1 \right\} \\ \quad + \sum_n \alpha_n \beta_n [ |c_{n,2}^0|^2 - |c_{n,1}^0|^2 ] \text{Re} \left\{ \bar{v} A_2 \right\} + \frac{1}{2} \sum_n \alpha_n \gamma_n [ |c_{n,2}^0|^2 - |c_{n,1}^0|^2 ] A_3. \\ b \approx \dot{a}_0(2\pi) + \sum_n \alpha_n \text{Im} \left\{ \bar{c}_{n,1}^0 c_{n,2}^0 B_1 \right\} \\ \quad + \sum_n \alpha_n \beta_n [ |c_{n,2}^0|^2 - |c_{n,1}^0|^2 ] \text{Re} \left\{ \bar{v} B_2 \right\} + \frac{1}{2} \sum_n \alpha_n \gamma_n [ |c_{n,2}^0|^2 - |c_{n,1}^0|^2 ] B_3. \end{array} \right. \quad (15.6)$$

**Remark 15.1.** i) The Poincaré map  $P$  is given by formulas (15.6) and (14.39).

ii) The second-order approximations (15.6) are obtained solving the equation (14.10) with  $k = 2$ , initial data (14.1), and the current (15.4). At the ‘‘ground states’’ (13.2), we have  $|c_{n,1}^0|^2 = 1$  and  $|c_{n,2}^0|^2 = 0$ , so the formula (15.4) takes the form

$$j^{(2)}(\tau) = - \sum_n \alpha_n \beta_n \tau \text{Im} \left\{ \bar{v} e^{-i\tau} \right\} - \frac{1}{2} \sum_n \alpha_n \gamma_n \tau \cos \tau. \quad (15.7)$$

## 16 Synchronisation of molecular currents

The key peculiarity of the perturbative calculations is the appearance of the summation

$$\mathbf{S} = \sum_n \alpha_n \beta_n \quad (16.1)$$

in the second-order approximation of the total molecular current (15.7). This appearance is due to the Taylor approximation (14.39). Also note that the same summation reappears in (20.15). It is of crucial importance that all terms of the sum  $\mathbf{S}$  are *nonnegative* since

$$\alpha_n \beta_n = \frac{2c}{\Omega_p} \mathbf{P}_n \mathbf{X}(x_n) \frac{\mathbf{P}_n \mathbf{X}(x_n)}{\hbar c} \geq 0, \quad (16.2)$$

and the sum is a non-negligible positive number, see (16.6) and (16.8) below. This means the synchronisation of all molecular currents in the total current (13.4) *independently of the phase of the Maxwell field* at the location of the molecules. This *synchronisation of molecular currents* is provisionally in line with the *superradiance* and with the role of the *stimulated emission* in the laser action. Note that the conjecture on appropriate kind of synchronisation in the laser action has been put forward repeatedly, see, e.g., [30].

We will consider two cases of amorphous and crystallin medium separately. In (12.8), all the parameters  $\mathbf{P}_n$ ,  $\mathbf{X}(x_n)$ ,  $\mathbf{a}_p(x_n)$  with  $n \in \bar{N}$  can be considered as *independent random values* on the event space  $\{1, \dots, N\}$ , and

$$\mathbb{E} \mathbf{X}(x_n) = \mathbb{E} \mathbf{a}_p(x_n) = 0. \quad (16.3)$$

**I. Amorphous medium.** By our assumption,  $|\mathbf{P}_n| = |\mathbf{P}|$  does not depend on  $n$ , so

$$(\mathbf{P}_n \cdot \mathbf{e}_1)^2 + (\mathbf{P}_n \cdot \mathbf{e}_2)^2 + (\mathbf{P}_n \cdot \mathbf{e}_3)^2 = |\mathbf{P}|^2. \quad (16.4)$$

Moreover, the dipole momenta  $\mathbf{P}_n$  are distributed uniformly over the angles, and hence,

$$\mathbb{E}(\mathbf{P}_n \cdot \mathbf{e}_1)^2 = \frac{1}{3} |\mathbf{P}|^2. \quad (16.5)$$

By Remark 12.1, we can apply the *law of large numbers* for weakly dependent random values [21] (recall that  $N \sim 10^{20}$ ), and obtain that  $\mathbf{S} \approx N\mathbb{E}\alpha_n\beta_n$ . Moreover,  $\mathbf{P}_n$  and  $\mathbf{X}(x_n)$  are independent. Hence, using (5.1), we obtain that

$$\begin{aligned} \mathbf{S} &\approx N\mathbb{E}\alpha_n\beta_n = N\frac{2}{\Omega_p\hbar}\mathbb{E}(\mathbf{P}_n\mathbf{X}(x_n))^2 = N\frac{2}{\Omega_p\hbar}\frac{\mathbb{E}|\mathbf{P}_n|^2}{3}\mathbb{E}\mathbf{X}^2(x_n) \\ &= \frac{2}{\Omega_p\hbar}\frac{|\mathbf{P}|^2}{3}\sum_n\mathbf{X}^2(x_n) \approx \frac{2|\mathbf{P}|^2}{3\Omega_p\hbar}\frac{N}{|V_a|}\int_{V_a}\mathbf{X}^2(x)dx = \frac{2|\mathbf{P}|^2}{3\Omega_p\hbar}d\frac{|V_a|}{|V|}, \quad d := \frac{N}{|V_a|} \end{aligned} \quad (16.6)$$

by the normalization of  $\mathbf{X}$ . Here the value  $|V_a|/|V|$  for the integral follows from Shnirelman's *Quantum Ergodic theorem* [34, 39, 40]. Note that this value obviously holds for the eigenmodes (7.6) with sufficiently large eigenvalues. For the case of ruby laser (12.13) we obtain  $d \approx 3 \times 10^{19}$  and

$$\mathbf{S} \sim 10^{-5}. \quad (16.7)$$

which is much larger than the terms  $|\alpha_n\beta_n| \sim 10^{-25}$  in the summation (16.1). Moreover,  $\mathbf{S}$  is a Gaussian random value since  $\alpha_n\beta_n$  are independent and identically distributed.

**II. Crystallin medium.** In this case the vectors  $\mathbf{P}_n = \mathbf{P}$  do not depend on  $n$ . We assume additionally that the mode  $\mathbf{X}(x)$  is linearly polarised. Now (16.6) changes to

$$\mathbf{S} \approx \frac{2}{\Omega_p\hbar}|\mathbf{P}|^2\cos^2\varphi\sum_n\mathbf{X}^2(x_n) \approx \frac{2|\mathbf{P}|^2\cos^2\varphi}{\Omega_p\hbar}d\frac{|V_a|}{|V|} \approx 32\frac{|V_a|}{|V|}10^{-5}\cos^2\varphi \sim 10^{-5}, \quad (16.8)$$

where  $\varphi$  is the angle between  $\mathbf{P}$  and  $\mathbf{X}(x)$ .

## 17 Factordynamics and the Hopf fibration

Let us denote by  $G = [U(1)]^N$  the symmetry gauge group of the system (13.3). The group acts on the phase space  $\mathbb{X} = \mathbb{R}^2 \oplus \mathbb{C}^{2N}$  by the formula

$$(e^{i\theta_1}, \dots, e^{i\theta_N})(a, b, (c_{11}, c_{12}), \dots, (c_{N,1}, c_{N,2})) = (a, b, (e^{i\theta_1}c_{11}, e^{i\theta_1}c_{12}), \dots, (e^{i\theta_N}c_{N,1}, e^{i\theta_N}c_{N,2})). \quad (17.1)$$

This action commutes with the dynamics (13.3), hence the latter induces the corresponding dynamics on the factorspace  $\mathbb{X}_* = \mathbb{X}/G$ . This factorspace is the base of the fibration  $\mathbb{X} \rightarrow \mathbb{X}/G$ .

The Poincaré map  $P$  commutes with the action (17.1). Moreover, this action does not change the Maxwell field. Accordingly, the instability and the parametric resonance (1.1) must be checked for the induced factordynamics only. More accurate justification is given in the next section: we reduce the problem to calculation of the differential  $DP_*(X_*^0)$ , where  $P_*$  denotes the action of  $P$  on the factorspace  $\mathbb{X}/G$ , and the point  $X_*^0$  corresponds to  $X^0$  in this factorization (see (13.1)). We show that the eigenvalues of  $DP(X^0)$  and  $DP_*(X_*^0)$  coincide up to  $\mu = 1$ , corresponding to the action (17.1).

To calculate  $DP_*(X_*^0)$ , we need suitable local coordinates on the factorspace. The charge conservation (12.11) means that

$$(c_{n,1}(\tau), c_{n,2}(\tau)) \in S^3, \quad (17.2)$$

where  $S^3$  is the unit sphere in  $\mathbb{C}^2$ . The sphere is the Hopf fibration with fibers

$$F = (c_1 e^{i\theta}, c_2 e^{i\theta}), \quad \theta \in \mathbb{R} \quad (17.3)$$

and the base diffeomorphic to the unit sphere  $S^2$ . Denote by  $(c_1, c_2)_* \in S^2$  the Hopf projection of a point  $(c_1, c_2) \in S^3$ . We choose the following invariant coordinate on the base:

$$z := \bar{c}_1 c_2. \quad (17.4)$$

Using (12.11), it is easy to check that

$$|c_1| + |c_2| = \sqrt{1 + 2|z|}, \quad |c_1| - |c_2| = \pm\sqrt{1 - 2|z|}. \quad (17.5)$$

Hence,

$$2|c_1| = \sqrt{1+2|z|} + \sqrt{1-2|z|}, \quad 2|c_2| = \sqrt{1+2|z|} - \sqrt{1-2|z|} \quad (17.6)$$

and

$$4|c_1|^2 = 2 + 2\sqrt{1-4|z|^2}, \quad 4|c_2|^2 = 2 - 2\sqrt{1-4|z|^2}. \quad (17.7)$$

In particular,  $|c_1|^2$  and  $|c_2|^2$  are smooth functions on the disk  $D := |z| < \frac{1}{2}$ , and

$$\nabla_z |c_l|^2|_{z=0} = 0, \quad (17.8)$$

where  $\nabla_z$  denotes the gradient with respect to  $\operatorname{Re} z$  and  $\operatorname{Im} z$ .

The coordinate  $z$  maps  $S^3$  onto the closed disk  $\bar{D}$ , and it is a local coordinate on the region of  $S^3$  outside the set  $|c_1|^2 = |c_2|^2 = 1/2$ . However, the inverse map  $D \rightarrow S^2$  is two-valued. Indeed, denote by  $(c_1, c_2)_* \in S^2$  the Hopf projection of a point  $(c_1, c_2) \in S^3$ . One branch of the inverse map is a diffeomorphism of  $D$  onto the neighborhood  $S^2_+ \subset S^2$  of the point  $(1, 0)_*$  (where  $z = 0$ ), and another branch is a diffeomorphism of  $D$  onto the neighborhood  $S^2_- \subset S^2$  of the point  $(0, 1)_*$  (where also  $z = 0$ ):

$$\left\{ \begin{array}{l} |c_1| > |c_2| \quad \text{for } (c_1, c_2)_* \in S^2_+ \\ |c_1| < |c_2| \quad \text{for } (c_1, c_2)_* \in S^2_- \end{array} \right\}. \quad (17.9)$$

The base  $S^2$  is the gluing of  $S^2_\pm$  along their boundaries, where  $|z| = \frac{1}{2}$  and  $|c_1|^2 = |c_2|^2 = \frac{1}{2}$ .

## 18 Reduction of spectral problem by the symmetry group

We choose coordinates (17.4) on the bases of the Hopf fibration for each active molecule, and denote

$$z_n^0 := \bar{c}_{n,1}^0 c_{n,2}^0 = \bar{c}_{n,1}(0) c_{n,2}(0), \quad z_n := \bar{c}_{n,1} c_{n,2} = \bar{c}_{n,1}(2\pi) c_{n,2}(2\pi). \quad (18.1)$$

Let us write  $z_n = z_{n,1} + iz_{n,2}$ ,  $z_n^0 = z_{n,1}^0 + iz_{n,2}^0$ , where  $z_{n,1}, z_{n,2}, z_{n,1}^0, z_{n,2}^0 \in \mathbb{R}$ , and

$$\mathbf{z}_n = \begin{pmatrix} z_{n,1} \\ z_{n,2} \end{pmatrix}, \quad \mathbf{z}_n^0 = (z_{n,1}^0, z_{n,2}^0), \quad \frac{\partial}{\partial \mathbf{z}_n^0} := \left( \frac{\partial}{\partial z_{n,1}^0}, \frac{\partial}{\partial z_{n,2}^0} \right), \quad n \in \bar{N}. \quad (18.2)$$

Denote

$$\mathbf{Z} = \begin{pmatrix} \mathbf{z}_1 \\ \vdots \\ \mathbf{z}_N \end{pmatrix}, \quad \mathbf{Z}^0 = (\mathbf{z}_1^0, \dots, \mathbf{z}_N^0). \quad (18.3)$$

Then the differential  $DP_*(X_*^0)$  of the Poincaré map  $P_*$  on the factorspace  $\mathbb{X}/G$  is represented by the matrix

$$DP_*(X_*^0) = \begin{pmatrix} \frac{\partial a}{\partial a^0} & \frac{\partial a}{\partial b^0} & \frac{\partial a}{\partial \mathbf{Z}^0} \\ \frac{\partial b}{\partial a^0} & \frac{\partial b}{\partial b^0} & \frac{\partial b}{\partial \mathbf{Z}^0} \\ \frac{\partial \mathbf{Z}}{\partial a^0} & \frac{\partial \mathbf{Z}}{\partial b^0} & \frac{\partial \mathbf{Z}}{\partial \mathbf{Z}^0} \end{pmatrix}. \quad (18.4)$$

The following lemma justifies the reduction to the eigenvalue problem for the factordynamics.

**Lemma 18.1.** *The eigenvalues of  $DP(X^0)$  and  $DP_*(X_*^0)$  coincide up to  $\mu = 1$ .*

*Proof.* The variables  $a^0, b^0, \mathbf{Z}^0$  are local coordinates on the factorspace  $\mathbb{X}_* = \mathbb{X}/G$ . This factorspace is the base of the fiberspace  $cX$  with the fibers, isomorphic to  $S^N$ , where  $S$  is the unit circle  $|\theta| = 1$ . Hence, in a small neighborhood of any fixed point  $X^0 \in \mathbb{X}$ , these coordinates can be completed by the phases  $\theta_n$  of the action (17.1). Now  $X^0 = (a^0, b^0, \mathbf{Z}^0, \Theta^0)$  with an arbitrary fixed  $\Theta^0 \in S^N$ , and the action of the symmetry group (17.1) becomes the shifts in  $\Theta$ :

$$(e^{i\theta_1}, \dots, e^{i\theta_N})(\tilde{a}^0, \tilde{b}^0, \tilde{\mathbf{Z}}^0, \tilde{\Theta}^0) = (\tilde{a}^0, \tilde{b}^0, \tilde{\mathbf{Z}}^0, \tilde{\Theta}^0 + \Theta), \quad \Theta = (\theta_1, \dots, \theta_N) \quad (18.5)$$

for  $(\tilde{a}^0, \tilde{b}^0, \tilde{\mathbf{Z}}^0, \tilde{\Theta}^0)$ , close to  $X^0$ , and small  $\Theta$ , since the Maxwell field and the variables  $z_n$  are invariant with respect to the action (17.1). Moreover,  $PX^0$  is close to  $X^0$ , and the Poincaré map  $P$  commutes with the shifts (18.5) with small  $\Theta$ . Hence, the differential  $DP(X^0)$  is represented by the matrix

$$DP(X^0) = \begin{pmatrix} \frac{\partial a}{\partial a^0} & \frac{\partial a}{\partial b^0} & \frac{\partial a}{\partial \mathbf{Z}^0} & \frac{\partial a}{\partial \Theta^0} \\ \frac{\partial b}{\partial a^0} & \frac{\partial b}{\partial b^0} & \frac{\partial b}{\partial \mathbf{Z}^0} & \frac{\partial b}{\partial \Theta^0} \\ \frac{\partial \mathbf{Z}}{\partial a^0} & \frac{\partial \mathbf{Z}}{\partial b^0} & \frac{\partial \mathbf{Z}}{\partial \mathbf{Z}^0} & \frac{\partial \mathbf{Z}}{\partial \Theta^0} \\ \frac{\partial \Theta}{\partial a^0} & \frac{\partial \Theta}{\partial b^0} & \frac{\partial \Theta}{\partial \mathbf{Z}^0} & \frac{\partial \Theta}{\partial \Theta^0} \end{pmatrix} = \begin{pmatrix} \frac{\partial a}{\partial a^0} & \frac{\partial a}{\partial b^0} & \frac{\partial a}{\partial \mathbf{Z}^0} & 0 \\ \frac{\partial b}{\partial a^0} & \frac{\partial b}{\partial b^0} & \frac{\partial b}{\partial \mathbf{Z}^0} & 0 \\ \frac{\partial \mathbf{Z}}{\partial a^0} & \frac{\partial \mathbf{Z}}{\partial b^0} & \frac{\partial \mathbf{Z}}{\partial \mathbf{Z}^0} & 0 \\ \frac{\partial \Theta}{\partial a^0} & \frac{\partial \Theta}{\partial b^0} & \frac{\partial \Theta}{\partial \mathbf{Z}^0} & I_N \end{pmatrix} = \begin{pmatrix} DP_*(X_*^0) & 0 \\ \dots & I_N \end{pmatrix}, \quad (18.6)$$

where  $I_N$  is the identity  $N \times N$ -matrix. Therefore, the eigenvalues of  $DP(X^0)$  and  $DP_*(X_*^0)$  coincide up to  $\mu = 1$ , which corresponds to the action (18.5).  $\square$

## 19 Differential of the Poincaré map on the factorspace

It remains to calculate the differential (18.4). Using (14.39) with  $\tau = 2\pi$ , we obtain that

$$\begin{aligned} z_n &= (\bar{c}_{n,1}^0 + i\bar{c}_{n,2}^0 2\pi\bar{\omega}_n^{(1)})(-ic_{n,1}^0 2\pi\bar{\omega}_n^{(1)} + c_{n,2}^0) + O(|\bar{\omega}_n^{(1)}|^2) \\ &= z_n^0 + 2\pi i\bar{\omega}_n^{(1)}[|c_{n,2}^0|^2 - |c_{n,1}^0|^2] + O(|\bar{\omega}_n^{(1)}|^2) \\ &= z_n^0 + 2\pi i(\beta_n \bar{\nabla} + \frac{\gamma_n}{2})[|c_{n,2}^0|^2 - |c_{n,1}^0|^2] + O(|\bar{\omega}_n^{(1)}|^2). \end{aligned} \quad (19.1)$$

We must calculate the matrix (18.4) at the ground states (13.2) which correspond to the single point

$$X_*^0 = (0, 0, 0). \quad (19.2)$$

The formulas (15.6) for the Poincaré map become

$$\left\{ \begin{array}{l} a \approx a_0(2\pi) + \sum_n \alpha_n \text{Im} \left\{ z_n^0 A_1 \right\} \\ \quad + \sum_n \alpha_n \beta_n [ |c_{n,2}^0|^2 - |c_{n,1}^0|^2 ] \text{Re} \left\{ \bar{\nabla} A_2 \right\} + \frac{1}{2} \sum_n \alpha_n \gamma_n [ |c_{n,2}^0|^2 - |c_{n,1}^0|^2 ] A_3 \\ b \approx \dot{a}_0(2\pi) + \sum_n \alpha_n \text{Im} \left\{ z_n^0 B_1 \right\} \\ \quad + \sum_n \alpha_n \beta_n [ |c_{n,2}^0|^2 - |c_{n,1}^0|^2 ] \text{Re} \left\{ \bar{\nabla} B_2 \right\} + \frac{1}{2} \sum_n \alpha_n \gamma_n [ |c_{n,2}^0|^2 - |c_{n,1}^0|^2 ] B_3 \end{array} \right. \quad (19.3)$$

Here  $a_0(2\pi)$  and  $\dot{a}_0(2\pi)$  are given by (14.12), so using (14.17) and neglecting errors  $O(\varkappa^2) = O(10^{-14})$ , we obtain

$$\left\{ \begin{array}{l} a_0(2\pi) = a^0 \dot{E}(2\pi) + [b^0 + 2\varkappa a^0] E(2\pi) \approx a^0(1 - 2\pi\varkappa) \\ \dot{a}_0(2\pi) = a^0 \ddot{E}(2\pi) + [b^0 + 2\varkappa a^0] \dot{E}(2\pi) \approx -2\varkappa a^0 + [b^0 + 2\varkappa a^0](1 - 2\pi\varkappa) \approx b^0(1 - 2\pi\varkappa) \end{array} \right. \quad (19.4)$$

At the point (19.2), we have by (17.8),

$$\frac{\partial}{\partial \mathbf{Z}_l^0} \Big|_{z_l^0=0} |c_{n,1}^0|^2 = \frac{\partial}{\partial \mathbf{Z}_l^0} \Big|_{z_l^0=0} |c_{n,2}^0|^2 = (0, 0). \quad (19.5)$$

Differentiating the formulas (14.40) at the point (19.2), we obtain that neglecting errors  $O(\varkappa^2) = O(10^{-14})$ , we have

$$\frac{\partial \bar{\nabla}}{\partial a^0} = -\frac{i}{2} + \frac{\varkappa}{4}(1 + 2\pi i), \quad \frac{\partial \bar{\nabla}}{\partial b^0} = \frac{1}{2} - \frac{\varkappa}{4}(2\pi - i) = i \frac{\partial \bar{\nabla}}{\partial a^0}. \quad (19.6)$$

Moreover, formulas (14.40) and (19.5) imply that at the point (19.2), we have

$$\frac{\partial \bar{v}}{\partial \mathbf{z}'_n} = (0, 0). \quad (19.7)$$

Now we can calculate the matrix (18.4) at the point (19.2). Differentiating the first formula of (19.3), and taking into account (19.4)–(19.7), we obtain that the first row of (18.4) is

$$\frac{\partial a}{\partial a^0} \approx 1 - 2\pi\kappa - \mathbf{S} \left[ \frac{A_{22}}{2} + \frac{\kappa}{4}(A_{21} - 2\pi A_{22}) \right], \quad \frac{\partial a}{\partial b^0} \approx -\mathbf{S} \left[ \frac{A_{21}}{2} - \frac{\kappa}{4}(2\pi A_{21} + A_{22}) \right], \quad \frac{\partial a}{\partial \mathbf{z}'_n} \approx \alpha_{n'}(A_{12}, A_{11}), \quad (19.8)$$

where  $\mathbf{S}$  is the summation (16.1). Similarly, using (B.7), we obtain the second row

$$\frac{\partial b}{\partial a^0} \approx -\mathbf{S} \left[ \frac{B_{22}}{2} + \frac{\kappa}{4}(B_{21} - 2\pi B_{22}) \right], \quad \frac{\partial b}{\partial b^0} \approx 1 - 2\pi\kappa - \mathbf{S} \left[ \frac{B_{21}}{2} - \frac{\kappa}{4}(2\pi B_{21} + B_{22}) \right], \quad \frac{\partial b}{\partial \mathbf{z}'_n} \approx \alpha_{n'}(B_{12}, B_{11}), \quad (19.9)$$

where  $B_{11} = \text{Re} B_1$ ,  $B_{12} = \text{Im} B_1$ , and similarly for  $B_{21}$  and  $B_{22}$ . Finally, differentiating (19.1) at the point (19.2), and using (19.5), (19.6), we get

$$\begin{aligned} \frac{\partial z_n}{\partial a^0} &\approx 2\pi i \beta_n \frac{\partial \bar{v}}{\partial a^0} = 2\pi i \beta_n \left[ -\frac{i}{2} + \frac{\kappa}{4}(1 + 2\pi i) \right] = \pi \beta_n \left[ 1 - \frac{\kappa}{2}(2\pi - i) \right] \\ \frac{\partial z_n}{\partial b^0} &\approx 2\pi i \beta_n \frac{\partial \bar{v}}{\partial b^0} = 2\pi i \beta_n \left[ \frac{1}{2} - \frac{\kappa}{4}(2\pi - i) \right] = \pi \beta_n \left[ i - \frac{\kappa}{2}(2\pi i + 1) \right] \end{aligned} \quad (19.10)$$

Similarly, using (19.5) and (19.7), we obtain

$$\frac{\partial z_n}{\partial \mathbf{z}'_{n'}} \approx \delta_{nn'}(1, i). \quad (19.11)$$

The expressions (19.10) and (19.11) can be written as vectors and matrices:

$$\frac{\partial \mathbf{z}_n}{\partial a^0} \approx \pi \beta_n \begin{pmatrix} 1 - \kappa\pi \\ \frac{\kappa}{2} \end{pmatrix}, \quad \frac{\partial \mathbf{z}_n}{\partial b^0} \approx \pi \beta_n \begin{pmatrix} -\frac{\kappa}{2} \\ 1 - \kappa\pi \end{pmatrix}, \quad \frac{\partial \mathbf{z}_n}{\partial \mathbf{z}'_{n'}} \approx \delta_{nn'} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (19.12)$$

## 20 Block-matrix approximation

All calculations below essentially rely on the constants (12.13) which correspond to the ruby laser. Let us denote the matrix

$$M = \begin{pmatrix} \frac{\partial a}{\partial a^0} & \frac{\partial a}{\partial b^0} \\ \frac{\partial b}{\partial a^0} & \frac{\partial b}{\partial b^0} \end{pmatrix}. \quad (20.1)$$

Recall that for the typical dipole moment (11.3) we have by (16.7), (16.8) and (12.13), that

$$\mathbf{S} = \sum_n \alpha_n \beta_n \sim 10^{-5}, \quad \gamma_n \approx 2 \times 10^{-7}, \quad \kappa \approx 10^{-7}. \quad (20.2)$$

Rewrite the formulas (19.8)–(19.9) substituting the expressions (19.6), and neglecting all terms containing  $\kappa \mathbf{S} \sim 10^{-12}$  and  $\kappa^2 \sim 10^{-14}$ :

$$\frac{\partial a}{\partial a^0} \approx 1 - 2\pi\kappa + \frac{1}{2}A_{22}\mathbf{S}, \quad \frac{\partial a}{\partial b^0} \approx -\frac{1}{2}A_{21}\mathbf{S}, \quad (20.3)$$

$$\frac{\partial b}{\partial a^0} \approx -\frac{1}{2}B_{22}\mathbf{S}, \quad \frac{\partial b}{\partial b^0} \approx 1 - 2\pi\kappa - \frac{1}{2}B_{21}\mathbf{S}. \quad (20.4)$$

Hence,

$$M = M(\kappa, \mathbf{S}) \approx \begin{pmatrix} 1 - 2\pi\kappa - \frac{1}{2}A_{22}\mathbf{S} & -\frac{1}{2}A_{21}\mathbf{S} \\ -\frac{1}{2}B_{22}\mathbf{S} & 1 - 2\pi\kappa - \frac{1}{2}B_{21}\mathbf{S} \end{pmatrix}. \quad (20.5)$$

Further, denote the matrices

$$D_n = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad V_n = \alpha_n \begin{pmatrix} A_{12} & A_{11} \\ B_{12} & B_{11} \end{pmatrix}, \quad W_n = \pi\beta_n \begin{pmatrix} 1 - \varkappa\pi & -\frac{\varkappa}{2} \\ \frac{\varkappa}{2} & 1 - \varkappa\pi \end{pmatrix}. \quad (20.6)$$

Using (B.7) and (B.4), we obtain that

$$V_n \approx \pi\alpha_n \begin{pmatrix} 1 - \varkappa\pi & \frac{\varkappa}{2} \\ -\frac{\varkappa}{2} & 1 - \varkappa\pi \end{pmatrix} \quad (20.7)$$

Now formulas (19.8), (19.9), (20.5), and (19.12) give the following block-matrix approximation for the differential (18.4):

$$DP_*(X_*^0) \approx \begin{pmatrix} M & V_1 & V_2 & V_3 & \dots & V_N \\ W_1 & D_1 & 0 & 0 & \dots & 0 \\ W_2 & 0 & D_2 & 0 & \dots & 0 \\ W_3 & 0 & 0 & D_3 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ W_N & 0 & 0 & 0 & \dots & D_N \end{pmatrix}, \quad (20.8)$$

which is not symmetric generally.

**Remark 20.1.** Since  $\alpha_n \sim 10^{-23}$ , the matrix is very close to its triangle approximation

$$DP_*(X_*^0) \approx \begin{pmatrix} M & 0 & 0 & 0 & \dots & 0 \\ W_1 & D_1 & 0 & 0 & \dots & 0 \\ W_2 & 0 & D_2 & 0 & \dots & 0 \\ W_3 & 0 & 0 & D_3 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ W_N & 0 & 0 & 0 & \dots & D_N \end{pmatrix}. \quad (20.9)$$

The corresponding eigenvalues coincide with the ones of the matrices  $M$  and  $D_n$ . In particular, for the eigenvalues of  $M$  we have  $|\mu - 1| \sim 10^{-5}$  by (20.5) and (16.7).

### 20.0.1 Reduction to polynomial equation

The spectral problem for the multipliers reads

$$DP_*(X_*^0)\mathbf{X} = \mu\mathbf{X}, \quad \mathbf{X} = \begin{pmatrix} \mathbf{x}_0 \\ \mathbf{x}_1 \\ \dots \\ \mathbf{x}_N \end{pmatrix}, \quad (20.10)$$

where all  $\mathbf{x}_n \in \mathbb{R}^2$  and  $\mathbf{X} \neq 0$ . We will reduce the spectral problem to a polynomial equation. Namely, substituting here the approximation (20.8), we obtain the equivalent system

$$(M - \mu)\mathbf{x}_0 + \sum_1^N V_n \mathbf{x}_n = 0, \quad W_n \mathbf{x}_0 + (D_n - \mu)\mathbf{x}_n = 0, \quad n \in \overline{N}. \quad (20.11)$$

Since we seek for eigenvalues with  $|\mu| > 1$ , we can write

$$\mathbf{x}_n = (\mu - D_n)^{-1} W_n \mathbf{x}_0 \approx \pi\beta_n \begin{pmatrix} \frac{1 - \varkappa\pi}{\mu - 1} & \frac{-\varkappa/2}{\mu - 1} \\ \frac{\varkappa/2}{\mu - 1 + 2\pi^2\gamma_n^2} & \frac{1 - \varkappa\pi}{\mu - 1 + 2\pi^2\gamma_n^2} \end{pmatrix} \mathbf{x}_0, \quad (20.12)$$

where  $\mathbf{x}_0 \neq 0$ . Substituting into the first equation of (20.11), we obtain that  $\mathcal{M}(\mu)\mathbf{x}_0 = 0$ , where

$$\begin{aligned} \mathcal{M}(\mu) &\approx (M-\mu) + \pi^2 \sum_1^N \left[ \alpha_n \beta_n \begin{pmatrix} 1 - \varkappa\pi & \frac{\varkappa}{2} \\ -\frac{\varkappa}{2} & 1 - \varkappa\pi \end{pmatrix} \begin{pmatrix} \frac{1 - \varkappa\pi}{\mu-1} & \frac{-\varkappa/2}{\mu-1} \\ \frac{\varkappa/2}{\mu-1+2\pi^2\gamma_n^2} & \frac{1 - \varkappa\pi}{\mu-1+2\pi^2\gamma_n^2} \end{pmatrix} \right] \\ &= (M-\mu) + \pi^2 \begin{pmatrix} 1 - \varkappa\pi & \frac{\varkappa}{2} \\ -\frac{\varkappa}{2} & 1 - \varkappa\pi \end{pmatrix} \begin{pmatrix} \frac{1 - \varkappa\pi}{\mu-1} \mathbf{S} & \frac{-\varkappa/2}{\mu-1} \mathbf{S} \\ \frac{\varkappa}{2} \sum_1^N \frac{\alpha_n \beta_n}{\mu-1+2\pi^2\gamma_n^2} & (1 - \varkappa\pi) \sum_1^N \frac{\alpha_n \beta_n}{\mu-1+2\pi^2\gamma_n^2} \end{pmatrix}. \end{aligned} \quad (20.13)$$

Thus, the characteristic equation for the matrix  $DP(X^0)$  is equivalent to

$$\det \mathcal{M}(\mu) = 0. \quad (20.14)$$

Let us consider the multipliers for which  $2\pi^2 \frac{\gamma_n^2}{\mu-1} \ll 1$ . The existence of such multipliers is suggested by Remark 20.1. For such multipliers, the last two sums in (20.13) can be calculated as follows. First,

$$\begin{aligned} \sum_1^N \frac{\alpha_n \beta_n}{\mu-1+2\pi^2\gamma_n^2} &= \frac{1}{\mu-1} \sum_1^N \frac{\alpha_n \beta_n}{1+2\pi^2 \frac{\gamma_n^2}{\mu-1}} \approx \frac{1}{\mu-1} \sum_1^N \alpha_n \beta_n (1 - 2\pi^2 \frac{\gamma_n^2}{\mu-1}) \\ &= \frac{\mathbf{S}}{\mu-1} - \frac{2\pi^2}{(\mu-1)^2} \sum_1^N \alpha_n \beta_n \gamma_n^2 = \frac{\mathbf{S}}{\mu-1} - \frac{2\pi^2}{(\mu-1)^2} \frac{2}{\Omega_p \hbar^2 c} \sum_1^N (\mathbf{P}_n \cdot \mathbf{X}(x_n))^2 (\mathbf{P}_n \cdot \mathbf{a}_p(x_n))^2. \end{aligned} \quad (20.15)$$

Second, by Remark 12.1, we can apply the *law of large numbers* for weakly dependent random values [21], and obtain that  $\mathbf{S} \approx \overline{N} \mathbb{E} \alpha_n \beta_n$ .

$$\sum_1^N (\mathbf{P}_n \cdot \mathbf{X}(x_n))^2 (\mathbf{P}_n \cdot \mathbf{a}_p(x_n))^2 = N \Sigma, \quad \Sigma := \mathbb{E} (\mathbf{P}_n \cdot \mathbf{X}(x_n))^2 (\mathbf{P}_n \cdot \mathbf{a}_p(x_n))^2. \quad (20.16)$$

The last expectation can be calculated using the fact that random values  $(\mathbf{P}^1, \mathbf{P}^2, \mathbf{P}^3) = \mathbf{P}_n$ ,  $(X^1, X^2, X^3) = \mathbf{X}(x_n)$  and  $(a^1, a^2, a^3) = \mathbf{a}_p(x_n)$  are independent, so

$$\Sigma = \mathbb{E} \sum_{i,l=1}^3 (\mathbf{P}^i X^i)^2 (\mathbf{P}^l X^l)^2 = \mathbb{E} \sum_{i,i',l,l'=1}^3 \mathbf{P}^i \mathbf{P}^{i'} \mathbf{P}^l \mathbf{P}^{l'} X^i X^{i'} a^l a^{l'} = \sum_{i,i',l,l'=1}^3 \mathbb{E} [\mathbf{P}^i \mathbf{P}^{i'} \mathbf{P}^l \mathbf{P}^{l'}] \mathbb{E} [X^i X^{i'}] \mathbb{E} [a^l a^{l'}] \quad (20.17)$$

By our assumption **H4**, the random vector  $\mathbf{a}_p(x_n)$  is distributed uniformly on the sphere of a radius  $a_p$ , so

$$\mathbb{E} [a^l a^{l'}] = \frac{1}{3} \delta^{ll'} a_p^2. \quad (20.18)$$

Further, *let us conjecture* that the distribution of the random vector  $\mathbf{X}(x_n)$  is close to the one of the eigenmodes (7.6) of rectangular cuboid  $V = [0, l_1] \times [0, l_2] \times [0, l_3]$ . In this case, similarly to (16.6),

$$\mathbb{E} [X^i X^{i'}] = \delta^{ii'} \mathbb{E} |X^i|^2 = \frac{\delta^{ii'}}{N} \sum_{n=1}^N |\mathbf{X}^i(x_n)|^2 \approx \frac{\delta^{ii'}}{|V_a|} \int_{V_a} |X^i|^2(x) dx = \frac{\delta^{ii'}}{3|V|} |a^i|^2, \quad (20.19)$$

where  $a^i = a_k^i$  are the amplitudes from (7.6). Substituting (20.18) and (20.19) into (20.17), we obtain that

$$\Sigma = \frac{1}{9|V|} |a_p|^2 \sum_{i,l=1}^3 \mathbb{E} [|\mathbf{P}^i|^2 |\mathbf{P}^l|^2] |a^i|^2 \quad (20.20)$$

**Remark 20.2.** According to Berry's conjecture, in the case of ergodic geodesic flow, the distribution of values of Laplacian eigenfunctions with sufficiently large eigenvalues can be Gaussian [6, 26], thus different from the "sinusoidal distribution" of eigenmodes (7.6).



Let us consider the following two cases separately.

**I. Polycrystalline medium.** Let us calculate  $\mathbb{E}[|\mathbf{P}^i|^2|\mathbf{P}^l|^2]$ . By our assumption **H1**, the random vector  $(\mathbf{P}^1, \mathbf{P}^2, \mathbf{P}^3)$  is distributed uniformly on the sphere of radius  $|\mathbf{P}|$ . Hence, integrating in spherical coordinates, we obtain

$$\mathbb{E}|\mathbf{P}^1|^2|\mathbf{P}^2|^2 = \frac{1}{15}|\mathbf{P}|^4, \quad \mathbb{E}(\mathbf{P}^1)^4 = \frac{1}{5}|\mathbf{P}|^4. \quad (20.21)$$

Hence,

$$\Sigma = \frac{1}{9|V|}|a_p|^2|\mathbf{P}|^4 \sum_{i=1}^3 \left(\frac{1}{5} + \frac{2}{15}\right)|a^i|^2 = \frac{1}{27|V|}|a_p|^2|\mathbf{P}|^4 \quad (20.22)$$

since  $a = (a^1, a^2, a^3)$  is the unit vector.

**II. Crystalline medium.** In this case the vectors  $\mathbf{P}_n = \mathbf{P}$  do not depend on  $n$ . Thus, (20.20) becomes

$$\Sigma = \frac{1}{9|V|}|a_p|^2 \sum_{i,l=1}^3 [|\mathbf{P}^i|^2|\mathbf{P}^l|^2]|a^i|^2 = \frac{a_p^2|\mathbf{P}|^2}{9|V|} \sum_{i=1}^3 |\mathbf{P}^i|^2|a^i|^2. \quad (20.23)$$

In both cases, (20.14) for  $\mu \neq 1$  is equivalent to the algebraic equation

$$p(\mu) = 0, \quad p(\mu) = \det[(\mu - 1)^2 \mathcal{M}(\mu)], \quad (20.24)$$

where  $p(\mu)$  is a polynomial of degree six.

**Remark 20.3.** i) The summation **S** reappears in the calculations (20.13) and (20.15).

ii) The degree six of the polynomial can be reduced to three using the approximate symmetry of type (4.7), see Remark 4.3.

**Remark 20.4.** For every nonzero eigenvector (20.10), its component  $\mathbf{x}_0$ , representing the Maxwell field, does not vanish. Hence, if (1.1) holds, then the magnitude of the Maxwell field provisionally increases exponentially for large (but bounded) times.

**Remark 20.5.** We expect that

- i) all the multipliers  $|\mu|$  are simple with probability one;
- ii) the multipliers form *clusters* which are close to the corresponding roots of the polynomial  $p(\mu)$ ;
- iii) the condition (1.1) holds for the roots of the polynomial  $p(\mu)$  when

$$a_p \geq d(\sigma_1), \quad (20.25)$$

where  $d(\sigma_1)$  is the corresponding threshold, which is an increasing function of  $\sigma_1$ . Physically, the condition (20.25) means that the amplitude of the pumping exceeds the damping losses.

## A On averaging of slow rotations

Here we justify the approximation of the Schrödinger equations from (13.3) by their averaged version (14.7) (“rotating wave approximation” [1, 3, 14, 28, 31, 33, 36, 42]). Let us show that this approximation is very accurate for small matrices  $\Omega_n(\tau)$ . Namely, consider the system

$$\dot{c}(\tau) = -i\Omega(\tau)c(\tau), \quad \tau \in [0, T], \quad (A.1)$$

where  $\Omega(\tau)$  is a bounded measurable  $2 \times 2$  complex matrix-function,

$$\varepsilon := \sup_{\tau \in [0, T]} |\Omega(\tau)| < \infty. \quad (A.2)$$

Denote the averaged matrix

$$\tilde{\Omega} := \frac{1}{T} \int_0^T \Omega(\tau) d\tau \quad (A.3)$$

and consider the corresponding averaged system

$$\dot{\hat{c}}(\tau) = -i\tilde{\Omega}\hat{c}(\tau), \quad \tau \in [0, T]. \quad (A.4)$$

**Lemma A.1.** For solutions of (A.1) and (A.4) with identical initial data  $\hat{c}(0) = c(0) = c_0$ , their final values  $c(T)$  and  $\hat{c}(T)$  are very close for small  $\varepsilon$ :

$$|c(T) - \hat{c}(T)| = O(\varepsilon^2), \quad \varepsilon \rightarrow 0. \quad (\text{A.5})$$

*Proof.* Rewrite (A.1) as

$$\dot{c}_\varepsilon(\tau) = -i\varepsilon\Omega_1(\tau)c_\varepsilon(\tau), \quad \tau \in [0, T]; \quad c_\varepsilon(0) = c_0, \quad (\text{A.6})$$

where  $\Omega_1(\tau) := \frac{1}{\varepsilon}\Omega(\tau)$ . The corresponding averaged system (A.4) now reads

$$\dot{\hat{c}}_\varepsilon(\tau) = -i\varepsilon\hat{\Omega}_1(\tau)\hat{c}_\varepsilon(\tau), \quad \tau \in [0, T]; \quad \hat{c}_\varepsilon(0) = c_0, \quad (\text{A.7})$$

where  $\tilde{\Omega}_1(\tau) := \frac{1}{T} \int_0^T \Omega_1(\tau) d\tau$ . It suffices to prove that

$$|c_\varepsilon(T) - \hat{c}_\varepsilon(T)| = O(\varepsilon^2), \quad \varepsilon \rightarrow 0. \quad (\text{A.8})$$

The Taylor formula gives

$$c_\varepsilon(\tau) = c_0 + \varepsilon[\partial_\varepsilon c_\varepsilon(\tau)]|_{\varepsilon=0} + O(\varepsilon^2), \quad \hat{c}_\varepsilon(\tau) = c_0 + \varepsilon[\partial_\varepsilon \hat{c}_\varepsilon(\tau)]|_{\varepsilon=0} + O(\varepsilon^2). \quad (\text{A.9})$$

Therefore, (A.8) will follow from the identity

$$[\partial_\varepsilon c_\varepsilon(T)]|_{\varepsilon=0} = [\partial_\varepsilon \hat{c}_\varepsilon(T)]|_{\varepsilon=0}. \quad (\text{A.10})$$

Indeed,  $\hat{c}_\varepsilon(T) = e^{-i\varepsilon\tilde{\Omega}_1 T} c_0$ , so the right-hand side of (A.10) is equal to

$$[\partial_\varepsilon \hat{c}_\varepsilon(T)]|_{\varepsilon=0} = -i\tilde{\Omega}_1 T c_0 = -i \int_0^T \Omega_1(\tau) c_0 d\tau. \quad (\text{A.11})$$

It remains to calculate the left-hand side. Denote

$$b_\varepsilon(\tau) := \partial_\varepsilon c_\varepsilon(\tau). \quad (\text{A.12})$$

Differentiating equation (A.6) in  $\varepsilon$ , we obtain

$$\dot{b}_\varepsilon(\tau) = -i\Omega_1(\tau)c_\varepsilon(\tau) - i\varepsilon\Omega_1(\tau)b_\varepsilon(\tau), \quad \tau \in [0, T]; \quad b_\varepsilon(0) = 0. \quad (\text{A.13})$$

In particular, for  $\varepsilon = 0$  we have

$$\dot{b}_0(\tau) = -i\Omega_1(\tau)c_0, \quad \tau \in [0, T]; \quad b_0(0) = 0. \quad (\text{A.14})$$

Hence,

$$b_0(T) = -i \int_0^T \Omega_1(\tau) c_0 d\tau. \quad (\text{A.15})$$

Now (A.10) is proved.  $\square$

**Remark A.2.** Equation (A.6) can be rewritten for  $\tilde{c}_\varepsilon(s) := c_\varepsilon(\frac{s}{\varepsilon})$  as

$$\partial_s \tilde{c}_\varepsilon(s) = -i\Omega_1(\frac{s}{\varepsilon})\tilde{c}_\varepsilon(s), \quad s \in [0, \varepsilon T], \quad (\text{A.16})$$

where  $s = \varepsilon\tau$  is the ‘slow time’. Thus, Lemma A.1 is a specific version of averaging principle.

## B Some integrals

### 2.1 Integrals (14.26)

First, (14.17) implies that

$$\begin{aligned}
I_1(\tau) &:= \int_0^\tau e^{-i\tau'} e^{-\varkappa(\tau-\tau')} \sin(\tau-\tau') d\tau' = \int_0^\tau e^{-i\tau'} [e^{(-\varkappa+i)(\tau-\tau')} - e^{(-\varkappa-i)(\tau-\tau')}] d\tau' \\
&= \frac{1}{2i} e^{(-\varkappa+i)\tau} \int_0^\tau e^{(\varkappa-2i)\tau'} d\tau' - \frac{1}{2i} e^{(-\varkappa-i)\tau} \int_0^\tau e^{\varkappa\tau'} d\tau' \\
&= \frac{1}{2i} e^{(-\varkappa+i)\tau} \frac{e^{(\varkappa-2i)\tau} - 1}{\varkappa - 2i} - \frac{1}{2i} e^{(-\varkappa-i)\tau} \frac{e^{\varkappa\tau} - 1}{\varkappa} \\
&= \frac{e^{-i\tau} - e^{(-\varkappa+i)\tau}}{2i\varkappa + 4} - \frac{e^{-i\tau} - e^{(-\varkappa-i)\tau}}{2i\varkappa} \\
&\approx -\frac{i}{2} \sin \tau - \frac{\tau}{2i} e^{-i\tau} + \varkappa \left[ \frac{-\sin \tau + \tau e^{i\tau}}{4} + \frac{\tau^2 e^{-i\tau}}{4i} \right], \tag{B.1}
\end{aligned}$$

where we have neglected errors  $O(\varkappa^2) = O(10^{-14})$ . Similarly,

$$\begin{aligned}
I_2(\tau) &:= \frac{1}{2} \int_0^\tau \tau' \cos \tau' e^{-\varkappa(\tau-\tau')} \sin(\tau-\tau') d\tau' = \frac{1}{4} \int_0^\tau \tau' e^{-\varkappa(\tau-\tau')} [\sin \tau + \sin(\tau-2\tau')] d\tau' \\
&= \frac{1}{4} e^{-\varkappa\tau} \left[ \sin \tau \int_0^\tau \tau' e^{\varkappa\tau'} d\tau' + \text{Im} \int_0^\tau \tau' e^{\varkappa\tau'} e^{i(\tau-2\tau')} d\tau' \right] \\
&= \frac{1}{4} e^{-\varkappa\tau} \left[ \sin \tau \left( \tau \frac{e^{\varkappa\tau}}{\varkappa} - \int_0^\tau \frac{e^{\varkappa\tau'}}{\varkappa} d\tau' \right) + \text{Im} \left( e^{i\tau} \tau \frac{e^{(\varkappa-2i)\tau}}{\varkappa-2i} - e^{i\tau} \int_0^\tau \frac{e^{(\varkappa-2i)\tau'}}{\varkappa-2i} d\tau' \right) \right] \\
&= \frac{1}{4} e^{-\varkappa\tau} \left[ \sin \tau \left( \tau \frac{e^{\varkappa\tau}}{\varkappa} - \frac{e^{\varkappa\tau} - 1}{\varkappa^2} \right) + \text{Im} \left( e^{i\tau} \tau \frac{e^{(\varkappa-2i)\tau}}{\varkappa-2i} - e^{i\tau} \frac{e^{(\varkappa-2i)\tau} - 1}{(\varkappa-2i)^2} \right) \right] \\
&= \frac{1}{4} e^{-\varkappa\tau} \left[ \sin \tau \left( \frac{\varkappa\tau e^{\varkappa\tau} - e^{\varkappa\tau} + 1}{\varkappa^2} \right) + \text{Im} \left( \tau \frac{e^{(\varkappa-i)\tau}(\varkappa+2i)}{\varkappa^2+4} - \frac{(e^{(\varkappa-i)\tau} - e^{i\tau})(\varkappa+2i)^2}{(\varkappa^2+4)^2} \right) \right] \\
&\approx \frac{1}{4} e^{-\varkappa\tau} \left[ \sin \tau \left( \frac{\tau^2}{2} + \frac{\varkappa\tau^3}{2} \right) + \tau \frac{e^{\varkappa\tau}}{4} \text{Im} (e^{-i\tau}(\varkappa+2i)) - \frac{1}{16} \text{Im} [(e^{(\varkappa-i)\tau} - e^{i\tau})(\varkappa+2i)^2] \right] \\
&\approx \frac{1}{4} e^{-\varkappa\tau} \left[ \sin \tau \left( \frac{\tau^2}{2} + \frac{\varkappa\tau^3}{2} \right) + \tau \frac{e^{\varkappa\tau}}{4} [2 \cos \tau - \varkappa \sin \tau] \right. \\
&\quad \left. - \frac{1}{16} [(e^{\varkappa\tau} \cos \tau - \cos \tau)4\varkappa - 4(-e^{\varkappa\tau} \sin \tau - \sin \tau)] \right] \\
&\approx \frac{1}{4} e^{-\varkappa\tau} \left[ \sin \tau \left( \frac{\tau^2}{2} + \frac{\varkappa\tau^3}{2} \right) + \frac{\tau}{4} [e^{\varkappa\tau} 2 \cos \tau - \varkappa \sin \tau] - \frac{1}{4} (e^{\varkappa\tau} + 1) \sin \tau \right] \\
&\approx \frac{1}{4} \left[ \sin \tau \frac{\tau^2}{2} + \frac{\tau}{4} [2 \cos \tau - \varkappa \sin \tau] - \frac{1}{4} (1 + e^{-\varkappa\tau}) \sin \tau \right] \\
&\approx \frac{1}{4} \left[ \sin \tau \frac{\tau^2}{2} + \frac{\tau}{4} [2 \cos \tau - \varkappa \sin \tau] - \frac{1}{4} (2 - \varkappa\tau) \sin \tau \right] \\
&= \frac{1}{8} \left[ \tau^2 \sin \tau + \tau \cos \tau - \sin \tau \right]. \tag{B.2}
\end{aligned}$$

## 2.2 Integrals (14.32)

Calculate integrals

$$\begin{aligned} \int_0^{2\pi} \tau e^{i\tau} d\tau &= -2\pi i \approx -6.28i, & \int_0^{2\pi} \tau \cos \tau d\tau &= 0, & \int_0^{2\pi} \tau e^{2i\tau} d\tau &= -\pi i, \\ \int_0^{2\pi} \tau^2 e^{i\tau} d\tau &= -4\pi^2 i + i \int_0^{2\pi} 2\tau e^{i\tau} d\tau = -4\pi^2 i + 4\pi, \\ \int_0^{2\pi} \tau^2 e^{2i\tau} d\tau &= -2\pi^2 i + i \int_0^{2\pi} \tau e^{2i\tau} d\tau = -2\pi^2 i + \pi, \\ \int_0^{2\pi} \tau^2 \sin \tau d\tau &= -4\pi^2 i. \end{aligned}$$

Then neglecting errors  $O(\varkappa^2) = O(10^{-14})$ , we can rewrite (14.32) as

$$\left\{ \begin{aligned} J_1 &:= \frac{1}{2\pi} \int_0^{2\pi} I_1'(\tau) e^{-i\tau} d\tau \approx \frac{1}{2\pi} \int_0^{2\pi} \left[ -\frac{i}{2} \sin \tau - \frac{\tau}{2i} e^{-i\tau} + \varkappa \left[ \frac{-\sin \tau + \tau e^{i\tau}}{4} + \frac{\tau^2 e^{-i\tau}}{4i} \right] \right]' e^{-i\tau} d\tau \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[ -\frac{i}{2} \cos \tau - \frac{e^{-i\tau} - i\tau e^{-i\tau}}{2i} + \varkappa \left[ \frac{-\cos \tau + e^{i\tau} + i\tau e^{i\tau}}{4} + \frac{2\tau e^{-i\tau} - i\tau^2 e^{-i\tau}}{4i} \right] \right] e^{-i\tau} d\tau \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[ -\frac{i}{2} \cos \tau e^{-i\tau} - \frac{e^{-2i\tau} - i\tau e^{-2i\tau}}{2i} + \varkappa \left[ \frac{-\cos \tau e^{-i\tau} + i\tau}{4} + \frac{2\tau e^{-2i\tau} - i\tau^2 e^{-2i\tau}}{4i} \right] \right] d\tau \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[ -\frac{i}{4} + \frac{\tau e^{-2i\tau}}{2} + \varkappa \left[ \frac{-\frac{1}{2} + i\tau}{4} + \frac{2\tau e^{-2i\tau} - i\tau^2 e^{-2i\tau}}{4i} \right] \right] d\tau \\ &= -\frac{i}{4} + \frac{\pi i}{4\pi} + \varkappa \left[ \frac{-\frac{1}{2} + i\pi}{4} + \frac{2\pi i - i(2\pi^2 i + \pi)}{8\pi i} \right] = 0, \\ J_2 &:= \frac{1}{2\pi} \int_0^{2\pi} I_2'(\tau) e^{-i\tau} d\tau \approx \frac{1}{16\pi} \int_0^{2\pi} \left[ \tau^2 \sin \tau + \tau \cos \tau - \sin \tau \right]' e^{-i\tau} d\tau \\ &= \frac{1}{16\pi} \int_0^{2\pi} \left[ \tau \sin \tau + \tau^2 \cos \tau \right] e^{-i\tau} d\tau = \frac{1}{32\pi} \int_0^{2\pi} \left[ -i\tau(1 - e^{-2i\tau}) + \tau^2(1 + e^{-2i\tau}) \right] d\tau \\ &= \frac{1}{32\pi} \left[ -i2\pi^2 - \pi + \frac{8\pi^3}{3} + 2\pi^2 i + \pi \right] = \frac{\pi^2}{12}. \end{aligned} \right. \quad (\text{B.3})$$

## 2.3 Integrals (15.5)

Neglecting errors  $O(\varkappa^2) = O(10^{-14})$ , we obtain

$$\begin{aligned} A_1 &\approx - \int_0^{2\pi} e^{-i\tau'} e^{-\varkappa(2\pi-\tau')} \sin \tau' d\tau' = -\frac{e^{-2\pi\varkappa}}{2i} \int_0^{2\pi} e^{-i\tau'} e^{\varkappa\tau'} [e^{i\tau'} - e^{-i\tau'}] d\tau' \\ &= -\frac{e^{-2\pi\varkappa}}{2i} \int_0^{2\pi} [e^{\varkappa\tau'} - e^{(\varkappa-2i)\tau'}] d\tau' = -\frac{e^{-2\pi\varkappa}}{2i} \left[ \frac{e^{2\pi\varkappa} - 1}{\varkappa} - \frac{e^{2\pi\varkappa} - 1}{\varkappa - 2i} \right] \\ &\approx -\frac{e^{-2\pi\varkappa}}{2i} \left[ \frac{2\pi\varkappa + \frac{1}{2}(2\pi\varkappa)^2}{\varkappa} - \frac{2\pi\varkappa(\varkappa + 2i)}{4} \right] \approx -\frac{e^{-2\pi\varkappa}}{2i} \left[ 2\pi + 2\varkappa\pi^2 - \varkappa\pi i \right] \\ &\approx (1 - 2\pi\varkappa) \left[ \pi i + \varkappa\pi^2 i + \frac{\varkappa\pi}{2} \right] \approx \pi i + \varkappa\pi^2 i + \frac{\varkappa\pi}{2} - \pi^2 2\varkappa i \\ &= \pi i + \varkappa[-\pi^2 i + \frac{\pi}{2}] = A_{11} + iA_{12}, \quad A_{11} = \varkappa\frac{\pi}{2}, \quad A_{12} = \pi - \varkappa\pi^2. \end{aligned} \quad (\text{B.4})$$

Similarly,

$$\begin{aligned}
A_2 &\approx -\frac{e^{-\pi\sigma_1}}{2i} \int_0^{2\pi} \tau' [e^{\varkappa\tau'} - e^{(\varkappa-2i)\tau'}] d\tau' \\
&= -\frac{e^{-\pi\sigma_1}}{2i} \left[ \tau \frac{e^{\varkappa\tau}}{\varkappa} - \frac{e^{\varkappa\tau}}{\varkappa^2} - \tau \frac{e^{\varkappa\tau}}{\varkappa-2i} + \frac{e^{\varkappa\tau}}{(\varkappa-2i)^2} \right] \Big|_0^{2\pi} = -\frac{e^{-\pi\sigma_1}}{2i} \left[ \frac{\tau\varkappa-1}{\varkappa^2} e^{\varkappa\tau} - \frac{\tau(\varkappa-2i)-1}{(\varkappa-2i)^2} e^{\varkappa\tau} \right] \Big|_0^{2\pi} \\
&\approx -\frac{e^{-\pi\sigma_1}}{2i} \left[ \frac{\tau\varkappa-1}{\varkappa^2} e^{\varkappa\tau} - \frac{\tau(\varkappa-2i)-1}{16} (\varkappa+2i)^2 e^{\varkappa\tau} \right] \Big|_0^{2\pi} \\
&= -\frac{e^{-\pi\sigma_1}}{2i} \left[ \frac{(\pi\sigma_1-1)e^{\pi\sigma_1}+1}{\varkappa^2} - \frac{(\varkappa+2i)^2}{16} [(\pi\sigma_1-4\pi i-1)e^{\pi\sigma_1}+1] \right] \\
&\approx -\frac{e^{-\pi\sigma_1}}{2i} \left[ \frac{(\pi\sigma_1-1)(1+\pi\sigma_1+\frac{1}{2}(\pi\sigma_1)^2+\frac{1}{6}(\pi\sigma_1)^3)+1}{(\varkappa)^2} - \pi i - \frac{\pi\sigma_1}{2} + \frac{1}{4}[\pi\sigma_1+(-4\pi i-1)\pi\sigma_1] \right] \\
&\approx -\frac{e^{-\pi\sigma_1}}{2i} \left[ \frac{(\pi\sigma_1)^2+(\pi\sigma_1-1)(\frac{1}{2}(\pi\sigma_1)^2+\frac{1}{6}(\pi\sigma_1)^3)}{\varkappa^2} - \pi i - \frac{\pi\sigma_1}{2} + \frac{1}{4}[\pi\sigma_1+(-4\pi i-1)\pi\sigma_1] \right] \\
&= -\frac{e^{-\pi\sigma_1}}{2i} \left[ (4\pi^2+(\pi\sigma_1-1)(2\pi^2+\frac{2}{3}\pi^3\sigma_1) - \pi i - \frac{\pi\sigma_1}{2} - \pi^2\sigma_1 i] \right] \\
&\approx -\frac{e^{-\pi\sigma_1}}{2i} \left[ 2\pi^2 - \pi i + \sigma_1 \left[ \frac{4}{3}\pi^3 - \frac{\pi}{2} - \pi^2 i \right] \right] = e^{-\pi\sigma_1} \left[ \pi^2 i + \frac{\pi}{2} + \sigma_1 \left[ \frac{2}{3}\pi^3 i - \frac{\pi}{4} i + \frac{\pi^2}{2} \right] \right] \\
&= \left[ \pi^2 i + \frac{\pi}{2} + \sigma_1 \left[ \frac{2}{3}\pi^3 i - \frac{\pi}{4} i + \frac{\pi^2}{2} \right] \right] - \pi\sigma_1 \left[ \pi^2 i + \frac{\pi}{2} \right] \\
&= \pi^2 i + \frac{\pi}{2} - \sigma_1 i \left[ \frac{\pi^3}{3} + \frac{\pi}{4} \right] = A_{21} + iA_{22}, \quad A_{21} = \frac{\pi}{2}, \quad A_{22} = \pi^2 - \sigma_1 \left[ \frac{\pi^3}{3} + \frac{\pi}{4} \right]. \tag{B.5}
\end{aligned}$$

Hence,

$$A_3 = \text{Re} A_2 \approx \pi^2 - \sigma_1 \left[ \frac{\pi^3}{3} + \frac{\pi}{4} \right]. \tag{B.6}$$

Finally,

$$\left\{ \begin{array}{l} B_1 = -i \int_0^{2\pi} e^{-i\tau'} E(2\pi - \tau') d\tau' = -iA_1. \\ B_2 = \int_0^{2\pi} [e^{-i\tau'} - i\tau' e^{-i\tau'}] E(2\pi - \tau') d\tau' = A_1 - iA_2 \\ B_3 = \text{Re} B_2 = A_{11} + A_{22} \end{array} \right. \tag{B.7}$$

## References

- [1] L. Allen, J.H. Eberly, *Optical Resonance and Two-Level Atoms*, Wiley, New York, 1975.
- [2] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer, New York, 1989.
- [3] S.H. Autler, C.H. Townes, Stark effect in rapidly varying fields, *Phys. Rev.* 100 (1955), no. 2, 703–722.
- [4] I. Bejenaru, D. Tataru, Global wellposedness in the energy space for the Maxwell–Schrödinger system, *Commun. Math. Phys.* **288** (2009), 145–198.
- [5] V. Benci, D. Fortunato, An eigenvalue problem for the Schrödinger–Maxwell equations, *Topological Methods in Nonlinear Analysis* **11** (1998), 283–293.
- [6] M.V. Berry, Regular and irregular semiclassical wavefunctions, *J. Phys. A: Math, Gen.* **10** (1977), no. 12, 2083–2091.

- [7] G. M. Coclite, V. Georgiev, Solitary waves for Maxwell–Schrödinger equations, *Electronic Journal of Differential Equations* 94 (2004), 1–31. [arXiv:math/0303142](https://arxiv.org/abs/math/0303142) [math.AP]
- [8] Ju.L. Daleckii, M.G. Krein, *Stability of Solutions of Differential Equations in Banach Space*, Amer Mathematical Society, Providence, Rhode Island, 1974. ISBN-10. 0821832387.
- [9] R.H. Dicke, Coherence in Spontaneous Radiation Processes, *Phys. Rev.* **93** (1954), no. 1, 99–110.
- [10] J. Demaison, W. Hüttner, *Dipole Moments and Related Constants of Diamagnetic Molecules*, Springer, Berlin, 1992.
- [11] A. Einstein, Quantentheorie der Strahlung (On the quantum theory of radiation) *Phys. Z.* **18** (1917), 121–128.
- [12] B. Garraway, The Dicke model in quantum optics: Dicke model revisited, *Phil. Trans. R. Soc. A* **369** (2011), 1137–1155.
- [13] Y. Guo, K. Nakamitsu, W. Strauss, Global finite-energy solutions of the Maxwell–Schrödinger system, *Comm. Math. Phys.* **170** (1995), no. 1, 181–196.
- [14] H. Haken, *Laser Theory*, Springer, Berlin, 1984.
- [15] K. Hepp, E.H. Lieb, On the superradiant phase transition for molecules in a quantised radiation field: The Dicke Maser model, *Ann. Physics* **76** (1973), 360–404.
- [16] K. Hepp, E.H. Lieb, Phase transitions in reservoir-driven open systems with applications to lasers and superconductors, *Helv. Phys. Acta* **46** (1973), 573–603.
- [17] K. Hepp, E.H. Lieb, Equilibrium statistical mechanics of matter interacting with the quantised radiation field, *Phys. Rev. A (3)* **8** (1973), 2517–2525.
- [18] K. Hepp, E.H. Lieb, The laser: a reversible quantum dynamical system with irreversible classical macroscopic motion. Dynamical systems, theory and applications (Rencontres, Battelle Res. Inst., Seattle, Wash., 1974), pp. 178–207 in: *Lecture Notes in Phys.*, Vol. 38, Springer, Berlin, 1975.
- [19] K. Hepp, E.H. Lieb, Laser models, pp 145–234 in: E. H. Lieb, *Condensed Matter Physics and Exactly Soluble Models*, Springer, Berlin, 2004.
- [20] R. D. Jackson, *Classical Electrodynamics*, Wiley, New York, 1999.
- [21] I. A. Ibragimov, Ju. V. Linnik, *Independent and Stationary Sequences of Random Variables*, Wolters-Noordhoff, Groningen, 1971.
- [22] A. Komech, *Quantum Mechanics: Genesis and Achievements*, Springer, Dordrecht, 2013.
- [23] A.I. Komech, *Lectures on Quantum Mechanics and Attractors*, World Scientific, Singapore, 2022.
- [24] A.I. Komech, On quantum jumps and attractors of the Maxwell–Schrödinger equations, *Annales mathématiques du Québec* **46** (2022), 139–159. [arXiv 1907.04297](https://arxiv.org/abs/1907.04297).
- [25] A. Komech, E. Kopylova, *Attractors of Hamiltonian Nonlinear Partial Differential Equations*, Cambridge University Press, Cambridge, 2021.
- [26] T. Letendre, H. Ueberschär, Random moments for the new eigenfunctions of point scatterers on rectangular flat tori, *Annales Henri Poincaré*, In press, [ff10.1007/s00023-021-01032-5](https://doi.org/10.1007/s00023-021-01032-5). [ffhal02308180v2](https://arxiv.org/abs/2008.180v2)
- [27] P. Mandel, *Theoretical Problems in Cavity Nonlinear Optics*, Cambridge University Press, Cambridge, 1997.
- [28] H. Nussenzveig, *Introduction to Quantum Optics*, Gordon and Breach, London, 1973.
- [29] K. Petersen, J. P. Solovej, Existence of travelling wave solutions to the Maxwell–Pauli and Maxwell–Schrödinger systems, Archive: [arXiv:1402.3936](https://arxiv.org/abs/1402.3936) .

- [30] A. Pikovsky, M. Rosenblum, J. Kurths, Synchronization. A Universal Concept in Nonlinear Sciences, Cambridge University Press, Cambridge, 2001.
- [31] I.I. Rabi, Space quantization in a gyrating magnetic field, *Phys. Rev.* **51** (1937), 652–654.
- [32] J.J. Sakurai, Advanced Quantum Mechanics, Pearson Education, Incorporated, 2006.
- [33] M. Sargent III, M.O. Scully, W.E. Lamb Jr, Laser Physics, Addison Wesley, Reading, 1978.
- [34] P. Sarnak, Recent progress on the quantum unique ergodicity conjecture, *Bull. AMS* **48**, no. 2, 211–228.
- [35] L.I. Schiff, Quantum Mechanics, McGraw-Hill, New York, 1955.
- [36] M. O. Scully, M. S. Zubairy, Quantum Optics, Cambridge University Press, Cambridge, 1997.
- [37] W.T. Silfvast, Laser Fundamentals, Cambridge University Press, Cambridge, 2012.
- [38] A. Shimomura, Modified wave operators for Maxwell–Schrödinger equations in three space dimensions, *Ann. Henri Poincaré* **4** (2003), 661–683.
- [39] A.I. Shnirelman, Ergodic properties of eigenfunctions, *Usp. Mat. Nauk* **29** (1974), no. 6, 181–182.
- [40] A.I. Shnirelman, On the asymptotic properties of eigenfunctions in the region of chaotic motion, addendum to V.F.Lazutkin, KAM theory and semiclassical approximations to eigenfunctions, Springer 1993.
- [41] V.A. Yakubovich, V.M. Starzhinskii, Linear Differential Equations with Periodic Coefficients, Wiley & Sons, New York, 1975.
- [42] O. Svelto, Principles of Lasers, Springer, New York, 2010.
- [43] G.K. Woodgate, Elementary Atomic Structure, Clarendon Press, Oxford, 2002.