Symmetry Classification of Scalar *nth* Order Ordinary Differential Equations

Said Waqas Shah¹, F. M. Mahomed² and H. Azad¹

¹Abdus Salam School Of Mathematical Sciences GC University, Lahore. 68-B, New Muslim Town, Lahore 54600, Pakistan Emails: waqas.shah@sms.edu.pk Hasan.Azad@sms.edu.pk

²DSI-NRF Centre of Excellence in Mathematical and Statistical Sciences, School of Computer Science and Applied Mathematics, University of the Witwatersrand, Johannesburg, Wits 2050, South Africa Email: Fazal.Mahomed@wits.ac.za

Abstract: We complete the Lie symmetry classification of scalar *n*th order, $n \ge 4$, ordinary differential equations by means of the symmetry Lie algebras they admit. It is known that there are three types of such equations depending upon the symmetry algebra they possess, viz. first-order equations which admit infinite dimensional Lie algebra of point symmetries, second-order equations possessing the maximum eight point symmetries and higher-order, $n \ge 3$, admitting the maximum n + 4 dimensional symmetry algebra. We show that *n*th order equations for $n \ge 4$ do not admit maximally an n + 3 dimensional Lie algebra except for n = 5 which can admit sl(3, R) algebra. Also, they can possess an n + 2 dimensional Lie algebra that gives rise to a nonlinear equation that is not linearizable via a point transformation. It is shown that for $n \ge 5$ there is only one such class of equations.

Keywords: Lie Classification, Symmetry Algebras, Fundamental Invariants and Lie Determinants.

1 Introduction

Symmetry Lie algebras of scalar *n*th order ordinary differential equations (ODEs) have been extensively studied over several years since the initial ground breaking works of Lie [1, 2]. Equations of order one are equivalent to each other via point transformation. For scalar higher order ODEs, Lie [1] proved that the maximum dimension of the point symmetry Lie algebra for a scalar second order ODE is eight dimensional and occurs for linear and linearizable by point transformation equations. Lie [2] obtained a complex classification of second order ODEs in terms of their Lie algebras. Mahomed and Leach [4] derived the real classification and showed that a second order equation can admit 0, 1, 2, 3 or the maximum 8 dimension real symmetry algebra. The original Lie classification and the classification in the real domain [4] for second order ODEs are, inter alia, compared in [5]. Algebraic linearizability criteria were initiated by Lie [1], who showed that such second order equations possessing a Lie algebra of dimension 2 and of rank one, are linearizable. This falls under Type II and Type IV canonical forms in Lie's classification. The Types I and III cases with focus on linearizability were achieved in [6], [7]. The reader is also referred to the survey article [8]. We furthermore deduce all the canonical forms for *n*th order, $n \ge 4$, equations that admit *n* dimensional symmetry algebras. The n = 2, 3 cases are well-known in the literature.

In the study of scalar linear ODEs of order $n, n \ge 3$, Mahomed and Leach [9] (see also [5], [8] as well as the contribution by Krause and Michel [11]) demonstrated that the point symmetry algebra can be n+1, n+2 or n+4. Thus, for $n \ge 3$, scalar linear ODEs are not necessarily equivalent to each other via point transformation. Moreover, for $n \ge 3$, there exist nonlinear as well as nonlinearizable ODEs with n+2 and n+3 symmetry algebras [9]. It is important to remark that second order ODEs are quite different to higher order equations $n \ge 3$ as per the symmetry algebras they admit. Apart, from the maximum dimension of the point symmetry algebra being 2+6 for second order ODEs and that for higher order, $n \ge 3$, equations n + 4 (see Lie [1, 2]), there are two more notable differences. Secondly, all linear second order ODEs are equivalent to the free particle equation whereas a linear higher order $n \ge 3$ ODE has been determined to have three equivalence classes depending upon whether it has n + 1, n + 2 or the maximum number n + 4 of point symmetries [9]. Thirdly, the complete or full algebra of symmetries of a second order ODE is a subalgebra of its maximum algebra s1(3, R), whereas the full algebra of a higher order $n \ge 3$ ODE is not necessarily a subalgebra of its maximum Lie algebra [9].

The symmetry Lie algebra classification of third order ODEs as well as linearization by point transformation have been investigated in a number of relevant publications (see [9, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21]). Furthermore, integrability and reductions for third order ODEs were looked at in [22].

Scalar fourth order ODEs were considered in recent works from the point of view of Lie classification in terms of four-dimensional algebras, canonical forms as well as integrability (see [23, 24, 25]. A complete Lie classification and algebraic linearization were also attempted in [26]. Linearization criteria, by point transformation, for such ODEs were found in [27].

This work is a continuation of our previous work on the classification of scalar fourth order ODEs [26]. Here we go on to complete the classification of *n*th order ODEs according to the symmetry algebras they admit. We emphasize that we use the same tools and scheme as we have for scalar fourth order ODEs but in general form as the order of the equations considered here are for any n > 4.

The first section deals with notation which we have taken from [26] as well as the established methods used in this paper. In the second and third sections we classify all equations of order n which admit (n + 1)- and (n + 2)-dimensional Lie algebras, respectively. Thereafter we find the (n - 1)- and n-dimensional fundamental invariants

of algebras of dimension n. In the last section we provide some applications and present concluding remarks.

2 Notation and Methods

By (m, n) we denote the type of algebra where m is the type of algebra in [28] and n is the dimension of that algebra. We denote a general vector field or generator as

$$X_i = \xi_i(x, y)\partial_x + \eta_i(x, y)\partial_y, \qquad i = 1, 2, ..., n$$

Here $(x, y) \in \mathbb{R}^2$ and ∂_x denotes $\frac{\partial}{\partial x}$. Note also that *n* is the dimension of a Lie algebra of which X_i s are the generators. Other notation will be mentioned as they arise.

2.1 Classification Scheme

Let L be an m-dimensional Lie subalgebra of vector fields. To find an invariant equation of order n we consider the normal form of an nth order ODE as

$$y^{(n)} = H(x, y, y', y'', \dots, y^{(n-1)}).$$
(2.1)

For a generator X to be a symmetry of (2.1) the following condition must hold:

$$pr_n(X)(y^{(n)} - H)|_{y^{(n)} = H} = 0$$
(2.2)

where pr_n denotes the *n*th prolongation and $|_{y^{(n)}=H}$ denotes the constraint. In this case the resulting *n*th order equation has the most general form of an *n*th order equation invariant under X. If the condition (2.1) holds for every X_i , i = 1, 2, ..., m, then the resulting *n*th order equation is called the invariant equation of L and L is called the symmetry algebra of such an equation. This will be more elaborated in the next sections.

2.2 Fundamental Invariants

Let L be an m-dimensional Lie subalgebra of vector fields defined on a subspace $\mathbb{D} \subset \mathbb{R}^2$. Then the nth order prolonged Lie algebra is defined on a subspace $\mathbb{D}^{(n)} \subset \mathbb{R}^{n+2}$. Suppose that r_0 is the rank of L in \mathbb{D} . Then r_n will be the rank of the prolonged L in $\mathbb{D}^{(n)}$. The rank here means the rank of the matrix whose rows are coefficients of the m generating vector fields of L.

Now let d_n be the number of differential invariants of order n. Then we have

$$d_n = n + 2 - r_n, \quad n \ge 0$$

Example 2.1. Consider $X = \partial_x$. Here n = 0 and $r_o = 1$. Thus $d_0 = 0 + 2 - 1 = 1$, so we have one zeroth order differential invariant which is u = y.

Example 2.2. Consider $X_1 = \partial_x$, $X_2 = \partial_y$. Again n = 0, $r_o = 2$ and $d_0 = 0 + 2 - 2 = 0$, and there is no zeroth order invariant. Now for the first prolongations of X_1 and X_2 , n = 1, $r_1 = 2$ and $d_1 = 1 + 2 - 2 = 1$. Thus there is a first order differential invariant u = y'.

A very important work for finding the fundamental differential invariants for n-th order equations is done by Neseterenko in [15].

2.3 Invariant Differentiation Operator

If u, v are invariants then by Lie's theorem, $\frac{D_x v}{D_x u}$ is also an invariant. This process is also called invariant differentiation [29]. Recall that D_x is the total differentiation operator. We can write this as

$$\frac{D_x v}{D_x u} = (D_x u)^{-1} D_x v$$
$$= \lambda D_x v$$

We call $\lambda D_x = D$ the invariant differentiation operator once we know λ .

Suppose we have an unknown $\lambda(x, y, y', ..., y^{(n)})$. We require $\lambda D_x v$ to be invariant, and therefore we need

$$pr_n(X)(\lambda D_x v) = 0 \tag{2.3}$$

For the time being we ignore the prolongation symbol and consider $pr_n(X)$ simply as X. We can write

$$X = \overline{X} + \xi D_x$$

where $\overline{X} = w\partial_y + D_x w\partial_{y'} + D_x^2 w\partial_{y''} + \dots + D_x^n w\partial_{y^{(n)}}$ is called the canonical operator and $w = \eta - \xi D_x$. The equation (2.3) is then written as

$$(\overline{X} + \xi D_x)(\lambda D_x v) = 0$$

$$\Rightarrow \xi D_x(\lambda D_x v) + \overline{X}(\lambda) D_x v + \lambda \overline{X}(D_x v) = 0$$

$$\Rightarrow \xi D_x \lambda D_x v + \xi \lambda D_x^2 v + (X\lambda - \xi D_x \lambda) D_x v + \lambda \overline{X}(D_x v) = 0$$

$$\Rightarrow \xi D_x \lambda D_x v + \lambda \xi D_x^2 v + X(\lambda) D_x v - \xi D_x \lambda D_x v + \lambda \overline{X}(D_x v) = 0$$

$$\Rightarrow \xi \lambda D_x^2 v + X(\lambda) D_x v + \lambda D_x (Xv - \xi D_x v) = 0$$

$$\Rightarrow (X(\lambda) - \lambda D_x \xi) D_x v = 0$$

This gives:

$$X(\lambda) = \lambda D_x \xi. \tag{2.4}$$

Hence, λ satisfies the non-homogenous linear PDE (2.4). One only requires one nontrivial solution for λ which can be a constant as well.

Example 2.3. Let $X_1 = \partial_x$ and $X_2 = \partial_y$. We know that u = y' is a first order differential invariant of X_1 and X_2 . Applying the condition (2.4) we have

$$X_1\lambda = 0$$
, $X_2\lambda = 0$

This clearly shows that we can set $\lambda = 1$ and therefore the invariant differentiation operator can be taken as $D = (1)D_x = D_x$. Thus

$$Dy' = D_x y' = y''$$

is a second order differential invariant.

2.4 Lie Determinants

Definition 2.1. Consider an m-dimensional Lie subalgebra of vector fields whose generators are given as

$$X_k = \xi_k \partial_x + \eta_k \partial_y, \ k = 1, 2, ..., m.$$

Then the determinant of the following matrix

$$\begin{pmatrix} \xi_1 & \eta_1 & \eta_1^{[1]} & \eta_1^{[2]} & \cdots & \eta_1^{[m-2]} \\ \xi_2 & \eta_2 & \eta_2^{[1]} & \eta_2^{[2]} & \cdots & \eta_2^{[m-2]} \\ \vdots & \vdots & \ddots & \vdots & & \\ \xi_m & \eta_m & \eta_m^{[1]} & \eta_m^{[2]} & \cdots & \eta_m^{[m-2]} \end{pmatrix}$$

is called the Lie determinant corresponds to the m-dimensional Lie algebra for $m \geq 2$. We denote the Lie determinant by Λ_L .

Lie proved that for an *m*-dimensional Lie subalgebra of vector fields L, the Lie determinant gives rise to all the invariant equations of order $\leq m - 2$ [3]. Similarly it can be noticed that the rank of the prolonged algebra is m, i.e. maximal unless the Lie determinant vanishes in which case the rank of L diminishes. Here the algebra is prolonged up to order m - 2. These invariant equations are called the singular invariant equations of L. The fundamental differential invariants which are not singular must be of order m - 1 and m and the higher order differential invariants can be then be found from the fundamental invariants.

Note that here, once we find the (m-1)th order differential invariant say u and the invariant differentiation operator $D = \lambda D_x$, we can determine the *m*th order differential invariant as it is none other than Du.

3 (n+1) Dimensional Algebras

3.1 Nonlinear Equations

It is easy to observe that (20, n + 1) is not admitted by an equation as there can only be n solution symmetries. The linearizable case (21, n + 1) is studied in the next subsection 3.2. Also, (22, n + 1) is not possessed as a maximal Lie algebra. We consider the other n + 1 dimensional algebras.

 $(24, n+1), r = n-2, n \ge 3: X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x + \alpha y\partial_y, X_4 = x\partial_y, X_5 = x^2\partial_y, \dots, X_{n+1} = x^{n-2}\partial_y.$

The generators, except X_3 , imply that a general *n*th order equation of the type (2.1) admitting these generators must be of the form:

$$y^{(n)} = H(y^{(n-1)}). ag{3.1}$$

The *n*th prolongation of X_3 is: $x\partial_x + \alpha y\partial_y + (\alpha - 1)y'\partial_{y'} + \dots + (\alpha - n)y^{(n)}\partial_{y^{(n)}}$.

Applying this to the equation (3.1) and solving we find the general form of an *n*th order equation admitting this algebra to be

$$y^{(n)} = K(y^{(n-1)})^{\frac{\alpha-n}{\alpha-n+1}}, \ K \neq 0,$$
(3.2)

where $K \neq 0$ is an arbitrary constant and $\alpha \neq n-1$. For K = 0 or $\alpha = n-1$, it is easy to see that the general form of such an equation admitting this algebra must be $y^{(n)} = 0$, though we know that such an equation admits the maximal (n+4)-dimensional algebra of which such an algebra is the subalgebra.

 $(25, n+1), r = n - 1, n \ge 2: X_1 = \partial_x, X_2 = \partial_y, X_3 = x \partial_y, X_4 = x^2 \partial_y, ..., X_n = x^{n-2} \partial_y, X_{n+1} = x \partial_x + (ry + x^r) \partial_y.$

The *n*th prolongation of X_{n+1} is: $x\partial_x + (ry + x^r)\partial_y + (rx^{r-1} + (r-1)y')\partial_{y'} + \dots + ((r(r-1)(r-n+1))x^{r-n} + (r-n)y^{(n)})\partial_{y^{(n)}}$.

The equation is:

$$y^{(n)} = K e^{\frac{-y^{(n-1)}}{(n-1)!}} K \neq 0.$$
(3.3)

 $(26, n+1), r = n-3, n \ge 4$: $X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_y, X_5 = x\partial_y, \dots, X_{n+1} = x^{n-3}\partial_y.$

Here the equation turns out to be

$$y^{(n)} = K \frac{(y^{(n-1)})^2}{y^{(n-2)}}, \ K \neq 0, n/(n-1).$$
 (3.4)

If K = n/(n-1), then there is one more symmetry $X = x^2 \partial x + (n-3)xy \partial y$ as discussed in Section 4.

$$(27, n+1), r = n-3, n \ge 4: X_1 = \partial_x, X_2 = \partial_y, X_3 = 2x\partial_x + ry\partial_y, X_4 = x^2\partial_x + rxy\partial_y, X_5 = x\partial_y, \dots, X_{n+1} = x^{n-3}\partial_y.$$

The *n*th prolongation of X_3 and X_4 are: $X_3 + \sum_{k=1}^n (r-2k)y^{(k)}\partial_{y^{(k)}}$ and $X_4 + \sum_{k=1}^n (k(r-k+1)y^{(k-1)} + x(r-2k)y^{(k)})\partial_{y^{(k)}}$, respectively.

The equation is

$$y^{(n)} = \frac{n}{n-1} \frac{(y^{(n-1)})^2}{y^{(n-2)}} + K(y^{(n-2)})^{\frac{n+3}{n-1}}, \ K \neq 0.$$
(3.5)

If K = 0, then there is one more symmetry $X = y \partial y$ as pursued in Section 4.

 $(28, n+1), r = n-4, n \ge 5: X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_y, X_5 = x^2\partial_x + (n-4)xy\partial_y, X_6 = x\partial_y, \dots, X_{n+1} = x^{n-4}\partial_y.$

The nth order invariant equation is:

$$y^{(n)} = (y^{(n-2)})^3 (y^{(n-3)})^{-2} \left[\frac{n(3(n-2)K_1 - 2n + 2)}{(n-2)^2} + K((n-2)K_1 - (n-1))^{\frac{3}{2}} \right], \quad (3.6)$$

where $K = e^{(n-3)e^{(n-1)}(e^{(n-2)})^{-2}}$

where $K_1 = y^{(n-3)}y^{(n-1)}(y^{(n-2)})^{-2}$.

3.2 Linearizable Equations for Higher Order

Here we consider linearization for higher order $n \ge 3$ equations. We demonstrate how one obtains the linear form. The reader is also referred to [9].

$$(21, n+1), r = n - 1: X_1 = \partial_y, X_2 = y \partial_y, X_3 = \xi_1(x) \partial_y, \dots, X_{n+1} = \xi_r(x) \partial_y.$$

By introducing coordinates: $\overline{x} = \xi_1(x)$, $\overline{y} = y$ and ignoring the bars, the generators of this algebra can be transformed to $X_1 = \partial_y$, $X_2 = y\partial_y$, $X_3 = x\partial_y$, $X_4 = \xi_2(x)\partial_y$,..., $X_{n+1} = \xi_r(x)\partial_y$.

Consider the general form of an *n*th order ODE as (2.1). For such an equation to admit X_1 and X_3 , H must be independent of y and y'. Invoking X_4 on this equation, the general form of H can be determined as

$$H = \frac{\xi_2^{(n)}}{\xi_2^{(n-1)}} y^{(n-1)} + H_1(x, \phi_1, \phi_2, ..., \phi_{n-3})$$
(3.7)

where

$$\phi_k = y^{(k+2)} - \frac{\xi_2^{(k+2)}}{\xi_2^{(k+1)}} y^{(k+1)}, \qquad k = 1, 2, ..., n-3.$$

Now imposing X_5 on this, the general form of H_1 is found to be

$$H_1 = \frac{\phi^0}{\phi^1} \phi_{n-3} + H_2(x, \psi^1, \psi^2, ..., \psi^{n-4})$$
(3.8)

where

$$\phi^k = \xi_3^{(n-k)} - \xi_3^{(n-k-1)} \frac{\xi_2^{(n-k)}}{\xi_2^{(n-k-1)}}, \qquad k = 0, 1, 2, ..., n-3.$$

and

$$\psi^k = \phi_{k+1} - \frac{\phi^{n-k-3}}{\phi^{n-k-2}}\phi_k, \qquad k = 1, 2, 3, ..., n-4$$

Similarly, for an equation to be invariant under each X_i , for $i \ge 5$, the functions H_{i-2} s that are arbitrary functions of their n-i-1 arguments, can be determined. As a result, we can write the general form of an *n*th order ODE invariant under X_1, X_3, \dots, X_{n+1} as

$$y^{(n)} = \sum_{i=2}^{n-1} A_i(x) y^{(i)} + H_{n-2}(x).$$
(3.9)

The invariance under X_2 now easily implies that H_{n-2} must vanish giving us the following homogenous equation whose symmetry algebra is (21, n+1):

$$y^{(n)} = \sum_{i=2}^{n-1} A_i(x) y^{(i)}.$$
(3.10)

One notices that since this equation is invariant under $X_1, X_3,..., X_{n+1}$, each ξ_i for i = 2, 3, ..., n - 1 must be an independent solution of equation (3.10). In general this proves the following proposition.

Proposition 3.1. (21, n+1) is a symmetry algebra of the nth order linear homogenous equation

$$y^{(n)} = \sum_{i=2}^{n-1} A_i(x) y^{(i)},$$

such that each ξ_i for i = 1, 2, ..., n-1 together with the number 1 form a set of independent solutions of this equation. Moreover, the functions A_i s are determined by solving the system of homogenous equations

$$\xi_k^{(n)} = \sum_{i=1}^{n-1} A_i(x)\xi_k^{(i)}, \qquad k = 1, 2, ..., n-1$$

4 (n+2)-Dimensional Algebras

4.1 Nonlinear Equations

For the n + 2 dimensional algebras, (20, n + 2), (21, n + 2) and (22, n + 2) are clearly not admissible algebras. The linearization case (23, n + 2) is looked at in 4.2.

$$(24, n+2), r = n - 1, n \ge 2: X_1 = \partial_x, X_2 = \partial_y, X_3 = x \partial_x + \alpha y \partial_y, X_4 = x \partial_y, X_5 = x^2 \partial_y, ..., X_{n+2} = x^{n-1} \partial_y. \Lambda_L = 1 \cdot 2! \cdot 3! ... (n-1)! (\alpha - n) y^{(n)}.$$

$$(4.1)$$

 $(25, n+2), r = n, n \ge 1: X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_y, X_4 = x^2\partial_y, ..., X_{n+1} = x^{n-1}\partial_y, X_{n+2} = x\partial_x + (ry + x^r)\partial_y.$

$$\Lambda_L = 1 \cdot 2! \cdot 3! \dots (n-2)! \cdot (n-1)! \cdot n!.$$
(4.2)

$$(26, n+2), r = n-2, n \ge 3: X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_y, X_5 = x\partial_y, ..., X_{n+2} = x^{n-2}\partial_y.$$

$$\Lambda_L = 1 \cdot 2! \cdot 3! ... (n-3)! \cdot (n-2)! y^{(n-1)} y^{(n)}.$$
(4.3)

 $(27, n+2), r = n-2, n \ge 3: X_1 = \partial_x, X_2 = \partial_y, X_3 = 2x\partial_x + ry\partial_y, X_4 = x^2\partial_x + rxy\partial_y, X_5 = x\partial_y, \dots, X_{n+2} = x^{n-2}\partial_y.$ $\Lambda_L = 1 \cdot 2! \cdot 3! \dots (n-2)! \cdot n^2 (y^{(n-1)})^2. \tag{4.4}$

$$(28, n+2), r = n-3, n \ge 4: X_1 = \partial_x, X_2 = x\partial_x, X_3 = y\partial_y, X_4 = x^2\partial_x + rxy\partial_y, X_5 = \partial_y, X_6 = x\partial_y, \dots, X_{n+2} = x^{n-3}\partial_y.$$

Equation:

$$y^{(n)} = \frac{n}{n-1} \frac{(y^{(n-1)})^2}{y^{(n-2)}}.$$
(4.5)

Here we have only one case, the last, which constitute an invariant equation with n + 2 dimensional symmetry algebra. The rest do not form an equation or are singular equations with maximal algebra.

Next we need to again deal with linearization.

4.2 Linearizable Higher Order Equations

Here we obtain the form for the reduced linear equation that results from linearizability (see also [9]).

$$(23, n+2), r = n$$
: $X_1 = \eta_1(x)\partial_y, X_2 = \eta_2(x)\partial_y, ..., X_n = \eta_n(x)\partial_y, X_{n+1} = y\partial_y, X_{n+2} = \partial_x.$

From Proposition 3.1, it is easy to show that the general form of an *n*th order equation invariant under the first n + 1 generators is the following linear homogenous equation:

$$y^{(n)} = \sum_{i=0}^{n-1} A_i(x) y^{(i)}.$$
(4.6)

For such an equation to admit X_{n+2} the coefficients A_i s must be constant for each i = 0, 1, 2, ..., n - 1. Note also that η_i s for i = 1, 2, ..., n form an independent set of solutions for this constant coefficient equation.

We can determine the A_i s as in Proposition 3.1 for known $\eta_i(x)$ s. However, note that these coefficients can also be determined by the roots of the corresponding characteristic polynomial equation.

We first consider the case when all the roots are real and distinct say, α_i , i = 1, 2, 3, ..., n.

The fundamental solutions of equation (4.6) are:

$$\eta_i(x) = e^{\alpha_i x}, \quad \text{for } i = 1, 2, 3, ..., n.$$

The constants A_i s which depend on α_i s can then be derived by solving the system

$$VX = B, (4.7)$$

where V is an $n \times n$ Vandermonde matrix in $(\alpha_1, \alpha_2, ..., \alpha_n)$,

$$X = [A_0, A_1, ..., A_{n-1}]^T$$
 and $B = [\alpha_1^n, \alpha_2^n, ..., \alpha_n^n]^T$.

The system given in (4.7) can be solved by Crammer's rule, i.e. we have by Crammer's rule,

$$A_i(\alpha_1, \alpha_2, ..., \alpha_n) = \frac{\det(f_i)}{\det(V)}, \qquad i = 0, 1, 2, ..., n - 1,$$

where f_i is the matrix obtained by replacing the (i + 1)th column in V by B. First, we state the following lemma whose proof is elementary and thus omitted.

Lemma 4.1. Let V be a Vandermonde matrix in $(\alpha_1, \alpha_2, ..., \alpha_n)$, then the determinant of V is

$$\det(V) = \prod_{1 \le i < j \le n} (\alpha_i - \alpha_j).$$

As an example, we find A_{n-1} . For this we need to find the determinant of f_n . We prove by induction on n that

$$A_{n-1}(\alpha_1, \alpha_2, \cdots, \alpha_n) = \sum_{k=1}^n \alpha_k.$$

Actually, we show that

$$\det(f_n) = A_{n-1} \det(V).$$

For this we write the matrix f_n as below:

$$f_n = \begin{pmatrix} 1 & \alpha_1 & \alpha_1^2 & \cdots & \alpha_1^{n-2} & \alpha_1^n \\ 1 & \alpha_2 & \alpha_2^2 & \cdots & \alpha_2^{n-2} & \alpha_2^n \\ \vdots & \vdots & \ddots & \vdots & & \\ 1 & \alpha_n & \alpha_n^2 & \cdots & \alpha_n^{n-2} & \alpha_n^n \end{pmatrix}$$

To find the determinant of this matrix we apply row operations to make all the entries in the first column zero except the first one.

 \Rightarrow

$$\det(f_n) = \begin{vmatrix} 1 & \alpha_1 & \alpha_1^2 & \cdots & \alpha_1^{n-2} & \alpha_1^n \\ 0 & \alpha_1 - \alpha_2 & \alpha_1^2 - \alpha_2^2 & \cdots & \alpha_1^{n-2} - \alpha_2^{n-2} & \alpha_1^n - \alpha_2^n \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \alpha_1 - \alpha_n & \alpha_1^2 - \alpha_n^2 & \cdots & \alpha_1^{n-2} - \alpha_n^{n-2} & \alpha_1^n - \alpha_n^n \end{vmatrix},$$

where by the vertical bars we denote the determinant of the matrix. \Rightarrow

$$\det(f_n) = \begin{vmatrix} \alpha_1 - \alpha_2 & \alpha_1^2 - \alpha_2^2 & \cdots & \alpha_1^{n-2} - \alpha_2^{n-2} & \alpha_1^n - \alpha_2^n \\ \alpha_1 - \alpha_3 & \alpha_1^2 - \alpha_3^2 & \cdots & \alpha_1^{n-2} - \alpha_3^{n-2} & \alpha_1^n - \alpha_3^n \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1 - \alpha_n & \alpha_1^2 - \alpha_n^2 & \cdots & \alpha_1^{n-2} - \alpha_n^{n-2} & \alpha_1^n - \alpha_n^n \end{vmatrix}$$

 \Rightarrow

 \Rightarrow

$$\prod_{k=2}^{n} (\alpha_1 - \alpha_k) \begin{vmatrix} 1 & \alpha_1 + \alpha_2 & \alpha_1(1) + \alpha_2^2 & \cdots & \alpha_1(1) + \alpha_2^{n-3} & \alpha_1(\alpha_1(1) + \alpha_2^{n-2}) + \alpha_2^{n-1} \\ 1 & \alpha_1 + \alpha_3 & \alpha_1(1) + \alpha_3^2 & \cdots & \alpha_1(1) + \alpha_3^{n-3} & \alpha_1(\alpha_1(1) + \alpha_2^{n-2}) + \alpha_3^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \alpha_1 + \alpha_n & \alpha_1(1) + \alpha_n^2 & \cdots & \alpha_1(1) + \alpha_n^{n-3} & \alpha_1(\alpha_1(1) + \alpha_n^{n-2}) + \alpha_n^{n-1} \end{vmatrix}$$

where in the brackets we have the corresponding entries of the previous columns in each successive step.

$$\det(f_n) = \prod_{k=2}^n (\alpha_1 - \alpha_k) \begin{pmatrix} 1 & \alpha_2 & \alpha_2^2 & \cdots & \alpha_2^{n-3} & \alpha_1(\alpha_2^{n-2}) + \alpha_2^{n-1} \\ 1 & \alpha_3 & \alpha_3^2 & \cdots & \alpha_3^{n-3} & \alpha_1(\alpha_2^{n-2}) + \alpha_3^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \alpha_n & \alpha_n^2 & \cdots & \alpha_n^{n-3} & \alpha_1(\alpha_n^{n-2}) + \alpha_n^{n-1} \end{pmatrix}$$
$$\det(f_n) = \prod_{k=2}^n (\alpha_1 - \alpha_k) \left(\alpha_1 \begin{vmatrix} 1 & \alpha_2 & \alpha_2^2 & \cdots & \alpha_n^{n-3} & \alpha_1^{n-2} \\ 1 & \alpha_3 & \alpha_3^2 & \cdots & \alpha_n^{n-3} & \alpha_2^{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \alpha_n & \alpha_n^2 & \cdots & \alpha_n^{n-3} & \alpha_n^{n-2} \end{vmatrix} + \begin{pmatrix} 1 & \alpha_2 & \alpha_2^2 & \cdots & \alpha_n^{n-3} & \alpha_2^{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \alpha_3 & \alpha_3^2 & \cdots & \alpha_3^{n-3} & \alpha_3^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \alpha_n & \alpha_n^2 & \cdots & \alpha_n^{n-3} & \alpha_n^{n-1} \end{pmatrix} \right)$$

The first matrix is the Vandermonde matrix in $(\alpha_2, \alpha_2, \dots, \alpha_n)$ and the second matrix is f_n in $(\alpha_2, \alpha_2, \dots, \alpha_n)$. By induction we now have

$$\det(f_n) = \prod_{k=1}^n (\alpha_1 - \alpha_k) \Big(\alpha_1 (\det(V(\alpha_2, \alpha_3, \cdots, \alpha_n))) + (\sum_{k=2}^n \alpha_k) (\det(V(\alpha_2, \alpha_3, \cdots, \alpha_n))) \Big)$$

$$\Rightarrow$$

$$\det(f_n) = \sum_{k=1}^n \alpha_k (\det(V(\alpha_1, \alpha_2, \cdots \alpha_n))),$$

which proves the assertion.

In general we have the following lemma:

Lemma 4.2. Let

 \Rightarrow

$$A_i = \frac{\det(f_i)}{\det(V)},$$

where V and f_i are defined as in the above discussion. Then A_i can be determined as

$$A_i = (-1)^i \sum_{1 \le k_1 < k_2 < \cdots < k_{i+1} \le n} \alpha_{k_1} \alpha_{k_2} \cdots \alpha_{k_{i+1}}.$$

The proof of this can be carried out by using induction on n and i simultaneously and therefore is omitted. The A_{n-1} above was derived by using the technique of the proof of this lemma.

Now we consider the case when 2k number of the roots are complex and the rest real, i.e. n-2k of them. If we show the complex roots as $\alpha_j = a_j + ib_j$ and $\overline{\alpha_j} = a_j - ib_j$ for $j = 1, 2, \dots k$ and the real roots as $\alpha_{2k+1}, \alpha_{2k+2} \dots \alpha_n$. Then the fundamental set of solutions of equation (4.6) with constant coefficients can be taken as $\eta_j(x) = e^{a_j(x)} \cos(b_j x)$ for $j = 1, 2, \dots k, \eta_j(x) = e^{a_j x} \sin(b_j x)$ for $j = k + 1, k + 2, \dots 2k$ and the rest as $e^{\alpha_j(x)}$ s which correspond to the real roots. The constants A_i s can be derived in exactly the same way. Thus, we actually prove the following proposition.

Proposition 4.1. The generators given in (23, n+2) form a Lie algebras if and only if the $\eta_i s$ form a fundamental set of solutions for the constant coefficient equation given as

$$y^{(n)} = \sum_{i=0}^{n-1} A_i y^{(i)}.$$
(4.8)

Moreover, the constants can be determined as in lemma 4.2.

5 Fundamental Invariants

In this section we now classify all the fundamental invariants for five and higher dimensions. Note that the algebras (20, n) and (21, n) are not admissible.

(5,5): The generators of this algebra are: $X_1 = \partial_x$, $X_2 = \partial_y$, $X_3 = x\partial_x - y\partial_y$, $X_4 = y\partial_x$, $X_5 = x\partial_y$.

This is a five-dimensional algebra, hence r = 5. We find the 3rd prolongations of the generators of this algebra. Then the Lie determinant, which is the determinant of the matrix

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & x & 1 & 0 & 0 \\ x & -y & -2y' & -3y'' & -4y''' \\ y & 0 & -y'^2 & -3y'y'' & -(3y''^2 + 4y'y'''), \end{pmatrix}$$

is then

$$\Lambda_L(M) = 9y''^2.$$

This shows that y'' = 0 is the only equation of order ≤ 3 invariant under this algebra.

The fundamental differential invariants can then be found in the usual way. For the given algebra we already know a fourth order invariant which is

$$\phi_1 = y^{(4)} y''^{\frac{-5}{3}} - \frac{5}{3} y'''^2 y''^{-8/3}$$

To find the 5th order differential invariant we can solve the system of linear partial differential equations

$$pr_5(X_i)(y^{(5)} - H)|_{eq} = 0, \quad \text{for } i = 1, 2, \cdots, 5,$$
 (5.1)

where $pr_n(X)$ is the *n*th prolongation of X.

Solving this system we derive the 5th order equation invariant under this algebra to be

$$y^{(5)} = \frac{-40}{9} y^{\prime\prime\prime3} y^{\prime\prime-2} + 5y^{\prime\prime\prime} y^{(4)} y^{\prime\prime-1} + y^{\prime\prime2} H(\phi_1),$$
(5.2)

where ϕ_1 is the fourth order invariant of this algebra and H is an arbitrary function of its argument. We can thus write the fifth order differential invariant to be

$$\phi_2 = (y''^2 y^{(5)} + \frac{40}{9} y'''^3 - 5y'' y''' y^{(4)}) / y''^4.$$

Here ϕ_1 and ϕ_2 are the fundamental differential invariants for this algebra. All higher order differential invariants can be deduced by Lie's invariant differentiation: $\phi_{n+1} = D_x(\phi_n)/D_x(\phi_{n-1})$ or by using invariant derivative operator $\mathcal{D} = y''^{-1/3}D_x$ so that higher order equations are given by

$$\mathcal{D}^{n-4}\phi_1 = H(\phi_1, \dots, \mathcal{D}^{n-5}\phi_1), \ n \ge 5,$$
 (5.3)

where D_x is the total derivative operator.

(15,5): $X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_y, X_5 = x^2\partial_x.$

The fundamental invariants are

$$\phi_1 = \frac{y'^2 y^{(4)} - 6(y'y''y''' - y''^3)}{(2y'y''' - 3y''^2)^{\frac{3}{2}}},$$

$$\phi_2 = \frac{3y'^3 y^{(5)} - 10y'^2 y''y^{(4)} - 50y'^2 y'''^2 + 240y'y''^2 y''' - 180y''^4}{(2y'y''' - 3y''^2)^2}$$

with the 5th order equation as $\phi_2 = H(\phi_1)$ and with $\mathcal{D} = y'(2y'y''' - 3y''^2)^{-1/2}D_x$ we can invoke (5.3) for higher order equations.

 $(6,6): X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_x, X_5 = x\partial_y, X_6 = y\partial_y.$

The generators X_1, X_2, X_5 and X_6 result in the invariants

$$K_1 = 3y''^{-1}y^{(4)} - 5y'''^2y''^{-2},$$

$$K_2 = 9y''^{-1}y^{(5)} - 45y''^{-2}y'''y^{(4)} + 40y''^{-3}y'''^3.$$

Writing X_4 in terms of K_1 and K_2 we find

$$X_4 = 2K_1\partial_{K_1} + 3K_2\partial_{K_2}$$

which provides the 5th order invariant of (6, 6)

$$= \frac{\phi_1 = K_1^{-3/2} K_2,}{(3y''y^{(4)} - 5y'''y^{'''})^{\frac{3}{2}}}.$$

The invariant differentiation operator is $\mathcal{D} = K_1^{-1/2} D_x$ and hence one can derive a sixth order invariant

$$\phi_2 = \frac{(9y''^3y^{(6)} - 63y''^2y'''y^{(5)} + 105y''y'''^2y^{(4)} - 35y'''^4)}{(3y''y^{(4)} - 5y'''^2)^2}.$$

The 5th order invariant equation is $\phi_1 = C$, C a constant and with $\mathcal{D} = y''(3y''y^{(4)} - 5y'''^2)^{-1/2}D_x$, we have

$$\mathcal{D}^{n-5}\phi_1 = H(\phi_1, \dots, \mathcal{D}^{n-6}\phi_1), \ n \ge 6,$$
(5.4)

for higher order equations. Here the 6th order equation is $\phi_2 = H(\phi_1)$ with ϕ_2 as above and higher order equations are deduced by invariant differentiation as stated.

Note that $K_1 = 0$ is the fourth order singular invariant equation with this algebra.

 $(16,6): X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_y, X_5 = x^2\partial_x, X_6 = y^2\partial_y.$

Here the invariants, using X_1 , X_2 , X_4 and X_6 , are

$$K_1 = y'''y^{-1} - \frac{3}{2}y'^{-2}y''^2,$$

$$K_2 = y'^{-1}y^{(4)} + 3y'^{-3}y''^3 - 4y'^{-2}y''y'''.$$

$$K_3 = y'^{-1}y^{(5)} + 5y'^{-3}y''^2y''' - 5y'^{-2}y''y^{(4)}.$$

Utilising X_5 in (K_1, K_2, K_3) space, we end up with

$$X_5 = 2xK_1\partial_{K_1} + (3xK_2 + 2K_1)\partial_{K_2} + (4xK_3 + 5K_2)\partial_{K_3}$$

Hence, a 5th order invariant transpires as

$$\phi_1 = K_1^{-3} (5K_2^2 - 4K_1K_3).$$

The invariant differentiation operator is $\mathcal{D} = K_1^{-1/2} D_x$ and higher order invariant equations are as in (5.4).

Here $K_1 = 0$ is the third order singular invariant equation admitting this algebra.

 $(7,6): X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x + y\partial_y, X_4 = y\partial_x - x\partial_y, X_5 = (x^2 - y^2)\partial_x + 2xy\partial_y, X_6 = 2xy\partial_x + (y^2 - x^2)\partial_y.$

The operators X_1 to X_4 gives rise to the invariants

$$K_{1} = (1 + y'^{2})y''^{-2}y''' - 3y',$$

$$K_{2} = (1 + y'^{2})^{2}y''^{-3}y^{(4)} + 15y'^{2} - 10y'y''^{-2}y'''(1 + y'^{2}),$$

$$K_{3} = (1 + y'^{2})^{3}y''^{-4}y^{(5)} - 15(1 + y'^{2})^{2}y'y''^{-3}y^{(4)} - 105y'^{3} + 105(1 + y'^{2})y'^{2}y''^{-2}y''' - 10(1 + y'^{2})^{2}y'y''^{-4}y'''^{2}.$$

Now X_5 in (K_1, K_2, K_3) space yields the two equations

$$-2K_1\partial_{K_2}F + (15 - 5K_2)\partial_{K_3}F = 0,$$

$$-2K_1\partial_{K_1}F + (9 - 3K_2)\partial_{K_2}F + (40K_1 - 4K_3)\partial_{K_3}F = 0.$$

The invariant deduced is of order five (which is admitted by X_6 as well since $[X_4, X_5] = X_6$):

$$\phi_1 = 30K_1^3K_2 - 5K_1^3K_2^2 + 4K_1^4K_3 + 45K_1^3 + 16K_1^5.$$

The operator of invariant differentiation is $\mathcal{D} = (1 + y'^2)y''^{-1}K_1^{-1/2}D_x$ and higher order equations are given by (5.4).

Here $K_1 = 0$ is the third order singular invariant equation.

 $(8,8): X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_y, X_4 = y\partial_y, X_5 = y\partial_x, X_6 = x\partial_x, X_7 = x^2\partial_x + xy\partial_y, X_8 = xy\partial_x + y^2\partial_y.$

The seventh order invariant is

$$\phi = K_2^{-5/3} D_x^2 K_2 - \frac{7}{6} K_2^{-8/3} (D_x K_2)^2 - y''^{-1} y''' K_2^{-5/3} D_x K_2$$
$$-y''^{-2} y'''^2 K_2^{2/3} - \frac{3}{2} y''^{-1} y^{(4)} K_2^{-2/3},$$

where $K_2 = 9y''^{-1}y^{(5)} - 45y''^{-2}y'''y^{(4)} + 40y''^{-3}y'''^{3}$ is the invariant of algebra (6, 6). The operator of invariant differentiation is $\mathcal{D} = K_2^{-1/3}D_x$ and higher order equations are obtained as

$$\mathcal{D}^{n-7}\phi = H(\phi, \dots, \mathcal{D}^{n-8}\phi), \ n \ge 8.$$
(5.5)

For this algebra the singular invariant equation is $K_2 = 0$ which is fifth order.

$$(22, n): r = n - 1, n \ge 2: X_1 = \partial_x, X_2 = \eta_1(x)\partial_y, \dots, X_n = \eta_{n-1}(x)\partial_y.$$

The nth order equation is

$$y^{(n)} + a_1 y^{(n-1)} + \dots + a_{n-1} y' = H(y^{(n-1)} + a_1 y^{(n-2)} + \dots + a_{n-1} y),$$

where the a_i s are constants and the η_i s satisfy the linear (n-1)th order equation

$$E_i \equiv \eta_i^{(n-1)} + a_1 \eta_i^{(n-2)} + \dots + a_{n-1} \eta_i = 0, i = 1, \dots, n-1$$

and its derived equation $D_x E_i = 0$.

$$(23, n): r = n - 2, n \ge 3: X_1 = \partial_x, X_2 = y \partial_y, X_3 = \eta_1(x) \partial_y, \dots, X_n = \eta_{n-2}(x) \partial_y.$$

The nth order equation that admits this algebra is

$$D_x^2 \ln |y^{(n-2)} + a_1 y^{(n-3)} + \dots + a_{n-2} y| = H(D_x \ln |y^{(n-2)} + a_1 y^{(n-3)} + \dots + a_{n-2} y|),$$

where the a_i s are constants and the η_i s satisfy the linear (n-1)th order equation

$$\eta_i^{(n-2)} + a_1 \eta_i^{(n-3)} + \dots + a_{n-2} \eta_i = 0, i = 1, \dots, n-2$$

 $(24,n): r = n - 3, n \ge 4: X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x + \alpha y\partial_y, X_4 = x\partial_y, \dots, X_n = x^{n-3}\partial_y.$

Fundamental Invariants, if $\alpha \neq n-1$, are

$$\phi_1 = (y^{(n-1)})^{\alpha-n+2} (y^{(n-2)})^{-(\alpha-n+1)}, \qquad \phi_2 = y^{(n)} (y^{(n-1)})^{-(\frac{\alpha-n}{\alpha-n+1})},$$

and for $\alpha = n - 1$

$$\phi_1 = y^{(n-1)}, \qquad \phi_2 = y^{(n)}y^{(n-2)}.$$

Thus the invariant equation for both cases is $\phi_2 = H(\phi_1)$.

If $\alpha = n - 1$, the invariant equation is

$$y^{(n)} = (y^{(n-2)})^{-1}H(y^{(n-1)}).$$

 $(25, n): r = n - 2, n \ge 3: X_1 = \partial_x, X_2 = \partial_y, X_3 = x \partial_y, X_4 = x^2 \partial_y, \cdots, X_{n-1} = x^{n-3} \partial_y, X_n = x \partial_x + ((n-2)y + x^{(n-2)}\partial_y.$

The fundamental differential invariants are

$$\phi_1 = y^{(n-1)} e^{\frac{y^{(n-2)}}{(n-2)!}}, \quad \phi_2 = y^{(n)} e^{\frac{2y^{(n-2)}}{(n-2)!}}$$

and invariant equation as before $\phi_2 = H(\phi_1)$.

 $(26, n), r = n - 4, n \ge 5: X_1 = \partial_x, X_2 = \partial_y, X_3 = x \partial_x, X_4 = y \partial_y, X_5 = x \partial_y, \cdots, X_n = x^{n-4} \partial_y.$

The fundamental differential invariants:

$$\phi_1 = \frac{y^{(n-3)}y^{(n-1)}}{(y^{(n-2)})^2}, \quad \phi_2 = \frac{(y^{(n-3)})^2 y^{(n)}}{(y^{(n-2)})^3}.$$

 $(27, n), r = n - 4, n \ge 5; X_1 = \partial_x, X_2 = \partial_y, X_3 = 2x\partial_x + y\partial_y, X_4 = x^2\partial_x + xy\partial_y, X_5 = x\partial_y, \cdots, X_n = x^{n-4}\partial_y.$

The fundamental invariants:

$$\phi_1 = y^{(n-1)} (y^{(n-3)})^{\frac{n+2}{2-n}} - \frac{n-1}{n-2} (y^{(n-2)})^2 (y^{(n-3)})^{\frac{-2n}{n-2}}$$

$$\phi_2 = y^{(n)} (y^{(n-3)})^{\frac{n+4}{2-n}} + \frac{2n(n-1)}{(n-2)^2} (y^{(n-2)})^3 (y^{(n-3)})^{\frac{3n}{2-n}} + \frac{3n}{2-n} y^{(n-1)} y^{(n-2)} (y^{(n-3)})^{\frac{2(n+1)}{2-n}}.$$

 $(28,n): r = n - 5, n \ge 6: X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_y, \cdots, X_{n-3} = x^{n-5}\partial_y, X_{n-2} = x\partial_x, X_{n-1} = y\partial_y, X_n = x^2\partial_x + rxy\partial_y.$

The fundamental invariants:

$$\phi_1 = \frac{(n-3)^2 K_2 - 3(n-1)(n-3)K_1 + 2(n-1)(n-2)}{((n-3)K_1 - (n-2))^{\frac{3}{2}}},$$

$$\phi_2 = K_3((n-3)K_1 - (n-2))^{-2} + ((n-3)K_1 - (n-2))^{-3}[-\frac{3}{2}(n-3)K_2^2 + \frac{1}{2}(5n-13)K_1K_2 - 3(n-1)K_1^3 + \frac{3}{2}(n-1)K_1^2 - 2(n-2)K_2],$$

where

$$K_i = \frac{y^{(n-3+i)}(y^{(n-4)})^i}{(y^{(n-3)})^{i+1}}, \ i = 1, 2, 3.$$

6 (n + 3)-Dimensional Algebras

Scalar first order equations admit infinite number of point symmetries. A second order equation does not admit a five dimensional symmetry algebra, as is well-known. The only third order equations that admit 6 dimensional algebras are

$$y''' = \frac{3y''^2}{2y'}$$
 and $y''' = \frac{3y'y''^2}{1+y'^2}$,

where the algebras are $\mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$ and $\mathfrak{so}(3,1)$, respectively.

In [26], it was shown that a fourth order equation either does not admit a 7 dimensional algebra or if it admits such an algebra, then it is not maximal.

For fifth order equations, the only equation which admits an 8 dimensional algebra is

$$y^{(5)} = \frac{5y'''y^{(4)}}{y''} - \frac{40y''^3}{9y''^2},$$
(6.1)

whose symmetry algebra is $\mathfrak{sl}(3)$ as stated in the previous section.

We check all the possible algebras of dimension n + 3 for $n \ge 5$. Since an *n*th order linear equation cannot have more than *n* independent solutions, the algebras (m, n + 3) for $m = 20, \dots, 25$ are not admitted. We discuss the remaining possible algebras:

$$(26, n+3): X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_y, X_5 = x\partial_y, \cdots, X_{n+3} = x^{n-1}\partial_y.$$

The generators X_1, X_2 and X_5, \dots, X_{n+1} implies that the equation must be of the form

$$y^{(n)} = K.$$

The X_4 then implies that K must vanish. Then X_3 is automatically admitted by this equation. However, the maximal symmetry algebra of this equation is n+4. In the same way it can be shown easily that the algebras (27, n+3) and (28, n+3) are admitted by an *n*th order equation if and only if the equation is equivalent to $y^{(n)} = 0$ and hence these algebras are admitted by the equation but are not maximal. This discussion proves the following proposition.

Proposition 6.1. For $n \ge 4$ an *n*th order ODE either does not admit an (n + 3)-dimensional algebra or it does admit it but is not maximal with the exception that for n = 5, $\mathfrak{sl}(3)$ is admitted by the equation given as (6.1).

In the next section we review n = 4 and discuss n = 5 at some length. Then we mention how one can obtain a classification for n > 5.

7 The Cases $n \ge 4$

In this section we review the main aspects on the classification of 4th order ODEs and then discuss the cases $n \ge 5$. We consider the 5th order case in detail as an example to show how one can obtain the classification of ODEs of any higher order.

In general we can classify scalar ODEs into 3 subclasses as follows:

Subclass (1): *n*th order equations admitting n + 1, n + 2, n + 3 and the maximal n + 4 dimensional algebras.

Subclass (2): *n*th order equations admitting *n* dimensional algebras. For this we have found the fundamental invariants of *n* dimensional algebras which are of order n-1 and n.

Subclass (3): nth order equations admitting algebras of dimension lower than n.

Subclasses (1) and (2) are completed in this work for $n \ge 5$ and the algebra of dimension n + 4 is already a well-known algebra since Lie with corresponding equation $y^{(n)} = 0$ that possesses this maximal dimension Lie algebra.

Subclass (3) can be performed in the following manner:

For algebras of dimension 1 and 2, this is easy and is given in Table 1 below.

The *n*th Order Equations Admitting 1 and 2 Dimensional Algebras for $n \ge 3$

For any 3 dimensional algebra we already know the fundamental invariants of orders 2, 3 and 4 from the works of Gat [14], Mahomed and Leach [10] as well as [26]. Then by

Algebra	Generators	Invariant Equation
(9, 1)	∂_x	$y^{(n)} = H(y, y', y'',, y^{(n-1)})$
(10, 2)	$\partial_x, \ x\partial_x$	$y^{(n)} = y'^n H(y, y''y'^{-2},, y^{(n-1)}y'^{1-n})$
(20, 2)	$\partial_y, x \partial_y$	$y^{(n)} = H(x, y'', \dots y^{(n-1)})$
(22, 2)	∂_x, ∂_y	$y^{(n)} = H(y', y'',, y^{(n-1)})$
$A_{2,1}$	$\partial_x, x\partial_x + y\partial_y$	$y^{(n)} = y^{1-n} H(y', y''y, y'''y^2,, y^{(n-1)}y^{n-2})$

Table 1: The *n*th Order Equations Admitting 1 and 2 Dimensional Algebras for $n \ge 3$

invariant differentiation we can find higher order invariants up to the required order, i.e. n. For this Lie's recursive formula

$$w_k = \frac{D_x(w_{k-1})}{D_x(w_{k-2})}, k = 5, 6, ..., n$$
(7.1)

can be utilised, where w_k is a kth order invariant. The general form of an nth order equation admitting this 3 dimensional algebra can be given as:

$$F(w_2, w_3, \dots, w_n) = 0 \tag{7.2}$$

or as

$$w_n = H(w_2, w_3, \dots, w_{n-1}). (7.3)$$

Another way is to use operator of invariant differentiation $\mathcal{D} = \lambda D_x$. For example, if we know a second order invariant ϕ_1 of a low dimensional algebra, then higher order equations for this algebra are of the form

$$\mathcal{D}^{n-2}\phi_1 = H(\phi_1, \dots, \mathcal{D}^{n-3}\phi_1), \quad n \ge 3.$$
 (7.4)

In a similar way, this can be done for algebras of higher dimensions.

We first briefly review fourth order equations that possess 4 dimensional algebras in a compact way than in [26]. This is presented in the following Table 2.

Algebra	λ	3rd order Invariant ϕ_1	4th Order ODE
(4, 4)	$(1+y'^2)y''^{-1}$	$y''^{-2}y'''(1+y'^2) - 3y'$	$y^{(4)} = y''^3 (1 + y'^2)^{-2} (15y'^2)$
			$+10\phi_1y'+H(\phi_1)$
(13, 4)	$y'y''^{-1}$	$y'y''^{-2}y'''$	$y^{(4)} = y'^{-2}y''^{3}H(\phi_1)$
(14, 4)	y'^{-1}	$y'''y'^{-3} - \frac{3}{2}y''^2y'^{-4}$	$y^{(4)} = y'^4 H(\phi_1) + 6y'^2 y'' \phi_1$
			$+3y'^{-2}y''^{3}$
(19, 4)	$y^{1/2}y''^{-1/2}$	$y^{1/2}y''^{-3/2}y''' + 3y'y^{-1/2}y''^{-1/2}$	$y^{(4)} = \frac{4}{3}y''^{-1}y'''^{2} + y^{-1}y''^{2}H(\phi_{1})$
(22,4)	1	$y''' + a_1 y'' + a_2 y' + a_3 y$	$D_x\phi_1 = H(\phi_1)$
		η_i solves $\phi_1 = 0$ for a_i s const.	
(23,4)	1	$D_x \ln E \equiv D_x \ln y'' + a_1 y' + a_2 y $	$D_x\phi_1 = H(\phi_1)$
		η_i satisfy $E = 0$ for a_i const.	
(24, 4)	$y^{\prime\prime 1/(\alpha-2)},$	$y^{\prime\prime\prime lpha-2}y^{\prime\prime 3-lpha}$	$y^{(4)} = y'''^{\frac{\alpha-4}{\alpha-3}} H(\phi_1), \alpha \neq 3$
	y''	$y^{\prime\prime\prime}$	$y^{(4)} = y''^{-1}H(y''')$
	y'''^{-1}	y''	$y^{(4)} = y'''^2 H(y'')$
(25,4)	$\exp(y''/2)$	$y''' \exp(y''/2)$	$y^{(4)} = \exp(-y'')H(\phi_1)$

Table 2: The 4th Order Equations Admitting 4 Dimensional Algebras

There are multiple cases of (22,4) in [26]. These are considered as one case in the Table 2. Note that invariant differentiation of the invariants in the above Table 2 will give fifth and higher order invariants.

We now consider fifth order invariant equations admitting algebras of dimension 5 and greater. These are given in Table 3 and 4 and follow from our deliberations in the previous sections.

The 5th order invariant ODEs that correspond to their symmetry algebras

5 D Algebra	5th order canonical equation
(5, 5)	$y^{(5)} = \frac{-40}{9} y^{\prime\prime\prime} y^{\prime\prime-2} + 5y^{\prime\prime\prime} y^{(4)} y^{\prime\prime-1} + y^{\prime\prime2} H(\phi_1)$
	$\phi_1 = y^{(4)} y''^{\frac{-5}{3}} - \frac{5}{3} y'''^2 y''^{-8/3}$
(15, 5)	$\frac{3y'^3y^{(5)} - 10y'^2y''y^{(4)} - 50y'^2y'''^2 + 240y'y''^2y''' - 180y''^4}{(2 + t''' - 2y'')^2} = H(\phi_1)$
(10, 0)	$(2y'y''' - 3y''^2)^2$
	$\phi_1 = \frac{y'^2 y^{(4)} - 6(y'y''y''' - y''^3)}{y''y''' - y''^3}$
	$\varphi_1 = {(2y'y''' - 3y''^2)^{\frac{3}{2}}}$
(22,5)	$D_x(E_4(y)) = H(E_4(y))$
	η_i is solution of $E_4(\eta_i) = 0, i = 1, \dots, 4$ for a_i consts.

5 D Algebra	5th order canonical equation
(23,5)	$D_x^2 \ln (E_3(y)) = H(D_x \ln E_3(y)) $
	η_i is solution of $E_3(\eta_i) = 0, i = 1, \dots, 3$ for a_i consts.
(24, 5)	$y^{(5)} = (y^{(4)})^{\frac{\alpha-5}{\alpha-4}} H((y^{(4)})^{\alpha-3}(y^{(3)})^{4-\alpha}), \ \alpha \neq 4$
	$y^{(5)} = (y''')^{-1}H(y^{(4)}), \alpha = 4$
(25, 5)	$y^{(5)} = \exp(-\frac{2y'''}{3!})H(y^{(4)}\exp(y'''/3!))$
(26, 5)	$y^{(5)}y''^2y'''^{-3} = H(y''y'''^{-2}y^{(4)})$
(27, 5)	$y^{(5)}y''^{-3} + \frac{40}{9}y'''^{3}y''^{-5} - 5y^{(4)}y'''y''^{-4} = H(\phi_1)$
	$\phi_1 = y^{(4)}y''^{-7/3} - \frac{4}{3}y'''^2y''^{-10/3}, H \neq K\phi_1^{3/2}, K \text{ const.}$

Table 3: The 5th order invariant ODES that correspond to their 5 dimensional symmetry algebras. Note that $E_n(z) \equiv z^{(n)} + a_1 z^{(n-1)} + \cdots + a_n z$.

Higher Algebra	5th order invariant equation
(6, 6)	$\frac{9y''^2y^{(5)} - 45y''y'''y^{(4)} + 40y'''^3}{(3y''y^{(4)} - 5y'''^2)^{\frac{3}{2}}} = K, K \text{ const.}$
(16, 6)	$K_1^{-3}(5K_2^2 - 4K_1K_3) = K, K \text{ const.}$
(7,6)	$K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2) = K, K \text{ const.}$
(21,6)	$y^{(5)} = \sum_{i=2}^{4} A_i(x) y^{(i)}, \xi_k \text{ satisfy } \xi_k^{(5)} = \sum_{i=2}^{4} A_i(x) \xi_k^{(i)}, k = 1, \dots, 4$
(24,6)	$y^{(5)} = K(y^{(4)})^{\frac{\alpha-5}{\alpha-4}}, \ \alpha \neq 4$
(25, 6)	$y^{(5)} = K \exp(-y^{(4)}/4!)$
(26, 6)	$y^{(5)} = K y^{\prime\prime\prime - 1} (y^{(4)})^2$
(27, 6)	$y^{(5)} = K(y^{(3)})^2 + \frac{5}{4}(y^{(4)})^2 y^{\prime\prime\prime - 1}$
(28, 6)	$\left(\frac{y^{(5)}y''^2}{y'''^3} - \frac{5(9K_1-8)}{9}\right)/(3K_1-4)^{3/2} = K, K \text{ const.}$
	$K_1 = y^{(4)} y'' y'''^{-2}$
(28, 7)	$y^{(5)} = \frac{5}{4} (y^{(4)})^2 y^{\prime\prime\prime - 1}$
(8,8)	$9y''^2y^{(5)} - 45y''y'''y^{(4)} + 40y'''^3 = 0$
(28, 9)	$y^{(5)} = 0$

Table 4: The 5th order invariant ODEs that correspond to their higher symmetry algebras

One can proceed to classify 6th order ODEs as follows. The algebras are (6,6) given by $\phi_2 = H(\phi_1, \text{ where } \phi_1 = K_1^{-3/2} K_2 \text{ and } \phi_2 = K_1^{-1/2} D_x \phi_1$; (16,6) with equation

$$K_1^{-1/2}D_x[K_1^{-3}(5K_2^2 - 4K_3K_1)] = H(K_1^{-3}(5K_2^2 - 4K_3K_1));$$

(7,6) with representative ODE

$$\mathcal{D}[K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)] = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_2^2 + 4K_1K_3 + 45 + 16K_1^2)) = H(K_1^3(30K_2 - 5K_1K_3 + 5K_$$

Also one can easily obtain the forms for the equations for the algebras (22,6) to (28,6). Altogether, there are hence 10 canonical forms for 6th order ODEs admitting 6 dimensional algebras. Moreover, for higher symmetries, there are 9 types given by (21,7), (24,7) to (28,7) as well as (23,8), (28,8) and (28,9).

Likewise, if we investigate 7th order ODEs, the number of canonical forms are 7 given by (22,7) to (28,7). Moreover, for higher symmetries, we end up with 10 types, viz. (21,8), (24,8) to (28,8), (23,9), (28,9), (28,11) and (8,8) given by the ODE

$$K_2^{-5/3} D_x^2 K_2 - \frac{7}{6} K_2^{-8/3} (D_x K_2)^2 - y''^{-1} y''' K_2^{-5/3} D_x K_2$$
$$-y''^{-2} y'''^2 K_2^{2/3} - \frac{3}{2} y''^{-1} y^{(4)} K_2^{-2/3} = K.$$

Now proceeding to 8th order ODEs, we have the 8 canonical forms (22,8) to (28,8) as well as (8,8) given by

$$\mathcal{D}w = K_2^{-13} D_x (K_2^{-5/3} D_x^2 K_2 - \frac{7}{6} K_2^{-8/3} (D_x K_2)^2 - y''^{-1} y''' K_2^{-5/3} D_x K_2$$
$$-y''^{-2} y'''^2 K_2^{2/3} - \frac{3}{2} y''^{-1} y^{(4)} K_2^{-2/3}) = H(w)$$

where $\mathcal{D} = K_2^{-13} D_x$.

There are also 9 types of higher symmetries, viz. (21,9), (24,9) to (28,9), (23,10), (28,10) and (28,12).

We have the generalization to order 9 and greater as follows:

For $n \ge 9$, there are 7 canonical forms for *n*th order ODEs admitting *n* symmetries. These are (22, n) to (28, n).

Also, for higher symmetries for nth order equations, $n \ge 9$, there are 9 types, viz. (21, n+1), (24, n+1) to (28, n+1), (23, n+2), (28, n+2) and (28, n+4) the maximal algebra.

We also remark here that the invariants above are verified by Maple code confirming that it is correct. Below is an example to show this.

Example 7.1. (28,6): $X_1 = \partial_x$, $X_2 = \partial_y$, $X_3 = x\partial_x$, $X_4 = x\partial_y$, $X_5 = x^2\partial_x + xy\partial_y$, $X_6 = y\partial_y$.

The fifth order differential invariant is:

$$\phi = \left(\frac{y^{(5)}y''^2}{y'''^3} - \frac{5(9K_1 - 8)}{9}\right) / (3K_1 - 4)^{\frac{3}{2}},$$
$$K_1 = \frac{y^{(4)}y''}{y'''^2}.$$

Using the Maple code we get the differential invariant :

$$\phi = (1/9)(9y''^2y^{(5)} - 45y''y'''y^{(4)} + 40y'''^3)/(3y''y^{(4)} - 4y'''^2)^{\frac{3}{2}}$$

which is equivalent to the above invariant computed by hand.

Appendix

A. Some Fourth Order Invariants in [26]

Here we provide the corrected invariants that had inadvertent errors in [26].

(1, 3):
$$X_1 = \partial_x, X_2 = \partial_y, X_3 = (\alpha x + y)\partial_x + (\alpha y - x)\partial_y$$

The second and third order invariants are:

$$w_2 = y''(1+y'^2)^{\frac{-3}{2}}e^{-\alpha \arctan(y')}, w_3 = \frac{y'''(1+y'^2)}{y''^2} - 3y'.$$

The fourth order invariant is

$$w = \lambda D_x(w_3)$$
 where $\lambda = (1 + y'^2)^{\frac{-1}{2}} e^{-\alpha \arctan(y')}$

and simplifies as w/w_2 which is

$$w_4 = (y'^2 + 1)^2 y''^{-3} y^{(4)} - 2y''^{-4} y'''^2 (y'^2 + 1)^2 + 2y' y''^{-2} y''' (y'^2 + 1) - 3(y'^2 + 1)$$

(2, 3): $X_1 = \partial_y, X_2 = x \partial_x + y \partial_y, X_3 = 2xy \partial_x + (y^2 - x^2) \partial_y$

We know the second and third order invariants from the works [10] and [14] which are

$$w_2 = xy''(1+y'^2)^{\frac{-3}{2}} - y'(1+y'^2)^{\frac{-1}{2}}, w_3 = \frac{x^2y'''}{(1+y'^2)^2} - \frac{-3x^2y'y''^2}{(1+y'^2)^3}$$

The fourth order invariant can then be found by invariant differentiation which is:

$$w_4 = \lambda D_x(w_3).$$

This can also be found once we know a suitable λ which in this case is $\lambda = x(1+y'^2)^{\frac{-1}{2}}$. The explicit form of this fourth order invariant is given below which is found using Maple. Moreover, the general form of a fourth order equation invariant under this algebra can also be written using the equation (7.3).

The fourth order invariant:

$$\frac{1}{(y'^2+1)^{(\frac{9}{2})}}((15y'^2y''^3-10y'y'''(y'^2+1)y''+y^{(4)}(y'^2+1)^2)x^3+(2(y'^2+1))(-\frac{15}{2})y'y''^2+y''(y'^2+1)x^2-27(y'^2+1)^2y''(y'^2+\frac{4}{3})x+33(y'^2+1)^3(y'^2+\frac{12}{11})y')$$

$$(3, 3): X_1 = y\partial_x - x\partial_y, X_2 = (x^2 - y^2 + 1)\partial_x + 2xy\partial_y, X_3 = 2xy\partial_x + (-x^2 + y^2 + 1)\partial_y$$

The second and third order invariants are:

$$w_2 = \frac{y''(1+x^2+y^2)}{(1+y'^2)^{\frac{3}{2}}} + \frac{2(y-xy')}{(1+y'^2)^{\frac{1}{2}}}, w_3 = \frac{y'''(1+x^2+y^2)^2}{(1+y'^2)^2} - \frac{3y'y''^2(1+x^2+y^2)^2}{(1+y'^2)^3}$$

The fourth order invariant is then:

$$w_4 = \lambda D_x(w_3)$$
 for $\lambda = (1 + x^2 + y^2)(1 + y'^2)^{\frac{-1}{2}}$

The fourth order invariant:

 $\frac{1}{(y'^2+1)^{\frac{9}{2}}} ((15y'^2y''^3-10y'y'''(y'^2+1)y''+y^{(4)}(y'^2+1)^2)x^6 + (4(-\frac{15}{2}y'y''^2+(y'^2+1)y'''-\frac{16}{2}y'y'')(y'^2+1)x^5 + (45y'^2(y^2+1)y''^3+(-12y'^4+6y'^2+18)yy''^2-30y'((y^2+1)y'''-\frac{6}{5}y'^3-\frac{6}{5}y')(y'^2+1)y''+(3*(\frac{4}{3}yy'y'''+y^{(4)}(y^2+1)))(y'^2+1)^2)x^4 + (8(y'^2+1))(-\frac{15}{2}y'(y^2+1)y'''-\frac{6}{5}y'^2-\frac{9y'y(y'^2+1)y''-(y'^2+1)y''-3y'^5-3y'^3)(y'^2+1)x^3+(45y'^2(y^2+1)^2y''^3-(24(y'^2-\frac{3}{2}))(y^2+1)y(y'^2+1)y''-(\frac{6}{5}((y'^2+1)y^2+y'^2))(y'^2+1)))(y'^2+1)y''+(3(\frac{8}{3}y'y(y^2+1)y'''-(\frac{6}{5}((y'^2+1)y^2+y'^2))(y'^2+1)))(y'^2+1)y''+(3(\frac{8}{3}y'y(y^2+1)y'''-\frac{18}{2}y'(y'^2+1)(y^2+1)y''+((y^2+1)^2y'''-(18y^2y'(y'^2+1))(y'^2+1)))(y'^2+1)x+15y'^2(y^2+1)y'''-(18y'y(y'^2+1)(y'^2+1)y''-(10(y'(y^2+1)^2y'''-(\frac{18}{5}(y'^2+1))y^2))(y^2+1)(y'^2+1)y''+(4y'y(y^2+1)^2y'''+y^{(4)}y^6+3y^{(4)}y^4+(24y'^2+24)y^3+3y^{(4)}y^2+y^{(4)})(y'^2+1)^2)$

(11,3): $X_1 = \partial_x, X_2 = x \partial_x, X_3 = x^2 \partial_x$ A zero and third order invariants are

$$w_0 = y, w_3 = \frac{2}{3}y'^{-3}y''' - y'^{-4}y''^2$$

A direct computation using $\mathcal{D} = y'^{-1}D_x$ is simple in this case and we get a fourth order invariant:

$$w_4 = y^{(4)}y'^{-4} + 6y''^3y'^{-6} - 6y''y'''y'^{-5}.$$
7, 3): $X_1 = \partial_y, X_2 = x\partial_x + y\partial_y, X_3 = 2xy\partial_x + (x^2 + y^2)\partial_y$

The second and third order invariants are:

$$w_2 = \frac{xy'' - y' + y'^3}{(1 - y'^2)^{\frac{3}{2}}}, w_3 = \frac{x^2 y'''}{(y'^2 - 1)^2} - \frac{3x^2 y' y''^2}{(y'^2 - 1)^3}$$

and the fourth order invariant is:

(1)

$$w_4 = \lambda D_x(w_3)$$
 where $\lambda = x(1 - y'^2)^{\frac{-1}{2}}$

B. The fundamental invariants of some algebras via Maple

Here we present the fundamental differential invariants of the algebras (7, 6), (8, 8) and (16, 6). We have computed them using Maple 18 as doing them by hand is a tedious job.

$$(7,6): X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x + y\partial_y, X_4 = y\partial_x - x\partial_y, X_5 = (x^2 - y^2)\partial_x + 2xy\partial_y, X_6 = 2xy\partial_x + (y^2 - x^2)\partial_y.$$

Fundamental Invariants:

$$\begin{split} \phi_1 &= (1/4) \frac{1}{(y'^2 y''' - 3y' y''^2 + y''')^3} ((135y'^4 - 270y'^2 - 45)y''^6 + (-180y'^5 + 180y')y''' y''^4 + \\ &\quad 30y^{(4)}(y'^2 + 1)^3 y''^3 - 12(y'^2 + 1)^2((-\frac{10}{3}y'^2 + \frac{20}{3})y'''^2 + y'y^{(5)}(y'^2 + 1))y''^2 + \\ &\quad 40y'y'''y^{(4)}(y'^2 + 1)^3y'' + 4(y'^2 + 1)^3(-10y'y'''^3 + y^{(5)}(y'^2 + 1)y''' - \frac{5}{4})(y^{(4)})^2(y'^2 + 1))). \end{split}$$

$$\begin{split} \phi_2 &= \frac{1}{(\sqrt{-y'^2y''' + 3y'y''^2 - y'''}(y'^2 y''' - 3y'y''^2 + y''')^4)} (y'^2 + 1)^3(-\frac{405}{4}) * y''^9 + \\ &\frac{405}{2}y'''y'y''^7 - \frac{135}{4}y^{(4)}(y'^2 - 3)y''^6 + ((-\frac{225}{2}y'^2 - \frac{405}{2})y'''^2 + \frac{27}{2}y'y^{(5)}(y'^2 - 3))y''^5 + (9(-\frac{5}{2}y'''(y'^2 - 7)y''(y'^2 + 1))y''(y''^4 + (-\frac{135}{4}(y'^2 + 1)^2(y'^4))^2 - \frac{153}{2}y'''((-\frac{40}{51}y'^3 + \frac{40}{51}y')y'''^2 + (y'^2 + 1)y^{(5)}(y'^2 - \frac{3}{17}))y''^3 - (6(y'^2 + 1))(((-\frac{175}{4}y'^2 - \frac{35}{4})y'''^2 - \frac{9}{4}y'y^{(5)}(y'^2 + 1))y''(y'' + (y''^2 + 1)y'''(-\frac{15}{8}y'^3 - \frac{15}{8}y')(y^{(4)})^2 + ((-\frac{85}{12}y'^2 - \frac{25}{12})y''''^2 + y''y^{(5)}(y'^2 + 1))y''' + (y'^2 + 1)^2((\frac{15}{4}y'^2 + \frac{15}{4})(y^{(4)})^3 - (\frac{9}{2}(-\frac{20}{9}y'y'''^2 + y^{(5)}(y'^2 + 1)))y'''y'' + y'''y^{(6)}(y'^2 + 1))). \end{split}$$

$$(16, 6): X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_y, X_5 = x^2\partial_x, X_6 = y^2\partial_y. \end{split}$$

Fundamental Invariants:

$$\phi_1 = (1/6) d \frac{y'^3 (12y'y'''y^{(5)} - 15y'(y^{(4)})^2 - 18y''^2y^{(5)} + 60y''y'''y^{(4)} - 40y'''^3)}{(2y'y''' - 3y''^2)^3}.$$

$$\phi_{2} = \frac{1}{(\sqrt{-2y'y'''+3y''^{2}}(2y'y'''-3y''^{2})^{4})} (4((y'''^{2}y'^{6}) - \frac{9}{2}y'''y'^{4}y'^{5}) + \frac{15}{4}(y'^{4})^{3})y'^{3} + (5y'''^{3}y'^{4}) + 12y''y'''^{2}y'^{5} + (-3y''^{2}y'^{6}) - \frac{45}{2}y''(y'^{4})^{2})y''' + \frac{27}{4}y''^{2}y'^{4}y'^{5})y'^{2} + \frac{9}{4}y''(y''^{3}y'^{6}) - 10y''^{2}y'''y'^{5} + \frac{70}{3}y''y'''^{2}y'^{4} - \frac{40}{3}y'''^{4})y' + \frac{27}{4}y''^{5}y'^{5} - \frac{45}{2}y'''^{4}y'''y'^{4} + 15y''^{3}y'''^{3})y'^{3}).$$

 $(8, 8): X_1 = \partial_x, X_2 = \partial_y, X_3 = x\partial_x, X_4 = y\partial_x, X_5 = x\partial_y, X_6 = y\partial_y, X_7 = x^2\partial_x + xy\partial_y, X_8 = xy\partial_x + y^2\partial_y.$

Fundamental Invariants:

$$\begin{split} & \phi_1 = (1/18) \frac{1}{(9y''^2y^{(5)} - 45y''y'''y^{(4)} + 40y''^3)^{8/3}} (162y''^6y^{(5)}y^{(7)} - 189y''^6(y^{(6)})^2 - \\ & 810y''^5y'''y^{(4)}y^{(7)} + 1134y''^5y'''y^{(5)}y^{(6)} + 1890y''^5(y^{(4)})^2y^{(6)} - 2835y''^5y^{(4)}(y^{(5)})^2 + \\ & 720y''^4y'''^3y^{(7)} - 3150y''^4y'''^2y^{(4)}y^{(6)} - 756y''^4y'''^2y^{(5)})^2 + 13230y''^4y'''(y^{(4)})^2y^{(5)} - \\ & 4725y''^4(y^{(4)})^4 - 12600y''^3y'''^3y^{(4)}y^{(5)} - 7875y''^3y'''^2(y^{(4)})^3 + 6720y''^2y'''^5y^{(5)} + \\ & 31500y''^2y'''^4(y^{(4)})^2 - 33600y''y'''^6y^{(4)} + 11200y'''^8). \end{split}$$
 $\phi_2 = (1/81) \frac{1}{(9y''^2y^{(5)} - 45y''y'''y^{(4)} + 40y'''^3)^4} (-2551500y''^6(y^{(4)})^6 + \\ & 7016625y''^5y'''^2(y^{(4)})^5 + (1530900(y''^3y^{(6)} + \frac{7}{2}y''^2y'''y^{(5)} + 45y'''^4))y''^4(y^{(4)})^4 + \\ & (-656100y''^7y'''y^{(7)} - 1760535y''^7(y^{(5)})^2 - 5868450y''^6y'''^2y^{(6)} - 73823400y''^5y'''^3y^{(5)} - \\ & 255150000y''^3y'''^6(y^{(4)})^3 + (131220(161y''^4y'''^2y^{(5)})^2 + y''^2(y''^4y^{(7)} + \frac{14}{3}y''^3y'''y^{(6)} + \\ & \frac{3640}{3}y'''^5)y^{(5)} - \frac{7}{3}y''^6(y^{(6)})^2 + \frac{5}{2}y''^5y''''y^{(8)} + \frac{160}{9}y''^4y'''^3y^{(7)} + \frac{1540}{24}y'''^3y'''y^{(6)} + \\ & \frac{70000}{3}y'''(y'^5y^{(8)} + \frac{40}{3}y''^4y'''y^{(7)} + \frac{80}{29}y''^3 x''''y^{(6)} + \frac{56000}{9}y'''^5)^{(5)} - 2y'''(y'''y^{(6)}y^{(7)} + \\ & \frac{35}{9}y''^6y'''(y^{(6)})^2 - \frac{20}{9}y''^5y'''^3y^{(8)} - \frac{440}{27}y'''y^{(4)}y''y^{(7)} - \frac{140}{21}y''^3y'''^5y^{(6)} - \frac{1120000}{729}y'''^9)))y''y^{(4)} + \\ & 653184y'''^6y''''(y^{(5)})^3 + 6561y''(y'''y^{(6)}y^{(7)} + 7y''^6y'''(y^{(6)})^2 - \frac{20}{9}y''^5y'''^3y^{(6)} + \\ & \frac{150080}{27}y'''^3y^{(6)} - \frac{1120000}{729}y'''^9)y^{(5)} + 20412y'''(y^{(6)})^3 - 116640y'''y'''^3y^{(6)}y^{(7)} + \\ & 129600y''^5y'''^6y^{(8)} + 518400y'''y'''^7 + 44800000y'''^1). \end{split}$

Acknowledgments. FM thanks the NRF and Wits for research support.

References

- [1] Lie, S., Differential Equations, Chelsea, New York, 1967 [in German].
- [2] Lie, S., "Vorlesungen iber Continuerliche Gruppen mit geometrischen und anderen Anwendungen," Bearbeitet und herausgegben von Dr. G. Scheffers, Tuebner, Leipzig, 1893.
- [3] Lie, S. (1888). Classification und Integration von gewöhnlichen Differentialgleichungen zwischen xy, die eine Gruppe von Transformationen gestatten. Mathematische Annalen, 32(2), 213-281.
- [4] Mahomed, F. M. and Leach, P. G. L., Lie algebras associated with second order ordinary differential equations, J. Math. Phys. 30 (1989), 277 0-2777.
- [5] Ibragimov, N. H. and Mahomed, F. M., Ordinary differential equations, in CRC Hanbook of Lie Group Analysis of Differential Equations, Vol. 3, N. H. Ibragimov (ed.), CRC Press, Boca Raton, FL, 1996, pp. 191–215.

- [6] Sarlet, W., Mahomed, F. M. and Leach, P. G. L., Symmetries of non-linear differential equations and linearisation, J. Phys. A: Math. Gen. 20 (1987), 277-292.
- [7] Mahomed, F. M. and Leach, P. G. L., The Lie algebra sl(3,R) and linearization, Queastiones Math. 12 (1989), 121-139.
- [8] Mahomed, F. M. 2007, Symmetry group classification of ordinary differential equations: Survey of some results, Math. Methods in the Applied Sciences, 30 (2007) 1995-2012.
- [9] Mahomed, F. M. and Leach, P. G. L., Symmetry Lie algebra of nth order ordinary differential equations, Journal of Mathematical Analysis Applications 151, 1990, 80–107.
- [10] Mahomed, F. M., and Leach, P. G. L. (1988, January). Normal forms for third order equations. In Proceedings of the Workshop on Finite Dimensional Integrable Nonlinear Dynamical Systems," World Scientific Pub (pp. 178-189).
- [11] Krause J., and Michel L. Classification of symmetries of ordinary differential equations. Lecture Notes in Physics 1991; 382:251.
- [12] Chern S. S. The geometry of the differential equation, Tensor N.S. 1940; 28: 173–176.
- [13] Mahomed, F. M, and Leach P. G. L. Normal forms for third order equations. Proceedings of the Workshop on Finite Dimensional Integrable Nonlinear Dynamical Systems, Johannesburg, Leach P. G. L., Steeb W. H. (eds). World Scientific: Singapore, January 1988; 178.
- [14] Gat O. Symmetries of third order differential equations. Journal of Mathematical Physics 1992; 33: 2966.
- [15] Nesterenko, M. O. (2006). Transformation groups on real plane and their differential invariants. International journal of mathematics and mathematical sciences, 2006.
- [16] Grebot G. The characterization of third order ordinary differential equations admitting a transitive fibre-preserving point symmetry group. Journal of Mathematical Analysis and Applications 1997; 206 (2): 364.
- [17] Ibragimov N. H, Meleshko S. V. Linearization of third-order ordinary differential equations by point and contact transformations. Journal of Mathematical Analysis and Applications 2005; 308 (1): 266.
- [18] Neut, S., and Petitot, M., La geometrie de equation y''' = f(x, y, y'), Comp Rend de Acad. des Sciences, Paris, Serie 1, 2002, 335: 515-518.

- [19] Al-Dweik Ahmad Y., Mustafa M. T. and Mahomed Fazal Mahmood Mahomed 2018, Invariant characterization of scalar third-order ODEs that admit the maximal point symmetry Lie algebra, Mathematical Methods in the Applied Sciences, Vol. 41 Issue 12, 4714-4723, https://doi.org/10.1002/mma.4923.
- [20] Al-Dweik Ahmad Y., Mustafa M. T. and Mahomed Fazal Mahmood Mahomed and Alassar Rajai S 2018, Linearization of third-order ordinary differential equations u''' = f(x; u; u'; u'') via point transformations, Mathematical Methods in the Applied Sciences, Vol. 41 Issue 16, Aug, 6017-7098, https://doi.org/10.1002/mma.5208.
- [21] Al-Dweik Ahmad Y., Mahomed F. M. and Mustafa M. T. 2019, Invariant characterization of third-order ordinary differential equations u''' = f(x; u; u'; u'') with five-dimensional point symmetry group, Communications in Nonlinear Science and Numerical Simulation, Vol 67, Feb, 627-636.
- [22] Ibragimov, Nail H., and Maria Clara Nucci. "Integration of third order ordinary differential equations by Lie's method: equations admitting three-dimensional Lie algebras." Lie Groups and Their Applications 1 (1994): 49-64.
- [23] Cerquetelli, T., Ciccoli N. and Nucci, M. C. "Four dimensional Lie symmetry algebras and fourth order ordinary differential equations." Journal of Nonlinear Mathematical Physics 9. Sup 2 (2002): 24-35.
- [24] Fatima, A., Ayub, M. and Mahomed, F. M. "A note on four-dimensional symmetry algebras and fourth-order ordinary differential equations." Journal of Applied Mathematics 2013 (2013).
- [25] Shah, Said Waqas, F. M. and H. Azad, A Note on the Integration of Scalar Fourth-Order Ordinary Differential Equations with Four-Dimensional Symmetry Algebras, Mathematical Problems in Engineering, Vol. 2021, Article ID 6619325, 7 pages https://doi.org/10.1155/2021/6619325
- [26] Shah, S. W., Mahomed, F. M., Azad, H. and Mustafa, M. T. (2021). Complete Classification of Scalar Fourth-Order Ordinary Differential Equations and Linearizing Algorithms. Dynamic Systems and Applications, 30 (2021) No.3, 519-534.
- [27] Ibragimov, Nail, H., Meleshko, Sergey V. and Suksern, Supaporn, Linearization of fourth-order ordinary differential equations by point transformations, J. Phys. A: Math. Theor. 41 (2008) 235206 (19pp) doi:10.1088/1751-8113/41/23/235206.
- [28] González-López, A., Kamran, N. and Olver, P. J. (1992). Lie algebras of vector fields in the real plane. Proceedings of the London Mathematical Society, 3(2), 339-368.
- [29] Ibragimov, N. H. (1999). Elementary Lie group analysis and ordinary differential equations (Vol. 197). New York: Wiley.