

Impact of $b \rightarrow c$ measurements on $\Lambda_b \rightarrow p\tau\bar{\nu}$ decay in U_1 leptoquark model

Neetu Raj Singh Chundawat,^{1,*} Shabana Khan,^{2,†} and Dinesh Kumar^{2,‡}

¹*Indian Institute of Technology Jodhpur, Jodhpur 342037, India*

²*Department of Physics, University of Rajasthan, Jaipur 302004, India*

The measurements of several lepton flavor universality (LFU) violating observables in the decays induced by the quark level transition $b \rightarrow c\tau\bar{\nu}$ provide an inkling of plausible physics beyond the standard model of electroweak interactions. Such new physics would also impact other sectors. In this work, we estimate the leverage of new physics in $b \rightarrow c\tau\bar{\nu}$ on $\Lambda_b \rightarrow p\tau\bar{\nu}$ decay in the context of U_1 leptoquark model. In this model, the new physics couplings in $b \rightarrow u\tau\bar{\nu}$ transition can be written in terms of $b \rightarrow c\tau\bar{\nu}$ couplings and hence the extent of allowed new physics in $\Lambda_b \rightarrow p\tau\bar{\nu}$ would be determined by $b \rightarrow c\tau\bar{\nu}$ transition. Using the new physics parameter space obtained by performing a fit to all $b \rightarrow c\tau\bar{\nu}$ data, we obtain predictions of several $\Lambda_b \rightarrow p\tau\bar{\nu}$ observables. We find that the current $b \rightarrow c\tau\bar{\nu}$ data allows an order of magnitude enhancement in the branching ratio as well as in the LFU ratio. The other observables such as convexity parameter, lepton forward-backward asymmetry, longitudinal polarization of final state baryon and tau lepton are consistent with the SM value.

arXiv:2208.11106v1 [hep-ph] 23 Aug 2022

* chundawat.1@iitj.ac.in

† shabana26k@gmail.com

‡ dineshsuman09@gmail.com

I. INTRODUCTION

The standard model (SM) of electroweak interactions can be considered as a baronial theory of fundamental interactions of nature. Ever since the discovery of weak neutral currents in 1973 in a neutrino scattering experiment in the Gargamelle bubble chamber at CERN, SM has been substantiated through a fecundity of experimental observations. The discovery of the Higgs Boson marks the culmination of the particle spectrum of the SM. Though SM may flare out to be an irrefutable theory, there are several other observations which propel us to clamor for physics beyond the SM. These include disappearance of anti-matter, existence of dark matter and dark energy. Further, gravity is excluded from the SM. Therefore the nonpareil theory of fundamental interactions of nature is still far away from the bay.

The evidence of physics beyond the SM has already started burgeoning at several fronts. These include observables related to the decays of B mesons. These anomalous discrepancies can be classified into two categories: decays induced by the charged current transition $b \rightarrow c\ell\nu$ ($\ell = e, \mu, \tau$) and neutral current transition $b \rightarrow s\ell\ell$ ($\ell = e, \mu$). In this work, we rivet on decays induced by the $b \rightarrow c\ell\nu$ transition which occurs at the tree level in the SM. A series of measurements by the Belle, BaBar and LHCb collaborations over the last decade have provided several enthralling hints of new physics in this sector.

The BaBar [1, 2], Belle [3–5] and LHCb [6] collaborations measured the following flavor ratios

$$R_{D^{(*)}} \equiv \frac{\Gamma(B \rightarrow D^{(*)} \tau \bar{\nu})}{\Gamma(B \rightarrow D^{(*)} (e, \mu) \bar{\nu})}. \quad (1)$$

The average values of these measurements differ from their respective SM predictions at the level of 3.9σ [7]. These deviations are inklings of lepton flavor universality violation. All of these experiments were based on methodologies where the τ lepton was identified through kinematical information rather than reconstruction. The reconstruction technique was emplaced by the LHCb collaboration using the 3π decay mode of the τ lepton [8]. This resulted in a distinct measurement of R_{D^*} . Including this measurement, the incongruence of the R_D - R_{D^*} data with SM predictions escalated to 4.1σ [9]. In 2019, Belle collaboration announced a new measurement of R_D and R_{D^*} [10], which is consistent with the SM prediction. By including these measurements, the discrepancy with SM reduced from 4.1σ to 3.1σ .

Apart from $R_{D^{(*)}}$, the LHCb collaboration measured the following ratio in $B_c \rightarrow J/\psi \ell \bar{\nu}$ decay modes

$$R_{J/\psi} = \frac{\Gamma(B_c \rightarrow J/\psi \tau \bar{\nu})}{\Gamma(B_c \rightarrow J/\psi \mu \bar{\nu})}. \quad (2)$$

These decays are generated by the same quark level transition which induces $R_{D^{(*)}}$. The measured value is 1.7σ higher than the SM prediction [11]. These dissension with the SM can be imputed to new physics in τ , μ or e sectors. However, in [12] it was shown that new physics only in μ or e sectors cannot accommodate these measurements. This is mainly due to measurements of the ratios $R_D^{\mu/e} = \Gamma(B \rightarrow D \mu \nu)/\Gamma(B \rightarrow D e \nu) = 0.995 \pm 0.022$ (stat.) ± 0.039 (syst.) and $R_{D^*}^{e/\mu} = \Gamma(B \rightarrow D^* e \nu)/\Gamma(B \rightarrow D^* \mu \nu) = 1.04 \pm 0.05$ (stat.) ± 0.01 (syst.) [13, 14]. The measured values of these ratios are in agreement with their SM predictions. Hence new physics only in $b \rightarrow c \mu \bar{\nu}$ or $b \rightarrow c e \bar{\nu}$ will blight this agreement. Therefore new physics in $b \rightarrow c \tau \bar{\nu}$ is imperative to accommodate the current measurements of flavor ratios in these sectors ¹.

In May 2022, the LHCb collaboration reported the first observation of the semileptonic b -baryon decay $\Lambda_b \rightarrow \Lambda_c^+ \tau^- \bar{\nu}_\tau$ with a significance of 6.1σ [15]. This was obtained by collecting a data sample corresponding to 3 fb^{-1} of integrated luminosity at centre-of-mass energies of 7 and 8 TeV. The LFU ratio $R(\Lambda_c)$ was measured to be [15]

$$R(\Lambda_c) = \frac{\Lambda_b \rightarrow \Lambda_c^+ \tau^- \bar{\nu}_\tau}{\Lambda_b \rightarrow \Lambda_c^+ \mu^- \bar{\nu}_\mu} = 0.242 \pm 0.026 \text{ (stat.)} \pm 0.040 \text{ (syst.)} \pm 0.059. \quad (3)$$

Here the last error is due to the external branching fraction uncertainty from the channel $\Lambda_b \rightarrow \Lambda_c^+ \mu^- \bar{\nu}_\mu$. The measured value is consistent with the SM prediction of 0.324 ± 0.004 [16].

Barring these LFU observables, we also have measurements of few angular observables. The Belle collaboration has measured the τ polarization, $P_\tau^{D^*}$, in $B \rightarrow D^* \tau \bar{\nu}$ decay. The measured value [5]

$$P_\tau^{D^*} = -0.38 \pm 0.51 \text{ (stat.)}_{-0.16}^{+0.21} \text{ (syst.)}, \quad (4)$$

is consistent with its SM prediction of -0.497 ± 0.013 [17]. In 2018, Belle collaboration reported the measurement of D^* longitudinal polarization fraction $F_L^{D^*}$ in the decay $B \rightarrow D^* \tau \bar{\nu}$. The measured value [18]

$$F_L^{D^*} = 0.60 \pm 0.08 \text{ (stat.)} \pm 0.04 \text{ (syst.)} \quad (5)$$

is 1.6σ higher than the SM prediction of 0.46 ± 0.04 [19].

The possible new physics effects in $b \rightarrow c \tau \bar{\nu}$ decay can be analyzed in a model independent way using the language of effective field theory. There are many such analyses, see for e.g, [20–33]. These analyses identified Lorentz structure of possible new physics. However, there are no unique

¹ In [34], it was shown that new physics only in muons can accommodate the entire $b \rightarrow c l \bar{\nu}$ data using a different set of combinations of new physics operators.

solutions. Depending upon the adopted methodology and assumptions, there are multiple new physics operators with specific values of corresponding WCs which can provide a good fit to data. A unique determination of the new Lorentz structure of new physics would require measurements of additional observables in $b \rightarrow c\tau\bar{\nu}$ sector [35].

The allowed model independent solutions can be realized in specific new physics models. There are a good number of such models. In context of some of these models it would be interesting to see whether some correlations exist between the observables in $b \rightarrow c$ sector and other sectors. In other words, what implications measurement in $b \rightarrow c$ sector have on other sectors. In this work we explore such correlations in $b \rightarrow u$ sector in the context of U_1 leptoquark (LQ) model. In particular, we study imprints of $b \rightarrow c$ measurements on several observables in $\Lambda_b \rightarrow p\tau\bar{\nu}$ decay mode. The baryonic decay mode $\Lambda_b \rightarrow p\tau\bar{\nu}$ is studied in the literature [36, 37].

The quark level transition $b \rightarrow u\tau\bar{\nu}$ induces decays such as $B^+ \rightarrow \tau\bar{\nu}$, $B \rightarrow \pi\tau\bar{\nu}$, $B \rightarrow \rho\tau\bar{\nu}$, $B \rightarrow \omega\tau\bar{\nu}$ and $\Lambda_b \rightarrow p\tau\bar{\nu}$. Out of these decays, currently, the only observed decay channel is the purely leptonic decay $B^+ \rightarrow \tau\bar{\nu}$ [38]. The measured value of its branching ratio is $(1.09 \pm 0.24) \times 10^{-5}$ which is consistent with the SM value $(8.80 \pm 0.73) \times 10^{-5}$. Further, the Belle collaboration provides an upper bound on the branching ratio of the semileptonic decay $B \rightarrow \pi\tau\bar{\nu}$. At 90% C.L., the branching ratio of $B \rightarrow \pi\tau\bar{\nu}$ can be as high as 2.5×10^{-4} [39]. Thus due to lack of enough measurements, any model independent analysis would allow a large new physics effects in some of the observables in $b \rightarrow u\tau\bar{\nu}$ transition. However, in the context of U_1 leptoquark model considered in this work, we show that the necessary couplings in $b \rightarrow u\tau\bar{\nu}$ decay are all related to the couplings in $b \rightarrow c\tau\bar{\nu}$ sector. Given the fact that we have relatively accurate measurements of number of observables in this sector, it would be interesting to see the extent up to which the new physics effects are allowed in $b \rightarrow u\tau\bar{\nu}$ sector. In particular, we study the impact of $b \rightarrow c\tau\bar{\nu}$ measurements on several observables in $\Lambda_b \rightarrow p\tau\bar{\nu}$ decay mode.

Plan of work is as follows. In Sec.II, we provide theoretical framework of this work. Starting with the effective Hamiltonian, we provide all necessary theoretical expressions in this section. This includes various observables in $b \rightarrow c\tau\bar{\nu}$ sector and $\Lambda_b \rightarrow p\tau\bar{\nu}$ decay. In the next section, we first provide constraints on $b \rightarrow c\tau\bar{\nu}$ couplings by performing a fit. Using the allowed parameter space of these couplings, we obtain predictions of several observables in $\Lambda_b \rightarrow p\tau\bar{\nu}$ decay. The conclusions are discussed in Sec.IV.

II. THEORETICAL FRAMEWORK

A. Effective Hamiltonian

Within the SM, the effective Hamiltonian for the quark level transition $b \rightarrow q \tau \bar{\nu}$ with $q = u, c$ is given by

$$H_{eff}^{\text{SM}} = \frac{4G_F}{\sqrt{2}} V_{qb} O_{V_L}, \quad (6)$$

where $O_{V_L} = (\bar{q}\gamma_\mu P_L b) (\bar{\tau}\gamma^\mu P_L \nu)$. In the presence of new physics, the effective Hamiltonian takes the form

$$H_{eff} = \frac{4G_F}{\sqrt{2}} V_{qb} [(1 + C_{V_L})O_{V_L} + C_{V_R}O_{V_R} + C_{S_L}O_{S_L} + C_{S_R}O_{S_R} + C_T O_T], \quad (7)$$

where

$$O_{V_R} = (\bar{q}\gamma_\mu P_R b) (\bar{\tau}\gamma^\mu P_L \nu), \quad (8)$$

$$O_{S_R} = (\bar{q}P_R b) (\bar{\tau}P_L \nu), \quad (9)$$

$$O_{S_L} = (\bar{q}P_L b) (\bar{\tau}P_L \nu), \quad (10)$$

$$O_T = (\bar{q}\sigma^{\mu\nu} P_L b) (\bar{\tau}\sigma_{\mu\nu} P_L \nu). \quad (11)$$

The interactions between the vector singlet U_1 LQ and the SM quarks and leptons can be written as [40, 41]

$$H_{eff}^{U_1} = h_{ij}^L \bar{Q}^i \gamma_\mu U_1^\mu P_L L^j + h_{ij}^R \bar{d}^i \gamma_\mu U_1^\mu P_R l_R^j + h.c., \quad (12)$$

where Q_i and L_j are the SM left-handed quark and lepton doublets and d_R^i and l_R^j are right handed quarks and leptons. Here h_{ij}^L and h_{ij}^R are the 3×3 matrices in the flavor space. This LQ contributes to $b \rightarrow c \tau \bar{\nu}$ at the tree level. As we only require $\bar{c} \nu U_1$ and $\bar{b} \tau U_1$ couplings to be non-zero, we have

$$h^L = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & h_{23}^L \\ 0 & 0 & h_{33}^L \end{pmatrix}, \quad h^R = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & h_{33}^R \end{pmatrix} \quad (13)$$

Assuming mixing in the up-type quark sector, the interaction Hamiltonian can be written as

$$H_{eff} = \left[(V_{us} h_L^{23} + V_{ub} h_{33}^L) \bar{u}_L \gamma_\mu \nu_L + (V_{cb} h_L^{33} + V_{cs} h_{23}^L) \bar{c}_L \gamma_\mu \nu_L \right. \\ \left. + h_L^{23} \bar{s}_L \gamma_\mu \tau_L + h_L^{33} \bar{b}_L \gamma_\mu \tau_L + h_R^{33} \bar{b}_R \gamma_\mu \tau_R \right] U_1^\mu. \quad (14)$$

It is ostensible from the above Lagrangian that only O_{V_L} and O_{S_R} contribute to $b \rightarrow c\tau\bar{\nu}$ and $b \rightarrow u\tau\bar{\nu}$ processes. Also, the same couplings appear in both decay modes. The relevant WCs for $b \rightarrow c\tau\bar{\nu}$ decay can be written as

$$C_{V_L}^{b \rightarrow c} = \frac{1}{2\sqrt{2}G_F V_{cb}} \frac{(V_{cb}h_{33}^L + V_{cs}h_{23}^L)h_{33}^L}{M_{U_1}^2}, \quad (15)$$

$$C_{S_R}^{b \rightarrow c} = -\frac{1}{\sqrt{2}G_F V_{cb}} \frac{(V_{cb}h_{33}^L + V_{cs}h_{23}^L)h_{33}^R}{M_{U_1}^2}. \quad (16)$$

The WCs for $b \rightarrow u\tau\bar{\nu}$ decay are

$$C_{V_L}^{b \rightarrow u} = \frac{1}{2\sqrt{2}G_F V_{ub}} \frac{(V_{ub}h_{33}^L + V_{us}h_{23}^L)h_{33}^L}{M_{U_1}^2}, \quad (17)$$

$$C_{S_R}^{b \rightarrow u} = -\frac{1}{\sqrt{2}G_F V_{ub}} \frac{(V_{ub}h_{33}^L + V_{us}h_{23}^L)h_{33}^R}{M_{U_1}^2}. \quad (18)$$

Thus we see that the $b \rightarrow c$ couplings can determine the new physics contributions to $b \rightarrow u$. Therefore we need to analyze observables in the $b \rightarrow c\tau\bar{\nu}$ sector. In the next section we provide theoretical expressions for $b \rightarrow c\tau\bar{\nu}$ observables used in our analysis to constrain the new physics parameter space.

B. Observables in $b \rightarrow c\tau\bar{\nu}$ sector

We consider following observables in our analysis:

- the flavor ratios R_D , R_{D^*} and $R(\Lambda_c)$,
- tau polarization in $B \rightarrow D^*\tau\bar{\nu}$ decays,
- D^* longitudinal polarization fraction in $B \rightarrow D^*\tau\bar{\nu}$ decay,
- branching ratio of $B_c \rightarrow \tau\bar{\nu}$.

We do not include the flavor ratio $R_{J/\psi}$ due to large theoretical errors. The theoretical expressions for R_D , R_{D^*} and R_{Λ_c} in terms of WCs are given as [42]

$$R_D^{th} \simeq R_D^{\text{SM}} \left\{ |1 + C_{V_L}|^2 + 1.54 \text{Re} [(1 + C_{V_L})C_{S_R}] + 1.09|C_{S_R}|^2 \right\}, \quad (19)$$

$$R_{D^*}^{th} \simeq R_{D^*}^{\text{SM}} \left\{ |1 + C_{V_L}|^2 + 0.13 \text{Re} [(1 + C_{V_L})C_{S_R}] + 0.05|C_{S_R}|^2 \right\}, \quad (20)$$

$$R_{\Lambda_c}^{th} \simeq R_{\Lambda_c}^{\text{SM}} \left\{ |1 + C_{V_L}|^2 + 0.50 \text{Re} [(1 + C_{V_L})C_{S_R}] + 0.33|C_{S_R}|^2 \right\}. \quad (21)$$

Tau polarization in $B \rightarrow D^*\tau\bar{\nu}$ decay, $P_\tau^{D^*}$, in the U_1 LQ model is given as [42]

$$P_\tau^{D^* th} \simeq \left(\frac{R_{D^*}^{th}}{R_{D^*}^{\text{SM}}} \right)^{-1} \left\{ -0.49|1 + C_{V_L}|^2 + 0.13 \text{Re} [(1 + C_{V_L})C_{S_R}] + 0.05|C_{S_R}|^2 \right\}. \quad (22)$$

The expression for D^* longitudinal polarization fraction, $f_L^{D^*}$, in $B \rightarrow D^* \tau \bar{\nu}$ decay is [42]

$$f_L^{D^* th} \simeq \left(\frac{R_{D^*}^{th}}{R_{D^*}^{SM}} \right)^{-1} \left\{ 0.46 |1 + C_{V_L}|^2 + 0.13 \operatorname{Re} [(1 + C_{V_L}) C_{S_R}] + 0.05 |C_{S_R}|^2 \right\}. \quad (23)$$

We also consider the constraints coming from the purely leptonic decay $B_c \rightarrow \tau \bar{\nu}$. This decay mode is not affected by the helicity suppression provided the transition is induced through the pseudo-scalar operators. The branching ratio of $B_c \rightarrow \tau \bar{\nu}$ in the U_1 LQ model can be written as

$$\mathcal{B}(B_c \rightarrow \tau \bar{\nu}) \simeq 0.02 \left(\frac{f_{B_c}}{0.43 \text{ GeV}} \right)^2 \left| 1 + C_{V_L} + 4.3 C_{S_R} \right|^2. \quad (24)$$

C. Observables in $\Lambda_b \rightarrow p \tau \bar{\nu}$ decay mode

In this section, we provide theoretical expressions for various $\Lambda_b \rightarrow p \tau \bar{\nu}$ observables used in our analysis. These observables can be defined with the help of angular differential decay distribution of this mode. The two-fold angular differential distribution for $\Lambda_b \rightarrow p \tau \bar{\nu}$ can be written in terms of q^2 and $\cos \theta_l$ where q^2 is the momentum transfer squared and θ_l is the angle between the daughter baryon and the lepton in the di-lepton rest frame. The two-fold angular differential distribution can be written as

$$\frac{d^2 \Gamma(\Lambda_b \rightarrow p \tau \bar{\nu})}{dq^2 d \cos \theta_l} = N \left(1 - \frac{m_l^2}{q^2} \right)^2 \left[A + \frac{m_l^2}{q^2} B + 2C + \frac{4m_l}{\sqrt{q^2}} D \right], \quad (25)$$

where

$$A = 2 \sin^2 \theta_l \left(H_{\frac{1}{2},0}^2 + H_{-\frac{1}{2},0}^2 \right) + \left(1 - \cos \theta_l \right)^2 H_{\frac{1}{2},1}^2 + \left(1 + \cos \theta_l \right)^2 H_{-\frac{1}{2},-1}^2, \quad (26)$$

$$B = 2 \cos^2 \theta_l \left(H_{\frac{1}{2},0}^2 + H_{-\frac{1}{2},0}^2 \right) + \sin^2 \theta_l \left(H_{\frac{1}{2},1}^2 + H_{-\frac{1}{2},-1}^2 \right) + 2 \left(H_{\frac{1}{2},t}^2 + H_{-\frac{1}{2},t}^2 \right) - 4 \cos \theta_l \left(H_{\frac{1}{2},t} H_{\frac{1}{2},0} + H_{-\frac{1}{2},t} H_{-\frac{1}{2},0} \right) \quad (27)$$

$$C = \left(H_{\frac{1}{2},0}^{SP} \right)^2 + \left(H_{-\frac{1}{2},0}^{SP} \right)^2, \quad (28)$$

$$D = -\cos \theta_l \left(H_{\frac{1}{2},0} H_{\frac{1}{2},0}^{SP} + H_{-\frac{1}{2},0} H_{-\frac{1}{2},0}^{SP} \right) + \left(H_{\frac{1}{2},t} H_{\frac{1}{2},0}^{SP} + H_{-\frac{1}{2},t} H_{-\frac{1}{2},0}^{SP} \right). \quad (29)$$

The differential decay rate for $\Lambda_b \rightarrow p \tau \bar{\nu}$ can be obtained after integrating out equation 25 over the $\cos \theta_l$ variable [43]

$$\frac{d\Gamma(\Lambda_b \rightarrow p \tau \bar{\nu})}{dq^2} = \frac{8N}{3} \left(1 - \frac{m_l^2}{q^2} \right)^2 \left[E + \frac{m_l^2}{2q^2} F + \frac{3}{2} G + \frac{3m_l}{\sqrt{q^2}} H \right]. \quad (30)$$

Here $N = \frac{G_F^2 |V_{ub}|^2 q^2 |p_{\vec{p}}|}{512 \pi^3 m_{\Lambda_b}^2}$, $|p_{\vec{p}}| = \sqrt{\lambda(m_{\Lambda_b}^2, m_p^2, q^2)}/(2m_{\Lambda_b})$ with $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2(ab + bc + ca)$.

Further,

$$E = H_{\frac{1}{2}0}^2 + H_{-\frac{1}{2}0}^2 + H_{\frac{1}{2}1}^2 + H_{-\frac{1}{2}-1}^2, \quad (31)$$

$$F = H_{\frac{1}{2}0}^2 + H_{-\frac{1}{2}0}^2 + H_{\frac{1}{2}1}^2 + H_{-\frac{1}{2}-1}^2 + 3(H_{\frac{1}{2}t}^2 + H_{-\frac{1}{2}t}^2), \quad (32)$$

$$G = (H_{\frac{1}{2}0}^{SP})^2 + (H_{-\frac{1}{2}0}^{SP})^2, \quad (33)$$

$$H = H_{\frac{1}{2}t} H_{\frac{1}{2}0}^{SP} + H_{-\frac{1}{2}t} H_{-\frac{1}{2}0}^{SP}. \quad (34)$$

The differential branching fraction can then be written as

$$\frac{d\mathcal{B}(\Lambda_b \rightarrow pl\bar{\nu})}{dq^2} = \tau_{\Lambda_b} \frac{d\Gamma}{dq^2}. \quad (35)$$

One can also define the following LFU ratios of the differential branching fractions as

$$R_p(q^2) = \frac{d\Gamma(\Lambda_b \rightarrow p\tau\bar{\nu})/dq^2}{d\Gamma(\Lambda_b \rightarrow p\mu\bar{\nu})/dq^2}, \quad (36)$$

The lepton forward-backward asymmetry is defined as

$$A_{FB} = \frac{\int_0^1 (d^2\Gamma/dq^2 d\cos\theta) d\cos\theta - \int_{-1}^0 (d^2\Gamma/dq^2 d\cos\theta) d\cos\theta}{\int_0^1 (d^2\Gamma/dq^2 d\cos\theta) d\cos\theta + \int_{-1}^0 (d^2\Gamma/dq^2 d\cos\theta) d\cos\theta}. \quad (37)$$

Moreover, the longitudinal polarization of final state baryon and τ lepton is given by

$$P_p^L = \frac{d\Gamma^{\lambda_p=1/2}/dq^2 - d\Gamma^{\lambda_p=-1/2}/dq^2}{d\Gamma^{\lambda_p=1/2}/dq^2 + d\Gamma^{\lambda_p=-1/2}/dq^2} \quad (38)$$

$$P_\tau^L = \frac{d\Gamma^{\lambda_\tau=1/2}/dq^2 - d\Gamma^{\lambda_\tau=-1/2}/dq^2}{d\Gamma^{\lambda_\tau=1/2}/dq^2 + d\Gamma^{\lambda_\tau=-1/2}/dq^2} \quad (39)$$

The convexity parameter, which is the measure of curvature of the $\cos\theta$ distribution, is defined as

$$C_F^l(q^2) = \frac{1}{\int d\cos\theta W(\theta)} \frac{d^2W(\theta)}{d(\cos\theta)^2} \quad (40)$$

with

$$W(\theta) = \frac{3}{8} \left[A + \frac{m_l^2}{q^2} B + 2C + \frac{4m_l}{\sqrt{q^2}} D \right].$$

The helicity amplitudes defined in terms of the form factors are given in Appendix V A.

III. RESULTS AND DISCUSSIONS

A. Fit results

From Sec II A, it is apparent that in the context of U_1 LQ model, the WCs in $b \rightarrow u$ transition can be written in terms of $b \rightarrow c$ couplings. Therefore the observables in $b \rightarrow u$ sector are expected

Observable	Experimental Values
R_D	$0.340 \pm 0.027 \pm 0.013$ [44]
R_{D^*}	$0.295 \pm 0.011 \pm 0.008$ [44]
R_{Λ_c}	0.242 ± 0.026 (stat.) ± 0.040 (syst.) ± 0.059 [15]
$P_\tau^{D^*}$	$-0.38 \pm 0.51^{+0.21}_{-0.16}$ [5]
$f_L^{D^*}$	0.60 ± 0.08 (stat.) ± 0.04 (syst.) [18, 45]

TABLE I. Experimental values of observables used in the fit. The third error in R_{Λ_c} is due to the external branching fractions measurements.

to have strong correlations with $b \rightarrow c$ observables. In other words, the extent up to which the new physics effects can be generated in $b \rightarrow u$ observables would be determined by the allowed parameter space of couplings by the current $b \rightarrow c$ data. Given the fact that we have relatively large number of measured observables in this sector and moreover, some of them are accurately measured and predicted fairly well within the SM, it would be interesting to see possible deviation in $\Lambda_b \rightarrow p\tau\bar{\nu}$ observables allowed by the $b \rightarrow c$ data.

The theoretical expressions of observables R_D , R_{D^*} , R_{Λ_c} , $P_\tau^{D^*}$ and $f_L^{D^*}$ as functions of the relevant WCs are given in Sec. II B. By fitting these expressions to the measured values of the observables, we obtain the values of WCs which are consistent with the data. The corresponding χ^2 is defined as

$$\chi^2(C_i^{\text{eff}}) = \sum_{m,n=R_D,R_{D^*}} \left(O^{th}(C_i) - O^{exp} \right)_m (V^{exp} + V^{SM})_{mn}^{-1} \left(O^{th}(C_i) - O^{exp} \right)_n + \frac{(R_{\Lambda_c}^{th}(C_i) - R_{\Lambda_c}^{exp})^2}{\sigma_{R_{\Lambda_c}}^2} + \frac{(P_\tau^{D^* th}(C_i) - P_\tau^{D^* exp})^2}{\sigma_{P_\tau}^2} + \frac{(f_L^{D^* th}(C_i) - f_L^{D^* exp})^2}{\sigma_{f_L}^2}. \quad (41)$$

Here $O^{th}(C_i^{\text{eff}})$ are the theoretical predictions for R_D and R_{D^*} whereas $R_{\Lambda_c}^{th}$, $P_\tau^{D^* th}$ and $f_L^{D^* th}$ are theoretical expressions for R_{Λ_c} , $P_\tau^{D^*}$ and $f_L^{D^*}$, respectively. These expressions depend upon the new physics WCs C_{V_L} and C_{S_R} which in turn are functions of h_{23}^L , h_{33}^L and h_{33}^R couplings. O^{exp} are the corresponding experimental measurements. V^{exp} and V^{SM} are the experimental and SM covariance matrices in the R_D , R_{D^*} space, respectively. The matrix V^{exp} includes the correlation in the combined experimental determination of R_D and R_{D^*} . In eq. (41), $\sigma_{R_{\Lambda_c}}$, σ_{P_τ} and σ_{f_L} are the uncertainties in the measurements of R_{Λ_c} , $P_\tau^{D^*}$ and $f_L^{D^*}$, respectively. The measured values are given in Table I. The fit results are shown in Table II. It is evident that the SM doesn't provide a good fit to the data as $\chi_{\min}^2 \sim 26.06$ whereas for the U_1 LQ model, the fit is significantly improved as indicated by that the χ_{\min}^2 value which is ~ 6.64 . The best fit of the new physics couplings are also shown in the Table II. While obtaining allowed parameter space of new physics couplings we

	Best fit value(s)	χ_{\min}^2
SM	$C_i = 0$	26.06
U_1 LQ	$h_{33}^L = 1.2 \pm 2.3, h_{23}^L = 0.05 \pm 0.2, h_{33}^R = 0.1 \pm 0.3$	6.64

TABLE II. Best fit values of new physics couplings by making use of data of $R_D, R_{D^*}, R_{\Lambda_c}, P_\tau^{D^*}$ and $f_L^{D^*}$ in the fit.

imposed additional constraint of $B(B_c \rightarrow \tau \bar{\nu}) < 0.3$ [46] by lifetime of B_c meson.

Using the allowed values of new physics couplings obtained in this section, in the next subsection, we predict several observables in the decay of $\Lambda_b \rightarrow pl\bar{\nu}$.

B. Predictions

We consider following $\Lambda_b \rightarrow pl\bar{\nu}$ observables in our analysis:

- differential branching ratio dB/dq^2 , defined in eq. (35)
- LFU ratio R_p , defined in defined in eq. (36)
- longitudinal polarization of final state baryon, defined in eq. (39)
- longitudinal polarization of τ , defined in eq. (39)
- lepton forward-backward asymmetry A_{FB} , defined in eq. (37)
- convexity parameter C , defined in eq. (40).

The SM prediction of these observables along with 1σ upper and lower new physics bounds are illustrated in Fig. 1. From the left panel of top figure, it is luculent that the new physics can ameliorate the branching ratio by an order of magnitude. Even the 1σ new physics lower bound is about three times the SM 1σ upper limit. Thus the current $b \rightarrow c$ data does allow a large enhancement in the branching ratio of $\Lambda_b \rightarrow pl\bar{\nu}$. This roseate feature is also toted to the LFU ratio $R(\Lambda_p)$ as can be seen from the right panel of the top figure. Here too an order of magnitude enhancement is allowed. This is due to the fact that a large enhancement was viable for the differential branching ratio. Therefore $\Lambda_b \rightarrow pl\bar{\nu}$ decay mode can serve as an important channel to probe LFU violation in the $b \rightarrow u$ sector.

The predictions of the longitudinal polarization of the final state baryon as well as the τ lepton in $\Lambda_b \rightarrow pl\bar{\nu}$ decay are shown in the left and right panels of Fig. 1. For $q^2 > 17 \text{ GeV}^2$, $P_p^L(q^2)$ is




FIG. 1. Predictions for various observables in $\Lambda_b \rightarrow pl\bar{\nu}$ decay. The band corresponds to the SM uncertainties.

consistent with the SM prediction whereas for $q^2 < 17 \text{ GeV}^2$, there is marginal deviation from the SM prediction. The deviation is more prominent for lower values of q^2 . On the other hand, the predictions of tau polarization is consistent with the SM prediction in the entire q^2 region. The same is true for lepton forward backward asymmetry and convexity parameter as can be seen from the left and right panels of bottom figure, respectively.

IV. CONCLUSIONS

In this work we anatomize new physics effects in $\Lambda_b \rightarrow p\tau\bar{\nu}$ decay in U_1 leptoquark model. This decay mode is induced by the quark level transition $b \rightarrow u\tau\bar{\nu}$. A model independent analysis of new physics in $b \rightarrow u\tau\bar{\nu}$ can lead to large effects due to the fact that, as of now, we only have one measurement in this sector. However, in the context of U_1 leptoquark model considered in this work, the new physics couplings in $b \rightarrow u\tau\bar{\nu}$ transition can be expressed in terms of couplings in $b \rightarrow c\tau\bar{\nu}$ decay along with a suitable combinations of elements of the CKM matrix. Therefore, one expects a strong correlations between these two sectors. Given the fact that, unlike $b \rightarrow u\tau\bar{\nu}$ sector, there are measurements of a number of observables in decays induced by $b \rightarrow c\tau\bar{\nu}$ transition, one expects that meaningful constraints on new physics parameter space can be obtained. It would then be interesting to see whether such constraints can allow for large enhancements in some of the observables in $\Lambda_b \rightarrow p\tau\bar{\nu}$ decay.

In order to obtain constraints on new physics couplings, we perform a fit to all $b \rightarrow c\tau\bar{\nu}$ data. For allowed parameter space of the couplings, we obtain predictions of the branching ratio, LFU ratio, the longitudinal polarization of final state baryon and τ lepton, lepton forward-backward asymmetry and in the decay of $\Lambda_b \rightarrow p\tau\bar{\nu}$. We find that

- The branching ratio as well as the LFU ratio can be enhanced by an order of magnitude over the SM value.
- There can be a marginal deviation from the SM in the longitudinal polarization of final state baryon in the low- q^2 region.
- The longitudinal polarization of τ , lepton forward-backward asymmetry as well as the convexity parameter are consistent with the SM.

Acknowledgements: The work of DK is supported by the SERB, India under the research grant no. SERB/EEQ/2021/000965.

V. APPENDIX

A. $\Lambda_b \rightarrow p$ transition form factors and Helicity amplitudes

The q^2 dependence of the helicity form factors in the lattice QCD calculations are defined as[47]:

$$f_i(q^2) = \frac{1}{1 - q^2/(m_{pole}^f)^2} [a_0^f + a_1^f z(q^2)], \quad (42)$$

where $i = +, \perp, 0$ and the expansion parameter is defined as

$$z(q^2) = \frac{\sqrt{t_+ - q^2} - \sqrt{t_+ - t_0}}{\sqrt{t_+ - q^2} + \sqrt{t_+ - t_0}}. \quad (43)$$

Here $t_+ = (m_{B_1} + m_{B_2})^2$ and $t_0 = (m_{B_1} - m_{B_2})^2$. The nominal form factor parameters $a_{0,1}^{f(g)}$ and m_{pole}^f for $\Lambda_b \rightarrow p$ are taken from [47].

The decay $\Lambda_b \rightarrow p l \bar{\nu}$ is considered to be through $\Lambda_b \rightarrow p W^*$ and the off-shell W^* decays to $l \bar{\nu}$. The helicity amplitudes for vector and axial-vector type current is defined by

$$H_{\lambda_p, \lambda_W} = H_{\lambda_p, \lambda_W}^V - H_{\lambda_p, \lambda_W}^A, \quad (44)$$

$$H_{\lambda_p, \lambda_W}^V = \epsilon^{\dagger\mu}(\lambda_W) \langle p, \lambda_p | \bar{c} \gamma_\mu b | \Lambda_b, \lambda_{\Lambda_b} \rangle, \quad (45)$$

$$H_{\lambda_p, \lambda_W}^A = \epsilon^{\dagger\mu}(\lambda_W) \langle p, \lambda_p | \bar{c} \gamma_\mu \gamma_5 b | \Lambda_b, \lambda_{\Lambda_b} \rangle. \quad (46)$$

Also, the helicity amplitudes for scalar and pseudo-scalar current is given by

$$H_{\lambda_p}^S = \langle p, \lambda_p | \bar{c} b | \Lambda_b, \lambda_{\Lambda_b} \rangle, \quad (47)$$

$$H_{\lambda_p}^P = \langle p, \lambda_p | \bar{c} \gamma_5 b | \Lambda_b, \lambda_{\Lambda_b} \rangle. \quad (48)$$

One can show from the parity argument or explicit calculation that $H_{-\lambda_p, -\lambda_W}^V = H_{\lambda_p, \lambda_W}^V$, $H_{-\lambda_p, -\lambda_W}^A = -H_{\lambda_p, \lambda_W}^A$, $H_{\lambda_p, \lambda_{NP}}^S = H_{-\lambda_p, -\lambda_{NP}}^S$ and $H_{\lambda_p, \lambda_{NP}}^P = -H_{-\lambda_p, -\lambda_{NP}}^P$.

The helicity amplitudes can be defined in terms of the helicity form factors as[43]:

$$H_{\frac{1}{2}, 0}^V = (1 + C_{V_L} + C_{V_R}) \frac{\sqrt{Q_-}}{\sqrt{q^2}} (m_{B_1} + m_{B_2}) f_+(q^2), \quad (49)$$

$$H_{\frac{1}{2}, 0}^A = (1 + C_{V_L} - C_{V_R}) \frac{\sqrt{Q_+}}{\sqrt{q^2}} (m_{B_1} - m_{B_2}) g_+(q^2), \quad (50)$$

$$H_{\frac{1}{2}, 1}^V = -(1 + C_{V_L} + C_{V_R}) \sqrt{2Q_-} f_\perp(q^2), \quad (51)$$

$$H_{\frac{1}{2}, 1}^A = -(1 + C_{V_L} - C_{V_R}) \sqrt{2Q_+} g_\perp(q^2), \quad (52)$$

$$H_{\frac{1}{2}, t}^V = (1 + C_{V_L} + C_{V_R}) \frac{\sqrt{Q_+}}{\sqrt{q^2}} (m_{B_1} - m_{B_2}) f_0(q^2), \quad (53)$$

$$H_{\frac{1}{2}, t}^A = (1 + C_{V_L} - C_{V_R}) \frac{\sqrt{Q_-}}{\sqrt{q^2}} (m_{B_1} + m_{B_2}) g_0(q^2), \quad (54)$$

where

$$Q_{\pm} = (m_{B_1} \pm m_{B_2})^2 - q^2. \quad (55)$$

The scalar and pseudo-scalar helicity amplitudes are defined as:

$$H_{\frac{1}{2},0}^{SP} = H_{\frac{1}{2},0}^S - H_{\frac{1}{2},0}^P, \quad (56)$$

$$H_{\frac{1}{2},0}^S = (C_{S_L} + C_{S_R}) \frac{\sqrt{Q_+}}{m_b - m_u} (m_{B_1} - m_{B_2}) f_0(q^2), \quad (57)$$

$$H_{\frac{1}{2},0}^P = (C_{S_L} - C_{S_R}) \frac{\sqrt{Q_-}}{m_b + m_u} (m_{B_1} + m_{B_2}) g_0(q^2). \quad (58)$$

The helicity-dependent differential decay rates are required to compute the longitudinal polarization asymmetry of final state baryon and τ and these decay rates are defined as

$$\frac{d\Gamma^{\lambda_p = \frac{1}{2}}}{dq^2} = \frac{m_l^2}{q^2} \left[\frac{4}{3} \left(H_{\frac{1}{2},1}^2 + H_{\frac{1}{2},0}^2 + 3H_{\frac{1}{2},t}^2 \right) \right] + \frac{8}{3} \left(H_{\frac{1}{2},0}^2 + H_{\frac{1}{2},1}^2 \right) + 4H_{\frac{1}{2},0}^{SP^2} + \frac{8m_l}{\sqrt{q^2}} H_{\frac{1}{2},t} H_{\frac{1}{2},0}^{SP} \quad (59)$$

$$\begin{aligned} \frac{d\Gamma^{\lambda_p = -\frac{1}{2}}}{dq^2} &= \frac{m_l^2}{q^2} \left[\frac{4}{3} \left(H_{-\frac{1}{2},1}^2 + H_{-\frac{1}{2},0}^2 + 3H_{-\frac{1}{2},t}^2 \right) \right] + \frac{8}{3} \left(H_{-\frac{1}{2},0}^2 + H_{-\frac{1}{2},-1}^2 \right) + 4H_{-\frac{1}{2},0}^{SP^2} \\ &\quad + \frac{8m_l}{\sqrt{q^2}} H_{-\frac{1}{2},t} H_{-\frac{1}{2},0}^{SP} \end{aligned} \quad (60)$$

$$\begin{aligned} \frac{d\Gamma^{\lambda_\tau = \frac{1}{2}}}{dq^2} &= \frac{m_l^2}{q^2} \left[\frac{4}{3} \left(H_{\frac{1}{2},1}^2 + H_{\frac{1}{2},0}^2 + H_{-\frac{1}{2},-1}^2 + H_{-\frac{1}{2},0}^2 \right) + 4 \left(H_{\frac{1}{2},t}^2 + H_{-\frac{1}{2},t}^2 \right) \right] + 4 \left(H_{\frac{1}{2},0}^{SP^2} + H_{-\frac{1}{2},0}^{SP^2} \right) \\ &\quad + \frac{8m_l}{\sqrt{q^2}} \left(H_{\frac{1}{2},t} H_{\frac{1}{2},0}^{SP} + H_{-\frac{1}{2},t} H_{-\frac{1}{2},0}^{SP} \right) \end{aligned} \quad (61)$$

$$\frac{d\Gamma^{\lambda_\tau = -\frac{1}{2}}}{dq^2} = \frac{8}{3} \left(H_{\frac{1}{2},1}^2 + H_{\frac{1}{2},0}^2 + H_{-\frac{1}{2},-1}^2 + H_{-\frac{1}{2},0}^2 \right) \quad (62)$$

- [1] J. P. Lees *et al.* [BaBar Collaboration], “Evidence for an excess of $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$ decays”, Phys. Rev. Lett. **109**, 101802 (2012) [arXiv:1205.5442 [hep-ex]].
- [2] J. P. Lees *et al.* [BaBar Collaboration], “Measurement of an Excess of $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$ Decays and Implications for Charged Higgs Bosons”, Phys. Rev. D **88**, no. 7, 072012 (2013) [arXiv:1303.0571 [hep-ex]].
- [3] M. Huschle *et al.* [Belle Collaboration], “Measurement of the branching ratio of $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$ relative to $\bar{B} \rightarrow D^{(*)} \ell^- \bar{\nu}_\ell$ decays with hadronic tagging at Belle”, Phys. Rev. D **92**, no. 7, 072014 (2015) [arXiv:1507.03233 [hep-ex]].
- [4] Y. Sato *et al.* [Belle Collaboration], “Measurement of the branching ratio of $\bar{B}^0 \rightarrow D^{*+} \tau^- \bar{\nu}_\tau$ relative to $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}_\ell$ decays with a semileptonic tagging method”, Phys. Rev. D **94**, no. 7, 072007 (2016) [arXiv:1607.07923 [hep-ex]].

- [5] S. Hirose *et al.* [Belle Collaboration], “Measurement of the τ lepton polarization and $R(D^*)$ in the decay $\bar{B} \rightarrow D^* \tau^- \bar{\nu}_\tau$ ”, Phys. Rev. Lett. **118**, no. 21, 211801 (2017) [arXiv:1612.00529 [hep-ex]].
- [6] R. Aaij *et al.* [LHCb Collaboration], “Measurement of the ratio of branching fractions $\mathcal{B}(\bar{B}^0 \rightarrow D^{*+} \tau^- \bar{\nu}_\tau) / \mathcal{B}(\bar{B}^0 \rightarrow D^{*+} \mu^- \bar{\nu}_\mu)$ ”, Phys. Rev. Lett. **115**, no. 11, 111803 (2015) [Phys. Rev. Lett. **115**, no. 15, 159901 (2015)] [arXiv:1506.08614 [hep-ex]].
- [7] <https://hflav-eos.web.cern.ch/hflav-eos/semi/summer16/html/RDsDsstar/RDRDs.html>
- [8] R. Aaij *et al.* [LHCb Collaboration], “Measurement of the ratio of the $B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$ and $B^0 \rightarrow D^{*-} \mu^+ \nu_\mu$ branching fractions using three-prong τ -lepton decays”, Phys. Rev. Lett. **120**, no. 17, 171802 (2018) [arXiv:1708.08856 [hep-ex]].
- [9] <http://www.slac.stanford.edu/xorg/hfag/semi/fpcp17/RDRDs.html>
- [10] A. Abdesselam *et al.* [Belle Collaboration], “Measurement of $\mathcal{R}(D)$ and $\mathcal{R}(D^*)$ with a semileptonic tagging method”, arXiv:1904.08794 [hep-ex].
- [11] R. Aaij *et al.* [LHCb Collaboration], “Measurement of the ratio of branching fractions $\mathcal{B}(B_c^+ \rightarrow J/\psi \tau^+ \nu_\tau) / \mathcal{B}(B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu)$ ”, Phys. Rev. Lett. **120** (2018) no.12, 121801 [arXiv:1711.05623 [hep-ex]].
- [12] A. K. Alok, D. Kumar, J. Kumar, S. Kumbhakar and S. U. Sankar, JHEP **09**, 152 (2018) [arXiv:1710.04127 [hep-ph]].
- [13] R. Glattauer *et al.* [Belle Collaboration], “Measurement of the decay $B \rightarrow D \ell \nu_\ell$ in fully reconstructed events and determination of the Cabibbo-Kobayashi-Maskawa matrix element $|V_{cb}|$ ”, Phys. Rev. D **93**, no. 3, 032006 (2016) [arXiv:1510.03657 [hep-ex]].
- [14] A. Abdesselam *et al.* [Belle Collaboration], “Precise determination of the CKM matrix element $|V_{cb}|$ with $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}_\ell$ decays with hadronic tagging at Belle”, arXiv:1702.01521 [hep-ex].
- [15] R. Aaij *et al.* [LHCb], “Observation of the decay $\Lambda_b^0 \rightarrow \Lambda_c^+ \tau^- \bar{\nu}_\tau$ ”, Phys. Rev. Lett. **128** (2022) no.19, 191803 [arXiv:2201.03497 [hep-ex]].
- [16] F. U. Bernlochner, Z. Ligeti, D. J. Robinson and W. L. Sutcliffe, “Precise predictions for $\Lambda_b \rightarrow \Lambda_c$ semileptonic decays,” Phys. Rev. D **99**, no.5, 055008 (2019) [arXiv:1812.07593 [hep-ph]].
- [17] M. Tanaka and R. Watanabe, “New physics in the weak interaction of $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}$,” Phys. Rev. D **87** (2013) no.3, 034028 [arXiv:1212.1878 [hep-ph]].
- [18] A. Abdesselam *et al.* [Belle Collaboration], “Measurement of the D^{*-} polarization in the decay $B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$,” arXiv:1903.03102 [hep-ex].
- [19] A. K. Alok, D. Kumar, S. Kumbhakar and S. U. Sankar, “ D^* polarization as a probe to discriminate new physics in $\bar{B} \rightarrow D^* \tau \bar{\nu}$,” Phys. Rev. D **95**, no. 11, 115038 (2017) [arXiv:1606.03164 [hep-ph]].
- [20] M. Freytsis, Z. Ligeti and J. T. Ruderman, “Flavor models for $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}$,” Phys. Rev. D **92**, no. 5, 054018 (2015) [arXiv:1506.08896 [hep-ph]].
- [21] M. Jung and D. M. Straub, “Constraining new physics in $b \rightarrow c \ell \nu$ transitions,” JHEP **01** (2019), 009 [arXiv:1801.01112 [hep-ph]].
- [22] S. Bhattacharya, S. Nandi and S. Kumar Patra, “ $b \rightarrow c \tau \nu_\tau$ Decays: a catalogue to compare, constrain,

- and correlate new physics effects,” *Eur. Phys. J. C* **79** (2019) no.3, 268 [arXiv:1805.08222 [hep-ph]].
- [23] Q. Y. Hu, X. Q. Li and Y. D. Yang, “ $b \rightarrow c\tau\nu$ transitions in the standard model effective field theory,” *Eur. Phys. J. C* **79** (2019) no.3, 264 [arXiv:1810.04939 [hep-ph]].
- [24] A. K. Alok, D. Kumar, S. Kumbhakar and S. Uma Sankar, “Solutions to $R_D-R_{D^*}$ in light of Belle 2019 data,” *Nucl. Phys. B* **953** (2020), 114957 [arXiv:1903.10486 [hep-ph]].
- [25] P. Asadi and D. Shih, “Maximizing the Impact of New Physics in $b \rightarrow c\tau\nu$ Anomalies,” *Phys. Rev. D* **100** (2019) no.11, 115013 [arXiv:1905.03311 [hep-ph]].
- [26] C. Murgui, A. Peñuelas, M. Jung and A. Pich, “Global fit to $b \rightarrow c\tau\nu$ transitions,” *JHEP* **09** (2019), 103 [arXiv:1904.09311 [hep-ph]].
- [27] D. Bardhan and D. Ghosh, “ B -meson charged current anomalies: The post-Moriond 2019 status,” *Phys. Rev. D* **100** (2019) no.1, 011701 [arXiv:1904.10432 [hep-ph]].
- [28] M. Blanke, A. Crivellin, T. Kitahara, M. Moscati, U. Nierste and I. Nišandžić, “Addendum to “Impact of polarization observables and $B_c \rightarrow \tau\nu$ on new physics explanations of the $b \rightarrow c\tau\nu$ anomaly”,” *Phys. Rev. D* **100** (2019) no.3, 035035 [arXiv:1905.08253 [hep-ph]].
- [29] R. X. Shi, L. S. Geng, B. Grinstein, S. Jäger and J. Martin Camalich, “Revisiting the new-physics interpretation of the $b \rightarrow c\tau\nu$ data,” *JHEP* **12** (2019), 065 [arXiv:1905.08498 [hep-ph]].
- [30] D. Bečirević, M. Fedele, I. Nišandžić and A. Tayduganov, “Lepton Flavor Universality tests through angular observables of $\bar{B} \rightarrow D^{(*)}\ell\bar{\nu}$ decay modes,” [arXiv:1907.02257 [hep-ph]].
- [31] S. Sahoo and R. Mohanta, “Investigating the role of new physics in $b \rightarrow c\tau\bar{\nu}_\tau$ transitions,” [arXiv:1910.09269 [hep-ph]].
- [32] K. Cheung, Z. R. Huang, H. D. Li, C. D. Lu, Y. N. Mao and R. Y. Tang, “Revisit to the $b \rightarrow c\tau\nu$ transition: in and beyond the SM,” [arXiv:2002.07272 [hep-ph]].
- [33] J. Cardozo, J. H. Muñoz, N. Quintero and E. Rojas, “Analysing the charged scalar boson contribution to the charged-current B meson anomalies,” [arXiv:2006.07751 [hep-ph]].
- [34] A. Carvunis, A. Crivellin, D. Guadagnoli and S. Gangal, “The Forward-Backward Asymmetry in $B \rightarrow D^*\ell\nu$: One more hint for Scalar Leptoquarks?,” *Phys. Rev. D* **105**, no.3, L031701 (2022) [arXiv:2106.09610 [hep-ph]].
- [35] A. K. Alok, D. Kumar, S. Kumbhakar and S. Uma Sankar, “Resolution of R_D/R_{D^*} puzzle,” *Phys. Lett. B* **784**, 16-20 (2018) [arXiv:1804.08078 [hep-ph]].
- [36] R. Dutta, *Phys. Rev. D* **93** (2016) no.5, 054003 doi:10.1103/PhysRevD.93.054003 [arXiv:1512.04034 [hep-ph]].
- [37] A. Ray, S. Sahoo and R. Mohanta, *Phys. Rev. D* **99** (2019) no.1, 015015 doi:10.1103/PhysRevD.99.015015 [arXiv:1812.08314 [hep-ph]].
- [38] R. L. Workman [Particle Data Group], *PTEP* **2022**, 083C01 (2022)
- [39] P. Hamer *et al.* [Belle], “Search for $B^0 \rightarrow \pi^- \tau^+ \nu_\tau$ with hadronic tagging at Belle,” *Phys. Rev. D* **93**, no.3, 032007 (2016) [arXiv:1509.06521 [hep-ex]].
- [40] I. Doršner, S. Fajfer, A. Greljo, J. F. Kamenik and N. Košnik, “Physics of leptoquarks in precision

- experiments and at particle colliders,” Phys. Rept. **641**, 1-68 (2016) [arXiv:1603.04993 [hep-ph]].
- [41] A. Bhaskar, D. Das, T. Mandal, S. Mitra and C. Neeraj, “Precise limits on the charge-2/3 U1 vector leptoquark,” Phys. Rev. D **104**, no.3, 035016 (2021) [arXiv:2101.12069 [hep-ph]].
- [42] M. Blanke, A. Crivellin, S. de Boer, T. Kitahara, M. Moscati, U. Nierste and I. Nišandžić, “Impact of polarization observables and $B_c \rightarrow \tau\nu$ on new physics explanations of the $b \rightarrow c\tau\nu$ anomaly,” Phys. Rev. D **99**, no.7, 075006 (2019) [arXiv:1811.09603 [hep-ph]].
- [43] S. Shivashankara, W. Wu and A. Datta, “ $\Lambda_b \rightarrow \Lambda_c \tau \bar{\nu}_\tau$ Decay in the Standard Model and with New Physics,” Phys. Rev. D **91** (2015) no.11, 115003 doi:10.1103/PhysRevD.91.115003 [arXiv:1502.07230 [hep-ph]].
- [44] <https://hflav-eos.web.cern.ch/hflav-eos/semi/spring19/html/RDsDsstar/RDRDs.html>
- [45] K. Adamczyk [Belle and Belle-II], “Semitauonic B decays at Belle/Belle II,” [arXiv:1901.06380 [hep-ex]].
- [46] R. Alonso, B. Grinstein and J. Martin Camalich, Phys. Rev. Lett. **118** (2017) no.8, 081802 doi:10.1103/PhysRevLett.118.081802 [arXiv:1611.06676 [hep-ph]].
- [47] W. Detmold, C. Lehner and S. Meinel, “ $\Lambda_b \rightarrow p \ell^- \bar{\nu}_\ell$ and $\Lambda_b \rightarrow \Lambda_c \ell^- \bar{\nu}_\ell$ form factors from lattice QCD with relativistic heavy quarks,” Phys. Rev. D **92** (2015) no.3, 034503 doi:10.1103/PhysRevD.92.034503 [arXiv:1503.01421 [hep-lat]].