

Banner appropriate to article type will appear here in typeset article

The high-Reynolds-number stratified wake of a slender body and its comparison with a bluff-body wake

Jose L. Ortiz-Tarin¹, Sheel Nidhan¹, Sutanu Sarkar¹†

¹Department of Mechanical and Aerospace Engineering, University of California San Diego, CA 92093, USA

(Received xx; revised xx; accepted xx)

The high-Reynolds number stratified wake of a slender body is studied using a high-resolution hybrid simulation. The wake generator is a 6:1 prolate spheroid with a tripped boundary layer, the diameter-based body Reynolds number is $Re = U_\infty D/\nu = 10^5$, and the body Froude numbers are $Fr = U_\infty/ND = \{2, 10, \infty\}$. The wake defect velocity (U_d) decays following three stages with different wake decay rates (Spedding 1997) as for a bluff body. However, the transition points among stages do not follow the expected $Nt = Nx/U_\infty$ values. Comparison with the wake of a circular disk in similar conditions (Chongsiripinyo & Sarkar 2020) quantifies the influence of the wake generator - bluff versus slender - in stratified flow. The strongly stratified $Fr = 2$ wake is in a resonant state. The steady lee waves strongly modulate the mean flow and, relative to the disk, the 6:1 spheroid (a high aspect ratio shape) wake at $Fr = 2$ shows an earlier transition from the non-equilibrium (NEQ) stage to the quasi two-dimensional (Q2D) stage. The NEQ-Q2D transition is followed by a sharp increase in the turbulent kinetic energy and horizontal wake meanders. At $Fr = 10$, the start of the NEQ stage is delayed for the spheroid. Transfers between kinetic energy and potential energy reservoirs (both mean and turbulence) are analyzed and the flows are compared in phase space (local Froude and Reynolds number as coordinates). Overall, the results of this study point to the difficulty of finding a universal framework for stratified wake evolution, independent of the features of the body, and provide insights into how buoyancy effects depend on the wake generator.

1. Introduction

Due to their low drag coefficients, slender bodies are extensively used in aerospace and naval applications. Multiple studies have described the flow around these bodies focusing on the drag force, the boundary layer, and the flow separation (Wang 1970; Costis *et al.* 1989; Wang *et al.* 1990; Chesnakas & Simpson 1994; Fu *et al.* 1994; Constantinescu *et al.* 2002; Wikström *et al.* 2004). However, despite their presence in many underwater applications, only a few works have looked into the wake of a slender body (Chevray 1968; Jiménez *et al.* 2010; Kumar & Mahesh 2018) and, only recently, the far wake of a slender body has been studied (Ortiz-Tarin *et al.* 2021).

The near wake of a slender body with a turbulent boundary layer (TBL) is characterized

† Email address for correspondence: sarkar@ucsd.edu

by having a small recirculation region. The recirculation region is surrounded by a ring of small scale turbulence that emerges from the boundary layer and does not show strong vortex shedding (Jiménez *et al.* 2010; Posa & Balaras 2016; Kumar & Mahesh 2018; Ortiz-Tarin *et al.* 2021). As a result, the wake is thin and develops slowly compared to the wake of bluff bodies. These particular features of the slender body high- Re near wake lead to interesting effects further downstream: (i) despite having a smaller drag coefficient than bluff bodies, the defect velocity ($U_d = U_\infty - U$) of the slender body wake can be larger than that of a bluff body for a long downstream distance, (ii) the turbulent kinetic energy of the wake shows an off-center radial peak at the location where the turbulent boundary layer separates – instead of a Gaussian profile with a central peak, and (iii) helical instabilities come into play only in the intermediate and far field of the wake. These particularities affect the scaling laws of the wake. In a domain spanning $80D$ the defect velocity, the kinetic energy, and the dissipation do not follow the classic high- Re scaling and they decay differently than bluff body wakes (Ortiz-Tarin *et al.* 2021) exhibiting a non-equilibrium scaling of dissipation (Vassilicos 2015; Dairay *et al.* 2015).

The few studies that look into slender body wakes assume that the body moves in an unstratified environment, where the density of the surrounding fluid is constant. However, in a realistic underwater marine environment the effect of density stratification due to salinity and temperature can become relevant. Density stratification suppresses vertical motions, triggers the formation and sustenance of coherent structures, and leads to the radiation of internal gravity waves. More importantly, in a stratified environment, the wake of a submersible lives longer than in an unstratified environment, i.e., it takes more time for the flow disturbance to die out (Spedding 2014). The study of stratified wakes has been nearly exclusively focused on the flow past bluff bodies (Lin & Pao 1979; Hanazaki 1988; Lin *et al.* 1992; Chomaz *et al.* 1992; Orr *et al.* 2015; Pal *et al.* 2017) and underwater topography (Drazin 1961; Castro *et al.* 1983; Baines 1998). Here, we study the influence of stratification on the high- Re wake of a prolate spheroid with a turbulent boundary layer.

The strength of ambient stratification is measured by the body-based Froude number $Fr = U_\infty / ND$. This is the ratio between the convective frequency of the flow, U_∞ / D – where U_∞ is the freestream velocity and D the diameter of the body – and the buoyancy frequency N . In the wake of ocean submersibles, $Fr \sim O(1 - 10^2)$. However, since the velocity deficit in the wake $U_d(x)$ decays with the streamwise distance and the wake width $L(x)$ increases, the Froude number defined with local variables $Fr_l = U_d / NL$ decreases as the flow evolves. Thus, even in a weakly stratified environment, eventually all wakes are affected by stratification.

Since the relative strength of stratification increases locally as the flow develops, the evolution of the stratified wake is multistage in nature. Based on the measurements of U_d and L in high- Fr (i.e. initially weak stratification) bluff body wakes, Spedding (1997) identified three regimes in stratified wake evolution based on the power-law decay rates of U_d . These regimes are generally identified by empirically fitting decay rates to U_d in different temporal (or spatial) regions and are as follows:

(i) Three-dimensional (3D) regime: Close to the generator, wake decay is similar to the unstratified wake of the corresponding body shape. This is the so-called 3D regime and lasts until the buoyancy time defined by $Nt = Nx/U = x/D \cdot 1/Fr$ approaches $O(1)$, equivalently until $x/D \sim Fr$.

(ii) Non-equilibrium (NEQ) regime: As the wake evolves, buoyancy effects become progressively stronger. The decay of U_d slows relative to the 3D regime and, furthermore, anisotropy between the vertical and horizontal velocity components increases. Spedding (1997) reported this non-equilibrium (NEQ) region to last for $Nt \approx 2 - 50$. Later, temporal simulations of Brucker & Sarkar (2010) and Diamessis *et al.* (2011) found an increase

in the span of the NEQ regime at higher Reynolds numbers. $U_d \sim x^{-0.25 \pm 0.04}$ during the NEQ regime according to Spedding (1997). However, there has been some variability in the observed NEQ decay rate in later studies. Bonnier & Eiff (2002) reported a NEQ regime with $U_d \sim x^{-0.38}$ for $1.5 < Fr < 5$. Brucker & Sarkar (2010) and Diamessis *et al.* (2011) reported $U_d \sim Nt^{-1/4}$ during the NEQ regime in their temporal simulations. Chongsiripinyo & Sarkar (2020) found that $U_d \sim x^{-0.18}$ during the NEQ regime of their $Fr = 2$ and 10 disk wakes. For wakes with $Fr \sim O(1)$, the NEQ decay rate is preceded by a pronounced oscillatory modulation in U_d which is linked to lee waves (Pal *et al.* 2017; Chongsiripinyo & Sarkar 2020; Ortiz-Tarin *et al.* 2019).

(iii) Quasi two-dimensional (Q2D) regime: After the NEQ regime, the stratified wake enters into the quasi two-dimensional regime (Q2D regime). The Q2D regime is characterized by transition in the U_d power law to a significantly increased decay rate, e.g. Spedding (1997) reports a transition to $U_d \sim x^{-3/4}$. In the Q2D regime, the wake progressively organizes into vortices that meander primarily in the horizontal plane (Gourlay *et al.* 2001; Dommermuth *et al.* 2002; Brucker & Sarkar 2010) and take the form of ‘pancakes’. Although the wake motion in this regime is primarily in the horizontal plane, there is a variability in the vertical direction in the form of layers (Spedding 2002), hence the prefix ‘quasi’.

In recent literature, stratified wakes have been characterized using turbulence features (Zhou & Diamessis 2019; Chongsiripinyo & Sarkar 2020) instead of the U_d -based criteria of Spedding (1997). These studies are motivated by an attempt to connect buoyancy-related wake transitions to the broader stratified turbulence field.

Notice that the arrival of the wake into each of the three stages in its evolution depends on the value of Nt , which is equivalent to a downstream distance of x/Fr from the wake generator. At high Froude number, the downstream distance required to reach the NEQ and Q2D regions can become very large. Consequently, the size of the computational domain required to access these regimes rapidly becomes computationally unfeasible. To circumvent these limitation temporal simulations were used in the study of Gourlay *et al.* (2001). Temporal simulations use a reference frame moving with the wake where time correlates with streamwise distance in a fixed reference frame. By assuming that the streamwise development of the flow is slow, periodic boundary conditions can be used and the equations are advanced in time without the need of introducing the wake generator. This reduces the computational cost significantly. Most of the studies that have contributed to our current understanding of stratified wakes use temporal simulations (Gourlay *et al.* 2001; Dommermuth *et al.* 2002; Brucker & Sarkar 2010; Diamessis *et al.* 2011; De Stadler & Sarkar 2012; Abdilghanie & Diamessis 2013; Redford *et al.* 2015; Rowe *et al.* 2020).

The main drawback of the temporal model is the influence of its initialization. Since the flow at the wake generator is not solved, the starting profiles of the mean and turbulence have to be assumed. These simulations lack some specific features that are generated due to the body, e.g., steady lee waves, near-wake buoyancy effects, and the vortical structures shed from the boundary layer. Even when it is tempting to assume that body specific features are lost far from the body, the universality of the wake decay has remained elusive to experiments (Bevilaqua & Lykoudis 1978; Wagnanski *et al.* 1986; Redford *et al.* 2012), even in unstratified wakes. An alternative to temporal simulations are body inclusive simulations that retain the wake generator dependent features at the expense of a higher computational cost and a limited domain size (Orr *et al.* 2015; Chongsiripinyo *et al.* 2017; Pal *et al.* 2017; Nidhan *et al.* 2019; More *et al.* 2021).

To the best of authors’ knowledge, Ortiz-Tarin *et al.* (2019) performed the first study of a stratified flow past a slender body that investigates the near and intermediate wake dynamics. Their analyses reveal that at $Fr \sim O(1)$ the type of separation and the subsequent wake establishment is strongly dependent on the characteristic frequency of the lee waves and

the aspect ratio of the body. When half the wavelength of the steady lee waves ($\lambda = 2\pi Fr$) matches the length of the slender body (L), the separation of the boundary layer is inhibited by buoyancy effects. Based on this condition, a critical Froude number can be defined $Fr_c = L/D\pi$. When $Fr > Fr_c$ stratification suppresses the generation of turbulence in the near wake, when $Fr \approx Fr_c$ buoyancy strongly limits the flow separation and can lead to a relaminarization of the wake at low Reynolds numbers. Finally, when $Fr < Fr_c$, the lee waves enlarge the separation region and there might be an increase in the turbulence intensities in the wake. When $Fr \approx Fr_c$ the wake is in a resonant state with both the separation and the wake dimensions being strongly modulated by the steady lee waves (Hunt & Snyder 1980; Chomaz *et al.* 1993; Ortiz-Tarin *et al.* 2019).

As mentioned before, the use of body inclusive simulations has one major limitation, i.e., the high computational cost. Due to the high resolution required to solve the boundary layer of the wake generator, the downstream domain is limited and thus the possibility of looking into the far wake gets significantly restricted, particularly at high Re . VanDine *et al.* (2018) presented a hybrid spatially-evolving model, which builds on the hybrid temporally-evolving model of Pasquetti (2011), and addresses most of the aforementioned problems. The hybrid method uses inflow conditions generated from a well-resolved body-inclusive simulation to perform a separate temporal simulation in the case of Pasquetti (2011) or spatially-evolving simulation in the work of VanDine *et al.* (2018) without including the body. By doing so, the amount of required points is substantially reduced since the flow near the body does not have to be resolved. This important reduction of the computational cost allows one to extend the domain farther downstream to gain insight in the far wake.

Here, we use a hybrid method that combines a body-inclusive simulation and a spatially evolving body-exclusive simulation to study the stratified high- Re far wake of a slender body for the first time. The Reynolds number is set to $Re = U_\infty D/\nu = 10^5$ and two levels of stratification are used, $Fr = U_\infty/ND = 2$ and 10. The simulation at $Fr = 10$ allows us to study the evolution of a weakly stratified wake in a domain that spans $80D$. Additionally, $Fr = 2$ is chosen because it is close to the critical Froude number for a 6:1 prolate spheroid $Fr_c = (L/D)/\pi = 6/\pi$. At the critical Froude number, the size of the separation region is strongly reduced by the lee waves (Ortiz-Tarin *et al.* 2019). These choices also allow us to compare our results with the findings of Chongsiripinyo & Sarkar (2020) (hereafter referred as CS20) regarding the stratified wake of a disk.

In CS20, the stratified wake of a disk at $Re = 5 \times 10^4$ is studied at $Fr = 2, 10, 50, \infty$. Apart from a detailed analysis of the decay rates of the mean and turbulent quantities, CS20 links the general evolution of stratified homogeneous turbulence (Brethouwer *et al.* 2007; de Bruyn Kops & Riley 2019) with the evolution of the wake turbulence. As the disk wake evolves, the influence of buoyancy is ‘felt’ by the turbulent motions at progressively smaller scales. First the mean flow and the large scales and later the r.m.s. velocities are affected by stratification. Simultaneously, the horizontal eddies start gaining energy. Based on the strength of these effects, three distinct stages can be identified: weakly, intermediate and strongly stratified turbulence. In CS20, the transition between these regimes is examined and parameterized using local Froude and Reynolds numbers. Zhou & Diamessis (2019) also examined these transitions and their link with the evolution of stratified homogeneous turbulence using temporal simulations.

The present work is the continuation of Ortiz-Tarin *et al.* (2021) – referred to as ONS21 – where the unstratified wake of a 6:1 prolate spheroid with a turbulent boundary layer was studied and compared with a large number of simulations and experiments. In our previous study we found that the particularities of the slender body wake, e.g., small recirculation region, low entrainment, large defect velocity, bimodal distribution of the turbulent kinetic energy, among others affect the wake decay significantly. In this study we are analyzing how

these features affect the evolution of the stratified wake. We also analyze the simulations of CS20 to closely compare our results with the stratified bluff body wake.

Some of the questions we want to answer are: do the stratified decay laws and their transition points depend on the shape of wake generator? how does the turbulence evolve in stratified slender body wakes and are there difference with bluff body wakes? how does the phase-space evolution of the stratified turbulence compares between bluff and slender body wakes? In broader terms, we attempt to find whether a turbulent stratified wake retains some imprint of the wake generator in the mean and turbulence evolution.

A description of the solver and the methodology is given in §2. The wakes are visualized in §3. The decay of the mean wake properties is analyzed in §4. Finally, the evolution of the turbulence and the phase-space analysis of the wake are presented in sections §5 and §6, respectively. The study is concluded in §7.

2. Methodology

To study the far wake of a slender body at a high Reynolds number we use a hybrid simulation. The hybrid model combines two simulations: body-inclusive (BI) that solves the flow past the wake generator and body-exclusive (BE) that resolves the intermediate and far wakes. Here, we use a spatially-evolving simulation following the procedure validated by VanDine *et al.* (2018). In the implementation, data from a selected cross-plane in the BI simulation is interpolated on to a new grid and used as an inlet boundary condition for the BE stage. This procedure allows us to alleviate the natural stiffness of the wake problem. Whereas the BI simulation is designed to capture the turbulent boundary layer and the flow separation, the BE simulation resolves the turbulence in the wake. Both the grid size and the time step required to solve the turbulent boundary layer are much smaller than those needed in the intermediate and far wakes. This method leads to significant savings in computational cost without compromising accuracy.

The setup and the solver here are the ones used in ONS2021 with the addition of stratification. Both simulations solve the three-dimensional Navier-Stokes with the Boussinesq approximation in cylindrical coordinates. The solver uses a third-order Runge-Kutta method combined with second-order Crank-Nicolson to advance the equations in time. Second-order-accurate central differences are used for the spatial derivatives in a staggered grid. A wall-adapting local eddy viscosity (WALE) is used to properly capture the turbulent boundary layer dynamics (Nicoud & Ducros 1999). Both the BI and BE simulations use Dirichlet boundary conditions at the inflow, convective outflow and Neumann at the radial boundary. Similarly to Ortiz-Tarin *et al.* (2019) a sponge layer is added to the boundaries to avoid the spurious reflection of gravity waves.

An immersed boundary method (Balaras 2004; Yang & Balaras 2006) is used to resolve the flow past a 6:1 prolate spheroid at zero angle of attack. A numerical bump is introduced on the surface of the body to accelerate the transition of the boundary layer to turbulence. The annular bump is located where the surface favorable pressure gradient is nearly zero. This location is found at approximately $0.5D$ from the nose. The radial extent of the bump is $0.002D$ (~ 15 wall units) and the streamwise extent is $0.1D$.

The stratification is set by a linear background density profile characterized by the Froude number, $Fr = U_\infty / ND$, where N is the buoyancy frequency. Three levels of stratification are simulated: $Fr = 2$, 10 and ∞ . $Fr = 2$ is close to the critical Froude number $Fr_c = 6/\pi$ for the 6:1 spheroid at which the suppression of turbulence in the wake by stratification is optimal (Ortiz-Tarin *et al.* 2019). $Fr = 10$ is a moderate level of stratification closer to oceanic values. Finally $Fr = \infty$ is the unstratified case which will be used as a reference (ONS21).

The cylindrical coordinate system is (x, r, θ) with the origin at the body center. For

Case	Re	Fr	L_r	L_θ	L_x^-	L_x^+	N_r	N_θ	N_x
1	10^5	∞	5	2π	8	15	746	512	2560
2	10^5	10	60	2π	20	30	848	512	3072
3	10^5	2	60	2π	20	30	848	512	3072

Table 1: Parameters of the body-inclusive simulation of prolate 6:1 spheroid. L_x^- and L_x^+ are the upstream and downstream distances from the wake generator.

Case	Re	Fr	L_r	L_θ	x_e	L_x	N_r	N_θ	N_x
1	10^5	∞	10	2π	6	80	479	256	4608
2	10^5	10	57	2π	9	90	619	256	4608
3	10^5	2	57	2π	9	90	619	256	4608

Table 2: Parameters of the body-exclusive simulations. x_e is the extraction location of the BI simulations that is fed as inlet to the BE simulations.

Case	Re	Fr	L_r	L_θ	L_x^-	L_x^+	N_r	N_θ	N_x
1	5×10^4	∞	15.14	2π	30.19	125.51	364	256	4608
2	5×10^4	10	80	2π	30.19	125.51	529	256	4608
3	5×10^4	2	80	2π	30.19	125.51	529	256	4608

Table 3: Parameters of the disk simulations (CS20).

convenience, the Cartesian coordinate system (x, y, z) will also be used, where z is the vertical direction aligned with gravity, y is the spanwise direction, and x is the streamwise direction.

The BI grid is designed to resolve the turbulent boundary layer and the small-scale wake turbulence. The turbulent boundary layer is resolved with $\Delta x^+ = 40$, $\Delta r^+ = 1$, and $r\Delta\theta^+ = 32$. There are 10 points in the viscous sublayer and 130 across the buffer and log layers. The mean velocities and turbulence intensities within the boundary layer were validated against existing studies (Posa & Balaras 2016; Kumar & Mahesh 2018) and the law of the wall. Additionally a grid refinement study was performed to guarantee the independence of the statistics to the grid choice.

In the wake, the peak ratio between the grid size and the Kolmogorov length $\eta = (\nu^3/\epsilon)^{1/4}$, in both BI and BE domains is $\max(\Delta x/\eta) = 7.5$, $\max(\Delta r/\eta) = 6$, and $\max(r\Delta\theta/\eta) = 5$. A figure showing the ratio between the Kolmogorov scale and the grid resolution can be found in ONS21 (figure 2). Additionally, the unstratified wake decay coincides with all the previous existing numerical and experimental works on slender body wakes (see figure 1 of ONS21).

The domain size in the stratified cases is large so that internal gravity waves are weak before reaching the sponge region near the walls. The total number of grid points across BI and BE domains is approximately 1.5 billion in the unstratified case and 2 billion in the stratified simulations. Tables 1 and 2 include the most relevant parameters of BI and BE simulations, respectively. Further details on the grid design can be found in section 2 of ONS21.




Figure 1: Instantaneous contours of streamwise velocity in the near wake for spheroid (left) and disk wakes (right) at $Fr = 2$ and $Fr = 10$ on center-vertical ($y = 0$) and center-horizontal ($z = 0$) planes. Red isolines show the limit of recirculation regions where the streamwise velocity is zero.

Once the flow has reached statistically steady state, the statistics are obtained by temporal averaging, denoted by $\langle \cdot \rangle$. Instantaneous quantities are written with lower case, mean quantities with upper case, and fluctuations with prime. In the stratified cases the average is performed over $270D/U_\infty$, approximately three flow-throughs. In the unstratified simulation flow statistics are obtained through the temporal (over $100D/U_\infty$) as well as azimuthal averaging. Apart from temporal averaging, some statistics are obtained from cross-wake area integration denoted by $\{ \cdot \}$. Unless otherwise indicated, the integral is performed over a cross-section of radius $4D$. All the flow statistics presented here die out well before they reach the limit of the integrated region.

Reported velocities and lengths are normalized with the free-stream velocity U_∞ and the body minor axis D , respectively. The normalized streamwise distance from the center of the body x is also measured as a function of the buoyancy frequency and the time. The time in the Nt axis refers to time measured by an observer attached to the mean flow that sees the body move at a speed $-U_\infty$. A Galilean transformation yields $x/Fr = Nt$.

To compare the stratified wake of the 6:1 spheroid with that of a bluff body we use the body-inclusive disk wake simulations of CS20. The solver used in CS20 is the same as the one used here although, instead of using the WALE closure model, CS20 uses a variant of dynamic Smagorinsky. The eddy-viscosity model was changed in the spheroid simulations since WALE was demonstrated to capture the behavior of the turbulent boundary layer with the resolution used in the present wall-resolved LES. Both sets of simulations are very well resolved and have a small subgrid contribution – see ONS21 and CS20 – hence the validity of the comparison. Further details of the simulations can be found in ONS21 and CS20. The main parameters of disk simulations are shown in table 3.

3. Visualizations

Figure 1 shows instantaneous snapshots of the near wake of a spheroid and a disk at $Fr = 2$ and $Fr = 10$. At both Fr , the near wake structure of the two bodies is very different. Compared to the spheroid with turbulent boundary layer (TBL), the disk wake has a large recirculation region ($\sim 2D$), as shown by the red isolines in figure 1. This large recirculation region oscillates (Rigas *et al.* 2014) and generates a vortex shedding structure that is advected downstream (Nidhan *et al.* 2020). In a spheroid with TBL, the recirculation region is very small ($\sim 0.1D$) and is surrounded by the small scale turbulence of the boundary layer. As a result, the near wake is highly organized and large scale oscillations are not observed in the near wake (Jiménez *et al.* 2010; Kumar & Mahesh 2018; Ortiz-Tarin *et al.* 2021). Only further downstream, does the wake begin to show a helical structure. This change in the structure of the slender body wake has been found to lead to a change in the decay rate and dissipation scaling in the unstratified wake (ONS21). In the following sections, we will analyze how the differences between the near wake of a disk and that of a spheroid lead to distinct trends of mean and turbulence evolution in a stratified environment. But first, let us describe different snapshots of the spheroid intermediate and far wakes. Snapshots of the disk intermediate and far wakes can be found in CS20.

Figure 2 shows an instantaneous visualization of the spheroid $Fr = 2$ wake in the center-vertical and center-horizontal planes. One of the distinctive features of the spheroid wake is that at $Fr \sim O(1)$ the separation of the boundary layer can be strongly modulated by the steady lee waves (Ortiz-Tarin *et al.* 2019). This interaction between the lee waves and the wake is particularly strong when the Froude number is close to a critical Froude number $Fr_c = AR/\pi$, where AR is the body aspect ratio. When $Fr \approx Fr_c$, half the wavelength of the lee wave ($\lambda/D = 2\pi Fr$) coincides with the length of the body and the size of the separation region is reduced. The flow is then in what is called resonant or saturated lee wave regime, (Hanazaki 1988; Chomaz *et al.* 1992). At low Reynolds numbers, this effect can lead to the relaminarization of the turbulent wake (Pal *et al.* 2016; Ortiz-Tarin *et al.* 2019). In the present case, figure 2(a) reveals that, even at $Re = 10^5$, the wake height is strongly modulated by the waves, although the wake is not relaminarized due to the high Re of the flow. For example, the wake height exhibits oscillations with a wavelength of $\lambda/D = 2\pi Fr = 4\pi$. The modulation of the wake by the waves leads to an unusual configuration in the intermediate wake ($x = 20 - 40$) where the wake width L_H is smaller than the wake height L_V , figure 2(a) and (c). In these figures, sinuous oscillations are observed only in the horizontal plane (figure 2d) due to strong stratification. These horizontal sinuous instabilities contrast with the lee wave induced varicose modulation in the vertical plane. As the wake evolves, the $L_H < L_V$ configuration transitions to the expected $L_H > L_V$. In this late region, the small-scale turbulence of the boundary layer has been dissipated and a layered-layer structure is observed in the vertical plane, figure 2(b). The qualitative trends of L_H and L_V discussed here are quantified in §4.

The main features of the $Fr = 10$ wake can be observed in the instantaneous snapshots of figure 3. The near wake, figure 3(a,c), is thin and carries the small scale turbulence generated in the boundary layer. Similar to the unstratified wake of ONS21, in the $x < 20$ region, it has a quasi-cylindrical structure. Only after $x \approx 20$, a helical structure develops. In the unstratified wake, the oscillation found at $x \approx 20$ is present until the end of the domain. Here, the $Fr = 10$ wake does not show major oscillations after $x \approx 30$. Stratification restrains the vertical motions in the wake and enhances the horizontal spread as can be seen in the visualization of the late wake, figure 3(b,d). Unlike the $Fr = 2$ wake, the horizontal and vertical $Fr = 10$ wake extent grow monotonically with increasing downstream distance.




Figure 2: Instantaneous contours of streamwise velocity of the spheroid $Fr = 2$ wake in center-vertical (a,b) and center-horizontal planes (c,d).

4. Evolution of the mean flow in spheroid and disk wakes

4.1. Evolution of the mean defect velocity (U_d)

The decay rate of the mean defect velocity $U_d = U_\infty - U$ shows the different stages in the evolution of a wake. In a stratified environment, wakes transverse the 3D, NEQ and Q2D regimes (Spedding 1997). Figure 4 compares the decay of U_d among the unstratified, $Fr = 10$, and $Fr = 2$ spheroid and disk wakes. To facilitate a one-to-one comparison, we present the disk data in the domain $3 \leq x \leq 80$, coinciding with the domain of the spheroid wake. Since x is measured from the center of the body, the location of $x = 3$ is in the near wake for the disk and is at the terminus of the body for the spheroid. The unstratified spheroid wake (figure 4a) shows a transition between the classical high- Re decay $U_d \sim x^{-2/3}$ to $U_d \sim x^{-6/5}$ at $x \approx 20$ coinciding with the development of a helical structure (ONS21). The $Fr = \infty$ disk wake decays as $U_d \sim x^{-0.9}$ from $10 \leq x \leq 65$ and transitions to the classical high- Re decay of $x^{-2/3}$ afterward (CS20), as shown in figure 4(b). Compared to the disk, the $Fr = 10$ and $Fr = \infty$ spheroid wakes have a higher value of U_d , owing to weaker near-wake entrainment and the slower development of slender body wakes.




Figure 3: Instantaneous contours of streamwise velocity of the spheroid $Fr = 10$ wake in center-vertical (a,b) and center-horizontal planes (c,d).

In the weakly stratified $Fr = 10$ regime, the defect velocity of the spheroid wake (figure 4a) evolves similarly to the unstratified wake until $Nt \approx 3.5$, when the decay rate slows down. However, at the same value of $Fr = 10$ but for the disk wake (figure 4b), U_d deviates from the unstratified case at $Nt \approx 1$. Based on U_d , the end of the 3D region and the beginning of the NEQ region of the spheroid wake occurs at $x \approx 30$ whereas in the disk it occurs at $x \approx 10$. We discuss the reason behind this delayed deviation of the spheroid-wake U_d from its unstratified counterpart in §5.

At $Fr = 2$, there are significant differences in U_d evolution between the disk and spheroid wakes. In the $Fr = 2$ spheroid wake, U_d shows an increased decay rate from the beginning. Although not shown here, the wake establishment is affected similarly to the $Fr = 1$ wake of the 4:1 spheroid of Ortiz-Tarin *et al.* (2019), where there was no 3D regime. The boundary layer evolution on the body and the separation are affected by stratification. At $Nt \approx \pi$, there is a sudden change in the decay rate due to the lee-wave-induced oscillatory modulation (Pal *et al.* 2017) observed in the $3 \lesssim x \lesssim 10$ region. This oscillatory modulation gets weaker downstream as the lee-wave amplitude decreases with the distance from the source.




Figure 4: Decay of the peak defect velocity in (a) spheroid and (b) disk. The red dashed line in (a) indicates the decay of the $Fr = 2$ centerline defect velocity. For all other cases, centerline and maximum U_d coincide. Note that the origin of the Nt scale is 1.5 for $Fr = 2$ and is 0.3 for $Fr = 10$.

At $Nt \approx \pi$, the wake transitions to the NEQ stage where U_d exhibits a slower decay compared to both the preceding stage and the following stage which commences at $Nt \approx 15$. Fitting a power law to the NEQ stage between $x = 6 - 25$ results in a decay with $x^{-0.266}$. This decay is close to the $-1/4$ decay in the NEQ regime found in the experiments of Spedding (1997) and later in numerical simulations (Diamessis *et al.* 2011; Brucker & Sarkar 2010; Redford *et al.* 2015; Pal *et al.* 2017). More details about the fitting strategy can be found in ONS21. At $Nt \approx 15$, the spheroid wake transitions to the Q2D regime with a sharper decay and a power-law fit between $x = 30 - 80$ results in $U_d \sim x^{-0.72}$, which is close to the $x^{-0.75}$ behavior established by Spedding (1997) for the Q2D regime. The $Fr = 2$ disk wake shows a very different behavior. Until at least $x = 125$ ($Nt = 62.5$) – the full extent of the computational domain – the disk wake exhibits no transition to the Q2D regime. Instead, after transitioning to the NEQ regime with a power law of $U_d \sim x^{-0.18}$, the disk wake stays in that regime.

Thus, the NEQ regime in the spheroid wake at $Fr = 2$ is significantly shortened compared to the disk wake, with it starting at $Nt \approx \pi$ and ending early at $Nt \approx 15$ when Q2D commences. In the experiments of Spedding (1997), the NEQ regime is reported to last until $Nt \approx 40$. Other temporal studies have found that the span of the NEQ regime depends on the Reynolds number. For example, in temporal simulations, Diamessis *et al.* (2011) found an increase of the NEQ duration to $Nt \approx 50$ when the Reynolds number increased to $Re = 10^5$. Brucker & Sarkar (2010) found a transition to a Q2D-type power law at $Nt \approx 100$. Only Redford *et al.* (2015) observed an earlier transition around $Nt \approx 25$. The reasons behind the early arrival of the Q2D regime in the spheroid $Fr = 2$ wake will be discussed in §5.

4.2. Evolution of the mean horizontal (L_H) and vertical (L_V) lengthscales

The evolution of the mean wake dimensions in the spheroid and disk wakes is shown in figure 5(a,c) and (b,d), respectively. L is defined such that $U_\infty - U(L) = \frac{1}{2}U_d$. The subscripts $\{V, H\}$ indicate that these measures have been taken in the vertical and horizontal planes so that they represent the half height and the half width.

The wake of a slender body is generally thinner than its bluff body counterpart. Compared to the disk wake of CS20, the present unstratified wake is smaller by a factor of about 3 – contrast black lines in figure 5(a) to (b). The difference in wake size stems from the different near-wake features. Here, the initial non-dimensional wake width is around 0.2




Figure 5: Wake dimensions measured using the mean defect velocity U_d for the spheroid (a,c) and disk (b,d) wakes in center-vertical (a,b) and center-horizontal (c,d) planes. The legends are same as in figure 4.




Figure 6: Instantaneous radial velocity contours of the $Fr = 2$ spheroid wake in the (a) center-vertical and (b) center-horizontal planes.

whereas its value for the disk is around 0.7. This observation agrees well with the scaling proposed by Tennekes & Lumley (1972) and used in stratified wake experiments by Meunier & Spedding (2004), where the wake dimensions behind a body with diameter D scale with the drag coefficient $\sqrt{C_D}$. We find that $C_D^{\text{disk}} \approx 1.11$ and $C_D^{\text{spheroid}} \approx 0.13$ resulting in $(C_D^{\text{disk}}/C_D^{\text{spheroid}})^{0.5} \approx 3.2$. Besides the initial dimensions, the near-wake growth rates of the spheroid and disk are also very different. Whereas in the $x = 3 - 20$ region the spheroid unstratified wake grows as $L \sim x^{0.2}$, the disk wake grows as $L \sim x^{0.45}$. Later, the growth rate of both wakes becomes comparable but the difference in size is already established and dictated by the near wake.

The evolution of the $Fr = 10$ spheroid wake height (L_V) is similar to its unstratified counterpart until $Nt \approx 3.5$ where the growth of L_V slows down. While L_V remains almost constant beyond $Nt \approx 4$, L_H keeps increasing with a growth rate of $\sim x^{1/3}$. In the $Fr = 10$ disk wake, the deviation from the $Fr = \infty$ case happens at $x \approx 20$ ($Nt \approx 2$). Interestingly,

after $Nt \approx 2$, the disk wake shows a continuous decrease in wake height. L_H of both spheroid and disk wakes at $Fr = 10$ closely follow the trend of the corresponding unstratified wake. See figures 5(c,d).

The wake dimensions at $Fr = 2$ for both disk and spheroid show oscillations with a wavelength of $\lambda/D = 2\pi Fr$. This reveals the influence of the steady lee waves especially on the wake height, see figure 5(a,b). From $Nt = 1 - 15$ the oscillations of L_V and L_H are consistent with the conservation of momentum deficit, i.e., to counteract the contraction of L_V caused by buoyancy, L_H is enhanced. Note that these initial oscillations are of similar amplitude in both disk and spheroid. However, the relative change over the initial wake dimensions is much more pronounced in the spheroid wake (~ 10 times) owing to its initial thinness. The influence of the lee waves on the spheroid $Fr = 2$ wake dimensions is illustrated by radial velocity contours in figure 6. The wake width contracts significantly in the region where the vertical velocity of the lee wave induces a rapid increase of wake height. Starting at $Nt = 20$, the width of the spheroid wake shows a rapid growth $L_H \sim x^{1.3}$ corresponding with: (i) the development of the horizontal wavy motions observed in figure 2(d) and (ii) the arrival of the Q2D stage with $U_d \sim x^{-3/4}$. The growth of L_V remains constant at a rate of $x^{0.25}$. Both, the very rapid growth of L_H and the sustained increase in L_V of the spheroid wake, are very different from the trends in the $Fr = 2$ disk wake. In the disk wake, we find that, instead of an increase, the vertical height exhibits a decrease ($L_V \sim x^{-0.18}$) at $x \gtrsim 20$. Furthermore, L_H grows at $x^{1/3}$, a moderate rate relative to its rapid growth rate in the disk wake.

4.3. Comparison of flow topology between stratified spheroid and disk wakes

The difference between the spheroid and disk with regards to the evolution of mean length scales (L_V and L_H), particularly at $Fr = 2$, points toward qualitative differences in the flow topology. To further characterize these differences in the $Fr = 2$ wake, figure 7 shows contours of mean streamwise velocity (U) at different streamwise locations for the spheroid (top two rows) and the disk (bottom row). In each panel of figure 7, the right half shows U_x and the left half shows turbulent kinetic energy (TKE), $E_T^K = (\langle u_x'^2 \rangle + \langle u_y'^2 \rangle + \langle u_z'^2 \rangle)/2$.

For the sake of brevity, we have not included contours of the $Fr = 10$ disk and spheroid wakes since their topology is similar – the mean can be well approximated by a vertically-squeezed two-dimensional Gaussian while the TKE evolves from a bimodal (off-center peaks) distribution in the radial direction to a Gaussian at intermediate to late streamwise distances. In the case of the disk, the TKE evolves as a two-dimensional Gaussian right beyond the recirculation region.

We first discuss the disk wake (bottom row). The mean shows a monotonic spread in the horizontal direction and resembles the shape of an ellipse or a two-dimensional Gaussian squeezed in the vertical direction. This shape does not change until the end of the computational domain at $x \approx 125$. The TKE for the disk wake also has a similar vertically squeezed appearance.

Turning to the spheroid wake, we find that its turbulence topology is different from that of the mean. In the region $5 \leq x \leq 15$, TKE shows two off-center peaks reminiscent of the TBL shedding from a slender body (Jiménez *et al.* 2010; Posa & Balaras 2016; Kumar & Mahesh 2018; Ortiz-Tarin *et al.* 2021) while mean U_x shows a single central peak. At $x = 20$, we see the start of a horizontal contraction of the mean velocity in the central region of the wake and the emergence of a ‘butterfly’ shape reminiscent of the separation and wake patterns observed in Ortiz-Tarin *et al.* (2019), where the $Fr_c = 4/\pi \approx 1$ wake of a 4:1 spheroid was studied. Note that in this stage, the wake is thinner than taller, i.e., $L_H < L_V$. At $x \approx 20$, TKE starts transitioning from a bimodal distribution to a single peak near the center-horizontal plane. In the next section, we will analyze how the horizontal contraction




Figure 7: $Fr = 2$ wake of spheroid (a-f) and disk (g-i) at different streamwise locations x . Contours of mean streamwise velocity in the right half and TKE in the left half of each contour. Contour limits are between the minimum (red) and maximum (white) values of the respective quantity at a given x with ten levels in between. Radial extent span till $r = 1$ and $r = 4$ for the spheroid and disk contours, respectively. The disk wake is larger than the spheroid wake as can be inferred from the $r = 1$ circle in the bottom row.

of the mean wake between $x = 20$ and $x = 40$ results in an increased horizontal mean shear, resulting in the maximum TKE being produced close to the center-horizontal plane. This leads to a transition in the TKE topology from bimodal distribution to a squeezed-Gaussian distribution at $x \gtrsim 40$. By $x \approx 60$, U has been organized into two distinct layers while TKE is sustained between these two vertically off-center layers. Note that the multi-layered mean flow structure at late x in the $Fr = 2$ spheroid wake is reminiscent of the layered structure of the Q2D regime Spedding (1997).

Previously, temporal simulations (Gourlay *et al.* 2001; Brucker & Sarkar 2010; Redford *et al.* 2015) have shown that the mean and the turbulence can evolve differently. Indeed, the effect of buoyancy is ‘felt’ very differently by the large and the small scales in the flow. The general trend is that, in the late wake, the turbulence occupies a smaller and smaller vertical fraction of the mean defect as time passes (Redford *et al.* 2014). Instead, the finding here for the spheroid wake is the combined effect of having a wake in saturated-lee-wave state and initial off-center peaks of TKE, established by the TBL separation. These characteristics of the flow have not been captured in temporal simulations since they have not accounted for the wake generator and the steady lee waves.




Figure 8: Comparison of evolution of TKE between spheroid (left) and disk (right) wakes. (a,b) total TKE, (c,d) streamwise TKE, (e,f) spanwise TKE, and (g,h) vertical TKE.

5. Evolution of the turbulent flow in spheroid and disk wakes

The energy of the flow is contrasted between spheroid and disk wake in this section. The turbulent kinetic energy (TKE also denoted as E_K^T), turbulent potential energy (TPE, E_P^T), mean kinetic energy (MKE, E_K^M) and mean potential energy (MPE, E_P^M) are defined as:

$$E_K^T = (\langle u_x'^2 \rangle + \langle u_y'^2 \rangle + \langle u_z'^2 \rangle)/2, \quad E_P^T = \gamma \langle \rho' \rho' \rangle / 2, \quad (5.1)$$

$$E_K^M = (U_d^2 + \langle u_y \rangle^2 + \langle u_z \rangle^2)/2, \quad E_P^M = \gamma \langle \rho_d \rangle^2 / 2, \quad (5.2)$$

where $\gamma = g^2 / \rho_0^2 N^2$. In what follows, trends of area-integrated values, denoted by $\{\cdot\}$, of these energy measures are reported. Area-integrated quantities are preferred because the peaks of mean and turbulence in stratified slender wakes are often times off-centered as can be seen in figure 7. The integration allows for a uniform comparison across cases. Also note that temporally averaged quantities are denoted by the angled brackets $\langle \cdot \rangle$.

5.1. Evolution of TKE, spectra and PE-to-KE ratios

The evolution of the area-integrated TKE of the disk and spheroid wakes is shown in figure 8. The most noticeable aspect is that, for all Fr numbers, $\{E_K^T\}$ in spheroid wakes is an order of magnitude smaller than in corresponding disk wakes. Although it is not shown here, we found a similar result for the area-averaged TKE (over an area of $r = 4D$ cross-section), instead of area-integrated TKE.

The decay of the unstratified wake is studied in more detail in ONS21 for the spheroid and

CS20 for the disk. In unstratified flow, $\{E_K^T\}$ decays following $\{E_K^T\} \sim x^{-2/5}$ for the spheroid and the disk wake follows $\{E_K^T\} \sim x^{-2/3}$. Note that these fits are empirical. Both decay rates are consistent with the decay of the peak TKE (k) and the growth of wake width (L) under the self-similarity framework, i.e., $\{E_K^T\} \sim kL^2$. Specifically, in the case of the spheroid $L \sim x^{3/5}$ and $k \sim x^{-8/5}$ (Ortiz-Tarin *et al.* 2021) and, in the case of the disk, $L \sim x^{1/3}$ and $k \sim x^{-4/3}$ (Chongsiripinyo & Sarkar 2020).

At $Fr = 10$ and for both the spheroid and the disk, $\{E_K^T\}$ deviates from the unstratified case at between $Nt \approx 1$ and $Nt \approx 3$, see figure 8(a,b). Interestingly, while the TKE is affected by stratification in the similar Nt range for both wake generators, the mean flow showed a different behavior. In the disk $Fr = 10$ wake, U_d deviated from its unstratified counterpart at $Nt \approx 1$ while, in the spheroid, this change occurred later at $Nt \approx 3$. The onset of deviations from the unstratified case will be explained in more detail in the following subsections.

At $Fr = 2$, there is a striking influence of the wake generator on the evolution of TKE. Whereas in the disk wake, the TKE decays monotonically, the far wake of the spheroid displays a rapid increase.

The disk wake shows a monotonic decay in $\{E_T^K\}$ and its individual components throughout $3 < x < 80$. Compared to the horizontal components, $\{E_{T_z}^K\}$ shows a sharper decay after $Nt \approx 10$ and turbulence anisotropy progressively increases. In the spheroid $Fr = 2$ wake, $\{E_T^K\}$ decays rapidly until $Nt \approx 10$. However, after $Nt \approx 10$, the decay slows down and is followed by a period of sustained growth starting at $Nt \approx 20$ and lasting until the end of the domain. The region of TKE growth coincides with the development of the large scale horizontal motions observed in figure 2(d) and the rapid growth of L_H shown in figure 5(c). It also coincides with the accelerated decay rate of U_d starting at $Nt \approx 20$.

Notice that right before the start of the rapid increase of TKE, the $Fr = 2$ wake has a configuration where $L_H < L_V$ (figure 7d). The horizontal response of the flow to the strong lee waves is what sustains this configuration. It is only after their strength subsides that the control on the wake is released to allow the horizontal wavy motion to develop. The rapid development of this motion coincides with the rapid increase in horizontal TKE, namely $\{E_{T_y}^T\}$ and $\{E_{K_x}^T\}$ as seen in figure 8(c,e). To the best of the authors' knowledge this is the first wake study, resolving the flow at the body, that observes an increase of fluctuation energy with downstream distance instead of its usual decrease.

To further quantify the horizontal wavy motions observed in figure 2(d), the energy spectra of the spanwise velocity are computed at the centerline. These spectra are compared between locations before (figure 9a) and after (figure 9b) the start of the the TKE increase associated with the Q2D regime. The spectra before $x < 40$ do not show preferential energization of the low frequencies. This finding is consistent with the visualizations in figure 2 where the intermediate wake does not show any sign of large scale motions. Beyond $x = 40$, however, spectra show a strong peak at Strouhal number $St = fD/U_\infty \approx 0.35$ (figure 9b). This value of Strouhal number agrees with the approximate wavelength of structures in figure 2(d), where the wavelength $\lambda/D \approx 1/St$. To summarize, the arrival of the Q2D regime in the $Fr = 2$ wake is accompanied by a strong increase in TKE (figure 8a) and the appearance of large scale motions in the center-horizontal plane (figure 2d and 9b).

In the $Fr = 2$ disk wake, the vortex shedding mode at $St = fD/U_\infty \approx 0.13 - 0.14$ is dominant throughout the whole domain. The horizontal meanders, which are prevalent in the spheroid wake, are absent in the disk wake at least until the end of the domain at $x/D = 125$. Since the vortex shedding mode, its long-lasting effect on the wake, and its internal wave field are described in detail by Nidhan *et al.* (2022), we do not discuss these aspects further.

The spheroid wake has significantly lower TKE content relative to the disk wake and also a different distribution of the mean momentum. The content of potential energy relative to




Figure 9: Energy spectra of the $Fr = 2$ spheroid wake computed with the spanwise velocity fluctuations at the centerline at (a) $x = 10, 30$ and (b) $x = 50, 70$.




Figure 10: Ratio of area-integrated (a) turbulent potential energy to turbulent kinetic energy and (b) mean potential energy to mean kinetic energy in stratified spheroid and disk wakes.

that of kinetic energy is also of interest. Figure 10 shows the ratio of area-integrated potential energy (PE) to kinetic energy (KE). Both fluctuating (10a) and mean (figure 10b) components are shown at $Fr = 2$ and 10 for both wake generators.

At $Fr = 10$, the turbulent PE-to-KE ratio increases steadily in both disk and spheroid cases indicating an increasing influence of buoyancy on turbulence (figure 10a). In the $Fr = 2$ wakes, the turbulent PE-to-KE ratio peak around $x \approx 30$ in both cases and decay afterward. By the end of the measurement region at $x = 80$, the turbulent PE-to-KE ratios are similar across $Fr = 2$ and 10 wakes. Thus, stratification and body shape do not qualitative affect the ability of turbulence to stir the density field in the intermediate and far wake. Quantitatively, the TPE-to-TKE ratios are somewhat higher for the disk relative to the spheroid for both wakes.

The mean PE-to-KE ratios in the $Fr = 10$ wakes (figure 10b) are minuscule compared to their turbulent counterparts. At $Fr = 2$, the mean-based ratio is much larger, particularly close

to the wake generators, pointing towards a strong influence of the steady lee waves. Both disk and spheroid mean-based ratios oscillate with a characteristic lengthscale corresponding to the wavelength of steady lee waves at $Fr = 2$. It is particularly revealing as to how much larger is the magnitude of $\{E_P^M\}/\{E_K^M\}$ in the spheroid wake compared to the disk, as it explains why the spheroid flow is so strongly modulated by the lee waves. It is worth noting that comparison of the absolute value of MPE between the disk and the spheroid reveals that it is the disk that has the larger MPE, an order of magnitude larger. The amplitude of the lee wave generated by the disk is larger than that of the spheroid by about a factor of 2.

5.2. Analyses of the spheroid TKE budget terms

To quantitatively understand the origin of TKE increase in the $Fr = 2$ spheroid wake, we look into the different terms of the TKE transport equation:

$$U_i \frac{\partial E_K^T}{\partial x_i} + \frac{\partial T_i}{\partial x_i} = P - \varepsilon + B, \quad (5.3)$$

where P is the turbulent production, ε is the turbulent dissipation and B denotes the turbulent buoyancy flux. These quantities are defined by:

$$P = -\langle u'_i u'_j \rangle \frac{\partial U_i}{\partial x_j}, \quad \varepsilon = 2\nu \langle s'_{ij} s'_{ij} \rangle - \langle \tau'^s_{ij} s'_{ij} \rangle, \quad B = -\frac{g}{\rho_0} \langle \rho' u'_z \rangle, \quad (5.4)$$

where $s_{ij} = (\partial_j u_i + \partial_i u_j)/2$ is the strain-rate tensor and $\tau'^s_{ij} = -2\nu_s s_{ij}$ is the subgrid stress tensor. The contribution of the subgrid term to the TKE transport equation is found to be small.

The turbulent production is a source of TKE and a sink in the MKE equation. The turbulent buoyancy flux transfers energy between TKE and TPE and the turbulent dissipation is a sink of TKE. Along with the sinks and sources of energy, there is a term responsible for the spatial redistribution of TKE, the turbulent transport,

$$T_i = \frac{1}{2} \langle u'_i u'_j u'_j \rangle + \langle u'_i p' \rangle - 2\nu \langle u'_j s'_{ij} \rangle - \langle u'_j \tau'^s_{ij} \rangle. \quad (5.5)$$

The turbulent transport redistributes energy, primarily in the y - z plane, and its contribution to the area-integrated budget is negligible.

The production term in equation (5.3) can be further expanded as follows:

$$P = -\langle u'_x u'_j \rangle \frac{\partial U_x}{\partial x_j} - \langle u'_y u'_j \rangle \frac{\partial U_y}{\partial x_j} - \langle u'_z u'_j \rangle \frac{\partial U_z}{\partial x_j}. \quad (5.6)$$

For the stratified wakes at hand, $U_y, U_z \ll U_x$. Hence equation (5.6) can be further simplified as below:

$$P = -\langle u'_x u'_x \rangle \frac{\partial U_x}{\partial x} - \langle u'_x u'_y \rangle \frac{\partial U_x}{\partial y} - \langle u'_x u'_z \rangle \frac{\partial U_x}{\partial z}. \quad (5.7)$$

In a wake developing in the x direction, P is primarily dominated by the transverse shear terms since $|\partial U_x/\partial y|, |\partial U_x/\partial z| \gg |\partial U_x/\partial x|$. Hence in the discussions on $\{P\}$ to follow, we focus on the transverse terms, namely $P_{xy} = \langle u'_x u'_y \rangle \partial_y U_x$ and $P_{xz} = \langle u'_x u'_z \rangle \partial_z U_x$.

Figure 11 shows the evolution of the area-integrated terms in the TKE transport equation of the spheroid wake. We do not present the TKE transport terms of the disk wakes here as they are presented and discussed in great detail in CS20. We first discuss the trends of the area-integrated production, $\{P\}$, and its components in $Fr = 2$ and 10 spheroid wakes. This




Figure 11: Spheroid wakes. (a) Area-integrated production. (b) Main components of the turbulent production. (c) Area-integrated dissipation. (d) Ratio between area-integrated production and dissipation. (e) Area-integrated buoyancy flux. (f) Maximum value of mean horizontal shear $\partial U_x / \partial y$.

is followed by a discussion of the area-integrated dissipation, $\{\varepsilon\}$. Thereafter, the variation of $\{P\}/\{\varepsilon\}$ sheds light on why an increase in TKE is observed in the $Fr = 2$ spheroid wake for $x > 30$.

$\{P\}$ in the $Fr = 10$ spheroid wake decays at a rate comparable to its unstratified counterpart until $Nt \approx 3$ when it starts to deviate (figure 11a). U_d in the $Fr = 10$ spheroid wake also starts deviating around $Nt \approx 3$ (figure 4a). At $Nt \approx 3$, the contribution of $P_{xz} = \langle u'_x u'_z \rangle \partial_z U_x$ starts declining due to a reduction in $\langle u'_x u'_z \rangle$ (not shown here for brevity). At $Nt \approx 4.5$, the contribution of the lateral production P_{xy} exceeds that of the vertical production P_{xz} .

The reduction of $\{P_{xz}\}$ in the $Fr = 10$ wake coincides with the transition of U_d to a slower decay rate. In figure 11(b), we note that $\{P_{xz}\}$ starts decaying faster than its unstratified counterpart between $3 < Nt < 4$ (or $30 < x < 40$). In figure 4(a), we see that this is the location where U_d of the $Fr = 10$ spheroid wake starts deviating from the $Fr = \infty$ spheroid wake. At the same location that $\{P_{xz}\}$ starts decaying rapidly, there is a maximum in the buoyancy flux $\{B\}$ (figure 11e) and a slowdown in the growth of L_V (figure 5a). Buoyancy is starting to affect the wake decay.

The initial value of $\{P\}$ in the $Fr = 2$ wake is higher than in the unstratified case. The distinct separation pattern and the vertical contraction of the mean flow (figure 5a) lead to

an increase in the vertical shear and hence $\{P_{xz}\}$, see red dashed lines in figure 11(b). This initially high value of $\{P_{xz}\}$ rapidly decays as L_V increases, leading to a reduction in the vertical shear. Note that the beginning of the U_d -based NEQ regime at $Nt \approx \pi$ also coincides with this reduction of $\{P_{xz}\}$. However, the mechanism at $Fr = 2$ is different from that of the $Fr = 10$ wake. Whereas at $Fr = 10$, the reduction of $\langle u'_x u'_z \rangle$ causes the decay of $\{P_{xz}\}$, at $Fr = 2$ the decay of $\langle u'_x u'_z \rangle$ is accompanied by a sudden reduction of $\partial_z U_x$ (not shown here). The reduction in vertical shear is driven by the expansion of L_V due to lee wave modulation in figure 5(a).

As the modulation of the wake by the lee waves continues, the horizontal production is enhanced by the strong reduction in L_H (figure 5c) in the NEQ region during $7 < x < 20$. The value of $\{P_{xy}\}$ overtakes $\{P_{xz}\}$ at $Nt \approx 5$. The magnitude of $\{P_{xy}\}$ remains nearly constant until the end of the domain. Figure 11(f) shows the maximum of the horizontal mean shear ($\partial U_x / \partial y$), i.e., $\max_{(r, \theta=0)} \partial U_x / \partial y$, in the central streamwise-horizontal plane for both wakes. The mean shear in the $Fr = 2$ wake increases in the region $10 < x < 30$, exactly in the region where the TKE decay starts plateauing (see figure 8a). The enhanced mean horizontal shear at $Fr = 2$ prevents the horizontal production from decaying monotonically unlike $Fr = 10$.

Figure 11(c) shows the evolution of $\{\varepsilon\}$ as a function of x for $Fr = \infty, 10$ and 2 spheroid wakes. In the $Fr = 10$ case, the decay of ε is similar to that of the unstratified wake until $Nt \approx 3$ after which the decay rate increases slightly. The $Fr = 2$ wake dissipation shows a sharper decay until $x \approx 20$. After $x \approx 20$, the decay rate appears to be closer to the $Fr = \infty$ decay rate of $x^{-7/5}$ (Ortiz-Tarin *et al.* 2021).

Since the turbulent dissipation keeps decreasing monotonically in the $Fr = 2$ spheroid wake (figure 11c), while the production tends to asymptote for $x > 30$ (figure 11a), the value of $\{P\}/\{\varepsilon\}$ becomes greater than 1 in the $Fr = 2$ spheroid wake, see figure 11(d), explaining the rapid increase of TKE beyond $x \approx 30$. Note that the $\{P\}/\{\varepsilon\}$ ratio oscillates with the characteristic wavelength of the lee waves revealing the strong influence of wave-related buoyancy effects on the energetics of the $Fr = 2$ wake.

One of the features of the arrival of the NEQ regime reported in previous studies is the radiation of internal gravity waves (Abdilghanie & Diamessis 2013; De Stadler & Sarkar 2012; Rowe *et al.* 2020). In these spheroid wakes, we find that the integrated wave flux remains negligible compared to the other terms in the TKE budget and hence is not shown here. The small magnitude of the turbulent wave flux is consistent with the findings of Meunier *et al.* (2018) who found that the magnitude of the wake-generated waves depends on the body drag coefficient – and the 6:1 spheroid has a very low drag compared to bluff bodies.

5.3. Early arrival of the Q2D regime in the $Fr = 2$ spheroid wake

Remember that the Q2D regime in stratified wakes is characterized by a faster decay of the mean defect velocity ($U_d \sim x^{-3/4}$) and the organization of wake into distinct layers in the vertical direction. Due to the strong effects of buoyancy, the fluid motions are primarily confined to the horizontal plane. However, the vertical variations in the velocity profiles still exist during the Q2D regime, primarily in the form of layered structures.

A key difference between the $Fr = 2$ disk and spheroid wakes is the early arrival of the Q2D regime in the spheroid wake. Whereas in the disk wake, CS20 did not observe a transition to the Q2D in a domain that extended up to $x = 125$ ($Nt = 62.5$), here we observe a transition at $x \approx 40$ ($Nt \approx 20$). The early transition in the spheroid wake is a consequence of the strong modulation of the intermediate wake by buoyancy, an effect that occurs for bodies with large aspect ratio (L/D) and, specifically, when Fr is near its critical value $Fr_c = L/D\pi$.

Figure 5(c) shows that L_H in the $Fr = 2$ spheroid wake contracts in the region $5 < x < 30$




Figure 12: Area-integrated (a) production and (b) buoyancy flux in the disk and spheroid $Fr = 10$ wakes. The unstratified wake ($Fr = \infty$) production is also shown in (a). The terms are normalized by the Lagrangian rate of change of their corresponding mean kinetic energy $\{\delta_t E_K^M\} = \{E_K^M\}U_\infty/x$.

as a response to the expansion of L_V (figure 5a) by steady lee waves. This phenomenon leads to the ‘butterfly’ shaped structure of mean U_x (figure 7(d,e)) which was also observed at a lower Reynolds number, $Re = 10^4$ in Ortiz-Tarin *et al.* (2019). In that study, which was performed at $Re = 10^4$, the spheroid wake at critical Fr_c relaminarized. Here, at a higher $Re = 10^5$, the flow response at the resonant state is quite different. The constriction of L_H leads to an enhancement in the mean horizontal shear shown in figure 11(f). This enhancement significantly slows down the decay of horizontal production (figure 11b). While ε continues to decay, $\{P\}/\{\varepsilon\}$ becomes > 1 leading to an increase of TKE for $x > 40$ (figure 8a). In figure 7(d-f), we also see the maximum TKE location moving to the wake axis from its off-center location in the near wake. It is worth noting that the enhanced $\{P\}$ that acts as a source of TKE is a sink for the mean energy. The sharp increase in TKE leads to a faster decay of U_d , and the $Fr = 2$ wake transitions to the Q2D regime early on, at $x \approx 40$.

In the disk wake, L_H is initially 3 – 4 times larger than in the spheroid wake. While the lee wave modulation of L_H is present in the disk wake as well (figure 5d), its amplitude relative the original value of L_H is quite small. Hence the horizontal mean shear (not shown here) and the horizontal production in the $Fr = 2$ disk wake continue to decay unlike in the spheroid wake.

The arrival of the Q2D regime in the spheroid wake is also accompanied by distinctive features of the Q2D regime reported in previous literature. Figure 2(d) shows lateral meanders in the late $Fr = 2$ wake similar to the lateral meanders in temporal simulations in the literature (Brucker & Sarkar 2010; Diamessis *et al.* 2011), albeit the temporal-simulation meanders occur much later in Nt units. As noted during the discussion of spectra, the waviness in the late wake has a characteristic frequency $St \approx 0.35$. The mean wake in the Q2D regime has a layered topology (figure 7f) as reported by Spedding (2002) and Chongsiripinyo *et al.* (2017). The turbulence state in the Q2D regime is characterized by weak vertical fluctuations $u'_z \ll u'_h$ (Spedding 1997) with $\{E_{Kz}^T\}/\{E_{Kh}^T\} < 0.1$ at $x > 60$ – where subscript h denotes the horizontal component of the fluctuations. Since the Q2D regime of the spheroid wake is in a relatively early phase, pancake vortices do not appear until the end of the simulation domain, $x = 80$.

5.4. Late transition to the NEQ regime in the $Fr = 10$ spheroid wake

At $Fr = 10$, the main difference between the disk and spheroid wakes is the location at which U_d deviates from the unstratified counterpart, i.e., the transition point to the NEQ regime. In the spheroid wake this transition occurs around $Nt \approx 3$, whereas in the disk it occurs at $Nt \approx 1$, compare the $Fr = 10$ curves between figure 4 (a) and (b). The beginning of the NEQ regime is marked by a slowdown in the decay rate of U_d in the stratified wake as compared to its unstratified counterpart. The NEQ regime is also characterized by the progressively increasing anisotropy between the horizontal and vertical components of TKE.

To understand better how this transition occurs we look into the mean kinetic energy (MKE) transport equation. The TKE transport equation was given in equation (5.3). In a similar fashion, the MKE transport equation is given by:

$$U_i \frac{\partial E_K^M}{\partial x_i} + \frac{\partial T_i^M}{\partial x_i} = -P - \varepsilon^M + B^M, \quad (5.8)$$

where superscript M denotes the mean counterparts of terms in equation (5.3).

Figure 12(a) shows the area-integrated production $\{P\}$ normalized by the Lagrangian change of the MKE ($\{E_K^M\}U_\infty/x$) in the disk and spheroid wakes at $Fr = \infty$ and 10. Likewise, figure 12(b) shows the normalized area-integrated mean buoyancy flux $\{B^M\}$. Normalization by the Lagrangian change of MKE allows us to quantify the individual importance of each budget term to the change of MKE.

Broadly, the mean and turbulence quantities in stratified wakes deviate from their unstratified counterparts as a result of buoyancy. In the MKE transport equation, buoyancy can affect (a) the turbulence production P as an indirect effect and (b) the MKE \leftrightarrow MPE transfer, through the mean buoyancy flux B^M . In both the disk and spheroid $Fr = 10$ wakes, we find that the contribution of the mean buoyancy flux to the MKE transport is significantly smaller than the contribution of the production, particularly for $x \lesssim 40$. Now turning to the area-integrated production in figure 12(a), we find that the production at $Fr = 10$ displays a strong reduction from the $Fr = \infty$ wake later in the spheroid at $x \approx 30$ ($Nt \approx 3$) compared to the disk where a similar strong reduction occurs at $x \approx 10$ ($Nt \approx 1$). This explains the late transition of the spheroid wake to the NEQ regime, i.e. U_d deviates from unstratified behavior later ($Nt \approx 3$) for the spheroid than the $Nt \approx 1$ transition point for the disk.

The decreased production in the $Fr = 10$ wakes of both bodies is a consequence of the decreased $\langle u'_x u'_z \rangle$ correlation (not shown here for brevity) in stratified turbulent shear flows (Jacobitz *et al.* 1997). In stratified wakes, this buoyancy effect has been observed experimentally by Spedding (2002) and numerically by Brucker & Sarkar (2010). The deviation between production of the $Fr = 10$ and ∞ wakes (figure 11a) is indeed caused by a reduction in its $\{P_{xz}\}$ component, see blue dashed lines in figure 11(b).

6. Evolution of the local flow state and its trajectory in the phase space

In previous sections, we showed how the spheroid and the disk wake do not transition between the 3D, NEQ and Q2D regimes at the same Nt . In this section, we examine the evolution of key local non-dimensional numbers describing the mean and fluctuating state to explore their roles. These non-dimensional numbers are local, streamwise-varying measures of stratification (Froude number) and the dynamical range of inertially dominated scales (Reynolds number). We also plot the trajectory of each wake in the Froude-Reynolds phase space.

The mean vertical Froude number (Fr_V) and the turbulent vertical and horizontal Froude numbers (Fr_V, Fr_h) are defined as




Figure 13: Evolution of the spheroid (solid lines) and disk (dashed lines) wakes at $Fr = 2$ (red) and $Fr = 10$ (blue) in the non-dimensional parameter space: (a) local vertical mean Froude number Fr_V , (b) local vertical turbulent Froude number Fr_V , (c) local horizontal turbulent Froude number Fr_h , and (d) local horizontal Reynolds number.




Figure 14: Description of the trajectories of spheroid (solid lines) and disk (dashed lines) wakes in $Fr_h - Re_h Fr_h^2$ phase space. Here, $Fr = 2$ and $Fr = 10$ are shown in red and blue, respectively. Dotted black line shows a $Re_h = \text{const.}$ line in phase space.

$$Fr_V = \frac{U_d}{2NL_V}, \quad Fr_v = \frac{u'_h}{Nl_v}, \quad Fr_h = \frac{u'_h}{NL_{Hk}}, \quad (6.1)$$

where $u'_h = (\langle u_x'^2 \rangle + \langle u_y'^2 \rangle)^{1/2}$ is the r.m.s. of the horizontal fluctuations and $l_v^2 = \langle u_x'^2 + u_y'^2 \rangle / \langle (\partial_z u_x')^2 + (\partial_z u_y')^2 \rangle$ is a vertical turbulent length scale (Riley & DeBruynKops 2003).

The mean Froude number is defined consistently with the global Froude number ($Fr = U_\infty/ND$), this is, with the wake full height $2L_V$. The turbulent Froude numbers are defined

with turbulence length scales in the energy-containing range. Since the horizontal turbulent integral length scale (l_h) is not easy to compute in a spatially evolving flow, we use the TKE-based horizontal half wake width (L_{Hk}) as a surrogate following CS20. L_{Hk} at a downstream location x is calculated by $E_K^T(y = L_{Hk}, z = 0, x) = 0.5E_K^T(y = 0, z = 0, x)$.

The mean vertical Froude number Fr_V (figure 13a) becomes $O(1)$ in both disk and spheroid wakes at the location at which the decay of U_d slows down signaling the beginning of the NEQ regime, marked by $Nt \approx 1$ for the disk and $Nt \approx 3$ for the spheroid. For the $Fr = 2$ cases, Fr_V in the spheroid wake starts dropping faster beyond $x \approx 40$ – the streamwise location where the wake transitions from NEQ to Q2D regime.

The turbulent Froude numbers play an analogous role to those based on the mean. When their values become $O(1)$, turbulence starts being affected by buoyancy. Fr_v is defined using l_v which has a significance to shear instability. Defined with l_v , Fr_v is proportional to $Ri^{-1/2}$, where Ri is the Richardson number of the fluctuating shear (Riley & DeBruynKops (2003), CS20). When Fr_v becomes $O(1)$ (figure 13b) is also when the spanwise and vertical components of the TKE start showing anisotropy of the turbulence stress tensor and $E_{Kz}^T > E_{Ky}^T$ as shown in figure 8 for the $Fr = 2$ wakes. See also figure 8 of CS20.

The avid reader will notice that the four $Fr_v = u'_h/Nl_v$ curves could collapse if plotted against Nt instead of x . Indeed, these curves collapse in the (Fr_v, Nt) plot for $1 < Nt < 10$ and the factor of 5 present between the curves in figure 13(b) is simply the ratio of buoyancy frequency N between $Fr = 2$ and $Fr = 10$. This result points to the fact that the spheroid and disk wakes are at a sufficiently high Re so that there is a range of small-scale wake turbulence with dynamics relatively unaffected by the large scales and, therefore, by neither the shape of the wake generator nor buoyancy. u'_h/l_v is a quantity that reflects such dynamics and thus, on dimensional grounds, evolves as $u'_h/l_v \sim t^{-1}$ so that $Fr_v = u'_h/Nl_v \sim (Nt)^{-1}$. We note that buoyancy does affect large scale wake features, which are affected by the body particularities – hence the lack of collapse in Fr_V . The values of the buoyancy Reynolds number $Re_b = \epsilon/\nu N^2$ and the Reynolds of the Taylor micro-scale $Re_\lambda = \sqrt{k}\lambda/\nu$ where $\lambda^2 = 15\nu u_x'^2/\epsilon$ (not included here for brevity) are very similar in magnitude for both disk and spheroid wakes at the two levels of stratification. This finding is consistent with the initial collapse of Fr_v when plotted against Nt .

In figure 13(c), the $Fr = 10$ wakes of both disk and spheroid reach $Fr_h \sim O(1)$ at $x \approx 10 - 20$, the location at which the $\{E_K^T\}$ starts deviating from its unstratified counterpart (figure 8(a,b)). In the $Fr = 2$ spheroid wake, $Fr_h < 1$ throughout the domain and $\{E_K^T\}$ deviates from the unstratified decay from the very beginning in both disk and spheroid wakes. Overall, we find that $Fr_h \sim O(1)$ is a good indicator of the deviation of $\{E_K^T\}$ from the unstratified counterpart. In contrast, $Fr_v \sim O(1)$ marks the location at which turbulence anisotropy between the vertical and the spanwise components starts growing. Figure 13(d) shows the evolution of turbulence Reynolds number, $Re_h = u'_h L_{Hk}/\nu$ with u'_h being the intensity of horizontal turbulent fluctuations. The value of Re_h (figure 13d) changes slowly as the wake evolves, remaining at $O(10^4)$ for the disk wake and at $O(10^3)$ for the spheroid wake.

A consolidated view of the evolution of the state of fluctuations is provided in phase space (Brethouwer *et al.* 2007; Zhou & Diamessis 2019; de Bruyn Kops & Riley 2019; Chongsiripinyo & Sarkar 2020). In the phase-space portrait, one axis measures the buoyancy effect on the large scales through the turbulent Froude number (Fr_h) and the other axis is a measure of Reynolds number that is not Re_h but one which accounts for buoyancy in addition to inertia. Ozmidov-scale eddies are the largest eddies unrestrained by buoyancy. The Ozmidov scales $l_o = (\epsilon/N^3)^{1/2}$ and $u_o = (\epsilon/N)^{1/2}$ lead to the definition of $Re_b = u_o l_o/\nu = \epsilon/\nu N^2$ as the so-called buoyancy Reynolds number. Another convenient measure

of Reynolds number, which does not require explicit computation of the turbulent dissipation rate, is the buoyancy-weighted Reynolds number, $Re_h Fr_h^2$ (Billant & Chomaz 2001; Riley & DeBruynKops 2003). The Reynolds numbers based on buoyancy tend to decrease with downstream distance as buoyancy progressively increases in importance and limits the range of scales which are susceptible to 3D turbulent motions. As long as $Re_h Fr_h^2 > O(1)$, viscous effects do not dominate.

Following CS20, figure 14 shows the evolution all four wakes in the $Re_h Fr_h^2 - Fr_h$ phase space, where $Re_h = u'_h L_{HK} / \nu$, u'_h being the intensity of horizontal turbulent fluctuations. The flow evolves in the direction of the arrow from a state where buoyancy effects are weak (almost negligible) to a region characterized by the presence of stratified turbulence. Within the state of stratified turbulence, three different regimes can be demarcated: weakly, intermediately, and strongly stratified turbulence (WST, IST, and SST). Unlike the case of freely decaying turbulence, the mean velocity is also important here. Therefore, CS20 elected to distinguish between WST (where buoyancy begins to affect the mean velocity) and IST (where the turbulence anisotropy begins to be affected) as we do here. The SST regime is one where buoyancy effects on the large scales is very strong ($Fr_h \ll 1$) but nevertheless the value of $Re_h Fr_h^2$ is sufficiently large so that viscous effects are not dominant at the large scales.

The slope in the $Re_h Fr_h^2 - Fr_h$ plane is similar for both disk and spheroid wakes in the weak buoyancy and WST stages. The slope almost follows a line of constant Re_h , marked by a dashed line, signaling the slow evolution of Re_h compared to that of Fr_h . However, there are significant differences in the way the spheroid wakes transverse the phase space. The spheroid wake starts out thinner than the disk wake, leading to larger local Froude numbers. Also, despite the body-based Re being larger in the spheroid wake by a factor of two, the turbulence intensity in the disk wake is higher, hence the Re_h difference observed in figure 13(d). This difference in Re_h can be seen in the horizontal offset between the disk and spheroid lines in the phase-space representation.

Despite the larger spheroid body-based Reynolds number – $Re = 10^5$ instead of $Re = 5 \times 10^4$, turbulence in the spheroid wake is unable to access either the the IST regime or the SST regime while the disk wake is able to access these regimes of stratified turbulence. Furthermore, the increase in TKE, which is not observed in the $Fr = 2$ disk wake, reverses the trajectory of the $Fr = 2$ spheroid wake. To the authors' best knowledge, a reversing trend of phase-space trajectory has not been seen before in the stratified turbulence literature. These differences reveal that the phase space evolution, at least for $Fr \leq 10$, depends on the features of the wake generator, e.g., its aspect ratio or the type of BL separation.

Based on the phase-space portrait alone, one may hastily conclude that stratified slender body wakes always experience a weaker buoyancy effect relative to bluff bodies. This is true for the $Fr = 10$ case, at least for the limited Nt simulation time, in terms of the relative amount of TPE (figure 10) and the vertical scale (L_V) deviation from the unstratified case – note that both are smaller in the spheroid wake than in the disk. However, we have also shown that for the $Fr = 2$ spheroid wake, compared to the disk wake, there is a much earlier onset of the Q2D regime. The mean and turbulence quantities are highly intertwined and the strong modulation of the mean flow by steady lee waves of the high aspect-ratio spheroid in the $Fr = 2$ case ultimately leads to an early entry into the Q2D regime. Hence, it is not entirely accurate to say that the slender body wakes are weakly affected by stratification. The present work calls for a need to generalize the parameter space of $\{Re, Fr\}$ in turbulent shear flows to account for the mean flow field (also possibly its instabilities) to build a more comprehensive understanding of buoyancy effects in shear flows. In the case of wakes, the shape of the body generator is brought into play through the mean flow as shown here.

7. Discussion and final remarks

The high-Reynolds number stratified wake of a slender body has been studied using a high-resolution hybrid simulation. The wake generator is a 6:1 prolate spheroid with a tripped turbulent boundary layer, the diameter-based Reynolds number is $Re = 10^5$ and the Froude numbers, namely $Fr = U_\infty/ND = \{2, 10, \infty\}$, take moderate to large values. By comparing the spheroid wake with the disk wake of Chongsiripinyo & Sarkar (2020) (referred to as CS20), we are able to study the influence of the wake generator - slender versus bluff - in the establishment and evolution of stratified wakes.

The near wake of the 6:1 prolate spheroid with a turbulent boundary layer is characterized by a small recirculation region ($\sim 0.1D$). The recirculation region is surrounded by small-scale turbulence that emerges from the boundary layer and the flow does not show strong vortex shedding at the body (Jiménez *et al.* 2010; Posa & Balaras 2016; Kumar & Mahesh 2018; Ortiz-Tarin *et al.* 2021). As a result, the wake is much thinner and develops slower than the wake of a bluff body like the disk, which has a large recirculation region ($\sim 2D$) and vortex shedding from the body (Nidhan *et al.* 2020). These body-dependent features of the near wake were recently shown to affect the decay of the far wake in environments *without* density stratification (Ortiz-Tarin *et al.* 2021). In the present stratified simulations also we find substantial differences in the decay of the disk and spheroid wake. Particularly, we find that the starting locations of the non-equilibrium (NEQ) and the following quasi-2D (Q2D) regions of wake deficit velocity depend on the wake generator.

At $Fr = 2 \approx (L/D)/\pi$, the wake of a 6:1 prolate spheroid is in a resonant state. The half wavelength of the lee waves is equal to the body length and, as a result, the flow separation and the wake are strongly modulated by the waves. Whereas previous works had described this regime in laminar-separation configurations of a sphere (Hanazaki 1988; Chomaz *et al.* 1992) and a 4:1 spheroid (Ortiz-Tarin *et al.* 2019), the present results show that the influence of the waves persists even at high Reynolds numbers and with the separation of a turbulent boundary layer. At $Fr = 2$, the flow and the turbulence in the spheroid wake evolve very differently from the disk wake. Both the lack of strong shedding in the near wake (Ortiz-Tarin *et al.* 2021) and the strong modulation of the mean flow by the lee waves, lead to a wake with vertical and horizontal profiles of mean velocity that depart strongly from Gaussian. These features are not observed in the disk wake at $Fr = 2$, which shows a vertically-squeezed Gaussian topology and a weak imprint of lee waves on the wake dimensions.

At $Fr = 2$, both disk and spheroid wakes transition to the NEQ regime at $Nt \approx \pi$. However the transition to the Q2D regime - with enhanced wake decay relative to the NEQ regime - is very different; whereas the spheroid wake transitions at $Nt \approx 15$, the disk wake does not access the Q2D regime in a domain that spans $Nt \approx 60$. Other bluff bodies, e.g., the towed sphere (Spedding 1997) show transition to the Q2D regime at $Nt \approx 50$, a location which is also delayed with respect to the spheroid wake. The early transition to the Q2D regime of the spheroid wake is driven by its strong modulation - horizontal contraction and expansion of the wake width - in response to the vertical contraction and expansion by the lee waves. This modulation has a particularly strong effect on the slender wake of a spheroid where the horizontal contraction is a large fraction of the wake width. The early start of the Q2D regime in the spheroid wake is accompanied by a sustained increase of turbulent kinetic energy (TKE), driven by an increase of the horizontal mean shear which acts on the turbulence of the separated boundary layer. The TKE increase is limited to the horizontal velocity with the spanwise component being strongest, having almost an order of magnitude larger energy than the vertical. Although coherent vortical structures and spanwise flapping are seen in the horizontal motion, pancake eddies are incipient and not fully formed at the end of the domain, $x/D = 80$.

At $Fr = 10$ also, there are differences between the disk and the spheroid wakes. Particularly in the spheroid wake, the beginning of the NEQ stage occurs later, at $Nt \approx 3$ instead of $Nt \approx 1$ ($x \approx 30$ instead of $x \approx 10$). The difference in the start of the NEQ can be attributed to the value of the local mean Froude number $Fr_V = U_d/2NL_v$. As noted previously, the spheroid wake is thinner than the disk wake, the mixing in the near wake is weaker, and as a result the defect velocity in the intermediate wake is larger. These features increase the value of the spheroid wake local Froude number and delay the onset of the buoyancy effect that gives rise to the NEQ regime. Additionally, the analysis of the mean kinetic energy (MKE) transport terms shows that the onset of buoyancy effect on the mean flow of both the disk and spheroid $Fr = 10$ wakes is associated with the decreased energy transfer from MKE to TKE.

Taking the unstratified case as a base line, the effect of buoyancy in the spheroid $Fr = 10$ wake is observed earlier (at $Nt \approx 1$) on the decay of the TKE than its effect (at $Nt \approx 3$) on the decay of U_d . In the spheroid wake at $Fr = 10$, the transfer from TKE to TPE is responsible for the enhanced decay of TKE at $Nt \approx 1$. On the other hand, the decrease in turbulent production at a farther downstream distance (compared to the disk $Fr = 10$ wake) in the spheroid $Fr = 10$ wake is responsible for the slowed decay of the mean defect velocity at $Nt \approx 3$. The decrease in the production is caused by a reduction in the $\langle u'_x u'_z \rangle$ correlation (Jacobitz *et al.* 1997; Spedding 2002; Brucker & Sarkar 2010).

Meunier & Spedding (2004) compared the evolution far into the stratified wake, up to $x \approx 8000$, among several body shapes that also included a 6:1 prolate spheroid and a circular disk. The body Reynolds number was $Re = 5000$ and their diameter based Froude numbers were $Fr = 4$ and 16. When normalized using $D_{\text{eff}} = D\sqrt{C_D/2}$ instead of D , the evolution of the peak defect velocity of different wake generators exhibited approximate collapse for $Nt \gtrsim 50$ (see their figure 5b) with a Q2D decay rate of $\sim x^{-0.75}$. Similarly, the wake width in the horizontal of different shapes approximately collapsed for $x/D_{\text{eff}} > 400$ to exhibit a growth rate of $\sim x^{0.35}$. Since their 6:1 spheroid data starts from $x/D \approx 100$ (see their figure 5a), a direct comparison is not possible with our simulations that end at $x/D = 80$.

In regard to the mean defect decay, a major difference between Meunier & Spedding (2004) and our results is that the transition to Q2D power-law behavior appears earlier, around $Nt \approx 15$, in the $Fr = 2$ spheroid wake relative to the the $Fr = 2$ disk wake which does not transition to the Q2D decay rate until the end of the domain at $Nt = 62.5$. The value of $Re = 10^5$ in the spheroid wake is larger here and it is possible that the features that we have linked to the early onset of the Q2D stage for the spheroid $Fr = O(1)$ wake, i.e., the instability that leads to horizontal meanders and also the enhanced TKE production, are inhibited by viscous damping at the lower Re of the experiments.

The present simulations, both of the disk and the spheroid, do not extend into the very far wake regime reached by their experiments. Future hybrid simulations or experimental work at higher Re that probe the very far wake would clearly be useful. At any given Re and Fr , it will also be of interest to look at how tripping affects the wake evolution for the different body shapes.

The simulations show that the buoyancy timescale Nt alone is not sufficient to determine the state of the wake decay for both generators. However, we find that the value of the local turbulent and mean Froude numbers can be a good proxy to describe some aspects of the wake state. For both disk and spheroid wakes, $Fr_V = U_d/2NL_v$ becomes $O(1)$ at the location at which the decay of U_d slows down; $Fr_h = u'_h/NL_{Hk} \sim O(1)$ marks the location at which the area-integrated TKE of the stratified wake starts deviating from the unstratified case; and $Fr_v = u'_h/Nl_v \sim O(1)$ signals the location at which anisotropy between the different TKE components starts growing.

The buoyancy-weighted Reynolds number ($Re_h Fr_h^2$) has been used widely in stratified

flow as a convenient surrogate for the buoyancy Reynolds number ($Re_b = \varepsilon/\nu N^2$) since it displays similar trends during the flow evolution and the two quantities can be shown to be proportional using classical inviscid scaling of the turbulent dissipation rate. The surrogacy is true for the stratified wakes considered here except for the spheroid $Fr = 2$ wake after its entry into the stage of Q2D wake decay. The horizontal fluctuation energy, therefore Fr_h , increases owing to horizontal meanders and flapping of the flow. However, ε continues to decrease, albeit at a reduced rate relative to the NEQ regime. The value of $Re_b = O(10)$ is not high in the Q2D regime realized here at $Fr = 2$. It remains to be seen if, in the Q2D regime at even higher body-based Reynolds number, the equivalence between $Re_h Fr_h^2$ and Re_b is recovered and whether the unusual upward trajectory seen here in $\{Fr_h, Re_h Fr_h^2\}$ phase space is also seen in $\{Fr_h, Re_b\}$ space. The duration of the upward trajectory in phase space until the eventual downward shift toward the viscous regime is also of interest.

The differences between bluff body (disk) and slender body (6:1 spheroid) wakes illustrate the difficulty of finding a universal scaling for the high- Re stratified wake. The initial magnitude of U_d for different wake generators and levels of stratification can be roughly scaled with the global Fr and the body drag coefficient (Meunier & Spedding 2004). However, the start and the duration of the NEQ regime cannot be assumed to be independent of the wake generator. We find that rather than a particular Nt , the local mean Froude number is a good proxy for the onset of the NEQ regime in the mean defect velocity and the values of local turbulent Froude number provide guidance for the behavior of TKE, e.g., the onset of buoyancy effect as well as the location at which the ratio of vertical to horizontal TKE starts decreasing. We are unable to connect Froude number to the Q2D regime transition of the wake. More numerical and experimental work spanning different wake generators, different sources of turbulence including freestream turbulence, and longer downstream distances will be instrumental in building a comprehensive picture of the effect of initial/boundary conditions on subsequent wake evolution.

Acknowledgments. Dr. K. Chongsiripinyo is thanked for useful discussions and for disk simulations. We thank the anonymous reviewers for their helpful comments.

Funding. We gratefully acknowledge the support of Office of Naval Research Grant N00014-20-1-2253.

Declaration of interests. The authors report no conflict of interest.

Author Contributions. J.L.O.-T. and S.N. contributed equally to this paper and are co-first authors.

Author ORCIDs.

J.L. Ortiz-Tarin <https://orcid.org/0000-0001-8325-9381>;

S. Nidhan <https://orcid.org/0000-0003-0433-6129>;

S. Sarkar <https://orcid.org/0000-0002-9006-3173>.

REFERENCES

- ABDILGHANIE, A. M. & DIAMESSIS, P. J. 2013 The internal gravity wave field emitted by a stably stratified turbulent wake. *J. Fluid Mech.* **720**, 104–139.
- BAINES, P. G. 1998 *Topographic Effects in Stratified Flows*. Cambridge University Press.
- BALARAS, E. 2004 Modeling complex boundaries using an external force field on fixed Cartesian grids in large-eddy simulations. *Comput. Fluids* **33**, 375–404.
- BEVILAQUA, P. M. & LYKODIS, P. S. 1978 Turbulence memory in self-preserving wakes. *J. Fluid Mech.* **89** (3), 589–606.
- BILLANT, P. & CHOMAZ, J. M. 2001 Self-similarity of strongly stratified inviscid flows. *Phys. Fluids* **13** (6), 1645–1651.
- BONNIER, M. & EIFF, O. 2002 Experimental investigation of the collapse of a turbulent wake in a stably stratified fluid. *Phys. Fluids* **14** (2), 791–801.

- BRETHOUWER, G., BILLANT, P., LINDBORG, E. & CHOMAZ, J.-M. 2007 Scaling analysis and simulation of strongly stratified turbulent flows. *J. Fluid Mech.* **585**, 343–368.
- BRUCKER, K. A. & SARKAR, S. 2010 A comparative study of self-propelled and towed wakes in a stratified fluid. *J. Fluid Mech.* **652**, 373–404.
- DE BRUYN KOPS, S. M. & RILEY, J. J. 2019 The effects of stable stratification on the decay of initially isotropic homogeneous turbulence. *J. Fluid Mech.* **860**, 787–821.
- CASTRO, I. P., SNYDER, W. H. & MARSH, G. L. 1983 Stratified flow over three-dimensional ridges. *J. Fluid Mech.* **135**, 261–282.
- CHESNAKAS, C. J. & SIMPSON, R. L. 1994 Full three-dimensional measurements of the cross-flow separation region of a 6:1 prolate spheroid. *Exp. Fluids* **17**, 68–74.
- CHEVRAY, R. 1968 The turbulent wake of a body of revolution. *J. Basic Eng* **90**, 275–284.
- CHOMAZ, J. M., BONNETON, P., BUTET, A., PERRIER, M. & HOPFINGER, E. J. 1992 Froude number dependence of the flow separation line on a sphere towed in a stratified fluid. *Phys. Fluids* **4** (2), 254–258.
- CHOMAZ, J. M., BONNETON, P. & HOPFINGER, E. J. 1993 The structure of the near wake of a sphere moving horizontally in a stratified fluid. *J. Fluid Mech.* **254** (II), 1–21.
- CHONGSIRIPINYO, K., PAL, A. & SARKAR, S. 2017 On the vortex dynamics of flow past a sphere at $Re = 3700$ in a uniformly stratified fluid. *Phys. Fluids* **29** (2), 020704.
- CHONGSIRIPINYO, K. & SARKAR, S. 2020 Decay of turbulent wakes behind a disk in homogeneous and stratified fluids. *J. Fluid Mech.* **885**.
- CONSTANTINESCU, G. S., PASINATO, H., WANG, Y. Q., FORSYTHE, J. R. & SQUIRES, K. D. 2002 Numerical investigation of flow past a prolate spheroid. *J. Fluids Eng.* **124** (4), 904–910.
- COSTIS, C. E., TELIONIS, D. P. & HOANG, N. T. 1989 Laminar separating flow over a prolate spheroid. *J. Aircr.* **26** (9), 810–816.
- DAIRAY, T., OBLIGADO, M. & VASSILICOS, J. C. 2015 Non-equilibrium scaling laws in axisymmetric turbulent wakes. *J. Fluid Mech.* **781**, 166–195.
- DE STADLER, M. B. & SARKAR, S. 2012 Simulation of a propelled wake with moderate excess momentum in a stratified fluid. *J. Fluid Mech.* **692**, 28–52.
- DIAMESSIS, P. J., SPEDDING, G. R. & DOMARADZKI, J. A. 2011 Similarity scaling and vorticity structure in high-Reynolds-number stably stratified turbulent wakes. *J. Fluid Mech.* **671**, 52–95.
- DOMMERMUTH, D. G., ROTTMAN, J. W., INNIS, G. E. & NOVIKOV, E. A. 2002 Numerical simulation of the wake of a towed sphere in a weakly stratified fluid. *J. Fluid Mech.* **473** (473), 83–101.
- DRAZIN, P. G. 1961 On the steady flow of a fluid of variable density past an obstacle. *Tellus* **13** (2), 239–251.
- FU, T. C., SHEKARRIZ, A., KATZ, J. & HUANG, T. T. 1994 The flow structure in the lee of an inclined 6:1 prolate spheroid. *J. Fluid Mech.* **269**, 79–106.
- GOURLAY, M. J., ARENDT, S. C., FRITTS, D. C. & WERNE, J. 2001 Numerical modeling of initially turbulent wakes with net momentum. *Phys. Fluids* **13** (12), 3783–3802.
- HANAZAKI, H. 1988 A numerical study of three-dimensional stratified flow past a sphere. *J. Fluid Mech.* **192** (1988), 393–419.
- HUNT, J. C. R. & SNYDER, W. H. 1980 Experiments on stably and neutrally stratified flow over a model three-dimensional hill. *J. Fluid Mech.* **96** (4), 671–704.
- JACOBITZ, F. G., SARKAR, S. & VANATTA, C. W. 1997 Direct numerical simulations of the turbulence evolution in a uniformly sheared and stably stratified flow. *J. Fluid Mech.* **342**, 231–261.
- JIMÉNEZ, J. M., HULTMARK, M. & SMITS, A. J. 2010 The intermediate wake of a body of revolution at high Reynolds numbers. *J. Fluid Mech.* **659**, 516–539.
- KUMAR, P. & MAHESH, K. 2018 Large-eddy simulation of flow over an axisymmetric body of revolution. *J. Fluid Mech.* **853**, 537–563.
- LIN, J. T. & PAO, Y. H. 1979 Wakes in stratified fluids. *Ann. Rev. Fluid Mech.* **11**, 317–338.
- LIN, Q., LINDBERG, W. R., BOYER, D. L. & FERNANDO, H. J. S. 1992 Stratified flow past a sphere. *J. Fluid Mech.* **240**, 315–354.
- MEUNIER, P., DIZÈS, S. L., REDEKOPP, L. & SPEDDING, G. R. 2018 Internal waves generated by a stratified wake : experiment and theory. *J. Fluid Mech.* pp. 752–788.
- MEUNIER, P. & SPEDDING, G. R. 2004 A loss of memory in stratified momentum wakes. *Phys. Fluids* **16** (2), 298–305.
- MORE, R. V., ARDEKANI, M. N., BRANDT, L. & ARDEKANI, A. M. 2021 Orientation instability of settling spheroids in a linearly density-stratified fluid. *J. Fluid Mech.* **929**.

- NICOUD, F. & DUCROS, F. 1999 Subgrid-scale stress modelling based on the square of the velocity gradient tensor. *Flow Turbul. Combust.* **62**, 183–200.
- NIDHAN, S., CHONGSIRIPINYO, K., SCHMIDT, O. T. & SARKAR, S. 2020 Spectral proper orthogonal decomposition analysis of the turbulent wake of a disk at $Re = 50\,000$. *Phys. Rev Fluids* **5** (12), 124606.
- NIDHAN, S., ORTIZ-TARIN, J. L., CHONGSIRIPINYO, K., SARKAR, S. & SCHMID, P. J. 2019 Dynamic Mode Decomposition of Stratified Wakes. In *AIAA Aviation 2019 Forum*, pp. 2019–3330.
- NIDHAN, S., SCHMIDT, O. T. & SARKAR, S. 2022 Analysis of coherence in turbulent stratified wakes using spectral proper orthogonal decomposition. *J. Fluid Mech.* **934**.
- ORR, T. S., DOMARADZKI, J. A., SPEDDING, G. R. & CONSTANTINESCU, G. S. 2015 Numerical simulations of the near wake of a sphere moving in a steady, horizontal motion through a linearly stratified fluid at $Re = 1000$. *Phys. Fluids* **27** (3), 035113.
- ORTIZ-TARIN, J. L., CHONGSIRIPINYO, K. C. & SARKAR, S. 2019 Stratified flow past a prolate spheroid. *Phys. Rev. Fluids* **094803**, 1–28.
- ORTIZ-TARIN, J. L., NIDHAN, S. & SARKAR, S. 2021 High-Reynolds number wake of a slender body. *J. Fluid Mech.* **261**, 333–374.
- PAL, A., SARKAR, S., POSA, A. & BALARAS, E. 2016 Regeneration of turbulent fluctuations in low-Froude number flow over a sphere at Reynolds number of 3700. *J. Fluid Mech.* **804 R2**, 1–11.
- PAL, A., SARKAR, S., POSA, A. & BALARAS, E. 2017 Direct numerical simulation of stratified flow past a sphere at a subcritical Reynolds number of 3700 and moderate Froude number. *J. Fluid Mech.* **826**, 5–31.
- PASQUETTI, R. 2011 Temporal/spatial simulation of the stratified far wake of a sphere. *Comput. Fluids* **40** (1), 179–187.
- POSA, A. & BALARAS, E. 2016 A numerical investigation of the wake of an axisymmetric body with appendages. *J. Fluid Mech.* **792**, 470–498.
- REDFORD, J. A., CASTRO, I. P. & COLEMAN, G. N. 2012 On the universality of turbulent axisymmetric wakes. *J. Fluid Mech.* **710**, 419–452.
- REDFORD, J. A., LUND, T. S. & COLEMAN, G. N. 2014 Direct Numerical Simulation of Weakly Stratified Turbulent Wake. *Nasa Technical Memorandum* **218523** (September), 568–609.
- REDFORD, J. A., LUND, T. S. & COLEMAN, G. N. 2015 A numerical study of a weakly stratified turbulent wake. *J. Fluid Mech.* **776**, 568–609.
- RIGAS, G., OXLADE, A. R., MORGANS, A. S. & MORRISON, J. F. 2014 Low-dimensional dynamics of a turbulent axisymmetric wake. *J. Fluid Mech.* **755**, R51–R511.
- RILEY, J. & DEBRUYNKOPS, S. M. 2003 Dynamics of turbulence strongly influenced by buoyancy. *Phys. Fluids* **15** (7), 2047–2059.
- ROWE, K. L., DIAMESSIS, P. J. & ZHOU, Q. 2020 Internal gravity wave radiation from a stratified turbulent wake. *J. Fluid Mech.* **888**, 1–26.
- SPEDDING, G. R. 1997 The evolution of initially turbulent bluff-body wakes at high internal Froude number. *J. Fluid Mech.* **337**, 283–301.
- SPEDDING, G. R. 2002 Vertical structure in stratified wakes with high initial froude number. *J. Fluid Mech.* **454**, 71–112.
- SPEDDING, G. R. 2014 Wake Signature Detection. *Annu. Rev. Fluid Mech.* **46** (1), 273–302.
- TENNEKES, H. & LUMLEY, J. L. 1972 *A First Course in Turbulence*. MIT Press.
- VANDINE, A., CHONGSIRIPINYO, K. & SARKAR, S. 2018 Hybrid spatially-evolving DNS model of flow past a sphere. *Comput. Fluids* **171**, 41–52.
- VASSILICOS, J. C. 2015 Dissipation in Turbulent Flows. *Annu. Rev. Fluid Mech.* **47** (1), 95–114.
- WANG, K. C. 1970 Three-dimensional boundary layer near the plane of symmetry of a spheroid at incidence. *J. Fluid Mech.* **43**, 187–209.
- WANG, K. C., ZHOU, H. C., HU, C. H. & HARRINGTON, S. 1990 Three-dimensional separated flow structure over prolate spheroids. *Proc. R. Soc. London, Ser. A: Mathematical and Physical Sciences* **429** (1876), 73–90.
- WIKSTRÖM, N., SVENNBERG, U., ALIN, N. & FUREBY, C. 2004 Large eddy simulation of the flow around an inclined prolate spheroid. *J. Turbul.* **5** (29).
- WYGNANSKI, W., CHAMPAGNE, F. & MARASLI, B. 1986 *On the large-scale structures in two-dimensional, small-deficit, turbulent wakes*, vol. 168.
- YANG, J. & BALARAS, E. 2006 An embedded-boundary formulation for large-eddy simulation of turbulent flows interacting with moving boundaries. *J. Comput. Phys.* **215** (1), 12–40.

ZHOU, Q. & DIAMESSIS, P. J. 2019 Large-scale characteristics of stratified wake turbulence at varying Reynolds number. *Physical Review Fluids* **4** (8), 1–30.