

Kaon production in high multiplicity events at the LHC

Yuri N. LIMA,^{1,*} André V. GIANNINI,^{2,†} and Victor P. GONÇALVES^{3,4,1,‡}

¹*Institute of Physics and Mathematics, Federal University of Pelotas,
Postal Code 354, 96010-900, Pelotas, RS, Brazil*

²*Instituto de Física Gleb Wataghin, Universidade Estadual de Campinas,
Rua Sérgio Buarque de Holanda 777, 13083-859 São Paulo, Brazil*

³*Institut für Theoretische Physik, Westfälische Wilhelms-Universität Münster,
Wilhelm-Klemm-Straße 9, D-48149 Münster, Germany*

⁴*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

The production of the K_S^0 meson in high multiplicity pp collisions at $\sqrt{s} = 13$ TeV is investigated considering the hybrid formalism and the solution of the running coupling Balitsky - Kovchegov equation. The associated cross section is estimated and compared with the experimental data for the transverse momentum spectrum. Moreover, we analyze the self-normalized yields of K_S^0 mesons as a function of the multiplicity of coproduced charged hadrons and demonstrate that a steep increasing is theoretically predicted. A comparison with the ALICE data is presented considering two distinct solutions of the BK equation.

PACS numbers:

Keywords: Particle production, Color Glass Condensate Framework, Hybrid factorization formalism

I. INTRODUCTION

In recent years, different experimental collaborations at the Relativistic Heavy Ion Collider (RHIC) and at the Large Hadron Collider (LHC) have found that the J/Ψ , D , K_S^0 and Λ yields observed in proton - proton (pp) collisions grow rapidly as a function of the multiplicities of co - produced charged particles [1–7]. Although the development of a theoretical framework for high multiplicity events has started several decades ago, the description of the current data using a unified approach remains a challenge (See e.g. Refs. [8–23]). One the main open questions is if the modification observed in high multiplicity events compared to the minimum bias case is due to either initial or final state effects or both. The similarity of multiplicity enhancements observed in the charm and strange sectors favours the interpretation that the behaviour is due to initial state effects, but models based on very distinct underlying assumptions and physical mechanisms are also able to describe the current data. Considering the recent restart of the LHC, we expect to have in the forthcoming years a larger amount of data for the production of different hadrons in high multiplicity events, which will allow us to improve our understanding of the mechanism that generates the multiplicity enhancement.

One of the promising frameworks to describe the particle production in low and high - multiplicity events at the LHC is the Color Glass Condensate (CGC) formalism [24], which is an effective field theory that states that a dense system of partons is produced in hadronic collisions. Such system is characterized by a new scale, the saturation scale Q_s , which increases as function of the energy, atomic number and multiplicity (See e.g. Refs. [25, 26]). In this framework, high multiplicity events are attributed to the presence of rare parton configurations (hot spots) in the hadrons that participate of the collision. Such highly occupied gluon states are characterized by larger saturation scales in comparison to the typical configurations present in minimum bias events. Naturally, in the CGC framework, low and high multiplicity approaches are expected to be described in a unified way, but with the scattering amplitude being estimated for different values of the saturation scale. Such assumption is the starting point of the studies performed in Refs. [16–23] and for the analysis that will be carried out in this paper, where we will focus on the production of the K_S^0 meson in high multiplicity events at the LHC. The strangeness enhancement has already been discussed in the literature in Ref. [20], where the authors have applied for the K production, the dipole approach developed for the calculation of open charm and bottom states. As a consequence, only the $gg \rightarrow s\bar{s}$ channel is taken

*Electronic address: limayuri.91@gmail.com

†Electronic address: giannini@ifi.unicamp.br

‡Electronic address: barros@ufpel.edu.br




FIG. 1: Representation of the K_S^0 meson production in the hybrid formalism. One has that a parton of the incident hadron interacts with the target and subsequently hadronizes in the K_S^0 meson.

into account and the presence of a strange (anti) quark in the wavefunction of the incident hadrons is disregarded. Moreover, these authors have assumed a phenomenological model for the description of the dipole - hadron interaction. In contrast, in this paper we will consider the hybrid formalism [27] to treat the K_S^0 production and the solution of the running coupling Balitsky - Kovchegov (BK) equation to describe the evolution of the dipole - hadron scattering amplitude [28–32]. One has that the hybrid formalism takes into account of the contribution associated to the gluon and quark - initiated channels and, over the last decade, has been extensively applied for the description of light particle production in hadronic colliders, with its predictions describing with reasonable success the RHIC and LHC data (See e.g. Refs. [33–48]). As we will demonstrate below, such approach is also able to describe the current minimum bias data for the transverse momentum spectrum of the K_S^0 meson, which motivates its application for high multiplicity events.

This paper is organized as follows. In the next Section, we will present a brief review of the hybrid formalism and will discuss the ingredients considered in our calculations. In Section III, we will present our predictions for the transverse momentum spectrum of the K_S^0 meson and the contribution of the gluon and quark - initiated processes will also be estimated. Moreover, we will analyze the dependence of the K_S^0 and charged particle yields in the value of the saturation scale, considering distinct transverse momentum ranges and two distinct solutions of the BK equation. Predictions for the multiplicity dependence of the K_S^0 meson production will be compared with the experimental data from ALICE. Finally, in Section IV we will summarize our main results and conclusions.

II. KAON PRODUCTION IN THE HYBRID FORMALISM

Over the last decade, the hybrid formalism has been developed, improved and applied for the description of hadron production in hadronic collisions at RHIC and LHC [33–48]. The basic assumption is that the cross section for particle production can be expressed as a convolution of the standard parton distributions, the scattering amplitude (which includes the high-density effects) and the parton fragmentation functions. In Fig. 1 we present the representation of the hybrid formalism for the K_S^0 meson production. Taking into account of the gluon and quark - initiated subprocesses, one has that the invariant yield for single-inclusive K_S^0 production in hadron-hadron processes will be described in the CGC formalism as follows [27]

$$\begin{aligned} \frac{dN_{K_S^0}}{dyd^2p_T} = & \frac{\mathcal{K}}{(2\pi)^2} \int_{x_F}^1 dx_1 \frac{x_1}{x_F} \left[\sum_{q=u,d,s} f_{q/p}(x_1, \mu^2) \tilde{\mathcal{N}}_F(k_T, x_2) D_{K_S^0/q} \left(\frac{x_F}{x_1}, \mu^2 \right) \right. \\ & \left. + f_{g/p}(x_1, \mu^2) \tilde{\mathcal{N}}_A(k_T, x_2) D_{K_S^0/g} \left(\frac{x_F}{x_1}, \mu^2 \right) \right]. \end{aligned} \quad (1)$$

One has that p_T , y and x_F are the transverse momentum, rapidity and the Feynman- x of the produced hadron, respectively. Moreover, x_1 denotes the momentum fraction of a projectile parton, $x_F = \frac{p_T}{\sqrt{s}} e^y$ and $k_T = \frac{x_1}{x_F} p_T$, with the momentum fraction of the target parton being given by $x_2 = x_1 e^{-2y}$. One also has that $f_{i/p}(x_1, \mu^2)$ are the projectile parton distribution functions and $D_{K_S^0/i}(z, \mu^2)$ are the parton fragmentation functions into the K_S^0 meson. As in previous studies [33–48], we will assume that the parton distribution and fragmentation functions evolve according to the DGLAP evolution equations [49] and obey the momentum sum-rule. In particular, we will consider the CT14 [50] and AKK08 [51] parametrizations for these quantities. The factorization scale μ^2 will be assumed as being $\mu^2 = \max(Q_s^2, p_T^2)$. Previous results derived using the hybrid formalism have pointed out that in order to




FIG. 2: (a) Dependence of the self-normalized yield for the K_S^0 -meson production on the transverse momentum (p_T) compared to experimental data provided by the ALICE collaboration [6]. (b) Contribution of the gluon and quark - initiated subprocesses for the K_S^0 -meson production derived using the UGD g1.119 model.

describe the data for different kinematical ranges one has to adjust the normalization by a \mathcal{K} -factor, which can be energy and rapidity dependent. Such factor is interpreted as taking account of higher order corrections and/or of other dynamical effects not included in the CGC formulation. Such uncertainty will not affect the analysis of high multiplicity events, since this factor is expected to be same in low and high multiplicity events and, consequently, it cancels in our prediction for the ratio between the results for high multiplicity and minimum bias events.

One has that the predictions for the K_s^0 yield are also strongly dependent on the description of the fundamental and adjoint representations of the forward dipole amplitude, $\tilde{\mathcal{N}}_F$ and $\tilde{\mathcal{N}}_A$, respectively. In the CGC framework, the forward dipole scattering amplitudes encodes all the information about the hadronic scattering, and thus about the non-linear and quantum effects in the hadron wave function. Such quantities can be expressed either in the momentum or in the position spaces, with the representations related by

$$\tilde{\mathcal{N}}_{A,F}(x, p_T) = \int d^2r e^{i\vec{p}_T \cdot \vec{r}} [1 - \mathcal{N}_{A,F}(x, r)] , \quad (2)$$

In our analysis we will consider the solutions of the running coupling Balitsky–Kovchegov equation for $\mathcal{N}_{A,F}(x, r)$ derived in Ref. [37] considering two distinct initial conditions characterized by an anomalous dimension larger than unity. In particular, we will consider the sets “ γ 1.101” and “ γ 1.119” from Ref. [37].

Following previous studies of the high multiplicity events using the CGC formalism, we will assume that the particle production mechanism is the same for low and high - multiplicity events, with the main difference being the saturation scale present in these two classes of events. In other words, we will assume that Eq. (1) is valid for both classes, and that the high multiplicity configurations can be approximated by increasing the value of Q_s as follows $Q_s^2(x, n) = n \cdot Q_s^2(x)$, where n characterizes the multiplicity. Such quantity is given approximately by the multiplicity of charged particles weighted by their minimum bias value: $n \approx dN_{ch}/d\eta / \langle dN_{ch}/d\eta \rangle$. As in Ref. [13], we will solve the BK equation considering multiples of initial saturation scale at $x = 0.01$, $Q_0^2 = 0.168 \text{ GeV}^2$, which has been determined from fits to the minimum bias ep HERA data. Such modification naturally implies larger values for the saturation scale probed in a given event, which depends on x and has its evolution determined by the BK equation.

III. RESULTS AND DISCUSSION

Initially, let’s compare our predictions for the transverse momentum distribution of the K_S^0 meson with the experimental data from the ALICE collaboration [6] for the self normalized yields $(1/N_{ev})d^2N/dydp_T$. We will consider the solutions of the BK equation derived assuming two distinct initial conditions [37], which will be denoted by “UGD g1.101” and “UGD g1.119” hereafter. As already emphasized in Ref. [20], the prediction of the normalization of the self normalized yields is strongly affected by non-perturbative effects, since this quantity is dominated by the contribution of kaon production at very low p_T , which cannot be evaluated reliably in our approach. As a consequence, in our analysis we will adjust the normalization of our predictions in order to describe the data. In Fig. 2(a) we present our results. One has that the shape of the spectrum can be quite well described by both models considered. The




FIG. 3: Relative multiplicity of (a) K_S^0 mesons and (b) charged particles as a function of $Q_{s,0}^2/Q_0^2$, considering different ranges of the transverse momentum p_T . (c) Ratio between the predictions for the production of K_S^0 mesons and charged particles as a function of $Q_{s,0}^2/Q_0^2$. Results derived using the UGD g1.119.

contribution of the gluon and quark - initiated channels for the K_S^0 production is presented separately in Fig. 2(b) for the UGD g1.119 model. One has verified that similar results are obtained for the g1.101 model. Our results indicate that the gluon channel is dominant in the kinematical range considered, which can explain why the approach used in Ref. [20] is also able to describe the ALICE data.

In order to estimate the impact of varying the initial saturation scale, in Figs. 3 (a) and (b) we present our results for the relative multiplicity of K_S^0 mesons and charged hadrons, respectively, for a pp collision at 13 TeV as a function of $Q_{s,0}^2/Q_0^2$, with $Q_0^2 = 0.168 \text{ GeV}^2$. We present our predictions, derived assuming the UGD g1.119 model, for different range of the transverse momentum p_T . For charged hadrons, we also have estimated the yield using the hybrid formalism, considering the contribution of charged pions, baryons and strange mesons. It is important to emphasize that we have verified that the corresponding predictions describe the current data for the inclusive hadron production at central rapidities in pp collisions at the LHC. One has that the predictions for K_S^0 mesons and charged hadrons are similar, but the increasing with $Q_{s,0}^2/Q_0^2$ is dependent of the p_T range considered. Such result is expected since the impact of the nonlinear effects are strongly dependent if Q_s^2 is larger or smaller than p_T^2 . Events where $Q_s^2 \gtrsim p_T^2$ are expected to be determined by the nonlinear QCD dynamics. Therefore, a larger value of the minimum value of p_T , implies a reduction of the number of events produced in the saturated regime. The differences between the predictions for K_S^0 mesons and charged hadrons can be quantified by calculating the ratio between the corresponding results,

$$R_{K_S^0,ch} = \frac{dN_{K_S^0}/dy / \langle dN_{K_S^0}/dy \rangle}{dN_{ch}/d\eta / \langle dN_{ch}/d\eta \rangle}, \quad (3)$$

with the results being presented in Fig. 3 (c). One has that when only large p_T values are considered, the ratio is consistent with unity, indicating that the production of K_S^0 mesons and charged hadrons are similarly affected by




FIG. 4: Correlation between the normalized K_S^0 and charged hadron yields in pp collisions at 13 TeV derived considering two distinct solutions of the BK equation. Data from the ALICE Collaboration [6].

the saturation effects. On the other hand, if events with lower transverse momentum are included, one has that the behavior with $Q_{s,0}^2/Q_0^2$ becomes final state dependent.

In Fig. 4 we present a comparison between our predictions for the multiplicity dependence of K_S^0 mesons and the experimental data from the ALICE Collaboration [6]. It is important to emphasize that these data were collected at central rapidities $|\eta| \leq 0.5$ and for transverse momentum bins in the $4.0 < p_T < 12$ GeV range. We present our predictions for the two initial conditions of the BK equation considered in our analysis. Our results indicate that the corresponding predictions are similar for low multiplicities, $dN_{ch}/d\eta / \langle dN_{ch}/d\eta \rangle \leq 2.0$, but differ for larger multiplicities. One has that the UGD g1.119 model provides a better description of the data, but also fails in the description of the data for the largest value of the multiplicity. A similar overshooting is also observed in Ref. [13] for the D meson production when events with larger transverse momentum are considered and in Ref. [20] for the K_S^0 meson production. One possible interpretation of this overestimation in the hybrid approach is that for high multiplicities, saturation effects cannot be neglected in the projectile. Another possible effect, not considered in our analysis, is the modification of the fragmentation function in high multiplicity events. Our results indicate that such possibilities should be analyzed in the future, which we intend to perform in a forthcoming study.

IV. CONCLUSIONS

The description of high multiplicity events observed in small collision systems still remains an open question. In recent years, new experimental data have renewed the interest in the topic, strongly motivating a large phenomenology based on models that take into account either initial or final state effects or both. In this paper we have focused on initial state effects, as described by the CGC framework, and applied the hybrid formalism for the description of the K_S^0 production in high multiplicity events present in pp collisions at $\sqrt{s} = 13$ TeV. Such formalism has been successfully applied for the description of the particle production in hadronic colliders and takes into account of the gluon and quark - initiated subprocesses. Moreover, its predictions can be derived using the solutions of the running coupling BK equation. It has been demonstrated that this formalism is also able to describe the current data for the transverse momentum spectrum and that the K_S^0 production is dominated by the gluon - initiated subprocess. Following the study performed in Ref. [13], which is able to describe the D meson and J/Ψ production at high multiplicities, we derived our predictions assuming that high multiplicity configurations can be approximated by increasing the value of the initial saturation scale in the BK evolution. The dependence of the relative multiplicity of K_S^0 mesons and charged particles with the initial condition for the BK evolution equation has been studied and we verified that they are slightly different, depending of the p_T range considered. Finally, we have compared our predictions with the ALICE data for the multiplicity dependence of K_S^0 mesons. Our results indicate that the hybrid formalism can describe the current data for values of multiplicity smaller than 2.5, but overestimate the data for larger multiplicities, which indicate that other effects and/or higher order corrections should be taken into account in this kinematical range. Surely, more data for the production of strange and charmed mesons in very high multiplicity

events will be very useful to improve our understanding of the dynamical effects present in the new kinematical range.

Acknowledgements

This work was partially supported by INCT-FNA (Process No. 464898/2014-5). V.P.G. was partially supported by the CAS President's International Fellowship Initiative (Grant No. 2021VMA0019) and by CNPq, CAPES and FAPERGS. Y.N.L. was partially financed by CAPES (process 001). The work of A.V.G. has been supported by FAPESP through grants 17/05685-2 and 21/04924-9.

-
- [1] J. Adam *et al.* [ALICE], JHEP **09**, 148 (2015)
- [2] D. Adamová *et al.* [ALICE], Phys. Lett. B **776**, 91-104 (2018)
- [3] J. Adam *et al.* [STAR], Phys. Lett. B **786**, 87-93 (2018)
- [4] S. Acharya *et al.* [ALICE], Phys. Lett. B **810**, 135758 (2020)
- [5] S. Acharya *et al.* [ALICE], JHEP **09**, 162 (2020)
- [6] ALICE Collaboration, Eur. Phys. J. C **80**, 167 (2020).
- [7] S. Acharya *et al.* [ALICE], JHEP **06**, 015 (2022)
- [8] E. G. Ferreira and C. Pajares, Phys. Rev. C **86**, 034903 (2012)
- [9] B. Z. Kopeliovich, H. J. Pirner, I. K. Potashnikova, K. Reygers and I. Schmidt, Phys. Rev. D **88**, no.11, 116002 (2013)
- [10] T. Lang and M. Bleicher, Phys. Rev. C **87**, no.2, 024907 (2013)
- [11] K. Werner, B. Guiot, I. Karpenko and T. Pierog, Phys. Rev. C **89**, no.6, 064903 (2014)
- [12] E. G. Ferreira and C. Pajares, [arXiv:1501.03381 [hep-ph]].
- [13] Y. Q. Ma, P. Tribedy, R. Venugopalan and K. Watanabe, Phys. Rev. D **98**, no.7, 074025 (2018)
- [14] A. Kovner and V. V. Skokov, Phys. Rev. D **98**, no.1, 014004 (2018)
- [15] S. G. Weber, A. Dubla, A. Andronic and A. Morsch, Eur. Phys. J. C **79**, no.1, 36 (2019)
- [16] E. Levin, I. Schmidt and M. Siddikov, Eur. Phys. J. C **80**, no.6, 560 (2020)
- [17] B. Z. Kopeliovich, H. J. Pirner, I. K. Potashnikova, K. Reygers and I. Schmidt, Phys. Rev. D **101**, no.5, 054023 (2020)
- [18] E. Gotsman and E. Levin, Eur. Phys. J. C **81**, no.2, 99 (2021)
- [19] M. Siddikov and I. Schmidt, Phys. Rev. D **104**, no.1, 016023 (2021)
- [20] M. Siddikov and I. Schmidt, Phys. Rev. D **104**, no.1, 016024 (2021)
- [21] M. Siddikov and I. Schmidt, Phys. Rev. D **104**, no.1, 016032 (2021)
- [22] T. Stebel and K. Watanabe, Phys. Rev. D **104**, no.3, 034004 (2021)
- [23] F. Salazar, B. Schenke and A. Soto-Ontoso, Phys. Lett. B **827**, 136952 (2022)
- [24] F. Gelis, E. Iancu, J. Jalilian-Marian and R. Venugopalan, Ann. Rev. Nucl. Part. Sci. **60**, 463-489 (2010)
- [25] Y. V. Kovchegov and E. Levin, "Quantum chromodynamics at high energy," Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **33**, 1-350 (2012) Cambridge University Press, 2012,
- [26] A. Morreale and F. Salazar, Universe **7**, no.8, 312 (2021)
- [27] A. Dumitru, A. Hayashigaki and J. Jalilian-Marian, Nucl. Phys. A **765**, 464 (2006); Nucl. Phys. A **770**, 57 (2006).
- [28] I. I. Balitsky, Nucl. Phys. **B463**, 99 (1996); Phys. Rev. Lett. **81**, 2024 (1998); Phys. Rev. D **60**, 014020 (1999); I. I. Balitsky, Phys. Lett. B **518**, 235 (2001); I.I. Balitsky and A.V. Belitsky, Nucl. Phys. **B629**, 290 (2002).
- [29] Y.V. Kovchegov, Phys. Rev. D **60**, 034008 (1999); Phys. Rev. D **61** 074018 (2000).
- [30] Y. V. Kovchegov and H. Weigert, Nucl. Phys. A **784**, 188 (2007); Nucl. Phys. A **789**, 260 (2007); Y. V. Kovchegov, J. Kuokkanen, K. Rummukainen and H. Weigert, Nucl. Phys. A **823**, 47 (2009).
- [31] J. L. Albacete and Y. V. Kovchegov, Phys. Rev. D **75**, 125021 (2007).
- [32] I. Balitsky, Phys. Rev. D **75**, 014001 (2007); I. Balitsky and G. A. Chirilli, Phys. Rev. D **77**, 014019 (2008).
- [33] D. Boer, A. Utermann and E. Wessels, Phys. Rev. D **77**, 054014 (2008)
- [34] M. A. Betemps and V. P. Goncalves, JHEP **09**, 019 (2008)
- [35] J. L. Albacete and C. Marquet, Phys. Lett. B **687**, 174-179 (2010)
- [36] T. Altinoluk and A. Kovner, Phys. Rev. D **83**, 105004 (2011)
- [37] J. L. Albacete, A. Dumitru, H. Fujii and Y. Nara, Nucl. Phys. A **897**, 1-27 (2013)
- [38] G. A. Chirilli, B. W. Xiao and F. Yuan, Phys. Rev. Lett. **108**, 122301 (2012)
- [39] Z. B. Kang, I. Vitev and H. Xing, Phys. Rev. Lett. **113**, 062002 (2014)
- [40] T. Altinoluk, N. Armesto, G. Beuf, A. Kovner and M. Lublinsky, Phys. Rev. D **91**, no.9, 094016 (2015)
- [41] E. Iancu, A. H. Mueller and D. N. Triantafyllopoulos, JHEP **12**, 041 (2016)
- [42] F. O. Durães, A. V. Giannini, V. P. Goncalves and F. S. Navarra, Phys. Rev. C **94**, no.2, 024917 (2016)
- [43] B. Ducloué, E. Iancu, T. Lappi, A. H. Mueller, G. Soyez, D. N. Triantafyllopoulos and Y. Zhu, Phys. Rev. D **97**, no.5, 054020 (2018)
- [44] B. Ducloué, T. Lappi and Y. Zhu, Phys. Rev. D **95**, no.11, 114007 (2017)
- [45] F. Carvalho, A. V. Giannini, V. P. Goncalves and F. S. Navarra, Phys. Rev. D **96**, no.9, 094002 (2017)

- [46] H. Y. Liu, Y. Q. Ma and K. T. Chao, Phys. Rev. D **100**, no.7, 071503 (2019)
- [47] E. Iancu and Y. Mulian, JHEP **03**, 005 (2021)
- [48] Y. Shi, L. Wang, S. Y. Wei and B. W. Xiao, Phys. Rev. Lett. **128**, no.20, 202302 (2022)
- [49] V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. **15**, 438 (1972); G. Altarelli and G. Parisi, Nucl. Phys. **B126**, 298 (1977); Yu.L. Dokshitzer, Sov. Phys. JETP **46**, 641 (1977).
- [50] S. Dulat *et al.*, Phys. Rev. D **93**, 033006 (2016).
- [51] S. Albino, B. A. Kniehl and G. Kramer, Nucl. Phys. B **803**, 42-104 (2008)