Kengo Matsumoto Department of Mathematics Joetsu University of Education Joetsu, 943-8512, Japan

October 13, 2022

Abstract

We introduce the notion of K-theoretic duality for extensions of separable unital nuclear C^* -algebras by using K-homology long exact sequence and cyclic six term exact sequence for K-theory groups of extensions. We then prove that the Toeplitz extension \mathcal{T}_A of a Cuntz-Krieger algebra \mathcal{O}_A is the K-theoretic dual of the Toeplitz extension \mathcal{T}_{A^t} of the Cuntz-Krieger algebra \mathcal{O}_A for the transposed matrix A^t of A. A pair of isomorphic Cuntz-Krieger algebras \mathcal{O}_A and \mathcal{O}_B does not necessarily yield the isomorphic pair of \mathcal{O}_{A^t} and \mathcal{O}_{B^t} . However, as an application, we may show that two Toeplitz algebras \mathcal{T}_A and \mathcal{T}_B are isomorphic as C^* -algebras if and only if the Toeplitz algebras \mathcal{T}_{A^t} of their transposed matrices are isomorphic.

Mathematics Subject Classification: Primary 19K33; Secondary 46L80.

Keywords and phrases: Extension, C^* -algebra, K-homology, K-group, Toeplitz algebra, Cuntz–Krieger algebra.

1 Introduction

Let \mathcal{A} be a separable unital C^* -algeba. Let us denote by $\mathcal{K}(H)$ the C^* -algebra of compact operators on a separable infinite dimensional Hilbert space H. An extension of a C^* algebra \mathcal{A} means a short exact sequence

$$0 \longrightarrow \mathcal{K}(H) \longrightarrow \mathcal{E} \longrightarrow \mathcal{A} \longrightarrow 0 \tag{1.1}$$

of C^* -algebras for which a unital C^* -algebra \mathcal{E} contains $\mathcal{K}(H)$ as an essential ideal. Classification of extensions of C^* -algebras have been developing from many sides, K-theory of C^* -algebras, non commutative geometry, structure theory of C^* -algebras, operator theory, homotopy theory in algebraic topology, etc. (see text books [2], [9], [16], etc.). In this paper, we will study 'duality' of extensions of C^* -algebras from the viewpoints of K-homology theory and K-group theory of C^* -algebras.

The notion of K-theoretic 'dual' of a C^* -algebra was first introduced by W. L. Paschke in [23] from the viewpoint of a duality of Ext-groups and K-groups. After Paschke, N. Higson refined the dual C^* -algebra written $\mathfrak{D}(\mathcal{A})$ of a C^* -algebra \mathcal{A} to define K-homology groups $K^*(\mathcal{A})$ for \mathcal{A} ([15], cf. [16]). By using the K-homology groups, Higson and Higson– Roe in [16] studied analytic K-homology theory for C^* -algebras comparing Kasparov's KK-theory ([20], [21]). Kaminker–Putnam in [17] introduced the notion of K-theoretic duality from the viewpoint of Spanier-Whitehead dual in topology and showed that the Cuntz–Krieger algebras \mathcal{O}_A and \mathcal{O}_{A^t} are the K-theoretic duality pair (cf. [18],[19]).

In this paper, we introduce the notion of K-theoretic duality for extensions of a separable unital nuclear C^* -algebras. Let us denote by $\widetilde{K}^*(\mathcal{A})$ and $K^*(\mathcal{A})$ the reduced Khomology and the unreduced K-homology of a C^* -algebra \mathcal{A} as in Higson–Roe's text book [16]. Since $\widetilde{K}^1(\mathcal{A})$ and $K^1(\mathcal{A})$ are identified with the strong extension group $\operatorname{Ext}_{s}(\mathcal{A})$ and the weak extension group $\operatorname{Ext}_{w}(\mathcal{A})$ for a separable unital nuclear C^* -algebra \mathcal{A} , the extension (1.1) of \mathcal{A} gives rise to an element $[\mathcal{E}]_s$ in $\widetilde{K}^1(\mathcal{A})$ and an element $[\mathcal{E}]_w$ in $K^1(\mathcal{A})$, respectively. Two extensions

$$0 \longrightarrow \mathcal{K}(H) \longrightarrow \mathcal{E} \longrightarrow \mathcal{A} \longrightarrow 0, \qquad 0 \longrightarrow \mathcal{K}(H) \longrightarrow \mathcal{E}' \longrightarrow \mathcal{A}' \longrightarrow 0$$
(1.2)

are said to be a K-theoretic duality pair (Definition 6.1) if the K-homology long exact sequence for \mathcal{A} (resp. \mathcal{A}')

$$\cdots \longrightarrow \widetilde{\mathrm{K}}^{p}(\mathcal{A}) \longrightarrow \mathrm{K}^{p}(\mathcal{A}) \longrightarrow \mathrm{K}^{p}(\mathbb{C}) \longrightarrow \widetilde{\mathrm{K}}^{p+1}(\mathcal{A}) \longrightarrow \mathrm{K}^{p+1}(\mathcal{A}) \longrightarrow \cdots$$
$$(\operatorname{resp.} \cdots \longrightarrow \widetilde{\mathrm{K}}^{p}(\mathcal{A}') \longrightarrow \mathrm{K}^{p}(\mathcal{A}') \longrightarrow \mathrm{K}^{p}(\mathbb{C}) \longrightarrow \widetilde{\mathrm{K}}^{p+1}(\mathcal{A}') \longrightarrow \mathrm{K}^{p+1}(\mathcal{A}') \longrightarrow \cdots)$$

is realized as the K-theory long exact sequence for the other extension

$$\cdots \longrightarrow \mathrm{K}_{p+1}(\mathcal{E}') \longrightarrow \mathrm{K}_{p+1}(\mathcal{A}') \longrightarrow \mathrm{K}_p(\mathcal{K}(H)) \longrightarrow \widetilde{\mathrm{K}}_p(\mathcal{E}') \longrightarrow \mathrm{K}_p(\mathcal{A}') \longrightarrow \cdots$$
$$(\mathrm{resp.} \cdots \longrightarrow \mathrm{K}_{p+1}(\mathcal{E}) \longrightarrow \mathrm{K}_{p+1}(\mathcal{A}) \longrightarrow \mathrm{K}_p(\mathcal{K}(H)) \longrightarrow \widetilde{\mathrm{K}}_p(\mathcal{E}) \longrightarrow \mathrm{K}_p(\mathcal{A}) \longrightarrow \cdots)$$

for $p = 0, 1 \pmod{2}$. If in particular, we may find the above realizations between $K^0(\mathbb{C})$ and $K_0(\mathcal{K}(H))$ in order preserving way, and between $\widetilde{K}^1(\mathcal{A})$ and $K_0(\mathcal{E}')$ (resp. between $\widetilde{K}^1(\mathcal{A}')$ and $K_0(\mathcal{E})$) in sending $[\mathcal{E}]_s$ to $-[1_{\mathcal{E}'}]$ (resp. $[\mathcal{E}']_s$ to $-[1_{\mathcal{E}}]$), then the K-theoretic duality pair is said to be *strong* K-*theoretic duality pair* (Definition 6.1), where $[1_{\mathcal{E}'}], [1_{\mathcal{E}}]$ denote the classes in $K_0(\mathcal{E}'), K_0(\mathcal{E})$ of the units of $\mathcal{E}', \mathcal{E}$, respectively.

For a Cuntz-Krieger algebra \mathcal{O}_A , there exists a specific extension

$$\tau_A: \qquad 0 \longrightarrow \mathcal{K}(H) \xrightarrow{\iota_A} \mathcal{T}_A \xrightarrow{\pi_A} \mathcal{O}_A \longrightarrow 0$$
(1.3)

called the Toeplitz extension of \mathcal{O}_A that was introduced in [12] and [14] (cf. [17]). We will prove that the K-homology long exact sequence for \mathcal{O}_A is realized as the K-theory cyclic six term exact sequence for the Toeplitz extension τ_{A^t} defined by the the transposed matrix A^t of A such that the class $[\mathcal{T}_A]_s \in \widetilde{K}(\mathcal{O}_A)$ of the Toeplitz extension τ_A exactly corresponds to the class $-[1_{\mathcal{T}_A t}] \in K_0(\mathcal{T}_A t)$ of the minus of the unit $1_{\mathcal{T}_A t}$ of $\mathcal{T}_A t$ under the identification between $K^0(\mathbb{C})$ and $K_0(\mathcal{K}(H))$ in order preserving way (Theorem 5.3). We then know that the Toeplitz extension of a Cuntz-Krieger algebra is the K-theoretic dual of the Toeplitz extension of the Cuntz-Krieger algebra for the transposed matrix in the following way.

Theorem 1.1. Let A be an irreducible non permutation matrix with entries in $\{0, 1\}$. The Toeplitz extensions τ_A and τ_{A^t} of the Cuntz-Krieger algebras \mathcal{O}_A and \mathcal{O}_{A^t} are strong K-theoretic duality pair. In general an isomorphic pair $\mathcal{O}_A \cong \mathcal{O}_B$ of Cuntz–Krieger algebras does not necessarily yield an isomorphic pair $\mathcal{O}_{A^t} \cong \mathcal{O}_{B^t}$ of the Cuntz–Krieger algebras of their transposed matrices. For example, the Cuntz–Krieger algebra \mathcal{O}_A for the matrix $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$ is isomorphic to \mathcal{O}_3 , whereas \mathcal{O}_{A^t} is not isomorphic to \mathcal{O}_3 (cf. [13], [27]). Eilers–Restroff– Ruiz in [10] (cf. [11]) showed that certain extensions of classifiable C^* -algebras are strongly classified by their associated cyclic six term exact sequences in K-groups together with the positive cones of the K₀-groups of the ideals and quotients. By applying their classification result in our setting, as an application of Theorem 1.1, we will prove the following theorem.

Theorem 1.2. Let A, B be irreducible non permutation matrices with entries in $\{0, 1\}$. The following are equivalent.

(i) There exist isomorphisms $\Phi : \mathcal{O}_A \longrightarrow \mathcal{O}_B$ and $\Psi : \mathcal{T}_A \longrightarrow \mathcal{T}_B$ of C^* -algebras such that the diagram

$$\begin{array}{cccc} \mathcal{T}_A & \xrightarrow{\pi_A} & \mathcal{O}_A \\ \Psi & & & \downarrow \Phi \\ \mathcal{T}_B & \xrightarrow{\pi_B} & \mathcal{O}_B \end{array} \tag{1.4}$$

commutes.

(ii) There exist isomorphisms $\Phi^t : \mathcal{O}_{A^t} \longrightarrow \mathcal{O}_{B^t}$ and $\Psi^t : \mathcal{T}_{A^t} \longrightarrow \mathcal{T}_{B^t}$ of C^* -algebras such that the diagram

commutes.

Since an isomorphism from \mathcal{T}_A to \mathcal{T}_B automatically induces the commutative diagram in Theorem 1.2 (i), we have the following corollary.

Corollary 1.3. Let A, B be irreducible non permutation matrices with entries in $\{0, 1\}$. the Toeplitz algebras \mathcal{T}_A and \mathcal{T}_B are isomorphic if and only if the Toeplitz algebras \mathcal{T}_{A^t} and \mathcal{T}_{B^t} defined by their transposed matrices are isomorphic.

We will finally present a classification result for the Toeplitz algebras (Proposition 7.4) with several examples (Section 8).

2 Preliminaries for Ext-groups and K-homology groups

In this section, we briefly give an introduction of K-homology groups based on [15] and [16], see [16] for detail. In what follows, \mathcal{A} stands for a separable unital nuclear C^* -algebra.

2.1 Ext-groups

It is well-known that an extension (1.1) bijectively corresponds to a unital *-monomorphism $\tau: \mathcal{A} \longrightarrow \mathcal{Q}(H)$ that is called a Busby invariant, where $\mathcal{Q}(H)$ denotes the Calkin algebra $\mathcal{B}(H)/\mathcal{K}(H)$. We also call such a *-monomorphism $\tau: \mathcal{A} \longrightarrow \mathcal{Q}(H)$ an extension of \mathcal{A} . Two extensions $\tau_i : \mathcal{A} \longrightarrow \mathcal{Q}(H), i = 1, 2$ are said to be weak equivalent if there exists a unitary $u \in \mathcal{Q}(H)$ such that $\tau_2(a) = u\tau_1(a)u^*, a \in \mathcal{A}$. If in particular the above unitary is taken as $u = \pi(U)$ for some unitary $U \in \mathcal{B}(H)$, where $\pi : \mathcal{B}(H) \longrightarrow \mathcal{Q}(H)$ denotes the natural quotient map, then the extensions $\tau_i : \mathcal{A} \longrightarrow \mathcal{Q}(H), i = 1, 2$ are said to be strong equivalent. Let us denote by $\operatorname{Ext}_{s}(\mathcal{A})$ and $\operatorname{Ext}_{w}(\mathcal{A})$ the strong equivalence classes and the weak equivalence classes of extensins of \mathcal{A} , respectively. The class of a unital *-monomorphism $\tau : \mathcal{A} \longrightarrow \mathcal{Q}(H)$ in $\operatorname{Ext}_{s}(\mathcal{A})$ is denoted by $[\tau]_{s}$, and similarly $[\tau]_{w}$ in $\operatorname{Ext}_{w}(\mathcal{A})$. Fix an identification $H \oplus H = H$ which induces an embleding $\mathcal{Q}(H) \oplus \mathcal{Q}(H) \hookrightarrow \mathcal{Q}(H)$. The sum of extensions $\tau_1 \oplus \tau_2$ are defined by a direct sum $\tau_1 \oplus \tau_2$ through the embedding $\mathcal{Q}(H) \oplus \mathcal{Q}(H) \hookrightarrow \mathcal{Q}(H)$. It is well-known that both $\operatorname{Ext}_{\mathrm{s}}(\mathcal{A})$ and $\operatorname{Ext}_{\mathrm{w}}(\mathcal{A})$ become abelian semigroups, and also they are abelian groups for nuclear C^* -algebra \mathcal{A} (cf. [1], [3], [5], [9], etc.). They are called the strong extension group for \mathcal{A} and the weak extension group for \mathcal{A} . Let us denote by $q_{\mathcal{A}} : \operatorname{Ext}_{s}(\mathcal{A}) \longrightarrow \operatorname{Ext}_{w}(\mathcal{A})$ the surjective homomorphism induced by a natural quotient map. As in [16] and [25], there exists a homomorphism $\iota_{\mathcal{A}}: \mathbb{Z} \longrightarrow \operatorname{Ext}_{\mathrm{s}}(\mathcal{A})$ such that the sequence

$$\mathbb{Z} \xrightarrow{\iota_{\mathcal{A}}} \operatorname{Ext}_{s}(\mathcal{A}) \xrightarrow{q_{\mathcal{A}}} \operatorname{Ext}_{w}(\mathcal{A})$$
(2.1)

is exact at the middle, and hence the quotient group $\operatorname{Ext}_{s}(\mathcal{A})/\iota_{\mathcal{A}}(\mathbb{Z})$ is isomorphic to $\operatorname{Ext}_{w}(\mathcal{A})$.

2.2 K-homology groups

Let \mathcal{A} be a separable unital C^* -algebra. Following [16], a non degenerate representation $\rho : \mathcal{A} \longrightarrow \mathcal{B}(H)$ on a separable infinite dimensional Hilbert space H is said to be *ample* if $\rho(a)$ never belongs to $\mathcal{K}(H)$ unless a = 0. Hence any non degenerate representation of a simple C^* -algebra is always ample if \mathcal{A} is of infinite dimensional. Let $\rho : \mathcal{A} \longrightarrow \mathcal{B}(H)$ be an ample representation. According to Paschke [23], Higson [15] and Higson-Roe [16], the dual C^* -algebra of \mathcal{A} means the C^* -subalgebra $\mathfrak{D}_{\rho}(\mathcal{A})$ of $\mathcal{B}(H)$ defined by

$$\mathfrak{D}_{\rho}(\mathcal{A}) = \{ T \in \mathcal{B}(H) \mid [T, \rho(a)] \in \mathcal{K}(H) \text{ for all } a \in \mathcal{A} \}.$$
(2.2)

The isomorphism type of the C^* -algebra $\mathfrak{D}_{\rho}(\mathcal{A})$ does not depend on the choice of ample representations, so that it is denoted by $\mathfrak{D}(\mathcal{A})$ ([15], [16], [23]). Let us denote by $\widetilde{\mathcal{A}}$ the unitization of \mathcal{A} which contains \mathcal{A} as a maximal two-sided ideal with codimension one. In Higson [15] and Higson–Roe [16], the reduced K-homology groups $\widetilde{K}^p(\mathcal{A})$ and the unreduced K-homology groups $K^p(\mathcal{A})$ for p = 0, 1 are defined by

$$\widetilde{\operatorname{K}}^p(\mathcal{A}) := \operatorname{K}_{1-p}(\mathfrak{D}(\mathcal{A})) \quad \text{and} \quad \operatorname{K}^p(\mathcal{A}) := \operatorname{K}_{1-p}(\mathfrak{D}(\widetilde{\mathcal{A}})) \quad \text{for } p = 0, 1.$$

Lemma 2.1 ([16]). For a separable unital nuclear C^* -algebra \mathcal{A} , we have

$$\widetilde{\mathrm{K}}^{1}(\mathcal{A}) = \mathrm{Ext}_{s}(\mathcal{A}), \qquad \mathrm{K}^{1}(\mathcal{A}) = \mathrm{Ext}_{s}(\widetilde{\mathcal{A}}) = \mathrm{Ext}_{w}(\mathcal{A})$$

As in [16, Section 5], the group $K_*(\mathfrak{D}(\widetilde{\mathbb{C}}))$ is naturally identified with $K_*(\mathfrak{D}(\mathbb{C})/\mathcal{K}(H))$. Since $\mathfrak{D}(\mathbb{C})$ is $\mathcal{B}(H)$, $K^0(\mathbb{C})$ is identified with $K_1(\mathcal{Q}(H))$. An element of $K_1(\mathcal{Q}(H))$ is the class [U] of a unitary $U \in \mathcal{Q}(H)$ whose Fredholm index $\mathrm{Ind}(U)$ takes its value in \mathbb{Z} . This gives an isomorphism between $K^0(\mathbb{C})$ and \mathbb{Z} . We fix this isomorphism and identify the group $K^0(\mathbb{C})$ with \mathbb{Z} to write $K^0(\mathbb{C}) = \mathbb{Z}$. By Lemma 2.1, the sequence (2.1) is rephrased as

$$\mathrm{K}^{0}(\mathbb{C}) \xrightarrow{\iota_{\mathcal{A}}} \widetilde{\mathrm{K}}^{1}(\mathcal{A}) \xrightarrow{j_{\mathcal{A}}^{1}} \mathrm{K}^{1}(\mathcal{A})$$

that is exact at the middle, where the map $q_{\mathcal{A}} : \operatorname{Ext}_{s}(\mathcal{A}) \longrightarrow \operatorname{Ext}_{w}(\mathcal{A})$ in (2.1) is written as $j^{1}_{\mathcal{A}}$. As K^{0} is a contravariant functor, the natural inclusion $\mathbb{C} \hookrightarrow \mathcal{A}$ induces a homomorphism $\iota^{*}_{\mathbb{C}} : \mathrm{K}^{0}(\mathcal{A}) \longrightarrow \mathrm{K}^{0}(\mathbb{C})$. As in [16, (5.2.8)], the algebra $\mathfrak{D}(\widetilde{\mathcal{A}})$ is identified with $\begin{bmatrix} \mathfrak{D}(\mathcal{A}) & \mathcal{K}(H) \\ \mathcal{K}(H) & \mathcal{B}(H) \end{bmatrix}$. The inclusion of $\mathfrak{D}(\mathcal{A})$ into the upper left corner of $\mathfrak{D}(\widetilde{\mathcal{A}})$ induces a homomorphism from $\widetilde{\mathrm{K}}^{0}(\mathcal{A})$ to $\mathrm{K}^{0}(\mathcal{A})$ that we denote by $j^{0}_{\mathcal{A}}$. We then have the following K-homology long exact sequence.

Lemma 2.2 ([16]). For a separable unital C^* -algebra \mathcal{A} , the K-homology cyclic six term exact sequence

holds.

For the group $K_0(\mathcal{K}(H))$, the correspondence from the class [e] of a projection e of rank one in $\mathcal{K}(H)$ to the positive generator 1 of the integer group \mathbb{Z} induces an isomorphism between $K_0(\mathcal{K}(H))$ and \mathbb{Z} . We fix this correspondence and write $K_0(\mathcal{K}(H)) = \mathbb{Z}$.

3 Cuntz–Krieger algebras and K-homology groups

3.1 Cuntz–Krieger algebras and Toeplitz algebras

Let $A = [A(i, j)]_{i,j=1}^N$ be an irreducible non permutation matrix with entries in $\{0, 1\}$. The Cuntz-Krieger algebra \mathcal{O}_A for the matrix A is defined by the universal C^* -algebra generated by partial isometries S_1, \ldots, S_N subject to the operator relations:

$$\sum_{j=1}^{N} S_j S_j^* = 1, \qquad S_i^* S_i = \sum_{j=1}^{N} A(i,j) S_j S_j^*, \quad i = 1, \dots, N \quad ([8]). \tag{3.1}$$

It is a separable unital nuclear simple purely infinite C^* -algebra satisfying UCT. Let \mathbb{C}^N be the *N*-dimensional Hilbert space with orthonormal basis $\{\xi_1, \ldots, \xi_N\}$. Let H_0 be the one-dimensional Hilbert space with unit vector v_0 . The full Fock space F_N is defined by the Hilbert space $H_0 \oplus (\bigoplus_{n=1}^{\infty} H^{\otimes n})$, where $H^{\otimes n}$ is the *n*-fold tensor product $\mathbb{C}^N \otimes \cdots \otimes \mathbb{C}^N$. The sub Fock space H_A associated to A is a subspace of F_N defined by the closed linear span of vectors

$$\{v_0\} \cup \{\xi_{k_1} \otimes \cdots \otimes \xi_{k_n} \in H^{\otimes n} \mid A(k_j, k_{j+1}) = 1, \ j = 1, \dots, n-1, \ n = 1, 2, \dots \}.$$

Let us define the creation operators T_1, \ldots, T_N on H_A by

$$T_i v_0 = \xi_i, \qquad T_i(\xi_{k_1} \otimes \dots \otimes \xi_{k_n}) = \begin{cases} \xi_i \otimes \xi_{k_1} \otimes \dots \otimes \xi_{k_n} & \text{if } A(i, k_1) = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Let us denote by P_0 the projection of rank one on H_A onto H_0 . The creation operators T_1, \ldots, T_N are partial isometries on H_A satisfying the operator relations:

$$\sum_{j=1}^{N} T_j T_j^* + P_0 = 1, \qquad T_i^* T_i = \sum_{j=1}^{N} A(i,j) T_j T_j^* + P_0, \quad i = 1, \dots, N \quad ([13], [14]). \quad (3.2)$$

The C^* -algebra $C^*(T_1, \ldots, T_N, P_0)$ generated by T_1, \ldots, T_N, P_0 is called the Toeplitz algebra for A, that is denoted by \mathcal{T}_A . Let $S_i, i = 1, \ldots, N$ be the generating partial isometries of \mathcal{O}_A satisfying (3.1). The correspondence $T_i \longrightarrow S_i$ and $P_0 \longrightarrow 0$ yields a surjective *-homomorphism $\mathcal{T}_A \longrightarrow \mathcal{O}_A$ which is written π_A . The kernel of the map $\pi_A : \mathcal{T}_A \longrightarrow \mathcal{O}_A$ is generated by the operators of the form $T_{\mu_1} \cdots T_{\mu_m} P_0 T^*_{\nu_n} \cdots T^*_{\nu_1}, \mu_1, \ldots, \mu_m, \nu_1, \ldots, \nu_n \in \{1, \ldots, N\}$ that generates a C^* -algebra isomorphic to $\mathcal{K}(H_A)$. Hence we have an extension

$$\tau_A: 0 \longrightarrow \mathcal{K}(H_A) \longrightarrow \mathcal{T}_A \longrightarrow \mathcal{O}_A \longrightarrow 0$$
(3.3)

called the Toeplitz extension for \mathcal{O}_A . It defines an element $[\mathcal{T}_A]_s$ in $\operatorname{Ext}_s(\mathcal{O}_A)$ and $[\mathcal{T}_A]_w$ in $\operatorname{Ext}_w(\mathcal{O}_A)$, respectively.

3.2 Cyclic exact sequence of $Ext_*(\mathcal{O}_A)$ -groups

Let $\sigma : \mathcal{O}_A \longrightarrow \mathcal{Q}(H)$ be an extension. Put $e_i = \sigma(S_i S_i^*), i = 1, \ldots N$. Take a *monomorphism $\tau_0 : \mathcal{O}_A \longrightarrow \mathcal{B}(H)$ such that $\pi(\tau_0(S_i S_i^*)) = e_i$ and consider the extension $\tau = \pi \circ \tau_0 : \mathcal{O}_A \longrightarrow \mathcal{Q}(H)$. Since the partial isometry $\sigma(S_i)\tau(S_i^*)$ commutes with e_i , its restriction $e_i\sigma(S_i)\tau(S_i^*)e_i$ becomes a unitary in $e_i\mathcal{Q}(H)e_i$, so we may define its Fredholm index $\operatorname{ind}_{e_i}\sigma(S_i)\tau(S_i^*)$ denoted by $d_i(\sigma,\tau) \in \mathbb{Z}, i = 1, \ldots, N$. Cuntz–Krieger in [8] proved that the correspondence $[\sigma]_w \in \operatorname{Ext}_w(\mathcal{O}_A) \longrightarrow [d_i(\sigma,\tau)]_{i=1}^N \in \mathbb{Z}^N/(I-A)\mathbb{Z}^N$ is well-defined and gives rise to an isomorphism that is denoted by $d_{\underline{w}}$, so that $\operatorname{Ext}_w(\mathcal{O}_A) \cong \mathbb{Z}^N/(I-A)\mathbb{Z}^N$.

For the matrix A, we define an $N \times N$ matrix \hat{A} by $\hat{A} = A + R_1 - AR_1$, where R_1 is the $N \times N$ matrix whose first row is $[1, \ldots, 1]$, and other rows are zero vectors. In [22], it was shown that the above family $d_i(\sigma, \tau) \in \mathbb{Z}, i = 1, \ldots, N$ also yields a homomorphism $[\sigma]_s \in \operatorname{Ext}_s(\mathcal{O}_A) \longrightarrow [d_i(\sigma, \tau)]_{i=1}^N \in \mathbb{Z}^N/(I - \hat{A})\mathbb{Z}^N$ that is actually an isomorphism, denoted by d_s . Since $I - \hat{A} = (I - A)(I + R_1)$, there is a natural quotient map from $\mathbb{Z}^N/(I - \hat{A})\mathbb{Z}^N$ to $\mathbb{Z}^N/(I - A)\mathbb{Z}^N$ that is denoted by \hat{q}_A .

As in [22], a homomorphism $\hat{\iota}_A : \mathbb{Z} \longrightarrow \mathbb{Z}^N / (I - \widehat{A})\mathbb{Z}^N$ is defined by $\hat{\iota}_A(m) = [(I - A)[k_i]_{i=1}^N]$ for $m = \sum_{i=1}^N k_i$. Then the following two lemmas were shown in [22].

Lemma 3.1 ([22]). There exist isomorphisms

$$d_w : \operatorname{Ext}_w(\mathcal{O}_A) \longrightarrow \mathbb{Z}^N / (I - A) \mathbb{Z}^N, \qquad d_s : \operatorname{Ext}_s(\mathcal{O}_A) \longrightarrow \mathbb{Z}^N / (I - \widehat{A}) \mathbb{Z}^N$$

of groups such that the diagram

commutes, and

$$d_w([\mathcal{T}_A]_w) = -[1_N] \quad in \quad \mathbb{Z}^N/(I-A)\mathbb{Z}^N,$$
(3.5)

$$d_s([\mathcal{T}_A]_s) = -\hat{\iota}_A(1) - [1_N] \quad in \quad \mathbb{Z}^N / (I - \widehat{A})\mathbb{Z}^N, \tag{3.6}$$

where $[1_N] = [(1, \ldots, 1)]$ means the class of the vector $(1, \ldots, 1) \in \mathbb{Z}^N$.

Let us denote by $\operatorname{Ker}(I - A)$, $\operatorname{Ker}(I - \widehat{A})$ the subgroups of \mathbb{Z}^N defined by the kernels in \mathbb{Z}^N of the matrices I - A and of $I - \widehat{A}$, respectively. Let

$$i_1: \mathbb{Z} \longrightarrow \operatorname{Ker}(I - \widehat{A}), \quad j_A: \operatorname{Ker}(I - \widehat{A}) \longrightarrow \operatorname{Ker}(I - A), \quad s_A: \operatorname{Ker}(I - A) \longrightarrow \mathbb{Z}$$

be homomorphisms of abelian groups defined by

$$i_1(m) = (m, 0, \dots, 0), \quad j_A([l_i]_{i=1}^N) = (-\sum_{i=2}^N l_i, l_2, \dots, l_N), \quad s_A([l_i]_{i=1}^N) = \sum_{i=1}^N l_i.$$
 (3.7)

Since $j_A \circ i_1 = 0$, the map j_A induces a homomorphism $\operatorname{Ker}(I - \widehat{A})/i_1(\mathbb{Z}) \longrightarrow \operatorname{Ker}(I - A)$ that is still denoted by j_A .

Lemma 3.2 ([22]). We have the following cyclic six term exact sequence.

3.3 K-homology long exact sequence for \mathcal{O}_A

In this subsection, we will compute the K-homology cyclic six term exact sequence (2.3) for Cuntz-Krieger algebra \mathcal{O}_A . The following lemma is an easy consequence of basic properties of K-theory of Cuntz-Krieger algebras.

Lemma 3.3. There exists an isomorphism $\phi^0 : \mathrm{K}^0(\mathcal{O}_A) \longrightarrow \mathrm{Ker}(I-A)$ such that the diagram

commutes, where $\iota_{\mathbb{C}}^* : \mathrm{K}^0(\mathcal{O}_A) \longrightarrow \mathrm{K}^0(\mathbb{C})$ is the homomorphism induced by the inclusion $\iota_{\mathbb{C}} : \mathbb{C} \hookrightarrow \mathcal{O}_A$, and $s_A : \mathrm{Ker}(I - A) \longrightarrow \mathbb{Z}$ is the homomorphism defined in (3.7).

Let us denote by $j_A^0 : \widetilde{K}^0(\mathcal{O}_A) \longrightarrow K^0(\mathcal{O}_A)$ the injective homomorphism $j_{\mathcal{O}_A}^0 : \widetilde{K}^0(\mathcal{O}_A) \longrightarrow K^0(\mathcal{O}_A)$ in the upper horizontal arrow in (2.3) for $\mathcal{A} = \mathcal{O}_A$. As $j_A : \operatorname{Ker}(I - \widehat{A})/i_1(\mathbb{Z}) \longrightarrow \operatorname{Ker}(I - A)$ is injective, we have the following lemma.

Lemma 3.4. There exists a unique isomorphism $\widetilde{\phi}^0 : \widetilde{K}^0(\mathcal{O}_A) \longrightarrow \operatorname{Ker}(I - \widehat{A})/i_1(\mathbb{Z})$ such that the diagram

$$\widetilde{\mathrm{K}}^{0}(\mathcal{O}_{A}) \xrightarrow{\widetilde{\phi}^{0}} \operatorname{Ker}(I - \widehat{A})/i_{1}(\mathbb{Z})$$

$$\begin{array}{ccc}
 j_{A}^{0} \downarrow & & \downarrow j_{A} \\
 \mathrm{K}^{0}(\mathcal{O}_{A}) \xrightarrow{\phi^{0}} & \operatorname{Ker}(I - A) \\
\end{array}$$
(3.10)

commutes.

Let us denote by $\iota_A : \mathrm{K}^0(\mathbb{C})(=\mathbb{Z}) \longrightarrow \widetilde{\mathrm{K}}^1(\mathcal{O}_A)$ the homomorphism $\iota_{\mathcal{O}_A} : \mathrm{K}^0(\mathbb{C}) \longrightarrow \widetilde{\mathrm{K}}^1(\mathcal{O}_A)$ in the lower horizontal arrow in (2.3) for $\mathcal{A} = \mathcal{O}_A$. Since $\widetilde{\mathrm{K}}^1(\mathcal{O}_A) = \mathrm{Ext}_s(\mathcal{O}_A)$ and $\mathrm{K}^1(\mathcal{O}_A) = \mathrm{Ext}_w(\mathcal{O}_A)$, by putting

$$\tilde{\phi}^1 = d_s : \tilde{\mathrm{K}}^1(\mathcal{O}_A) \longrightarrow \mathbb{Z}^N / (I - \hat{A})\mathbb{Z}^N, \qquad \phi^1 = d_w : \mathrm{K}^1(\mathcal{O}_A) \longrightarrow \mathbb{Z}^N / (I - A)\mathbb{Z}^N,$$

we have the following proposition.

Proposition 3.5. The K-homology cyclic six term exact sequence (2.3) for \mathcal{O}_A

is computed to be the cyclic six term exact sequence (3.8), such that there is a commutative diagram

satisfying $\widetilde{\phi}^1([\mathcal{T}_A]_s) = -\hat{\iota}_A(1) - [1_N]$, where $[\mathcal{T}_A]_s \in \widetilde{\mathrm{K}}^1(\mathcal{O}_A)$ is the strong equivalence class of the Toeplitz extension and $[1_N] = [(1, \ldots, 1)]$ is the class of the vector $(1, \ldots, 1)$ in \mathbb{Z}^N .

Proof. By Lemma 3.1, Lemma 3.3 and Lemma 3.4, we have the commutative diagram (3.12). By (3.6), we know that $\tilde{\phi}^1([\mathcal{T}_A]_s) = -\hat{\iota}_A(1) - [1_N]$.

4 K-theory for Toeplitz algebra \mathcal{T}_{A^t}

In this section, we will compute the K-theory cyclic six term exact sequence for the Toeplitz extension (3.3) for the transposed matrix A^t of A. Let us consider the Toeplitz algebra \mathcal{T}_{A^t} for the transposed matrix A^t . Its generating partial isometries and projection are denoted by t_1, \ldots, t_N and p_0 that satisfy the operator relations

$$\sum_{j=1}^{N} t_j t_j^* + p_0 = 1, \qquad t_i^* t_i = \sum_{j=1}^{N} A^t(i,j) t_j t_j^* + p_0 \quad \text{for } i = 1, \dots, N \quad (\text{see } (3.2)).$$
(4.1)

Let $A_{\mathcal{T}}^t$ be the $(N+1) \times (N+1)$ matrix defined by $A_{\mathcal{T}}^t = \begin{bmatrix} 0 & \cdots & 0 \\ \vdots & A^t \end{bmatrix}$. The C^* -algebra \mathcal{T}_{A^t} is then regarded as the graph algebra $C^*(G_{A_{\mathcal{T}}^t})$ for the directed graph $G_{A_{\mathcal{T}}^t}$ associated to the matrix $A_{\mathcal{T}}^t$. The graph is obtained by adjoining a sink to the original graph G_{A^t} for the matrix A^t . Let A_1 be the $(N+1) \times (N+1)$ matrix defined by $A_1 = \begin{bmatrix} 1 & \cdots & 1 \\ 0 & A \end{bmatrix}$. The general K-theory formula due to Raeburn–Szymański [26] for graph C^* -algebras tells us the following lemma.

Lemma 4.1 (Raeburn–Szymański [26]). Keep the above notation.

- (i) $K_0(\mathcal{T}_{A^t}) = \mathbb{Z}^{N+1}/(I A_1)\mathbb{Z}^{N+1}$
- (ii) $\operatorname{K}_1(\mathcal{T}_{A^t}) = \operatorname{Ker}(s_A : \operatorname{Ker}(I A) \longrightarrow \mathbb{Z}).$

Let us define homomorphisms

$$\iota_{A_1} : \mathbb{Z} \longrightarrow \mathbb{Z}^{N+1}(I - A_1)\mathbb{Z}^{N+1}$$
 and $q_{A_1} : \mathbb{Z}^{N+1}/(I - A_1)\mathbb{Z}^{N+1} \longrightarrow \mathbb{Z}^N/(I - A)\mathbb{Z}^N$

by setting $\iota_{A_1}(m) = [0, \hat{\iota}_A(m)]$ for $m \in \mathbb{Z}$ and $q_{A_1}([(z_0, z_1, \dots, z_N)]) = [(z_1, \dots, z_N)]$ for $[(z_0, z_1, \dots, z_N)] \in \mathbb{Z}^{N+1}/(I - A_1)\mathbb{Z}^{N+1}$. Let

$$\iota_{s_A} : \operatorname{Ker}(s_A : \operatorname{Ker}(I - A) \longrightarrow \mathbb{Z}) \longrightarrow \operatorname{Ker}(I - A)$$

be the natural inclusion map. It is direct to see the following lemma.

Lemma 4.2. We have the following cyclic six term exact sequence.

Let $s_i, i = 1, ..., N$ be the generating partial isometries of the Cuntz-Krieger algebra \mathcal{O}_{A^t} for the transposed matrix A^t , that satisfy the operator relations

$$\sum_{j=1}^{N} s_j s_j^* = 1, \qquad s_i^* s_i = \sum_{j=1}^{N} A^t(i,j) s_j s_j^* \quad \text{for } i = 1, \dots, N \quad (\text{see } (3.1)).$$
(4.3)

As in (3.3), the correspondence $t_i \longrightarrow s_i$ and $p_0 \longrightarrow 0$ yields a surjective *-homomorphism $\mathcal{T}_{A^t} \longrightarrow \mathcal{O}_{A^t}$ which is written π_{A^t} , so that we have an extension

$$\tau_{A^t}: \quad 0 \longrightarrow \mathcal{K}(H) \longrightarrow \mathcal{T}_{A^t} \xrightarrow{\pi_{A^t}} \mathcal{O}_{A^t} \longrightarrow 0.$$
(4.4)

Lemma 4.3. Define homomorphisms

$$\eta_0^{\mathcal{T}} : \mathrm{K}_0(\mathcal{T}_{A^t}) \longrightarrow \mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1} \quad and \qquad \eta_0^{\mathcal{O}} : \mathrm{K}_0(\mathcal{O}_{A^t}) \longrightarrow \mathbb{Z}^N/(I-A)\mathbb{Z}^N$$

 $by \eta_0^{\mathcal{T}}([t_i t_i^*]) = [e_i], \ \eta_0^{\mathcal{T}}([p_0]) = [e_0] \ and \ \eta_0^{\mathcal{O}}([s_i s_i^*]) = [f_i], \ where \ e_i = (0, \dots, 0, \stackrel{i+1}{1}, 0, \dots, 0) \in \mathbb{Z}^{N+1}, i = 0, 1, \dots, N, \ and \ f_i = (0, \dots, 0, \stackrel{i}{1}, 0, \dots, 0) \in \mathbb{Z}^N, i = 1, \dots, N. \ Then \ both \ \eta_0^{\mathcal{T}} : K_0(\mathcal{T}_{A^t}) \longrightarrow \mathbb{Z}^{N+1}/(I - A_1)\mathbb{Z}^{N+1} \ and \ \eta_0^{\mathcal{O}} : K_0(\mathcal{O}_{A^t}) \longrightarrow \mathbb{Z}^N/(I - A)\mathbb{Z}^N \ are \ isomorphisms \ such \ that \ the \ diagram$

commutes, and $\eta_0^{\mathcal{T}}([1_{\mathcal{T}_{A^t}}]) = [1_{N+1}]$ holds, where $[1_{\mathcal{T}_{A^t}}]$ denotes the class in $\mathcal{K}_0(\mathcal{T}_{A^t})$ of the unit $1_{\mathcal{T}_{A^t}}$ of \mathcal{T}_{A^t} and $[1_{N+1}] = [(1, \ldots, 1)]$ denotes the class in $\mathbb{Z}^{N+1}/(I - A_1)\mathbb{Z}^{N+1}$ of the vector $(1, \ldots, 1) \in \mathbb{Z}^{N+1}$.

Proof. As $\eta_0^{\mathcal{T}} : \mathrm{K}_0(\mathcal{T}_{A^t}) \longrightarrow \mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1}$ and $\eta_0^{\mathcal{O}} : \mathrm{K}_0(\mathcal{O}_{A^t}) \longrightarrow \mathbb{Z}^N/(I-A)\mathbb{Z}^N$ are both isomorphisms, it suffices to show that the commutativity of the diagram. The commutativity $q_{A_1} \circ \eta_0^{\mathcal{T}} = \eta_0^{\mathcal{O}} \circ \pi_{A^{t_*}}$ is immediate by definition of π_{A^t} . In the short exact sequence (4.4), the class $[t_1p_0t_1^*]$ in $\mathrm{K}_0(\mathcal{K}(H))$ corresponds to the positive generator of \mathbb{Z} . As $[t_1p_0t_1^*] = [(t_1p_0)(t_1p_0)^*] = [(t_1p_0)^*(t_1p_0)] = [p_0]$ in $\mathrm{K}_0(\mathcal{T}_{A^t})$, we have

$$\eta_0^{\mathcal{T}}(\iota_*([t_1p_0t_1^*])) = \eta_0^{\mathcal{T}}([p_0]) = [e_0].$$

On the other hand, as

$$\hat{\iota}_A(1) = (I - A) \begin{bmatrix} 1\\0\\\vdots\\0 \end{bmatrix} = \begin{bmatrix} 1 - A(1,1)\\-A(2,1)\\\vdots\\-A(N,1) \end{bmatrix} \text{ and hence } \iota_{A_1}(1) = \begin{bmatrix} 0\\1 - A(1,1)\\-A(2,1)\\\vdots\\-A(N,1) \end{bmatrix}$$

so that

$$\iota_{A_1}(1) - [e_0] = \begin{bmatrix} -1\\ 1 - A(1, 1)\\ -A(2, 1)\\ \vdots\\ -A(N, 1) \end{bmatrix} = (I - A_1) \begin{bmatrix} 0\\ 1\\ 0\\ \vdots\\ 0 \end{bmatrix}.$$

This shows that $\iota_{A_1}(1) = [e_0]$ in $\mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1}$ and $\eta_0^{\mathcal{T}}(\iota_*([t_1p_0t_1^*])) = \iota_{A_1}(1)$, proving that the diagram $K_0(\mathcal{K}(H)) = \mathbb{Z}$

$$\begin{array}{ccc} \mathbb{X}_{0}(\mathcal{K}(H)) & = & \mathbb{Z} \\ & & & \downarrow^{\iota_{A_{1}}} \\ \mathbb{K}_{0}(\mathcal{T}_{A^{t}}) & \xrightarrow{\eta_{0}^{\mathcal{T}}} \mathbb{Z}^{N+1}/(I-A_{1})\mathbb{Z}^{N+1} \end{array}$$

commutes. The equality $\eta_0^{\mathcal{T}}([1_{\mathcal{T}_{A^t}}]) = [1_{N+1}]$ is obvious.

Let $\partial : K_1(\mathcal{O}_{A^t}) \longrightarrow K_0(\mathcal{K}(H))$ be the connecting map of the K-theory cyclic six term exact sequence for the extension (4.4).

Lemma 4.4. There are isomorphisms

$$\eta_1^{\mathcal{T}} : \mathrm{K}_1(\mathcal{T}_{A^t}) \longrightarrow \mathrm{Ker}(s_A : \mathrm{Ker}(I - A) \longrightarrow \mathbb{Z}) \quad and \quad \eta_1^{\mathcal{O}} : \mathrm{K}_1(\mathcal{O}_{A^t}) \longrightarrow \mathrm{Ker}(I - A)$$

such that the diagram

commutes.

Proof. Let $\zeta_1^{\mathcal{O}} : \mathrm{K}_1(\mathcal{O}_{A^t}) \ni [U(x)] \longrightarrow x = (x_i)_{i=1}^N \in \mathrm{Ker}(I-A)$ be the isomorphism defined by the inverse of the isomorphism from $\mathrm{Ker}(I-A)$ to $\mathrm{K}_1(\mathcal{O}_{A^t})$ explicitly defined in [27, P. 33]. As in Lemma 3.1, the class $[\mathcal{T}_{A^t}]_w$ in $\mathrm{Ext}_w(\mathcal{O}_{A^t})(=\mathrm{K}^1(\mathcal{O}_{A^t}))$ corresponds to the class $[(-1,\ldots,-1)]$ of $(-1,\ldots,-1)$ in $\mathbb{Z}^N/(I-A^t)\mathbb{Z}^N$. Then the connecting map $\partial : \mathrm{K}_1(\mathcal{O}_{A^t}) \longrightarrow \mathrm{K}_0(\mathcal{K}(H)) = \mathbb{Z}$ is given by the pairing

$$\partial([U]) = \langle [-1, \dots, -1] \mid \zeta_1^{\mathcal{O}}([U]) \rangle, \qquad [U] \in \mathcal{K}_1(\mathcal{O}_{A^t})$$

$$(4.6)$$

where the right hand side above is the inner product. For $[U(x)] \in K_1(\mathcal{O}_{A^t})$, we have

$$\partial([U(x)]) = < [-1, ..., -1] \mid x > = -\sum_{j=1}^{N} x_j \text{ for } x \in \operatorname{Ker}(I - A).$$

By setting $\eta_1^{\mathcal{O}}([U]) = \zeta_1^{\mathcal{O}}([U^*])$ for $[U] \in \mathcal{K}_1(\mathcal{O}_{A^t})$, we have

$$\partial([U(x)]) = < [1, ..., 1] \mid \eta_1^{\mathcal{O}}([U(x)]) > = \sum_{j=1}^N x_j = (s_A \circ \eta_1^{\mathcal{O}})([U(x)]),$$

proving $\partial = s_A \circ \eta_1^{\mathcal{O}}$.

Since both the homomorphisms $\pi_{A^{t_*}}$: $K_1(\mathcal{T}_{A^t}) \longrightarrow K_1(\mathcal{O}_{A^t})$ and ι_{s_A} : $\operatorname{Ker}(s_A : \operatorname{Ker}(I - A) \longrightarrow \mathbb{Z}) \longrightarrow \operatorname{Ker}(I - A)$ are injective, there exists a unique isomorphism $\eta_1^{\mathcal{T}}: K_1(\mathcal{T}_{A^t}) \longrightarrow \operatorname{Ker}(s_A : \operatorname{Ker}(I - A) \longrightarrow \mathbb{Z})$ satisfying $\iota_{s_A} \circ \eta_1^{\mathcal{T}} = \eta_1^{\mathcal{O}} \circ \pi_{A^{t_*}}$, showing the commutative diagram (4.5).

We thus have

Proposition 4.5. The K-theory cyclic six term exact sequence

for the Toeplitz extension (4.4) is computed to be the cyclic six term exact sequence (4.2), such that there is a commutative diagram

where the horizontal arrows $\eta_1^{\mathcal{T}}, \eta_1^{\mathcal{O}}, \eta_0^{\mathcal{T}}, \eta_0^{\mathcal{O}}$ are all isomorphisms satisfying $\eta_0^{\mathcal{T}}([1_{\mathcal{T}_{A^t}}]) = [1_{N+1}]$, and $[1_{\mathcal{T}_{A^t}}]$ denotes the class in $K_0(\mathcal{T}_{A^t})$ of the unit $1_{\mathcal{T}_{A^t}}$ of \mathcal{T}_{A^t} .

5 K-theoretic duality for Toeplitz extensions

We have four kinds of cyclic six term exact sequences (3.8), (3.11), (4.2) and (4.7). The first two are isomorphic by (3.12), and the second two are isomorphic by (4.8). In this section, we will show that (3.11) and (4.2) are isomorphic to prove that all of the four cyclic six term exact sequences are isomorphic.

Lemma 5.1. There exists an isomorphism $\xi^0: \mathbb{Z}^N/(I-\widehat{A})\mathbb{Z}^N \longrightarrow \mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1}$ satisfying $\xi^0(\hat{\iota}_A(1) + [1_N]) = [1_{N+1}]$ such that the diagram

commutes.

Proof. Let U, V be the $(N+1) \times (N+1)$ matrices defined by

$$U = \begin{bmatrix} 1 & 0 & \cdots & \cdots & 0 \\ 1 - A(1,1) & 1 & \ddots & & \vdots \\ -A(2,1) & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ -A(N,1) & 0 & \cdots & 0 & 1 \end{bmatrix}, \qquad V = \begin{bmatrix} 0 & 1 & 0 & \cdots & \cdots & 0 \\ -1 & 0 & -1 & \cdots & \cdots & -1 \\ 0 & 0 & 1 & 0 & \cdots & 0 \\ \vdots & & \ddots & \ddots & \ddots & \vdots \\ \vdots & & & \ddots & \ddots & \ddots & \vdots \\ \vdots & & & \ddots & \ddots & 0 \\ 0 & \cdots & \cdots & 0 & 1 \end{bmatrix}.$$

For $(x_1, ..., x_N) \in \mathbb{Z}^N$, define $\xi^0(x_1, ..., x_N) = (0, x_1, ..., x_N) \in \mathbb{Z}^{N+1}$. Note that $U^{-1}(0, x_1, ..., x_N) = (0, x_1, ..., x_N) \in \mathbb{Z}^{N+1}$ holds. As $U(I - A_1)V = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & I - \widehat{A} \end{bmatrix}$, we

have

$$(I - A_1)V[0 \oplus \mathbb{Z}^N] = U^{-1}[0 \oplus (I - \widehat{A})\mathbb{Z}^N] = \xi^0((I - \widehat{A})\mathbb{Z}^N)$$

so that $\xi^0((I-\widehat{A})\mathbb{Z}^N) \subset (I-A_1)\mathbb{Z}^{N+1}$. Hence $\xi^0: \mathbb{Z}^N \longrightarrow \mathbb{Z}^{N+1}$ induces a homomorphism $\mathbb{Z}^N/(I-\widehat{A})\mathbb{Z}^N \longrightarrow \mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1}$ that is still denoted by ξ^0 . It gives rise to an isomorphism because

$$\mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1} = U\mathbb{Z}^{N+1}/U(I-A_1)\mathbb{Z}^{N+1} = \mathbb{Z}^{N+1}/(\mathbb{Z}\oplus(I-\widehat{A})\mathbb{Z}^N) = \mathbb{Z}^N/(I-\widehat{A})\mathbb{Z}^N.$$

Since

$$\hat{\iota}_A(1) + [1_N] = (I - A) \begin{bmatrix} 1\\0\\\vdots\\0 \end{bmatrix} + \begin{bmatrix} 1\\1\\\vdots\\1 \end{bmatrix} = \begin{bmatrix} 1 - A(1,1) + 1\\-A(2,1) + 1\\\vdots\\-A(N,1) + 1 \end{bmatrix},$$

we have

$$\xi^{0}(\hat{\iota}_{A}(1) + [1_{N}]) - [1_{N+1}] = \begin{bmatrix} -1\\1 - A(1,1)\\-A(2,1)\\\vdots\\-A(N,1) \end{bmatrix} = (I - A_{1}) \begin{bmatrix} 0\\1\\0\\\vdots\\0 \end{bmatrix}$$

so that we have $\xi^0(\hat{\iota}_A(1) + [1_N]) = [1_{N+1}]$ in $\mathbb{Z}^{N+1}/(I - A_1)\mathbb{Z}^{N+1}$. The equalities $q_{\widehat{A}} = q_{A_1} \circ \xi^0$ and $\iota_{A_1} = \xi^0 \circ \iota_{\widehat{A}}$ are obvious.

Recall that a homomorphism $j_A : \operatorname{Ker}(I - \widehat{A})/\iota_1(\mathbb{Z}) \longrightarrow \operatorname{Ker}(I - A)$ is defined in (3.7), and $\iota_{s_A} : \operatorname{Ker}(s_A : \operatorname{Ker}(I - A) \longrightarrow \mathbb{Z}) \longrightarrow \operatorname{Ker}(I - A)$ stands for the inclusion map. Define $\tilde{\xi}^0 : \operatorname{Ker}(I - \widehat{A})/\iota_1(\mathbb{Z}) \longrightarrow \operatorname{Ker}(s_A : \operatorname{Ker}(I - A) \longrightarrow \mathbb{Z})$ by j_A . Since the diagram

$$\begin{array}{cccc} \operatorname{Ker}(I - \widehat{A})/\iota_{1}(\mathbb{Z}) & \xrightarrow{\widehat{\xi}^{0}} & \operatorname{Ker}(s_{A} : \operatorname{Ker}(I - A) \longrightarrow \mathbb{Z}) \\ & & & \downarrow^{\iota_{s_{A}}} & & \downarrow^{\iota_{s_{A}}} \\ & & & \operatorname{Ker}(I - A) & \underbrace{} & & \operatorname{Ker}(I - A) \end{array} \tag{5.1}$$

commutes, we have the following proposition.

Proposition 5.2. We have a commutative diagram

such that $\xi^0(\hat{\iota}_A(1) + [1_N]) = [1_{N+1}].$

Let us define the following four isomorphisms through (3.12), (4.8) and (5.2) by

$$\begin{split} \widetilde{\Phi}^{0}_{A} : \widetilde{\mathrm{K}}^{0}(\mathcal{O}_{A}) & \xrightarrow{\widetilde{\phi}^{0}} & \mathrm{Ker}(I - \widehat{A})/\iota_{1}(\mathbb{Z}) & \xrightarrow{\widetilde{\xi}^{0}} & \mathrm{Ker}(s_{A} : \mathrm{Ker}(I - A) \longrightarrow \mathbb{Z}) & \xrightarrow{\eta_{1}^{\tau-1}} & \mathrm{K}_{1}(\mathcal{T}_{A^{t}}), \\ \Phi^{0}_{A} : \mathrm{K}^{0}(\mathcal{O}_{A}) & \xrightarrow{\phi^{0}} & \mathrm{Ker}(I - A) & \underbrace{\qquad} & \mathrm{Ker}(I - A) & \xrightarrow{\eta_{1}^{\mathcal{O}-1}} & \mathrm{K}_{1}(\mathcal{O}_{A^{t}}), \\ \widetilde{\Phi}^{1}_{A} : \widetilde{\mathrm{K}}^{1}(\mathcal{O}_{A}) & \xrightarrow{\widetilde{\phi}^{1}} & \mathbb{Z}^{N}/(I - \widehat{A})\mathbb{Z}^{N} & \xrightarrow{\xi^{0}} & \mathbb{Z}^{N+1}/(I - A_{1})\mathbb{Z}^{N+1} & \xrightarrow{\eta_{0}^{\tau-1}} & \mathrm{K}_{0}(\mathcal{T}_{A^{t}}), \\ \Phi^{1}_{A} : \mathrm{K}^{1}(\mathcal{O}_{A}) & \xrightarrow{\phi^{1}} & \mathbb{Z}^{N}/(I - A)\mathbb{Z}^{N} & \underbrace{\qquad} & \mathbb{Z}^{N}/(I - A)\mathbb{Z}^{N} & \xrightarrow{\eta_{0}^{\mathcal{O}-1}} & \mathrm{K}_{0}(\mathcal{O}_{A^{t}}). \end{split}$$

Therefore we reach the following theorem.

Theorem 5.3. We have a commutative diagram

where two vertical sequences are exact and all of the horizontal arrows are isomorphisms such that

$$\widetilde{\Phi}^1_A([\mathcal{T}_A]_s) = -[1_{\mathcal{T}_A t}] \qquad in \quad \mathrm{K}_0(\mathcal{T}_{A^t}).$$

Proof. By combining the three commutative diagrams (3.12), (4.8) and (5.2), we get the disired commutative diagram from Proposition 3.5, Proposition 4.5 and Proposition 5.2. The equalities $\tilde{\Phi}^1_A([\mathcal{T}_A]_s) = (\eta_0^{\mathcal{T}-1} \circ \xi^0 \circ \tilde{\phi}^1)([\mathcal{T}_A]_s) = -[1_{\mathcal{T}_A t}]$ in $K_0(\mathcal{T}_A t)$ are obvious. \Box

6 K-theoretic duality for extensions

We now reach the following definition.

Definition 6.1. Let \mathcal{A} and \mathcal{A}' be separable unital nuclear C^* -algebras. Two extensions

$$\tau: 0 \longrightarrow \mathcal{K}(H) \xrightarrow{\iota} \mathcal{E} \xrightarrow{q} \mathcal{A} \longrightarrow 0 \quad \text{and} \quad \tau': 0 \longrightarrow \mathcal{K}(H) \xrightarrow{\iota'} \mathcal{E}' \xrightarrow{q'} \mathcal{A}' \longrightarrow 0$$

are said to be a K-theoretic duality pair if there are horizontal arrows of isomorphisms in the diagrams below making them commutative

If in particular $\Phi^0_{\mathbb{C}} = \Phi'^0_{\mathbb{C}} = \text{id}$ under the identification $\mathcal{K}^0(\mathbb{C}) = \mathcal{K}_0(\mathcal{K}(H)) = \mathbb{Z}$, and

$$\widetilde{\Phi}^{1}([\tau]_{s}) = -[1_{\mathcal{E}'}] \text{ in } \mathrm{K}_{0}(\mathcal{E}'), \qquad \widetilde{\Phi}'^{1}([\tau']_{s}) = -[1_{\mathcal{E}}] \text{ in } \mathrm{K}_{0}(\mathcal{E}), \tag{6.1}$$

then they are said to be strong K-theoretic duality pair, where $[1_{\mathcal{E}'}], [1_{\mathcal{E}}]$ denote the classes in $K_0(\mathcal{E}'), K_0(\mathcal{E})$ of the units of $\mathcal{E}', \mathcal{E}$, respectively.

For the strong K-theoretic duality pair, the above commutative diagrams say that the equalities (6.1) imply

$$\Phi^{1}([\tau]_{w}) = -[1_{\mathcal{A}'}] \text{ in } \mathrm{K}_{0}(\mathcal{A}'), \qquad \Phi'^{1}([\tau']_{w}) = -[1_{\mathcal{A}}] \text{ in } \mathrm{K}_{0}(\mathcal{A}).$$
(6.2)

By virtue of the introduction of the notion of K-theoretic duality for extensions of unital nuclear C^* -algebras, Theorem 5.3 implies Theorem 1.1. We note the following proposition.

Proposition 6.2. Let \mathcal{A} and \mathcal{A}' be separable unital nuclear C^* -algebras. Let

$$\tau_i: \ 0 \longrightarrow \mathcal{K}(H) \longrightarrow \mathcal{E}_i \longrightarrow \mathcal{A} \longrightarrow 0 \quad and \quad \tau'_i: \ 0 \longrightarrow \mathcal{K}(H) \longrightarrow \mathcal{E}'_i \longrightarrow \mathcal{A}' \longrightarrow 0$$

be extensions for i = 1, 2. Suppose that $[\tau_1]_s = [\tau_2]_s$ in $\operatorname{Ext}_s(\mathcal{A})$ and $[\tau'_1]_s = [\tau'_2]_s$ in $\operatorname{Ext}_s(\mathcal{A}')$. If τ_1 and τ'_1 are K-theoretic duality pair (resp. strong K-theoretic duality pair),

then τ_2 and τ'_2 are K-theoretic duality pair (resp. strong K-theoretic duality pair). Hence the (strong) K-theoretic duality pair depends only on the strong equivalence classes of extensions.

Proof. Assume that $[\tau_1]_s = [\tau_2]_s$ in $\operatorname{Ext}_s(\mathcal{A})$ and $[\tau'_1]_s = [\tau'_2]_s$ in $\operatorname{Ext}_s(\mathcal{A})$. We may find unitaries $U, U' \in \mathcal{B}(H)$ and isomorphisms $\gamma : \mathcal{E}_1 \longrightarrow \mathcal{E}_2, \ \gamma' : \mathcal{E}'_1 \longrightarrow \mathcal{E}'_2$ such that the diagrams

and

commute. Hence it is a direct consequence that τ_1 and τ'_1 are K-theoretic duality pair implies that τ_2 and τ'_2 are K-theoretic duality pair.

Suppose further that τ_1 and τ'_1 are strong K-theoretic pair. Let $\widetilde{\Phi}_1^1 : \widetilde{K}^1(\mathcal{A}_1) \longrightarrow K_0(\mathcal{E}_1')$ and $\widetilde{\Phi}_1'^1 : \widetilde{K}^1(\mathcal{A}_1') \longrightarrow K_0(\mathcal{E}_1)$ be isomorphisms giving rise to the strong K-theoretic duality pair for τ_1 and τ'_1 such that $\widetilde{\Phi}_1^1([\tau_1]_s) = -[1_{\mathcal{E}_1'}]$ in $K_0(\mathcal{E}_1')$ and $\widetilde{\Phi}_1'^1([\tau'_1]_s) = -[1_{\mathcal{E}_1}]$ in $K_0(\mathcal{E}_1)$. As $[\tau_1]_s = [\tau_2]_s$ and $[\tau'_1]_s = [\tau'_2]_s$, we have $(\gamma'_* \circ \widetilde{\Phi}_1^1)([\tau_2]_s) = -[\gamma'_*(1_{\mathcal{E}_1'})] = -[1_{\mathcal{E}_2'}]$, and symmetrically $(\gamma_* \circ \widetilde{\Phi}_1'^1)([\tau'_2]_s) = -[1_{\mathcal{E}_2}]$. Since $\operatorname{Ad}(U)_* = \operatorname{Ad}(U')_* = \operatorname{id}$ on $K_0(\mathcal{K}(H))$, the isomorphisms $\widetilde{\Phi}_2^1 := \gamma'_* \circ \widetilde{\Phi}_1^1 : \widetilde{K}^1(\mathcal{A}_2) \longrightarrow K_0(\mathcal{E}_2')$ and $\widetilde{\Phi}_2'^1 := \gamma_* \circ \widetilde{\Phi}_1'^1 : \widetilde{K}^1(\mathcal{A}_2') \longrightarrow K_0(\mathcal{E}_2)$ yield that τ_2 and τ'_2 are strong K-theoretic pair.

7 Duality for Toeplitz extensions

We will first show the following proposition, that is a sort of duality of the matrix A and its transpose A^t from the viewpoint of Cuntz–Krieger algebras and extension groups.

Proposition 7.1. Let A, B be irreducible non permutation matrices with entries in $\{0, 1\}$. The following are equivalent.

- (i) The Cuntz-Krieger algebras \mathcal{O}_A and \mathcal{O}_B are isomorphic, and there exists an isomorphism $\varphi : \operatorname{Ext}_w(\mathcal{O}_A) \longrightarrow \operatorname{Ext}_w(\mathcal{O}_B)$ such that $\varphi([\mathcal{T}_A]_w) = [\mathcal{T}_B]_w$.
- (ii) The Cuntz-Krieger algebras \mathcal{O}_{A^t} and \mathcal{O}_{B^t} are isomorphic, and there exists an isomorphism $\phi : \operatorname{Ext}_w(\mathcal{O}_{A^t}) \longrightarrow \operatorname{Ext}_w(\mathcal{O}_{B^t})$ such that $\phi([\mathcal{T}_{A^t}]_w) = [\mathcal{T}_{B^t}]_w$.

Proof. Under the identification between $K_0(\mathcal{O}_A)$ and $Ext_w(\mathcal{O}_{A^t})$, the class $[1_{\mathcal{O}_A}]$ of the unit $1_{\mathcal{O}_A}$ of \mathcal{O}_A correspondes to the class $-[\mathcal{T}_{A^t}]_w$ of the minus of the Toeplitz extensioon of \mathcal{O}_{A^t} . Hence the equivalence between (i) and (ii) is immediate from Rørdam's classification result of the Cuntz-Krieger algebras ([27]).

The following lemma is elementary, so we omit its proof.

Lemma 7.2. Let $\tau_i : \mathcal{A}_i \longrightarrow \mathcal{Q}(H)$ be the Busby invariants for extensions

$$0 \longrightarrow \mathcal{K}(H) \longrightarrow \mathcal{E}_i \longrightarrow \mathcal{A}_i \longrightarrow 0, \qquad i = 1, 2.$$

Suppose that there exist isomorphisms $\alpha : \mathcal{K}(H) \longrightarrow \mathcal{K}(H), \ \beta : \mathcal{E}_1 \longrightarrow \mathcal{E}_2, \ \gamma : \mathcal{A}_1 \longrightarrow \mathcal{A}_2$ of C^* -algebras such that the diagarm

commutes. Then $\gamma : \mathcal{A}_1 \longrightarrow \mathcal{A}_2$ induces isomorphisms $\gamma_s^* : \operatorname{Ext}_s(\mathcal{A}_2) \longrightarrow \operatorname{Ext}_s(\mathcal{A}_1)$ and $\gamma_w^* : \operatorname{Ext}_w(\mathcal{A}_2) \longrightarrow \operatorname{Ext}_w(\mathcal{A}_1)$ such that $\gamma_s^*([\tau_2]_s) = [\tau_1]_s, \ \gamma_w^*([\tau_2]_w) = [\tau_1]_w$ and the diagram

$$\mathbb{Z} \xrightarrow{\iota_{\mathcal{A}_{2}}} \operatorname{Ext}_{s}(\mathcal{A}_{2}) \xrightarrow{q_{2}} \operatorname{Ext}_{w}(\mathcal{A}_{2})$$

$$\| \qquad \gamma_{s}^{*} \downarrow \qquad \downarrow \gamma_{w}^{*}$$

$$\mathbb{Z} \xrightarrow{\iota_{\mathcal{A}_{1}}} \operatorname{Ext}_{s}(\mathcal{A}_{1}) \xrightarrow{q_{1}} \operatorname{Ext}_{w}(\mathcal{A}_{1})$$

commutes.

We thus have the following proposition.

Proposition 7.3. The following are equivalent.

- (i) There exists an isomorphism $\Phi : \mathcal{O}_A \longrightarrow \mathcal{O}_B$ of C^* -algebras such that the induced isomorphism $\Phi_s^* : \operatorname{Ext}_{\mathrm{s}}(\mathcal{O}_B) \longrightarrow \operatorname{Ext}_{\mathrm{s}}(\mathcal{O}_A)$ satisfies $\Phi_s^*([\mathcal{T}_A]_s) = [\mathcal{T}_B]_s$.
- (ii) There exist isomorphisms $\Phi : \mathcal{O}_A \longrightarrow \mathcal{O}_B$ and $\Psi : \mathcal{T}_A \longrightarrow \mathcal{T}_B$ of C^* -algebras such that the diagram

$$\begin{array}{cccc} \mathcal{T}_{A} & \xrightarrow{\pi_{A}} & \mathcal{O}_{A} \\ \Psi & & & \downarrow \Phi \\ \mathcal{T}_{B} & \xrightarrow{\pi_{B}} & \mathcal{O}_{B} \end{array} \tag{7.1}$$

commutes.

Proof. Let us denote by $\tau_A : \mathcal{O}_A \longrightarrow \mathcal{Q}(H)$ and $\tau_B : \mathcal{O}_B \longrightarrow \mathcal{Q}(H)$ the Busby invariants for the Toeplitz extensions \mathcal{T}_A and \mathcal{T}_B , respectively.

(i) \implies (ii): Since $\Phi_s^*([\tau_B]_s) = [\tau_A]_s$, there exists a unitary $U \in \mathcal{B}(H)$ such that $\tau_B(\Phi(X)) = \pi(U)\tau_A(X)\pi(U)^*, X \in \mathcal{O}_A$. As

$$\mathcal{T}_A = \{ (T, X) \in \mathcal{B}(H) \oplus \mathcal{O}_A \mid \pi(T) = \tau_A(X) \}, \mathcal{T}_B = \{ (T', X') \in \mathcal{B}(H) \oplus \mathcal{O}_B \mid \pi(T') = \tau_B(X') \},$$

we may define the map $\Psi : \mathcal{T}_A \longrightarrow \mathcal{T}_B$ by setting $\Psi(T, X) = (UTU^*, \Phi(X))$. It is easy to see that $\Psi(T, X) \in \mathcal{T}_B$ for $(T, X) \in \mathcal{T}_A$ and $\Psi : \mathcal{T}_A \longrightarrow \mathcal{T}_B$ gives rise to an isomorphism such that the diagram (7.1) commutes.

(ii) \implies (i): This implication follows from Lemma 7.2.

Now we give a proof of Theorem 1.2.

Proof of Theorem 1.2. (i) \Longrightarrow (ii): Assume that there exist isomorphisms $\Phi : \mathcal{O}_A \longrightarrow \mathcal{O}_B$ and $\Psi : \mathcal{T}_A \longrightarrow \mathcal{T}_B$ of C^* -algebras such that the diagram (1.4) commutes. Since \mathcal{T}_A has exactly one non-trivial ideal, which is the kernel of the surjection $\pi_A : \mathcal{T}_A \longrightarrow \mathcal{O}_A$. It is isomorphic to the C^* -algebra $\mathcal{K}(H_A)$ of compact operators on the sub Fock space H_A . Hence the isomorphism $\Psi : \mathcal{T}_A \longrightarrow \mathcal{T}_B$ satisfies $\Psi(\mathcal{K}(H_A)) = \mathcal{K}(H_B)$. Put $\alpha =$ $\Psi|_{\mathcal{K}(H_A)} : \mathcal{K}(H_A) \longrightarrow \mathcal{K}(H_B)$ the restriction of Ψ to $\mathcal{K}(H_A)$. It induces an isomorphism from $K_0(\mathcal{K}(H_A))$ to $K_0(\mathcal{K}(H_B))$, that maps \mathbb{Z} to \mathbb{Z} as the identity. Therefore we have a commutative diagram:

such that $\alpha_* = \operatorname{id} : \operatorname{K}_0(\mathcal{K}(H_A)) = \mathbb{Z} \longrightarrow \operatorname{K}_0(\mathcal{K}(H_B)) = \mathbb{Z}$. In the general exact sequence (2.1), the homomorphism $\iota_{\mathcal{A}} : \mathbb{Z} \longrightarrow \operatorname{Ext}_{\mathrm{s}}(\mathcal{A})$ is given by $\iota_{\mathcal{A}} : m \in \mathbb{Z} \longrightarrow [\sigma_m] \in \operatorname{Ext}_{\mathrm{s}}(\mathcal{A})$ where $\sigma_m = \operatorname{Ad}(u_m) \circ \tau : \mathcal{A} \longrightarrow \mathcal{Q}(H)$ for a trivial extension $\tau : \mathcal{A} \longrightarrow \mathcal{Q}(H)$ and a unitary $u_m \in \mathcal{Q}(H)$ of Fredholm index m. We denote by ι_A, ι_B the homomorphisms $\iota_{\mathcal{O}_A}, \iota_{\mathcal{O}_B}$ in (2.1) for $\mathcal{A} = \mathcal{O}_A$, respectively. For the isomorphism $\Phi : \mathcal{O}_A \longrightarrow \mathcal{O}_B$, define isomorphisms $\Phi_s^* : \operatorname{Ext}_{\mathrm{s}}((\mathcal{O}_B) \longrightarrow \operatorname{Ext}_{\mathrm{s}}(\mathcal{O}_A)$ and $\Phi_w^* : \operatorname{Ext}_{\mathrm{w}}(\mathcal{O}_B) \longrightarrow \operatorname{Ext}_{\mathrm{w}}(\mathcal{O}_A)$ by setting $\Phi_s^*([\sigma]_s) = [\sigma \circ \Phi]_s$ and $\Phi_w^*([\sigma]_w) = [\sigma \circ \Phi]_w$, respectively, for a unital *-monomorphism $\sigma : \mathcal{O}_B \longrightarrow \mathcal{Q}(H)$. As the equality $\iota_A(m) = \Phi_s^* \circ \iota_B(m), m \in \mathbb{Z}$ holds, we have the commutative diagram:

By virtue of Proposition 7.3, the commutative diagram (7.2) tells us that $\Phi_s^*([\mathcal{T}_B]_s) = [\mathcal{T}_A]_s$ in $\operatorname{Ext}_s(\mathcal{O}_A)$ and $\Phi_w^*([\mathcal{T}_B]_w) = [\mathcal{T}_A]_w$ in $\operatorname{Ext}_w(\mathcal{O}_A)$ so that $\Phi_s^*(\iota_B(1) + [\mathcal{T}_B]_s) = \iota_A(1) + [\mathcal{T}_A]_s$. Under the identifications $\widetilde{K}^1(\cdot) = \operatorname{Ext}_s(\cdot)$ and $K^1(\cdot) = \operatorname{Ext}_w(\cdot)$, the fact that the Toeplitz extensions are strong K-theoretic duality pair gives the following commutative diagram

By putting

$$\Phi_0^{\mathcal{T}} := \widetilde{\Phi}_A^1 \circ \Phi_s^* \circ \widetilde{\Phi}_B^{1-1} : \mathrm{K}_0(\mathcal{T}_{B^t}) \longrightarrow \mathrm{K}_0(\mathcal{T}_{A^t}),$$

$$\Phi_0^{\mathcal{O}} := \Phi_A^1 \circ \Phi_w^* \circ \Phi_B^{1-1} : \mathrm{K}_0(\mathcal{O}_{B^t}) \longrightarrow \mathrm{K}_0(\mathcal{O}_{A^t}),$$

we have a commutative diagram

such that $\Phi_0^{\mathcal{T}}([1_{\mathcal{T}_{B^t}}]) = [1_{\mathcal{T}_{A^t}}]$ and $\Phi_0^{\mathcal{O}}([1_{\mathcal{O}_{B^t}}]) = [1_{\mathcal{O}_{A^t}}]$. The the groups $K_0(\mathcal{T}_{A^t})$ and $K_0(\mathcal{T}_{B^t})$ consist of the equivalence classes of projections in \mathcal{T}_{A^t} and \mathcal{T}_{B^t} , respectively, Thanks to the classification result [10, Theorem 5.3] (cf. [11]) proved by Eilers–Restroff–Ruiz, we may find an isomorphism $\varphi : \mathcal{T}_{B^t} \longrightarrow \mathcal{T}_{A^t}$ such that $\varphi_* = \Phi_0^{\mathcal{T}} : K_0(\mathcal{T}_{B^t}) \longrightarrow K_0(\mathcal{T}_{A^t})$. Since the isomorphism $\varphi : \mathcal{T}_{B^t} \longrightarrow \mathcal{T}_{A^t}$ preserves their ideals of compact operators, we see that $\varphi(\mathcal{K}(H_{B^t})) = \mathcal{K}(H_{A^t})$, so that the isomorphism $\Psi^t := \varphi^{-1} : \mathcal{T}_{A^t} \longrightarrow \mathcal{T}_{B^t}$ induces an isomorphism $\Phi^t : \mathcal{O}_{A^t} \longrightarrow \mathcal{O}_{B^t}$ such that the diagram (7.5) commutes.

We finally present several equivalence conditions under which two Toeplitz algebras are isomorphic.

Proposition 7.4. Let $A = [A(i,j)]_{i,j=1}^N$, $B = [B(i,j)]_{i,j=1}^M$ be irreducible non permutation matrices with entries in $\{0,1\}$. Then the following four assertions are isomorphic.

- (i) \mathcal{T}_A is isomorphic to \mathcal{T}_B .
- (ii) There exist isomorphisms $\Phi : \mathcal{O}_A \longrightarrow \mathcal{O}_B$ and $\Psi : \mathcal{T}_A \longrightarrow \mathcal{T}_B$ of C^* -algebras such that the diagram

commutes.

(iii) There exists an isomorphism $\varphi : \mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1} \longrightarrow \mathbb{Z}^{M+1}/(I-B_1)\mathbb{Z}^{M+1}$ of groups such that $\varphi([(1,\ldots,1)]) = [(1,\ldots,1)]$ and $\varphi([(1,0,\ldots,0)]) = [(1,0,\ldots,0)]$.

(iv) There exists an isomorphism $\Phi : \mathcal{O}_A \longrightarrow \mathcal{O}_B$ of C^* -algebras such that $\Phi_s^*([\mathcal{T}_B]_s) = [\mathcal{T}_A]_s$ in $\operatorname{Ext}_s(\mathcal{O}_A)$

where A_1 is the $(N+1) \times (N+1)$ matrix and B_1 is the $(M+1) \times (M+1)$ matrix defined by $A_1 = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ 0 & & \\$

Proof. (i) \iff (ii) is clear.

(ii) \iff (iv) is seen in Proposition 7.3.

(ii) \implies (iii): Assume the condition (ii), so that we have a commutative diagram (7.2). Put

$$\varphi := \Psi_* : \mathrm{K}_0(\mathcal{T}_A) = \mathbb{Z}^{N+1} / (I - A_1) \mathbb{Z}^{N+1} \longrightarrow \mathrm{K}_0(\mathcal{O}_B) = \mathbb{Z}^{M+1} / (I - B_1) \mathbb{Z}^{M+1}.$$

Since the position $[1_{\mathcal{T}_A}]$ of the unit of \mathcal{T}_A in $K_0(\mathcal{T}_A)$ is the class $[(1,\ldots,1)]$ of the vector $(1,\ldots,1)$ in the group $\mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1}$, we have $\varphi([(1,\ldots,1)]) = [(1,\ldots,1)]$. Since $\Psi(\mathcal{T}_A) = \mathcal{T}_B$ and the class of a minimal projection is the class $[(1,0,\ldots,0)]$ of the vector $(1,0,\ldots,0)$ in $K_0(\mathcal{T}_A) = \mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1}$, and similarly in $K_0(\mathcal{T}_B) = \mathbb{Z}^{M+1}/(I-B_1)\mathbb{Z}^{M+1}$, we have $\varphi([(1,0,\ldots,0)]) = [(1,0,\ldots,0)]$.

(iii) \Longrightarrow (i): Assume (iii). Since the class $[(1,0,\ldots,0)]$ of the vector $(1,0,\ldots,0)$ is $[e_0]$ in the notation of Lemma 4.3, the condition $\varphi([(1,0,\ldots,0)]) = [(1,0,\ldots,0)]$ implies that $\varphi : Z^{N+1}/(I-A_1)\mathbb{Z}^{N+1} \longrightarrow \mathbb{Z}^{M+1}/(I-B_1)\mathbb{Z}^{M+1}$ induces an isomorphism $\varphi_0 : Z^N/(I-A)\mathbb{Z}^N \longrightarrow \mathbb{Z}^M/(I-B)\mathbb{Z}^M$ which sends $[(1,\ldots,1)]$ in $Z^N/(I-A)\mathbb{Z}^N$ to $[(1,\ldots,1)]$ in $Z^M/(I-B)\mathbb{Z}^M$. Hence there exists an isomorphism $\Phi : \mathcal{O}_{A^t} \longrightarrow \mathcal{O}_{B^t}$ such that $\Phi_* = \varphi_0 : K_0(\mathcal{O}_{A^t}) \longrightarrow K_0(\mathcal{O}_{B^t})$ by [27]. It also induces an isomorphism $\Phi_* : K_1(\mathcal{O}_{A^t}) = \operatorname{Ker}(I-A) \longrightarrow K_1(\mathcal{O}_{B^t}) = \operatorname{Ker}(I-B)$ written φ_1 . Let $\zeta_1^{\mathcal{O}} : K_0(\mathcal{O}_{A^t}) \longrightarrow \operatorname{Ker}(I-A)$ be the isomorphism defined in the proof of Lemma 4.4. Let $\partial_{A^t} : K_1(\mathcal{O}_{A^t}) \longrightarrow K_0(\mathcal{K}(H))$ denote the connectiong map ∂ in the diagram (4.5). For $[U] \in K_1(\mathcal{O}_{B^t})$, by (4.6) we have

$$\partial_{A^{t}}(\Phi_{*}([U])) = \langle [-1, \dots, -1] \mid \zeta_{1}^{\mathcal{O}}([\Phi(U)]) \rangle = \langle [-1, \dots, -1] \mid \Phi_{*}(\zeta_{1}^{\mathcal{O}}([U]) \rangle \\ = \langle \varphi([-1, \dots, -1]) \mid \zeta_{1}^{\mathcal{O}}([U]) \rangle = \langle [-1, \dots, -1]) \mid \zeta_{1}^{\mathcal{O}}([U]) \rangle = \partial_{B^{t}}([U])$$

because of the condition $\varphi([-1,\ldots,-1]) = [-1,\ldots,-1]$. The map $\iota_{A_1} : \mathbb{Z} \longrightarrow Z^{N+1}/(I - A_1)\mathbb{Z}^{N+1}$ is given by by $m \in \mathbb{Z} \longrightarrow m[e_0] \in Z^{N+1}/(I - A_1)\mathbb{Z}^{N+1}$. Hence we have a commutative diagram

We thus have a commutative diagram

where the horizontal arrows in (7.6) are all isomorphisms. The the groups $K_0(\mathcal{T}_{A^t})$ and $K_0(\mathcal{T}_{B^t})$ consist of the equivalence classes of projections in \mathcal{T}_{A^t} and \mathcal{T}_{B^t} , respectively, Since $\varphi([1, \ldots, 1]) = [1, \ldots, 1]$ in $K_0(\mathcal{T}_{B^t})$ and $\varphi_0([1, \ldots, 1]) = [1, \ldots, 1]$ in $K_0(\mathcal{O}_{B^t})$, by the classification result [10, Theorem 5.3] (cf. [11]) proved by Eilers–Restroff–Ruiz, we may find an isomorphism $\varphi : \mathcal{T}_{B^t} \longrightarrow \mathcal{T}_{A^t}$, because of Proposition 4.5. Therefore we conclude that \mathcal{T}_A is isomorphic to \mathcal{T}_B by Corollary 1.3.

8 Examples

We present examples of the triplet $(\mathbb{Z}^{N+1}/(I-A_1)\mathbb{Z}^{N+1}, [(1,0,\ldots,0)], [(1,\ldots,1)])$ that shows $(K_0(\mathcal{T}_{A^t}), \iota_*([1]), [1_{\mathcal{T}_{A^t}}])$ for several matrices, computed without difficulty by hand. **1.**

•
$$B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$
, $(\mathbb{Z}^3/(I - B_1)\mathbb{Z}^3, [(1,0,0)], [(1,1,1)]) \cong (\mathbb{Z}, 1, 0)$.
• $B_- = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix}$, $(\mathbb{Z}^5/(I - B_{-1})\mathbb{Z}^5, [(1,0,0,0,0)], [(1,1,1,1,1)]) \cong (\mathbb{Z}, 1, -1)$.
• $F = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$, $(\mathbb{Z}^3/(I - F_1)\mathbb{Z}^3, [(1,0,0)], [(1,1,1)]) \cong (\mathbb{Z}, 1, -2)$.
• $C = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}$, $(\mathbb{Z}^4/(I - C_1)\mathbb{Z}^4, [(1,0,0,0)], [(1,1,1,1)]) \cong (\mathbb{Z}, 1, -1)$.

•
$$D = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$
, $(\mathbb{Z}^4/(I - D_1)\mathbb{Z}^4, [(1, 0, 0, 0)], [(1, 1, 1, 1)]) \cong (\mathbb{Z}, 1, 0).$

Hence we have $\mathcal{T}_B \cong \mathcal{T}_{D^t}$ and $\mathcal{T}_{B_-} \cong \mathcal{T}_{C^t}$, whereas \mathcal{T}_F is not isomorphic to any of $\mathcal{T}_B, \mathcal{T}_{B_-}, \mathcal{T}_{C^t}, \mathcal{T}_{D^t}$. Since $\mathrm{K}_0(\mathcal{O}_{A^t}) \cong \mathrm{K}_0(\mathcal{T}_{A^t})/\mathbb{Z}_{\ell_*}([1])$, we have $\mathcal{O}_B \cong \mathcal{O}_{B_-} \cong \mathcal{O}_F \cong \mathcal{O}_{C^t} \cong \mathcal{O}_{D^t}$, that are all isomorphic to \mathcal{O}_2 .

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \qquad B = A^t = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix},$$
$$(\mathbb{Z}^4/(I - A_1)\mathbb{Z}^4, [(1, 0, 0, 0)], [(1, 1, 1, 1)]) \cong (\mathbb{Z}, 2, -2),$$

$$(\mathbb{Z}^4/(I-B_1)\mathbb{Z}^4, [(1,0,0,0)], [(1,1,1,1)]) \cong (\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}, 1 \oplus 0, (-1) \oplus 1).$$

Hence we have \mathcal{T}_A is not isomorphic to \mathcal{T}_B , and \mathcal{O}_A is not isomorphic to \mathcal{O}_B . We indded see that \mathcal{O}_A is isomorphic to \mathcal{O}_3 , and \mathcal{O}_B is isomorphic to $\mathcal{O}_3 \otimes M_2(\mathbb{C})$ (cf. [13], [24]).

Acknowledgment: This work was supported by JSPS KAKENHI Grant Number 19K03537.

References

2.

- W. B. ARVESON, Notes on extensions of C^{*}-algebras, Duke Math. J. 44(1977), pp. 329–335.
- [2] B. E. BLACKADAR, K-theory for operator algebras, MSRI Publications 5, Springer-Verlag, Berlin, Heidelberg and New York, 1986.
- [3] L. G. BROWN, R. G. DOUGLAS AND P. A. FILLMORE, Extensions of C^{*}-algebras and K-homology, Ann. Math. 105(1977), pp. 265–324.
- [4] R. C. BUSBY, Double centralizers and extensions of C*-algebras, Trans. Amer. Math. Soc. 132(1968), 79–99.
- [5] M. D. CHOI AND E. G. EFFROS, The completely positive lifting problem for C^{*}algebras, Ann. Math. 104(1976), 585–609.
- [6] J. CUNTZ, A class of C*-algebras and topological Markov chains II: reducible chains and the Ext-functor for C*-algebras, Invent. Math. 63(1980), pp. 25–40.
- [7] J. CUNTZ, K-theory for certain simple C*-algebras, Ann. Math. 117(1981), pp. 181– 197.
- [8] J. CUNTZ AND W. KRIEGER, A class of C^{*}-algebras and topological Markov chains, Invent. Math. 56(1980), pp. 251–268.
- R. G. DOUGLAS, C^{*}-algebra extensions and K-homology, Princeton University Press, Princeton, New Jersey, 1980.

- [10] S. EILERS, G. RESTROFF AND E. RUIZ, Strong classification of extensions of classifiable C^{*}-algebras, Bull. Korean Math. Soc. 59(2022), pp. 567–608.
- [11] S. EILERS, G. RESTROFF, E. RUIZ AND A. P. W. SØRENSEN, The complete classification of unital graph C*-algebras: Geometric and strong, Duke Math. J. 170(2021), pp. 2421–2517.
- [12] M. ENOMOTO, M. FUJII AND Y. WATATANI, Tensor algebras on the sub Fock space associated with \mathcal{O}_A , Math. Japon. 24(1979), pp. 463–468.
- [13] M. ENOMOTO, M. FUJII AND Y. WATATANI, K₀-groups and classifications of Cuntz-Krieger algebras, Math. Japon. 26(1981), pp. 443–460.
- [14] D. EVANS, Gauge actions on \mathcal{O}_A , J. Operator theory 7(1982), pp. 79–100.
- [15] N. HIGSON, C^{*}-algebra extension theory and duality, Jour. Funct. Anal. 129(1995), pp. 349–363.
- [16] N. HIGSON AND J. ROE, Analytic K-homology, Oxford Mathematical Monographs, Oxford Science Publications, Oxford University Press, Oxford, 2000
- [17] J. KAMINKER AND I. PUTNAM, K-Theoretic duality for shifts of finite type, Commun. Math. Phys. 187(1997), pp. 509–522.
- [18] J. KAMINKER, I.F. PUTNAM AND M. WHITTAKER, K-Theoretic duality for hyperbolic dynamical systems, J. Reine Angew. Math. 730(2017), pp. 263–299.
- [19] J. KAMINKER AND C. L. SCHOCHET, Spanier-Whitehead K-duality for C*-algebras, J. Topol. Anal. 11(2019), pp. 21–52.
- [20] G. G. KASPAROV, Topological invariants of elliptic operators I : K-homology, Math. USSR Izvestijia 9(1975), pp. 751–792.
- [21] G. G. KASPAROV, The operator K-functor and extensions of C^{*}-algebras, Math. USSR Izvestijia 16(1981), pp. 513–572.
- [22] K. MATSUMOTO, On strong extension groups of Cuntz-Krieger algebras, preprint, arXiv:2208.08001 [math.OA].
- [23] W. L. PASCHKE, K-theory for commutants in the Calkin algebra, Pacific J. Math. 95(1981), pp. 427–437.
- [24] W. L. PASCHKE AND N. SALINAS, *Matrix algebras over* \mathcal{O}_n , Michigan Math. J. **26**(1979), pp. 3–12.
- [25] M. PIMSNER AND S. POPA, The Ext-groups of some C*-algebras considered by J. Cuntz, Rev. Roum. Math. Pures et Appl. 23(1978), pp. 1069–1076.
- [26] I. RAEBURN AND W. SZYMAŃSKI, Cuntz-Krieger algebras of infinite graphs and matrices, Trans. Amer. Math. Soc. 356(2004), 39–59.
- [27] M. RØRDAM, Classification of Cuntz-Krieger algebras, K-theory 9(1995), pp. 31–58.