

Edge universality of random normal matrices generalizing to higher dimensions

L. D. Molag*

Faculty of Mathematics and Faculty of Physics, Bielefeld University, P.O. Box 100131, D-33501 Bielefeld, Germany

(Dated: August 31, 2022)

As recently proved in generality by Hedenmalm and Wennman, it is a universal behavior of complex random normal matrix models that one finds a complementary error function behavior at the boundary (also called edge) of the droplet as the matrix size increases. Such behavior is seen both in the density, and in the off-diagonal case, where the Faddeeva plasma kernel emerges. These results are neatly expressed with the help of the outward unit normal vector on the edge. We prove that such universal behaviors transcend this class of random normal matrices, being also valid in higher dimensional determinantal point processes, defined on \mathbb{C}^d . The models under consideration concern higher dimensional generalizations of the determinantal point processes describing the eigenvalues of the complex Ginibre ensemble and the complex elliptic Ginibre ensemble. These models describe a system of particles in \mathbb{C}^d with mutual repulsion, that are confined to the origin by an external field $V(z) = |z|^2 - \tau \operatorname{Re}(z_1^2 + \dots + z_d^2)$, where $0 \leq \tau < 1$. Their average density of particles converges to a uniform law on a $2d$ -dimensional ellipsoidal region. It is on the boundary of this region that we find a complementary error function behavior and the Faddeeva plasma kernel. To the best of our knowledge, this is the first instance of the Faddeeva plasma kernel emerging in a higher dimensional model.

I. INTRODUCTION

In 1965, Ginibre introduced a complex $n \times n$ Gaussian matrix with i.i.d. entries of mean zero and unit variance, that we now call the complex Ginibre ensemble [18], along with a real and symplectic version. In this random matrix ensemble, one considers $n \times n$ complex matrices M that are distributed as

$$\frac{1}{Z_n} e^{-\operatorname{Tr}(M^*M)} dM, \quad (1)$$

where dM is the standard Lebesgue measure on this space of matrices, and Z_n is some normalization constant. A complex random normal matrix model is any random matrix model described by (1), with M^*M replaced by a general external field $V(M)$ (satisfying some normalization requirement), on the space of $n \times n$ complex normal matrices. One major difference with, e.g., the general unitary ensembles, is that the eigenvalues of such matrices are not necessarily real. Such models are therefore called non-Hermitian. The eigenvalues z_1, \dots, z_n of the Ginibre ensemble are distributed as

$$\frac{1}{c_n} \prod_{1 \leq i < j \leq n} |z_i - z_j|^2 \prod_{j=1}^n e^{-|z_j|^2} d^2 z_j, \quad z_1, \dots, z_n \in \mathbb{C},$$

where c_n is some normalization constant. In deriving this JPDF, Ginibre assumed that all eigenvalues are distinct, as the remaining set of matrices has measure 0. A less restrictive, but still sufficient condition would be to demand that the matrices are normal. The eigenvalues form a determinantal point process (DPP) in the plane, i.e., the k -point correlation functions can be expressed as

$$\rho_n^{(k)}(z_1, \dots, z_k) = \det(\mathcal{K}_n(z_i, z_j))_{1 \leq i, j \leq k},$$

where $\mathcal{K}_n : \mathbb{C}^2 \rightarrow \mathbb{C}$ is a function called the correlation kernel. For the Ginibre ensemble it may be expressed in the particularly simple form

$$\mathcal{K}_n(z, w) = \frac{1}{\pi} e^{-\frac{|z|^2 + |w|^2}{2}} \sum_{j=0}^{n-1} \frac{(z\bar{w})^j}{j!}. \quad (2)$$

Ginibre proved that the average density of particles $\rho_n^{(1)}(z) = \frac{1}{n} \mathcal{K}_n(z, z)$ converges to a uniform distribution on the unit disc under a proper scaling [18]. Namely, we have

$$\lim_{n \rightarrow \infty} n \rho_n^{(1)}(\sqrt{n}z) = \begin{cases} \frac{1}{\pi}, & |z| < 1, \\ \frac{1}{2\pi}, & |z| = 1, \\ 0, & |z| > 1. \end{cases} \quad (3)$$

* lmolag@math.uni-bielefeld.de

(The case $|z| = 1$ was not treated in [18] though.) The region $|z| \leq 1$, where the limiting distribution lives, is called the *droplet*, the interior $|z| < 1$ is often called the *bulk*. The boundary of the droplet, $|z| = 1$, is called the *edge*. The edge will be our main focus in this paper (albeit for different but related models). The pointwise limit for $|z| = 1$ in (3) is actually a direct consequence of a more general result concerning edge scaling limits. For any z in the unit circle, and any $u, v \in \mathbb{C}$, we have

$$\lim_{n \rightarrow \infty} \mathcal{K}_n(\sqrt{n}z + u, \sqrt{n}z + v) = \frac{1}{2\pi} e^{u\bar{v} - \frac{|u|^2 + |v|^2}{2}} \operatorname{erfc}\left(\frac{\bar{v}z + u\bar{z}}{\sqrt{2}}\right), \quad (4)$$

where erfc is the complementary error function, given by

$$\operatorname{erfc}(z) = 1 - \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt.$$

Alternatively, we may write (4) as

$$\lim_{n \rightarrow \infty} \mathcal{K}_n((\sqrt{n} - u)z, (\sqrt{n} - v)z) = \frac{1}{2\pi} e^{u\bar{v} - \frac{|u|^2 + |v|^2}{2}} \operatorname{erfc}\left(\frac{u + \bar{v}}{\sqrt{2}}\right). \quad (5)$$

The limiting kernel on the RHS in (5) is known as the Faddeeva plasma kernel [12, 20] (note that Hedenmalm and Wenmann use a different definition for the error function), and seems to have been first derived as a scaling limit in [13]. This edge behavior is known to be universal, i.e., it arises as a scaling limit in a large class of other models. For example, Tao and Vu proved that it holds in the vague topology for random normal matrix models (real or complex), that match up to four moments with the Ginibre ensemble [26]. Ameur, Kang, and Makarov proved that it holds for random normal matrix models with radially symmetric external field [5]. Very recently, it was proved by Hedenmalm and Wennman that the behavior is universal in random normal matrix models, under weak assumptions on the external field [20] (being “1-admissible”), essentially bringing the questions about edge universality of these models to their logical conclusion. One model that falls in this category is the elliptic Ginibre ensemble of $n \times n$ complex matrices with parameter τ . These are distributed according to

$$\frac{1}{Z_n} e^{-\frac{1}{1-\tau^2}(M^*M - \frac{\tau}{2}(M^2 + (M^*)^2))}.$$

The model describes random matrices of the form $M = \sqrt{1+\tau}G_1 + i\sqrt{1-\tau}G_2$, where G_1, G_2 are $n \times n$ complex Hermitian matrices picked from the GUE. As such, the elliptic Ginibre ensemble interpolates between the Ginibre ensemble ($\tau = 0$) and the GUE ($\tau = 1$). After a scaling $M \rightarrow (1 - \tau^2)^{-\frac{1}{2}}M$, and restricting to normal matrices, the corresponding eigenvalues are distributed according to

$$\frac{1}{c_n} \prod_{1 \leq i < j \leq n} |z_i - z_j|^2 \prod_{j=1}^n e^{-|z_j|^2 + \tau \operatorname{Re}(z_j^2)} d^2 z_j, \quad z_1, \dots, z_n \in \mathbb{C}.$$

Again, these eigenvalues form a DPP, where now the correlation kernel is given by

$$\mathcal{K}_n(z, w) = \frac{\sqrt{1-\tau^2}}{\pi} \sqrt{\omega(z)\omega(w)} \sum_{j=0}^{n-1} \frac{1}{j!} \left(\frac{\tau}{2}\right)^j H_j\left(\sqrt{\frac{1-\tau^2}{2\tau}}z\right) \overline{H_j\left(\sqrt{\frac{1-\tau^2}{2\tau}}w\right)}, \quad (6)$$

where

$$\omega(z) = e^{-|z|^2 + \tau \operatorname{Re}(z^2)},$$

and $H_j(z) = (-1)^j e^{z^2} \frac{d^j}{dz^j} e^{-z^2}$ is the Hermite polynomial of degree j . The asymptotic behavior for large n of such a kernel is generally harder to derive, because there does not exist a Christoffel-Darboux formula for planar orthogonal polynomials, and tools such as the Deift-Zhou steepest descent method [11, 14] for Riemann-Hilbert problems cannot be applied unless there happens to be some additional structure. The average density of particles also converges to a uniform distribution, where now the bulk is not given by the unit disc, but by the elliptic domain

$$\mathcal{E}_\tau = \left\{ z \in \mathbb{C} : \frac{1-\tau}{1+\tau} \operatorname{Re}(z)^2 + \frac{1+\tau}{1-\tau} \operatorname{Im}(z)^2 < 1 \right\}.$$

Lee and Riser were able to obtain fine asymptotics for the density $\rho_n^{(1)}$ of the Ginibre ensemble and the elliptic Ginibre ensemble [23]. Picking $z \in \partial E_\tau$, and letting \mathbf{n} denote the outward normal vector on ∂E_τ in z , they showed that

$$\begin{aligned} n\rho_n^{(1)}(\sqrt{n}z + \lambda\mathbf{n}) &= \frac{1}{2\pi} \operatorname{erfc}(\sqrt{2}\lambda) + \frac{\kappa}{\sqrt{n}} \frac{1}{3\sqrt{2\pi^3}} (\lambda^2 - 1) e^{-2\lambda^2} \\ &\quad + \frac{1}{n} \frac{e^{-2\lambda^2}}{\sqrt{2\pi^3}} \left(\kappa^2 \frac{2\lambda^5 - 8\lambda^3 + 3\lambda}{18} + \left(\frac{(\partial_s \kappa)^2}{9\kappa^2} - \frac{\partial_s^2 \kappa}{12\kappa} \right) \lambda \right) + \mathcal{O}(n^{-\frac{3}{2} + 9\nu}), \quad (7) \end{aligned}$$

where κ is the curvature of the ellipse $\partial\mathcal{E}_\tau$ in z , and we have any fixed $0 < \nu < \frac{1}{6}$ (∂_s denotes the derivative with respect to the arclength parameter, see [23] for details). The error bound is uniform for $\lambda \in \mathbb{R}$ with $\lambda = \mathcal{O}(n^\nu)$ and $z \in \partial\mathcal{E}_\tau$. The first treatment of the off-diagonal case for the elliptic Ginibre ensemble can be found in Riser's thesis [24], and its error bounds were recently refined by Byun and Ebke in a paper about the symplectic elliptic Ginibre ensemble [9]. Namely, for some explicit unimodular factors $c_n : \partial\mathcal{E}_\tau \times \mathbb{C} \rightarrow \mathbb{T}$ (where $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$), and any $\varepsilon > 0$, we have

$$c_n(z, u) \overline{c_n(z, v)} \mathcal{K}_n(\sqrt{n}z + u\mathbf{n}, \sqrt{n}z + v\mathbf{n}) = \frac{1}{2\pi} \exp\left(u\bar{v} - \frac{|u|^2 + |v|^2}{2}\right) \operatorname{erfc}\left(\frac{u + \bar{v}}{\sqrt{2}}\right) + \frac{1}{\sqrt{n}} \exp\left(-\frac{|u|^2 + u^2 + |v|^2 + \bar{v}^2}{2}\right) \kappa \frac{u^2 + \bar{v}^2 - u\bar{v} - 1}{3\sqrt{2}\pi^3} + \mathcal{O}(n^{-1+\varepsilon}), \quad (8)$$

as $n \rightarrow \infty$, uniformly for $z \in \partial\mathcal{E}_\tau$, and $u, v \in \mathbb{C}$ bounded (in fact, when $0 < \varepsilon < \frac{1}{2}$, the condition $u, v = \mathcal{O}(n^{\varepsilon/3})$ is sufficient for the error bound to be uniform). Here κ is the curvature of $\partial\mathcal{E}_\tau$ in z . Note that these unimodular factors drop out when calculating determinants, and are thus irrelevant for the k -point correlation functions.

The goal of this paper is to show that edge scaling limits such as (7) and (8) hold more generally for higher dimensional DPPs with correlation kernel of the form

$$\mathcal{K}_n(z, w) = \sum_{|j| < n} \prod_{k=1}^d \sqrt{\omega(z_k) \omega(w_k)} P_{j_k}(z_k) \overline{P_{j_k}(w_k)}, \quad z, w \in \mathbb{C}^d, \quad (9)$$

for some planar weight $\omega : \mathbb{C} \rightarrow [0, \infty)$ and polynomials P_j that are orthonormal with respect to the weight, and the summation is over all multi-indices (j_1, \dots, j_d) such that $|j| = j_1 + \dots + j_d < n$. These weights are assumed to vanish sufficiently fast at ∞ , otherwise there is no bulk, end thus no edge. One motivation for studying such higher dimensional processes comes from [2]. There, a DPP on \mathbb{C}^d , essentially a higher dimensional generalization of the elliptic Ginibre ensemble, was considered, with correlation kernel

$$\mathcal{K}_n(z, w) = \left(\frac{\sqrt{1-\tau^2}}{\pi}\right)^d \sum_{|j| < n} \frac{\left(\frac{\tau}{2}\right)^j}{j_1! \dots j_d!} \prod_{k=1}^d \sqrt{\omega(z_k) \omega(w_k)} H_{j_k}\left(\sqrt{\frac{1-\tau^2}{2\tau}} z_k\right) \overline{H_{j_k}\left(\sqrt{\frac{1-\tau^2}{2\tau}} w_k\right)}, \quad z, w \in \mathbb{C}^d, \quad (10)$$

where $\omega(z) = e^{-|z|^2 + \tau \operatorname{Re}(z^2)}$, and $0 < \tau < 1$. The model describes a system of particles in \mathbb{C}^d that repel each other (due to the determinantal structure), which are confined to the origin due to the external field $V(z) = |z|^2 - \tau \operatorname{Re}(z_1^2 + \dots + z_d^2)$. Kernels in higher dimensions of the type (9) have been studied by Berman [6] (for more general multivariate weights in fact). In particular, (9), without the factor of weights, is the Bergman kernel of the Hilbert space of all polynomials $H(z_1, \dots, z_n)$ of total degree $< n$ with weighted norm

$$\|H\| = \int_{\mathbb{C}^d} |H(z_1, \dots, z_n)|^2 \prod_{k=1}^d \omega(z_k) d^2 z_k.$$

For $d = 1$, the kernel (10) coincides with the kernel in (6). In the limit $\tau \rightarrow 1$, this model corresponds to spinless free fermions with quantum harmonic oscillator in \mathbb{R}^d [10]. One major reason to investigate the model for $d > 1$, was that it allowed to probe a weak non-Hermiticity regime [15–17]. For $d = 1$ it is known that interpolating kernels are found in the bulk and on the edge in the weak non-Hermiticity regime [1, 3, 7]. Recently, an interpolating kernel was also found for the rightmost eigenvalue of the elliptic Ginibre ensemble in the weak non-Hermiticity regime [8]. Higher dimensional versions of such interpolating kernels for the bulk and edge were indeed found for $d > 1$ in [2]. For $d = 1$, the limit $\tau \rightarrow 0$ corresponds to spinless free Fermions in two dimensions in a rotating trap [22], with angular speed near some critical value, but it is not clear if such a physical interpretation exists for $d > 1$. We can however, simply write down the corresponding kernel for $d \geq 1$ and $\tau = 0$, namely

$$\mathcal{K}_n(z, w) = \frac{1}{\pi^d} e^{-\frac{|z|^2 + |w|^2}{2}} \sum_{|j| < n} \prod_{k=1}^d \frac{(z_k \bar{w}_k)^{j_k}}{j_k!}, \quad z, w \in \mathbb{C}^d. \quad (11)$$

Indeed, one obtains this expression by taking the limit $\tau \rightarrow 0$ in (10). Note that this is yet another instance of (9), where now $\omega(z) = e^{-|z|^2}$. Henceforth, we agree that \mathcal{K}_n is determined by (10) when $0 < \tau < 1$, and by (11) when $\tau = 0$ (equivalently, \mathcal{K}_n is determined by $\omega(z) = e^{-|z|^2 + \tau \operatorname{Re}(z^2)}$ via (9) for any $0 \leq \tau < 1$). The corresponding DPP can be interpreted as a higher dimensional version of the complex Ginibre ensemble. At least formally, the approach in [22] can be extended to $2d$ dimensions, by generalizing the Hamiltonian in a straightforward way, the model defined via (11) then describing spinless free Fermions in a rotating $2d$ -dimensional space (with angular speed close to some critical value).

Unlike the $d = 1$ case, it is considerably more complicated to write down the JPDF associated to DPPs with kernel (9). For example, in what is probably the simplest case, i.e., the kernel (11), writing down the JPDF merely for $n = 2$ gives

$$\rho_2^{(2)}(z, w) = (|z - w|^2 + |z|^2|w|^2 - |z \cdot w|^2) \frac{e^{-|z|^2 - |w|^2}}{\pi^{2d}}, \quad z, w \in \mathbb{C}^d.$$

Here $z \cdot w = z_1 \bar{w}_1 + \dots + z_d \bar{w}_d$ denotes the dot product. A Vandermonde type expression, with factors expressing the mutual distances between the points, is therefore not going to work, and we have little hope that a closed form expression can be derived for general n . Consequently, it is not easy to find an explicit equilibrium problem for measures on \mathbb{C}^d corresponding to the DPP. An associated equilibrium problem for measures on \mathbb{C}^d can be found in [6], although its characterisation is somewhat abstract. For the particular cases of (10) and (11), it is implied by [6, Theorem 3.4] that we should find a uniform law, given by the so-called Monge-Ampere measure [21], although it is not a priori clear what the support is. Nevertheless, it is possible to determine the support [2]. Under a scaling $(z, w) \mapsto \sqrt{n}(z, w)$, the average density of particles for both (10) and (11) converges to a uniform distribution on a $2d$ -dimensional ellipsoidal region, given by

$$\mathcal{E}_\tau^d = \left\{ z \in \mathbb{C}^d : \frac{1 - \tau}{1 + \tau} |\operatorname{Re} z|^2 + \frac{1 + \tau}{1 - \tau} |\operatorname{Im} z|^2 < 1 \right\}. \quad (12)$$

(For $\tau = 0$, we prove this in Corollary II.4). In the $2d$ -dimensional bulk, defined via (12), there is a local scaling limit in the form of a factorization in Ginibre kernels [2]. That is, there exist unimodular functions $c_n : \partial \mathcal{E}_\tau^d \times \mathbb{C}^d \rightarrow \mathbb{T}$ such that

$$c_n(z, u) \overline{c_n(z, v)} \mathcal{K}_n(\sqrt{n}z + u, \sqrt{n}z + v) = \frac{1}{\pi^d} \exp\left(u \cdot v - \frac{|u|^2 + |v|^2}{2}\right),$$

as $n \rightarrow \infty$, uniformly on compact sets of $z \in \mathcal{E}_\tau^d$ and $u, v \in \mathbb{C}$ (for $\tau = 0$, see Corollary IV.4). This bulk scaling limit is well-known to be universal for random normal matrices under certain conditions ($d = 1$) [4]. A scaling limit at the edge $\partial \mathcal{E}_\tau^d$ was not obtained in [2]. The aim of this paper is to prove that we find scaling limits, similar to (7) and (8), for the DPP with kernel (10) or (11). Indeed, we find higher dimensional generalizations. We start with a result for the average density of points $\rho_n^{(1)}(z)$ (for (10) when $0 < \tau < 1$, and (11) when $\tau = 0$).

Theorem I.1. *Let d be a positive integer, let $0 \leq \tau < 1$, and let $0 < \nu < \frac{1}{6}$. Let $z \in \partial \mathcal{E}_\tau^d$, and denote by \mathbf{n} the outward unit normal vector on $\partial \mathcal{E}_\tau^d$ in z . Then we have as $n \rightarrow \infty$ that*

$$n^d \rho_n^{(1)}(\sqrt{n}z + \lambda \mathbf{n}) = \frac{d!}{2\pi^d} \operatorname{erfc}(\sqrt{2}\lambda) + \frac{\kappa}{\sqrt{n}} \frac{d!}{3\pi^d \sqrt{2\pi}} \left(\lambda^2 - 1 + \mathbf{1}_{\tau \neq 0} \frac{d-1}{(2\kappa^2)^{\frac{1}{3}}} + \mathcal{O}(n^{-\frac{1}{2}+3\nu}) \right) e^{-2\lambda^2}, \quad (13)$$

uniformly for $z \in \partial \mathcal{E}_\tau^d$, and $\lambda \in \mathbb{R}$ such that $\lambda = \mathcal{O}(n^\nu)$, where $\kappa = \kappa(z)$ is defined by

$$\kappa(z) = \frac{1}{\left((|\operatorname{Re} z|^2 - |\operatorname{Im} z|^2 - \frac{4\tau}{1-\tau^2})^2 + 4|\operatorname{Re} z|^2 |\operatorname{Im} z|^2 \right)^{\frac{3}{4}}}. \quad (14)$$

This result should be seen as an extension of (7) to higher dimensions (although we give fewer terms in the expansion). This is not entirely surprising. The asymptotic behavior in terms of the complementary error function was also found near the edge in higher dimensional models investigated by Ross-Singer [25] and Zelditch-Zhou [27], although we believe that our models defined via (10) and (11) fall in a different category. It is an interesting question whether the universality class contains other models of the form (9). One possible direction to pursue is as follows. It is known that a version of the Mehler kernel exists for generalized Laguerre polynomials, expressed via the so-called Hardy-Hill formula, and an adaptation of the approach of the current paper and [2] will likely work here. Perhaps the universality class can be extended to multivariate (exponentially varying) weights and their corresponding multivariate orthogonal polynomials. At the moment, it is not known whether, analogous to the $d = 1$ case, the expression κ in (14) has a geometric interpretation pertaining to $\partial \mathcal{E}_\tau^d$ (although obviously, it yields the curvature of $\partial \mathcal{E}_\tau$ in $|\operatorname{Re} z| + i|\operatorname{Im} z|$).

We also derive a scaling limit, essentially a higher dimensional analogue of the Faddeeva plasma kernel, for the general, i.e., diagonal and off-diagonal, case.

Theorem I.2. *Let d be a positive integer, let $0 \leq \tau < 1$, and let $0 < \nu < \frac{1}{6}$. Let $z \in \partial \mathcal{E}_\tau^d$, and denote by \mathbf{n} the outward unit normal vector on $\partial \mathcal{E}_\tau^d$ in z . Then there exist continuous unimodular functions $c_n : \partial \mathcal{E}_\tau^d \times \mathbb{C}^d \rightarrow \mathbb{T}$ such that*

$$c_n(z, u) \overline{c_n(z, v)} \pi^d \exp\left(\frac{|u|^2 + |v|^2}{2} - u \cdot v\right) \mathcal{K}_n(\sqrt{n}z + u, \sqrt{n}z + v) = \frac{1}{2} \operatorname{erfc}\left(\frac{u \cdot \mathbf{n} + \mathbf{n} \cdot v}{\sqrt{2}}\right) + \exp\left(-\frac{(u \cdot \mathbf{n} + \mathbf{n} \cdot v)^2}{2}\right) \frac{\mathcal{O}(1 + |u|^2 + |v|^2)}{\sqrt{n}},$$

as $n \rightarrow \infty$, uniformly for $z \in \partial \mathcal{E}_\tau^d$, and $u, v \in \mathbb{C}^d$ such that $|u|, |v| = \mathcal{O}(n^\nu)$.

Indeed, this result shows that the universality of (4) is not limited to the eigenvalues of random normal matrices, but its universality class includes higher dimensional models as well. Again, we wonder if such edge behavior also holds in models described by (9) with other weights ω , or more general multivariate weights. When u and v are picked from a bounded set, we infer that

$$\lim_{n \rightarrow \infty} c_n(z, u) \overline{c_n(z, v)} \mathcal{K}_n(\sqrt{n}z + u, \sqrt{n}z + v) = \frac{1}{2\pi^d} \exp\left(u \cdot v - \frac{|u|^2 + |v|^2}{2}\right) \operatorname{erfc}\left(\frac{u \cdot \mathbf{n} + \mathbf{n} \cdot v}{\sqrt{2}}\right).$$

We obtain the Faddeeva plasma kernel after a substitution $u \rightarrow u\mathbf{n}, v \rightarrow v\mathbf{n}$, where now $u, v \in \mathbb{C}$. To the best of our knowledge, a scaling limit of this form (excluding the diagonal case) has not yet appeared in the literature for higher dimensional models. For $d = 1$, we can be more explicit about the error, and indeed, we manage to rederive a slightly stronger version of (8) (see Proposition V.1). We consider our derivation of this result of independent interest.

The paper is built up as follows. In Section II we recap some results from [2], that were used for a steepest descent analysis to derive the asymptotic behavior of (10). The reason that the edge limit was not treated in [2], is that it corresponds to a more complicated situation where a saddle point and pole coalesce in the limit $n \rightarrow \infty$. In Section III we clarify how this situation can be treated. Some preparatory relevant identities and estimates are derived in Section IV, and finally, in Section V we prove the main theorems.

ACKNOWLEDGMENTS

The author is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – SFB 1283/2 2021 – 317210226 "Taming uncertainty and profiting from randomness and low regularity in analysis, stochastics and their applications". The author thanks Gernot Akemann, Sungsoo Byun, Maurice Duits and Markus Ebke for useful discussions.

II. APPROACH AND SET UP

A. Single integral representation

Our primary tool to prove the main results will be a steepest descent analysis. In [2] it was shown, using a formula for the Mehler kernel, that (10) (with a different scaling) admits a single integral representation of the form

$$\mathcal{K}_n(\sqrt{n}z, \sqrt{n}w) = \left(\frac{\sqrt{1-\tau^2}}{\pi}\right)^d \sqrt{\omega(\sqrt{n}z_+) \omega(\sqrt{n}z_-)} I_{n,\tau}^d(z_\pm), \quad (15)$$

valid for any $z, w \in \mathbb{C}$, where

$$z_\pm = \frac{\sqrt{\sinh 2\xi_\tau}}{2} \left(\sqrt{\sum_{j=1}^d (z_j + \bar{w}_j)^2} \pm \sqrt{\sum_{j=1}^d (z_j - \bar{w}_j)^2} \right), \quad (16)$$

and, with γ_0 a small positively oriented loop around $s = 0$, the integral $I_{n,\tau}^d(z_\pm)$ is given by

$$I_{n,\tau}^d(z_\pm) = -\frac{1}{2\pi i} \oint_{\gamma_0} \frac{e^{nF(s)}}{s-\tau} \frac{ds}{(1-s^2)^{\frac{d}{2}}} \quad (17)$$

and $F(s) = F_\tau(z_\pm; s)$ is given by

$$F(s) = \frac{s}{1+s} \frac{(z_+ + z_-)^2}{2} - \frac{s}{1-s} \frac{(z_+ - z_-)^2}{2} - \log s + \log \tau. \quad (18)$$

We do not have to be explicit about the choice of branch of the square roots in (16), because the choice is irrelevant for (18), and thus for $\mathcal{K}_n(\sqrt{n}z, \sqrt{n}w)$. The same holds for the branch of the logarithm in (18). Nevertheless, let us use the convention that

logarithms and power functions will be defined as $\log z = \log |z| + i \arg z$ and $z^\alpha = |z|^\alpha e^{i\alpha \arg z}$, where $\arg z \in (-\pi, \pi]$. The saddle points of $F(s)$ have a surprisingly simple form in elliptic coordinates. We write

$$z_\pm = \sqrt{2} \cosh(\xi_\pm + i\eta_\pm),$$

where $\xi_\pm \geq 0$ and $\eta_\pm \in (-\pi, \pi]$ when $\xi_\pm > 0$, while $\eta_\pm \in [0, \pi]$ when $\xi_\pm = 0$. Note that any constant value of ξ_\pm corresponds to an ellipse with vertex $\sqrt{2} \cosh \xi_\pm$ and co-vertex $\sqrt{2} \sinh \xi_\pm$. The edge, i.e., the ellipse forming the boundary of the droplet, is described by the particular choice $\xi_\pm = \xi_\tau$, where $\xi_\tau = \frac{1}{2} \log \frac{1}{\tau}$. We define

$$a = e^{\xi_+ + \xi_- + i(\eta_+ + \eta_-)}, \quad \text{and} \quad b = e^{\xi_+ - \xi_- + i(\eta_+ - \eta_-)}. \quad (19)$$

With these notations, the following result was derived in [2].

Proposition II.1 ($0 < \tau < 1$).

If $z_+, z_- \in \mathbb{C} \setminus \{-\sqrt{2}, \sqrt{2}\}$, then the saddle points of $s \mapsto F_\tau(z_\pm; s)$ are simple, and we have the following:

- (i) When $z_+ \neq \pm z_-$, there are exactly four saddle points given by a, a^{-1}, b and b^{-1} .
- (ii) When $z_+ = \pm z_-$ and $z_+ \neq 0$, there are exactly two saddle points, which are given by a and a^{-1} .

If $z_+ \in \{-\sqrt{2}, \sqrt{2}\}$ or $z_- \in \{-\sqrt{2}, \sqrt{2}\}$, then all saddle points have order two and we have the following:

- (iii) When $z_\pm \in \{-\sqrt{2}, \sqrt{2}\}$ and $z_\mp \notin \{-\sqrt{2}, \sqrt{2}\}$, then we have two saddle points $a = b^{\mp 1}$ and $a^{-1} = b^{\pm 1}$.
- (iv) When $z_+ = \pm z_- \in \{-\sqrt{2}, \sqrt{2}\}$, then we have one saddle point $a^{-1} = a = b = b^{-1} = \pm 1$.

Finally, when $z_+ = z_- = 0$ there are no saddle points.

Furthermore, in order to understand what deformations of γ_0 were allowed, the following theorem was proved. Notice in particular that one can deform γ_0 to the circle $|s| = |a|^{-1}$ when $\xi_+, \xi_- > 0$.

Theorem II.2 ($0 < \tau < 1$). With the notations as above, we have the inequality

$$\operatorname{Re} F(s) \leq \operatorname{Re} F(a^{-1}), \quad |s| = |a|^{-1}. \quad (20)$$

- (i) When $\xi_+ > 0$ and $\xi_- > 0$, we have equality if and only if $s = a^{-1}$.
- (ii) When $\xi_+ > 0$ and $\xi_- = 0$, we have equality if and only if $s = a^{-1}$ or $s = b^{-1}$.
- (iii) When $\xi_+ = 0$ and $\xi_- > 0$, we have equality if and only if $s = a^{-1}$ or $s = b$.
- (iv) When $\xi_+ = 0$ and $\xi_- = 0$, we have equality for all s .

So far, these results describe \mathcal{K}_n for $0 < \tau < 1$ (as defined in (10)). For $\tau = 0$, we can significantly simplify \mathcal{K}_n (as defined in (11)).

Proposition II.3 ($\tau = 0$). Let $z, w \in \mathbb{C}^d$. Then we may write

$$\mathcal{K}_n(\sqrt{n}z, \sqrt{n}w) = \frac{1}{\pi^d} e^{-n \frac{|z|^2 + |w|^2}{2}} \sum_{j=0}^{n-1} \frac{(nz \cdot w)^j}{j!}. \quad (21)$$

Proof. We start by noticing that

$$e^{(z \cdot w)s} = \prod_{k=1}^d e^{(z_k \bar{w}_k)s} = \prod_{k=1}^d \sum_{j_k=0}^{\infty} \frac{(z_k \bar{w}_k s)^{j_k}}{j_k!} = \sum_{m=0}^{\infty} \sum_{|j|=m} s^m \prod_{k=1}^d \frac{(z_k \bar{w}_k)^{j_k}}{j_k!}.$$

Then by applying the residue theorem in two directions, we have

$$\sum_{|j| < n} \prod_{k=1}^d \frac{(z_k \bar{w}_k)^{j_k}}{j_k!} = \frac{1}{2\pi i} \oint_{\gamma_0} e^{(z \cdot w)s} \left(1 + \frac{1}{s} + \dots + \frac{1}{s^{n-1}} \right) \frac{ds}{s} = \sum_{j=0}^{n-1} \frac{(z \cdot w)^j}{j!}. \quad (22)$$

Now substituting $(z, w) \mapsto \sqrt{n}(z, w)$, and multiplying with the remaining factors in (11), we arrive at (21). \square

Comparing with (2), we infer that \mathcal{K}_n for $d \geq 1$ and $\tau = 0$ is closely related to the Ginibre ensemble ($d = 1$). In particular, some properties are immediately inherited from the Ginibre ensemble. For instance, the model has a bulk, which is given by the $2d$ -dimensional unit ball in \mathbb{C}^d .

Corollary II.4 ($\tau = 0$). *The average density of particles converges to a uniform law on the unit ball in \mathbb{C}^d . More precisely, we have*

$$\lim_{n \rightarrow \infty} n^d \rho_n^{(1)}(\sqrt{n}z) = \begin{cases} \frac{d!}{\pi^d}, & |z| < 1, \\ \frac{d!}{2\pi^d}, & |z| = 1, \\ 0, & |z| > 1. \end{cases} \quad (23)$$

The convergence is uniform on compact subsets such that $|z| \neq 1$.

Proof. By some straightforward combinatorial arguments, the number of points of the DPP defined via (11) is given by the binomial coefficient $\binom{n+d-1}{d}$, which behaves as $\frac{n^d}{d!}(1 + \mathcal{O}(1/n))$ for large n . For the average density of particles, one takes $z = w$. Then (21) turns into

$$\mathcal{K}_n(\sqrt{n}z, \sqrt{n}z) = \frac{1}{\pi^d} e^{-n|z|^2} \sum_{j=0}^{n-1} \frac{(n|z|^2)^j}{j!}.$$

This, apart from a factor π^{d-1} , is simply the (rescaled) average density of points of the Ginibre ensemble ($d = 1$) in $|z|$ (or any rotation in the plane thereof), and we obtain the result directly from (3). \square

Though Proposition II.3 gives an insightful relation between the $d > 1$ and $d = 1$ model for $\tau = 0$, it will turn out to be both instructive and beneficial to find a single integral representation for this model as well. We have the following result.

Proposition II.5 ($\tau = 0$). *Let $z, w \in \mathbb{C}^d$. We may write*

$$\mathcal{K}_n(\sqrt{n}z, \sqrt{n}w) = \frac{1}{\pi^d} e^{-n \frac{|z|^2 + |w|^2}{2}} I_{n,0}^d(z \cdot w), \quad (24)$$

where, for $\zeta \in \mathbb{C}$, we have

$$I_{n,0}^d(\zeta) = -\frac{1}{2\pi i} \oint_{\gamma_0} \frac{e^{nF(s)}}{s-1} ds, \quad (25)$$

with γ_0 a small positively oriented loop around 0, and $F(s) = F(\zeta; s)$ is defined by

$$F(s) = \zeta s - \log s. \quad (26)$$

Proof. Since $s = 1$ is not enclosed by γ_0 (which is assumed to be small), the residue theorem implies that (22) can alternatively be written as

$$\sum_{|j| < n} \prod_{k=1}^d \frac{(z_k \bar{w}_k)^{j_k}}{j_k!} = \frac{1}{2\pi i} \oint_{\gamma_0} e^{(z \cdot w)s} \frac{1-s^{-n}}{1-s^{-1}} \frac{ds}{s} = -\frac{1}{2\pi i} \oint_{\gamma_0} e^{(z \cdot w)s} s^{-n} \frac{ds}{s-1}. \quad (27)$$

Now substituting $(z, w) \mapsto \sqrt{n}(z, w)$, writing $s^{-n} = e^{-n \log s}$, and multiplying with the remaining factors in (11), we arrive at (24), with $I_{n,0}^d$ and F as defined in (25) and (26) respectively. \square

We state the following proposition. The proof is trivial, and is therefor omitted.

Proposition II.6 ($\tau = 0$). *Let $\zeta \in \mathbb{C}$, and let $F(s) = F(\zeta; s)$ be as in Proposition II.5, i.e., $F(s) = \zeta s - \log s$.*

(i) *When $\zeta \neq 0$, there is only one saddle point $s_0 = \zeta^{-1}$, which is simple.*

In this case $F(s_0) = 1 + \log \zeta$ and $F''(s_0) = \zeta^2$.

(ii) *When $\zeta = 0$, there are no saddle points.*

B. Set up

We shall need the results of the previous subsection, tailored to a specific situation. In our present case, we investigate the asymptotic behavior of $\mathcal{K}_n(\sqrt{n}z + u, \sqrt{n}z + v)$, where $z \in \partial\mathcal{E}_\tau^d$, and $u, v \in \mathbb{C}^d$ satisfy $u, v = \mathcal{O}(n^\nu)$ as $n \rightarrow \infty$, for some fixed $0 < \nu < \frac{1}{6}$. First, we focus on the case $0 < \tau < 1$. Using the definition in (16), we have that

$$\frac{z_\pm}{\sqrt{\sinh 2\xi_\tau}} = \sqrt{|\operatorname{Re}z|^2 + \frac{u+\bar{v}}{\sqrt{n}} \cdot (\operatorname{Re}z) + \frac{(u+\bar{v})^2}{4n}} \pm i\sqrt{|\operatorname{Im}z|^2 + \frac{u-\bar{v}}{\sqrt{n}} \cdot (i\operatorname{Im}z) - \frac{(u-\bar{v})^2}{4n}}, \quad (28)$$

where we use a short-hand notation $\zeta^2 = \zeta_1^2 + \dots + \zeta_d^2$ for any $\zeta \in \mathbb{C}^d$. In particular, z_\pm depend piecewise continuously on z and $\frac{u}{\sqrt{n}}, \frac{v}{\sqrt{n}}$, and we have uniformly for $z \in \partial\mathcal{E}_\tau^d$ and $u, v = \mathcal{O}(n^\nu)$ that

$$\lim_{n \rightarrow \infty} \frac{z_\pm}{\sqrt{\sinh 2\xi_\tau}} = \frac{\hat{z}_\pm}{\sqrt{\sinh 2\xi_\tau}} := |\operatorname{Re}z| + i|\operatorname{Im}z| \in \partial\mathcal{E}_\tau.$$

In particular, there is an $\eta \in [0, \frac{\pi}{2}]$ such that

$$\hat{z}_\pm = \sqrt{2} \cosh(\xi_\tau \pm i\eta).$$

When $d = 1$, we simply have $z_+ = \sqrt{\sinh 2\xi_\tau}(z + u)$ and $z_- = \sqrt{\sinh 2\xi_\tau}(z + v)$. When $d > 1$, there is an essential difference though, and this is caused by the branch-cut of the square roots in (28). In the end, these branch-cuts will somehow drop out (at least to the dominant order), since $I_{n,\tau}^d(z_\pm)$ depends only on $(z_+ \pm z_-)^2$, which have no branch-cut. However, in our derivation we shall have to take the presence of these branch-cuts into account. When z is in a subset of $\partial\mathcal{E}_\tau^d$ such that $\operatorname{Re}z, \operatorname{Im}z \neq 0$, we have

$$\frac{z_\pm}{\sqrt{\sinh 2\xi_\tau}} = \frac{\hat{z}_\pm}{\sqrt{\sinh 2\xi_\tau}} + \frac{u+\bar{v}}{\sqrt{n}} \cdot \frac{\operatorname{Re}z}{|\operatorname{Re}z|} \pm \frac{u-\bar{v}}{\sqrt{n}} \cdot \frac{\operatorname{Im}z}{|\operatorname{Im}z|} + \mathcal{O}\left(\frac{n^{-1+2\nu}}{|\operatorname{Re}z|} + \frac{n^{-1+2\nu}}{|\operatorname{Im}z|}\right) \quad (29)$$

uniformly for $u, v = \mathcal{O}(n^\nu)$ as $n \rightarrow \infty$. On the other hand, when $\operatorname{Re}z = 0$, we have

$$\frac{z_\pm}{\sqrt{\sinh 2\xi_\tau}} = \frac{\hat{z}_\pm}{\sqrt{\sinh 2\xi_\tau}} + \frac{\sqrt{(u+\bar{v})^2}}{\sqrt{n}} \pm \frac{u-\bar{v}}{\sqrt{n}} \cdot \frac{\operatorname{Im}z}{|\operatorname{Im}z|} + \mathcal{O}(n^{-1+2\nu}), \quad (30)$$

while the case $\operatorname{Im}z = 0$ yields

$$\frac{z_\pm}{\sqrt{\sinh 2\xi_\tau}} = \frac{\hat{z}_\pm}{\sqrt{\sinh 2\xi_\tau}} + \frac{u+\bar{v}}{\sqrt{n}} \cdot \frac{\operatorname{Re}z}{|\operatorname{Re}z|} \pm \frac{\sqrt{(u-\bar{v})^2}}{\sqrt{n}} + \mathcal{O}(n^{-1+2\nu}), \quad (31)$$

uniformly for $u, v = \mathcal{O}(n^\nu)$ as $n \rightarrow \infty$. Although we have $z_\pm = \hat{z}_\pm + \mathcal{O}(n^{-\frac{1}{2}+\nu})$ in all cases, it is not a priori clear that the constant implied by this \mathcal{O} term can be taken uniformly for $z \in \partial\mathcal{E}_\tau^d$. Nevertheless, this does indeed hold.

Lemma II.7 ($0 < \tau < 1$). *We have $z_\pm - \hat{z}_\pm = \mathcal{O}(n^{-\frac{1}{2}+\nu})$ as $n \rightarrow \infty$, uniformly for $z \in \partial\mathcal{E}_\tau^d$ and $u, v = \mathcal{O}(n^\nu)$.*

Proof. One simply has to divide into the case $|\operatorname{Re}z|, |\operatorname{Im}z| > n^{-\frac{1}{2}+\nu}$, the case $|\operatorname{Re}z| \leq n^{-\frac{1}{2}+\nu}$, and the case $|\operatorname{Im}z| \leq n^{-\frac{1}{2}+\nu}$. In the first case, we have

$$\begin{aligned} \left| \sqrt{|\operatorname{Re}z|^2 + \frac{u+\bar{v}}{\sqrt{n}} \cdot (\operatorname{Re}z) + \frac{(u+\bar{v})^2}{4n}} - |\operatorname{Re}z| \right| &= \left| \frac{\frac{u+\bar{v}}{\sqrt{n}} \cdot \frac{\operatorname{Re}z}{|\operatorname{Re}z|} + \frac{(u+\bar{v})^2}{4|\operatorname{Re}z|n}}{1 + \sqrt{1 + \frac{u+\bar{v}}{\sqrt{n}} \cdot \frac{\operatorname{Re}z}{|\operatorname{Re}z|} + \frac{(u+\bar{v})^2}{4|\operatorname{Re}z|n}}} \right| \leq \frac{|u+\bar{v}|}{\sqrt{n}} + \frac{|u+\bar{v}|^2}{4n^{\frac{1}{2}+\nu}}, \\ \left| \sqrt{|\operatorname{Im}z|^2 + \frac{u-\bar{v}}{\sqrt{n}} \cdot (i\operatorname{Im}z) - \frac{(u-\bar{v})^2}{4n}} - |\operatorname{Im}z| \right| &= \left| \frac{\frac{u-\bar{v}}{\sqrt{n}} \cdot \frac{i\operatorname{Im}z}{|\operatorname{Im}z|} - \frac{(u-\bar{v})^2}{4|\operatorname{Im}z|n}}{1 + \sqrt{1 + \frac{u+\bar{v}}{\sqrt{n}} \cdot \frac{\operatorname{Re}z}{|\operatorname{Re}z|} + \frac{(u+\bar{v})^2}{4|\operatorname{Re}z|n}}} \right| \leq \frac{|u-\bar{v}|}{\sqrt{n}} + \frac{|u-\bar{v}|^2}{4n^{\frac{1}{2}+\nu}}. \end{aligned} \quad (32)$$

We used here that for any $\zeta \in \mathbb{C}$ we have $\operatorname{Re} \sqrt{\zeta} \geq 0$, and thus for all $\zeta \in \mathbb{C} \cup \{\infty\}$ one has

$$\left| \frac{1}{1 + \sqrt{\zeta}} \right| \leq 1.$$

When we are in the second case, we have

$$\left| \sqrt{|\operatorname{Re} z|^2 + \frac{u + \bar{v}}{\sqrt{n}} \cdot (\operatorname{Re} z) + \frac{(u + \bar{v})^2}{4n}} - |\operatorname{Re} z| \right| \leq \sqrt{n^{-1+2v} + |u + \bar{v}|n^{-1+v} + \frac{|u + \bar{v}|^2}{4n}} + n^{-\frac{1}{2}+v} = n^{-\frac{1}{2}+v} \left(2 + \frac{|u + \bar{v}|}{2n^v} \right),$$

while we still have the same estimate (32). The third case is analogous. Obviously, we can find a constant $C > 0$ such that $|z_{\pm} - \hat{z}_{\pm}| \leq Cn^{-\frac{1}{2}+v}$ for the three cases simultaneously. \square

It will turn out to be convenient to introduce

$$\Delta_{\pm} = \frac{z_{\pm} - \hat{z}_{\pm}}{\sqrt{\hat{z}_{\pm}^2 - 2}} = \frac{\cosh(\xi_{\pm} + i\eta_{\pm}) - \cosh(\xi_{\tau} + i\eta)}{\sinh(\xi_{\tau} + i\eta)}, \quad (33)$$

where, as before, $z_{\pm} = \sqrt{2} \cosh(\xi_{\pm} + i\eta_{\pm})$ and $\hat{z}_{\pm} = \sqrt{2} \cosh(\xi_{\tau} \pm i\eta)$. Rather than working with $u, v = \mathcal{O}(n^v)$ we shall simply consider $z_{\pm} = \hat{z}_{\pm} + \sqrt{\hat{z}_{\pm}^2 - 2} \Delta_{\pm}$, and demand that $\Delta_{\pm} = \mathcal{O}(n^{-\frac{1}{2}+v})$ as $n \rightarrow \infty$. An incidental advantage of this perspective, is that we only need to consider limits that are uniform in $\hat{z}_{\pm} \in \sqrt{\sinh 2\xi_{\tau}} \partial \mathcal{E}_{\tau} = \{\sinh(2\xi_{\tau}) w : w \in \partial \mathcal{E}_{\tau}\}$. For notational convenience, we introduce

$$\hat{z} = \frac{z_+}{\sqrt{\sinh 2\xi_{\tau}}}.$$

Summarizing, our aim is to understand the asymptotic behavior of $I_{n,\tau}^d(z_{\pm})$, where

$$z_{\pm} = \hat{z}_{\pm} + \sqrt{\hat{z}_{\pm}^2 - 2} \Delta_{\pm},$$

uniformly for $\hat{z} \in \partial \mathcal{E}_{\tau}$, under the assumption that $\Delta_{\pm} = \mathcal{O}(n^{-\frac{1}{2}+v})$ as $n \rightarrow \infty$.

Lemma II.8 ($0 < \tau < 1$). *Let a and b be defined as in (19). Then we have*

$$a^{\pm 1} = \tau^{\mp 1} \pm \tau^{\mp 1} (\Delta_+ + \Delta_-) + \mathcal{O}(n^{-1+2v}). \quad (34)$$

$$b^{\pm 1} = e^{\pm 2i\eta} \pm e^{\pm 2i\eta} (\Delta_+ - \Delta_-) + \mathcal{O}(n^{-1+2v}). \quad (35)$$

as $n \rightarrow \infty$, uniformly for $\hat{z} \in \partial \mathcal{E}_{\tau}$. Furthermore, we have

$$\tau a - 1 = \Delta_+ + \Delta_- + \frac{(\Delta_+ + \Delta_-)^2}{2} - \coth(\xi_{\tau} + i\eta) \frac{\Delta_+^2}{2} - \coth(\xi_{\tau} - i\eta) \frac{\Delta_-^2}{2} + \mathcal{O}(n^{-\frac{3}{2}+3v}) \quad (36)$$

as $n \rightarrow \infty$, uniformly for $\hat{z} \in \partial \mathcal{E}_{\tau}$.

Proof. As shown in [2], we can alternatively write the saddle points as

$$a^{\pm 1} = \frac{1}{2} \left(z_+ \pm \sqrt{z_+^2 - 2} \right) \left(z_- \pm \sqrt{z_-^2 - 2} \right) \quad (37)$$

$$b^{\pm 1} = \frac{1}{2} \left(z_+ \pm \sqrt{z_+^2 - 2} \right) \left(z_- \mp \sqrt{z_-^2 - 2} \right). \quad (38)$$

We observe by Taylor series expansion that

$$\begin{aligned} \sqrt{z_{\pm}^2 - 2} &= \sqrt{\hat{z}_{\pm}^2 - 2} + \frac{\hat{z}_{\pm}}{\sqrt{\hat{z}_{\pm}^2 - 2}} (z_{\pm} - \hat{z}_{\pm}) - \frac{1}{(\hat{z}_{\pm}^2 - 2)^{\frac{3}{2}}} (z_{\pm} - \hat{z}_{\pm})^2 + \mathcal{O}(\Delta_{\pm}^3) \\ &= \sqrt{\hat{z}_{\pm}^2 - 2} + \hat{z}_{\pm} \Delta_{\pm} - \frac{\Delta_{\pm}^2}{\sqrt{\hat{z}_{\pm}^2 - 2}} + \mathcal{O}(\Delta_{\pm}^3). \end{aligned}$$

Henceforth

$$\frac{z_{\pm} + \sqrt{z_{\pm}^2 - 2}}{\sqrt{2}} = e^{\xi_{\tau} \pm i\eta} + e^{\xi_{\tau} \pm i\eta} \Delta_{\pm} - \frac{1}{\sinh(\xi_{\tau} \pm i\eta)} \frac{\Delta_{\pm}^2}{2} + \mathcal{O}(\Delta_{\pm}^3). \quad (39)$$

Plugging these in (37), we obtain

$$a^{\pm 1} = \tau^{\mp 1} \pm \tau^{\mp 1}(\Delta_+ + \Delta_-) + \mathcal{O}(n^{-1+2\nu}). \quad (40)$$

Analogously, we find

$$b^{\pm 1} = e^{\pm 2i\eta} \pm e^{\pm 2i\eta}(\Delta_+ - \Delta_-) + \mathcal{O}(n^{-1+2\nu}). \quad (41)$$

Using (39) to one order higher, we find that

$$\tau a - 1 = \Delta_+ + \Delta_- + \Delta_+ \Delta_- - \frac{e^{-\xi_\tau - i\eta}}{\sinh(\xi_\tau + i\eta)} \frac{\Delta_+^2}{2} - \frac{e^{-\xi_\tau + i\eta}}{\sinh(\xi_\tau - i\eta)} \frac{\Delta_-^2}{2} + \mathcal{O}(n^{-\frac{3}{2}+3\nu}).$$

□

Lemma II.8 shows in particular that a, b and b^{-1} lie outside the circle $|s| = |a|^{-1}$ for n big enough. Following Lemma II.2, we intend to deform the integration contour γ_0 to the circle $|s| = |a|^{-1}$, and we should only take the saddle point $s = a^{-1}$ into account. What makes this situation complicated, is that the saddle point a^{-1} and the pole at τ of the integrand of $I_{n,\tau}^d(\hat{z}_\pm + \sqrt{\hat{z}_\pm^2 - 2\Delta_\pm})$ coalesce in the limit $n \rightarrow \infty$.

Since a^{-1} is the only saddle point that will give a contribution, we shall need to know the values of F and F'' only in a^{-1} . By [2], these are given by

$$\begin{aligned} F(a^{-1}) &= 1 + \log \tau + \xi_+ + \xi_- + i(\eta_+ + \eta_-) + \frac{1}{2}e^{-2(\xi_+ + i\eta_+)} + \frac{1}{2}e^{-2(\xi_- + i\eta_-)}, \\ F''(a^{-1}) &= 2a^2 \frac{\sinh(\xi_+ + i\eta_+) \sinh(\xi_- + i\eta_-)}{\sinh(\xi_+ + i\eta_+ + \xi_- + i\eta_-)}. \end{aligned} \quad (42)$$

We would rather express $F''(a^{-1})$ in terms of Δ_\pm however, hence the following lemma. As it turns out, we need not be precise about the behavior of $F(a^{-1})$.

Lemma II.9 ($0 < \tau < 1$). *Uniformly for $\hat{z} \in \partial \mathcal{E}_\tau$, we have as $n \rightarrow \infty$ that*

$$\frac{1}{2}a^{-2}F''(a^{-1}) = \frac{|\sinh(\xi_\tau + i\eta)|^2}{\sinh(2\xi_\tau)} (1 + \coth(\xi_\tau + i\eta)\Delta_+ + \coth(\xi_\tau - i\eta)\Delta_- - \coth(2\xi_\tau)(\Delta_+ + \Delta_-)) + \mathcal{O}(n^{-1+2\nu}). \quad (43)$$

Proof. As observed in [2], we can write $F''(a^{-1})$ entirely in terms of the saddle points. Namely

$$F''(a^{-1}) = \frac{(a-b)(a-b^{-1})}{1-a^{-2}}.$$

Plugging in (40) and (41), we find that

$$\begin{aligned} a - b^{\pm 1} &= (\tau^{-1} - e^{\pm 2i\eta})(1 + \Delta_\pm + \coth(\xi_\tau \mp i\eta)\Delta_\mp) + \mathcal{O}(n^{-1+2\nu}), \\ a^2 - 1 &= (\tau^{-2} - 1) \left(1 + 2\frac{\Delta_+ + \Delta_-}{1 - \tau^2} \right) + \mathcal{O}(n^{-1+2\nu}). \end{aligned}$$

We thus have

$$a^{-2}F''(a^{-1}) = \frac{|\tau^{-1} - e^{2i\eta}|^2}{\tau^{-2} - 1} (1 + \coth(\xi_\tau + i\eta)\Delta_+ + \coth(\xi_\tau - i\eta)\Delta_- - \coth(2\xi_\tau)(\Delta_+ + \Delta_-)) + \mathcal{O}(n^{-1+2\nu}).$$

□

So far, we set up everything for the case $0 < \tau < 1$. The case $\tau = 0$ is considerably easier, and there is not much to set up. Substituting $\sqrt{n}(z, w) \mapsto (\sqrt{n}z + u, \sqrt{n}z + v)$ in Proposition II.6, we find that there is exactly one saddle point (for n large enough), which is simple, and given by

$$s_0 = \frac{1}{\left(z + \frac{u}{\sqrt{n}}\right) \cdot \left(z + \frac{v}{\sqrt{n}}\right)} = \left(1 + \frac{u \cdot z + z \cdot v}{\sqrt{n}} + \frac{u \cdot v}{n}\right)^{-1} = 1 - \frac{u \cdot z + z \cdot v}{\sqrt{n}} + \frac{(u \cdot z + z \cdot v)^2 - u \cdot v}{n} + \mathcal{O}(n^{-\frac{3}{2}+3\nu}) \quad (44)$$

as $n \rightarrow \infty$, uniformly for $|z| = 1$ and $u, v \in \mathbb{C}^d$ with $u, v = \mathcal{O}(n^\nu)$. Note that, in this case, just as in the case $0 < \tau < 1$, the saddle point ($s = s_0$) coalesces with the pole ($s = 1$) in the limit $n \rightarrow \infty$. We also find that

$$\begin{aligned} F(s_0) &= 1 - \log s_0 = 1 + \log \left(1 + \frac{u \cdot z + z \cdot v}{\sqrt{n}} + \frac{u \cdot v}{n} \right) = 1 + \frac{u \cdot z + z \cdot v}{\sqrt{n}} - \frac{(u \cdot z + z \cdot v)^2 - u \cdot v}{n} + \mathcal{O}(n^{-\frac{3}{2}+3\nu}), \\ F''(s_0) &= \frac{1}{s_0^2} = \left(1 + \frac{u \cdot z + z \cdot v}{\sqrt{n}} + \frac{u \cdot v}{n} \right)^2, \end{aligned} \quad (45)$$

as $n \rightarrow \infty$, uniformly for $|z| = 1$ and $u, v \in \mathbb{C}^d$ with $u, v = \mathcal{O}(n^\nu)$. As before, rather than using u, v , we shall consider the asymptotic behavior of $I_{n,0}^d(1 + \Delta)$, where we assume that $\Delta = \mathcal{O}(n^{-\frac{1}{2}+\nu})$ as $n \rightarrow \infty$. In the end, we then have to substitute

$$\Delta = \frac{u \cdot z + z \cdot v}{\sqrt{n}} + \frac{u \cdot v}{n}. \quad (46)$$

Having prepared all the ingredients necessary for the steepest descent analysis, we shall explicitly derive the asymptotic behavior of $I_{n,\tau}^d$ in the next section.

III. STEEPEST DESCENT ANALYSIS

For $0 < \tau < 1$, our goal is to derive the asymptotic behavior of $I_{n,\tau}^d(\hat{z}_\pm + \sqrt{\hat{z}_\pm^2 - 2\Delta_\pm})$ with a steepest descent analysis, under the assumption that $\Delta_\pm = \mathcal{O}(n^{-\frac{1}{2}+\nu})$ as $n \rightarrow \infty$. We deform the integration contour γ_0 to the circle $|s| = |a|^{-1} = e^{-\xi_+ - \xi_-}$. We need to understand the behavior of $F(s)$ around the saddle point $s = a^{-1}$ and the pole (of the integrand) $s = \tau$.

When $\tau = 0$, the situation is comparable. Here we attempt to find the large n behavior of $I_{n,0}^d(1 + \Delta)$, under the condition that $\Delta = \mathcal{O}(n^{-\frac{1}{2}+\nu})$ as $n \rightarrow \infty$. The saddle point will coalesce with the pole $s = 1$ in the limit $n \rightarrow \infty$.

A. A saddle point coalescing with a pole

Under proper conditions, it is possible to perform a steepest descent analysis where a saddle point and a pole coalesce. Although our arguments can be extended to more general situations, we consider only the quadratic case on the real line. After the right preparations, Proposition III.1 turns out to be enough for our purposes.

Proposition III.1. *Let $\ell_1 < 0 < \ell_2$ and $\delta > 0$. Uniformly for $p \in \mathbb{C}$ with $\ell_1 + \delta < \operatorname{Re} p < \ell_2 - \delta$, we have as $n \rightarrow \infty$ that*

$$\int_{\ell_1}^{\ell_2} e^{-nt^2} \frac{dt}{t-p} = -\pi i e^{-np^2} \operatorname{erfc}(i\sqrt{n}p) + \mathcal{O}\left(\frac{e^{-n\ell_1^2} + e^{-n\ell_2^2}}{n}\right),$$

where the path from ℓ_1 to ℓ_2 is such that p is to the right of the path.

Proof. Without loss of generality, we assume that $\operatorname{Im} p < 0$ and that the curve from ℓ_1 to ℓ_2 is a line segment. First we note that

$$\begin{aligned} \int_{-\infty}^{\ell_1} e^{-nt^2} \frac{dt}{t-p} + \int_{\ell_2}^{\infty} e^{-nt^2} \frac{dt}{t-p} &\leq \frac{1}{\operatorname{Re} p - \ell_1} \sqrt{\frac{\pi}{n}} \operatorname{erfc}(\sqrt{n}\ell_1) + \frac{1}{\ell_2 - \operatorname{Re} p} \sqrt{\frac{\pi}{n}} \operatorname{erfc}(\sqrt{n}\ell_2) \\ &\leq \frac{1}{\delta} \frac{e^{-n\ell_1^2}}{\pi n \ell_1} + \frac{1}{\delta} \frac{e^{-n\ell_2^2}}{\pi n \ell_2}, \end{aligned}$$

as $n \rightarrow \infty$. Hence, we may just as well replace ℓ_1 by $-T$ and ℓ_2 by T for some large $T > 0$, and integrate over the line segment $[-T, T]$ in what follows. We create a closed integration contour by adding the line segments $[\operatorname{Im} p - T, p - \varepsilon]$, $[p + \varepsilon, \operatorname{Im} p + T]$ and $\pm T + i[\operatorname{Im} p, 0]$, and the upper semicircle from $p + \varepsilon$ to $p - \varepsilon$ for some small $\varepsilon > 0$ (see Figure 1). Performing the corresponding contour integration, and letting $T \rightarrow \infty$ and $\varepsilon \rightarrow 0$, we infer that

$$\int_{-\infty}^{\infty} e^{-nt^2} \frac{dt}{t-p} = \pi i e^{-np^2} - \text{p.v.} \int_{-\infty}^{\infty} e^{-n(t+p)^2} \frac{dt}{t}.$$

We can rewrite the principle value integral as




FIG. 1. The integration contour used in the proof of Proposition III.1.

$$\begin{aligned}
 \text{p. v. } \int_{-\infty}^{\infty} e^{-n(t+p)^2} \frac{dt}{t} &= e^{-np^2} \int_{-\infty}^{\infty} e^{-nt^2} \sum_{k=0}^{\infty} \frac{(-2npt)^{2k+1}}{(2k+1)!} \frac{dt}{t} \\
 &= \frac{e^{-np^2}}{\sqrt{n}} \sum_{k=1}^{\infty} \frac{(-2\sqrt{np})^{2k+1}}{(2k+1)!} \Gamma(k+1/2) \\
 &= 2e^{-np^2} \sqrt{\pi} \sum_{k=0}^{\infty} \frac{(-\sqrt{np})^{2k+1}}{k!(2k+1)} = -\pi i e^{-np^2} \operatorname{erf}(i\sqrt{np})
 \end{aligned}$$

where we used the Legendre duplication formula $\Gamma(s)\Gamma(s+1/2) = 2^{1-2s}\sqrt{\pi}\Gamma(2s)$ for $s = k+1/2$, and the Taylor series expansion for the error function. Together with the half residue from before, this yields the complementary error function. \square

B. Conformal map

Let us consider $\tau = 0$. By (45) we know that $F''(s_0)$ is approximately equal to 1. Hence the steepest descent path going through s_0 is almost a vertical line segment in a neighborhood of $s = s_0$ (and $s = 1$). There is a conformal map transforming this steepest descent path to a real line segment, and this, after some rewriting, will allow the application of Proposition III.1. Then by a simple Taylor series expansion of $F(s) = (1+\Delta)s - \log s$, we have

$$F(s) = F(s_0) + \sum_{k=2}^{\infty} \frac{(-1)^k}{k} \left(\frac{s-s_0}{s_0} \right)^k.$$

The convergence radius of this series is given by $|s_0| = 1 + \mathcal{O}(n^{-\frac{1}{2}+\nu})$. For n large enough, we may assume that the convergence radius ρ is at least, say, $\frac{1}{2}$. Clearly then, we can find a conformal map $\phi : D(s_0; \rho) \rightarrow \mathbb{C}$ such that

$$F(s) - F(s_0) = -\phi(s)^2.$$

Here $D(s_0; \rho)$ denotes the disc with radius ρ , centered at s_0 . There are two such maps, but let us take the one with expansion

$$\phi(s) = -\frac{i}{\sqrt{2}} \frac{s-s_0}{s_0} + \frac{i}{3\sqrt{2}} \left(\frac{s-s_0}{s_0} \right)^2 + \mathcal{O}((s-s_0)^3). \quad (47)$$

Since the saddle point converges to 1, we want to find out how ϕ behaves there. To this end, we have the following lemma.

Lemma III.2 ($\tau = 0$). *Let ϕ be as defined above. Then we have as $n \rightarrow \infty$ that*

$$i\phi(1) = \frac{\Delta}{\sqrt{2}} - \frac{\Delta^2}{3\sqrt{2}} + \mathcal{O}(n^{-\frac{3}{2}+3\nu}). \quad (48)$$

Proof. From (49) and (44) we have

$$i\phi(1) = \frac{1}{\sqrt{2}}(s_0^{-1} - 1) - \frac{1}{3\sqrt{2}}(s_0^{-1} - 1)^2 + \mathcal{O}((s_0^{-1} - 1)^3) = \frac{\Delta}{\sqrt{2}} - \frac{\Delta^2}{3\sqrt{2}} + \mathcal{O}(n^{-\frac{3}{2}+3\nu}),$$

as $n \rightarrow \infty$, where we used that $s_0^{-1} = 1 + \Delta$. \square

Let us move to the case $0 < \tau < 1$, and derive an analogous result. As n becomes large, by (43), we have that

$$F''(a^{-1}) = 2\tau^{-2} \frac{|\sinh(\xi_\tau + i\eta)|^2}{\sinh(2\xi_\tau)} + \mathcal{O}(n^{-\frac{1}{2}+\nu})$$

as $n \rightarrow \infty$. This is close to being a positive real number, hence the steepest descent direction is almost a vertical line in a neighborhood of $s = a^{-1}$ (and $s = \tau$). Again, there is a conformal map transforming this steepest descent path to a real line segment. Explicitly, we can write

$$F(s) - F(a^{-1}) = \sum_{k=2}^{\infty} \left(\frac{(z_+ + z_-)^2}{2} \frac{(-1)^k}{(1 + a^{-1})^{k+1}} - \frac{(z_+ - z_-)^2}{2} \frac{1}{(1 - a^{-1})^{k+1}} + \frac{(-a)^k}{k} \right) (s - a^{-1})^k$$

The convergence radius of this series is given by $\min(\tau, 1 - \tau) + \mathcal{O}(n^{-\frac{1}{2}+\nu})$. In particular, for n large enough, we may assume that it has convergence radius at least $\rho := \frac{1}{2} \min(\tau, 1 - \tau)$, uniformly for $\hat{z} \in \partial\mathcal{E}_\tau$. Thus, for n large enough, we have

$$F(s) - F(a^{-1}) = -\phi(s)^2,$$

for some conformal map $\phi : D(a^{-1}, \rho) \rightarrow \mathbb{C}$, that depends on n . There are two such conformal maps, but we choose the one with expansion

$$\phi(s) = -i\sqrt{\frac{1}{2}F''(a^{-1})(s - a^{-1})} - \frac{i}{6\sqrt{2}} \frac{F'''(a^{-1})}{\sqrt{F''(a^{-1})}} (s - a^{-1})^2 + \mathcal{O}((s - a^{-1})^3). \quad (49)$$

By possibly taking ρ smaller, the reader may convince oneself that the length of the real line segment, corresponding to the inverse image of the steepest descent contour, can be arranged to be bounded, both from below and from above, by a positive constant, uniformly for n and $\hat{z} \in \partial\mathcal{E}_\tau$. In particular, we will consider a single line segment $[-r, r]$, with $r > 0$, for all $\hat{z} \in \partial\mathcal{E}_\tau$ and all n , for which $\phi^{-1}([-r, r]) \subset D(a^{-1}, \rho)$ lies on the steepest descent contour through $s = a^{-1}$.

Lemma III.3 ($0 < \tau < 1$). *We have*

$$i\phi(\tau) = \frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} (\Delta_+ + \Delta_-) - \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \frac{\Delta_+^2 - \Delta_+ \Delta_- + \Delta_-^2}{6} + \mathcal{O}(n^{-\frac{3}{2}+3\nu}), \quad (50)$$

as $n \rightarrow \infty$, uniformly for $\hat{z} \in \partial\mathcal{E}_\tau$.

Proof. Using (43) and (36), we infer that

$$\begin{aligned} \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \sqrt{\frac{1}{2}F''(a^{-1})(\tau - a^{-1})} &= \Delta_+ + \Delta_- + \frac{(\Delta_+ + \Delta_-)^2}{2} - \coth(\xi_\tau + i\eta) \frac{\Delta_+^2}{2} - \coth(\xi_\tau - i\eta) \frac{\Delta_-^2}{2} \\ &+ \frac{1}{2}(\Delta_+ + \Delta_-)(\coth(\xi_\tau + i\eta)\Delta_+ + \coth(\xi_\tau - i\eta)\Delta_- - \coth(2\xi_\tau)(\Delta_+ + \Delta_-)) + \mathcal{O}(n^{-\frac{3}{2}+3\nu}) \\ &= \Delta_+ + \Delta_- - \frac{\tau^2}{1 - \tau^2}(\Delta_+ + \Delta_-)^2 + \frac{\sinh(2\xi_\tau)}{|\sinh(\xi_\tau + i\eta)|^2} \frac{\Delta_+ \Delta_-}{2} + \mathcal{O}(n^{-\frac{3}{2}+3\nu}). \end{aligned} \quad (51)$$

To arrive at the last line, we used the identities

$$\begin{aligned} \coth(2\xi_\tau) - 1 &= \frac{2\tau^2}{1 - \tau^2}, \\ \coth(\xi_\tau + i\eta) + \coth(\xi_\tau - i\eta) &= \frac{\sinh(2\xi_\tau)}{|\sinh(\xi_\tau + i\eta)|^2}. \end{aligned}$$

Also, we have

$$\frac{i}{6\sqrt{2}} \frac{F'''(a^{-1})}{\sqrt{F''(a^{-1})}} (\tau - a^{-1})^2 = i\tau \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \left(\frac{(\operatorname{Re} \hat{z}_+)^2}{(1+\tau)^4} + \frac{(\operatorname{Im} \hat{z}_+)^2}{(1-\tau)^4} - \frac{1}{6\tau^3} \right) \tau^2 (\Delta_+ + \Delta_-)^2 + \mathcal{O}(n^{-\frac{3}{2}+3\nu}). \quad (52)$$

This we can simplify by noticing that

$$\frac{(\operatorname{Re} \hat{z}_+)^2}{(1+\tau)^4} + \frac{(\operatorname{Im} \hat{z}_+)^2}{(1-\tau)^4} = \frac{\cos^2 \eta}{2\tau(1+\tau)^2} + \frac{\sin^2 \eta}{2\tau(1-\tau)^2} = \frac{1+\tau^2-2\tau\cos 2\eta}{2\tau(1-\tau^2)^2} = \frac{|\sinh(\xi_\tau + i\eta)|^2}{\tau \sinh(2\xi_\tau)(1-\tau^2)}. \quad (53)$$

Combining (51) with (52) and (53), and the defining relation (49) for ϕ , we conclude that

$$\begin{aligned} i\phi(\tau) &= \frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} (\Delta_+ + \Delta_-) + \frac{1}{2} \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \Delta_+ \Delta_- \\ &+ \left(-\frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} \frac{\tau^2}{1-\tau^2} + \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \left(\frac{\tau^2 |\sinh(\xi_\tau + i\eta)|^2}{\sinh(2\xi_\tau)(1-\tau^2)} - \frac{1}{6} \right) \right) (\Delta_+ + \Delta_-)^2 + \mathcal{O}(n^{-\frac{3}{2}+3\nu}) \\ &= \frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} (\Delta_+ + \Delta_-) + \frac{1}{2} \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \Delta_+ \Delta_- - \frac{1}{6} \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} (\Delta_+^2 + 2\Delta_+ \Delta_- + \Delta_-^2) + \mathcal{O}(n^{-\frac{3}{2}+3\nu}), \end{aligned}$$

as $n \rightarrow \infty$, uniformly for $\hat{z} \in \partial \mathcal{E}_\tau$, and this yields the result. \square

C. Asymptotic behavior of $I_{n,\tau}^d$

In this section, we perform a steepest descent analysis to obtain the asymptotic behavior of $I_{n,\tau}^d$, both for $0 < \tau < 1$ and $\tau = 0$. Let us start with the latter, $\tau = 0$. Here, we simply deform γ_0 to the steepest descent path through $s = s_0$. This is allowed because the steepest descent path ends in $-\infty$ as $n \rightarrow \infty$, from both directions (to see this, one should solve the contour lines $\operatorname{Im}(s_0^{-1}s - \log s) = \arg s_0$). We may then write

$$\begin{aligned} I_{n,0}^d(z \cdot w) &= -\frac{1}{2\pi i} \oint_{\gamma_0} e^{nF(s)} \frac{1}{s-1} \left(1 - \frac{\phi'(s)(s-1)}{\phi(s) - \phi(\tau)} \right) ds - \frac{e^{nF(s_0)}}{2\pi i} \oint_{\gamma_0} \frac{e^{-n\phi(s)^2}}{s-1} \frac{\phi'(s)(s-1)}{\phi(s) - \phi(\tau)} ds \\ &= -\frac{1}{2\pi i} \oint_{\gamma_0} e^{nF(s)} \frac{1}{s-1} \left(1 - \frac{\phi'(s)(s-1)}{\phi(s) - \phi(\tau)} \right) ds - \frac{1}{2\pi i} e^{nF(s_0)} \int_{-\infty}^{\infty} \frac{e^{-ns^2}}{s - \phi(1)} ds. \end{aligned}$$

The first integral can be done with a standard steepest descent procedure, where we rescale variables locally around $s = s_0$ (with a factor $n^{-\frac{1}{2}+\nu}$). The reader may verify that the assumption that $0 < \nu < \frac{1}{6}$ is necessary for this procedure to work (the cubic term has to be small). We see that

$$\frac{1}{s-1} \left(1 - \frac{\phi'(s)(s-1)}{\phi(s) - \phi(1)} \right) = \frac{\phi(s) - \phi(1) + \phi'(s)(1-s)}{(s-1)(\phi(s) - \phi(1))} = -\frac{1}{2} \frac{\phi''(1)}{\phi'(1)} + \mathcal{O}(s-1) = \frac{1}{3} + \mathcal{O}(\Delta) + \mathcal{O}(s-1).$$

Performing the steepest descent analysis and invoking Proposition III.1, we get

$$I_{n,0}^d(z \cdot w) = \frac{1}{2} e^{nF(1)} \operatorname{erfc}(i\sqrt{n}\phi(1)) + \left(-\frac{1}{3} + \mathcal{O}(n^{-1+2\nu}) \right) \frac{e^{nF(s_0)}}{\sqrt{2\pi F''(s_0)n}}. \quad (54)$$

We thus get the following proposition.

Proposition III.4 ($\tau = 0$). *We have as $n \rightarrow \infty$ that*

$$e^{-nF(1)} I_{n,0}^d(1 + \Delta) = \frac{1}{2} \operatorname{erfc} \left(\sqrt{n} \frac{\Delta}{\sqrt{2}} \right) + \frac{1}{\sqrt{2\pi n}} \exp \left(-n \frac{\Delta^2}{2} \right) \left(\frac{n\Delta^2 - 1}{3} + \mathcal{O}(n^{-\frac{1}{2}+3\nu}) \right). \quad (55)$$

Proof. By the preceding, we have

$$e^{-nF(1)} I_{n,0}^d(1 + \Delta) = \frac{1}{2} \operatorname{erfc}(i\sqrt{n}\phi(1)) + \left(-\frac{1}{3} + \mathcal{O}(n^{-1+2\nu}) \right) \frac{e^{-n\phi(1)^2}}{\sqrt{2\pi F''(s_0)n}}. \quad (56)$$

By Taylor expanding and Proposition III.2, we have

$$\operatorname{erfc}(i\sqrt{n}\phi(1)) = \operatorname{erfc}\left(\sqrt{n}\frac{\Delta}{\sqrt{2}}\right) - \frac{2}{\sqrt{\pi}} \exp\left(-n\frac{\Delta^2}{2}\right) \left(-\sqrt{n}\frac{\Delta^2}{3\sqrt{2}} + \mathcal{O}(n^{-1+3\nu})\right), \quad (57)$$

and similarly

$$e^{-n\phi(1)^2} = \exp\left(-n\frac{\Delta^2}{2}\right) (1 + \mathcal{O}(n^{-\frac{1}{2}+3\nu})). \quad (58)$$

Plugging (57) and (58) in (56), and using that $F''(s_0) = s_0^2$ and $s_0^{-1} = 1 + \Delta$, we arrive at the result. \square

Now we turn to our integral (17) for the case $0 < \tau < 1$, where we deform γ_0 to the circle $|s| = |a|^{-1}$. As the reader may verify, similar to what we did for $\tau = 0$, we can rewrite our integral as

$$I_{n,\tau}^d(z_{\pm}) = -\frac{1}{2\pi i} \oint_{\gamma_0} \frac{e^{nF(s)}}{s-\tau} \left(\frac{1}{(1-s^2)^{\frac{d}{2}}} - \frac{1}{(1-\tau^2)^{\frac{d}{2}}} \right) ds - \frac{1}{2\pi i} \frac{1}{(1-\tau^2)^{\frac{d}{2}}} \int_{\phi(-r)}^{\phi(r)} \frac{e^{nF(s)}}{s-\tau} \left(1 - \frac{\phi'(s)(s-\tau)}{\phi(s)-\phi(\tau)} \right) ds - \frac{1}{2\pi i} \frac{e^{nF(a^{-1})}}{(1-\tau^2)^{\frac{d}{2}}} \int_{-r}^r \frac{e^{-ns^2}}{s-\phi(\tau)} ds + \mathcal{O}(e^{nF(a^{-1})} e^{-cn}) \quad (59)$$

uniformly for $\hat{z} \in \partial\mathcal{E}_\tau$ as $n \rightarrow \infty$, for some constant $c > 0$. The first two integrals on the RHS of (59) can be asymptotically solved with a standard steepest descent procedure, where we rescale variables locally around $s = a^{-1}$ (with a factor $n^{-\frac{1}{2}+\nu}$). Again, the reader may verify that the assumption that $0 < \nu < \frac{1}{6}$ is necessary for this procedure to work. We note that

$$\frac{1}{s-\tau} \left(1 - \frac{\phi'(s)(s-\tau)}{\phi(s)-\phi(\tau)} \right) = \frac{\phi(s)-\phi(\tau)+\phi'(s)(\tau-s)}{(s-\tau)(\phi(s)-\phi(\tau))} = -\frac{1}{2} \frac{\phi''(\tau)}{\phi'(\tau)} + \mathcal{O}(s-\tau).$$

Hence, executing the steepest descent method for the first two integrals, and applying Proposition III.1, we get

$$I_{n,\tau}^d(z_{\pm}) = \frac{1}{2} \frac{e^{nF(\tau)}}{(1-\tau^2)^{\frac{d}{2}}} \operatorname{erfc}(i\sqrt{n}\phi(\tau)) - \frac{1}{\sqrt{\pi n}} \frac{\tau \sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \frac{e^{nF(a^{-1})}}{(1-\tau^2)^{\frac{d}{2}}} \left(\frac{d\tau}{1-\tau^2} - \frac{1}{2} \frac{\phi''(\tau)}{\phi'(\tau)} + \mathcal{O}(n^{-\frac{1}{2}+3\nu}) \right).$$

We used here that it did not matter whether $\phi(\tau)$ was to the right or the left of the contour, by application of the residue theorem, since we started with a closed contour. In particular, we have

$$(1-\tau^2)^{\frac{d}{2}} e^{-nF(\tau)} I_{n,\tau}^d(z_{\pm}) = \frac{1}{2} \operatorname{erfc}(i\sqrt{n}\phi(\tau)) - \frac{1}{\sqrt{\pi n}} \frac{\tau \sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} e^{n\phi(\tau)^2} \left(\frac{d\tau}{1-\tau^2} - \frac{1}{2} \frac{\phi''(\tau)}{\phi'(\tau)} + \mathcal{O}(n^{-\frac{1}{2}+3\nu}) \right).$$

as $n \rightarrow \infty$, uniformly for $\hat{z} \in \partial\mathcal{E}_\tau$.

Summarizing, we have the following theorem.

Proposition III.5 ($0 < \tau < 1$). *Let $\hat{z} \in \partial\mathcal{E}_\tau$. Let p be as in Lemma III.3. Then we have*

$$(1-\tau^2)^{\frac{d}{2}} e^{-nF(\tau)} I_{n,\tau}^d(z_{\pm}) = \frac{1}{2} \operatorname{erfc}\left(\frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} \sqrt{n}(\Delta_+ + \Delta_-)\right) + \frac{1}{\sqrt{\pi n}} \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \exp\left(-\frac{|\sinh(\xi_\tau + i\eta)|^2}{\sinh 2\xi_\tau} n(\Delta_+ + \Delta_-)^2\right) \times \left(n \frac{\Delta_+^2 - \Delta_+ \Delta_- + \Delta_-^2}{3} - \frac{1}{6} \frac{\sinh 2\xi_\tau}{|\sinh(\xi_\tau + i\eta)|^2} - \tau^2 \frac{d-1}{1-\tau^2} + \mathcal{O}(n^{-\frac{1}{2}+3\nu}) \right), \quad (60)$$

as $n \rightarrow \infty$, uniformly for $\hat{z} \in \partial\mathcal{E}_\tau$.

Proof. This follows from the formulae above, and

$$\begin{aligned} \operatorname{erfc}(i\sqrt{n}\phi(\tau)) &= \operatorname{erfc}\left(\frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} \sqrt{n}(\Delta_+ + \Delta_-)\right) \\ &\quad - \frac{2}{\sqrt{\pi}} \exp\left(-\frac{|\sinh(\xi_\tau + i\eta)|^2}{\sinh 2\xi_\tau} n(\Delta_+ + \Delta_-)^2\right) \left(-\frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \sqrt{n} \frac{\Delta_+^2 - \Delta_+ \Delta_- + \Delta_-^2}{6} + \mathcal{O}(n^{-1+3\nu}) \right), \\ e^{n\phi(\tau)^2} &= \exp\left(-\frac{|\sinh(\xi_\tau + i\eta)|^2}{\sinh 2\xi_\tau} n(\Delta_+ + \Delta_-)^2\right) (1 + \mathcal{O}(n^{-\frac{1}{2}+3\nu})), \end{aligned}$$

which are obtained by taking a Taylor series and using Lemma III.3. Finally, with arguments similar to those in the proof of Lemma III.3, we have

$$\frac{d\tau}{1-\tau^2} - \frac{1}{2} \frac{\phi''(\tau)}{\phi'(\tau)} = \frac{(d-1)\tau}{1-\tau^2} + \frac{1}{6\tau} \frac{\sinh 2\xi_\tau}{|\sinh(\xi_\tau + i\eta)|^2} + \mathcal{O}(n^{-\frac{1}{2}+v}).$$

□

IV. EXPLICIT FORM OF RELEVANT EXPRESSIONS

To prove the main theorems we need to start plugging u, v and z into our formulae. Doing so, we can state Lemma III.2 and Lemma III.3 differently.

Lemma IV.1 ($\tau = 0$). Fix $0 < v < \frac{1}{6}$. For $z \in \mathbb{C}^d$ with $|z| = 1$, denote by \mathbf{n} the outward unit normal vector on the $2d$ -dimensional unit sphere ∂E_0^d in z . We have

$$i\sqrt{n}\phi(1) = \frac{u \cdot \mathbf{n} + \mathbf{n} \cdot v}{\sqrt{2}} + \frac{3u \cdot v - (u \cdot \mathbf{n} + \mathbf{n} \cdot v)^2}{3\sqrt{2n}} + \mathcal{O}(n^{-1+3v}), \quad (61)$$

as $n \rightarrow \infty$, uniformly for $|z| = 1$ and $u, v \in \mathbb{C}^d$ with $u, v = \mathcal{O}(n^v)$. Furthermore, if $u = v = \lambda \mathbf{n}$ with $\lambda \in \mathbb{R}$, then we have

$$i\sqrt{n}\phi(1) = \sqrt{2}\lambda - \frac{\lambda^2}{3\sqrt{2n}} + \mathcal{O}(n^{-1+3v}),$$

as $n \rightarrow \infty$, uniformly for $|z| = 1$ and $\lambda = \mathcal{O}(n^v)$.

Proof. It is well-known that the unit normal vector on the boundary of the $2d$ -dimensional unit ball in z is given simply by z . Substituting (46) in (48), we find that

$$\sqrt{2ni}\phi(1) = u \cdot z + z \cdot v + \frac{u \cdot v}{\sqrt{n}} - \frac{(u \cdot z + z \cdot v)^2}{3} + \mathcal{O}(n^{-1+3v}).$$

Identifying $\mathbf{n} = z$ in this equation yields the first part of the lemma. The second part follows trivially from the first part. □

With a little more effort, we find an analogous statement for the case $0 < \tau < 1$.

Lemma IV.2 ($0 < \tau < 1$). Fix $0 < v < \frac{1}{6}$. For $z \in \partial \mathcal{E}_\tau^d$, denote by \mathbf{n} the outward unit normal vector on $\partial \mathcal{E}_\tau^d$ in z . We have

$$i\sqrt{n}\phi(\tau) = \frac{u \cdot \mathbf{n} + \mathbf{n} \cdot v}{\sqrt{2}} + \frac{\mathcal{O}(|u|^2 + |v|^2)}{\sqrt{n}} + \mathcal{O}(n^{-1+3v}), \quad (62)$$

as $n \rightarrow \infty$, uniformly for $z \in \partial \mathcal{E}_\tau$ and $u, v \in \mathbb{C}^d$ with $u, v = \mathcal{O}(n^v)$. Furthermore, if $u = v = \lambda \mathbf{n}$ with $\lambda \in \mathbb{R}$, then we have

$$i\sqrt{n}\phi(\tau) = \sqrt{2}\lambda + \frac{1}{6} \left(\frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \right)^3 \frac{\lambda^2}{\sqrt{n}} + \mathcal{O}(n^{-1+3v}),$$

as $n \rightarrow \infty$, uniformly for $z \in \partial \mathcal{E}_\tau^d$ and $\lambda = \mathcal{O}(n^v)$.

Proof. First, we need to describe the outward unit normal vector \mathbf{n} . The ellipsoid $\partial \mathcal{E}_\tau^d$ can be written as a $(2d-1)$ -dimensional hypersurface $f(z) = 0$, where

$$f(z) = \frac{1-\tau}{1+\tau} |\operatorname{Re} z|^2 + \frac{1+\tau}{1-\tau} |\operatorname{Im} z|^2 - 1.$$

A normal vector is then given by the gradient of f , which, when normalized, gives

$$\mathbf{n} = \frac{(1-\tau)^2 \operatorname{Re} z + i(1+\tau)^2 \operatorname{Im} z}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}}. \quad (63)$$

This is the outward normal vector on $\partial \mathcal{E}_\tau^d$. Now let us investigate $\Delta_+ + \Delta_-$, as defined via (33). We notice that

$$\begin{aligned} \frac{1}{\sinh(\xi_\tau + i\eta)} + \frac{1}{\sinh(\xi_\tau - i\eta)} &= \frac{\sinh(\xi_\tau) \cos \eta}{|\sinh(\xi_\tau + i\eta)|^2} = \frac{1}{|\sinh(\xi_\tau + i\eta)|} \frac{(1-\tau)^2 |\operatorname{Re} z|}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}}, \\ \frac{1}{\sinh(\xi_\tau + i\eta)} - \frac{1}{\sinh(\xi_\tau - i\eta)} &= i \frac{\cosh(\xi_\tau) \sin \eta}{|\sinh(\xi_\tau + i\eta)|^2} = \frac{1}{|\sinh(\xi_\tau + i\eta)|} \frac{(1+\tau)^2 i |\operatorname{Im} z|}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}}. \end{aligned}$$

Hence we have

$$\begin{aligned} &\sqrt{2} \sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2} \frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} (\Delta_+ + \Delta_-) \\ &= (1-\tau)^2 |\operatorname{Re} z| \left(\sqrt{|\operatorname{Re} z|^2 + \frac{u+\bar{v}}{\sqrt{n}} \cdot (\operatorname{Re} z) + \frac{(u+\bar{v})^2}{4n}} - |\operatorname{Re}(z)| \right) \\ &\quad + (1+\tau)^2 |\operatorname{Im} z| \left(\sqrt{|\operatorname{Im} z|^2 + \frac{u-\bar{v}}{\sqrt{n}} \cdot (i\operatorname{Im} z) - \frac{(u-\bar{v})^2}{4n}} - |\operatorname{Im} z| \right). \end{aligned} \quad (64)$$

This, for $|\operatorname{Re} z|, |\operatorname{Im} z| > \frac{|u+v|}{\sqrt{n}}$, we can rewrite as

$$\begin{aligned} &\frac{(1-\tau)^2}{2} \frac{u+\bar{v}}{\sqrt{n}} \cdot (\operatorname{Re} z) + \frac{(1+\tau)^2}{2} \frac{u-\bar{v}}{\sqrt{n}} \cdot (i\operatorname{Im} z) \\ &+ \frac{(u+\bar{v})^2 - ((u+\bar{v}) \cdot \frac{\operatorname{Re} z}{|\operatorname{Re} z|})^2}{4n} \frac{|\operatorname{Re} z| + \frac{1}{2} \frac{u+\bar{v}}{\sqrt{n}} \cdot (\operatorname{Re} z)}{|\operatorname{Re} z| + \frac{1}{2} \frac{u+\bar{v}}{\sqrt{n}} \cdot (\operatorname{Re} z) + \sqrt{|\operatorname{Re} z|^2 + \frac{u+\bar{v}}{\sqrt{n}} \cdot (\operatorname{Re} z) + \frac{(u+\bar{v})^2}{4n}}} \\ &- \frac{(u-\bar{v})^2 - ((u-\bar{v}) \cdot \operatorname{Im} z)^2}{4n} \frac{|\operatorname{Im} z| + \frac{1}{2} \frac{u-\bar{v}}{\sqrt{n}} \cdot (i\operatorname{Im} z)}{|\operatorname{Im} z| + \frac{1}{2} \frac{u-\bar{v}}{\sqrt{n}} \cdot (i\operatorname{Im} z) + \sqrt{|\operatorname{Im} z|^2 + \frac{u-\bar{v}}{\sqrt{n}} \cdot (i\operatorname{Im} z) - \frac{(u-\bar{v})^2}{4n}}}. \end{aligned}$$

Since $\operatorname{Re} \sqrt{\zeta} \geq 0$ for all $\zeta \in \mathbb{C}$, we have for all $\zeta \in \mathbb{C} \cup \{\infty\}$ that

$$\left| \frac{1}{1 + \sqrt{\zeta}} \right| \leq 1.$$

We conclude that, when $|\operatorname{Re} z|, |\operatorname{Im} z| > \frac{|u+v|}{\sqrt{n}}$, we have

$$\left| \sqrt{2} \frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} (\Delta_+ + \Delta_-) - \frac{u \cdot \mathbf{n} + \mathbf{n} \cdot v}{2\sqrt{n}} \right| \leq \frac{1}{2n} \frac{|u+\bar{v}|^2 + |u-\bar{v}|^2}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}}.$$

The reader may check by explicit calculation that this estimate is still valid for $|\operatorname{Re} z| \leq \frac{|u+v|}{\sqrt{n}}$ or $|\operatorname{Im} z| \leq \frac{|u+v|}{\sqrt{n}}$. Plugging our estimate into (50), we arrive at the first statement of the lemma.

When $u = v = \lambda \mathbf{n}$, equation (64) simplifies considerably. We have

$$\begin{aligned} (u+\bar{v}) \cdot (\operatorname{Re} z) &= 2\lambda (\operatorname{Re} \mathbf{n}) \cdot (\operatorname{Re} z) = \frac{2\lambda (1-\tau)^2}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}} |\operatorname{Re} z|^2, \\ (u-\bar{v}) \cdot (i\operatorname{Im} z) &= 2\lambda (\operatorname{Im} \mathbf{n}) \cdot (\operatorname{Im} z) = \frac{2\lambda (1+\tau)^2}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}} |\operatorname{Im} z|^2, \end{aligned}$$

and

$$\begin{aligned} \frac{(u+\bar{v})^2}{4} &= \lambda^2 |\operatorname{Re} \mathbf{n}|^2 = \lambda^2 \frac{(1-\tau)^4}{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2} |\operatorname{Re} z|^2, \\ -\frac{(u-\bar{v})^2}{4} &= \lambda^2 |\operatorname{Im} \mathbf{n}|^2 = \lambda^2 \frac{(1+\tau)^4}{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2} |\operatorname{Im} z|^2. \end{aligned}$$

Hence we find that

$$\begin{aligned} & \sqrt{2} \frac{|\sinh(\xi_\tau + i\eta)|}{\sqrt{\sinh 2\xi_\tau}} (\Delta_+ + \Delta_-) \\ &= \frac{(1-\tau)^2 |\operatorname{Re} z|}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}} \left(|\operatorname{Re} z| \left(1 + \frac{\lambda(1-\tau)^2}{\sqrt{n} \sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}} \right) - |\operatorname{Re}(z)| \right) \\ &+ \frac{(1+\tau)^2 |\operatorname{Im} z|}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}} \left(|\operatorname{Im} z| \left(1 + \frac{\lambda(1+\tau)^2}{\sqrt{n} \sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}} \right) - |\operatorname{Im} z| \right) = \frac{\lambda}{\sqrt{n}}. \end{aligned}$$

In particular, we have

$$i\sqrt{n}\phi(\tau) = \sqrt{2}\lambda - \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \frac{\Delta_+^2 - \Delta_+ \Delta_- + \Delta_-^2}{6\sqrt{n}} + \mathcal{O}(n^{-1+3\nu}). \quad (65)$$

Similar to before, we can show that

$$\sqrt{n}\Delta_\pm = \frac{\sqrt{\sinh 2\xi_\tau}}{\sinh(\xi_\tau \pm i\eta)} \frac{(1-\tau)^2 |\operatorname{Re} z| \pm i(1+\tau)^2 |\operatorname{Im} z|}{\sqrt{(1-\tau)^4 |\operatorname{Re} z|^2 + (1+\tau)^4 |\operatorname{Im} z|^2}} \lambda = \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \lambda.$$

Plugging these into (65) finishes the proof. \square

Lemma IV.3. *Let $\tau = 0$. Then we have*

$$\exp\left(-\frac{|\sqrt{n}z+u|^2 + |\sqrt{n}z+v|^2}{2}\right) e^{nF(1)} = \exp\left(u \cdot v - \frac{|u|^2 + |v|^2}{2}\right). \quad (66)$$

Now let $0 < \tau < 1$. There exist unimodular functions $c_n : \partial \mathcal{E}_\tau^d \times \mathbb{C} \rightarrow \mathbb{T}$ such that

$$c_n(z, u) \overline{c_n(z, v)} \sqrt{\omega(\sqrt{n}z_+) \omega(\sqrt{n}z_-)} e^{nF(\tau)} = \exp\left(u \cdot v - \frac{|u|^2 + |v|^2}{2}\right). \quad (67)$$

Proof. Let us start with the case $\tau = 0$. We see that

$$|\sqrt{n}z+u|^2 + |\sqrt{n}z+v|^2 = 2n + |u|^2 + |v|^2 + 2\sqrt{n}(u+v) \cdot \bar{z}.$$

It readily follows from the definition of F for $\tau = 0$ that

$$F(1) = \left(z + \frac{u}{\sqrt{n}}\right) \cdot \left(z + \frac{v}{\sqrt{n}}\right) = 1 + \frac{u \cdot v}{n} + \frac{(u+v) \cdot \bar{z}}{\sqrt{n}}.$$

Combining these two identities, we obtain (66). Let move on to the case $0 < \tau < 1$. We know that

$$\begin{aligned} & \sqrt{\omega(\sqrt{n}z_+) \omega(\sqrt{n}z_-)} = \prod_{j=1}^d \sqrt{\omega(\sqrt{n}z_j + u_j) \omega(\sqrt{n}z_j + v_j)} \\ &= \exp\left(-\frac{1}{2} \sum_{j=1}^d (|\sqrt{n}z_j + u_j|^2 - \frac{\tau}{2}((\sqrt{n}z_j + u_j)^2 + \overline{(\sqrt{n}z_j + u_j)^2}) + |\sqrt{n}z_j + v_j|^2 - \frac{\tau}{2}((\sqrt{n}z_j + v_j)^2 + \overline{(\sqrt{n}z_j + v_j)^2}))\right). \quad (68) \end{aligned}$$

We notice that

$$\sum_{j=1}^d (|\sqrt{n}z_j + u_j|^2 - \frac{\tau}{2}((\sqrt{n}z_j + u_j)^2 + \overline{(\sqrt{n}z_j + u_j)^2})) = n(|z|^2 - \tau \operatorname{Re}(z^2)) + |u|^2 - \tau \operatorname{Re}(u^2) + \sqrt{n}(u \cdot (\bar{z} - \tau z) + (z - \tau \bar{z}) \cdot u), \quad (69)$$

and, similarly with u replaced by v . On the other hand, we have

$$\begin{aligned} F(\tau) &= \frac{\tau(z_+ + z_-)^2}{2(1+\tau)} - \frac{\tau(z_+ - z_-)^2}{2(1-\tau)} \\ &= \frac{1-\tau}{4} \sum_{j=1}^d \left(2\operatorname{Re} z_j + \frac{u_j + \bar{v}_j}{\sqrt{n}}\right)^2 - \frac{1+\tau}{4} \sum_{j=1}^d \left(2i\operatorname{Im} z_j + \frac{u_j - \bar{v}_j}{\sqrt{n}}\right)^2 \\ &= |z|^2 - \tau \operatorname{Re}(z^2) + \frac{u \cdot v - \frac{\tau}{2}(u^2 + v^2)}{n} + \frac{u \cdot (\bar{z} - \tau z) + (z - \tau \bar{z}) \cdot v}{\sqrt{n}}. \quad (70) \end{aligned}$$

Plugging the identities (69) (also for u replaced by v) and (70) into (68), we find that

$$\sqrt{\omega(\sqrt{n}z_+)\omega(\sqrt{n}z_-)}e^{nF(\tau)} = \exp\left(u \cdot v - \frac{|u|^2 + |v|^2}{2}\right) \exp(-i\tau \text{Im}(u^2 - v^2)) \exp(i\tau\sqrt{n} \text{Im}((z - \tau\bar{z}) \cdot (u - v))).$$

Clearly then, to get (67), we should define the unimodular functions by

$$c_n(z, u) = \exp(i\tau \text{Im}(u^2)) \exp(-i\tau\sqrt{n} \text{Im}((z - \tau\bar{z}) \cdot u)).$$

□

The following is more or less a direct consequence of Proposition II.3 and Lemma IV.3, we omit a proof. The result means that we find the higher dimensional analogue of the Ginibre kernel, i.e., a factorisation in d Ginibre kernels, as a scaling limit in the bulk. In [2], this was proved for $0 < \tau < 1$.

Corollary IV.4 ($\tau = 0$). *Take $|z| < 1$ and $u, v \in \mathbb{C}^d$. Then we have*

$$\lim_{n \rightarrow \infty} \mathcal{K}_n(\sqrt{n}z + u, \sqrt{n}z + v) = \frac{1}{\pi^d} \exp\left(u \cdot v - \frac{|u|^2 + |v|^2}{2}\right).$$

The convergence is uniform on compact sets of $|z| < 1$ and u, v .

V. PROOF OF THE MAIN THEOREMS

We now have all the ingredients necessary to prove the main results.

Proof of Theorem I.2. We first treat the case $\tau = 0$. Substituting (61) in (54), and using Taylor expansions in the same way as in the proof of Proposition III.4, we find that

$$e^{-nF(1)} I_{n,0}^d \left(\left(z + \frac{u}{\sqrt{n}} \right) \cdot \left(z + \frac{v}{\sqrt{n}} \right) \right) = \frac{1}{2} \text{erfc} \left(\frac{u \cdot \mathbf{n} + \mathbf{n} \cdot v}{\sqrt{2}} \right) + \frac{1}{\sqrt{2\pi n}} \exp \left(\frac{(u \cdot \mathbf{n} + \mathbf{n} \cdot v)^2}{2} \right) \left(\frac{(u \cdot \mathbf{n} + \mathbf{n} \cdot v)^2 - 3u \cdot v - 1}{3} + \mathcal{O}(n^{-\frac{1}{2}+3v}) \right), \quad (71)$$

as $n \rightarrow \infty$, uniformly for $|z| = 1$ and $u, v \in \mathbb{C}^d$ such that $u, v = \mathcal{O}(n^v)$. Multiplying by the remaining factors in (11), and using Lemma IV.3, we obtain the statement of Theorem I.2 for $\tau = 0$. Next, we treat the case $0 < \tau < 1$. Starting from the formula (15), we simply combine Lemma IV.2 and Lemma IV.3 with Proposition III.5 (which is valid with $\Delta_+^2 - \Delta_+ \Delta_- + \Delta_-^2$ replaced by $\mathcal{O}(|u|^2 + |v|^2)$). □

As the proof shows, we can be more precise about the error when $\tau = 0$. In the case of the elliptic Ginibre ensemble, i.e., the case $d = 1$ and $0 < \tau < 1$, we can also be more precise about the error. The following result (but slightly weaker) was proved by Byun and Ebke in [9]. We consider our novel derivation of this result of independent interest.

Proposition V.1. *Pick $0 < v < \frac{1}{6}$. Let \mathcal{K}_n be the kernel of the eigenvalues of the elliptic Ginibre ensemble with parameter $0 < \tau < 1$, as defined in (6). Let $z \in \partial \mathcal{E}_\tau$, and denote by \mathbf{n} and κ , respectively, the outward unit normal vector and the curvature, of $\partial \mathcal{E}_\tau$ in z . Then there exist continuous unimodular functions $c_n : \partial \mathcal{E}_\tau \times \mathbb{C} \rightarrow \mathbb{T}$ such that*

$$c_n(z, u) \overline{c_n(z, v)} \mathcal{K}_n(\sqrt{n}z + u\mathbf{n}, \sqrt{n}z + v\mathbf{n}) = \frac{1}{2\pi} \exp\left(u\bar{v} - \frac{|u|^2 + |v|^2}{2}\right) \text{erfc}\left(\frac{u + \bar{v}}{\sqrt{2}}\right) + \frac{\kappa}{\sqrt{n}} \exp\left(-\frac{|u|^2 + u^2 + |v|^2 + \bar{v}^2}{2}\right) \left(\frac{u^2 + \bar{v}^2 - u\bar{v} - 1}{3\sqrt{2\pi^3}} + \mathcal{O}(n^{-\frac{1}{2}+3v})\right), \quad (72)$$

as $n \rightarrow \infty$, uniformly for $z \in \partial \mathcal{E}_\tau$, and $u, v \in \mathbb{C}$ such that $u, v = \mathcal{O}(n^v)$.

Proof. In this case we have $z_+ = \sqrt{\sinh(2\xi_\tau)}(z + \frac{u}{\sqrt{n}}\mathbf{n})$ and $z_- = \sqrt{\sinh(2\xi_\tau)}(z + \frac{v}{\sqrt{n}}\mathbf{n})$. The outward unit normal vector is now given simply by

$$\mathbf{n} = \frac{\sinh(\xi_\tau + i\eta)}{|\sinh(\xi_\tau + i\eta)|}.$$

Hence, we have

$$\Delta_+ = \frac{1}{\sqrt{n}} \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \frac{u}{\sqrt{2}}, \quad \text{and} \quad \Delta_- = \frac{1}{\sqrt{n}} \frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \frac{\bar{v}}{\sqrt{2}}.$$

Plugging these in (60) yields

$$(1 - \tau^2)^{\frac{d}{2}} e^{-nF(\tau)} J_{n,\tau}^d(z_\pm) = \frac{1}{2} \operatorname{erfc} \left(\frac{u + \bar{v}}{\sqrt{2}} \right) + \frac{1}{6\sqrt{\pi n}} \left(\frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \right)^3 \exp \left(-\frac{(u + \bar{v})^2}{2} \right) \left(u^2 - u\bar{v} + \bar{v}^2 - 1 + \mathcal{O}(n^{-\frac{1}{2}+3v}) \right), \quad (73)$$

as $n \rightarrow \infty$, uniformly for $z \in \partial\mathcal{E}_\tau$ and $u, v = \mathcal{O}(n^v)$. As shown in [23], the factor in the second line of (73) can be identified with the curvature κ of $\partial\mathcal{E}_\tau$ in z (up to an explicit constant factor). In particular, according to [23, C13] we have

$$\kappa = \frac{(1 - \tau^2)^{\frac{3}{2}}}{(1 + \tau^2 - 2\tau \cos 2\eta)^{\frac{3}{2}}} = \left(\frac{2\tau \sinh 2\xi_\tau}{4\tau |\sinh(\xi_\tau + i\eta)|^2} \right)^{\frac{3}{2}} = \frac{1}{2\sqrt{2}} \left(\frac{\sqrt{\sinh 2\xi_\tau}}{|\sinh(\xi_\tau + i\eta)|} \right)^3. \quad (74)$$

Finally, reinstating the weight factors as in (15) and applying Lemma IV.3, we obtain the result. \square

Proof of Theorem I.1. As the reader may verify with some straightforward combinatorial arguments, the number of points of the DPP defined via (10) is given by the binomial coefficient $\binom{n+d-1}{d}$, which behaves as $\frac{n^d}{d!} (1 + \mathcal{O}(1/n))$ for large n . For $\tau = 0$, the result is more or less a direct consequence of Theorem I.2, where, according to (71), we are allowed to replace $\mathcal{O}(1 + |u|^2 + |v|^2)$ by

$$\frac{(u \cdot \mathbf{n} + \mathbf{n} \cdot v)^2 - 3u \cdot v - 1}{3} + \mathcal{O}(n^{-\frac{1}{2}+3v}).$$

Taking $u = v = \lambda \mathbf{n}$, we find the result (note that κ reduces to the value 1 here). Let us move to the case $0 < \tau < 1$. Note that the unimodular factors from Lemma IV.3 cancel each other when $u = v$. We can rewrite (74) as

$$\kappa = \left(\frac{\sqrt{\sinh 2\xi_\tau}}{|z_+^2 - 2|^{\frac{1}{2}}} \right)^3 = \frac{1}{(|\operatorname{Re} z| + i|\operatorname{Im} z|)^2 - \frac{4\tau}{1-\tau^2}}^{\frac{3}{2}} = \frac{1}{\left((|\operatorname{Re} z|^2 - |\operatorname{Im} z|^2 - \frac{4\tau}{1-\tau^2})^2 + 4|\operatorname{Re} z|^2 |\operatorname{Im} z|^2 \right)^{\frac{3}{4}}}.$$

The statement is now a direct consequence of Proposition III.5, and (the second part of) Lemma IV.2. \square

-
- [1] G. Akemann and M. Bender, "Interpolation between Airy and Poisson statistics for unitary chiral non-Hermitian random matrix ensembles." J. Math. Phys. 51 (2010) 103524.
- [2] G. Akemann, M. Duits, and L. D. Molag, "The Elliptic Ginibre Ensemble: A Unifying Approach to Local and Global Statistics for Higher Dimensions". arXiv preprint: [arXiv:2203.00287](https://arxiv.org/abs/2203.00287).
- [3] Y. Ameur and S. Byun, "Almost-Hermitian random matrices and bandlimited point processes." Preprint [arXiv:2101.03832](https://arxiv.org/abs/2101.03832) (2021).
- [4] Y. Ameur, H. Hedenmalm and N. Makarov, "Fluctuations of eigenvalues of random normal matrices." Duke Math. J., 159 (2011), 31-81.
- [5] Y. Ameur, N.-G. Kang, and N. Makarov, "Rescaling Ward identities in the random normal matrix model". Constr Approx 50, 63-127 (2019).
- [6] R. Bergman, "Bergman kernels for weighted polynomials and weighted equilibrium measures of \mathbb{C}^d ." Indiana University Mathematics Journal Vol. 58, No. 4 (2009), pp. 1921-1946.
- [7] M. Bender, "Edge scaling limits for a family of non-Hermitian random matrix ensembles". Probab. Theory Relat Fields, 147 (2010) 241.
- [8] T. Bothner and A. Little, "The complex elliptic Ginibre ensemble at weak non-Hermiticity." Preprint [arXiv:2208.04684](https://arxiv.org/abs/2208.04684) (2022).
- [9] S. Byun and M. Ebke, "Universal scaling limits of the symplectic elliptic Ginibre ensemble." Preprint [arXiv:2108.05541](https://arxiv.org/abs/2108.05541) (2021).
- [10] D. S. Dean, P. Le Doussal, S. N. Majumdar, and G. Schehr, "Noninteracting fermions in a trap and random matrix theory". J. Phys. A: Math. Theor. 52 (2019) 144006.
- [11] P. Deift and X. Zhou, "A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the MKdV equation" Ann. Math. 137 (1993), 295-368.
- [12] V. N. Faddeyeva and N. M. Terent'ev, "Tables of Values of the Function $w(z) = e^{-z^2} (1 + 2i\pi^{-1/2} \int_0^z e^{t^2} dt)$ for Complex Argument." Mathematical Tables Series, 11. Pergamon Press, Oxford-London-New York-Paris, 1961.

- [13] P. J. Forrester and G. Honner, “*Exact statistical properties of the zeros of complex random polynomials.*” J. Phys. A: Math. Gen. 32 (1999), 2961.
- [14] A. S. Fokas, A. R. Its and A. V. Kitaev, “*The isomonodromy approach to matrix models in 2d quantum gravity.*” Comm. Math. Phys., 142 (1991), 313-344.
- [15] Y. V. Fyodorov, B. A. Khoruzhenko, and H.-J. Sommers. “*Almost Hermitian random matrices: crossover from Wigner-Dyson to Ginibre eigenvalue statistics*”. Phys. Rev. Lett. 79 (1997) 557.
- [16] Y. V. Fyodorov, B. A. Khoruzhenko, and H.-J. Sommers. “*Almost-Hermitian random matrices: eigenvalue density in the complex plane*”. Phys. Lett. A 226 (1997) 46.
- [17] Y. V. Fyodorov, B. A. Khoruzhenko, and H.-J. Sommers. “*Universality in the random matrix spectra in the regime of weak non-Hermiticity*”. Ann. Inst. H. Poincaré Phys. Théor. 68 (1998) 449.
- [18] J. Ginibre, “*Statistical ensembles of complex, quaternion, and real matrices*”. J. Math. Phys. 6 (1965) 440.
- [19] V. L. Girko, “*Elliptic law*”. Theory Probab. Appl. 30 (1986) 677.
- [20] H. Hedenmalm and A. Wennman, “*Planar orthogonal polynomials and boundary universality in the random normal matrix model*”. Acta Math., 227 (2021), 309-406.
- [21] S. Kolodziej, “*The complex Monge-Ampere equation and pluripotential theory.*” Mem. Amer. Math. So. 178 (2005), no. 840.
- [22] B. Lacroix-A-Chéz-Toine, S. N. Majumdar, and G. Schehr, “*Rotating trapped fermions in two dimensions and the complex Ginibre ensemble: Exact results for the entanglement entropy and number variance*”. Phys. Rev. A 99 (2019) 021602(R).
- [23] S. Lee and R. Riser, “*Fine asymptotic behavior for eigenvalues of random normal matrices: Ellipse case*”. J. Math. Phys. 57 (2016) 023302.
- [24] R. Riser, “*Universality in Gaussian random normal matrices*”. PhD Thesis, ETH Zurich, [arXiv:1312.0068](https://arxiv.org/abs/1312.0068), 2013.
- [25] J. Ross and M. Singer, “*Asymptotics of partial density functions for divisors.*” J. Geom. Anal., 27 (2017), 1803-1854.
- [26] T. Tao, V. Vu, “*Random matrices: universality of local spectral statistics of non-Hermitian matrices*”. Ann. Prob. 43 (2015) 782.
- [27] S. Zelditch and P. Zhou, “*Central limit theorem for spectral partial Bergman kernels.*” Geom. Topol., 23 (2004), 1961-2004.