NEARLY PARALLEL G₂-MANIFOLDS: FORMALITY AND ASSOCIATIVE SUBMANIFOLDS

MARISA FERNÁNDEZ, ANNA FINO, ALEXEI KOVALEV, AND VICENTE MUÑOZ

ABSTRACT. We construct new examples of non-formal simply connected compact Sasaki–Einstein 7-manifolds. We determine the minimal model of the total space of any fibre bundle over \mathbb{CP}^2 with fibre $S^1 \times S^2$ or S^3/\mathbb{Z}_p (p > 0), and we apply this to conclude that the Aloff–Wallach spaces are formal. We also find examples of formal manifolds and non-formal manifolds, which are locally conformal parallel Spin(7)-manifolds.

On the other hand, we construct associative minimal submanifolds in the Aloff– Wallach spaces and in any regular Sasaki–Einstein 7-manifold; in particular, in the space $Q(1,1,1) = (SU(2) \times SU(2) \times SU(2))/(U(1) \times U(1))$ with the natural S^{1} family of nearly parallel G₂-structures induced by the Sasaki–Einstein structure. In each of those cases, we obtain a family of non-trivial associative deformations.

1. INTRODUCTION

A nearly parallel G₂-structure on a 7-manifold P is given by a positive 3-form φ satisfying the differential equation $d\varphi = \tau_0 \star \varphi$, for some non-zero constant τ_0 , where \star is the Hodge star operator determined by the induced metric by φ . Nearly parallel G₂-manifolds were introduced as manifolds with weak holonomy G₂ by Gray in [33].

In [9], it was shown that there is a 1-1 correspondence between nearly parallel G₂manifolds and Killing spinors. As a consequence, the metric g induced by the 3-form φ has to be Einstein with positive scalar curvature $\operatorname{scal}(g) = \frac{21}{8}\tau_0^2$. Thus, if (P,g) is complete, then P is a compact manifold with finite fundamental group due to Myers' theorem.

Another equivalent description of nearly parallel G₂-structures is in terms of the metric cone ($\hat{P} = P \times \mathbb{R}^+, g^c = r^2g + dr^2$), which has to have holonomy contained in Spin(7), viewed as subgroup of SO(8). If (P, g) is simply connected and not isometric to the standard sphere, then there are three possible cases: the holonomy of (\hat{P}, g^c) is Sp(2) or, equivalently, (P, g) is a 3-Sasakian manifold; the holonomy of (\hat{P}, g^c) can be SU(4) if and only if (P, g) is a Sasaki–Einstein manifold; or the holonomy of (\hat{P}, g^c)

Date: April 19, 2023.

²⁰¹⁰ Mathematics Subject Classification. 53C25, 53C30, 53B25, 53C40.

Key words and phrases. nearly parallel G₂-structures, formality, associative 3-folds, mapping tori.

coincides with Spin(7), in which case the G₂-structure is called *proper*. We recall that these three cases correspond to the existence of a space of Killing spinors of dimension 3, 2 or 1, respectively (see [29]). In particular, by [1], a Sasaki–Einstein 7-manifold has a canonical S^1 -family of nearly parallel G₂-structures inducing the Sasaki–Einstein metric. Moreover, by [30], a 3-Sasakian 7-manifold carries a second Einstein metric induced by a proper nearly parallel G₂-structure (see Section 2 for details).

A celebrated result of Deligne, Griffiths, Morgan and Sullivan states that any compact Kähler manifold is formal [21]. In the same spirit, formality of a manifold is related to the existence of suitable geometric structures on the manifold (see Section 3 for details about formality).

Nearly parallel G₂-manifolds are in various ways rather similar to nearly Kähler 6manifolds. Such a manifold is also Einstein, it has a Killing spinor, and its metric cone has holonomy contained in G₂. Simply connected compact manifolds of dimension ≤ 6 are always formal [49]. Even more, any simply connected compact nearly Kähler manifold is formal by [3]. In contrast, an example of a non-formal simply connected compact nearly parallel G₂-manifold is given by the simply connected compact Sasaki– Einstein manifold $Q(1, 1, 1) = (SU(2) \times SU(2) \times SU(2))/(U(1) \times U(1))$ (see [13, Theorem 3.2]). The space Q(1, 1, 1) is the only example, known in the literature, of a nearly parallel G₂-manifold satisfying all those properties.

The goal of this paper is two-fold. Our first purpose is to study the formality of nearly parallel G_2 -manifolds. For simply connected compact 3-Sasakian 7-manifolds, that study was already previously considered in [25], so that we focus on 7-dimensional Sasaki–Einstein manifolds and proper nearly parallel G_2 -manifolds.

In Section 4 (Theorem 4.2), we construct new examples of non-formal simply connected compact Sasaki–Einstein 7-manifolds. Such a manifold S_k ($3 \le k \le 8$) is obtained as the total space of a principal circle bundle over the Sasaki–Einstein manifold $X_k = P_k \times S^2$, where $P_k = \mathbb{CP}^2 \# k \mathbb{CP}^2$ is a del Pezzo surface. We prove that the manifold S_k is non-formal because it has a non-zero triple Massey product.

In Section 5, we study the formality of the Berger space and of Aloff-Wallach spaces considering both as fibrations. More precisely, in Theorem 5.1 we find the minimal model of the total space of any principal S^3 -bundle over S^4 and show that such a space is formal. We use this result to prove that the Berger space and S^7 have the same minimal model, and so the Berger space is formal. For the Aloff-Wallach spaces $W_{k,l}$, by [34], we know that they are the total space of a F-fiber bundle over \mathbb{CP}^2 , where $F = S^3/\mathbb{Z}_p$ with p > 0 if $p = k + l \neq 0$, or $F = S^1 \times S^2$ if $k + l \neq 0$. We determine the minimal model of the total space of such a fiber bundle, and we prove that such a space is formal. Moreover, we show that $W_{k,l}$ and the product manifold $S^2 \times S^5$ have the same minimal model (Theorem 5.2 and Theorem 5.3). The formality of the Berger space and Aloff-Wallach spaces was previously proved by a different method [42, 43]. In fact, in [42] it is proved that the Berger space is formal because it is geometrically formal (that is, it has a Riemannian metric such that all wedge products of harmonic forms are harmonic). In [43], it is proved that the spaces $W_{k,l}$ are formal since they are homogeneous spaces of Cartan type.

On the other hand, if (P, φ) is a nearly parallel G_2 -manifold, then the torsion-free Spin(7)-structure on the metric cone (\hat{P}, g^c) gives rise to a canonical closed 4-form which defines a calibration on \hat{P} . A 3-dimensional submanifold Y in (P, φ) is called *associative* if $\varphi|_Y = \operatorname{vol}_Y$ or, equivalently, if its cone \hat{Y} in \hat{P} is calibrated by the 4-form (see [40, Lemma 2.10]). By [5], associative submanifolds in nearly parallel G₂-manifolds P are minimal and their infinitesimal deformations were considered by Kawai in [40].

Associative 3-folds have been studied by Lotay [46] when P is the standard 7-sphere, by Kawai [39] when P is the squashed 7-sphere, and by Ball and Madnick when Pis the Berger space [6], the squashed 7-sphere and the squashed exceptional Aloff– Wallach space $W_{1,1}$ [7]. However, nothing seems to be known for other nearly-parallel G₂-manifolds.

Noting the above, the second goal of the paper is to provide constructions of associative submanifolds in a regular Sasaki–Einstein manifold and in the Aloff–Wallach spaces. In Section 7, we study such submanifolds in the case of a regular Sasaki– Einstein manifold (Propositions 7.1 and Theorem 7.4). In particular, we consider the case of the compact Sasaki–Einstein space Q(1, 1, 1), which is the total space of a principal S^1 -bundle over $S^2 \times S^2 \times S^2$. We first make explicit the canonical S^1 -family φ_t of nearly parallel G₂-structures associated to the Sasaki–Einstein structure on Q(1, 1, 1) (see Section 6). Then, considering the Calabi–Yau cone over Q(1, 1, 1), we construct minimal associative 3-folds in Q(1, 1, 1), and we determine a family of nontrivial associative deformations (Propositions 7.5 and 7.6). We also study minimal associative 3-folds in the Aloff–Wallach spaces (Theorem 8.1).

Given a nearly parallel G_2 -manifold P, the product manifold $P \times S^1$ and, more generally, the mapping torus of P by a diffeomorphism preserving the nearly parallel G_2 -structure, carries a natural locally conformal parallel Spin(7)-structure. In Section 9, we use the mapping torus construction to the nearly-parallel G_2 -manifolds $(Q(1,1,1), \varphi_t)$, the Berger space and the Aloff–Wallach spaces, and we construct examples of formal and non-formal locally conformal parallel Spin(7)-manifolds (Propositions 9.2 – Proposition 9.5).

The paper is organized as follows. In Section 2 and Section 3, we briefly review properties on nearly parallel G₂-structures and minimal models, respectively, that we need for our results. Then, in Section 4 and Section 5, we examine the formality of nearly parallel G₂-manifolds. In Section 6, we apply the construction of the canonical S^1 -family of nearly parallel G₂-structures to the Sasaki–Einstein manifold Q(1, 1, 1). Section 7 is devoted to the construction of minimal associative 3-folds in 7-dimensional regular Sasaki–Einstein manifolds and, in particular, in Q(1, 1, 1). Deformations of minimal associative 3-folds in any 7-dimensional regular Sasaki–Einstein manifold are also considered in Section 7. In Section 8 we study the associative 3-dimensional submanifolds of the Aloff–Wallach spaces. Finally, Section 9 contains the construction of locally conformal parallel Spin(7)-manifolds as mapping tori of each of the following spaces: $(Q(1, 1, 1), \varphi_t)$, the Berger space and the Aloff–Wallach spaces.

2. Nearly parallel G₂-structures

In this section, we recall the canonical S^1 -family of nearly parallel G₂-structures that exists on any Sasaki–Einstein 7-manifold [1, 15, 28], and the proper nearly parallel G₂-structure on any 3-Sasakian 7-manifold [29, 30] (see [17] for more details).

A 7-manifold P has a G₂-structure if there is a reduction of the structure group of its frame bundle from $GL(7, \mathbb{R})$ to the exceptional Lie group G₂. By [32], a 7-manifold P carries G₂-structures if and only if P is orientable and spin.

The presence of a G₂-structure is equivalent to the existence of a differential 3-form φ on P, which can be described locally as

$$\varphi = e^{127} + e^{347} + e^{567} + e^{135} - e^{146} - e^{236} - e^{245},$$

with respect to some basis $\{e^1, \ldots, e^7\}$ of the (local) 1-forms on P. Here, e^{127} stands for $e^1 \wedge e^2 \wedge e^7$, and so on.

Since $G_2 \subset SO(7)$, a G_2 -structure φ on P determines a Riemannian metric $g = g_{\varphi}$ and an orientation on P such that

$$g(U,V) \text{ vol} = \frac{1}{6} (U \lrcorner \varphi) \land (V \lrcorner \varphi) \land \varphi, \qquad (2.1)$$

for any vector fields U, V on P, where vol is the volume form on P.

A G₂-structure φ on a 7-manifold P is said to be *nearly parallel* if there exists a non-zero real number τ_0 such that

$$d\varphi = \tau_0 \star \varphi, \tag{2.2}$$

where \star is the Hodge star operator determined by the metric g_{φ} and the volume form on P induced by φ . A 7-manifold P with a nearly parallel G₂-structure is called a *nearly parallel* G₂-manifold.

Sasaki–Einstein and 3-Sasakian manifolds. Let us recall that a Riemannian manifold (S, g) of dimension 2n+1 is called *Sasakian* if its cone $(S \times \mathbb{R}^+, g^c = r^2g + dr^2)$ is Kähler, that is the cone metric g^c admits a compatible integrable almost complex structure J so that $(S \times \mathbb{R}^+, g^c, J)$ is Kähler. In this case the Reeb vector field $\xi = J\partial_r$ on S is a unit Killing vector field. The corresponding 1-form η defined by

 $\eta(U) = g(\xi, U)$, for any vector field U on S, is a *contact form*, meaning $\eta \wedge (d\eta)^n \neq 0$ at every point of S. The Kähler form on the cone can be expressed as

$$\omega^c = \frac{1}{2}d(r^2\eta),\tag{2.3}$$

where we extended η (and ξ) to $S \times \mathbb{R}^+$ by $\eta(U) = \frac{1}{r^2}g^c(J(r\partial_r), U)$. Let ∇ be the Levi–Civita connection of g. The endomorphism Φ of the tangent bundle TS of S, given by $\Phi U = \nabla_U \xi$, satisfies the following identities

$$\Phi^{2} = -\operatorname{Id} + \eta \otimes \xi, \ g(\Phi U, \Phi V) = g(U, V) - \eta(U)\eta(V), \ d\eta(U, V) = 2g(\Phi U, V), \ (2.4)$$

for any vector fields U, V on S. If the integral curves of the Reeb vector field ξ are closed, hence circles, then ξ integrates to a locally free isometric action of S^1 on (S, g). A Sasakian manifold is called *regular* when this latter S^1 -action is free.

A Sasakian manifold (S, g) of dimension 2n + 1 is said to be *Sasaki–Einstein* if the cone metric $g^c = r^2g + dr^2$ on $S \times \mathbb{R}^+$ is Kähler and Ricci-flat or, equivalently, the restricted holonomy group of g^c is contained in SU(n + 1). Thus, the Sasakian metric g is Einstein with Einstein constant 2n.

Compact regular Sasaki–Einstein manifolds are principal circle bundles over compact Kähler–Einstein manifolds with positive scalar curvature, and whose Kähler form defines a cohomology class proportional to an integral cohomology class [16, 28, 41] (therefore, the base manifold is a simply connected projective algebraic variety, more precisely a Fano variety). The converse also holds. In fact, the following result is proved in [16, Theorem 2.8]. Let (X, g_X) be a compact Kähler–Einstein manifold with positive scalar curvature, and whose Kähler form ω_X defines an integral cohomology class, and let S be the total space of the circle bundle

$$S^1 \hookrightarrow S \xrightarrow{\pi} X$$
 (2.5)

with Euler class $[\omega_X] \in H^2(X, \mathbb{Z})$. Then, S with the metric $g = \pi^* g_X + \eta \otimes \eta$ is a (regular) Sasaki–Einstein manifold, whose contact form η satisfies the equation $d\eta = 2\pi^* \omega_X$, where π is the projection in (2.5).

Let (S, g) be a Sasaki–Einstein manifold of dimension 7 with contact form η . Then, according to [1, pp. 723–724], S has an S¹-family of nearly parallel G₂-structures φ_t , which are given by

$$\varphi_t = \Omega \wedge \eta + \cos t \,\Psi_+ + \sin t \,\Psi_-. \tag{2.6}$$

Here Ω is the horizontal Kähler form related to the Ricci-flat Kähler form ω^c on the cone $S \times \mathbb{R}^+$ via $\Omega \wedge \eta = \frac{1}{2} \partial_r \lrcorner (\omega^c \wedge \omega^c)|_{r=1}$, equivalently $\Omega = \pi^* \omega_X$ in the case when (S,g) is regular and $\pi \colon S \to X$ is the projection of the principal circle bundle. Further, $\Psi = \Psi_+ + i\Psi_- = \partial_r \lrcorner \widehat{\Psi}|_{r=1}$ is a horizontal complex volume form, where $\widehat{\Psi}$ denotes a holomorphic 4-form of unit length on $S \times \mathbb{R}^+$. Now, a straightforward computation using (2.1) shows that every φ_t induces the Sasaki–Einstein metric g on S. **Remark 2.1.** Note that the expression for φ_t given in [1] (where the authors write σ_t instead of φ_t) is

$$\varphi_t = -\Omega \wedge \eta + \cos t \, \Psi_+ + \sin t \, \Psi_-,$$

with η the contact form of the Sasakian structure on S. The change that we made of the first term on the right-hand side of (2.6) is due to the following. By (2.4), we have $d\eta(U, V) = 2g(\Phi U, V)$, while in [1] the authors consider $d\eta(U, V) = 2g(U, \Phi V)$, for all vector fields U, V on S.

Let us recall that a 3-Sasakian structure is a collection of three Sasakian structures $(\phi_i, \xi_i, \eta_i, g)$ $(i \in \{1, 2, 3\})$ on a (4n + 3)-dimensional Riemannian manifold satisfying quaternionic-like identities. More precisely, a Riemannian manifold (S, g) of dimension 4n + 3 is called 3-Sasakian if its cone $(S \times \mathbb{R}^+, g^c = r^2g + dr^2)$ is hyperkähler, or equivalently, the holonomy group of the cone metric g^c is a subgroup of $\operatorname{Sp}(n + 1)$. In this case, the Reeb vector fields $\xi_i = J_i \partial_t$ (i = 1, 2, 3) satisfy the relations: $g(\xi_i, \xi_j) = \delta_{ij}$ and $[\xi_i, \xi_j] = 2\xi_k$ for (i, j, k) a cyclic permutation of (1, 2, 3).

If (S, g) is a 7-dimensional 3-Sasakian manifold, then S carries a second nearly parallel G₂-structure whose underlying Einstein metric is such that its cone metric has holonomy equal to Spin(7) (see [29, 30, 48]). This second Einstein metric \tilde{g} on S is given by

$$\widetilde{g} = \frac{1}{\sqrt{5}} g|_{\mathcal{H}} + g|_{\mathcal{V}}, \qquad (2.7)$$

where \mathcal{V} is the 3-dimensional distribution $\mathcal{V} = \text{span}\{\xi_1, \xi_2, \xi_3\}$, and $\mathcal{H} = \mathcal{V}^{\perp}$. The explicit expression of the second nearly parallel G₂-structure inducing the metric \tilde{g} is given in [30, Prop. 2.4] (see also [48]).

3. MINIMAL MODELS AND FORMALITY

In order to analyze the property of formality of nearly parallel G_2 -manifolds (Section 4 and Section 5) and of 8-manifolds with a locally conformal parallel Spin(7)-structure (see Section 9), we review here concepts about minimal models and formality (see [21, 23, 26] for more details).

A differential graded algebra (or DGA) over the real numbers \mathbb{R} , is a pair (\mathcal{A}, d) consisting of a graded commutative algebra $\mathcal{A} = \bigoplus_{k \ge 0} \mathcal{A}^k$ over \mathbb{R} (that is, it is positively graded), and a differential d of degree 1, satisfying the Leibnitz rule $d(a \cdot b) = (da) \cdot b + (-1)^{|a|} a \cdot (db)$, where |a| is the degree of a.

The cohomology $H^*(\mathcal{A})$ of a differential graded algebra (\mathcal{A}, d) is naturally a DGA with the product inherited from that on \mathcal{A} and with the differential being identically zero. The DGA (\mathcal{A}, d) is *connected* if $H^0(\mathcal{A}) = \mathbb{R}$, and \mathcal{A} is 1-connected if, in addition, $H^1(\mathcal{A}) = 0$. Henceforth we shall assume that all our DGAs are connected. In our context, the main example of DGA is the de Rham complex $(\Omega^*(M), d)$ of a connected differentiable manifold M, where d is the exterior differential.

Morphisms between DGAs are required to preserve the degree and to commute with the differential. A morphism $f : (\mathcal{A}, d) \to (\mathcal{B}, d)$ is a *quasi-isomorphism* if the map induced in cohomology $f^* : H^*(\mathcal{A}, d) \to H^*(\mathcal{B}, d)$ is an isomorphism.

A DGA (\mathcal{M}, d) is said to be *minimal* if

- (1) \mathcal{M} is free as an algebra, that is, \mathcal{M} is the free algebra $\bigwedge V$ over a graded vector space $V = \bigoplus_{i>1} V^i$, and
- (2) there is a collection of generators $\{x_{\tau}\}_{\tau \in I}$ indexed by some well ordered set I, such that $|x_{\mu}| \leq |x_{\tau}|$ if $\mu < \tau$, and each dx_{τ} is expressed in terms of preceding $x_{\mu}, \mu < \tau$. This in particular implies that dx_{τ} is always decomposable, i.e. dx_{τ} has no linear part.

We say that $(\bigwedge V, d)$ is a minimal model of the differential graded commutative algebra (\mathcal{A}, d) if $(\bigwedge V, d)$ is minimal and there exists a quasi-isomorphism $f: (\bigwedge V, d) \longrightarrow (\mathcal{A}, d)$. In [35], it is proved that any connected DGA has a minimal model unique up to isomorphism. For 1-connected DGAs, a similar result was proved earlier in [21].

A minimal model of a connected differentiable manifold M is a minimal model $(\bigwedge V, d)$ for $(\Omega^*(M), d)$. If M is simply connected, then the dual of the real homotopy vector space $\pi_i(M) \otimes \mathbb{R}$ is isomorphic to V^i , for any i (see [21]).

We say that a DGA (\mathcal{A}, d) is a *model* of a manifold M if (\mathcal{A}, d) and M have the same minimal model. Thus, if $(\bigwedge V, d)$ is the minimal model of M, we have

$$(\mathcal{A}, d) \xleftarrow{g} (\bigwedge V, d) \xrightarrow{f} (\Omega^*(M), d),$$

where f and g are quasi-isomorphisms.

A minimal algebra $(\bigwedge V, d)$ is called *formal* if there exists a morphism of differential algebras $f: (\bigwedge V, d) \longrightarrow (H^*(\bigwedge V), 0)$ inducing the identity map on cohomology. Also a differentiable manifold M is called *formal* if its minimal model is formal.

Simply connected compact manifolds of dimension ≤ 6 are always formal [49], so dimension 7 is the lowest dimension in which formality is an issue.

The formality of a minimal algebra $(\bigwedge V, d)$ is characterized by the condition that V can be decomposed into a direct sum $V = C \oplus N$ with d(C) = 0 with d injective on N and such that every closed element in the ideal I(N) in $\bigwedge V$ generated by N is exact [21]. This characterization can be weakened using the concept of s-formality introduced in [26, Definition 2.1] as follows.

Definition 3.1. A minimal algebra $(\bigwedge V, d)$ is *s*-formal (s > 0) if for each $i \leq s$ the space V^i of generators of degree *i* decomposes as a direct sum $V^i = C^i \oplus N^i$, where the spaces C^i and N^i satisfy the three following conditions:

MARISA FERNÁNDEZ, ANNA FINO, ALEXEI KOVALEV, AND VICENTE MUÑOZ

- (1) $d(C^i) = 0$,
- (2) the differential map $d: N^i \longrightarrow \bigwedge V$ is injective, and
- (3) any closed element in the ideal $I_s = I(\bigoplus_{i \le s} N^i)$, generated by the space $\bigoplus_{i \le s} N^i$

in the free algebra $\bigwedge (\bigoplus_{i \leq s} V^i)$, is exact in $\bigwedge V$.

A differentiable manifold M is *s*-formal if its minimal model is *s*-formal. Clearly, if M is formal then M is *s*-formal, for any s > 0. The main result of [26] shows that sometimes the weaker condition of *s*-formality implies formality.

Theorem 3.2. [26, Theorem 3.1] Let M be a connected and orientable compact differentiable manifold of dimension 2n or (2n-1). Then M is formal if and only if it is (n-1)-formal.

We will use also the following property, whose proof is exactly the same as the one given in [25, Lemma 2.4] for 7-dimensional simply connected compact manifolds with $b_2 \leq 1$.

Lemma 3.3. Let M be a 7-dimensional compact manifold with $b_1(M) = 0$ and $b_2(M) \leq 1$. Then, M is 3-formal and so formal.

In order to detect non-formality, instead of computing the minimal model, which is usually a lengthy process, one can use (triple) Massey products, which are obstructions to formality [21] and they are defined in the following way. Let (\mathcal{A}, d) be a DGA and suppose that there are $[a_i] \in H^{p_i}(\mathcal{A}), p_i > 0, 1 \leq i \leq 3$, such that $a_1 \cdot a_2$ and $a_2 \cdot a_3$ are exact. Write $a_1 \cdot a_2 = da_{1,2}$ and $a_2 \cdot a_3 = da_{2,3}$. The *(triple) Massey product* of the classes $[a_i]$ is defined as

$$\langle [a_1], [a_2], [a_3] \rangle = [a_1 \cdot a_{2,3} + (-1)^{p_1 + 1} a_{1,2} \cdot a_3] \in \frac{H^{p_1 + p_2 + p_3 - 1}(\mathcal{A})}{[a_1] \cdot H^{p_2 + p_3 - 1}(\mathcal{A}) + [a_3] \cdot H^{p_1 + p_2 - 1}(\mathcal{A})}$$

Note that a Massey product $\langle [a_1], [a_2], [a_3] \rangle$ on $(\mathcal{A}, d_{\mathcal{A}})$ is zero (or trivial) if and only if there exist $\tilde{x}, \tilde{y} \in \mathcal{A}$ such that $a_1 \cdot a_2 = d_{\mathcal{A}} \tilde{x}, a_2 \cdot a_3 = d_{\mathcal{A}} \tilde{y}$ and

$$0 = [a_1 \cdot \widetilde{y} + (-1)^{p_1 + 1} \widetilde{x} \cdot a_3] \in H^{p_1 + p_2 + p_3 - 1}(\mathcal{A}).$$

Models of fibrations. Let $F \to E \to B$ be a fibration of simply connected spaces. Let (\mathcal{A}_B, d_B) be a model (not necessarily minimal) of the base B, and let $(\bigwedge V_F, d_F)$ be a minimal model of the fiber F. By [23, section 15], a model of E is the *KS*-extension $(\mathcal{A}_B \otimes \bigwedge V_F, D)$, where D is defined as $Db = d_Bb$, for $b \in \mathcal{A}_B$, and $Dx = d_Fx + \Theta(x)$, for $x \in V_F$, and where

$$\Theta: V_F \to \mathcal{A}_B \oplus \left(\bigwedge^+ V_F \otimes \mathcal{A}_B^+\right).$$

Note that $\Theta(x)$ is written in terms of the previous generators of $\bigwedge V_F$, by minimality. The component of Θ in the first summand $\tau : V_F \to \mathcal{A}_B$ is the transgression map.

8

This is also true in the case that F, E and B are nilpotent spaces and the fibration is nilpotent, that is $\pi_1(B)$ acts nilpotently in the homotopy groups $\pi_i(F)$ of the fiber.

We will need two cases. In the case that E and B are simply connected and $F = SU(2) = S^3$ or $F = S^3/\mathbb{Z}_r$, with r > 0, the fibration is nilpotent. The minimal model of S^3 is $(\bigwedge a, d)$, with |a| = 3 and da = 0. Both spaces S^3 and S^3/\mathbb{Z}_r are rationally homotopy equivalent, because the quotient map $S^3 \to S^3/\mathbb{Z}_r$ is an isomorphism on cohomology and the fundamental group of S^3/\mathbb{Z}_r is trivial after rationalization. Therefore a fibration $S^3/\mathbb{Z}_r \to E \to B$ is a rational S^3 -fibration (that is, after rationalization of the spaces, it becomes a fibration). The map Θ is such that $\Theta(a) \in \mathcal{A}^4_B$ is a closed element of degree 4 defining the (rational) Euler class e(E) of the fibration.

The second case that we will need is for fibrations with $F = S^1 \times S^2$. Then the minimal model of F is $(\bigwedge(b, c, x), d)$, where |b| = 1, |c| = 2, |x| = 3, and the differential map is defined by db = 0, dc = 0 and $dx = c^2$. If B is simply connected then the fibration is nilpotent, and a model of E is given as $(\mathcal{A}_B \otimes \bigwedge(b, c, x), D)$, where the differential D is of the form

$$Db = \tau(b), \ Dc = b a'_2 + \tau(c), \ Dx = c^2 + c a_2 + b a_3 + \tau(x),$$

where τ is the transgression map, and $a_2, a'_2 \in \mathcal{A}^2_B, a_3 \in \mathcal{A}^3_B$. Here $\tau(b)$ is a closed element of degree 2 defining the Chern class of the S^1 -bundle induced from E via the fiberwise projection $S^1 \times S^2 \to S^1$.

4. On the formality of nearly parallel G_2 -manifolds

Let P be a simply connected compact 7-manifold with a nearly parallel G₂-structure φ inducing a metric g such that (P, g) is not isometric to the standard sphere S^7 . As already mentioned in the introduction, (P, g) belongs to one of the following three classes: a *proper* nearly parallel G₂-manifold, a Sasaki–Einstein manifold or a 3-Sasakian manifold.

Formality of simply connected compact 3-Sasakian manifolds, of dimension 7, was studied in [25], showing that a simply connected compact 3-Sasakian manifold (S, g) of dimension 7 is formal if and only if its second Betti number $b_2(S) \leq 1$.

Regarding the formality of Sasaki–Einstein manifolds, by [13, Theorem 3.2], if $\omega = \omega_1 + \omega_2 + \omega_3$ is the Kähler form on $S^2 \times S^2 \times S^2$, where $[\omega_1]$, $[\omega_2]$ and $[\omega_3]$ are the generators of the integral cohomology group of each of the S^2 -factors on $S^2 \times S^2 \times S^2$, then, the total space of the principal S^1 -bundle

$$S^1 \hookrightarrow Q(1,1,1) \longrightarrow S^2 \times S^2 \times S^2$$
,

with Euler class $[\omega] \in H^2(S^2 \times S^2 \times S^2, \mathbb{Z})$ is non-formal.

The above is a consequence of the following more general result.

Proposition 4.1. Consider the principal S^1 -bundle

 $S^1 \hookrightarrow Q \longrightarrow S^2 \times S^2 \times S^2$,

with Euler class $e_1a_1 + e_2a_2 + e_3a_3$, where $e_1, e_2, e_3 \in \mathbb{Z}$, and a_i is the generator of $H^2(S^2, \mathbb{Z})$ for the *i*-th copy of S^2 , i = 1, 2, 3. Then, Q is formal if and only if $e_1e_2e_3 = 0$.

Proof. We will determine a model of the 7-manifold Q. A minimal model of $S^2 \times S^2 \times S^2$ is the differential algebra $(\bigwedge(a_1, a_2, a_3, x_1, x_2, x_3), d)$, where $|a_i| = 2$ while $|x_i| = 3$ with $1 \le i \le 3$, and the differential d is defined by $da_i = 0$ and $dx_i = a_i^2$. Therefore, a model of the total space of a fiber bundle

$$S^1 \hookrightarrow Q \longrightarrow S^2 \times S^2 \times S^2$$

is the differential algebra over the vector space V generated by the elements y of degree 1, a_1, a_2, a_3 of degree 2, and x_1, x_2, x_3 of degree 3, and the differential d is given by

$$da_i = 0$$
, $dx_i = a_i^2$, $1 \le i \le 3$, $dy = e_1a_1 + e_2a_2 + e_3a_3$,

where $e_1a_1 + e_2a_2 + e_3a_3 \in H^2(S^2 \times S^2 \times S^2, \mathbb{Z})$ is the Euler class of the S^1 -bundle.

If all $e_1 = e_2 = e_3 = 0$, then $(\bigwedge V, d)$ is a minimal DGA, and so it is the minimal model of Q. But $(\bigwedge V, d)$ is also the minimal model of $S^1 \times S^2 \times S^2 \times S^2$, which is formal being the product of formal manifolds. Thus Q is formal because its minimal model is so.

If not all e_i are zero, we can assume $e_1 \neq 0$ (up to reordering). Then $e_1a_1 = dy - e_2a_2 - e_3a_3$, so

$$e_1^2 dx_1 = e_1^2 a_1^2 = (dy - e_2 a_2 - e_3 a_3)^2$$

= $d(y \cdot dy) - 2 \sum_{i=2}^3 e_i d(y \cdot a_i) + \sum_{i=2}^3 e_i^2 dx_i + 2e_2 e_3 a_2 \cdot a_3$.

Thus, letting

$$\widetilde{x}_1 = x_1 - e_1^{-2} \left(y \cdot dy - 2 \sum_{i=2}^3 e_i y \cdot a_i + \sum_{i=2}^3 e_i^2 x_i \right),$$

we have

 $d\widetilde{x}_1 = 2e_1^{-2}e_2e_3a_2 \cdot a_3$.

Then, the differential algebra $(\bigwedge(a_2, a_3, \tilde{x}_1, x_2, x_3), d)$ is a model of Q. In fact, the map $f: (\bigwedge(a_2, a_3, \tilde{x}_1, x_2, x_3), d) \to (\bigwedge(a_1, a_2, a_3, x_1, x_2, x_3, y), d)$ defined by $f(a_i) = a_i, f(x_i) = x_i \ (i = 2, 3), \text{ and } f(\tilde{x}_1) = x_1 - e_1^{-2} \left(y \cdot dy - 2 \sum_{i=2}^3 e_i y \cdot a_i + \sum_{i=2}^3 e_i^2 x_i \right)$ is a quasi-isomorphism.

Let us assume that $e_2e_3 = 0$. In this case, $d\tilde{x}_1 = 0$, and $(\bigwedge(a_2, a_3, x_2, x_3, \tilde{x}_1), d)$ is a minimal differential graded algebra. So $(\bigwedge(a_2, a_3, x_2, x_3, \tilde{x}_1), d)$ is the minimal model

of Q. Hence Q is formal since $(\bigwedge(a_2, a_3, x_2, x_3, \widetilde{x}_1), d)$ is also the minimal model of $S^2 \times S^2 \times S^3$, which is formal being the product of formal manifolds.

Finally, if $e_2e_3 \neq 0$, then

$$a_2 \cdot a_3 = \frac{e_1^2}{2e_2e_3} d\widetilde{x}_1.$$

We are going to show that Q is non-formal because there exists a non-zero Massey product on Q. By [10] we know that Massey products on a manifold can be computed by using any model for the manifold. Since $(\bigwedge(a_2, a_3, x_2, x_3, \widetilde{x}_1), d)$ is a model of Q, we have $H^*(Q) \cong H^*(\Lambda(a_2, a_3, x_2, x_3, \tilde{x}_1), d)$, so that $H^1(Q) = 0 = H^6(Q)$, $H^2(Q) \cong \langle [a_2], [a_3] \rangle, H^3(Q) = 0$, and by Poincaré duality for the 7-manifold Q, $H^4(Q) = 0$ and $H^5(Q)$ has dimension 2. Moreover, the Massey product $\langle [a_2], [a_2], [a_3] \rangle$ is defined and

$$\langle [a_2], [a_2], [a_3] \rangle = \left[\frac{e_1^2}{2e_2e_3} a_2 \cdot \widetilde{x}_1 - x_2 \cdot a_3 \right].$$

This element in $H^5(Q)$ cannot be exact since there is no non-zero element $x \in$ $\bigwedge^4(a_2, a_3, x_2, x_3, \widetilde{x}_1)$ such that $dx = \frac{e_1^2}{2e_2e_3}a_2 \cdot \widetilde{x}_1 - x_2 \cdot a_3$. Moreover, the indeterminacy of the Massey product is zero because $H^3(Q) = 0$. So $\langle [a_2], [a_2], [a_3] \rangle \neq 0$, and hence Q is non-formal.

In order to exhibit further examples of non-formal simply connected Sasaki–Einstein manifolds, we consider a del Pezzo surface P_k , for $3 \le k \le 8$, that is the blow-up of the complex projective space \mathbb{CP}^2 at k points,

$$P_k = \mathbb{CP}^2 \# k \overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# \overline{\mathbb{CP}}^2 \# \overbrace{\cdots}^k \# \overline{\mathbb{CP}}^2, \quad 3 \le k \le 8$$

where $\overline{\mathbb{CP}}^2$ is \mathbb{CP}^2 with the opposite of the standard orientation. Then the de Rham cohomology of P_k is

- $H^0(P_k) = \langle 1 \rangle$, $H^1(P_k) = 0$,
- $H^{2}(P_{k}) = \langle a, a_{1}, \dots, a_{k} \rangle,$ $H^{3}(P_{k}) = 0,$ $H^{4}(P_{k}) = \langle \nu \rangle,$

where $\nu = a^2$ is the volume form, and a is the integral cohomology class defined by the Kähler form on \mathbb{CP}^2 . Among these cohomology classes, the following relations are satisfied

 $a^{2} = -a_{i}^{2} = \nu$, for $1 \le i \le k$, $a \cdot a_{i} = 0 = a_{i} \cdot a_{j}$, for $1 \le i, j \le k$ and $i \ne j$. Now consider the 6-manifold $X_k = P_k \times S^2$, $3 \le k \le 8$.

Theorem 4.2. Let S_k be the total space of the circle bundle $S^1 \longrightarrow S_k \longrightarrow X_k = P_k \times S^2$, with Euler class $N(a - \sum_{i=1}^k \epsilon_i a_i + b)$, for $\epsilon_i > 0$ such that $\sum \epsilon_i < 1$, where

b is the generator of $H^2(S^2, \mathbb{Z})$, and N is a large integer satisfying that $N\epsilon_i \in \mathbb{Z}$ for all i. Then, for $3 \leq k \leq 8$ and some choice of Euler class as above, S_k is a simply connected compact Sasaki–Einstein manifold, with second Betti number $b_2 = k + 1$, which is non-formal.

Proof. It is standard that $a - \sum_{i=1}^{k} \epsilon_i a_i$ is a Kähler class for $P_k = \mathbb{CP}^2 \# k \overline{\mathbb{CP}}^2$, when $\epsilon_i > 0$ are small enough, since a is the Kähler class of \mathbb{CP}^2 and a_i represent the classes of the exceptional divisors of the blow-up. The effective cone of P_k is generated by $a - a_i$ and a_i , $i = 1, \ldots, k$. Hence the Kähler cone, which is its dual, is the open convex hull of a and $a - a_i$, $i = 1, \ldots, k$. So $a - \sum_{i=1}^{k} \epsilon_i a_i$ is a Kähler class for $\sum \epsilon_i < 1$, and $a - \sum_{i=1}^{k} \epsilon_i a_i + b$ is a Kähler class for $X_k = P_k \times S^2$. Now we choose N so that $N\epsilon_i \in \mathbb{Z}$, hence $N(a - \sum_{i=1}^{k} \epsilon_i a_i + b)$ is an integral cohomology class defined by a Kähler form on the product complex manifold $X_k = P_k \times S^2$. Therefore, there is a circle bundle $S_k \longrightarrow X_k = P_k \times S^2$ with Euler class equal to $N(a - \sum_{i=1}^{k} \epsilon_i a_i + b)$, where b is the generator of $H^2(S^2, \mathbb{Z})$.

Clearly S_k is a 7-dimensional simply connected, compact manifold, with second Betti number $b_2 = k + 1$. For some choice of Kähler form, S_k is Sasaki–Einstein. Indeed, Tian and Yau in [55], proved that there are Kähler–Einstein structures with $c_1 > 0$ on any manifold $P_k = \mathbb{CP}^2 \# k \overline{\mathbb{CP}}^2$, for $3 \le k \le 8$. Then, there exists a Kähler–Einstein metric on any manifold $X_k = P_k \times S^2$, for $3 \le k \le 8$ [29]. Thus, according to Section 2, S_k is a Sasaki–Einstein manifold.

By [21], if S_k has a non-zero Massey product, then S_k is non-formal. By [10], we know that Massey products on a manifold can be computed using any model for the manifold. Since X_k is a compact Kähler manifold, X_k is formal. Thus, a model of X_k is $(H^*(X_k), 0)$, where $H^*(X_k)$ is the de Rham cohomology algebra of X_k , that is

$$H^{0}(X_{k}) = \langle 1 \rangle,$$

$$H^{1}(X_{k}) = H^{3}(X_{k}) = H^{5}(X_{k}) = 0,$$

$$H^{2}(X_{k}) = \langle a, a_{1}, \dots, a_{k}, b \rangle,$$

$$H^{4}(X_{k}) = \langle a^{2}, a \cdot b, a_{1} \cdot b, \dots, a_{k} \cdot b \rangle,$$

$$H^{6}(X_{k}) = \langle a^{2} \cdot b \rangle.$$

Then, a model of S_k is the differential graded algebra (\mathcal{A}, d) , where $\mathcal{A} = H^*(X_k) \otimes \bigwedge(y)$, with |y| = 1, $d(H^*(X_k)) = 0$ and $dy = N(a - \sum_{i=1}^k \epsilon_i a_i + b)$. Write $\tilde{y} = \frac{1}{N}y$. Then, $H^1(\mathcal{A}, d) = H^3(\mathcal{A}, d) = 0$, $H^2(\mathcal{A}, d) = \langle [a], [a_1], \ldots, [a_k] \rangle$.

Using this model, we compute the Massey product $\langle [a], [a], [a_1] \rangle$. In this model $a \cdot a_i = 0$ $(1 \leq i \leq k)$, and

$$(a - \sum_{i=1}^{k} \epsilon_i a_i + b) \cdot (a - \sum_{i=1}^{k} \epsilon_i a_i - b) = (1 - \sum_{i=1}^{k} \epsilon_i^2) a^2,$$

since $b^2 = 0$. Then, $a \cdot a = (1 - \sum_{i=1}^k \epsilon_i^2)^{-1} d((a - \sum_{i=1}^k \epsilon_i a_i - b) \cdot \widetilde{y})$. So the Massey product $\langle [a], [a], [a_1] \rangle$ is defined, and

$$\langle [a], [a], [a_1] \rangle = \left[-(1 - \sum_{i=1}^k \epsilon_i^2)^{-1} \left((a - \sum_{i=1}^k \epsilon_i a_i - b) \cdot \widetilde{y} \right) \cdot a_1 \right]$$
$$= \left[-(1 - \sum_{i=1}^k \epsilon_i^2)^{-1} (\epsilon_1 \nu - b \cdot a_1) \cdot \widetilde{y} \right],$$

which is non-zero in $H^5(S_k)$. Therefore, S_k is non-formal. Note that there is no indeterminacy of this Massey product, since it lives in $[a] \cdot H^3(S_k) + [a_1] \cdot H^3(S_k)$, and we know that $H^3(S_k) \cong H^3(\mathcal{A}, d) = 0$.

Note that in [25] there is an example of a 7-dimensional regular simply connected Sasaki–Einstein manifold, with second Betti number $b_2 \geq 2$, which is formal, and so it does not admit any 3-Sasakian structure by [25]. Such a manifold is the total space of an S^1 -bundle over the blow-up of the complex projective space \mathbb{CP}^3 at four points. Other examples of simply connected formal Sasaki–Einstein manifolds, of dimension 7, are the total space of a circle bundle over the Kähler–Einstein manifold $\mathbb{CP}^2 \times S^2$ [10], and the space $W_{1,1}$ which, as Sasaki–Einstein manifold, is the total space of a circle bundle over the flag manifold F(1,2) [29].

5. Formality of proper nearly parallel G_2 -manifolds

According to [29], the only examples of proper nearly parallel G₂-manifolds, whose underlying metric is homogeneous, are the squashed 7-sphere S_{sq}^7 , the Berger space $\mathcal{B} = \mathrm{SO}(5)/\mathrm{SO}(3)$ and the Aloff–Wallach spaces $W_{k,l} = \mathrm{SU}(3)/S_{k,l}^1$. The only compact non-homogeneous examples of proper nearly parallel G₂-manifolds, known in the literature, are 7-dimensional compact non-homogeneous 3-Sasakian manifolds (S, g)with the canonical variation metric \tilde{g} of g given by (2.7). (Examples of such 3-Sasakian manifolds are given in [18].)

In this section we determine the minimal models of appropriate fibre bundles over \mathbb{CP}^2 and show that \mathcal{B} and $W_{k,l}$ are both formal. In particular, we show that the Aloff–Wallach spaces and $S^2 \times S^5$ have the same minimal model. It was previously known that as \mathcal{B} is a rational homology sphere it is geometrically formal for any choice of Riemannian metric, i.e. have the property that wedge products of harmonic forms are harmonic [42]. The homogeneous spaces $W_{k,l}$ are of Cartan type and therefore are formal by [43]. Thus the result that \mathcal{B} and $W_{k,l}$ are formal is not new but the proofs are new.

5.1. The Berger space. Consider the usual action of SO(3) on $\mathbb{R}^3 = \operatorname{span}\{x, y, z\}$. This action extends to an action of SO(3) on the polynomial ring $\mathbb{R}[x, y, z]$. Let $V_n \subset \mathbb{R}[x, y, z]$ be the SO(3)-submodule of homogeneous polynomials of degree n, and let $\mathcal{H}_n \subset V_n$ denote the SO(3)-submodule of harmonic polynomials of degree n, an irreducible SO(3)-module of dimension 2n+1. Every finite dimensional irreducible SO(3)-module is isomorphic to \mathcal{H}_n , for some n. The irreducible representation \mathcal{H}_2 of SO(3) has dimension 5, and so defines a non-standard embedding SO(3) \subset SO(5). The *Berger space* is the compact homogeneous space $\mathcal{B} = SO(5)/SO(3)$, given by the quotient of SO(5) by the copy of SO(3) embedded in SO(5) via the irreducible representation \mathcal{H}_2 of SO(3). The space \mathcal{B} has a metric such that the holonomy of its cone metric is Spin(7) [19]. The proper nearly parallel G₂-structure on \mathcal{B} is given explicitly in [6, Subsection 2.4.1].

Berger [12] proved that \mathcal{B} is a rational homology sphere with $H^4(\mathcal{B}, \mathbb{Z}) = \mathbb{Z}_{10}$, and it has the cohomology ring of an S^3 -bundle over S^4 . Therefore, \mathcal{B} and the sphere S^7 have the same minimal model (and in particular, it is formal). In the following theorem, we determine the minimal model of the total space of a principal S^3 -bundle over S^4 , and we show that such a space is formal. We apply this to conclude again that the space \mathcal{B} and S^7 have the same minimal model, and hence \mathcal{B} is formal.

Theorem 5.1. Consider an S^3 -bundle $S^3 \to P \to S^4$. Then P is formal. In particular, the Berger space $\mathcal{B} = SO(5)/SO(3)$ is formal.

Proof. Let $e[\omega]$ be the Euler class of the bundle, where $e \in \mathbb{Z}$, and $[\omega]$ is the generator of the integral cohomology group $H^4(S^4, \mathbb{Z})$. The minimal model of S^4 is the differential graded algebra $(\bigwedge(a, u), d)$, with |a| = 4, |u| = 7, da = 0 and $du = a^2$. The minimal model of S^3 is $(\bigwedge b, d)$, where |b| = 3 and db = 0. Then, according to Section 3, a model of P is $(\bigwedge(a, u, b), D)$, with

$$Da = 0$$
, $Du = a^2$, $Db = ea$.

If e = 0, then the DGA ($\bigwedge(a, u, b), d$) is minimal, and so it is the minimal model of P. But ($\bigwedge(a, u, b), D$) is the minimal model of $S^3 \times S^4$, which is formal being the product of two formal manifolds. Therefore, P is formal.

Suppose now that $e \neq 0$. In this case we have $du = a^2 = e^{-1}D(b \cdot a)$. So the element $\tilde{u} = u - e^{-1}b \cdot a$ has degree 7 and $D\tilde{u} = 0$. Then the DGA $(\bigwedge(a, \tilde{u}, b), D)$ is also a model of P, because it is quasi-isomorphic to $(\bigwedge(a, u, b), D)$. Moreover, $(\bigwedge \tilde{u}, 0)$ is the minimal model of P. In fact, $(\bigwedge \tilde{u}, 0)$ is a minimal DGA, and a model of P because, taking into account that Db = ea, the map $f: (\bigwedge \tilde{u}, 0) \to (\bigwedge(a, \tilde{u}, b), D)$ given by $f(\tilde{u}) = \tilde{u}$ is a quasi-isomorphism. Therefore, P is formal since $(\bigwedge \tilde{u}, 0)$ is the minimal model of the sphere S^7 , which is formal.

In [31] it is proved that the Berger space is diffeomorphic to the S^3 -bundle over S^4 with Euler class $-10 \, [\omega]$. Thus \mathcal{B} and S^7 have the same minimal model, and consequently \mathcal{B} is formal.

5.2. The Aloff–Wallach spaces. Let $k, l \in \mathbb{Z}$ be non-zero, co-prime integers, and $S_{k,l}^1$ be a circle subgroup of SU(3) consisting of elements of the form

$$\left(\begin{array}{ccc} e^{ik\theta} & 0 & 0\\ 0 & e^{il\theta} & 0\\ 0 & 0 & e^{im\theta} \end{array}\right),$$

where k + l + m = 0. The Aloff–Wallach space $W_{k,l} = SU(3)/S_{k,l}^1$ is the quotient of SU(3) by this circle subgroup [2]. Note that there are examples of different pairs (k, l) such that the corresponding Aloff–Wallach spaces are homeomorphic but not diffeomorphic [44]. The spaces $W_{k,l}$ are called generic if $\{k, l, -(k+l)\}$ is different from $\{1, 1, -2\}$ or $\{1, -1, 0\}$. We will denote by $W_{1,1}$ and $W_{1,-1}$ the two exceptional Aloff–Wallach spaces.

By [51] (see also [58]) all the spaces $W_{k,l}$ admit two homogeneous Einstein metrics. If (k, l) = (1, 1) one of those metrics is the 3-Sasakian structure on the space $W_{1,1}$ mentioned in Section 4, and the other is induced by a proper homogeneous nearly parallel G₂-structure. If (k, l) = (1, -1), the space $W_{1,-1}$ admits only one proper homogeneous nearly parallel G₂-structure, up to homotheties [51]. On the generic Aloff-Wallach spaces the two metrics are induced by proper homogeneous nearly parallel G₂-structures [9], which by [51, 52] are only two, up to homotheties. The expressions of those two G₂-structures are given in [8, 20].

The manifold $W_{k,l}$ is simply connected with $H^2(W_{k,l},\mathbb{Z}) \cong \mathbb{Z}$ and $H^3(W_{k,l},\mathbb{Z}) = 0$ (see [44]). Thus, $b_1(W_{k,l}) = b_3(W_{k,l}) = 0$ and $b_2(W_{k,l}) = 1$. Hence, $W_{k,l}$ is formal by Lemma 3.3.

In [34] (see also [8]) it is shown that there is a canonical fibration

$$\pi: W_{k,l} \to \mathbb{CP}^2,$$

whose fibers are the lens spaces $S^3/\mathbb{Z}_{|k+l|}$ if $k+l \neq 0$, or $S^1 \times S^2$ if k+l = 0. In the two following theorems, we determine the minimal model of the total space of a F-fiber bundle over \mathbb{CP}^2 , where $F = S^1 \times S^2$, or $F = S^3/\mathbb{Z}_p$ with p > 0, and we prove that such a space is formal. In particular, we show that $W_{k,l}$ and $S^5 \times S^2$ have the same minimal model.

Theorem 5.2. Let P be the total space of an S^3/\mathbb{Z}_p -bundle $S^3/\mathbb{Z}_p \to P \to \mathbb{CP}^2$ with p > 0. Then P is formal. In particular, if $k + l \neq 0$, the Aloff–Wallach space $W_{k,l}$ is formal, and $W_{k,l}$ and the product manifold $S^2 \times S^5$ have the same minimal model.

Proof. We will determine the minimal model of P and show that it is formal. According to Section 3, the fibre bundle $S^3/\mathbb{Z}_p \to P \to \mathbb{CP}^2$ is a rational S^3 -fibration. Let $e a^2$ be its (rational) Euler class, where $e \in \mathbb{Q}$ and $a = [\omega]$ is the generator of the integral cohomology group $H^2(\mathbb{CP}^2, \mathbb{Z})$. By [23], if $(\mathcal{A}, d_{\mathcal{A}})$ is a model of \mathbb{CP}^2 , we have that $(\mathcal{A} \otimes \bigwedge u, d)$, with |u| = 3, $d|_{\mathcal{A}} = d_{\mathcal{A}}$ and $du = e a^2$, is a model of P.

The minimal model of \mathbb{CP}^2 is the differential graded algebra $(\bigwedge(a, x), d)$, where |a| = 2, |x| = 5, da = 0 and $dx = a^3$. Therefore, the KS-model of P is $(\bigwedge(a, x, u), d)$, with $du = e a^2$.

If e = 0 then the differential graded algebra $(\bigwedge(a, x, u), d)$ is minimal, and so it is the minimal model of P. Moreover, $(\bigwedge(a, x, u), d)$ is the minimal model of $S^3 \times \mathbb{CP}^2$, which is formal being the product of two formal manifolds. Thus, P is formal.

Suppose now that $e \neq 0$. In this case we have $a^2 = e^{-1}du$, and so the element of degree five $\tilde{x} = x - e^{-1}a \cdot u$ is such that $d\tilde{x} = 0$. Clearly $(\bigwedge(a, \tilde{x}, u), d)$ is a minimal DGA, and a model of P because the map $f: (\bigwedge(a, \tilde{x}, u), d) \to (\bigwedge(a, x, u), d)$ given by $f(a) = a, f(\tilde{x}) = x - e^{-1}a \cdot u$ and f(u) = u is a quasi-isomorphism. Therefore, $(\bigwedge(a, \tilde{x}, u), d)$ is the minimal model of P. Thus, P is formal since $(\bigwedge(a, \tilde{x}, u), d) = (\bigwedge\tilde{x} \otimes \bigwedge(a, u), d)$ is the minimal model of $S^5 \times S^2$, which is formal.

All this shows not only that P is formal but also the minimal model of P. Indeed, the minimal model of P is either the minimal model of $S^3 \times \mathbb{CP}^2$ or the minimal model of $S^5 \times S^2$. Thus, if the third Betti number of P is $b_3(P) = 1$, then P and $S^3 \times \mathbb{CP}^2$ have the same minimal model, while if $b_3(P) = 0$, then the minimal model of P is the minimal model of $S^5 \times S^2$. Therefore, for $k + l \neq 0$, the space $W_{k,l}$ and $S^5 \times S^2$ have the same minimal model since $b_3(W_{k,l}) = 0$.

Theorem 5.3. The total space of an $S^1 \times S^2$ -bundle $S^1 \times S^2 \to P \to \mathbb{CP}^2$ is formal. In particular, for $k \neq 0$, the Aloff–Wallach space $W_{k,-k}$ is formal, and $W_{k,-k}$ and the product manifold $S^2 \times S^5$ have the same minimal model.

Proof. We know that the minimal model of \mathbb{CP}^2 is the differential graded algebra $(\bigwedge(a, x), d)$, where |a| = 2, |x| = 5, da = 0 and $dx = a^3$. Moreover, the minimal model of $S^1 \times S^2$ is $(\bigwedge(b, c, y), d)$, where |b| = 1, |c| = 2, |y| = 3, and the differential map is defined by db = 0, dc = 0 and $dy = c^2$. Hence the KS-model of P is of the form

$$(\bigwedge (a, x, b, c, y), D), Da = 0, Dx = a^3, Db = e a, Dc = g ab, Dy = c^2 + f a^2 + h ac,$$

for some $e, f, g, h \in \mathbb{Z}$. First note that taking $\tilde{c} = c + \frac{1}{2}h a$, we have $D\tilde{c} = g ab$ and $Dy = \tilde{c}^2 + \tilde{f}a^2$, for some \tilde{f} . Therefore we can assume in the above that h = 0. We do that without changing the names of the variables. By noting that $D^2(y) = 2g abc = 0$, we get g = 0. Altogether we get the model of P as

$$\left(\bigwedge (a, x, b, c, y), D\right), Da = 0, Dx = a^3, Db = e a, Dc = 0, Dy = c^2 + f a^2.$$

If e = f = 0 then $(\bigwedge (a, x, b, c, y), D)$ is the minimal model of $S^1 \times S^2 \times \mathbb{CP}^2$, which is formal being the product of three formal manifolds. Hence P is formal. Suppose now that $e \neq 0$ and f = 0. In this case, $a = e^{-1}Db$. Then the element \tilde{x} of degree 5 given by $\tilde{x} = x - e^{-1}a \cdot u$ is such that $D\tilde{x} = 0$. Thus, $(\bigwedge(a, \tilde{x}, b, c, y), D)$ is a model of P because this DGA is quasi-isomorphic to $(\bigwedge(a, x, b, c, y), D)$. Moreover, $(\bigwedge(\tilde{x}, c, y), D)$ is a minimal DGA, and a model of P. In fact, taking into account that $a = e^{-1}Db$, one can check that the map $f: (\bigwedge(\tilde{x}, c, y), D) \to (\bigwedge(a, \tilde{x}, b, c, y), D)$ given by $f(\tilde{x}) = \tilde{x}, f(c) = c$ and f(y) = y is a quasi-isomorphism. Therefore, $(\bigwedge(\tilde{x}, c, y), D)$ is the minimal model of P. Thus, P is formal since $(\bigwedge(\tilde{x}, c, y), D)$ is the minimal model of $S^5 \times S^2$, which is formal.

If $e \neq 0$ and $f \neq 0$, as before we determine the minimal model of P. Take the elements $\tilde{x} = x - e^{-1}b \cdot a^2$ and $\tilde{y} = y - fe^{-1}a \cdot b$ of degree 5 and 3, respectively. Then, we get the model $(\bigwedge(a, \tilde{x}, b, c, \tilde{y}), D)$ of P with Da = 0, $d\tilde{x} = 0$, Db = ea, Dc = 0 and $D\tilde{y} = c^2$. Consider the differential graded algebra $(\bigwedge(\tilde{x}, c, \tilde{y}), D)$, which is a minimal DGA, and a model of P. In fact, the map $f: (\bigwedge(\tilde{x}, c, \tilde{y}), D) \to (\bigwedge(a, \tilde{x}, b, c, \tilde{y}), D)$ given by $f(\tilde{x}) = \tilde{x}$, f(c) = c and $f(\tilde{y}) = \tilde{y}$ is a quasi-isomorphism. Therefore, $(\bigwedge(\tilde{x}, c, \tilde{y}), D)$ is the minimal model of P. Thus, P is formal since $(\bigwedge(\tilde{x}, c, \tilde{y}), D)$ is the minimal model of $S^5 \times S^2$, which is formal.

Finally, suppose that e = 0 and $f \neq 0$. Then the model $(\bigwedge(a, x, b, c, y), D)$ of P is such that Da = 0, $Dx = a^3$, Db = 0, Dc = 0 and $Dy = c^2 + f a^2$. This implies that the differential graded algebra $(\bigwedge(a, x, b, c, y), D)$ is minimal, and so it is the minimal model of P. To show that P is formal we proceed as follows. First note that $\bigwedge(a, x, b, c, y) = \bigwedge b \otimes \bigwedge(a, x, c, y)$. Clearly, $(\bigwedge b, 0)$ is the minimal model of S^1 , which is formal, and $(\bigwedge(a, x, c, y), D)$ is the minimal model of the total space X of an S^2 -bundle

$$S^2 \to X \to \mathbb{CP}^2.$$

The de Rham cohomology of X is $H^1(X) = H^3(X) = H^5(X) = 0$, $H^2(X) = \langle [a], [c] \rangle$, $H^4(X) = \langle [a]^2, [a] \cdot [c] \rangle$ and $H^6(X) = \langle [a]^2 \cdot [c] \rangle$. Thus, the 6-manifold X is formal since $b_1(X) = 0$. In fact, denote by V the graded vector space generated by the elements a, x, c and y. Because V is a graded vector space, we can consider the space V^i of generators of degree i, which decomposes as a direct sum $V^i = C^i \oplus N^i$, with $C^1 = N^1 = 0$, $C^2 = \langle a, c \rangle$ and $N^2 = 0$. Thus, according to Definition 3.1, the manifold X is 2-formal and, by Theorem 3.2, X is formal.

Therefore, the minimal model $(\bigwedge(a, x, b, c, y), D)$ of P is the minimal model of the product manifold $S^1 \times X$, which is formal being the product of two formal manifolds. Hence P is formal.

Thus, if P is the total space of an $S^1 \times S^2$ -bundle over \mathbb{CP}^2 , then P and $S^5 \times S^2$ have the same minimal model, or the minimal model of P is the minimal model of the manifold M, where $M = S^1 \times S^2 \times \mathbb{CP}^2$, or $M = S^1 \times X$. So, for $k \neq 0$, $W_{k,-k}$ and $S^5 \times S^2$ have the same minimal model since $b_1(W_{k,-k}) = 0$.

Remark 5.4. In [22] it is proved that, for k, l co-prime integers, the space $W_{k,l} = SU(3)/S_{k,l}^1$ can be described as follows. Let X be the total space of the S^2 -bundle $S^2 \longrightarrow X \xrightarrow{\pi} \mathbb{CP}^2$ with Pontryagin class $p_1 = -3$ and Stiefel-Whitney class $w_2 \neq 0$. Then, $W_{k,l}$ is the total space of the circle bundle

$$S^1 \longrightarrow W_{k,l} \xrightarrow{\varpi} X$$
 (5.1)

determined by the Euler class ka+lb, where a, b are the generators of $H^2(X, \mathbb{Z}) \cong \mathbb{Z}^2$, and the base space $X = \mathrm{SU}(3)/T^2$.

Note that the space $W_{k,l}$ thus defined can be also considered as the aforementioned F-fiber bundle $F \longrightarrow W_{k,l} \longrightarrow \mathbb{CP}^2$, where $F = S^1 \times S^2$, or $F = S^3/\mathbb{Z}_p$ with $p \neq 0$. Indeed, for $z \in \mathbb{CP}^2$, consider $X_z = \pi^{-1}(z)$ and $W_{k,l}(z) = \mathrm{pr}^{-1}(z)$, where $\mathrm{pr} = \pi \circ \varpi$. If we restrict to each X_z , this circle bundle $S^1 \longrightarrow W_{k,l}(z) \longrightarrow X_z$ must be $W_{k,l}(z) = S^1 \times S^2$ if the Euler class is e = 0, or a lens space S^3/\mathbb{Z}_e if the Euler class is $e \neq 0$. Varying over all $z \in \mathbb{CP}^2$, we have a fiber bundle $F \longrightarrow W_{k,l} \longrightarrow \mathbb{CP}^2$, where $F = S^1 \times S^2$ or $F = S^3/\mathbb{Z}_e$.

6. The nearly parallel G_2 -manifold Q(1, 1, 1)

In this section, we consider the regular Sasaki–Einstein 7-manifold Q(1,1,1) and we work out explicitly, in coordinates, the S^1 -family of nearly parallel G_2 -structures on it, which we will use in Section 7 and Section 9.

We start from the Kähler manifold $X = S^2 \times S^2 \times S^2$ with Kähler form

$$\omega = \omega_1 + \omega_2 + \omega_3, \qquad (6.1)$$

where ω_1 , ω_2 and ω_3 are the generators of the integral cohomology group for each of the S^2 -factors on $S^2 \times S^2 \times S^2$. Let M be the total space of the principal S^1 -bundle

$$S^1 \hookrightarrow M \xrightarrow{\pi_M} X = S^2 \times S^2 \times S^2,$$
 (6.2)

with Euler class $[\omega] \in H^2(X, \mathbb{Z})$. Then, by application of [16, cf. Theorem 2.8], M is a simply connected compact Sasaki–Einstein manifold, with contact form η such that $d\eta = 2\pi_M^*(\omega)$. From [9, §4.2] (see also [28, 29]), we know that M is the homogeneous space

$$M = Q(1, 1, 1) = \left(\,\mathrm{SU}(2) \times \mathrm{SU}(2) \times \mathrm{SU}(2) \right) / \left(\mathrm{U}(1) \times \mathrm{U}(1) \right). \tag{6.3}$$

Throughout this section, the notation M and X will always mean the manifolds defined in (6.2) and (6.3).

We apply the Kobayashi construction [41] to determine the contact form for the Sasaki–Einstein metric of the principal circle bundle M. Let $\theta_j \in (0, \pi)$ and $\phi_j \in (0, 2\pi)$, j = 1, 2, 3, be the standard spherical coordinates on each of the S^2 -factors in X and $z_j = \cot(\theta_j/2)e^{i\phi_j} \in \mathbb{C}$ a complex coordinate defined via the stereographic projection. The Fubini–Study metric of volume 1 on the j-th S^2 factor has the Kähler

form $\omega_j = \frac{1}{4\pi} dd^c \log(1 + z_j \bar{z}_j) = \alpha_{2j-1} \wedge \alpha_{2j}$, where we used an orthonormal co-frame field

$$\alpha_{2j-1} = \frac{-1}{2\sqrt{\pi}} \, d\theta_j, \qquad \alpha_{2j} = \frac{1}{2\sqrt{\pi}} \, \sin\theta_j \, d\phi_j, \qquad j = 1, 2, 3, \tag{6.4}$$

on the open dense region $U_X \subset X$ defined by $z_j \neq 0$ for all j. The Kähler form (6.1) on X restricts to an exact form on U_X

$$\omega = \alpha_1 \wedge \alpha_2 + \alpha_3 \wedge \alpha_4 + \alpha_5 \wedge \alpha_6 = \frac{1}{4\pi} \sum_{j=1}^3 d(\cos\theta_j \, d\phi_j). \tag{6.5}$$

The principal S^1 -bundle M trivializes over U_X , $\pi_M^{-1}(U_X) \cong U_X \times S^1$. Let $s \in (0, 4\pi)$ denote a coordinate on the fibre S^1 and consider on $\pi_M^{-1}(U_X)$ a 1-form

$$\eta = \frac{1}{2\pi} \left(-ds + \sum_{j=1}^{3} \cos \theta_j \, d\phi_j \right). \tag{6.6}$$

Noting that $[\omega] \in H^2(X,\mathbb{Z})$ is an integral cohomology class, it can be checked by following the general construction in [41, §2] that η extends smoothly from $\pi_M^{-1}(U_X)$ to a well-defined connection 1-form on all of M (still denoted by η). Since

$$d\eta = 2\pi_M^*(\omega),\tag{6.7}$$

it also follows that η is a well-defined contact form on M (cf. Section 2).

We shall slightly abuse the notation by writing α_i also for the lifting to M via π_M^* of the local 1-forms (6.4). Then $\alpha_1 + i\alpha_2$, $\alpha_3 + i\alpha_4$, $\alpha_5 + i\alpha_6$ can be considered as point-wise orthonormal (1,0)-forms on the complex cone $\pi^{-1}(U_X) \times \mathbb{R}^+$ as π_M is a Riemannian submersion and the associated bundle projection $M \times \mathbb{R}^+ \to X$ is holomorphic.

The Riemannian cone $M \times \mathbb{R}^+$ is simply-connected and Ricci-flat Kähler. Therefore, $M \times \mathbb{R}^+$ has a non-vanishing holomorphic (4, 0)-form $\widehat{\Psi}$, which is parallel with respect to the Kähler metric. In local coordinates, the Kähler form on $M \times \mathbb{R}^+$ is given by $rdr \wedge \eta + r^2 \sum_{j=1}^3 \alpha_{2j-1} \wedge \alpha_{2j}$ (cf. (2.3)) and $\widehat{\Psi}$ can be written as

$$\widehat{\Psi} = r^3 e^{i\mu} \left(dr + i\frac{\pi}{2} r\eta \right) \wedge (\alpha_1 + i\alpha_2) \wedge (\alpha_3 + i\alpha_4) \wedge (\alpha_5 + i\alpha_6), \tag{6.8}$$

where a smooth real function μ on M is determined, up to adding a constant, by the condition $d\widehat{\Psi} = 0$. Setting

$$\Psi = e^{-is}(\alpha_1 + i\alpha_2) \wedge (\alpha_3 + i\alpha_4) \wedge (\alpha_5 + i\alpha_6), \tag{6.9}$$

it is straightforward to check that

$$d\Psi = -2\pi i \,\Psi \wedge \eta,\tag{6.10}$$

thus, noting also that $\Psi \wedge d\eta = 0$, one can take $\mu = -s$ in (6.8). The holomorphic 4-form $\widehat{\Psi}$ extends from $\pi^{-1}(U_X) \times \mathbb{R}^+$ to all of $M \times \mathbb{R}^+$ (e.g. by patching with a different choice of spherical coordinate neighbourhood in (6.4)). Respectively, Ψ extends to a well-defined horizontal complex volume form on M.

We can now write a coordinate expression for the S^1 -family of nearly parallel G₂structures φ_t on M induced by the Sasaki–Einstein structure, cf. (2.6),

$$\varphi_t = \frac{\pi}{2}\omega \wedge \eta + \cos(s+t)\Psi_+ + \sin(s+t)\Psi_- = \frac{\pi}{2}\omega \wedge \eta + \operatorname{Re}(e^{-it}\Psi)$$
(6.11)

where the real 3-forms Ψ_+ , Ψ_- in (6.11) are determined by $\Psi = e^{-is}(\Psi_+ + i\Psi_-)$ and can be written in coordinates using (6.9). The respective G₂ 4-forms are

$$\star_{\varphi_t}\varphi_t = \frac{1}{2}\omega \wedge \omega + (\cos(s+t)\Psi_- - \sin(s+t)\Psi_+) \wedge \eta = \frac{1}{2}\omega \wedge \omega + \operatorname{Im}(e^{-it}\Psi) \wedge \eta,$$

and from (6.7) and (6.10) it follows that

$$d\varphi_t = 2\pi \star \varphi_t \,, \tag{6.12}$$

for each t. The induced metric $g = g_{\varphi_t}$ on M does not depend on t and is given by

$$g = \frac{1}{16} \left(ds - \sum_{j=1}^{3} \cos \theta_j \, d\phi_j \right)^2 + \frac{1}{4\pi} \sum_{j=1}^{3} \left(d\theta_i^2 + \sin^2 \theta_j \, d\phi_j^2 \right),$$

so the local coframe of 1-forms $\{\alpha_1, \ldots, \alpha_6, \alpha_7 = \eta\}$ is orthonormal.

7. Associative 3-folds in Sasaki–Einstein 7-manifolds

We now turn to consider minimal associative 3-folds in nearly parallel G₂-manifolds. In this section, we deal with the case when the nearly parallel G₂-structure arises from a Sasaki–Einstein structure on a 7-manifold, with particular attention to the regular Sasaki–Einstein manifolds S_k and Q(1,1,1) discussed in Section 4 and Section 6, respectively.

Firstly, let us recall that if (P, φ) is a nearly-parallel G₂-manifold, then an *associa*tive 3-fold is an oriented 3-dimensional submanifold $X \subset P$ such that

$$\varphi|_X = \operatorname{vol}_X.$$

In particular, associative 3-folds in P are the links of Cayley cones in the metric cone over P, and hence are also minimal submanifolds of P. Even more, Ball and Madnick in [5, Theorem 2.18] prove that the largest torsion class of G₂-structures for which every associative 3-fold is minimal is given by the nearly parallel G_2 -structures.

Let (S, g_S) be a Sasaki–Einstein 7-manifold and φ_t the corresponding S^1 -family of nearly parallel G₂-structures on S defined in (2.6). Assume that S is simplyconnected, so the metric cone $S \times \mathbb{R}^+$ has holonomy in SU(4), and denote by ω^c the Kähler form and by $\widehat{\Psi}$ a non-vanishing holomorphic 4-form on $S \times \mathbb{R}^+$. Then $S\times \mathbb{R}^+$ has a 1-parameter family of torsion-free Spin(7)-structures induced by the closed 4-forms

$$\widehat{\Phi}_t = \frac{1}{2}\omega^c \wedge \omega^c + \operatorname{Re}(e^{-it}\widehat{\Psi}), \quad t \in \mathbb{R}.$$
(7.1)

cf. [38, Prop. 13.1.4]. Recall that we identify the 7-manifold S as a submanifold $S \times \{1\}$ of the cone. The nearly parallel G₂-structures φ_t on S are then related to the Spin(7)-structures (7.1) by

$$\varphi_t = \partial_r \lrcorner \Phi_t |_{r=1}, \quad \text{for each } t,$$

cf. [1, pp. 723–724]. If a 4-dimensional submanifold Z of $S \times \mathbb{R}^+$ is calibrated by $\widehat{\Phi}$ (i.e. Z is a Cayley submanifold) and $Z = Y \times \mathbb{R}^+$ for some submanifold $Y \subset S$, then $(\varphi_t|_Y) \wedge dr$ is the volume form of the conical metric $r^2g_S + dr^2$ on $Y \times \mathbb{R}^+$ as ∂_r defines a unit normal vector field along each $Y \times \{r\}$. It follows that Y is an associative 3-fold in S. Conversely, if $Y \subset S$ is an associative 3-fold then $Z = Y \times \mathbb{R}^+ \subset S \times \mathbb{R}^+$ is Cayley.

Examples of Cayley submanifolds, in the case when the Spin(7)-structure is of the form (7.1) induced by an SU(4)-structure, include complex surfaces and special Lagrangian submanifolds. We shall consider these two possibilities in order.

If $Z = Y \times \mathbb{R}^+$ is a complex surface in $S \times \mathbb{R}^+$, so the tangent spaces of Z are preserved by the (integrable) almost complex structure J on $S \times \mathbb{R}^+$, then Y is an 'invariant' submanifold for the contact structure on S as defined in [14, §8.1]. More explicitly, the endomorphism Φ discussed in Section 2 maps each tangent space of Y into itself and the Reeb vector field ξ is tangent to Y.

Proposition 7.1. Let S be a regular Sasaki–Einstein 7-manifold with contact form η arising from a principal S¹-bundle $\pi : S \to X$ with Euler class $c_1 = [\omega]$, where X is a projective complex 3-fold with Kähler form ω and $d\eta = \pi^*(\omega)$. Let φ_t be the corresponding 1-parameter family of nearly parallel G₂-forms defined in (2.6).

Then, for each complex curve Σ in X, the preimage $Y_{\Sigma} = \pi^{-1}(\Sigma) \subset S$ is an invariant submanifold for the contact structure η and a (minimal) associative 3-fold with respect to φ_t for each t.

In particular, if S is Q(1,1,1) or S_k , so $X = \mathbb{CP}^1 \times P$, with $P = \mathbb{CP}^1 \times \mathbb{CP}^1$ or $P = P_k$ ($3 \le k \le 8$) a del Pezzo surface, and $\omega = \omega_1 + \omega_P$, where $[\omega_1]$ is the (positive) generator of $H^2(\mathbb{CP}^1,\mathbb{Z})$, $[\omega_P] \in H^2(P,\mathbb{Z})$ and $\Sigma = \mathbb{CP}^1 \times \{p\}$, $p \in P$, then the minimal associative Y_{Σ} is diffeomorphic to the sphere S^3 .

Proof. Recall from (2.3) that the Kähler form on the cone $S \times \mathbb{R}^+$ is

$$\omega^c = r^2 d\eta + 2r \, dr \wedge \eta = r^2 \pi^* \omega + 2r \, dr \wedge \eta, \tag{7.2}$$

where $r \in \mathbb{R}^+$ and we extended π to a projection $\pi : S \times \mathbb{R}^+ \to X$ independent of the \mathbb{R}^+ factor. Pulling back to submanifolds $Y_{\Sigma} \times \mathbb{R}^+$ and $\Sigma \subset Y_{\Sigma} \times \{1\}$ we find that

$$\frac{1}{2}(\omega^c \wedge \omega^c)|_{Y_{\Sigma} \times \mathbb{R}^+} = \pi^*(\omega|_{\Sigma \times \{1\}}) \wedge 2\eta \wedge dr = \operatorname{vol}_{Y_{\Sigma} \times \mathbb{R}^+}$$

noting also that π is a Riemannian submersion and 2η has unit length in the metric induced by φ_t . Thus, $Y_{\Sigma} \times \mathbb{R}^+$ is calibrated by $\frac{1}{2}\omega^c \wedge \omega^c$ and must be a complex surface in $S \times \mathbb{R}^+$ by the Wirtinger inequality. The link Y_{Σ} is therefore associative.

For the last part, we can consider $S \times \mathbb{R}^+$ as the total space of a holomorphic line bundle over X. The restriction of this bundle to a projective line Σ is isomorphic to the hyperplane bundle $\mathcal{O}(1)$ over \mathbb{CP}^1 . The total space $Y_{\Sigma} \times \mathbb{R}$ is then biholomorphic to $\mathbb{CP}^2 \setminus \{(0:0:1)\}$. We obtain that the associative 3-fold Y_{Σ} is diffeomorphic to the sphere S^3 , noting that the fibres of $Y_{\Sigma} \times \mathbb{R}$ correspond to projective lines through (0:0:1), so the zero section Σ can be identified as a projective line in \mathbb{CP}^2 avoiding (0:0:1).

Remarks.

- (i) It is not difficult to see that every minimal associative Y_{Σ} in Proposition 7.1 is invariant under the (isometric) S^1 -action on the principal bundle S and every deformation of the holomorphic curve Σ in X induces an associative deformation of Y_{Σ} .
- (ii) We obtain from (2.6) that $\varphi_t|_{Y_{\Sigma}} = 2\eta \wedge \Omega_{\Sigma}$, where $\Omega_{\Sigma} = \pi^*(\omega|_{\Sigma})$. As π is a Riemannian submersion, the form Ω_{Σ} at each point in Y_{Σ} can be written as a wedge product of two unit-length covectors which are orthogonal to η in the metric induced by φ_t . (If S = Q(1, 1, 1) and $\Sigma = \mathbb{CP}^1 \times \{(p_1, p_2)\}$, then $\Omega_{\Sigma} = \alpha_1 \wedge \alpha_2$ in the notation of (6.4).)
- (iii) In the last part of Proposition 7.1, one can more generally take Σ to be the graph of a holomorphic embedding $\mathbb{CP}^1 \to P$. For S = Q(1,1,1), the ambiguity of taking such Σ corresponds to a generic choice of two rational functions of one complex variable.

Concerning deformations of minimal associative 3-folds for nearly parallel G_2 -structures of this type, we more generally have:

Proposition 7.2. Let S be a regular Sasaki–Einstein 7-manifold with contact form η arising from a principal S¹-bundle $\pi : S \to X$ and let φ_t be the induced S¹-family (2.6) of nearly parallel G₂-structures on S.

If $Y \subset S$ is an associative 3-fold with respect to φ_{t_0} for some fixed t_0 and $Y \neq \pi^{-1}(\Sigma)$ for any real 2-dimensional submanifold $\Sigma \subset X$, then the free S^1 -action on the principal bundle S induces non-trivial φ_{t_0} -associative deformations of Y (in particular, Y is not rigid).

Proof. The free S^1 -action on S preserves the family φ_t and all φ_t induce the same (Sasaki–Einstein) metric on S. Therefore, the S^1 -orbit of Y consists of volume minimizing submanifolds of S which are in general distinct from Y by the hypothesis.

The result now follows by application of [36, Theorem II.4.2].

We now turn to consider associative 3-folds arising as links of Cayley submanifolds $Z = Y \times \mathbb{R}^+$ which are a special Lagrangian in $S \times \mathbb{R}^+$, so $\omega^c|_Z = 0$ and $\operatorname{Re}(e^{-it}\widehat{\Psi})|_Z = 0$ for some fixed t. Then (and only then) the cross-section Y is, by definition, a *special Legendrian* submanifold of the Sasaki–Einstein manifold S. Equivalently, $\eta|_Y = 0$ and $\operatorname{Re} \Psi_t|_Y = 0$, where $\Psi_t = \partial_r \lrcorner (e^{-it}\widehat{\Psi})|_{r=1}$ is the horizontal volume form, cf. [47, Prop. 3.3]. (As before, η is the contact form on S.) Since Y is associative with respect to the G₂-structure φ_t , Y is a minimal submanifold for the Sasaki–Einstein metric on S. Conversely, it is known that an oriented Legendrian submanifold $Y \subset S$ is minimal if and only if Y is special Legendrian [47, Prop. 3.2].

The following result will be useful for producing examples of associative 3-folds.

Proposition 7.3 ([47, Prop. 3.4]). Let S be a Sasaki–Einstein (2n+1)-manifold and let $\tau_S : S \times \mathbb{R}^* \to S \times \mathbb{R}^*$ be an anti-holomorphic involution for the complex structure of the Kähler–Einstein cone $S \times \mathbb{R}^*$ fixing the coordinate $r \in \mathbb{R}^+$. If the fixed point set C_{τ} of τ is not empty, then the link $C_{\tau} \cap (S \times \{1\})$ is a special Legendrian submanifold of S.

In the case of a regular Sasaki–Einstein manifold S we have the following.

Theorem 7.4. Let S be a regular Sasaki–Einstein 7-manifold with contact form η arising from a principal S¹-bundle $\pi : S \to X$ with Euler class $c_1 = [\omega]$, where X is a Kähler–Einstein Fano 3-fold with Kähler form ω and $d\eta = \pi^*(\omega)$. Let φ_t be the corresponding 1-parameter family of nearly parallel G₂-forms defined in (2.6).

Then for each compact special Legendrian submanifold $Y \subset S$, the restriction $\pi|_Y : Y \to Y_X$ is a finite covering of a Lagrangian submanifold $Y_X \subset X$.

If $Y_X \subset X$ is a compact simply-connected Lagrangian submanifold, thus a Lagrangian 3-sphere, then Y_X lifts to an S¹-family of Legendrian submanifolds $Y_s \subset S$ such that $\pi(Y_s) = Y_X$ for each $s \in S^1$.

Assume that $\tau : X \to X$ is an isometric anti-holomorphic involution. If the fixed point set $Y_X \subset X$ of τ is non-empty, then Y_X is Lagrangian and diffeomorphically lifts to a special Legendrian (hence associative) submanifold of S.

Proof. If a submanifold Y is Legendrian, i.e. $\eta|_Y = 0$, then Y cannot be tangent to any fibre in the principal circle bundle S and π maps Y locally diffeomorphically onto the image $\pi(Y) = Y_X$ which is a submanifold of X. Since Y, hence also Y_X , are compact we obtain that $\pi|_Y$ is a finite cover. Considering (7.2) we find that Y_X is Lagrangian submanifold of X, $\omega|_{Y_X} = 0$.

24 MARISA FERNÁNDEZ, ANNA FINO, ALEXEI KOVALEV, AND VICENTE MUÑOZ

For the second claim, since $c_1 = [\omega]$ vanishes on Y_X and $d\eta = \pi^* \omega$, the connection η on the S^1 -bundle S restricts to a flat connection over Y_X . Furthermore, $\eta|_{Y_X}$ must be a trivial product connection as Y_X is simply connected. So $\pi^{-1}(Y_X) \cong Y_X \times S^1$ with $\eta|_{Y_X}$ corresponding to $ds, s \in S^1$, and $Y_s = Y_X \times \{s\}$ for each s, is a Legendrian submanifold of S.

The principal bundle S is associated to the anticanonical bundle K_X^{-1} and $S \times \mathbb{R}^*$ is biholomorphic to the complement of the zero section. The hypotheses on τ imply that $\tau^* \omega_X = -\omega_X$ and $\tau^* \overline{K_X^{-1}}$ is a holomorphic line bundle isomorphic to K_X^{-1} . We find that τ^* defines a lift of τ to an antiholomorphic involution on K_X^{-1} and (by restriction) on $S \times \mathbb{R}^+$ with $\tau^* \eta = -\eta$ and $\tau \circ r = r$. Thus τ^* preserves the Kähler-Einstein metric on $S \times \mathbb{R}^+$ and the fixed point set of τ^* is an oriented (hence trivial) real line bundle over Y_X . The desired special Legendrian is obtained as the intersection with $S = S \times \{1\}$ by application of Proposition 7.3.

If the Fano 3-fold X also appears as a smooth fibre in a Lefschetz fibration $\lambda : E \to \Delta$ over a disc with $0 \in \Delta$ the only critical value, then the vanishing cycles in X (the cycles that degenerate to a point as X is deformed to a fibre over 0) are represented by Lagrangian 3-spheres (see [4] or [53, § 4]).

In the case when the nearly parallel G_2 -manifold S is Q(1, 1, 1) or S_3 (see Theorem 4.2) we can say more.

Proposition 7.5. Let $\pi_M : M = Q(1, 1, 1) \to X = S^2 \times S^2 \times S^2$ be the principal S^1 bundle (6.2) and φ_t the 1-dimensional family of nearly parallel G₂-structures (6.11) on M. Let $L \subset X$ be a 3-torus defined by $\theta_j = \pi/2$, j = 1, 2, 3, in the spherical coordinates ϕ_j, θ_j on X (see Section 6).

Then L lifts via π_M to a family of minimal Legendrian 3-tori $L_s \subset M$, $s \in \mathbb{R}/2\pi\mathbb{Z}$. For each s, the 3-torus L_s is associative with respect to φ_t for all t.

As X and hence Q(1,1,1) are toric manifolds, the existence of a compact special Legendrian submanifold in Q(1,1,1) follows from the main result in [47]. However, the explicit examples of special Legendrians in Q(1,1,1) are not considered in [47].

Proof of Proposition 7.5. It follows at once from the expression (6.5) for the Kähler form that the 3-torus L is Lagrangian in X. We know from (6.7) that ω is proportional to the curvature of the principal S^1 -bundle M, thus the restriction of this bundle to L has first Chern class zero and is a trivial bundle. Furthermore, with $\theta_j = \pi/2$ the 1-form η in (6.6) defines a trivial product connection on $\pi_M^{-1}(L)$. Thus L lifts via π to a family of horizontal 3-tori $L_s \subset M$, parametrised by $s \in S^1$.

When $s + t = \pi/2$, from (6.11) and (6.9) we obtain that $\varphi_t|_{L_s} = \Psi_-|_{L_s} = -\alpha_2 \wedge \alpha_4 \wedge \alpha_6$, whence L_s (with appropriate orientation) is φ_t -associative and minimal Legendrian. Then L_s is φ_t -associative, for all t, by application of Proposition 7.2. \Box

The Fano 3-fold $X_3 = P_3 \times \mathbb{CP}^1$ is toric because P_3 is so. Recall that the del Pezzo surface is the blow-up $P_3 = \mathbb{CP}^2 \# 3\mathbb{CP}^2$ and is, up to isomorphism, independent of the choice of three non-colinear points in \mathbb{CP}^2 . Also, P_3 admits a Kähler–Einstein metric ([54] or [55]) which can be taken to be invariant under the torus action. (The induced nearly parallel G₂-structure φ_t of (2.6) on S_3 therefore is also torus-invariant for all t.)

We construct a special Legendrian associative 3-fold in S_3 as a fixed point set of an anti-holomorphic involution. The existence of this special Legendrian is of course again a special case of the main result in [47], but is rather simpler than in a general case.

The complex surface P_3 can be identified as the simultaneous solution of two complex bilinear equations

$$\{(z,w) \in \mathbb{CP}^2 \times \mathbb{CP}^2 : z_0 w_0 = z_1 w_1 = z_2 w_2\}$$
(7.3)

and the blow-up $P_3 \to \mathbb{CP}^2$ is the restriction of the first projection. The effective torus action on P_3 is given by $(z_0, z_1, z_2, w_0, w_1, w_2) \mapsto (\xi_0 z_0, \xi_1 z_1, \xi_2 z_2, \xi_0^{-1} w_0, \xi_1^{-1} w_1, \xi_2^{-1} w_2)$, where $\xi_i \in \mathbb{C}^*$, $\xi_0 \xi_1 \xi_2 = 1$, and this can be interpreted as an embedding of $(\mathbb{C}^*)^2$ in P_3 as an open dense subset. We deduce that in the notation of (7.3) the map $\tau_3(z_i, w_i) = (\bar{w}_i, \bar{z}_i)$ induces an antiholomorphic isometry of the Kähler–Einstein P_3 . The fixed point set of this involution is diffeomorphic to the 2-torus [57].

Extending to an antiholomorphic involution of the Fano 3-fold X_3 with the complex conjugation on the \mathbb{CP}^1 -factor, we obtain, by application of Theorem 7.4.

Proposition 7.6. There exists an associative 3-torus in the nearly parallel G_2 -manifold (S_3, φ_t) .

8. Associative 3-folds in the Aloff–Wallach spaces

We now revisit the Aloff–Wallach spaces $W_{k,l}$, discussed in Subsection 5.2, and find examples of associative submanifolds in $W_{k,l}$. Recall that k, l are non-zero, co-prime integers. We begin by briefly recalling from [8, 20] the construction of proper nearly parallel G₂-structures on $W_{k,l}$.

Let $\{e_i\}_{i=0,\ldots,7}$ be a basis of left-invariant vector fields on SU(3) such that e_0 is everywhere tangent to the orbits of the $S_{k,l}^1$ action. We shall interchangeably think of e_i as elements of the Lie algebra $\mathfrak{su}(3)$ with $e_0 \in \mathfrak{s}_{k,l}^1$. Choose $\{e_i\}$ so that writing $\{e^i\}$ for the dual basis of $\mathfrak{su}(3)^*$ (or the left-invariant 1-forms) the Maurer-Cartan form $\Omega = \sum_i e_i e^i$ on SU(3) with values in $\mathfrak{su}(3)$ is expressed as

$$\Omega = \frac{1}{\sqrt{2}} \begin{pmatrix} i(\frac{k}{\sqrt{3}s}e^0 + \frac{l-m}{3s}e^4) & e^1 + ie^5 & -e^3 + ie^7 \\ -e^1 + ie^5 & i(\frac{l}{\sqrt{3}s}e^0 + \frac{m-k}{3s}e^4) & e^2 + ie^6 \\ e^3 + ie^7 & -e^2 + ie^6 & i(\frac{m}{\sqrt{3}s}e^0 + \frac{k-l}{3s}e^4) \end{pmatrix},$$
(8.1)

where m = -k - l and $s = \sqrt{(k^2 + l^2 + m^2)/6}$.

Then for any choice of non-zero constants A, B, C, D the 3-form

$$\varphi_W = ABC(e^{123} - e^{167} + e^{257} - e^{356}) - D(A^2e^{15} + B^2e^{26} + C^2e^{37}) \wedge e^4$$
(8.2)

descends to a well-defined positive 3-form giving a coclosed G₂-structure on $W_{k,l}$. The induced orientation corresponds to a non-vanishing 7-form $De^{1234567}$ and the coframe $Ae^1, Be^2, Ce^3, De^4, Ae^5, Be^6, Ce^7$ of 1-forms is orthonormal in the induced metric g_{φ_W} . Note that we may assume without loss of generality that A > 0 and B > 0. From the results in [20] it follows that if $\{k, l, m\}$ are not $\{1, 1, -2\}$ or $\{1, -1, 0\}$, then every homogeneous, coclosed G₂-structure on $W_{k,l}$ is of the form (8.2). Furthermore, by using the Maurer–Cartan equation $d\Omega = -\frac{1}{2}[\Omega, \Omega]$, it was proved that exactly two such G₂-structures are nearly parallel, more precisely, these are proper nearly parallel.

Recall from Subsection 5.2 that each generic Aloff–Wallach space $W_{k,l}$ has a structure of a smooth fibre bundle $\pi_{k,l} : W_{k,l} \to \mathbb{CP}^2$ with typical fibre the spherical space form $S^3/\mathbb{Z}_{|k+l|}$. This fibration is not unique. The Weyl group of SU(3) contains an element of order 3 which induces a diffeomorphism $\upsilon : W_{k,l} \to W_{l,m}$. The composition $\pi_{l,m} \circ \upsilon$ defines a different fibration, in general by different spherical space forms.

Theorem 8.1. Let φ_W be a nearly parallel G₂-structure of the form (8.2) on an Aloff– Wallach space $W_{k,l}$, with k, l non-zero and co-prime integers such that $\{k, l, m\}$ are not $\{1, 1, -2\}$ or $\{1, -1, 0\}$. Then the fibres of $\pi_{k,l}$ are embedded minimal associative 3-folds with respect to φ_W .

Furthermore, for suitably 'generic' k, l, the Aloff–Wallach space $W_{k,l}$ has three different 4-dimensional deformation families of minimal associative spherical space forms.

Proof. The fibre bundle $\pi_{k,l}$ can be obtained by considering the embedding of U(2) as a Lie subgroup of SU(3) consisting of the block-diagonal matrices with blocks $Ae^{i\theta}$ and $1/\det(Ae^{i\theta}) = e^{-2i\theta}$, for all $A \in SU(2)$ and $\theta \in \mathbb{R}$. Then

$$\pi_{k,l}: W_{k,l} = SU(3)\S^1_{k,l} \to SU(3)/U(2) \cong \mathbb{CP}^2$$

between respective cosets and the fibres is $U(2)/S_{k,l}^1 \cong S^3/\mathbb{Z}_{|k+l|}$.

Comparing the tangent spaces to the cosets of U(2) in SU(3) with (8.1), we find that the vertical spaces of the principal U(2)-bundle SU(3) $\rightarrow \mathbb{CP}^2$ are defined by the vanishing of e^2, e^3, e^6, e^7 . Thus φ_W restricts on each fibre of $\pi_{k,l}$ to the volume form $A^2 D e^{145}$ of the metric induced by g_{φ_W} , thus the fibres are associative.

We next show that the fibres of $\pi_{l,m} \circ \upsilon$ are also minimal associative for the same G_2 -structure φ_W .

To this end, we further note that $\pi_{k,l}$ factors through the flag manifold $F(1,2) \cong$ SU(3)/ T^2 (where $T^2 \subset$ SU(3) is the subgroup of diagonal matrices),

$$W_{k,l} \to F(1,2) \to \mathbb{CP}^2.$$
 (8.3)

Here the first map is a principal S^1 -bundle and the second map is a \mathbb{CP}^1 -bundle associated to the principal U(2)-bundle SU(3) $\rightarrow \mathbb{CP}^2$ discussed above. In particular, the vertical space of the S^1 -bundle is spanned by e_4 .

It is well-known that F(1, 2) is a complex manifold with a Kähler structure defined by the Kirillov–Kostant–Souriau symplectic form. We can consider a different choice of the \mathbb{CP}^1 -bundle $h_3 : F(1,2) \to \mathbb{CP}^2$ corresponding to a different embedding of U(2) as a Lie subgroup of SU(3), with the tangent space at the identity now spanned by $e_0, e_3, e_4, e_7 \in \mathfrak{su}(3)$ (rather than e_0, e_1, e_4, e_5 chosen above). Let Y be a fibre of the map $W_{k,l} \to F(1,2) \xrightarrow{h_3} \mathbb{CP}^2$. Then the forms e^1, e^5, e^2, e^6 vanish on Y and so $\varphi_W|_Y = C^2 D e^{347}$. Thus Y is an embedded minimal associative 3-fold with respect to a G₂-structure φ_W of the form (8.2) on $W_{k,l}$.

The third family of associative 3-folds follows by a similar argument, replacing the \mathbb{CP}^1 -bundle h_3 with $h_2: F(1,2) \to \mathbb{CP}^2$ where the vertical spaces are now spanned by e_0, e_2, e_4, e_6 .

Remark 8.2. The exceptional Aloff–Wallach space $W_{1,-1}$ has, up to homotheties, only one homogeneous nearly parallel G₂-structure, which is of the form (8.2). The argument and result of Theorem 8.1 applies to this latter G₂-structure noting that the fibres of $\pi_{1,-1}$ are now $S^2 \times S^1$ as $S^1_{1,-1}$ is a subgroup of SU(2).

When the G_2 -structure φ_W in Theorem 8.1 is *not* nearly parallel, the fibres of $\pi_{k,l}$ are associative 3-folds but need not be minimal.

Remark 8.3. For the exceptional Aloff–Wallach space $W_{1,1}$ there are still exactly two homogeneous Einstein metrics [50]. However, one of these metrics is induced by a proper nearly parallel G_2 -structure on $W_{1,1}$ which is *not* of the form (8.2), whereas the other metric is induced by a 3-Sasakian structure on $W_{1,1}$. For this latter 3-Sasakian manifold, the above construction of minimal associative 3-folds based on (8.3) remains valid. On the other hand, very recently Ball and Madnick constructed in $W_{1,1}$ examples of associative 3-folds diffeomorphic to an S^1 -bundle over a genus g surface for all $g \ge 0$ [7, Theorem 5.9].

9. LOCALLY CONFORMAL PARALLEL Spin(7)-STRUCTURES

In this section we give examples of formal compact 8-manifolds, with a locally conformal parallel Spin(7)-structure.

An 8-dimensional manifold N has a Spin(7)-structure if there is a reduction of the structure group of its frame bundle from $GL(8, \mathbb{R})$ to the exceptional Lie group

27

Spin(7). In opposite to the existence of G_2 -structures on any orientable and spin manifold of dimension 7, it happens that not every 8-dimensional spin manifold N admits a Spin(7)-structure. In fact, in [45] it is proved that N has a Spin(7)-structure if and only if

$$p_1^2(N) - 4 p_2(N) + 8\chi(N) = 0,$$

for an appropriate choice of the orientation, where $p_1(N)$, $p_2(N)$ and $\chi(N)$ are the first Pontryagin class, the second Pontryagin class and the Euler characteristic of N, respectively.

The presence of a Spin(7)-structure is equivalent to the existence of a nowhere vanishing global differential 4-form Ω on the 8-manifold N, which can be written locally as

$$\Omega = \left(e^{127} + e^{347} + e^{567} + e^{135} - e^{146} - e^{236} - e^{245}\right) \wedge e^{8} + e^{1234} + e^{1256} + e^{1367} + e^{1457} + e^{2357} + e^{2467} + e^{3456},$$

with respect to some (local) basis $\{e^1, \ldots, e^8\}$ of the (local) 1-forms on N.

Since $\text{Spin}(7) \subset \text{SO}(8)$, a Spin(7)-structure Ω on N determines a Riemannian metric g_{Ω} and an orientation on N such that

$$g_{\Omega}(U,V)$$
 vol $= \frac{1}{7}(U \lrcorner \Omega) \land \star_8(V \lrcorner \Omega),$

for any vector fields U, V on N, where vol is the volume form on N, and \star_8 is the Hodge star operator determined by g_{Ω} . Note that if Ω is a Spin(7)-structure on N, then Ω is a self-dual 4-form, i.e. $\star_8 \Omega = \Omega$.

Let us recall that a Spin(7)-structure Ω on a 8-manifold N is said to be *parallel* if the induced metric by Ω has holonomy contained in Spin(7). This is equivalent to say that the 4-form Ω is parallel with respect to the Levi–Civita connection of the metric g_{Ω} , which happens if and only if $d\Omega = 0$ [24].

Definition 9.1. A Spin(7)-structure Ω on a 8-manifold N is said to be *locally con*formal parallel if

$$d\Omega = \Omega \wedge \Theta, \tag{9.1}$$

for a closed non-vanishing 1-form Θ , which is known as the *Lee form* of the Spin(7)structure. A manifold endowed with such a structure is called *locally conformal parallel* Spin(7)-manifold.

Let φ be a nearly parallel G₂-structure on a 7-manifold P, so that $d\varphi = \tau_0 \star \varphi$ by (2.2). Then, the product manifold $P \times S^1$ carries a natural locally conformal parallel Spin(7)-structure Ω defined by

$$\Omega = \varphi \wedge \theta + \star \varphi, \tag{9.2}$$

where θ is the volume form on S^1 , and \star is the Hodge star operator determined by the induced metric by φ on P. Note that the Lee form Θ is given by $\Theta = \tau_0 \theta$. In fact, using (9.2), we have $d\Omega = \tau_0 \star \varphi \wedge \theta = \tau_0 \Omega \wedge \theta$, that is $d\Omega = \Omega \wedge \Theta$.

By [13, Theorem 3.2], we know that Q(1,1,1) is non-formal. Hence the product manifold $Q(1,1,1) \times S^1$ is non-formal by [21]. Then, if we consider the S^1 -family of nearly parallel G₂-structures φ_t defined in (6.11), we have:

Proposition 9.2. The product manifold $Q(1,1,1) \times S^1$ is non-formal and has an S^1 -family of locally conformal parallel Spin(7)-structures $\Omega_t = \varphi_t \wedge \theta + \star \varphi_t$, with Lee form $\Theta = 4 \theta$.

Let P be a differentiable manifold and let $\rho : P \to P$ be a diffeomorphism. The mapping torus P_{ρ} of ρ is the manifold obtained from $P \times [0, 1]$ by identifying the ends with ρ , that is

$$P_{\rho} = \frac{P \times [0, 1]}{(x, 0) \sim (\rho(x), 1)}.$$

The natural map $\pi: P_{\rho} \to S^1$ defined by $\pi(x,t) = e^{2\pi i t}$ is the projection of a locally trivial fiber bundle (here we think of S^1 as the interval [0,1] with identified end points). Thus, any ρ -invariant form β on P defines a form on P_{ρ} , since the pullback of β to $P \times \mathbb{R}$ is invariant under the diffeomorphism $(x,t) \mapsto (\rho(x), t+1)$. For the same reason, the 1-form dt on \mathbb{R} , where t is the coordinate on \mathbb{R} , induces a closed 1-form ν on P_{ρ} .

A theorem of Tischler [56] asserts that a compact manifold is a mapping torus if and only if it admits a non-vanishing closed 1-form. This result was extended to locally conformal parallel Spin(7) manifolds in [37, Theorem B]. There, it is proved that there exists a fibre bundle $N \to S^1$ with abstract fibre P/Γ , where P is a compact simply connected nearly parallel G₂-manifold and Γ is a finite subgroup of isometries of P acting freely. Moreover, the cone of P/Γ covers N with cyclic infinite covering transformation group.

Notice that if P is a 7-dimensional compact manifold endowed with a nearly parallel G_2 -structure φ , and $\rho : P \to P$ is a diffeomorphism such that $\rho^* \varphi = \varphi$, then ρ preserves the orientation and the metric on P induced by the G_2 -structure φ . So, $\rho^*(\star \varphi) = \star \varphi$, and the mapping torus $N = P_{\rho}$ has a locally conformal parallel Spin(7)-structure Ω given by $\Omega = \varphi \wedge \nu + \star \varphi$.

Let us recall that if P is a differentiable manifold and $\rho : P \to P$ is a diffeomorphism, then the cohomology of the mapping torus $N = P_{\rho}$ of ρ sits in an exact sequence [11, Lemma 12]

$$0 \to C^{r-1} \to H^r(N) \to K^r \to 0,$$

where K^r is the kernel of $\varphi^* - \mathrm{Id} \colon H^r(P) \to H^r(P)$, and C^r is its cokernel. Thus,

$$H^{r}(P_{\rho}) \cong \ker \left(\varphi^{*} - \operatorname{Id} : H^{r}(P) \to H^{r}(P)\right)$$
$$\oplus \left[\nu\right] \wedge \frac{H^{r-1}(P)}{\operatorname{Im}\left(\varphi^{*} - \operatorname{Id} : H^{r-1}(P) \to H^{r-1}(P)\right)}$$

Now we consider the manifold M = Q(1, 1, 1) with the family of nearly parallel G₂-structures φ_t defined by (6.11). Let ρ be the diffeomorphism of Q(1, 1, 1) given by

$$\rho(\theta_1, \phi_1, \theta_2, \phi_2, \theta_3, \phi_3, s) = (\theta_2, \phi_2, -\theta_1, \phi_1, \theta_3, \phi_3, s)$$

Then, the diffeomorphism ρ on the 1-forms $\{\alpha_1, \ldots, \alpha_6, \alpha_7 = \zeta\}$ on Q(1, 1, 1) is given by

$$\rho^* \alpha_1 = \alpha_3, \quad \rho^* \alpha_2 = \alpha_4, \quad \rho^* \alpha_3 = -\alpha_1, \quad \rho^* \alpha_4 = -\alpha_2, \quad \rho^* \alpha_i = \alpha_i, \ i = 5, 6, 7.$$

Proposition 9.3. The mapping torus $M_{\rho} = Q(1,1,1)_{\rho}$ is formal and it has an S^1 family of locally conformal parallel Spin(7)-structures.

Proof. Clearly the diffeomorphism ρ preserves the nearly parallel G₂-structures φ_t defined in (6.11) and, taking into account (6.12), ρ preserves also $\star \varphi_t$. Then, by [37, Theorem B], we know that $M_{\rho} = Q(1, 1, 1)_{\rho}$ carries a S¹-family of locally conformal parallel Spin(7)-structures.

In order to prove that $M_{\rho} = Q(1,1,1)_{\rho}$ is formal we proceed as follows. As in Section 6, we write with the same symbol the lifting to M of the 1-forms α_i , $1 \le i \le 6$, and of the Kähler forms ω_j , $1 \le j \le 3$, where $[\omega_1]$, $[\omega_2]$ and $[\omega_3]$ are the generators of the integral cohomology group of each of the S^2 -factors on $S^2 \times S^2 \times S^2$. Now, for simplicity of notation we write $a_1 = [\omega_1]$, $a_2 = [\omega_2]$ and $a_3 = [\omega_3]$. Since M = Q(1, 1, 1)is the principal S^1 -bundle

$$S^1 \longrightarrow M = Q(1, 1, 1) \longrightarrow S^2 \times S^2 \times S^2$$

with first Chern class equal to $a_1 + a_2 + a_3$, the Gysin sequence gives that

$$\begin{aligned} H^{0}(M,\mathbb{Z}) &= H^{7}(M,\mathbb{Z}) = \mathbb{Z}, \\ H^{1}(M,\mathbb{Z}) &= H^{3}(M,\mathbb{Z}) = H^{6}(M,\mathbb{Z}) = 0, \\ H^{2}(M,\mathbb{Z}) &= H^{5}(M,\mathbb{Z}) = \mathbb{Z}^{2}, \\ H^{4}(M,\mathbb{Z}) &= \mathbb{Z}\langle a_{1}a_{2}, a_{1}a_{3}, a_{2}a_{3} \rangle / \langle a_{1}a_{2} + a_{1}a_{3}, a_{2}a_{1} + a_{2}a_{3}, a_{3}a_{1} + a_{3}a_{2} \rangle = \mathbb{Z}_{2}. \end{aligned}$$

So, the de Rham cohomology groups of M = Q(1, 1, 1) up to the degree 4 are

$$H^{0}(M) = \langle 1 \rangle, \quad H^{1}(M) = H^{3}(M) = H^{4}(M) = 0, \quad H^{2}(M) = \langle a_{1}, a_{2} \rangle.$$

Since $\rho^*\omega_1 = \omega_2$ and $\rho^*\omega_2 = \omega_1$, the de Rham cohomology groups of the mapping torus $M_{\rho} = Q(1, 1, 1)_{\rho}$ up to the degree 4 are

$$H^{0}(M_{\rho}) = \langle 1 \rangle, \quad H^{1}(M_{\rho}) = \langle [\nu] \rangle, \quad H^{2}(M_{\rho}) = \langle a_{1} + a_{2} \rangle,$$
$$H^{3}(M_{\rho}) = \langle (a_{1} + a_{2}) \wedge [\nu] \rangle, \quad H^{4}(M_{\rho}) = 0.$$

Therefore, the minimal model of M_{ρ} must be a differential graded algebra $(\bigwedge V, d)$, being $\bigwedge V$ the free algebra of the form $\bigwedge V = \bigwedge (a, b, x) \otimes \bigwedge V^{\geq 4}$, where |a| = 1, |b| = 2, |x| = 3, and d is defined by da = 0 = db, $dx = b^2$. According to Definition 3.1, we get $N^j = 0$ for j = 1, 2, thus M_{ρ} is 2-formal. Moreover, M_{ρ} is 3-formal. In fact, take $z \in I(N^{\leq 3})$ a closed element in $\bigwedge V$. As $H^*(\bigwedge V) = H^*(M_{\rho})$ has non-zero cohomology in degrees 0, 1, 2, 3, 5, 6, 7, 8, it must be deg z = 5, 6, 7, 8. If deg z = 5then $z = b \cdot x$ (up to non-zero scalar), which is not closed because $d(b \cdot x) = b^3 \neq 0$. If deg z = 6 then $z = a \cdot b \cdot x$ which is not closed because $d(a \cdot b \cdot x) = -a \cdot b^3 \neq 0$. If deg z = 7 then $z = b^2 \cdot x$ which is not closed, and if deg z = 8 then $z = a \cdot b^2 \cdot x$ which is not closed either. Thus, according to Definition 3.1, M_{ρ} is 3-formal and, by Theorem 3.2, M_{ρ} is formal.

Next, for $M = \mathcal{B}$ or $M = W_{k,l}$, where $\mathcal{B} = SO(5)/SO(3)$ is the Berger space and $W_{k,l} = SU(3)/S_{k,l}^1$ are the the Aloff–Wallach spaces, we study the formality of the mapping torus of a diffeomorphism of M preserving a nearly-parallel G₂-structure on M (see subsections 5.1 and 5.2).

Proposition 9.4. Let $\rho : \mathbb{B} \to \mathbb{B}$ be a diffeomorphism preserving a nearly-parallel G_2 -structure on \mathbb{B} . Then, the mapping torus \mathbb{B}_{ρ} is formal and it has a locally conformal parallel Spin(7)-structure.

Proof. We consider a nearly parallel G_2 -structure on \mathcal{B} , which is preserved by the diffeomorphism $\rho : \mathcal{B} \to \mathcal{B}$. (As we already mentioned in subsection 5.1, the only known nearly parallel G_2 -structure on \mathcal{B} is proper and is explicitly given in [6].) Theorem B in [37] implies that \mathcal{B}_{ρ} has a locally conformal parallel Spin(7)-structure.

According to the proof of Theorem 5.1, we know that \mathcal{B} and the 7-sphere S^7 have the same minimal model. Hence, the de Rham cohomology groups of \mathcal{B} up to the degree 6 are

$$H^0(\mathfrak{B}) = \langle 1 \rangle, \quad H^k(\mathfrak{B}) = 0, \quad 1 \le k \le 6.$$

So, the de Rham cohomology groups of the mapping torus \mathcal{B}_{ρ} up to the degree 6 are

$$H^0(\mathcal{B}_{\rho}) = \langle 1 \rangle, \quad H^1(\mathcal{B}_{\rho}) = \langle [\nu] \rangle, \quad H^k(\mathcal{B}_{\rho}) = 0, \quad 2 \le k \le 6.$$

Therefore, the minimal model of \mathcal{B}_{ρ} must be a differential graded algebra $(\bigwedge V, d)$, being $\bigwedge V$ the free algebra of the form $\bigwedge V = \bigwedge (a) \otimes \bigwedge V^{\geq 7}$, with |a| = 1, and d is given by da = 0. Now, we have $C^1 = \langle a \rangle$, $C^i = 0$ for $2 \leq i \leq 4$, and $N^j = 0$ for $1 \leq j \leq 6$. Thus, Definition 3.1 implies that \mathcal{B}_{ρ} is 3-formal and, by Theorem 3.2, \mathcal{B}_{ρ} is formal. **Proposition 9.5.** Let $\rho: W_{k,l} \to W_{k,l}$ be a diffeomorphism preserving a nearly parallel G₂-structure (not necessarily proper) on $W_{k,l}$. Then, the mapping torus $(W_{k,l})_{\rho}$ is formal and it has a locally conformal parallel Spin(7)-structure.

Proof. If $\rho: W_{k,l} \to W_{k,l}$ is a diffeomorphism preserving a nearly parallel G₂-structure on $W_{k,l}$, the mapping torus $(W_{k,l})_{\rho}$ has a locally conformal parallel Spin(7)-structure by [37, Theorem B].

In order to prove that $(W_{k,l})_{\rho}$ is formal we proceed as follows. By Theorem 5.2 and Theorem 5.3, we know that $W_{k,l}$ and the product manifold $S^2 \times S^5$ have the same minimal model, and hence the same de Rham cohomology. So, the de Rham cohomology groups of $W_{k,l}$ are:

$$H^{0}(W_{k,l}) = \langle 1 \rangle, \quad H^{1}(W_{k,l}) = H^{3}(W_{k,l}) = H^{4}(W_{k,l}) = H^{6}(W_{k,l}) = 0,$$

$$H^{2}(W_{k,l}) = \langle \xi \rangle, \quad H^{5}(W_{k,l}) = \langle \tau \rangle, \quad H^{7}(W_{k,l}) = \langle \xi \wedge \tau \rangle.$$

Let us consider the map $\rho^* : H^2(W_{k,l}) \to H^2(W_{k,l})$. It is clear that either $\rho^*(\xi) \neq \xi$ or $\rho^*(\xi) = \xi$.

We deal first with the possibility that $\rho^*(\xi) \neq \xi$. Then, the cohomology of $(W_{k,l})_{\rho}$ up to the degree 4 is

$$H^{0}((W_{k,l})_{\rho}) = \langle 1 \rangle, \qquad H^{1}((W_{k,l})_{\rho}) = \langle [\nu] \rangle, \qquad H^{i}((W_{k,l})_{\rho}) = 0, \ 2 \le i \le 4.$$

Therefore, the minimal model of $(W_{k,l})_{\rho}$ must be a differential graded algebra $(\bigwedge V, d)$, being $\bigwedge V$ the free algebra of the form $\bigwedge V = \bigwedge(a) \otimes \bigwedge V^{\geq 5}$, where |a| = 1, and dis defined by da = 0. According to Definition 3.1, we get $C^1 = \langle a \rangle$, $C^i = 0$ for $2 \leq i \leq 4$, and $N^j = 0$ for $1 \leq j \leq 4$. Hence, by Definition 3.1, $(W_{k,l})_{\rho}$ is 3-formal and, by Theorem 3.2, $(W_{k,l})_{\rho}$ is formal.

Suppose now that $\rho^*(\xi) = \xi$. In this case, the cohomology of $(W_{k,l})_{\rho}$ up to the degree 4 is

$$H^{0}((W_{k,l})_{\rho}) = \langle 1 \rangle, \qquad H^{1}((W_{k,l})_{\rho}) = \langle [\nu] \rangle, \qquad H^{2}((W_{k,l})_{\rho}) = \langle \xi \rangle,$$

$$H^{3}((W_{k,l})_{\rho}) = \langle [\nu] \wedge \xi \rangle, \qquad H^{4}((W_{k,l})_{\rho}) = 0.$$

Thus, the minimal model of $(W_{k,l})_{\rho}$ must be a differential graded algebra $(\bigwedge V, d)$, being $\bigwedge V$ the free algebra of the form $\bigwedge V = \bigwedge (a, b, x) \otimes \bigwedge V^{\geq 4}$, where |a| = 1, |b| = 2, |x| = 3, and d is defined by da = db = 0 and $dx = b^2$. According to Definition 3.1, we get $N^j = 0$ for $j \leq 2$, thus $(W_{k,l})_{\rho}$ is 2-formal. Moreover, $(W_{k,l})_{\rho}$ is 3-formal. In fact, take $\alpha \in I(N^{\leq 3})$ a closed element in $\bigwedge V$. As $H^*(\bigwedge V) = H^*((W_{k,l})_{\rho})$ has cohomology in all the degrees except in degree 4, and since $|\alpha| \geq 4$, it must be $|\alpha| = 5, 6, 7, 8$. If $|\alpha| = 5$, then $\alpha = b \cdot x$ (up to non-zero scalar), which is not closed. If $|\alpha| = 6$, then $\alpha = a \cdot b \cdot x$, which is not closed either. If $|\alpha| = 7$, then $\alpha = b^2 \cdot x$, and if $|\alpha| = 8$, then $\alpha = a \cdot b^2 \cdot x$, but α is not closed in either case. So, according to Definition 3.1, $(W_{k,l})_{\rho}$ is 3-formal and, by Theorem 3.2, $(W_{k,l})_{\rho}$ is formal.

Acknowledgements. We are very grateful to Dieter Kotschick and the anonymous referees for useful comments. The first author was partially supported by the Basque Government Grant IT1094-16 and by the Grant PGC2018-098409-B-100 of the Ministerio de Ciencia, Innovación y Universidades of Spain. The second author is supported by the project PRIN "Real and complex manifolds: Topology, Geometry and Holomorphic Dynamics", by GNSAGA of INdAM and by a grant from the Simons Foundation (#944448). The fourth author was partially supported by Project MINECO (Spain) PID2020-118452GB-I00.

References

- B, Alexandrov, U. Semmelmann, Deformations of nearly parallel G₂-structures, Asian J. Math. 16 (2012), no. 4, 713–744.
- [2] S. Aloff, N. Wallach, An infinite family of distinct 7-manifolds admitting positively curved Riemannian structures, Bull. Amer. Math. Soc. 81 (1975), 93–97
- [3] M. Amann, I. Taimanov, On the formality of nearly Kähler manifolds and of Joyce's examples in G₂ holonomy. Preprint arXiv:2012.10915 [math.DG].
- [4] V. Arnol'd, Some remarks on symplectic monodromy of Milnor fibrations. In The Floer memorial volume, volume 133 of Progr. Math., pages 99–103. Birkhäuser, Basel, 1995.
- [5] G. Ball, J. Madnick, The Mean Curvature of First-Order Submanifolds in Exceptional Geometries with Torsion, Ann. Global Anal. and Geom. 59 (2021), no. 1, 27–67.
- [6] G. Ball, J. Madnick, Associative submanifolds of the Berger space. Preprint arXiv: 2003.13169 [math.DG], to appear in Comm. Anal. Geom..
- [7] G. Ball, J. Madnick, Associative Submanifolds of Squashed 3-Sasakian Manifolds. Preprint arXiv: 2208.10622 [math.DG].
- [8] G. Ball, G. Oliveira, Gauge theory on Aloff-Wallach spaces, Geom. Topol. 23 (2019), no. 2, 685–743.
- [9] H. Baum, T. Friedrich, R. Grunewald, I. Kath, Twistors and Killing spinors on Riemannian manifolds, Seminarberichte Nr. 108, Humboldt-Universität zu Berlin, 1990.
- [10] G. Bazzoni, I. Biswas, M. Fernández, V. Muñoz, A. Tralle, *Homotopic properties of Kähler orbifolds*, Special Metrics and Group Actions in Geometry (Rome, 2015), 23–58, Springer INdAM 23, 2017.
- G. Bazzoni, M. Fernández, V. Muñoz, Non-formal co-symplectic manifolds, Trans. Amer. Math. Soc. 367 (2015), 4459-4481.
- [12] M. Berger, Les variétés riemanniennes homogènes normales simplement connexes à courbure strictement positive, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) 15 (1961), 179–246.
- [13] I. Biswas, M. Fernández, V. Muñoz, A. Tralle, On formality of Sasakian manifolds, J. Topol. 9 (2016), no. 1, 161–180.
- [14] D. Blair, Riemannian geometry of contact and symplectic manifolds, Progress in Mathematics 203, Birkhäuser Boston, Inc., Boston, MA, 2002.
- [15] C. Boyer, K. Galicki, 3-Sasakian manifolds, Surveys Diff. Geom. 6 (1999), 123–184.
- [16] C. Boyer, K. Galicki, On Sasakian-Einstein geometry, Internat. J. Math. 11 (2000), 873–909.
- [17] C. Boyer, K. Galicki, Sasakian Geometry, Oxford Univ. Press, Oxford, 2007.

- [18] C. Boyer, K. Galicki, B. Mann, E. Rees, Compact 3-Sasakian 7-manifolds with arbitrary second Betti number, Invent. Math. 131 (1998), 321–344.
- [19] R. Bryant, Metrics with exceptional holonomy, Ann Math. 126 (1987), 525–576.
- [20] F. Cabrera, M. Monar, A. Swann, *Classification of G₂-structures*, J. London Math. Soc. (2) 53 (1996), no. 2, 407–416.
- [21] P. Deligne, P. Griffiths, J. Morgan, D. Sullivan, Real homotopy theory of Kähler manifolds, Invent. Math. 29 (1975), 245–274.
- [22] C. Escher, W. Ziller, Topology of non-negatively curved manifolds, Ann. Global Anal. Geom. 46 (2014), no. 1, 23-55.
- [23] Y. Félix, S. Halperin, J.-C. Thomas, Rational Homotopy Theory, Springer, 2002.
- [24] M. Fernández, A classification of Riemannian manifolds with structure group Spin(7), Ann. Mat. Pura Appl. 143 (1986), 101–122.
- [25] M. Fernández, S. Ivanov, V. Muñoz, Formality of 7-dimensional 3-Sasakian manifolds, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5) 19 (2019), no. 1, 297–309.
- [26] M. Fernández, V. Muñoz, Formality of Donaldson submanifolds, Math. Z. 250 (2005), 149–175.
- [27] M. Fernández, V. Muñoz, J. Sánchez, On SO(3)-bundles over the Wolf spaces, Rev. Mat. Iberoamericana 36 (2020), 159–193.
- [28] T. Friedrich, I. Kath, 7-Dimensional compact Riemannian manifolds with Killing spinors, Comm. Math. Phys. 133 (1990), 543–561.
- [29] T. Friedrich, I. Kath, A. Moroianu, U. Semmelmann, On nearly parallel G₂-structures, J. Geom. Phys. 23 (1997), 259–286.
- [30] K. Galicki, S. Salamon, On Betti Numbers of 3-Sasakian Manifolds, Geom. Dedicata 63 (1996), 45–68.
- [31] S. Goette, N. Kitchloo, K. Skankar, Diffeomorphism type of the Berger space SO(5)/SO(3), Amer. J. Math. 126 (2004), 395–416.
- [32] A. Gray, Vector cross products on manifolds, Trans. Amer. Math. Soc. 141 (1969), 465–504.
- [33] A. Gray, Weak holonomy groups, Math. Z. 123 (1971), 290–300.
- [34] S. Gukov, J. Sparks, *M*-Theory on Spin(7)-manifolds, Nuclear Phys. B 625 (2002), no. 1-2, 3–69.
- [35] S. Halperin, Lectures on minimal models, Mém. Soc. Math. France 230, 1983.
- [36] R. Harvey, H. Lawson, Calibrated geometries, Acta Math. 148 (1982) 47–157.
- [37] S. Ivanov, M. Parton, P. Piccinni, Locally conformal parallel G₂ and Spin(7) manifolds, Math. Res. Lett. 13 (2006), 167–177.
- [38] D. Joyce, Compact manifolds with special holonomy, Oxford University Press, 2000.
- [39] K. Kawai, Some associative submanifolds of the squashed 7-sphere, Q. J. Math. 66 (2015), 861–893.
- [40] K. Kawai, Deformations of homogeneous associative submanifolds in nearly parallel G₂manifolds, Asian J. Math. 21 (2017), 429–462.
- [41] S. Kobayashi, Principal fibre bundles with the 1-dimensional toroidal group, Tohoku Math. J. (2), 8 (1956), 29–45.
- [42] D. Kotschick, On products of harmonic forms, Duke Math. J. 107 (2001), 521–531.
- [43] D. Kotschick, S. Terzić, On formality of generalised symmetric spaces, Math. Proc. Cam. Phil. Soc. 134 (2003), 491–505.
- [44] M. Kreck, S. Stolz, Some nondiffeomorphic homeomorphic homogeneous 7-manifolds with positive sectional curvature, J. Differ. Geom. 33 (1991), 465–486. A correction on Some nondiffeomorphic homeomorphic homogeneous 7-manifolds with positive sectional curvature, J. Differ. Geom. 49 (1998), 203–204.
- [45] B. Lawson, M. Michelsohn, Spin Geometry, Princeton University Press, 1989.

- [46] J. Lotay, Associative submanifolds of the 7-sphere, Proc. Lond. Math. Soc. 105 (2012), 1183– 1214.
- [47] T. Moriyama, Special Legendrian submanifolds in toric Sasaki-Einstein manifolds, New York J. Math. 21 (2015), 465-484.
- [48] P. Nagy, U. Semmelmann, The G₂ geometry of 3-Sasaki structures. Preprint arXiv: 2101.04494 [math.DG].
- [49] J. Neisendorfer, T. Miller, Formal and coformal spaces, Illinois J. Math. 22 (1978), 565–580.
- [50] Y. Nikonorov, Compact homogeneous Einstein 7-manifolds, Geom. Dedicata 109 (2004), 7–30.
- [51] D. Page, C. Pope, New squashed solutions of d = 11 supergravity, Phys. Lett. B 147 (1984) 55–60.
- [52] F. Reidegeld, Spaces admitting homogeneous G₂-structures, Differential Geom. Appl. 28 (2010), no. 3, 301–312.
- [53] P. Seidel, Floer homology and the symplectic isotopy problem. D.Phil. Thesis, Oxford 1997.
- [54] Y-T. Siu The existence of Kähler-Enstein metrics on manifolds with positive anticanonical line bundle and a suitable finite symmetry group, Ann. of Math. 127 (1988), 585–627.
- [55] G. Tian, S-T. Yau, Kähler–Einstein metrics on complex surfaces with $c_1 > 0$, Comm. Math. Phys. **112** (1987), 175–203.
- [56] D. Tischler, On fibering certain foliated manifolds over S^1 , Topology 9 (1970), 153–154.
- [57] C. Wall, Real forms of smooth del Pezzo surfaces, J. Reine Angew. Math. 375/376 (1987) 47-66.
- [58] M. Wang, Einstein metrics from symmetry and bundle constructions: a sequel, Differential geometry, 253–309, Adv. Lect. Math. 22, Int. Press, Somerville, MA, 2012.

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIA Y TECNOLOGÍA, UNIVERSIDAD DEL PAÍS VASCO (UPV/EHU), APARTADO 644, 48080 BILBAO, SPAIN

Email address: marisa.fernandez@ehu.es

DIPARTIMENTO DI MATEMATICA "GIUSEPPE PEANO, UNIVERSITÀ DI TORINO, VIA CARLO AL-BERTO 10, 10123 TORINO, ITALY, & DEPARTMENT OF MATHEMATICS AND STATISTICS, FLORIDA INTERNATIONAL UNIVERSITY, MIAMI 33199, USA

Email address: annamaria.fino@unito.it, afino@fiu.edu

DPMMS, University of Cambridge, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WB, UK

Email address: A.G.Kovalev@dpmms.cam.ac.uk

FACULTAD DE CIENCIAS MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, PLAZA CIENCIAS 3, CIUDAD UNIVERSITARIA, 28040 MADRID, SPAIN

Email address: vicente.munoz@ucm.es