

Compact leaves of the foliation defined by the kernel of a T^2 -invariant presymplectic form

Asuka Hagiwara *

Abstract

We investigate the foliation defined by the kernel of an exact presymplectic form $d\alpha$ of rank $2n$ on a $(2n + r)$ -dimensional closed manifold M . For $r = 2$, we prove that the foliation has at least two leaves which are homeomorphic to a 2-dimensional torus, if M admits a locally free T^2 -action which preserves $d\alpha$ and satisfies that the function $\alpha(Z_2)$ is constant, where Z_1, Z_2 are the infinitesimal generators of the T^2 -action. We also give its generalization for $r \geq 1$.

1 Introduction

Let n and r be positive integers. A smooth manifold M of dimension $2n + r$ is said to be *presymplectic* if it carries a closed 2-form ω such that the dimension of the kernel of $\omega(x)$ is r for all $x \in M$, that is, ω is of constant rank $2n$. The 2-form ω is called a *presymplectic form* on M . We denote the kernel of ω by $\ker \omega$, which is an involutive distribution of dimension r . Hence, by Frobenius' theorem, $\ker \omega$ defines an r -dimensional foliation \mathcal{F}_ω . The leaves of \mathcal{F}_ω are integral manifolds of the distribution.

A special case of presymplectic manifolds is when a manifold M is $(2n + 1)$ -dimensional and carries a contact form α , that is, α is a 1-form on M such that $\alpha \wedge (d\alpha)^n \neq 0$. In this case, the 2-form $\omega := d\alpha$ is a presymplectic form on M and the foliation \mathcal{F}_ω is 1-dimensional. The leaves of \mathcal{F}_ω are integral curves of $\ker \omega$, which are called *characteristics*. The Weinstein conjecture [9] asserts that there exists at least one closed characteristic on a closed contact manifold, which is one of the fundamental problems concerning dynamics on contact manifolds. This conjecture is still open, but has been proved in several cases. For instance, Viterbo [8] proved it for compact contact manifolds which are hypersurfaces of contact type in $(\mathbb{R}^{2n}, \omega_0)$, where $\omega_0 = \sum_{j=1}^n dy_j \wedge dx_j$ is the standard symplectic form. Hofer [4] proved this conjecture

2020 *Mathematics Subject Classification*. Primary 37C86, Secondary 37C85, 53D10

Key Words and Phrases. presymplectic form, foliation, torus action, generalized Weinstein conjecture

*Graduate School of Science and Engineering, Ibaraki University, Bunkyo 2-1-1, Mito, Ibaraki, 310-8512, Japan.

for S^3 and for overtwisted 3-dimensional contact manifolds, and after that Taubes [7] solved it affirmatively in dimension 3.

In [1], Banyaga and Rukimbira raised the following question for presymplectic manifolds (M, ω) , which generalizes the Weinstein conjecture.

Question ([1, p.3902]). When does the foliation \mathcal{F}_ω admits a compact leaf?

In addition, they focused their attention on the case where $\dim \mathcal{F}_\omega = 1$ and proved the following theorem.

Theorem 1 (Banyaga-Rukimbira, [1]). *Let M be a $(2n + 1)$ -dimensional oriented closed C^∞ -manifold with a 1-form α such that the 2-form $\omega := d\alpha$ has constant rank $2n$ everywhere. If there exists a locally free circle-action on M which preserves ω , then \mathcal{F}_ω has at least two closed leaves.*

In particular, they pointed out that this result guarantees the existence of periodic orbits of the Hamiltonian system on hypersurfaces in \mathbb{R}^{2n} which are not necessarily of contact type. However, they did not mention the case where $\dim \mathcal{F}_\omega \geq 2$. For such a direction, it seems that there are few works which assume that (M, ω) is r -contact (see Remark in Section 3, and [3] for details).

In this paper, we consider a higher dimensional version of Theorem 1 without assuming that (M, ω) is r -contact. We show the following result for the case where $\dim \mathcal{F}_\omega = 2$.

Theorem 2. *Let M be a $(2n + 2)$ -dimensional oriented closed C^∞ -manifold with an exact presymplectic form ω , that is, there exists a 1-form $\alpha \in \Omega^1(M)$ such that the 2-form $\omega = d\alpha$ has rank $2n$ everywhere. Assume that M admits a locally free T^2 -action with the following conditions:*

- (i) *The T^2 -action preserves ω ,*
- (ii) *The function $\alpha(Z_2)$ is constant on M ,*

where Z_1, Z_2 denote the infinitesimal generators of the T^2 -action. Then

- (1) *the 2-dimensional C^∞ -foliation \mathcal{F}_ω has at least two leaves which are homeomorphic to a 2-dimensional torus.*
- (2) *Moreover, if the function $\alpha(Z_1)$ is also constant, then \mathcal{F}_ω coincides with the foliation of the T^2 -action and hence, all leaves of \mathcal{F}_ω are homeomorphic to a 2-dimensional torus.*

We prove Theorem 2 in the next section. In Section 3 we provide examples of Theorem 2 and mention that a similar result holds for the general case where $\dim \mathcal{F}_\omega = r$.

2 Proof of Theorem 2

Let $\rho : M \times T^2 \rightarrow M$ be a locally free T^2 -action on M with the conditions (i) and (ii) above, and let $\mathcal{D} : M \ni q \mapsto \mathcal{D}_q = \text{span}\{Z_1(q), Z_2(q)\} \subset T_q M$ be the distribution determined by ρ , where Z_1, Z_2 denote the infinitesimal generators of ρ . The distribution \mathcal{D} is involutive and of dimension 2 since ρ is locally free. Therefore, now M has two foliations of codimension $2n$ which are determined by $\ker \omega$ and \mathcal{D} . Note that, by definition, every leaf of \mathcal{D} is homeomorphic to a 2-dimensional torus T^2 .

We define the diffeomorphism $s : M \rightarrow M$ by $s(x) := \rho(x, s)$ for $s \in T^2$ and the 1-form

$$\alpha_0 := \int_{T^2} (s^* \alpha) d\sigma,$$

where σ is the Haar measure on T^2 . Then α_0 is T^2 -invariant. Thus,

$$L_{Z_i} \alpha_0 = \lim_{t \rightarrow 0} \frac{(\varphi_i^t)^* \alpha_0 - \alpha_0}{t} = 0, \quad i = 1, 2,$$

where $\varphi_i^t = \rho(\cdot, \exp t Z_i)$ denotes the flow of Z_i , respectively. Moreover, define smooth functions $S_i : M \rightarrow \mathbb{R}$ by

$$S_i(x) := -\alpha_0(x)(Z_i(x)) = -i_{Z_i} \alpha_0(x), \quad i = 1, 2.$$

By the condition (i), we have

$$d\alpha_0 = \int_{T^2} d(s^* \alpha) d\sigma = \int_{T^2} (s^* d\alpha) d\sigma = \omega,$$

and hence, due to the Cartan formula,

$$dS_i = -di_{Z_i} \alpha_0 = -L_{Z_i} \alpha_0 + i_{Z_i} d\alpha_0 = i_{Z_i} \omega. \quad (1)$$

Since the group T^2 is commutative, we have $s \circ \varphi_2^t(x) = \varphi_2^t \circ s(x)$ for all $x \in M$. Differentiating this equation in t , we obtain $ds(x)Z_2(x) = Z_2(s(x))$, so that

$$S_2(x) = - \int_{T^2} s^* \alpha(x)(Z_2(x)) d\sigma = - \int_{T^2} \alpha(s(x))(Z_2(s(x))) d\sigma.$$

Thus, by the condition (ii) the function S_2 is constant and so $dS_2 = 0$. Hence,

$$Z_2(q) \in \ker \omega(q), \quad \forall q \in M \quad (2)$$

by (1) for $i = 2$. To investigate the relation between Z_1 and $\ker \omega$, we shall take a special chart. Since $\ker \omega$ is involutive, by using Frobenius' theorem [2, p.89, Theorem 1], for any point $p \in M$ we can choose the following chart (φ, U) around p : for $q \in U$ we write

$$\varphi(q) = (x(q), z_1(q), z_2(q)), \quad x(q) = (x_1(q), \dots, x_{2n}(q)),$$

then this chart satisfies that $\varphi(p) = (0, 0, 0)$ and

$$\xi_i := \frac{\partial}{\partial z_i}, \quad i = 1, 2$$

form a local frame of $\ker \omega$ on U . For $q \in U$ we put

$$Z_1(q) = \sum_{j=1}^{2n} X_j(x, z_1, z_2) \frac{\partial}{\partial x_j} + u_1(x, z_1, z_2) \frac{\partial}{\partial z_1} + u_2(x, z_1, z_2) \frac{\partial}{\partial z_2}$$

and prove the following

Lemma 3. *The function X_j is independent of z_1, z_2 for any $j = 1, \dots, 2n$.*

Proof. In U , since $\xi_i \in \ker \omega$, we have $L_{\xi_i} \omega = i_{\xi_i} d\omega + di_{\xi_i} \omega = 0$ and

$$\begin{aligned} L_{\xi_i}(i_{Z_j} \omega) &= i_{\xi_i} d(i_{Z_j} \omega) + di_{\xi_i}(i_{Z_j} \omega) \\ &= i_{\xi_i} ddS_j - di_{Z_j} i_{\xi_i} \omega = 0 \quad (\because (1)). \end{aligned}$$

Therefore,

$$i_{[\xi_i, Z_j]} \omega = L_{\xi_i} i_{Z_j} \omega - i_{Z_j} L_{\xi_i} \omega = 0$$

and namely, there exist functions $\lambda_i^1, \lambda_i^2 : U \rightarrow \mathbb{R}$ such that

$$[\xi_i, Z_1] = \lambda_i^1 \xi_1 + \lambda_i^2 \xi_2.$$

On the other hand,

$$\begin{aligned} [\xi_i, Z_1] &= \left[\frac{\partial}{\partial z_i}, Z_1 \right] = \left[\frac{\partial}{\partial z_i}, \sum_{j=1}^{2n} X_j \frac{\partial}{\partial x_j} \right] + \left[\frac{\partial}{\partial z_i}, u_1 \frac{\partial}{\partial z_1} \right] + \left[\frac{\partial}{\partial z_i}, u_2 \frac{\partial}{\partial z_2} \right] \\ &= \sum_{j=1}^{2n} \frac{\partial X_j}{\partial z_i} \frac{\partial}{\partial x_j} + \frac{\partial u_1}{\partial z_i} \frac{\partial}{\partial z_1} + \frac{\partial u_2}{\partial z_i} \frac{\partial}{\partial z_2}. \end{aligned}$$

Therefore, $\partial X_j / \partial z_i = 0$. □

By this lemma, we have

$$Z_1(q) = \sum_{j=1}^{2n} X_j(x) \frac{\partial}{\partial x_j} + u_1(x, z_1, z_2) \frac{\partial}{\partial z_1} + u_2(x, z_1, z_2) \frac{\partial}{\partial z_2} \quad \text{for } q \in U.$$

Now, we assume that $p \in M$ is a critical point of S_1 . By (1) for $i = 1$, we have $Z_1(p) \in \ker \omega(p)$. Because $\varphi(p) = (0, 0, 0)$, we have

$$Z_1(p) = \sum_{j=1}^{2n} X_j(0) \frac{\partial}{\partial x_j} + u_1(0, 0, 0) \frac{\partial}{\partial z_1} + u_2(0, 0, 0) \frac{\partial}{\partial z_2},$$

so that

$$X_j(0) = 0, \quad j = 1, \dots, 2n. \quad (3)$$

We denote by $\mathcal{F}_\omega(p)$ the leaf of \mathcal{F}_ω passing through the point p . Let $q \in \mathcal{F}_\omega(p) \cap U$. Then there exist $t_1, t_2 \in \mathbb{R}$ such that $\varphi(q) = (0, t_1, t_2)$, since

$$T_q(\mathcal{F}_\omega(p)) = \ker \omega(q) = \text{span} \left\{ \left(\frac{\partial}{\partial z_1} \right)_q, \left(\frac{\partial}{\partial z_2} \right)_q \right\}.$$

Thus, by (3), we obtain

$$\begin{aligned} Z_1(q) &= \sum_{j=1}^{2n} X_j(0) \frac{\partial}{\partial x_j} + u_1(0, t_1, t_2) \frac{\partial}{\partial z_1} + u_2(0, t_1, t_2) \frac{\partial}{\partial z_2} \\ &= u_1(0, t_1, t_2) \frac{\partial}{\partial z_1} + u_2(0, t_1, t_2) \frac{\partial}{\partial z_2} \in \ker \omega(q). \end{aligned} \quad (4)$$

It follows from (2) that

$$\ker \omega(q) = \mathcal{D}_q, \quad \forall q \in \mathcal{F}_\omega(p) \cap U, \quad (5)$$

since the dimensions of $\ker \omega$ and \mathcal{D} are the same. In order to verify that $\ker \omega$ coincides with \mathcal{D} on the leaf $\mathcal{F}_\omega(p)$, we cover $\mathcal{F}_\omega(p)$ by a set of charts $\{(U_a, \varphi_a)\}_{a \in A}$, where each (U_a, φ_a) is a chart as we chose above. For $(p \neq) \hat{p} \in \mathcal{F}_\omega(p)$ we take a chart $(\hat{U}, \hat{\varphi}) \in \{(U_a, \varphi_a)\}_{a \in A}$ around \hat{p} which satisfies $\mathcal{F}_\omega(p) \cap U \cap \hat{U} \neq \emptyset$. We write

$$\hat{\varphi}(q) = (y(q), w_1(q), w_2(q)), \quad y(q) = (y_1(q), \dots, y_{2n}(q))$$

for $q \in \hat{U}$, then $\hat{\varphi}(\hat{p}) = (0, 0, 0)$ and $\partial/\partial w_1, \partial/\partial w_2$ form a local frame of $\ker \omega$ on \hat{U} . With this setting, by applying Lemma 3, we have

$$Z_1(q) = \sum_{j=1}^{2n} \hat{X}_j(y) \frac{\partial}{\partial y_j} + \hat{u}_1(y, w_1, w_2) \frac{\partial}{\partial w_1} + \hat{u}_2(y, w_1, w_2) \frac{\partial}{\partial w_2}$$

in \hat{U} as well. Since $T_q(\mathcal{F}_\omega(p)) = \ker \omega(q)$, we have $\hat{\varphi}(q) = (0, w_1(q), w_2(q))$ for all $q \in \mathcal{F}_\omega(p) \cap \hat{U}$. Therefore, for $q \in \mathcal{F}_\omega(p) \cap U \cap \hat{U}$,

$$Z_1(q) = \sum_{j=1}^{2n} \hat{X}_j(0) \frac{\partial}{\partial y_j} + \hat{u}_1(0, w_1(q), w_2(q)) \frac{\partial}{\partial w_1} + \hat{u}_2(0, w_1(q), w_2(q)) \frac{\partial}{\partial w_2}$$

holds. From (5), we obtain $\hat{X}_j(0) = 0$, $j = 1, \dots, 2n$. Thus, for all $q \in \mathcal{F}_\omega(p) \cap \hat{U}$ we have

$$Z_1(q) = \hat{u}_1(0, w_1(q), w_2(q)) \frac{\partial}{\partial w_1} + \hat{u}_2(0, w_1(q), w_2(q)) \frac{\partial}{\partial w_2} \in \ker \omega(q),$$

so that $\ker \omega(q) = \mathcal{D}_q$. Repeating this argument we see that

$$\ker \omega(q) = \mathcal{D}_q, \quad \forall q \in \mathcal{F}_\omega(p). \quad (6)$$

Therefore, $\mathcal{F}_\omega(p)$ is not only an integral manifold of $\ker \omega$, but also that of \mathcal{D} passing through p . On the other hand, the T^2 -orbit $T^2(p)$ of p is nothing but the maximal connected integral manifold of \mathcal{D} passing through p . Hence, by [5, p.172, Theorem 1], $\mathcal{F}_\omega(p)$ is an open submanifold in $T^2(p)$. Actually, we claim the following

Lemma 4. $\mathcal{F}_\omega(p) = T^2(p)$ holds. In particular, $\mathcal{F}_\omega(p)$ is homeomorphic to T^2 .

Proof. Since $\mathcal{F}_\omega(p)$ is a nonempty open set in a connected space $T^2(p)$, it suffices to show that $\mathcal{F}_\omega(p)$ is a closed set in $T^2(p)$. We denote by $\overline{\mathcal{F}_\omega(p)}$ the closure of $\mathcal{F}_\omega(p)$ in $T^2(p)$. Let $q_0 \in \overline{\mathcal{F}_\omega(p)}$. We take the following chart (U, φ) around q_0 as we chose above: for $q \in U$ we write

$$\varphi(q) = (x(q), z_1(q), z_2(q)), \quad x(q) = (x_1(q), \dots, x_{2n}(q)),$$

then $\varphi(q_0) = (0, 0, 0)$, and $\partial/\partial z_1, \partial/\partial z_2$ form a local frame of $\ker \omega$ on U . By Frobenius' theorem [2, p.89, Theorem 1], the slice

$$S_0 := \{q \in U \mid x_1(q) = 0, \dots, x_{2n}(q) = 0\}$$

is a connected integral manifold of $\ker \omega$ passing through q_0 .

Claim. The slice S_0 is an open submanifold in $T^2(p)$ which contains q_0 .

Proof of Claim. For $q \in U$, by Lemma 3, we have

$$Z_1(q) = \sum_{j=1}^{2n} X_j(x) \frac{\partial}{\partial x_j} + u_1(x, z_1, z_2) \frac{\partial}{\partial z_1} + u_2(x, z_1, z_2) \frac{\partial}{\partial z_2}.$$

First we prove that $Z_1(q_0) \in \ker \omega(q_0)$. Arguing by contradiction we assume that $Z_1(q_0) \notin \ker \omega(q_0)$, that is, $X_j(x(q_0)) = X_j(0) \neq 0$ for some $j \in \{1, \dots, 2n\}$. Since the function X_j is smooth on U , there exists a neighborhood $U' \subset U$ of q_0 such that $X_j(x(q)) \neq 0$ for any $q \in U'$. Because $q_0 \in \overline{\mathcal{F}_\omega(p)}$, the intersection $\mathcal{F}_\omega(p) \cap U'$ is nonempty, so that for $q \in \mathcal{F}_\omega(p) \cap U'$ it holds that $X_j(x(q)) \neq 0$. This is a contradiction to (6). Hence, we have $X_j(0) = 0$ for all $j = 1, \dots, 2n$.

Therefore, for all $q \in S_0$, we can write $\varphi(q) = (0, z_1(q), z_2(q))$ and we have

$$Z_1(q) = u_1(0, z_1(q), z_2(q)) \frac{\partial}{\partial z_1} + u_2(0, z_1(q), z_2(q)) \frac{\partial}{\partial z_2} \in \ker \omega(q).$$

Thus, by (2) we obtain $\mathcal{D}_q = \ker \omega(q)$ for all $q \in S_0$. It follows from $T_q S_0 = \ker \omega(q)$ that S_0 is a connected integral manifold of \mathcal{D} through q_0 . On the other hand, $T^2(p)$ is the maximal connected integral manifold of \mathcal{D} which contains q_0 . Therefore, by [5, p.172, Theorem 1], we see that S_0 is an open submanifold in $T^2(p)$ which contains q_0 . \square

By this claim, S_0 is an open neighborhood of q_0 in $T^2(p)$. Because $q_0 \in \overline{\mathcal{F}_\omega(p)}$, the intersection $S_0 \cap \mathcal{F}_\omega(p)$ is nonempty. Thus, $\mathcal{F}_\omega(p)$ is the leaf of $\ker \omega$ passing through a point $p_0 \in S_0 \cap \mathcal{F}_\omega(p)$ as well. Due to Frobenius' theorem [2, p.89, Theorem 1], S_0 is a connected integral manifold of $\ker \omega$ which contains the point p_0 . Hence, by [5, p.172, Theorem 1], S_0 is also an open submanifold in $\mathcal{F}_\omega(p)$. Consequently, $q_0 \in S_0 \subset \mathcal{F}_\omega(p)$ and so we have $\overline{\mathcal{F}_\omega(p)} = \mathcal{F}_\omega(p)$. \square

Finally, we check the existence of critical points of the function S_1 on M . If S_1 is not constant, then it has at least two critical points $p_{\max}, p_{\min} \in M$ corresponding to a maximum and a minimum, respectively, since M is closed. Moreover, p_{\max} and p_{\min} are on different leaves of \mathcal{F}_ω . This follows from the following. By (1) and (2), we have

$$L_{Z_1} S_1 = i_{Z_1} dS_1 + di_{Z_1} S_1 = i_{Z_1} i_{Z_1} \omega = 0$$

and

$$L_{Z_2} S_1 = i_{Z_2} dS_1 + di_{Z_2} S_1 = i_{Z_2} i_{Z_1} \omega = -i_{Z_1} i_{Z_2} \omega = 0,$$

so that S_1 is constant along $T^2(p) = \{\varphi_1^s \circ \varphi_2^t(p) \mid s, t \in \mathbb{R}\}$. Therefore, $T^2(p_{\max}) \cap T^2(p_{\min}) = \emptyset$ and hence, by Lemma 4, we obtain $\mathcal{F}_\omega(p_{\max}) \cap \mathcal{F}_\omega(p_{\min}) = \emptyset$. If S_1 is constant on M , then $dS_1(p) = 0$ for all $p \in M$, so that $\ker \omega = \mathcal{D}$ on M . Hence, from the uniqueness of foliations, \mathcal{F}_ω coincides with the foliation defined by \mathcal{D} and therefore, each leaf of \mathcal{F}_ω is homeomorphic to T^2 . Consequently, in any case, we see that \mathcal{F}_ω has at least two leaves which are homeomorphic to T^2 . In particular, if $\alpha(Z_1)$ is constant, then, by a similar calculation for S_2 , we see that S_1 is constant on M . Thus, we complete the proof of Theorem 2.

3 Examples and Remarks

We first provide an example of Theorem 2.

Example 5. We consider the case of a submanifold of codimension 2 of a symplectic manifold $(\mathbb{R}^6, d\lambda)$, where

$$\lambda = \frac{1}{2} \sum_{j=1}^3 (y_j dx_j - x_j dy_j)$$

is the Liouville form. Then $d\lambda = \omega_0$ is the standard symplectic form on \mathbb{R}^6 . We define functions $G_i : \mathbb{R}^6 \rightarrow \mathbb{R}$, $i = 1, 2$, by

$$G_1 := (x_1^2 + y_1^2)^2 + x_2^2 + y_2^2, \quad G_2 := x_2^2 + y_2^2 + x_3^2 + y_3^2$$

and for real numbers $c_1 > c_2 > 0$ we put

$$M := \{(x, y) \in \mathbb{R}^6 \mid G_1 = c_1, G_2 = c_2\}.$$

Since the gradients

$$\nabla G_1 = \begin{pmatrix} 4x_1(x_1^2 + y_1^2) \\ 4y_1(x_1^2 + y_1^2) \\ 2x_2 \\ 2y_2 \\ 0 \\ 0 \end{pmatrix}, \quad \nabla G_2 = \begin{pmatrix} 0 \\ 0 \\ 2x_2 \\ 2y_2 \\ 2x_3 \\ 2y_3 \end{pmatrix}$$

are linearly independent on M , we see that M is a 4-dimensional closed submanifold of \mathbb{R}^6 . Because $\{G_1, G_2\} = 0$, we deduce that $\dim(\ker(\omega_0|_M)) = 2$ and $\omega_0|_M$ is a presymplectic form of constant rank 2 (see [6, p.27]). We define a T^2 -action ρ on M as follows. We put $z_j = x_j + \sqrt{-1}y_j$ and for $(e^{\sqrt{-1}2\pi s_1}, e^{\sqrt{-1}2\pi s_2}) \in T^2$, $s_1, s_2 \in \mathbb{R}$, we set

$$f_{s_1} : \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix} \mapsto \begin{pmatrix} e^{\sqrt{-1}2\pi s_1} & 0 & 0 \\ 0 & e^{\sqrt{-1}2\pi s_1} & 0 \\ 0 & 0 & e^{\sqrt{-1}2\pi s_1} \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix},$$

$$g_{s_2} : \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{\sqrt{-1}2\pi s_2} & 0 \\ 0 & 0 & e^{\sqrt{-1}2\pi s_2} \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix}$$

and define

$$\rho^{(e^{\sqrt{-1}2\pi s_1}, e^{\sqrt{-1}2\pi s_2})}(z) := f_{s_1} \circ g_{s_2}(z) = g_{s_2} \circ f_{s_1}(z), \quad z = (z_1, z_2, z_3) \in M.$$

Then $\rho^{(e^{\sqrt{-1}2\pi s_1}, e^{\sqrt{-1}2\pi s_2})}(z) \in M$ and we can easily check that

$$\rho^{(e^{\sqrt{-1}2\pi s_1}, e^{\sqrt{-1}2\pi s_2})^*} \lambda = \lambda \tag{7}$$

and ρ is a free action preserving $d\lambda$. The infinitesimal generators of ρ are given by

$$Z_1 = \left. \frac{d}{ds_1} f_{s_1}(x, y) \right|_{s_1=0} = \sum_{j=1}^3 \left(-2\pi y_j \frac{\partial}{\partial x_j} + 2\pi x_j \frac{\partial}{\partial y_j} \right),$$

$$Z_2 = \left. \frac{d}{ds_2} g_{s_2}(x, y) \right|_{s_2=0} = \sum_{j=2}^3 \left(-2\pi y_j \frac{\partial}{\partial x_j} + 2\pi x_j \frac{\partial}{\partial y_j} \right),$$

and therefore, the function

$$\lambda(Z_1) = -\pi \sum_{j=1}^3 (x_j^2 + y_j^2) = -\pi(x_1^2 + y_1^2 + c_2)$$

is nonconstant on M and the function

$$\lambda(Z_2) = -\pi(x_2^2 + y_2^2 + x_3^2 + y_3^2) = -\pi c_2$$

is constant on M . It follows that $(M, \omega_0|_M)$ satisfies the conditions (i), (ii) of Theorem 2, but not the additional condition in (2) of Theorem 2. Thus, the foliation $\mathcal{F}_{\omega_0|_M}$ has at least two leaves which are homeomorphic to T^2 .

In this case, by (7), the function S_1 defined in the proof of Theorem 2 is given by

$$S_1 = - \int_{T^2} \left(\rho^{(e^{\sqrt{-1}2\pi s_1}, e^{\sqrt{-1}2\pi s_2})^*} \lambda \right) (Z_1) d\sigma = - \int_{T^2} \lambda(Z_1) d\sigma = -\lambda(Z_1).$$

Due to the compactness of M , there exist $p_{\max}, p_{\min} \in M$ such that $S_1(p_{\max})$ is a maximal value and $S_1(p_{\min})$ is a minimal value of S_1 . By the former part of the last paragraph of the proof of Theorem 2, the function S_1 is constant along the leaves $\mathcal{F}_{\omega_0|_M}(p_{\max})$ and $\mathcal{F}_{\omega_0|_M}(p_{\min})$. In fact, by using the method of Lagrange multipliers, we see that S_1 has only two critical submanifolds and S_1 takes the maximal value $S_1(p_{\max}) = \pi(\sqrt{c_1} + c_2)$ on

$$\mathcal{F}_{\omega_0|_M}(p_{\max}) = \{(x, y) \in M \mid x_1^2 + y_1^2 = \sqrt{c_1}, x_2^2 = y_2^2 = 0, x_3^2 + y_3^2 = c_2\}$$

and the minimal value $S_1(p_{\min}) = \pi(\sqrt{c_1 - c_2} + c_2)$ on

$$\mathcal{F}_{\omega_0|_M}(p_{\min}) = \{(x, y) \in M \mid x_1^2 + y_1^2 = \sqrt{c_1 - c_2}, x_2^2 + y_2^2 = c_2, x_3 = y_3 = 0\}.$$

Moreover, we can easily check that

$$Z_1(q) \notin \ker \omega(q) \quad \text{for } q \in M \setminus (\mathcal{F}_{\omega_0|_M}(p_{\max}) \cup \mathcal{F}_{\omega_0|_M}(p_{\min})). \quad (8)$$

Another example can be found in [6, p.24], which satisfies the condition in (2) of Theorem 2.

Example 6. We also consider $(\mathbb{R}^6, \omega_0 = d\lambda)$, where λ is the Liouville form. Define functions on \mathbb{R}^6 by

$$G_1 := x_1^2 + y_1^2, \quad G_2 := \sum_{j=1}^3 (x_j^2 + y_j^2).$$

For real numbers $c_2 > c_1 > 0$, we set

$$M := \{(x, y) \in \mathbb{R}^6 \mid G_1 = c_1, G_2 = c_2\}$$

then, as in Example 5, M is a 4-dimensional closed manifold with a presymplectic form $\omega_0|_M$ of constant rank 2. We shall define a T^2 -action on M . Put $z_j = x_j + \sqrt{-1}y_j$. For $(e^{\sqrt{-1}2\pi s_1}, e^{\sqrt{-1}2\pi s_2}) \in T^2$, $s_1, s_2 \in \mathbb{R}$, we set

$$f_{s_1} : \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix} \mapsto \begin{pmatrix} e^{\sqrt{-1}2\pi s_1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix},$$

$$g_{s_2} : \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix} \mapsto \begin{pmatrix} e^{\sqrt{-1}2\pi s_2} & 0 & 0 \\ 0 & e^{\sqrt{-1}2\pi s_2} & 0 \\ 0 & 0 & e^{\sqrt{-1}2\pi s_2} \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix}$$

and define

$$\rho^{(e^{\sqrt{-1}2\pi s_1}, e^{\sqrt{-1}2\pi s_2})}(z) := f_{s_1} \circ g_{s_2}(z) = g_{s_2} \circ f_{s_1}(z), \quad z = (z_1, z_2, z_3) \in M.$$

We see that this action is free and preserves ω_0 . Similar to Example 5, we denote the infinitesimal generators of ρ by Z_1, Z_2 . Then the functions $\lambda(Z_1), \lambda(Z_2)$ are constant on M . Thus, the foliation $\mathcal{F}_{\omega_0|_M}$ coincides with the foliation of the T^2 -action ρ and therefore, all leaves of $\mathcal{F}_{\omega_0|_M}$ are homeomorphic to T^2 .

Similar to the proof of Theorem 2, we have the following result for arbitrary $r \geq 1$.

Theorem 7. *Let M be a $(2n + r)$ -dimensional oriented closed C^∞ -manifold with an exact presymplectic form ω , that is, there exists a 1-form $\alpha \in \Omega^1(M)$ such that the 2-form $\omega = d\alpha$ has rank $2n$ everywhere. Assume that M admits a locally free T^r -action with the following conditions:*

- (i) *The T^r -action preserves ω ,*
- (ii) *The functions $\alpha(Z_i)$, $i = 2, \dots, r$, are constant on M ,*

where Z_1, Z_2, \dots, Z_r denote the infinitesimal generators of the T^r -action. Then the r -dimensional C^∞ -foliation \mathcal{F}_ω has at least two leaves which are homeomorphic to an r -dimensional torus. Moreover, if the function $\alpha(Z_1)$ is also constant, then \mathcal{F}_ω coincides with the foliation of the T^r -action and hence, all leaves of \mathcal{F}_ω are homeomorphic to an r -dimensional torus.

Theorem 7 also gives a partial answer to Question in Section 1. If $r = 1$, then Theorem 7 agrees with Theorem 1.

Remark. We emphasize that in Theorem 7 (and 2) we do not assume that (M, ω) is r -contact, which means that M carries r linearly independent non-vanishing 1-forms $\alpha_1, \dots, \alpha_r$ with a splitting $TM = \mathcal{R} \oplus (\cap_i^r \ker \alpha_i)$ satisfying that $d\alpha_i|_{\cap_i^r \ker \alpha_i}$ is non-degenerate and $\ker d\alpha_i = \mathcal{R}$ for every i . In our context, \mathcal{R} corresponds to $\ker \omega$. Finamore [3, Theorem 3.23] proved that if a closed presymplectic manifold (M, ω) is r -contact with a special metric, then the r -dimensional foliation defined by \mathcal{R} has at least two leaves which are homeomorphic to an r -dimensional torus. In the case where M is r -contact, by definition, M admits a locally free \mathbb{R}^r -action with the infinitesimal generators $R_1, \dots, R_r \in \mathcal{R}$. On the other hand, in Theorem 7, M does not always satisfy that $Z_1 \in \ker \omega$, as (8) in Example 5 shows. Thus, [3, Theorem 3.23] and Theorem 7 are independent results.

Acknowledgement

This paper is a part of the author's Master thesis presented in February 2022 at Ibaraki University. I would like to thank my supervisor, Prof. Hiroshi Iriyeh for suggesting the problem, enlightening discussions and helpful comments.

References

- [1] A. Banyaga, P. Rukimbira, *On characteristics of circle invariant presymplectic forms*, Proc. Amer. Math. Soc. 123 (1995), no. 12, 3901–3906. doi:10.2307/2161922
- [2] C. Chevalley, *Theory of Lie groups*, Princeton, 1946.
- [3] D. Finamore, *Contact foliations and generalised Weinstein conjectures*, doi:10.48550/arXiv.2202.07622
- [4] H. Hofer, *Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjecture in dimension three*, Invent. Math. 114 (1993), 515–563. doi:10.1007/BF01232679
- [5] Y. Matsushima, *Differentiable manifolds*, Translated from the Japanese by E. T. Kobayashi. Pure and Applied Mathematics, 9. Marcel Dekker, Inc., New York, 1972. vii+303 pp.
- [6] J. Moser, *A fixed point theorem in symplectic geometry*, Acta Math. 141 (1978), no. 1-2, 17–34. doi:10.1007/BF02545741
- [7] C. Taubes, *The Seiberg-Witten equations and the Weinstein conjecture*, Geom. Topol. 11(4) (2007), 2117–2202. doi:10.2140/gt.2007.11.2117
- [8] C. Viterbo, *A proof of the Weinstein conjecture in \mathbb{R}^{2n}* , Ann. Inst. H. Poincaré, Anal. Non linéaire, 4(4) (1987), 337–357. doi:10.1016/S0294-1449(16)30363-8
- [9] A. Weinstein, *On the hypothesis of Rabinowitz’s periodic orbit theorems*, J. Diff. Equ. 33 (1979), 353–358. doi:10.1016/0022-0396(79)90070-6

Present Address:

Numata Girls’ High School,
Higashikurauchimachi 753-3, Numata, Gumma, 378-0043, Japan.
e-mail: hagiwara-asuka@edu-g.gsn.ed.jp