

Fermionic contribution to the anomalous dimension
of twist-2 operators in $\mathcal{N} = 4$ SYM theory,
critical indices and integrability

V.N. VELIZHANIN

*Theoretical Physics Division
NRC “Kurchatov Institute”
Petersburg Nuclear Physics Institute
Orlova Roscha, Gatchina
188300 St. Petersburg, Russia
velizh@thd.pnpi.spb.ru*

Abstract

We compute the contribution to the anomalous dimension of the twist-2 operators in $\mathcal{N} = 4$ SYM theory, which is proportional to the number of fermion loops inside Feynman diagrams or, formally, to the number of fermions. The result was obtained by the method based on the calculation of critical indices at the critical point by analogy with previous similar computations in scalar theories and in QCD. The obtained result is much simpler with compare to analogous results in QCD and almost satisfies the maximal transcendentality principle. A possible relation between the obtained result and integrability is discussed.

The anomalous dimension of twist-2 operators is intensively studied in $\mathcal{N} = 4$ SYM theory due to the relation with integrability in the framework of AdS/CFT-correspondence [1–3]. A number of exact results were obtained with the help of integrability, such as the all-loop asymptotic Bethe ansatz [4–16], the TBA-equations and the Y-system [17–21], and finally the quantum spectral curve approach [22, 23], which solve the spectral problem for composite operators in $\mathcal{N} = 4$ SYM theory.

From the point of view of applicability in realistic models, first of all QCD, the most interesting results obtained in $\mathcal{N} = 4$ SYM theory give the most complicated (more precisely, the most transcendental) contribution according to the maximal transcendentality principle [24]. In fact, the maximal transcendentality principle was proposed to obtain the results in $\mathcal{N} = 4$ SYM theory from the available results in QCD. Initially, an indication of this was obtained during the computations of the eigenvalues of the Balitsky-Fadin-Kuraev-Lipatov (BFKL) equations [25–27] in $\mathcal{N} = 4$ SYM theory [28] based on the original computations in QCD [29]. It turned out that if we assign to each term in the general expression for this quantity its transcendentality, which is the sum of absolute values of indices of multiple zeta-values, that are obtained from the function as its argument tends to infinity, then the result in the given order of perturbative theory will contain only functions with maximal transcendentality, equal to $2\ell - 1$ for the ℓ -th order. To prove this assumption, we calculated the two-loop anomalous dimension of twist-2 operators in $\mathcal{N} = 4$ SYM theory [30] and found that it actually holds if we put into the corresponding QCD results [31–34] the following identifications for Casimir operators: $C_F = N_c$ and $C_A = N_c$.

However, in QCD there are also some exact (all-loop) results for twist-2 operators. The most famous result is the eigenvalue of the BFKL equation in the leading and next-to-leading logarithmic approximations [25–27, 29], from which one can obtain an expression for the anomalous dimension of the gluon operator in any order of perturbative theory near $j = 1 + \omega$, where it has poles in ω . Another similar exact result is the double-logarithmic equation for the flavour non-singlet quark twist-2 operator [35–37], which gives information about the corresponding anomalous dimension in any order of perturbative theory near $j = 0 + \omega$, where it has poles in ω .

The most interesting all-loop result related to the subject of this paper is the expansion of the non-singlet anomalous dimension in the number of quark flavors or the number of quark loops inside the Feynman diagrams. This result was obtained from the calculation of the critical indices at the critical point within dimensional regularization in Ref. [38], following the methods proposed in Refs. [39, 40].

In this paper, we compute the contribution proportional to the number of fermionic loops to the anomalous dimension of the special twist-2 operator in $\mathcal{N} = 4$ SYM theory, which is similar to the flavor non-singlet operator in QCD. The operator we will consider belong to the representation **15** of the $SU(4)$ group of internal symmetry of $\mathcal{N} = 4$ SYM




Figure 1: One-loop diagrams contributing to anomalous dimension of $\tilde{\mathcal{O}}_{15}^{qq}$.

theory and can formally be written in two-componet notation as [41]

$$\left[\tilde{\mathcal{O}}_{jl}^{qq,15} \right]_A^B = \sigma_j \text{tr} [P_{15}]_{AD}^{BC} \bar{\lambda}_{\dot{\alpha}C} \sigma^{+\dot{\alpha}\beta} (i\partial^+)^l C_j^{3/2} \left(\overleftrightarrow{\mathcal{D}}^+ / \partial^+ \right) \lambda_\beta^D, \quad (1)$$

$$[P_{15}]_{AD}^{BC} = \delta_A^C \delta_D^B - \frac{1}{4} \delta_A^B \delta_D^C, \quad \sigma_j \equiv 1 - (-1)^j, \quad (2)$$

where $C_j^{3/2}$ is the Gegenbauer polynomial. In the notation close to usual QCD form it can be written as

$$\tilde{\mathcal{O}}_{15}^{qq}(n) = \bar{\psi}^I \gamma_{\mu_1} \gamma_5 \mathcal{D}_{\mu_2} \dots \mathcal{D}_{\mu_n} \psi^K \quad (3)$$

for even n .

In the leading order, there are three diagrams Fig.1 that contribute to the anomalous dimension of this operator. The diagram in Fig.1.a is the same as in QCD [31] and together with its mirror image gives

$$\gamma_{\text{Fig.1.a+m.d.}}^{(0)} = 2 \times (2 S_1(n) - 2). \quad (4)$$

The diagram in Fig.1.b is also the same as in QCD [31] with the following result

$$\gamma_{\text{Fig.1.b}}^{(0)} = -\frac{2}{n(n+1)}. \quad (5)$$

The diagram in Fig.1.c contains scalar or pseudoscalar fields and the result is the same in both cases, and the overall result is

$$\gamma_{\text{Fig.1.c}}^{(0)} = 2 \times \frac{1}{n(n+1)}. \quad (6)$$

Putting it all together, we get

$$\gamma_{\tilde{\mathcal{O}}_{15}^{qq}}^{(0)}(n) = 4 S_1(n), \quad (7)$$

where the constant in eq.(4) cancels out when the external fermionic field is renormalised.

In what follows, we compute the contribution from insertions of fermion and, in general, scalar loops into gauge and scalar propagators in all order of perturbative theory. Our task is to study the maximal transcendentality principle for the contribution proportional to the number of fermions in order to have additional information that can be used for the reconstruction of the corresponding result in QCD.

For such calculations we use the method proposed in Refs. [39, 40] and applied to the computations of the anomalous dimension of non-singlet twist-2 operator in QCD in Ref. [38]. The main idea is that we have to compute the diagrams in Fig.1 with propagators with critical exponents related with critical indices, which can be calculated at the critical point through the expansion in the number of quark (scalar) flavors or the number of quark (scalar) loops. So, we have to use propagators in the following form [38–40, 42–45]

$$\psi = \frac{A \not{k}}{(k^2)^{\mu-\alpha}}, \quad \mathcal{A}_{\mu\nu} = \frac{B g_{\mu\nu}}{(k^2)^{\mu-\beta}}, \quad \phi = \frac{C}{(k^2)^{\mu-\gamma}}, \quad (8)$$

where A , B and C are momentum independent amplitudes, $\mu = 2 - \epsilon$ and α , β and γ are the critical exponents. These amplitudes or their combinations can be fixed by considering the skeleton Dyson equations with dressed propagators [39, 40], which can be represented graphically in the case of gauge and scalar propagators with insertions of fermionic fields as [38, 42–45]

$$0 = \mathcal{A}_{\mu\nu}^{-1} + \text{diagram} \quad (9)$$

$$0 = \phi^{-1} + \text{diagram} \quad (10)$$

Strictly speaking, eq.(9) is correct only for the gauge field propagator in the Landau gauge $\mathcal{A}_{\mu\nu} = (g_{\mu\nu} - k_\mu k_\nu / k^2) / (k^2)^{\mu-\beta}$, since only the transverse part of the gauge field propagator ensures the correct behaviour of the computed quantities [46, 47], but for the computations of the gauge-invariant quantities, such as anomalous dimension, for the leading N_f contribution one can choose for simplicity the Feynman gauge, as was done in [38]. After substituting the propagators from eq.(8) we obtain the sum of general one-loop integrals, which can be computed using the well-known Tkachov formula [48]

$$\int \frac{d^D P}{(2\pi)^D} \frac{\mathcal{P}(P)}{P^{2\alpha} (P-Q)^{2\beta}} = \frac{1}{(4\pi)^2} (Q^2)^{D/2-\alpha-\beta} \sum_{\sigma \geq 0}^{[n/2]} G(\alpha, \beta, n, \sigma) Q^{2\sigma} \left\{ \frac{1}{\sigma!} \left(\frac{\square}{4} \right)^\sigma \mathcal{P}_n(P) \right\}_{P=Q} \quad (11)$$

where

$$\mathcal{P}_n(P) = P_{\mu_1} P_{\mu_2} \cdots P_{\mu_n}, \quad \square = \frac{\partial^2}{\partial P_\mu \partial P^\mu}, \quad (12)$$

D is the dimension of space-time $D = 4 - 2\epsilon$ and G can be expressed in terms of Γ -functions:

$$G(\alpha, \beta, n, \sigma) = (4\pi)^\epsilon \frac{\Gamma(\alpha + \beta - \sigma - D/2) \Gamma(D/2 - \alpha + n - \sigma) \Gamma(D/2 - \beta + \sigma)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(D - \alpha - \beta + n)}. \quad (13)$$

We use the Feynman rules from Refs. [49, 50], which we used for all our previous computations. The calculations of eqs.(9) and (10) after substitution of expressions for critical

indices in the leading approximation $\alpha = \mu - 1$ and $\beta = 1$ fixes the combinations of amplitudes A^2B and A^2C as [38, 42–45]

$$A^2B = \frac{\Gamma[2\mu]}{2 N_f \Gamma[2 - \mu] \Gamma[\mu]^2}, \quad (14)$$

$$A^2C = \frac{\Gamma[2\mu - 1]}{N_f \Gamma[1 - \mu] \Gamma[\mu]^2}. \quad (15)$$

These results will be used to calculate the diagrams in Fig.1 using the expressions for the propagators from eq.(8).

We now turn to the calculation of the diagrams in Fig.1 using the expressions for critical propagators from eq.(8). However, if we compute the obtained integrals and expand them after the substitution of the critical indices $\alpha = \mu - 1 + \eta/2$ and $\beta = 1 - \eta - \chi_A$ and $\gamma = 1 - \eta - \chi_\phi$, where η is the anomalous dimension of the fermions and χ_A and χ_ϕ are the anomalous dimensions of the respective vertices*, we will find that the integral diverges and therefore requires regularisation. This can be done by shifting the exponents of the gauge and scalar fields by an infinitesimal quantity δ , $\beta \rightarrow \beta - \delta$ [38, 42, 45, 51]. With such regularisation, all diagrams in Fig.1 will have the following formal structure after expanding near $\delta \rightarrow 0$ [38, 44, 51, 52]

$$\frac{P}{\delta} + Q + R \ln p^2 + \mathcal{O}(\delta), \quad (16)$$

where P , Q and R depend on μ and p is the external momentum following through the fermions. The simple pole is absorbed by conventional (critical point) renormalisation [51, 52]. The δ -finite Green's function must resum to the structure $(p^2)^{\gamma_{\mathcal{O}}^{(n)}/2}$ [51, 52], where $\gamma_{\mathcal{O}}^{(n)}$ is related to the exponent we are interested in, $\eta_1^{(n)}$, through [38, 44, 51, 52]:

$$\eta^{(n)} = \eta + \gamma_{\mathcal{O}}^{(n)} \quad (17)$$

and is equal to R from eq.(16) or $(-P)$ in the leading N_f order.

The diagrams Fig.1.a and Fig.1.b are the same as in QCD with the following already computed results [38]:

$$\eta_{\text{Fig.1.a+m.d.}}^{(n)} = \frac{\mu(\mu - 1)\Gamma[2\mu]}{\Gamma[\mu]^2\Gamma[\mu + 1]\Gamma[2 - \mu]N_f} \sum_{j=2}^n \frac{1}{\mu + j - 2}, \quad (18)$$

$$\eta_{\text{Fig.1.b}}^{(n)} = -\frac{\mu(\mu - 1)^3\Gamma[2\mu]}{2(\mu + n - 1)(\mu + n - 2)\Gamma[\mu]^2\Gamma[\mu + 1]\Gamma[2 - \mu]N_f}. \quad (19)$$

Expanding the sum of these results in ϵ in the critical point \hat{a}_s where the one-loop QCD β -function is equal to zero in the leading N_f order ($a_s = \alpha_s/(4\pi)$)

$$\beta(a_s) = -\epsilon a_s + \left(\frac{2}{3}N_f - \frac{11}{3}C_A\right)a_s^2 + \mathcal{O}(a_s^3), \quad \hat{a}_s = \frac{3\epsilon}{2N_f}, \quad \epsilon = \frac{2\hat{a}_s N_f}{3}, \quad (20)$$

*In the leading N_f order they do not give contributions into result, see Refs. [38, 44, 45]




Figure 2: Two-loop diagrams with fermion and scalar loops insertions.

we obtain the results for the leading N_f contribution [38], which coincide in the first four orders of perturbative theory with the corresponding results from Refs. [53–55]. Note that in the Landau gauge, only eq.(19) gets the constant $(-1/2)$ multiplied by n -independent factor from eq.(19).

The diagram Fig.1.c contains the scalar field and together with the result for A^2C from eq.(15) gives

$$\eta_{\text{Fig.1.c}}^{(n)} = \frac{\mu(\mu-1)^3\Gamma[2\mu-1]}{(\mu+n-1)(\mu+n-2)\Gamma[\mu]^2\Gamma[\mu+1]\Gamma[2-\mu]N_f}. \quad (21)$$

Moreover, we must take into account the contribution from the so-called ϵ -scalars, that appear in the framework of the dimensional reduction scheme [56–59]. Their contribution is the same as contribution of the scalars (21), but multiplied by $\epsilon = 2 - \mu$, which together with eq.(21) gives

$$\eta_{\text{Fig.1.c}+\epsilon\text{-sc}}^{(n)} = \frac{\mu(\mu-1)^4\Gamma[2\mu-1]}{(\mu+n-1)(\mu+n-2)\Gamma[\mu]^2\Gamma[\mu+1]\Gamma[2-\mu]N_f}. \quad (22)$$

One can see, that the results (19) and (22) for the diagrams Fig.1.b and Fig.1.c have different structures and their sum will produce terms proportional to $(1/n/(n+1))^i$ after expansion in ϵ , that is, in higher orders of perturbative theory, while we expect such terms to be absent, as in the expression for the universal anomalous dimension of twist-2 operators in $\mathcal{N} = 4$ SYM theory [30, 60, 61] (see eq.(7)).

To analyse this situation, we used our previous calculations for the two-loop anomalous dimension of twist-2 operators in $\mathcal{N}=4$ SYM theory [30]. During that calculations we obtained the results for individual diagrams for an arbitrary Lorentz spin of operators. Using these results, we tried to find a combination of diagrams that do not have such terms as $1/n/(n+1)$ in the sum. We found, that it is enough to add to the corresponding two-loops diagrams Fig.2.a and 2.b, whose contributions are respectively

$$\gamma_{\text{Fig.2.a}}^{(1)} = N_f \left(-\frac{4}{3n^2} + \frac{4}{3(n+1)^2} + \frac{44}{9n} - \frac{44}{9(n+1)} \right), \quad (23)$$

$$\gamma_{\text{Fig.2.b}}^{(1)} = N_f \left(\frac{2}{n^2} - \frac{2}{(n+1)^2} - \frac{6}{n} + \frac{6}{n+1} \right), \quad (24)$$

only one diagram Fig.2.c with the scalar loop in the gauge field, with the following result

$$\gamma_{\text{Fig.2.c}}^{(1)} = N_s \left(-\frac{1}{3n^2} + \frac{1}{3(n+1)^2} + \frac{14}{9n} - \frac{14}{9(n+1)} \right), \quad (25)$$

and taking into account the contribution of two ϵ -scalars that must be added within the dimensional reduction scheme [56–58] and is given by

$$\gamma_{\text{Fig.2.b} \times 2\epsilon}^{(1)} = N_f \left(-\frac{2}{n} + \frac{2}{n+1} \right), \quad (26)$$

but if we relate the number of scalar fields N_s with the number of fermion fields N_f . In $\mathcal{N} = 1$, $\mathcal{N} = 2$ and $\mathcal{N} = 4$ SYM theories, the number of scalar fields N_s is related to the number of fermion fields N_f through $N_s = 2N_f - 2$. Thus, if we express N_s through N_f in Fig.2.c, then the sum of all three diagrams in Fig.2 will give zero for the contribution proportional to N_f . This cancellation occurs in higher loops as well, as we checked in three loops by explicit computations with MINCER-package [62] for FORM [63] the relevant three-loop diagrams, which can be obtained by inserting scalar and fermion loops into the gauge propagators in diagrams Fig.2.a and Fig.2.c and inserting the fermion loops into the scalar propagators in diagram Fig.2.b and in three, four and five loops by the direct computations with Tkachov formula (11). We consider only such minimal gauge-invariant subclass of diagrams, since it corresponds to the leading N_f computations in the original paper [39] and in QED [42–44] or QCD [38, 45], while we want to get results close to the result in QCD. The inclusion of other diagrams will violate the cancellation of $1/n$ terms.

For calculations in the general case with critical exponents, one should additionally consider in eq.(9) the diagram with the scalar loop

$$0 = \mathcal{A}_{\mu\nu}^{-1} + \text{diagram with fermion loop in gauge field} + \text{diagram with scalar loop in gauge field} \quad (27)$$

and do not take into account N_f expansion for the scalar propagator here. The result for eq.(27) has the following form

$$0 = 1 - 2A^2B N_f (1 - \mu) \frac{\Gamma[1 - \mu] \Gamma[\mu]^2}{\Gamma[2\mu]} + C^2B N_s \frac{\Gamma[1 - \mu] \Gamma[\mu]^2}{2\Gamma[2\mu]} \quad (28)$$

Substituting the expression for A^2B into eq.(19) divided by eq.(14) we fix C^2B and then A^2B demanding the cancellation with eq.(22)

$$C^2B N_s = 2 \frac{(\mu - 1)\Gamma[2\mu - 1]}{\Gamma[2 - \mu]\Gamma[\mu]^2}, \quad A^2B N_f = \frac{(\mu - 1)\Gamma[2\mu - 1]}{\Gamma[2 - \mu]\Gamma[\mu]^2}. \quad (29)$$

The results for A^2B and C^2B can be obtained by the direct computations of the critical indices for the fermion and scalar fields and using the Slavnov-Taylor identity between the fermion-fermion-gauge field and scalar-scalar-gauge field vertices, similar to the computations of the critical indices for gluon and ghost fields in QCD in Refs. [45, 47], but this requires additional consideration of the relevant vertex diagrams.

So, we are left with one diagram Fig.1.a, which contributes to the critical index for the anomalous dimension. Its contribution is as follows (with A^2B from eq.(29))

$$\eta_{\text{Fig.1.a+m.d.}}^{(n)} = 2 A^2 B \frac{\mu(\mu-1)}{\Gamma[\mu+1]} \sum_{j=2}^n \frac{1}{\mu+j-2} = 2 \frac{\mu(\mu-1)^2}{N_f \Gamma[\mu+1]} \frac{\Gamma[2\mu-1]}{\Gamma[2-\mu] \Gamma[\mu]^2} \sum_{j=2}^n \frac{1}{\mu+j-2}. \quad (30)$$

Remember now the expression for the one-loop beta function in $\mathcal{N} = 1$, $\mathcal{N} = 2$ and $\mathcal{N} = 4$ SYM theories [31,64]

$$\beta_{\text{SUSY}}(g) = -\epsilon g^2 + \left(\frac{2}{3} N_f + \frac{1}{6} N_s - \frac{11}{3} \right) g^4 \quad (31)$$

and substituting the above relation between the number of fermionic and scalar fields $N_s = 2N_f - 2$, we obtain at the critical point g_c , where β_{SUSY} equals zero

$$\beta_{\text{SUSY}}(g) = -\epsilon g^2 + (N_f - 4) g^4, \quad g_c^2 = \frac{\epsilon}{N_f}, \quad \epsilon = N_f g_c^2. \quad (32)$$

Expanding the expression for $\eta^{(n)}$ from eq.(30) in ϵ and using the relation between ϵ and g^2 at the critical point (32) one can obtain the expression for the anomalous dimension of $\tilde{\mathcal{O}}_{15}^{qq}$ operator in the number of fermionic loops[†]

$$\begin{aligned} \hat{\gamma}_{N_f}^{(n)} &= 4 S_1 g^2 + 4(S_2 - 2 S_1) g^4 N_f + 4(S_3 - 2 S_2) g^6 N_f^2 + 4(S_4 - 2 S_3 + 2 \zeta_3 S_1) g^8 N_f^3 \\ &\quad + 4(S_5 - 2 S_4 + 2 \zeta_3 S_2 - 4 \zeta_3 S_1 + 3 \zeta_4 S_1) g^{10} N_f^4 + \dots \end{aligned} \quad (33)$$

where the nested harmonic sums $S_{\vec{a}} = S_{\vec{a}}(n)$ are defined as [65]

$$S_i(M) = \sum_{j=1}^M \frac{\text{sign}(i)^j}{j^{|i|}}, \quad S_{i_1, i_2, \dots, i_k}(M) = \sum_{j=1}^M \frac{\text{sign}(i_1)^j}{j^{|i_1|}} S_{i_2, \dots, i_k}(j). \quad (34)$$

The explicit calculations with MINCER [62] give the same results for the first two N_f terms in eq.(33). This expression can be compared with the analogies expression for the leading N_f (the number of fermionic loop) contribution in QCD [38], which looks like:

$$\begin{aligned} \hat{\gamma}_{\text{NS}}^{(n)} &= 4 S_1 C_F \alpha_s + \left(\frac{8}{3} S_2 - \frac{40}{9} S_1 \right) C_F n_f \alpha_s^2 + \left(\frac{16}{9} S_3 - \frac{80}{27} S_2 - \frac{16}{27} S_1 \right) C_F n_f^2 \alpha_s^3 \\ &\quad + \left(\frac{32}{27} S_4 - \frac{160}{81} S_3 - \frac{32}{81} S_2 - \frac{32}{81} S_1 + \frac{64}{27} \zeta_3 S_1 \right) C_F n_f^3 \alpha_s^4 + \dots, \end{aligned} \quad (35)$$

where we left only the terms proportional to S_i .

Let us now try to relate the obtained result to integrability. Namely, we will try to modify the results of integrability in $\mathcal{N} = 4$ SYM theory to include the presented results (33). According to the relation between the computation of the energy of integrable

[†]The renormalisation of the external fermions will change the summation over j from $j = 2$ to $j = 1$.

spin chain and the anomalous dimension of composite operators, we should consider only one-loop diagrams with fermionic and scalar loops insertions into the gauge propagator in Fig.1.a and Fig.1.b and fermionin loops insertions into scalar propagator on Fig.1.c. The most simple way is to consider the solution of the Baxter's equation [66–68] and modify it in such a way, that the final results match our result (33). The Baxter function for twist-2 operators in leading order is expressed through the hypergeometric function

$$Q^{(0)}(u, M) = {}_3F_2 \left(\begin{matrix} -M, M+1, \frac{1}{2} + iu \\ 1, 1 \end{matrix} \middle| 1 \right), \quad i \left(\partial_u Q^{(0)}(u, M) \right) \Big|_{u=-i/2}^{u=i/2} = 4 S_1(M) \quad (36)$$

and the anomalous dimension in the leading order, equal to $4S_1(M)$, can be found in the standard way by differentiating with respect to u and setting u equals to $i/2$. Its modifications for higher order were studied in Ref. [69, 70] and we assume that since our two-loop result (33) gives a part of the full result in the second order, therefore, the next-to-leading order modification of the Baxter function has to contain its in some way. We found that the following δ -modification of the Baxter function from Ref. [69, 70]

$$Q_\delta^{(0)}(u, M) = {}_3F_2 \left(\begin{matrix} -M, M+1+2\delta, \frac{1}{2} + iu \\ 1+\delta, 1 \end{matrix} \middle| 1 \right) \quad (37)$$

will give the necessary contributions to the anomalous dimension according to the general formula

$$i \left(\partial_u \left(\partial_\delta^k Q_\delta^{(0)}(u, M) \right) \Big|_{\delta=0} \right) \Big|_{u=-i/2}^{u=i/2} = 2 (-1)^k k! S_{k+1} \left(\frac{M}{2} \right). \quad (38)$$

The modified Baxter function that will give our result (33) can be written as:

$$Q_{N_f}^{\text{LA}}(u, M) = \frac{\mu(\mu-1)^2}{N_f \Gamma[\mu+1]} \frac{\Gamma[2\mu-1]}{\Gamma[2-\mu] \Gamma[\mu]^2} (2 Q_{-\epsilon}^{(0)}(u, 2M) - 2 Q^{(0)}(u, 2M) + Q^{(0)}(u, M)), \quad \mu = 2 - \epsilon. \quad (39)$$

Expanding the above equation in ϵ (32) and using eq.(36) we reproduce our result (33).

However, if we try to go with such minimal modification to the subleading order in N_f expansion, starting with the two-loop Baxter function and its modification similar to eq.(39)

$$Q_{N_f}^{(1)}(u, M) = \frac{\mu(\mu-1)^2}{N_f \Gamma[\mu+1]} \frac{\Gamma[2\mu-1]}{\Gamma[2-\mu] \Gamma[\mu]^2} Q_\epsilon^{(1)}(u, M), \quad (40)$$

with $Q_\delta^{(1)}(u, M)$ from Ref. [69, 70] we will not obtain the leading transcendental structures from available results in QCD, which can be extracted from Refs. [53–55] and have the

following form:

$$\hat{\gamma}_{\text{NS}}^{(1)} = 8C_F(C_A - 2C_F)(2S_{1,-2} - S_{-3} + S_3), \quad (41)$$

$$\begin{aligned} \hat{\gamma}_{\text{NS}}^{(2),n_f} &= \frac{16}{3}n_f C_F \left[(C_A - 2C_F)(4S_{1,-2,1} - 2S_{1,-3} - 2S_{-3,1} + S_{-4} + 3S_{1,3} - S_{3,1} - S_4) \right. \\ &\quad \left. + C_F(2S_{1,3} - 2S_{3,1}) \right], \end{aligned} \quad (42)$$

$$\begin{aligned} \hat{\gamma}_{\text{NS}}^{(3),n_f^2} &= \frac{32}{9}n_f^2 C_F \left[(C_A - 2C_F)(8S_{1,-2,1,1} - 4S_{1,-2,2} - 4S_{1,-3,1} - 4S_{-3,1,1} + 2S_{1,-4} \right. \\ &\quad \left. + 2S_{-4,1} + 2S_{-3,2} - S_{-5} + 4S_{1,-2,-2} - 2S_{-3,-2} - S_{1,4} - 2S_{1,3,1} + S_{4,1} + S_5) \right. \\ &\quad \left. - 2C_F(2S_{1,3,1} - S_{1,4} - S_{4,1}) \right]. \end{aligned} \quad (43)$$

Nevertheless, we observed that if we pass from the usual anomalous dimension to the reciprocity-respecting anomalous dimension [71, 72] the above equations change to

$$\hat{\gamma}_{\text{NS,RR}}^{(1)} = C_F(2C_F - C_A)(4\mathbb{B}_{1,2} - 4\mathbb{B}_{2,1}), \quad (44)$$

$$\hat{\gamma}_{\text{NS,RR}}^{(2),n_f} = n_f C_F \left[(2C_F - C_A) \left(\frac{8}{3}\mathbb{B}_{1,3} - 4\mathbb{B}_{1,2,1} + \frac{4}{3}\mathbb{B}_{2,1,1} \right) + C_F \left(\frac{8}{3}\mathbb{B}_{1,2,1} - \frac{8}{3}\mathbb{B}_{2,1,1} \right) \right], \quad (45)$$

$$\begin{aligned} \hat{\gamma}_{\text{NS,RR}}^{(3),n_f^2} &= n_f^2 C_F \left[(2C_F - C_A) \left(\frac{16}{9}\mathbb{B}_{1,4} - \frac{16}{9}\mathbb{B}_{1,3,1} - \frac{8}{9}\mathbb{B}_{2,2,1} + \frac{8}{9}\mathbb{B}_{1,2,1,1} \right) \right. \\ &\quad \left. + C_F^2 \left(\frac{16}{9}\mathbb{B}_{2,2,1} - \frac{16}{9}\mathbb{B}_{1,2,1,1} \right) \right], \end{aligned} \quad (46)$$

where the binomial harmonic sums $\mathbb{B}_{\vec{a}} = \mathbb{B}_{\vec{a}}(M)$ are defined as [65]

$$\mathbb{B}_{\vec{a}}(M) = \sum_{k=1}^M \binom{M}{k} \binom{M+k}{k} S_{\vec{a}}(k). \quad (47)$$

The first $\mathbb{B}_{\vec{a}}$ in eqs.(44)-(46) that correspond to the terms in eqs.(41)-(43) with the maximal number of indices can be generated by the following sequence of the binomial sums:

$$\mathbb{B}_{1,2} \quad \mapsto \quad \mathbb{B}_{1,3} \quad \mapsto \quad \mathbb{B}_{1,4}, \quad (48)$$

that is, from $\mathbb{B}_{1,2}$ formally differentiated by the last index with exactly the same relative coefficients as for the leading n_f contributions (35). It seems, that the most complicated part for the subleading N_f contribution can be generated using an expression similar to eq.(18) but with the usual harmonic sums changed to binomial harmonic sums. Additional information can be obtained from the relevant direct diagrammatic calculations, which are in progress both in QCD and in $\mathcal{N} = 4$ SYM theory.

We expect that in higher loops there is some set of diagrams that gives the most transcendent terms to N_f contribution, and this set of diagrams contains QCD diagrams and diagrams in which gauge fields are replaced by scalar fields, canceling the ‘‘simple’’

terms, especially with $1/n$, as discussed here. The number of terms in the final result will decrease due to such cancellations. This will be very important for the four-loop computations, because so far the only way to get the full result for the anomalous dimension of twist-2 operators in four loops is to compute the results for the first few fixed moments of operators and reconstruct the general result using number theory, assume that the final results will consist of nested harmonic sums[‡]. Any additional information, in particular, that can be obtained by extending the consideration presented in this paper, will allow us finally to find the general result for n_f contribution to the four-loop anomalous dimension of the non-singlet twist-2 operator in QCD.

Acknowledgments

The research was supported by a grant from the Russian Science Foundation No. 22-22-00803, <https://rscf.ru/en/project/22-22-00803/>.

References

- [1] J.M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [[hep-th/9711200](#)].
- [2] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)].
- [3] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)].
- [4] J.A. Minahan and K. Zarembo, *The Bethe ansatz for N=4 superYang-Mills*, *JHEP* **03** (2003) 013 [[hep-th/0212208](#)].
- [5] N. Beisert, C. Kristjansen and M. Staudacher, *The Dilatation operator of conformal N=4 superYang-Mills theory*, *Nucl. Phys. B* **664** (2003) 131 [[hep-th/0303060](#)].
- [6] N. Beisert and M. Staudacher, *The N=4 SYM integrable super spin chain*, *Nucl. Phys. B* **670** (2003) 439 [[hep-th/0307042](#)].
- [7] I. Bena, J. Polchinski and R. Roiban, *Hidden symmetries of the AdS(5) x S**5 superstring*, *Phys. Rev. D* **69** (2004) 046002 [[hep-th/0305116](#)].
- [8] V.A. Kazakov, A. Marshakov, J.A. Minahan and K. Zarembo, *Classical/quantum integrability in AdS/CFT*, *JHEP* **05** (2004) 024 [[hep-th/0402207](#)].

[‡]Such method, proposed by us in Ref. [73], was successfully applied for the computation at four loops for the non-planar contribution to the universal anomalous dimension in $\mathcal{N} = 4$ SYM theory [74, 75], n_f^2 contribution to the non-singlet anomalous dimension in QCD [55] and planar limit of the non-singlet anomalous dimension in QCD [76].

- [9] N. Beisert, V. Dippel and M. Staudacher, *A Novel long range spin chain and planar $N=4$ super Yang-Mills*, *JHEP* **07** (2004) 075 [[hep-th/0405001](#)].
- [10] G. Arutyunov, S. Frolov and M. Staudacher, *Bethe ansatz for quantum strings*, *JHEP* **10** (2004) 016 [[hep-th/0406256](#)].
- [11] M. Staudacher, *The Factorized S-matrix of CFT/AdS*, *JHEP* **05** (2005) 054 [[hep-th/0412188](#)].
- [12] N. Beisert and M. Staudacher, *Long-range $psu(2,2-4)$ Bethe Ansatz for gauge theory and strings*, *Nucl. Phys. B* **727** (2005) 1 [[hep-th/0504190](#)].
- [13] R.A. Janik, *The AdS(5) x S**5 superstring worldsheet S-matrix and crossing symmetry*, *Phys. Rev. D* **73** (2006) 086006 [[hep-th/0603038](#)].
- [14] B. Eden and M. Staudacher, *Integrability and transcendentality*, *J. Stat. Mech.* **0611** (2006) P11014 [[hep-th/0603157](#)].
- [15] N. Beisert, B. Eden and M. Staudacher, *Transcendentality and Crossing*, *J. Stat. Mech.* **0701** (2007) P01021 [[hep-th/0610251](#)].
- [16] N. Beisert et al., *Review of AdS/CFT Integrability: An Overview*, *Lett. Math. Phys.* **99** (2012) 3 [[1012.3982](#)].
- [17] G. Arutyunov and S. Frolov, *String hypothesis for the AdS(5) x S**5 mirror*, *JHEP* **03** (2009) 152 [[0901.1417](#)].
- [18] N. Gromov, V. Kazakov and P. Vieira, *Exact spectrum of anomalous dimensions of planar $N=4$ supersymmetric Yang-Mills theory*, *Phys. Rev. Lett.* **103** (2009) 131601 [[0901.3753](#)].
- [19] D. Bombardelli, D. Fioravanti and R. Tateo, *Thermodynamic Bethe Ansatz for planar AdS/CFT: A Proposal*, *J. Phys. A* **42** (2009) 375401 [[0902.3930](#)].
- [20] N. Gromov, V. Kazakov, A. Kozak and P. Vieira, *Exact spectrum of anomalous dimensions of planar $N = 4$ supersymmetric Yang-Mills theory: TBA and excited states*, *Lett. Math. Phys.* **91** (2010) 265 [[0902.4458](#)].
- [21] G. Arutyunov and S. Frolov, *Thermodynamic Bethe Ansatz for the AdS(5) x S(5) Mirror Model*, *JHEP* **05** (2009) 068 [[0903.0141](#)].
- [22] N. Gromov, V. Kazakov, S. Leurent and D. Volin, *Quantum spectral curve for planar $\mathcal{N} = 4$ Super-Yang-Mills theory*, *Phys. Rev. Lett.* **112** (2014) 011602 [[1305.1939](#)].
- [23] N. Gromov, V. Kazakov, S. Leurent and D. Volin, *Quantum spectral curve for arbitrary state/operator in AdS₅/CFT₄*, *JHEP* **09** (2015) 187 [[1405.4857](#)].
- [24] A.V. Kotikov and L.N. Lipatov, *DGLAP and BFKL equations in the $N = 4$ supersymmetric gauge theory*, *Nucl. Phys. B* **661** (2003) 19 [[hep-ph/0208220](#)].
- [25] L.N. Lipatov, *Reggeization of the vector meson and the vacuum singularity in non-abelian gauge theories*, *Sov. J. Nucl. Phys.* **23** (1976) 338.

- [26] E.A. Kuraev, L.N. Lipatov and V.S. Fadin, *The Pomeron singularity in non-abelian gauge theories*, *Sov. Phys. JETP* **45** (1977) 199.
- [27] I.I. Balitsky and L.N. Lipatov, *The Pomeron singularity in quantum chromodynamics*, *Sov. J. Nucl. Phys.* **28** (1978) 822.
- [28] A.V. Kotikov and L.N. Lipatov, *NLO corrections to the BFKL equation in QCD and in supersymmetric gauge theories*, *Nucl. Phys. B* **582** (2000) 19 [[hep-ph/0004008](#)].
- [29] V.S. Fadin and L.N. Lipatov, *BFKL pomeron in the next-to-leading approximation*, *Phys. Lett. B* **429** (1998) 127 [[hep-ph/9802290](#)].
- [30] A.V. Kotikov, L.N. Lipatov and V.N. Velizhanin, *Anomalous dimensions of Wilson operators in $N=4$ SYM theory*, *Phys. Lett. B* **557** (2003) 114 [[hep-ph/0301021](#)].
- [31] D.J. Gross and F. Wilczek, *Asymptotically free gauge theories - I*, *Phys. Rev. D* **8** (1973) 3633.
- [32] H. Georgi and H.D. Politzer, *Electroproduction scaling in an asymptotically free theory of strong interactions*, *Phys. Rev. D* **9** (1974) 416.
- [33] E.G. Floratos, D.A. Ross and C.T. Sachrajda, *Higher order effects in asymptotically free gauge theories: the anomalous dimensions of Wilson operators*, *Nucl. Phys. B* **129** (1977) 66.
- [34] A. Gonzalez-Arroyo, C. Lopez and F.J. Yndurain, *Second order contributions to the structure functions in deep inelastic scattering. 1. Theoretical calculations*, *Nucl. Phys. B* **153** (1979) 161.
- [35] R. Kirschner and L.N. Lipatov, *Double logarithmic asymptotics of quark scattering amplitudes with flavor exchange*, *Phys. Rev. D* **26** (1982) 1202.
- [36] R. Kirschner and L.n. Lipatov, *Double logarithmic asymptotics and Regge singularities of quark amplitudes with flavor exchange*, *Nucl. Phys. B* **213** (1983) 122.
- [37] V.N. Velizhanin, *Generalised double-logarithmic equation in QCD*, *Mod. Phys. Lett. A* **32** (2017) 1750213 [[1412.7143](#)].
- [38] J.A. Gracey, *Anomalous dimension of nonsinglet Wilson operators at $O(1/N(f))$ in deep inelastic scattering*, *Phys. Lett. B* **322** (1994) 141 [[hep-ph/9401214](#)].
- [39] A.N. Vasiliev, Y.M. Pismak and Y.R. Khonkonen, *Simple method of calculating the critical indices in the $1/N$ expansion*, *Theor. Math. Phys.* **46** (1981) 104.
- [40] A.N. Vasiliev, Y.M. Pismak and Y.R. Khonkonen, *$1/N$ expansion: calculation of the exponents η and Nu in the order $1/N^2$ for arbitrary number of dimensions*, *Theor. Math. Phys.* **47** (1981) 465.
- [41] A.V. Belitsky, S.E. Derkachov, G.P. Korchemsky and A.N. Manashov, *Superconformal operators in $N=4$ superYang-Mills theory*, *Phys. Rev. D* **70** (2004) 045021 [[hep-th/0311104](#)].

- [42] J.A. Gracey, *Quantum electrodynamics at $O(1/N_f(2))$ in arbitrary dimensions*, *Mod. Phys. Lett. A* **7** (1992) 1945.
- [43] J.A. Gracey, *Algorithm for computing the beta function of quantum electrodynamics in the large $N(f)$ expansion*, *Int. J. Mod. Phys. A* **8** (1993) 2465 [[hep-th/9301123](#)].
- [44] J.A. Gracey, *Electron mass anomalous dimension at $O(1/(N_f(2)))$ in quantum electrodynamics*, *Phys. Lett. B* **317** (1993) 415 [[hep-th/9309092](#)].
- [45] J.A. Gracey, *Quark, gluon and ghost anomalous dimensions at $O(1/N(f))$ in quantum chromodynamics*, *Phys. Lett. B* **318** (1993) 177 [[hep-th/9310063](#)].
- [46] M. Ciuchini, S.E. Derkachov, J.A. Gracey and A.N. Manashov, *Quark mass anomalous dimension at $O(1/N(f)^{**2})$ in QCD*, *Phys. Lett. B* **458** (1999) 117 [[hep-ph/9903410](#)].
- [47] M. Ciuchini, S.E. Derkachov, J.A. Gracey and A.N. Manashov, *Computation of quark mass anomalous dimension at $O(1 / N^{**2}(f))$ in quantum chromodynamics*, *Nucl. Phys. B* **579** (2000) 56 [[hep-ph/9912221](#)].
- [48] F.V. Tkachov, *A theorem on analytical calculability of four loop renormalization group functions*, *Phys. Lett. B* **100** (1981) 65.
- [49] L. Brink, J.H. Schwarz and J. Scherk, *Supersymmetric Yang-Mills theories*, *Nucl. Phys. B* **121** (1977) 77.
- [50] F. Gliozzi, J. Scherk and D.I. Olive, *Supersymmetry, supergravity theories and the dual spinor model*, *Nucl. Phys. B* **122** (1977) 253.
- [51] A.N. Vasiliev and M.Y. Nalimov, *Analog of dimensional regularization for calculation of the renormalization group functions in the $1/n$ expansion for arbitrary dimension of space*, *Theor. Math. Phys.* **55** (1983) 423.
- [52] A.N. Vasiliev and M.Y. Nalimov, *The $CP^{*(n-1)}$ model: calculation of anomalous dimensions and the mixing matrices in the order $1/N$* , *Theor. Math. Phys.* **56** (1983) 643.
- [53] S. Moch, J.A.M. Vermaseren and A. Vogt, *Nonsinglet structure functions at three loops: Fermionic contributions*, *Nucl. Phys. B* **646** (2002) 181 [[hep-ph/0209100](#)].
- [54] S. Moch, J.A.M. Vermaseren and A. Vogt, *The three loop splitting functions in QCD: The nonsinglet case*, *Nucl. Phys. B* **688** (2004) 101 [[hep-ph/0403192](#)].
- [55] J. Davies, A. Vogt, B. Ruijl, T. Ueda and J.A.M. Vermaseren, *Large- n_f contributions to the four-loop splitting functions in QCD*, *Nucl. Phys. B* **915** (2017) 335 [[1610.07477](#)].
- [56] W. Siegel, *Supersymmetric dimensional regularization via dimensional reduction*, *Phys. Lett. B* **84** (1979) 193.

- [57] P.K. Townsend and P. van Nieuwenhuizen, *Dimensional regularization and supersymmetry at the two loop level*, *Phys. Rev. D* **20** (1979) 1832.
- [58] E. Sezgin, *Dimensional regularization and the massive Wess-Zumino model*, *Nucl. Phys. B* **162** (1980) 1.
- [59] D.M. Capper, D.R.T. Jones and P. van Nieuwenhuizen, *Regularization by Dimensional Reduction of Supersymmetric and Nonsupersymmetric Gauge Theories*, *Nucl. Phys. B* **167** (1980) 479.
- [60] L.N. Lipatov, *Next-to-leading corrections to the BFKL equation and the effective action for high energy processes in QCD*, *Nucl. Phys. B Proc. Suppl.* **99** (2001) 175.
- [61] A.V. Kotikov, L.N. Lipatov, A.I. Onishchenko and V.N. Velizhanin, *Three loop universal anomalous dimension of the Wilson operators in $N = 4$ SUSY Yang-Mills model*, *Phys. Lett. B* **595** (2004) 521 [[hep-th/0404092](#)].
- [62] S.G. Gorishnii, S.A. Larin, L.R. Surguladze and F.V. Tkachov, *Mincer: program for multiloop calculations in quantum field theory for the Schoonschip system*, *Comput. Phys. Commun.* **55** (1989) 381.
- [63] J.A.M. Vermaseren, *New features of FORM*, [math-ph/0010025](#).
- [64] D.R.T. Jones, *Asymptotic behavior of supersymmetric Yang-Mills theories in the two loop approximation*, *Nucl. Phys. B* **87** (1975) 127.
- [65] J.A.M. Vermaseren, *Harmonic sums, Mellin transforms and integrals*, *Int. J. Mod. Phys. A* **14** (1999) 2037 [[hep-ph/9806280](#)].
- [66] R.J. Baxter, *One-dimensional anisotropic Heisenberg chain*, *Annals Phys.* **70** (1972) 323.
- [67] R.J. Baxter, *Eight vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain. 1. Eigenvectors of the transfer matrix and Hamiltonian*, *Annals Phys.* **76** (1973) 48.
- [68] E.K. Sklyanin, *Bäcklund transformations and Baxter's Q-operator*, *CRM Proceedings and Lecture Notes, Amer. Math. Soc., Providence, RI* **26** (2000) 227 [[nlin/0009009](#)].
- [69] A.V. Kotikov, A. Rej and S. Zieme, *Analytic three-loop solutions for $N=4$ SYM twist operators*, *Nucl. Phys. B* **813** (2009) 460 [[0810.0691](#)].
- [70] M. Beccaria, A.V. Belitsky, A.V. Kotikov and S. Zieme, *Analytic solution of the multiloop Baxter equation*, *Nucl. Phys. B* **827** (2010) 565 [[0908.0520](#)].
- [71] Y.L. Dokshitzer, G. Marchesini and G.P. Salam, *Revisiting parton evolution and the large- x limit*, *Phys. Lett. B* **634** (2006) 504 [[hep-ph/0511302](#)].
- [72] Y.L. Dokshitzer and G. Marchesini, *$N=4$ SUSY Yang-Mills: three loops made simple(r)*, *Phys. Lett. B* **646** (2007) 189 [[hep-th/0612248](#)].

- [73] V.N. Velizhanin, *Six-Loop Anomalous Dimension of Twist-Three Operators in $N=4$ SYM*, *JHEP* **11** (2010) 129 [[1003.4717](#)].
- [74] B.A. Kniehl and V.N. Velizhanin, *Nonplanar Cusp and Transcendental Anomalous Dimension at Four Loops in $\mathcal{N}=4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **126** (2021) 061603 [[2010.13772](#)].
- [75] B.A. Kniehl and V.N. Velizhanin, *Non-planar universal anomalous dimension of twist-two operators with general Lorentz spin at four loops in $N = 4$ SYM theory*, *Nucl. Phys. B* **968** (2021) 115429 [[2103.16420](#)].
- [76] S. Moch, B. Ruijl, T. Ueda, J.A.M. Vermaseren and A. Vogt, *Four-Loop Non-Singlet Splitting Functions in the Planar Limit and Beyond*, *JHEP* **10** (2017) 041 [[1707.08315](#)].