

CONFORMAL AND EXTRINSIC UPPER BOUNDS FOR THE HARMONIC MEAN OF NEUMANN AND STEKLOV EIGENVALUES

HANG CHEN

ABSTRACT. Let M be an m -dimensional compact Riemannian manifold with boundary. We obtain the upper bound of the harmonic mean of the first m nonzero Neumann eigenvalue and Steklov eigenvalues involving the conformal volume and relative conformal volume, respectively. We also give an optimal sharp extrinsic upper bound for closed submanifolds in the space forms. These extend the previous related results for the first nonzero eigenvalue.

1. INTRODUCTION

The study of eigenvalues on manifolds is an important topic in Riemannian geometry, and there have been lots of results, including various estimates bounded from below or above in different situations. In this paper, we study the upper bounds of the harmonic mean of the lower Neumann and Steklov eigenvalues.

1.1. Conformal upper bounds for Neumann eigenvalues. Let (M, g) be an m -dimensional compact Riemannian manifold. It is well known that the eigenvalues of Laplacian on M (with Neumann boundary condition if $\partial M \neq \emptyset$) are discrete and satisfies

$$0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \rightarrow +\infty.$$

In 1982, Li and Yau [30] introduced the concept of the conformal volume in order to study the Willmore conjecture and the first eigenvalue of compact surfaces.

Definition 1.1 (cf. [30]). Let M be an m -dimensional compact Riemannian manifold with boundary that admits a conformal map $\phi : M \rightarrow \mathbb{S}^n$. Define

$$V_c(M, n, \phi) := \sup_{\gamma \in \text{Conf}(\mathbb{S}^n)} V(\gamma(\phi(M))), \quad (1.1)$$

$$V_c(M, n) := \inf_{\phi} V_c(M, n, \phi), \quad (1.2)$$

$$V_c(M) := \lim_{n \rightarrow +\infty} V_c(M, n). \quad (1.3)$$

Here $\text{Conf}(\mathbb{S}^n)$ is the group of conformal diffeomorphisms of \mathbb{S}^n , and the infimum is over all non-degenerate conformal maps $\phi : M \rightarrow \mathbb{S}^n$.

We call $V_c(M, n, \phi)$, $V_c(M, n)$ and $V_c(M)$ the n -conformal volume of ϕ , the n -conformal volume of M , and the conformal volume M , respectively.

We remark that, the conformal map ϕ exists for n big enough because of the Nash embedding theorem (via the stereographic projection). And the limit in (1.3) exists since $V_c(M, n) \geq V_c(M, n+1)$ (cf. [30, Fact 4]).

2020 *Mathematics Subject Classification.* 58C40, 53C42, 35P15.

Key words and phrases. Neumann eigenvalues, Steklov eigenvalues, conformal volume, Reilly inequality.

We set the following condition for convenience.

(C1) (M, g) admits, up to a homothety, a minimal isometric immersion in \mathbb{S}^n given by a subspace of the first eigenfunctions.

Then the first eigenvalue λ_1 can be bounded by the conformal volume from above. Precisely, we have

Theorem 1.2 ([14, 30]). *Let (M, g) be an $m(\geq 2)$ -dimensional compact Riemannian manifold. Then the first nonzero eigenvalue λ_1 of the Laplacian (with Neumann boundary condition if $\partial M \neq \emptyset$) satisfies*

$$\lambda_1 \leq m \left(\frac{V_c(M, n)}{V(M, g)} \right)^{2/m}$$

for all n for which $V_c(M, n)$ is defined.

Equality implies (C1). If M is closed, (C1) is also sufficient for equality holding.

This estimate was firstly obtained by Li and Yau [30] for $m = 2$, and then El Soufi and Ilias [14] for general $n \geq 3$. Matei [32] generalized it to the p -Laplacian. For higher eigenvalues of a closed manifold, Kokarev [27] very recently proved that there is a constant $C(n, m)$ depending only on m and n such that

$$\lambda_k \leq C(n, m) \left(\frac{V_c(M, n)}{V(M, g)} \right)^{2/m} k^{2/m} \quad (1.4)$$

holds for any $k \geq 1$, which is compatible with the famous Weyl asymptotic formula:

$$\lambda_k V(M, g)^{2/m} \sim \frac{4\pi^2}{\omega_m^{2/m}} k^{2/m} \text{ as } k \rightarrow \infty, \quad (1.5)$$

where ω_m is the volume of the unit ball in \mathbb{R}^m . This can be viewed as an improvement of Korevaar's result [28], which says that

$$\lambda_k \leq C k^{2/m} \quad (1.6)$$

for some constant C depending on $[g]$ in a rather implicit way. In the same paper [27], the author also proved a version of the Reilly inequality for higher eigenvalues, that is,

$$\lambda_k \leq C(n, m) \left(\frac{1}{V(M)} \int_M (c + |\mathbf{H}|^2) \right) k \quad (1.7)$$

for a closed submanifold M^m of $\mathbb{R}^n(c)$.

In this paper, we consider the harmonic mean of the first m nonzero eigenvalues. We write the harmonic mean of k positive numbers a_1, \dots, a_k as

$$\mathfrak{H}(a_1, \dots, a_k) = \left(\frac{a_1^{-1} + \dots + a_k^{-1}}{k} \right)^{-1}. \quad (1.8)$$

If $a_1 \leq \dots \leq a_k$, then one can easily check that

$$a_1 = \mathfrak{H}(a_1) \leq \mathfrak{H}(a_1, a_2) \leq \dots \leq \mathfrak{H}(a_1, \dots, a_{k-1}) \leq \mathfrak{H}(a_1, \dots, a_k). \quad (1.9)$$

The first main result in this paper is the following estimate involving the conformal volume for $\mathfrak{H}(\lambda_1, \dots, \lambda_m)$.

Theorem 1.3. *Let M be an $m(\geq 2)$ -dimensional compact Riemannian manifold. Then the first m nonzero eigenvalues of the Laplacian (with Neumann boundary condition if $\partial M \neq \emptyset$) satisfy*

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \leq m \left(\frac{V_c(M, n)}{V(M, g)} \right)^{2/m} \quad (1.10)$$

for all n for which $V_c(M, n)$ is defined.

Equality implies (C1). If M is closed, (C1) is also sufficient for equality holding.

Remark 1.4. For some particular manifolds, such as compact symmetric spaces of rank 1 and the minimal Clifford torus with their canonical metric can, we have $V_c(M, n) = V(M, \frac{\lambda_1}{m} \text{can})$ for $n+1$ greater or equal to the multiplicity of λ_1 (cf. [14, pp. 266–267]). In these cases the upper bound is explicit.

Clearly, Theorem 1.3 is stronger than Theorem 1.2 due to (1.9). For $m = 2$, the upper bound can be controlled by the topology of the surface.

Corollary 1.5. Let (M, g) be a closed surface¹. Then

$$\mathfrak{H}(\lambda_1, \lambda_2) \leq \frac{8\pi}{V(M, g)} \left[\frac{\text{genus}(M) + 3}{2} \right], \quad (1.11)$$

where $[\cdot]$ denotes the integer part. In particular, for a 2-sphere we have

$$\mathfrak{H}(\lambda_1, \lambda_2) \leq \frac{8\pi}{V(\mathbb{S}^2, g)}, \quad (1.12)$$

This corollary is directly obtained from Theorem 1.3 since Li and Yau [30] pointed out that

$$V_c(M, g) \leq 4\pi(\text{genus}(M) + 1), \quad (1.13)$$

and then El Soufi and Ilias [13] remark that the factor $\text{genus}(M) + 1$ in (1.13) can be improved to $[\frac{\text{genus}(M)+3}{2}]$,

Let's review some previous related results for $m = 2$. Obviously, Corollary 1.5 implies

$$\lambda_1 \leq \frac{8\pi}{V(M, g)}(\text{genus}(M) + 1), \quad (1.14)$$

which was originally proved by Hersch [24] for $M = S^2$ (i.e., $\text{genus}(M) = 0$) and Yang-Yau [47] for $\text{genus}(M) \geq 1$. Actually, Yang-Yau [47] proved a stronger result

$$\mathfrak{H}(\lambda_1, \lambda_2, \lambda_3) \leq \frac{8\pi}{V(M, g)}(\text{genus}(M) + 1). \quad (1.15)$$

By (1.13), Li and Yau [30] gave a simpler proof of (1.14); they also proved that ([30, Corollary 2])

$$\mathfrak{H}(\lambda_1, \dots, \lambda_k) \leq n \frac{V_c(M, n-1)}{V(M)} \text{ for any } n \geq k, \quad (1.16)$$

but the constant is not sharp.

Another remarkable estimate for the higher eigenvalues of a 2-sphere (\mathbb{S}^2, g) is

$$\lambda_k(\mathbb{S}^2, g) \leq \frac{8\pi k}{V(\mathbb{S}^2, g)}, \text{ for any } k \geq 1. \quad (1.17)$$

The metric g is smooth outside a finite number of conical singularities. For $k = 1$, this is just Hersch's result [24] and the equality in (1.17) is attained if and only if g is a round metric. For $k \geq 2$, the inequality in (1.17) is strict, and the equality can be attained in the limit by a sequence of metrics degenerating to a union of k touching identical round spheres. This was proved by Nadirashvili [34] (and Petrides [37] with a different argument)

¹In this article, we always consider orientable surfaces except special declaration.

for $k = 2$, Nadirashvili and Sire [35] for $k = 3$, and Karpukhin et al. [25] for any $k \geq 2$ very recently. From (1.17) we have

$$\mathfrak{H}(\lambda_1, \lambda_2) \leq \frac{4}{3} \frac{8\pi}{V(\mathbb{S}^2, g)}, \quad (1.18)$$

which is weaker than (1.12). Our result (1.12) means that if λ_2 is close to $\frac{8\pi \times 2}{V(\mathbb{S}^2, g)}$, then λ_1 is not close to $\frac{8\pi}{V(\mathbb{S}^2, g)}$.

1.2. Conformal upper bounds for Steklov eigenvalues. Beside Neumann eigenvalues, Steklov eigenvalues are intensively studied for a compact Riemannian manifold (M, g) with nonempty boundary ∂M . A real number σ is called a Steklov eigenvalue if there exists a non-zero function u (which is called the Steklov eigenfunction corresponding to σ) on M satisfying

$$\begin{cases} \Delta u = 0, & \text{on } M; \\ \frac{\partial u}{\partial \nu} = \sigma u, & \text{on } \partial M, \end{cases} \quad (1.19)$$

where ν is the outward normal on ∂M .

The Steklov problem has a physical background, and it can be traced back to the turn of the 20th century, when Steklov studied liquid sloshing [40, 41]. The readers can refer [29] for the history and a very recent survey [10] for researches in this area. There are also the very recent developments and various results (e.g., [11, 12, 18, 19, 44, 46] and the references therein).

The Steklov eigenvalues can be interpreted as the eigenvalues of the Dirichlet-to-Neumann operator L , which is defined as follows:

$$\begin{aligned} L : C^\infty(\partial M) &\rightarrow C^\infty(\partial M) \\ Lu &= \frac{\partial \hat{u}}{\partial \nu}, \end{aligned} \quad (1.20)$$

where u is a function on ∂M and \hat{u} is the harmonic extension of u , i.e.,

$$\begin{cases} \Delta \hat{u} = 0, & \text{on } M; \\ \hat{u} = u, & \text{on } \partial M. \end{cases} \quad (1.21)$$

L is a nonnegative self-adjoint operator with discrete spectrum

$$0 = \sigma_0 < \sigma_1 \leq \sigma_2 \leq \cdots \rightarrow +\infty.$$

Fraser and Schoen [17] gave an upper bound involved the *relative conformal volume* for the first nonzero Steklov eigenvalue, which is analogous to Theorem 1.2 for the first Neumann eigenvalue of the Laplacian.

Theorem 1.6 ([17, Theorem 6.2]). *Let (M, g) be a compact m -dimensional Riemannian manifold with nonempty boundary. Then the first nonzero Steklov eigenvalue σ_1 satisfies*

$$\sigma_1 V(\partial M) V(M)^{\frac{2-m}{m}} \leq m V_{rc}(M, n)^{2/m} \quad (1.22)$$

for all n for which $V_{rc}(M, n)$ is defined.

Equality implies that there exists a conformal harmonic map $\phi : M \rightarrow B^n$ which (after rescaling the metric g) is an isometry on ∂M , with $\phi(\partial M) \subset \partial B^n$ and such that $\phi(M)$ meets ∂B^n orthogonally along $\phi(\partial M)$. For $m > 2$, this map is an isometric minimal immersion of M to its image. Moreover, the immersion is given by a subspace of the first eigenspace.

We explain $V_{rc}(M, n)$ in the above theorem.

Definition 1.7 (cf. [17]). Let M be an m -dimensional compact Riemannian manifold with boundary that admits a conformal map $\phi : M \rightarrow B^n$ with $\phi(\partial M) \subset \partial B^n$. Define

$$V_{rc}(M, n, \phi) := \sup_{\gamma \in \text{Conf}(B^n)} V(\gamma(\phi(M))), \quad (1.23)$$

$$V_{rc}(M, n) := \inf_{\phi} V_{rc}(M, n, \phi), \quad (1.24)$$

$$V_{rc}(M) := \lim_{n \rightarrow +\infty} V_{rc}(M, n). \quad (1.25)$$

Here $\text{Conf}(B^n)$ is the group of conformal diffeomorphisms of B^n , and the infimum is over all non-degenerate conformal maps $\phi : M \rightarrow B^n$ with $\phi(\partial M) \subset \partial B^n$.

We call $V_{rc}(M, n, \phi)$, $V_{rc}(M, n)$ and $V_{rc}(M)$ the relative n -conformal volume of ϕ , the relative n -conformal volume of M , and the relative conformal volume of M , respectively. The existence of the limit in (1.25) is from the monotonicity of $V_{rc}(M, n)$ (cf. [17, Lemma 5.7]).

We extend Theorem 1.6 to the harmonic mean of the first m nonzero eigenvalues.

Theorem 1.8. *Let (M, g) be a compact m -dimensional Riemannian manifold with nonempty boundary. Then the first m nonzero Steklov eigenvalues satisfy*

$$\mathfrak{H}(\sigma_1, \dots, \sigma_m) V(\partial M) V(M)^{\frac{2-m}{m}} \leq m V_{rc}(M, n)^{2/m} \quad (1.26)$$

for all n for which $V_{rc}(M, n)$ is defined.

Equality implies that there exists a conformal harmonic map $\phi : M \rightarrow B^n$ with $\phi(\partial M) \subset \partial B^n$. For $m > 2$, this map (after rescaling the metric g) is an isometric minimal immersion of M to its image. Moreover, each coordinate functions of this immersion belongs to some eigenspace E_{σ_i} ($1 \leq i \leq m$) (counted with multiplicity).

For compact surfaces, we have

Theorem 1.9. *Let (M, g) be a compact surface with k boundary components. Then*

$$\mathfrak{H}(\sigma_1, \sigma_2) V(\partial M) \leq 2(\text{genus}(M) + k)\pi. \quad (1.27)$$

In particular, for a compact simply-connected surface with boundary, we have

$$\mathfrak{H}(\sigma_1, \sigma_2) V(\partial M) \leq 2\pi, \quad (1.28)$$

Moreover, with equality if and only if there is a conformal map from M to the unit disk which is an isometry on the boundary.

This theorem extends the estimates of Fraser-Schoen ([17, Theorem 2.3]) which says

$$\sigma_1 V(\partial M) \leq 2(\text{genus}(M) + k)\pi \quad (1.29)$$

under the same assumptions. When $\text{genus}(M) = 0$, (1.29) was obtained by Weinstock [43] for $k = 1$ and it is sharp; however, it is not sharp for $k \geq 2$ (see [17, Theorem 2.3]). These estimates for Steklov eigenvalues are analogous to the estimates (1.11) and (1.14) for eigenvalues of the Laplacian. We also have

Theorem 1.10. *Let (M, g) be a compact surface with boundary. Then*

$$\mathfrak{H}(\sigma_1, \sigma_2) V(\partial M) < 8\pi(\text{genus}(M) + 1). \quad (1.30)$$

In particular, for a compact surface with boundary of genus 0, we have

$$\mathfrak{H}(\sigma_1, \sigma_2) V(\partial M) < 8\pi. \quad (1.31)$$

This extends some corresponding estimates for σ_1 (cf. [26, Theorem A₁] and [10, Theorem 3.6]).

1.3. Reilly inequalities in space forms. The conformal upper bounds are proved by finding suitable test functions via conformal maps. This technique also allows us to obtain some extrinsic upper bounds for closed submanifolds in space forms.

Let $(\mathbb{R}^n(c), h_c)$ be the n -dimensional simply-connected space form of constant curvature c equality with the canonical metric h_c , namely, $\mathbb{R}^n(c)$ represents the Euclidean space \mathbb{R}^n , the hyperbolic space $\mathbb{H}^n(-1)$ and the unit sphere \mathbb{S}^n for $c = 0, -1$ and 1 , respectively. When M is a closed submanifold of $\mathbb{R}^n(c)$, the first nonzero eigenvalue can be bounded by the mean curvature \mathbf{H} from above.

Theorem 1.11 ([15, 38]). *Let M be an $m(\geq 2)$ -dimensional closed submanifold in $\mathbb{R}^n(c)$. Then the first nonzero eigenvalue of Laplacian satisfies*

$$\lambda_1 \leq \frac{m}{V(M)} \int_M (c + |\mathbf{H}|^2). \quad (1.32)$$

If we exclude the case that M is minimal in \mathbb{S}^n for $c = 1$, then the equality holds if and only if M is minimally immersed in a geodesic sphere Σ_c of $\mathbb{R}^n(c)$, where the geodesic radius r_c of Σ_c is given by

$$r_0 = \left(\frac{m}{\lambda_1} \right)^{1/2}, \quad r_1 = \arcsin r_0, \quad r_{-1} = \operatorname{arsinh} r_0.$$

Remark 1.12. *We use the following convention:*

When M is a closed hypersurface of $\mathbb{R}^n(c)$ (i.e., $n - m = 1$), “ M lies in a geodesic sphere Σ_c of $\mathbb{R}^n(c)$ ” means “ $M = \Sigma_c$ ”.

Theorem 1.11 is usually referred “Reilly inequality” as it was first proved by Reilly [38] for $c = 0$ by using the coordinate functions as the test functions after translating the center of mass of M to the origin. By embedding the sphere $\mathbb{S} \rightarrow \mathbb{R}^{n+1}$, one can reduce the case $c = 1$ to the case $c = 0$. But this method fails for $c = -1$. The case $c = -1$ was proved by El Soufi and Ilias [15] via the technique of conformal maps in [30].

There are lots of generalizations for Reilly inequality. For example, one can consider the case that the ambient space is a manifold whose sectional curvature bounded above by a constant, even can have a weighted measure; one can consider other operators, such as so-called L_T operator and p -Laplacian, instead of the Laplacian. The readers can refer [6, 23, 31, 36, 39] and the references therein. In this paper, we prove

Theorem 1.13. *Let M be an $m(\geq 2)$ -dimensional closed submanifold in $\mathbb{R}^n(c)$. Then the first m nonzero eigenvalues of the Laplacian satisfy:*

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \leq \frac{m}{V(M)} \int_M (c + |\mathbf{H}|^2). \quad (1.33)$$

Moreover, equality in (1.33) holds if and only if either

(1) *M is minimal in \mathbb{S}^n and the immersion is given by a subspace of the first eigenfunctions (this occurs only when $c = 1$), or*

(2) *M is minimal in a geodesic sphere Σ_c of $\mathbb{R}^n(c)$, where the geodesic radius r_c of Σ_c is given by*

$$r_0 = \left(\frac{m}{\lambda_1} \right)^{1/2}, \quad r_1 = \arcsin r_0, \quad r_{-1} = \operatorname{arsinh} r_0.$$

1.4. Extrinsic estimates for submanifolds of \mathbb{R}^n . When M is a submanifold of \mathbb{R}^n , we can directly use the coordinate functions as the test functions without conformal maps. This is just the original idea in [38], which allows us to get more general extrinsic upper bounds, including both closed eigenvalues (i.e., eigenvalues for closed manifolds) and Steklov eigenvalues. In fact, the original Reilly's inequalities in [38] involve not only the mean curvature, but also the higher order mean curvatures. By this approach, we can prove the following results. First, considering the closed eigenvalues, we have

Theorem 1.14. *Let M be an $m(\geq 2)$ -dimensional closed submanifold of \mathbb{R}^n . Let T be a symmetric, divergence-free $(1,1)$ -tensors on M . Then the first m nonzero eigenvalues satisfy*

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \left(\int_M \operatorname{tr} T \right)^2 \leq mV(M) \int_M |H_T|^2. \quad (1.34)$$

Assume that $H_T \not\equiv 0$, then equality holds if and only if the following conditions hold:

- (1) $\operatorname{tr} T$ is constant;
- (2) M is minimal and T -minimal in a hypersphere of \mathbb{R}^n with the radius $\sqrt{m/\lambda_1}$.

Theorem 1.15. *Let M be an $m(\geq 2)$ -dimensional closed submanifold of \mathbb{S}^n . Let T be a symmetric, divergence-free $(1,1)$ -tensors on M . Then the first m nonzero eigenvalues satisfy*

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \left(\int_M \operatorname{tr} T \right)^2 \leq mV(M) \int_M (|H_T|^2 + (\operatorname{tr} T)^2). \quad (1.35)$$

Assume that $H_T \not\equiv 0$, then equality holds if and only if the following conditions hold:

- (1) $\operatorname{tr} T$ are constant;
- (2) M is minimal and T -minimal in \mathbb{S}^n and the immersion is given by a subspace of the first eigenfunctions; or M is minimal and T -minimal in a geodesic sphere of \mathbb{S}^n with the geodesic radius $\arcsin(\sqrt{m/\lambda_1})$;

We will introduce the definitions of H_T and T -minimality in Sect. 2.3. The above theorems imply [21, Proposition 2.1]. We can take $T = T_r$, the Newton transformation of the second fundamental form, then H_T and $\operatorname{tr} T$ are corresponding to the $(r+1)$ -th and r -th mean curvature, respectively.

Corollary 1.16. *Let M be an $m(\geq 2)$ -dimensional closed submanifold of $\mathbb{R}^n(c)$, $c = 0, 1$. Then the first m nonzero eigenvalues satisfy:*

- (1) if $n > m + 1$ and $r \in \{0, \dots, m-1\}$ is an even integer, then

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \left(\int_M H_r \right)^2 \leq mV(M) \int_M (|\mathbf{H}_{r+1}|^2 + cH_r^2),$$

- (2) if $n = m + 1$ and $r \in \{0, \dots, m-1\}$ is any integer, then

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \left(\int_M H_r \right)^2 \leq mV(M) \int_M (H_{r+1}^2 + cH_r^2),$$

where \mathbf{H}_{r+1} is the $(r+1)$ -th mean curvature vector of M .

Corollary 1.16 implies [38, Theorem A] and [21, Theorem 2.1]. In particular, we again obtain Theorem 1.13 for $c = 0, 1$ when taking $r = 0$.

For the Steklov eigenvalues, we have

Theorem 1.17. *Let M be an $m(\geq 2)$ -dimensional compact submanifold with non-empty boundary ∂M of \mathbb{R}^n . Let T be a symmetric, divergence-free $(1,1)$ -tensors on ∂M . Then the first m nonzero Steklov eigenvalues satisfy*

$$\mathfrak{H}(\sigma_1, \dots, \sigma_m) \left(\int_{\partial M} \operatorname{tr} T \right)^2 \leq mV(M) \int_{\partial M} |H_T|^2. \quad (1.36)$$

In particular, when $T = I$, we obtain

$$\mathfrak{H}(\sigma_1, \dots, \sigma_m) \leq \frac{mV(M)}{(V(\partial M))^2} \int_{\partial M} |\mathbf{H}^{\partial M}|^2,$$

where $\mathbf{H}^{\partial M}$ denotes the mean curvature of ∂M in \mathbb{R}^n .

This paper is organized as follows. In Sect. 2 we recall some formulae about conformal maps and submanifolds. Then we give the proofs of Theorems 1.3 and 1.8 in Sect. 3 and Sect. 4, respectively. we give In Sect. 5 we prove the Reilly-type inequality Theorem 1.13; a general version Theorem 5.1 is also obtained. In the last section, we give the proofs of results mentioned in Sect. 1.4. The choice of test functions is analogous in the proofs, but discussions on equalities are different. Unlike the estimates for the first eigenvalues, we need more careful analysis when studying the equality cases.

2. PRELIMINARIES AND NOTATIONS

In this section, we give the relations between some geometric quantities of M^m as a submanifold of (N^n, g_N) and their corresponding quantities of M as a submanifold in (N^n, \tilde{g}_N) , where \tilde{g}_N is conformal to g_N . Although the relations are well-known in the literature (cf. [4, 5]), we give a brief proof of these relations for the reader's convenience. We also recall some basic formulas for submanifolds in space forms.

We use the following convention on the ranges of indices except special declaration:

$$1 \leq i, j, k, \dots \leq m; \quad m+1 \leq \alpha, \beta, \gamma, \dots \leq n; \quad 1 \leq A, B, C, \dots \leq n.$$

2.1. Conformal relations: the ambient manifold. Let (N, g_N) be an n -dimensional n -dimensional Riemannian manifold.

Let $\{e_A\}_{A=1}^n$ be an orthonormal frame of (N, g_N) , where $\{e_i\}_{i=1}^m$ are tangent to M^2 and $\{e_\alpha\}_{\alpha=m+1}^n$ are normal to M . Let $\{\omega_A\}_{A=1}^n$ be the dual coframe of $\{e_A\}_{A=1}^n$. Then the structure equations of (N, g_N) are (see [9]):

$$\begin{cases} d\omega_A = \sum_B \omega_{AB} \wedge \omega_B, \\ \omega_{AB} + \omega_{BA} = 0, \end{cases} \quad (2.1)$$

where $\{\omega_{AB}\}$ are the connection forms of (N, g_N) .

Let \tilde{g}_N be a metric on N conforming to g_N . We assume $\tilde{g}_N = e^{2\rho} g_N$ for some $\rho \in C^\infty(N)$. Then $\{\tilde{e}_A = e^{-\rho} e_A\}$ is an orthonormal frame of (N, \tilde{g}_N) , and $\{\tilde{\omega}_A = e^\rho \omega_A\}$ is the dual coframe of $\{\tilde{e}_A\}$. The structure equations of (N, \tilde{g}_N) are given by

$$\begin{cases} d\tilde{\omega}_A = \sum_B \tilde{\omega}_{AB} \wedge \tilde{\omega}_B, \\ \tilde{\omega}_{AB} + \tilde{\omega}_{BA} = 0, \end{cases} \quad (2.2)$$

²Here we identify e_i with $\phi_*(e_i)$

where $\{\tilde{\omega}_{AB}\}$ are the connection forms of (N, \tilde{g}_N) .

From (2.1) and (2.2), we can derive that

$$\tilde{\omega}_{AB} = \omega_{AB} + \rho_A \omega_B - \rho_B \omega_A, \quad (2.3)$$

where ρ_A means the covariant derivative of ρ with respect to e_A .

Let $\bar{\nabla}$ (or $\tilde{\nabla}$ resp.) denote Levi-Civita connections on N with respect to metric g_N (or \tilde{g}_N resp.). In general, for any $F \in C^\infty(N)$, we compute the gradient

$$\begin{aligned} dF &= \sum_{A=1}^n F_A \omega_A = \sum_{A=1}^n \bar{\nabla}_A F \omega_A \\ &= \sum_{A=1}^n \tilde{F}_A \tilde{\omega}_A = \sum_{A=1}^n \tilde{\nabla}_A F \tilde{\omega}_A, \end{aligned}$$

which gives the relation

$$\tilde{F}_A = \tilde{\nabla}_A F = e^{-\rho} \bar{\nabla}_A F = e^{-\rho} F_A, \quad \text{for } A = 1, \dots, n. \quad (2.4)$$

2.2. Conformal relations: the submanifold. Let M be an $m(\geq 2)$ -dimensional submanifold of N immersed by ϕ .

When consider the metric g_N , M has an induced metric $g_M = \phi^* g_N$. Denote $\phi^* \omega_A = \theta_A$, $\phi^* \omega_{AB} = \theta_{AB}$, then restricted to (M, g_M) , we have (see [9])

$$\theta_\alpha = 0, \quad \theta_{i\alpha} = \sum_j h_{ij}^\alpha \theta_j. \quad (2.5)$$

and

$$\begin{cases} d\theta_i = \sum_j \theta_{ij} \wedge \theta_j, & \theta_{ij} + \theta_{ji} = 0, \\ d\theta_{ij} - \sum_k \theta_{ik} \wedge \theta_{kj} = -\frac{1}{2} \sum_{k,l} R_{ijkl} \theta_k \wedge \theta_l, \end{cases} \quad (2.6)$$

where R_{ijkl} are components of the curvature tensor of (M, g_M) and h_{ij}^α are components of the second fundamental form of (M, g_M) in (N, g_N) .

We denote the mean curvature vector of M (precisely, of ϕ) by

$$\mathbf{H}_\phi = \frac{1}{n} \sum_\alpha \left(\sum_i h_{ii}^\alpha \right) e_\alpha = \sum_\alpha H^\alpha e_\alpha. \quad (2.7)$$

We sometimes omit the subscript ϕ and just write \mathbf{H} when the immersion is clear.

Similarly, we consider the induced metric $\tilde{g}_M = \phi^* \tilde{g}_N$, and denote $\phi^* \tilde{\omega}_A = \tilde{\theta}_A$, $\phi^* \tilde{\omega}_{AB} = \tilde{\theta}_{AB}$. Then restricted to (M, \tilde{g}_M) , we have

$$\tilde{\theta}_\alpha = 0, \quad \tilde{\theta}_{i\alpha} = \sum_j \tilde{h}_{ij}^\alpha \tilde{\theta}_j, \quad (2.8)$$

and

$$\begin{cases} d\tilde{\theta}_i = \sum_j \tilde{\theta}_{ij} \wedge \tilde{\theta}_j, & \tilde{\theta}_{ij} + \tilde{\theta}_{ji} = 0, \\ d\tilde{\theta}_{ij} - \sum_k \tilde{\theta}_{ik} \wedge \tilde{\theta}_{kj} = -\frac{1}{2} \sum_{k,l} \tilde{R}_{ijkl} \tilde{\theta}_k \wedge \tilde{\theta}_l, \end{cases} \quad (2.9)$$

where \tilde{R}_{ijkl} are components of the curvature tensor of (M, \tilde{g}_M) and \tilde{h}_{ij}^α are components of the second fundamental form of (M, \tilde{g}_M) in (N, \tilde{g}_N) .

By pulling back (2.3) to M by ϕ and using (2.5) and (2.8), we obtain the following relation.

$$\tilde{h}_{ij}^\alpha = e^{-\rho}(h_{ij}^\alpha - \rho_\alpha \delta_{ij}), \quad \tilde{H}^\alpha = e^{-\rho}(H^\alpha - \rho_\alpha). \quad (2.10)$$

Combining (2.3), (2.4), (2.6), and (2.9), we can obtain the following relation:

$$\begin{aligned} e^{2\rho} \tilde{R}_{ijkl} = & R_{ijkl} - (\rho_{ik} \delta_{jl} + \rho_{jl} \delta_{ik} - \rho_{il} \delta_{jk} - \rho_{jk} \delta_{il}) \\ & + (\rho_i \rho_k \delta_{jl} + \rho_j \rho_l \delta_{ik} - \rho_j \rho_k \delta_{il} - \rho_i \rho_l \delta_{jk}) \\ & - \left(\sum_m \rho_m^2 \right) (\delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}), \end{aligned} \quad (2.11)$$

$$e^{2\rho} \tilde{R}_{ij} = R_{ij} + (n-2)(\rho_i \rho_j - \rho_{ij} - |\nabla \rho|^2 \delta_{ij}) - (\Delta \rho) \delta_{ij}, \quad (2.12)$$

where R_{ij} (or \tilde{R}_{ij} resp.) are components of Ricci curvature with respect to g_M (or \tilde{g}_M resp.).

The following proposition gives the relation between the conformal fact and the mean curvature.

Proposition 2.1 (cf. Proposition 3.3 and Remark 3.4 in [8]). *Let (N, g_N) be an n -dimensional Riemannian manifold (possibly not complete) which admits a conformal immersion γ in the sphere (\mathbb{S}^n, h_1) . Assume $\gamma^* h_1 = e^{2\rho} g_N$.*

Let M be an $m(\geq 2)$ -dimensional submanifold immersed in (N, g_N) by ϕ . Then we have

$$e^{2\rho} = \left(|\mathbf{H}_\phi|^2 + \bar{\mathcal{R}}_\phi \right) - \frac{2}{m} \Delta \rho - \frac{m-2}{m} |\nabla \rho|^2 - |(\bar{\nabla} \rho)^\perp - \mathbf{H}_\phi|^2, \quad (2.13)$$

where

$$\bar{\mathcal{R}}_\phi = \frac{1}{m(m-1)} \sum_{i,j} K_N(e_i, e_j) \quad (2.14)$$

for a local orthogonal tangent frame $\{e_i\}$ on M , and K_N is the sectional curvature of (N, g_N) .

In particular, when $(N, g_N) = (\mathbb{R}^n(c), h_c)$, we have

$$e^{2\rho} = (|\mathbf{H}|^2 + c) - \frac{2}{m} \Delta \rho - \frac{m-2}{m} |\nabla \rho|^2 - |(\bar{\nabla} \rho)^\perp - \mathbf{H}|^2. \quad (2.15)$$

Proof. We denote the immersion of M to (N, g_N) by ϕ . Consider two metric $g_M = \phi^* g_N$ and $\tilde{g}_M = \phi^*(\gamma^* h_1) = (\gamma \circ \phi)^* h_1$ on M . The Gauss equations for the immersions ϕ and $\gamma \circ \phi$ imply:

$$R_{ij} = \sum_k \bar{R}_{ikjk} + \sum_\alpha m H^\alpha h_{ij}^\alpha - \sum_{k,\alpha} h_{ik}^\alpha h_{kj}^\alpha, \quad (2.16)$$

$$\tilde{R}_{ij} = (m-1)\delta_{ij} + \sum_\alpha m \tilde{H}^\alpha \tilde{h}_{ij}^\alpha - \sum_{k,\alpha} \tilde{h}_{ik}^\alpha \tilde{h}_{kj}^\alpha. \quad (2.17)$$

From (2.10), (2.12), (2.16) and (2.17), we obtain

$$\begin{aligned} (m-2)(\rho_i \rho_j - \rho_{ij} - |\nabla \rho|^2 \delta_{ij}) - (\Delta \rho) \delta_{ij} &= e^{2\rho} \tilde{R}_{ij} - R_{ij} \\ &= (m-1)(e^{2\rho} + \sum_\alpha \rho_\alpha^2) \delta_{ij} - \sum_k \bar{R}_{ikjk} - (m-2) \sum_\alpha \rho_\alpha h_{ij}^\alpha - \sum_\alpha m H^\alpha \rho_\alpha \delta_{ij}. \end{aligned} \quad (2.18)$$

Contracting (2.18) with δ_{ij} , we obtain

$$-2\Delta \rho = (m-2)|\nabla \rho|^2 + m(e^{2\rho} + \sum_\alpha \rho_\alpha^2) - 2 \sum_\alpha m H^\alpha \rho_\alpha - \frac{1}{m-1} \sum_{i \neq k} \bar{R}_{ikik}, \quad (2.19)$$

which is equivalent to (2.13). \square

When $(N, g_N) = (\mathbb{R}^n(c), h_c)$, we usually denote the immersion of M to $\mathbb{R}^n(c)$ by x instead of ϕ , and x can be viewed as the *position vector*. Then we have (cf. [3, 9, 42]):

$$dx = \sum_i \theta_i e_i, \quad de_i = \sum_j \theta_{ij} e_j + \sum_j h_{ij}^\alpha \theta_j e_\alpha - c\theta_i x, \quad de_\alpha = -\sum_{i,j} h_{ij}^\alpha \theta_j e_i + \sum_\beta \theta_{\alpha\beta} e_\beta,$$

which implies

$$x_i = e_i, \quad x_{ij} = \sum_\alpha h_{ij}^\alpha e_\alpha - c\delta_{ij}x, \quad \Delta x = m\mathbf{H} - mcx. \quad (2.20)$$

2.3. The generalization of the mean curvature. Given a symmetric $(0, 2)$ -tensor $T = (T_{ij})$ on a (closed) manifold M , we can define a normal vector field H_T associated with T by

$$H_T = \sum_{i=1}^m A(Te_i, e_i) = \sum_{\substack{m+1 \leq \alpha \leq n \\ 1 \leq i, j \leq m}} h_{ij}^\alpha T_{ij} e_\alpha.$$

We call M is T -minimal in N if $H_T \equiv 0$.

If we additionally assume that T is divergence-free, then we can define a self-adjoint operator L_T associated to T by

$$L_T u = \operatorname{div}(T\nabla u), \quad u \in C^\infty(M).$$

where T is regarded as a $(1, 1)$ -tensor.

Now H_T is a generalization of the mean curvature and the higher order mean curvature, while L_T is a generalization of the Laplacian. Indeed, $H_T = m\mathbf{H}$, $L_T = \Delta$ when $T = I$ the identity.

Both H_T and L_T have been involved a lot of results (e.g., [7, 22]).

We have (cf. [21, Lemma 2.3])

$$L_T x = H_T, \quad \frac{1}{2} L_T |x|^2 = \langle x, H_T \rangle + \operatorname{tr} T. \quad (2.21)$$

For a closed submanifold M of \mathbb{R}^n , we obtain the following generalized Hsiung–Minkowski formulas

$$0 = \frac{1}{2} \int_M L_T |x|^2 = \int_M \langle x, H_T \rangle + \int_M \operatorname{tr} T \quad (2.22)$$

by integrating the second equation of (2.21).

3. PROOFS FOR THE NEUMANN EIGENVALUES

In this section, we prove Theorem 1.3. We prove the inequality (1.10) in Sect. 3.1 and then analyze the equality carefully in Sect. 3.2. A corollary is given to end this section.

3.1. The inequality in (1.10). To construct suitable test functions, we recall the following lemma.

Lemma 3.1 (see [16, 30]). *Let $\phi : (M^m, g) \rightarrow (\mathbb{S}^n, h_1)$ be a conformal map. Then there exists $\gamma \in \operatorname{Conf}(\mathbb{S}^n)$ such that the immersion $\Phi = \gamma \circ \phi = (\Phi^1, \dots, \Phi^{n+1})$ satisfies that*

$$\int_M \Phi^A = 0, \quad A = 1, \dots, n+1. \quad (3.1)$$

It is well known that the variational characterization of $\lambda_i (i \geq 1)$

$$\lambda_i = \inf_{u \in H^1(M) \setminus \{0\}} \left\{ \frac{\int_M |\nabla u|^2}{\int_M u^2} \mid \int_M uu_j = 0, j = 0, \dots, i-1 \right\}, \quad (3.2)$$

where $\{u_i\}_{i \geq 0}$ is an orthonormal set of eigenfunctions satisfying $\Delta u_i = -\lambda_i u_i$.

Now we can start the proof. (3.1) means that Φ^A is L^2 -orthogonal to the first eigenfunction u_0 (a constant). By using the QR-decomposition via the Gram–Schmidt process, we can further assume that $\{\Phi^1, \dots, \Phi^{n+1}\}$ satisfy

$$\int_M \Phi^A u_B = 0, \text{ for } 1 \leq B < A \leq n+1. \quad (3.3)$$

Indeed, we denote

$$d_{AB} = \int_M \Phi^A u_B, \text{ for } 1 \leq A, B \leq n+1, \quad (3.4)$$

then the matrix $D = (d_{AB}) = QR$, where $Q = (q_{AB}) \in O(n+1)$ and R is an upper triangular matrix. This is equivalent to $R = Q^T D$. Hence, if we replace the old orthonormal basis (E_1, \dots, E_n) of \mathbb{R}^{n+1} by the new one $(E'_1, \dots, E'_{n+1}) = (E_1, \dots, E_n)Q^T$, then these new coordinate functions, still denoted by Φ^A , satisfy (3.3).

Hence, by the variational characterization (3.2), for each $A = 1, \dots, n+1$, we have

$$\lambda_A \int_M (\Phi^A)^2 \leq \int_M |\nabla \Phi^A|^2. \quad (3.5)$$

On the other hand, if we denote $\tilde{g} = \Phi^* h_1 = fg$, then

$$0 \leq |\tilde{\nabla} \Phi^A|^2 \leq 1 - (\Phi^A)^2, \quad \sum_{A=1}^{n+1} |\tilde{\nabla} \Phi^A|^2 = m, \quad (3.6)$$

and

$$mf = \sum_{A=1}^{n+1} |\nabla \Phi^A|^2. \quad (3.7)$$

where $\tilde{\nabla}$ is the gradient operator on (M, \tilde{g}) . Noting that $\sum_{A=1}^{n+1} \Phi_i^2 = 1$, we obtain

$$V(M, g) = \int_M dv_g \leq \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \sum_{A=m+1}^{n+1} \frac{1}{\lambda_A} \int_M f |\tilde{\nabla} \Phi^A|^2 \quad (3.8)$$

$$\leq \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \frac{1}{\lambda_m} \int_M f \sum_{A=m+1}^{n+1} |\tilde{\nabla} \Phi^A|^2 \quad (3.9)$$

$$= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \frac{1}{\lambda_m} \int_M f (m - \sum_{A=1}^m |\tilde{\nabla} \Phi^A|^2)$$

$$= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \frac{1}{\lambda_m} \int_M f \sum_{A=1}^m (1 - |\tilde{\nabla} \Phi^A|^2)$$

$$\leq \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f (1 - |\tilde{\nabla} \Phi^A|^2) \quad (3.10)$$

$$= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f. \quad (3.11)$$

By using the Hölder inequalities, we have

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \leq \frac{m}{V(M, g)} \int_M f dv_g \quad (3.12)$$

$$\leq \frac{m}{V(M, g)} \left(\int_M f^{m/2} dv_g \right)^{2/m} (V(M))^{m-2} \quad (3.13)$$

$$= m \left(\frac{V(M, \tilde{g})}{V(M, g)} \right)^{2/m} \quad (3.14)$$

$$\leq m \left(\frac{\sup_{\gamma \in \text{Conf}(\mathbb{S}^n)} V(M, (\gamma \circ \phi)^* h_1)}{V(M, g)} \right)^{2/m}. \quad (3.15)$$

Taking the infimum over all $\phi \in I_n(M, [g])$, we obtain the desired inequality.

3.2. Equality in (1.10). Now we discuss the equality. Choose a sequence of conformal maps $\phi_j : M \rightarrow \mathbb{S}^n$ such that

$$\lim_{j \rightarrow \infty} V_c(M, n, \phi_j) = V_c(M, n), \quad (3.16)$$

and by composing with a conformal transformation of the sphere we may assume

$$\int_M \phi_j^A = 0, \quad 1 \leq A \leq n+1, \quad (3.17)$$

$$\int_M \phi_j^A u_B = 0, \quad \text{for } 1 \leq B < A \leq n+1. \quad (3.18)$$

We denote $mf_j = \sum_{A=1}^{n+1} |\nabla \phi_j^A|^2$ and write $\mathfrak{H}(\lambda_1, \dots, \lambda_m)$ as \mathfrak{H} for short.

We reproduce the previous steps from (3.8) to (3.15) for each ϕ_j and then take the limit. It follows that all the inequalities must be sharp. Indeed, noticing (3.8) and (3.11), we have

$$V(M, g) = \int_M \sum_{A=1}^{n+1} (\phi_j^A)^2 \leq \mathfrak{H}^{-1} \int_M mf_j \quad (3.19)$$

$$\leq \mathfrak{H}^{-1} m \left(\int_M f_j^{m/2} dv_g \right)^{2/m} (V(M))^{m-2} \quad (3.20)$$

$$\leq \mathfrak{H}^{-1} m (V_c(M, n, \phi_j))^{2/m} (V(M))^{m-2}. \quad (3.21)$$

Letting $j \rightarrow \infty$ and using $\mathfrak{H} = m \left(\frac{V_c(M, n)}{V(M, g)} \right)^{2/m}$, we obtained

$$V(M, g) = \lim_{j \rightarrow \infty} \int_M \sum_{A=1}^{n+1} (\phi_j^A)^2 = \mathfrak{H}^{-1} \lim_{j \rightarrow \infty} \int_M \sum_{A=1}^{n+1} |\nabla \phi_j^A|^2 \quad (3.22)$$

$$= \mathfrak{H}^{-1} \lim_{j \rightarrow \infty} \left(\int_M \left(\sum_{A=1}^{n+1} |\nabla \phi_j^A|^2 \right)^{m/2} dv_g \right)^{2/m} (V(M))^{m-2} \quad (3.23)$$

$$= V(M, g). \quad (3.24)$$

Hence, for any fixed A , $\{\phi_j^A\}$ is a bounded sequence in $W^{1,m}(M)$. Due to the compact inclusion $W^{1,m}(M) \hookrightarrow L^2(M)$, by passing to a subsequence we can assume that $\{\phi_j^A\}$ converges weakly in $W^{1,m}(M)$, strongly in $L^2(M)$, and pointwise a.e., to a map $\psi^A :$

$M \rightarrow \mathbb{R}$. We have

$$\sum_{A=1}^{n+1} (\psi^A)^2 = 1, \text{ a.e. on } M; \quad (3.25)$$

$$\lim_{j \rightarrow \infty} \int_M (\phi_j^A)^2 = \lim_{j \rightarrow \infty} \frac{1}{\lambda_A} \int_M |\nabla \phi_j^A|^2, \quad (3.26)$$

We give two claims.

Claim 1. For each A fixed, ψ^A belongs to the eigenspace E_{λ_A} corresponding to the eigenvalue λ_A , and then ψ^A is C^∞ .

Proof. Let π be the orthogonal projection on E_{λ_A} and π^\perp the projection on $E_{\lambda_A}^\perp$, then

$$\|\phi_j^A\|_{L^2}^2 = \|\pi \phi_j^A\|_{L^2}^2 + \|\pi^\perp \phi_j^A\|_{L^2}^2, \quad (3.27)$$

$$\|\nabla \phi_j^A\|_{L^2}^2 = \|\nabla \pi \phi_j^A\|_{L^2}^2 + \|\nabla \pi^\perp \phi_j^A\|_{L^2}^2. \quad (3.28)$$

Combining with (3.26), we derive that

$$\lim_{j \rightarrow \infty} \|\nabla \pi^\perp \phi_j^A\|_{L^2}^2 = \lim_{j \rightarrow \infty} \lambda_A \|\pi^\perp \phi_j^A\|_{L^2}^2. \quad (3.29)$$

Noticing (3.18), we have

$$\|\nabla \pi^\perp \phi_j^A\|_{L^2}^2 \geq \lambda_B \|\pi^\perp \phi_j^A\|_{L^2}^2, \quad (3.30)$$

where $\lambda_B (> \lambda_A)$ is the next distinct eigenvalue after λ_A . Hence,

$$0 \geq \lim_{j \rightarrow \infty} (\lambda_A - \lambda_B) \|\pi^\perp \phi_j^A\|_{L^2}^2 \geq 0, \quad (3.31)$$

which implies $\lim_{j \rightarrow \infty} \|\pi^\perp \phi_j^A\|_{L^2}^2 = 0$ and then $\pi^\perp \psi^A = 0$. This proves Claim 1. \square

Claim 2. $\{\phi_j^A\}$ converges to ψ^A strongly in $W^{1,2}(M)$.

Proof. By using Claim 1, we have

$$\|\nabla \phi_j^A - \nabla \psi^A\|_{L^2}^2 = \|\nabla \phi_j^A\|_{L^2}^2 + \|\nabla \psi^A\|_{L^2}^2 - 2\langle \nabla \phi_j^A, \nabla \psi^A \rangle_{L^2} \quad (3.32)$$

$$= \|\nabla \phi_j^A\|_{L^2}^2 + \lambda_A \|\psi^A\|_{L^2}^2 - 2\lambda_A \langle \phi_j^A, \psi^A \rangle_{L^2}, \quad (3.33)$$

By using (3.18), we derive

$$\lim_{j \rightarrow \infty} \|\nabla \phi_j^A - \nabla \psi^A\|_{L^2}^2 = 0. \quad (3.34)$$

This proves Claim 2. \square

Now we can continue the proof. Denote $mf = \sum_{A=1}^m |\nabla \psi^A|^2$. By taking the limit, all the inequalities from (3.8) to (3.15) for ϕ becomes the equalities, that is,

$$\begin{aligned} V(M, g) &= \int_M \sum_{A=1}^m |\psi^A|^2 = \sum_{A=1}^m \frac{1}{\lambda_A} \int_M |\nabla \psi^A|^2 + \sum_{A=m+1}^{n+1} \frac{1}{\lambda_A} \int_M |\nabla \psi^A|^2 \\ &= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M |\nabla \psi^A|^2 + \frac{1}{\lambda_m} \int_M \sum_{A=m+1}^{n+1} |\nabla \psi^A|^2 \end{aligned} \quad (3.35)$$

$$\begin{aligned} &= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M |\nabla \psi^A|^2 + \frac{1}{\lambda_m} \int_M f \sum_{A=1}^m \left(1 - \frac{1}{f} |\nabla \psi^A|^2\right) \\ &= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M |\nabla \psi^A|^2 + \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f \left(1 - \frac{1}{f} |\nabla \psi^A|^2\right) \end{aligned} \quad (3.36)$$

$$\begin{aligned}
 &= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M f \\
 &= \mathfrak{H}^{-1} m \left(\int_M f^{m/2} dv_g \right)^{2/m} (V(M))^{\frac{m-2}{m}} \\
 &= \mathfrak{H}^{-1} m \left(V_c(M, n) \right)^{2/m} (V(M))^{\frac{m-2}{m}} = V(M, g).
 \end{aligned}$$

The strong convergence in $W^{1,2}(M)$ proves that

$$\psi^* h_1 = \lim_{j \rightarrow \infty} \phi_j^* h_1 = \lim_{j \rightarrow \infty} f_j g = fg,$$

which means that $\psi : M \rightarrow \mathbb{S}^n$ is a conformal map with the conformal factor f .

Next, we prove the following lemma.

Lemma 3.2. *If the equality holds in (1.33), then $\lambda_1 = \dots = \lambda_m$ and $f = \lambda_1/m$.*

Proof. Recall Claim 1, we have

$$\Delta \psi^A = -\lambda_A \psi^A, \quad (3.37)$$

then

$$mf = \sum_{A=1}^{n+1} |\nabla \psi^A|^2 = \sum_{A=1}^{n+1} \left(\frac{1}{2} \Delta (\psi^A)^2 - \psi^A \Delta \psi^A \right) = \sum_{A=1}^{n+1} \lambda_A (\psi^A)^2. \quad (3.38)$$

We point out that $|\nabla \psi^A|^2/f \equiv 1$ is impossible. Otherwise, $1 \equiv |\nabla \psi^A|^2/f = |\tilde{\nabla} \psi^A|^2 \leq 1 - (\psi^A)^2$, which means $\psi^A \equiv 0$ and then $|\nabla \psi^A|^2/f = 0$, a contradiction. Hence, the equality in (3.36) gives $\lambda_t = \lambda_m$ for all $t \leq m-1$.

If $|\nabla \psi^A|^2 \equiv 0$, then ψ^A is constant on M , which follows that $\psi^A \equiv 0$ by using (3.37). If $\psi^s \not\equiv 0$ for some $s \geq m+1$, then the equality in (3.35) gives $\lambda_m = \lambda_s$. Hence, we conclude from (3.38) that

$$mf = \lambda_m \sum_{\substack{1 \leq A \leq n+1 \\ \psi^A \neq 0}} (\psi^A)^2 = \lambda_m = \lambda_1.$$

We complete the proof. \square

By scaling we may assume $\lambda_1 = m$. Then $\psi : M \rightarrow \mathbb{S}^n$ is an isometric immersion. Since each $\psi^A \in E_{\lambda_1}$, we know that ψ is minimal by Takahashi theorem.

Conversely, the sufficiency for the equality is directly from the following theorem when M is closed.

Theorem 3.3 ([15, Theorem 1.1]). *Let (M, g) be an $m(\geq 2)$ -dimensional closed Riemannian manifold. Suppose there exists a minimal isometric immersion ϕ of (M, g) in \mathbb{S}^n . Then*

$$V(M, g) = V_c(M, n, \phi) \geq V_c(M, n). \quad (3.39)$$

Moreover, if (M, g) is not isometric to (\mathbb{S}^m, h_1) , then $V(M, g) > V(M, (\gamma \circ \phi)^* \text{can})$ for all $\gamma \in G \setminus O(n+1)$.

Remark 3.4. *For each $a \in B^{n+1}$, we can define a conformal map γ_a on $\mathbb{S}^n(1)$ (cf. [33]):*

$$\gamma_a(x) = \frac{x + (\mu f + \lambda)a}{\lambda(1+f)}, \quad \forall x \in \mathbb{S}^n(1), \quad (3.40)$$

where B^{n+1} is the open unit ball in \mathbb{R}^{n+1} , x is the position vector of $\mathbb{S}^n(1)$, and

$$\lambda = (1 - |a|^2)^{-1/2}, \quad \mu = (\lambda - 1)|a|^{-2}, \quad f(x) = \langle x, a \rangle.$$

When $a = 0$, we set $\lambda = 1, \mu = 0, \gamma_0(x) = x$.

The proof of Theorem 3.3 is used the following fact, which allows us to replace $\text{Conf}(\mathbb{S}^n)$ by the subgroup $G = \{\gamma_a | a \in B^{n+1}\}$ when taking the supremum in the definition of $V(M, n, \phi)$. (see [14, pp. 259]):

For any $\gamma \in \text{Conf}(\mathbb{S}^n)$, there exists $a \in B^{n+1}$ and $r \in O(n+1)$ such that $\gamma = r \circ \gamma_a$.

3.3. A corollary.

Corollary 3.5. (1) Let (M, g) be a closed $m(\geq 2)$ -dimensional Riemannian manifold which can be minimally immersed into \mathbb{S}^n by ϕ . If (M, g) is not isometric to (\mathbb{S}^m, h_1) , then for any metric $\tilde{g} \in [g]$, we have

$$\mathfrak{H}(\tilde{g})V(M, \tilde{g})^{2/m} \leq mV(M, g)^{2/m}, \quad (3.41)$$

where $[g]$ is the conformal class of g , and $\mathfrak{H}(\tilde{g})$ is the harmonic mean of the first m nonzero eigenvalues with respect to \tilde{g} .

Moreover, equality holds if and only if the minimal immersion ϕ is given by a subspace of the first eigenfunctions and $\tilde{g} = kg$ for some constant $k > 0$.

(2) For any metric g on $\mathbb{S}^m, g \in [h_1]$, we have

$$\mathfrak{H}(g)V(\mathbb{S}^m, g)^{2/m} \leq \mathfrak{H}(h_1)V(\mathbb{S}^m, h_1)^{2/m}, \quad (3.42)$$

where equality holds if and only if $g = kh_1$ for some constant $k > 0$.

Proof. (1) The inequality (3.41) is from Theorem 1.3 and Theorem 3.3. Now assume that the equality holds. After rescaling, we can assume that $\mathfrak{H}(\tilde{g}) = m$. Then $V(M, \tilde{g}) = V(M, g)$ and there is a minimal immersion $\psi : (M, \tilde{g}) \rightarrow \mathbb{S}^n$ with $\lambda_1(\tilde{g}) = \cdots = \lambda_m(\tilde{g}) = m$ by the case of equality in Theorem 1.3.

By reviewing the proof of Theorem 1.3, we can assume $\psi = \gamma_a \circ \phi$ for some $a \in B^{n+1}$. Since (M, g) is not isometric to (\mathbb{S}^m, h_1) , γ_a must be the identity map by Theorem 3.3 (and Remark 3.4). This follows that $\tilde{g} = \psi^*h_1 = \phi^*h_1 = g$.

(2) This assertion follows from Theorem 1.3, by noting that $V_c(\mathbb{S}^m, g) = V_c(\mathbb{S}^m, h_1) = V(\mathbb{S}^m, h_1)$ and $\mathfrak{H}(h_1) = \lambda_1(h_1) = m$. \square

4. PROOFS FOR THE STEKLOV EIGENVALUES

In this section, we prove Theorems 1.8, 1.9 and 1.10. Since the main approaches are the same as in Sect. 3, we will omit some same details but emphasize the differences.

4.1. The inequality in (1.26). Recall that the variational characterization of the Steklov eigenvalue $\sigma_i (i \geq 1)$ (cf. [45, 46])

$$\sigma_i = \inf_{\substack{u \in H^1(M) \setminus \{0\} \\ u|_{\partial M} \neq 0}} \left\{ \frac{\int_M |\nabla u|^2}{\int_{\partial M} u^2} \mid \int_{\partial M} uv_j = 0, j = 0, \dots, i-1 \right\}, \quad (4.1)$$

where $\{v_i\}_{i \geq 0}$ is an orthonormal set of Steklov eigenfunctions.

Let $\phi : (M, g) \rightarrow B^n$ be a conformal map with $\phi(\partial M) \subset \partial B^n$. By analogous arguments in Sect. 3.1, there exists $\gamma \in \text{Conf}(B^n)$ such that $\Phi = \gamma \circ \phi = (\Phi_1, \dots, \Phi_n)$ satisfies that

$$\int_{\partial M} \Phi^A = 0, \quad A = 1, \dots, n. \quad (4.2)$$

$$\int_{\partial M} \Phi^A v_B = 0, \text{ for } 1 \leq B < A \leq n. \quad (4.3)$$

Let $\hat{\Phi}^A$ be a harmonic extension of $\Phi^A|_{\partial M}$. Now for each $A = 1, \dots, n$, we have

$$\sigma_A \int_{\partial M} (\Phi^A)^2 \leq \int_M |\nabla \hat{\Phi}^A|^2 \leq \int_M |\nabla \Phi^A|^2. \quad (4.4)$$

On the other hand, if we denote $\tilde{g} = \Phi^* g_{B^n} = fg$, then

$$0 \leq |\tilde{\nabla} \Phi^A|^2 \leq 1, \quad \sum_{A=1}^n |\tilde{\nabla} \Phi^A|^2 = m, \quad (4.5)$$

and

$$mf = \sum_{A=1}^n |\nabla \Phi^A|^2. \quad (4.6)$$

where $\tilde{\nabla}$ is the gradient operator on (M, \tilde{g}) .

An analogous computation as in Sect. 3.1 shows that

$$\begin{aligned} V(\partial M) &= \int_{\partial M} 1 \leq \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \sum_{A=m}^n \frac{1}{\sigma_A} \int_M f |\tilde{\nabla} \Phi^A|^2 \\ &\leq \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \frac{1}{\sigma_m} \int_M f \sum_{A=m}^n |\tilde{\nabla} \Phi^A|^2 \end{aligned} \quad (4.7)$$

$$\begin{aligned} &= \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \frac{1}{\sigma_m} \int_M f (m - \sum_{A=1}^{m-1} |\tilde{\nabla} \Phi^A|^2) \\ &= \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \frac{1}{\sigma_m} \int_M f \sum_{A=1}^{m-1} (1 - |\tilde{\nabla} \Phi^A|^2) + \frac{1}{\sigma_m} \int_M f \\ &\leq \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M f |\tilde{\nabla} \Phi^A|^2 + \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M f (1 - |\tilde{\nabla} \Phi^A|^2) + \frac{1}{\sigma_m} \int_M f \end{aligned} \quad (4.8)$$

$$= \sum_{A=1}^m \frac{1}{\sigma_A} \int_M f. \quad (4.9)$$

Hence, by using the Hölder inequalities, we have

$$\mathfrak{H}(\sigma_1, \dots, \sigma_m) V(\partial M) \leq m \int_M f dv_g \quad (4.10)$$

$$\leq m \left(\int_M f^{m/2} dv_g \right)^{2/m} (V(M))^{\frac{m-2}{m}} \quad (4.11)$$

$$= m \left(V(M, \tilde{g}) \right)^{2/m} (V(M))^{\frac{m-2}{m}} \quad (4.12)$$

$$\leq m \left(\sup_{\gamma \in \text{Conf}(B^n)} V(M, (\gamma \circ \phi)^* g_{B^n}) \right)^{2/m} (V(M))^{\frac{m-2}{m}}. \quad (4.13)$$

Taking the infimum over all non-degenerate conformal maps $\phi : M \rightarrow B^n$ with $\phi(\partial M) \subset \partial B^n$, we obtain the desired inequality.

4.2. **Equality in (1.26).** Now, we assume that the equality holds.

Choose a sequence of conformal maps $\phi_j : M \rightarrow B^n$ with $\phi_j(\partial M) \subset \partial B^n$ such that

$$\lim_{j \rightarrow \infty} V_{rc}(M, n, \phi_j) = V_{rc}(M, n), \quad (4.14)$$

and by composing with a conformal transformation of the sphere we may assume

$$\int_{\partial M} \phi_j^A = 0, \quad 1 \leq A \leq n. \quad (4.15)$$

$$\int_{\partial M} \phi_j^A v_B = 0, \quad \text{for } 1 \leq B < A \leq n \quad (4.16)$$

Let $\hat{\phi}_j^A$ be a harmonic extension of $\phi_j^A|_{\partial M}$ as before. Denote $mf_j = \sum_{A=1}^n |\nabla \phi_j^A|^2$ and write $\mathfrak{H}(\sigma_1, \dots, \sigma_m)$ as \mathfrak{H} for short.

Like the arguments in Sect. 3.2, We reproduce the previous steps in Sect. 4.1 for each ϕ_j and then take the limit. It follows that all the inequalities must be sharp.

$$V(\partial M) = \lim_{j \rightarrow \infty} \int_M \sum_{A=1}^n (\phi_j^A)^2 = \mathfrak{H}^{-1} \lim_{j \rightarrow \infty} \int_M \sum_{A=1}^n |\nabla \phi_j^A|^2 \quad (4.17)$$

$$= \mathfrak{H}^{-1} \lim_{j \rightarrow \infty} \left(\int_M \left(\sum_{A=1}^n |\nabla \phi_j^A|^2 \right)^{m/2} dv_g \right)^{2/m} (V(M))^{\frac{m-2}{m}} \quad (4.18)$$

$$= V(\partial M). \quad (4.19)$$

Hence, by passing to a subsequence we can assume that $\{\phi_j^A\}$ converges weakly in $W^{1,m}(M)$, strongly in $L^2(M)$, and pointwise a.e., to a map $\psi^A : M \rightarrow \mathbb{R}$. We have

$$\sum_{A=1}^n (\psi^A)^2 = 1, \quad \text{a.e. on } \partial M; \quad (4.20)$$

$$\sum_{A=1}^n (\psi^A)^2 \leq 1, \quad \text{a.e. on } M; \quad (4.21)$$

$$\int_{\partial M} (\phi_j^A)^2 \leq \frac{1}{\sigma_A} \int_M |\nabla \phi_j^A|^2, \quad \text{for } A = 1, \dots, n; \quad (4.22)$$

$$\lim_{j \rightarrow \infty} \sum_{A=1}^n \int_{\partial M} (\phi_j^A)^2 = \lim_{j \rightarrow \infty} \sum_{A=1}^n \frac{1}{\sigma_A} \int_M |\nabla \phi_j^A|^2, \quad (4.23)$$

Claim 3. For each A fixed, $\{\phi_j^A\}$ converges to ψ^A strongly in $W^{1,2}(M)$; ψ^A belongs to the eigenspace E_{σ_A} corresponding to the eigenvalue σ_A , and ψ^A is harmonic.

Proof. It follows from (4.22) and (4.23) that

$$\lim_{j \rightarrow \infty} \frac{1}{\sigma_A} \int_M |\nabla \phi_j^A|^2 = \lim_{j \rightarrow \infty} \int_{\partial M} (\phi_j^A)^2 = \int_{\partial M} (\psi^A)^2 \leq \frac{1}{\sigma_A} \int_M |\nabla \psi^A|^2, \quad (4.24)$$

On the other hand, $\phi_j^A \rightarrow \psi^A$ weakly in $W^{1,m}(M)$ implies

$$\int_M |\nabla \psi^A|^2 \leq \lim_{j \rightarrow \infty} \int_M |\nabla \phi_j^A|^2. \quad (4.25)$$

Therefore, we must have

$$\lim_{j \rightarrow \infty} \int_M |\nabla \phi_j^A|^2 = \int_M |\nabla \psi^A|^2 \quad (4.26)$$

and

$$\sigma_A \int_{\partial M} (\psi^A)^2 = \int_M |\nabla \psi^A|^2. \quad (4.27)$$

These equalities proves Claim 3. \square

Now we continue the proof. Denote $mf = \sum_{A=1}^n |\nabla \psi^A|^2$. Claim 3 implies $\psi : M \rightarrow B^n$ is a conformal map with the conformal factor f . Also, ψ is harmonic with $\psi(\partial M) \subset \partial B^n$. By taking the limit, we have

$$V(\partial M) = \int_{\partial M} \sum_{A=1}^n |\psi^A|^2 = \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M |\nabla \psi^A|^2 + \sum_{A=m}^n \frac{1}{\sigma_A} \int_M |\nabla \psi^A|^2 \quad (4.28)$$

$$= \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M |\nabla \psi^A|^2 + \frac{1}{\sigma_m} \int_M \sum_{A=m}^n |\nabla \psi^A|^2 \quad (4.29)$$

$$= \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M |\nabla \psi^A|^2 + \frac{1}{\sigma_m} \int_M f \sum_{A=1}^{m-1} \left(1 - \frac{1}{f}\right) |\nabla \psi^A|^2 + \frac{1}{\sigma_m} \int_M f$$

$$= \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M |\nabla \psi^A|^2 + \sum_{A=1}^{m-1} \frac{1}{\sigma_A} \int_M f \left(1 - \frac{1}{f}\right) |\nabla \psi^A|^2 + \frac{1}{\sigma_m} \int_M f \quad (4.30)$$

$$= \sum_{A=1}^m \frac{1}{\sigma_A} \int_M f \quad (4.31)$$

$$= \mathfrak{H}^{-1} m \left(\int_M f^{m/2} dv_g \right)^{2/m} (V(M))^{\frac{m-2}{m}} \quad (4.32)$$

$$= \mathfrak{H}^{-1} m \left(V_{rc}(M, n) \right)^{2/m} (V(M))^{\frac{m-2}{m}} = V(M, g). \quad (4.33)$$

The strong convergence in $W^{1,2}(M)$ proves that

$$\tilde{g} := \psi^* g_{B^n} = \lim_{j \rightarrow \infty} \phi_j^* g_{B^n} = \lim_{j \rightarrow \infty} f_j g = fg, \quad (4.34)$$

which means that $\psi : M \rightarrow B^n$ is a conformal map with the conformal factor f .

For $s \geq m+1$, from (4.29) we know that if $\nabla \psi^s \not\equiv 0$ on M , then $\sigma_s = \sigma_m$; if $\nabla \psi^s \equiv 0$ on M , then we still have $\sigma_s \int_{\partial M} |\psi|^2 = \int_M |\nabla \psi^s|^2 = 0$. Therefore, we always have $\sigma_s \in E_{\sigma_m}$ for any $s \geq m+1$.

When $m > 2$, equality in the Hölder inequality (4.32) implies f is constant. Hence, ψ is an isomeric immersion after rescaling ($f \equiv 1$ and $\sum_{A=1}^m \frac{1}{\sigma_A} = m$), and then it is a minimal immersion since each coordinate function is harmonic.

4.3. Proof of Theorems 1.9 and 1.10. As pointed out in the proof of [17, Theorem 2.3], for any compact surface M with boundary, there exists a conformal branched cover $\phi : M \rightarrow D$ with $\deg(\phi) \leq \text{genus}(M) + k$ (cf. [1, 20]).

Since ϕ is proper we have $\phi(\partial M) \subset \partial D$. For the same reason as in Sect. 4.1, we may assume that the coordinate functions ϕ^1, ϕ^2 can be used as the test functions. Hence, we

have (cf. (4.10))

$$\mathfrak{H}(\sigma_1, \sigma_2)V(\partial M) \leq \int_M \sum_{i=1}^2 |\nabla \phi^i|^2 dv_g \quad (4.35)$$

Since ϕ is conformal and $\dim M = 2$, the right hand of (4.35) equals to $2V(\phi(M))$ and bounded by $2(\text{genus}(M) + k)\pi$ from above. This proves Theorem 1.9.

To prove Theorem 1.10, we just notice that the stereographic projection from \mathbb{R}^2 to (\mathbb{S}^2, h_1) is conformal and then consider the conformal map $\psi : M \rightarrow (\mathbb{S}^2, h_1)$. By using the coordinate functions ψ^1, ψ^2, ψ^3 (in \mathbb{R}^3) as the test functions, we have

$$\begin{aligned} \mathfrak{H}(\sigma_1, \sigma_2)V(\partial M) &\leq \int_M \sum_{i=1}^3 |\nabla \psi^i|^2 dv_g = 2V(\psi(M)) \\ &< 2V(\mathbb{S}^2, h_1) \deg(\psi) \leq 8\pi(\text{genus}(M) + 1). \end{aligned}$$

Hence, we complete the proof.

5. PROOFS FOR THE REILLY-TYPE INEQUALITIES

In this section, we prove Theorem 1.13 and give a general version.

5.1. The inequality in (1.33). For the immersion $x : M \rightarrow \mathbb{R}^n(c)$, by analogous arguments in Sect. 3.1, there exists a regular conformal map $\Gamma : \mathbb{R}^n(c) \rightarrow \mathbb{S}^n \subset \mathbb{R}^{n+1}$ such that the immersion $\Phi = \Gamma \circ x = (\Phi^1, \dots, \Phi^{n+1})$ satisfies that

$$\begin{aligned} \int_M \Phi^A &= 0, \quad A = 1, \dots, n+1. \\ \int_M \Phi^A u_B &= 0, \quad \text{for } 1 \leq B < A \leq n+1. \end{aligned}$$

Set $\Gamma^*h_1 = e^{2\rho}h_c$, then $g_M = x^*h_c$, $\tilde{g}_M = (\Gamma \circ x)^*h_1$.

By the variational characterization (3.2), for each $A = 1, \dots, n+1$, we have

$$\lambda_A \int_M (\Phi^A)^2 \leq \int_M |\nabla \Phi^A|^2 = \int_M e^{2\rho} |\tilde{\nabla} \Phi^A|^2, \quad (5.1)$$

An analogous computation as in Sect. 3.1 shows that

$$\begin{aligned} V(M) &= \int_M 1 \leq \sum_{A=1}^m \frac{1}{\lambda_A} \int_M e^{2\rho} |\tilde{\nabla} \Phi^A|^2 + \sum_{A=m+1}^{n+1} \frac{1}{\lambda_A} \int_M e^{2\rho} |\tilde{\nabla} \Phi^A|^2 \\ &\leq \sum_{A=1}^m \frac{1}{\lambda_A} \int_M e^{2\rho} |\tilde{\nabla} \Phi^A|^2 + \frac{1}{\lambda_m} \int_M e^{2\rho} \sum_{A=m+1}^{n+1} |\tilde{\nabla} \Phi^A|^2 \\ &= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M e^{2\rho} |\tilde{\nabla} \Phi^A|^2 + \frac{1}{\lambda_m} \int_M e^{2\rho} \sum_{A=1}^m (1 - |\tilde{\nabla} \Phi^A|^2) \\ &\leq \sum_{A=1}^m \frac{1}{\lambda_A} \int_M e^{2\rho} |\tilde{\nabla} \Phi^A|^2 + \sum_{A=1}^m \frac{1}{\lambda_A} \int_M e^{2\rho} (1 - |\tilde{\nabla} \Phi^A|^2) \quad (5.2) \end{aligned}$$

$$= \sum_{A=1}^m \frac{1}{\lambda_A} \int_M e^{2\rho}. \quad (5.3)$$

Hence, by using (2.15) we derive that

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) = \frac{m}{\sum_{i=1}^m \frac{1}{\lambda_i}} \leq \frac{m}{V(M)} \int_M e^{2\rho} \leq \frac{m}{V(M)} \int_M (c + |\mathbf{H}|^2).$$

5.2. Equality in (1.33). Firstly, we discuss the necessary conditions for equality in (1.33).

When M is minimally immersed in \mathbb{S}^n , we know that $\lambda = m$ is an eigenvalue and its multiplicity is not less than m since $\Delta x = -mx$. Hence, $\mathfrak{H}(\lambda_1, \dots, \lambda_m) = \frac{m}{V(M)} \int_M (c + |\mathbf{H}|^2) = m$ implies $\lambda_1 = \dots = \lambda_m = m$, i.e., the immersion x is given by a subspace of the first eigenfunctions.

When $c \neq 1$ or M is not minimal in \mathbb{S}^n , following the proof of Lemma 3.2, equality in (5.2) implies $\lambda_1 = \dots = \lambda_m$. This makes us reduce the equality case in (1.33) to the equality case in (1.32) and complete the proof for the necessity.

Next, we discuss the sufficient conditions for equality in (1.33). We already have

$$\lambda_1 \leq \mathfrak{H}(\lambda_1, \dots, \lambda_m) \leq \frac{m}{V(M)} \int_M (c + |\mathbf{H}|^2),$$

Noting Theorem 1.11, from either case (1) or case (2) we can derive

$$\lambda_1 = \frac{m}{V(M)} \int_M (c + |\mathbf{H}|^2),$$

so the equality in (1.33) is attained.

5.3. A general version. When the ambient is not the space form, we can prove the following theorem, which implies [15, Theorem 2].

Theorem 5.1. *Let (N, g_N) be a Riemannian manifold of dimension n (possibly not complete) which admits a conformal immersion in the sphere (\mathbb{S}^n, h_1) . Then, for any $m(\geq 2)$ -dimensional closed submanifold M immersed in (N, g_N) by ϕ , we have*

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \leq \frac{m}{V(M)} \int_M (|\mathbf{H}_\phi|^2 + \bar{\mathcal{R}}_\phi), \quad (5.4)$$

where K_N is the sectional curvature of (N, g_N) , and

$$\bar{\mathcal{R}}_\phi = \frac{1}{m(m-1)} \sum_{i \neq j} K_N(e_i, e_j) \quad (5.5)$$

for a local orthonormal tangent frame $\{e_i\}$ on M .

Moreover, equality holds if and only if $\lambda_1 = \dots = \lambda_m$ and $|\mathbf{H}_\phi|^2 + \bar{\mathcal{R}}_\phi$ equals the constant λ_1/m ,

Proof. The inequality follows from the same arguments as in Sect. 5.1 (just using (2.13) instead of (2.15) in the last step). The sufficiency for the equality is obvious. The necessary for the equality is due to Lemma 3.2. We omit the details. \square

6. PROOFS OF THEOREMS 1.14 AND 1.17

Proof of Theorem 1.14. When M is a submanifold of \mathbb{R}^n , without lost of generality, we assume that the coordinate functions satisfy

$$\int_M x^A = 0 \text{ for } A = 1, \dots, n.$$

We can assume further

$$\int_M x^A u_B = 0 \text{ for } 1 \leq B < A \leq n$$

by using the QR-decomposition. We claim

$$\mathfrak{H}(\lambda_1, \dots, \lambda_m) \int_M |x|^2 \leq mV(M). \quad (6.1)$$

Then (6.1) is obtained by a similar computation (cf. (5.3)) as in Sect. 5.1, just noting that the conformal factor $e^{2\rho} \equiv 1$ since we don't need the conformal transformation, and replacing the left side $V(M) = \int_M 1$ by $\int_M |x|^2$ since we don't have $M \subset \mathbb{S}^{n+1}$.

Multiplying the both sides of (6.1) by $\int_M |H_T|^2$, we have

$$\begin{aligned} mV(M) \int_M |H_T|^2 &\geq \mathfrak{H}(\lambda_1, \dots, \lambda_m) \int_M |x|^2 \int_M |H_T|^2 \\ &\geq \mathfrak{H}(\lambda_1, \dots, \lambda_m) \left(\int_M |x| \cdot |H_T| \right)^2 \end{aligned} \quad (6.2)$$

$$\geq \mathfrak{H}(\lambda_1, \dots, \lambda_m) \left(\int_M \langle x, H_T \rangle \right)^2 \quad (6.3)$$

$$= \mathfrak{H}(\lambda_1, \dots, \lambda_m) \left(\int_M \text{tr } T \right)^2, \quad (6.4)$$

where we used the Hölder inequality, the Cauchy-Schwarz inequalities, and the Hsiung-Minkowski formula (2.22).

Equality case: necessity. If the equality in (1.34) holds, then (6.1) is sharp, which implies $\lambda_1 = \dots = \lambda_m$ (cf. Lemma 3.2). Moreover, we have $\Delta x = -\lambda_1 x$. Hence, by the well-known Takahashi theorem ([42, Theorem 3]), we conclude that M is actually a minimal submanifold in a hypersphere $\mathbb{S}^{n-1}(r)$ with the radius $r = \sqrt{m/\lambda_1}$.

Equalities in (6.2) and (6.3) imply H_T is proportional to x . Suppose $H_T = kx$ for some constant $k \neq 0$, then $|H_T| = |k||x| = |k|r$. From (2.21) we derive $\text{tr } T = -\langle x, H_T \rangle = -kr^2$ is also constant.

Let $\{e_{m+2}, \dots, e_n\}$ be the normal frame of the immersion from M to $\mathbb{S}^{n-1}(r)$, then $\{e_{m+1} = \frac{x}{r}, e_{m+2}, \dots, e_n\}$ forms a local normal frame of M in \mathbb{R}^n . Since $H_T = kx$ and

$$H_T = \sum_{i,j,\alpha} T_{ij} h_{ij}^\alpha e_\alpha = \sum_{i,j} T_{ij} h_{ij}^{m+1} e_{m+1} + \sum_{i,j;\alpha \geq m+2} T_{ij} h_{ij}^\alpha e_\alpha, \quad (6.5)$$

we obtain $H'_T := \sum_{i,j;\alpha \geq m+2} T_{ij} h_{ij}^\alpha e_\alpha$ must vanish, i.e., M is T -minimal in $\mathbb{S}^{n-1}(r)$.

Equality case: sufficiency. Conversely, if M is minimal and T -minimal in $\mathbb{S}^{n-1}(r)$, then $\lambda_1 = \dots = \lambda_m$ and (6.5) becomes

$$H_T = \sum_{i,j} T_{ij} h_{ij}^{m+1} e_{m+1} = - \sum_{i,j} T_{ij} \frac{1}{r} \delta_{ij} e_{m+1} = - \text{tr } T \frac{1}{r^2} x,$$

hence, $|H_T| = \frac{1}{r} \text{tr } T$. Since $r^2 = m/\lambda_1$ and $\text{tr } T$ is constant, one can easily check that equality in (1.34) holds. \square

Proof of Theorem 1.15. Since $x : M \rightarrow \mathbb{S}^n \subset \mathbb{R}^{n+1}$, we consider M as a submanifold of \mathbb{R}^{n+1} . Let H_T and H'_T be associated to $M \rightarrow \mathbb{S}^n$ and $M \rightarrow \mathbb{R}^{n+1}$, respectively. Then we can apply Theorem 1.14 to obtain (1.35), just noticing the following relations:

$$H'_T = H_T - (\text{tr } T)x, \quad |H'_T|^2 = |H_T|^2 + (\text{tr } T)^2.$$

Equality case: The sufficiency can be verified by using the Takahashi's theorems, we omit the details. Next, we check the necessity.

Since $\text{tr } T \neq 0$ implies $H'_T \neq 0$, equality in (1.35) implies $\text{tr } T$ is constant, and M is minimal and T -minimal in a hypersphere S' of \mathbb{R}^{n+1} by Theorem 1.14.

If $S'(r)$ is just \mathbb{S}^n , then the immersion is given by a subspace of the first eigenfunctions. This is the first case.

Otherwise, M lies in $\Sigma_1 := S' \cap \mathbb{S}^n$. We know that Σ_1 is a geodesic sphere of \mathbb{S}^n . Since M is minimal and T -minimal in S' , M is also minimal and T -minimal in Σ_1 . The geodesic radius of Σ_1 can be determined by the Gauss equation. This is the second case. \square

Proof of Theorem 1.17. The proof is similar with the proof of Theorem 1.14, but there are two main differences: one is that we should apply the Hsiung-Minkowski formula for ∂M instead of M since ∂M is closed; the other one is that $m = n$ is allowed when considering Steklov eigenvalues.

We assume that the coordinate functions satisfy

$$\int_{\partial M} x^A = 0 \text{ for } A = 1, \dots, n.$$

We claim

$$\mathfrak{H}(\sigma_1, \dots, \sigma_m) \int_{\partial M} |x|^2 \leq mV(M). \quad (6.6)$$

When $n > m$, we assume further

$$\int_{\partial M} x^A v_B = 0 \text{ for } 1 \leq B < A \leq n$$

by using the QR-decomposition. Then (6.6) is obtained by a similar computation as in Sect. 4.1 (cf. (4.10)), just noting that $f \equiv 1$ since we don't need the conformal transformation, and replacing $V(\partial M) = \int_{\partial M} 1$ by $\int_{\partial M} |x|^2$ since we don't have $\partial M \subset \partial B^n = \mathbb{S}^{n-1}$.

When $n = m$, this means M is a domain of \mathbb{R}^m . We use the following characterization for the inverse trace of the Steklov eigenvalues (see [2, pp. 99]):

$$\sum_{i=1}^m \frac{1}{\sigma_i} = \max \left\{ \sum_{i=1}^m \int_{\partial M} u_i^2 \left| \begin{array}{l} u_i \in W^{1,2}(M), \int_M \nabla u_i \nabla u_j = \delta_{ij}, \int_M \nabla u_i \nabla v_0 = 0, \\ \int_{\partial M} u_i = 0 \text{ for } 1 \leq i, j \leq m \end{array} \right. \right\}.$$

Noticing that the eigenfunction v_0 is constant, we take $u_i = x^i / \sqrt{V(M)}$. Then we have

$$\sum_{i=1}^m \frac{1}{\sigma_i} \geq \sum_{i=1}^m \int_{\partial M} \left(\frac{x^i}{\sqrt{V(M)}} \right)^2 = \frac{1}{V(M)} \int_{\partial M} |x|^2,$$

which is equivalent to (6.6).

Multiplying the both sides of (6.6) by $\int_{\partial M} |H_T|^2$, we have

$$\begin{aligned} mV(M) \int_{\partial M} |H_T|^2 &\geq \mathfrak{H}(\sigma_1, \dots, \sigma_m) \int_{\partial M} |x|^2 \int_{\partial M} |H_T|^2 \\ &\geq \mathfrak{H}(\sigma_1, \dots, \sigma_m) \left(\int_{\partial M} |x| \cdot |H_T| \right)^2 \end{aligned}$$

$$\begin{aligned} &\geq \mathfrak{H}(\sigma_1, \dots, \sigma_m) \left(\int_{\partial M} \langle x, H_T \rangle \right)^2 \\ &= \mathfrak{H}(\sigma_1, \dots, \sigma_m) \left(\int_{\partial M} \operatorname{tr} T \right)^2, \end{aligned}$$

□

REFERENCES

- [1] Lars V. Ahlfors, *Open Riemann surfaces and extremal problems on compact subregions*, Comment. Math. Helv. **24** (1950), 100–134. MR36318
- [2] Catherine Bandle, *Isoperimetric inequalities and applications*, Monographs and Studies in Mathematics, vol. 7, Pitman (Advanced Publishing Program), Boston, Mass.-London, 1980. MR572958
- [3] Linfen Cao and Haizhong Li, *r-minimal submanifolds in space forms*, Ann. Global Anal. Geom. **32** (2007), no. 4, 311–341. MR2346221
- [4] Bang-Yen Chen, *Geometry of submanifolds*, Marcel Dekker, Inc., New York, 1973. Pure and Applied Mathematics, No. 22. MR0353212
- [5] ———, *Some conformal invariants of submanifolds and their applications*, Boll. Un. Mat. Ital. (4) **10** (1974), 380–385. MR0370436
- [6] Hang Chen and Xudong Gui, *Reilly-type inequalities for submanifolds in Cartan-Hadamard manifolds*, arXiv:2206.11164 (2022).
- [7] Hang Chen and Xianfeng Wang, *Sharp Reilly-type inequalities for a class of elliptic operators on submanifolds*, Differential Geom. Appl. **63** (2019), 1–29. MR3896192
- [8] Hang Chen and Guofang Wei, *Reilly-type inequalities for p-Laplacian on submanifolds in space forms*, Nonlinear Anal. **184** (2019), 210–217. MR3921104
- [9] Shiing-Shen Chern, *Minimal submanifolds in a Riemannian manifold*, University of Kansas, Department of Mathematics Technical Report 19 (New Series), Univ. of Kansas, Lawrence, Kan., 1968. MR0248648
- [10] Bruno Colbois, Alexandre Girouard, Carolyn Gordon, and David Sher, *Some recent developments on the Steklov eigenvalue problem*, arXiv:2212.12528 (2022).
- [11] Bruno Colbois, Alexandre Girouard, and Asma Hassannezhad, *The Steklov and Laplacian spectra of Riemannian manifolds with boundary*, J. Funct. Anal. **278** (2020), no. 6, 108409, 38. MR4054110
- [12] Bruno Colbois and Sheela Verma, *Sharp Steklov upper bound for submanifolds of revolution*, J. Geom. Anal. **31** (2021), no. 11, 11214–11225. MR4310168
- [13] A. El Soufi and S. Ilias, *Le volume conforme et ses applications d’après Li et Yau*, Séminaire de Théorie Spectrale et Géométrie, Année 1983–1984, 1984, pp. VII.1–VII.15. MR1046044
- [14] ———, *Immersiones minimales, première valeur propre du laplacien et volume conforme*, Math. Ann. **275** (1986), no. 2, 257–267. MR854009
- [15] Ahmad El Soufi and Saïd Ilias, *Une inégalité du type “Reilly” pour les sous-variétés de l’espace hyperbolique*, Comment. Math. Helv. **67** (1992), no. 2, 167–181. MR1161279
- [16] ———, *Second eigenvalue of Schrödinger operators and mean curvature*, Comm. Math. Phys. **208** (2000), no. 3, 761–770. MR1736334
- [17] Ailana Fraser and Richard Schoen, *The first Steklov eigenvalue, conformal geometry, and minimal surfaces*, Adv. Math. **226** (2011), no. 5, 4011–4030. MR2770439
- [18] ———, *Sharp eigenvalue bounds and minimal surfaces in the ball*, Invent. Math. **203** (2016), no. 3, 823–890. MR3461367
- [19] ———, *Shape optimization for the Steklov problem in higher dimensions*, Adv. Math. **348** (2019), 146–162. MR3925929
- [20] Alexandre Gabard, *Sur la représentation conforme des surfaces de Riemann à bord et une caractérisation des courbes séparantes*, Comment. Math. Helv. **81** (2006), no. 4, 945–964. MR2271230
- [21] Jean-François Grosjean, *Upper bounds for the first eigenvalue of the Laplacian on compact submanifolds*, Pacific J. Math. **206** (2002), no. 1, 93–112. MR1924820
- [22] ———, *Extrinsic upper bounds for the first eigenvalue of elliptic operators*, Hokkaido Math. J. **33** (2004), no. 2, 319–339. MR2073001
- [23] Ernst Heintze, *Extrinsic upper bounds for λ_1* , Math. Ann. **280** (1988), no. 3, 389–402. MR936318
- [24] Joseph Hersch, *Quatre propriétés isopérimétriques de membranes sphériques homogènes*, C. R. Acad. Sci. Paris Sér. A-B **270** (1970), A1645–A1648. MR292357

- [25] Mikhail Karpukhin, Nikolai Nadirashvili, Alexei V. Penskoi, and Iosif Polterovich, *An isoperimetric inequality for Laplace eigenvalues on the sphere*, J. Differential Geom. **118** (2021), no. 2, 313–333. MR4278696
- [26] Gerasim Kokarev, *Variational aspects of Laplace eigenvalues on Riemannian surfaces*, Adv. Math. **258** (2014), 191–239. MR3190427
- [27] ———, *Conformal volume and eigenvalue problems*, Indiana Univ. Math. J. **69** (2020), no. 6, 1975–2003. MR4170085
- [28] Nicholas Korevaar, *Upper bounds for eigenvalues of conformal metrics*, J. Differential Geom. **37** (1993), no. 1, 73–93. MR1198600
- [29] Nikolay Kuznetsov, Tadeusz Kulczycki, Mateusz Kwaśnicki, Alexander Nazarov, Sergey Poborchi, Iosif Polterovich, and B. Siudeja, *The legacy of Vladimir Andreevich Steklov*, Notices Amer. Math. Soc. **61** (2014), no. 1, 9–22. MR3137253
- [30] Peter Li and Shing Tung Yau, *A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces*, Invent. Math. **69** (1982), no. 2, 269–291. MR674407
- [31] Fernando Manfio, Julien Roth, and Abhitosh Upadhyay, *Extrinsic eigenvalues upper bounds for submanifolds in weighted manifolds*, Annals of Global Analysis and Geometry (2022).
- [32] Ana-Maria Matei, *Conformal bounds for the first eigenvalue of the p -Laplacian*, Nonlinear Anal. **80** (2013), 88–95. MR3010757
- [33] Sebastián Montiel and Antonio Ros, *Minimal immersions of surfaces by the first eigenfunctions and conformal area*, Invent. Math. **83** (1986), no. 1, 153–166. MR813585
- [34] Nikolai Nadirashvili, *Isoperimetric inequality for the second eigenvalue of a sphere*, J. Differential Geom. **61** (2002), no. 2, 335–340. MR1972149
- [35] Nikolai Nadirashvili and Yannick Sire, *Isoperimetric inequality for the third eigenvalue of the Laplace-Beltrami operator on S^2* , J. Differential Geom. **107** (2017), no. 3, 561–571. MR3715349
- [36] Yanyan Niu and Shicheng Xu, *Total squared mean curvature of immersed submanifolds in a negatively curved space*, arXiv:2106.01912v1 (2021).
- [37] Romain Petrides, *Maximization of the second conformal eigenvalue of spheres*, Proc. Amer. Math. Soc. **142** (2014), no. 7, 2385–2394. MR3195761
- [38] Robert C. Reilly, *On the first eigenvalue of the Laplacian for compact submanifolds of Euclidean space*, Comment. Math. Helv. **52** (1977), no. 4, 525–533. MR0482597
- [39] Julien Roth, *General Reilly-type inequalities for submanifolds of weighted Euclidean spaces*, Colloq. Math. **144** (2016), no. 1, 127–136. MR3485886
- [40] W. Stekloff, *Sur les problèmes fondamentaux de la physique mathématique*, Ann. Sci. École Norm. Sup. (3) **19** (1902), 191–259. MR1509012
- [41] ———, *Sur les problèmes fondamentaux de la physique mathématique (suite et fin)*, Ann. Sci. École Norm. Sup. (3) **19** (1902), 455–490. MR1509018
- [42] Tsunero Takahashi, *Minimal immersions of Riemannian manifolds*, J. Math. Soc. Japan **18** (1966), 380–385. MR0198393
- [43] Robert Weinstock, *Inequalities for a classical eigenvalue problem*, J. Rational Mech. Anal. **3** (1954), 745–753. MR64989
- [44] C. Xia and C. Xiong, *Escobar’s conjecture on a sharp lower bound for the first nonzero Steklov eigenvalue*, arXiv:1907.07340 (2019).
- [45] Changwei Xiong, *Optimal estimates for Steklov eigenvalue gaps and ratios on warped product manifolds*, Int. Math. Res. Not. IMRN **22** (2021), 16938–16962. MR4345819
- [46] ———, *On the spectra of three Steklov eigenvalue problems on warped product manifolds*, J. Geom. Anal. **32** (2022), no. 5, Paper No. 153, 35. MR4386421
- [47] Paul C. Yang and Shing Tung Yau, *Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **7** (1980), no. 1, 55–63. MR577325

SCHOOL OF MATHEMATICS AND STATISTICS, NORTHWESTERN POLYTECHNICAL UNIVERSITY, XI’ AN 710129, P. R. CHINA

Email address: chenhang86@nwpu.edu.cn