

Neutrino propagation in the neutron star with uncertainties from nuclear, hadron, and particle physics

Parada T. P. Hutaauruk,^{1,2,*} Hana Gil,^{3,†} Seung-il Nam,^{1,3,4,‡} and Chang Ho Hyun^{2,3,§}

¹*Department of Physics, Pukyong National University (PKNU), Busan 48513, Korea*

²*Department of Physics Education, Daegu University, Gyeongsan 38453, Korea*

³*Center for Extreme Nuclear Matters (CENuM),
Korea University, Seoul 02841, Korea*

⁴*Asia Pacific Center for Theoretical Physics (APCTP), Pohang 37673, Korea*

(Dated: March 14, 2023)

Abstract

In the present work, we investigate the neutral-current neutrino-nucleon scattering in the nuclear medium using various energy-density functional (EDF) models such as the KIDS (Korea-IBS-Daegu-SKKU) and SLy4, together with the quark-meson coupling (QMC) model for the nucleon form factors at finite density. The differential cross section (DCS) and neutrino mean free path (NMFP) are computed numerically, considering the density-dependent nucleon form factors (DDFF) and neutrino structural properties such as the neutrino magnetic moment (NMM) and its electric charge radius (NCR). It turns out that the DDFF decreases the scattering cross-section, while the NCR increases it considerably. The effect of the NMM turns out to be almost negligible. We also observe that the value of the neutron effective mass is of importance in the neutron-star cooling process, indicating that for the neutron effective mass larger than the mass in free space, the neutrino can interact with matter at densities $\rho \gtrsim 1.5\rho_0$ in the neutron star with radius 13 km.

*Electronic address: phutaauruk@gmail.com

†Electronic address: khn1219@gmail.com

‡Electronic address: sinam@pknu.ac.kr

§Electronic address: hch@daegu.ac.kr

I. INTRODUCTION

In the recent work, we investigated the weak interaction of neutrinos in the homogeneous neutron-star (NS) matter within the framework of Korea-IBS-Daegu-SKKU (KIDS) density functional [1]. The work focused on the effect of uncertainties and/or corrections of the nuclear matter equation of state (EoS), i.e., symmetry energy and nucleon effective mass, to the neutrino mean free path (NMFP) within the NS systematically. We found that the NMFP depends strongly on these uncertainties and/or corrections. Compared with the NS radius, the NMFP could be as short as about half of the NS radius, but also could be larger than the NS radius. Such wide-range values of NMFP can lead to a different result in the cooling behavior of the NS. Overall, the work of Ref. [1] demonstrated the importance of the accurate determination of nuclear matter EoS in the neutrino-weak interaction at finite density and zero temperature. However, it is worth noting that in Ref. [1], the nucleons are treated as point particles, while, in fact, the nucleon has a structure such as the electromagnetic form factor in the transverse momentum and parton distribution functions in the longitudinal momentum. Such structure of the nucleon has been confirmed not only theoretically [2], but also experimentally [3–6]. Moreover, in the past, many studies have been done in investigating the neutral and charged current weak interaction of a neutrino with the matter by considering the electromagnetic form factor in the free space [7–12]. In the present work, we consider the structure of the nucleon—which is originally from non-perturbative quantum chromodynamic (QCD) aspect of hadron and particle physics in both free space and medium. It is well known that the electromagnetic form factors are expected to be modified in the medium [13–18, 20]. Such medium effect can be non-negligible in the model [21, 22], providing another source of the corrections that can affect the NMFP.

In most standard studies, neutrinos are assumed to be elementary particles. However, several experiments show the non-zero values for the magnetic moment and charge radius of the neutrino [23–30], although the evidence is not firmly established yet. More theoretical studies and experiments using modern facilities like CONUS [31], DUNE [32], NOMAD [33], MiniBooNe [34], MINERvA [35], Hyper-Kamiokande, and other reactor or accelerator experiments as well as galactic or atmospheric neutrinos are really needed to collect more data in order to establish the properties of the neutrino. If the neutrinos have an internal structure, this evidence of neutrino moment magnetic (NMM) and charge radius (NCR) will signifi-

cantly impact elementary particle physics, nuclear physics, astrophysics, and cosmology in the standard model calculation.

In this study, we take into account the aforementioned uncertainties and/or corrections from nuclear physics, hadron physics, elementary particle physics, i.e., nucleon form factor (free space and medium), and NMM and NCR in the description of neutrino electro-weak interaction at finite density and zero temperature. The values of the NMM and NCR are obtained from experimental constraints [26, 29, 30]. We then calculate the differential cross-section (DCS) for the neutrino-nucleon scattering and NMFP of the neutrinos in the core of the NS. The roles of the nucleon and neutrino corrections in the neutrino propagation inside NS are explored in further detail. We find that the in-medium corrections of the nucleon form factor and NCR give a significant impact on the DCS and NMFP. Compared to the vacuum nucleon form factor (VFF), the density dependence of the nucleon form factor (DDFF) decreases the DCS, implying an increase in the NMFP. The increase of the NMFP is more significant at higher matter density. It means that the neutrino with the DDFF will more freely escape from the core of NS. The DCS does not change significantly by taking an NMM value from the experimental constraints [26]. This indicates that the NMM plays a small role in the neutrino emission from the NS core. However, it is drastically changed when we consider a finite NCR value obtained from the experiment constraint [29, 30]. It increases the DCS significantly in comparison with other scenarios without the NCR. This shows that the neutrinos are more strongly interacting with the matter if the NCR contribution is taken into account appropriately.

The present work is organized as follows: Section II is devoted to a brief description of the theoretical framework. The numerical results and related discussions are given in Section III. The final Section is devoted to the summary.

II. NEUTRINO-NUCLEON INTERACTION

Here we adopt four non-relativistic energy density functional (EDF) models: KIDS0, KIDS-A, KIDS0-m*87, and SLy4 [1]. All the models have yielded identical values of the saturation density $\rho_0 = 0.16 \text{ fm}^{-3}$ and the binding energy per nucleon $E_B = 16 \text{ MeV}$. However, they have distinctive behavior for the EoS at densities below and above the saturation.

	K_0	J	L	K_{sym}	m_s^*/M	m_v^*/M
KIDS0	240	32.8	49.1	-156.7	1.0	0.8
KIDS-A	230	33	66	-139.5	1.0	0.8
KIDS0-m*87	240	32.8	49.1	-156.7	0.8	0.7
SLy4	229.9	32	45.9	-119.7	0.7	0.8

TABLE I: Nuclear matter parameters of the models. K_0 , J , L and K_{sym} are in the units of MeV. m_s^* , m_v^* and M denote the isoscalar effective mass, isovector effective mass, and nucleon mass in free space, respectively.

A conventional expansion of the energy per nucleon can be written as

$$\mathcal{E}(\rho, \delta) = E(\rho) + S(\rho)\delta^2 + O(\delta^3), \quad (1)$$

$$E(\rho) = E_B + \frac{1}{2}K_0x^2 + O(x^3), \quad (2)$$

$$S(\rho) = J + Lx + \frac{1}{2}K_{\text{sym}}x^2 + O(x^3), \quad x = \frac{\rho - \rho_0}{3\rho_0}, \quad \delta = \frac{\rho_n - \rho_p}{\rho}. \quad (3)$$

The parameters that characterize the density-dependence of EoS are summarized in Tab. I. The KIDS0, KIDS0-m*87, and SLy4 models have similar density dependencies for the symmetry energy but are very different in the effective mass. On the other hand, the KIDS0 and KIDS-A models have similar effective masses, whereas the symmetry energy for the KIDS-A model is much stiffer than that for the KIDS0 model. Comparison among these models will show the role of the effective mass and symmetry energy in the neutrino-nucleon interaction in the nuclear medium.

The neutral-current weak and electromagnetic (EM) interactions for the neutrino-nucleon scattering in the nuclear medium can be described in terms of the following effective Lagrangian:

$$\mathcal{L}_{\text{int}}^N = \frac{G_F}{\sqrt{2}} (\bar{\nu}\Gamma_W^\mu\nu) (\bar{N}J_\mu^W N) + \frac{4\pi\alpha_{\text{EM}}}{q^2} (\bar{\nu}\Gamma_{\text{EM}}^\mu\nu) (\bar{N}J_\mu^{\text{EM}} N), \quad (4)$$

where the ν and $N = (n, p)$ denote the neutrino and nucleon fields, respectively. The weak

Target	G_A	F_1^W	F_2^W	F_1^{EM}	F_2^{EM}
n	$-\frac{g_A}{2}$	-0.5	$-\frac{1}{2}(\kappa_p - \kappa_n) - 2 \sin^2 \theta_w \kappa_n$	0	κ_n
p	$\frac{g_A}{2}$	$0.5 - 2 \sin^2 \theta_w$	$\frac{1}{2}(\kappa_p - \kappa_n) - 2 \sin^2 \theta_w \kappa_n$	1	κ_p

TABLE II: Vacuum form factor values at $q^2 = 0$. In the numerical calculation, we use $\sin^2 \theta_w = 0.231$, $g_A = 1.260$, $\kappa_p = 1.793$ and $\kappa_n = -1.913$.

and EM currents for the nucleon, J_μ^W and J_μ^{EM} are defined by

$$\begin{aligned}
J_\mu^W &= F_1^W(q^2)\gamma_\mu - G_A(q^2)\gamma_\mu\gamma^5 + iF_2^W(q^2)\frac{\sigma_{\mu\nu}q^\nu}{2M} + \frac{G_p(q^2)}{2M}q_\mu\gamma^5, \\
J_\mu^{\text{EM}} &= F_1^{\text{EM}}(q^2)\gamma_\mu + iF_2^{\text{EM}}(q^2)\frac{\sigma_{\mu\nu}q^\nu}{2M}.
\end{aligned} \tag{5}$$

The values of the nucleon form factors G_A and $F_{1,2}^W$ at $q^2 = 0$ in vacuum are summarized in Tab. II.

Since we are interested in the density effects of the neutrino-nucleon scattering, the nucleon form factors should be described as functions of density. For this purpose, the DDFFs are calculated in the quark-meson coupling (QMC) model [18]. In the QMC model, the standard values of the current quark mass $m_u = 5$ MeV and nucleon bag radius in free space $R_N = 0.8$ fm are used. For more details on the model and related calculations, refer to Ref. [18] and references therein.

The weak-interaction vertex of the Dirac neutrino in Eq. (4) can be written in the standard $V - A$ form as follows:

$$\Gamma_W^\mu = \gamma^\mu(1 - \gamma^5), \tag{6}$$

while the EM-interaction vertex is constructed in terms of the four independent form factors in general

$$\Gamma_{\text{EM}}^\mu = f_1\gamma^\mu - \frac{i}{2m_e}f_2\sigma^{\mu\nu}q_\nu + g_1\left(g^{\mu\nu} - \frac{q^\mu q^\nu}{q^2}\right)\gamma_\nu\gamma^5 - \frac{i}{2m_e}g_2\sigma^{\mu\nu}q_\nu\gamma^5. \tag{7}$$

Here, f_1 , g_1 , f_2 , and g_2 are called the Dirac, anapole, magnetic, and electric **dipole** form factors as functions of q^2 , respectively. The NCR is simply defined by

$$\langle R_\nu^2 \rangle \equiv \langle R_V^2 \rangle + \langle R_A^2 \rangle, \tag{8}$$

where R_V and R_A are the vector and axial-vector charge radii, which are defined by

$$\langle R_V^2 \rangle = 6 \frac{df_1(q^2)}{dq^2} \Big|_{q^2=0}, \quad \langle R_A^2 \rangle = 6 \frac{dg_1(q^2)}{dq^2} \Big|_{q^2=0}. \quad (9)$$

By doing this, we can explore the effects of the spatial extension of the neutrino in the medium. In the Breit frame with $q_0 = 0$, we can use the approximate relation:

$$f_1(q^2) \simeq -\frac{1}{6} \langle R_V^2 \rangle \mathbf{q}^2, \quad g_1(q^2) \simeq -\frac{1}{6} \langle R_A^2 \rangle \mathbf{q}^2. \quad (10)$$

At $q^2 = 0$, $f_2(q^2)$ and $g_2(q^2)$ define the NMM and charge-parity violating electric dipole moment as

$$\mu_\nu^m = f_2(0)\mu_B \quad \text{and} \quad \mu_\nu^e = g_2(0)\mu_B, \quad (11)$$

from which we can define the effective NMM as

$$\mu_\nu^2 \equiv (\mu_\nu^m)^2 + (\mu_\nu^e)^2 \quad (12)$$

with the Bohr magneton $\mu_B = \frac{e}{2m_e}$, where e and m_e is the electron unit charge and mass, respectively.

The DCS density for the neutrino-nucleon scattering in the weak and EM neutral-current interactions is given by

$$\begin{aligned} \frac{1}{V} \frac{d^3\sigma}{dE'_\nu d^2\Omega} &= -\frac{1}{16\pi^2} \frac{E'_\nu}{E_\nu} \left[\frac{G_F^2}{2} (L_\nu^{\alpha\beta} \Pi_{\alpha\beta}^{\text{Im}})_{\text{W}} + \frac{16\pi^2 \alpha_{\text{EM}}^2}{q^4} (L_\nu^{\alpha\beta} \Pi_{\alpha\beta}^{\text{Im}})_{\text{EM}} \right. \\ &\quad \left. + \frac{8\pi G_F \alpha_{\text{EM}}}{q^2 \sqrt{2}} (L_\nu^{\alpha\beta} \Pi_{\alpha\beta}^{\text{Im}})_{\text{INT}} \right] \end{aligned} \quad (13)$$

where E'_ν and E_ν are respectively the final and initial neutrino energies. The detailed forms of L_ν and Π^{Im} for the weak, EM, and interference terms are given in Refs. [8, 11, 18].

The inverse of the neutrino mean free path (NMFP) is determined by integrating the DCS in Eq. (13) over q_0 and $|\mathbf{q}|$, resulting in

$$\lambda^{-1}(E_\nu) = 2\pi \int_{q_0}^{(2E_\nu - q_0)} d|\mathbf{q}| \int_0^{2E_\nu} dq_0 \frac{|\mathbf{q}|}{E_\nu E'_\nu} \left[\frac{1}{V} \frac{d^3\sigma}{dE'_\nu d^2\Omega} \right]. \quad (14)$$

Here, $E'_\nu = E_\nu - q_0$.




FIG. 1: Normalized density-dependent weak form factors (DDFF) for the nucleon as functions of $Q^2 \leq 0.1 \text{ GeV}^2$ and $\rho/\rho_0 \leq 3$ from the QMC model [19]: (a) $G_A(Q^2, \rho)/G_A(0, 0)$, (b) $F_1^W(Q^2, \rho)/F_1^W(0, 0)$, and (c) $F_2^W(Q^2, \rho)/F_2^W(0, 0)$.

III. NUMERICAL RESULT AND DISCUSSION

In this Section, we present the numerical results with detailed discussions for the DDDFF, DCR, and NMFP.

A. Nucleon DDDFF

As mentioned already, since we are interested in the neutrino-nucleon scattering inside the nuclear medium, density effects should be taken into account carefully for the nucleon form factors. For this purpose, we employ the QMC model [19]. In Fig. 1, we depict the numerical results for the normalized density-dependent weak form factors for the nucleon as functions of Q^2 and ρ/ρ_0 , showing (a) $G_A(Q^2, \rho)/G_A(0, 0)$, (b) $F_1^W(Q^2, \rho)/F_1^W(0, 0)$, and (c) $F_2^W(Q^2, \rho)/F_2^W(0, 0)$ for the proton. Note that the F_1^W is almost independent of the density in the low Q^2 region, in which we are interested mostly, whereas the G_A decreases with respect to the density, as shown in panel (a) of Fig. 1. This decreasing behavior of the G_A can be understood by the lower component of the quark spinor $\mathcal{L}(r) \propto \mathcal{O}(1/M_q^*)$ is enhanced more than its upper component $\mathcal{U}(r) \propto \mathcal{O}(1)$ as functions of the density when it is calculated using the three-point quark operator [19, 37]: $G_A \equiv \int d^3r [\mathcal{U}^2(r) - \mathcal{L}^2(r)] > 1$. On the contrary, the F_2^W increases with respect to the density, and this tendency is originated from $F_2^W \equiv \int d^3r \mathcal{U}(r)\mathcal{L}(r)$. From the numerical calculations, we verified that the contributions from the G_A are the most dominant to describe the scattering cross-section. Hence, with the DDDFF, the scattering cross-section will be reduced with respect to the density in comparison




FIG. 2: Triple isotropic differential cross-section (DCS) of the neutrino-nucleon scattering as a function of the transferred energy q_0 for $\rho = (1.0 - 3.0) \rho_0$ for the (a) KIDS0, (b) KIDS-A, (c) KIDS0-m*87, and (d) SLy4 models. The numerical results are given with the DCS+VFF (dotted), DCS+DDFF (dashed), DCS+DDFF+NMM (solid), DCS+DDFF+NCR1 (long-dashed), DCS+DDFF+NCR2 (dot-dashed), and DCS+DDFF+NCR3 (dot-long-dashed). Neutrino charge radii are $R_\nu = 3.5, 1.28$ and 2.48 in units of 10^{-5} MeV^{-1} for NCR1, NCR2 and NCR3, respectively.

to that without it. It is worth mentioning that the result for the G_A at saturation density, for instance, turns out to be consistent with other theoretical calculations [36–38]. Although we do not show the EM form factors here, we verify that their density dependencies are not significant in the cross-section.

B. Neutrino-nucleon scattering DCS

The temperature of the proto-neutron star that is formed after the bounce of material in the supernova explosion is the order of 10 MeV. It falls down quickly by the deleptonization process and reaches a few MeV in a few minutes. The energy of the neutrino in the β -

equilibrium is at the order of thermal fluctuation, so we choose $E_\nu = 5$ MeV for the neutrinos in the NS throughout the calculation.

The numerical results for the DCS are depicted as a function of q_0 for the densities $\rho = \rho_0$ (left), $2\rho_0$ (middle), and $3\rho_0$ (right) in Fig. 2. In each row, we compute the DCS using different EDF models, i.e., KIDS0, KIDS-A, KIDS0-m*87, and SLy4 from top to bottom. In order to explore the effects of the new ingredients DDFF, NMM, and NCR, we separately show the results with the vacuum form factor (VFF), DDFF, DDFF+NMM, and DDFF+NCR. Note that the VFF indicates the DDFF at $\rho = 0$.

First, we explain the overall tendency depending on the different EDF models. It is obvious that the DCS from KIDS0 and KIDS-A are qualitatively larger than those from KIDS0-m*87 and SLy4, although quantitative differences are still shown. As discussed in detail in Ref. [1], the larger DCS is originated from the effective neutron mass following the condition $M_n^* \gtrsim M_n$, where M_n is the neutron mass in a vacuum, as in KIDS0 and KIDS-A. This observation can be basically explained by that the DCS is proportional to the $\mathcal{O}(M_N^*)$ and $\mathcal{O}(M_N^{*2})$ terms from the spin summation over the nucleonic tensor given in Eq. (13). Physically, the increasing nucleon mass in the scattering process results in decreasing energy transfer Δq_0 in the t channel at a certain \sqrt{s} value. In turn, the interaction time increases, resulting in a larger cross-section from $\Delta q_0 \Delta \tau \gtrsim \hbar/s$, where the τ stands for the interaction time. The difference between the models becomes more obvious as the density increases as also reported previously in Ref. [1]. The different endpoints of each cross-section are determined by the effective nucleon masses depending on the models.

As for the effects of the DDFF, it turns out that the inclusion of the DDFF provides a non-negligible reduction of the cross-section in comparison to that of VFF. Moreover, the reduction becomes more significant as the density increases. If we compare the maximum values with the DCS+DDFF with those of the DCS+VFF, the reduction rates are 0.83, 0.77, and 0.71 at $\rho = \rho_0$, $2\rho_0$, and $3\rho_0$, respectively, and it is weakly dependent on the EDF models as shown in the figure. Hence, the decreasing behavior of the DCS can be understood in terms of the in-medium behavior of the DDFF, especially G_A , as already discussed in the previous subsection.

Now we are in a position to discuss the effects of NMM in the DCS. Here, we used $\mu_\nu = 2.9 \times 10^{-11} \mu_B$ for the numerical calculations. This value is constrained from the experiment [26] and close to the astronomical observation [39]. By comparing the DCS+DDFF

(dashed) and DCS+DDFF+NMM (solid) in Fig. 2, it is obvious that the NMM gives negligible contributions to the scattering process. Note that the neutrino magnetic tensor currents including the NMM are proportional to q_0 as shown in Eq. (7). Hence, the effects of the NMM in the DCS are considerably suppressed in the low q_0 region as shown in Fig. 2, in addition to its extremely small value $\propto 10^{-11} \mu_B$, in comparison to other scales such as μ_n and μ_p in the scattering process.

The NCR relates to the neutrino electric vector current as in Eq. (7) and indicates the EM structure of the particle, although the neutrino has been believed to be a point-like particle in general. In the LAMPF experiment for the measurement $\nu_e e^- \rightarrow \nu_e e^-$ [29], the NCR was estimated by $R_\nu = 3.5 \times 10^{-5} \text{ MeV}^{-1}$. In Ref. [30] from the new analyses based on the COHERENT elastic neutrino-nucleus scattering, the NCR is given by the upper limit $R_\nu = 1.28 \times 10^{-5} \text{ MeV}^{-1}$. To explore the tendency of the NCR effect, we also try a middle value $R_\nu = 2.48 \times 10^{-5} \text{ MeV}^{-1}$. For convenience, we assigned these three NCR values the acronyms NCR1, NCR2, and NCR3, respectively.

Using these values in the numerical calculations, the results are given in the long-dashed (NCR1), dot-dashed (NCR2), and dot-long-dashed (NCR3) lines in the figure. Very interestingly, all DCSs increase sizably with the NCR1 [29] and NCR3 (between those of Ref. [29] and [30]) commonly for the different EDF models, while the NCR2 [30] only makes negligible changes. This observation can be easily understood by that the effects of the NCR2 are about 4 – 8 times smaller than NCR1 and NCR3 in the cross-section qualitatively. The effects of the NCR are especially profound for the SLy4 model. We also note that the increase of the DCS due to the NCR is less sensitive to the density than the DDFF since the neutrino EM current is not dependent on the density. The drastic changes observed in the DCS due to the NCR can be understood physically by the spatial extension of the neutrino wave function, increasing the overlap with that for the nucleon, resulting in a rapidly growing DCS.

C. NMFP

Finally, we present the numerical results for the NMFP in Fig. 3 in the same manner as that for the DCS given above. The NMFP is one of the critical quantities that affect the cooling rate of the NS. It is proportional by construction to the inverse of the DCS, so




FIG. 3: Neutrino mean free path (NMFP) as a function of density for the (a) KIDS0, (b) KIDS-A, (c) KIDS0-m*87, and (d) SLy4 modes. The numerical results are given with the vacuum form factor NMFP+VFF (dotted), NMFP+DDFF (dashed), NMFP+DDFF+NMM (solid), NMFP+DDFF+NCR1 (long-dashed), NMFP+DDFF+NCR2 (dot-dashed), and NMFP+DDFF+NCR3 (dot-long-dashed). Model dependence is compared with NMFP+DDFF+NCR1 in panel (e). The shaded area stands for the region $R_{NS} = 13$ km, an upper bound for the radius of $1.4M_{\odot}$ mass NS.

one can easily expect that shorter NMFP at a certain density for $M_n^* \gtrsim M_n$ in the KIDS0 and KIDS-A models in comparison to the others, as shown in Fig. 3. In common for all the cases, the NMFP is a decreasing function of the density, except for the NMFP+DDFF, NMFP+DDFF+NMM, and NMFP+DDFF+NCR2 in the SLy4 model, being consistent with the consideration that the neutrino-nucleon interaction rate increases in a dense matter.

In the low-density region, where $VFF \sim DDFF$, the NMFP+VFF, and NMFP+DDFF are close to each other as expected, and the difference gets larger as the density grows for all the cases. Similar to the DCS, the effects of the NCR turn out to be critical throughout the densities with the NCR1 and NCR3 values, decreasing the NMFP by 3 – 5 km on average. Again, the NNM does not make any significant contributions to the NMFP, being consistent with the DCS results. Taking into account that the radius of the NS whose mass is $1.4M_\odot$ is $R_{1.4M_\odot} \lesssim 13$ km [40], indicated by the horizontal shaded line, as for the results from the KIDS0 and KIDS-A models, the neutrino can not escape freely from the NS if it is emitted at the center of NS where the density is beyond $\rho \approx 1.5\rho_0$. In contrast, the neutrino seldom experiences weak and EM interactions until it escapes from the NS for the KIDS0-m*87 and SLy4 models. For instance, there is almost no delay of the neutrino emission for $R_{NS} \approx 10$ km because NMFP is always larger than R_{NS} .

Interestingly, the NMFP+DDFF for the SLy4 model shows an increasing curve beyond $\rho \approx \rho_0$, being different from others. That obvious difference shown in the SLy4 model is caused by the smallest neutron-effective mass among the EDF models [1] in addition to the considerable DDFF effects, which make the DCS reduced. From this observation, the importance of the effective mass is crucial in analyzing the cooling processes of the NS via the neutrinos. Since the range of the NMFP predicted by the different EDF models is very wide, the application of the present results to the calculation of the thermal evolution of the NS should be necessarily followed up.

Here is a discussion on the effect of symmetry energy. A comparison of the KIDS0 and KIDS-A models clearly shows the role of the symmetry energy because the two models differ evidently in the stiffness of symmetry energy but have similar effective masses. In the result of both DCS and NMFP, the two models show similar behavior up to $3\rho_0$, so the stiffness of the symmetry energy appears to be insignificant. Symmetry energy directly affects the particle fraction, giving a larger proton fraction with stiffer symmetry energy. A large proton fraction can ignite the direct URCA process which leads to super fast cooling of the NS. In

addition, it's been shown in [1] that the particle fraction shows sizable dependence on the symmetry energy at densities higher than $3\rho_0$. Therefore, the effect of the symmetry energy could be probed correctly when the NS cooling is considered explicitly.

IV. SUMMARY AND CONCLUSION

In the present work, we have investigated the neutral-current neutrino-nucleon scattering in the nuclear medium using the quark-meson coupling (QMC) model together with the four different energy-density functional (EDF) models, i.e., KIDS0, KIDS-A, KIDS0-m*87, and SLy4. The nucleon density-dependent form factor (DDFF), differential cross-section (DCS), and neutrino mean free path (NMFP) were computed numerically at various densities. In addition, we also explored the effects of the finite neutrino magnetic moment (NMM) and its EM size via the charge radius (NCR). Below, we list relevant observations found in the present work:

- Among the weak DDFFs, G_A is a decreasing function of the density, and vice versa for F_2^W , whereas F_1^W is almost insensitive to the density in the small Q^2 region. These opposite behaviors between G_A and F_2^W can be understood by the different combinations of the density-dependent lower and upper components of the quark spinor in the QMC model. We also find out that the lower part is more sensitive to density and increases with respect to it. The density dependencies in the EM DDFFs turn out to be weak.
- The DCS increases with respect to the density in general and is larger in the KIDS0 and KIDS-A models which have $M_n^* \gtrsim M_n$ in comparison to other EDF models. The dominant contribution among the weak DDFFs turns out to be G_A , which reduces the DCS with respect to the density as understood by the above discussions. The effect of the NMM is almost negligible since it is highly suppressed in the small q_0 region. The finite NCR indicates a larger overlap with the nucleon wave function, resulting in the increase of the DCS in general. The range of increase is, however, strongly dependent on the the magnitude of NCR.
- The NMFP which is inversely proportional to the DCS is scrutinized in the same manner as the DCS. The weak DDFF G_A makes the NMFP increase as understood by

its density dependence. The inclusion of the NCR can drastically decrease the NMFP by about 5 km when $R_\nu = 3.5 \times 10^{-5} \text{ MeV}^{-1}$. If we take $R_{\text{NS}} \approx 13 \text{ km}$ for instance in the present theoretical framework, the neutrino escapes from the NS almost without interactions up to $\rho \approx 3\rho_0$ for $M_n^* \lesssim M_n$. On the contrary for $M_n^* \gtrsim M_n$, DCS increases as density increases, and it leads to NMFP shorter than the NS radius. A decrease in NMFP implies that the interaction of the neutrino with NS matter becomes more probable, and it can impose a non-negligible effect on the thermal evolution of the NS.

As discussed previously, there are considerable theoretical uncertainties in the present theoretical framework, depending on the EDF models, density-dependent form factors, and neutrino properties. Differential cross sections and neutrino mean free paths are highly sensitive to the magnitude of NCR, and the effect of NCR is closely coupled to the nuclear physics input, i.e. the effective mass of the nucleon. Hence, it is necessary to investigate the NS thermal evolution to construct a more realistic theoretical model and determine model parameters consistently. Coherent elastic neutrino-nucleus scattering ($\text{CE}\nu\text{NS}$) is expected to provide brand-new constraints to the beyond-standard-model physics that includes NCR and NMM from the laboratory experiment. Recently bounds on NCR and NMM are extracted from the COHERENT data [41]. Uncertainty diagnosis in $\text{CE}\nu\text{NS}$ can open a way to obtain more stringent constraints on the relevant quantities and more realistic physical results from the experiment. The effect of effective mass and NCR to $\text{CE}\nu\text{NS}$ cross-section is being investigated. A combination of various independent phenomena such as NS cooling, $\text{CE}\nu\text{NS}$, quasielastic and deep inelastic scatterings will help reduce the range of uncertainty. Related works will appear elsewhere.

Acknowledgments

This work was supported by the National Research Foundation of Korea (NRF) Grants No. 2018R1A5A1025563, No. 2020R1F1A1052495, and No. 2022R1A2C1003964.

-
- [1] P. T. P. Hutaaruk, H. Gil, S. i. Nam and C. H. Hyun, *Phys. Rev. C* **106**, no.3, 035802 (2022).
 - [2] G. A. Miller, E. Piassetzky and G. Ron, *Phys. Rev. Lett.* **101**, 082002 (2008).
 - [3] C. E. Hyde and K. de Jager, *Ann. Rev. Nucl. Part. Sci.* **54**, 217-267 (2004).

- [4] L. Andivahis *et al.*, Phys. Rev. D **50**, 5491-5517 (1994).
- [5] J. Litt *et al.*, Phys. Lett. B **31**, 40-44 (1970).
- [6] B. Mueller *et al.* [SAMPLE], Phys. Rev. Lett. **78**, 3824-3827 (1997).
- [7] C. J. Horowitz and M. A. Perez-Garcia, Phys. Rev. C **68**, 025803 (2003).
- [8] S. Reddy, M. Prakash, J. M. Lattimer and J. A. Pons, Phys. Rev. C **59**, 2888-2918 (1999).
- [9] A. Sulaksono, P. T. P. Hutaauruk and T. Mart, Phys. Rev. C **72**, 065801 (2005).
- [10] G. Guo, G. Martínez-Pinedo, A. Lohs and T. Fischer, Phys. Rev. D **102**, no.2, 023037 (2020).
- [11] A. Sulaksono *et al.*, Phys. Rev. C **73**, 025803 (2006).
- [12] P. T. P. Hutaauruk, A. Sulaksono and T. Mart, Nucl. Phys. A **782**, 400-405 (2007).
- [13] I. C. Cloet, G. A. Miller, E. Piassetzky and G. Ron, Phys. Rev. Lett. **103**, 082301 (2009).
- [14] D. H. Lu, K. Tsushima *et al.*, Phys. Rev. C **60**, 068201 (1999).
- [15] D. F. Geesaman, K. Saito and A. W. Thomas, Ann. Rev. Nucl. Part. Sci. **45**, 337-390 (1995).
- [16] J. J. Aubert *et al.* [European Muon], Phys. Lett. B **123**, 275-278 (1983).
- [17] S. Malace *et al.* [Jefferson Lab Hall A], AIP Conf. Proc. **1056**, no.1, 141-147 (2008).
- [18] P. T. P. Hutaauruk, Y. Oh and K. Tsushima, Phys. Rev. D **98**, no.1, 013009 (2018).
- [19] K. Saito, K. Tsushima and A. W. Thomas, Prog. Part. Nucl. Phys. **58**, 1-167 (2007).
- [20] P. T. P. Hutaauruk, A. Sulaksono and K. Tsushima, Nucl. Phys. A **1017**, 122356 (2022).
- [21] C.Y. Ryu, C.H. Hyun, T.-S. Park, and S.W. Hong, Phys. Lett. B **674**, 122 (2009).
- [22] C.Y. Ryu, C.H. Hyun, and M.-K. Cheoun, J. Phys. G **37**, 105002 (2010).
- [23] D. W. Liu *et al.* [Super-Kamiokande], Phys. Rev. Lett. **93**, 021802 (2004).
- [24] H. B. Li *et al.* [TEXONO], Phys. Rev. Lett. **90**, 131802 (2003).
- [25] Z. Daraktchieva *et al.* [MUNU], Phys. Lett. B **564**, 190-198 (2003).
- [26] A. G. Beda *et al.*, Phys. Part. Nucl. Lett. **10**, 139-143 (2013).
- [27] E. Aprile *et al.* [XENON], Phys. Rev. D **102**, no.7, 072004 (2020).
- [28] M. Agostini *et al.* [Borexino], Phys. Rev. D **96**, no.9, 091103 (2017).
- [29] R. C. Allen *et al.*, Phys. Rev. D **47**, 11-28 (1993).
- [30] M. Cadeddu, C. Giunti, K. A. Kouzakov, Y. F. Li, Y. Y. Zhang and A. I. Studenikin, Phys. Rev. D **98**, no.11, 113010 (2018) [erratum: Phys. Rev. D **101**, no.5, 059902 (2020)].
- [31] H. Bonet *et al.* [CONUS], [arXiv:2201.12257 [hep-ex]].
- [32] S. Jana, Y. P. Porto-Silva and M. Sen, [arXiv:2203.01950 [hep-ph]].
- [33] V. Lyubushkin *et al.* [NOMAD], Eur. Phys. J. C **63**, 355-381 (2009).

- [34] A. A. Aguilar-Arevalo *et al.* [MiniBooNE], Phys. Rev. D **82**, 092005 (2010).
- [35] G. A. Fiorentini *et al.* [MINERvA], Phys. Rev. Lett. **111**, 022502 (2013).
- [36] P. T. P. Hutaeruk, Y. Oh and K. Tsushima, Phys. Rev. C **99**, no.1, 015202 (2019).
- [37] D. H. Lu, A. W. Thomas and K. Tsushima, [arXiv:nucl-th/0112001 [nucl-th]].
- [38] A. M. Rakhimov *et al.*, Nucl. Phys. A **643**, 383-401 (1998).
- [39] G. G. Raffelt, Phys. Rept. **320**, 319-327 (1999).
- [40] M. C. Miller *et al.*, Astrophys. J. **918**, L28 (2021).
- [41] M. A. Corona *et al.*, JHEP **09**, 164 (2022).