

# UV asymptotics of $n$ -point correlators of twist-2 operators in $SU(N)$ Yang-Mills theory

---

Marco Bochicchio<sup>a</sup> Mauro Papinutto<sup>b,a</sup> Francesco Scardino<sup>b,a</sup>

<sup>a</sup>*Physics Department, INFN Roma1,  
Piazzale A. Moro 2, Roma, I-00185, Italy*

<sup>b</sup>*Physics Department, Sapienza University,  
Piazzale A. Moro 2, Roma, I-00185, Italy*

*E-mail:* [marco.bochicchio@roma1.infn.it](mailto:marco.bochicchio@roma1.infn.it), [mauro.papinutto@roma1.infn.it](mailto:mauro.papinutto@roma1.infn.it),  
[francesco.scardino@roma1.infn.it](mailto:francesco.scardino@roma1.infn.it)

ABSTRACT: Extending our previous work [1], we compute – directly from its path-integral definition as a Gaussian integral – the generating functional of the conformal correlators to the lowest perturbative order of all the twist-2 operators with maximal spin along the  $p_+$  direction in  $SU(N)$  Yang-Mills theory, both in Minkowskian and Euclidean space-time. We thus provide a purely perturbative explanation as to why the aforementioned generating functional has the structure of the logarithm of a functional determinant. Besides, by means of a careful choice of the renormalization scheme that reduces the mixing of the above operators to the multiplicatively renormalizable case to all orders of perturbation theory, we lift the generating functional of the Euclidean conformal correlators to the generating functional of the corresponding renormalization-group (RG) improved correlators that inherits the very same structure of the logarithm of a functional determinant. For large  $N$  the RG-improved generating functional admits a decomposition into planar and leading-order (LO) nonplanar contributions. Remarkably, according to the prediction in [2] the RG-improved LO nonplanar generating functional – being asymptotic in the ultraviolet (UV) to the sum of the glueball one-loop diagrams – admits the structure of the logarithm of a functional determinant as well. Hence, the computation above sets strong UV asymptotic constraints on the nonperturbative solution of large- $N$  YM theory and it may be a pivotal guide for the search of such a solution.

---

## Contents

<b>1</b>	<b>Introduction, physics motivations and conclusions</b>	<b>1</b>
<b>2</b>	<b>Plan of the paper</b>	<b>3</b>
<b>3</b>	<b>Generating functional <math>\mathcal{W}_{conf}</math> of the Minkowskian conformal correlators from the YM path integral</b>	<b>4</b>
3.1	Standard basis	5
<b>4</b>	<b>Connection of <math>\mathcal{W}_{conf}</math> with the generating functionals <math>\Gamma_{conf}</math> in [1]</b>	<b>8</b>
4.1	An example of the dictionary	10
4.2	Dictionary in the general case	12
4.3	Generating functional $\Gamma_{conf}$	13
<b>5</b>	<b>Generating functional in the momentum representation</b>	<b>15</b>
5.1	Standard basis	15
<b>6</b>	<b>Mixed 3- and 4-point conformal correlators</b>	<b>18</b>
6.1	3-point correlators in Minkowskian space-time	18
6.2	4-point correlators in Minkowskian space-time	20
6.3	4-point correlators in Euclidean space-time	24
<b>7</b>	<b>UV asymptotics of Euclidean <math>n</math>-point correlators</b>	<b>25</b>
7.1	Operator mixing	25
7.2	Canonical nonresonant diagonal renormalization scheme	26
7.3	Anomalous dimensions of twist-2 operators	28
<b>8</b>	<b>Generating functional of Euclidean RG-improved correlators</b>	<b>29</b>
<b>A</b>	<b>Determinants of block matrices</b>	<b>31</b>
<b>B</b>	<b>Conformal properties of the standard basis</b>	<b>31</b>

---

## 1 Introduction, physics motivations and conclusions

Recently, we have computed [1] to the lowest perturbative order in  $SU(N)$  Yang-Mills (YM) theory  $n$ -point conformal correlators in the coordinate,  $G_{conf}^{(n)}(x_1, \dots, x_n)$ , and momentum,  $G_{conf}^{(n)}(p_1, \dots, p_n)$ , representation of the gauge-invariant twist-2 operators with maximal spin along the  $p_+$  direction, both in Minkowskian and – by analytic continuation – Euclidean space-time.

Specifically, we have calculated the  $n$ -point conformal correlators of the balanced and unbalanced<sup>1</sup> operators with collinear twist 2 [1] in their separate sectors and the 3-point correlators in the mixed sector as well.

We have also reconstructed [1] from the  $n$ -point conformal correlators in each separate sector the corresponding generating functional, both in the coordinate and momentum representation, that turns out to have the structure of the logarithm of a functional determinant.

We have pointed out [1] that the rationale behind the generating functional being the logarithm of a functional determinant is the prediction [2] that, by fundamental principles of the 't Hooft large- $N$  expansion in  $SU(N)$  YM theory, the leading-order (LO) nonplanar contribution to the nonperturbative large- $N$  YM effective action should have the structure of the logarithm of a functional determinant that sums the glueball one-loop diagrams. In relation to the above computations, the aim of the present paper is fourfold.

First, as opposed to the aforementioned nonperturbative argument, since the computation [1] of the  $n$ -point conformal correlators is of purely perturbative nature, we find a perturbative reason as to why the corresponding generating functional is the logarithm of a functional determinant: As all the aforementioned collinear twist-2 operators are quadratic in the gauge field in the light-cone gauge to the lowest order of perturbation theory, the functional integral that defines their generating functional is Gaussian and leads to a functional determinant involving the corresponding sources.

Then, once some technical difficulties are properly understood, a tedious but straightforward computation shows that the functional determinants worked out in [1] actually arise from their very definition as Gaussian functional integrals in the present paper.

Second, our new method to compute the generating functional allows us to produce a vast generalization of our previous computations. For example, we calculate the generating functional in the mixed balanced/unbalanced sector in  $SU(N)$  YM theory, verify by means of it our previous computation of the generating functionals in the separate sectors and of the mixed 3-point correlators [1] and compute the mixed 4-point correlators.

Besides, the same technique will allow us to compute [5] – as the logarithm of a functional (super-)determinant – the generating functional of the lowest-order conformal correlators of the collinear twist-2 operators in  $SU(N)$  QCD,  $\mathcal{N} = 1$  SUSY YM theory,  $\mathcal{N} = 1$  SUSY QCD and  $\mathcal{N} = 2, 4$  SUSY YM theory – the super-determinant arising in the super-symmetric cases.

Remarkably, the very same technique also applies [5] to the higher-twist operators that are quadratic [3] in the elementary fields in the light-cone gauge to the lowest order of perturbation theory in all the above theories.

Third, we work out the ultraviolet (UV) asymptotics of the Euclidean nonperturbative  $n$ -point correlators in a certain renormalization scheme described below. The computation is based both on our lowest-order calculation of the Euclidean conformal correlators and its improvement by means of the renormalization group (RG).

---

<sup>1</sup>In our terminology 'unbalanced' and 'balanced' refers to either the different or the equal number of dotted and undotted indices that the aforementioned operators respectively possess in the spinorial representation. Unbalanced operators are referred to as 'asymmetric' in [3] and 'anisotropic' in [4].

Generally, collinear twist-2 operators  $\mathcal{O}_{s_i}$  with maximal spin projection  $s_i \geq 2$  mix under renormalization with total derivatives of the same operators with lower spins, so that an explicit evaluation of the RG-improved asymptotics of their Euclidean  $n$ -point correlators in the  $\overline{MS}$  renormalization scheme would require the asymptotic estimate – that seems practically impossible – of the product of the sum of  $O(s_i)$  terms for each  $\mathcal{O}_{s_i}$  occurring in the  $n$ -point correlators because of the triangular nature of their renormalized mixing matrix  $Z(\lambda)$  in the  $\overline{MS}$  scheme, not to mention that in general  $Z(\lambda)$  is actually only known to two loops [6–8] in perturbation theory and not in its RG-improved form.

Recently, a differential-geometric approach to operator mixing in massless QCD-like theories has been proposed [9] precisely to overcome the above difficulty. Specifically, it has been determined under which conditions a renormalization scheme exists where  $Z(\lambda)$  may be set in a diagonal canonical form that is one-loop exact to all perturbative orders – according to the nonresonant diagonalizable  $\frac{\gamma_0}{\beta_0}$  case (I) [9]. Moreover, it has been verified [10] that the balanced collinear twist-2 operators belong to the case (I) and we show in the present paper that this is also the case for the unbalanced ones <sup>2</sup>.

Remarkably, in such a scheme the needed terms for the asymptotic evaluation of the  $n$ -point correlators reduce to just one, since in this scheme the operators are multiplicatively renormalizable and in general their correlators do not vanish to the lowest order of perturbation theory. The nonresonant diagonal scheme may also apply [5] to the collinear twist-2 operators in QCD and its supersymmetric extensions above.

Fourth, we employ the UV asymptotics of the Euclidean  $n$ -point correlators in the nonresonant diagonal scheme to work out the corresponding generating functional that also turns out to be the logarithm of a functional determinant. In fact, after a suitable rescaling of the operators, it may be decomposed for large  $N$  into the sum of the generating functionals of the planar and LO nonplanar asymptotic correlators, which has the structure of the logarithm of a functional determinant as well.

This is our last – and perhaps most remarkable – step. Indeed, as already pointed out [1] according to the prediction in [2], the generating functional of the LO nonplanar asymptotic correlators must be asymptotic in the UV to the corresponding nonperturbative glueball one-loop generating functional that is therefore asymptotically known by means of our computation.

In a forthcoming paper we will further investigate the above nonperturbative interpretation. In any case the generating functionals of planar and LO nonplanar asymptotic correlators set strong UV asymptotic constraints on the nonperturbative solution of large- $N$  YM theory and may be a pivotal guide for the search of such a solution.

## 2 Plan of the paper

In section 3 we compute the generating functional in the coordinate representation of all the correlators of collinear twist-2 operators (appendix B) to the lowest order of perturbation theory from the YM path integral in Minkowskian space-time.

---

<sup>2</sup>In both cases numerically up to  $s = 10^4$ .

In section 4 we connect the aforementioned path-integral computation of the generating functional – that includes the mixed balanced/unbalanced sector – with the generating functionals in the separate balanced and unbalanced sectors [1] finding perfect agreement.

In section 5 we work out the generating functional in the momentum representation.

In section 6 we calculate the 3- and 4-point correlators in the mixed sector from the generating functional. The result agrees with our previous computation [1] of the mixed 3-point correlators.

In section 7 we recall some results on operator mixing and apply them to work out the UV asymptotics in the coordinate representation of Euclidean  $n$ -point correlators of collinear twist-2 operators in the canonical nonresonant diagonal scheme.

In section 8 we compute the generating functional in the coordinate representation of the RG-improved correlators in Euclidean space-time.

In appendix A we compute the functional determinant that leads to the generating functional.

In appendix B we recall the definition of the standard basis [1] of collinear twist-2 operators and describe their conformal properties to the leading and next-to-leading perturbative order.

### 3 Generating functional $\mathcal{W}_{conf}$ of the Minkowskian conformal correlators from the YM path integral

The Minkowskian YM action is:

$$S_{YM} = -\frac{1}{2} \int d^4x \operatorname{Tr} F_{\mu\nu} F^{\mu\nu} \quad (3.1)$$

where:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + i \frac{g}{\sqrt{N}} [A_\mu, A_\nu] \quad (3.2)$$

and:

$$A_\mu = A_\mu^a T^a \quad (3.3)$$

with the Hermitian generators of the  $SU(N)$  Lie algebra:

$$[T^a, T^b] = i f^{abc} T^c \quad (3.4)$$

in the fundamental representation normalized as:

$$\operatorname{Tr}(T^a T^b) = \frac{\delta^{ab}}{2} \quad (3.5)$$

and  $g^2 = N g_{YM}^2$  the 't Hooft coupling [11]. We set:

$$\begin{aligned} V_+ &= \frac{V_0 + V_3}{\sqrt{2}} & V_- &= \frac{V_0 - V_3}{\sqrt{2}} \\ V &= \frac{V_1 + iV_2}{\sqrt{2}} & \bar{V} &= \frac{V_1 - iV_2}{\sqrt{2}} \end{aligned} \quad (3.6)$$

for any vector  $V_\mu$ . We choose the light-cone gauge:

$$A_+ = 0 \quad (3.7)$$

After integrating out  $A_-$ , the YM action in the light-cone gauge reads [12]:

$$\begin{aligned} S_{YM}(A, \bar{A}) = & - \int \bar{A}^a \square A^a + 2 \frac{g}{\sqrt{N}} f^{abc} (A^a \partial_+ \bar{A}^b \bar{\partial}_+^{-1} A^c + \bar{A}^a \partial_+ A^b \partial_+^{-1} \bar{A}^c) \\ & + 2 \frac{g^2}{N} f^{abc} f^{ade} \partial_+^{-1} (A^b \partial_+ \bar{A}^c) \partial_+^{-1} (\bar{A}^d \partial_+ A^e) d^4x \end{aligned} \quad (3.8)$$

The v.e.v. of a product of local gauge-invariant operators  $\mathcal{O}_i(A)$  that do not depend on  $A_-$  reads:

$$\langle \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \rangle = \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} e^{iS_{YM}(A, \bar{A})} \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \quad (3.9)$$

To the leading perturbative order it reduces to:

$$\langle \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \rangle = \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} e^{-i \int d^4x \bar{A}^a \square A^a} \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \quad (3.10)$$

with:

$$\square = g^{\mu\nu} \partial_\mu \partial_\nu = \partial_0^2 - \sum_{i=1}^3 \partial_i^2 \quad (3.11)$$

where we employ the mostly minus metric  $g^{\mu\nu}$  in Minkowskian space-time [1]. The corresponding generating functional reads to the leading order:

$$\mathcal{Z}_{conf}[J\mathcal{O}] = \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} e^{-i \int d^4x \bar{A}^a \square A^a} \exp \left( \int d^4x \sum_i J_{\mathcal{O}_i} \mathcal{O}_i \right) \quad (3.12)$$

### 3.1 Standard basis

The collinear twist-2 operators in the standard basis [1] are (appendix B):

$$\begin{aligned} \mathbb{O}_s &= \frac{1}{2} \bar{A}^a(x) \overleftarrow{\partial}_+ (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \overrightarrow{\partial}_+ A^a(x) \\ &= \frac{1}{2} \bar{A}^a(x) \mathcal{Y}_{s-2}^{\frac{5}{2}}(\overrightarrow{\partial}_+, \overleftarrow{\partial}_+) A^a(x) \quad s = 2, 4, 6, \dots \\ \tilde{\mathbb{O}}_s &= \frac{1}{2} \bar{A}^a(x) \overleftarrow{\partial}_+ (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \overrightarrow{\partial}_+ A^a(x) \\ &= \frac{1}{2} \bar{A}^a(x) \mathcal{H}_{s-2}^{\frac{5}{2}}(\overrightarrow{\partial}_+, \overleftarrow{\partial}_+) A^a(x) \quad s = 3, 5, 7, \dots \\ \mathbb{S}_s &= \frac{1}{2\sqrt{2}} \bar{A}^a(x) \overleftarrow{\partial}_+ (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \overrightarrow{\partial}_+ \bar{A}^a(x) \\ &= \frac{1}{2\sqrt{2}} \bar{A}^a(x) \mathcal{Y}_{s-2}^{\frac{5}{2}}(\overrightarrow{\partial}_+, \overleftarrow{\partial}_+) \bar{A}^a(x) \quad s = 2, 4, 6, \dots \\ \bar{\mathbb{S}}_s &= \frac{1}{2\sqrt{2}} A^a(x) \overleftarrow{\partial}_+ (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \overrightarrow{\partial}_+ A^a(x) \\ &= \frac{1}{2\sqrt{2}} A^a(x) \mathcal{Y}_{s-2}^{\frac{5}{2}}(\overrightarrow{\partial}_+, \overleftarrow{\partial}_+) A^a(x) \quad s = 2, 4, 6, \dots \end{aligned} \quad (3.13)$$

where the sum over repeated color indices is understood, and [1]:

$$\begin{aligned}\mathcal{Y}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &= \overleftarrow{\partial}_+(i\vec{\partial}_+ + i\overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \vec{\partial}_+ \\ &= \frac{\Gamma(3)\Gamma(s+3)}{\Gamma(5)\Gamma(s+1)} i^{s-2} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-1)^{s-k} \overleftarrow{\partial}_+^{s-k-1} \vec{\partial}_+^{k+1}\end{aligned}\quad (3.14)$$

for even spin, and:

$$\begin{aligned}\mathcal{H}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &= \overleftarrow{\partial}_+(i\vec{\partial}_+ + i\overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \vec{\partial}_+ \\ &= \frac{\Gamma(3)\Gamma(s+3)}{\Gamma(5)\Gamma(s+1)} i^{s-2} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-1)^{s-k} \overleftarrow{\partial}_+^{s-k-1} \vec{\partial}_+^{k+1}\end{aligned}\quad (3.15)$$

for odd spin. The corresponding generating functional of conformal correlators to the leading order reads:

$$\begin{aligned}\mathcal{Z}_{conf}[J_{\mathbb{O}}, J_{\tilde{\mathbb{O}}}, J_{\mathbb{S}}, J_{\tilde{\mathbb{S}}}] \\ = \frac{1}{Z} \int \mathcal{D}AD\bar{A} e^{-i \int d^4x \bar{A}^a \square A^a} \exp \left( \int d^4x \sum_s J_{\mathbb{O}_s} \mathbb{O}_s + J_{\tilde{\mathbb{O}}_s} \tilde{\mathbb{O}}_s + J_{\mathbb{S}_s} \mathbb{S}_s + J_{\tilde{\mathbb{S}}_s} \tilde{\mathbb{S}}_s \right)\end{aligned}\quad (3.16)$$

where the currents  $J_{\mathbb{O}_s}, J_{\tilde{\mathbb{O}}_s}, J_{\mathbb{S}_s}, J_{\tilde{\mathbb{S}}_s}$  are defined to be zero for  $s$  different from the spin of the corresponding operator. Explicitly:

$$\begin{aligned}\mathcal{Z}_{conf}[J_{\mathbb{O}}, J_{\tilde{\mathbb{O}}}, J_{\mathbb{S}}, J_{\tilde{\mathbb{S}}}] \\ = \frac{1}{Z} \int \mathcal{D}AD\bar{A} e^{-i \int d^4x \bar{A}^a \square A^a} \exp \left( \frac{1}{2} \int d^4x \sum_s J_{\mathbb{O}_s} \bar{A}^a(x) \mathcal{Y}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) A^a(x) \right. \\ \left. + J_{\tilde{\mathbb{O}}_s} \bar{A}^a(x) \mathcal{H}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) A^a(x) \right. \\ \left. + J_{\mathbb{S}_s} \frac{1}{\sqrt{2}} \bar{A}^a(x) \mathcal{Y}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) \bar{A}^a(x) + J_{\tilde{\mathbb{S}}_s} \frac{1}{\sqrt{2}} A^a(x) \mathcal{Y}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) A^a(x) \right)\end{aligned}\quad (3.17)$$

The above functional integral is quadratic in the elementary fields and therefore it may be computed exactly. Employing the symmetry properties of the Gegenbauer polynomials (appendix B):

$$\begin{aligned}\mathbb{O}_s &= \frac{1}{4} \left( \bar{A}^a(x) \mathcal{Y}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) A^a(x) + A^a(x) \mathcal{Y}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) \bar{A}^a(x) \right) \\ \tilde{\mathbb{O}}_s &= \frac{1}{4} \left( \bar{A}^a(x) \mathcal{H}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) A^a(x) - A^a(x) \mathcal{H}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) \bar{A}^a(x) \right)\end{aligned}\quad (3.18)$$

we obtain:

$$\mathcal{Z}_{conf}[J_{\mathbb{O}}, J_{\tilde{\mathbb{O}}}, J_{\mathbb{S}}, J_{\tilde{\mathbb{S}}}] = \frac{1}{Z} \int \mathcal{D}AD\bar{A} e^{-\frac{i}{2} \int d^4x (A^a(x) \bar{A}^a(x)) \mathcal{M}^{ab} \begin{pmatrix} \bar{A}^b(x) \\ A^b(x) \end{pmatrix}}\quad (3.19)$$

with:

$$\mathcal{M}^{ab} = \delta^{ab} \begin{pmatrix} \square + \frac{i}{2} \sum_s J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{i}{2} \sum_s J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} & \frac{i}{\sqrt{2}} \sum_s J_{\mathbb{S}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} \\ \frac{i}{\sqrt{2}} \sum_s J_{\mathbb{S}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} & \square + \frac{i}{2} \sum_s J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{i}{2} \sum_s J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \end{pmatrix} \quad (3.20)$$

where we have introduced the symbol  $\otimes$  to imply that the right and left derivatives do not act on the sources  $J$ . Hence, performing the Gaussian integral we get:

$$\mathcal{Z}_{conf}[J_{\mathbb{0}}, J_{\tilde{\mathbb{0}}}, J_{\mathbb{S}}, J_{\mathbb{S}}] = \text{Det}^{-\frac{1}{2}}(\mathcal{M}) \quad (3.21)$$

where the above determinant – up to a trivial normalization factor – is computed in eq. (A.5). Hence, the generating functional of the connected correlators reads:

$$\begin{aligned} \mathcal{W}_{conf}[J_{\mathbb{0}}, J_{\tilde{\mathbb{0}}}, J_{\mathbb{S}}, J_{\mathbb{S}}] &= \log \mathcal{Z}_{conf}[J_{\mathbb{0}}, J_{\tilde{\mathbb{0}}}, J_{\mathbb{S}}, J_{\mathbb{S}}] \\ &= -\frac{1}{2} \log \text{Det} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{1}{2} i \square^{-1} J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \\ &\quad - \frac{1}{2} \log \text{Det} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2} i \square^{-1} J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \\ &\quad - \frac{1}{2} \log \text{Det} \left[ \mathcal{I} - \frac{1}{2} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2} i \square^{-1} J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right)^{-1} \right. \\ &\quad \left. i \square^{-1} J_{\mathbb{S}_{s_1}} \otimes \mathcal{Y}_{s_1-2}^{\frac{5}{2}} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{0_{s_2}} \otimes \mathcal{Y}_{s_2-2}^{\frac{5}{2}} + \frac{1}{2} i \square^{-1} J_{\tilde{0}_{s_2}} \otimes \mathcal{H}_{s_2-2}^{\frac{5}{2}} \right)^{-1} \right. \\ &\quad \left. i \square^{-1} J_{\mathbb{S}_{s_3}} \otimes \mathcal{Y}_{s_3-2}^{\frac{5}{2}} \right] \end{aligned} \quad (3.22)$$

where  $\mathcal{I}$  is the identity in both color and space-time and the sum over repeated spin indices is understood. After rescaling the operators:

$$\mathcal{O}'_s(x) = \frac{1}{N} \frac{2\Gamma(5)\Gamma(s+1)}{\Gamma(3)\Gamma(s+3)} \mathcal{O}_s(x) \quad (3.23)$$



so that their 2-point correlators are of order 1 for large  $N$ , we obtain more explicitly:

$$\begin{aligned}
\mathcal{W}_{conf}[J_{\mathbb{O}'}, J_{\tilde{\mathbb{O}'}}, J_{\mathbb{S}'}, J_{\tilde{\mathbb{S}'}}] = & \\
& - \frac{1}{2} \log \text{Det} \left( \mathcal{I} + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i\vec{\partial}_+)^{s-k-1} i\Box^{-1} (J_{\mathbb{O}'_s} + J_{\tilde{\mathbb{O}'_s}}) (i\vec{\partial}_+)^{k+1} \right) \\
& - \frac{1}{2} \log \text{Det} \left( \mathcal{I} + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i\vec{\partial}_+)^{s-k-1} i\Box^{-1} (J_{\mathbb{O}'_s} - J_{\tilde{\mathbb{O}'_s}}) (i\vec{\partial}_+)^{k+1} \right) \\
& - \frac{1}{2} \log \text{Det} \left[ \mathcal{I} - \frac{2}{N^2} \left( \mathcal{I} + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i\vec{\partial}_+)^{s-k-1} i\Box^{-1} (J_{\mathbb{O}'_s} - J_{\tilde{\mathbb{O}'_s}}) (i\vec{\partial}_+)^{k+1} \right)^{-1} \right. \\
& \quad \sum_{k_1=0}^{s_1-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} (i\vec{\partial}_+)^{s_1-k_1-1} i\Box^{-1} J_{\tilde{\mathbb{S}'_{s_1}}} (i\vec{\partial}_+)^{k_1+1} \\
& \quad \left. \left( \mathcal{I} + \frac{1}{N} \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (i\vec{\partial}_+)^{s_2-k_2-1} i\Box^{-1} (J_{\mathbb{O}'_{s_2}} + J_{\tilde{\mathbb{O}'_{s_2}}}) (i\vec{\partial}_+)^{k_2+1} \right)^{-1} \right. \\
& \quad \left. \sum_{k_3=0}^{s_3-2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} (i\vec{\partial}_+)^{s_3-k_3-1} i\Box^{-1} J_{\tilde{\mathbb{S}'_{s_3}}} (i\vec{\partial}_+)^{k_3+1} \right] \tag{3.24}
\end{aligned}$$

where we have employed the definitions in eqs. (3.14) and (3.15) and the identity:

$$i\Box^{-1} \overleftarrow{\partial}_+^{s-k-1} = (-1)^{s-k-1} \overrightarrow{\partial}_+^{s-k-1} i\Box^{-1} \tag{3.25}$$

that follows from the propagator in the coordinate representation [1]:

$$\frac{1}{4\pi^2} \frac{1}{|x-y|^2 - i\epsilon} = i\Box^{-1}(x-y) \tag{3.26}$$

#### 4 Connection of $\mathcal{W}_{conf}$ with the generating functionals $\Gamma_{conf}$ in [1]

Remarkably,  $\mathcal{W}_{conf}[J_{\mathbb{O}}, J_{\tilde{\mathbb{O}}}, J_{\mathbb{S}}, J_{\tilde{\mathbb{S}}}]$  is the generating functional of all the Minkowskian conformal correlators that extends the generating functionals in the separate balanced and unbalanced sectors obtained [1] by guessing their structure from the perturbative computation of the corresponding  $n$ -point correlators. However, even restricting to the separate balanced and unbalanced sectors, the connection with the approach in [1] is not immediately visible. Indeed, the connected generating functionals in the separate sectors read [1]:

$$\begin{aligned}
\Gamma_{conf}[j_{\mathbb{O}}, j_{\tilde{\mathbb{O}}}] = & - \frac{N^2 - 1}{2} \log \text{Det} \left( \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} + \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \right) \\
& - \frac{N^2 - 1}{2} \log \text{Det} \left( \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} - \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \right) \\
\Gamma_{conf}[j_{\mathbb{S}}, j_{\tilde{\mathbb{S}}}] = & - \frac{N^2 - 1}{2} \log \text{Det} \left( \mathbb{I} - 2\mathcal{D}^{-1} j_{\tilde{\mathbb{S}}} \mathcal{D}^{-1} j_{\mathbb{S}} \right) \tag{4.1}
\end{aligned}$$

in Minkowskian space-time, where  $\mathbb{I}$  is the identity in both space-time and the discrete indices defined below. By making manifest the space-time and discrete-indices dependence in the kernels of the integral operators above, eq. (4.1) reads in the coordinate representation [1]:

$$\begin{aligned}
& \Gamma_{conf}[j_{\mathbb{O}}, j_{\tilde{\mathbb{O}}}] \\
&= -\frac{N^2-1}{2} \log \text{Det} \left( \delta_{s_1 k_1, s_2 k_2} \delta^{(4)}(x-y) + \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x-y) (j_{\mathbb{O}_{s_2 k_2}}(y) + j_{\tilde{\mathbb{O}}_{s_2 k_2}}(y)) \right) \\
&\quad - \frac{N^2-1}{2} \log \text{Det} \left( \delta_{s_1 k_1, s_2 k_2} \delta^{(4)}(x-y) + \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x-y) (j_{\mathbb{O}_{s_2 k_2}}(y) - j_{\tilde{\mathbb{O}}_{s_2 k_2}}(y)) \right) \\
& \\
& \Gamma_{conf}[j_{\mathbb{S}}, j_{\tilde{\mathbb{S}}}] \\
&= -\frac{N^2-1}{2} \log \text{Det} \left( \delta_{s_1 k_1, s_2 k_2} \delta^{(4)}(x-y) \right. \\
&\quad \left. - 2 \int d^4 z \sum_{sk} \mathcal{D}_{s_1 k_1, sk}^{-1}(x-z) j_{\tilde{\mathbb{S}}_{sk}}(z) \mathcal{D}_{sk, s_2 k_2}^{-1}(z-y) j_{\mathbb{S}_{s_2 k_2}}(y) \right) \tag{4.2}
\end{aligned}$$

where, by a slight abuse of notation, we have displayed as arguments of the determinants the corresponding integral kernels, with [1]:

$$\begin{aligned}
\mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x-y) &= \frac{i^{s_1+1} \Gamma(3) \Gamma(s_1+3)}{2 \Gamma(5) \Gamma(s_1+1)} \binom{s_1}{k_1} \binom{s_2}{k_2+2} (-\partial_+)^{s_1-k_1+k_2} \square^{-1}(x-y) \\
&= \frac{i^{s_1} \Gamma(3) \Gamma(s_1+3)}{8\pi^2 \Gamma(5) \Gamma(s_1+1)} \binom{s_1}{k_1} \binom{s_2}{k_2+2} (-\partial_+)^{s_1-k_1+k_2} \frac{1}{|x-y|^2 - i\epsilon} \tag{4.3}
\end{aligned}$$

The currents  $j_{\mathcal{O}_{sk}}$  are dual to the component operators  $\mathcal{O}_{sk}$  that are employed to construct the conformal operators  $\mathcal{O}_s$  [1]:

$$\mathcal{O}_s = \sum_{k=0}^l \mathcal{O}_{sk} \tag{4.4}$$

with  $l = s - 2$  for the standard basis. Consequently, the  $n$ -point correlators satisfy [1]:

$$\langle \mathcal{O}_{s_1}(x_1) \dots \mathcal{O}_{s_n}(x_n) \rangle = \sum_{k_1=0}^{l_1} \frac{\delta}{\delta j_{\mathcal{O}_{s_1 k_1}}(x_1)} \dots \sum_{k_n=0}^{l_n} \frac{\delta}{\delta j_{\mathcal{O}_{s_n k_n}}(x_n)} \Gamma_{conf}[j_{\mathcal{O}}] \tag{4.5}$$

Hence, we should demonstrate that:

$$\sum_{k_1=0}^{l_1} \frac{\delta}{\delta j_{\mathcal{O}_{s_1 k_1}}(x_1)} \dots \sum_{k_n=0}^{l_n} \frac{\delta}{\delta j_{\mathcal{O}_{s_n k_n}}(x_n)} \Gamma_{conf}[j_{\mathcal{O}}] = \frac{\delta}{\delta J_{\mathcal{O}_{s_1}}(x_1)} \dots \frac{\delta}{\delta J_{\mathcal{O}_{s_n}}(x_n)} \mathcal{W}_{conf}[J_{\mathcal{O}}] \tag{4.6}$$

with the functional derivatives computed at  $j_{\mathcal{O}_{sk}} = 0$  and  $J_{\mathcal{O}_s} = 0$  respectively. While  $\Gamma_{conf}[j_{\mathcal{O}}]$  involves determinants of integral operators formally of Fredholm type, the path-integral computation of  $\mathcal{W}_{conf}[J_{\mathcal{O}}]$  involves determinants of quadratic forms that originally arise from their very definition in eq. (3.22) by employing both left and right derivatives in eqs. (3.14) and (3.15). This is the source of some technical complications that we resolve momentarily creating a dictionary that relates  $\mathcal{W}_{conf}[J_{\mathcal{O}}]$  to  $\Gamma_{conf}[j_{\mathcal{O}}]$ .

#### 4.1 An example of the dictionary

We choose as example the generating functional  $\mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0]$  restricted to the balanced operators with even spin in the standard basis. We formally expand it:

$$\begin{aligned}
\mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0] &= -\log \text{Det} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} \right) \\
&= -\log \text{Det} \left( \delta^{ab} \delta^{(4)}(x-y) + \frac{1}{2} \delta^{ab} i \square^{-1}(x-y) J_{\mathbb{O}_s}(y) \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} \right) \\
&= (N^2 - 1) \sum_{k=1}^{\infty} \frac{(-1)^k}{k} \frac{1}{2^k} \int d^4 x_1 \dots d^4 x_k \sum_{s_1} \dots \sum_{s_k} \\
&\quad i \square^{-1}(x_1 - x_2) J_{\mathbb{O}_{s_2}}(x_2) \otimes \mathcal{Y}_{s_2-2}^{\frac{5}{2}}(\vec{\partial}_{x_2^+}, \overleftarrow{\partial}_{x_2^+}) \\
&\quad \dots i \square^{-1}(x_k - x_1) J_{\mathbb{O}_{s_1}}(x_1) \otimes \mathcal{Y}_{s_1-2}^{\frac{5}{2}}(\vec{\partial}_{x_1^+}, \overleftarrow{\partial}_{x_1^+})
\end{aligned} \tag{4.7}$$

where the factor  $N^2 - 1$  arises by performing the color trace. By definition:

$$\begin{aligned}
\langle \mathbb{O}_{s_1}(x_1) \dots \mathbb{O}_{s_n}(x_n) \rangle &= \frac{\delta}{\delta J_{\mathbb{O}_{s_1}}(x_1)} \dots \frac{\delta}{\delta J_{\mathbb{O}_{s_n}}(x_n)} \mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0] \\
&= (N^2 - 1) \frac{(-1)^n}{n} \frac{1}{2^n} \sum_{\sigma \in P_n} i \square^{-1}(x_{\sigma(1)} - x_{\sigma(2)}) \mathcal{Y}_{s_{\sigma(2)}-2}^{\frac{5}{2}}(\vec{\partial}_{x_{\sigma(2)}^+}, \overleftarrow{\partial}_{x_{\sigma(2)}^+}) \\
&\quad \dots i \square^{-1}(x_{\sigma(n)} - x_{\sigma(1)}) \mathcal{Y}_{s_{\sigma(1)}-2}^{\frac{5}{2}}(\vec{\partial}_{x_{\sigma(1)}^+}, \overleftarrow{\partial}_{x_{\sigma(1)}^+})
\end{aligned} \tag{4.8}$$

We relabel the permutations in the sum,  $\sigma(n) \rightarrow \sigma(2)$ ,  $\sigma(n-1) \rightarrow \sigma(3)$ ,  $\sigma(n-2) \rightarrow \sigma(4), \dots$ , so that:

$$\begin{aligned}
\langle \mathbb{O}_{s_1}(x_1) \dots \mathbb{O}_{s_n}(x_n) \rangle &= \frac{\delta}{\delta J_{\mathbb{O}_{s_1}}(x_1)} \dots \frac{\delta}{\delta J_{\mathbb{O}_{s_n}}(x_n)} \mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0] \\
&= (N^2 - 1) \frac{(-1)^n}{n} \frac{1}{2^n} \sum_{\sigma \in P_n} i \square^{-1}(x_{\sigma(1)} - x_{\sigma(n)}) \mathcal{Y}_{s_{\sigma(n)}-2}^{\frac{5}{2}}(\vec{\partial}_{x_{\sigma(n)}^+}, \overleftarrow{\partial}_{x_{\sigma(n)}^+}) \\
&\quad \dots i \square^{-1}(x_{\sigma(2)} - x_{\sigma(1)}) \mathcal{Y}_{s_{\sigma(1)}-2}^{\frac{5}{2}}(\vec{\partial}_{x_{\sigma(1)}^+}, \overleftarrow{\partial}_{x_{\sigma(1)}^+})
\end{aligned} \tag{4.9}$$

Employing the definition in eq. (3.14), we get:

$$\begin{aligned}
\langle \mathbb{O}_{s_1}(x_1) \dots \mathbb{O}_{s_n}(x_n) \rangle &= \frac{\delta}{\delta J_{\mathbb{O}_{s_1}}(x_1)} \dots \frac{\delta}{\delta J_{\mathbb{O}_{s_n}}(x_n)} \mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0] \\
&= (N^2 - 1) \frac{(-1)^n}{n} \frac{1}{2^n} i^{\sum_{i=1}^n s_i} (-1)^n \frac{\Gamma(3)\Gamma(s_1+3)}{\Gamma(5)\Gamma(s_1+1)} \dots \frac{\Gamma(3)\Gamma(s_n+3)}{\Gamma(5)\Gamma(s_n+1)} \\
&\quad \sum_{k_1=0}^{s_1-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} (-1)^{s_1-k_1} \dots \sum_{k_n=0}^{s_n-2} \binom{s_n}{k_n} \binom{s_n}{k_n+2} (-1)^{s_n-k_n} \\
&\quad \sum_{\sigma \in P_n} i \square^{-1}(x_{\sigma(1)} - x_{\sigma(n)}) \overleftarrow{\partial}_{x_{\sigma(n)}^+}^{s_{\sigma(n)}-k_{\sigma(n)}-1} \overrightarrow{\partial}_{x_{\sigma(n)}^+}^{k_{\sigma(n)}+1} \\
&\quad \dots i \square^{-1}(x_{\sigma(2)} - x_{\sigma(1)}) \overleftarrow{\partial}_{x_{\sigma(1)}^+}^{s_{\sigma(1)}-k_{\sigma(1)}-1} \overrightarrow{\partial}_{x_{\sigma(1)}^+}^{k_{\sigma(1)}+1}
\end{aligned} \tag{4.10}$$

We arrange the left and right derivatives into a single object:

$$\begin{aligned} & \overrightarrow{\partial}_{x_{\sigma(a)}^+}^{k_{\sigma(a)}+1} i \square^{-1} (x_{\sigma(a)} - x_{\sigma(b)}) \overleftarrow{\partial}_{x_{\sigma(b)}^+}^{s_{\sigma(b)} - k_{\sigma(b)} - 1} \\ &= (-1)^{s_{\sigma(b)} - k_{\sigma(b)} - 1} \overrightarrow{\partial}_{x_{\sigma(a)}^+}^{s_{\sigma(b)} - k_{\sigma(b)} + k_{\sigma(a)}} i \square^{-1} (x_{\sigma(a)} - x_{\sigma(b)}) \end{aligned} \quad (4.11)$$

so that, employing eq. (3.26) and rearranging the terms in eq. (4.10) from the last to the first, we obtain:

$$\begin{aligned} \langle \mathbb{O}_{s_1}(x_1) \dots \mathbb{O}_{s_n}(x_n) \rangle &= \frac{\delta}{\delta J_{\mathbb{O}_{s_1}}(x_1)} \dots \frac{\delta}{\delta J_{\mathbb{O}_{s_n}}(x_n)} \mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0] \\ &= \frac{1}{(4\pi^2)^n} \frac{N^2 - 1}{2^n} i^{\sum_{i=1}^n s_i} \frac{\Gamma(3)\Gamma(s_1 + 3)}{\Gamma(5)\Gamma(s_1 + 1)} \dots \frac{\Gamma(3)\Gamma(s_n + 3)}{\Gamma(5)\Gamma(s_n + 1)} \\ &\quad \sum_{k_1=0}^{s_1-2} \binom{s_1}{k_1} \binom{s_1}{k_1 + 2} \dots \sum_{k_n=0}^{s_n-2} \binom{s_n}{k_n} \binom{s_n}{k_n + 2} \\ &\quad \frac{(-1)^n}{n} \sum_{\sigma \in P_n} \partial_{x_{\sigma(1)}^+}^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}} \frac{1}{|x_{\sigma(2)} - x_{\sigma(1)}|^2} \\ &\quad \partial_{x_{\sigma(2)}^+}^{s_{\sigma(2)} - k_{\sigma(2)} + k_{\sigma(3)}} \frac{1}{|x_{\sigma(3)} - x_{\sigma(2)}|^2} \\ &\quad \dots \partial_{x_{\sigma(n)}^+}^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}} \frac{1}{|x_{\sigma(1)} - x_{\sigma(n)}|^2} \end{aligned} \quad (4.12)$$

After switching the position of the coordinates inside the propagators that introduces a minus sign in the derivatives, we get:

$$\begin{aligned} \langle \mathbb{O}_{s_1}(x_1) \dots \mathbb{O}_{s_n}(x_n) \rangle &= \frac{\delta}{\delta J_{\mathbb{O}_{s_1}}(x_1)} \dots \frac{\delta}{\delta J_{\mathbb{O}_{s_n}}(x_n)} \mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0] \\ &= \frac{1}{(4\pi^2)^n} \frac{N^2 - 1}{2^n} i^{\sum_{i=1}^n s_i} \frac{\Gamma(3)\Gamma(s_1 + 3)}{\Gamma(5)\Gamma(s_1 + 1)} \dots \frac{\Gamma(3)\Gamma(s_n + 3)}{\Gamma(5)\Gamma(s_n + 1)} \\ &\quad \sum_{k_1=0}^{s_1-2} \binom{s_1}{k_1} \binom{s_1}{k_1 + 2} \dots \sum_{k_n=0}^{s_n-2} \binom{s_n}{k_n} \binom{s_n}{k_n + 2} \\ &\quad \frac{(-1)^n}{n} \sum_{\sigma \in P_n} (-\partial_{x_{\sigma(1)}^+})^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}} \frac{1}{|x_{\sigma(1)} - x_{\sigma(2)}|^2} \\ &\quad (-\partial_{x_{\sigma(2)}^+})^{s_{\sigma(2)} - k_{\sigma(2)} + k_{\sigma(3)}} \frac{1}{|x_{\sigma(2)} - x_{\sigma(3)}|^2} \\ &\quad \dots (-\partial_{x_{\sigma(n)}^+})^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}} \frac{1}{|x_{\sigma(n)} - x_{\sigma(1)}|^2} \end{aligned} \quad (4.13)$$

that perfectly agrees with [1]. We may rewrite the above equation according with eq. (4.3):

$$\begin{aligned} \langle \mathbb{O}_{s_1}(x_1) \dots \mathbb{O}_{s_n}(x_n) \rangle &= \frac{\delta}{\delta J_{\mathbb{O}_{s_1}}(x_1)} \dots \frac{\delta}{\delta J_{\mathbb{O}_{s_n}}(x_n)} \mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0] \\ &= (N^2 - 1) \sum_{k_1=0}^{s_1-2} \dots \sum_{k_n=0}^{s_n-2} \frac{(-1)^n}{n} \sum_{\sigma \in P_n} \mathcal{D}_{s_{\sigma(1)} k_{\sigma(1)}, s_{\sigma(2)} k_{\sigma(2)}}^{-1} (x_{\sigma(1)} - x_{\sigma(2)}) \\ &\quad \dots \mathcal{D}_{s_{\sigma(n)} k_{\sigma(n)}, s_{\sigma(1)} k_{\sigma(1)}}^{-1} (x_{\sigma(n)} - x_{\sigma(1)}) \end{aligned} \quad (4.14)$$

Hence, introducing the new currents  $j_{\mathbb{O}_{s_k}}$  as above, we obtain according with eq. (4.14):

$$\begin{aligned} & \frac{\delta}{\delta J_{\mathbb{O}_{s_1}}(x_1)} \cdots \frac{\delta}{\delta J_{\mathbb{O}_{s_n}}(x_n)} \mathcal{W}_{conf}[J_{\mathbb{O}}, 0, 0, 0] \\ &= \sum_{k_1=0}^{s_1-1} \frac{\delta}{\delta j_{\mathbb{O}_{s_1 k_1}}(x_1)} \cdots \sum_{k_n=0}^{s_n-1} \frac{\delta}{\delta j_{\mathbb{O}_{s_n k_n}}(x_n)} \Gamma_{conf}[j_{\mathbb{O}}, 0, 0, 0] \end{aligned} \quad (4.15)$$

where [1]:

$$\begin{aligned} \Gamma_{conf}[j_{\mathbb{O}}, 0, 0, 0] &= - (N^2 - 1) \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \int d^4 x_1 \cdots d^4 x_n \sum_{s_1 k_1} \cdots \sum_{s_n k_n} \\ & \quad \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x_1 - x_2) j_{\mathbb{O}_{s_2 k_2}}(x_2) \cdots \mathcal{D}_{s_n k_n, s_1 k_1}^{-1}(x_n - x_1) j_{\mathbb{O}_{s_1 k_1}}(x_1) \\ &= - (N^2 - 1) \log \text{Det} (\mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}}) \end{aligned} \quad (4.16)$$

according with eq. (4.1).

## 4.2 Dictionary in the general case

The dictionary in the general case is constructed as follows. We expand the traces of the logarithms in eq. (3.22):

$$\begin{aligned} & \mathcal{W}_{conf}[J_{\mathbb{O}}, J_{\tilde{\mathbb{O}}}, J_{\mathbb{S}}, J_{\bar{\mathbb{S}}}] \\ &= \frac{N^2 - 1}{2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr} \left( \frac{1}{2} i \square^{-1} (J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + J_{\tilde{\mathbb{O}}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}}) \right)^n \\ &+ \frac{N^2 - 1}{2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr} \left( \frac{1}{2} i \square^{-1} (J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - J_{\tilde{\mathbb{O}}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}}) \right)^n \\ &+ \frac{N^2 - 1}{2} \sum_{n=1}^{\infty} \frac{1}{n} \text{Tr} \left[ \frac{1}{2} \left( I + \frac{1}{2} i \square^{-1} (J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - J_{\tilde{\mathbb{O}}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}}) \right)^{-1} i \square^{-1} J_{\bar{\mathbb{S}}_{s_1}} \otimes \mathcal{Y}_{s_1-2}^{\frac{5}{2}} \right. \\ & \quad \left. \left( I + \frac{1}{2} i \square^{-1} (J_{\mathbb{O}_{s_2}} \otimes \mathcal{Y}_{s_2-2}^{\frac{5}{2}} + J_{\tilde{\mathbb{O}}_{s_2}} \otimes \mathcal{H}_{s_2-2}^{\frac{5}{2}}) \right)^{-1} i \square^{-1} J_{\mathbb{S}_{s_3}} \otimes \mathcal{Y}_{s_3-2}^{\frac{5}{2}} \right]^n \end{aligned} \quad (4.17)$$

where  $I$  is the identity in space-time and we have already performed the color trace. Expanding:

$$\begin{aligned} & \left( I + \frac{1}{2} i \square^{-1} (J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} \pm J_{\tilde{\mathbb{O}}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}}) \right)^{-1} \\ &= \sum_{n=0}^{\infty} (-1)^n \left( \frac{1}{2} i \square^{-1} (J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} \pm J_{\tilde{\mathbb{O}}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}}) \right)^n \end{aligned} \quad (4.18)$$

we obtain:

$$\begin{aligned}
& \mathcal{W}_{conf}[J_{\mathbb{O}}, J_{\tilde{\mathbb{O}}}, J_{\mathbb{S}}, J_{\tilde{\mathbb{S}}}] \\
&= \frac{N^2 - 1}{2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr} \left( \frac{1}{2} i^{\square-1} (J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + J_{\tilde{\mathbb{O}}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}}) \right)^n \\
&+ \frac{N^2 - 1}{2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr} \left( \frac{1}{2} i^{\square-1} (J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - J_{\tilde{\mathbb{O}}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}}) \right)^n \\
&+ \frac{N^2 - 1}{2} \sum_{n=1}^{\infty} \frac{1}{n} \\
&\text{Tr} \left[ \frac{1}{2} \sum_{m_1=0}^{\infty} (-1)^{m_1} \left( \frac{1}{2} i^{\square-1} (J_{\mathbb{O}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - J_{\tilde{\mathbb{O}}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}}) \right)^{m_1} i^{\square-1} J_{\tilde{\mathbb{S}}_{s_1}} \otimes \mathcal{Y}_{s_1-2}^{\frac{5}{2}} \right. \\
&\left. \sum_{m_2=0}^{\infty} (-1)^{m_2} \left( \frac{1}{2} i^{\square-1} (J_{\mathbb{O}_{s_2}} \otimes \mathcal{Y}_{s_2-2}^{\frac{5}{2}} + J_{\tilde{\mathbb{O}}_{s_2}} \otimes \mathcal{Y}_{s_2-2}^{\frac{5}{2}}) \right)^{m_2} i^{\square-1} J_{\mathbb{S}_{s_3}} \otimes \mathcal{Y}_{s_3-2}^{\frac{5}{2}} \right]^n \quad (4.19)
\end{aligned}$$

Though the above expansion looks complicated, we do not actually need to explicitly perform the functional derivatives. It suffices to observe that a generic term of the expansion has the structure:

$$\text{Tr} i^{\square-1} J \mathcal{Y} i^{\square-1} J \mathcal{Y} i^{\square-1} J \mathcal{Y} i^{\square-1} \dots i^{\square-1} J \mathcal{H} i^{\square-1} J \mathcal{H} i^{\square-1} J \mathcal{H} i^{\square-1} \dots i^{\square-1} J \mathcal{Y} i^{\square-1} J \mathcal{Y} \quad (4.20)$$

where the trace includes all the objects to its right. Therefore, by performing the functional derivatives, the products of  $i^{\square-1} \mathcal{Y} i^{\square-1}$  or  $i^{\square-1} \mathcal{H} i^{\square-1}$  may be rearranged thanks to the cyclicity of the trace to build the corresponding kernels of integral operators by acting with the right or left derivatives on the same  $i^{\square-1}$ , as in the previous example of the dictionary.

Accordingly, the generating functional may be rewritten in terms of the kernel  $\mathcal{D}^{-1}$  and the new currents  $j_{\mathcal{O}_{sk}}$  dual to the component operators  $\mathcal{O}_{sk}$ .

### 4.3 Generating functional $\Gamma_{conf}$

Hence, the construction above may be condensed into the single formula:

$$\Gamma_{conf}[j_{\mathbb{O}}, j_{\tilde{\mathbb{O}}}, j_{\mathbb{S}}, j_{\tilde{\mathbb{S}}}] = -\frac{N^2 - 1}{2} \log \text{Det } \mathbf{M} \quad (4.21)$$

with:

$$\mathbf{M} = \begin{pmatrix} \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} - \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} & \sqrt{2} \mathcal{D}^{-1} j_{\tilde{\mathbb{S}}} \\ \sqrt{2} \mathcal{D}^{-1} j_{\mathbb{S}} & \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} + \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \end{pmatrix} \quad (4.22)$$

and:

$$\mathcal{D}_{s_1 k_1, s_2 k_2}^{-1} = \frac{i^{s_1} \Gamma(3) \Gamma(s_1 + 3)}{2 \Gamma(5) \Gamma(s_1 + 1)} \binom{s_1}{k_1} \binom{s_2}{k_2 + 2} (-\partial_+)^{s_1 - k_1 + k_2} i^{\square-1} \quad (4.23)$$

The above determinant reads (appendix A):

$$\begin{aligned}
& \Gamma_{conf}[j_{\mathbb{O}}, j_{\tilde{\mathbb{O}}}, j_{\mathbb{S}}, j_{\tilde{\mathbb{S}}}] \\
&= -\frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} + \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \right] \\
&\quad - \frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} - \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \right] \\
&\quad - \frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} - 2 \left( \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} - \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \right)^{-1} \mathcal{D}^{-1} j_{\tilde{\mathbb{S}}} \left( \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} + \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \right)^{-1} \mathcal{D}^{-1} j_{\mathbb{S}} \right]
\end{aligned} \tag{4.24}$$

The generating functional immediately reduces – by setting the corresponding sources to zero – to the generating functionals in the separate balanced and unbalanced sectors [1]:

$$\begin{aligned}
& \Gamma_{conf}[j_{\mathbb{O}}, j_{\tilde{\mathbb{O}}}] = \Gamma_{conf}[j_{\mathbb{O}}, j_{\tilde{\mathbb{O}}}, 0, 0] \\
&= -\frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} - \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \right] - \frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} + \mathcal{D}^{-1} j_{\mathbb{O}} + \mathcal{D}^{-1} j_{\tilde{\mathbb{O}}} \right] \\
& \Gamma_{conf}[j_{\mathbb{S}}, j_{\tilde{\mathbb{S}}}] = \Gamma_{conf}[0, 0, j_{\mathbb{S}}, j_{\tilde{\mathbb{S}}}] = -\frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} - 2 \mathcal{D}^{-1} j_{\tilde{\mathbb{S}}} \mathcal{D}^{-1} j_{\mathbb{S}} \right]
\end{aligned} \tag{4.25}$$

Moreover, the generating functional of the Euclidean correlators reads:

$$\begin{aligned}
& \Gamma_{conf}^E[j_{\mathbb{O}^E}, j_{\tilde{\mathbb{O}}^E}, j_{\mathbb{S}^E}, j_{\tilde{\mathbb{S}}^E}] \\
&= -\frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} + \mathcal{D}_E^{-1} j_{\mathbb{O}^E} + \mathcal{D}_E^{-1} j_{\tilde{\mathbb{O}}^E} \right] \\
&\quad - \frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} + \mathcal{D}_E^{-1} j_{\mathbb{O}^E} - \mathcal{D}_E^{-1} j_{\tilde{\mathbb{O}}^E} \right] \\
&\quad - \frac{N^2-1}{2} \text{Tr log} \left[ \mathbb{I} - 2 \left( \mathbb{I} + \mathcal{D}_E^{-1} j_{\mathbb{O}^E} - \mathcal{D}_E^{-1} j_{\tilde{\mathbb{O}}^E} \right)^{-1} \mathcal{D}_E^{-1} j_{\tilde{\mathbb{S}}^E} \right. \\
&\quad \quad \left. \left( \mathbb{I} + \mathcal{D}_E^{-1} j_{\mathbb{O}^E} + \mathcal{D}_E^{-1} j_{\tilde{\mathbb{O}}^E} \right)^{-1} \mathcal{D}_E^{-1} j_{\mathbb{S}^E} \right]
\end{aligned} \tag{4.26}$$

with:

$$\mathcal{D}_E^{-1}{}_{s_1 k_1, s_2 k_2} = \frac{(-i)^{k_1-k_2}}{2} \frac{\Gamma(3)\Gamma(s_1+3)}{\Gamma(5)\Gamma(s_1+1)} \binom{s_1}{k_1} \binom{s_2}{k_2+2} \partial_z^{s_1-k_1+k_2} \Delta^{-1}$$

obtained by Wick rotation [1], where:

$$\Delta = \delta_{\mu\nu} \partial_\mu \partial_\nu = \partial_4^2 + \sum_{i=1}^3 \partial_i^2 \tag{4.27}$$

and [1]:

$$\Delta^{-1}(x-y) = -\frac{1}{4\pi^2} \frac{1}{(x-y)^2} \tag{4.28}$$

## 5 Generating functional in the momentum representation

The generating functional in the momentum representation is defined by the path integral:

$$\mathcal{Z}_{conf}[J_{\mathcal{O}}] = \int \mathcal{D}A \mathcal{D}\bar{A} e^{-i \int d^4x \bar{A}^a \square A^a} \exp \left( \int \frac{d^4p}{(2\pi)^4} \sum_i J_{\mathcal{O}_i}(-p) \mathcal{O}_i(p) \right) \quad (5.1)$$

Correspondingly, the correlators in the momentum representation read:

$$\langle \mathcal{O}_{s_1}(p_1) \dots \mathcal{O}_{s_n}(p_n) \rangle = (2\pi)^4 \frac{\delta}{\delta J_{\mathcal{O}_{s_1}}(-p_1)} \dots (2\pi)^4 \frac{\delta}{\delta J_{\mathcal{O}_{s_n}}(-p_n)} \mathcal{W}_{conf}[J_{\mathcal{O}}] \quad (5.2)$$

Employing the dictionary, we get as well:

$$\begin{aligned} & \langle \mathcal{O}_{s_1}(p_1) \dots \mathcal{O}_{s_n}(p_n) \rangle \\ &= \sum_{k_1=0}^{l_1} (2\pi)^4 \frac{\delta}{\delta j_{\mathcal{O}_{s_1 k_1}}(-p_1)} \dots \sum_{k_n=0}^{l_n} (2\pi)^4 \frac{\delta}{\delta j_{\mathcal{O}_{s_n k_n}}(-p_n)} \Gamma_{conf}[j_{\mathcal{O}}] \end{aligned} \quad (5.3)$$

### 5.1 Standard basis

To find the explicit form of the generating functional in the momentum representation we choose again as example:

$$\Gamma_{conf}[j_{\mathcal{O}}] = -(N^2 - 1) \log \text{Det} \left( \delta_{s_1 k_1, s_2 k_2} \delta^{(4)}(x - y) + \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x - y) j_{\mathbb{O}_{s_2 k_2}}(y) \right) \quad (5.4)$$

where the argument of the determinant is the kernel:

$$K_{s_1 k_1, s_2 k_2}(x, y) = \delta_{s_1 k_1, s_2 k_2} \delta^{(4)}(x - y) + \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x - y) j_{\mathbb{O}_{s_2 k_2}}(y) \quad (5.5)$$

of the integral operator:

$$\psi_{s_1 k_1}(x) = \sum_{s_2 k_2} \int K_{s_1 k_1, s_2 k_2}(x, y) \phi_{s_2 k_2}(y) d^4 y \quad (5.6)$$

In order to obtain the kernel in the momentum representation [1] we perform the Fourier transform of the lhs:

$$\psi_{s_1 k_1}(q) = \int \psi_{s_1 k_1}(x) e^{-iqx} d^4 x = \sum_{s_2 k_2} \int K_{s_1 k_1, s_2 k_2}(x, y) \phi_{s_2 k_2}(y) e^{-iqx} d^4 x d^4 y \quad (5.7)$$

and write the rhs in terms of the Fourier-transformed fields:

$$\psi_{s_1 k_1}(q) = \sum_{s_2 k_2} \int K_{s_1 k_1, s_2 k_2}(x, y) \phi_{s_2 k_2}(p) e^{ipy} e^{-iqx} d^4 x d^4 y \frac{d^4 p}{(2\pi)^4} \quad (5.8)$$

Substituting the kernel in eq. (5.5):

$$\begin{aligned} \psi_{s_1 k_1}(q) &= \sum_{s_2 k_2} \int \delta_{s_1 k_1, s_2 k_2} \delta^{(4)}(x - y) \phi_{s_2 k_2}(p) e^{ipy} e^{-iqx} \\ &\quad + \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x - y) j_{\mathbb{O}_{s_2 k_2}}(y) \phi_{s_2 k_2}(p) e^{ipy} e^{-iqx} d^4 x d^4 y \frac{d^4 p}{(2\pi)^4} \end{aligned} \quad (5.9)$$



we get:

$$\begin{aligned}\psi_{s_1 k_1}(q) &= \sum_{s_2 k_2} \int \delta_{s_1 k_1, s_2 k_2} (2\pi)^4 \delta^{(4)}(p - q) \phi_{s_2 k_2}(p) \frac{d^4 p}{(2\pi)^4} \\ &\quad + \sum_{s_2 k_2} \int \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x - y) j_{\mathbb{O}_{s_2 k_2}}(y) \phi_{s_2 k_2}(p) e^{ipy} e^{-iqx} d^4 x d^4 y \frac{d^4 p}{(2\pi)^4}\end{aligned}\quad (5.10)$$

The second line in the above equation becomes:

$$\sum_{s_2 k_2} \int \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(k_1) j_{\mathbb{O}_{s_2 k_2}}(k_2) \phi_{s_2 k_2}(p) e^{ik_1(x-y)} e^{ik_2 y} e^{ipy} e^{-iqx} d^4 x d^4 y \frac{d^4 p}{(2\pi)^4} \frac{d^4 k_1}{(2\pi)^4} \frac{d^4 k_2}{(2\pi)^4}\quad (5.11)$$

that reduces to:

$$\sum_{s_2 k_2} \int \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(q) j_{\mathbb{O}_{s_2 k_2}}(q - p) \phi_{s_2 k_2}(p) \frac{d^4 p}{(2\pi)^4}\quad (5.12)$$

Therefore, the kernel in the momentum representation is:

$$K_{s_1 k_1, s_2 k_2}(q_1, q_2) = \delta_{s_1 k_1, s_2 k_2} (2\pi)^4 \delta^{(4)}(q_1 - q_2) + \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(q_1) j_{\mathbb{O}_{s_2 k_2}}(q_1 - q_2)\quad (5.13)$$

that defines the integral operator:

$$\psi_{s_1 k_1}(q_1) = \sum_{s_2 k_2} \int K_{s_1 k_1, s_2 k_2}(q_1, q_2) \phi_{s_2 k_2}(q_2) \frac{d^4 q_2}{(2\pi)^4}\quad (5.14)$$

with:

$$\mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(p) = \frac{i^{s_1} \Gamma(3) \Gamma(s_1 + 3)}{2 \Gamma(5) \Gamma(s_1 + 1)} \binom{s_1}{k_1} \binom{s_2}{k_2 + 2} (-ip_+)^{s_1 - k_1 + k_2} \frac{-i}{|p|^2 + i\epsilon}\quad (5.15)$$

Hence, the generating functional in the momentum representation reads [1]:

$$\Gamma_{conf}[j_{\mathbb{O}}] = -(N^2 - 1) \log \text{Det} \left( \delta_{s_1 k_1, s_2 k_2} (2\pi)^4 \delta^{(4)}(q_1 - q_2) + \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(q_1) j_{\mathbb{O}_{s_2 k_2}}(q_1 - q_2) \right)\quad (5.16)$$

In order to obtain the correlators we expand the generating functional:

$$\begin{aligned}\Gamma_{conf}[j_{\mathbb{O}}] &= -(N^2 - 1) \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \int \frac{d^4 q_1}{(2\pi)^4} \cdots \frac{d^4 q_n}{(2\pi)^4} \sum_{s_1 k_1} \cdots \sum_{s_n k_n} \\ &\quad \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(q_1) j_{\mathbb{O}_{s_2 k_2}}(q_1 - q_2) \mathcal{D}_{s_2 k_2, s_3 k_3}^{-1}(q_2) j_{\mathbb{O}_{s_3 k_3}}(q_2 - q_3) \\ &\quad \cdots \mathcal{D}_{s_n k_n, s_1 k_1}^{-1}(q_n) j_{\mathbb{O}_{s_1 k_1}}(q_n - q_1)\end{aligned}\quad (5.17)$$

We change variables defining  $p_i = q_i - q_{i+1}$  with  $q_{n+1} \equiv q_1$  that automatically ensures  $\sum_{i=1}^n p_i = 0$ :

$$\begin{aligned}
q_1 &= q_n - p_n \\
q_2 &= q_1 - p_1 = q_n - p_n - p_1 \\
q_3 &= q_2 - p_2 = q_n - p_n - p_2 - p_1 \\
&\dots \\
q_{n-1} &= q_n - p_n - \sum_{i=1}^{n-2} p_i \\
q_n &= q_{n-1} - p_{n-1}
\end{aligned} \tag{5.18}$$

so that, setting  $q_n \equiv q$  and inserting in the rhs of eq. (5.17) the integral  $\int \frac{d^4 p_n}{(2\pi)^4} (2\pi)^4 \delta^{(4)}(p_1 + p_2 + \dots + p_n)$  to keep  $p_n$  while enforcing  $\sum_{i=1}^n p_i = 0$ , we obtain:

$$\begin{aligned}
\Gamma_{conf}[j_{\mathbb{O}}] &= - (N^2 - 1) \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \int \frac{d^4 p_1}{(2\pi)^4} \dots \frac{d^4 p_n}{(2\pi)^4} (2\pi)^4 \delta^{(4)}(p_1 + p_2 + \dots + p_n) \\
&\int \frac{d^4 q}{(2\pi)^4} \sum_{s_1 k_1} \dots \sum_{s_n k_n} \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(q - p_n) j_{\mathbb{O}_{s_2 k_2}}(p_1) \\
&\mathcal{D}_{s_2 k_2, s_3 k_3}^{-1}(q - p_1 - p_n) j_{\mathbb{O}_{s_3 k_3}}(p_2) \dots \mathcal{D}_{s_n k_n, s_1 k_1}^{-1}(q) j_{\mathbb{O}_{s_1 k_1}}(p_n)
\end{aligned} \tag{5.19}$$

Moreover, making the substitution  $p_i \rightarrow -p_i$ , we get:

$$\begin{aligned}
\Gamma_{conf}[j_{\mathbb{O}}] &= - (N^2 - 1) \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \int \frac{d^4 p_1}{(2\pi)^4} \dots \frac{d^4 p_n}{(2\pi)^4} (2\pi)^4 \delta^{(4)}(p_1 + p_2 + \dots + p_n) \\
&\int \frac{d^4 q}{(2\pi)^4} \sum_{s_1 k_1} \dots \sum_{s_n k_n} \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(q + p_n) j_{\mathbb{O}_{s_2 k_2}}(-p_1) \\
&\mathcal{D}_{s_2 k_2, s_3 k_3}^{-1}(q + p_1 + p_n) j_{\mathbb{O}_{s_3 k_3}}(-p_2) \dots \mathcal{D}_{s_n k_n, s_1 k_1}^{-1}(q) j_{\mathbb{O}_{s_1 k_1}}(-p_n)
\end{aligned} \tag{5.20}$$

We employ eq. (5.3) to obtain the correlators:

$$\begin{aligned}
\langle \mathbb{O}_{s_1}(p_1) \dots \mathbb{O}_{s_n}(p_n) \rangle &= \frac{N^2 - 1}{2^n} \frac{(-1)^n}{n} (2\pi)^4 \delta^{(4)}(p_1 + p_2 + \dots + p_n) \\
&\sum_{\sigma \in P_n} \int \frac{d^4 q}{(2\pi)^4} \mathcal{D}_{s_{\sigma(1)} k_{\sigma(1)}, s_{\sigma(2)} k_{\sigma(2)}}^{-1}(q + p_{\sigma(n)}) \\
&\dots \mathcal{D}_{s_{\sigma(n)} k_{\sigma(n)}, s_{\sigma(1)} k_{\sigma(1)}}^{-1}(q)
\end{aligned} \tag{5.21}$$

Explicitly, by eq. (4.3):

$$\begin{aligned}
& \left( \frac{2\Gamma(5)\Gamma(s_1+1)}{\Gamma(3)\Gamma(s_1+3)} \right) \cdots \left( \frac{2\Gamma(5)\Gamma(s_n+1)}{\Gamma(3)\Gamma(s_n+3)} \right) \\
& \langle \mathbb{O}_{s_1}(p_1) \cdots \mathbb{O}_{s_n}(p_n) \rangle \\
& = (N^2 - 1)(2\pi)^4 i^n \delta^{(4)}(p_1 + \cdots + p_n) \\
& \quad \sum_{k_1=0}^{s_1-2} \cdots \sum_{k_n=0}^{s_n-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \cdots \binom{s_n}{k_n} \binom{s_n}{k_n+2} \\
& \quad \frac{1}{n} \sum_{\sigma \in P_n} \int \frac{d^4 q}{(2\pi)^4} \frac{(p_{\sigma(1)} + q)_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{|p_{\sigma(1)} + q|^2} \frac{(p_{\sigma(1)} + p_{\sigma(2)} + q)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_{\sigma(3)}}}{|p_{\sigma(1)} + p_{\sigma(2)} + q|^2} \\
& \quad \cdots \frac{(\sum_{l=1}^{n-1} p_{\sigma(l)} + q)_+^{s_{\sigma(n-1)} - k_{\sigma(n-1)} + k_{\sigma(n)}}}{|\sum_{l=1}^{n-1} p_{\sigma(l)} + q|^2} \frac{(q)_+^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}}}{|q|^2} \tag{5.22}
\end{aligned}$$

that coincides with the computation in [1]. More generally, the generating functional in the momentum representation reads from eqs. (3.24) and (4.24) with the propagator in the momentum representation:

$$i\Box^{-1}(p) = \frac{-i}{|p|^2 + i\epsilon} \tag{5.23}$$

and correspondingly:

$$(i\partial_+)^{s-k_1+k_2} i\Box^{-1}(p) = p_+^{s-k_1+k_2} i\Box^{-1}(p) \tag{5.24}$$

## 6 Mixed 3- and 4-point conformal correlators

We employ the generating functional to calculate the 3- and 4-point correlators in the mixed balanced/unbalanced sector.

### 6.1 3-point correlators in Minkowskian space-time

The generating functional of the mixed  $\mathbb{O}$ ,  $\mathbb{S}$  and  $\bar{\mathbb{S}}$  correlators reads:

$$\Gamma_{conf}[j_{\mathbb{O}}, 0, j_{\mathbb{S}}, j_{\bar{\mathbb{S}}}] = -\frac{N^2 - 1}{2} \text{Tr} \log \left[ \mathbb{I} - 2(\mathbb{I} + \mathcal{D}^{-1}j_{\mathbb{O}})^{-1} \mathcal{D}^{-1}j_{\bar{\mathbb{S}}} (\mathbb{I} + \mathcal{D}^{-1}j_{\mathbb{O}})^{-1} \mathcal{D}^{-1}j_{\mathbb{S}} \right] \tag{6.1}$$

Only two terms contribute to the mixed 3-point correlator  $\langle \mathbb{O}_{s_1}(x) \mathbb{S}_{s_2}(y) \bar{\mathbb{S}}_{s_3}(z) \rangle$ :

$$\begin{aligned}
& \Gamma_{conf}[j_{\mathbb{O}}, 0, j_{\mathbb{S}}, j_{\bar{\mathbb{S}}}] \\
& = \frac{N^2 - 1}{2} 2 \text{Tr} \left( (\mathbb{I} + \mathcal{D}^{-1}j_{\mathbb{O}})^{-1} \mathcal{D}^{-1}j_{\bar{\mathbb{S}}} (\mathbb{I} + \mathcal{D}^{-1}j_{\mathbb{O}})^{-1} \mathcal{D}^{-1}j_{\mathbb{S}} \right) + \cdots \\
& = \frac{N^2 - 1}{2} 2 \text{Tr} \left( (\mathbb{I} - \mathcal{D}^{-1}j_{\mathbb{O}}) \mathcal{D}^{-1}j_{\bar{\mathbb{S}}} (\mathbb{I} - \mathcal{D}^{-1}j_{\mathbb{O}}) \mathcal{D}^{-1}j_{\mathbb{S}} \right) + \cdots \\
& = -\frac{N^2 - 1}{2} 2 \text{Tr} \left( \mathcal{D}^{-1}j_{\mathbb{O}} \mathcal{D}^{-1}j_{\bar{\mathbb{S}}} \mathcal{D}^{-1}j_{\mathbb{S}} \right) - \frac{N^2 - 1}{2} 2 \text{Tr} \left( \mathcal{D}^{-1}j_{\bar{\mathbb{S}}} \mathcal{D}^{-1}j_{\mathbb{O}} \mathcal{D}^{-1}j_{\mathbb{S}} \right) + \cdots \tag{6.2}
\end{aligned}$$

Hence:

$$\begin{aligned}
& \langle \mathbb{O}_{s_1}(x) \mathbb{S}_{s_2}(y) \bar{\mathbb{S}}_{s_3}(z) \rangle \\
&= \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \frac{\delta}{\delta j_{\mathbb{O}_{s_1 k_1}}(x)} \frac{\delta}{\delta j_{\mathbb{S}_{s_2 k_2}}(y)} \frac{\delta}{\delta j_{\bar{\mathbb{S}}_{s_3 k_3}}(z)} \Gamma_{conf}[j_{\mathbb{O}}, j_{\bar{\mathbb{O}}}, j_{\mathbb{S}}, j_{\bar{\mathbb{S}}}] \\
&= -\frac{N^2-1}{2} 2 \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \frac{\delta}{\delta j_{\mathbb{O}_{s_1 k_1}}(x)} \frac{\delta}{\delta j_{\mathbb{S}_{s_2 k_2}}(y)} \frac{\delta}{\delta j_{\bar{\mathbb{S}}_{s_3 k_3}}(z)} \\
&\quad \int d^4 x_1 d^4 x_2 d^4 x_3 \sum_{k'_1 s'_1} \sum_{k'_2 s'_2} \sum_{k'_3 s'_3} \left( \mathcal{D}_{s'_1 k'_1, s'_2 k'_2}^{-1}(x_1 - x_2) j_{\mathbb{O}_{s'_2 k'_2}}(x_2) \right. \\
&\quad \mathcal{D}_{s'_2 k'_2, s'_3 k'_3}^{-1}(x_2 - x_3) j_{\mathbb{S}_{s'_3 k'_3}}(x_3) \mathcal{D}_{s'_3 k'_3, s'_1 k'_1}^{-1}(x_3 - x_1) j_{\bar{\mathbb{S}}_{s'_1 k'_1}}(x_1) \\
&\quad + \mathcal{D}_{s'_1 k'_1, s'_2 k'_2}^{-1}(x_1 - x_2) j_{\bar{\mathbb{S}}_{s'_2 k'_2}}(x_2) \mathcal{D}_{s'_2 k'_2, s'_3 k'_3}^{-1}(x_2 - x_3) j_{\mathbb{O}_{s'_3 k'_3}}(x_3) \\
&\quad \left. \mathcal{D}_{s'_3 k'_3, s'_1 k'_1}^{-1}(x_3 - x_1) j_{\mathbb{S}_{s'_1 k'_1}}(x_1) \right) \\
&= -(N^2-1) \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \left( \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x-y) \mathcal{D}_{s_2 k_2, s_3 k_3}^{-1}(y-z) \mathcal{D}_{s_3 k_3, s_1 k_1}^{-1}(z-x) \right. \\
&\quad \left. + \mathcal{D}_{s_2 k_2, s_3 k_3}^{-1}(y-z) \mathcal{D}_{s_3 k_3, s_1 k_1}^{-1}(z-x) \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x-y) \right) \\
&= -2(N^2-1) \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \mathcal{D}_{s_1 k_1, s_2 k_2}^{-1}(x-y) \mathcal{D}_{s_2 k_2, s_3 k_3}^{-1}(y-z) \mathcal{D}_{s_3 k_3, s_1 k_1}^{-1}(z-x) \quad (6.3)
\end{aligned}$$

Employing eq. (4.3), we obtain:

$$\begin{aligned}
\langle \mathbb{O}_{s_1}(x) \mathbb{S}_{s_2}(y) \bar{\mathbb{S}}_{s_3}(z) \rangle &= -\frac{1}{(4\pi^2)^3} 2 \left( \frac{2}{4!} \right)^3 \frac{N^2-1}{8} i^{s_1+s_2+s_3} 2^{s_1+s_2+s_3} \\
&\quad (s_1+1)(s_1+2)(s_2+1)(s_2+2)(s_3+1)(s_3+2) \\
&\quad \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} \\
&\quad (s_1 - k_1 + k_2)! (s_2 - k_2 + k_3)! (s_3 - k_3 + k_1)! \\
&\quad \frac{(x-y)_+^{s_1-k_1+k_2}}{(|x-y|^2)^{s_1+1-k_1+k_2}} \frac{(y-z)_+^{s_2-k_2+k_3}}{(|y-z|^2)^{s_2+1-k_2+k_3}} \frac{(z-x)_+^{s_3-k_3+k_1}}{(|z-x|^2)^{s_3+1-k_3+k_1}} \\
&\hspace{15em} (6.4)
\end{aligned}$$

that agrees with [1]. We also infer from the generating functional that [1]:

$$\langle \tilde{\mathbb{O}}_{s_1}(x) \mathbb{S}_{s_2}(y) \bar{\mathbb{S}}_{s_3}(z) \rangle = 0 \quad (6.5)$$

Indeed:

$$\Gamma_{conf}[0, j_{\bar{\mathbb{O}}}, j_{\mathbb{S}}, j_{\bar{\mathbb{S}}}] = -\frac{N^2-1}{2} \text{Tr} \log \left[ \mathbb{I} - 2(\mathbb{I} - \mathcal{D}^{-1} j_{\bar{\mathbb{O}}})^{-1} \mathcal{D}^{-1} j_{\mathbb{S}} \left( \mathbb{I} + \mathcal{D}^{-1} j_{\bar{\mathbb{O}}} \right)^{-1} \mathcal{D}^{-1} j_{\bar{\mathbb{S}}} \right] \quad (6.6)$$

contributes two terms linear in  $j_{\bar{\mathbb{O}}}$  with opposite sign that cancel each other.

## 6.2 4-point correlators in Minkowskian space-time

Only three terms contribute to the mixed 4-point correlator  $\langle \mathbb{O}_{s_1}(x_1) \mathbb{O}_{s_2}(x_2) \mathbb{S}_{s_3}(x_3) \bar{\mathbb{S}}_{s_4}(x_4) \rangle$ :

$$\begin{aligned}
& \Gamma_{conf}[j_{\mathbb{O}}, 0, j_{\mathbb{S}}, j_{\bar{\mathbb{S}}}] \\
&= \frac{N^2 - 1}{2} 2 \operatorname{Tr} \left( (\mathbb{I} - \mathcal{D}^{-1} j_{\mathbb{O}} + \mathcal{D}^{-1} j_{\mathbb{O}} \mathcal{D}^{-1} j_{\mathbb{O}}) \mathcal{D}^{-1} j_{\bar{\mathbb{S}}} (\mathbb{I} - \mathcal{D}^{-1} j_{\mathbb{O}} + \mathcal{D}^{-1} j_{\mathbb{O}} \mathcal{D}^{-1} j_{\mathbb{O}}) \mathcal{D}^{-1} j_{\mathbb{S}} \right) + \dots \\
&= \frac{N^2 - 1}{2} 2 \operatorname{Tr} (\mathcal{D}^{-1} j_{\mathbb{O}} \mathcal{D}^{-1} j_{\bar{\mathbb{S}}} \mathcal{D}^{-1} j_{\mathbb{O}} \mathcal{D}^{-1} j_{\mathbb{S}}) + \frac{N^2 - 1}{2} 2 \operatorname{Tr} (\mathcal{D}^{-1} j_{\mathbb{O}} \mathcal{D}^{-1} j_{\mathbb{O}} \mathcal{D}^{-1} j_{\bar{\mathbb{S}}} \mathcal{D}^{-1} j_{\mathbb{S}}) \\
&\quad + \frac{N^2 - 1}{2} 2 \operatorname{Tr} (\mathcal{D}^{-1} j_{\bar{\mathbb{S}}} \mathcal{D}^{-1} j_{\mathbb{O}} \mathcal{D}^{-1} j_{\mathbb{O}} \mathcal{D}^{-1} j_{\mathbb{S}}) + \dots \tag{6.7}
\end{aligned}$$

Hence:

$$\begin{aligned}
& \langle \mathbb{O}_{s_1}(x_1) \mathbb{O}_{s_2}(x_2) \mathbb{S}_{s_3}(x_3) \bar{\mathbb{S}}_{s_4}(x_4) \rangle \\
&= \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \frac{\delta}{\delta j_{\mathbb{O}_{s_1 k_1}}(x_1)} \frac{\delta}{\delta j_{\mathbb{O}_{s_2 k_2}}(x_2)} \frac{\delta}{\delta j_{\mathbb{S}_{s_3 k_3}}(x_3)} \frac{\delta}{\delta j_{\bar{\mathbb{S}}_{s_4 k_4}}(x_4)} \Gamma_{conf}[j_{\mathbb{O}}, j_{\bar{\mathbb{O}}}, j_{\mathbb{S}}, j_{\bar{\mathbb{S}}}] \\
&= \frac{N^2 - 1}{2} 2 \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \frac{\delta}{\delta j_{\mathbb{O}_{s_1 k_1}}(x_1)} \frac{\delta}{\delta j_{\mathbb{O}_{s_2 k_2}}(x_2)} \frac{\delta}{\delta j_{\mathbb{S}_{s_3 k_3}}(x_3)} \frac{\delta}{\delta j_{\bar{\mathbb{S}}_{s_4 k_4}}(x_4)} \\
&\quad \int d^4 y_1 d^4 y_2 d^4 y_3 d^4 y_4 \sum_{k'_1 s'_1} \sum_{k'_2 s'_2} \sum_{k'_3 s'_3} \sum_{k'_4 s'_4} \\
&\quad \left( \mathcal{D}_{s'_1 k'_1, s'_2 k'_2}^{-1}(y_1 - y_2) j_{\mathbb{O}_{s'_2 k'_2}}(y_2) \mathcal{D}_{s'_2 k'_2, s'_3 k'_3}^{-1}(y_2 - y_3) j_{\bar{\mathbb{S}}_{s'_3 k'_3}}(y_3) \right. \\
&\quad \quad \mathcal{D}_{s'_3 k'_3, s'_4 k'_4}^{-1}(y_3 - y_4) j_{\mathbb{O}_{s'_4 k'_4}}(y_4) \mathcal{D}_{s'_4 k'_4, s'_1 k'_1}^{-1}(y_4 - y_1) j_{\bar{\mathbb{S}}_{s'_1 k'_1}}(y_1) \\
&\quad + \mathcal{D}_{s'_1 k'_1, s'_2 k'_2}^{-1}(y_1 - y_2) j_{\mathbb{O}_{s'_2 k'_2}}(y_2) \mathcal{D}_{s'_2 k'_2, s'_3 k'_3}^{-1}(y_2 - y_3) j_{\mathbb{O}_{s'_3 k'_3}}(y_3) \\
&\quad \quad \mathcal{D}_{s'_3 k'_3, s'_4 k'_4}^{-1}(y_3 - y_4) j_{\bar{\mathbb{S}}_{s'_4 k'_4}}(y_4) \mathcal{D}_{s'_4 k'_4, s'_1 k'_1}^{-1}(y_4 - y_1) j_{\bar{\mathbb{S}}_{s'_1 k'_1}}(y_1) \\
&\quad + \mathcal{D}_{s'_1 k'_1, s'_2 k'_2}^{-1}(y_1 - y_2) j_{\mathbb{O}_{s'_2 k'_2}}(y_2) \mathcal{D}_{s'_2 k'_2, s'_3 k'_3}^{-1}(y_2 - y_3) j_{\mathbb{O}_{s'_3 k'_3}}(y_3) \\
&\quad \quad \left. \mathcal{D}_{s'_3 k'_3, s'_4 k'_4}^{-1}(y_3 - y_4) j_{\bar{\mathbb{S}}_{s'_4 k'_4}}(y_4) \mathcal{D}_{s'_4 k'_4, s'_1 k'_1}^{-1}(y_4 - y_1) j_{\bar{\mathbb{S}}_{s'_1 k'_1}}(y_1) \right) \tag{6.8}
\end{aligned}$$

Performing the functional derivatives and employing the symmetry properties of the Gegenbauer polynomials (appendix B), we get:

$$\begin{aligned}
& \langle \mathbb{O}_{s_1}(x_1) \mathbb{O}_{s_2}(x_2) \mathbb{S}_{s_3}(x_3) \bar{\mathbb{S}}_{s_4}(x_4) \rangle \\
&= \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \frac{\delta}{\delta j_{\mathbb{O}_{s_1 k_1}}(x_1)} \frac{\delta}{\delta j_{\mathbb{O}_{s_2 k_2}}(x_2)} \frac{\delta}{\delta j_{\mathbb{S}_{s_3 k_3}}(x_3)} \frac{\delta}{\delta j_{\bar{\mathbb{S}}_{s_4 k_4}}(x_4)} \Gamma_{conf}[j_{\mathbb{O}}, j_{\bar{\mathbb{O}}}, j_{\mathbb{S}}, j_{\bar{\mathbb{S}}}] \\
&= (N^2 - 1) \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \sum_{\sigma \in P_2} \\
&\quad \left( \mathcal{D}_{s_3 k_3, s_{\sigma(1)} k_{\sigma(1)}}^{-1}(x_3 - x_{\sigma(1)}) \mathcal{D}_{s_{\sigma(1)} k_{\sigma(1)}, s_4 k_4}^{-1}(x_{\sigma(1)} - x_4) \right. \\
&\quad \mathcal{D}_{s_4 k_4, s_{\sigma(2)} k_{\sigma(2)}}^{-1}(x_4 - x_{\sigma(2)}) \mathcal{D}_{s_{\sigma(2)} k_{\sigma(2)}, s_3 k_3}^{-1}(x_{\sigma(2)} - x_3) \\
&\quad + 2 \mathcal{D}_{s_3 k_3, s_{\sigma(1)} k_{\sigma(1)}}^{-1}(x_3 - x_{\sigma(1)}) \mathcal{D}_{s_{\sigma(1)} k_{\sigma(1)}, s_{\sigma(2)} k_{\sigma(2)}}^{-1}(x_{\sigma(1)} - x_{\sigma(2)}) \\
&\quad \left. \mathcal{D}_{s_{\sigma(2)} k_{\sigma(2)}, s_4 k_4}^{-1}(x_{\sigma(2)} - x_4) \mathcal{D}_{s_4 k_4, s_3 k_3}^{-1}(x_4 - x_3) \right) \tag{6.9}
\end{aligned}$$

Explicitly:

$$\begin{aligned}
& 2^4 \frac{\Gamma(5)\Gamma(s_1+1)}{\Gamma(3)\Gamma(s_1+3)} \frac{\Gamma(5)\Gamma(s_2+1)}{\Gamma(3)\Gamma(s_2+3)} \frac{\Gamma(5)\Gamma(s_3+1)}{\Gamma(3)\Gamma(s_3+3)} \frac{\Gamma(5)\Gamma(s_4+1)}{\Gamma(3)\Gamma(s_4+3)} \\
& \langle \mathbb{O}_{s_1}(x_1) \mathbb{O}_{s_2}(x_2) \mathbb{S}_{s_3}(x_3) \bar{\mathbb{S}}_{s_4}(x_4) \rangle \\
&= (N^2 - 1) \frac{1}{(4\pi^2)^4} 2^{\sum_{i=1}^4 s_i} i^{\sum_{i=1}^4 s_i} \\
&\quad \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} \binom{s_4}{k_4} \binom{s_4}{k_4+2} \\
&\quad \sum_{\sigma \in P_2} \left[ (s_3 - k_3 + k_{\sigma(1)})! (s_{\sigma(1)} - k_{\sigma(1)} + k_4)! (s_4 - k_4 + k_{\sigma(2)})! (s_{\sigma(2)} - k_{\sigma(2)} + k_3)! \right. \\
&\quad \frac{(x_3 - x_{\sigma(1)})_+^{s_3 - k_3 + k_{\sigma(1)}}}{(|x_3 - x_{\sigma(1)}|^2)^{s_3 - k_3 + k_{\sigma(1)} + 1}} \frac{(x_{\sigma(1)} - x_4)_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_4}}{(|x_{\sigma(1)} - x_4|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_4 + 1}} \\
&\quad \frac{(x_4 - x_{\sigma(2)})_+^{s_4 - k_4 + k_{\sigma(2)}}}{(|x_4 - x_{\sigma(2)}|^2)^{s_4 - k_4 + k_{\sigma(2)} + 1}} \frac{(x_{\sigma(2)} - x_3)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_3}}{(|x_4 - x_1|^2)^{s_4 - k_4 + k_1 + 1}} \\
&\quad + 2 (s_3 - k_3 + k_{\sigma(1)})! (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! (s_{\sigma(2)} - k_{\sigma(2)} + k_4)! (s_4 - k_4 + k_3)! \\
&\quad \frac{(x_3 - x_{\sigma(1)})_+^{s_3 - k_3 + k_{\sigma(1)}}}{(|x_3 - x_{\sigma(1)}|^2)^{s_3 - k_3 + k_{\sigma(1)} + 1}} \frac{(x_{\sigma(1)} - x_{\sigma(2)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{(|x_{\sigma(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \\
&\quad \left. \frac{(x_{\sigma(2)} - x_4)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_4}}{(|x_{\sigma(2)} - x_4|^2)^{s_{\sigma(2)} - k_{\sigma(2)} + k_4 + 1}} \frac{(x_4 - x_3)_+^{s_4 - k_4 + k_3}}{(|x_4 - x_3|^2)^{s_4 - k_4 + k_3 + 1}} \right] \tag{6.10}
\end{aligned}$$

that in the momentum representation reads:

$$\begin{aligned}
& 2^4 \frac{\Gamma(5)\Gamma(s_1+1)}{\Gamma(3)\Gamma(s_1+3)} \frac{\Gamma(5)\Gamma(s_2+1)}{\Gamma(3)\Gamma(s_2+3)} \frac{\Gamma(5)\Gamma(s_3+1)}{\Gamma(3)\Gamma(s_3+3)} \frac{\Gamma(5)\Gamma(s_4+1)}{\Gamma(3)\Gamma(s_4+3)} \\
& \langle \mathbb{O}_{s_1}(p_1) \mathbb{O}_{s_2}(p_2) \mathbb{S}_{s_3}(p_3) \bar{\mathbb{S}}_{s_4}(p_4) \rangle \\
& = (N^2 - 1) (2\pi)^4 i^4 \delta^{(4)}(p_1 + p_2 + p_3 + p_4) \\
& \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} \binom{s_4}{k_4} \binom{s_4}{k_4+2} \\
& \sum_{\sigma \in P_2} \int \frac{d^4 q}{(2\pi)^4} \left[ \frac{(p_3 + q)_+^{s_3 - k_3 + k_{\sigma(1)}}}{|p_3 + q|^2 + i\epsilon} \frac{(p_3 + p_{\sigma(1)} + q)_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_4}}{|p_3 + p_{\sigma(1)} + q|^2 + i\epsilon} \right. \\
& \frac{(p_3 + p_{\sigma(1)} + p_4 + q)_+^{s_{\sigma(4)} - k_{\sigma(4)} + k_{\sigma(2)}}}{|p_3 + p_{\sigma(1)} + p_4 + q|^2} \frac{(q)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_3}}{|q|^2 + i\epsilon} \\
& \left. + 2 \frac{(p_3 + q)_+^{s_3 - k_3 + k_{\sigma(1)}}}{|p_3 + q|^2 + i\epsilon} \frac{(p_3 + p_{\sigma(1)} + q)_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{|p_3 + p_{\sigma(1)} + q|^2 + i\epsilon} \right. \\
& \left. \frac{(p_3 + p_{\sigma(1)} + p_{\sigma(2)} + q)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_4}}{|p_3 + p_{\sigma(1)} + p_{\sigma(2)} + q|^2 + i\epsilon} \frac{(q)_+^{s_4 - k_4 + k_3}}{|q|^2 + i\epsilon} \right] \tag{6.11}
\end{aligned}$$

Similarly, we obtain:

$$\begin{aligned}
& \langle \tilde{\mathbb{O}}_{s_1}(x_1) \tilde{\mathbb{O}}_{s_2}(x_2) \mathbb{S}_{s_3}(x_3) \bar{\mathbb{S}}_{s_4}(x_4) \rangle \\
& = \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \frac{\delta}{\delta j_{\tilde{\mathbb{O}}_{s_1 k_1}}(x_1)} \frac{\delta}{\delta j_{\tilde{\mathbb{O}}_{s_2 k_2}}(x_2)} \frac{\delta}{\delta j_{\mathbb{S}_{s_3 k_3}}(x_3)} \frac{\delta}{\delta j_{\bar{\mathbb{S}}_{s_4 k_4}}(x_4)} \Gamma_{conf}[j_{\mathbb{O}}, j_{\tilde{\mathbb{O}}}, j_{\mathbb{S}}, j_{\bar{\mathbb{S}}}] \\
& = (N^2 - 1) \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \sum_{\sigma \in P_2} \\
& \left( - \mathcal{D}_{s_3 k_3, s_{\sigma(1)} k_{\sigma(1)}}^{-1}(x_3 - x_{\sigma(1)}) \mathcal{D}_{s_{\sigma(1)} k_{\sigma(1)}, s_4 k_4}^{-1}(x_{\sigma(1)} - x_4) \right. \\
& \mathcal{D}_{s_4 k_4, s_{\sigma(2)} k_{\sigma(2)}}^{-1}(x_4 - x_{\sigma(2)}) \mathcal{D}_{s_{\sigma(2)} k_{\sigma(2)}, s_3 k_3}^{-1}(x_{\sigma(2)} - x_3) \\
& \left. + 2 \mathcal{D}_{s_3 k_3, s_{\sigma(1)} k_{\sigma(1)}}^{-1}(x_3 - x_{\sigma(1)}) \mathcal{D}_{s_{\sigma(1)} k_{\sigma(1)}, s_{\sigma(2)} k_{\sigma(2)}}^{-1}(x_{\sigma(1)} - x_{\sigma(2)}) \right. \\
& \left. \mathcal{D}_{s_{\sigma(2)} k_{\sigma(2)}, s_4 k_4}^{-1}(x_{\sigma(2)} - x_4) \mathcal{D}_{s_4 k_4, s_3 k_3}^{-1}(x_4 - x_3) \right) \tag{6.12}
\end{aligned}$$

Explicitly, in the coordinate:

$$\begin{aligned}
& 2^4 \frac{\Gamma(5)\Gamma(s_1+1)}{\Gamma(3)\Gamma(s_1+3)} \frac{\Gamma(5)\Gamma(s_2+1)}{\Gamma(3)\Gamma(s_2+3)} \frac{\Gamma(5)\Gamma(s_3+1)}{\Gamma(3)\Gamma(s_3+3)} \frac{\Gamma(5)\Gamma(s_4+1)}{\Gamma(3)\Gamma(s_4+3)} \\
& \langle \tilde{\mathcal{O}}_{s_1}(x_1) \tilde{\mathcal{O}}_{s_2}(x_2) \mathbb{S}_{s_3}(x_3) \bar{\mathbb{S}}_{s_4}(x_4) \rangle \\
& = (N^2 - 1) \frac{1}{(4\pi^2)^4} 2^{\sum_{i=1}^4 s_i} i^{\sum_{i=1}^4 s_i} \\
& \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} \binom{s_4}{k_4} \binom{s_4}{k_4+2} \\
& \sum_{\sigma \in P_2} \left[ - (s_3 - k_3 + k_{\sigma(1)})! (s_{\sigma(1)} - k_{\sigma(1)} + k_4)! (s_4 - k_4 + k_{\sigma(2)})! (s_{\sigma(2)} - k_{\sigma(2)} + k_3)! \right. \\
& \frac{(x_3 - x_{\sigma(1)})_+^{s_3 - k_3 + k_{\sigma(1)}}}{(|x_3 - x_{\sigma(1)}|^2)^{s_3 - k_3 + k_{\sigma(1)} + 1}} \frac{(x_{\sigma(1)} - x_4)_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_4}}{(|x_{\sigma(1)} - x_4|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_4 + 1}} \\
& \frac{(x_4 - x_{\sigma(2)})_+^{s_4 - k_4 + k_{\sigma(2)}}}{(|x_4 - x_{\sigma(2)}|^2)^{s_4 - k_4 + k_{\sigma(2)} + 1}} \frac{(x_{\sigma(2)} - x_3)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_3}}{(|x_4 - x_1|^2)^{s_4 - k_4 + k_1 + 1}} \\
& + 2(s_3 - k_3 + k_{\sigma(1)})! (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! (s_{\sigma(2)} - k_{\sigma(2)} + k_4)! (s_4 - k_4 + k_3)! \\
& \frac{(x_3 - x_{\sigma(1)})_+^{s_3 - k_3 + k_{\sigma(1)}}}{(|x_3 - x_{\sigma(1)}|^2)^{s_3 - k_3 + k_{\sigma(1)} + 1}} \frac{(x_{\sigma(1)} - x_{\sigma(2)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{(|x_{\sigma(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \\
& \left. \frac{(x_{\sigma(2)} - x_4)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_4}}{(|x_{\sigma(2)} - x_4|^2)^{s_{\sigma(2)} - k_{\sigma(2)} + k_4 + 1}} \frac{(x_4 - x_3)_+^{s_4 - k_4 + k_3}}{(|x_4 - x_3|^2)^{s_4 - k_4 + k_3 + 1}} \right] \tag{6.13}
\end{aligned}$$

and momentum representations:

$$\begin{aligned}
& 2^4 \frac{\Gamma(5)\Gamma(s_1+1)}{\Gamma(3)\Gamma(s_1+3)} \frac{\Gamma(5)\Gamma(s_2+1)}{\Gamma(3)\Gamma(s_2+3)} \frac{\Gamma(5)\Gamma(s_3+1)}{\Gamma(3)\Gamma(s_3+3)} \frac{\Gamma(5)\Gamma(s_4+1)}{\Gamma(3)\Gamma(s_4+3)} \\
& \langle \tilde{\mathcal{O}}_{s_1}(p_1) \tilde{\mathcal{O}}_{s_2}(p_2) \mathbb{S}_{s_3}(p_3) \bar{\mathbb{S}}_{s_4}(p_4) \rangle \\
& = (N^2 - 1) (2\pi)^4 i^4 \delta^{(4)}(p_1 + p_2 + p_3 + p_4) \\
& \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} \binom{s_4}{k_4} \binom{s_4}{k_4+2} \\
& \sum_{\sigma \in P_2} \int \frac{d^4 q}{(2\pi)^4} \left[ - \frac{(p_3 + q)_+^{s_3 - k_3 + k_{\sigma(1)}}}{|p_3 + q|^2 + i\epsilon} \frac{(p_3 + p_{\sigma(1)} + q)_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_4}}{|p_3 + p_{\sigma(1)} + q|^2 + i\epsilon} \right. \\
& \frac{(p_3 + p_{\sigma(1)} + p_4 + q)_+^{s_{\sigma(4)} - k_{\sigma(4)} + k_{\sigma(2)}}}{|p_3 + p_{\sigma(1)} + p_4 + q|^2} \frac{(q)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_3}}{|q|^2 + i\epsilon} \\
& + 2 \frac{(p_3 + q)_+^{s_3 - k_3 + k_{\sigma(1)}}}{|p_3 + q|^2 + i\epsilon} \frac{(p_3 + p_{\sigma(1)} + q)_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{|p_3 + p_{\sigma(1)} + q|^2 + i\epsilon} \\
& \left. \frac{(p_3 + p_{\sigma(1)} + p_{\sigma(2)} + q)_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_4}}{|p_3 + p_{\sigma(1)} + p_{\sigma(2)} + q|^2 + i\epsilon} \frac{(q)_+^{s_4 - k_4 + k_3}}{|q|^2 + i\epsilon} \right] \tag{6.14}
\end{aligned}$$



### 6.3 4-point correlators in Euclidean space-time

In Euclidean space-time we obtain as well:

$$\begin{aligned}
& 2^4 \frac{\Gamma(5)\Gamma(s_1+1)}{\Gamma(3)\Gamma(s_1+3)} \frac{\Gamma(5)\Gamma(s_2+1)}{\Gamma(3)\Gamma(s_2+3)} \frac{\Gamma(5)\Gamma(s_3+1)}{\Gamma(3)\Gamma(s_3+3)} \frac{\Gamma(5)\Gamma(s_4+1)}{\Gamma(3)\Gamma(s_4+3)} \\
& \langle \mathbb{O}_{s_1}^E(x_1) \mathbb{O}_{s_2}^E(x_2) \mathbb{S}_{s_3}^E(x_3) \bar{\mathbb{S}}_{s_4}^E(x_4) \rangle \\
& = (N^2 - 1) \frac{1}{(4\pi^2)^4} 2^{\sum_{l=1}^4 s_l} (-1)^{\sum_{l=1}^4 s_l} \\
& \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} \binom{s_4}{k_4} \binom{s_4}{k_4+2} \\
& \sum_{\sigma \in P_2} \left[ (s_3 - k_3 + k_{\sigma(1)})! (s_{\sigma(1)} - k_{\sigma(1)} + k_4)! (s_4 - k_4 + k_{\sigma(2)})! (s_{\sigma(2)} - k_{\sigma(2)} + k_3)! \right. \\
& \frac{(x_3 - x_{\sigma(1)})_z^{s_3 - k_3 + k_{\sigma(1)}}}{((x_3 - x_{\sigma(1)})^2)^{s_3 - k_3 + k_{\sigma(1)} + 1}} \frac{(x_{\sigma(1)} - x_4)_z^{s_{\sigma(1)} - k_{\sigma(1)} + k_4}}{((x_{\sigma(1)} - x_4)^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_4 + 1}} \\
& \frac{(x_4 - x_{\sigma(2)})_z^{s_4 - k_4 + k_{\sigma(2)}}}{((x_4 - x_{\sigma(2)})^2)^{s_4 - k_4 + k_{\sigma(2)} + 1}} \frac{(x_{\sigma(2)} - x_3)_z^{s_{\sigma(2)} - k_{\sigma(2)} + k_3}}{((x_4 - x_1)^2)^{s_4 - k_4 + k_1 + 1}} \\
& + 2(s_3 - k_3 + k_{\sigma(1)})! (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! (s_{\sigma(2)} - k_{\sigma(2)} + k_4)! (s_4 - k_4 + k_3)! \\
& \frac{(x_3 - x_{\sigma(1)})_z^{s_3 - k_3 + k_{\sigma(1)}}}{((x_3 - x_{\sigma(1)})^2)^{s_3 - k_3 + k_{\sigma(1)} + 1}} \frac{(x_{\sigma(1)} - x_{\sigma(2)})_z^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{((x_{\sigma(1)} - x_{\sigma(2)})^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \\
& \left. \frac{(x_{\sigma(2)} - x_4)_z^{s_{\sigma(2)} - k_{\sigma(2)} + k_4}}{((x_{\sigma(2)} - x_4)^2)^{s_{\sigma(2)} - k_{\sigma(2)} + k_4 + 1}} \frac{(x_4 - x_3)_z^{s_4 - k_4 + k_3}}{((x_4 - x_3)^2)^{s_4 - k_4 + k_3 + 1}} \right] \tag{6.15}
\end{aligned}$$

and:

$$\begin{aligned}
& 2^4 \frac{\Gamma(5)\Gamma(s_1+1)}{\Gamma(3)\Gamma(s_1+3)} \frac{\Gamma(5)\Gamma(s_2+1)}{\Gamma(3)\Gamma(s_2+3)} \frac{\Gamma(5)\Gamma(s_3+1)}{\Gamma(3)\Gamma(s_3+3)} \frac{\Gamma(5)\Gamma(s_4+1)}{\Gamma(3)\Gamma(s_4+3)} \\
& \langle \tilde{\mathcal{O}}_{s_1}^E(x_1) \tilde{\mathcal{O}}_{s_2}^E(x_2) \mathbb{S}_{s_3}^E(x_3) \bar{\mathbb{S}}_{s_4}^E(x_4) \rangle \\
& = (N^2 - 1) \frac{1}{(4\pi^2)^4} 2^{\sum_{i=1}^4 s_i} (-1)^{\sum_{i=1}^4 s_i} \\
& \sum_{k_1=0}^{s_1-2} \sum_{k_2=0}^{s_2-2} \sum_{k_3=0}^{s_3-2} \sum_{k_4=0}^{s_4-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} \binom{s_4}{k_4} \binom{s_4}{k_4+2} \\
& \sum_{\sigma \in P_2} \left[ - (s_3 - k_3 + k_{\sigma(1)})! (s_{\sigma(1)} - k_{\sigma(1)} + k_4)! (s_4 - k_4 + k_{\sigma(2)})! (s_{\sigma(2)} - k_{\sigma(2)} + k_3)! \right. \\
& \frac{(x_3 - x_{\sigma(1)})_z^{s_3 - k_3 + k_{\sigma(1)}}}{((x_3 - x_{\sigma(1)})^2)^{s_3 - k_3 + k_{\sigma(1)} + 1}} \frac{(x_{\sigma(1)} - x_4)_z^{s_{\sigma(1)} - k_{\sigma(1)} + k_4}}{((x_{\sigma(1)} - x_4)^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_4 + 1}} \\
& \frac{(x_4 - x_{\sigma(2)})_z^{s_4 - k_4 + k_{\sigma(2)}}}{((x_4 - x_{\sigma(2)})^2)^{s_4 - k_4 + k_{\sigma(2)} + 1}} \frac{(x_{\sigma(2)} - x_3)_z^{s_{\sigma(2)} - k_{\sigma(2)} + k_3}}{((x_4 - x_1)^2)^{s_4 - k_4 + k_1 + 1}} \\
& + 2 (s_3 - k_3 + k_{\sigma(1)})! (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! (s_{\sigma(2)} - k_{\sigma(2)} + k_4)! (s_4 - k_4 + k_3)! \\
& \frac{(x_3 - x_{\sigma(1)})_z^{s_3 - k_3 + k_{\sigma(1)}}}{((x_3 - x_{\sigma(1)})^2)^{s_3 - k_3 + k_{\sigma(1)} + 1}} \frac{(x_{\sigma(1)} - x_{\sigma(2)})_z^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{((x_{\sigma(1)} - x_{\sigma(2)})^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \\
& \left. \frac{(x_{\sigma(2)} - x_4)_z^{s_{\sigma(2)} - k_{\sigma(2)} + k_4}}{((x_{\sigma(2)} - x_4)^2)^{s_{\sigma(2)} - k_{\sigma(2)} + k_4 + 1}} \frac{(x_4 - x_3)_z^{s_4 - k_4 + k_3}}{((x_4 - x_3)^2)^{s_4 - k_4 + k_3 + 1}} \right] \tag{6.16}
\end{aligned}$$

## 7 UV asymptotics of Euclidean $n$ -point correlators

### 7.1 Operator mixing

The Euclidean  $n$ -point correlator of local renormalized operators  $\mathcal{O}_i(x)$  that mix under renormalization:

$$\langle \mathcal{O}_{k_1}(x_1) \dots \mathcal{O}_{k_n}(x_n) \rangle = G_{k_1 \dots k_n}^{(n)}(x_1, \dots, x_n; \mu, g(\mu)) \tag{7.1}$$

satisfies the Callan-Symanzik equation:

$$\begin{aligned}
& \left( \sum_{\alpha=1}^n x_\alpha \cdot \frac{\partial}{\partial x_\alpha} + \beta(g) \frac{\partial}{\partial g} + \sum_{\alpha=1}^n D_{\mathcal{O}_\alpha} \right) G_{k_1 \dots k_n}^{(n)} + \\
& \sum_a \left( \gamma_{k_1 a}(g) G_{a k_2 \dots k_n}^{(n)} + \gamma_{k_2 a}(g) G_{k_1 a k_3 \dots k_n}^{(n)} \dots + \gamma_{k_n a}(g) G_{k_1 \dots a}^{(n)} \right) = 0
\end{aligned} \tag{7.2}$$

where  $D_{\mathcal{O}_i}$  is the canonical dimension of  $\mathcal{O}_i(x)$  and  $\gamma_{ij}(g)$  the matrix of the anomalous dimensions. Its solution satisfies:

$$\begin{aligned} & G_{k_1 \dots k_n}^{(n)}(\lambda x_1, \dots, \lambda x_n; \mu, g(\mu)) \\ &= \sum_{j_1 \dots j_n} Z_{k_1 j_1}(\lambda) \dots Z_{k_n j_n}(\lambda) \lambda^{-\sum_{i=1}^n D_{\mathcal{O}_i}} G_{j_1 \dots j_n}^{(n)}(x_1, \dots, x_n; \mu, g\left(\frac{\mu}{\lambda}\right)) \end{aligned} \quad (7.3)$$

where in matrix notation:

$$\left( \frac{\partial}{\partial g} + \frac{\gamma(g)}{\beta(g)} \right) Z(\lambda) = 0 \quad (7.4)$$

with  $g \equiv g(\mu)$  and:

$$Z(\lambda) = P \exp \left( \int_{g(\mu)}^{g(\frac{\mu}{\lambda})} \frac{\gamma(g')}{\beta(g')} dg' \right) \quad (7.5)$$

Hence, the corresponding UV asymptotics as  $\lambda \rightarrow 0$  reads:

$$\begin{aligned} & G_{k_1 \dots k_n}^{(n)}(\lambda x_1, \dots, \lambda x_n; \mu, g(\mu)) \\ & \sim \sum_{j_1 \dots j_n} Z_{k_1 j_1}(\lambda) \dots Z_{k_n j_n}(\lambda) \lambda^{-\sum_{i=1}^n D_{\mathcal{O}_i}} G_{conf\ j_1 \dots j_n}^{(n)}(x_1, \dots, x_n) \end{aligned} \quad (7.6)$$

provided that  $G_{conf\ j_1 \dots j_n}^{(n)}(x_1, \dots, x_n)$  – the conformal correlators to the lowest order of perturbation theory – do not vanish.

The evaluation of the asymptotics above involves the estimate of each term in eq. (7.6) that in turn involves the computation of the path-ordered exponential defining  $Z_{ij}(\lambda)$  in eq. (7.5). Both the computations are technically challenging, even in the special case where  $\gamma(g)$  is triangular so that the expansion of the path-ordered exponential in eq. (7.5) terminates to a finite order [13].

Therefore, it is of the utmost importance to establish whether a renormalization scheme exists where  $Z(\lambda)$  is diagonalizable to all perturbative orders. Indeed, in such a scheme the sum in eq. (7.6) would reduce to just one term and the path-ordered exponential to just the ordinary exponential, as in the multiplicatively renormalizable case.

## 7.2 Canonical nonresonant diagonal renormalization scheme

The question above has been addressed on the basis of the differential-geometric interpretation [9] of the operator mixing in massless QCD-like theories to all orders of perturbation theory that we summarize in the following. We interpret a finite change of the basis of renormalized operators in matrix notation:

$$\mathcal{O}'(x) = S(g)\mathcal{O}(x) \quad (7.7)$$

as a real-analytic invertible gauge transformation  $S(g)$  that depends on  $g$  [9]. Under the action of the aforementioned gauge transformation, the matrix:

$$A(g) = -\frac{\gamma(g)}{\beta(g)} = \frac{1}{g} \left( \frac{\gamma_0}{\beta_0} + \dots \right) \quad (7.8)$$

associated to the differential equation for  $Z(\lambda)$ :

$$\left(\frac{\partial}{\partial g} - A(g)\right)Z(\lambda) = 0 \quad (7.9)$$

defines a connection  $A(g)$  with a regular singularity at  $g = 0$  that transform as:

$$A'(g) = S(g)A(g)S^{-1}(g) + \frac{\partial S(g)}{\partial g}S^{-1}(g) \quad (7.10)$$

with:

$$\mathcal{D} = \frac{\partial}{\partial g} - A(g) \quad (7.11)$$

the corresponding covariant derivative. Consequently,  $Z(\lambda)$  is interpreted as a Wilson line that transforms as:

$$Z'(\lambda) = S(g(\mu))Z(\lambda)S^{-1}\left(g\left(\frac{\mu}{\lambda}\right)\right) \quad (7.12)$$

for the gauge transformation  $S(g)$ . Moreover, everything that we have mentioned also applies by allowing the coupling  $g$  to be complex valued. It follows from the Poincaré-Dulac theorem that [9], if any two eigenvalues  $\lambda_1, \lambda_2, \dots$  of the matrix  $\frac{\gamma_0}{\beta_0}$ , in nonincreasing order  $\lambda_1 \geq \lambda_2 \geq \dots$ , do not differ by a positive even integer:

$$\lambda_i - \lambda_j - 2k \neq 0 \quad (7.13)$$

for  $i \leq j$  and  $k$  a positive integer, then a formal holomorphic gauge transformation exists that sets  $A(g)$  in the canonical nonresonant form [9]:

$$A'(g) = \frac{\gamma_0}{\beta_0} \frac{1}{g} \quad (7.14)$$

that is one-loop exact to all orders of perturbation theory. As a consequence, if in addition  $\frac{\gamma_0}{\beta_0}$  is diagonalizable,  $Z(\lambda)$  is diagonalizable as well and its eigenvalues  $Z_{\mathcal{O}_i}(\lambda)$  are computed by [9]:

$$Z_{\mathcal{O}_i}(\lambda) = \left(\frac{g\left(\frac{\mu}{\lambda}\right)}{g(\mu)}\right)^{\frac{\gamma_{0\mathcal{O}_i}}{\beta_0}} \quad (7.15)$$

with  $\gamma_{0\mathcal{O}_i}$  the eigenvalues of  $\gamma_0$ . Correspondingly, if the nonresonant diagonal scheme exists:

$$\langle \mathcal{O}_{j_1}(\lambda x_1) \dots \mathcal{O}_{j_n}(\lambda x_n) \rangle_{conn} \sim \frac{Z_{\mathcal{O}_{j_1}}(\lambda) \dots Z_{\mathcal{O}_{j_n}}(\lambda)}{\lambda^{D_{\mathcal{O}_1} + \dots + D_{\mathcal{O}_n}}} G_{conf\ j_1 \dots j_n}^{(n)}(x_1, \dots, x_n) \quad (7.16)$$

provided that  $G_{conf\ j_1 \dots j_n}^{(n)}(x_1, \dots, x_n)$  does not vanish, with:

$$g^2\left(\frac{\mu}{\lambda}\right) \sim \frac{1}{\beta_0 \log\left(\frac{1}{\lambda^2}\right)} \left(1 - \frac{\beta_1}{\beta_0^2} \frac{\log \log\left(\frac{1}{\lambda^2}\right)}{\log\left(\frac{1}{\lambda^2}\right)}\right) \quad (7.17)$$

as  $\lambda \rightarrow 0$ . Actually, the nonresonant diagonal scheme where eq. (7.14) holds is constructed inductively order by order in perturbation theory starting from any other scheme by means of the Poincaré-Dulac theorem [9]. However, for the purposes of the present paper we only need to know whether it exists for the collinear twist-2 operators.

### 7.3 Anomalous dimensions of twist-2 operators

We define the bare operators with  $s \geq 2$  and  $k \geq 0$  [6]:

$$\mathcal{O}_{Bs}^{(k)} = (i\partial_+)^k \mathcal{O}_{Bs} \quad (7.18)$$

that, to the leading order of perturbation theory, for  $k > 0$  are conformal descendants [14] of the corresponding primary conformal operator  $\mathcal{O}_{Bs}^{(0)} = \mathcal{O}_{Bs}$  in the standard basis. As a consequence of the operator mixing we obtain [6, 14] for the renormalized operators:

$$\mathcal{O}_s^{(k)} = \sum_i^s Z_{si} \mathcal{O}_{Bi}^{(k+s-i)} \quad (7.19)$$

with the mixing matrix  $Z$  and the matrix of the anomalous dimensions:

$$\gamma(g) = -\frac{\partial Z}{\partial \log \mu} Z^{-1} = \sum_{j=0}^{\infty} \gamma_j g^{2j+2} \quad (7.20)$$

lower triangular and  $\gamma_0$  diagonal in the  $\overline{MS}$  scheme [6, 14]. In our notation the eigenvalues of  $\gamma_0$  are given by [6]:

$$\gamma_{0s} = \frac{2}{(4\pi)^2} \left( 4\psi(s+1) - 4\psi(1) - \frac{11}{3} - 8 \frac{s^2 + s + 1}{(s-1)s(s+1)(s+2)} \right) \quad (7.21)$$

for  $\mathcal{O}_s^{(k)}$  with even  $s \geq 2$ , and:

$$\gamma_{0s} = \frac{2}{(4\pi)^2} \left( 4\psi(s+1) - 4\psi(1) - \frac{11}{3} - 8 \frac{s^2 + s - 2}{(s-1)s(s+1)(s+2)} \right) \quad (7.22)$$

for  $\tilde{\mathcal{O}}_s^{(k)}$  with odd  $s \geq 3$ .  $\gamma_{02} = 0$  consistently with the conservation of the stress-energy tensor. The eigenvalues of  $\gamma_0$  for  $\mathbb{S}_s^{(k)}$  and  $\bar{\mathbb{S}}_s^{(k)}$  are given by [12]:

$$\gamma_{0s} = \frac{2}{(4\pi)^2} \left( 4\psi(s+1) - 4\psi(1) - \frac{11}{3} \right) \quad (7.23)$$

with even  $s \geq 2$ .

Interestingly, from the above equations it follows that in  $SU(N)$  YM theory  $\gamma_0$  is independent of  $N$  and thus has no nonplanar contribution [10].

Moreover, we have verified numerically that the nonresonant condition in eq. (7.13) is satisfied for all the collinear twist-2 operators in the standard basis up to  $s = 10^4$ .

Hence, the nonresonant diagonal basis exists for the collinear twist-2 operators and restricts to the lowest order of perturbation theory to the standard basis because  $\gamma_0$  is diagonal in this basis.

Therefore, the standard basis may be employed to compute the conformal correlators in eq. (7.6).

## 8 Generating functional of Euclidean RG-improved correlators

From the vantage point of view of the 't Hooft large- $N$  expansion it is convenient to employ the rescaled operators in eqs. (3.23) in Minkowskian and Euclidean space-time respectively, so that the 2-point correlators are of order 1 for large  $N$ . Choosing again the operators  $\mathbb{O}'_s(x)$  as example, the corresponding correlators acquire the simpler form in Minkowskian:

$$\begin{aligned}
& \langle \mathbb{O}'_{s_1}(x_1) \dots \mathbb{O}'_{s_n}(x_n) \rangle_{conn} \\
&= (N^{2-n} - N^{-n}) \frac{1}{(4\pi^2)^n} 2^{\sum_{l=1}^n s_l} i^{\sum_{l=1}^n s_l} \sum_{k_1=0}^{s_1-2} \dots \sum_{k_n=0}^{s_n-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \dots \binom{s_n}{k_n} \binom{s_n}{k_n+2} \\
& \frac{(-1)^n}{n} \sum_{\sigma \in P_n} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! \dots (s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)})! \\
& \frac{(x_{\sigma(1)} - x_{\sigma(2)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{(|x_{\sigma(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \\
& \dots \frac{(x_{\sigma(n)} - x_{\sigma(1)})_+^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}}}{(|x_{\sigma(n)} - x_{\sigma(1)}|^2)^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)} + 1}} \tag{8.1}
\end{aligned}$$

and Euclidean space-time:

$$\begin{aligned}
& \langle \mathbb{O}'_{s_1}{}^E(x_1) \dots \mathbb{O}'_{s_n}{}^E(x_n) \rangle_{conn} \\
&= (N^{2-n} - N^{-n}) \frac{1}{(4\pi^2)^n} 2^{\sum_{l=1}^n s_l} (-1)^{\sum_{l=1}^n s_l} \sum_{k_1=0}^{s_1-2} \dots \sum_{k_n=0}^{s_n-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \dots \binom{s_n}{k_n} \binom{s_n}{k_n+2} \\
& \frac{1}{n} \sum_{\sigma \in P_n} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! \dots (s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)})! \\
& \frac{(x_{\sigma(1)} - x_{\sigma(2)})_z^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{((x_{\sigma(1)} - x_{\sigma(2)})^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \\
& \dots \frac{(x_{\sigma(n)} - x_{\sigma(1)})_z^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}}}{((x_{\sigma(n)} - x_{\sigma(1)})^2)^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)} + 1}} \tag{8.2}
\end{aligned}$$

Correspondingly, the generating functional of the Euclidean RG-improved  $n$ -point correlators in the canonical nonresonant diagonal scheme decomposes into its planar  $\mathcal{W}_{Sphere\ asym}^E$  and LO nonplanar  $\mathcal{W}_{Torus\ asym}^E$  contributions:

$$\mathcal{W}_{asym}^E[j_{\mathcal{O}'^E}, \lambda] = \mathcal{W}_{Sphere\ asym}^E[j_{\mathcal{O}'^E}, \lambda] + \mathcal{W}_{Torus\ asym}^E[j_{\mathcal{O}'^E}, \lambda] \tag{8.3}$$

with:

$$\mathcal{W}_{Sphere\ asym}^E[j_{\mathcal{O}'^E}, \lambda] = -N^2 \mathcal{W}_{Torus\ asym}^E[j_{\mathcal{O}'^E}, \lambda] \tag{8.4}$$

and as  $\lambda \rightarrow 0$ :

$$\begin{aligned}
& \mathcal{W}_{Torus\ asym}^E[J_{\mathbb{O}'E}, J_{\tilde{\mathbb{O}}'E}, J_{\mathbb{S}'E}, J_{\tilde{\mathbb{S}}'E}, \lambda] = \\
& -\frac{1}{2} \log \text{Det} \left( \mathcal{I} + \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i\vec{\partial}_+)^{s-k-1} \Delta^{-1} \frac{(Z_{\mathbb{O}'s}(\lambda)J_{\mathbb{O}'E} + Z_{\tilde{\mathbb{O}}'s}(\lambda)J_{\tilde{\mathbb{O}}'E})}{N\lambda^{2+s}} (i\vec{\partial}_+)^{k+1} \right) \\
& -\frac{1}{2} \log \text{Det} \left( \mathcal{I} + \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i\vec{\partial}_+)^{s-k-1} \Delta^{-1} \frac{(Z_{\mathbb{O}'s}(\lambda)J_{\mathbb{O}'E} - Z_{\tilde{\mathbb{O}}'s}(\lambda)J_{\tilde{\mathbb{O}}'E})}{N\lambda^{2+s}} (i\vec{\partial}_+)^{k+1} \right) \\
& -\frac{1}{2} \log \text{Det} \left[ \mathcal{I} \right. \\
& -2 \left( \mathcal{I} + \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i\vec{\partial}_+)^{s-k-1} \Delta^{-1} \frac{(Z_{\mathbb{O}'s}(\lambda)J_{\mathbb{O}'E} - Z_{\tilde{\mathbb{O}}'s}(\lambda)J_{\tilde{\mathbb{O}}'E})}{N\lambda^{2+s}} (i\vec{\partial}_+)^{k+1} \right)^{-1} \\
& \sum_{k_1=0}^{s_1-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} (i\vec{\partial}_+)^{s_1-k_1-1} \Delta^{-1} \frac{Z_{\mathbb{S}'s_1}(\lambda)}{N\lambda^{2+s_1}} J_{\mathbb{S}'E} (i\vec{\partial}_+)^{k_1+1} \\
& \left. \left( \mathcal{I} + \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (i\vec{\partial}_+)^{s_2-k_2-1} \Delta^{-1} \frac{(Z_{\mathbb{O}'s_2}(\lambda)J_{\mathbb{O}'E} - Z_{\tilde{\mathbb{O}}'s_2}(\lambda)J_{\tilde{\mathbb{O}}'E})}{N\lambda^{2+s_2}} (i\vec{\partial}_+)^{k_2+1} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-2} \binom{s_3}{k_3} \binom{s_3}{k_3+2} (i\vec{\partial}_+)^{s_3-k_3-1} \Delta^{-1} \frac{Z_{\mathbb{S}'s_3}(\lambda)}{N\lambda^{2+s_3}} J_{\mathbb{S}'E} (i\vec{\partial}_+)^{k_3+1} \right] \quad (8.5)
\end{aligned}$$

in the coordinate representation.

By the above computation we verify the prediction [2] that the generating functional  $\mathcal{W}_{Torus\ asym}^E$  of the LO nonplanar Euclidean RG-improved collinear twist-2 correlators should inherit the very same loop structure of the corresponding nonperturbative object that involves the sum of the glueball one-loop diagrams.

Hence, the aforementioned asymptotic result strongly constrains the nonperturbative solution of the large- $N$  YM theory and it may actually provide an essential guide to find it.

## A Determinants of block matrices

The matrix in eq. (3.20) has a block structure with noncommutative entries:

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \quad (\text{A.1})$$

where  $A, B, C, D$  are operators. Correspondingly, the determinant of  $M$  reads [15]:

$$\text{Det } M = \text{Det}(D) \text{Det}(A - BD^{-1}C) \quad (\text{A.2})$$

that may be rewritten as:

$$\text{Det } M = \text{Det}(D) \text{Det}(A) \text{Det}(1 - A^{-1}BD^{-1}C) \quad (\text{A.3})$$

In our case:

$$\mathcal{M} = \delta^{ab} \begin{pmatrix} \square + \frac{i}{2} \sum_s J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{i}{2} \sum_s J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} & \frac{i}{\sqrt{2}} \sum_s J_{\mathbb{S}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} \\ \frac{i}{\sqrt{2}} \sum_s J_{\mathbb{S}_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} & \square + \frac{i}{2} \sum_s J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{i}{2} \sum_s J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \end{pmatrix} \quad (\text{A.4})$$

Hence, up to a trivial constant normalization, we get:

$$\begin{aligned} \text{Det } \mathcal{M} &= \text{Det} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{1}{2} i \square^{-1} J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \\ &\quad \text{Det} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2} i \square^{-1} J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \\ &\quad \text{Det} \left[ \mathcal{I} - \frac{1}{2} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{0_s} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2} i \square^{-1} J_{\tilde{0}_s} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right)^{-1} \right. \\ &\quad \quad \left. i \square^{-1} J_{\mathbb{S}_{s_1}} \otimes \mathcal{Y}_{s_1-2}^{\frac{5}{2}} \left( \mathcal{I} + \frac{1}{2} i \square^{-1} J_{0_{s_2}} \otimes \mathcal{Y}_{s_2-2}^{\frac{5}{2}} + \frac{1}{2} i \square^{-1} J_{\tilde{0}_{s_2}} \otimes \mathcal{H}_{s_2-2}^{\frac{5}{2}} \right)^{-1} \right. \\ &\quad \quad \left. i \square^{-1} J_{\mathbb{S}_{s_3}} \otimes \mathcal{Y}_{s_3-2}^{\frac{5}{2}} \right] \quad (\text{A.5}) \end{aligned}$$

where the sum over repeated spin indices is understood.

## B Conformal properties of the standard basis

The gauge-invariant collinear twist-2 operators in the light-cone gauge in the standard basis read [1]:

$$\begin{aligned} \mathbb{O}_s &= \text{Tr} \partial_+ \bar{A}(x) (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ A(x) \\ \tilde{\mathbb{O}}_s &= \text{Tr} \partial_+ \bar{A}(x) (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ A(x) \\ \mathbb{S}_s &= \frac{1}{\sqrt{2}} \text{Tr} \partial_+ \bar{A}(x) (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ \bar{A}(x) \\ \bar{\mathbb{S}}_s &= \frac{1}{\sqrt{2}} \text{Tr} \partial_+ A(x) (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ A(x) \quad (\text{B.1}) \end{aligned}$$



where  $C_l^{\alpha'}(x)$  are the Gegenbauer polynomials that are a special case of the Jacobi polynomials [1]:

$$C_l^{\alpha'}(x) = \frac{\Gamma(l+2\alpha')\Gamma(\alpha'+\frac{1}{2})}{\Gamma(2\alpha')\Gamma(l+\alpha'+\frac{1}{2})} P_l^{(\alpha'-\frac{1}{2}, \alpha'-\frac{1}{2})}(x) \quad (\text{B.2})$$

with the symmetry properties:

$$C_l^{\alpha'}(-x) = (-1)^l C_l^{\alpha'}(x) \quad (\text{B.3})$$

They are the restriction, up to perhaps normalization and linear combinations [1], to the component with maximal-spin projection  $s$  along the  $p_+$  direction of the balanced,  $\mathbb{O}_s^{\mathcal{T}=2}$ ,  $\tilde{\mathbb{O}}_s^{\mathcal{T}=2}$ , and unbalanced,  $\mathbb{S}_s^{\mathcal{T}=2}$ , twist-2 operators that to the leading and next-to-leading order of perturbation theory transform as primary operators with respect to the conformal group [3]:

$$\begin{aligned} \mathbb{O}_s^{\mathcal{T}=2} &= \text{Tr } F_{(\rho_1}^\mu \overleftrightarrow{D}_{\rho_2} \dots \overleftrightarrow{D}_{\rho_{s-1}} F_{\rho_s)\mu} - \text{traces} \\ \tilde{\mathbb{O}}_s^{\mathcal{T}=2} &= \text{Tr } \tilde{F}_{(\rho_1}^\mu \overleftrightarrow{D}_{\rho_2} \dots \overleftrightarrow{D}_{\rho_{s-1}} F_{\rho_s)\mu} - \text{traces} \\ \mathbb{S}_s^{\mathcal{T}=2} &= \text{Tr } (F_{\mu(\nu} + i\tilde{F}_{\mu(\nu}) \overleftrightarrow{D}_{\rho_1} \dots \overleftrightarrow{D}_{\rho_{s-2}} (F_{\lambda)\sigma} + i\tilde{F}_{\lambda)\sigma}) - \text{traces} \end{aligned} \quad (\text{B.4})$$

where the parentheses stand for symmetrization of all the indices in between and the subtraction of the traces ensures that the contraction of any two indices is zero.

The above statement about the conformal properties needs some refinements. In a conformal field theory the propagator of a vector field  $V_\mu(x)$  with conformal dimension 1 is purely longitudinal [16]:

$$\langle V_\mu(x)V_\nu(0) \rangle = \left( g_{\mu\nu} - 2\frac{x_\mu x_\nu}{x^2} \right) \frac{1}{x^2} = \frac{1}{2} \partial_\mu \partial_\nu \log x^2 \quad (\text{B.5})$$

Therefore, there is no Lorentz-covariant gauge where the field  $A_\mu$  may be primary with respect to the conformal group to the lowest order of perturbation theory. Yet,  $F_{\mu\nu}$  decomposes into the  $(1, 0) \oplus (0, 1)$  representations of the Lorentz group [1]:

$$F_{a\dot{a}b\dot{b}} = 2(f_{ab}\epsilon_{\dot{a}\dot{b}} - \epsilon_{ab}\dot{f}_{\dot{a}\dot{b}}) \quad (\text{B.6})$$

where:

$$F_{a\dot{a}b\dot{b}} = \sigma_{\dot{a}\dot{b}}^\mu \sigma_{ab}^\nu F_{\mu\nu} \quad (\text{B.7})$$

and:

$$\begin{aligned} f_{ab} &= \frac{i}{2} (\sigma^{\mu\nu})_{ab} F_{\mu\nu} \\ \dot{f}_{\dot{a}\dot{b}} &= -\frac{i}{2} (\bar{\sigma}^{\mu\nu})_{\dot{a}\dot{b}} F_{\mu\nu} \end{aligned} \quad (\text{B.8})$$

with:

$$\bar{f}_{ab} = \dot{f}_{\dot{a}\dot{b}} \quad (\text{B.9})$$

It turns out that to the leading and next-to-leading order of perturbation theory  $f_{ab}$  and  $f_{\dot{a}\dot{b}}$  are local primary fields with respect to the conformal group [3]. Consequently, it follows the construction [3] in eq. (B.4) of the local gauge-invariant operators that are conformal primaries to the leading and next-to-leading order, with the caveat above that the corresponding representation of the conformal group does not extend outside the local gauge-invariant sector by including the field  $A_\mu$  in a covariant gauge.

Besides, by projecting on the maximal-spin component along the  $p_+$  direction the aforementioned gauge-invariant operators restrict to [1]:

$$\begin{aligned}
\mathbb{O}_s &= \text{Tr } f_{11}(x) (i\vec{D}_+ + i\overleftarrow{D}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{D}_+ - \overleftarrow{D}_+}{\vec{D}_+ + \overleftarrow{D}_+} \right) f_{\dot{1}\dot{1}}(x) & s = 2, 4, 6, \dots \\
\tilde{\mathbb{O}}_s &= \text{Tr } f_{11}(x) (i\vec{D}_+ + i\overleftarrow{D}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{D}_+ - \overleftarrow{D}_+}{\vec{D}_+ + \overleftarrow{D}_+} \right) f_{\dot{1}\dot{1}}(x) & s = 3, 5, 7, \dots \\
\mathbb{S}_s &= \frac{1}{\sqrt{2}} \text{Tr } f_{11}(x) (i\vec{D}_+ + i\overleftarrow{D}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{D}_+ - \overleftarrow{D}_+}{\vec{D}_+ + \overleftarrow{D}_+} \right) f_{11}(x) & s = 2, 4, 6, \dots \\
\bar{\mathbb{S}}_s &= \frac{1}{\sqrt{2}} \text{Tr } f_{\dot{1}\dot{1}}(x) (i\vec{D}_+ + i\overleftarrow{D}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left( \frac{\vec{D}_+ - \overleftarrow{D}_+}{\vec{D}_+ + \overleftarrow{D}_+} \right) f_{\dot{1}\dot{1}}(x) & s = 2, 4, 6, \dots \quad (\text{B.10})
\end{aligned}$$

that are primaries with respect to the collinear conformal subgroup  $SL(2, R)$  [6] and reduce in the light-cone gauge to the operators in eq. (B.1) with  $f_{11} = -\partial_+ \bar{A}$ .

In fact, the fields  $f \equiv f_{11}$  and  $\bar{f} \equiv f_{\dot{1}\dot{1}}$  may be realized as primary operators for the collinear conformal subgroup to the leading and next-to-leading order of perturbation theory in a noncanonical path-integral quantization of the YM theory in the light-cone gauge by a suitable change of variables [12]:

$$\begin{aligned}
S_{YM}(f, \bar{f}) &= \int \bar{f}^a \partial_+^{-2} \square f^a + 2 \frac{g}{\sqrt{N}} f^{abc} (\partial_+^{-1} \bar{f}^a f^b \bar{\partial} \partial_+^{-2} \bar{f}^c + \partial_+^{-1} f^a \bar{f}^b \partial \partial_+^{-2} f^c) \\
&\quad - 2 \frac{g^2}{N} f^{abc} f^{ade} \partial_+^{-1} (\partial_+^{-1} \bar{f}^b f^c) \partial_+^{-1} (\partial_+^{-1} f^d \bar{f}^e) d^4 x \quad (\text{B.11})
\end{aligned}$$

where:

$$\langle \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \rangle = \frac{1}{Z} \int \mathcal{D}f \mathcal{D}\bar{f} e^{iS_{YM}(f, \bar{f})} \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \quad (\text{B.12})$$

## References

- [1] M. Bochicchio, M. Papinutto and F. Scardino, *n-point correlators of twist-2 operators in  $SU(N)$  Yang-Mills theory to the lowest perturbative order*, *JHEP* **08** (2021) 142, [[2104.13163](#)].
- [2] M. Bochicchio, *An asymptotic solution of Large- $N$  QCD, for the glueball and meson spectrum and the collinear S-matrix*, *AIP Conf. Proc.* **1735** (2016) 030004.
- [3] N. Beisert, G. Ferretti, R. Heise and K. Zarembo, *One-loop QCD spin chain and its spectrum*, *Nucl. Phys. B* **717** (2005) 137–189, [[hep-th/0412029](#)].
- [4] D. G. Robertson and F. Wilczek, *Anomalous dimensions of anisotropic gauge theory operators*, *Phys. Lett. B* **251** (1990) 434–438.

- [5] To appear in arXiv.
- [6] A. V. Belitsky and D. Mueller, *Broken conformal invariance and spectrum of anomalous dimensions in QCD*, *Nucl. Phys. B* **537** (1999) 397–442, [[hep-ph/9804379](#)].
- [7] V. M. Braun, A. N. Manashov, S. Moch and M. Strohmaier, *Three-loop evolution equation for flavor-nonsinglet operators in off-forward kinematics*, *JHEP* **06** (2017) 037, [[1703.09532](#)].
- [8] V. M. Braun, Y. Ji and A. N. Manashov, *Two-loop evolution equation for the B-meson distribution amplitude*, *Phys. Rev. D* **100** (2019) 014023, [[1905.04498](#)].
- [9] M. Bochicchio, *On the geometry of operator mixing in massless QCD-like theories*, *Eur. Phys. J. C* **81** (2021) 749, [[2103.15527](#)].
- [10] U. Aglietti, M. Becchetti, M. Bochicchio, M. Papinutto and F. Scardino, *Operator mixing, UV asymptotics of nonplanar/planar 2-point correlators, and nonperturbative large-N expansion of QCD-like theories*, [2105.11262](#).
- [11] G. 't Hooft, *A Planar Diagram Theory for Strong Interactions*, *Nucl. Phys. B* **72** (1974) 461.
- [12] A. V. Belitsky, S. E. Derkachov, G. P. Korchemsky and A. N. Manashov, *Dilatation operator in (super-)Yang-Mills theories on the light-cone*, *Nucl. Phys. B* **708** (2005) 115–193, [[hep-th/0409120](#)].
- [13] M. Becchetti and M. Bochicchio, *Operator mixing in massless QCD-like theories and Poincaré-Dulac theorem*, *Eur. Phys. J. C* , [[2103.16220](#)].
- [14] V. M. Braun, G. P. Korchemsky and D. Mueller, *The Uses of conformal symmetry in QCD*, *Prog. Part. Nucl. Phys.* **51** (2003) 311–398, [[hep-ph/0306057](#)].
- [15] P. D. Powell, *Calculating Determinants of Block Matrices*, [1112.4379](#).
- [16] Y. M. Makeenko, *On conformal operators in quantum chromodynamics*, tech. rep., USSR, 1980.