

THE DERIVED AUSLANDER–IYAMA CORRESPONDENCE

GUSTAVO JASSO AND FERNANDO MURO

With an appendix by Bernhard Keller

ABSTRACT. We work over a perfect field. Recent work of the second-named author established a Derived Auslander Correspondence that relates finite-dimensional self-injective algebras that are twisted 3-periodic to algebraic triangulated categories of finite type. Moreover, the aforementioned work also shows that the latter triangulated categories admit a unique differential graded enhancement. In this article we prove a higher-dimensional version of this result that, given an integer $d \geq 1$, relates twisted $(d + 2)$ -periodic algebras to algebraic triangulated categories with a $d\mathbb{Z}$ -cluster tilting object. We also show that the latter triangulated categories admit a unique differential graded enhancement. Our result yields recognition theorems for interesting algebraic triangulated categories, such as the Amiot cluster category of a self-injective quiver with potential in the sense of Herschend and Iyama and, more generally, the Amiot–Guo–Keller cluster category associated with a d -representation finite algebra in the sense of Iyama and Oppermann. As an application of our result, we obtain infinitely many triangulated categories with a unique differential graded enhancement that is not strongly unique. In the appendix, B. Keller explains how—combined with crucial results of August and Hua–Keller—our main result yields the last key ingredient to prove the Donovan–Wemyss Conjecture in the context of the Homological Minimal Model Program for threefolds.

CONTENTS

Introduction	1
1. Cluster tilting subcategories	13
2. $(d + 2)$ -angulated categories	19
3. Enhanced $(d + 2)$ -angulated categories	29
4. Higher structures and $(d + 2)$ -angulations	35
5. Existence and uniqueness of enhancements	77
6. Recognition theorems	106
Appendix A. The Donovan–Wemyss conjecture, by Bernhard Keller	113
References	115

INTRODUCTION

We work over a field \mathbf{k} . Triangulated categories, introduced by Verdier in the late 1960s [Ver96] (see also Puppe’s [Pup62]) are central objects of study in homological algebra [HJR10]. Typical examples of triangulated categories include derived categories of rings and schemes, stable categories of representations of finite groups, and the stable homotopy category (of spectra). Despite their ubiquity and usefulness, it has long been recognised that triangulated categories lack a number of desirable

2020 *Mathematics Subject Classification*. Primary: 18G80; secondary: 18N40.

Key words and phrases. Triangulated categories; differential graded categories; A_∞ -categories; cluster tilting; twisted periodicity; Massey products; Hochschild cohomology; Toda brackets.

formal properties. For example, limits of triangulated categories in general do not exist, nor is there a reasonable tensor product operation on triangulated categories; moreover, most triangulated categories do not admit interesting (co)limits, such as pushouts and pullbacks, see [Toë11, Sec. 2.2] for a more detailed discussion of these shortcomings.

In the seminal article [BK90], it is proposed to address the above issues by endowing the triangulated categories that appear in practice (at least in algebra and geometry) with what nowadays is called a ‘differential graded enhancement.’ Recall that a differential graded (=DG) category [Kel65] is a category-like structure in which the morphisms between two given objects form a cochain complex, and whose composition law is compatible with the differentials in a suitable sense. A DG category \mathcal{A} has an associated homotopy category $H^0(\mathcal{A})$, obtained by passing to the 0-th cohomology of its morphism complexes; when the DG category \mathcal{A} has the property of being *pre-triangulated* (Definition 3.1.1), the category $H^0(\mathcal{A})$ is canonically a triangulated category. Given a triangulated category \mathcal{T} , we say that a pre-triangulated DG category \mathcal{A} is a *DG enhancement of \mathcal{T}* if there is an equivalence of triangulated categories $\mathcal{T} \simeq H^0(\mathcal{A})$. Interpreting these in the appropriate manner, pre-triangulated DG categories have all the expected formal properties that triangulated categories lack and, moreover, all triangulated categories that arise naturally in algebra and geometry (but, in general, not in topology) admit a DG enhancement, see [Kel06] for details. For completeness, we mention that there are alternative frameworks for enhancing triangulated categories, such as (stable) Quillen model categories [HPS97, Hov99], spectral categories [SS03], triangulated A_∞ -categories [LH, BLM08], stable ∞ -categories [Lur17], etc.

Given a triangulated category \mathcal{T} , two natural questions arise: Does \mathcal{T} admit a DG enhancement? and if it does, is the enhancement unique up to the appropriate notion of equivalence? In general, the first question has a negative answer, for the stable homotopy category does not admit a DG enhancement [Sch10], and there are even triangulated categories that are linear over a field and do not admit a DG enhancement [RVdB20] (there also exist triangulated categories that do not even admit ‘topological’ enhancements [MSS07]). The second question is also rather delicate. Indeed, there exist triangulated categories that admit inequivalent DG enhancements [DS07, Sch02], and there are even such triangulated categories that are linear over a field [Kaj13, RVdB19]. Notwithstanding, there are several results that show that important classes of triangulated categories, such as derived categories of modules or of quasi-coherent sheaves, admit a unique DG enhancement [LO10, CS18, CNS22]. We note that there are also several important results on the uniqueness of ‘topological’ enhancements [Ant, Sch01, Sch07, SS02].

On the other hand, recent work of the second-named author established an existence and uniqueness result for a quite different class of triangulated categories. We say that a triangulated category \mathcal{T} that is linear over a field is of *finite type* if it has finite-dimensional morphism spaces and split idempotents and, moreover, there exists an object $c \in \mathcal{T}$ such that every other object of \mathcal{T} is a direct summand of a finite direct sum of copies of c . This means, in particular, that there are only finitely many indecomposable objects in \mathcal{T} up to isomorphism (in fact, the underlying additive category of \mathcal{T} is equivalent to the category of finitely-generated projective $\mathcal{T}(c, c)$ -modules). We also need the following terminology: Given an algebra Λ , recall that a Λ -bimodule I is *invertible* if there exists a Λ -bimodule I^{-1} such that there are isomorphisms

$$I \otimes_{\Lambda} I^{-1} \cong \Lambda \cong I^{-1} \otimes_{\Lambda} I$$

of Λ -bimodules. A finite-dimensional basic self-injective algebra Λ is *twisted n -periodic* if the n -th syzygy $\Omega_{\Lambda^e}^n(\Lambda)$ of the diagonal Λ -bimodule is stably isomorphic

to an invertible Λ -bimodule, where Λ^e denotes the enveloping algebra of Λ . Finally, we denote the category of finitely-generated projective (right) Λ -modules by $\text{proj}(\Lambda)$. The following theorem summarises the main results in [Mur22]. Recall that a field is perfect if it has characteristic 0 or if it has characteristic $p > 0$ and every element has a p -th root.

Theorem (Derived Auslander Correspondence). *Suppose that \mathbf{k} is a perfect field. There is a bijective correspondence between the following:*

- (1) *Quasi-isomorphism classes of DG algebras A such that the perfect derived category $\mathcal{D}^c(A)$ is of finite type.*
- (2) *Equivalence classes of algebraic triangulated categories \mathcal{T} of finite type.*
- (3) *Equivalence classes of pairs (Λ, I) consisting of*
 - *a finite-dimensional basic self-injective algebra Λ that is twisted 3-periodic and*
 - *an invertible Λ -bimodule I such that $\Omega_{\Lambda^e}^3(\Lambda) \cong I$ in the stable category of Λ -bimodules.*

The correspondences are given by

$$A \longmapsto (\mathcal{D}^c(A), A) \quad \text{and} \quad (\mathcal{T}, c) \longmapsto (\mathcal{T}(c, c), \mathcal{T}(c, c[-1])).$$

Moreover, the algebraic triangulated categories in (2) admit a unique DG enhancement.

In the context of the above theorem, two pairs (Λ, I) and (Λ', I') are *equivalent* if the pairs $(\text{proj}(\Lambda), - \otimes_{\Lambda} I)$ and $(\text{proj}(\Lambda'), - \otimes_{\Lambda'} I')$ are equivalent as pairs in the natural sense (Definition 2.1.10). In particular, if the pairs (Λ, I) and (Λ', I') are equivalent, then the algebras Λ and Λ' are Morita equivalent (even isomorphic, since they are basic). The restriction to basic algebras is technically convenient since, under these assumptions, the Picard group of invertible Λ -bimodules and the group of outer automorphisms of Λ are isomorphic. In particular, up to isomorphism, every invertible Λ -bimodule is of the form ${}_{\sigma}\Lambda_1$ for some algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$, where ${}_{\sigma}\Lambda_1$ is the Λ -bimodule with underlying right Λ -module Λ_{Λ} but with left action twisted by σ . We also observe that if the algebra Λ is connected and non-separable, then the invertible Λ -bimodule I is uniquely determined up to bimodule isomorphism by the algebra Λ , see [Mur22, Prop. 9.8] (this observation generalises to arbitrary periods, see Proposition 2.2.6).

We have stated the main results in [Mur22] in the above form in order to emphasise the analogy with a classical result in representation theory of algebras: the famous Auslander correspondence [Aus71]. Recall that a finite-dimensional algebra A is said to have finite representation type if it admits an additive generator, that is a finite-dimensional A -module M such that every other finite-dimensional A -module is a direct summand of a finite direct sum of copies of M .

Theorem (Auslander Correspondence). *Let \mathbf{k} be a field. There is a bijective correspondence between the following:*

- (1) *Morita equivalence classes of finite-dimensional algebras A of finite representation type and*
- (2) *Morita equivalence classes of Auslander algebras, that is finite-dimensional algebras Γ with global dimension at most 2 and dominant dimension [Tac64] at least 2.*

The correspondence is given by the formula $A \mapsto \text{Hom}_A(M, M)$, where M is an additive generator of the category of finite-dimensional A -modules.

Thus, the Derived Auslander Correspondence can be interpreted both as a result concerning the general theory of triangulated categories and their DG enhancements, and as a result in representation theory of finite-dimensional algebras.

Notice, moreover, the following parallel between the triangulated and the classical Auslander Correspondence: in both cases, the second class of algebras that is considered (twisted 3-periodic algebras in the triangulated case and Auslander algebras in the classical case) is part of an infinite hierarchy of finite-dimensional algebras. Indeed, as part of his foundational work on higher-dimensional Auslander–Reiten theory, Iyama introduced in [Iya07a] the class of d -Auslander algebras, obtained by replacing the number 2 by the number $d + 1$ in the definition of Auslander algebra. Remarkably, Iyama was also able to extend the classical Auslander Correspondence to d -Auslander algebras by introducing the important concept of a d -cluster tilting module [Iya07b] (see Definition 6.2.2). We state Iyama’s higher-dimensional Auslander Correspondence below as it helps us position the main result in this article in an appropriate context.

Theorem (Auslander–Iyama correspondence). *Let \mathbf{k} be a field and $d \geq 1$ an integer. There is a bijective correspondence between the following:*

- (1) *Equivalence classes of pairs (A, M) consisting of*
 - *a finite-dimensional algebra A and*
 - *a d -cluster tilting A -module M .*
- (2) *Morita equivalence classes of d -Auslander algebras.*

The correspondence is given by the formula $(A, M) \mapsto \text{Hom}_A(M, M)$.

In the context of the previous theorem, two pairs (A, M) and (A', M') are *equivalent* if there exists an equivalence

$$\text{mod}(A) \xrightarrow{\simeq} \text{mod}(A')$$

between the corresponding categories of finite-dimensional modules that restricts to an equivalence between the closures of M and M' under finite direct sums and direct summands. In particular, if the pairs (A, M) and (A', M') are equivalent, then the algebras A and A' are Morita equivalent.

At this stage, it is worth pointing out that a 1-cluster tilting module is simply an additive generator of the ambient module category. In particular, for $d = 1$, the Auslander–Iyama Correspondence specialises to the classical Auslander Correspondence. We also mention that there are other results in representation theory of algebras that follow a similar pattern, such as the Auslander–Iyama–Solberg Correspondence [IS18] that extends the correspondence in [AS93]. If $d = 2$, other than in higher Auslander–Reiten theory, 2-cluster tilting modules (and their triangulated counter parts [IY08]) play a crucial role in the categorification of Fomin–Zelevinsky cluster algebras [FZ02] by means of 2-Calabi–Yau triangulated categories [Kel08], see for example [Ami09, BMR⁺06, BIRS09, DK08, GLS06, Kel13, Pal08] among many others. Cluster tilting modules also appear in commutative algebra, (possibly non-commutative) algebraic geometry and symplectic geometry, see for example [AIR15, DJL21, HIMO, IT13, IW14].

Our main result. Inspired by the above pattern, the aim of this article is to prove a suitable higher-dimensional extension of the Derived Auslander Correspondence established in [Mur22]. For the definition of $d\mathbb{Z}$ -cluster tilting object, see Definition 1.3.1 (and compare with Remark 1.3.5); here we only mention that such objects are in particular d -step generators of the ambient category in the sense of Rouquier [Rou08], but they generate in a stronger sense (see Theorem 1.3.7).

Theorem A (Derived Auslander–Iyama Correspondence). *Suppose that \mathbf{k} is a perfect field and $d \geq 1$ an integer. There are bijective correspondences between the following:*

- (1) *Quasi-isomorphism classes of DG algebras A that satisfy the following:*

- (a) The ordinary algebra $H^0(A)$ is a basic finite-dimensional algebra.
- (b) The free DG A -module $A \in \mathcal{D}^c(A)$ is a $d\mathbb{Z}$ -cluster tilting object, that is A is a d -cluster tilting object that satisfies $A \cong A[d]$.
- (2) Equivalence classes of pairs (\mathcal{T}, c) consisting of
 - (a) an algebraic triangulated category \mathcal{T} with finite-dimensional morphism spaces and split idempotents and
 - (b) a basic $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$, that is a d -cluster tilting object such that $\mathcal{T}(c, c[i]) = 0$ for all $i \notin d\mathbb{Z}$.
- (3) Equivalence classes of pairs (Λ, I) consisting of
 - (a) a basic finite-dimensional self-injective algebra Λ that is twisted $(d+2)$ -periodic and
 - (b) an invertible Λ -bimodule I such that $\Omega_{\Lambda^e}^{d+2}(\Lambda) \cong I$ in the stable category of Λ -bimodules.

The correspondences are given by

$$A \longmapsto (\mathcal{D}^c(A), A) \quad \text{and} \quad (\mathcal{T}, c) \longmapsto (\mathcal{T}(c, c), \mathcal{T}(c, c[-d])).$$

Moreover, the algebraic triangulated categories in (2) admit a unique DG enhancement.

In the context of **Theorem A**, two pairs (\mathcal{T}, c) and (\mathcal{T}', c') as in (2) are equivalent if there exists an equivalence of triangulated categories

$$F: \mathcal{T} \xrightarrow{\sim} \mathcal{T}'$$

such that $F(c) \cong c'$. Similarly, two pairs (Λ, I) and (Λ', I') as in (3) are equivalent if there exists an algebra isomorphism $\varphi: \Lambda \xrightarrow{\sim} \Lambda'$ such that $\varphi^*(I') \cong I$. Notice also that a devissage argument shows that $\mathcal{D}^c(A)$ has finite-dimensional morphism spaces if and only if for all $i \in \mathbb{Z}$ the vector space $H^i(A) \cong \text{Hom}_{\mathcal{D}(A)}(A, A[i])$ is finite dimensional, which is the case for the DG algebras considered in the theorem. We also mention that the compact derived category of a non-contractible DG algebra satisfying the conditions in **Theorem A** cannot be equivalent to the derived category of an abelian or exact category: Indeed, the finite-dimensionality of the morphism spaces implies that we would be dealing with a bounded derived category, and such categories do not admit non-zero objects that are isomorphic to any of its non-zero shifts. In particular, the results in [CNS22] cannot be used to establish uniqueness of DG enhancements for the triangulated categories in **Theorem A**(2).

As before, we note that a $1\mathbb{Z}$ -cluster tilting object is simply an additive generator of the ambient triangulated category (in particular, the notions of 1- and $1\mathbb{Z}$ -cluster tilting object coincide). Hence, for $d = 1$, **Theorem A** specialises to the Derived Auslander Correspondence from [Mur22]. In fact, our proof of **Theorem A** shares some methods with the second-named author's proof of the Derived Auslander Correspondence; in particular, we rely crucially on techniques of homotopy theory as well as on the obstruction theory developed by the second-named author in [Mur20b]. As we explain in more detail later in this introduction, the central problem that arises is to interpret the property of an object being $d\mathbb{Z}$ -cluster tilting in terms of the Hochschild(–Tate) cohomology of a certain graded algebra.

Theorem A shows a deep connection between $d\mathbb{Z}$ -cluster tilting objects and twisted $(d+2)$ -periodic algebras. Indeed, the fact that the endomorphism algebra of a $d\mathbb{Z}$ -cluster tilting object in an triangulated category is twisted $(d+2)$ -periodic was observed first by Dugas [Dug12] in the algebraic case and in general by Chan, Darpö, Iyama and Marczinzik [CDIM] where they also investigate the connection between twisted periodicity and higher Auslander–Reiten theory focusing on the fundamental class of trivial extensions of finite-dimensional algebras. From the viewpoint promoted in this article, this is a consequence of the relationship

between $d\mathbb{Z}$ -cluster tilting objects and $(d+2)$ -angulated categories and a general characterisation of twisted periodicity [GSS03, Han20], see [Proposition 2.2.9](#). For more information on (twisted) periodic algebras and their importance in representation theory and elsewhere in mathematics, we refer the reader to [ES08, CDIM] and the references therein.

We wish to highlight that [Theorem A](#) is closely related to the theory of $(d+2)$ -angulated categories in the sense of Geiss, Keller and Oppermann [GKO13] and their DG enhancements (which we introduce in this article). Indeed, by the main result in [GKO13], the additive closure of a $d\mathbb{Z}$ -cluster tilting object in a triangulated category inherits the structure of a $(d+2)$ -angulated category (we recall this result as [Theorem 2.2.1](#)). One of the main results in this article ([Theorem 5.1.3](#)) establishes the existence and uniqueness of enhancements for a class of $(d+2)$ -angulated categories introduced by Amiot [Ami07] in the case $d = 1$ and by Lin [Lin19] in the general case $d \geq 1$. From this perspective, [Theorem A](#) can be regarded as a contribution to the emerging subject of higher-dimensional homological algebra, where $(d+2)$ -angulated categories and their exact/abelian variants [Jas16] play a central role.

Notice that [Theorem A](#) cannot be generalised to triangulated categories with d -cluster tilting objects that are not $d\mathbb{Z}$ -cluster tilting, at least not without further assumptions. On the one hand, for $d > 2$, Ladkani [Lad] has shown that *every* basic finite-dimensional algebra A over an algebraically closed field arises as the endomorphism algebra of a distinguished d -cluster tilting object in an algebraic triangulated category constructed explicitly from a presentation of A in terms of a quiver with relations. This means that the endomorphism algebras of d -cluster tilting objects do not admit any useful characterisation (homological or otherwise) in this case. Moreover, still for $d > 2$, Ladkani also has shown that there are *non-equivalent* algebraic triangulated categories with d -cluster tilting objects whose endomorphism algebras are isomorphic. In both instances, Ladkani's constructions involve d -cluster tilting objects that are not $d\mathbb{Z}$ -cluster tilting, see [Remark 6.1.7](#) for details and precise references. On the other hand, for $d = 2$, Keller and Lie have announced [KL23] a general (and stronger) result for algebraic 2-Calabi–Yau triangulated categories that requires additional assumptions on the enhancements which are necessary to apply a result of Van den Bergh [VdB15] in the course of the proof, see [Remark 6.3.4](#) for details. We also mention recent work of Hanihara [Han22] that deals with ‘hereditary’ d -Calabi–Yau triangulated categories with a d -cluster tilting object.

Existence and uniqueness of A_∞ -structures. Our approach to the proof of [Theorem A](#) suggests a further interpretation in terms of the existence and uniqueness of A_∞ -structures on certain graded algebras that are d -sparse in the sense that their non-zero components are concentrated in degrees that are multiples of d (see [Theorem 5.1.2](#) for a precise statement). Indeed, to prove that the correspondence in [Theorem A](#) is surjective we construct, given a basic twisted $(d+2)$ -periodic algebra Λ with respect to an algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$, a minimal A_∞ -structure on the d -sparse graded algebra $\Lambda(\sigma, d)$ with non-zero components

$$\Lambda(\sigma, d)^{di} := {}_{\sigma^i} \Lambda_1, \quad i \in \mathbb{Z},$$

with the property that its associated perfect derived category admits a $d\mathbb{Z}$ -cluster tilting object with graded endomorphism algebra isomorphic to $\Lambda(\sigma, d)$. As explained previously in this introduction, we establish the existence of such an A_∞ -algebra structure by leveraging the obstruction theory developed in [Mur20a].

On the other hand, the uniqueness of enhancements for the triangulated categories in [Theorem A\(2\)](#) (which is essentially equivalent to the injectivity of the

correspondences) can be explained in terms of Hochschild cohomology of d -sparse graded algebras. For the sake of comparison, recall that a graded algebra A is intrinsically formal if every DG algebra (or, more generally, A_∞ -algebra) B such that $H^\bullet(B) \cong A$ as graded algebras is already quasi-isomorphic to A (equipped with the trivial differential and vanishing higher operations). The following criterion for intrinsic formality, due to Kadeishvili, is well known.

Theorem ([Kad88], see also [ST01, Thm. 4.7]). *Let A be a graded algebra such that*

$$\mathrm{HH}^{p+2, -p}(A, A) = 0, \quad p \geq 1.$$

Then, the following statements hold:

- (1) *The graded algebra A is intrinsically formal.*
- (2) *If B is a DG algebra with $H^\bullet(B) = A$, then B is quasi-isomorphic to A (viewed as a DG algebra with vanishing differential) via a quasi-isomorphism that is the identity in cohomology.*

Kadeishvili’s Theorem can be applied, for example, to show that the algebra of Laurent polynomials $\mathbf{k}[\iota^{\pm 1}]$ on a variable of cohomological degree $-d$ is intrinsically formal (more generally, Kadeishvili’s Theorem can be used to prove that the Laurent polynomial algebra $K[\iota^{\pm 1}] = K \otimes_{\mathbf{k}} \mathbf{k}[\iota^{\pm 1}]$ is intrinsically formal if K is a finite-dimensional algebra of projective dimension at most d as a K -bimodule [Sai20, Cor. 4.2]). At this juncture it is interesting to observe that the d -sparse graded algebra $\Lambda(\sigma, d)$ introduced above can be identified with the twisted Laurent polynomial algebra

$$\Lambda(\sigma, d) \cong \frac{\Lambda\langle \iota^{\pm 1} \rangle}{(\iota x - \sigma(x)\iota)_{x \in \Lambda}}, \quad |\iota| = -d.$$

However, in general $\Lambda(\sigma, d)$ is *far from being intrinsically formal* and, indeed, has a very rich Hochschild cohomology.

Nonetheless, we leverage a key insight (already present in [Mur20a, Mur22] in the case $d = 1$): Given a d -sparse Massey algebra, which is a pair (A, m) consisting d -sparse graded algebra A and a class $m \in \mathrm{HH}^{d+2, -d}(A)$ whose Gerstenhaber square $\mathrm{Sq}(m)$ vanishes (when the characteristic of \mathbf{k} is different from 2, this is equivalent to the familiar Maurer–Cartan equation

$$\frac{1}{2}[m, m] = 0$$

with respect to the usual Lie bracket on $\mathrm{HH}^{\bullet, *}(A)$), there is a bigraded cochain complex $\mathbf{C}^{\bullet, *}(A, m)$ whose vector spaces of cochains are given by the Hochschild cohomology $\mathrm{HH}^{\bullet, *}(A)$ of the graded algebra A and whose differential of bidegree $(d+1, -d)$ is given by

$$\partial: x \mapsto \pm[m, x]$$

almost everywhere (Definition 5.2.5). The Hochschild cohomology of the d -sparse Massey algebra (A, m) is the cohomology of this complex, denoted by $\mathrm{HH}^{\bullet, *}(A, m)$. The uniqueness of enhancements for the triangulated categories in Theorem A(2) is closely related to the following generalisation of Kadeishvili’s intrinsic formality theorem, which is of independent interest (see the proofs of Theorem 5.3.3(1) and Theorem 5.1.2(2)).

Theorem B. *Let (A, m) be a d -sparse Massey algebra A . Suppose that*

$$\mathrm{HH}^{p+2, -p}(A, m) = 0, \quad p > d.$$

If B and C are minimal A_∞ -algebras with $H^\bullet(B) = H^\bullet(C) = A$ as graded algebras such that

$$\{m_{d+2}^B\} = \{m_{d+2}^C\} = m \in \mathrm{HH}^{d+2, -d}(A),$$

then B and C are A_∞ -isomorphic through an A_∞ -isomorphism with identity linear part.

Kadeishvili's theorem is an immediate consequence of the case $d = 1$ of [Theorem B](#), see [Corollary 5.2.8](#). Although [Theorem B](#) likely has a direct proof, we deduce it from the vanishing of certain terms in an extended spectral sequence of Bousfield–Kan type that is the main technical tool in the obstruction theory developed in [[Mur20a](#)]. Results similar to [Theorem B](#) have appeared in other contexts; for example, in symplectic geometry, Lekili and Perutz have shown that the A_∞ -structures on certain graded algebras arising from the Fukaya category of the punctured torus are controlled by a pair of Hochschild cohomology classes, see [[LP11](#)] for details.

Applications. Although [Theorem A](#) has a theoretical character, it has interesting consequences and useful applications. First of all, notice that a twisted $(d + 2)$ -periodic algebra Λ is also twisted $(d + 2)k$ -periodic for every $k \geq 1$. Choose an invertible Λ -bimodule I such that $\Omega_{\Lambda^e}^{d+2}(\Lambda) \cong I$ in the stable category of Λ -bimodules and, consequently, $\Omega_{\Lambda^e}^{(d+2)k}(\Lambda) \cong I^k$ in the stable category of Λ -bimodules for every $k \geq 1$. Thus, we may apply [Theorem A](#) to the infinite sequence of pairs

$$(\Lambda, I), (\Lambda, I^{\otimes 2}), (\Lambda, I^{\otimes 3}), \dots$$

to obtain an infinite sequence of algebraic triangulated categories

$$\mathcal{T}_1, \mathcal{T}_2, \mathcal{T}_3, \dots$$

equipped with an $n_k\mathbb{Z}$ -cluster tilting object $c_k \in \mathcal{T}_k$ such that $\mathcal{T}_k(c_k, c_k) \cong \Lambda$, where $n_k = dk + 2k - 2$. It would be interesting to investigate whether there is a relationship between these triangulated categories (see also [Remark 3.2.10](#) and [Theorem 5.1.3](#)).

Our approach to the proof of [Theorem A](#) yields a more precise statement concerning the equivalence classes of strong enhancements of the (algebraic) $(d + 2)$ -angulated categories we consider ([Theorem 5.4.10](#)). As a consequence we obtain, to our knowledge, the first examples of triangulated categories that admit a unique enhancement but not a unique strong enhancement (that is, an enhancement that is not ‘strongly unique’ in the sense of [[LO10](#)]), see [Corollary 5.4.12](#).

Recognition theorems. As for further applications, the bijectivity of the correspondence in [Theorem A](#) yields recognition theorems for interesting classes of algebraic triangulated categories that admit $d\mathbb{Z}$ -cluster tilting objects, from which we single out the following (we refer the reader to [Section 6](#) for details and definitions). Below, we restrict to connected non-separable algebras for simplicity.

Following Iyama and Oppermann [[IO11](#)], we say that a finite-dimensional algebra A is (*d-hereditary*) *d-representation finite* if A has global dimension at most d and there exists a d -cluster tilting A -module (which turns out to be unique up to multiplicity of its indecomposable direct summands [[Iya11](#)]). For example, if $d = 1$, then A is a hereditary algebra of finite representation type and, if the ground field is moreover algebraically closed, then A is Morita equivalent to the path algebra of a Dynkin quiver. Recall that the category of finitely-generated projective modules over the Gelfand–Ponomarev preprojective algebra [[GP79](#)] of a Dynkin quiver admits an algebraic (1-Calabi–Yau) triangulated structure [[AR96](#)] that is essentially unique by the main result in [[Mur22](#)]. [Theorem A](#) yields the following analogous recognition theorem.

Theorem ([Theorem 6.2.7](#)). *Suppose that \mathbf{k} is a perfect field. The d -Calabi–Yau Amiot–Guo–Keller cluster category of a connected, non-separable d -representation*

finite algebra has a unique enhancement and is uniquely characterised among algebraic triangulated categories by the existence of a $d\mathbb{Z}$ -cluster tilting object whose endomorphism algebra is isomorphic to the $(d + 1)$ -preprojective algebra $\Pi_{d+1}(A)$ of A in the sense of [IO13].

In the case $d = 1$ of the above theorem, the algebra $\Pi_{d+1}(A) = \Pi_2(A)$ is the preprojective algebra of A as defined in [BGL87].

As a further application of **Theorem A**, we obtain the following recognition theorem for the Amiot cluster category of a self-injective quiver with potential. This result complements the Recognition Theorem for acyclic cluster categories of Keller and Reiten [KR08], which deals with 2-Calabi–Yau categories with a cluster tilting object whose endomorphism algebra has acyclic Gabriel quiver.

Theorem (Theorem 6.3.1). *Let \mathbf{k} be an arbitrary field. The 2-Calabi–Yau Amiot cluster category of a connected, non-separable self-injective quiver with potential (Q, W) has a unique enhancement and is uniquely characterised among algebraic triangulated categories by the existence of a $2\mathbb{Z}$ -cluster tilting object whose endomorphism algebra is isomorphic to the (completed) Jacobian algebra of (Q, W) .*

Both of the above results are special cases of a more general recognition theorem for the Amiot–Guo–Keller cluster category of a (pseudo-compact) homologically smooth DG algebra that satisfies a number of technical conditions that guarantee, in particular, the existence of a $d\mathbb{Z}$ -cluster tilting object, see **Theorem 6.1.5**.

Proof of the Donovan–Wemyss Conjecture. As a final application, in the appendix, B. Keller explains how—combined with crucial results of August [Aug20] and Hua–Keller [HK]—our main result yields the last key ingredient to prove the Donovan–Wemyss Conjecture ([DW16, Conj. 1.4] and [Aug20, Conj. 1.3]) in the context of the Homological Minimal Model Program for threefolds [Wem18].

Let $X = \text{Spec } R$ be the formal neighbourhood of an isolated compound Du Val (=cDV) singularity over the field of complex numbers, that is R is a complete local three-dimensional hypersurface with an isolated singularity whose generic hyperplane section is a Du Val/Kleinian singularity. The class of cDV singularities was introduced by Reid [Rei83] who was motivated by their natural occurrence in the minimal model program for three-folds [KM98]. Suppose now that X admits a crepant resolution (=smooth minimal model) $p: \tilde{X} \rightarrow X$. To this geometric setup, Donovan and Wemyss [DW16] associate a basic finite-dimensional algebra $\Lambda = \Lambda(p)$ – the contraction algebra – that controls the non-commutative deformations of the exceptional fibre of p , see [Wem23] for a survey of the state of the art. Donovan and Wemyss made the following remarkable conjecture [DW16, Aug20]:

Let R_1 and R_2 be isolated cDV singularities that admit crepant resolutions $p_1: \tilde{X}_1 \rightarrow X_1$ and $p_2: \tilde{X}_2 \rightarrow X_2$ with corresponding contraction algebras $\Lambda(p_1)$ and $\Lambda(p_2)$. Then, the \mathbb{C} -algebras R_1 and R_2 are isomorphic if and only if the contraction algebras $\Lambda(p_1)$ and $\Lambda(p_2)$ are derived equivalent.

The Donovan–Wemyss Conjecture is considered the most important open problem in the context of the Homological Minimal Model Program [Wem18]. It is known that derived contraction algebras of a given isolated compound Du Val singularity are derived equivalent [Wem18, Dug10]. In the appendix, Keller explains how to reduce the remaining implication of the Donovan–Wemyss Conjecture to the validity of the following statement:

Let R be an isolated cDV singularity that admits a crepant resolution $p: \tilde{X} \rightarrow X$ and $\text{sg}(R) = \text{D}^b(\text{mod } R)/\text{K}^b(\text{proj } R)$ its singularity category with its canonical DG enhancement $\text{sg}_{dg}(R)$. Then, up to

quasi-equivalence of DG categories, $\text{sg}_{dg}(R)$ is uniquely determined by the following properties:

- The algebraic triangulated category $H^0(\text{sg}_{dg}R) = \text{sg}(R)$ has finite-dimensional morphism spaces and split idempotents.
- There exists a $2\mathbb{Z}$ -cluster tilting object in $\text{sg}(R)$ whose graded endomorphism algebra is isomorphic to the algebra of Laurent polynomials $\Lambda(p)[i^{\pm 1}]$, $|i| = -2$.

The above statement is a special case of [Theorem A](#), and hence the remaining implication of the Donovan–Wemyss Conjecture follows, see the appendix for details.

Outline of the proof of [Theorem A](#) and structure of the article. The proof of [Theorem A](#) is involved and occupies the vast majority of the article. For this reason, we provide a rough outline of the argument that we hope will help the reader have a clear overview of the proof and of the article as a whole.

In [Section 1](#) we recall the definition and basic aspects related to d -cluster tilting subcategories. Most of the material in this section is well known to specialists, but [Theorem 1.3.9](#) (a characterisation of $d\mathbb{Z}$ -cluster tilting subcategories) is new.

In [Section 2](#) we sum up the essentials of the theory of $(d+2)$ -angulated categories. Again, most of the material in this section is known to experts, but [Theorem 2.2.4](#) (essentially a reformulation of [Theorem 1.3.9](#) in terms of $(d+2)$ -angulated structures) is new. In particular, [Proposition 2.2.10](#) shows that the correspondence in [Theorem A](#) is well defined. Moreover, we recall the construction of $(d+2)$ -angulated structures on the category of finitely-generated projective modules over a twisted $(d+2)$ -periodic algebra Λ given by Amiot [[Ami07](#)] in the case $d = 1$ and Lin [[Lin19](#)] in the general case $d \geq 1$. In a nutshell, a choice of an exact sequence of Λ -bimodules

$$\eta: \quad 0 \rightarrow {}_1\Lambda_\sigma \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

with projective-injective middle terms exhibiting the twisted periodicity of Λ defines a class of $(d+2)$ -angles \triangleleft_η on $\text{proj}(\Lambda)$ that, up to equivalence, depends only on the class of σ in the outer automorphism group of Λ and not on the choice of η (see [Proposition 2.2.18](#), which is also new). We call these structures Amiot–Lin (AL) $(d+2)$ -angulations. In [Section 3](#) we review the theory of DG categories and their derived categories, and introduce DG enhancements of $(d+2)$ -angulated categories ([Definition 3.2.5](#)) as straightforward generalisations of DG enhancements of triangulated categories.

We prove [Theorem A](#) by providing an inverse to the map $(\mathcal{T}, c) \mapsto (\Lambda, \sigma)$ that associates to an algebraic triangulated category (with the relevant finiteness assumptions) with a $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$ the corresponding twisted $(d+2)$ -periodic algebra $\Lambda = \mathcal{T}(c, c)$ together with the algebra automorphism σ induced by the action of the d -fold shift of \mathcal{T} . Thus, given a pair (Λ, σ) consisting of a twisted $(d+2)$ -periodic algebra with respect to the algebra automorphism σ , we need to construct an *algebraic* triangulated category \mathcal{T} with a $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$ together with a fully faithful functor

$$\begin{aligned} \text{proj}(\Lambda) &\longrightarrow \mathcal{T} \\ \Lambda &\longmapsto c \end{aligned}$$

that is compatible with the actions of the functors $- \otimes_\Lambda \sigma \Lambda_1$ and $[d]$ on the source and the target category, respectively. For this, we observe that in fact \mathcal{T} must be equivalent to the perfect derived category $\text{D}^c(A)$ of a DG algebra A whose cohomology is (isomorphic to) the d -sparse graded algebra

$$\Lambda(\sigma, d) = \bigoplus_{di \in \mathbb{Z}} \sigma^i \Lambda_1, \quad a \cdot b = \sigma^j(a)b, \quad |b| = dj.$$

Thus, we need to construct a DG algebra A such that $H^\bullet(A) \cong \Lambda(\sigma, d)$ and $A \in \mathcal{D}^c(A)$ is a $d\mathbb{Z}$ -cluster tilting object. Equivalently, by Kadeishvili’s Homotopy Transfer Theorem, we need to endow $\Lambda(\sigma, d)$ with a minimal A_∞ -algebra structure

$$A = (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, m_{3d+2}, \dots),$$

where $m_{i+2} = 0$ if $i \notin d\mathbb{Z}$ due to the fact that $\Lambda(\sigma, d)$ is d -sparse ([Proposition 4.3.4](#)), in such a way that $A \in \mathcal{D}^c(A)$ is a $d\mathbb{Z}$ -cluster tilting object. Moreover, in order to obtain an inverse to the correspondence in [Theorem A](#), such a minimal A_∞ -algebra A should be unique up to A_∞ -quasi-isomorphism. In particular, we need to translate the property ‘ $A \in \mathcal{D}^c(A)$ is a $d\mathbb{Z}$ -cluster tilting object’ into a property of the minimal A_∞ -algebra that can be leveraged using techniques of homological and homotopical algebra.

Suppose first that A is a minimal A_∞ -algebra with $H^\bullet(A) = \Lambda(\sigma, d)$ and observe that, since the latter graded algebra is d -sparse, the first non-trivial higher operation

$$m_{d+2} \in \mathcal{C}^{d+2, -d}(\Lambda(\sigma, d), \Lambda(\sigma, d))$$

is in fact a Hochschild *cocycle* ([Proposition 4.4.4](#)). Thus, associated to A there is a well-defined class in Hochschild cohomology

$$\{m_{d+2}\} \in \mathrm{HH}^{d+2, -d}(\Lambda(\sigma, d), \Lambda(\sigma, d))$$

that we call its *universal Massey product of length $d + 2$* . As it turns out, most relevant to our purpose is the so-called *restricted universal Massey product*

$$j^* \{m_{d+2}\} \in \mathrm{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d)) \cong \mathrm{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$$

obtained by restricting $\{m_{d+2}\}$ along the morphism

$$j^*: \mathrm{HH}^{\bullet, *}\Lambda(\sigma, d), \Lambda(\sigma, d) \longrightarrow \mathrm{HH}^{\bullet, *}\Lambda, \Lambda(\sigma, d)$$

induced by the inclusion $j: \Lambda \rightarrow \Lambda(\sigma, d)$ of the degree 0 part. [Corollary 4.5.17](#) then establishes the following remarkable fact: $A \in \mathcal{D}^c(A)$ is a $d\mathbb{Z}$ -cluster tilting object if and only if the restricted universal Massey product $j^* \{m_{d+2}\}$ is represented by an extension all of whose middle terms are projective-injective Λ -bimodules. Furthermore, the latter property is equivalent to the class $j^* \{m_{d+2}\}$ being a *unit* in the Hochschild–Tate cohomology $\underline{\mathrm{HH}}^{\bullet, *}\Lambda, \Lambda(\sigma, d)$, see [Remark 4.5.16](#).

The proof of [Corollary 4.5.17](#), which is at the core of the argument, relies on a careful analysis of the relationship between Toda brackets, Massey products, minimal A_∞ -structures, Hochschild cohomology and $(d + 2)$ -angulations; we study these relationships in [Section 4](#), the results of which can be regarded as $(d + 2)$ -angulated analogues of results in [[Mur20b](#), [Mur22](#)] and of classical results for triangulated categories. To understand why this technical excursion is essential in our approach, notice that if the restricted universal Massey product $j^* \{m_{d+2}\} \in \mathrm{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$ is represented by an extension with projective-injective middle terms, then there is a well-defined AL $(d + 2)$ -angulation $\diamond_{j^* \{m_{d+2}\}}$ on $\mathrm{proj}(\Lambda)$. On the other hand, a general theorem of Geiss, Keller and Oppermann ([Theorem 2.2.1](#)) shows that

$$\mathrm{proj}(\Lambda) \simeq \mathrm{add}(A) \subseteq \mathcal{D}^c(A)$$

is endowed with the structure of $(d + 2)$ -angulated category—which we call standard—that is induced by the canonical triangulation of $\mathcal{D}^c(A)$, provided that $A \in \mathcal{D}^c(A)$ is a $d\mathbb{Z}$ -cluster tilting object. The proof of [Corollary 4.5.17](#) relies on the fact that the AL $(d + 2)$ -angulation and the standard $(d + 2)$ -angulation of $\mathrm{proj}(\Lambda)$ coincide up to sign; we demonstrate this in [Theorem 4.5.8](#). Toda brackets enter the proof of [Theorem 4.5.8](#) as these can be used to detect standard $(d + 2)$ -angles ([Theorem 4.1.12](#)), and a certain agreement between Toda brackets and Massey products on $H^\bullet(A) = \Lambda(\sigma, d)$ ([Theorem 4.2.6](#)) turns out to be essential for computations. Due to its technical nature, the reader may want to skip [Section 4](#) on

a first reading and refer to it as necessary throughout their reading of [Section 5](#) (familiarity with [Definitions 4.4.5](#) and [4.5.4](#) will largely suffice).

Having established all the necessary technical preliminaries, in [Section 5](#) we prove [Theorem A](#) and [Theorem B](#). We deduce [Theorem A](#) from a more precise statement, [Theorem 5.1.2](#), which allows us to construct the inverse of the correspondence: There exists an essentially unique minimal A_∞ -algebra structure A on $\Lambda(\sigma, d)$ whose restricted universal Massey product is a unit in the corresponding Hochschild–Tate cohomology, that is such that $A \in \mathcal{D}^c(A)$ is a $d\mathbb{Z}$ -cluster tilting object. The proofs of [Theorem B](#) and [Theorem 5.1.2](#) utilise the same homotopical techniques (which we discuss in the relevant subsections) leveraged in [\[Mur22\]](#) to establish [Theorem A](#) in the case $d = 1$ of triangulated categories of finite type; in particular the obstruction theory developed in [\[Mur20a\]](#) by the second-named author plays a crucial role in this part of the article. In [Section 5.5](#) we illustrate the necessity for the hypotheses in [Theorem A](#) and [Theorem 5.1.3](#) by means of examples and, in particular, show that there exist triangulated categories with a $d\mathbb{Z}$ -cluster tilting object that admit inequivalent topological enhancements although they admit a unique (\mathbf{k} -linear) DG enhancement (see [Section 5.5.4](#)). Finally, [Section 6](#) includes several recognition theorems that we deduce from [Theorem A](#).

Conventions. We fix a positive integer $d \geq 1$ once and for all (we occasionally let $d = -1, 0$, but this is always clear from the context). All algebras and categories in this article are assumed to be linear over a ground field \mathbf{k} , except for a few obvious examples; unless explicitly noted otherwise, all modules are right modules. The Jacobson radical of an algebra Λ is denoted by J_Λ . Our main results require \mathbf{k} to be a perfect field, and this assumption is indicated whenever relevant. Unless noted otherwise, all (ordinary) categories considered in this article are additive; accordingly, all subcategories are assumed to be (strictly) full and closed under the formation of finite direct sums in the ambient category. Given a collection X of objects in a category \mathcal{C} with split idempotents (=Karoubian), we denote by $\mathbf{add}(X)$ the smallest (additive) subcategory of \mathcal{C} that contains X and is closed under direct summands; if $X = \{x\}$ consists of a single object, we write $\mathbf{add}(x) = \mathbf{add}(X)$. We say that an object c in a category in which the Krull–Remak–Schmidt Theorem holds (for example, in an additive category with finite-dimensional morphism spaces and split idempotents [\[Kra15, Cor. 4.4\]](#)) is *basic* if in any decomposition $c = c_1 \oplus c_2 \oplus \cdots \oplus c_n$ into indecomposable objects, there is an isomorphism $c_i \cong c_j$ if and only if $i = j$. A finite-dimensional algebra A is *basic* if the regular representation of A is basic as an object of its category of finite-dimensional (projective) modules. We usually denote the shift/suspension functor in a triangulated category by $[1]$, although the notation Σ is used when dealing with general $(d + 2)$ -angulated categories. If \mathcal{T} is a triangulated category with split idempotents and X is a class of objects in \mathcal{T} , we denote by $\mathbf{thick}(X)$ the smallest triangulated subcategory of \mathcal{T} that contains X and is closed under direct summands; if $X = \{x\}$ consists of a single object, we write $\mathbf{thick}(x) = \mathbf{thick}(X)$. Finally, we compose morphisms in a category as functions: the composite of morphisms $f: x \rightarrow y$ and $g: y \rightarrow z$ is the morphism $gf = g \circ f: x \rightarrow z$.

Acknowledgements. The authors thank Peter Jørgensen, Zhengfang Wang and the Uppsala Representation Theory Group for comments on a previous versions of this article, as well as Yankı Lekili for bringing references [\[Sei15, LP11\]](#) to our attention. They are particularly grateful to Julian Külshammer for his numerous comments and suggestions. The authors also wish to thank A. Lorenzin for a question that lead to the inclusion of [Section 5.4](#). Finally, the authors thank Martin Herschend for discussions in relation to extending the scope of our main results, see [Remark 2.2.19](#).

Financial support. G. J. was partially supported by the Deutsche Forschungsgemeinschaft (German Research Foundation) under Germany’s Excellence Strategy – GZ 2047/1, Projekt- ID 390685813 as well by the Swedish Research Council (Vetenskapsrådet) Research Project Grant ‘Higher structures in higher-dimensional homological algebra.’ F. M. was partially supported by grants PID2020-117971GB-C21 funded by MCIN/AEI/10.13039/501100011033, US-1263032 (US/JUNTA/FEDER, UE), and P20_01109 (JUNTA/FEDER, UE).

1. CLUSTER TILTING SUBCATEGORIES

Setting 1.0.1. We fix a triangulated category \mathcal{T} with finite-dimensional morphism spaces and split idempotents as well as a subcategory $\mathcal{C} \subseteq \mathcal{T}$ that satisfies $\mathcal{C} = \text{add}(\mathcal{C})$.

1.1. Functorially finite subcategories. We begin by recalling a classical definition due to Auslander and Smalø, and independently to Enochs.

Definition 1.1.1 ([AS80, AS81] and [Eno81]). The subcategory $\mathcal{C} \subseteq \mathcal{T}$ is *contravariantly finite in \mathcal{T}* if the following condition is satisfied:

- (1) For each object $x \in \mathcal{T}$ there exist an object $c \in \mathcal{C}$ and a morphism $f: c \rightarrow x$ with the following property: For each morphism $g: c' \rightarrow x$ with $c' \in \mathcal{C}$, there exists a (not necessarily unique) morphism $h: c' \rightarrow c$ such that $g = f \circ h$.

$$\begin{array}{ccc} & & c' \\ & \exists h \swarrow & \downarrow \forall g \\ c & \xrightarrow{f} & x \end{array}$$

Equivalently, the induced linear map

$$\mathcal{T}(c', c) \longrightarrow \mathcal{T}(c', x), \quad h \longmapsto f \circ h,$$

is surjective (but not necessarily injective). Such a morphism f is called a *right \mathcal{C} -approximation of x* .

Dually, the subcategory $\mathcal{C} \subseteq \mathcal{T}$ is *covariantly finite in \mathcal{T}* if the following condition is satisfied:

- (1) For each object $x \in \mathcal{T}$ there exist an object $c \in \mathcal{C}$ and a morphism $f: x \rightarrow c$ with the following property: For each morphism $g: x \rightarrow c'$ with $c' \in \mathcal{C}$, there exists a (not necessarily unique) morphism $h: c \rightarrow c'$ such that $g = h \circ f$.

$$\begin{array}{ccc} x & \xrightarrow{f} & c \\ & \searrow \forall g & \downarrow \exists h \\ & & c' \end{array}$$

Equivalently, the induced linear map

$$\mathcal{T}(c, c') \longrightarrow \mathcal{T}(x, c'), \quad h \longmapsto h \circ f,$$

is surjective (but not necessarily injective). Such a morphism f is called a *left \mathcal{C} -approximation of x* .

Finally, the subcategory $\mathcal{C} \subseteq \mathcal{T}$ is *functorially finite in \mathcal{T}* if it is both contravariantly finite and covariantly finite in \mathcal{T} .

Remark 1.1.2. Contravariantly finite subcategories are also called ‘pre-covering’ and right approximations are also called ‘pre-covers’. Similarly, covariantly finite subcategories are also called ‘pre-enveloping’ and left approximations are also called ‘pre-envelopes,’ see [Eno81].

Example 1.1.3. Suppose that \mathcal{C} is a right admissible subcategory of \mathcal{T} , that is the inclusion $\mathcal{C} \subseteq \mathcal{T}$ admits a right adjoint $R: \mathcal{T} \rightarrow \mathcal{C}$. Let $x \in \mathcal{T}$ and $c = R(x)$. The co-unit $\varepsilon: c \rightarrow x$ is a right \mathcal{C} -approximation of x , for it induces natural isomorphisms

$$\mathcal{T}(c', c) = \mathcal{T}(c', R(x)) \xrightarrow{\cong} \mathcal{T}(c', x), \quad c' \in \mathcal{C}.$$

In particular, one may think of contravariantly finite subcategories as mild generalisations of right admissible subcategories. Similarly, covariantly finite subcategories are mild generalisations of left admissible subcategories, that is subcategories such that the inclusion admits a left adjoint.

The following basic example of a functorially finite subcategory is the most relevant for the purposes of this article.

Example 1.1.4. Let $c \in \mathcal{T}$ and $\mathcal{C} = \mathbf{add}(c)$. Then, the subcategory \mathcal{C} is functorially finite in \mathcal{T} . Indeed, \mathcal{C} -approximations of an object $x \in \mathcal{T}$ are easily constructed from a decomposition $c = c_1 \oplus \cdots \oplus c_k$ into indecomposable objects and bases of the (finite-dimensional) vector spaces $\mathcal{T}(c_i, x)$ and $\mathcal{T}(x, c_i)$, $1 \leq i \leq k$.

Remark 1.1.5. Suppose that \mathcal{C} is contravariantly finite in \mathcal{T} . For each object $x \in \mathcal{T}$ there exists a commutative diagram, constructed inductively, of the form

$$\begin{array}{ccccccc} \cdots & c_2 & \longrightarrow & c_1 & \longrightarrow & c_0 & \\ & \searrow & & \nearrow & & \searrow & \\ \cdots & & & x_2 & \xleftarrow{+1} & x_1 & \xleftarrow{+1} & x_0 = x \end{array}$$

in which each of the morphisms $c_i \rightarrow x_i$ is a right \mathcal{C} -approximation and, as is customary, the triangles

$$\begin{array}{ccc} & c_i & \\ \nearrow & & \searrow \\ x_{i+1} & \xleftarrow{+1} & x_i \end{array}$$

represent exact triangles

$$x_{i+1} \longrightarrow c_i \longrightarrow x_i \longrightarrow x_{i+1}[1].$$

Dually, if \mathcal{C} is covariantly finite in \mathcal{T} , then for each object x of \mathcal{T} there exists a commutative diagram of the form

$$\begin{array}{ccccccc} & & c^0 & \longrightarrow & c^1 & \longrightarrow & c^2 & \cdots \\ & \nearrow & & \searrow & \nearrow & \searrow & \nearrow & \\ x = x^0 & \xleftarrow{+1} & x^1 & \xleftarrow{+1} & x^2 & \cdots & \end{array}$$

in which each of the morphisms $x^i \rightarrow c^i$ is a left \mathcal{C} -approximation and with corresponding exact triangles

$$x^i \longrightarrow c^i \longrightarrow x^{i+1} \longrightarrow x^i[1].$$

From the point of view of this article, the existence of such ‘ \mathcal{C} -(co)resolutions’ is one of the main motivations for considering functorially finite subcategories.

1.2. The Verdier product. The following elementary operation on subcategories of a triangulated category is needed in the proof of [Theorem 1.3.9](#) below. All subcategories $\mathcal{X} \subseteq \mathcal{T}$ considered in this section are assumed to satisfy $\mathbf{add}(\mathcal{X}) = \mathcal{X}$.

Definition 1.2.1. Let $\mathcal{X}, \mathcal{Y} \subseteq \mathcal{T}$ be subcategories. The *Verdier product of \mathcal{X} with \mathcal{Y}* is the full subcategory $\mathcal{X} * \mathcal{Y}$ of \mathcal{T} spanned by the objects $t \in \mathcal{T}$ such that there exists an exact triangle

$$x \longrightarrow t \longrightarrow y \longrightarrow x[1]$$

with $x \in \mathcal{X}$ and $y \in \mathcal{Y}$.

Notation 1.2.2. Given subcategories $\mathcal{X}, \mathcal{Y} \subseteq \mathcal{T}$, we let $\mathcal{X} \vee \mathcal{Y}$ be the smallest additive subcategory of \mathcal{T} containing \mathcal{X} and \mathcal{Y} . Obviously, $\mathcal{X} \vee \mathcal{Y} \subseteq \mathcal{X} * \mathcal{Y}$ and $\mathcal{X} \vee \mathcal{Y} = \mathcal{Y} \vee \mathcal{X}$.

Lemma 1.2.3. *Let $\mathcal{X}, \mathcal{Y}, \mathcal{Z} \subseteq \mathcal{T}$ be subcategories. The following statements hold:*

- (1) *The Verdier product is associative: $(\mathcal{X} * \mathcal{Y}) * \mathcal{Z} = \mathcal{X} * (\mathcal{Y} * \mathcal{Z})$.*
- (2) *Suppose that*

$$\forall x \in \mathcal{X}, \forall y \in \mathcal{Y}, \quad \mathcal{T}(y, x[1]) = 0.$$

*Then, $\mathcal{X} * \mathcal{Y} = \mathcal{X} \vee \mathcal{Y} \subseteq \mathcal{Y} * \mathcal{X}$.*

- (3) *If*

$$\forall x \in \mathcal{X}, \forall y \in \mathcal{Y}, \quad \mathcal{T}(x, y) = 0,$$

*then $\mathcal{X} * \mathcal{Y}$ is closed under direct summands.*

Proof. (1) This is an immediate consequence of the octahedral axiom.

- (2) The equality $\mathcal{X} * \mathcal{Y} = \mathcal{X} \vee \mathcal{Y}$ follows from the fact that every triangle

$$x \rightarrow z \rightarrow y \rightarrow x[1]$$

with $x \in \mathcal{X}$ and $y \in \mathcal{Y}$ splits, for the morphism $y \rightarrow x[1]$ vanishes by assumption. The inclusion $\mathcal{X} \vee \mathcal{Y} = \mathcal{Y} \vee \mathcal{X} \subseteq \mathcal{Y} * \mathcal{X}$ is obvious.

- (3) See [IY08, Prop. 2.1(1)]. □

Example 1.2.4. Let $\mathcal{C} \subseteq \mathcal{T}$ be a subcategory. The reader can easily verify, using the rotation axiom, that the Verdier product $\mathcal{C} * \mathcal{C}[1] * \cdots * \mathcal{C}[d-1]$ coincides with the full subcategory of \mathcal{T} spanned by those objects $x \in \mathcal{T}$ for which there exists a commutative diagram of the form

$$\begin{array}{ccccccc} & & c_{d-2} & & \cdots & & c_1 & \xrightarrow{\quad} & c_0 \\ & \nearrow & & \searrow & & & \nearrow & & \searrow \\ c_{d-1} & \xleftarrow{+1} & & x_{d-2} & \cdots & & x_2 & \xleftarrow{+1} & x_1 & \xleftarrow{+1} & x_0 = x \end{array}$$

in which $c_i \in \mathcal{C}$, $0 \leq i < d$. In other words, $\mathcal{C} * \mathcal{C}[1] * \cdots * \mathcal{C}[d-1]$ is the full subcategory of \mathcal{T} spanned by those objects that admit a \mathcal{C} -resolution of length (at most) d .

1.3. Cluster tilting subcategories. The following remarkable class of subcategories of a triangulated category, introduced by Iyama and Yoshino (see also [Iya07a, Iya07b, IJ17] for the case of abelian categories), is the main focus of this article.

Definition 1.3.1 ([IY08, Sec. 3]). The subcategory \mathcal{C} is *d-cluster tilting* in \mathcal{T} if the following three conditions are satisfied:

- (1) The subcategory \mathcal{C} is functorially finite in \mathcal{T} .
- (2) An object $x \in \mathcal{T}$ lies in \mathcal{C} if and only if

$$\forall c \in \mathcal{C}, \forall 0 < i < d, \quad \mathcal{T}(x, c[i]) = 0.$$

- (3) An object $x \in \mathcal{T}$ lies in \mathcal{C} if and only if

$$\forall c \in \mathcal{C}, \forall 0 < i < d, \quad \mathcal{T}(c, x[i]) = 0.$$

The subcategory \mathcal{C} is *d \mathbb{Z} -cluster tilting* in \mathcal{T} if, in addition to the three conditions above, \mathcal{C} satisfies the following condition:

- (4) The equality $\mathcal{C} = \mathcal{C}[d]$ holds.

We say that an object $c \in \mathcal{T}$ is *d- or d \mathbb{Z} -cluster-tilting* in \mathcal{T} , if the subcategory $\mathcal{C} = \text{add}(c)$ is *d- or d \mathbb{Z} -cluster tilting* in \mathcal{T} , respectively.

Remark 1.3.2. Let $d = 1$. By vacuity, a subcategory $\mathcal{C} \subseteq \mathcal{T}$ is 1-cluster tilting if and only if $\mathcal{C} = \mathcal{T}$. Consequently, the notions of 1-cluster tilting and $1\mathbb{Z}$ -cluster tilting coincide. In particular, $c \in \mathcal{T}$ is a 1-cluster tilting object (= $1\mathbb{Z}$ -cluster tilting object) if and only if $\text{add}(c) = \mathcal{T}$, that is c generates \mathcal{T} under finite direct sums and direct summands.

Definition 1.3.3. A subcategory $\mathcal{C} \subseteq \mathcal{T}$ is *d-rigid* if

$$\forall c, c' \in \mathcal{C}, \forall 0 < i < d, \quad \mathcal{T}(c, c'[i]) = 0.$$

We say that \mathcal{C} is *d \mathbb{Z} -rigid* if, moreover, for each $c, c' \in \mathcal{C}$ and each $i \in \mathbb{Z}$,

$$i \notin d\mathbb{Z} \quad \implies \quad \mathcal{T}(c, c'[i]) = 0.$$

Remark 1.3.4. By definition, d -cluster tilting subcategories are d -rigid.

Remark 1.3.5. Let $\mathcal{C} \subseteq \mathcal{T}$ be a d -cluster tilting subcategory. It follows immediately from the definition of d -cluster tilting subcategory that the following conditions are equivalent:

- (1) The equality $\mathcal{C} = \mathcal{C}[d]$ holds.
- (2) The subcategory \mathcal{C} is $d\mathbb{Z}$ -rigid.

In other words, the $d\mathbb{Z}$ -cluster tilting subcategories are precisely the d -cluster tilting subcategories that are $d\mathbb{Z}$ -rigid.

Lemma 1.3.6. *Let $\mathcal{C} \subseteq \mathcal{T}$ be a d -rigid subcategory. For all $0 \leq j < i < d$ such that $(j, i) \neq (0, d-1)$, the inclusion*

$$\mathcal{C}[i] * \mathcal{C}[j] \subseteq \mathcal{C}[j] * \mathcal{C}[i]$$

holds.

Proof. Given that $\mathcal{C} \subseteq \mathcal{T}$ is a d -rigid subcategory, the assumption on i and j implies that

$$\forall c, c' \in \mathcal{C}, \quad \mathcal{T}(c, c'[i-j+1]) = 0.$$

Thus, by [Lemma 1.2.3\(2\)](#) the inclusion $\mathcal{C}[i-j] * \mathcal{C} \subseteq \mathcal{C} * \mathcal{C}[i-j]$ holds. Since

$$\mathcal{C}[i] * \mathcal{C}[j] \subseteq \mathcal{C}[j] * \mathcal{C}[i] \quad \iff \quad \mathcal{C}[i-j] * \mathcal{C} \subseteq \mathcal{C} * \mathcal{C}[i-j],$$

the claim follows. \square

The class of d -cluster tilting subcategories admits the following useful characterisation, due to Iyama and Yoshino ((1) \implies (2)) and to Beligiannis ((2) \implies (1)).

Theorem 1.3.7 ([IY08, Thm. 3.1], [Bel15, Thm. 5.3]). *Let $\mathcal{C} \subseteq \mathcal{T}$ be a subcategory that satisfies $\mathcal{C} = \text{add}(\mathcal{C})$. The following statements are equivalent:*

- (1) *The subcategory \mathcal{C} is d -cluster tilting in \mathcal{T} .*
- (2) *The following conditions are satisfied:*
 - (a) *The subcategory \mathcal{C} is d -rigid.*
 - (b) *The equality $\mathcal{T} = \mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1]$ holds.*

Proof. The implication (1) \implies (2) is proven in [IY08, Thm. 3.1] while the converse implication (2) \implies (1) is part of [Bel15, Thm. 5.3], which contains many more equivalent conditions and requires the subcategory $\mathcal{C} \subseteq \mathcal{T}$ to be contravariantly finite. We prove that the latter property in fact follows from the assumptions in part (2) of the statement theorem. The argument is well known to experts (compare with [GKO13, Lemma 4.3]); we include it for the sake of completeness.

Suppose that the conditions in part (2) of the statement of the theorem are satisfied. If $d = 1$, then $\mathcal{C} = \mathcal{T}$ and there is nothing to prove; thus, we may assume that $d \geq 2$. By [Example 1.2.4](#), for every object $x \in \mathcal{T}$ there exists a commutative diagram of the form

$$\begin{array}{ccccccc}
& & c_{d-2} & \cdots & & c_1 & \longrightarrow & c_0 \\
& \nearrow & & \searrow & & \nearrow & & \searrow \\
c_{d-1} & \longleftarrow +1 & \longrightarrow & x_{d-2} & \cdots & x_2 & \longleftarrow +1 & \longrightarrow & x_1 & \longleftarrow +1 & \longrightarrow & x = x_0
\end{array}$$

in which $c_i \in \mathcal{C}$, $0 \leq i < d$. We claim that

$$\forall c \in \mathcal{C}, \forall 0 < i \leq j \leq d-2, \quad \mathcal{T}(c, x_j[i]) = 0.$$

We prove the claim by reverse induction on j . If $j = d-2$, then for each $0 < i \leq d-2$ and each $c \in \mathcal{C}$ there is an induced exact sequence

$$0 = \mathcal{T}(c, c_{d-2}[i]) \longrightarrow \mathcal{T}(c, x_{d-2}[i]) \longrightarrow \mathcal{T}(c, c_{d-1}[i+1]) = 0$$

in which the outer terms vanish since the subcategory \mathcal{C} is d -rigid; this proves the claim in this case. Suppose now that the claim holds for some $1 < j \leq d-2$. Then, for each $0 < i \leq j-1$ and each $c \in \mathcal{C}$ there is an induced exact sequence

$$0 = \mathcal{T}(c, c_{j-1}[i]) \longrightarrow \mathcal{T}(c, x_{j-1}[i]) \longrightarrow \mathcal{T}(c, x_j[i+1]) = 0$$

in which the right-most term vanishes by the inductive hypothesis and the left-most terms since the subcategory \mathcal{C} is d -rigid; this finishes the induction step. We claim now that the morphism $c_0 \rightarrow x_0$ is in fact a right \mathcal{C} -approximation of $x = x_0$. Indeed, there is an induced exact sequence

$$\mathcal{T}(c, c_0) \longrightarrow \mathcal{T}(c, x_0) \longrightarrow \mathcal{T}(c, x_1[1]) = 0$$

in which the term on the right vanishes, as shown above. This proves the claim. Since the object $x \in \mathcal{T}$ was chosen arbitrarily, this shows that \mathcal{C} is contravariantly finite in \mathcal{T} , which is what we needed to prove. \square

Remark 1.3.8. Let $\mathcal{C} \subseteq \mathcal{T}$ be a d -cluster tilting subcategory. Then, \mathcal{C} generates \mathcal{T} as a triangulated category: indeed, $\mathcal{T} = \mathcal{C} * \mathcal{C}[1] * \cdots * \mathcal{C}[d-1]$ and the right-hand side is of course contained in the triangulated closure of \mathcal{C} . Thus, in particular, \mathcal{C} is a d -step generator in the sense of [Rou08]. However, it is remarkable that every object in \mathcal{T} admits a \mathcal{C} -resolution of length at most d (not involving any shifts).

The following characterisation of $d\mathbb{Z}$ -cluster tilting subcategories is quite natural from the point of view of $(d+2)$ -angulated categories, whose definition we recall in [Definition 2.1.1](#).

Theorem 1.3.9. *Let $\mathcal{C} \subseteq \mathcal{T}$ be a subcategory of \mathcal{T} that satisfies $\mathcal{C} = \text{add}(\mathcal{C})$. The following statements are equivalent:*

- (1) *The subcategory \mathcal{C} is $d\mathbb{Z}$ -cluster tilting in \mathcal{T} .*
- (2) *The following conditions are satisfied:*
 - (a) *The subcategory \mathcal{C} is d -rigid.*
 - (b) *The equality $\mathcal{C} = \mathcal{C}[d]$ holds.*
 - (c) *The inclusion $(\mathcal{C}[-1] * \mathcal{C}) \subseteq (\mathcal{C} * \mathcal{C}[1] * \cdots * \mathcal{C}[d-1])$ holds.*
 - (d) *The subcategory \mathcal{C} satisfies $\text{thick}(\mathcal{C}) = \mathcal{T}$.*

Proof. The implication (1) \Rightarrow (2) follows immediately from the definition of $d\mathbb{Z}$ -cluster tilting subcategory and [Theorem 1.3.7](#).

(2) \Rightarrow (1) Notice that the case $d = 1$ is trivial, for assumptions (2b) and (2c) imply that \mathcal{C} is closed under (co)cones and, in view of assumption (2d), we must have $\mathcal{C} = \text{add}(\mathcal{C}) = \mathcal{T}$ (assumption (2a) is vacuous in this case).

Suppose now that $d \geq 2$. Since $\mathcal{C} = \mathcal{C}[d]$ by assumption, we only need to prove that \mathcal{C} is a d -cluster tilting subcategory of \mathcal{T} ; for this, we shall verify the remaining condition in [Theorem 1.3.7\(2\)](#), namely that

$$\mathcal{T} = \mathcal{C} * \mathcal{C}[1] * \cdots * \mathcal{C}[d-1].$$

Let $\text{thick}(\mathcal{C})$ be the smallest thick subcategory of \mathcal{T} containing \mathcal{C} . Clearly [AI12, Lemma 2.15(a)]

$$\mathcal{T} = \text{thick}(\mathcal{C}) = \bigcup_{\ell > 0, n_1, \dots, n_\ell \in \mathbb{Z}} \text{smd}(\mathcal{C}[n_1] * \dots * \mathcal{C}[n_\ell]),$$

where $\mathcal{X} \mapsto \text{smd } \mathcal{X}$ denotes the passage to the closure under direct summands (recall that $\text{thick}(\mathcal{C}) = \mathcal{T}$ by assumption (2d)). We make the following observations:

- Since $\mathcal{C} = \mathcal{C}[d]$, for each $n \in \mathbb{Z}$ there exists an integer $0 \leq i < d$ such that $\mathcal{C}[n] = \mathcal{C}[i]$. Consequently,

$$\mathcal{T} = \text{thick}(\mathcal{C}) = \bigcup_{\ell > 0, 0 \leq n_1, \dots, n_\ell < d} \text{smd}(\mathcal{C}[n_1] * \dots * \mathcal{C}[n_\ell]).$$

- Since \mathcal{C} is a d -rigid subcategory of \mathcal{T} , Lemma 1.2.3(3) readily implies that

$$\mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1] = \text{smd}(\mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1]).$$

Thus, it is enough to prove that, given $\ell > 0$ and $0 \leq n_1, \dots, n_\ell < d$, the inclusion

$$\mathcal{C}[n_1] * \dots * \mathcal{C}[n_\ell] \subseteq \mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1]$$

holds. We prove this claim by induction on ℓ . For $\ell = 1$ the claim is obvious, so suppose that the claim holds for some $\ell > 0$. Let $0 \leq n_1, \dots, n_\ell, n < d$ and notice that, by the inductive hypothesis, the inclusion

$$(\mathcal{C}[n_1] * \dots * \mathcal{C}[n_\ell]) * \mathcal{C}[n] \subseteq (\mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1]) * \mathcal{C}[n].$$

holds. We distinguish two cases:

$n = 0$: Since $\mathcal{C}[d-1] = (\mathcal{C}[d])[-1] = \mathcal{C}[-1]$, the equality

$$\mathcal{C}[d-1] * \mathcal{C}[n] = \mathcal{C}[-1] * \mathcal{C}$$

holds. Moreover, by assumption,

$$(\mathcal{C}[-1] * \mathcal{C}) \subseteq (\mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1]).$$

Consequently,

$$\begin{aligned} (\mathcal{C}[n_1] * \dots * \mathcal{C}[n_\ell]) * \mathcal{C}[n] &\subseteq (\mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1]) * \mathcal{C}[n] \\ &= (\mathcal{C} * \dots * \mathcal{C}[d-2]) * (\mathcal{C}[d-1] * \mathcal{C}[n]) \\ &= (\mathcal{C} * \dots * \mathcal{C}[d-2]) * (\mathcal{C}[-1] * \mathcal{C}) \\ &\subseteq (\mathcal{C} * \dots * \mathcal{C}[d-2]) * (\mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1]) \\ &\subseteq (\mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-2]) * \mathcal{C}[d-1], \end{aligned}$$

where the last inclusion is obtained from the inclusions

$$\mathcal{C}[i] * \mathcal{C}[j] \subseteq \mathcal{C}[j] * \mathcal{C}[i]$$

for $0 \leq j < i < d$ with $(j, i) \neq (0, d-1)$ (see Lemma 1.3.6) and the equality

$$\mathcal{C}[i] * \mathcal{C}[i] = \mathcal{C}[i] \vee \mathcal{C}[i] = \mathcal{C}[i],$$

(see Lemma 1.2.3(2)) which stems from the fact that there are no non-trivial self-extensions between the objects of $\mathcal{C}[i]$ (recall that $d \geq 2$ by assumption). This proves the claim in this case.

$n \neq 0$: A simpler version of the argument used in the case $n = 0$ shows that there are inclusions

$$\begin{aligned} \mathcal{C}[n_1] * \dots * \mathcal{C}[n_\ell] * \mathcal{C}[n] &\subseteq (\mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1]) * \mathcal{C}[n] \\ &\subseteq \mathcal{C} * \mathcal{C}[1] * \dots * \mathcal{C}[d-1], \end{aligned}$$

where in the last inclusion we use Lemma 1.3.6 and Lemma 1.2.3(2) and the assumption that $n \neq 0$.

In view of [Theorem 1.3.7](#), this finishes the proof of the theorem. \square

Remark 1.3.10. In the context of [Theorem 1.3.9](#), condition (2d) is not essential: We may replace \mathcal{T} by its full subcategory $\text{thick}(\mathcal{C})$, noticing that conditions (2a)–(2c) only depend on the latter subcategory.

2. $(d + 2)$ -ANGULATED CATEGORIES

2.1. Definition and basic properties. Although we do not use most axioms in the definition (at least not explicitly), we include the complete definition of a $(d + 2)$ -angulated category since it plays an important conceptual role in our main results. Recall that $d \geq 1$ is a fixed integer.

Definition 2.1.1 ([\[GKO13, Def. 2.1\]](#) and [\[BT13, Thm. 4.2\]](#)). Let (\mathcal{F}, Σ) be a pair consisting of an additive category \mathcal{F} and an automorphism¹ $\Sigma: \mathcal{F} \xrightarrow{\sim} \mathcal{F}$. A $(d + 2)$ -angulation of (\mathcal{F}, Σ) is a class \diamond of sequences

$$x_1 \rightarrow x_2 \rightarrow \cdots \rightarrow x_{d+2} \rightarrow \Sigma(x_1)$$

that satisfy the following axioms:

- (*d*-TR1) (a) The class \diamond is closed under direct sums and direct summands.
 (b) For each object $x \in \mathcal{F}$, the sequence

$$x \xrightarrow{1} x \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \Sigma(x)$$

(*d* consecutive zeroes) lies in \diamond .

- (c) Every morphism $x \rightarrow y$ in \mathcal{F} is the *leftmost* morphism in a sequence that lies in \diamond .

- (*d*-TR2) A sequence

$$x_1 \xrightarrow{f} x_2 \rightarrow \cdots \rightarrow x_{d+1} \rightarrow \Sigma(x_1)$$

lies in \diamond if and only if its *left rotation*

$$x_2 \rightarrow x_3 \rightarrow \cdots \rightarrow x_{d+2} \rightarrow \Sigma(x_1) \xrightarrow{(-1)^d \Sigma(f)} \Sigma(x_2)$$

lies in \diamond (notice that the sign in the rotation depends on *d*).

- (*d*-TR3) Given a solid diagram in \mathcal{F} of the form

$$\begin{array}{ccccccccccc} x_1 & \rightarrow & x_2 & \rightarrow & x_3 & \rightarrow & \cdots & \rightarrow & x_{d+1} & \rightarrow & x_{d+2} & \rightarrow & \Sigma(x_1) \\ \downarrow u & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \Sigma(u) \\ y_1 & \rightarrow & y_2 & \rightarrow & y_3 & \rightarrow & \cdots & \rightarrow & y_{d+1} & \rightarrow & y_{d+2} & \rightarrow & \Sigma(y_1), \end{array}$$

in which the leftmost square commutes and both rows lie in \diamond , there exist morphisms $x_i \rightarrow y_i$, $3 \leq i \leq d + 2$, rendering the diagram commutative.

- (*d*-TR4) Given the solid part of the diagram

$$\begin{array}{ccccccccccc} x_1 & \rightarrow & x_2 & \xrightarrow{u} & x_3 & \rightarrow & x_4 & \rightarrow & \cdots & \rightarrow & x_{d+1} & \rightarrow & x_{d+2} & \rightarrow & \Sigma(x_1) \\ \parallel & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \parallel \\ x_1 & \rightarrow & y_2 & \rightarrow & y_3 & \rightarrow & y_4 & \rightarrow & \cdots & \rightarrow & y_{d+1} & \rightarrow & y_{d+2} & \rightarrow & \Sigma(x_1) \\ \downarrow f & & \parallel & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \Sigma(f) \\ x_1 & \rightarrow & y_2 & \rightarrow & z_3 & \rightarrow & z_4 & \rightarrow & \cdots & \rightarrow & z_{d+1} & \rightarrow & z_{d+2} & \xrightarrow{v} & \Sigma(x_1) \end{array}$$

¹As in the case of triangulated categories, one may assume instead that Σ is merely an auto-equivalence, but every autoequivalence can be strictified by replacing \mathcal{F} by an equivalent category, see for example [\[KV87, Sec. 2\]](#). We mostly ignore this distinction as it is not crucial to our results.

whose rows lie in \diamond , a collection of dotted morphisms rendering the diagram commutative exists, and with the further property that the natural complex (see [BT13, Sec. 4] for the precise signs)

$$\begin{aligned} x_3 &\rightarrow x_4 \oplus y_3 \rightarrow x_5 \oplus y_4 \oplus z_3 \rightarrow \cdots \\ \cdots &\rightarrow x_{d+2} \oplus y_{d+1} \oplus z_d \rightarrow y_{d+2} \oplus z_{d+1} \rightarrow z_{d+2} \xrightarrow{\Sigma(u)_{\text{ov}}} \Sigma(x_3) \end{aligned}$$

lies in \diamond .

In this case, the triple $(\mathcal{F}, \Sigma, \diamond)$ is called a $(d+2)$ -angulated category. Similarly, we say that \diamond is a *pre- $(d+2)$ -angulation* if it satisfies axioms $(d\text{-TR1})$ – $(d\text{-TR3})$, in which case the triple $(\mathcal{F}, \Sigma, \diamond)$ is a *pre- $(d+2)$ -angulated category*.

Notation 2.1.2. Let $(\mathcal{F}, \Sigma, \diamond)$ be a (pre-) $(d+2)$ -angulated category. If there is no risk of confusion, we often abuse the notation and identify the triple $(\mathcal{F}, \Sigma, \diamond)$ with its underlying additive category \mathcal{F} .

Remark 2.1.3. In [Definition 2.1.1](#) we use the equivalent axioms given by Bergh and Thaule [BT13], that is, we use a suitable analogue of octahedral axiom instead of a higher analogue of Neeman’s mapping cone axiom for triangulated categories [Nee01, Def. 1.3.13] used in the original definition of Geiss, Keller and Oppermann. The higher analogue of Neeman’s mapping cone axiom is used in [Example 2.1.9](#) below, so we include it here:

$(d\text{-TR4}')$ In the situation of $(d\text{-TR3})$, the morphisms $x_i \rightarrow y_i$, $3 \leq i \leq d+2$, can be chosen such that the usual mapping cone (see [GKO13, 2.1] for the precise form of the morphisms)

$$x_2 \oplus y_1 \rightarrow x_3 \oplus y_2 \rightarrow \cdots \rightarrow \Sigma x_1 \oplus y_{d+2} \rightarrow \Sigma(x_2 \oplus y_1)$$

lies in \diamond .

Remark 2.1.4. A 3-angulated category is nothing else but a triangulated category. Indeed, for $d = 1$, axioms $(d\text{-TR1})$ – $(d\text{-TR4})$ are identical to axioms (TR1)–(TR4) in the usual definition of a triangulated category.

Remark 2.1.5. We do not know examples of pre- $(d+2)$ -angulated categories that do not satisfy axiom $(d\text{-TR4})$ for any $d \geq 1$. In fact, no such examples are known even in the classical case of triangulated categories.

Remark 2.1.6. Given a $(d+2)$ -angulated category $(\mathcal{F}, \Sigma, \diamond)$ and a unit $u \in \mathbf{k}^\times$ in the ground field, we can form a new $(d+2)$ -angulation $u \cdot \diamond$ of (\mathcal{F}, Σ) as follows:

$$x_1 \xrightarrow{f} x_2 \rightarrow \cdots x_{d+1} \rightarrow \Sigma(x_1)$$

belongs to \diamond if and only if

$$x_1 \xrightarrow{u^{-1} \cdot f} x_2 \rightarrow \cdots x_{d+1} \rightarrow \Sigma(x_1)$$

belongs to $u \cdot \diamond$ (see [Bal02] for the case $d = 1$ of ordinary triangulations).

Remark 2.1.7. For the purposes of this article, it is enough to consider only $(d+2)$ -angulated categories whose underlying additive category has split idempotents. This is not a serious restriction, since $(d+2)$ -angulations can be extended to idempotent completions in an essentially unique way, see [BS01] for the case $d = 1$ of triangulated categories and [Lin21] for the general case.

The following result is a $(d+2)$ -angulated analogue of a classical result of Freyd [Fre66] and of Heller’s parametrisation of the pre-triangulations on a pre-triangulated category [Hel68].

Proposition 2.1.8 ([GKO13, Prop. 2.5 and Prop. 3.4]). *Let (\mathcal{F}, Σ) be a pair consisting of an additive category and an automorphism $\Sigma: \mathcal{F} \xrightarrow{\sim} \mathcal{F}$. Suppose that (\mathcal{F}, Σ) admits a pre- $(d+2)$ -angulation. Then, the following statements hold:*

- (1) *The category $\mathbf{mod}(\mathcal{F})$ of finitely-presented functors $\mathcal{F}^{\text{op}} \rightarrow \mathbf{Mod} \mathbf{k}$ to vector spaces is a Frobenius abelian category. Moreover, if idempotents split in \mathcal{F} , then the Yoneda embedding $x \mapsto \mathcal{F}(-, x)$ induces an equivalence between \mathcal{F} and the full subcategory of $\mathbf{mod}(\mathcal{F})$ spanned by the projective-injective objects.*
- (2) *The automorphism of $\Sigma: \mathbf{mod}(\mathcal{F}) \xrightarrow{\sim} \mathbf{mod}(\mathcal{F})$ induced by $\Sigma: \mathcal{F} \xrightarrow{\sim} \mathcal{F}$ is an exact functor. In particular, there is an induced natural isomorphism*

$$\sigma: \Sigma \Omega_{\mathcal{F}}^{-1} \xrightarrow{\sim} \Omega_{\mathcal{F}}^{-1} \Sigma,$$

where $\Omega_{\mathcal{F}}^{-1}: \mathbf{mod}(\mathcal{F}) \xrightarrow{\sim} \mathbf{mod}(\mathcal{F})$ is a choice of cosyzygy functor on the stable category, that makes the pair (Σ, σ) into an exact functor on $\mathbf{mod}(\mathcal{F})$.

- (3) *There is a (canonical) bijective map from the class of pre- $(d+2)$ -angulations on (\mathcal{F}, Σ) to the class of isomorphisms of exact functors*

$$(\Sigma, \sigma) \xrightarrow{\sim} (\Omega_{\mathcal{F}}^{-(d+2)}, (-1)^d \mathbf{1}_{\Omega_{\mathcal{F}}^{-(d+2)-1}})$$

on $\mathbf{mod}(\mathcal{F})$. In particular, if the pair (\mathcal{F}, Σ) admits a pre- $(d+2)$ -angulation, then there are natural isomorphisms

$$\Sigma \cong \Omega_{\mathcal{F}}^{-(d+2)} \quad \text{and} \quad \Sigma^{-1} \cong \Omega_{\mathcal{F}}^{d+2}$$

of exact functors on $\mathbf{mod}(\mathcal{F})$.

The following example, that to our knowledge has not yet appeared in the literature, provides a simple method to constructing ‘longer’ angulations from ‘shorter’ ones.

Example 2.1.9. Let $(\mathcal{F}, \Sigma, \diamond)$ be a $(d+2)$ -angulated category and $n \geq 1$ an integer. The pair (\mathcal{F}, Σ^n) is endowed with an $n(d+2)$ -angulation \diamond^n consisting of the sequences

$$\begin{array}{ccccccc} x_1 & \longrightarrow & x_2 & \longrightarrow & \cdots & \longrightarrow & x_{d+2} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Sigma(x_1) & \longrightarrow & \Sigma(x_2) & \longrightarrow & \cdots & \longrightarrow & \Sigma(x_{d+2}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \cdots & & \cdots & & \cdots & & \cdots \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Sigma^{n-1}(x_1) & \longrightarrow & \Sigma^{n-1}(x_2) & \longrightarrow & \cdots & \longrightarrow & \Sigma^{n-1}(x_{d+2}) \longrightarrow \Sigma^n(x_1) \end{array}$$

in which the first sequence

$$x_1 \rightarrow x_2 \rightarrow \cdots \rightarrow x_{d+2} \rightarrow \Sigma(x_1)$$

lies in \diamond and the subsequent rows of the diagram are obtained from the first one by iterated application of the suspension functor (without any additional signs). Utilising the fact that \diamond is a $(d+2)$ -angulation of (\mathcal{F}, Σ) , it is straightforward to verify that \diamond^n is an $n(d+2)$ -angulation of (\mathcal{F}, Σ^n) . Indeed, in terms of **Proposition 2.1.8(3)**, if the class \diamond is classified by an isomorphism $\Sigma \cong \Omega_{\mathcal{F}}^{d+2}$ of exact functors on $\mathbf{mod}(\mathcal{F})$, then the class \diamond^n is the pre- $n(d+2)$ -angulation of (\mathcal{F}, Σ^n) classified by the induced isomorphism $\Sigma^n \cong \Omega_{\mathcal{F}}^{n(d+2)}$. Thus, to prove that \diamond^n is a $(d+2)$ -angulation, it suffices to verify that the higher analogue of Neeman’s mapping cone axiom is satisfied (**Remark 2.1.3**). We leave the details of this verification to the reader.

We record the apparent notion of equivalence between (pre-) $(d+2)$ -angulated categories for later use. For this, we also introduce the following auxiliary definition.

Definition 2.1.10. Let $(\mathcal{C}, \Phi_{\mathcal{C}})$ and $(\mathcal{D}, \Phi_{\mathcal{D}})$ be pairs consisting of categories \mathcal{C} and \mathcal{D} and automorphisms $\Phi_{\mathcal{C}}: \mathcal{C} \xrightarrow{\sim} \mathcal{C}$ and $\Phi_{\mathcal{D}}: \mathcal{D} \xrightarrow{\sim} \mathcal{D}$. A *functor of pairs* $(\mathcal{C}, \Phi_{\mathcal{C}}) \rightarrow (\mathcal{D}, \Phi_{\mathcal{D}})$ is a pair (F, F^{\sharp}) consisting of a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ and a natural isomorphism $F^{\sharp}: F\Phi_{\mathcal{C}} \Rightarrow \Phi_{\mathcal{D}}F$. A functor of pairs is an *equivalence* if its underlying functor is an equivalence of categories.

Remark 2.1.11. Observe that the collection of pairs (as in [Definition 2.1.10](#)) form a 2-category in a natural way. In particular, the notion of equivalence of pairs is simply the corresponding notion of equivalence in this 2-category.

Notation 2.1.12. In the context of [Definition 2.1.10](#), we always identify a functor of pairs with the functor between the corresponding underlying categories.

Definition 2.1.13. Let $(\mathcal{F}, \Sigma, \diamond)$ and $(\mathcal{F}', \Sigma', \diamond')$ be two pre- $(d+2)$ -angulated categories. A *pre- $(d+2)$ -angulated equivalence* between $(\mathcal{F}, \Sigma, \diamond)$ and $(\mathcal{F}', \Sigma', \diamond')$ is an equivalence of pairs

$$F: (\mathcal{F}, \Sigma) \xrightarrow{\sim} (\mathcal{F}', \Sigma')$$

such that, for each $(d+2)$ -angle

$$x_1 \rightarrow x_2 \rightarrow \cdots \rightarrow x_{d+2} \xrightarrow{\eta} \Sigma(x_0)$$

in \diamond , the sequence

$$F(x_1) \rightarrow F(x_2) \rightarrow \cdots \rightarrow F(x_{d+2}) \xrightarrow{\eta'} \Sigma(F(x_0))$$

lies in \diamond' , where $\eta' = F^{\sharp}_x F(\eta)$. If \diamond and \diamond' are in fact $(d+2)$ -angulations, we say that F is a *$(d+2)$ -angulated equivalence* or an *equivalence of $(d+2)$ -angulated categories*.

2.2. Constructions of $(d+2)$ -angulated categories. We recall the two constructions of $(d+2)$ -angulated categories that are most relevant for the purposes of this article.

2.2.1. Standard $(d+2)$ -angulated categories. The following theorem of Geiss, Keller and Oppermann is one of the main sources of $(d+2)$ -angulated categories; in fact, this theorem was the main motivation for introducing the notion of $(d+2)$ -angulated category, and it also motivated the definition of a $d\mathbb{Z}$ -cluster tilting subcategory ([Definition 1.3.1](#)).

Theorem 2.2.1 ([GKO13, Thm. 1]). *Let $\mathcal{C} \subseteq \mathcal{T}$ be a $d\mathbb{Z}$ -cluster tilting subcategory of \mathcal{T} . Then, the pair $(\mathcal{C}, [d])$ is endowed with a $(d+2)$ -angulation whose $(d+2)$ -angles are those sequences*

$$c_{d+2} \rightarrow c_{d+1} \rightarrow \cdots \rightarrow c_2 \rightarrow c_1 \rightarrow c_{d+2}[d]$$

in \mathcal{C} which fit as the spine of a commutative diagram of the form

$$\begin{array}{ccccccc} & & c_{d+1} & \cdots & & c_3 & \xrightarrow{\quad} & c_2 \\ & \nearrow & & & \searrow & \nearrow & & \searrow \\ c_{d+2} & \xleftarrow{+1} & x_{d,5} & \cdots & x_{3,5} & \xleftarrow{+1} & x_{2,5} & \xleftarrow{+1} & c_1 \end{array}$$

in which the oriented triangles are exact triangles in \mathcal{T} and such that the connecting morphism $c_0 \rightarrow c_{d+2}[d]$ is given by the obvious (shifted) composite along the bottom row of the diagram.

Definition 2.2.2. We call a $(d+2)$ -angulated category $(\mathcal{F}, \Sigma, \diamond_{\mathcal{F}})$ *standard* if there exists a fully faithful functor of pairs $\iota: (\mathcal{F}, \Sigma) \hookrightarrow (\mathcal{T}, [d])$ with the following properties:

- (1) The category \mathcal{T} is triangulated with suspension functor $[1]$.
- (2) The essential image $\iota(\mathcal{F})$ of ι is a d -rigid subcategory of \mathcal{T} .

- (3) The image of the $(d+2)$ -angulation of \mathcal{F} under ι coincides with the class of sequences considered in [Theorem 2.2.1](#). More precisely, the closure under isomorphisms of the class of sequences

$$F(x_1) \rightarrow F(x_2) \rightarrow \cdots \rightarrow F(x_{d+2}) \xrightarrow{F_{x_1}^\# F(\eta)} \Sigma(F(x_1))$$

in $\iota(\mathcal{F})$ that are the image of a $(d+2)$ -angle

$$x_1 \rightarrow x_2 \rightarrow \cdots \rightarrow x_{d+2} \xrightarrow{\eta} \Sigma(x_1)$$

that lies in the class $\diamond_{\mathcal{F}}$ coincides with the class of sequences considered in [Theorem 2.2.1](#).

In this case, we say that $\iota(\mathcal{F}) \subseteq \mathcal{T}$ is a *standard $(d+2)$ -angulated subcategory* and call the class $\diamond_{\mathcal{F}}$ a *standard $(d+2)$ -angulation*.

Remark 2.2.3. Let $(\mathcal{F}, \Sigma, \diamond)$ be a standard $(d+2)$ -angulated category with respect to a fully faithful functor $\iota: \mathcal{F} \hookrightarrow \mathcal{T}$ into a triangulated category \mathcal{T} . Since ι is a functor of pairs $\iota(\mathcal{F})[d] = \iota(\mathcal{F})$ and since moreover $\iota(\mathcal{F})$ is d -rigid then $\iota(\mathcal{F})$ is actually a $d\mathbb{Z}$ -rigid subcategory of \mathcal{T} .

[Theorems 1.3.9](#) and [2.2.1](#) combine into the following result.

Theorem 2.2.4. *Suppose that \mathcal{T} has finite-dimensional morphism spaces and split idempotents. Let $\mathcal{C} \subseteq \mathcal{T}$ be a subcategory that satisfies $\mathbf{add}(\mathcal{C}) = \mathcal{C}$ and $\iota: \mathcal{C} \hookrightarrow \mathcal{T}$ the inclusion functor. Then, the following statements are equivalent:*

- (1) *The subcategory \mathcal{C} is a $d\mathbb{Z}$ -cluster tilting subcategory of \mathcal{T} .*
- (2) *$\mathbf{thick}(\mathcal{C}) = \mathcal{T}$ and the functor ι exhibits \mathcal{C} as a standard $(d+2)$ -angulated category; in other words, the pair $(\mathcal{C}, [d])$ admits a $(d+2)$ -angulation whose $(d+2)$ -angles are given as in [Theorem 2.2.1](#).*

Proof. The implication (1) \Rightarrow (2) is precisely [Theorem 2.2.1](#). We prove the converse implication (2) \Rightarrow (1). By assumption, \mathcal{C} is a $d\mathbb{Z}$ -rigid subcategory such that $\mathbf{thick}(\mathcal{C}) = \mathcal{T}$ and $\mathcal{C} = \mathcal{C}[d]$. In view of [Theorem 1.3.9](#), it is enough to prove that the inclusion

$$(\mathcal{C}[-1] * \mathcal{C}) \subseteq (\mathcal{C} * \mathcal{C}[1] * \cdots * \mathcal{C}[d-1])$$

holds. Indeed, let $x \in \mathcal{C}[-1] * \mathcal{C}$ and choose an exact triangle in \mathcal{T} of the form

$$x \rightarrow c_2 \xrightarrow{f} c_1 \rightarrow x[1]$$

with $c_1, c_2 \in \mathcal{C}$. By axioms [\(d-TR1\)](#) and [\(d-TR2\)](#), there exists a $(d+2)$ -angle in \mathcal{C} of the form

$$c_{d+2} \rightarrow c_{d+1} \rightarrow \cdots \rightarrow c_2 \xrightarrow{f} c_1 \rightarrow c_{d+2}[d].$$

Thus, since ι exhibits \mathcal{C} as a standard $(d+2)$ -angulated category, there must exist a commutative diagram of the form

$$\begin{array}{ccccccc} & & c_{d+1} & & \cdots & & c_3 & \xrightarrow{\quad} & c_2 & & \\ & \nearrow & & \searrow & & \nearrow & & \searrow & \nearrow & \searrow & \\ c_{d+2} & \xleftarrow{+1} & & \xrightarrow{+1} & x_{d,5} & \cdots & x_{3,5} & \xleftarrow{+1} & & \xrightarrow{+1} & x_{2,5} & \xleftarrow{+1} & & c_1 \end{array}$$

where $x = x_{2,5}$. This shows that $x \in \mathcal{C} * \mathcal{C}[1] * \cdots * \mathcal{C}[d-1]$, as required, see [Example 1.2.4](#). This finishes the proof of the theorem. \square

Remark 2.2.5. In the context of [Theorem 2.2.4](#), the condition

$$\mathbf{thick}(\mathcal{C}) = \mathcal{T}$$

is not essential: We may replace \mathcal{T} by its full subcategory $\mathbf{thick}(\mathcal{C})$, noticing that statement (2) only depends on the latter subcategory (compare with [Theorem 1.3.9\(2a\)](#)).

2.2.2. *Amiot–Lin $(d+2)$ -angulations.* In addition to [Theorem 2.2.1](#), an important source of $(d+2)$ -angulated categories is the following ‘higher-dimensional’ version of a theorem of Amiot, who proved the result in the case $d = 1$. We need some preliminaries to state the result.

Given an algebra Λ , we let $\Lambda^e = \Lambda \otimes \Lambda^{\text{op}}$ be its enveloping algebra and identify the category of Λ -bimodules with that of (right) Λ^e -modules in the canonical way. Recall that a finite-dimensional self-injective algebra Λ is *twisted n -periodic* if there exist an algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$ and an isomorphism

$$\Omega_{\Lambda^e}^n(\Lambda) \cong {}_1\Lambda_\sigma$$

in the stable category of Λ -bimodules; here, ${}_1\Lambda_\sigma$ denotes the diagonal Λ -bimodule with right action twisted by σ . Equivalently, Λ is twisted n -periodic if there exists an exact sequence of Λ -bimodules

$$0 \rightarrow {}_1\Lambda_\sigma \rightarrow P_n \rightarrow \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow \Lambda \rightarrow 0$$

whose middle terms are projective Λ -bimodules. Similarly, Λ is *n -periodic* if it is twisted n -periodic with $\sigma = \mathbf{1}_\Lambda$.

On the other hand, let Λ be a basic self-injective algebra. By [\[Bol84, Prop. 3.8\]](#), the map

$$\text{Out}(\Lambda) \longrightarrow \text{Pic}(\Lambda), \quad [\sigma] \longmapsto [{}_1\Lambda_\sigma],$$

yields an isomorphism between the outer automorphism group $\text{Out}(\Lambda)$ of Λ and the Picard group $\text{Pic}(\Lambda)$ of invertible Λ -bimodules. In particular, every autoequivalence of the additive category $\text{proj}(\Lambda)$ is of the form

$$- \otimes_{\Lambda} {}_1\Lambda_\sigma: \text{proj}(\Lambda) \xrightarrow{\sim} \text{proj}(\Lambda)$$

for some automorphism σ of Λ ; notice also that the above autoequivalence is isomorphic to the restriction of scalars

$$(-)_\sigma: P \longmapsto P_\sigma$$

along σ . In particular, Λ is twisted n -periodic if and only if the Λ -bimodule $\Omega_{\Lambda^e}^n(\Lambda)$ is stably isomorphic to an invertible Λ -bimodule.

We record the following observation for later use. Recall that the syzygy $\Omega(M)$ of a module M is the kernel of its projective cover.

Proposition 2.2.6. *Let Λ be a finite-dimensional self-injective algebra that is twisted $(d+2)$ -periodic. Suppose that Λ is connected and non-separable. Then, the following statements hold:*

- (1) *Every invertible Λ -bimodule I that is isomorphic to $\Omega_{\Lambda^e}^{d+2}(\Lambda)$ in $\underline{\text{mod}}(\Lambda^e)$ is already isomorphic to $\Omega_{\Lambda^e}^{d+2}(\Lambda)$ as a Λ -bimodule.*
- (2) *The $(d+2)$ -syzygy $\Omega_{\Lambda^e}^{d+2}(\Lambda)$ is an invertible Λ -bimodule.*

Proof. The proof of [\[Mur22, Prop. 9.8\]](#) applies verbatim. \square

Recall that a finite-dimensional algebra Λ is *separable* if it is isomorphic to a finite product of matrix algebras over skew fields, whose centres are separable field extensions of the ground field \mathbf{k} ; in particular, if \mathbf{k} is perfect, then Λ is separable if and only if it is semisimple (see for example [\[Wei94, Sec. 9.2.1\]](#)). The following theorem of Green, Snashall and Solberg (with some simplifications due to Hanihara) gives a convenient characterisation of the class of twisted periodic algebras, see also the comments in the proof of [\[CDIM, Prop. 3.5\]](#).

Theorem 2.2.7 ([\[GSS03, Thm. 1.4\]](#) and [\[Han20, Cor. 2.2\]](#)). *Let Λ be a finite-dimensional self-injective algebra and $n \geq 1$ an integer; suppose, moreover, that Λ/J_Λ is separable. The following statements are equivalent:*

- (1) *The algebra Λ is twisted n -periodic.*

(2) There exist an algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$ and a natural isomorphism

$$\Omega_{\Lambda}^n \cong (-)_{\sigma}$$

of exact functors $\underline{\text{mod}}(\Lambda) \xrightarrow{\sim} \underline{\text{mod}}(\Lambda)$.

Remark 2.2.8. Suppose that the equivalent conditions in [Theorem 2.2.7](#) are satisfied and that Λ is connected and non-separable. [Proposition 2.2.6](#) shows that $\Omega_{\Lambda^e}^{d+2}(\Lambda)$ is an invertible Λ -bimodule but it need not be isomorphic to ${}_1\Lambda_{\sigma}$. Although these bimodules induce isomorphic functors on $\underline{\text{mod}}\Lambda$, this only permits us to conclude that $\Omega_{\Lambda^e}^{d+2}(\Lambda) \cong {}_1\Lambda_{\sigma\gamma}$ where γ is a *stable* inner automorphism, that is an algebra automorphism such that the restriction of scalars along γ is isomorphic to the identity functor of the stable category $\underline{\text{mod}}\Lambda$ (compare with [IV14, Thm. 1.8] for example).

Our interest in twisted periodic algebras in the context of $(d+2)$ -angulated categories stems from the following observation, which is essentially a reformulation of [Proposition 2.1.8](#).

Proposition 2.2.9. *Let $(\mathcal{F}, \Sigma, \diamond)$ be a $(d+2)$ -angulated category with finite-dimensional morphism spaces and split idempotents. Suppose that there exists a basic object $x \in \mathcal{F}$ such that $\text{add}(x) = \mathcal{F}$ and set $\Lambda := \mathcal{F}(x, x)$. If Λ/J_{Λ} is separable, then Λ is self-injective and twisted $(d+2)$ -periodic.*

Proof. The proof is an adaptation of the proof of [Han20, Prop. 3.1], which treats the case $d = 1$ of triangulated categories. By assumption, the functor

$$\mathcal{F}(x, -): \mathcal{F} \longrightarrow \text{proj}(\Lambda), \quad y \longmapsto \mathcal{F}(x, y)$$

is an equivalence of categories; hence the finite-dimensional algebra Λ is self-injective by [Proposition 2.1.8\(1\)](#). In particular, there are commutative diagrams of equivalences of categories

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{\sim} & \text{proj}(\Lambda) \\ \downarrow \Sigma^{-1} & & \downarrow (-)_{\sigma} \\ \mathcal{F} & \xrightarrow{\sim} & \text{proj}(\Lambda) \end{array} \quad \text{and} \quad \begin{array}{ccc} \underline{\text{mod}}(\mathcal{F}) & \xrightarrow{\sim} & \underline{\text{mod}}(\Lambda) \\ \downarrow \Sigma^{-1} & & \downarrow (-)_{\sigma} \\ \underline{\text{mod}}(\mathcal{F}) & \xrightarrow{\sim} & \underline{\text{mod}}(\Lambda), \end{array}$$

where $\sigma: \Lambda \xrightarrow{\sim} \Lambda$ is an algebra automorphism (compare [Bol84, Prop. 3.8]) and the functors in the diagram on the right are exact with respect to the induced triangulated structures. [Proposition 2.1.8\(3\)](#) yields the existence of a natural isomorphism $\Sigma^{-1} \cong \Omega_{\mathcal{F}}^{d+2}$ of exact functors on $\underline{\text{mod}}(\mathcal{F})$ and, consequently, there is also a natural isomorphism $(-)_{\sigma} \cong \Omega_{\Lambda}^{d+2}$ of exact functors on $\underline{\text{mod}}(\Lambda)$. [Theorem 2.2.7](#) then implies that Λ is twisted $(d+2)$ -periodic, which is what we needed to prove. \square

Combining [Theorem 2.2.1](#) and [Proposition 2.2.9](#), we obtain an alternative proof of the following result that emphasises the role of $(d+2)$ -angulations in this context.

Proposition 2.2.10 ([Dug12], [CDIM, Prop. 8.5(a)]). *Let \mathcal{T} be a triangulated category with finite-dimensional morphism spaces and split idempotents and such that there exists a $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$. Set $\Lambda := \mathcal{T}(c, c)$. If Λ/J_{Λ} is separable, then Λ is self-injective and twisted $(d+2)$ -periodic.*

Proof. Indeed, [Theorem 2.2.1](#) shows that the additive category $\mathcal{C} := \text{add}(c) \subseteq \mathcal{T}$ is endowed with the structure of a $(d+2)$ -angulated category with suspension functor $[d]: \mathcal{C} \xrightarrow{\sim} \mathcal{C}$. Both claims then follows from [Proposition 2.2.9](#). \square

Remark 2.2.11. In [CDIM, Prop. 8.5(a)], the authors assume that the ambient triangulated category admits a Serre functor in the sense of [BK89], but this fact is not used in their proof.

Remark 2.2.12. **Proposition 2.2.10** shows that the correspondence in **Theorem A** is well defined.

Construction 2.2.13. Let Λ be a basic finite-dimensional self-injective algebra and σ an automorphism. Notice that Λ and Λ^{op} are Frobenius algebras [Zim14, Prop. 4.5.7] and therefore so is their tensor product Λ^e ; in particular the projective Λ -bimodules are also injective Λ -bimodules. Consider an exact sequence of Λ -bimodules

$$\eta: 0 \rightarrow {}_1\Lambda_\sigma \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

with P_i projective-injective as Λ -bimodules for $0 \leq i < d+1$. In particular η , regarded as a complex of left Λ -modules, is contractible². We will be mainly interested in the case that Λ is twisted $(d+2)$ -periodic with respect to σ and P_{d+1} is also a projective-injective Λ -bimodule, so that η exhibits the existence of an isomorphism $\Omega_{\Lambda^e}^{d+2}(\Lambda) \cong {}_1\Lambda_\sigma$ in $\underline{\text{mod}}(\Lambda^e)$. Let

$$\Sigma := - \otimes_{\Lambda} \sigma \Lambda_1: \text{proj}(\Lambda) \xrightarrow{\sim} \text{proj}(\Lambda)$$

and

$$\Sigma^{-1} := - \otimes_{\Lambda} {}_1\Lambda_\sigma: \text{proj}(\Lambda) \xrightarrow{\sim} \text{proj}(\Lambda)$$

its quasi-inverse. We define a class \diamond_η of *exact $(d+2)$ -angles* in $\text{proj}(\Lambda)$ as follows: A complex of finitely generated projective-injective Λ -modules

$$Q_{d+2} \xrightarrow{f} Q_{d+1} \rightarrow \cdots \rightarrow Q_1 \xrightarrow{g} \Sigma Q_{d+2}$$

lies in \diamond_η if it has the following two properties:

- (1) The extended complex

$$Q_{d+2} \xrightarrow{f} Q_{d+1} \rightarrow \cdots \rightarrow Q_1 \xrightarrow{g} \Sigma Q_{d+2} \xrightarrow{\Sigma f} \Sigma Q_{d+1}$$

is exact (notice that this condition only depends on the pair $(\text{proj}(\Lambda), \Sigma)$).

- (2) Denote the Λ -module $\text{coker } g$ by N . By construction, there is an exact sequence

$$0 \rightarrow \Sigma^{-1}N \xrightarrow{i} Q_{d+1} \rightarrow \cdots \rightarrow Q_1 \xrightarrow{g} \Sigma Q_{d+2} \xrightarrow{p} N \rightarrow 0$$

such that $f = i \circ \Sigma^{-1}p$. We require the above exact sequence to be equivalent to the exact sequence

$$N \otimes_{\Lambda} (0 \rightarrow {}_1\Lambda_\sigma \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0)$$

in the extension space $\text{Ext}_{\Lambda^e}^{d+2}(N, {}_1\Lambda_\sigma)$. The latter sequence is exact because, as we have pointed out above, η is contractible as a complex of left Λ -modules.

Remark 2.2.14. In connection with **Remark 2.1.6**, given an exact sequence of Λ -bimodules

$$\eta: 0 \rightarrow {}_1\Lambda_\sigma \xrightarrow{\iota} P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

and a unit $u \in \mathbf{k}^\times$ in the ground field the class

$$u \cdot [\eta] \in \text{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$$

is represented by the exact sequence of Λ -bimodules

$$u \cdot \eta: 0 \rightarrow {}_1\Lambda_\sigma \xrightarrow{u^{-1} \cdot \iota} P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0.$$

Therefore, the class $\diamond_{u \cdot \eta}$ given by **Construction 2.2.13** coincides with $u \cdot \diamond_\eta$.

²It suffices to assume that P_i is left projective for $0 \leq i < d+1$ (as a consequence P_{d+1} too), so that the tensored sequence in (2) remains exact. This hypothesis is missing in [Mur20b, Prop. 5.6 and Rmk. 5.7]. Nevertheless, any extension has representatives satisfying this assumption, such as the representative considered in the proof of [Mur20b, Prop. 5.6].

We are ready to state the theorem(s) of Amiot and Lin.

Theorem 2.2.15 ([Ami07, Thm. 8.1] and [Lin19, Thm. 1.3]). *Let Λ be a basic finite-dimensional algebra that is twisted $(d+2)$ -periodic with respect to an algebra automorphism σ and let*

$$\Sigma := - \otimes_{\Lambda} \sigma \Lambda_1 : \text{proj}(\Lambda) \xrightarrow{\sim} \text{proj}(\Lambda).$$

Then, the class \diamond_{η} of exact $(d+2)$ -angles from [Construction 2.2.13](#) associated to an extension η with projective-injective middle Λ -bimodules endows the pair $(\text{proj}(\Lambda), \Sigma)$ with the structure of a $(d+2)$ -angulated category.

Remark 2.2.16. Notice that [Theorem 2.2.15](#) is *a priori* quite different from [Theorem 2.2.1](#). Indeed, the former result does not require the existence of an ambient triangulated category in which to embed $\text{proj}(\Lambda)$. However, [Theorem A](#) and [Theorem 4.5.8](#) show that, if \mathbf{k} is perfect, $\text{proj}(\Lambda)$ embeds as a $d\mathbb{Z}$ -cluster tilting subcategory in an essentially unique algebraic triangulated category in such a way that the $(d+2)$ -angulated structure from [Theorem 2.2.15](#) coincides with that from [Theorem 2.2.1](#).

Definition 2.2.17. In the context of [Theorem 2.2.15](#), we call the class \diamond_{η} an *Amiot–Lin (AL) $(d+2)$ -angulation* of the pair $(\text{proj}(\Lambda), \Sigma)$.

The following result shows that the $(d+2)$ -angulated structure from [Theorem 2.2.15](#) is independent of the choice of (truncated) projective resolution of the diagonal bimodule.

Proposition 2.2.18. *Let Λ be a basic finite-dimensional algebra that is twisted $(d+2)$ -periodic with respect to an algebra automorphism σ and let*

$$\Sigma := - \otimes_{\Lambda} \sigma \Lambda_1 : \text{proj}(\Lambda) \xrightarrow{\sim} \text{proj}(\Lambda).$$

Choose exact sequences

$$\eta: \quad 0 \rightarrow {}_1\Lambda_{\sigma} \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

and

$$\eta': \quad 0 \rightarrow {}_1\Lambda_{\sigma} \rightarrow Q_{d+1} \rightarrow \cdots \rightarrow Q_1 \rightarrow Q_0 \rightarrow \Lambda \rightarrow 0$$

of Λ -bimodules with projective-injective middle terms. Then, there exists a morphism of exact sequences

$$\begin{array}{ccccccccccc} \eta: & 0 & \rightarrow & {}_1\Lambda_{\sigma} & \rightarrow & P_{d+1} & \rightarrow & \cdots & \rightarrow & P_1 & \rightarrow & P_0 & \rightarrow & \Lambda & \rightarrow & 0 \\ & & & \downarrow u & & \downarrow & & & & \downarrow & & \downarrow & & \parallel & & \\ \eta': & 0 & \rightarrow & {}_1\Lambda_{\sigma} & \rightarrow & Q_{d+1} & \rightarrow & \cdots & \rightarrow & Q_1 & \rightarrow & Q_0 & \rightarrow & \Lambda & \rightarrow & 0 \end{array}$$

where $u \in Z(\Lambda)^{\times}$ is a unit in the centre of Λ . In particular, the $(d+2)$ -angulations \diamond_{η} and $\diamond_{\eta'}$ of $(\text{proj}(\Lambda), \Sigma)$ are equivalent in the sense of [Definition 2.1.13](#).

Proof. The Comparison Lemma for projective resolutions yields a morphism of complexes

$$\begin{array}{ccccccccccc} \eta: & 0 & \rightarrow & {}_1\Lambda_{\sigma} & \rightarrow & P_{d+1} & \rightarrow & \cdots & \rightarrow & P_1 & \rightarrow & P_0 & \rightarrow & \Lambda & \rightarrow & 0 \\ & & & \downarrow \varphi & & \downarrow & & & & \downarrow & & \downarrow & & \parallel & & \\ \eta': & 0 & \rightarrow & {}_1\Lambda_{\sigma} & \rightarrow & Q_{d+1} & \rightarrow & \cdots & \rightarrow & Q_1 & \rightarrow & Q_0 & \rightarrow & \Lambda & \rightarrow & 0 \end{array}$$

in which the morphism φ is an isomorphism in the stable category of Λ -bimodules. Since the algebra Λ is finite-dimensional, the morphism φ can be replaced by an isomorphism in the category of Λ -bimodules without modifying the rows and the other vertical morphisms in the diagram (except for possibly $P_{d+1} \rightarrow Q_{d+1}$), see [[Che21](#), Cor. 2.3]. Also, it is easily verified that there are isomorphisms of algebras

$$\text{End}_{\Lambda^e}({}_1\Lambda_{\sigma}) = \text{End}_{\Lambda^e}(\Lambda) \cong Z(\Lambda),$$

where the rightmost isomorphism is classical. In particular, the isomorphism φ is given by right multiplication by a unit $u \in Z(\Lambda)^\times$, which we interpret as a natural isomorphism $u: \Sigma \Rightarrow \Sigma$ in the usual way. It readily follows that the pair $(\mathbf{1}_{\text{proj}(\Lambda)}, u)$ yields an equivalence between the $(d+2)$ -angulations $\diamond_{\eta'}$ and \diamond_{η} on the pair $(\text{proj}(\Lambda), \Sigma)$. \square

Remark 2.2.19. We mention that [Proposition 2.2.18](#) holds, more generally, for locally bounded categories (or even more general objects) for which one can lift stable isomorphisms between twists of the diagonal bimodule to actual bimodule isomorphisms. We thank Martin Herschend for informing us that the latter property holds for Krull–Schmidt categories whose quiver of irreducible maps is connected, for such categories have local centre.

The following theorem is a $(d+2)$ -angulated analogue of [\[Han20, Thm. 1.2\]](#), which treats the case $d = 1$.

Theorem 2.2.20. *Let Λ be a basic finite-dimensional algebra such that Λ/J_Λ is separable. The following statements are equivalent:*

- (1) *The algebra Λ is twisted $(d+2)$ -periodic.*
- (2) *The additive category $\text{proj}(\Lambda)$ admits the structure of a $(d+2)$ -angulated category.*

Proof. The proof is a straightforward adaptation of the proof of [\[Han20, Thm. 1.2\]](#). If Λ is twisted $(d+2)$ -periodic, then [Theorem 2.2.15](#) shows that the additive category $\text{proj}(\Lambda)$ admits the structure of a $(d+2)$ -angulated category. The converse is precisely [Proposition 2.2.9](#). \square

2.2.3. Twisted periodic algebras and higher Auslander–Reiten Theory. Although not needed in the sequel, the following discussion provides further motivation for the study of twisted periodic algebras from the perspective of Iyama’s higher Auslander–Reiten theory, see the introduction to [\[Iya07b\]](#) for the analogous discussion in the (d) -abelian case.

Let Λ be a basic Frobenius algebra; for simplicity, assume that Λ is connected and not semisimple. The *Nakayama functor*

$$\nu := - \otimes_\Lambda D\Lambda: \text{proj } \Lambda \xrightarrow{\sim} \text{proj } \Lambda$$

is a Serre functor in the sense that there are natural isomorphisms

$$\text{Hom}_\Lambda(Y, \nu X) \xrightarrow{\sim} D\text{Hom}_\Lambda(X, Y), \quad X, Y \in \text{proj } \Lambda,$$

where $V \mapsto DV$ denotes the passage to the \mathbf{k} -linear dual, see [\[BO01, Def. 1.2\]](#).

Suppose now that Λ is twisted $(d+2)$ -periodic with respect to an algebra automorphism σ and let

$$\Sigma := - \otimes_\Lambda \sigma \Lambda_1: \text{proj}(\Lambda) \xrightarrow{\sim} \text{proj}(\Lambda).$$

Fix an arbitrary Amiot–Lin $(d+2)$ -angulation on the pair $(\text{proj } \Lambda, \Sigma)$. Following [\[IY08\]](#), we set

$$\nu_d := \Sigma^{-d} \nu: \text{proj } \Lambda \xrightarrow{\sim} \text{proj } \Lambda.$$

Let P^0 be an indecomposable projective Λ -module. Extend the map $f: P^0 \rightarrow \nu P^0$ that maps the simple top of P^0 to the simple socle of νP^0 to a $(d+2)$ -angle

$$\nu_d P^0 \rightarrow P^{-d} \rightarrow \cdots \rightarrow P^{-1} \rightarrow P^0 \xrightarrow{f} \nu P^0,$$

which we may assume is minimal in the sense that all the morphisms lie in the radical of $\text{proj } \Lambda$. Since $(d+2)$ -angles are exact [\[GKO13, Prop. 2.5\(a\)\]](#), the above $(d+2)$ -angle is the first part of a minimal projective resolution of the simple top S of P^0 and, simultaneously, of a minimal injective co-resolution of the simple socle

$\Omega_{\Lambda}^{d+2}(S) \cong \sigma^*(S)$ of $\nu_d P^0$. Following the paradigm of classical Auslander–Reiten theory [AR75, AR78], the above $(d+2)$ -angle can be regarded as the prototypical example of an *almost-split $(d+2)$ -angle* in the sense of [IY08, 3.8], see also [Fed19, Def. 5.1]. For the equivalence between the existence of a Serre functor and the existence of almost-split $(d+2)$ -angles see [RVdB02, Thm. I.2.4] for the case $d=1$ and [Zho22, Thm. 1.5] for the general case (see also [Ami07, Prop. 1.2.] for a related statement).

3. ENHANCED $(d+2)$ -ANGULATED CATEGORIES

3.1. Pre-triangulated differential graded categories. We begin by recalling the notion of a pre-triangulated differential graded category in the sense of [BK90]. We need a few preliminaries.

3.1.1. Reminder on differential graded categories. We recall basic aspects of the theory of differential graded categories that are needed in the sequel; we refer the reader to [Kel06] and the references therein for details.

A *differential graded (=DG) category* is a category enriched in the symmetric monoidal category $\mathbf{C}(\mathbf{k}) = \mathbf{C}(\text{Mod}(\mathbf{k}))$ of cochain complexes of \mathbf{k} -modules (endowed with the usual tensor product of cochain complexes). Thus, the morphisms between two objects in a DG category form a cochain complex and the composition law satisfies the graded Leibniz rule. A *DG functor* is simply an enriched functor between DG categories. A *graded category* is a DG category whose cochain complexes of morphisms have vanishing differential, and a *graded functor* is a DG functor between graded categories. For example, given a (\mathbf{k} -linear) additive category \mathcal{C} , there is a DG category $\mathbf{C}_{\text{dg}}(\mathcal{C})$ whose objects are the cochain complexes in \mathcal{C} and for $X, Y \in \mathbf{C}_{\text{dg}}(\mathcal{C})$ we let $\text{hom}_{\mathcal{C}}^i(X, Y)$ be the complex whose component of degree i is the \mathbf{k} -module

$$\text{hom}_{\mathcal{C}}^i(X, Y) := \prod_{k \in \mathbb{Z}} \mathcal{C}(X^k, Y^{k+i})$$

of degree i morphisms of graded objects $X \rightarrow Y$ and differential

$$\partial: f \mapsto d_Y f - (-1)^i f d_X, \quad f \in \text{hom}_{\mathcal{C}}^i(X, Y).$$

Note also that the usual shift of cochain complexes

$$(X, d_X) \mapsto X[1] := (X(1), -d_X),$$

where $X \mapsto X(1)$ denotes the shift of graded objects, can be promoted to a DG functor

$$[1]: \mathbf{C}_{\text{dg}}(\mathcal{C}) \longrightarrow \mathbf{C}_{\text{dg}}(\mathcal{C}), \quad X \mapsto X[1]$$

whose action on morphisms is given by

$$[1]: f \mapsto (-1)^i f, \quad f \in \text{hom}_{\mathcal{C}}^i(X, Y).$$

A DG category \mathcal{A} has two associated categories whose objects are the same as those of \mathcal{A} : the *underlying category* $Z^0(\mathcal{A})$ and the *0-th cohomology category* $H^0(\mathcal{A})$. As the notation suggests, the morphisms in $Z^0(\mathcal{A})$ are the \mathbf{k} -modules of cocycles

$$Z^0(\mathcal{A})(x, y) := Z^0(\mathcal{A}(x, y)), \quad x, y \in Z^0(\mathcal{A}),$$

while the morphisms in $H^0(\mathcal{A})$ are the cohomology \mathbf{k} -modules

$$H^0(\mathcal{A})(x, y) := H^0(\mathcal{A}(x, y)), \quad x, y \in H^0(\mathcal{A}).$$

Similarly, \mathcal{A} has an associated graded category $H^\bullet(\mathcal{A})$ whose objects are the same as those of \mathcal{A} and with graded \mathbf{k} -modules of morphisms

$$H^\bullet(\mathcal{A})(x, y) := H^\bullet(\mathcal{A}(x, y)), \quad x, y \in H^\bullet(\mathcal{A}).$$

A DG functor $F: \mathcal{A} \rightarrow \mathcal{B}$ induces apparent functors

$$Z^0(F): Z^0(\mathcal{A}) \longrightarrow Z^0(\mathcal{B}) \quad \text{and} \quad H^0(F): H^0(\mathcal{A}) \longrightarrow H^0(\mathcal{B})$$

as well as a graded functor

$$H^\bullet(F): H^\bullet(\mathcal{A}) \longrightarrow H^\bullet(\mathcal{B}).$$

A DG functor F is a *quasi-equivalence* if the induced functor $H^\bullet(F)$ is an equivalence of graded categories.

For example, if \mathcal{C} is an additive category then

$$Z^0(\mathbf{C}_{\text{dg}}(\mathcal{C})) = \mathbf{C}(\mathcal{C})$$

while

$$H^0(\mathbf{C}_{\text{dg}}(\mathcal{C})) = \mathbf{K}(\mathcal{C})$$

is the homotopy category of cochain complexes in \mathcal{C} . Similarly, the graded category $H^\bullet(\mathbf{C}_{\text{dg}}(\mathcal{C}))$ has graded \mathbf{k} -modules of morphisms

$$H^i(\text{hom}_{\mathcal{C}}^\bullet(X, Y)) = \mathbf{K}(\mathcal{C})(X, Y[i]), \quad X, Y \in \mathbf{C}(\mathcal{C}), \quad i \in \mathbb{Z}.$$

Under these identifications, the shift (DG) functor on $\mathbf{C}_{\text{dg}}(\mathcal{C})$ induces the usual shift functors on $\mathbf{C}(\mathcal{C})$ and $\mathbf{K}(\mathcal{C})$.

3.1.2. The homotopy category of small DG categories. As with quasi-isomorphisms between cochain complexes, quasi-equivalences between DG categories need not admit a quasi-inverse given by a DG functor. However, the category $\mathbf{dgc}\mathbf{at}$ of small DG categories admits a cofibrantly generated Quillen model category structure, called the *Tabuada model structure*, whose weak equivalences are the quasi-equivalences (we do not need to recall here what are the classes of (co)fibrations) [Tab05]. We denote the homotopy category of the Tabuada model structure by $\mathbf{Ho}(\mathbf{dgc}\mathbf{at})$, which is, by the general theory of Quillen model categories, equivalent to the localisation of $\mathbf{dgc}\mathbf{at}$ at the class of quasi-equivalences. In particular, two small DG categories \mathcal{A} and \mathcal{B} are isomorphic in $\mathbf{Ho}(\mathbf{dgc}\mathbf{at})$ if and only if they are connected by a finite zig-zag of quasi-equivalences. Furthermore, the set of morphisms $\mathcal{A} \rightarrow \mathcal{B}$ in $\mathbf{Ho}(\mathbf{dgc}\mathbf{at})$ is in bijection with the set of quasi-isomorphism classes of DG \mathcal{A} - \mathcal{B} -bimodules X such that, for each $a \in \mathcal{A}$, the DG \mathcal{B} -module $X(-, a)$ is quasi-isomorphic to a representable DG \mathcal{B} -module, see [Toë11, Cor. 4.8].

3.1.3. The derived category of a DG category. We recall the construction of the derived category of a (small) DG category. We refer the reader [Kel94] for details. Let \mathcal{A} be a small DG category, that is the objects in \mathcal{A} form a set. The *opposite DG category* \mathcal{A}^{op} of \mathcal{A} is the DG category with the same objects as \mathcal{A} and graded morphism spaces

$$\mathcal{A}^{\text{op}}(x, y) := \mathcal{A}(y, x), \quad x, y \in \mathcal{A}^{\text{op}};$$

the composition law in \mathcal{A}^{op} is given in terms of the composition law in \mathcal{A} by the formula

$$g \circ^{\text{op}} f := (-1)^{ij} f \circ g$$

whenever f and g are homogeneous of degree i and j , respectively. The *DG category of (right) DG \mathcal{A} -modules* is the DG category

$$\mathbf{dgMod}_{\text{dg}}(\mathcal{A}) := \mathbf{Fun}_{\text{dg}}(\mathcal{A}^{\text{op}}, \mathbf{C}_{\text{dg}}(\mathbf{k}))$$

of DG functors $\mathcal{A}^{\text{op}} \rightarrow \mathbf{C}_{\text{dg}}(\mathbf{k}) = \mathbf{C}_{\text{dg}}(\mathbf{Mod} \mathbf{k})$. We denote the morphism complexes in $\mathbf{dgMod}_{\text{dg}}(\mathcal{A})$ by

$$\text{hom}_{\mathcal{A}}(M, N), \quad M, N \in \mathbf{dgMod}_{\text{dg}}(\mathcal{A}).$$

The DG Yoneda embedding

$$\mathbf{h}: \mathcal{A} \longrightarrow \mathbf{dgMod}_{\text{dg}}(\mathcal{A}), \quad x \longmapsto \mathcal{A}(-, x).$$

identifies \mathcal{A} with full DG subcategory of $\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A})$.

The underlying category

$$\mathbf{dgMod}(\mathcal{A}) := Z^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$$

is a Grothendieck abelian category. Moreover, $\mathbf{dgMod}(\mathcal{A})$ is a Frobenius exact category in which a short exact sequence

$$0 \rightarrow L \xrightarrow{i} M \rightarrow N \rightarrow 0$$

of DG \mathcal{A} -modules is admissible (=a conflation) if there exists a morphism $p: M \rightarrow L$ of graded (!) \mathcal{A} -modules such that $pi = \mathbf{1}_L$; a DG \mathcal{A} -module is projective-injective in this exact structure if and only if it is *contractible*, that is if it is a zero object in the homotopy category $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$. In particular, the stable category of the Frobenius exact category $\mathbf{dgMod}(\mathcal{A})$ equals $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$ and is a triangulated category with suspension functor induced by the shift functor on $\mathbf{C}_{\mathbf{dg}}(\mathbf{k})$ by means of the formula

$$[1]: M \longmapsto M[1] := [1] \circ M, \quad M \in \mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}),$$

keeping in mind that a DG \mathcal{A} -module is a DG functor $\mathcal{A}^{\text{op}} \rightarrow \mathbf{C}_{\mathbf{dg}}(\mathbf{k})$. By construction, there are canonical isomorphisms

$$H^i(\mathbf{hom}_{\mathcal{A}}(M, N)) \cong H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))(M, N[i]), \quad M, N \in \mathbf{dgMod}(\mathcal{A})$$

and, as a consequence of the DG Yoneda Lemma,

$$H^i(\mathbf{hom}_{\mathcal{A}}(\mathbf{h}_x, M)) \cong H^i(M_x), \quad x \in \mathcal{A}, \quad M \in \mathbf{dgMod}(\mathcal{A}).$$

A DG \mathcal{A} -module N is *acyclic* if for each $x \in \mathcal{A}$ the cochain complex of \mathbf{k} -modules N_x is acyclic. The *derived category* $\mathbf{D}(\mathcal{A})$ of \mathcal{A} is the full subcategory of the triangulated category $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$ spanned by the DG \mathcal{A} -modules that are *DG projective*, that is the DG \mathcal{A} -modules P such that every epimorphism $M \rightarrow P$ with acyclic kernel is split. The derived category of \mathcal{A} is a triangulated subcategory of $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$ that is closed under small coproducts and retracts and is compactly generated by the *free DG \mathcal{A} -modules*

$$\mathbf{h}_x = \mathcal{A}(-, x): y \longmapsto \mathcal{A}(y, x), \quad x \in \mathcal{A}.$$

The *perfect derived category* $\mathbf{D}^c(\mathcal{A})$ of \mathcal{A} is the full subcategory of $\mathbf{D}(\mathcal{A})$ spanned by the compact objects; equivalently, $\mathbf{D}^c(\mathcal{A})$ is the thick subcategory of $\mathbf{D}(\mathcal{A})$ generated by the free DG \mathcal{A} -modules [Kel94, Thm. 5.3]. In particular, $\mathbf{D}^c(\mathcal{A})$ is a triangulated category in which idempotents split.

Let $F: \mathcal{A} \rightarrow \mathcal{B}$ be a DG functor. The restriction of scalars

$$F^*: \mathbf{dgMod}_{\mathbf{dg}}(\mathcal{B}) \longrightarrow \mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A})$$

induces an exact functor

$$F^*: \mathbf{D}(\mathcal{B}) \longrightarrow \mathbf{D}(\mathcal{A})$$

that admits an exact left adjoint

$$\mathbb{L}F_!: \mathbf{D}(\mathcal{A}) \longrightarrow \mathbf{D}(\mathcal{B})$$

that preserves small co-products (constructed explicitly as a left derived functor). Moreover, the functor $\mathbb{L}F_!$ preserves compact objects (since it preserves the free DG modules) and hence restricts to an exact functor

$$\mathbb{L}F_!: \mathbf{D}^c(\mathcal{A}) \longrightarrow \mathbf{D}^c(\mathcal{B})$$

between the corresponding perfect derived categories.

3.1.4. *Pre-triangulated DG categories.* The following definitions motivate our definition of DG enhancement of a $(d + 2)$ -angulated category (see [Definitions 3.2.1](#) and [3.2.5](#)).

Definition 3.1.1 ([\[BK90\]](#)). A small DG category \mathcal{A} is *pre-triangulated* if the fully faithful functor

$$H^0(\mathbf{h}): H^0(\mathcal{A}) \longrightarrow \mathbf{D}^c(\mathcal{A})$$

identifies $H^0(\mathcal{A})$ with a triangulated subcategory of $\mathbf{D}^c(\mathcal{A})$. Similarly, \mathcal{A} is *Karoubian pre-triangulated* if the above functor is an equivalence of categories.

Remark 3.1.2. The term *pre-triangulated* is also used for triangulated categories which need not satisfy the octahedral axiom [[Nee01](#), Def. 1.1.2], see also [Definition 2.1.1](#) and [Remark 2.1.5](#). Despite the clash of terminology, it is always clear from the context which notion is in use. In particular, notice that the 0-th cohomology category of a pre-triangulated DG category is a triangulated category (and therefore satisfies the octahedral axiom).

Definition 3.1.3 ([\[BK90\]](#)). Let \mathcal{T} be a triangulated category.

- (1) A *(DG) enhancement* of \mathcal{T} is a pre-triangulated DG category \mathcal{A} such that there exists an equivalence of triangulated categories

$$\mathcal{T} \simeq H^0(\mathcal{A})$$

(notice that we only require the existence of such an equivalence, and not the datum of a preferred such).

- (2) We say that \mathcal{T} *admits a unique (DG) enhancement* if it admits an enhancement and any two enhancements of \mathcal{T} are quasi-equivalent (that is, isomorphic in $\mathbf{Ho}(\mathbf{dgcats})$).

3.1.5. *Morita equivalences.* The category of small DG categories and DG functors between them admits a cofibrantly generated Quillen model category structure, called the Morita model structure [[Tab05](#), Thm. 5.3], see also [[Tab06](#), [Tab07a](#)], whose weak equivalences are the *Morita equivalences*, that is the DG functors $F: \mathcal{A} \rightarrow \mathcal{B}$ such that the induced exact functor

$$\mathbb{L}F_1: \mathbf{D}^c(\mathcal{A}) \longrightarrow \mathbf{D}^c(\mathcal{B})$$

is an equivalence of triangulated categories. A DG category is fibrant in the Morita model structure if and only if it is Karoubian pre-triangulated; more generally, the DG Yoneda embedding

$$\mathcal{A} \longrightarrow \mathbf{D}_{\mathbf{dg}}^c(\mathcal{A})$$

is a fibrant replacement (that is a Morita equivalence with fibrant target) for every small DG category \mathcal{A} , where $\mathbf{D}_{\mathbf{dg}}^c(\mathcal{A})$ is the *perfect derived DG category* of \mathcal{A} , that is the full DG subcategory of $\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A})$ spanned by the DG \mathcal{A} -modules that are DG projective and compact in $\mathbf{D}(\mathcal{A})$. The *Morita category of small DG categories* is the homotopy category \mathbf{Hmo} associated to the Morita model structure. Two DG categories are *Morita equivalent* if they are isomorphic in \mathbf{Hmo} . In particular, every small DG category is Morita equivalent to its perfect derived DG category.

Remark 3.1.4. If \mathcal{A} and \mathcal{B} are small DG categories that are Morita equivalent, then the perfect derived categories $\mathbf{D}^c(\mathcal{A})$ and $\mathbf{D}^c(\mathcal{B})$ are equivalent as triangulated categories. The converse, however, is false [[DS07](#), [Sch02](#), [Kaj13](#), [RVdB19](#)].

Remark 3.1.5. Every quasi-equivalence between small DG categories is also a Morita equivalence. Closely related to this is the fact that \mathbf{Hmo} is equivalent to the full subcategory of $\mathbf{Ho}(\mathbf{dgcats})$ spanned by the Karoubian pre-triangulated DG categories. In particular, two Karoubian pre-triangulated DG categories are quasi-equivalent if and only if they are Morita equivalent.

Theorem 3.1.6 ([Kel06, Thm. 3.8]). *Let \mathcal{X} be a Karoubian pre-triangulated DG category and $\mathcal{A} \subseteq \mathcal{X}$ a full DG subcategory such that $\text{thick}(H^0(\mathcal{A})) = H^0(\mathcal{X})$. Then, the inclusion functor $\mathcal{A} \hookrightarrow \mathcal{X}$ is a Morita equivalence and, in particular, there are canonical equivalences of triangulated categories*

$$\mathrm{D}^c(\mathcal{A}) \xrightarrow{\sim} \mathrm{D}^c(\mathcal{X}) \xleftarrow{\sim} H^0(\mathcal{X}).$$

Example 3.1.7. Given a DG algebra A , there are canonical morphisms of DG algebras (=DG functors between DG categories with a single object)

$$H^0(A) \longleftarrow \tau^{\leq 0} A \longrightarrow A,$$

where

$$(\tau^{\leq 0} A)^i := \begin{cases} A^i & i < 0, \\ Z^0(A) & i = 0, \\ 0 & i > 0. \end{cases}$$

By construction, the map $\tau^{\leq 0} A \rightarrow A$ is a quasi-isomorphism if and only if the cohomology of A is concentrated in non-positive degrees. In particular, if $H^\bullet(A) = H^0(A)$ is concentrated in degree 0, then both of the above maps are quasi-isomorphisms of DG algebras and thus also Morita equivalences. Combining this observation with [Theorem 3.1.6](#), we conclude that the (compact) derived category of an ordinary algebra admits a unique enhancement in the sense of [Definition 3.1.3](#).

Remark 3.1.8. In contrast to the situation described in [Example 3.1.7](#) for the derived category of an ordinary algebra, establishing the uniqueness of enhancements for more general triangulated categories is a subtle endeavour. Moreover, the majority of results of this kind that are available rely on the existence of t -structures in a crucial way, see for example [LO10, CS18, CNS22]. Similar to the situation in [Mur22], in this article we consider triangulated categories that need not admit any non-trivial t -structures (such as 2-Calabi–Yau triangulated categories with a 2-cluster tilting object, see [Theorem 6.1.5](#) and compare with [ZZ, Thm. 4.1]).

3.2. Pre- $(d+2)$ -angulated differential graded categories. The following are straightforward extensions of [Definitions 3.1.1](#) and [3.1.3](#) to the context of $(d+2)$ -angulated categories.

Definition 3.2.1. A small DG category \mathcal{A} is *pre- $(d+2)$ -angulated* if the fully faithful functor

$$H^0(\mathbf{h}): H^0(\mathcal{A}) \longrightarrow \mathrm{D}^c(\mathcal{A})$$

identifies $H^0(\mathcal{A})$ with a standard $(d+2)$ -angulated subcategory of $\mathrm{D}^c(\mathcal{A})$ (see [Definition 2.2.2](#)). Similarly, \mathcal{A} is *Karoubian pre- $(d+2)$ -angulated* if it is pre- $(d+2)$ -angulated and idempotents split in $H^0(\mathcal{A})$.

Remark 3.2.2. A (Karoubian) pre-3-angulated DG category is precisely a (Karoubian) pre-triangulated DG category.

Remark 3.2.3. Let \mathcal{A} be a pre- $(d+2)$ -angulated DG category. Since, by definition, $H^0(\mathcal{A})$ is equivalent to a d -rigid subcategory of $\mathrm{D}^c(\mathcal{A})$ closed under the action of the d -fold suspension and its inverse, the standard isomorphisms

$$\mathrm{Hom}_{\mathrm{D}(\mathcal{A})}(\mathbf{h}_x, \mathbf{h}_y[i]) \cong H^i(\mathcal{A}(x, y)), \quad x, y \in \mathcal{A},$$

imply that

$$\forall i \notin d\mathbb{Z}, \quad H^i(\mathcal{A}(x, y)) = 0.$$

This is a strong restriction on the cohomology of \mathcal{A} (see also [Definition 4.3.2](#)).

Proposition 3.2.4. *Let \mathcal{A} be a small DG category such that $H^0(\mathcal{A})$ is an additive category with finite-dimensional morphism spaces and split idempotents. The following statements are equivalent:*

- (1) The DG category \mathcal{A} is Karoubian pre- $(d+2)$ -angulated.
- (2) The fully faithful functor

$$H^0(\mathbf{h}): H^0(\mathcal{A}) \longrightarrow D^c(\mathcal{A})$$

identifies $H^0(\mathcal{A})$ with a $d\mathbb{Z}$ -cluster tilting subcategory of $D^c(\mathcal{A})$.

Proof. Notice that, since $D^c(\mathcal{A})$ is generated as a triangulated category with split idempotents by the free DG \mathcal{A} -modules, the finiteness assumption on $H^0(\mathcal{A})$ implies that $D^c(\mathcal{A})$ has finite-dimensional morphism spaces. The equivalence between the two conditions then follows from [Theorem 2.2.4](#). \square

Definition 3.2.5. Let \mathcal{F} be a $(d+2)$ -angulated category.

- (1) A (DG) enhancement of \mathcal{F} is a pre- $(d+2)$ -angulated DG category \mathcal{A} such that there exists an equivalence of $(d+2)$ -angulated categories

$$\mathcal{F} \simeq H^0(\mathcal{A}),$$

where the $(d+2)$ -angulation on the right-hand side is induced by the triangulation on $D^c(\mathcal{A})$ as in [Theorem 2.2.1](#). If \mathcal{F} has an enhancement we say that it is *algebraic*.

- (2) We say that \mathcal{F} admits a unique (DG) enhancement if it admits an enhancement and any two enhancements of \mathcal{F} are quasi-equivalent (that is isomorphic in $\mathbf{Ho}(\mathbf{dgcatt})$).

The following two questions are natural.

Question 3.2.6. Let \mathcal{F} be a $(d+2)$ -angulated category. Does \mathcal{F} admit an enhancement? If it does, under which conditions is the enhancement unique?

Definition 3.2.7. Let (\mathcal{C}, Σ) be a pair consisting of an additive category \mathcal{C} and an autoequivalence $\Sigma: \mathcal{C} \xrightarrow{\sim} \mathcal{C}$. We say that a pre- $(d+2)$ -angulated DG category \mathcal{A} induces a (DG) enhanced $(d+2)$ -angulated structure on (\mathcal{C}, Σ) if there exists an equivalence $F: \mathcal{C} \xrightarrow{\sim} H^0(\mathcal{A})$ such that the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & H^0(\mathcal{A}) \\ \Sigma \downarrow & & \downarrow [d] \\ \mathcal{C} & \xrightarrow{F} & H^0(\mathcal{A}) \end{array}$$

commutes up to natural isomorphism. In this case, we say that the pair (\mathcal{C}, Σ) admits a (DG) enhanced $(d+2)$ -angulated structure.

Question 3.2.8. Let (\mathcal{C}, Σ) be a pair consisting of an additive category \mathcal{C} and an automorphism $\Sigma: \mathcal{C} \xrightarrow{\sim} \mathcal{C}$. Does the pair admit an enhanced $(d+2)$ -angulated structure? If it does, is the enhancement of the underlying $(d+2)$ -angulated category unique?

Remark 3.2.9. [Theorems 4.5.8](#) and [5.1.3](#) provide a complete answer to [Questions 3.2.6](#) and [3.2.8](#) in the setting of [Theorems 2.2.15](#) and [2.2.20](#).

Remark 3.2.10. Let $(\mathcal{F}, \Sigma, \diamond)$ be a $(d+2)$ -angulated category and $n \geq 1$ an integer. Recall from [Example 2.1.9](#) that the pair (\mathcal{F}, Σ^n) is endowed with an $n(d+2)$ -angulation \diamond^n obtained by extending the sequences in \diamond periodically up to iterated applications of Σ . In general, we do not know if the existence of an enhancement of $(\mathcal{F}, \Sigma, \diamond)$ implies the existence of an enhancement of $(\mathcal{F}, \Sigma^n, \diamond^n)$. Notwithstanding, as explained in the introduction, [Theorem 5.1.3](#) provides a positive answer when $\mathcal{F} = \mathbf{proj}(\Lambda)$ for some twisted $(d+2)$ -periodic algebra with respect to an algebra

automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$ and $\diamond = \diamond_\eta$ is the AL $(d+2)$ -angulation associated to an exact sequence of Λ -bimodules

$$\eta: \quad 0 \rightarrow {}_1\Lambda_\sigma \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

with projective-injective middle terms. Indeed, the algebra Λ is twisted $n(d+2)$ -periodic with respect to σ^n and the class \diamond^n is precisely the AL $n(d+2)$ -angulation associated to the exact sequence obtained by splicing together the exact sequences

$$\eta, \Sigma(\eta), \dots, \Sigma^{n-1}(\eta).$$

Therefore, in this case, the $n(d+2)$ -angulated category $(\mathcal{F}, \Sigma^n, \diamond^n)$ admits an enhancement by [Theorem 5.1.3](#). However, the precise relationship between enhancements of $(\mathcal{F}, \Sigma, \diamond)$ and $(\mathcal{F}, \Sigma^n, \diamond^n)$ is not clear (even if the semisimple algebra Λ/J_Λ is separable, so that we can guarantee that enhancements are unique). For example, can one construct—explicitly—an enhancement of $(\mathcal{F}, \Sigma^n, \diamond^n)$ from a given enhancement of $(\mathcal{F}, \Sigma, \diamond)$?

4. HIGHER STRUCTURES AND $(d+2)$ -ANGULATIONS

4.1. Toda brackets and standard $(d+2)$ -angulations.

Setting 4.1.1. We fix a triangulated category \mathcal{T} with shift functor $[1]$ in this subsection.

We aim to characterise standard $(d+2)$ -angles ([Definition 2.2.2](#)) in terms of higher Toda brackets. We begin by recalling the necessary definitions.

Definition 4.1.2. Suppose given a sequence of morphisms in \mathcal{T} of the form

$$x_3 \xrightarrow{f_3} x_2 \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0.$$

The *Toda family* $\Upsilon(f_1, f_2, f_3)$ is the set of pairs (β, α) such that there exists a commutative diagram of the form

$$(4.1.3) \quad \begin{array}{ccccccc} x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{f_1} & x_0 \\ \alpha \downarrow & & \parallel & & \parallel & & \uparrow \beta[1] \\ c & \xrightarrow{u} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{v} & c[1] \end{array}$$

in which the horizontal row is an exact triangle in \mathcal{T} .

Remark 4.1.4. In the setting of [Definition 4.1.2](#), notice that the Toda family $\Upsilon(f_1, f_2, f_3)$ is non-empty if and only if $f_1 f_2 = 0$ and $f_2 f_3 = 0$.

Remark 4.1.5. An alternative (perhaps more suggestive) way to depict the condition for the pair (β, α) to lie in $\Upsilon(f_1, f_2, f_3)$ is as a commutative diagram

$$\begin{array}{ccccc} & & x_2 & \xrightarrow{f_2} & x_1 \\ & f_3 \nearrow & & \nwarrow & \nearrow f_1 \\ x_3 & \xrightarrow{\alpha} & c & \xrightarrow{\beta} & x_0 \\ & & & \searrow -1 & \\ & & & & \end{array}$$

in which β is a morphism of degree -1 . Here, the oriented triangle is an exact triangle in \mathcal{T} and the un-oriented triangles commute. In particular, the pair (β, α) yields the well-defined composite

$$x_3[1] \xrightarrow{\alpha[1]} c[1] \xrightarrow{\beta[1]} x_0.$$

Definition 4.1.6. Suppose given a sequence of morphisms in \mathcal{T} of the form

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0.$$

The *Toda bracket* $\langle f_1, \dots, f_{d+2} \rangle$ is the subset of $\mathcal{T}(x_{d+2}[d], x_0)$ defined inductively as follows:

- (1) If $d = 0$, then $\langle f_1, f_2 \rangle := \{f_1 \circ f_2\}$.
- (2) If $d \geq 1$, then

$$\langle f_1, \dots, f_{d+2} \rangle := \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} \langle \beta[1], \alpha[1], f_4[1], \dots, f_{d+2}[1] \rangle$$

is the union of Toda brackets of length $d + 1$.

Remark 4.1.7. **Definition 4.1.6** is equivalent to that in [Shi02], which is itself based on that in [Coh68], see [CF17, Prop. 5.8]. We also note that **Definition 4.1.2** is equivalent to the desuspension of that in [CF17, Sec. 5].

Remark 4.1.8. The Toda bracket is not quite a $(d + 2)$ -ary operation as its output is a *set* of morphisms rather than a single map (compare with **Proposition 4.1.31**).

Example 4.1.9. The Toda bracket of a triple of composable morphisms

$$x_3 \xrightarrow{f_3} x_2 \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0.$$

is the subset of $\mathcal{T}(x_3[1], x_0)$ given by

$$\langle f_1, f_2, f_3 \rangle = \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} \langle \beta[1], \alpha[1] \rangle = \{(\beta \circ \alpha)[1] \mid (\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)\}.$$

Example 4.1.10. The Toda bracket of the morphisms in an exact triangle

$$x_3 \xrightarrow{f_3} x_2 \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_3[1]$$

satisfies $\mathbf{1}_{x_3[1]} \in \langle f_1, f_2, f_3 \rangle$. Indeed, in view of **Example 4.1.9**, it is enough to observe that the tautological diagram

$$\begin{array}{ccccccc} x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{f_1} & x_3[1] \\ \mathbf{1}_{x_3} \downarrow & & \parallel & & \parallel & & \uparrow \mathbf{1}_{x_3[1]} \\ x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{f_1} & x_3[1] \end{array}$$

exhibits the membership $(\mathbf{1}_{x_3}, \mathbf{1}_{x_3}) \in \mathbb{T}(f_1, f_2, f_3)$.

The following classical result, which is a refinement of **Example 4.1.10**, illustrates the usefulness of Toda brackets.

Theorem 4.1.11 ([Hel68, Thm. 13.2], see also [CF17, Thm. B.1]). *Suppose given a sequence of morphisms in \mathcal{T} of the form*

$$x_3 \xrightarrow{f_3} x_2 \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_3[1].$$

The following statements are equivalent:

- (1) *The above sequence is an exact triangle in \mathcal{T} .*
- (2) *The following two conditions are satisfied:*
 - (a) *For each $y \in \mathcal{T}$, The induced sequence of \mathbf{k} -modules*

$$\mathcal{T}(y, x_3) \rightarrow \mathcal{T}(y, x_2) \rightarrow \mathcal{T}(y, x_1) \rightarrow \mathcal{T}(y, x_3[1]) \rightarrow \mathcal{T}(y, x_2[1])$$

is exact.

- (b) *We have $\mathbf{1}_{x_3[1]} \in \langle f_1, f_2, f_3 \rangle$.*

The objective of this section is to prove the following $(d + 2)$ -angulated analogue of **Theorem 4.1.11** (compare also with **Theorem 2.2.1**). As is customary, given a subset X of an abelian group, we let

$$-X := \{-x \mid x \in X\}.$$

Theorem 4.1.12. *Let $\mathcal{C} \subseteq \mathcal{T}$ be a d -rigid subcategory such that $\mathcal{C} = \text{add}(c)$ for some basic object c , $\mathcal{C}[d] = \mathcal{C}$ and $\text{thick}(\mathcal{C}) = \mathcal{T}$. Suppose given a sequence of morphisms in \mathcal{C} of the form*

$$c_{d+2} \xrightarrow{f_{d+2}} c_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_3} c_2 \xrightarrow{f_2} c_1 \xrightarrow{f_1} c_{d+2}[d],$$

with $d \geq 1$. The following statements are equivalent:

(1) *The above sequence fits as the spine of a commutative diagram of the form*

$$\begin{array}{ccccccc} & & c_{d+1} & & \cdots & & c_3 & \xrightarrow{\quad} & c_2 & & \\ & \nearrow & & \searrow & & \nearrow & & & \searrow & & \\ c_{d+2} & \xleftarrow{+1} & & \xrightarrow{+1} & x_{d,5} & \cdots & x_{3,5} & \xleftarrow{+1} & & \xrightarrow{+1} & x_{2,5} & \xleftarrow{+1} & & c_1 \end{array}$$

in which the oriented triangles are exact triangles in \mathcal{T} and such that the connecting morphism $c_0 \rightarrow c_{d+2}[d]$ is given by the obvious (shifted) composite along the bottom row of the diagram.

(2) *The following two conditions are satisfied:*

(a) *For each $y \in \mathcal{C}$, the induced sequence of vector spaces*

$$\mathcal{T}(y, c_{d+2}) \rightarrow \cdots \rightarrow \mathcal{T}(y, c_2) \rightarrow \mathcal{T}(y, c_1) \rightarrow \mathcal{T}(y, c_{d+2}[d]) \rightarrow \mathcal{T}(y, c_{d+1}[d])$$

is exact.

(b) *We have*

$$\mathbf{1}_{c_{d+2}[d]} \in (-1)^{1+\sum_{i=1}^{d+1} i} \langle f_1, f_2, \dots, f_{d+2} \rangle.$$

Computing higher Toda brackets can be rather intricate. We establish a few computation rules that are needed for the proof of [Theorem 4.1.12](#) as well as elsewhere in the article.

Lemma 4.1.13. *Suppose given a sequence of morphisms in \mathcal{T} of the form*

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0$$

and let $\varphi: x_1 \rightarrow x'_1$ be an isomorphism. Then, there is an equality of Toda brackets

$$\langle f_1, f_2, \dots, f_{d+2} \rangle = \langle f_1 \varphi^{-1}, \varphi f_2, \dots, f_{d+2} \rangle.$$

Proof. The claim is obvious for $d = 0$ for we have

$$\langle f_1, f_2 \rangle = \{f_1 \circ f_2\} = \{(f_1 \varphi^{-1}) \circ (\varphi \circ f_2)\} = \langle f_1 \varphi^{-1}, \varphi f_2 \rangle.$$

To prove the claim for $d \geq 1$ it is enough to show that

$$\mathbb{T}(f_1, f_2, f_3) = \mathbb{T}(f_1 \varphi^{-1}, \varphi f_2, f_3).$$

Moreover, it is enough to prove that

$$\mathbb{T}(f_1, f_2, f_3) \subseteq \mathbb{T}(f_1 \varphi^{-1}, \varphi f_2, f_3)$$

for the reverse inclusion is obtained by replacing f_1 and f_2 by $f_1 \varphi^{-1}$ and φf_2 , respectively, and φ by its inverse. Indeed, given a pair $(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)$ there is a commutative diagram in \mathcal{T} of the form

$$\begin{array}{ccccccc} & & x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{f_1} & x_0 \\ & \swarrow \alpha & \parallel & & \parallel & & \parallel & & \parallel \\ c & \xrightarrow{u} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{v} & c[1] & & \beta[1] \\ & \parallel & \parallel & & \parallel & & \parallel & & \parallel \\ & \swarrow \alpha & x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{\varphi f_2} & x'_1 & \xrightarrow{f_1 \varphi^{-1}} & x_0 \\ c & \xrightarrow{u} & x_2 & \xrightarrow{\varphi f_2} & x'_1 & \xrightarrow{v \varphi^{-1}} & c[1] & & \beta[1] \end{array}$$

in which the rows of the front face are exact triangles in \mathcal{T} (the top layer exists by assumption). The bottom face of the diagram exhibits the membership $(\beta, \alpha) \in \mathbb{T}(f_1\varphi^{-1}, \varphi f_2, f_3)$, which is what we needed to prove. \square

Notation 4.1.14. Suppose given a sequence of morphisms in \mathcal{T} of the form

$$x_3 \xrightarrow{f_3} x_2 \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0$$

and let S be a set of exact triangles of the form

$$c \rightarrow x_2 \xrightarrow{f_2} x_1 \rightarrow c[1].$$

We let $\mathbb{T}(f_1, f_2, f_3)_S \subseteq \mathbb{T}(f_1, f_2, f_3)$ be the subset of all pairs (β, α) that are part of a commutative diagram of the form (4.1.3) in which the bottom horizontal row belongs to the set S .

Lemma 4.1.15. *Let $d \geq 1$ and suppose given a sequence of morphisms in \mathcal{T} of the form*

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \dots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0.$$

Let S be a non-empty class of triangles with middle morphism f_2 as in Notation 4.1.14. Then, in the definition of the Toda bracket $\langle f_1, \dots, f_{d+2} \rangle$, we can replace $\mathbb{T}(f_1, f_2, f_3)$ by its subset $\mathbb{T}(f_1, f_2, f_3)_S$.

Proof. Given an arbitrary exact triangle in \mathcal{T} of the form

$$c' \xrightarrow{u'} x_2 \xrightarrow{f_2} x_1 \xrightarrow{v'} c'[1],$$

there exists an isomorphism of exact triangles

$$\begin{array}{ccccccc} c' & \xrightarrow{u'} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{v'} & c'[1] \\ \downarrow \varphi & & \parallel & & \parallel & & \downarrow \varphi[1] \\ c & \xrightarrow{u} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{v} & c[1] \end{array}$$

where the bottom row is an exact triangle in S (chosen arbitrarily). Thus, given a commutative diagram in \mathcal{T} of the form

$$\begin{array}{ccccccc} x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{f_1} & x_0 \\ \alpha \downarrow & & \parallel & & \parallel & & \uparrow \beta[1] \\ c' & \xrightarrow{u'} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{v'} & c'[1] \end{array}$$

there exists an extended commutative diagram

$$\begin{array}{ccccccccccc} & & & & x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{f_1} & x_0 \\ & & & & \parallel & & \parallel & & \parallel & & \parallel \\ & & & & \alpha \swarrow & & \parallel & & \parallel & & \beta[1] \swarrow \\ c' & \xrightarrow{u'} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{v'} & c'[1] & & & & \\ \downarrow \varphi & & \parallel & & \parallel & & \parallel & & \parallel & & \downarrow \varphi[1] \\ & & & & \parallel & & \parallel & & \parallel & & \parallel \\ & & & & \varphi \swarrow & & \parallel & & \parallel & & (\beta\varphi^{-1})[1] \swarrow \\ c & \xrightarrow{u} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{v} & c[1] & & & & \end{array}$$

inductive hypothesis,

$$\begin{aligned}
\langle f_1, -f_2, f_3, \dots, f_n, f_{n+1} \rangle &= \bigcup_{(\beta, \gamma) \in \mathbb{T}(f_1, -f_2, f_3)} \langle \beta[1], \gamma[1], f_4[1], \dots, f_n[1], f_{n+1}[1] \rangle \\
&= \bigcup_{(\beta, \gamma) \in -\mathbb{T}(f_1, f_2, f_3)} \langle \beta[1], \gamma[1], f_4[1], \dots, f_n[1], f_{n+1}[1] \rangle \\
&= \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} \langle \beta[1], -\alpha[1], f_4[1], \dots, f_n[1], f_{n+1}[1] \rangle \\
&= \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} -\langle \beta[1], \alpha[1], f_4[1], \dots, f_n[1], f_{n+1}[1] \rangle \\
&= -\langle f_1, f_2, f_3, \dots, f_n, f_{n+1} \rangle.
\end{aligned}$$

The claim follows. \square

Notation 4.1.19. Given a set X of morphisms in \mathcal{T} , we let

$$X[1] := \{f[1] \mid f \in X\},$$

and similarly for sets of pairs of morphisms in \mathcal{T} (we are thinking of Toda families as in [Definition 4.1.2](#)).

Lemma 4.1.20 ([\[CF17, Lemma 5.12\]](#)). *Suppose given a sequence of morphisms in \mathcal{T} of the form*

$$x_3 \xrightarrow{f_3} x_2 \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0.$$

Then, there is an equality

$$\mathbb{T}(f_1[1], f_2[1], f_3[1]) = -\mathbb{T}(f_1, f_2, f_3)[1].$$

Lemma 4.1.21. *Suppose given a sequence of morphisms in \mathcal{T} of the form*

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \dots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0.$$

Then, the following relation between Toda brackets holds:

$$\langle f_1, f_2, \dots, f_{d+2} \rangle[1] = (-1)^d \langle f_1[1], f_2[1], f_3[1], \dots, f_{d+2}[1] \rangle.$$

Proof. We proceed by induction on the number n of morphisms in the sequence. If $n = 2$, then the claim follows from the functoriality of the shift:

$$\langle f_1, f_2 \rangle[1] = \{f_1 \circ f_2\}[1] = \{(f_1 \circ f_2)[1]\} = \{f_1[1] \circ f_2[1]\} = \langle f_1[1], f_2[1] \rangle.$$

Suppose that the claim holds for all sequences of n composable morphisms in \mathcal{T} for some $n \geq 2$. Let f_1, \dots, f_n, f_{n+1} be a sequence of composable morphisms in \mathcal{T} . Then, in view of [Lemmas 4.1.18](#) and [4.1.20](#) and the inductive hypothesis, the Toda bracket

$$\langle f_1, f_2, \dots, f_n, f_{n+1} \rangle[1]$$

equals

$$\begin{aligned}
& \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} \langle \beta[1], \alpha[1], f_4[1], \dots, f_n[1], f_{n+1}[1] \rangle[1] \\
&= \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} (-1)^{n-2} \langle \beta[2], \alpha[2], f_4[2], \dots, f_n[2], f_{n+1}[2] \rangle \\
&= \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} (-1)^{n-1} \langle \beta[2], -\alpha[2], f_4[2], \dots, f_n[2], f_{n+1}[2] \rangle \\
&= \bigcup_{(\beta, -\alpha) \in -\mathbb{T}(f_1, f_2, f_3)} (-1)^{n-1} \langle \beta[2], -\alpha[2], f_4[2], \dots, f_n[2], f_{n+1}[2] \rangle \\
&= \bigcup_{(\beta[1], -\alpha[1]) \in -\mathbb{T}(f_1, f_2, f_3)[1]} (-1)^{n-1} \langle \beta[2], -\alpha[2], f_4[2], \dots, f_n[2], f_{n+1}[2] \rangle \\
&= \bigcup_{(\beta[1], -\alpha[1]) \in \mathbb{T}(f_1[1], f_2[1], f_3[1])} (-1)^{n-1} \langle \beta[2], -\alpha[2], f_4[2], \dots, f_n[2], f_{n+1}[2] \rangle \\
&= (-1)^{n-1} \langle f_1[1], \dots, f_n[1], f_{n+1}[1] \rangle.
\end{aligned}$$

The claim follows. \square

We combine the above results into the following equivalent way to compute the Toda bracket of a sequence of morphisms.

Proposition 4.1.22. *Let $d \geq 1$ and suppose given a sequence of morphisms in \mathcal{T} of the form*

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \dots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0.$$

Then, there is an equality

$$\langle f_1, \dots, f_{d+2} \rangle = \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} (-1)^{d-1} \langle \beta, \alpha, f_4, \dots, f_{d+2} \rangle[1].$$

Proof. According to [Definition 4.1.6](#) and [Lemma 4.1.21](#), there are equalities

$$\begin{aligned}
\langle f_1, \dots, f_{d+2} \rangle &= \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} \langle \beta[1], \alpha[1], f_4[1], \dots, f_{d+2}[1] \rangle \\
&= \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} (-1)^{d-1} \langle \beta, \alpha, f_4, \dots, f_{d+2} \rangle[1]. \quad \square
\end{aligned}$$

Proposition 4.1.23. *Let $d \geq 1$ and suppose given a sequence of morphisms in \mathcal{T} of the form*

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \dots \xrightarrow{f_3} x_2 \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_{d+2}[d].$$

Suppose that the above sequence fits as the spine of a commutative diagram of the form

$$\begin{array}{ccccccc}
& & x_{d+1} & \cdots & & x_3 & \longrightarrow & x_2 \\
& \nearrow & & & & \nearrow & & \nearrow \\
x_{d+2} & \longleftarrow +1 & \longrightarrow & x_{d.5} & \cdots & x_{3.5} & \longleftarrow +1 & \longrightarrow & x_{2.5} & \longleftarrow +1 & \longrightarrow & x_1 \\
& \searrow & & & & \searrow & & \searrow
\end{array}$$

in which the oriented triangles are exact triangles in \mathcal{T} and the morphism f_1 is the apparent (shifted) composite along the bottom row of the diagram. Then,

$$\mathbf{1}_{x_{d+2}[d]} \in (-1)^{1+\sum_{i=1}^{d+1} i} \langle f_1, f_2, \dots, f_{d+2} \rangle.$$

Proof. We proceed by induction on d . The case $d = 1$ follows from [Theorem 4.1.11](#). Suppose then that the claim holds for $d \geq 1$ and consider a sequence of morphisms $f_1, f_2, \dots, f_{d+2}, f_{d+3}$ that satisfies the conditions in the statement of the proposition. Thus, there are exact triangles

$$x_{i+1.5} \xrightarrow{h_{i+1}} x_{i+1} \xrightarrow{g_{i+1}} x_{i.5} \xrightarrow{u_i} x_{i+1.5}[1], \quad i = 1, \dots, d+1,$$

such that the triangles

$$\begin{array}{ccc} x_{i+1} & \xrightarrow{f_{i+1}} & x_i \\ & \searrow^{g_{i+1}} & \nearrow^{h_i} \\ & & x_{i.5} \end{array}, \quad i = 2, \dots, d+1,$$

commute, where we set $x_{1.5} := x_1$ and $x_{d+2.5} := x_{d+3}$, so that also $g_2 = f_2$ and $h_{d+2} = f_{d+3}$. We also set

$$f_1^{(d)} = (u_{d+1}[d-1]) \circ \dots \circ (u_3[1]) \circ u_2$$

and notice that $f_1^{(d)}: x_{2.5} \rightarrow x_{d+3}[d]$ and $f_1 = (f_1^{(d)}[1]) \circ u_1$. Having fixed the necessary notation, the induction hypothesis implies that

$$\mathbf{1}_{x_{d+3}[d]} \in (-1)^{1+\sum_{i=1}^{d+1} i} \langle f_1^{(d)}, g_3, f_4, \dots, f_{d+2}, f_{d+3} \rangle.$$

We claim that $(f_1^{(d)}, g_3) \in \mathbb{T}(f_1, f_2, f_3)$ and therefore, according to [Proposition 4.1.22](#),

$$\begin{aligned} \langle f_1, f_2, \dots, f_{d+2}, f_{d+3} \rangle &= \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} (-1)^d \langle \beta, \alpha, f_3, \dots, f_{d+2}, f_{d+3} \rangle[1] \\ &\supseteq (-1)^d \langle f_1^{(d)}, g_3, f_4, \dots, f_{d+2}, f_{d+3} \rangle[1] \\ &\supseteq (-1)^{d+2} ((-1)^{1+\sum_{i=1}^{d+1} i} \mathbf{1}_{x_{d+3}[d]})[1] \\ &= (-1)^{1+\sum_{i=1}^{d+2} i} \mathbf{1}_{x_{d+3}[d+1]}. \end{aligned}$$

To prove the claim, it suffices observe that the following diagram commutes:

$$\begin{array}{ccccccc} x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{f_1} & x_{d+3}[d+1] \\ \downarrow^{g_3} & & \parallel & & \parallel & & \uparrow^{f_1^{(d)}[1]} \\ x_{2.5} & \xrightarrow{h_2} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{u_1} & x_{2.5}[1]. \end{array}$$

This finishes the proof. \square

We also need the following lemma.

Lemma 4.1.24. *Suppose given a sequence of morphisms in \mathcal{T} of the form*

$$x_3 \xrightarrow{f_3} x_2 \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_3[1].$$

The following statements are equivalent:

- (1) *We have $\mathbf{1}_{x_3[1]} \in \langle f_1, f_2, f_3 \rangle$.*
- (2) *There exist an object $y \in \mathcal{T}$ and morphisms $g: y \rightarrow x_2$ and $h: x_1 \rightarrow y[1]$ such that the sequence*

$$x_3 \oplus y \xrightarrow{(f_3 \ g)} x_2 \xrightarrow{f_2} x_1 \xrightarrow{\begin{pmatrix} f_1 \\ h \end{pmatrix}} (x_3 \oplus y)[1]$$

is an exact triangle in \mathcal{T} .

Proof. By definition, $\mathbf{1}_{x_2[1]} \in \langle f_0, f_1, f_2 \rangle$ if and only if there exists a commutative diagram in \mathcal{T} of the form

$$\begin{array}{ccccccc}
x_3 & \xrightarrow{f_3} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{f_1} & x_3[1] \\
\alpha \downarrow & & \parallel & & \parallel & & \uparrow \beta[1] \\
c & \xrightarrow{u} & x_2 & \xrightarrow{f_2} & x_1 & \xrightarrow{v} & c[1]
\end{array}$$

in which the bottom row is an exact triangle in \mathcal{T} and $(\beta\alpha)[1] = \mathbf{1}_{x_3[1]}$ or, equivalently $\beta\alpha = \mathbf{1}_{x_3}$. In particular, $c \cong x_3 \oplus y$ for some object $y \in \mathcal{T}$. If we identify c with $x_3 \oplus y$ we see that the morphism u has components $f_3: x_3 \rightarrow x_2$ and $g: y \rightarrow x_2$ for some morphism g . Similarly, under the identification $c[1] = (x_3 \oplus y)[1] = x_3[1] \oplus y[1]$ the morphism v has components $f_1: x_1 \rightarrow x_3[1]$ and $h: x_1 \rightarrow y[1]$ for some morphism h . The claim follows. \square

In the proof of [Theorem 4.1.12](#) we will use elementary properties of Poincaré polynomials of periodic graded modules.

Definition 4.1.25. Let A be a graded algebra with a degree d unit. The *Poincaré polynomial* of a degree-wise finite-dimensional A -module M is

$$p(M) = \sum_{i=0}^{d-1} \dim_{\mathbf{k}} M^i \cdot t^i \in \mathbb{Z}[t]/(t^d - 1).$$

We now show some elementary properties of these Poincaré polynomials. All proofs are simple, albeit somewhat tedious, exercises, so we only sketch them.

Proposition 4.1.26. *Given a graded algebra A with a degree d unit $u \in A^d$, the Poincaré polynomials of degree-wise finite-dimensional A -modules satisfy the following properties:*

- (1) $p(M \oplus N) = p(M) + p(N)$.
- (2) *Given a short exact sequence $0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$, $p(N) = p(M) + p(P)$.*
- (3) $p(M) = 0 \Leftrightarrow M = 0$.
- (4) $p(M) \in \mathbb{Z}$ if and only if $M^i = 0$ for $i \notin d\mathbb{Z}$.
- (5) If $p(M) \in \mathbb{Z}$ and $N \subset M$ then $p(N) \in \mathbb{Z}$.
- (6) $p(M(-1)) = t \cdot p(M)$.
- (7) $p(M(d)) = p(M)$.
- (8) *Given an exact sequence*

$$\begin{array}{ccc}
& & N \\
& \nearrow f & \\
M & & P \\
& \longleftarrow h & \longleftarrow +1
\end{array}$$

if $K = \ker g = \operatorname{im} f$ then

$$p(M) + tp(N) - tp(P) = (t+1)p(K).$$

Proof. The first five properties are obvious. For (6), we use the following three facts:

- $M(-1)^i = M^{i-1}$,
- multiplication by u induces an isomorphism $M(-1)^0 = M^{-1} \cong M^{d-1}$,
- $t^d = 1$ in $\mathbb{Z}[t]/(t^d - 1)$.

Property (7) is a direct consequence of (6). For (8), we break the exact sequence

$$0 \rightarrow K(-1) \rightarrow N(-1) \xrightarrow{g} P(-1) \xrightarrow{h} M \rightarrow K \rightarrow 0$$

into three short exact sequences and apply (2) and (6). \square

Proposition 4.1.27. *Let $d \geq 1$. Given a graded algebra A with a degree d unit and an exact sequence of degree-wise finite-dimensional A -modules,*

$$\begin{array}{ccccc}
& & C_{d+1} & \xrightarrow{f_{d+1}} & \cdots & \xrightarrow{f_3} & C_2 & & \\
& & \nearrow^{f_{d+2}} & & & & \searrow^{f_2} & & \\
C_{d+2} & \xleftarrow{f_{d+2}} & & & & & & \xrightarrow{+d} & C_1
\end{array}$$

if $K = \ker f_2$ then the following relation between Poincaré polynomials holds,

$$(-1)^d (p(C_2) - p(C_1)) + \sum_{i=3}^{d+2} (-1)^i p(C_i) = ((-1)^d - 1) p(K).$$

Proof. Consider $K_i = \ker f_i$ and the short exact sequences

$$\begin{array}{l}
0 \rightarrow K_i \rightarrow C_i \rightarrow K_{i-1} \rightarrow 0, \quad 2 \leq i \leq d+2, \\
0 \rightarrow K_1 \rightarrow C_1 \rightarrow K_{d+2}(d) \rightarrow 0.
\end{array}$$

Now it is just a matter of applying (2) and (7) from [Proposition 4.1.26](#). \square

Proposition 4.1.28. *Let $d \geq 1$. Given a graded algebra A with a degree d unit and exact sequences of degree-wise finite-dimensional A -modules,*

$$\begin{array}{ccccccc}
& & M_{d+1} & & M_d & \cdots & & M_2 & \\
& & \nearrow & & \nearrow & & & \nearrow & \\
M_{d+2} & \xleftarrow{+1} & X_{d.5} & \xleftarrow{+1} & X_{d-1.5} & \cdots & & X_{2.5} & \xleftarrow{+1} & M_1
\end{array}$$

if $K_i = \ker g_i$ then

$$\begin{aligned}
p(M_2) + p(M_{d+2}) - p(M_1) + \sum_{i=3}^{d+1} p(M_i) t^{d+2-i} = \\
p(K_2) + p(K_{d+1}) + \sum_{i=3}^{d+1} (p(K_i) + p(K_{i-1})) t^{d+2-i}
\end{aligned}$$

Proof. This follows from [Proposition 4.1.26](#) (8). \square

Corollary 4.1.29. *In the setting of [Proposition 4.1.28](#), if the M_i are concentrated in degrees $d\mathbb{Z}$ then*

$$(-1)^d (p(M_2) - p(M_1)) + \sum_{i=3}^{d+2} (-1)^i p(M_i) = ((-1)^d - 1) p(K_2).$$

Proof. By [Proposition 4.1.26](#) (4) and (5), $p(M_i), p(K_i) \in \mathbb{Z}$. In the polynomial equation of [Proposition 4.1.28](#), both polynomials must have the same coefficients in the same degrees. This translates into d equations involving $p(M_i), p(K_i) \in \mathbb{Z}$. From these equations, it is easy to deduce the equation in the statement. \square

We are ready to prove [Theorem 4.1.12](#).

Proof of [Theorem 4.1.12](#). Let $\mathcal{C} \subseteq \mathcal{T}$ be a d -rigid subcategory such that $\text{add}(\mathcal{C}) = \mathcal{C}$ and $\mathcal{C}[d] = \mathcal{C}$, and

$$(4.1.30) \quad c_{d+2} \xrightarrow{f_{d+2}} c_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_3} c_2 \xrightarrow{f_2} c_1 \xrightarrow{f_1} c_{d+2}[d]$$

a sequence of morphisms in \mathcal{C} with $d \geq 1$.

(1) \Rightarrow (2) Suppose that the sequence (4.1.30) satisfies the conditions in [Theorem 4.1.12](#)(1). The fact, for each $y \in \mathcal{C}$, the induced sequence of vector spaces

$$\mathcal{T}(y, c_{d+2}) \rightarrow \cdots \rightarrow \mathcal{T}(y, c_2) \rightarrow \mathcal{T}(y, c_1) \rightarrow \mathcal{T}(y, c_{d+2}[d]) \rightarrow \mathcal{T}(y, c_{d+1}[d])$$

is exact follows from standard arguments using that $\mathcal{C} \subseteq \mathcal{T}$ is a d -rigid subcategory, see for example the first paragraph in the proof of [Lin19, Lemma 4.6]. That

$$\mathbf{1}_{c_{d+2}[d]} \in (-1)^{1+\sum_{i=1}^{d+1}} \langle f_1, f_2, \dots, f_{d+2} \rangle$$

follows from **Proposition 4.1.23**.

(2) \Rightarrow (1) If $d = 1$, then we are precisely in the setting of **Theorem 4.1.11**; hence, we may assume that $d > 1$. Inductively, we shall construct a commutative diagram

$$\begin{array}{ccc} & c_{d+1} & \cdots \\ f_{d+2} \nearrow & & \searrow g_{d+1} \\ c_{d+2} & \xleftarrow{u_d} +1 - & x_{d.5} \end{array} \quad \cdots \quad \begin{array}{ccccc} & c_3 & \xrightarrow{f_3} & c_2 & \\ h_3 \nearrow & & \searrow g_3 & h_2 \nearrow & \searrow f_2 \\ x_{3.5} & \xleftarrow{u_2} +1 - & x_{2.5} & \xleftarrow{u_1} +1 - & c_1 \end{array}$$

with the following properties:

- The triangles

$$x_{k.5} \xrightarrow{h_k} c_k \xrightarrow{g_k} x_{k-1.5} \xrightarrow{u_{k-1}} x_{k.5}[1], \quad 2 \leq k \leq d,$$

are exact, where $x_{1.5} = c_1$ and $p_2 = f_2$;

- We have $\mathbf{1}_{c_{d+2}[1]} \in \langle u_d, g_{d+1}, f_{d+2} \rangle$, although the triangle

$$c_{d+2} \xrightarrow{f_{d+2}} c_{d+1} \xrightarrow{g_{d+1}} x_{d.5} \xrightarrow{u_d} c_{d+2}[1]$$

is not (yet) known to be exact.

- There are equalities $f_k = h_{k-1} \circ g_k$, for all $3 \leq k \leq d+1$.
- There is an equality $f_1 = u_d[d-1] \circ \cdots \circ u_2[1] \circ u_1$.

In the first step we notice that, since the morphism $(-1)^{1+\sum_{i=1}^{d+1} i} \mathbf{1}_{c_{d+2}[d]}$ lies in

$$\langle f_1, f_2, \dots, f_{d+2} \rangle = \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)} (-1)^{d-1} \langle \beta, \alpha, f_4, \dots, f_{d+2} \rangle[1],$$

there exist a pair of morphisms $(f_1^{(d)}, g_3) \in \mathbb{T}(f_1, f_2, f_3)$ such that

$$(-1)^{1+\sum_{i=1}^{d+1} i} \mathbf{1}_{c_{d+2}[d]} \in (-1)^{d-1} \langle f_1^{(d)}, g_3, f_4, \dots, f_{d+2} \rangle[1]$$

or, equivalently,

$$(-1)^{1+\sum_{i=1}^d i} \mathbf{1}_{c_{d+2}[d-1]} \in \langle f_1^{(d)}, g_3, f_4, \dots, f_{d+2} \rangle.$$

In particular, there exists a diagram of the form

$$\begin{array}{ccccc} & c_2 & \xrightarrow{f_2} & c_1 & \\ f_3 \nearrow & & \searrow h_2 & u_1 \nearrow & +1 \\ c_3 & \xrightarrow{g_3} & x_{2.5} & \xleftarrow{-1} & c_{d+2}[d] \end{array} \quad \begin{array}{c} \searrow f_1 \\ \downarrow f_1^{(d)} \end{array}$$

in which the oriented triangle is exact. Thus, we obtain a diagram of the form

$$\begin{array}{ccccccc} & c_{d+1} & \xrightarrow{f_{d+1}} & \cdots & \xrightarrow{f_4} & c_3 & \xrightarrow{f_3} & c_2 \\ f_{d+2} \nearrow & & & & & \searrow g_3 & h_2 \nearrow & \searrow f_2 \\ c_{d+2} & \xleftarrow{f_1^{(d)}} & + (d-1) & \xleftarrow{-1} & x_{2.5} & \xleftarrow{u_1} & +1 - & c_1 \end{array}$$

We may then continue the inductive procedure with the sequence $f_1^{(d)}, g_3, f_4, \dots, f_{d+2}$

$$(-1)^{1+(1+2)} \mathbf{1}_{c_{d+2}[1]} = \mathbf{1}_{c_{d+2}[1]} \in \langle u_d, g_{d+1}, f_{d+2} \rangle,$$

as required). It remains to prove that the sequence

$$c_{d+2} \xrightarrow{f_{d+2}} c_{d+1} \xrightarrow{g_{d+1}} x_{d.5} \xrightarrow{u_d} c_{d+2}[1]$$

is an exact triangle. Since $\mathbf{1}_{c_{d+2}[1]} \in \langle u_d, g_{d+1}, f_{d+2} \rangle$, [Lemma 4.1.24](#) shows that there exists an object $y \in \mathcal{T}$ and an exact triangle of the form

$$y \oplus c_{d+2} \xrightarrow{\begin{pmatrix} f_{d+2} & g \end{pmatrix}} c_{d+1} \xrightarrow{g_{d+1}} x_{d.5} \xrightarrow{\begin{pmatrix} u_d \\ h \end{pmatrix}} (y \oplus c_{d+2})[1].$$

We need to prove that $y = 0$; since $\text{thick}(c) = \text{thick}(\mathcal{C}) = \mathcal{T}$ it suffices to prove that $\mathcal{T}(c[i], y) = 0$ for all $i \in \mathbb{Z}$. Moreover, since $\mathcal{C}[d] = \mathcal{C} = \text{add}(c)$ and c is basic, $c[d] \cong c$ and the graded endomorphism algebra A of c in \mathcal{T} is of the form $\Lambda(\sigma, d)$ in [Section 4.5.1](#), so it has a degree d unit. Moreover, $M_x = \bigoplus_{i \in \mathbb{Z}} \mathcal{T}(c[i], x)$ is a degree-wise finite-dimensional A -module for any $x \in \mathcal{T}$. We have to show that $M_y = 0$. We will denote $p(x) = p(M_x)$, and we will show that $p(y) = 0$, which suffices by [Proposition 4.1.26 \(3\)](#).

Applying $\bigoplus_{i \in \mathbb{Z}} \mathcal{T}(c[i], -)$ to

$$\begin{array}{ccccc} & & c_{d+1} & \xrightarrow{f_{d+1}} & \cdots & \xrightarrow{f_3} & c_2 & & \\ & f_{d+2} \nearrow & & & & & & & f_2 \searrow \\ c_{d+2} & & & & & & & & c_1 \\ & & & \xleftarrow{f_{d+2}} & +d & \xrightarrow{\quad} & & & \end{array}$$

and

$$\begin{array}{ccccccc} & & c_{d+1} & & c_d & & \cdots & & c_2 & & \\ & (f_{d+2} \ g) \nearrow & & \xrightarrow{g_{d+1}} & \xrightarrow{h_d} & & & & & \xrightarrow{h_2} & \searrow f_2 \\ y \oplus c_{d+2} & & & \xleftarrow{\begin{pmatrix} u_{d+2} \\ h \end{pmatrix}} +1 - & x_{d.5} & \xleftarrow{u_{d-1}} +1 - & x_{d-1.5} & \cdots & x_{2.5} & \xleftarrow{u_1} +1 - & c_1 \end{array}$$

we obtain diagrams of A -modules as in [Proposition 4.1.27](#) and [Proposition 4.1.28](#). By [Proposition 4.1.27](#), if $K = \ker \bigoplus_{i \in \mathbb{Z}} \mathcal{T}(c[i], f_2)$,

$$(-1)^d (p(c_2) - p(c_1)) + \sum_{i=3}^{d+2} (-1)^i p(c_i) = ((-1)^d - 1) p(K).$$

Moreover, since $c_i \in \mathcal{C}$ and \mathcal{C} is d -rigid, M_{c_i} is concentrated in $d\mathbb{Z}$. Hence, by [Corollary 4.1.29](#),

$$(-1)^d (p(c_2) - p(c_1)) + \sum_{i=3}^{d+1} (-1)^i p(c_i) + p(y) + p(c_{d+2}) = ((-1)^d - 1) p(K).$$

Both equations are identical, except for the fact that the second one has an extra term, $p(y)$. Therefore, $p(y) = 0$, as we wanted to prove. \square

We conclude this section with the following result.

Proposition 4.1.31 ([\[Sag08, Prop. 4.10\]](#)). *Let $\mathcal{C} \subseteq \mathcal{T}$ a d -rigid subcategory such that $\text{add}(\mathcal{C}) = \mathcal{C}$ and $\mathcal{C}[d] = \mathcal{C}$. Suppose given a sequence of morphisms in \mathcal{C} of the form*

$$c_{d+2} \xrightarrow{f_{d+2}} c_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} c_1 \xrightarrow{f_1} c_0$$

Then, the Toda bracket $\langle f_1, \dots, f_{d+2} \rangle$ is an element of the quotient

$$\text{Toda}(f_{d+2}, f_1) := \frac{\mathcal{T}(c_{d+2}[d], c_0)}{f_1 \cdot \mathcal{T}(c_{d+2}[d], c_1) + \mathcal{T}(c_{d+1}[d], c_0) \cdot f_{d+2}[d]}.$$

4.2. Massey products versus Toda brackets. Let \mathcal{A} be a DG category. We recall that, similarly to the cohomology of a space, the graded category $H^\bullet(\mathcal{A})$ is endowed with so-called Massey products [BK90, Sec. 5.A]. Massey products are closely related to Toda brackets. In fact, it is stated in *loc. cit.* that

“It can be verified that for an enhanced triangulated category the Massey products coincide with the Toda brackets.”

As it turns out, this statement indeed holds, but only up to a suitable sign. Since this agreement is crucial to our main results, we include a proof for the convenience of the reader (see also [Bod14] for an alternative proof that involves the language of twisted complexes).

We begin by recalling the definition of the Massey products, see for example [May69].

Definition 4.2.1. Let \mathcal{A} be a DG category and suppose given a sequence in $H^\bullet(\mathcal{A})$ of the form

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0$$

such that f_i is homogeneous of degree $|f_i|$, where $d \geq 0$.

- (1) A *defining system* (for the above sequence) is a set

$$\{g_{ij} : x_j \rightarrow x_i \mid 0 \leq i < j \leq d+2, j-i < d+2\}$$

of morphisms in \mathcal{A} that are homogeneous of degree

$$|g_{ij}| = i - j + 1 + \sum_{i < k \leq j} |f_k|$$

with the following two properties:

- (a) The equality

$$\partial(g_{ij}) = \sum_{i < k < j} (-1)^{|g_{ik}|-1} g_{ik} g_{kj}$$

holds (notice that $|g_{ik}| - 1 = i - k + \sum_{i < \ell \leq k} |f_\ell|$). In particular,

$$\partial(g_{i-1,i}) = 0$$

so that $g_{i-1,i}$ is a cocycle of degree $|f_i|$.

- (b) The morphism $g_{i-1,i}$ represents f_i in $H^\bullet(\mathcal{A})$.

- (2) The *Massey product* is the subset of

$$\langle\langle f_1, \dots, f_{d+2} \rangle\rangle \subseteq H^{-d + \sum_{i=1}^{d+2} |f_i|}(\mathcal{A}(x_{d+2}, x_0))$$

consisting of the classes of the cocycles

$$\sum_{0 < k < d+2} (-1)^{|g_{0k}|-1} g_{0k} g_{k,d+2} = \sum_{0 < k < d+2} (-1)^{-k + \sum_{0 < \ell \leq k} |f_\ell|} g_{0k} g_{k,d+2}$$

for all possible defining systems.

Remark 4.2.2. In the setting of **Definition 4.2.1**, the degree of the Massey product $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$ is more recognisable by letting $n = d+2$ so that $\langle\langle f_1, \dots, f_n \rangle\rangle$ is a subset of the cohomology of $\mathcal{A}(x_n, x_0)$ in degree $2 - n + \sum_i |f_i|$.

Remark 4.2.3. The Massey product $\langle\langle f_1, f_2 \rangle\rangle$ of a pair of (homogeneous) composable morphisms in $H^\bullet(\mathcal{A})$ is always defined and is given by the singleton

$$\langle\langle f_1, f_2 \rangle\rangle = \left\{ (-1)^{|f_1|-1} f_1 \circ f_2 \right\}.$$

We see that, if $|f_1| = 0$, the Massey product of f_1 and f_2 is the sign-twisted composite $-f_1 \circ f_2$ (compare with **Definition 4.1.6** and **Theorem 4.2.6**).

Remark 4.2.4. In the setting of [Definition 4.2.1](#), if $d \geq 1$, then the Massey product is defined (i.e. non-empty) if and only if

$$\forall 0 \leq i < j \leq d+2, j-i < d+2, \quad 0 \in \langle\langle f_i, \dots, f_j \rangle\rangle.$$

Remark 4.2.5. Let \mathcal{A} be a small DG category. Suppose given a sequence of degree 0 morphisms

$$M_{d+2} \xrightarrow{f_{d+2}} M_{d+1} \xrightarrow{f_{d+1}} \dots \xrightarrow{f_2} M_1 \xrightarrow{f_1} M_0$$

in the triangulated category $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$. By definition, their Massey product satisfies

$$\langle\langle f_1, \dots, f_{d+2} \rangle\rangle \subseteq H^{-d}(\mathbf{hom}_{\mathcal{A}}(M_{d+2}, M_0)) \cong H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))(M_{d+2}, M_0[-d]),$$

while their Toda bracket satisfies

$$\langle f_1, \dots, f_{d+2} \rangle \subseteq H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))(M_{d+2}[d], M_0).$$

Of course, we may compare both brackets by means of the isomorphism

$$H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))(M_{d+2}, M_0[-d]) \cong H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))(M_{d+2}[d], M_0)$$

induced by the d -fold shift functor.

We shall prove the following agreement result for Massey products and Toda brackets.

Theorem 4.2.6. *Let \mathcal{A} be a small DG category. Suppose given a sequence of degree 0 morphisms*

$$M_{d+2} \xrightarrow{f_{d+2}} M_{d+1} \xrightarrow{f_{d+1}} \dots \xrightarrow{f_2} M_1 \xrightarrow{f_1} M_0$$

in the triangulated category $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$. Then, their Massey product and their Toda bracket are related by the formula

$$\langle\langle f_1, \dots, f_{d+2} \rangle\rangle[d] = (-1)^{\sum_{i=1}^{d+1} i} \langle f_1, \dots, f_{d+2} \rangle.$$

Remark 4.2.7. Let \mathcal{A} be a small DG category. Since $\mathbf{D}(\mathcal{A}) \subseteq H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$ is a full triangulated subcategory, [Theorem 4.2.6](#) also establishes a relationship between Massey products and Toda brackets in $\mathbf{D}(\mathcal{A})$.

For the proof of [Theorem 4.2.6](#), we need the following observation. Given a small DG category \mathcal{A} and a morphism $f: M \rightarrow N$ in $\mathbf{dgMod}(\mathcal{A})$, recall that the *cone of f* , denoted by $\mathbf{cone}(f)$, is the DG \mathcal{A} -module whose underlying graded \mathcal{A} -module is $M(1) \oplus N$ equipped with the differential

$$d_{\mathbf{cone}(f)} := d_{M[1]} \oplus d_N + \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix};$$

above, $M \mapsto M(1)$ denotes the shift of the underlying graded \mathcal{A} -module of M . The definition above ensures that the canonical split short exact sequence of graded \mathcal{A} -modules

$$0 \rightarrow N \xrightarrow{\iota} \mathbf{cone}(f) \xrightarrow{\pi} M[1] \rightarrow 0$$

is in fact a (not necessarily split) short exact sequence of DG \mathcal{A} -modules. The above sequence yields an exact triangle

$$M \xrightarrow{f} N \xrightarrow{\iota} \mathbf{cone}(f) \xrightarrow{\pi} M[1]$$

in the triangulated category $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$.

Proposition 4.2.8. *Let \mathcal{A} be a small DG category. Suppose given a sequence of degree 0 morphisms*

$$M_{d+2} \xrightarrow{f_{d+2}} M_{d+1} \xrightarrow{f_{d+1}} \dots \xrightarrow{f_2} M_1 \xrightarrow{f_1} M_0$$

in the triangulated category $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$, with $d \geq 1$. Let S be the set consisting of the exact triangles of the precise form

$$\mathbf{cone}(g_{12})[-1] \xrightarrow{\pi[-1]} M_2 \xrightarrow{f_2} M_1 \xrightarrow{-\iota} \mathbf{cone}(g_{12}),$$

where g_{12} ranges over all degree 0 cocycles $M_2 \rightarrow M_1$ in $\mathbf{dgMod}(\mathcal{A})$ that represent the morphism f_2 . Then, the Massey product $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$ satisfies the formula

$$\langle\langle f_1, \dots, f_{d+2} \rangle\rangle = \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)_S} \langle\langle \beta, \alpha, f_4, \dots, f_{d+2} \rangle\rangle.$$

Proof. Let

$$\{g_{ij} : M_j \rightarrow M_i \mid 0 \leq i < j \leq d+2, j-i < d+2\}$$

be a defining system for the Massey product $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$. Define the morphisms in $\mathbf{dgMod}(\mathcal{A})$

$$\alpha := \begin{pmatrix} g_{23} \\ g_{13} \end{pmatrix} : M_3 \longrightarrow \mathbf{cone}(g_{12})[-1],$$

$$\beta := (g_{02} \quad -g_{01}) : \mathbf{cone}(g_{12})[-1] \longrightarrow M_0[-1].$$

Notice that $|\alpha| = 0 = |\beta|$. We claim that the pair formed by the cohomology classes of these morphisms $(\{\beta\}, \{\alpha\}) \in \mathbb{T}(f_1, f_2, f_3)_S$. Notice first that α and β are indeed cocycles since, in view of the equations satisfied by a defining system, we have

$$\begin{aligned} \partial(\alpha) &= \begin{pmatrix} d_{M_2} & 0 \\ -g_{12} & -d_{M_1} \end{pmatrix} \circ \begin{pmatrix} g_{23} \\ g_{13} \end{pmatrix} - \begin{pmatrix} g_{23} \\ g_{13} \end{pmatrix} \circ d_{M_3} \\ &= \begin{pmatrix} d_{M_2} \circ g_{23} - g_{23} \circ d_{M_3} \\ -g_{12}g_{23} - (d_{M_1} \circ g_{13} + g_{13} \circ d_{M_3}) \end{pmatrix} \\ &= \begin{pmatrix} \partial(g_{23}) \\ \partial(g_{13}) - \partial(g_{13}) \end{pmatrix} = 0 \end{aligned}$$

and

$$\begin{aligned} \partial(\beta) &= -d_{M_0} \circ (g_{02} \quad -g_{01}) - (g_{02} \quad -g_{01}) \circ \begin{pmatrix} d_{M_2} & 0 \\ -g_{12} & -d_{M_1} \end{pmatrix} \\ &= (-(d_{M_0} \circ g_{02} + g_{02} \circ d_{M_2}) - g_{01}g_{12} \quad d_{M_0} \circ g_{01} - g_{01} \circ d_{M_1}) \\ &= (-\partial(g_{02}) + \partial(g_{02}) \quad \partial(g_{01})) = 0. \end{aligned}$$

Here we use that $d_{M_0[-1]} = -d_{M_0}$. Thus, to exhibit the membership relation $(\{\beta\}, \{\alpha\}) \in \mathbb{T}(f_1, f_2, f_3)$, it suffices to observe that the diagram (see [Remark 4.1.5](#))

$$\begin{array}{ccccc} & & M_2 & \xrightarrow{f_2} & M_1 & & \\ & & \swarrow & & \searrow & & \\ & & & & & & \\ f_3 \nearrow & & & & & & \\ & & M_3 & \xrightarrow{\{\alpha\}} & \mathbf{cone}(g_{12})[-1] & \xrightarrow{\{\beta\}} & -1 \longrightarrow & M_0 \\ & & & & \swarrow & & \nearrow & \\ & & & & & & & \end{array}$$

commutes in $H^\bullet(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$, for g_{01} and g_{23} represent the morphisms f_1 and f_3 and $\pi[-1]$ and ι are the apparent projection and inclusion into the corresponding direct summands.

We wish to use the given defining system for the Massey product $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$ to construct a defining system for the Massey product $\langle\langle \{\beta\}, \{\alpha\}, f_4, \dots, f_{d+2} \rangle\rangle$. We claim that the collection of morphisms

$$\bar{g}_{01} := \beta = (g_{02} \quad -g_{01}) : \mathbf{cone}(g_{12})[-1] \longrightarrow M_0;$$

$$\bar{g}_{0j} := g_{0,j+1} : M_{j+1} \longrightarrow M_0[-1], \quad d+2 > j \geq 2;$$

$$\bar{g}_{1j} := \begin{pmatrix} g_{2,j+1} \\ g_{1,j+1} \end{pmatrix} : M_{j+1} \longrightarrow \mathbf{cone}(g_{12})[-1], \quad d+2 > j \geq 2;$$

$$\bar{g}_{ij} := g_{i+1,j+1} : M_{j+1} \longrightarrow M_{i+1}, \quad d+2 > j > i \geq 2;$$

is such a defining system. Notice that

- $\bar{g}_{01} = \beta$ represents $\{\beta\}$;
- $\bar{g}_{12} = \alpha$ represents $\{\alpha\}$;
- $\bar{g}_{i-1,i} = g_{i,i+1}: M_{i+1} \rightarrow M_i$ represents f_{i+1} .

We now analyse the differentials $\partial(\bar{g}_{i,j})$ in the four cases:

- We have already shown that $\partial(\bar{g}_{01}) = \partial(\beta) = 0$.
- If $j \geq 2$, then

$$\begin{aligned}
\partial(\bar{g}_{0j}) &= -\partial(g_{0,j+1}) \\
&= - \sum_{0 < \ell < j+1} (-1)^{-\ell} g_{0\ell} g_{\ell,j+1} \\
&= - (g_{02} \quad -g_{01}) \begin{pmatrix} g_{2,j+1} \\ g_{1,j+1} \end{pmatrix} + \sum_{2 < \ell < j+1} (-1)^{1-\ell} g_{0\ell} g_{\ell,j+1} \\
&= -\bar{g}_{01} \bar{g}_{1j} + \sum_{2 < \ell < j+1} (-1)^{1-\ell} \bar{g}_{0,\ell-1} \bar{g}_{\ell-1,j} \\
&= \sum_{0 < k < j} (-1)^{-k} \bar{g}_{0k} \bar{g}_{kj},
\end{aligned}$$

where the equality $\partial(\bar{g}_{0j}) = -\partial(g_{0,j+1})$ stems from the fact that the target of \bar{g}_{0j} is the DG \mathcal{A} -module $M_0[-1]$, which has the differential $-d_{M_0}$.

- If $i = 1$ and $j \geq 2$, then

$$\begin{aligned}
\partial(\bar{g}_{1j}) &= \begin{pmatrix} d_{M_2} & 0 \\ -g_{12} & -d_{M_1} \end{pmatrix} \circ \begin{pmatrix} g_{2,j+1} \\ g_{1,j+1} \end{pmatrix} - (-1)^{|\bar{g}_{1j}|} \begin{pmatrix} g_{2,j+1} \\ g_{1,j+1} \end{pmatrix} \circ d_{M_{j+1}} \\
&= \begin{pmatrix} d_{M_2} \circ g_{2,j+1} - (-1)^{|g_{2,j+1}|} g_{2,j+1} \circ d_{M_{j+1}} \\ -g_{12} g_{2,j+1} - (d_{M_1} \circ g_{1,j+1} - (-1)^{|g_{1,j+1}|} g_{1,j+1} \circ d_{M_{j+1}}) \end{pmatrix} \\
&= \begin{pmatrix} \partial(g_{2,j+1}) \\ -g_{12} g_{2,j+1} - \partial(g_{1,j+1}) \end{pmatrix} \\
&= \begin{pmatrix} \sum_{2 < \ell < j+1} (-1)^{2-\ell} g_{2\ell} g_{\ell,j+1} \\ -\sum_{2 < \ell < j+1} (-1)^{1-\ell} g_{1\ell} g_{\ell,j+1} \end{pmatrix} \\
&= \sum_{2 < \ell < j+1} (-1)^{2-\ell} \begin{pmatrix} g_{2\ell} \\ g_{1\ell} \end{pmatrix} g_{\ell,j+1} \\
&= \sum_{2 < \ell < j+1} (-1)^{2-\ell} \bar{g}_{1,\ell-1} \bar{g}_{\ell-1,j} \\
&= \sum_{1 < k < j} (-1)^{1-k} \bar{g}_{1k} \bar{g}_{kj}.
\end{aligned}$$

- If $i \geq 2$, then

$$\begin{aligned}
\partial(\bar{g}_{ij}) &= \partial(g_{i+1,j+1}) = \sum_{i+1 < k+1 < j+1} (-1)^{(i+1)-(k+1)} g_{i+1,k+1} g_{k+1,j+1} \\
&= \sum_{i < k < j} (-1)^{i-k} \bar{g}_{ik} \bar{g}_{kj}.
\end{aligned}$$

This shows that the collection of morphisms $\{\bar{g}_{ij}\}$ is indeed a defining system for the Massey product $\langle\langle \{\beta\}, \{\alpha\}, f_4, \dots, f_{d+2} \rangle\rangle$.

Finally, the above process is clearly reversible: Given $(\{\beta\}, \{\alpha\}) \in \mathbb{T}(f_1, f_2, f_3)_S$ and a defining system for the Massey product $\langle\langle \{\beta\}, \{\alpha\}, f_4, \dots, f_{d+2} \rangle\rangle$, one obtains a defining system for the Massey product $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$ by means of the above formulas. This finishes the proof. \square

We are ready to prove [Theorem 4.2.6](#).

Proof of [Theorem 4.2.6](#). Given a sequence of degree 0 morphisms

$$M_{d+2} \xrightarrow{f_{d+2}} M_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} M_1 \xrightarrow{f_1} M_0$$

in the triangulated category $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$, for \mathcal{A} a small DG category, we need to prove that

$$\langle\langle f_1, \dots, f_{d+2} \rangle\rangle[d] = (-1)^{\sum_{i=1}^{d+1} i} \langle f_1, \dots, f_{d+2} \rangle.$$

If $d = 0$, then the claim follows from the discussion in [Remark 4.2.3](#). For $d \geq 1$, the claim follows from a straightforward induction using [Propositions 4.1.22](#) and [4.2.8](#) and [Lemma 4.1.15](#). Indeed, suppose that the claim holds for all sequences of n composable morphisms for some $n \geq 2$. Let f_1, \dots, f_n, f_{n+1} be a sequence of composable homogeneous morphisms in $H^0(\mathbf{dgMod}_{\mathbf{dg}}(\mathcal{A}))$. Then, the $(n-1)$ -fold shifted Massey product

$$\langle\langle f_1, \dots, f_n, f_{n+1} \rangle\rangle[n-1]$$

equals

$$\begin{aligned} & \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)_S} \langle\langle \beta, \alpha, f_4, \dots, f_n, f_{n+1} \rangle\rangle[n-1] \\ &= \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)_S} (-1)^{\sum_{i=1}^{n-1} i} \langle \beta, \alpha, f_4, \dots, f_n, f_{n+1} \rangle[1] \\ &= (-1)^{\sum_{i=1}^{n-1} i} (-1)^n \bigcup_{(\beta, \alpha) \in \mathbb{T}(f_1, f_2, f_3)_S} (-1)^{n-2} \langle \beta, \alpha, f_4, \dots, f_n, f_{n+1} \rangle[1] \\ &= (-1)^{\sum_{i=1}^n i} \langle f_1, \dots, f_n, f_{n+1} \rangle, \end{aligned}$$

where S is as in [Proposition 4.2.8](#). The claim follows. \square

4.3. Massey products and minimal A_∞ -structures. In this subsection we relate (under some conditions) higher operations in a minimal A_∞ -category and Massey products. We begin by recalling the necessary definitions.

4.3.1. Reminder on A_∞ -categories. We recall the most basic aspects of the theory of A_∞ -categories and its relationship to that of DG categories. We refer the reader to [\[Kel01, LH\]](#) for details.

An A_∞ -category is a categorical structure \mathcal{A} consisting of a class of objects and graded vector spaces of morphisms

$$\mathcal{A}(x, y), \quad x, y \in \mathcal{A}.$$

Moreover, \mathcal{A} is equipped with higher composition operations

$$m_n: \mathcal{A}(x_1, x_0) \otimes \mathcal{A}(x_2, x_1) \otimes \cdots \otimes \mathcal{A}(x_n, x_{n-1}) \longrightarrow \mathcal{A}(x_n, x_0), \quad n \geq 1$$

given by graded morphisms of degree $2-n$ that must satisfy a certain infinite system of equations that we do not recall here. We only mention that, in particular, these equations imply the following:

- For each $x, y \in \mathcal{A}$, the pair $(\mathcal{A}(x, y), m_1)$ is a cochain complex, that is $m_1 \circ m_1 = 0$.
- The differential m_1 is a graded derivation with respect to the composition operation

$$m_2: \mathcal{A}(y, z) \otimes \mathcal{A}(x, y) \longrightarrow \mathcal{A}(x, z),$$

(that is, it satisfies the graded Leibniz rule).

- Although the composition operation m_2 is not necessarily associative, the failure of the associativity is controlled by the ternary operation m_3 in a precise sense. In particular, as in the case of DG algebras, \mathcal{A} has an associated graded category $H^\bullet(\mathcal{A})$ as well as an ordinary category $H^0(\mathcal{A})$, whose (associative) composition laws are induced by m_2 .

We also note that DG categories can be identified with the A_∞ -categories with $m_n = 0$ for all $n \geq 3$; in particular, the composition operation m_2 is associative in this case. An A_∞ -category \mathcal{A} is *minimal* if $m_1 = 0$; in this case the composition operation is also associative.

Let \mathcal{A} and \mathcal{B} be A_∞ -categories. An A_∞ -functor $F: \mathcal{A} \rightarrow \mathcal{B}$ consists of the following data:

- A map $F_0: \text{ob}(\mathcal{A}) \rightarrow \text{ob}(\mathcal{B})$ between the classes of objects of \mathcal{A} and \mathcal{B} .
- Graded morphisms

$$F_n: \mathcal{A}(x_1, x_0) \otimes \mathcal{A}(x_2, x_1) \otimes \cdots \otimes \mathcal{A}(x_n, x_{n-1}) \longrightarrow \mathcal{B}(F_0(x_n), F_0(x_0)),$$

for $n \geq 1$, of degree $1 - n$.

Moreover, the above morphisms are required to satisfy a further infinite system of equations that imply that the morphisms

$$F_1: \mathcal{A}(x, y) \longrightarrow \mathcal{B}(F_0(x), F_0(y)), \quad x, y \in \mathcal{A},$$

are cochain maps and, moreover, F induces a graded functor

$$H^\bullet(F): H^\bullet(\mathcal{A}) \longrightarrow H^\bullet(\mathcal{B})$$

via the apparent formulas

$$x \longmapsto F_0(x), \quad x \in \text{ob}(\mathcal{A}), \quad \text{and} \quad \{f\} \longmapsto \{F_1(f)\}, \quad f \in Z(\mathcal{A}(x, y)),$$

and in particular an ordinary functor

$$H^0(F): H^0(\mathcal{A}) \longrightarrow H^0(\mathcal{B}).$$

An A_∞ -functor $F: \mathcal{A} \rightarrow \mathcal{B}$ is a *quasi-equivalence* if the induced graded functor

$$H^\bullet(F): H^\bullet(\mathcal{A}) \longrightarrow H^\bullet(\mathcal{B})$$

is an equivalence of graded categories.

Remark 4.3.1. There are different notions of ‘unitality’ for the composition operation in an A_∞ -category. First of all, the existence of units (identities) may not be required, in which case the corresponding A_∞ -categories are termed *non-unital*. Second, one may require that the graded category $H^\bullet(\mathcal{A})$ is unital, in which case one speaks of *cohomologically unital* A_∞ -categories. Finally, one may require that the (higher) composition operations m_n are *strictly unital* in a precise sense. All these variants of unitality extend to A_∞ -functors. We do not wish to elaborate on how the various notions of unitality relate to each other, but only remark that strict unitality implies cohomological unitality, and that every cohomologically unital A_∞ -category is quasi-equivalent to a strictly unital A_∞ -category, see for example [Sei08, Sec. I.2a]. The A_∞ -categories in this article are cohomologically unital.

Let \mathcal{A} be a DG category. By the Homotopy Transfer Theorem [Kad80], the graded category $H^\bullet(\mathcal{A})$ admits a (non-unique) minimal A_∞ -structure

$$(H^\bullet(\mathcal{A}), m_3, m_4, \dots)$$

whose composition operation m_2 is the composition law in the graded category $H^\bullet(\mathcal{A})$ (and is implicit in the notation), and such that there is a quasi-equivalence of A_∞ -categories

$$F: (H^\bullet(\mathcal{A}), m_3, m_4, \dots) \xrightarrow{\sim} \mathcal{A}$$

that is the identity on the objects: $F_0 = \mathbf{1}$. In this case, $(H^\bullet(\mathcal{A}), m_3, m_4, \dots)$ is called a *minimal A_∞ -model of \mathcal{A}* . Up to quasi-equivalence of A_∞ -categories, the DG category structure on \mathcal{A} is determined by the additional structure on its cohomology $H^\bullet(\mathcal{A})$. This is the main motivation for the use of A_∞ -categories in this article (see the proof of [Theorem 5.1.3](#) to see why this is important for us).

4.3.2. Massey products in DG categories with sparse cohomology. Let \mathcal{A} be a DG category. It was a longstanding belief that the higher Massey products in the graded category $H^\bullet(\mathcal{A})$ were related to the higher operations in a minimal model for \mathcal{A} via the formula

$$\pm m_n(f_1, \dots, f_n) \in \langle\langle f_1, \dots, f_n \rangle\rangle.$$

However, counter-examples were found recently by Buijs, Moreno-Fernández and Murillo in [\[BMFM20\]](#). There, the authors give sufficient conditions for the above formula to hold; however, these conditions are not suitable for our purposes. Below we show that the above relationship between Massey products and higher operations holds under a different sufficient condition that is better adapted to our framework.

The following definition is motivated by [Remark 3.2.3](#).

Definition 4.3.2. A graded category \mathcal{C} is *d-sparse* if

$$\forall x, y \in \mathcal{C}, \forall i \notin d\mathbb{Z}, \quad \mathcal{C}(x, y)^i = 0.$$

Similarly, a DG category \mathcal{A} has *d-sparse cohomology* or is *cohomologically d-sparse* if the graded category $H^\bullet(\mathcal{A})$ is *d-sparse*. Explicitly, \mathcal{A} is cohomologically *d-sparse* if

$$\forall x, y \in \mathcal{A}, \quad \forall i \notin d\mathbb{Z}, \quad H^i(\mathcal{A}(x, y)) = 0.$$

Remark 4.3.3. For $d = 1$, the sparseness condition on a (DG) category is, of course, vacuous.

Proposition 4.3.4. *Let \mathcal{A} be DG category with d-sparse cohomology, for some $d \geq 1$. Suppose given a sequence of composable morphisms in $H^\bullet(\mathcal{A})$ of the form*

$$x_{k+2} \xrightarrow{f_{k+2}} x_{k+1} \xrightarrow{f_{k+1}} \dots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0$$

with $k \notin d\mathbb{Z}$. Let $(H^\bullet(\mathcal{A}), m_3, m_4, \dots)$ be a minimal A_∞ -model of \mathcal{A} . Then,

$$m_{k+2}(f_1, f_2, \dots, f_{k+2}) = 0.$$

In other words, the higher operation m_{k+2} is identically 0.

Proof. By definition, m_{k+2} is a homogeneous morphism of degree $2 - (k + 2) = -k$. Since \mathcal{A} is cohomologically *d-sparse* by assumption and

$$|m_{k+2}(f_1, f_2, \dots, f_{k+2})| = -k + \sum_{i=1}^{k+2} |f_i|,$$

we must have $m_{k+2}(f_1, f_2, \dots, f_{k+2}) = 0$, because either some $|f_i| \notin d\mathbb{Z}$ and hence $f_i = 0$ or all $|f_i| \in d\mathbb{Z}$ and then $-k + \sum_{i=1}^{k+2} |f_i| \notin d\mathbb{Z}$. \square

Notation 4.3.5. Let $(\mathcal{A}, m_3, m_4, \dots)$ be a minimal A_∞ -category whose underlying graded category $\mathcal{A} = H^\bullet(\mathcal{A})$ is *d-sparse*. In view of [Proposition 4.3.4](#), we write

$$(\mathcal{A}, m_{d+2}, m_{2d+2}, \dots) = (\mathcal{A}, m_3, m_4, \dots)$$

since the higher operations m_{i+2} with $i \notin d\mathbb{Z}$ must vanish.

The following proposition shows that, in a DG category with *d-sparse* cohomology, the lowest non-trivial higher operation of any minimal A_∞ -model, that is m_{d+2} , determines the Massey products of the corresponding length.

Proposition 4.3.6. *Let \mathcal{A} be DG category with d -sparse cohomology, for some $d \geq 1$, and suppose given a sequence of composable morphisms in $H^\bullet(\mathcal{A})$ of the form*

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0$$

and such that $f_i f_{i+1} = 0$ for each $0 < i < d+2$. Then, for every minimal A_∞ -model $(H^\bullet(\mathcal{A}), m_{d+2}, m_{2d+2}, \dots)$ of \mathcal{A} , we have

$$-(-1)^{\sum_{0 < k \leq d+2} (d+2-k)|f_k|} m_{d+2}(f_1, \dots, f_{d+2}) \in \langle\langle f_1, \dots, f_{d+2} \rangle\rangle.$$

Proof. Let $(H^\bullet(\mathcal{A}), m_{d+2}, m_{2d+2}, \dots)$ be a minimal A_∞ -model of \mathcal{A} and choose a quasi-equivalence

$$F: (H^\bullet(\mathcal{A}), m_{d+2}, m_{2d+2}, \dots) \xrightarrow{\sim} \mathcal{A}$$

with $F_0 = \mathbf{1}$. For $0 \leq i < j \leq d+2$ and $j-i < d+2$, we define

$$g_{ij} := (-1)^{\sum_{i < k \leq j} (j-k)|f_k|} F_{j-i}(f_{i+1}, \dots, f_j).$$

We claim that the g_{ij} 's form a defining system for computing the Massey product $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$. Indeed, in view of the equations satisfied by the A_∞ -functor F [Kel01, Sec. 3.4], the morphism $d(F_{j-i}(f_{i+1}, \dots, f_j))$ equals

$$- \sum_{i < k < j} (-1)^{(k-i-1)+(j-k-1)\sum_{i < \ell \leq k} |f_\ell|} F_{k-i}(f_{i+1}, \dots, f_k) F_{j-k}(f_{k+1}, \dots, f_j)$$

and the claim follows by multiplying by the sign $(-1)^{\sum_{i < k \leq j} (j-k)|f_k|}$ in the definition of g_{ij} .

Finally, the morphism $d(F_{d+2}(f_1, \dots, f_{d+2}))$ equals

$$F_1(m_{d+2}(f_1, \dots, f_{d+2})) - \sum_{0 < k < d+2} (-1)^{(k-1)+(d+1-k)\sum_{0 < \ell \leq k} |f_\ell|} F_k(f_1, \dots, f_k) F_{d+2-k}(f_{k+1}, \dots, f_{d+2}).$$

From this, it follows that the morphism

$$-(-1)^{\sum_{0 < k \leq d+2} (d+2-k)|f_k|} m_{d+2}(f_1, \dots, f_{d+2})$$

represents the same cohomology class in $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$ as the one induced by the above defining system. \square

4.4. The universal Massey product. In this subsection we introduce the universal Massey product of length $d+2$ associated to a cohomologically d -sparse DG category (Definition 4.4.5).

4.4.1. *Reminder on Hochschild cohomology.* We refer the reader to [Mur22, Sec. 1] for more details on the material in this section. Let \mathcal{C} be a category and $\mathcal{C}^e := \mathcal{C} \otimes \mathcal{C}^{\text{op}}$ its enveloping category. As usual, we identify the abelian category of \mathcal{C} -bimodules with that of (right) \mathcal{C}^e -modules; in particular, the category \mathcal{C} yields the *diagonal \mathcal{C} -bimodule*

$$(x, y) \longmapsto \mathcal{C}(x, y), \quad x, y \in \mathcal{C},$$

that, with some abuse of notation, we also denote by \mathcal{C} . Recall that the *Hochschild cohomology of \mathcal{C} with coefficients in a \mathcal{C} -bimodule M* , also known as the *Hochschild–Mitchell cohomology* [Mit72], is the graded vector space

$$\text{HH}^\bullet(\mathcal{C}, M) := \text{Ext}_{\mathcal{C}^e}^\bullet(\mathcal{C}, M).$$

In particular, the Hochschild cohomology $\text{HH}^n(\mathcal{C}, M)$ vanishes if n is negative. If $M = \mathcal{C}$ the diagonal \mathcal{C} -bimodule, we write

$$\text{HH}^\bullet(\mathcal{C}) := \text{HH}^\bullet(\mathcal{C}, \mathcal{C}).$$

If \mathcal{C} is a graded category, then the category of \mathcal{C} -bimodules is also graded and the Hochschild cohomology is then a bi-graded vector space $\mathrm{HH}^{\bullet,*}(\mathcal{C}, M)$ and

$$\mathrm{HH}^p(\mathcal{C}, M) = \prod_{q \in \mathbb{Z}} \mathrm{HH}^{p+q}(\mathcal{C}, M), \quad p \geq 0.$$

Here, p is the *Hochschild (=horizontal) degree* and q is the *internal (=vertical) degree*. Hochschild cohomology is Morita invariant and suitably functorial with respect to graded functors in the first variable and bimodule morphisms in the second variable, see for example [Mur06].

The Hochschild cohomology of a graded category can be computed by means of the *bar complex*, which is the projective resolution of the diagonal bimodule whose component in degree n is

$$B_n(\mathcal{C}) := \bigoplus_{x_0, x_1, \dots, x_n} \mathcal{C}(x_0, -) \otimes \mathcal{C}(x_1, x_0) \otimes \mathcal{C}(x_2, x_1) \otimes \cdots \otimes \mathcal{C}(x_n, x_{n-1}) \otimes \mathcal{C}(-, x_n)$$

and whose differential $d: B_n(\mathcal{C}) \rightarrow B_{n-1}(\mathcal{C})$ and augmentation $d: B_0(\mathcal{C}) \rightarrow \mathcal{C}$ act on an elementary tensor by the formula

$$d(f_0 \otimes f_1 \otimes \cdots \otimes f_n \otimes f_{n+1}) = \sum_{i=0}^n (-1)^i f_0 \otimes f_1 \otimes \cdots \otimes f_i f_{i+1} \otimes \cdots \otimes f_n \otimes f_{n+1}$$

that uses the composition operation in \mathcal{C} . The *normalised bar complex* $\bar{B}_\bullet(\mathcal{C})$, consisting of replacing $\mathcal{C}(x, x)$ with $\mathcal{C}(x, x)/\mathbf{k} \cdot \mathbf{1}_x$ above whenever $x_{i+1} = x_i = x \in \mathcal{C}$, is also a projective resolution of the diagonal bimodule. The *Hochschild cochain complex* of \mathcal{C} is the the cochain complex

$$\mathbf{C}^{\bullet,*}(\mathcal{C}, M) := \mathrm{Hom}_{\mathcal{C}^e}^*(B_\bullet(\mathcal{C}), M).$$

Explicitly, in Hochschild degree n ,

$$\mathbf{C}^{n,*}(\mathcal{C}, M) = \prod_{x_0, x_1, \dots, x_n} \mathrm{Hom}_{\mathbf{k}}^*(\mathcal{C}(x_1, x_0) \otimes \mathcal{C}(x_2, x_1) \otimes \cdots \otimes \mathcal{C}(x_n, x_{n-1}), M(x_n, x_0))$$

and the differential $\varphi \mapsto d(\varphi)$, of bidegree $(1, 0)$, is given by the formula

$$\begin{aligned} d(\varphi)(f_1 \otimes \cdots \otimes f_{n+1}) &= (-1)^{|\varphi|_v |f_1|} f_1 \cdot \varphi(f_2 \otimes \cdots \otimes f_{n+1}) \\ &\quad + \sum_{i=1}^{n-1} (-1)^i \varphi(f_1 \otimes \cdots \otimes f_i f_{i+1} \otimes \cdots \otimes f_n \otimes f_{n+1}) \\ &\quad + (-1)^n \varphi(f_1 \otimes \cdots \otimes f_n) \cdot f_{n+1}, \end{aligned}$$

where $|\varphi|_v$ denotes the vertical degree. We can alternatively use the *normalised Hochschild cochain complex*,

$$\bar{\mathbf{C}}^{\bullet,*}(\mathcal{C}, M) := \mathrm{Hom}_{\mathcal{C}^e}^*(\bar{B}_\bullet(\mathcal{C}), M),$$

which is the subcomplex formed by the Hochschild cochains which vanish when at least one entry is an identity morphism in \mathcal{C} . If $M = \mathcal{C}$ the diagonal \mathcal{C} -bimodule, we also write

$$\mathbf{C}^\bullet(\mathcal{C}) := \mathbf{C}^\bullet(\mathcal{C}, \mathcal{C}).$$

Essentially by definition, the Hochschild cohomology

$$\mathrm{HH}^{0,*}(\mathcal{C}, M) = \mathrm{Hom}_{\mathcal{C}^e}^*(\mathcal{C}, M)$$

is isomorphic to the so-called *end* of the bimodule M , see [Kel05b, Sec. 2.1] for the definition. In particular,

$$\mathrm{HH}^{0,*}(\mathcal{C}) = \mathrm{Hom}_{\mathcal{C}^e}^*(\mathcal{C}, \mathcal{C}) \cong Z(\mathcal{C})$$

is isomorphic to the end of the diagonal bimodule, that is the *graded centre* of \mathcal{C} .

If \mathcal{C} is an ungraded category (=graded category concentrated in degree 0), then the Hochschild cohomology of \mathcal{C} with coefficients in a graded (!) \mathcal{C} -bimodule can be computed in terms of ungraded Hochschild cohomology by means of the formula

$$\mathrm{HH}^{p,q}(\mathcal{C}, M) \cong \mathrm{HH}^p(\mathcal{C}, M^q), \quad p \geq 0, \quad q \in \mathbb{Z}.$$

The Hochschild cohomology

$$\mathrm{HH}^{\bullet,*}(\mathcal{C}) = \mathrm{Ext}_{\mathcal{C}^e}^{\bullet,*}(\mathcal{C}, \mathcal{C})$$

of a graded category has the structure of a *Gerstenhaber algebra* [Ger63] and [Mur20a, Sec. 1]. Thus, $\mathrm{HH}^{\bullet,*}(\mathcal{C})$ is equipped with a graded commutative product

$$\mathrm{HH}^{p,q}(\mathcal{C}) \otimes \mathrm{HH}^{r,s}(\mathcal{C}) \longrightarrow \mathrm{HH}^{p+r,q+s}(\mathcal{C}), \quad a \otimes b \longmapsto a \cdot b,$$

called the *cup product*; up to sign, it agrees with the Yoneda product on the corresponding extension spaces. Thus, the cup product satisfies the identities

$$\begin{aligned} (a \cdot b) \cdot c &= a \cdot (b \cdot c), \\ a \cdot b &= (-1)^{|a||b|} b \cdot a. \end{aligned}$$

Here $|x| = p + q$ is the *total degree* of an element $x \in \mathrm{HH}^{p,q}(\mathcal{C})$. Moreover, there is a degree -1 Lie bracket

$$\mathrm{HH}^{p,q}(\mathcal{C}) \otimes \mathrm{HH}^{r,s}(\mathcal{C}) \longrightarrow \mathrm{HH}^{p+r-1,q+s}(\mathcal{C}), \quad a \otimes b \longmapsto [a, b],$$

that satisfies the following graded variants of the usual Lie algebra identities:

$$\begin{aligned} [a, a] &= 0, \quad |a| \text{ is odd}, \\ [a, b] &= -(-1)^{(|a|-1)(|b|-1)} [b, a], \\ [a, [b, c]] &= [[a, b], c] + (-1)^{(|a|-1)(|b|-1)} [b, [a, c]], \\ [a, [a, a]] &= 0, \quad |a| \text{ is even}. \end{aligned}$$

The cup product and the Lie bracket are compatible in the sense that

$$[a, b \cdot c] = [a, b] \cdot c + (-1)^{(|a|-1)|b|} b \cdot [a, c].$$

Finally, there is a *Gerstenhaber squaring operation* or *Gerstenhaber square*

$$\mathrm{HH}^{p,q}(\mathcal{C}) \longrightarrow \mathrm{HH}^{2p-1,2q}(\mathcal{C}), \quad a \longmapsto \mathrm{Sq}(a),$$

defined whenever $p + q$ is even or $\mathrm{char}(\mathbf{k}) = 2$. The Gerstenhaber square satisfies the following compatibility relations with respect to the cup product and the Lie bracket:

$$\begin{aligned} \mathrm{Sq}(a + b) &= \mathrm{Sq}(a) + \mathrm{Sq}(b) + [a, b], \\ \mathrm{Sq}(a \cdot b) &= \mathrm{Sq}(a) \cdot b^2 + a \cdot [a, b] \cdot b + a^2 \cdot \mathrm{Sq}(b), \\ [\mathrm{Sq}(a), b] &= [a, [a, b]]. \end{aligned}$$

If $\mathrm{char}(\mathbf{k}) \neq 2$, then the Gerstenhaber square is in fact determined by the Lie bracket according to the formula

$$\mathrm{Sq}(a) = \frac{1}{2}[a, a],$$

which follows from the previous relations. In particular, if $\mathrm{char}(\mathbf{k}) \neq 2$, then the relations that involve the Gerstenhaber square are redundant.

Remark 4.4.1. For a graded algebra A (that we can regard as a graded category with only one object), the automorphism group $\mathrm{Aut}(A)$ acts on the right of the cochain complex $\mathbf{C}^{\bullet,*}(A)$ by conjugation,

$$\varphi^f(x_1, \dots, x_n) = f^{-1}\varphi(f(x_1), \dots, f(x_n)), \quad \varphi \in \mathbf{C}^{n,*}(A), \quad f \in \mathrm{Aut}(A).$$

This induces a right action of $\text{Aut}(A)$ on $\text{HH}^{\bullet,*}(A)$ which is determined by

$$x^f = (f^{-1})_* f^*(x).$$

Here $f^*: \text{HH}^{\bullet,*}(A, A) \rightarrow \text{HH}^{\bullet,*}(A, {}_f A_f)$ is induced by the algebra automorphism f in the first variable and $f_*^{-1}: \text{HH}^{\bullet,*}(A, {}_f A_f) \rightarrow \text{HH}^{\bullet,*}(A, A)$ is induced by the A -bimodule morphism $f^{-1}: {}_f A_f \rightarrow A$ in the second variable.

The following result is probably well known but we have not found any reference.

Proposition 4.4.2. *The right action of $\text{Aut}(A)$ on $\text{HH}^{\bullet,*}(A)$ (Remark 4.4.1) factors through the outer automorphism group $\text{Out}(A)$.*

Proof. We have to show that the inner automorphism $f_u: A \xrightarrow{\sim} A$, $f_u(x) := (-1)^{|u||x|} u x u^{-1}$, induced by a unit $u \in A$ acts trivially on $\text{HH}^{\bullet,*}(A)$. The sign in the definition of the outer automorphism comes from the fact that $\mathbf{1}_A$ and f_u can be regarded as endofunctors of the graded category with one object A and multiplication by u can then be regarded as a degree $|u|$ natural isomorphism $u: \mathbf{1}_A \Rightarrow f_u$. We will produce a null-homotopy for the map

$$\begin{aligned} \mathbf{C}^{\bullet,*}(A) &\longrightarrow \mathbf{C}^{\bullet,*}(A), \\ \varphi &\longmapsto \varphi^{f_u}. \end{aligned}$$

The null-homotopy is the map

$$h: \mathbf{C}^{\bullet,*}(A) \longrightarrow \mathbf{C}^{\bullet-1,*}(A)$$

defined by

$$h(\varphi)(x_1, \dots, x_{n-1}) = \sum_{i=0}^{n-1} (-1)^{i+|u||\varphi|v} u^{-1} \varphi(u x_1 u^{-1}, \dots, u x_i u^{-1}, u, x_{i+1}, \dots, x_{n-1}).$$

A tedious but straightforward computation shows that

$$dh + hd = \mathbf{1}_{\mathbf{C}^{\bullet,*}(A)} - (-)^{f_u}. \quad \square$$

Remark 4.4.3. Proposition 4.4.2 can be generalized from algebras A to categories \mathcal{C} . In theory, the role of $\text{Aut}(A)$ should be replaced by the self-equivalences of \mathcal{C} , but these do not form a group (taking pseudo-inverses is not well defined). Nevertheless, $\text{Out}(A)$ can be replaced by the group of natural isomorphism classes of self-equivalences of \mathcal{C} , which happens to act on $\text{HH}^{\bullet,*}(\mathcal{C})$ by the same formulas as above. This can be checked by using the functoriality properties of Hochschild cohomology of categories, compare [Mur06].

4.4.2. *The universal Massey product of length $d+2$.* Let $(\mathcal{A}, m_3, m_4, \dots)$ be a minimal A_∞ -category. We are interested in the Hochschild cohomology of the graded category $\mathcal{A} = H^\bullet(\mathcal{A})$ (recall that $m_1 = 0$ by definition and m_2 is the composition law in \mathcal{A}). Firstly, notice that by definition the higher operations are Hochschild cochains:

$$m_n \in \mathbf{C}^{n, 2-n}(\mathcal{A}), \quad n \geq 3.$$

Similarly, if

$$F: (\mathcal{A}, m_3, m_4, \dots) \longrightarrow (\mathcal{B}, m_3, m_4, \dots)$$

is an A_∞ -functor between minimal A_∞ -categories, then F_0 and F_1 define a graded functor

$$F_{0,1}: \mathcal{A} \longrightarrow \mathcal{B}.$$

between the underlying graded categories, and the rest of components are Hochschild cochains:

$$F_n \in \mathbf{C}^{n, 1-n}(\mathcal{A}, \mathcal{B}(F_{0,1}, F_{0,1})), \quad n \geq 2.$$

The following result is a straightforward consequence of results of Lefèvre-Hasegawa [LH].

Proposition 4.4.4. *Let $(\mathcal{A}, m_{d+2}, m_{2d+2}, \dots)$ be a minimal A_∞ -category whose underlying graded category \mathcal{A} is d -sparse. The following statements hold:*

- (1) *The first non-trivial higher operation (see Proposition 4.3.4)*

$$m_{d+2} \in \mathbf{C}^{d+2, -d}(\mathcal{A})$$

is a Hochschild cocycle.

- (2) *Let $(\mathcal{A}, m'_{d+2}, m'_{2d+2}, \dots)$ be another minimal A_∞ -category with the same underlying graded category \mathcal{A} and*

$$F: (\mathcal{A}, m_{d+2}, m_{2d+2}, \dots) \longrightarrow (\mathcal{A}, m'_{d+2}, m'_{2d+2}, \dots)$$

an A_∞ -functor with $F_0 = \mathbf{1}$ and $F_1 = \mathbf{1}$. Then,

$$\{m_{d+2}\} = \{m'_{d+2}\} \quad \text{in} \quad \mathbf{HH}^{d+2, -d}(\mathcal{A}).$$

Proof. (1) Since \mathcal{A} is d -sparse, the higher operations m_{i+2} and m'_{i+2} vanish whenever $i \notin d\mathbb{Z}$ (Proposition 4.3.4). The claim follows from [LH, Lemma B.4.1], which shows that the differential $d(m_{d+2})$ of m_{d+2} in the Hochschild complex $\mathbf{C}^{\bullet, *}(A)$ equals an expression of the form

$$\sum \pm m_{i+2}(\dots),$$

where in particular $0 < i < d$, and hence the above expression must vanish.

(2) Similarly as in the previous statement, since \mathcal{A} is d -sparse, the higher components F_{i+1} of the A_∞ -functor F must vanish whenever $i \notin d\mathbb{Z}$ as, by definition, $|F_{i+1}| = -i$. The claim follows from [LH, Lemma B.4.2], which shows that the differential $d(F_{d+1})$ of F_{d+1} in the Hochschild complex $\mathbf{C}^{\bullet, *}(A)$ equals an expression of the form

$$m_{d+2} - m'_{d+2} + \sum \pm F_{i+1}(\dots) - \sum \pm m'_{i+2}(\dots),$$

where in particular $0 < i < d$, and hence the two right-most terms in the expression must vanish. This finishes the proof. \square

Definition 4.4.5. Let \mathcal{A} be a cohomologically d -sparse DG category and $H^\bullet(\mathcal{A})$ is graded cohomology category. The *universal Massey product of length $d+2$* is the Hochschild cohomology class

$$\{m_{d+2}\} \in \mathbf{HH}^{d+2, -d}(H^\bullet(\mathcal{A}))$$

represented by the operation m_{d+2} in any minimal A_∞ -model of \mathcal{A} .

Remark 4.4.6. Given a cohomologically d -sparse DG category, Proposition 4.4.4 shows that the universal Massey product of length $d+2$ is well defined and, moreover, is natural with respect to quasi-equivalences: if $F: \mathcal{A} \rightarrow \mathcal{B}$ is a quasi-equivalence between cohomologically d -sparse DG categories, then the induced isomorphism

$$\mathbf{HH}^{d+2, -d}(H^\bullet(\mathcal{A})) \cong \mathbf{HH}^{d+2, -d}(H^\bullet(\mathcal{B}))$$

takes the universal Massey product of length $d+2$ of \mathcal{A} to that of \mathcal{B} . The universal Massey product of length $d+2$ is also considered in [Sei15, Ch. 3]. For $d=1$, it has been investigated for example in [BKS04, Kad82] and plays a crucial role in [Mur22] as well.

Remark 4.4.7. The formula in Proposition 4.3.6 still holds if we replace m_{d+2} by any other Hochschild cochain m of bidegree $(d+2, -d)$ representing the universal

Massey product of length $d+2$ of \mathcal{A} , i.e. given a sequence of composable morphisms in $H^\bullet(\mathcal{A})$ of the form

$$x_{d+2} \xrightarrow{f_{d+2}} x_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} x_1 \xrightarrow{f_1} x_0$$

and such that $f_i f_{i+1} = 0$ for each $0 < i < d+2$, then

$$-(-1)^{\sum_{0 < k \leq d+2} (d+2-k)|f_k|} m(f_1, \dots, f_{d+2}) \in \langle\langle f_1, \dots, f_{d+2} \rangle\rangle.$$

This is straightforward to check, see [BKS04, Lem. 5.14] for the case $d = 1$. Compare also [BD89, Thm. 3.3] in topology.

Remark 4.4.8. Occasionally, we consider, more generally, the universal Massey product

$$\{m_{d+2}\} \in \mathrm{HH}^{d+2, -d}(\mathcal{A})$$

associated to a minimal A_∞ -category $\mathcal{A} = H^\bullet(\mathcal{A})$ whose underlying graded category is d -sparse (here, $\mathrm{HH}^{d+2, -d}(\mathcal{A})$ denotes the Hochschild cohomology of the underlying graded category of \mathcal{A}).

4.5. The restricted universal Massey product. In this subsection we relate universal Massey products to standard and Amiot–Lin $(d+2)$ -angulations ([Theorem 4.5.8](#)). The key ingredient for establishing the desired relationship is the restricted universal Massey product ([Definition 4.5.4](#)).

Setting 4.5.1. We fix a small DG category \mathcal{A} with the following properties:

- (1) The DG category \mathcal{A} is cohomologically d -sparse. We fix a minimal A_∞ -model

$$(H^\bullet(\mathcal{A}), m_{d+2}, m_{2d+2}, \dots)$$

of \mathcal{A} (see [Notation 4.3.5](#)).

- (2) The essential image of $H^0(\mathcal{A})$ under the canonical fully faithful functor

$$H^0(\mathbf{h}): H^0(\mathcal{A}) \longrightarrow \mathrm{D}^c(\mathcal{A})$$

is closed under the action of the d -fold shift and its inverse. In particular, there is a commutative diagram

$$\begin{array}{ccc} H^0(\mathcal{A}) & \longrightarrow & \mathrm{D}^c(\mathcal{A}) \\ [d] \downarrow & & \downarrow [d] \\ H^0(\mathcal{A}) & \longrightarrow & \mathrm{D}^c(\mathcal{A}) \end{array}$$

where $[d]: H^0(\mathcal{A}) \xrightarrow{\sim} H^0(\mathcal{A})$ is an autoequivalence such that

$$\forall x, y \in \mathcal{A}, \quad H^0(\mathcal{A}(x, y[d])) \cong H^0(\mathcal{A}(x, y)[d]) \cong H^d(\mathcal{A}(x, y))$$

as vector spaces (functorially on x and y).

- (3) The category $H^0(\mathcal{A})$ is additive and has split idempotents and finite-dimensional morphism spaces.
- (4) There exists a basic object $c \in H^0(\mathcal{A})$ such that $\mathrm{add}(c) = H^0(\mathcal{A})$. In particular, the Yoneda functor

$$H^0(\mathcal{A}) \longrightarrow \mathrm{mod}(\Lambda), \quad x \longmapsto H^0(\mathcal{A})(c, x)$$

restricts to an equivalence of categories

$$H^0(\mathcal{A}) \xrightarrow{\sim} \mathrm{proj}(\Lambda),$$

where $\Lambda := H^0(\mathcal{A}(c, c))$.

Notation 4.5.2. Since the finite-dimensional algebra Λ is basic, there exists an algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$ (unique up to inner automorphisms) such that the diagram

$$\begin{array}{ccc} H^0(\mathcal{A}) & \xrightarrow{\sim} & \text{proj}(\Lambda) \\ [d] \downarrow & & \downarrow -\otimes_{\Lambda} \sigma \Lambda_1 \\ H^0(\mathcal{A}) & \xrightarrow{\sim} & \text{proj}(\Lambda) \end{array}$$

commutes up to natural isomorphism. Explicitly, we fix an isomorphism $\varphi: c \xrightarrow{\sim} c[d]$ in $H^0(\mathcal{A})$ (which exists since c is a basic additive generator of $H^0(\mathcal{A})$, which is closed under $[d]$). The algebra automorphism

$$\sigma = \sigma_\varphi: H^0(\mathcal{A})(c, c) \xrightarrow{\sim} H^0(\mathcal{A})(c, c), \quad a \mapsto \varphi^{-1}(a[d])\varphi,$$

has the desired property. In particular, the autoequivalence

$$\Sigma := - \otimes_{\Lambda} \sigma \Lambda_1: \text{proj}(\Lambda) \xrightarrow{\sim} \text{proj}(\Lambda)$$

can be identified with the restriction of scalars $P \mapsto P_{\sigma^{-1}}$ along σ^{-1} .

4.5.1. *The d -sparse graded algebra $\Lambda(\sigma, d)$.* The isomorphism $\varphi: c \xrightarrow{\sim} c[d]$ ([Notation 4.5.2](#)) can be interpreted as an invertible cohomology class

$$\varphi \in H^0(\mathcal{A})(c, c[d]) \cong H^d(\mathcal{A}(c, c));$$

under this identification, the algebra automorphism $\sigma = \sigma_\varphi$ of Λ acts by conjugation $a \mapsto \varphi^{-1}a\varphi$. Notice also that φ induces isomorphisms of vector spaces

$$H^0(\mathcal{A}(c, c)) \xrightarrow{\sim} H^{di}(\mathcal{A}(c, c)), \quad b \mapsto \varphi^i b,$$

for each $i \in \mathbb{Z}$. Moreover, as a bimodule over $\Lambda = H^0(\mathcal{A}(c, c))$, the cohomology space $H^{di}(\mathcal{A}(c, c))$ identifies with the twisted Λ -bimodule ${}_{\sigma^i} \Lambda_1$. Indeed, for $a \in H^0(\mathcal{A}(c, c))$ and

$$b \in H^0(\mathcal{A}(c, c)) \xrightarrow{\sim} H^{di}(\mathcal{A}(c, c))$$

we have

$$b \cdot a = \varphi^{-i}((\varphi^i b)a) = ba \quad \text{and} \quad a \cdot b = \varphi^{-i}(a(\varphi^i b)) = \sigma^i(a)b.$$

More generally, we introduce the d -sparse graded algebra with underlying graded vector space

$$\Lambda(\sigma, d) := \bigoplus_{di \in d\mathbb{Z}} {}_{\sigma^i} \Lambda_1$$

and multiplication law

$$a \cdot b := \sigma^j(a)b \in \Lambda(\sigma, d)^{d(i+j)}, \quad a \in {}_{\sigma^i} \Lambda_1, \quad b \in {}_{\sigma^j} \Lambda_1.$$

It is straightforward to verify that the linear map

$$H^\bullet(\mathcal{A})(c, c) \xrightarrow{\sim} \Lambda(\sigma, d), \quad a \mapsto \varphi^{-i}a, \quad a \in H^{di}(\mathcal{A})(c, c)$$

is an isomorphism of graded algebras and, therefore, we obtain an induced isomorphism of Gerstenhaber algebras

$$\text{HH}^{\bullet,*}(H^\bullet(\mathcal{A})(c, c)) \xrightarrow{\sim} \text{HH}^{\bullet,*}(\Lambda(\sigma, d)),$$

which we treat as an identification in what follows. Finally, since the object $c \in H^0(\mathcal{A})$ is an additive generator there is also a canonical isomorphism of Gerstenhaber algebras

$$\text{HH}^{\bullet,*}(H^\bullet(\mathcal{A})(c, c)) \xrightarrow{\sim} \text{HH}^{\bullet,*}(H^\bullet(\mathcal{A}))$$

and, consequently, a further isomorphism of Gerstenhaber algebras

$$\text{HH}^{\bullet,*}(H^\bullet(\mathcal{A})) \xrightarrow{\sim} \text{HH}^{\bullet,*}(\Lambda(\sigma, d)).$$

Remark 4.5.3. There is a further isomorphism of (d -sparse) graded algebras

$$\Lambda(\sigma, d) \cong \frac{\Lambda\langle \iota^{\pm 1} \rangle}{(\iota a - \sigma(a)\iota)_{a \in \Lambda}}, \quad |\iota| = -d,$$

induced by the Λ -linear map

$$\Lambda\langle \iota^{\pm 1} \rangle \longrightarrow \Lambda(\sigma, d), \quad \iota \longmapsto 1 \in \Lambda(\sigma, d)^{-d} = {}_{\sigma^{-1}}\Lambda_1.$$

Indeed, for $a \in \Lambda = \Lambda(\sigma, d)^0$ and $1 \in \Lambda(\sigma, d)^{-d}$ we have

$$\sigma(a) \cdot 1 = \sigma^{-1}(\sigma(a))1 = a = 1a = \sigma^0(1)a = 1 \cdot a.$$

We treat this isomorphism as an identification simply for the purpose of removing the ambiguities that arise when considering homogeneous elements of the graded algebra $\Lambda(\sigma, d)$ (whose non-zero components are all equal to Λ as vector spaces).

4.5.2. The restricted universal Massey product. The inclusion $j: \Lambda \hookrightarrow \Lambda(\sigma, d)$ of the degree 0 part induces a restriction morphism

$$j^*: \mathrm{HH}^{\bullet,*}(\Lambda(\sigma, d), \Lambda(\sigma, d)) \longrightarrow \mathrm{HH}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$$

that, in bidegree $(d+2, -d)$, yields a morphism

$$j^*: \mathrm{HH}^{d+2,-d}(\Lambda(\sigma, d), \Lambda(\sigma, d)) \longrightarrow \mathrm{HH}^{d+2,-d}(\Lambda, \Lambda(\sigma, d)) \cong \mathrm{HH}^{d+2}(\Lambda, {}_1\Lambda_\sigma),$$

since

$$\Lambda(\sigma, d)^{-d} = {}_{\sigma^{-1}}\Lambda_1 \cong {}_1\Lambda_\sigma$$

is the degree $-d$ part of $\Lambda(\sigma, d)$ and

$$\mathrm{HH}^{d+2,-d}(\Lambda, \Lambda(\sigma, d)) = \mathrm{HH}^{d+2}(\Lambda, \Lambda(\sigma, d)^{-d}) = \mathrm{HH}^{d+2}(\Lambda, {}_{\sigma^{-1}}\Lambda_1);$$

notice also that the target of j^* is (isomorphic to) the extension space

$$\mathrm{HH}^{d+2}(\Lambda, {}_1\Lambda_\sigma) = \mathrm{Ext}_{\Lambda_e}^{d+2}(\Lambda, {}_1\Lambda_\sigma).$$

Using the graded Morita invariance of Hochschild cohomology, we may identify j^* with the restriction morphism

$$j^*: \mathrm{HH}^{d+2,-d}(H^\bullet(\mathcal{A}), H^\bullet(\mathcal{A})) \longrightarrow \mathrm{HH}^{d+2}(H^0(\mathcal{A}), H^{-d}(\mathcal{A})),$$

where $H^{-d}(\mathcal{A})$ is the ungraded $H^0(\mathcal{A})$ -bimodule

$$H^{-d}(\mathcal{A}): (x, y) \longmapsto H^{-d}(\mathcal{A}(x, y)).$$

Definition 4.5.4. The *restricted universal Massey product of length $d+2$* is the class

$$j^* \{m_{d+2}\} \in \mathrm{HH}^{d+2}(H^0(\mathcal{A}), H^{-d}(\mathcal{A})),$$

where

$$\{m_{d+2}\} \in \mathrm{HH}^{\bullet,*}(H^\bullet(\mathcal{A}), H^\bullet(\mathcal{A}))$$

is the universal Massey product of length $d+2$ associated to the minimal A_∞ -model

$$(H^\bullet(\mathcal{A}), m_{d+2}, m_{2d+2}, \dots)$$

of the d -sparse DG category \mathcal{A} .

Remark 4.5.5. In view of the isomorphism

$$\mathrm{HH}^{\bullet,*}(H^\bullet(\mathcal{A})) \xrightarrow{\sim} \mathrm{HH}^{\bullet,*}(\Lambda(\sigma, d)),$$

the universal Massey product $\{m_{d+2}\}$ can be identified with a class

$$\{m_{d+2}\} \in \mathrm{HH}^{\bullet,*}(\Lambda(\sigma, d), \Lambda(\sigma, d))$$

and therefore

$$j^* \{m_{d+2}\} \in \mathrm{HH}^{d+2,-d}(\Lambda, \Lambda(\sigma, d)) \cong \mathrm{Ext}_{\Lambda_e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$$

can be represented by an exact sequence of Λ -bimodules.

Remark 4.5.6. Occasionally, we consider, more generally, the restricted universal Massey product

$$j^* \{m_{d+2}\} \in \mathrm{HH}^{d+2, -d}(\mathcal{A}^0, \mathcal{A}) = \mathrm{HH}^{d+2}(\mathcal{A}^0, \mathcal{A}^{-d})$$

associated to a minimal A_∞ -category $\mathcal{A} = H^\bullet(\mathcal{A})$ whose underlying graded category is d -sparse, where \mathcal{A}^0 is the degree 0 part of \mathcal{A} (in particular, $\mathrm{HH}^{d+2, -d}(\mathcal{A}^0, \mathcal{A})$ denotes the Hochschild cohomology of the ungraded category \mathcal{A}^0 with coefficients in the graded \mathcal{A}^0 -bimodule \mathcal{A}).

The cohomology class investigated in [BD89] is a topological version of the restricted universal Massey products considered here in the case $d = 1$. It computes Toda brackets, which are the topological counterpart of Massey products. The connection is explained in [Mur20b].

4.5.3. Standard $(d+2)$ -angles versus Amiot–Lin $(d+2)$ -angulations.

Notation 4.5.7. Choose an exact sequence

$$\eta: \quad 0 \rightarrow {}_1\Lambda_\sigma \xrightarrow{l} P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

that represents $j^* \{m_{d+2}\}$ in $\mathrm{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$; we may and we will assume that P_0, P_1, \dots, P_d are projective Λ -bimodules. We let

$$\diamond_{j^* \{m_{d+2}\}} := \diamond_\eta$$

be the class of $(d+2)$ -angles in $\mathrm{proj}(\Lambda)$ obtained as in [Construction 2.2.13](#). Note, however, that we do not claim that $j^* \{m_{d+2}\}$ can be represented by an extension in which *all* middle terms are projective Λ -bimodules, and therefore *a priori* the class $\diamond_{j^* \{m_{d+2}\}}$ is not known to be a $(d+2)$ -angulation of $(\mathrm{proj}(\Lambda), \Sigma)$.

We aim to prove the following theorem. We remind the reader of our standing assumptions ([Setting 4.5.1](#)).

Theorem 4.5.8. *Suppose that the DG category \mathcal{A} is pre- $(d+2)$ -angulated. Then, the following statements hold:*

- (1) *The class of standard $(d+2)$ -angles on*

$$(\mathrm{proj}(\Lambda), \Sigma) \simeq (H^0(\mathcal{A}), [d])$$

coincides with the class $(-1)^{\sum_{i=1}^{d+2} i} \diamond_{j^ \{m_{d+2}\}}$ from [Notation 4.5.7](#) (see also [Remark 2.1.6](#)). In particular, the class $(-1)^{\sum_{i=1}^{d+2} i} \diamond_{j^* \{m_{d+2}\}}$ does not depend on the choice of an exact sequence representing $j^* \{m_{d+2}\}$ in $\mathrm{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$.*

- (2) *Suppose that Λ/J_Λ is separable. Let*

$$0 \rightarrow {}_1\Lambda_\sigma \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

be an exact sequence that represents $j^ \{m_{d+2}\}$ in $\mathrm{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$ such that the Λ -bimodules P_i , $0 \leq i < d+1$, are projective-injective Λ -bimodules. Then, P_{d+1} is also a projective-injective Λ -bimodule.*

In particular, if Λ/J_Λ is separable, the standard $(d+2)$ -angulation and the AL $(d+2)$ -angulation of $(\mathrm{proj}(\Lambda), \Sigma)$ coincide.

Notation 4.5.9. To alleviate the notation, we denote by $\mathcal{C} \subseteq \mathrm{D}^c(\mathcal{A})$ be the essential image of $H^0(\mathcal{A})$ under the canonical fully faithful functor

$$H^0(\mathbf{h}): H^0(\mathcal{A}) \longrightarrow \mathrm{D}^c(\mathcal{A}).$$

Under the induced equivalence of categories $H^0(\mathcal{A}) \simeq \mathcal{C}$, the $H^0(\mathcal{A})$ -bimodule

$$H^{di}(\mathcal{A}): (x, y) \longmapsto H^{di}(\mathcal{A}(x, y)) \cong H^0(\mathcal{A})(x, y[di]), \quad i \in \mathbb{Z},$$

identifies with the twisted \mathcal{C} -bimodule

$$[di]\mathcal{C}_1: (M, N) \mapsto \mathcal{C}(M, N[di]), \quad i \in \mathbb{Z}.$$

Similarly, the autoequivalence $[d]: H^0(\mathcal{A}) \xrightarrow{\sim} H^0(\mathcal{A})$ induces the autoequivalence

$$\Sigma := - \otimes_{\mathcal{C}} [d]\mathcal{C}_1: \text{Mod}(\mathcal{C}) \xrightarrow{\sim} \text{Mod}(\mathcal{C})$$

with quasi-inverse

$$\Sigma^{-1} := - \otimes_{\mathcal{C}} [-d]\mathcal{C}_1: \text{Mod}(\mathcal{C}) \xrightarrow{\sim} \text{Mod}(\mathcal{C}).$$

Working towards the proof of [Theorem 4.5.8](#), we first investigate the relationship between the restricted universal Massey product and extensions. The reader might find it useful to compare the discussion below with [Construction 2.2.13](#). We begin with a general observation. Let \mathcal{X} be a category and consider an exact sequence of \mathcal{X} -bimodules of the form

$$0 \rightarrow M \rightarrow E_{d+1} \rightarrow \cdots \rightarrow E_1 \rightarrow E_0 \rightarrow \mathcal{X} \rightarrow 0,$$

where E_0, E_1, \dots, E_d are projective \mathcal{X} -bimodules. Notice that for each $x \in \mathcal{X}$ the sequence of *left* \mathcal{X} -modules

$$0 \rightarrow M(x, -) \rightarrow E_{d+1}(x, -) \rightarrow \cdots \rightarrow E_1(x, -) \rightarrow E_0(x, -) \rightarrow \mathcal{X}(x, -) \rightarrow 0$$

is split-exact (=contractible as a complex), for the representable left \mathcal{X} -module $\mathcal{X}(x, -)$ as well as $E_i(x, -)$, $i = 0, 1, \dots, d$, are projective as left \mathcal{X} -modules. From this follows that, for every (right) \mathcal{X} -module N , the sequence of (right) \mathcal{X} -modules

$$N \otimes_{\mathcal{X}} (0 \rightarrow M \rightarrow E_{d+1} \rightarrow \cdots \rightarrow E_1 \rightarrow E_0 \rightarrow \mathcal{X} \rightarrow 0)$$

is exact. Thus, in view of the canonical isomorphism $N \otimes_{\mathcal{X}} \mathcal{X} \cong N$, there is a well-defined morphism

$$\Phi_N: \text{Ext}_{\mathcal{X}^e}^{\bullet}(\mathcal{X}, M) \longrightarrow \text{Ext}_{\mathcal{X}}^{\bullet}(N, N \otimes_{\mathcal{X}} M),$$

which is an algebra morphism with respect to the Yoneda product if $M = \mathcal{X}$ the diagonal \mathcal{X} -bimodule.

Returning to the context of [Setting 4.5.1](#), if we let $\mathcal{X} = \mathcal{C}$ and $M = [q]\mathcal{C}_1$ in the previous discussion, for each \mathcal{C} -module N we obtain a morphism

$$\Phi_N: \text{Ext}_{\mathcal{C}^e}^{d+2}(\mathcal{C}, [q]\mathcal{C}_1) \longrightarrow \text{Ext}_{\mathcal{C}}^{d+2}(N, N \otimes_{\mathcal{C}} [q]\mathcal{C}_1).$$

In particular, if $q = -d$, we may consider the class

$$\Phi_N(j^* \{m_{d+2}\}) \in \text{Ext}_{\mathcal{C}}^{d+2}(N, N \otimes_{\mathcal{C}} [-d]\mathcal{C}_1) \cong \text{Ext}_{\mathcal{C}}^{d+2}(N, \Sigma^{-1}N)$$

induced by the restricted universal Massey product of length $d+2$. Thus, given an exact sequence of \mathcal{C} -bimodules

$$\eta: \quad 0 \rightarrow [-d]\mathcal{C}_1 \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \mathcal{C} \rightarrow 0$$

that represents the restricted universal Massey product $j^* \{m_{d+2}\}$, we see that the class $\Phi_N(j^* \{m_{d+2}\})$ is represented by the exact sequence

$$N \otimes_{\mathcal{C}} (0 \rightarrow [-d]\mathcal{C}_1 \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \mathcal{C} \rightarrow 0).$$

Suppose given a sequence of composable morphisms in \mathcal{C} of the form

$$M_{d+2} \xrightarrow{f_{d+2}} M_{f+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} M_1 \xrightarrow{f_1} M_0$$

and such that $f_i f_{i+1} = 0$ for all $0 \leq i < d+2$, so that the Massey product $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$ in $H^{\bullet}(\mathcal{A})$ is non-empty (see [Remark 4.2.4](#)). Let

$$N := \text{coker } \mathcal{C}(-, f_1): \mathcal{C}^{\text{op}} \longrightarrow \text{mod } \mathbf{k}.$$

In view of [Proposition 4.1.31](#) and [Theorem 4.2.6](#), the Massey product is an element of the vector space $\text{Massey}(f_{d+2}, f_1)$ defined as the quotient

$$\frac{\text{Hom}_{\mathbb{D}^c(\mathcal{A})}(M_{d+2}, M_0[-d])}{f_1[-d] \cdot \text{Hom}_{\mathbb{D}^c(\mathcal{A})}(M_{d+2}, M_1[-d]) + \text{Hom}_{\mathbb{D}^c(\mathcal{A})}(M_{d+1}, M_0) \cdot f_{d+2}}.$$

We wish to construct a further morphism

$$\Psi: \text{Ext}_{\mathbb{C}}^{d+2}(N, \Sigma^{-1}N) \longrightarrow \text{Massey}(f_{d+2}, f_1).$$

For this, let $B_{\bullet}(\mathbb{C})$ be the bar complex of \mathbb{C} , so that $N \otimes_{\mathbb{C}} B_{\bullet}(\mathbb{C})$ is a projective resolution of N that we use to compute the extension space $\text{Ext}_{\mathbb{C}}^{d+2}(N, \Sigma^{-1}N)$ as

$$\text{Ext}_{\mathbb{C}}^{d+2}(N, \Sigma^{-1}N) = H^{d+2}(\text{Hom}_{\mathbb{C}}(N \otimes_{\mathbb{C}} B_{\bullet}(\mathbb{C}), \Sigma^{-1}N)).$$

Following essentially the same arguments as in [\[Mur20b, Sec. 5\]](#), one can check that under the above isomorphism the class $\Phi_N(j^* \{m_{d+2}\})$ is represented by the class of the morphism

$$N \otimes_{\mathbb{C}} B_{d+2}(\mathbb{C}) \longrightarrow \Sigma^{-1}N$$

that corresponds, under the Yoneda embedding, to the morphism

$$\bigoplus_{L_0, L_1, \dots, L_{d+2}} N(L_0) \otimes \mathbb{C}(L_1, L_0) \otimes \cdots \otimes \mathbb{C}(L_{d+2}, L_{d+1}) \longrightarrow \Sigma^{-1}N(L_{d+2})$$

given by

$$(4.5.10) \quad [h] \otimes g_1 \otimes \cdots \otimes g_{d+2} \longmapsto [(h[-d]) \circ m_{d+2}(g_1, \dots, g_{d+2})],$$

where we use that

$$m_{d+2}(g_1, \dots, g_{d+2}) \in \text{Hom}_{\mathbb{D}^c(\mathcal{A})}(L_{d+2}, L_0[-d])$$

and therefore the composite

$$L_{d+2} \xrightarrow{m_{d+2}(g_1, \dots, g_{d+2})} L_0[-d] \xrightarrow{h[-d]} N[-d] = \Sigma^{-1}N$$

indeed represents an element in $\Sigma^{-1}N(L_{d+2})$. We refer the reader to the last paragraph of the proof of [\[Mur20b, Prop. 5.2\]](#) as well as [\[Mur20b, Prop. 5.6 and Rmks. 5.4 and 5.7\]](#) for details.

Continuing with the construction of the morphism Ψ , consider the morphism of augmented chain complexes

$$\begin{array}{ccccccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & \mathbb{C}(-, M_{d+2}) & \longrightarrow & \cdots & \longrightarrow & \mathbb{C}(-, M_1) & \longrightarrow & \mathbb{C}(-, M_0) & \xrightarrow{p} & N \\ & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \parallel \\ \cdots & \rightarrow & N \otimes_{\mathbb{C}} B_{d+3}(\mathbb{C}) & \rightarrow & N \otimes_{\mathbb{C}} B_{d+2}(\mathbb{C}) & \rightarrow & \cdots & \rightarrow & N \otimes_{\mathbb{C}} B_1(\mathbb{C}) & \rightarrow & N \otimes_{\mathbb{C}} B_0(\mathbb{C}) & \rightarrow & N. \end{array}$$

that corresponds, under the Yoneda embedding, to the elements

$$(4.5.11) \quad (-1)^{\sum_{j=1}^i j} p \otimes f_1 \otimes \cdots \otimes f_i \otimes \mathbf{1}_{M_i} \in (N \otimes_{\mathbb{C}} B_i(\mathbb{C}))(M_i),$$

compare with the proof of [\[Mur20b, Prop. 5.9\]](#). Here p is the natural projection onto the cokernel. The morphism Ψ is obtained by applying the functor $\text{Hom}_{\mathbb{C}}(-, \Sigma^{-1}N)$ to the above morphism of chain complexes and passing to cohomology in degree $d+2$, while keeping in mind the isomorphisms

$$\text{Hom}_{\mathbb{C}}(\mathbb{C}(-, M_{d+2}), \Sigma^{-1}N) \cong \Sigma^{-1}N(M_{d+2}) \cong \frac{\mathbb{C}(M_{d+2}, M_0[-d])}{(f_1[-d]) \cdot \mathbb{C}(M_{d+2}, M_1[-d])}.$$

As a consequence of the above discussion, we obtain the following relationship between the restricted universal Massey product $j^* \{m_{d+2}\}$ and the Massey product $\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$.

Proposition 4.5.12. *Suppose given a sequence of composable morphisms in \mathcal{C} of the form*

$$M_{d+2} \xrightarrow{f_{d+2}} M_{f+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} M_1 \xrightarrow{f_1} M_0$$

and such that $f_i f_{i+1} = 0$ for all $0 \leq i < d+2$. Set $N = \text{coker } \mathcal{C}(-, f_1)$. Then,

$$(\Psi \circ \Phi_N)(j^* \{m_{d+2}\}) = -(-1)^{\sum_{i=1}^{d+2} i} \langle\langle f_1, \dots, f_{d+2} \rangle\rangle$$

in Massey (f_{d+2}, f_1) .

Proof. In view of (4.5.10) and (4.5.11),

$$(\Psi \circ \Phi_N)(j^* \{m_{d+2}\}) = (-1)^{\sum_{i=1}^{d+2} i} \{m_{d+2}(f_1, \dots, f_{d+2})\} \in \text{Massey}(f_{d+2}, f_1).$$

The claim then follows from Proposition 4.3.6, which in this case says that

$$-m_{d+2}(f_1, \dots, f_{d+2}) \in \langle\langle f_1, \dots, f_{d+2} \rangle\rangle. \quad \square$$

Remark 4.5.13. In view of Theorem 4.2.6, under the assumptions of Proposition 4.5.12,

$$(\Psi \circ \Phi_N)(j^* \{m_{d+2}\})[d] = (-1)^{d+1} \langle f_1, \dots, f_{d+2} \rangle \in \text{Toda}(f_{d+2}, f_1).$$

The following proposition shows that the standard $(d+2)$ -angles in \mathcal{C} are detected by the restricted universal Massey product of length $d+2$. In the statement we do not assume \mathcal{A} to be pre- $(d+2)$ -angulated.

Proposition 4.5.14. *Let $(H^\bullet(\mathcal{A}), m_{d+2}, m_{2d+2}, \dots)$ be a minimal A_∞ -model of \mathcal{A} . Suppose given a sequence of composable morphisms*

$$(4.5.15) \quad M_{d+2} \xrightarrow{f_{d+2}} M_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} M_1 \xrightarrow{f_1} \Sigma M_{d+2}.$$

in $(\mathcal{C}, \Sigma) \simeq (H^0(\mathcal{A}), [d])$. Then, the following statements are equivalent:

- (1) *The above sequence satisfies the conditions in Theorem 4.1.12.*
- (2) *The above sequence lies in $(-1)^{\sum_{i=1}^{d+2} i} \langle j^* \{m_{d+2}\} \rangle$ (see Notation 4.5.7 and Remark 2.2.14).*

Proof. For simplicity, we identify (4.5.15) with a sequence of projective Λ -modules. In view of Theorem 4.2.6, statement (1) is equivalent to the following two conditions:

- The sequence

$$M_{d+2} \xrightarrow{f_{d+2}} M_{d+1} \xrightarrow{f_{d+1}} \cdots \xrightarrow{f_2} M_1 \xrightarrow{f_1} \Sigma M_{d+2} \xrightarrow{\Sigma f_{d+2}} \Sigma M_{d+1}$$

is exact.

- We have $-\mathbf{1}_{M_{d+2}} \in \langle\langle f_1, \dots, f_{d+2} \rangle\rangle$.

The first condition is precisely the first condition that a sequence must satisfy to be a member of the class $(-1)^{\sum_{i=1}^{d+2} i} \langle j^* \{m_{d+2}\} \rangle$.

On the other hand, the second (and last) condition that a sequence must satisfy to be a member of the class $(-1)^{\sum_{i=1}^{d+2} i} \langle j^* \{m_{d+2}\} \rangle$ says the following: Let $N = \text{coker } f_1$, so that there is an exact sequence

$$0 \rightarrow \Sigma^{-1} N \xrightarrow{i} M_{d+1} \rightarrow \cdots \rightarrow M_1 \xrightarrow{f_1} \Sigma M_{d+2} \xrightarrow{p} N \rightarrow 0$$

with $f_{d+2} = i \circ \Sigma^{-1} p$. Then, the above sequence must represent the class

$$\Phi_N((-1)^{\sum_{i=1}^{d+2} i} j^* \{m_{d+2}\}) \in \text{Ext}_{\Lambda^e}^{d+2}(N, \Sigma^{-1} N)$$

that, by definition, is represented by the exact sequence

$$N \otimes_{\Lambda} (0 \rightarrow {}_1\Lambda_{\sigma} \xrightarrow{(-1)^{\sum_{i=1}^{d+2} i} \iota} P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0)$$

(the Λ -bimodule extension on the right of the tensor product represents the class $(-1)^{\sum_{i=1}^{d+2} i} j^* \{m_{d+2}\}$, see [Notation 4.5.7](#) and [Remark 2.2.14](#)). The extension space can be computed by means of the projective resolution

$$\cdots \rightarrow \Sigma^{-1}M_1 \xrightarrow{\Sigma^{-1}f_1} M_{d+2} \rightarrow M_{d+1} \rightarrow \cdots \rightarrow M_1 \xrightarrow{f_1} \Sigma M_{d+2} \xrightarrow{p} N \rightarrow 0$$

and, in these terms, we see that the former sequence is represented by the class

$$[\Sigma^{-1}p] \in \frac{\ker(? \circ \Sigma^{-1}f_1)}{\mathrm{Hom}_\Lambda(M_{d+1}, \Sigma^{-1}N) \cdot f_{d+2}} \cong \mathrm{Ext}_\Lambda^{d+2}(N, \Sigma^{-1}N).$$

Here

$$? \circ \Sigma^{-1}f_1: \mathrm{Hom}_\Lambda(M_{d+2}, \Sigma^{-1}N) \longrightarrow \mathrm{Hom}_\Lambda(\Sigma^{-1}M_1, \Sigma^{-1}N)$$

is the morphism given by pre-composition with $\Sigma^{-1}f_1$. Thus, to summarise, the second condition for membership in the class $(-1)^{\sum_{i=1}^{d+2} i} \diamond_{j^*} \{m_{d+2}\}$ says that

$$[\Sigma^{-1}p] = \Phi_N((-1)^{\sum_{i=1}^{d+2} i} j^* \{m_{d+2}\})$$

in $\mathrm{Ext}_{\Lambda^e}^{d+2}(N, \Sigma^{-1}N)$.

Now, the morphism Ψ is injective by construction and identifies the extension space $\mathrm{Ext}_{\Lambda^e}^{d+2}(N, \Sigma^{-1}N)$ with the subspace

$$\frac{\ker(? \circ \Sigma^{-1}f_1)}{\mathrm{Hom}_\Lambda(M_{d+1}, \Sigma^{-1}N) \cdot f_{d+2}} \subseteq \frac{\mathrm{Hom}_\Lambda(M_{d+2}, \Sigma^{-1}N)}{\mathrm{Hom}_\Lambda(M_{d+1}, \Sigma^{-1}N) \cdot f_{d+2}} \cong \mathrm{Massey}(f_{d+2}, f_1).$$

Thus, to conclude the proof it is enough to prove that

$$\Psi(\Phi_N((-1)^{\sum_{i=1}^{d+2} i} j^* \{m_{d+2}\})) = \Psi([\Sigma^{-1}p])$$

if and only if $-\mathbf{1}_{M_{d+2}} \in \langle\langle f_1, \dots, f_{d+2} \rangle\rangle$. Indeed,

$$\Psi: [\Sigma^{-1}p] \mapsto [\mathbf{1}_{M_{d+2}}] \in \mathrm{Massey}(f_{d+2}, f_1).$$

and

$$\Psi: \Phi_N((-1)^{\sum_{i=1}^{d+2} i} j^* \{m_{d+2}\}) \mapsto -\langle\langle f_1, \dots, f_{d+2} \rangle\rangle$$

by [Proposition 4.5.12](#). The claim follows. \square

The graded algebra structure on $\Lambda(\sigma, d)$ endows the Hochschild cohomology

$$\mathrm{HH}^{\bullet,*}(\Lambda, \Lambda(\sigma, d)) = \mathrm{Ext}_{\Lambda^e}^{\bullet}(\Lambda, \Lambda(\sigma, d))$$

with the structure of a bigraded algebra, where the internal grading is induced by the grading of $\Lambda(\sigma, d)$. In particular,

$$\mathrm{HH}^{\bullet,0}(\Lambda, \Lambda(\sigma, d)) = \mathrm{Ext}_{\Lambda^e}^{\bullet}(\Lambda, \Lambda(\sigma, d)^0) = \mathrm{Ext}_{\Lambda^e}^{\bullet}(\Lambda, \Lambda)$$

is the Yoneda algebra of the diagonal Λ -bimodule. Similarly, since Λ^e is also self-injective, we may consider the *Hochschild–Tate cohomology*

$$\underline{\mathrm{HH}}^{\bullet,*}(\Lambda, \Lambda(\sigma, d)) := \underline{\mathrm{Ext}}_{\Lambda^e}^{\bullet,*}(\Lambda, \Lambda(\sigma, d)),$$

which is again a bigraded algebra, and note that there are isomorphisms

$$\underline{\mathrm{HH}}^{p,q}(\Lambda, \Lambda(\sigma, d)) \cong \mathrm{HH}^{p,q}(\Lambda, \Lambda(\sigma, d)) = \mathrm{Ext}_{\Lambda^e}^p(\Lambda, \sigma^q \Lambda_1), \quad p > 0, \quad q \in \mathbb{Z}.$$

We refer the reader to [[Mur22](#), Sec. 5] for details.

Remark 4.5.16. An extension

$$0 \rightarrow \sigma \Lambda_1 \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

of Λ -bimodules such that P_i , $0 \leq i < d+1$, is a projective-injective Λ -bimodule represents a unit in the Hochschild–Tate cohomology $\underline{\mathrm{HH}}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$ if and only if P_{d+1} is also a projective-injective Λ -bimodule, see [[Mur22](#), Rmk. 5.9].

We are finally ready to prove [Theorem 4.5.8](#).

Proof of Theorem 4.5.8. Statement (1) follows from Proposition 4.5.14. We prove statement (2). Choose an exact sequence

$$\eta: \quad 0 \rightarrow {}_1\Lambda_\sigma \xrightarrow{\iota} P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

that represents the restricted universal Massey product

$$j^* \{m_{d+2}\} \in \text{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$$

and such that P_i , $0 \leq i < d+1$, is a projective-injective Λ -bimodule (such a representative always exists). We need to prove that the Λ -bimodule P_{d+1} is also projective-injective. It is enough to prove that $N \otimes P_{d+1}$ is projective for every finite-dimensional Λ -module N , see [AR91, Thm. 3.1] and notice that the proof remains valid under the assumption of Λ/J_Λ being separable, since in this case the finite-dimensional algebra $(\Lambda/J_\Lambda) \otimes (\Lambda/J_\Lambda)^{\text{op}}$ is semisimple [ERZ57, Cor. 18].

Thus, let N be a finite-dimensional Λ -module and choose a projective presentation

$$Q_1 \xrightarrow{f_1} \Sigma Q_{d+2} \rightarrow N \rightarrow 0.$$

Using axioms (*d-TR1*) and (*d-TR2*), complete the morphism f_1 to a standard $(d+2)$ -angle

$$Q_{d+2} \rightarrow Q_{d+1} \rightarrow \cdots \rightarrow Q_1 \xrightarrow{f_1} \Sigma Q_{d+2}.$$

In view of the validity of statement (1), the above $(d+2)$ -angle lies in the class $\pm \diamond_{j^* \{m_{d+2}\}}$. In particular, the exact sequence

$$N \otimes_\Lambda (0 \rightarrow {}_1\Lambda_\sigma \xrightarrow{\pm \iota} P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0)$$

is equivalent to an exact sequence of the form

$$0 \rightarrow \Sigma^{-1}N \rightarrow Q_{d+1} \rightarrow \cdots \rightarrow Q_1 \rightarrow \Sigma Q_{d+2} \rightarrow N \rightarrow 0.$$

Notice that all of the middle terms in the latter exact sequence are projective-injective Λ -bimodules and, therefore, the class of this sequence in the Hochschild–Tate cohomology $\underline{\text{HH}}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$ is a unit. Consequently, the former (equivalent) sequence also represents a unit in $\underline{\text{HH}}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$, and hence the Λ -module $N \otimes_\Lambda P_{d+1}$ must be projective-injective, which is what we needed to prove. This finishes the proof of the theorem. \square

Theorem 4.5.8 and Proposition 4.5.14 yield the following important corollaries.

Corollary 4.5.17. *Let \mathcal{A} be a small DG category satisfying the assumptions in Setting 4.5.1. Consider the following statements:*

- (1) *The DG category \mathcal{A} is pre- $(d+2)$ -angulated.*
- (2) *The canonical fully faithful functor*

$$H^0(\mathbf{h}): H^0(\mathcal{A}) \longrightarrow \text{D}^c(\mathcal{A})$$

identifies $H^0(\mathcal{A})$ with a $d\mathbb{Z}$ -cluster tilting subcategory of $\text{D}^c(\mathcal{A})$.

- (3) *The restricted universal Massey product*

$$j^* \{m_{d+2}\} \in \text{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$$

is represented by an extension all of whose middle terms are projective-injective Λ -bimodules.

In general (1) \Leftrightarrow (2) and (1) \Leftarrow (3). If Λ/J_Λ is separable, then (1) \Rightarrow (3).

Proof. The equivalence (1) \Leftrightarrow (2) is Proposition 3.2.4.

(1) \Rightarrow (3) This is precisely Theorem 4.5.8(2), which requires the algebra Λ/J_Λ to be separable.

(3) \Rightarrow (1) By Proposition 4.5.14 and Remark 2.2.14, the Amiot–Lin $(d+2)$ -angulation associated to a representative of $(-1)^{\sum_{i=1}^{d+2} i} j^* \{m_{d+2}\}$ coincides with the

class of sequences that satisfy the conditions in [Theorem 4.1.12](#), which therefore forms a $(d + 2)$ -angulation. Consequently, the canonical fully faithful functor

$$H^0(\mathbf{h}): H^0(\mathcal{A}) \longrightarrow \mathcal{D}^c(\mathcal{A})$$

identifies the left-hand side with a standard $(d + 2)$ -angulated subcategory of the right-hand side, which is precisely the definition of \mathcal{A} being pre- $(d + 2)$ -angulated. \square

Remark 4.5.18. Suppose for simplicity that the field k is perfect. In the context of [Corollary 4.5.17](#), it follows from [[Dug12](#), Thm. 3.2] that there is a stable bimodule isomorphism $\Omega_{\Lambda^e}^{d+2}(\Lambda) \cong {}_1\Lambda_\sigma$. An important aspect of [Corollary 4.5.17](#) is that this stable bimodule isomorphism is witnessed by the restricted universal Massey product of \mathcal{A} .

Recall that the Nakayama automorphism η of a Frobenius algebra Λ is characterised (up to inner automorphisms) by the existence of an isomorphism of Λ -bimodules $D\Lambda \cong {}_1\Lambda_\eta$.

Corollary 4.5.19 ([[Dug12](#), Prop. 3.3], [[GKO13](#), Sec. 5.4], [[IO13](#), Cor. 4.6]). *Let \mathcal{A} be a small DG category satisfying the assumptions in [Setting 4.5.1](#) and such that the canonical fully faithful functor*

$$H^0(\mathbf{h}): H^0(\mathcal{A}) \longrightarrow \mathcal{D}^c(\mathcal{A})$$

identifies $H^0(\mathcal{A})$ with a $d\mathbb{Z}$ -cluster tilting subcategory of $\mathcal{D}^c(\mathcal{A})$. If $H^0(\mathcal{A})$ is ds -Calabi–Yau, $\ell \in \mathbb{Z}$, in the sense that there are functorial isomorphisms

$$H^0(\mathcal{A})(y, x[d\ell]) \xrightarrow{\sim} DH^0(\mathcal{A})(x, y), \quad x, y \in H^0(\mathcal{A}),$$

then $\sigma^{-\ell}$ is a Nakayama automorphism for Λ . In particular, the stable module category $\underline{\text{mod}} \Lambda$ is $((d + 2)\ell - 1)$ -Calabi–Yau.

Proof. The argument is standard; we reproduce it for the convenience of the reader. The functorial isomorphism

$$H^0(\mathcal{A})(c, c[d\ell]) \xrightarrow{\sim} DH^0(\mathcal{A})(c, c)$$

can be interpreted as an isomorphism of Λ -bimodules ${}_1\Lambda_{\sigma^{-\ell}} \cong D\Lambda$ that exhibits $\sigma^{-\ell}$ as a Nakayama automorphism for Λ . This proves the first claim. To prove the second claim, observe that [Corollary 4.5.17](#) yields the existence of an isomorphism

$$\Omega_{\Lambda^e}^{d+2}(\Lambda) \cong {}_1\Lambda_\sigma,$$

in $\underline{\text{mod}} \Lambda^e$ that in turn induces an isomorphism of functors

$$\Omega_\Lambda^{(d+2)\ell} \cong \nu^{-1}$$

on $\underline{\text{mod}} \Lambda$, where $\nu = - \otimes_\Lambda D\Lambda$ is the Nakayama functor. The claim follows from the fact that $\nu\Omega_\Lambda$ is a Serre functor on $\underline{\text{mod}} \Lambda$, see [[ES06](#), Prop. 1.2 and Cor. 1.3] and compare also with [[IV14](#), Thm. 1.8]. This finishes the proof. \square

Corollary 4.5.20. *Let Λ be a finite-dimensional self-injective algebra that is twisted $(d + 2)$ -periodic with respect to an algebra automorphism σ and let*

$$\Sigma := - \otimes_\Lambda \sigma \Lambda_1: \text{proj}(\Lambda) \xrightarrow{\sim} \text{proj}(\Lambda).$$

If Λ/J_Λ is separable, then, up to equivalence, the pair $(\text{proj}(\Lambda), \Sigma)$ admits at most one algebraic $(d + 2)$ -angulation.

Proof. Indeed, [Theorem 4.5.8](#) shows that any two algebraic $(d + 2)$ -angulations of $(\text{proj}(\Lambda), \Sigma)$ coincide with the AL $(d + 2)$ -angulations given by any choice of exact sequences with projective-middle terms in $\text{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$ that represent the corresponding restricted universal Massey products of the enhancements, up to a sign depending on d , see also [Remark 2.2.14](#). But such AL $(d + 2)$ -angulations are unique up to equivalence by [Proposition 2.2.18](#). \square

Remark 4.5.21. Notice that [Corollary 4.5.20](#) does not prove that any two enhancements are equivalent, provided at least one exists. This is part of the content of [Theorem A](#).

4.6. Example of a universal Massey product. In this subsection we compute the universal Massey product of a specific Karoubian pre- $(d + 2)$ -angulated DG category \mathcal{A} or, rather, the first non-zero higher operation in a minimal A_∞ -model of \mathcal{A} . Let $A = A_3$ be the path algebra of the quiver $1 \rightarrow 2 \rightarrow 3$ and consider its compact derived category $D^c(A)$. It is well known and easy to verify that A is DG Morita equivalent to the factor algebra $B = A/J_A^2$ (an explicit tilting object in $D^c(A)$ with endomorphism algebra isomorphic to B is given by the direct sum of the simple A -modules at the vertices 1 and 3 together with the unique indecomposable projective-injective A -module). Thus, we may and we will identify minimal A_∞ -models of $D^c(A)$ with those of $D^c(B)$ (we do this to minimise the number of projective resolutions needed in the computation below).

The algebra B has global dimension 2 and $D^c(B)$ admits a $2\mathbb{Z}$ -cluster tilting subcategory given by

$$\mathcal{U} = \mathcal{U}(B) := \text{add} \{(B \oplus S_1)[2i] \mid i \in \mathbb{Z}\} \subset D^c(B),$$

where B is the regular representation and S_1 is the simple B -module concentrated at the vertex 1, see [\[Iya11, Cor. 1.15 and Thm. 1.21\]](#) (the claim can also be verified directly using the explicit description of the Auslander–Reiten quiver of $D^c(A) \simeq D^c(B)$ available in [\[Hap88, Sec. I.5\]](#) for example). By [Theorem 2.2.1](#) the pair $(\mathcal{U}, [2])$ has an induced standard 4-angulation; it is also easy to verify that all 4-angles in \mathcal{U} are finite direct sums of rotations of trivial 4-angles as in [\(d-TR1b\)](#) and the apparent 4-angle

$$P_3 \xrightarrow{a} P_2 \xrightarrow{b} P_1 \rightarrow S_1 \rightarrow P_3[2],$$

where P_i denotes the indecomposable projective B -module at the vertex i and the connecting morphism $S_1 \rightarrow P_3[2]$ classifies the (essentially unique) non-zero class in $\text{Ext}_B^2(S_1, P_3)$; thus, the above 4-angle consists of the minimal projective resolution of S_1 together with its canonical augmentation. [Theorems 4.1.12](#) and [4.2.6](#) imply that the Toda bracket/Massey product of the morphisms in the above 4-angle contains $\pm 1_{P_3}$. By [Proposition 4.1.31](#), this Massey product is an element of the quotient

$$\frac{\text{hom}_B(P_3, P_3)}{\text{hom}_B(P_2, P_3) \cdot a},$$

hence it cannot contain 0 since it contains $\pm 1_{P_3}$ and the morphism a is not a split monomorphism. Therefore, any minimal A_∞ -model of the canonical DG enhancement of $D^c(B)$ must have non-vanishing m_4 by [Proposition 4.3.6](#). For the convenience of the reader, below we provide an explicit computation of the action of m_4 on the above 4-angle by means of Kadeishvili’s Homotopy Transfer Theorem.

The minimal projective resolution Q of S_1 is the apparent complex

$$Q: \quad P_3 \xrightarrow{a} P_2 \xrightarrow{b} P_1.$$

As a graded quiver with relations, the full DG subcategory of $\text{C}_{\text{dg}}(\text{Mod } B)$ spanned by Q and the P_i ’s the following:

$$\begin{array}{ccccc}
P_3 & \xrightarrow{a} & P_2 & \xrightarrow{b} & P_1 \\
u' \uparrow \downarrow u & & v' \uparrow \downarrow v & & w' \uparrow \downarrow w \\
(P_3 & \xrightarrow{a} & P_2 & \xrightarrow{b} & P_1)
\end{array}$$

The bottom row is the complex Q and the degrees of the arrows and the relations are as follows:

$$|a| = |b| = |w| = |w'| = 0 \quad -|v| = |v'| = 1 \quad -|u| = |u'| = 2$$

$$\begin{aligned}
u'u &= \mathbf{1}_{P_3} & v'v &= \mathbf{1}_{P_2} & w'w &= \mathbf{1}_{P_1} & uu' + vv' + ww' &= \mathbf{1}_Q & ba &= 0 \\
u'v &= 0 & v'u &= 0 & v'w &= 0 & w'v &= 0 & u'w &= 0 & w'u &= 0.
\end{aligned}$$

The non-zero differentials of the generating morphisms are given by

$$\partial(u) = va \quad \partial(v) = wb \quad \partial(w') = -bv' \quad \partial(v') = -au';$$

the remaining differentials are uniquely determined by the graded Leibniz rule. Similarly, as a graded quiver with relations, the graded cohomology category is

$$\begin{array}{ccccc}
P_3 & \xrightarrow{a} & P_2 & \xrightarrow{b} & P_1 \\
\underline{u}' \uparrow & & & & \downarrow \underline{w} \\
(P_3 & \xrightarrow{a} & P_2 & \xrightarrow{b} & P_1)
\end{array}$$

The latter quiver is subject to the relation that the composite of any two consecutive arrows vanishes.

We claim that³

$$m_4(\underline{u}', \underline{w}, \underline{b}, \underline{a}) = \pm \mathbf{1}_{P_3}.$$

The part of the necessary homotopy retract that is relevant to our computation is the following:

$$\begin{aligned}
i(\underline{a}) &= a & i(\underline{b}) &= b & i(\underline{w}) &= w & i(\underline{u}') &= u' \\
p(u'u) &= \mathbf{1}_{P_3} & h(va) &= u & h(wb) &= v.
\end{aligned}$$

Recall that m_4 is defined combinatorially in terms of planar binary rooted trees by interpolating the original composition operation with the homotopy h , see for example [Val14, p. 233]. Since the composites $u'w$ and ba vanish on the nose, the summands of $m_4(\underline{u}', \underline{w}, \underline{b}, \underline{a})$ that involve first composing (after applying i) either the leftmost two or the rightmost two morphisms vanish. This leaves only the two summands that involve first composing w with b and then applying h . Finally, given that the composite $u'h(wb) = u'v = 0$ vanishes,

$$\begin{array}{ccccccc}
u' = i(\underline{u}') & & w = i(\underline{w}) & & b = i(\underline{b}) & & a = i(\underline{a}) \\
& & \searrow & & \swarrow & & \\
& & & v = h(wb) & & & \\
& & & \searrow & & \swarrow & \\
& & & & u = h(va) & & \\
& \searrow & & \swarrow & & & \\
& & \mathbf{1}_{P_3} = p(u'u) & & & &
\end{array}$$

is the only non-zero summand of $m_4(\underline{u}', \underline{w}, \underline{b}, \underline{a})$, which is what we had to prove.

³This computation goes back to Kontsevich, see [Sei08, Rmk. 3.11] and compare with [HKK17, Eq. 3.18].

Remark 4.6.1. The above example shows that [Her16, Thm. D] cannot hold as stated. Indeed, [Her16, Thm. D] claims that the compact derived category of an acyclic quiver has a minimal A_∞ -model with only one non-zero higher operation, namely m_3 ; the previous example shows that there are higher non-vanishing operations in general. Notice also that [Her16] would contradict our [Corollary 4.5.17](#), as the latter implies that the Hochschild class of m_4 in the above example must be non-zero. The error in the proof of [Her16, Thm. D] occurs in the proof of Prop. 3.8.1 therein, where it is claimed that the higher operations m_n , $n > 3$, vanish for the minimal A_∞ -models of bounded derived categories of hereditary abelian categories with enough projectives. There, the author argues that the higher composite of a sequence of n morphisms in the graded cohomology category always represents a morphism in the $(n - 2)$ -nd extension space between two complexes that lie in the heart of the derived category, and these spaces indeed vanish for $n > 3$ due to the hereditary assumption. However, as the example above shows, the sequence

$$P_3 \xrightarrow{a} P_2 \xrightarrow{b} P_1 \xrightarrow{w} Q \xrightarrow{w'} P_3$$

has total degree 2 and its higher composite yields a non-zero morphism $P_3 \rightarrow P_3$ of degree 0 (notice that m_4 is a morphism of degree -2).

4.7. Hochschild cohomology of sparse graded algebras.

Setting 4.7.1. We fix a finite-dimensional self-injective algebra Λ and an algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$.

In this technical subsection we study the Hochschild cohomology of the d -sparse graded algebra

$$\Lambda(\sigma, d) := \frac{\Lambda\langle \iota^{\pm 1} \rangle}{(\iota x - \sigma(x)\iota)_{x \in \Lambda}}, \quad |\iota| = -d,$$

(compare with [Remark 4.5.3](#)). The results in this section extend results obtained in [Mur22] in the case $d = 1$.

We begin with some general observations that hold for an arbitrary sparse graded algebra.

Definition 4.7.2. Let A be a d -sparse graded algebra. The *fractional Euler derivation* $\delta_{/d}: A \rightarrow A$ is the degree 0 map

$$\delta_{/d}(a) = \frac{|a|}{d} \cdot a,$$

where the coefficient is indeed an integer in view of the sparseness of A .

Remark 4.7.3. Let A be a d -sparse graded algebra. The (usual) Euler derivation is $\delta = \delta_{/1}$. If d is invertible in the ground field, then $\delta_{/d} = \frac{1}{d}\delta$.

Proposition 4.7.4. Let A be a d -sparse graded algebra and $\varphi \in \mathbb{C}^{p,q}(A)$ a Hochschild cochain. Then,

$$[\delta_{/d}, \varphi] = \frac{q}{d} \cdot \varphi.$$

Moreover, $\delta_{/d}$ is a Hochschild cocycle.

Proof. The proof is almost identical to those of [Mur22, Prop. 3.2 and Cor. 3.3] and is left to the reader. \square

Notation 4.7.5. Let A be a d -sparse graded algebra. The Hochschild cohomology class of the fractional Euler derivation, called *fractional Euler class*, is denoted

$$\{\delta_{/d}\} \in \mathrm{HH}^{1,0}(A).$$

Proposition 4.7.6. *The square of the fractional Euler class vanishes:*

$$\{\delta_{/d}\}^2 = 0 \in \mathrm{HH}^{2,0}(A).$$

Proof. The proof of [Mur22, Prop. 3.8] applies almost verbatim, only that one needs to replace the cochain $\beta \in \mathbb{C}^{1,0}(A)$ in *loc. cit.* by the cochain

$$\beta: x \mapsto \frac{\frac{|x|}{d} \left(1 - \frac{|x|}{d}\right)}{2} \cdot x. \quad \square$$

Proposition 4.7.7. *If $\mathrm{char}(\mathbf{k}) = 2$, then $\mathrm{Sq}(\{\delta_{/d}\}) = \{\delta_{/d}\}$.*

Proof. Indeed, the claim follows from the fact that every integer is congruent to its square modulo 2 (compare with [Mur22, Prop. 3.5]). \square

Straightforward computations that combine the above propositions with the laws of a Gerstenhaber algebra yield the following identities (compare with [Mur22, Props. 3.6, 3.9, 3.10]). We leave the proof to the reader.

Proposition 4.7.8. *Let A be a d -sparse graded algebra. The following identities hold for $x \in \mathrm{HH}^{p,dq}(A)$ and $y \in \mathrm{HH}^{s,dt}(A)$:*

$$[\{\delta_{/d}\} \cdot x, \{\delta_{/d}\} \cdot y] = (t - q) \cdot \{\delta_{/d}\} \cdot x \cdot y.$$

$$\mathrm{Sq}(\{\delta_{/d}\} \cdot x) = 0 \quad \text{if } \mathrm{char}(\mathbf{k}) \neq 2 \text{ and } p + dq \text{ is odd.}$$

$$\mathrm{Sq}(\{\delta_{/d}\} \cdot x) = (q + 1) \cdot \{\delta_{/d}\} \cdot x^2 \quad \text{if } \mathrm{char}(\mathbf{k}) = 2.$$

$$y \cdot x = [y, \{\delta_{/d}\} \cdot x] + \{\delta_{/d}\} \cdot [y, x] \quad \text{if } t = -1 \text{ and } d - s \text{ is even.}$$

We return to our object of interest in this subsection, which is the Hochschild cohomology of the d -sparse graded algebra $\Lambda(\sigma, d)$ (see [Setting 4.7.1](#)).

Remark 4.7.9. Recall that, by definition, the following identity holds in $\Lambda(\sigma, d)$:

$$\iota x \iota^{-1} = \sigma(x), \quad x \in \Lambda.$$

Now, the automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$ extends to $\Lambda(\sigma, d)$ via the formula

$$\sigma(\iota) := (-1)^{|\iota|} \iota = (-1)^{-d} \iota.$$

Since $\Lambda(\sigma, d)^{dk} = \{\iota^{-k} x \mid x \in \Lambda\}$, we see that the equality

$$\iota x \iota^{-1} = (-1)^{|x|} \sigma(x)$$

holds for all homogeneous elements $x \in \Lambda(\sigma, d)$.

Remark 4.7.10. Observe that the group $\mathrm{Aut}(\Lambda(\sigma, d))$ of graded algebra automorphisms of $\Lambda(\sigma, d)$ acts naturally on the right by conjugation on the bigraded algebra $\mathrm{HH}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$ as in [Remark 4.4.1](#), since $\Lambda = \Lambda(\sigma, d)^0$ is the degree 0 part. It also acts on $\underline{\mathrm{HH}}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$ since Hochschild–Tate cohomology satisfies the same functoriality properties as Hochschild cohomology. In particular the latter action preserves units. The canonical comparison map

$$\mathrm{HH}^{\bullet,*}(\Lambda, \Lambda(\sigma, d)) \longrightarrow \underline{\mathrm{HH}}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$$

is $\mathrm{Aut}(\Lambda(\sigma, d))$ -equivariant. Hence the action of $\mathrm{Aut}(\Lambda(\sigma, d))$ on $\mathrm{HH}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$ preserves edge units.

A unit $u \in Z(\Lambda)^\times$ in the centre of Λ induces a graded algebra automorphism $g_u: \Lambda(\sigma, d) \xrightarrow{\sim} \Lambda(\sigma, d)$ that is uniquely determined by the requirements

$$g_u(x) = x, \quad x \in \Lambda = \Lambda(\sigma, d)^0, \quad \text{and} \quad g_u(\iota) = \iota u^{-1}.$$

Indeed, these formulas define g_u on the graded algebra generators of $\Lambda(\sigma, d)$. In order to check compatibility with the defining relations of $\Lambda(\sigma, d)$ in [Setting 4.7.1](#),

We have a short exact sequence of $\Lambda(\sigma, d)$ -bimodules

$$\Lambda(\sigma, d) \otimes_{\Lambda} \Lambda(\sigma, d) \xrightarrow{1-\Gamma} \Lambda(\sigma, d) \otimes_{\Lambda} \Lambda(\sigma, d) \xrightarrow{\mu} \Lambda(\sigma, d)$$

where μ is the product in $\Lambda(\sigma, d) \otimes_{\Lambda} \Lambda(\sigma, d)$ and Γ is defined as

$$\Gamma(a \otimes b) = av^{-1} \otimes ib.$$

Indeed, using the isomorphisms of Λ -bimodules

$$\begin{aligned} \sigma^i \Lambda_1 \otimes_{\Lambda} \sigma^j \Lambda_1 &\xrightarrow{\cong} \sigma^{i+j} \Lambda_1, \\ a \otimes b &\longmapsto \sigma^j(a)b, \end{aligned}$$

we see that, as Λ -bimodules,

$$(\Lambda(\sigma, d) \otimes_{\Lambda} \Lambda(\sigma, d))^k \cong \bigoplus_{i \in \mathbb{Z}} \sigma^k \Lambda_1.$$

Through this decomposition, μ maps all copies of $\sigma^k \Lambda_1$ to itself identically, and Γ maps the i^{th} direct summand to the $(i+1)^{\text{st}}$ direct summand identically. Hence, the previous sequence is degree-wise a well-known short exact sequence.

The $\Lambda(\sigma, d)$ -bimodule $\Lambda(\sigma, d) \otimes_{\Lambda} \Lambda(\sigma, d)$ is the extension of scalars of Λ along $j^{\otimes 2}: \Lambda^e \rightarrow \Lambda(\sigma, d)^e$. The long exact sequence in the statement will be the one associated to the previous short exact sequence and the derived functors of $\text{hom}_{\Lambda(\sigma, d)^e}^*(-, \Lambda(\sigma, d))$. We just have to check that the following square is commutative

$$\begin{array}{ccc} \text{hom}_{\Lambda(\sigma, d)^e}^*(\Lambda(\sigma, d) \otimes_{\Lambda} \Lambda(\sigma, d), \Lambda(\sigma, d)) & \xrightarrow{\cong} & \text{hom}_{\Lambda^e}^*(\Lambda, \Lambda(\sigma, d)) \\ \text{hom}_{\Lambda(\sigma, d)^e}^*(\Gamma, \Lambda(\sigma, d)) \downarrow & & \downarrow \sigma_*^{-1} \sigma^* \\ \text{hom}_{\Lambda(\sigma, d)^e}^*(\Lambda(\sigma, d) \otimes_{\Lambda} \Lambda(\sigma, d), \Lambda(\sigma, d)) & \xrightarrow{\cong} & \text{hom}_{\Lambda^e}^*(\Lambda, \Lambda(\sigma, d)) \end{array}$$

Here, the horizontal isomorphisms are the adjunction isomorphisms. Let us check this. Let $f: \Lambda(\sigma, d) \otimes_{\Lambda} \Lambda(\sigma, d) \rightarrow \Lambda(\sigma, d)$ be a $\Lambda(\sigma, d)$ -bimodule morphism. Since source and target are d -sparse, we can suppose that f has degree kd for some $k \in \mathbb{Z}$. The image of f through the upper right corner is the Λ -bimodule morphism $g: \Lambda \rightarrow \Lambda(\sigma, d)$ given by $g(x) = \sigma^{-1}f(\sigma(x) \otimes 1)$. The image through the lower left corner is the Λ -bimodule morphism $h: \Lambda \rightarrow \Lambda(\sigma, d)$ given by

$$\begin{aligned} h(x) &= f\Gamma(x \otimes 1) \\ &= f(xv^{-1} \otimes v) \\ &= f(v^{-1}xv \otimes v) \\ &= (-1)^{kd^2} v^{-1}f(vxv^{-1} \otimes v)v \\ &= \sigma^{-1}f(\sigma(x) \otimes 1) \\ &= g(x). \end{aligned}$$

Here we use that f is a $\Lambda(\sigma, d)$ -bimodule morphism and that $d^2 \equiv d \pmod{2}$.

The morphisms j^* and $\sigma_*^{-1} \sigma^*$ in the statement are algebra maps by functoriality properties of Hochschild cohomology. The multiplicative properties of ∂ require a deeper analysis.

The $\Lambda(\sigma, d)$ -bimodule endomorphism Γ lifts to the projective resolution $\Lambda(\sigma, d) \otimes_{\Lambda} B_{\bullet}(\Lambda) \otimes_{\Lambda} \Lambda(\sigma, d)$ in the following way,

$$\begin{aligned} \Gamma: \Lambda(\sigma, d) \otimes_{\Lambda} B_k(\Lambda) \otimes_{\Lambda} \Lambda(\sigma, d) &\longrightarrow \Lambda(\sigma, d) \otimes_{\Lambda} B_k(\Lambda) \otimes_{\Lambda} \Lambda(\sigma, d), \\ a \otimes x_1 \otimes \cdots \otimes x_k \otimes b &\longmapsto av^{-1} \otimes \sigma(x_1) \otimes \cdots \otimes \sigma(x_k) \otimes ib. \end{aligned}$$

Therefore, the mapping cone of the endomorphism $\mathbf{1} - \Gamma$ of $\Lambda(\sigma, d) \otimes_{\Lambda} B_{\bullet}(\Lambda) \otimes_{\Lambda} \Lambda(\sigma, d)$ is a projective resolution of $\Lambda(\sigma, d)$ as a $\Lambda(\sigma, d)$ -bimodule. This mapping cone is

$$C_{\mathbf{1}-\Gamma} = \Lambda(\sigma, d) \otimes_{\Lambda} B_{\bullet}(\Lambda) \otimes_{\Lambda} \Lambda(\sigma, d) \oplus \Lambda(\sigma, d) \otimes_{\Lambda} B_{\bullet-1}(\Lambda) \otimes_{\Lambda} \Lambda(\sigma, d)$$

with differential

$$\begin{pmatrix} d_B & 0 \\ \mathbf{1} - \Gamma & -d_B \end{pmatrix}.$$

Here d_B stands for the bar complex differential.

Consider the maps

$$\Lambda(\sigma, d) \otimes_{\Lambda} B_{\bullet}(\Lambda) \otimes_{\Lambda} \Lambda(\sigma, d) \xrightarrow{\mathbf{1}-\Gamma} \Lambda(\sigma, d) \otimes_{\Lambda} B_{\bullet}(\Lambda) \otimes_{\Lambda} \Lambda(\sigma, d) \xrightarrow{j_*} B_{\bullet}(\Lambda(\sigma, d))$$

where j_* is defined by

$$j_*(a \otimes x_1 \otimes \cdots \otimes x_j \otimes b) = a \otimes j(x_1) \otimes \cdots \otimes j(x_j) \otimes b.$$

We could omit the j 's on the right because it is just the inclusion $\Lambda \subset \Lambda(\sigma, d)$. The composite is null-homotopic, with explicit homotopy

$$\begin{aligned} h(a \otimes x_1 \otimes \cdots \otimes x_j \otimes b) = \\ \sum_{k=0}^j (-1)^k a \iota^{-1} \otimes \sigma(x_1) \otimes \cdots \otimes \sigma(x_k) \otimes \iota \otimes x_{k+1} \otimes \cdots \otimes x_j \otimes b. \end{aligned}$$

Indeed, it is easy to check that

$$j_*(\mathbf{1} - \Gamma) = d_B h + h d_B.$$

This gives rise to an explicit map of resolutions

$$\begin{aligned} C_{\mathbf{1}-\Gamma} &\longrightarrow B_{\bullet}(\Lambda(\sigma, d)), \\ (x, y) &\longmapsto j_*(x) + h(y). \end{aligned}$$

Taking $\mathrm{hom}_{\Lambda(\sigma, d)^e}^*$, we obtain a quasi-isomorphism

$$\mathbf{C}^{\bullet,*}(\Lambda(\sigma, d), \Lambda(\sigma, d)) \xrightarrow{\sim} \mathrm{hom}_{\Lambda(\sigma, d)^e}^*(C_{\mathbf{1}-\Gamma}, \Lambda(\sigma, d)).$$

The Hochschild complex (source) is a differential graded algebra. We can endow the target with a differential graded algebra structure turning this map into an algebra map. It is the following square-zero extension

$$\mathrm{hom}_{\Lambda(\sigma, d)^e}^*(C_{\mathbf{1}-\Gamma}, \Lambda(\sigma, d)) = \mathbf{C}^{\bullet,*}(\Lambda, \Lambda(\sigma, d)) \oplus_{\sigma_*^{-1}\sigma^*} \mathbf{C}^{\bullet-1,*}(\Lambda, \Lambda(\sigma, d))_1$$

of the differential graded algebra $\mathbf{C}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$ by the twisted and then shifted bimodule ${}_{\sigma_*^{-1}\sigma^*} \mathbf{C}^{\bullet-1,*}(\Lambda, \Lambda(\sigma, d))_1$. It is straightforward to check that the differential satisfies the Leibniz rule with respect to the product. Moreover, the quasi-isomorphism is an algebra map because, given $f, g \in \mathbf{C}^{\bullet,*}(\Lambda(\sigma, d), \Lambda(\sigma, d))$,

$$h^*(f \cdot g) = h^*(f) \cdot j^*(g) + (-1)^{|f|} \sigma_*^{-1} \sigma^* j^*(f) \cdot h^*(g).$$

This is tedious but straightforward to check.

The long exact sequence in the statement is induced by the square-zero extension

$${}_{\sigma_*^{-1}\sigma^*} \mathbf{C}^{\bullet-1,*}(\Lambda, \Lambda(\sigma, d))_1 \xrightarrow{i_2} \mathrm{hom}_{\Lambda(\sigma, d)^e}^*(C_{\mathbf{1}-\Gamma}, \Lambda(\sigma, d)) \xrightarrow{p_1} \mathbf{C}^{\bullet,*}(\Lambda, \Lambda(\sigma, d))$$

where p_1 is an algebra map and i_2 is a $\mathrm{hom}_{\Lambda(\sigma, d)^e}^*(C_{\mathbf{1}-\Gamma}, \Lambda(\sigma, d))$ -bimodule morphism. This proves that ∂ , which is induced by $-i_2$, is a bimodule morphism. In cohomology, we can remove the left twisting by $\sigma_*^{-1}\sigma^*$ since $\mathrm{HH}^{\bullet,*}(\Lambda(\sigma, d), \Lambda(\sigma, d))$ maps to the kernel of $\mathbf{1} - \sigma_*^{-1}\sigma^*$.

The coordinates of the image of the fractional Euler derivation under the quasi-isomorphism are

$$j^*(\delta/d) = 0,$$

because Λ is ungraded, and

$$h^*(\delta_{/d}) = \iota^{-1}\delta_{/d}(\iota) = \frac{|\iota|}{d}\iota^{-1}\iota = -1,$$

hence it corresponds to $(0, -1) \in \text{hom}_{\Lambda(\sigma, d)^e}^*(C_{\mathbf{1}-\Gamma}, \Lambda(\sigma, d))$. The composite ∂j^* is induced by $-i_2 p_1$ and, in $\text{hom}_{\Lambda(\sigma, d)^e}^*(C_{\mathbf{1}-\Gamma}, \Lambda(\sigma, d))$,

$$(0, -1) \cdot (f, g) = (0, -f) = -i_2 p_1(f, g).$$

This proves the last claim. \square

We record the following immediate corollary of [Proposition 4.7.11](#). Below, we denote by $\langle \sigma \rangle \subseteq \text{Aut } \Lambda(\sigma, d)$ be the subgroup generated by the extension of σ to $\Lambda(\sigma, d)$

Corollary 4.7.12. *In the setting of [Proposition 4.7.11](#), the inclusion $j : \Lambda \rightarrow \Lambda(\sigma, d)$ induces a square-zero extension of bigraded algebras*

$$\text{HH}^{\bullet-1, *}(\Lambda, \Lambda(\sigma, d))_{\langle \sigma \rangle} \hookrightarrow \text{HH}^{\bullet, *}(\Lambda(\sigma, d), \Lambda(\sigma, d)) \twoheadrightarrow \text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))^{\langle \sigma \rangle},$$

where $\text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))^{\langle \sigma \rangle}$ and $\text{HH}^{\bullet-1, *}(\Lambda, \Lambda(\sigma, d))_{\langle \sigma \rangle}$ denote the bigraded algebra of invariants and the shifted bimodule of coinvariants of the action of $\langle \sigma \rangle \subset \text{Aut}(\Lambda(\sigma, d))$ on $\text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))$, respectively (see [Remark 4.7.10](#)).

Remark 4.7.13. [Corollary 4.7.12](#) implies that $\text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))^{\langle \sigma \rangle}$ is graded commutative with respect to the total degree as is $\text{HH}^{\bullet, *}(\Lambda(\sigma, d), \Lambda(\sigma, d))$. Moreover $\text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))_{\langle \sigma \rangle}$ is a symmetric $\text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))^{\langle \sigma \rangle}$ -bimodule, that is a module in the commutative sense. Furthermore, when the obvious map

$$\text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))^{\langle \sigma \rangle} \longrightarrow \text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))_{\langle \sigma \rangle}$$

is an isomorphism, [Proposition 4.7.11](#) identifies the kernel of the square-zero extension with the ideal generated by $\{\delta_{/d}\}$. In particular, in this case, multiplication by $\{\delta_{/d}\}$ in $\text{HH}^{\bullet, *}(\Lambda(\sigma, d), \Lambda(\sigma, d))$ induces a splitting of the square-zero extension. A sufficient condition is that the order of σ as an automorphism of $\Lambda(\sigma, d)$ is finite (compare with the Periodicity Conjecture of Erdmann and Skoworński [[ES08](#)]) and invertible in the ground field \mathbf{k} . Notice that the order of σ as an automorphism of $\Lambda(\sigma, d)$ is twice the order of σ as an automorphism of Λ if d is odd and \mathbf{k} has odd characteristic, and they coincide otherwise, see [Remark 4.7.9](#).

Recall that we have a canonical comparison map

$$\text{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d)) \longrightarrow \underline{\text{HH}}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))$$

from Hochschild cohomology to Hochschild–Tate cohomology, that is an isomorphism in positive horizontal degree and an epimorphism in horizontal degree 0. In particular, we may identify $\text{HH}^{>0, *}(\Lambda, \Lambda(\sigma, d))$ with the (horizontal) positive part of $\underline{\text{HH}}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))$.

We introduce the following auxiliary definition, see [[Mur22](#), Sec. 6] for justification for the terminology (in particular [[Mur22](#), Prop. 6.5 and Cor. 6.7]).

Definition 4.7.14. Let $p > 0$.

- (1) A class

$$x \in \text{HH}^{p, q}(\Lambda, \Lambda(\sigma, d)) = \text{Ext}_{\Lambda^e}^p(\Lambda, \sigma^q \Lambda_1)$$

is an *edge unit* if it is a unit $\underline{\text{HH}}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))$.

- (2) Let $j : \Lambda \hookrightarrow \Lambda(\sigma, d)$ be the inclusion of Λ as the degree 0 part. A class $x \in \text{HH}^{p, q}(\Lambda(\sigma, d), \Lambda(\sigma, d))$ is an *edge unit* if the image of x under the map

$$j^* : \text{HH}^{p, q}(\Lambda(\sigma, d), \Lambda(\sigma, d)) \longrightarrow \text{HH}^{p, q}(\Lambda, \Lambda(\sigma, d))$$

is an edge unit.

The following property of edge units is key for later computations.

Proposition 4.7.15. *Let $x \in \mathrm{HH}^{d+2,-d}(\Lambda(\sigma, d))$ be an edge unit. Then, the left-multiplication map*

$$\mathrm{HH}^{p,*}(\Lambda(\sigma, d)) \longrightarrow \mathrm{HH}^{p+d+2,*-d}(\Lambda(\sigma, d)), \quad y \longmapsto x \cdot y,$$

is bijective for $p \geq 2$ and surjective for $p = 1$.

Proof. Using [Proposition 4.7.11](#), both claims can be proved with the same argument used in the proof of [\[Mur22, Lemma 6.8\]](#). \square

The following result—a direct consequence of [Proposition 2.2.18](#)—is responsible for the *uniqueness* of enhancements in [Theorem 5.1.2](#) (compare with [\[Mur22, Prop. 9.4\]](#)⁴). In particular, it is here that it is crucial for us to work with finite-dimensional algebras and not with more general objects such as locally-finite categories (see the proof of [Proposition 2.2.18](#), in particular the use of [\[Che21, Cor. 2.3\]](#) and compare with [Remark 2.2.19](#)).

Proposition 4.7.16. *Let Λ be a finite-dimensional self-injective algebra that is twisted $(d+2)$ -periodic with respect to an algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$, $d \geq 1$. Then, the set formed by the edge units in $\mathrm{HH}^{d+2,-d}(\Lambda, \Lambda(\sigma, d))$ is an orbit for the action of $\mathrm{Aut}(\Lambda(\sigma, d))$.*

Proof. By [Remark 4.7.10](#), the orbit of an edge unit consists of edge units. Let x and x' be edge units in $\mathrm{HH}^{d+2,-d}(\Lambda, {}_1\Lambda_\sigma)$ and choose exact sequences

$$\eta: \quad 0 \rightarrow {}_1\Lambda_\sigma \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

and

$$\eta': \quad 0 \rightarrow {}_1\Lambda_\sigma \rightarrow Q_{d+1} \rightarrow \cdots \rightarrow Q_1 \rightarrow Q_0 \rightarrow \Lambda \rightarrow 0$$

of Λ -bimodules with projective-injective middle terms that represent x and y , respectively (see [Remark 4.5.16](#)). [Proposition 2.2.18](#) yields a morphism of exact sequences

$$\begin{array}{ccccccccccc} \eta: & 0 & \rightarrow & {}_1\Lambda_\sigma & \rightarrow & P_{d+1} & \rightarrow & \cdots & \rightarrow & P_1 & \rightarrow & P_0 & \rightarrow & \Lambda & \rightarrow & 0 \\ & & & \downarrow u & & \downarrow & & & & \downarrow & & \downarrow & & \parallel & & \\ \eta': & 0 & \rightarrow & {}_1\Lambda_\sigma & \rightarrow & Q_{d+1} & \rightarrow & \cdots & \rightarrow & Q_1 & \rightarrow & Q_0 & \rightarrow & \Lambda & \rightarrow & 0 \end{array}$$

where $u \in Z(\Lambda)^\times$, which implies that

$$x' = u \cdot x = x^{g_u},$$

see again [Remark 4.7.10](#). \square

5. EXISTENCE AND UNIQUENESS OF ENHANCEMENTS

Setting 5.0.1. We fix a finite-dimensional basic self-injective algebra Λ that is twisted $(d+2)$ -periodic for some $d \geq 1$ with respect to an algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$ and let

$$\Sigma := - \otimes_{\Lambda} \sigma \Lambda_1: \mathrm{proj}(\Lambda) \xrightarrow{\sim} \mathrm{proj}(\Lambda).$$

As in [Remark 4.5.3](#), we consider the d -sparse graded algebra

$$\Lambda(\sigma, d) := \frac{\Lambda\langle \iota^\pm \rangle}{(\iota x - \sigma(x)\iota)_{x \in \Lambda}}, \quad |\iota| = -d.$$

⁴The formula for g_u in the proof of [\[Mur22, Prop. 9.4\]](#) (called $g(x)$ therein for $x = u \in Z(\Lambda)^\times$), where the case $d = 1$ is considered, is wrong in degrees $\neq 0, -1$ unless $\sigma(u) = u$, since otherwise u is not a central element in $\Lambda(\sigma, d)$ and hence $g(x)$ does not preserve products. The correct definition is the one in [Remark 4.7.10](#) in terms of generators and relations, which yields different degree-wise formulas. The rest of the argument in the proof of [\[Mur22, Prop. 9.4\]](#) is nevertheless valid since it only uses the explicit formulas in degrees 0 and -1 .

5.1. **Proof Theorem A.** In this subsection we give the proof **Theorem A**, modulo the proof of the following main result (**Theorem 5.1.2**) that we postpone until **Section 5.3**.

Notation 5.1.1. Let $g \in \text{Aut}(\Lambda(\sigma, d))$ be a graded algebra automorphism and

$$(\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, m_{3d+2}, \dots)$$

a minimal A_∞ -algebra structure on $\Lambda(\sigma, d)$. We let

$$(\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) * g := (\Lambda(\sigma, d), m_{d+2}^g, m_{2d+2}^g, \dots),$$

where $m_i^g := g^{-1}m_i g^{\otimes i}$, and note that this is also a minimal A_∞ -algebra structure on the d -sparse graded algebra $\Lambda(\sigma, d)$.

Theorem 5.1.2. *The following statements hold:*

(1) *There exists a minimal A_∞ -algebra structure*

$$(\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots)$$

on the d -sparse graded algebra $\Lambda(\sigma, d)$ whose restricted universal Massey product

$$j^* \{m_{d+2}\} \in \text{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d)) = \text{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$$

*is a unit in the Hochschild–Tate cohomology $\underline{\text{HH}}^{\bullet, *}$ ($\Lambda, \Lambda(\sigma, d)$) (equivalently, the class $j^* \{m_{d+2}\}$ is represented by an extension all of whose middle terms are projective-injective Λ -bimodules).*

(2) *Let*

$$(\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) \quad \text{and} \quad (\Lambda(\sigma, d), m'_{d+2}, m'_{2d+2}, \dots)$$

be minimal A_∞ -algebra structures on $\Lambda(\sigma, d)$ whose corresponding restricted universal Massey products $j^ \{m_{d+2}\}$ and $j^* \{m'_{d+2}\}$ are units in the Hochschild–Tate cohomology $\underline{\text{HH}}^{\bullet, *}$ ($\Lambda, \Lambda(\sigma, d)$). Then, there exist a graded algebra automorphism $g \in \text{Aut}(\Lambda(\sigma, d))$ and a quasi-isomorphism of minimal A_∞ -algebras*

$$F: (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) \xrightarrow{\cong} (\Lambda(\sigma, d), m'_{d+2}, m'_{2d+2}, \dots) * g$$

such that $F_0 = \mathbf{1}$ and $F_1 = \mathbf{1}$.

We need some recollections on the derived category of an A_∞ -algebra. Given an A_∞ -algebra A , we denote by $A_\infty\text{-Mod}(A)$ the DG (!) category of (*right*) A_∞ -modules over A , that is the DG category of A_∞ -functors $A^{\text{op}} \rightarrow \text{C}_{\text{dg}}(\mathbf{k})$ from the opposite A_∞ -algebra of A (viewed as an A_∞ -category with a single object $*$) to the DG category of cochain complexes of vector spaces, see for example [Sei08, Sec. I.1j]. The A_∞ -categorical Yoneda embedding

$$A \longrightarrow A_\infty\text{-Mod}(A), \quad * \longmapsto A_A$$

yields a quasi-isomorphism of A_∞ -algebras

$$A \xrightarrow{\sim} \overline{A} := \text{hom}_A(A_A, A_A)$$

where \overline{A} is the DG (!) algebra of endomorphisms of the free A_∞ -module A_A , see for example [Sei08, Sec. I.2g] (to our knowledge, this important fact was first observed by Kontsevich and Soibelman [KS06, Sec. 6.2], but see also [LH, Sec. 7.5]). In particular, there is a canonical isomorphism of graded algebras

$$H^\bullet(A) \xrightarrow{\sim} H^\bullet(\overline{A}).$$

Finally, the derived category of A is defined as

$$\text{D}(A) := H^0(A_\infty\text{-Mod}(A));$$

this is a sensible definition since quasi-isomorphisms of A_∞ -modules are also homotopy equivalences, see [Kel01, Sec. 4] or [LH, Thm. 4.1.3.1] while keeping in

mind that we work over a field. By construction, there is a canonical equivalence of triangulated categories

$$\mathrm{D}(\overline{A}) \xrightarrow{\sim} \mathrm{D}(A),$$

which is in fact induced by the quasi-equivalence of DG categories

$$A_\infty\text{-Mod}(\overline{A}) \xrightarrow{\sim} A_\infty\text{-Mod}(A)$$

given by restriction of scalars along the quasi-isomorphism $A \xrightarrow{\sim} \overline{A}$, see for example [Sei08, Lemma I.2.10]. The following result can be regarded as a $(d+2)$ -angulated analogue of one of the main theorems in [Mur22].

Theorem 5.1.3. *Let Λ be a finite-dimensional self-injective algebra that is twisted $(d+2)$ -periodic with respect to an algebra automorphism σ and let*

$$\Sigma := - \otimes_{\Lambda} \sigma \Lambda_1 : \mathrm{proj}(\Lambda) \xrightarrow{\sim} \mathrm{proj}(\Lambda).$$

Then, the AL $(d+2)$ -angulation \diamond_{η} of $(\mathrm{proj}(\Lambda), \Sigma)$ induced by any exact sequence of Λ -bimodules

$$\eta: \quad 0 \rightarrow {}_1\Lambda_\sigma \rightarrow P_{d+1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \Lambda \rightarrow 0$$

with projective-injective middle terms (Theorem 2.2.15) admits an enhancement. Moreover, if Λ/J_Λ is separable, then the enhancement is unique.

Proof. Recall from Proposition 2.2.18 that, up to equivalence, the AL $(d+2)$ -angulation is independent of the choice of the exact sequence η (as long as it satisfies the conditions in the statement of the theorem).

We prove the first claim, that the AL $(d+2)$ -angulation on $(\mathrm{proj}(\Lambda), \Sigma)$ admits an enhancement. Firstly, Theorem 5.1.2(1) yields a minimal A_∞ -algebra structure

$$A := (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots)$$

on the d -sparse graded algebra $\Lambda(\sigma, d)$ whose restricted universal Massey product

$$j^* \{m_{d+2}\} \in \mathrm{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d)) = \mathrm{Ext}_{\Lambda^e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$$

is represented by an exact sequence $\overline{\mathcal{A}}$ of whose middle terms are projective-injective Λ -bimodules. Secondly, let $\overline{\mathcal{A}} \subseteq \mathrm{D}(A)$ be the smallest subcategory containing A that is closed under finite direct sums and direct summands, and $\mathcal{A} \subseteq A_\infty\text{-Mod}(A)$ the full DG subcategory spanned by the objects of $\overline{\mathcal{A}}$. By construction,

- $H^\bullet(\mathcal{A})$ and $\Lambda(\sigma, d)$ are graded Morita equivalent (indeed, the graded category $H^\bullet(\mathcal{A})$ identifies with that of projective graded $\Lambda(\sigma, d)$ -modules) and
- the induced diagram of equivalences of categories

$$\begin{array}{ccc} H^0(\mathcal{A}) & \xrightarrow{\sim} & \mathrm{proj}(\Lambda) \\ [d] \downarrow & & \downarrow - \otimes_{\Lambda} \sigma \Lambda_1 \\ H^0(\mathcal{A}) & \xrightarrow{\sim} & \mathrm{proj}(\Lambda) \end{array}$$

commutes up to natural isomorphism. It follows that there is a commutative diagram

$$\begin{array}{ccc} \mathrm{HH}^{\bullet, *}(H^\bullet(\mathcal{A}), H^\bullet(\mathcal{A})) & \xrightarrow{j^*} & \mathrm{HH}^{\bullet, *}(H^0(\mathcal{A}), H^\bullet(\mathcal{A})) \\ \downarrow \sim & & \downarrow \sim \\ \mathrm{HH}^{\bullet, *}(\Lambda(\sigma, d), \Lambda(\sigma, d)) & \xrightarrow{j^*} & \mathrm{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d)) \end{array}$$

where the vertical maps are isomorphisms of bigraded algebras and the horizontal maps are induced by the corresponding inclusion of the degree 0 part. In particular, the restricted universal Massey product

$$j^* \{m_{d+2}\} \in \mathrm{HH}^{d+2, -d}(H^0(\mathcal{A}), H^\bullet(\mathcal{A}))$$

is also unit. Consequently, the DG category \mathcal{A} is Karoubian pre- $(d+2)$ -angulated ([Corollary 4.5.17 \(1\) \$\Leftarrow\$ \(3\)](#)) and the standard $(d+2)$ -angulation on $(H^0(\mathcal{A}), [d])$ coincides with the AL $(d+2)$ -angulation induced by any representative of $(-1)^{\sum_{i=1}^{d+2} i} j^* \{m_{d+2}\}$ with projective-injective middle terms ([Theorem 4.5.8\(1\)](#)). Finally, by construction, \mathcal{A} is an enhancement of the AL $(d+2)$ -angulation of the pair $(\mathrm{proj}(\Lambda), \Sigma)$, which proves the first claim.

We prove the second claim, that any two enhancements of the AL $(d+2)$ -angulation on $(\mathrm{proj}(\Lambda), \Sigma)$ are equivalent under the separability hypothesis. Indeed, suppose that \mathcal{A} and \mathcal{B} are two such enhancements. By assumption, we may identify the graded endomorphism algebras of basic additive generators $c \in H^0(\mathcal{A})$ and $c' \in H^0(\mathcal{B})$ with the d -sparse graded algebra $\Lambda(\sigma, d)$. It follows that any choice of minimal A_∞ -models on the graded categories $H^\bullet(\mathcal{A})$ and $H^\bullet(\mathcal{B})$ yield minimal A_∞ -algebra structures

$$(\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) \quad \text{and} \quad (\Lambda(\sigma, d), m'_{d+2}, m'_{2d+2}, \dots)$$

on $\Lambda(\sigma, d)$. Moreover, as a consequence of [Corollary 4.5.17 \(1\) \$\Rightarrow\$ \(3\)](#), which requires the separability hypothesis on Λ/J_Λ , we see that the restricted universal Massey products $j^* \{m_{d+2}\}$ and $j^* \{m'_{d+2}\}$ are units in $\mathrm{HH}^{\bullet, *}(\Lambda, \Lambda(\sigma, d))$. Thus, by [Theorem 5.1.2\(2\)](#), there exist a graded algebra automorphism $g \in \mathrm{Aut}(\Lambda(\sigma, d))$ and a quasi-isomorphism of A_∞ -algebras

$$F: (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) \xrightarrow{\sim} (\Lambda(\sigma, d), m'_{d+2}, m'_{2d+2}, \dots) * g$$

such that $F_1 = \mathbf{1}$. It is straightforward to verify that the tuple

$$(gF_1, gF_2, gF_3, \dots): (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) \xrightarrow{\sim} (\Lambda(\sigma, d), m'_{d+2}, m'_{2d+2}, \dots)$$

is also quasi-isomorphism of A_∞ -algebras. By well-known rectification results for A_∞ -algebras [[LV12](#), Sec. 11.4.3], the existence of the latter quasi-isomorphism of A_∞ -algebras implies that the DG algebras $A = \mathcal{A}(c, c)$ and $B = \mathcal{B}(c', c')$ are quasi-isomorphic. However the quasi-isomorphism might not be direct, in the sense that it might be given by zig-zag through intermediate DG algebras,

$$A = A^{(0)} \xleftarrow{\sim} A^{(1)} \xrightarrow{\sim} \dots \xleftarrow{\sim} A^{(n-1)} \xrightarrow{\sim} A^{(n)} = B.$$

Moreover, the quasi-isomorphisms in the zig-zag might not preserve the unit, and the intermediate DG algebras might not even be unital. Notwithstanding, since both A and B are unital DG algebras, everything can be made unital using the results in [[Mur14](#), Prop. 6.2]. Finally, any unital quasi-isomorphism between A and B can be extended to a quasi-equivalence between \mathcal{A} and \mathcal{B} in an essentially unique way (this is a well-know property of the homotopy Karoubi envelope of a DG algebra, see for example [[GHW21](#), Sec. 4]),

$$\mathcal{A} = \mathcal{A}^{(0)} \xleftarrow{\sim} \mathcal{A}^{(1)} \xrightarrow{\sim} \dots \xleftarrow{\sim} \mathcal{A}^{(n-1)} \xrightarrow{\sim} \mathcal{A}^{(n)} = \mathcal{B}.$$

This finishes the proof of the theorem. \square

Remark 5.1.4. [Theorems 4.5.8](#) and [5.1.3](#) show that if Λ is a finite-dimensional self-injective algebra that its twisted $(d+2)$ -periodic and such that Λ/J_Λ is separable, then the algebraic and the AL $(d+2)$ -angulated structures on $\mathrm{proj}(\Lambda)$ coincide, see also [Remark 2.2.14](#).

Remark 5.1.5. The use of derived categories of A_∞ -algebras in the proof of [Theorem 5.1.3](#) is not essential; these are only used for constructing an explicit DG algebra that is quasi-isomorphic to a given A_∞ -algebra, and any other choice of such a DG algebra would suffice.

Theorem 5.1.6. *Let $d \geq 1$. For $i = 1, 2$, let \mathcal{T}_i be an algebraic triangulated category with finite-dimensional morphism spaces and split idempotents, and $c_i \in \mathcal{T}_i$ a basic $d\mathbb{Z}$ -cluster tilting object. Assume that the algebras $\mathcal{T}(c_1, c_1)$ and $\mathcal{T}(c_2, c_2)$ are isomorphic and write Λ for a representative of their isomorphism class. Suppose, moreover, that the algebra Λ/J_Λ is separable and that there exists a Λ -bimodule I such the diagram*

$$\begin{array}{ccc} \mathcal{T}_i \supset \mathbf{add}(c_i) & \xrightarrow{\mathcal{T}(c_i, -)} & \mathbf{proj}(\Lambda) \\ \downarrow [-d] & & \downarrow -\otimes_\Lambda I \\ \mathcal{T}_i \supset \mathbf{add}(c_i) & \xrightarrow{\mathcal{T}(c_i, -)} & \mathbf{proj}(\Lambda) \end{array}$$

commutes up to natural isomorphism, for $i = 1, 2$. Then, there exists an equivalence of triangulated categories

$$F: \mathcal{T}_1 \xrightarrow{\sim} \mathcal{T}_2$$

that restricts to an equivalence

$$F: \mathbf{add}(c_1) \xrightarrow{\sim} \mathbf{add}(c_2).$$

Proof. The algebra Λ is finite-dimensional, self-injective and twisted $(d+2)$ -periodic by [Proposition 2.2.10](#), and the Λ -bimodule I is invertible because the left vertical and the horizontal arrows in the commutative square of the statement are equivalences. Moreover, Λ is basic since the c_i are, so $I \cong {}_1\Lambda_\sigma$ for a some automorphism σ of Λ by [[Bol84](#), Prop. 3.8]. Furthermore, σ realises the $(d+2)$ -periodicity of Λ by [Theorem 2.2.1](#) and [Proposition 2.1.8\(3\)](#). Compare the hypotheses of [Theorem 5.1.3](#).

Let \mathcal{B}_i , $i = 1, 2$, be pre-triangulated enhancements of \mathcal{T}_i , $i = 1, 2$, so that there are equivalences of triangulated categories

$$\mathcal{T}_i \simeq H^0(\mathcal{B}_i) \quad i = 1, 2.$$

Set $\mathcal{C}_i := \mathbf{add}(c_i)$, $i = 1, 2$, and let $\mathcal{A}_i \subseteq \mathcal{B}_i$, $i = 1, 2$, be the full DG subcategories spanned by the objects in \mathcal{C}_i , $i = 1, 2$. In particular, since $\mathbf{thick}(\mathcal{C}_i) = \mathcal{T}_i$ (see [Theorem 1.3.7](#)), the inclusions $\mathcal{A}_i \subseteq \mathcal{B}_i$, $i = 1, 2$, are Morita equivalences of DG categories and, consequently, there are canonical equivalences of triangulated categories

$$\mathbf{D}^c(\mathcal{A}_i) \xrightarrow{\sim} \mathbf{D}^c(\mathcal{B}_i) \xleftarrow{\sim} H^0(\mathcal{B}_i) \quad i = 1, 2,$$

see [Theorem 3.1.6](#). Next, by construction, \mathcal{A}_i , $i = 1, 2$, yields a pre- $(d+2)$ -angulated enhancements of

$$(\mathcal{C}_i, [d]_{\mathcal{T}_i}) \simeq (\mathbf{proj}(\Lambda), -\otimes_\Lambda I^{-1}), \quad i = 1, 2,$$

where the pair of the left-hand side is equipped with the standard $(d+2)$ -angulation as in [Theorem 2.2.1](#). We may apply [Theorem 5.1.3](#) (see also [Remark 5.1.4](#)) to deduce that \mathcal{A}_1 and \mathcal{A}_2 are isomorphic as objects of the homotopy category $\mathbf{Ho}(\mathbf{dgc})$. Thus, \mathcal{A}_1 and \mathcal{A}_2 are connected by a finite zig-zag of quasi-equivalences of DG categories

$$(5.1.7) \quad \mathcal{A}_1 = \mathcal{A}^{(0)} \xleftarrow{\sim} \mathcal{A}^{(1)} \xrightarrow{\sim} \dots \xleftarrow{\sim} \mathcal{A}^{(n-1)} \xrightarrow{\sim} \mathcal{A}^{(n)} = \mathcal{A}_2$$

that extends to a finite zig-zag of equivalences of triangulated categories ([Theorem 3.1.6](#))

$$\mathbf{D}^c(\mathcal{A}_1) = \mathbf{D}^c(\mathcal{A}^{(0)}) \xleftarrow{\sim} \mathbf{D}^c(\mathcal{A}^{(1)}) \xrightarrow{\sim} \dots \xleftarrow{\sim} \mathbf{D}^c(\mathcal{A}^{(n-1)}) \xrightarrow{\sim} \mathbf{D}^c(\mathcal{A}^{(n)}) = \mathbf{D}^c(\mathcal{A}_2).$$

Choosing quasi-inverses of the left-pointing functors in the latter zig-zag yields an equivalence of triangulated categories

$$H^0(\mathcal{B}_1) \simeq D^c(\mathcal{A}_1) \xrightarrow{\sim} D^c(\mathcal{A}_2) \simeq H^0(\mathcal{B}_2)$$

which restricts to an equivalence

$$\mathcal{C}_1 \simeq H^0(\mathcal{A}_1) \xrightarrow{\sim} H^0(\mathcal{A}_2) \simeq \mathcal{C}_2,$$

which is what we needed to prove. \square

Theorem 5.1.8. *Let \mathcal{T} be an algebraic triangulated category with finite-dimensional morphism spaces and split idempotents, and $c \in \mathcal{T}$ a basic $d\mathbb{Z}$ -cluster tilting object. Set $\Lambda = \mathcal{T}(c, c)$ and suppose that the algebra Λ/J_Λ is separable. Then \mathcal{T} has a unique enhancement.*

Proof. The argument is identical to the proof of [Theorem 5.1.6](#), but choosing enhancements \mathcal{B}_i of $\mathcal{T}_i = \mathcal{T}$, $i = 1, 2$ and with $c_1 = c_2 = c$. Indeed, letting $\mathcal{A}_i \subseteq \mathcal{B}_i$, $i = 1, 2$, as above, by [Remark 3.1.5](#) the zig-zag of quasi-equivalences [\(5.1.7\)](#) shows that \mathcal{A}_1 and \mathcal{A}_2 are Morita equivalent DG categories, but then so are the pre-triangulated DG categories \mathcal{B}_1 and \mathcal{B}_2 . This finishes the proof of the theorem. \square

Theorem 5.1.9. *Let $d \geq 1$. Let Λ be a basic finite-dimensional algebra that is twisted $(d + 2)$ -periodic with respect to an algebra automorphism σ . Then, there exists an algebraic triangulated category \mathcal{T} with finite-dimensional morphism spaces, split idempotents, and a $d\mathbb{Z}$ -cluster tilting object such that $\mathcal{T}(c, c) \cong \Lambda$ and the square*

$$\begin{array}{ccc} \mathcal{T} \supset \mathbf{add}(c) & \xrightarrow{\mathcal{T}(c, -)} & \mathbf{proj}(\Lambda) \\ \downarrow [-d] & & \downarrow -\otimes_{\Lambda} \sigma \\ \mathcal{T} \supset \mathbf{add}(c) & \xrightarrow{\mathcal{T}(c, -)} & \mathbf{proj}(\Lambda) \end{array}$$

commutes up to natural isomorphism.

Proof. [Theorem 2.2.15](#) shows that the pair $(\mathbf{proj}(\Lambda), -\otimes_{\Lambda} \sigma \Lambda_1)$ can be endowed with an AL $(d + 2)$ -angulation, and the latter admits an enhancement \mathcal{A} by [Theorem 5.1.3](#). Let $\mathcal{T} = D^c(\mathcal{A})$, which is an algebraic triangulated category with split idempotents, and let $c \in \mathcal{T}$ be the image of Λ under the identification

$$\mathbf{proj}(\Lambda) \simeq H^0(\mathcal{A}),$$

so that $\mathcal{T}(c, c) \cong \Lambda$. Since the canonical inclusion functor identifies $H^0(\mathcal{A})$ with a $d\mathbb{Z}$ -cluster tilting subcategory $\mathcal{C} \subseteq D^c(\mathcal{A})$ ([Proposition 3.2.4](#)), it follows that \mathcal{T} has finite-dimensional morphism spaces and $\mathbf{proj}(\Lambda) \simeq H^0(\mathcal{A})$ is equivalent to the standard $(d + 2)$ -angulated category $\mathcal{C} \subseteq D^c(\mathcal{A})$, so the square in the statement commutes. \square

We now use [Theorems 5.1.6](#), [5.1.8](#) and [5.1.9](#) to prove [Theorem A](#).

Proof of Theorem A. We analyse each correspondence separately.

The map $(\mathcal{T}, c) \mapsto (\mathcal{T}(c, c), \mathcal{T}(c, c[-d]))$ is well defined. Let (\mathcal{T}, c) be a pair satisfying the conditions in [Theorem A\(2\)](#), so that $c \in \mathcal{T}$ is a $d\mathbb{Z}$ -cluster tilting object with finite-dimensional endomorphism algebra. Then, [Proposition 2.2.10](#) shows that $\Lambda = \mathcal{T}(c, c)$ is self-injective and twisted $(d+2)$ -periodic (here we use the assumption of the ground field \mathbf{k} being perfect). In particular, as noted in [Remark 2.2.12](#), the map

$$(\mathcal{T}, c) \longmapsto (\mathcal{T}(c, c), \mathcal{T}(c, c[-d]))$$

in [Theorem A](#) is well defined since the Λ -bimodule $I = \mathcal{T}(c, c[-d])$ renders the diagram

$$\begin{array}{ccc} \mathrm{add}(c) & \xrightarrow{\mathcal{T}(c, -)} & \mathrm{proj}(\Lambda) \\ [-d] \downarrow & & \downarrow -\otimes_{\Lambda} I \\ \mathrm{add}(c) & \xrightarrow{\mathcal{T}(c, -)} & \mathrm{proj}(\Lambda) \end{array}$$

commutative up to natural isomorphism (compare with the proof of [Proposition 2.2.9](#)).

The map $(\mathcal{T}, c) \mapsto (\mathcal{T}(c, c), \mathcal{T}(c, c[-d]))$ is surjective. Let Λ be a basic finite-dimensional algebra that is twisted $(d+2)$ -periodic and I an invertible Λ -bimodule isomorphic to $\Omega_{\Lambda^e}^{d+2}(\Lambda)$ in the stable category of Λ -bimodules. Then $I \cong {}_1\Lambda_{\sigma}$ for a certain automorphism σ of Λ by [[Bol84](#), Prop. 3.8]. Furthermore, σ realises the $(d+2)$ -periodicity of Λ by the stable isomorphism $I \cong \Omega_{\Lambda^e}^{d+2}(\Lambda)$. Hence [Theorem 5.1.9](#) applies.

The map $(\mathcal{T}, c) \mapsto (\mathcal{T}(c, c), \mathcal{T}(c, c[-d]))$ is injective. This follows from [Theorem 5.1.6](#). Here Λ/J_{Λ} is separable for $\Lambda = \mathcal{T}(c, c)$ because the ground field \mathbf{k} is perfect.

The map $A \mapsto (\mathrm{D}^c(A), A)$ is surjective. Indeed, given an algebraic triangulated category \mathcal{T} with a basic $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$, any choice of enhancement \mathcal{A} of \mathcal{T} yields a DG algebra A together with an equivalence of triangulated categories

$$\mathrm{D}^c(A) \xrightarrow{\sim} \mathcal{T}, \quad A \mapsto c.$$

Explicitly, by assumption there exists an equivalence of triangulated categories $\mathcal{T} \simeq H^0(\mathcal{A})$; identify \mathcal{T} with $H^0(\mathcal{A})$ along this equivalence and let $A := \mathcal{A}(c, c)$ be the DG algebra of endomorphisms of c viewed as an object of \mathcal{A} . Since $d\mathbb{Z}$ -cluster tilting objects are in particular classical generators of the ambient triangulated category ([Theorem 1.3.7](#)), [Theorem 3.1.6](#) yields an equivalence of triangulated categories

$$\mathrm{D}^c(A) \xrightarrow{\sim} H^0(\mathcal{A}), \quad A \mapsto c,$$

and the claim follows.

The map $A \mapsto (\mathrm{D}^c(A), A)$ is injective. The injectivity of the map is a direct consequence of [Theorem 5.1.2\(2\)](#). Indeed, let A and B be DG algebras satisfying the conditions in (1) and suppose that the pairs $(\mathrm{D}^c(A), A)$ and $(\mathrm{D}^c(B), B)$ are equivalent; explicitly, this means that there exists an equivalence of triangulated categories

$$\mathrm{D}^c(A) \xrightarrow{\sim} \mathrm{D}^c(B), \quad A \mapsto B;$$

here we use that A and B are assumed to be basic objects of $\mathrm{D}^c(A)$. In particular, we may and we will identify the graded algebras $H^{\bullet}(A)$ and $H^{\bullet}(B)$ via this equivalence. Choose minimal A_{∞} -models

$$(H^{\bullet}(A), m_{d+2}^A, m_{2d+2}^A, \dots) \text{ and } (H^{\bullet}(B), m_{d+2}^B, m_{2d+2}^B, \dots).$$

of A and B , respectively. By [Corollary 4.5.17](#), that the restricted universal Massey products $j^*\{m_{d+2}^A\}$ and $j^*\{m_{d+2}^B\}$ are edge units, and [Theorem 5.1.2\(2\)](#) shows that, up to quasi-isomorphism of A_{∞} -algebras, there is a unique A_{∞} -structure on $H^{\bullet}(A) = H^{\bullet}(B)$ with this property. Consequently, A and B are quasi-isomorphic DG algebras (compare with the proof of [Theorem 5.1.3](#)). This finishes the proof of the theorem.

Uniqueness of enhancements. Finally, we need to show that the triangulated categories that satisfy the conditions in [Theorem 5.1.3\(2\)](#) admit a unique enhancement. This follows from [Theorem 5.1.8](#). \square

5.2. Proof of Theorem B. Let A be a graded vector space. Essentially by definition, a minimal A_∞ -algebra structure on A can be identified with a point in the mapping space (a simplicial set)

$$\mathrm{Map}(A_\infty, \mathrm{End}(A)) = \mathrm{Map}_{\mathrm{dgOp}}(A_\infty, \mathrm{End}(A))$$

of operadic maps from the A_∞ -operad A_∞ to the endomorphism operad $\mathrm{End}(A)$ of A viewed as a cochain complex with vanishing differential, computed in the model category dgOp of differential graded (non-symmetric) operads [Lyu11, Mur11]. By [Fre09, Thm. 5.2.1] or [Mur16, Cor. 2.3], two points in the space $\mathrm{Map}(A_\infty, \mathrm{End}(A))$ lie in the same path-connected component if and only if the corresponding minimal A_∞ -structures are related by a quasi-isomorphism of A_∞ -algebras F with $F_0 = \mathbf{1}$ and $F_1 = \mathbf{1}$ (compare with Theorem B). Moreover, the mapping space $\mathrm{Map}(A_\infty, \mathrm{End}(A))$ is the homotopy limit of the tower of fibrations

$$\cdots \rightarrow \mathrm{Map}(A_{i+2}, \mathrm{End}(A)) \rightarrow \cdots \rightarrow \mathrm{Map}(A_3, \mathrm{End}(A)) \rightarrow \mathrm{Map}(A_2, \mathrm{End}(A)),$$

where A_n is the operad of A_n -algebra structures.⁵ Thus, a minimal A_∞ -algebra structure $x_\infty \in \mathrm{Map}(A_\infty, \mathrm{End}(A))$ on A yields minimal A_{i+2} -algebra structures

$$x_i \in \mathrm{Map}(A_{i+2}, \mathrm{End}(A)), \quad i \geq 0,$$

by restriction. To simplify the notation, we write

$$X_i := \mathrm{Map}(A_{i+2}, \mathrm{End}(A)), \quad i \geq 0,$$

so that we have a tower of *pointed* fibrations

$$\cdots \rightarrow X_{i+2} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0.$$

The above perspective on A_∞ -structures is advantageous in that it enables us to apply robust techniques from homotopy theory to compute the set of path-connected components of the homotopy limit $\mathrm{Map}(A_\infty, \mathrm{End}(A))$. Indeed, suppose that A is a d -sparse graded algebra. In this case, the results in [Mur20a] provide an extension of the classical Bousfield–Kan (fringed) spectral sequence [BK72, Ch. IX, Sec. 4] with useful computational properties, of which we highlight the following:

- The classical Bousfield–Kan spectral sequence is only defined on the upper-half of the bisection of the first quadrant, that is $E_r^{s,t}$ is defined for $t \geq s \geq 0$. In contrast, the extension of the Bousfield–Kan spectral sequence is defined in most of the right half-plane; however, the extended region of definition ‘converges’ to that of the classical Bousfield–Kan spectral sequence as $r \rightarrow \infty$, see Figure 1 and [Mur20a] for details.
- The extended Bousfield–Kan spectral sequence is partially-defined even if we only have a minimal A_{i+2} -structure on A , so just the bottom part of the tower of fibrations is pointed. More precisely, the extended spectral sequence is defined up to the terms of page $\lfloor \frac{i+3}{2} \rfloor$, see [Mur20a, Def. 5.1].
- Suppose given a minimal A_{i+2} -algebra structure on A , that is a base point

$$x_i \in X_i = \mathrm{Map}(A_{i+2}, \mathrm{End}(A)).$$

The obstruction to the existence of a minimal A_{i+3} -algebra structure

$$x_{i+1} \in x_{i+1} = \mathrm{Map}(A_{i+3}, \mathrm{End}(A))$$

⁵Recall that an A_n -algebra has operations

$$m_i: A^{\otimes i} \rightarrow A(2-i), \quad 1 \leq i \leq n,$$

that satisfy all the A_∞ -equations that involve only these operations; in particular, an A_∞ -algebra yields an A_n -algebra by discarding the operations m_i , $i > n$. Notice also that there is no condition of the n -ary multiplication in an A_n -algebra when $m_1 = 0$, that is when A is minimal. For example, a minimal A_3 -algebra consists of an *associative* graded algebra with an *arbitrary* ternary operation.




FIGURE 1. Range of definition of the extended Bousfield–Kan spectral sequence ($r = 5$). The red region is where the classical Bousfield–Kan spectral sequence is defined; the extended spectral sequence is defined also in the blue region.

such that x_i and x_{i+1} have the same underlying minimal A_{i-r+3} -algebra structure

$$x_{i-r+1} \in X_{i-r+1} = \text{Map}(A_{i-r+3}, \text{End}(A))$$

lies in the term $E_r^{i+1, i}$, $1 \leq r \leq \lfloor \frac{i+3}{2} \rfloor$. This obstruction vanishes precisely when such a minimal A_{i+3} -algebra structure exists, see [Mur20a, Prop. 5.4].

- The $E_2^{\bullet, *}$ -terms of the extended Bousfield–Kan spectral sequence are given by [Mur20a, Cor. 6.3]

$$E_2^{s, t} = \begin{cases} \text{HH}^{s+2, -t}(A) & s > 0, t \in \mathbb{Z}; \\ \text{Z}^{2, -t}(\mathbf{C}^{\bullet, *}(A)) & s = 0, t > 0. \end{cases}$$

In particular, since the graded algebra A is d -sparse, $E_2^{s, t} = 0$ if $t \notin d\mathbb{Z}$. Therefore, the spectral sequence differentials $\partial_i = 0$ are necessarily trivial for $2 \leq i \leq d$ and $E_{d+1}^{s, t} = E_2^{s, t}$ wherever they are defined.

Remark 5.2.1. By construction, the spectral sequence described above is related to the homotopy groups of the space $\text{Map}(A_\infty, \text{End}(A))$, which turn out to be computed in terms of Hochschild cohomology of A_∞ -algebras, see [Mur20a, Prop. 6.9] for a precise statement. In particular, notice that the spectral sequence above is different from those usually considered in the context of deformation problems, such as the one discussed in [Sei15, Ch. 3]. Seidel’s spectral sequence is a particular case of the third sequence in Elmendorf–Kriz–Mandell–May’s [EKMM07, Thm. 1.6] in the context of ‘brave new algebra.’ For the relation between these spectral sequences and ours, see [Ang11, Sec. 2] and [Mur20a, Sec. 7].

A minimal A_{i+2} -algebra structure on A has an m_{d+2} operation for any $i \geq d$, but we need $i > d$ to ensure that m_{d+2} is a cocycle, whose cohomology class

$$\{m_{d+2}\} \in \text{HH}^{d+2, -d}(A)$$

also deserves to be called *universal Massey product of length $d + 2$* , and $i > 2d$ so that

$$\mathrm{Sq}(\{m_{d+2}\}) = 0.$$

Proposition 5.2.2. *Let A be a d -sparse graded algebra equipped with a minimal A_{2d+3} -algebra structure, so the spectral sequence is defined up to the $E_{d+2}^{\bullet,*}$ terms. The spectral sequence differential*

$$\partial_{d+1}: E_{d+1}^{s,t} \longrightarrow E_{d+1}^{s+d+1,t+d}$$

is given by

$$\partial_{d+1}(x) = \pm[\{m_{d+2}\}, x]$$

wherever it is defined, except for $(s, t) = (d - 1, d)$ in $\mathrm{char}(\mathbf{k}) = 2$, where it is given by

$$\partial_{d+1}(x) = [\{m_{d+2}\}, x] + x^2.$$

Remark 5.2.3. The differential defined in [Proposition 5.2.2](#) is the first possibly non-trivial spectral sequence differential beyond page 1. Notice also that $x^2 = 0$ in $\mathrm{char}(\mathbf{k}) \neq 2$ by graded commutativity.

Proof of [Proposition 5.2.2](#). The general formula follows from [[Mur20a](#), Lem. 6.1] by the same argument as in the proof of [[Mur20a](#), Thm. 6.5], which is the case $d = 1$. Condition $m \leq s + 1$ in the statement of [[Mur20a](#), Lem. 6.1] is actually unnecessary. Indeed, as indicated therein, this lemma follows directly from [[Mur20a](#), Prop. 3.5] where no upper bound is required. In our current context, where A is d -sparse, [[Mur20a](#), Prop. 3.5] is used for $r = s + 1$, $s = d + 1$ and $t = 1$ and only the first and last summands ($p = 2, d + 2$) produce possibly non-vanishing elements since $m_{i+2} = 0$ for $0 < i < d$. Hence the proof goes through as for [[Mur20a](#), Thm. 6.5].

In the special case $(s, t) = (d - 1, d)$ we must invoke [[Mur20a](#), Cors. 3.9,3.10] instead, as in the proof of [[Mur20a](#), Thm. 6.5], here for $r = 2d + 1$ and $s = 1$. \square

Remark 5.2.4. For $s = 0$ and $t > 0$, $x \in E_2^{0,t} = Z^{2,-t}(\mathbf{C}^{\bullet,*}(A))$ is a cocycle. Hence, on the right hand side of the formula for $\partial_{d+1}(x)$ in [Proposition 5.2.2](#), we understand that x stands for its Hochschild cohomology class.

Definition 5.2.5. A *d -sparse Massey algebra (A, m)* is a pair given by a d -sparse graded algebra A and a Hochschild cohomology class

$$m \in \mathrm{HH}^{d+2,-d}(A)$$

with trivial Gerstenhaber square

$$\mathrm{Sq}(m) = 0.$$

The *Hochschild complex* of a d -sparse Massey algebra is the bigraded vector space $\mathbf{C}^{\bullet,*}(A, m)$ defined as $\mathrm{HH}^{\bullet,*}(A)$ in horizontal degrees $\bullet \geq 2$ and zero otherwise, with bidegree $(d + 1, -d)$ differential

$$\begin{aligned} \mathrm{HH}^{s,t}(A) &\longrightarrow \mathrm{HH}^{s+d+1,t-d}(A), \\ x &\longmapsto [m, x], \end{aligned}$$

except in bidegree $(d + 1, -d)$, where it is given by

$$\begin{aligned} \mathrm{HH}^{d+1,-d}(A) &\longrightarrow \mathrm{HH}^{2(d+1),-2d}(A), \\ x &\longmapsto [m, x] + x^2. \end{aligned}$$

The cohomology of this complex is the *Hochschild cohomology* of the d -sparse Massey algebra, and it is denoted by $\mathrm{HH}^{\bullet,*}(A, m)$.

Remark 5.2.6. The Hochschild complex of a d -sparse Massey algebra is indeed a complex since

$$[m, [m, x]] = [\text{Sq}(m), x] = 0$$

by the Gerstenhaber algebra relations. Moreover, if x is in bidegree $(d+1, -d)$

$$[m, [m, x] + x^2] = [m, [m, x]] + [m, x^2] = [\text{Sq}(m), x] + [m, x] \cdot x - x \cdot [m, x] = 0.$$

Notice also that $x^2 = 0$ in $\text{char}(k) \neq 2$ by graded commutativity.

Remark 5.2.7. Let A be a d -sparse graded algebra equipped with a minimal A_{2d+3} -algebra structure. The terms $E_{d+1}^{s,t}$ of the (truncated) extended Bousfield–Kan spectral sequence constructed in [Mur20a] are defined for $t \geq s$ and for $s \geq 2d-1$ and $t \in \mathbb{Z}$. Moreover, the differential

$$\partial_{d+1}: E_{d+1}^{s,t} \longrightarrow E_{d+1}^{s+d+1,t+d}$$

is defined except for $0 \leq s = t \leq d$. Furthermore, $E_{d+2}^{s,t}$ is given by the cohomology of $(E_{d+1}^{\bullet,*}, \partial_{d+1})$ whenever the incoming and outgoing differentials are defined on $E_{d+1}^{s,t}$. Notice that the assumptions on A imply that $(A, \{m_{d+2}\})$ is a d -sparse Massey algebra. Moreover, [Proposition 5.2.2](#) shows that $(E_{d+1}^{\bullet,*}, \partial_{d+1})$ surjects onto $\mathbf{C}^{\bullet+2,-*}(A, \{m_{d+2}\})$ whenever the source is defined, and this surjection is actually an isomorphism for $\bullet > 0$. Therefore, we deduce that

$$E_{d+2}^{s,t} \cong \text{HH}^{s+2,-t}(A, \{m_{d+2}\})$$

in the following cases:

- $t > s$,
- $t = s > d$,
- $s \geq 3d, t \in \mathbb{Z}$.

We are ready to prove [Theorem B](#).

Proof of Theorem B. Let A be a d -sparse graded algebra and

$$B = (A, m_{d+2}^B, m_{2d+2}^B, m_{3d+2}^B, \dots) \quad \text{and} \quad C = (A, m_{d+2}^C, m_{2d+2}^C, m_{3d+2}^C, \dots)$$

minimal A_∞ -algebra structures on A , which we identify with points

$$x_\infty, y_\infty \in X_\infty = \text{Map}(A_\infty, \text{End}(A)),$$

respectively. Suppose that there is an equality of universal Massey products

$$m = \{m_{d+2}^B\} = \{m_{d+2}^C\} \in \text{HH}^{d+2,-d}(A)$$

and that

$$\text{HH}^{p+2,-p}(A, m) = 0, \quad p > d.$$

As explained at the beginning of this subsection, we need to prove that x_∞ and y_∞ lie in the same connected component of X_∞ . We take $x_\infty \in X_\infty$ as a base point for the definition of the spectral sequence in [Mur20a]. By [Remark 5.2.7](#)

$$E_{d+2}^{p,p} = \text{HH}^{p+2,-p}(A, m) = 0, \quad p > d,$$

therefore

$$E_r^{p,p} = 0, \quad p > d, \quad d+2 \leq r \leq \infty.$$

Denote by $x_i, y_i \in X_i$ the images of $x_\infty, y_\infty \in X_\infty$ along the restriction map $X_\infty \rightarrow X_i$. We have chosen x_i as the base point of X_i , $0 \leq i < \infty$, hence $\pi_0(X_i)$ is a set pointed by $[x_i]$. We will inductively show that $[x_i] = [y_i] \in \pi_0(X_i)$ for $0 \leq i < \infty$.

($i = 0$) Since A has trivial differential, the set $\pi_0(X_0)$ is in bijection with the set of degree 0 maps of graded vector spaces $A \otimes A \rightarrow A$, and the class $[x_0]$ corresponds to the product operation of A ; the class $[y_0]$ also corresponds to the product operation

of A since A is the underlying graded algebra of both minimal A_∞ -algebra structures B and C . We distinguish between the various natural cases:

($0 < i < d$) Since the graded algebra A is d -sparse, in this range the set $\pi_0(X_i)$ is in bijection with the set of graded associative algebra structures on the underlying graded vector space of A , and both $[x_i]$ and $[y_i]$ correspond the product of A by the same reason as above.

($i = d$) The previous case shows, in particular, that $[x_{d-1}] = [y_{d-1}]$. Therefore the classes $[x_d]$ and $[y_d]$ lie in

$$\ker(\pi_0 X_d \rightarrow \pi_0 X_{d-1}) \cap \text{im}(\pi_0 X_{2d} \rightarrow \pi_0 X_d)$$

and, according to [Mur20a, Eqn. 4.8], this intersection is in fact equal to

$$E_{d+1}^{d,d} = E_2^{d,d} = \text{HH}^{d+2,-d}(A).$$

The elements in Hochschild cohomology corresponding to the classes $[x_d]$ and $[y_d]$ are the universal Massey products $\{m_{d+2}^B\} = \{m_{d+2}^C\}$, which are equal by hypothesis and hence we obtain the required equality $[x_d] = [y_d]$.

($d < i < \infty$) As in the previous case, the classes $[x_i]$ and $[y_i]$ lie in the intersection

$$\ker(\pi_0 X_i \rightarrow \pi_0 X_{i-1}) \cap \text{im}(\pi_0 X_{2i} \rightarrow \pi_0 X_i) = E_{i+1}^{i,i} = 0,$$

see [Mur20a, Eqn. 4.8], therefore $[x_i] = [y_i]$.

($i = \infty$) We deduce the desired equality $[x_\infty] = [y_\infty]$ from the following two facts:

- [BK72, Sec. IX.3.1] There is a Milnor short exact sequence of pointed sets

$$* \longrightarrow \varprojlim^1 \pi_1(X_i) \longrightarrow \pi_0(X_\infty) \longrightarrow \varprojlim \pi_0(X_i) \longrightarrow *$$

and, by the previous discussion, the classes $[x_\infty]$ and $[y_\infty]$ have the same image under the canonical map $\pi_0(X_\infty) \rightarrow \varprojlim \pi_0(X_i)$, namely the base point of $\varprojlim \pi_0(X_i)$.

- [BK72, Sec. IX.5.4] For every $s \geq 0$, the term $E_r^{s,s+1}$ stabilises for $r \geq \max\{d+2, s+1\}$ since the source of the incoming differential is $E_r^{s-r,s-r+1} = 0$ being in the left half-plane ($s-r \leq s-(s+1) < 0$), and the target of the outgoing differential is $E_r^{s+r,s+r} = 0$ since $s+r \geq r \geq d+2 > d$. Therefore $\varprojlim^1 \pi_1(X_i) = 0$. This implies that the inverse image of the base point along the natural map of pointed sets $\pi_0(X_\infty) \rightarrow \varprojlim \pi_0(X_i)$ consists just of the source's base point, hence $[x_\infty] = [y_\infty]$ by the last observation in the previous item.

This finishes the proof of the theorem. \square

For the sake of completeness, we observe that **Theorem B** implies Kadeishvili's intrinsic formality theorem, which we state in its version for minimal A_∞ -algebras.

Corollary 5.2.8 ([Kad88], see also [ST01, Thm. 4.7]). *Let A be a graded algebra such that*

$$\text{HH}^{p+2,-p}(A, A) = 0, \quad p \geq 1.$$

If B and C are minimal A_∞ -algebras with $H^\bullet(B) = H^\bullet(C) = A$ as graded algebras, then B and C are A_∞ -isomorphic through an A_∞ -isomorphism with identity linear part. In particular, the graded algebra A is intrinsically formal.

Proof. Endow the graded algebra A with the trivial 1-sparse Massey algebra structure $(A, 0)$ and notice that, in fact, this is the only possible such structure since $\text{HH}^{3,-1}(A, A) = 0$ by assumption. We will apply **Theorem B** to $(A, 0)$. By definition, the cochain complex $C^{\bullet,*}(A, 0)$ has vanishing differential and therefore the




FIGURE 2. The $E_{d+2}^{\bullet,*}$ -page of the extended Bousfield–Kan spectral sequence can be non-trivial only in the green part ($d = 4$), see [Proposition 5.2.9](#).

Hochschild cohomology of the 1-sparse Massey algebra $(A, 0)$ coincides with the (ordinary) Hochschild cohomology of the graded algebra A . In particular,

$$\mathrm{HH}^{p+2, -p}(A, 0) = \mathrm{HH}^{p+2, -p}(A) = 0, \quad p > 1,$$

by assumption. Thus, we may apply [Theorem B](#) to the 1-sparse Massey algebra $(A, 0)$ and the first claim follows. The second claim follows by taking $C = A$ viewed as a minimal A_∞ -algebra with vanishing higher operations. \square

We record the following vanishing properties that are crucial to the proof of [Theorem 5.3.3](#) below, compare with the hypothesis in [Remark 5.2.7](#) and the assumptions in [Theorem B](#).

Proposition 5.2.9. *Let $A = \Lambda(\sigma, d)$ as in [Setting 5.0.1](#). Assume we have at least a minimal A_{2d+3} -algebra structure on A such that $\{m_{d+2}\}$ is an edge unit. Then the terms $E_{d+2}^{s,t}$ in the extended Bousfield–Kan spectral sequence associated to the tower of fibrations*

$$\cdots \rightarrow X_i \rightarrow \cdots \rightarrow X_1 \rightarrow X_0$$

can be non-zero only in the following cases ([Figure 2](#)):

- $s = 0$ and $t = 0$.
- $t = kd$, $k \geq 1$, and $0 \leq s \leq d$.
- $t = kd$, $k \leq 2$, and $2d + 1 \leq s \leq 3d - 1$.

Proof. We are going to prove that

$$\mathrm{HH}^{s,t}(A, \{m_{d+2}\}) = 0, \quad s > d + 2.$$

Hence, the proposition follows from the relation between $E_{d+2}^{\bullet,*}$ and the Hochschild cohomology of the d -sparse Massey algebra $(A, \{m_{d+2}\})$ in [Remark 5.2.7](#), and from the fact that $\mathrm{HH}^{s,t}(A, \{m_{d+2}\})$ is concentrated in $t \in d\mathbb{Z}$ since A is d -sparse.

Consider the degree $(d+2, -d)$ endomorphism of $\mathbf{C}^{\bullet,*}(A, \{m_{d+2}\})$ given by the isomorphisms

$$\begin{aligned} \mathrm{HH}^{s,t}(A) &\xrightarrow{\cong} \mathrm{HH}^{s+d+2,t-d}(A), \\ x &\longmapsto \{m_{d+2}\} \cdot x, \end{aligned}$$

if $(s, t) \neq (d+1, -d)$, see [Proposition 4.7.15](#), and by

$$\begin{aligned} \mathrm{HH}^{d+1,-d}(A) &\xrightarrow{\cong} \mathrm{HH}^{2d+3,-2d}(A), \\ x &\longmapsto \{m_{d+2}\} \cdot x + \{\delta_{/d}\} \cdot x^2. \end{aligned}$$

The latter map is also bijective. This can be checked as in the proof of [[Mur22](#), Thm. 9.1]. The remaining components of this endomorphism are maps from 0 to possibly non-trivial targets, so it not a point-wise isomorphism everywhere.

Indeed, the previous maps assemble to a cochain map because, by the Gerstenhaber algebra relations,

$$\begin{aligned} [\{m_{d+2}\}, \{m_{d+2}\} \cdot x] &= [\{m_{d+2}\}, \{m_{d+2}\}] \cdot x + \{m_{d+2}\} \cdot [\{m_{d+2}\}, x] \\ &= 2\mathrm{Sq}(\{m_{d+2}\}) \cdot x + \{m_{d+2}\} \cdot [\{m_{d+2}\}, x] \\ &= \{m_{d+2}\} \cdot [\{m_{d+2}\}, x], \end{aligned}$$

and if x has bidegree $(d+1, -1)$ then

$$\begin{aligned} [\{m_{d+2}\}, \{m_{d+2}\} \cdot x + \{\delta_{/d}\} \cdot x^2] &= [\{m_{d+2}\}, \{m_{d+2}\} \cdot x] + [\{m_{d+2}\}, \{\delta_{/d}\} \cdot x^2] \\ &= \{m_{d+2}\} \cdot [\{m_{d+2}\}, x] + [\{m_{d+2}\}, \{\delta_{/d}\}] \cdot x^2 \\ &= \{m_{d+2}\} \cdot [\{m_{d+2}\}, x] + \{m_{d+2}\} \cdot x^2 \\ &= \{m_{d+2}\} \cdot ([\{m_{d+2}\}, x] + x^2), \end{aligned}$$

see [Proposition 4.7.4](#).

Since the previous cochain endomorphism is mostly given by isomorphisms, it induces isomorphisms in most of the cohomology. Namely,

$$\mathrm{HH}^{s,t}(A, \{m_{d+2}\}) \cong \mathrm{HH}^{s+d+2,t-d}(A, \{m_{d+2}\}), \quad s > d+2,$$

because in these bidegrees the endomorphism of $\mathbf{C}^{\bullet,*}(A, \{m_{d+2}\})$ is an isomorphism on the source and target of the incoming and outgoing differentials.

Strikingly, the previous endomorphism of the complex $\mathbf{C}^{\bullet,*}(A, \{m_{d+2}\})$ is null-homotopic. The chain homotopy is given by the bidegree $(1, 0)$ maps

$$\begin{aligned} \mathrm{HH}^{s,t}(A) &\xrightarrow{\cong} \mathrm{HH}^{s+1,t}(A), \\ x &\longmapsto \{\delta_{/d}\} \cdot x. \end{aligned}$$

Indeed, by [Proposition 4.7.8](#),

$$\{m_{d+2}\} \cdot x = [\{m_{d+2}\}, \{\delta_{/d}\} \cdot x] + \{\delta_{/d}\} \cdot [\{m_{d+2}\}, x],$$

in particular if x has bidegree $(d+1, -d)$,

$$\{m_{d+2}\} \cdot x + \{\delta_{/d}\} \cdot x^2 = [\{m_{d+2}\}, \{\delta_{/d}\} \cdot x] + \{\delta_{/d}\} \cdot ([\{m_{d+2}\}, x] + x^2),$$

and if $d > 1$ and x has bidegree $(d, -d)$,

$$\begin{aligned} \{m_{d+2}\} \cdot x &= [\{m_{d+2}\}, \{\delta_{/d}\} \cdot x] + (\{\delta_{/d}\} \cdot x)^2 + \{\delta_{/d}\} \cdot [\{m_{d+2}\}, x] \\ &= [\{m_{d+2}\}, \{\delta_{/d}\} \cdot x] + \{\delta_{/d}\} \cdot [\{m_{d+2}\}, x], \end{aligned}$$

since the product is graded commutative and $\{\delta_{/d}\}^2 = 0$ by [Proposition 4.7.6](#).

Therefore, the previous isomorphisms on the Hochschild cohomology of the d -sparse Massey algebra $(A, \{m_{d+2}\})$ are trivial maps. This implies that

$$\mathrm{HH}^{s,t}(A, \{m_{d+2}\}) = 0, \quad s > d+2,$$

as desired. \square

5.3. Proof of Theorem 5.1.2. In this subsection we complete the proof of [Theorem A](#) by providing a proof of [Theorem 5.1.2](#). We remind the reader of our standing assumptions for this section ([Setting 5.0.1](#)).

Notation 5.3.1. We denote by $S_d(\Lambda, \sigma)$ the set of (minimal) A_∞ -algebra structures

$$(\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, m_{3d+2}, \dots)$$

on the d -sparse graded algebra $\Lambda(\sigma, d)$ whose restricted universal Massey product

$$j^* \{m_{d+2}\} \in \mathrm{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d)) = \mathrm{Ext}_{\Lambda_e}^{d+2}(\Lambda, {}_1\Lambda_\sigma)$$

is an edge unit. Similarly, we denote by $S_d^\sim(\Lambda, \sigma)$ the quotient of $S_d(\Lambda, \sigma)$ by the equivalence relation given by the existence of a quasi-isomorphism of A_∞ -algebras F with $F_1 = \mathbf{1}$.

Remark 5.3.2. The group $\mathrm{Aut}(\Lambda(\sigma, d))$ of graded algebra automorphisms of $\Lambda(\sigma, d)$ acts on the right on the set $S_d(\Lambda, \sigma)$ via the formula in [Notation 5.1.1](#), and this action descends to the quotient $S_d^\sim(\Lambda, \sigma)$ (compare with [Remarks 4.4.1](#) and [4.7.10](#)). Moreover, the set of path-connected components of the space $\mathrm{Map}(\mathbf{A}_\infty, \mathrm{End}(\Lambda(\sigma, d)))$ of minimal A_∞ -algebra structures on $\Lambda(\sigma, d)$ contains the quotient set $S_d^\sim(\Lambda, \sigma)$, see [[Mur20a](#), Prop. 6.9].

We shall prove the following theorem.

Theorem 5.3.3. *Suppose that Λ/J_Λ is separable. The following statements hold:*

(1) *The $\mathrm{Aut}(\Lambda(\sigma, d))$ -equivariant map*

$$\begin{aligned} \Phi: S_d^\sim(\Lambda, \sigma) &\longrightarrow \mathrm{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d)) \\ (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) &\longmapsto j^* \{m_{d+2}\} \end{aligned}$$

is injective.

(2) *The image of Φ consists precisely of the edge units in $\mathrm{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d))$.*

(3) *The set of orbits*

$$S_d^\sim(\Lambda, \sigma) / \mathrm{Aut}(\Lambda(\sigma, d))$$

is a singleton.

Proof. The map

$$\begin{aligned} \Phi: S_d^\sim(\Lambda, \sigma) &\longrightarrow \mathrm{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d)) \\ (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) &\longmapsto j^* \{m_{d+2}\} \end{aligned}$$

is clearly $\mathrm{Aut}(\Lambda(\sigma, d))$ -equivariant and, by definition, its image consists of units in the Hochschild–Tate cohomology $\mathrm{HH}^{\bullet, *}_{\bullet}(\Lambda, \Lambda(\sigma, d))$ (the set of Hochschild–Tate units in bidegree $(d+2, -d)$ is non-empty since the algebra Λ is twisted $(d+2)$ -periodic with respect to the algebra automorphism $\sigma: \Lambda \xrightarrow{\sim} \Lambda$, see [Setting 5.0.1](#) and [Remark 4.5.16](#)).

Notice also that statement (3) follows immediately from statements (1) and (2) and [Proposition 4.7.16](#): statements (1) and (2) imply that we may identify $S_d^\sim(\Lambda, \sigma) / \mathrm{Aut}(\Lambda(\sigma, d))$ with the set of orbits of edge units in $\mathrm{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d))$, and the latter set is a singleton by [Proposition 4.7.16](#). Recall that

$$X_i = \mathrm{Map}(\mathbf{A}_{i+2}, \mathrm{End}(\Lambda(\sigma, d))), \quad i \geq 0,$$

is the space of minimal A_{i+2} -algebra structures on $\Lambda(\sigma, d)$, and that we have a tower of fibrations

$$\cdots \rightarrow X_{i+2} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0$$

as well as a distinguished base point $x_0 \in X_0$ given by multiplication on $\Lambda(\sigma, d)$. As in [Section 5.2](#), we identify

$$X_\infty = \text{Map}(A_\infty, \text{End}(\Lambda(\sigma, d))),$$

with the (homotopy) limit of the above tower. In particular, we have restriction/truncation maps $X_\infty \rightarrow X_i$ for all $i \geq 0$.

Proof of statement (2): We have to prove that every edge unit is in the image of Φ . Let $x \in \text{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d))$ be an edge unit. We need to prove that there exists an A_∞ -algebra structure

$$x_\infty \in X_\infty = \text{Map}(A_\infty, \text{End}(\Lambda(\sigma, d))), \quad x_\infty \mapsto x_0,$$

whose underlying graded algebra structure is the given one ([Remark 4.5.3](#)) and whose restricted universal Massey product of length $d + 2$ is x . We make the following observations:

- The class x lies in the image of the restriction map

$$j^*: \text{HH}^{d+2, -d}(\Lambda(\sigma, d), \Lambda(\sigma, d)) \longrightarrow \text{HH}^{d+2, -d}(\Lambda, \Lambda(\sigma, d))$$

induced by the inclusion $j: \Lambda \hookrightarrow \Lambda(\sigma, d)$ of the degree 0 part. Indeed, in view of the long exact sequence described in [Proposition 4.7.11](#), it is enough to check that $\sigma_*^{-1}\sigma^*(x) = x$; this can be shown exactly as in the proof of [\[Mur22, Cor. 7.7\]](#), that is using [\[Mur22, Prop. 7.6\]](#) for $n = d + 2$, replacing σ by σ^{-1} and keeping in mind that the extension of σ^{-1} acts on $\Lambda(\sigma, d)^{-d} = \sigma^{-1}\Lambda_1$ by

$$(-1)^{-d}\sigma^{-1} = (-1)^{(d+2)^2}\sigma^{-1}.$$

- There exists a unique $u \in \text{HH}^{d+2, -d}(\Lambda(\sigma, d), \Lambda(\sigma, d))$ such that $j^*(u) = x$ and $\text{Sq}(u) = 0$. To prove the existence of such a class u , notice that if $j^*(u) = x$, then also

$$j^*(u + \{\delta_{/d}\} \cdot y) = x$$

for arbitrary $y \in \text{HH}^{d+1, -d}(\Lambda(\sigma, d), \Lambda(\sigma, d))$; this follows from the fact that Λ is concentrated in degree 0 and therefore $j^*\{\delta_{/d}\} = 0$. Arguing as in the proof of [\[Mur22, Prop. 7.9\]](#), one shows that one can choose an appropriate y such that $\text{Sq}(u + \{\delta_{/d}\} \cdot y) = 0$ (in the argument, one must replace [\[Mur22, Props. 3.6, 3.9 and 3.10\]](#) by [Proposition 4.7.8](#) and [\[Mur22, Lemma 6.8\]](#) by [Proposition 4.7.15](#)). The proof that such a pre-image u is unique is completely analogous to that of [\[Mur22, Prop. 7.10\]](#), replacing [\[Mur22, Prop. 3.6\]](#) by the last equation in [Proposition 4.7.8](#) and [\[Mur22, Lemma 6.8\]](#) by [Proposition 4.7.15](#).

- Since the graded algebra $\Lambda(\sigma, d)$ is d -sparse, tautologically, a minimal A_{d+2} -algebra structure $x_d \in X_d$ is simply a Hochschild cochain

$$m_{d+2} \in \mathbb{C}^{d+2, -d}(\Lambda(\sigma, d), \Lambda(\sigma, d)),$$

which we choose to be a representing cocycle of the unique class

$$u \in \text{HH}^{d+2, -d}(\Lambda(\sigma, d), \Lambda(\sigma, d)) \quad \text{with} \quad \text{Sq}(u) = 0 \quad \text{and} \quad j^*(u) = x$$

(compare with [Proposition 4.4.4](#)). The fact that m_{d+2} is a cocycle implies that it extends to an A_{d+3} -algebra structure on $\Lambda(\sigma, d)$ choosing m_{d+3} arbitrarily, e.g. $m_{d+3} = 0$ (this is actually the only possible choice if $d > 1$). This is immediate from the definition of an A_∞ -algebra, see [\[Kel01, Sec. 3.1\]](#). Thus, there exists a point $x_{d+1} \in X_{d+1}$ that maps to x_d under the fibration $X_{d+1} \twoheadrightarrow X_d$. Moreover, the same argument shows that we can extend the A_{d+3} -algebra structure to an A_{2d+2} -algebra structure, necessarily setting $m_{i+2} = 0$ for $d < i < 2d$ and choosing m_{2d+2}

arbitrarily, for example $m_{2d+2} = 0$. Hence we have $x_{2d} \in X_{2d}$ that maps to x_d under the fibration $X_{2d} \rightarrow X_d$. In other words, the Hochschild cocycle m_{d+2} with $\{m_{d+2}\} = u$ gives rise to a d -sparse minimal A_{d+2} -algebra $(\Lambda(\sigma, d), m_{d+2})$ which can be extended to a d -sparse minimal A_{2d+2} -algebra $(\Lambda(\sigma, d), m_{d+2}, 0)$, with all other operations being trivial due to the fact that $\Lambda(\sigma, d)$ is d -sparse. Notice, however, that $(\Lambda(\sigma, d), m_{d+2}, 0, 0, \dots)$ need not be an A_∞ -algebra since, although $\mathrm{Sq}(\{m_{d+2}\}) = \mathrm{Sq}(u) = 0$, the Gerstenhaber square of m_{d+2} need not vanish at the cochain level.

- The condition $\mathrm{Sq}(u) = 0$ is equivalent to the existence of an A_{2d+3} -algebra structure on $\Lambda(\sigma, d)$ with the same underlying A_{d+2} -algebra as above. In other words, $\mathrm{Sq}(u) = 0$ if and only if there exists a point $y_{2d+1} \in X_{2d+1}$ such that $y_{2d+1} \mapsto x_d$. This can be shown exactly as in the proof of [Mur20a, Prop. 6.7], since the relevant results [Mur20a, Props. 3.4 and 5.4] are not particular to the case $d = 1$. Indeed, in the notation of [Mur20a, Prop. 5.4] we have $k = 2d$ and $l = 2d - 1$, the recipient of the obstruction is $E_{k+1-l}^{k+1,k} = E_2^{2d+1,2d} = \mathrm{HH}^{2d+3,-2d}(\Lambda(\sigma, d))$, and the obstruction is $\mathrm{Sq}(u)$ by [Mur20a, Prop. 3.4] (here, only the last summand counts, since the other ones correspond to the elements $m_{i+2} = 0$, $d < i < 2d$, which vanish since $\Lambda(\sigma, d)$ is d -sparse, as remarked in the previous item). In other words, we have obtained a d -sparse minimal A_{2d+3} -algebra $(\Lambda(\sigma, d), m_{d+2}, m_{2d+2})$ with $\{m_{d+2}\} = u$ and whose operation of top arity $2d+3$ must vanish since $\Lambda(\sigma, d)$ is d -sparse.

Let $j \geq d$ and suppose that we have constructed points $x_i \in X_i$, $d \leq i \leq j$, as well as an auxiliary point $y_{j+d+1} \in X_{j+d+1}$ that map to one another under the bonding maps in the tower:

$$\begin{array}{ccccccccccc} \cdots & \twoheadrightarrow & X_{j+d+2} & \twoheadrightarrow & X_{j+d+1} & \twoheadrightarrow & \cdots & \twoheadrightarrow & X_{j+1} & \twoheadrightarrow & X_j & \twoheadrightarrow & \cdots & \twoheadrightarrow & X_d & \twoheadrightarrow & \cdots & \twoheadrightarrow & X_0 \\ & & \exists y_{j+d+2} & & y_{j+d+1} & \xrightarrow{\quad\quad\quad} & & & x_j & \mapsto & \cdots & \mapsto & x_d & \xrightarrow{\quad\quad\quad} & & & & & x_0 \end{array}$$

(the base of the induction $j = d$ was discussed above). Since $E_{d+2}^{j+d+2,j+d+1} = 0$ by [Proposition 5.2.9](#), the obstruction for the existence of a point

$$y_{j+d+2} \in X_{j+d+2}, \quad y_{j+d+2} \mapsto x_j,$$

vanishes; therefore such a y_{j+d+2} exists and we define $x_{j+1} \in X_{j+1}$ as its image. Here we use [Mur20a, Prop. 5.4] for $k = j + d + 1$ and $l = j$. This process results in a sequence $\{x_i \in X_i \mid i \geq d\}$ that defines a point $x_\infty \in X_\infty$ in the (homotopy) limit with $x_\infty \mapsto x_d$, that is a minimal A_∞ -algebra structure on the graded algebra $\Lambda(\sigma, d)$ whose restricted universal Massey product is $j^* \{m_{d+2}\} = x$. This finishes the proof of statement (2).

Proof of statement (1): Let x_∞ and y_∞ be A_∞ -algebra structures on $\Lambda(\sigma, d)$ with universal Massey products $\{m_{d+2}^{x_\infty}\}$ and $\{m_{d+2}^{y_\infty}\}$, respectively. We need to prove that, if there is an equality of restricted universal Massey products

$$j^* \{m_{d+2}^{y_\infty}\} = j^* \{m_{d+2}^{x_\infty}\} \in \mathrm{HH}^{d+2,-d}(\Lambda, \Lambda(\sigma, d)),$$

then x_∞ and y_∞ lie in the same connected component of X_∞ . The claim follows from [Theorem B](#) since we have

$$\mathrm{Sq}(\{m_{d+2}^{x_\infty}\}) = 0 = \mathrm{Sq}(\{m_{d+2}^{y_\infty}\})$$

and therefore $\{m_{d+2}^{y_\infty}\} = \{m_{d+2}^{x_\infty}\}$ (as explained in the proof of statement (2)). This finishes the proof of statement (1). \square

We are finally ready to prove [Theorem 5.1.2](#), the last logical step for establishing [Theorem A](#).

Proof of [Theorem 5.1.2](#). The theorem is a direct consequence of [Theorem 5.3.3\(3\)](#). Indeed, [Theorem 5.1.2\(1\)](#), that is the existence of an A_∞ -algebra structure on the d -sparse graded algebra $\Lambda(\sigma, d)$ whose restricted universal Massey product is a unit in Hochschild–Tate cohomology, follows immediately from the fact that the quotient $S_d^\simeq(\Lambda, \sigma)/\text{Aut}(\Lambda(\sigma, d))$ is non-empty. Finally, [Theorem 5.1.2\(2\)](#), that is the essential uniqueness of such A_∞ -algebra structures, is readily seen to be equivalent to the fact that the quotient $S_d^\simeq(\Lambda, \sigma)/\text{Aut}(\Lambda(\sigma, d))$ has exactly one element. \square

5.4. Strong differential graded enhancements. The following definition extends that of a ‘strong enhancement’ of a triangulated category (the case $d = 1$) introduced by Lunts and Orlov in [\[LO10\]](#) to the setting of $(d + 2)$ -angulated categories.

Definition 5.4.1. Let \mathcal{F} be a $(d + 2)$ -angulated category.

- (1) A *strong (DG) enhancement* of \mathcal{F} is a pair (\mathcal{A}, φ) consisting of a pre- $(d + 2)$ -angulated DG category \mathcal{A} and a \mathbf{k} -linear equivalence of $(d + 2)$ -angulated categories

$$\varphi: H^0(\mathcal{A}) \xrightarrow{\sim} \mathcal{F}.$$

- (2) Two strong enhancements (\mathcal{A}, φ) and (\mathcal{B}, ψ) of \mathcal{F} are *equivalent* if there exists a quasi-equivalence of DG categories $F: \mathcal{A} \xrightarrow{\sim} \mathcal{B}$ such that the diagram

$$\begin{array}{ccc} H^0(\mathcal{A}) & & \\ \downarrow H^0(F) & \searrow \varphi & \\ H^0(\mathcal{B}) & & \mathcal{F} \end{array} \begin{array}{c} \nearrow \psi \\ \end{array}$$

commutes up to (unspecified) natural isomorphism.

We say that \mathcal{F} *admits a unique strong (DG) enhancement*⁶ if any two strong enhancements of \mathcal{F} are equivalent.

In the case $d = 1$ of triangulated categories (with split idempotents), [Definition 5.4.1](#) admits the following equivalent formulation.

Definition 5.4.2. Let \mathcal{T} be a triangulated category with split idempotents.

- (1) A *strong (DG) enhancement* of \mathcal{T} is a pair (\mathcal{A}, φ) consisting of a DG category \mathcal{A} and a \mathbf{k} -linear equivalence of triangulated categories

$$\varphi: D^c(\mathcal{A}) \xrightarrow{\sim} \mathcal{T}.$$

- (2) Two strong enhancements (\mathcal{A}, φ) and (\mathcal{B}, ψ) of \mathcal{T} are *equivalent* if there exists a Morita equivalence of DG categories $F: \mathcal{A} \xrightarrow{\sim} \mathcal{B}$ such that the diagram

$$\begin{array}{ccc} D^c(\mathcal{A}) & & \\ \downarrow \mathbb{L}F_1 & \searrow \varphi & \\ D^c(\mathcal{B}) & & \mathcal{T} \end{array} \begin{array}{c} \nearrow \psi \\ \end{array}$$

commutes up to (unspecified) natural isomorphism.

In order to investigate strong enhancements of $(d + 2)$ -angulated categories, we extend the notion to pairs.

⁶In this case one also says that \mathcal{T} ‘admits a strongly unique enhancement.’

Definition 5.4.3. Let (\mathcal{C}, Σ) be a pair consisting of an additive category and a self-equivalence, that we often omit from the notation.

- (1) A *strong $(d+2)$ -angulated (DG) enhancement* of \mathcal{C} is a pair (\mathcal{A}, φ) consisting of a pre- $(d+2)$ -angulated DG category \mathcal{A} and a \mathbf{k} -linear equivalence of pairs

$$\varphi: H^0(\mathcal{A}) \xrightarrow{\sim} \mathcal{C}.$$

- (2) Two strong enhancements (\mathcal{A}, φ) and (\mathcal{B}, ψ) of \mathcal{C} are *equivalent* if there exists a quasi-equivalence of DG categories $F: \mathcal{A} \xrightarrow{\sim} \mathcal{B}$ such that the diagram

$$\begin{array}{ccc} H^0(\mathcal{A}) & & \\ H^0(F) \downarrow & \searrow \varphi & \\ H^0(\mathcal{B}) & \nearrow \psi & \mathcal{C} \end{array}$$

commutes up to (unspecified) natural isomorphism.

Remark 5.4.4. For $d = 1$, [Definition 5.4.3](#) also admits an equivalent formulation in terms of perfect derived categories, compare [Definitions 5.4.1](#) and [5.4.2](#).

Remark 5.4.5. A strong $(d+2)$ -angulated enhancement of (\mathcal{C}, Σ) induces a $(d+2)$ -angulated structure and two equivalent enhancements induce *identical* $(d+2)$ -angulated structures, not just equivalent ones. Therefore, the set of equivalence classes of strong enhancements of a $(d+2)$ -angulated category $(\mathcal{C}, \Sigma, \diamond)$ is a subset of the set of equivalence classes of strong $(d+2)$ -angulated enhancements of (\mathcal{C}, Σ) . Moreover, the latter can be partitioned by subsets of the former form as \diamond runs over the set of $(d+2)$ -angulated structures on the pair (\mathcal{C}, Σ) .

Recall the standing assumptions of this section, [Setting 5.0.1](#). We endow $\text{proj}(\Lambda)$ with any of the equivalent algebraic, that is AL $(d+2)$ -angulated structures \diamond with suspension functor $\Sigma = - \otimes_{\Lambda} \sigma \Lambda_1$, which we fix, see [Definition 2.2.17](#), [Proposition 2.2.18](#) and [Remark 5.1.4](#). Our aim is to compute the set of equivalence classes of strong enhancements of the $(d+2)$ -angulated category $(\text{proj}(\Lambda), \Sigma, \diamond)$. For this, we need some auxiliary definitions.

Definition 5.4.6. The *stable centre* $\underline{Z}(\Lambda)$ of Λ is the quotient of the centre $Z(\Lambda)$ by the ideal formed by the elements $a \in Z(\Lambda)$ such that the Λ -bimodule morphism

$$\Lambda \rightarrow \Lambda, \quad x \mapsto xa,$$

factors through a projective-injective Λ -bimodule.

Remark 5.4.7. In the same way that the centre of Λ can be interpreted in terms of Hochschild cohomology as $Z(\Lambda) = \text{HH}^0(\Lambda)$, the stable centre is interpreted in terms of Hochschild–Tate cohomology as

$$\underline{Z}(\Lambda) = \underline{\text{HH}}^0(\Lambda) = \underline{\text{Hom}}_{\Lambda^e}(\Lambda, \Lambda).$$

We recall the following classical definition.

Definition 5.4.8. The *centre* $Z(\mathcal{D})$ of a \mathbf{k} -linear category \mathcal{D} is the endomorphism algebra of the identity functor $\mathbf{1}_{\mathcal{D}}$.

Remark 5.4.9. There is a commutative square of algebra maps

$$\begin{array}{ccc} Z(\Lambda) & \xrightarrow{\cong} & Z(\text{mod } \Lambda) \\ \rho \downarrow & & \downarrow \\ \underline{Z}(\Lambda) & \xrightarrow{\zeta} & Z(\underline{\text{mod}} \Lambda) \end{array}$$

where the top arrow is the well-known isomorphism that sends $a \in Z(\Lambda)$ to the natural transformation with components

$$M \longrightarrow M, \quad x \mapsto xa, \quad M \in \mathbf{mod}(\Lambda),$$

the left vertical arrow is the quotient map, and the right vertical arrow is induced by the ideal quotient

$$\mathbf{mod}(\Lambda) \twoheadrightarrow \underline{\mathbf{mod}}(\Lambda) = \mathbf{mod}(\Lambda)/[\mathbf{proj}(\Lambda)].$$

We denote by

$$\zeta^\times : \underline{Z}(\Lambda)^\times \longrightarrow Z(\underline{\mathbf{mod}}(\Lambda))^\times$$

the morphism induced by ζ between the corresponding groups of units.

Theorem 5.4.10. *Suppose that the algebra Λ/J_Λ is separable and endow the pair $(\mathbf{proj}(\Lambda), \Sigma)$ with an algebraic $(d+2)$ -angulation \diamond . The following statements hold:*

- (1) *The set of equivalence classes of strong $(d+2)$ -angulated enhancements of the pair $(\mathbf{proj}(\Lambda), \Sigma)$ is in bijection with the set $\underline{Z}(\Lambda)^\times$ of units in the stable centre of Λ .*
- (2) *The subset of equivalence classes of strong enhancements of the $(d+2)$ -angulated category $(\mathbf{proj}(\Lambda), \Sigma, \diamond)$ is in bijection with $\ker \zeta^\times$.*

Proof. Let (\mathcal{A}, φ) be a strong enhancement of $(\mathbf{proj}(\Lambda), \Sigma, \diamond)$, which is in particular a strong $(d+2)$ -angulated enhancement of $(\mathbf{proj}(\Lambda), \Sigma)$. If (\mathcal{B}, ψ) is another strong $(d+2)$ -angulated enhancement of $(\mathbf{proj}(\Lambda), \Sigma)$, then \mathcal{A} and \mathcal{B} are quasi-equivalent, were we use [Proposition 2.2.18](#), [Theorem 5.1.3](#) and the separability hypothesis. Therefore, without loss of generality, we may replace \mathcal{B} by \mathcal{A} and consider only strong enhancements of the form (\mathcal{A}, ψ) .

A strong $(d+2)$ -angulated enhancement of the form (\mathcal{A}, ψ) is equivalent to $(\mathcal{A}, \xi\varphi)$ for $\xi = \psi\varphi^{-1}$, where φ^{-1} is a choice of quasi-inverse of the equivalence of triangulated categories φ ; indeed, the diagram

$$\begin{array}{ccc} H^0(\mathcal{A}) & & \\ \mathbf{1}_{H^0(\mathcal{A})} = H^0(\mathbf{1}_{\mathcal{A}}) \downarrow & \begin{array}{c} \nearrow \xi\varphi \\ \searrow \psi \end{array} & \mathcal{C} \\ H^0(\mathcal{A}) & & \end{array}$$

commutes up to natural isomorphism. Moreover, given two self-equivalences ξ_1, ξ_2 of $\mathbf{proj}(\Lambda)$, the strong $(d+2)$ -angulated enhancements $(\mathcal{A}, \xi_1\varphi)$ and $(\mathcal{A}, \xi_2\varphi)$ are equivalent if and only if there exists an automorphism F of \mathcal{A} in $\mathbf{Ho}(\mathbf{dgcats})$ such that the following square commutes up to natural isomorphism

$$\begin{array}{ccc} H^0(\mathcal{A}) & \xrightarrow{\varphi} & \mathbf{proj}(\Lambda) \\ H^0(F) \downarrow & & \downarrow \xi_2^{-1}\xi_1 \\ H^0(\mathcal{A}) & \xrightarrow{\varphi} & \mathbf{proj}(\Lambda) \end{array}$$

In this context, talking about the self-equivalence of $H^0(\mathcal{A})$ induced by an automorphism F of \mathcal{A} in $\mathbf{Ho}(\mathbf{dgcats})$ makes sense because any such automorphism is represented by a zig-zag of quasi-equivalences

$$\mathcal{A} = \mathcal{A}^{(0)} \xleftarrow{\sim} \mathcal{A}^{(1)} \xrightarrow{\sim} \dots \xleftarrow{\sim} \mathcal{A}^{(n-1)} \xrightarrow{\sim} \mathcal{A}^{(n)} = \mathcal{A}.$$

Here, taking cohomology we obtain a zig-zag of equivalences of pairs

$$H^0(\mathcal{A}) = H^0(\mathcal{A}^{(0)}) \xleftarrow{\sim} H^0(\mathcal{A}^{(1)}) \xrightarrow{\sim} \dots \xleftarrow{\sim} H^0(\mathcal{A}^{(n-1)}) \xrightarrow{\sim} H^0(\mathcal{A}^{(n)}) = H^0(\mathcal{A}).$$

We can then take quasi-inverses of arrows pointing backwards

$$H^0(\mathcal{A}) = H^0(\mathcal{A}^{(0)}) \xrightarrow{\sim} H^0(\mathcal{A}^{(1)}) \xrightarrow{\sim} \dots \xrightarrow{\sim} H^0(\mathcal{A}^{(n-1)}) \xrightarrow{\sim} H^0(\mathcal{A}^{(n)}) = H^0(\mathcal{A})$$

and the composite is well defined up to natural isomorphism. We conclude that the set of equivalence classes of strong $(d + 2)$ -angulated enhancements of $(\text{proj}(\Lambda), \Sigma)$ is in bijection with the set of left cosets of the group of natural isomorphism classes of self-equivalences of $(\text{proj}(\Lambda), \Sigma)$ modulo the subgroup defined by the self-equivalences induced by an automorphism of \mathcal{A} in $\text{Ho}(\text{dgcats})$, see [Section 3.1.2](#).

Recall that the group of self-equivalences of the category $\text{proj}(\Lambda)$ modulo natural isomorphisms—the Picard group of Λ —identifies with the outer automorphism group $\text{Out}(\Lambda)$, see [Section 2.2.2](#). Similarly, the group of self-equivalences of the pair $(\text{proj}(\Lambda), \Sigma)$ modulo natural isomorphisms identifies with the outer automorphism group $\text{Out}(\Lambda(\sigma, d))$ of the graded algebra $\Lambda(\sigma, d)$. We now investigate the self-equivalences of \mathcal{A} . We use homotopical techniques that have also been used in [\[Mur22, Sec. 8\]](#) with similar purposes.

Let $\mathcal{B} \subset \mathcal{A}$ be the full DG subcategory spanned by the objects which are basic additive generators of $H^0(\mathcal{A}) \simeq \text{proj}(\Lambda)$. Denote by $\text{dgcats}_{\mathcal{A}}, \text{dgcats}_{\mathcal{B}} \subset \text{dgcats}$ the non-full (!) subcategories whose objects are the DG categories which are quasi-isomorphic to \mathcal{A} and \mathcal{B} , respectively, and whose morphisms are quasi-equivalences.

The following argument is based in the proof of [\[Mur22, Prop. 8.5\]](#). Consider the functors

$$\text{dgcats}_{\mathcal{B}} \begin{array}{c} \xrightarrow{R} \\ \xleftarrow{j} \end{array} \text{dgcats}_{\mathcal{A}}$$

defined as follows:

- The functor R is the homotopy Karoubi envelope (defined for example in [\[GHW21, Sec. 4\]](#)), which is an endofunctor of dgcats .
- The functor j takes a DG category \mathcal{D} quasi-equivalent to \mathcal{A} to its full DG subcategory spanned by the objects that are basic additive generators in $H^0(\mathcal{D}) \simeq H^0(\mathcal{A}) \simeq \text{proj}(\Lambda)$.

For an arbitrary DG category \mathcal{D} , there is a natural functor $\mathcal{D} \rightarrow R(\mathcal{D})$ that is a quasi-equivalence if and only if $H^0(\mathcal{D})$ is idempotent complete. If \mathcal{D} is quasi-equivalent to \mathcal{B} , then the previous canonical functor corestricts to a natural quasi-equivalence $\mathcal{D} \xrightarrow{\sim} jR(\mathcal{D})$. If \mathcal{D} is quasi-equivalent to \mathcal{A} , then there is a zig-zag of natural quasi-equivalences

$$Rj(\mathcal{D}) \xrightarrow{\sim} R(\mathcal{D}) \xleftarrow{\sim} \mathcal{D}.$$

This proves that R and j induce inverse homotopy equivalences between the classifying spaces of the categories $\text{dgcats}_{\mathcal{A}}$ and $\text{dgcats}_{\mathcal{B}}$,

$$|\text{dgcats}_{\mathcal{B}}| \begin{array}{c} \xrightarrow{|R|} \\ \xleftarrow{|j|} \end{array} |\text{dgcats}_{\mathcal{A}}|.$$

Taking fundamental groups (these spaces are connected by construction) we obtain an isomorphism between the automorphism groups of \mathcal{A} and \mathcal{B} in $\text{Ho}(\text{dgcats})$.

Let $c \in \mathcal{A}$ be an object such that $\varphi(c) \cong \Lambda$ in $\text{proj}(\Lambda)$; in particular, $c \in \mathcal{B}$. Since all objects in $H^0(\mathcal{B})$ are isomorphic, the homotopy fiber sequence in the proof of [\[TV08, Prop. 2.3.3.5\]](#) gives an isomorphism between the automorphism group of \mathcal{B} in $\text{Ho}(\text{dgcats})$ and the automorphism group of $\mathcal{B}(c, c)$ in $\text{Ho}(\text{dgalg})$, the homotopy category of the model category of DG algebras with quasi-isomorphisms as weak equivalences and surjections as fibrations.

A choice of isomorphism $\varphi(c) \cong \Lambda$ in $\text{proj}(\Lambda)$ induces an isomorphism of graded algebras $H^\bullet(\mathcal{B}(c, c)) \cong \Lambda(\sigma, d)$. Fix a minimal model

$$(\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots)$$

of the DG algebra $\mathcal{B}(c, c)$. The automorphism group of $\mathcal{B}(c, c)$ in $\text{Ho}(\text{dgalg})$ is isomorphic to the group of homotopy classes of A_∞ -automorphisms of this minimal

model (an A_∞ -isomorphism of minimal A_∞ -algebras is an A_∞ -morphism F such that F_1 is an isomorphism); this follows from [LH, Cor. 1.3.1.3] (despite the fact that only non-unital DG algebras are considered in [LH], the claim is still valid by [Mur14, Prop. 6.2]).

An A_∞ -automorphism F of $(\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots)$ is the same as an A_∞ -isomorphism

$$\begin{aligned} G: (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) &\longrightarrow (\Lambda(\sigma, d), m_{d+2}, m_{2d+2}, \dots) * F_1 \\ &= (\Lambda(\sigma, d), m_{d+2}^{F_1}, m_{2d+2}^{F_1}, \dots) \end{aligned}$$

with $G_1 = \mathbf{1}$ and $G_n = F_1^{-1}F_n$, see [Notation 5.1.1](#) and compare with the proof of [Theorem 5.1.3](#). By [Theorem 5.3.3\(1\)](#), such an A_∞ -isomorphism exists if and only if there is an equality of restricted universal Massey products

$$j^*\{m_{d+2}\} = j^*\{m_{d+2}^{F_1}\} \in \mathbb{H}^{d+2, -d}(\Lambda, \Lambda(\sigma, d)).$$

Moreover, using the right conjugation action of $\text{Aut}(\Lambda(\sigma, d))$ on $\mathbb{H}^{d+2, -d}(\Lambda, \Lambda(\sigma, d))$ that appeared in [Remark 4.7.10](#),

$$j^*\{m_{d+2}^{F_1}\} = j^*\{m_{d+2}\}^{F_1}.$$

Hence, if $G \subset \text{Aut}(\Lambda(\sigma, d))$ is now the isotropy group of this action at $j^*\{m_{d+2}\}$ and $H \subset \text{Out}(\Lambda(\sigma, d))$ is the image of G along the quotient map

$$\text{Aut}(\Lambda(\sigma, d)) \twoheadrightarrow \text{Out}(\Lambda(\sigma, d)),$$

the set of cosets we wish to describe is $\text{Out}(\Lambda(\sigma, d))/H$.

The right conjugation action of $\text{Aut}(\Lambda(\sigma, d))$ on $\mathbb{H}^{\bullet, *}$ $(\Lambda(\sigma, d), \Lambda(\sigma, d))$ factors through $\text{Out}(\Lambda(\sigma, d))$ ([Proposition 4.4.2](#)). Since $j^*\{m_{d+2}\}$ is in the image of the $\text{Aut}(\Lambda(\sigma, d))$ -equivariant morphism

$$j^*: \mathbb{H}^{d+2, -d}(\Lambda(\sigma, d), \Lambda(\sigma, d)) \longrightarrow \mathbb{H}^{d+2, -d}(\Lambda, \Lambda(\sigma, d)),$$

we deduce that G contains all inner automorphisms of $\Lambda(\sigma, d)$ and, therefore, $\text{Out}(\Lambda(\sigma, d))/H$ is in bijection with $\text{Aut}(\Lambda(\sigma, d))/G$.

By [Proposition 4.7.16](#), $\text{Aut}(\Lambda(\sigma, d))/G$ is in bijection with the edge units of $\mathbb{H}^{d+2, -d}(\Lambda, \Lambda(\sigma, d))$, that is with the Hochschild cohomology classes in bidegree $(d+2, -d)$ that become units in the Hochschild–Tate cohomology $\underline{\mathbb{H}}^{\bullet, *}$ $(\Lambda, \Lambda(\sigma, d))$. These classes are in (non-canonical) bijection with the units in

$$\underline{\mathbb{H}}^{0,0}(\Lambda, \Lambda(\sigma, d)) = \underline{\mathbb{Z}}(\Lambda),$$

since the latter group of units acts freely and transitively on the set of edge units of bidegree $(d+2, -d)$. An explicit bijection between $\text{Aut}(\Lambda(\sigma, d))/G$ and $\underline{\mathbb{Z}}(\Lambda)^\times$ is induced by the principal crossed homomorphism

$$\begin{aligned} \chi: \text{Aut}(\Lambda(\sigma, d)) &\longrightarrow \underline{\mathbb{Z}}(\Lambda)^\times, \\ g &\longmapsto j^*\{m_{d+2}\}^g j^*\{m_{d+2}\}^{-1} \end{aligned}$$

whose kernel is $\ker \chi = G$. This proves statement [\(1\)](#).

To prove statement [\(2\)](#) we have to identify the image along χ of the subgroup $K \subset \text{Aut}(\Lambda(\sigma, d))$ corresponding to the self-equivalences of $(\text{proj}(\Lambda), \Sigma)$ which preserve the $(d+2)$ -angulated structure \diamond .

The subgroup K contains G since the latter is the subgroup of $\text{Aut}(\Lambda(\sigma, d))$ formed by the automorphisms of $\Lambda(\sigma, d)$ induced by an automorphism of the pre- $(d+2)$ -angulated DG category \mathcal{A} in $\text{Ho}(\text{dgcats})$. Since \mathcal{A} is an enhancement of $(\text{proj}(\Lambda), \Sigma, \diamond)$, the self-equivalences of the pair $H^0(\mathcal{A}) \simeq (\text{proj}(\Lambda), \Sigma)$ induced by the automorphisms of \mathcal{A} also preserve the $(d+2)$ -angulated structure \diamond , that is $G \subset K$. Since $G = \ker \chi$ we deduce that $K = \chi^{-1}(\chi(K))$ (this is well known for group homomorphisms and easy to deduce for crossed homomorphisms).

Recall that the automorphisms of $\Lambda(\sigma, d)$ introduced in [Remark 4.7.10](#) define a group homomorphism

$$\begin{aligned} \gamma: Z(\Lambda)^\times &\longrightarrow \text{Aut}(\Lambda(\sigma, d)), \\ u &\longmapsto g_u. \end{aligned}$$

The composite $\chi\gamma$ is the surjective group homomorphism $Z(\Lambda)^\times \rightarrow \underline{Z}(\Lambda)^\times$ induced by the natural projection $Z(\Lambda) \rightarrow \underline{Z}(\Lambda)$ (the former homomorphism is surjective by [\[Che21, Cor. 2.3\]](#)). Since $K = \chi^{-1}(\chi(K))$ and $\chi\gamma$ is surjective,

$$\chi(K \cap \gamma(Z(\Lambda)^\times)) = \chi(K) \cap \chi\gamma(Z(\Lambda)^\times) = \chi(K).$$

Hence, $\chi(K) \subset \underline{Z}(\Lambda)^\times$ consists of the classes of those units $u \in Z(\Lambda)^\times$ such that g_u preserves the $(d+2)$ -angulation \diamond . The self-equivalence of $(\text{proj}(\Lambda), \Sigma)$ induced by g_u , $u \in Z(\Lambda)^\times$, is $(\mathbf{1}_{\text{proj}(\Lambda)}, u)$ (considered already in the proof of [Proposition 2.2.18](#)). Therefore, in order to finish the proof we have to check that $(\mathbf{1}_{\text{proj}(\Lambda)}, u)$ preserves \diamond if and only if $[u] \in \ker \zeta^\times$.

First, assume $[u] \in \ker \zeta^\times$. We have to show that, for every $(d+2)$ -angle

$$x_1 \xrightarrow{f_1} x_2 \xrightarrow{f_2} x_3 \rightarrow \cdots \rightarrow x_{d+2} \xrightarrow{f_{d+2}} \Sigma(x_1)$$

in \diamond , the sequence

$$x_1 \xrightarrow{f_1} x_2 \xrightarrow{f_2} x_3 \rightarrow \cdots \rightarrow x_{d+2} \xrightarrow{uf_{d+2}} \Sigma(x_1)$$

is also in \diamond ; notice that we can in fact place the multiplication by u in any given arrow, since the resulting sequences are isomorphic. We therefore consider the sequence

$$x_1 \xrightarrow{uf_1} x_2 \xrightarrow{f_2} x_3 \rightarrow \cdots \rightarrow x_{d+2} \xrightarrow{f_{d+2}} \Sigma(x_1)$$

and show that it is isomorphic to the first diagram. This will suffice by [\(*d*-TR1\)](#). We factor f_1 in $\text{mod}(\Lambda)$ as

$$x_1 \xrightarrow{p} M \xleftarrow{i} x_2$$

and consider the following commutative square of solid arrows:

$$\begin{array}{ccc} M & \xleftarrow{i} & x_2 \\ \downarrow u & \swarrow \alpha & \downarrow u \\ M & \xleftarrow{i} & x_2 \end{array}$$

Since $[u] \in \ker \zeta^\times$, multiplication by u induces the identity in the stable module category $\underline{\text{mod}}(\Lambda)$ for every finite-dimensional Λ -module. This implies the existence of a dashed arrow α such that

$$\alpha i = u - \mathbf{1}_M.$$

An easy computation shows that the following diagram commutes:

$$\begin{array}{ccccccc} x_1 & \xrightarrow{f_1} & x_2 & \xrightarrow{f_2} & x_3 & \longrightarrow & \cdots & \longrightarrow & x_{d+1} & \xrightarrow{f_{d+1}} & x_{d+2} & \xrightarrow{f_{d+2}} & \Sigma(x_1) \\ \parallel & & \downarrow \mathbf{1}_{x_2} + i\alpha & & \parallel & & & & \parallel & & \parallel & & \parallel \\ x_1 & \xrightarrow{uf_1} & x_2 & \xrightarrow{f_2} & x_3 & \longrightarrow & \cdots & \longrightarrow & x_{d+1} & \xrightarrow{f_{d+1}} & x_{d+2} & \xrightarrow{f_{d+2}} & \Sigma(x_1). \end{array}$$

The arrow $\mathbf{1}_{x_2} + i\alpha$ is an isomorphism by the five lemma, which is valid in the setting of $(d+2)$ -angulated categories by [\(*d*-TR2\)](#) and [\[GKO13, Prop. 2.5\(a\)\]](#).

Assume now that $(\mathbf{1}_{\text{proj}(\Lambda)}, u)$ preserves the $(d+2)$ -angulation \diamond . Let M be a finite-dimensional Λ -module. Recall that $\Sigma := - \otimes_{\Lambda} \sigma \Lambda_1$. There exists a $(d+2)$ -angle in \diamond

$$x_1 \xrightarrow{f_1} x_2 \xrightarrow{f_2} x_3 \rightarrow \cdots \rightarrow x_{d+2} \xrightarrow{f_{d+2}} \Sigma(x_1)$$

such that f_{d+2} factors as

$$x_{d+2} \xrightarrow{p} \Sigma M \xrightarrow{\Sigma(i)} \Sigma(x_1).$$

in $\text{mod}(\Lambda)$. Indeed, choose an epimorphism $p: x_{d+2} \rightarrow \Sigma M$ with projective source and a monomorphism $i: M \hookrightarrow x_1$ with injective target; define $f_{d+2} = \Sigma(i)p$ and then apply [\(d-TR1\)](#) and [\(d-TR2\)](#). The previous factorisation and $(d+2)$ -angle induce an isomorphism in $\underline{\text{mod}}(\Lambda)$,

$$\omega: \Sigma M \xrightarrow{\sim} \Omega^{-(d+2)} M.$$

Since $(\mathbf{1}_{\text{proj}(\Lambda)}, u)$ preserves \diamond , the following diagram is also in \diamond

$$x_1 \xrightarrow{f_1} x_2 \xrightarrow{f_2} x_3 \rightarrow \cdots \rightarrow x_{d+2} \xrightarrow{u f_{d+2}} \Sigma(x_1).$$

We can factor f_{d+2} as

$$x_{d+2} \xrightarrow{up} \Sigma M \xrightarrow{\Sigma(i)} \Sigma(x_1).$$

The natural isomorphism in $\underline{\text{mod}}(\Lambda)$ induced by this factorisation and the previous diagram is

$$u\omega: \Sigma M \xrightarrow{\sim} \Omega^{-(d+2)} M.$$

By [Proposition 2.1.8\(3\)](#) both isomorphisms in $\underline{\text{mod}}(\Lambda)$ must coincide:

$$\omega = u\omega.$$

Therefore, the isomorphism $u: M \xrightarrow{\sim} M$ in $\underline{\text{mod}}(\Lambda)$ represented by multiplication by u must be the identity for every $M \in \text{mod}(\Lambda)$. This shows that $[u] \in \ker \zeta^\times$, as required. This finishes the proof of the theorem. \square

Remark 5.4.11. It is an interesting problem to study the set of equivalence classes of strong enhancements of an algebraic triangulated category \mathcal{T} with finite-dimensional morphism spaces, split idempotents, and a $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$. [Theorem 5.4.10](#) provides a complete answer in the case $d = 1$ when $\text{add}(c) = \mathcal{T}$. In general, there is an apparent map (given by restriction) from the set of equivalence classes of strong enhancements of the ambient triangulated category \mathcal{T} to the set of equivalence classes of strong enhancements of the standard $(d+2)$ -angulated category $(\text{add}(c), [d], \diamond)$. We do not know, however, whether this map is injective or surjective.

The following result provides, in particular, the simplest example of an algebraic triangulated category with a unique enhancement but with non-unique strong enhancements.

Corollary 5.4.12. *Let $\Lambda = \mathbf{k}[\varepsilon]/(\varepsilon^2)$ be the algebra of dual numbers. The following statements hold:*

- (1) *The algebra Λ is connected and twisted $(d+2)$ -periodic for every $d \geq -1$ and Λ/J_Λ is separable, hence $\text{proj}(\Lambda)$ is algebraic $(d+2)$ -angulated for every $d \geq 1$ and there is only one choice of suspension functor.*
- (2) *If $\text{char}(\mathbf{k}) \neq 2$ and $d \geq 1$, then every algebraic $(d+2)$ -angulated structure on $\text{proj}(\Lambda)$ admits a unique strong enhancement.*

- (3) If $\text{char}(\mathbf{k}) = 2$ and $d \geq 1$, then for every algebraic $(d+2)$ -angulated structure on $\text{proj}(\Lambda)$ the set of equivalence classes of strong enhancements is in bijection with the elements of the ground field \mathbf{k} , and therefore $\text{proj}(\Lambda)$ does not admit a unique strong enhancement.

Proof. (1) The algebra Λ is obviously connected and $\Lambda/J_\Lambda = \Lambda/(\varepsilon) = \mathbf{k}$ is separable. The algebra Λ is also twisted 1-periodic, as witnessed by the short exact sequence of Λ -bimodules

$$0 \longrightarrow {}_\sigma\Lambda_1 \xrightarrow{\nu} \Lambda \otimes \Lambda \xrightarrow{\mu} \Lambda \longrightarrow 0,$$

where μ is the multiplication map, the automorphism σ is given by $\sigma(\varepsilon) = -\varepsilon$, and

$$\nu(1) = 1 \otimes \varepsilon - \varepsilon \otimes 1.$$

Therefore, Λ is twisted $(d+2)$ -periodic for every $d \geq -1$. By [Theorem 2.2.20](#), the category $\text{proj}(\Lambda)$ is $(d+2)$ -angulated for any $d \geq 1$. Indeed, by [Theorem 2.2.15](#), [Proposition 2.2.6](#) and [Corollaries 4.5.17](#) and [4.5.20](#) it has a unique AL $(d+2)$ -angulated structure, up to equivalence, and the only choice of suspension functor is $\Sigma = - \otimes_\Lambda {}_\sigma\Lambda_1$. Algebraic and AL $(d+2)$ -angulated structures are the same thing ([Remark 5.1.4](#)). Now, the first claim follows.

(2) and (3) In view of [Theorem 5.4.10](#), we need a set-theoretic description of the kernel of the canonical morphism

$$\zeta^\times : \underline{Z}(\Lambda)^\times \longrightarrow Z(\underline{\text{mod}}(\Lambda))^\times.$$

Clearly $Z(\Lambda) = \Lambda$ and the stable category

$$\underline{\text{mod}}(\Lambda) \simeq \underline{\text{mod}}(\mathbf{k})$$

has centre isomorphic to the ground field \mathbf{k} . Notice that all Λ -bimodule morphisms $\Lambda \rightarrow \Lambda \otimes \Lambda$ are of the form

$$1 \mapsto \alpha(1 \otimes \varepsilon + \varepsilon \otimes 1) + \beta(\varepsilon \otimes \varepsilon), \quad \alpha, \beta \in \mathbf{k}.$$

The composition of such a Λ -bimodule morphism with μ satisfies

$$1 \mapsto 2\alpha\varepsilon.$$

Since μ is a surjection from a projective Λ -bimodule, the stable centre is the quotient of $Z(\Lambda)$ by the ideal (2ε) . Hence, if $\text{char}(\mathbf{k}) \neq 2$, then ζ is an isomorphism. On the other hand, if $\text{char}(\mathbf{k}) = 2$ then $\zeta: \Lambda \rightarrow \mathbf{k}$ is the natural projection and

$$\ker \zeta^\times = 1 + (\varepsilon) = \{1 + \alpha\varepsilon \mid \alpha \in \mathbf{k}\}.$$

The two last claims now follow from [Theorem 5.4.10](#). \square

Example 5.4.13. Let $\text{char}(\mathbf{k}) = 2$ and $\Lambda = \mathbf{k}[\varepsilon]/(\varepsilon^2)$ the algebra of dual numbers. Following the proof of [Theorem 5.4.10](#) and [Corollary 5.4.12](#), we give an explicit description of the strong enhancements of the triangulated category $\text{proj}(\Lambda)$ (for any triangulated structure, all of them are algebraic).

Since $\text{char}(\mathbf{k}) = 2$ the suspension functor must be $\Sigma = \mathbf{1}_{\text{proj}(\Lambda)}$ (see the proof of [Corollary 5.4.12](#)). Moreover, the stable category $\underline{\text{mod}}(\Lambda)$ is equivalent to $\underline{\text{mod}}(\mathbf{k})$ and $\Omega = \Sigma = \mathbf{1}_{\underline{\text{mod}}(\mathbf{k})}$. [Proposition 2.1.8](#) establishes a bijection between the triangulated structures on $(\text{proj}(\Lambda), \Sigma)$ and the set \mathbf{k}^\times of units of the ground field. It is straightforward to verify that the triangulated structure \diamond_α corresponding to $\alpha \in \mathbf{k}^\times$ is characterised by the fact that the triangle

$$\Lambda \xrightarrow{\varepsilon} \Lambda \xrightarrow{\varepsilon} \Lambda \xrightarrow{\alpha\varepsilon} \Lambda$$

belongs to \diamond_α , compare with [[MSS07](#), Rmk. 8].

Consider the DG algebra with underlying graded algebra

$$A = \frac{k[u, v^{\pm 1}]\langle a \rangle}{(a^2, av + va, au + ua + 1)}, \quad |u| = 0, \quad |v| = 1, \quad |a| = 0,$$

and differential

$$d(u) = 0, \quad d(v) = 0, \quad d(a) = u^2v.$$

It is proved in [MSS07, Rmk. 8] that there is an equivalence of pairs

$$\begin{aligned} \varphi: \mathbf{D}^c(A) &\xrightarrow{\sim} \mathbf{proj}(\Lambda), \\ M &\longmapsto H^0(M), \end{aligned}$$

which is a triangulated equivalence if we endow the target with the triangulated structure \diamond_1 . For any $\alpha \in \mathbf{k}^\times$, we have a triangulated equivalence

$$(\mathbf{1}_{\mathbf{proj}(\Lambda)}, \alpha): (\mathbf{proj}(\Lambda), \Sigma, \diamond_1) \longrightarrow (\mathbf{proj}(\Lambda), \Sigma, \diamond_\alpha)$$

of the kind considered in the proofs of [Proposition 2.2.18](#) and [Corollary 5.4.12](#). Given $\beta \in \mathbf{k}$ we consider the self-equivalence $(\mathbf{1}_{\mathbf{proj}(\Lambda)}, 1 + \beta\varepsilon)$ and define

$$\varphi_{\alpha,\beta} = (\mathbf{1}_{\mathbf{proj}(\Lambda)}, 1 + \beta\varepsilon)(\mathbf{1}_{\mathbf{proj}(\Lambda)}, \alpha)\varphi = (\mathbf{1}_{\mathbf{proj}(\Lambda)}, (1 + \beta\varepsilon)\alpha)\varphi.$$

The map

$$\beta \longmapsto (A, \varphi_{\alpha,\beta})$$

realises the bijection between \mathbf{k} and the set of equivalence classes of strong enhancements of $(\mathbf{proj}(\Lambda), \Sigma, \diamond_\alpha)$ given by [Corollary 5.4.12](#).

5.5. Comments on the (non-)uniqueness of enhancements. We wish to illustrate the necessity of the assumptions in [Theorems 5.1.3](#), [5.1.8](#) and [5.4.10](#) by analysing the case when $\Lambda = \mathbf{k}$ is an arbitrary field, which is the most basic example of a periodic algebra of any period $d \geq 1$. Thus, let $d \geq 1$ and $\sigma = \mathbf{1}_\Lambda$ so that

$$\Lambda(\sigma, d) = \mathbf{k}\langle t^\pm \rangle, \quad |t| = -d,$$

is the algebra of Laurent polynomials in a single variable of degree $-d$, which we view as a DG algebra with vanishing differential (this algebra is considered implicitly in [[Lad](#), Ex. 3.30] and explicitly in [[Lor](#), Ex. 6.7], see also [Remark 5.5.2](#)).

By definition, a DG $\Lambda(\sigma, d)$ -module $M = (V, \varphi)$ consists of a cochain complex of vector spaces V equipped with an invertible degree $-d$ cocycle $\varphi \in Z^{-d}(\mathbf{hom}_{\mathbf{k}}(V, V))$ that we interpret as an isomorphism $\varphi: V \xrightarrow{\sim} V[-d]$. Similarly, a morphism of DG $\Lambda(\sigma, d)$ -modules $f: (V, \varphi) \rightarrow (W, \psi)$ is a morphism of cochain complexes of vector spaces $f: V \rightarrow W$ such that the diagram

$$\begin{array}{ccc} V & \xrightarrow{\varphi} & V[-d] \\ \downarrow f & & \downarrow f[-d] \\ W & \xrightarrow{\psi} & W[-d] \end{array}$$

commutes. In particular, such an f is a quasi-isomorphism if and only if the induced linear map

$$H^i(f): H^i(V) \longrightarrow H^i(W)$$

is an isomorphism for all $0 \leq i < d$. We claim that functor

$$\mathbf{D}(\Lambda(\sigma, d)) \longrightarrow \prod_{i=0}^{d-1} \mathbf{Mod}(\mathbf{k}), \quad M \longmapsto (H^0(M), H^1(M), \dots, H^{d-1}(M)),$$

is an equivalence of (ordinary) categories (although this fact is well known to experts, we give a proof for the convenience of the reader). For this, notice that there is an apparent fully faithful functor

$$\prod_{i=0}^{d-1} \mathbf{Mod}(\mathbf{k}) \longrightarrow \mathbf{dgMod}(\Lambda(\sigma, d))$$

which sends a tuple $(V^0, V^1, \dots, V^{d-1})$ to the DG $\Lambda(\sigma, d)$ -module with trivial differential $(V, \mathbf{1}_V)$ where

$$V: \quad \dots \quad V^{d-1} \quad V^0 \quad V^1 \quad \dots \quad V^{d-1} \quad V^0 \quad \dots$$

Notice that the composite

$$\prod_{i=0}^{d-1} \text{Mod}(\mathbf{k}) \longrightarrow \text{dgMod}(\Lambda(\sigma, d)) \longrightarrow H^0(\text{dgMod}_{\text{dg}}(\Lambda(\sigma, d)))$$

is also fully faithful. To complete the proof, it is enough to observe the following:

- Every DG $\Lambda(\sigma, d)$ -module is homotopy equivalent to its cohomology (viewed as a DG $\Lambda(\sigma, d)$ -module in the obvious way).
- A morphism of DG $\Lambda(\sigma, d)$ -modules is a quasi-isomorphism if and only if it is a homotopy equivalence and therefore

$$H^0(\text{dgMod}_{\text{dg}}(\Lambda(\sigma, d))) = \text{D}(\Lambda(\sigma, d)).$$

- The composite

$$\prod_{i=0}^{d-1} \text{Mod}(\mathbf{k}) \longrightarrow \text{D}(\Lambda(\sigma, d)) \longrightarrow \prod_{i=0}^{d-1} \text{Mod}(\mathbf{k})$$

is isomorphic to the identity functor.

The first claim is obvious and the second and third claims can be shown exactly as in the case of cochain complexes of vector spaces, observing that the category $\text{dgMod}(\Lambda(\sigma, d))$ is equivalent to the category of d -periodic cochain complexes of vector spaces. Under the above equivalence, the shift functor on $\text{D}(\Lambda(\sigma, d))$ corresponds to the automorphism

$$(V^0, V^1, \dots, V^{d-1}) \longmapsto (V^1, \dots, V^{d-1}, V^0).$$

of $\prod_{i=0}^{d-1} \text{Mod}(\mathbf{k})$. Since the graded algebra $H^\bullet(\Lambda(\sigma, d)) = \Lambda(\sigma, d)$ is finite-dimensional in each degree and

$$H^i(\Lambda(\sigma, d)[-j]) = \Lambda(\sigma, d)^{i-j} = \begin{cases} \mathbf{k} & i = j, \\ 0 & i \neq j, \end{cases} \quad 0 \leq i, j < d,$$

it is clear that the functor

$$\text{D}^c(\Lambda(\sigma, d)) \longrightarrow \prod_{i=0}^{d-1} \text{mod}(\mathbf{k}), \quad M \longmapsto (H^0(M), H^1(M), \dots, H^{d-1}(M)),$$

is also an equivalence of (ordinary) categories. Finally, the free DG module $\Lambda(\sigma, d) \in \text{D}^c(\Lambda(\sigma, d))$ is a $d\mathbb{Z}$ -cluster tilting object, since this is obviously the case for its image

$$(\mathbf{k}, 0, \dots, 0) \in \prod_{i=0}^{d-1} \text{mod}(\mathbf{k}).$$

Remark 5.5.1. According to [Theorem 2.2.1](#), the additive category

$$\text{mod}(\mathbf{k}) \simeq \text{add}(\Lambda(\sigma, d)) \subseteq \text{D}^c(\Lambda(\sigma, d))$$

admits a $(d+2)$ -angulated structure with suspension functor $\Sigma = \mathbf{1}$ (up to natural isomorphism, the only autoequivalence of $\text{mod}(\mathbf{k})$ whose $(d+2)$ -angles are induced by the triangles in $\text{D}^c(\Lambda(\sigma, d))$). In fact, $\text{mod}(\mathbf{k})$ admits a unique $(d+2)$ -angulation: A $(d+2)$ -angle

$$V_1 \rightarrow V_2 \rightarrow \dots \rightarrow V_{d+2} \rightarrow V_1$$

in $\text{mod}(\mathbf{k})$ must yield a $(d+2)$ -periodic exact sequence

$$\dots \rightarrow V_{d+2} \rightarrow V_1 \rightarrow V_2 \rightarrow \dots \rightarrow V_{d+2} \rightarrow V_1 \rightarrow \dots$$

(see [GKO13, Prop. 2.5]) and, conversely, every $(d + 2)$ -angulation must contain all such sequences for these are isomorphic to a finite direct sum of rotations of the trivial sequence

$$\mathbf{k} \xrightarrow{1} \mathbf{k} \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \mathbf{k}$$

with d zeroes. Moreover, $\text{mod}(\mathbf{k})$ has a unique DG-enhancement, namely the graded category $\text{add}(\Lambda(\sigma, d))$ endowed with the trivial differential.

Remark 5.5.2. The above discussion and **Theorem 5.1.6** show that the triangulated category $\text{D}^c(\Lambda(\sigma, d))$ is equivalent to the d -cluster category of \mathbf{k} , that is the orbit category [Kel05a]


$$\mathcal{C}_d(\mathbf{k}) := \text{D}^c(\text{mod}(\mathbf{k}))/[d].$$

Indeed, $\mathcal{C}_d(\mathbf{k})$ is an algebraic triangulated category with $[d] = \mathbf{1}$, finite-dimensional morphism spaces, split idempotents and $\mathbf{k} \in \mathcal{C}_d(\mathbf{k})$ is a $d\mathbb{Z}$ -cluster tilting object with endomorphism algebra isomorphic to \mathbf{k} (alternatively, one can also invoke the Recognition Theorem of Keller and Reiten [KR08]).

Remark 5.5.3. We record the following observation for the sake of completeness. The triangulated category $\text{D}^c(\Lambda(\sigma, d))$ admits an additive generator that, under the identification $\text{D}^c(\Lambda(\sigma, d)) \simeq \prod_{i=0}^{d-1} \text{mod}(\mathbf{k})$, is given by the direct sum $\bigoplus_{i=0}^{d-1} \mathbf{k}(i)$, where

$$\mathbf{k}(i) := (0, \dots, 0, \mathbf{k}, 0, \dots, 0)$$

with \mathbf{k} placed in the i -th coordinate. It is straightforward to verify that the derived endomorphism algebra of this additive generator is quasi-isomorphic to the graded algebra $\mathbf{k}Q/I$ where Q is the graded quiver



with arrows $u: i \rightarrow i + 1$ of degree -1 and $v: i + 1 \rightarrow i$ of degree 1 , and I is the graded ideal generated by the relations $vu = e_i$ and $uv = e_{i+1}$ (with all indices modulo d). We leave the details to the reader.

5.5.1. *Non-unique enhancements in the Hom-infinite case.* Let ℓ be a field and $\mathbf{k} = \ell(x_0, x_1, \dots, x_{d-1})$; notice that \mathbf{k} is perfect if ℓ has characteristic 0. Rizzardo and Van den Bergh [RVdB19] show that $\text{D}^c(\Lambda(\sigma, d))$, viewed as an ℓ -linear (!) algebraic triangulated category, admits two inequivalent enhancements. This does not contradict **Theorem A**, **Theorem 5.1.8** or **Theorem 5.4.10**: as an ℓ -linear category, the morphism spaces in $\text{D}^c(\Lambda(\sigma, d))$ are infinite dimensional and therefore the hypotheses in **Theorem A** and **Theorems 5.1.8** and **5.4.10** are not satisfied. Thus, this example explains why it is necessary to restrict to triangulated categories with finite-dimensional morphism spaces.

5.5.2. *Non-unique enhancements for non-separable algebras.* Although **Theorem A** requires the ground field \mathbf{k} to be perfect, in **Theorems 5.1.3**, **5.1.6** and **5.4.10** it is enough to assume that Λ/J_Λ is separable. We shall see that the latter assumption is essential. Indeed, let $p > 0$ be a prime number and $\mathbf{k} = \mathbb{F}_p(x)$; thus, \mathbf{k} is not perfect. Let

$$\Lambda := \mathbb{F}_p(x^{\frac{1}{p}}) = \frac{\mathbf{k}[X]}{(X^p - x)},$$

which is a purely inseparable field extension of \mathbf{k} of degree p with trivial Galois group; thus, in particular, $\Lambda = \Lambda/J_\Lambda$ is not separable over \mathbf{k} . Notice also that the algebras \mathbf{k} and Λ are isomorphic over \mathbb{F}_p , but not over \mathbf{k} . Moreover, there is an isomorphism of \mathbf{k} -algebras

$$\Lambda^e = \Lambda \otimes_{\mathbf{k}} \Lambda^{\text{op}} \cong \frac{\mathbf{k}[X, Y]}{(X^p - x, Y^p - x)} \cong \frac{\Lambda[Y]}{(Y - X)^p};$$

we treat the above isomorphism as an identification. Over Λ , we have seen that the semisimple category $\text{mod}(\Lambda)$ admits a unique $(d + 2)$ -angulated structure (Remark 5.5.1), and the latter admits a unique enhancement, represented by the graded category $\text{add}(\Lambda(\sigma, d))$ with trivial differential. We shall see that $\text{mod}(\Lambda)$ admits two inequivalent enhancements over the field \mathbf{k} . Indeed, notice that the diagonal Λ -bimodule has infinite projective dimension and admits a 2-periodic projective resolution

$$\dots \xrightarrow{(Y-X)^{p-1}} \Lambda^e \xrightarrow{Y-X} \Lambda^e \xrightarrow{(Y-X)^{p-1}} \Lambda^e \xrightarrow{Y-X} \Lambda^e \longrightarrow 0$$

that is in fact 1-periodic if $p = 2$. Thus, for even d (or every $d \geq 1$ if $p = 2$), we obtain a non-trivial extension of length $d + 2$

$$0 \longrightarrow \Lambda \longrightarrow \Lambda^e \xrightarrow{Y-X} \dots \xrightarrow{(Y-X)^{p-1}} \Lambda^e \xrightarrow{Y-X} \Lambda^e \longrightarrow \Lambda \longrightarrow 0$$

and we may consider the AL $(d + 2)$ -angulation associated to this extension (Theorem 2.2.15). By Theorem 5.1.3, this AL $(d + 2)$ -angulation admits an enhancement \mathcal{B} whose restricted universal Massey product is represented, up to sign, by the class of the above non-trivial extension; in particular, \mathcal{B} cannot be quasi-isomorphic to $H^\bullet(\mathcal{B})$ as DG categories over \mathbf{k} and, therefore, \mathcal{B} cannot be quasi-equivalent to the enhancement $\text{add}(\Lambda(\sigma, d))$ of $\text{mod}(\Lambda)$. This does not contradict Theorem 5.1.3: the algebra $\Lambda = \Lambda/J_\Lambda$ is not separable as a \mathbf{k} -algebra, and hence the hypotheses in Theorem 5.1.3 are not satisfied.

Moreover, just as in [RVdB20], $D^c(\mathcal{B})$ is equivalent to $D^c(\Lambda(\sigma, d))$ as triangulated categories, but the DG enhancements \mathcal{B} and $\text{add}(\Lambda(\sigma, d))$ cannot be Morita equivalent. Indeed, if they were, the arguments in the proof of Theorem 5.4.10 would show that they also have to be quasi-equivalent, and we have already argued that they are not. This proves that the separability hypothesis is necessary in Theorems 5.1.8 and 5.4.10.

5.5.3. *$d\mathbb{Z}$ -cluster tilting does not guarantee twisted $(d + 2)$ -periodicity.* According to Proposition 2.2.10, the endomorphism algebra Λ of a $d\mathbb{Z}$ -cluster tilting object in a triangulated category with finite-dimensional morphism spaces is twisted $(d + 2)$ -periodic, provided that Λ/J_Λ is separable. This need not be the case if the algebra Λ/J_Λ is not separable, even if the ambient triangulated category is algebraic. Indeed, as in Section 5.5.2 above, let $p > 0$ be a prime number, $\mathbf{k} = \mathbb{F}_p(x)$ and $\Lambda = \mathbb{F}_p(x^{\frac{1}{p}})$. As explained in Section 5.5, the triangulated category

$$D^c(\Lambda\langle i^\pm \rangle), \quad |i| = -d,$$

is algebraic and admits a $d\mathbb{Z}$ -cluster tilting object with endomorphism algebra isomorphic to Λ (since Λ is itself a field). However, if $p \neq 2$ and d is odd (in particular if $d = 1$), then Λ is not twisted $(d + 2)$ -periodic as a \mathbf{k} -algebra and, therefore, the restricted universal Massey product associated to the canonical \mathbf{k} -linear (!) enhancement of $D^c(\Lambda\langle i^\pm \rangle)$ cannot be represented by a unit in the Hochschild–Tate cohomology of Λ (viewed as a \mathbf{k} -algebra). This does not contradict Proposition 2.2.10 nor the implication (1) \Rightarrow (3) in Corollary 4.5.17, since $\Lambda = \Lambda/J_\Lambda$ is not separable as a \mathbf{k} -algebra.

5.5.4. *Differential graded enhancements vs topological enhancements.* Our results do not give any information on the (non-)existence of exotic *topological* enhancements, that is enhancements given in terms of spectral categories, say. We illustrate this point with the following example from [SS03, Rmk. 3.2.1]. Fix an arbitrary prime number p and an integer $n \geq 1$. We consider the perfect field $\mathbf{k} = \mathbb{F}_p$ and set $d = 2p^n - 2$ and $\Lambda = \mathbf{k}$. The graded algebra

$$\Lambda\langle i^{\pm 1} \rangle = \mathbf{k}\langle i^{\pm 1} \rangle = \mathbb{F}_p[i^{\pm 1}], \quad |i| = -d = -(2p^n - 2),$$

is isomorphic to the graded homotopy ring $\pi_*(K(n))$ of the Morava K -theory spectrum $K(n)$ of height n at the prime p . Moreover, there is an equivalence of triangulated categories

$$D(K(n)) \simeq D(\mathbb{F}_p[i^{\pm 1}]),$$

where $D(K(n))$ is the derived category of $K(n)$ -module spectra; the above equivalence of triangulated categories implies that $K(n) \in D^c(K(n)) \simeq D^c(\mathbb{F}_p[i^{\pm 1}])$ is a $d\mathbb{Z}$ -cluster tilting object in the compact derived category of $K(n)$. By construction, the triangulated category on the left-hand side is equipped with a topological enhancement while the triangulated category on the right-hand side is equipped with a DG enhancement; however, identifying DG \mathbf{k} -algebras with algebra spectra over the Eilenberg–MacLane commutative ring spectrum $H\mathbf{k}$ of \mathbf{k} [Shi07], these two enhancements are inequivalent (the reason being that the spectrum $K(n)$ is not weakly equivalent to the Eilenberg–MacLane spectrum of $\mathbb{F}_p[i^{\pm 1}]$). This does not contradict **Theorem A** since the theorem only pertains to \mathbf{k} -linear DG enhancements and DG \mathbf{k} -algebras (equivalently, spectral enhancements over $H\mathbf{k}$ and $H\mathbf{k}$ -algebra spectra).

6. RECOGNITION THEOREMS

In this section we discuss several recognition theorems for algebraic triangulated categories of interest in representation theory and algebraic and symplectic geometry that are immediate consequences of **Theorem A** (given the existent knowledge about these categories). Except in special cases, the triangulated categories considered below do not admit an additive generator and therefore the results in [Mur22] generally do not apply to them.

Setting 6.0.1. For simplicity, we assume that \mathbf{k} is a perfect field throughout this section.

Excluding **Theorem 6.4.3**, all of the applications of **Theorem A** that we discuss in this section hold, more generally, when the endomorphism algebra Λ of the (basic) $d\mathbb{Z}$ -cluster tilting objects under consideration is such that the quotient Λ/J_Λ is a separable algebra over the ground field. This assumption is satisfied automatically in the following cases:

- When the ground field is perfect (for example if it is algebraically closed or it has characteristic 0).
- When the ground field is arbitrary and $\Lambda \cong \mathbf{k}Q/I$ is (isomorphic to) the bounded path algebra of a finite quiver. Indeed, in this case there is an isomorphism $\Lambda/J_\Lambda \cong \mathbf{k}^{Q_0}$, where Q_0 denotes the set of vertices of Q . The analogous statement holds when the finite-dimensional algebra Λ is the (complete) Jacobian algebra of a quiver with potential.

Under this more general assumption, references to **Theorem A** should be replaced by references to **Theorems 5.1.6** and **5.1.8**. Similarly, non-separable algebras should be considered in place of non-semisimple algebras when appropriate (since these notions coincide for perfect fields but not for arbitrary fields).

6.1. Amiot–Guo–Keller cluster categories. We begin by recalling the following theorem, proven by Amiot in the case $d = 2$ and by Guo in the general case, both using results of Keller [Kel05a, Kel11] in an essential way.

Recall that a triangulated category \mathcal{T} with finite-dimensional morphism spaces is *n-Calabi–Yau* if there exists a natural isomorphism

$$\mathcal{T}(y, x[n]) \xrightarrow{\cong} D\mathcal{T}(x, y), \quad x, y \in \mathcal{T}.$$

where $V \mapsto DV$ denotes the passage to the \mathbf{k} -linear dual.

Given a DG algebra Γ , we denote by $D^{\text{fd}}(\Gamma)$ the full triangulated subcategory of $D(\Gamma)$ spanned by the DG Γ -modules M such that

$$\sum_{i \in \mathbb{Z}} \dim H^i(M) < \infty.$$

Following Kontsevich, we say that Γ is *homologically smooth* if the diagonal bimodule Γ is perfect as a DG Γ -bimodule, that is $\Gamma \in D^c(\Gamma^e)$. Also, given an integer n , the DG algebra Γ is *bimodule n-Calabi–Yau* if there is an isomorphism

$$\text{RHom}_{\Gamma^e}(\Gamma, \Gamma^e) \cong \Gamma[-n]$$

in the derived category of DG Γ -bimodules. If Γ is homologically smooth and bimodule n -Calabi–Yau, then the triangulated category $D^{\text{fd}}(\Gamma)$ is *n-Calabi–Yau*, see [Kel08, Lemma 4.1].

Theorem 6.1.1 ([Ami09, Thm. 2.1] and [Guo11, Thm. 2.2], see also [IY18, Thm. 5.8] for the case $d = 1$). *Let $d \geq 1$ and Γ a DG algebra that satisfies the following properties:*

- *The DG algebra Γ is homologically smooth.*
- *The cohomology of Γ is concentrated in non-positive degrees.*
- *The (ordinary) algebra $H^0(\Gamma)$ is finite-dimensional.*
- *The DG algebra Γ is bimodule $(d + 1)$ -Calabi–Yau.*

Then, the following statements hold:

- (1) *The Verdier quotient*

$$\mathcal{C}(\Gamma) = D^c(\Gamma) / D^{\text{fd}}(\Gamma)$$

is well defined and is an (algebraic) d-Calabi–Yau triangulated category with split idempotents. In particular, $\mathcal{C}(\Gamma)$ has finite-dimensional morphism spaces.

- (2) [Lad, Lemma 2.7] *There are isomorphisms of vector spaces*

$$\mathcal{C}(\Gamma)(\Gamma[i], \Gamma) = H^{-i}(\Gamma), \quad 0 \leq i < d.$$

- (3) *The image of Γ in $\mathcal{C}(\Gamma)$ is a d-cluster tilting object whose endomorphism algebra is isomorphic to $H^0(\Gamma)$.*

Definition 6.1.2. Let Γ be a DG algebra that satisfies the conditions in **Theorem 6.1.1** for some $d \geq 1$. Following [IY18], we call $\mathcal{C}(\Gamma)$ the *Amiot–Guo–Keller (AGK) cluster category* of Γ .

In the context of **Theorem 6.1.1**, the following result of Iyama and Oppermann gives necessary and sufficient conditions for the image of Γ in $\mathcal{C}(\Gamma)$ to be a $d\mathbb{Z}$ -cluster tilting object. We introduce the following definition, simply as a matter of convenience, compare with the ‘vosnex property’ in [IO13, Not. 3.5].

Definition 6.1.3. Let Γ be a DG algebra and $d \geq 1$. We say that the cohomology of Γ *vanishes in small negative degrees (relative to d)*, if

$$\forall 0 < i < d - 1, \quad H^{-i}(\Gamma) = 0.$$

(notice that this condition is in fact vacuous for $d = 1, 2$).

Proposition 6.1.4 ([IO11, Prop. 3.6]). *Let Γ be a DG algebra that satisfies the conditions in [Theorem 6.1.1](#) for some $d \geq 1$. The following statements are equivalent:*

- (1) *The image of Γ in $\mathcal{C}(\Gamma)$ is a $d\mathbb{Z}$ -cluster tilting object.*
- (2) *The following conditions are satisfied:*
 - *The finite-dimensional algebra $H^0(\Gamma)$ is self-injective.*
 - *The cohomology of Γ vanishes in small negative degrees (relative to d).*

Our main result yields the following recognition theorem for the AGK cluster category of a DG algebra with vanishing cohomology in small negative degrees.

Theorem 6.1.5. *Let Γ be a DG algebra that satisfies the conditions in [Theorem 6.1.1](#) for some $d \geq 1$. Assume that*

- *The finite-dimensional algebra $H^0(\Gamma)$ is self-injective.*
- *The cohomology of Γ vanishes in small negative degrees (relative to d).*

(equivalently, the image of Γ in $\mathcal{C}(\Gamma)$ is a $d\mathbb{Z}$ -cluster tilting object). Then, the following statements hold:

- (1) *The d -Calabi–Yau AGK cluster category $\mathcal{C}(\Gamma)$ admits a unique DG enhancement.*
- (2) *Suppose, moreover, that the algebra $H^0(\Gamma)$ is connected and non-semisimple. Let \mathcal{T} be an algebraic triangulated category with finite-dimensional morphism spaces and split idempotents. If there exists a $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$ such that the algebras $\mathcal{T}(c, c)$ and $H^0(\Gamma)$ are isomorphic, then*

$$\mathcal{T} \simeq \mathcal{C}(\Gamma)$$

as triangulated categories.

Proof. In view of [Theorem 6.1.1](#) and [Proposition 6.1.4](#), both claims are immediate consequences of [Theorem A](#) and [Proposition 2.2.6](#). \square

Remark 6.1.6. In the literature, there are several theorems (both in the Calabi–Yau case and in the general case) that exhibit an algebraic triangulated category with a d -cluster tilting object (or even a more general type of generator) as the AGK cluster category of a homologically smooth DG algebra, see [[Tab07b](#)], [[KY](#), Thm.4.1] and [[Han22](#), Thm. 3.3]. These theorems rely on the choice of an enhancement and do not address the question of *uniqueness* of the enhancement. Notice, however, that [Theorem 6.1.5](#) includes a stringent assumption on the DG algebra being considered: Namely, its cohomology is required to vanish in small negative degrees, see also [Remark 6.1.7](#) below.

Remark 6.1.7. Suppose that \mathbf{k} is an algebraically closed field and let $d > 2$. Given an *arbitrary* basic finite-dimensional algebra A , Ladkani constructs a DG algebra Γ with $H^0(\Gamma) \cong A$ and that satisfies the conditions in [Theorem 6.1.1](#), see [[Lad](#), Thm. 3.26]. In fact, Ladkani’s methods permit to construct, for certain algebras A , DG algebras Γ and Γ' such that

$$H^0(\Gamma) \cong A \cong H^0(\Gamma'),$$

but such that $\mathcal{C}(\Gamma)$ and $\mathcal{C}(\Gamma')$ are *not* equivalent as triangulated categories, see [[Lad](#), Ex. 3.30 and Prop. 3.31]. However, Ladkani also shows that the cohomology of Γ vanishes in small negative degrees only when A is hereditary and, as it turns out, $\mathcal{C}(\Gamma)$ is equivalent as a triangulated category to the d -Calabi–Yau cluster category of A [[Kel05a](#), Sec. 6] when this is the case, see [[Lad](#), Prop. 3.18 and Ex. 3.30]. In particular, in view of the equivalent conditions in [Theorem 6.1.1](#), Ladkani’s methods yield a $d\mathbb{Z}$ -cluster tilting object $\Gamma \in \mathcal{C}(\Gamma)$ only when A is semisimple (since A must be both hereditary and self-injective).

6.2. The AGK cluster category of a d -representation finite algebra.

Setting 6.2.1. We assume that \mathbf{k} is a perfect field throughout this subsection.

We highlight a class of DG algebras that satisfy the conditions in [Theorem 6.1.5](#), and hence their associated AGK cluster category is essentially unique. These DG algebras arise naturally in the context of Iyama’s higher-dimensional Auslander–Reiten theory [[Iya07a](#), [Iya07b](#)].

The following class of algebras was introduced by Iyama and Oppermann. Our choice of terminology is inspired by [[HIO14](#), Def. 3.2 and Thm. 3.4].

Definition 6.2.2 ([[IO11](#), Def. 2.2]). Let A be a finite-dimensional algebra. We say that A is *d -representation-finite* (*d -hereditary*) if the following conditions are satisfied:

- (1) The algebra A has global dimension at most d .
- (2) There exists a *d -cluster tilting A -module* $M \in \text{mod } A$, that is such that the following conditions are satisfied:

- (a) An A -module L lies in $\text{add}(M)$ if and only if

$$\forall 0 < i < d, \quad \text{Ext}_A^i(L, M) = 0.$$

- (b) An A -module N lies in $\text{add}(M)$ if and only if

$$\forall 0 < i < d, \quad \text{Ext}_A^i(M, N) = 0.$$

Remark 6.2.3. Notice that the 1-representation finite algebras are precisely the hereditary finite-dimensional algebras of finite representation type. Thus, d -representation finite algebras can be regarded as ‘higher-dimensional analogues’ of the latter class of algebras.

Remark 6.2.4. In the context of [Definition 6.2.2](#), the d -cluster tilting module turns out to be unique, up to multiplicity of its indecomposable direct summands [[Iya11](#), Prop. 1.3].

Following Keller [[Kel11](#)], given a homologically smooth DG algebra A and an integer n , we consider its *derived n -preprojective algebra* $\mathbf{\Pi}_n(A)$, which is defined as the tensor DG algebra of the shifted inverse dualising DG A -bimodule

$$\text{RHom}_{A^e}(A, A^e)[n-1].$$

Crucially, the DG algebra $\mathbf{\Pi}_n(A)$ is homologically smooth and bimodule n -Calabi–Yau [[Kel11](#), Thm. 4.8].

Notice that we may apply the above construction when A is a finite-dimensional algebra of global dimension at most d , viewed as a DG algebra concentrated in degree 0 (recall that we assume \mathbf{k} to be a perfect field). In this case, we consider the derived $(d+1)$ -preprojective algebra $\mathbf{\Pi}_{d+1}(A)$ and, following Iyama and Oppermann, we also consider the (non-derived) *$(d+1)$ -preprojective algebra*

$$\Pi_{d+1}(A) := H^0(\mathbf{\Pi}_{d+1}(A)),$$

compare with [[IO13](#), Def. 2.11 and Rmk. 2.12]. As explained in the paragraph before [[Guo11](#), Cor. 4.8], the cohomology of $\mathbf{\Pi}_{d+1}(A)$ is concentrated in non-positive degrees.

Remark 6.2.5. Let A be a finite-dimensional algebra of global dimension at most d and $\mathbf{\Pi}_{d+1}(A)$ its derived $(d+1)$ -preprojective algebra. Then, the $(d+1)$ -preprojective algebra $\Pi_{d+1}(A)$ is isomorphic to the tensor algebra of the A -bimodule $\text{Ext}_A^d(DA, A)$, see [[IO11](#), Prop. 2.12] and compare with [[Guo11](#), Lemma 2.7] and the discussion afterwards.

In the context of this article, our interest in d -representation finite algebras stems from the following theorem, see also the proof of [CDIM, Prop. 8.6].

Theorem 6.2.6 ([IO13, Lemma 2.13 and Cor. 3.7]). *Let A be a d -representation finite algebra. Then, its derived $(d+1)$ -preprojective algebra $\mathbf{\Pi}_{d+1}(A)$ satisfies the equivalent conditions in [Proposition 6.1.4](#).*

We obtain the following recognition theorem for the AGK cluster category of a d -representation finite algebra.

Theorem 6.2.7. *Let A be a d -representation finite algebra and $\mathbf{\Pi}_{d+1}(A)$ its derived $(d+1)$ -preprojective algebra. Then, the following statements hold:*

- (1) *The d -Calabi–Yau AGK cluster category $\mathcal{C}(\mathbf{\Pi}_{d+1}(A))$ admits a unique DG enhancement.*
- (2) *Suppose, moreover, that the algebra $\mathbf{\Pi}_{d+1}(A)$ is connected and non-semisimple. Let \mathcal{T} be an algebraic triangulated category with finite-dimensional morphism spaces and split idempotents. If there exists a $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$ such that the algebra $\mathcal{T}(c, c)$ is isomorphic to the (non-derived) $(d+1)$ -preprojective algebra $\mathbf{\Pi}_{d+1}(A)$, then*

$$\mathcal{T} \simeq \mathcal{C}(\mathbf{\Pi}_{d+1}(A))$$

as triangulated categories.

Proof. In view of [Theorem 6.2.6](#), the claims is immediate from [Theorem 6.1.5](#) and [Proposition 2.2.10](#). \square

6.3. The Amiot cluster category of a self-injective quiver with potential.

Recall that a *quiver with potential* is a pair (Q, W) consisting of a finite quiver and a possibly-infinite linear combination W of (oriented) cycles in Q . To these data, one associates its (completed) *Jacobian algebra* $J(Q, W)$, see [DWZ08, Def. 3.1] for the precise definition. Following Herschend and Iyama [HI11], we say that (Q, W) is *self-injective* if its completed Jacobian algebra is a finite-dimensional self-injective algebra.

Given a quiver with potential (Q, W) , we are also interested in the *completed (3-Calabi–Yau) Ginzburg DG algebra* $\Gamma(Q, W)$ of (Q, W) , see [Gin] and [KY11, Sec. 2.6]. Replacing DG algebras and DG modules by their pseudo-compact counter-parts, [Theorem 6.1.1](#) holds *mutatis mutandis* for the pseudo-compact DG algebra $\Gamma(Q, W)$, see [KY11, Thm. A.27]. In particular, associated to (Q, W) there is a well-defined 2-Calabi–Yau triangulated category

$$\mathcal{C}(Q, W) = \mathcal{C}(\Gamma(Q, W)),$$

which we call the *Amiot cluster category of (Q, W)* , and the image of Γ in $\mathcal{C}(\Gamma)$ is a 2-cluster tilting object whose endomorphism algebra is isomorphic to the Jacobian algebra of (Q, W) , see also [Ami09]. Consequently, we obtain the following recognition theorem for the Amiot cluster category of a self-injective quiver with potential.

Theorem 6.3.1. *Let \mathbf{k} be an arbitrary field. Let (Q, W) be a quiver with potential whose Jacobian algebra is finite-dimensional and self-injective. Then, the following statements hold:*

- (1) *The (2-Calabi–Yau) Amiot cluster category $\mathcal{C}(Q, W)$ admits a unique DG enhancement.*
- (2) *Suppose, moreover, that the Jacobian algebra $J(Q, W)$ is connected and non-semisimple. Let \mathcal{T} be an algebraic triangulated category with finite-dimensional morphism spaces and split idempotents. If there exists a $2\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$ such that the algebra $\mathcal{T}(c, c)$ and $J(Q, W)$ are*

isomorphic, then

$$\mathcal{T} \simeq \mathcal{C}(Q, W)$$

as triangulated categories.

Proof. In view of the previous discussion, both claims are immediate consequences of [Theorems 5.1.6](#) and [5.1.8](#) and [Proposition 2.2.6](#), keeping in mind that, for $d = 2$, the condition on the cohomology of Γ vanishing in small negative degrees is vacuous. \square

Remark 6.3.2. Suppose that \mathbf{k} is an algebraically closed field. In this case, [Theorem 6.3.1](#) can be seen as an instance of [Theorem 6.2.7](#) for completed derived higher preprojective algebras. Indeed, the completed Ginzburg DG algebra of a self-injective quiver with potential is quasi-isomorphic to the completed derived 3-preprojective algebra of some 2-representation finite algebra, see the proof of [[HI11](#), Thm. 3.11(a)] and [[HI11](#), Prop. 2.4].

Remark 6.3.3. Classifying all quivers with potential whose Jacobian algebra is self-injective seems to be an intractable problem. Families of such quivers with potential are constructed in [[HI11](#), [Pas20](#)], see also [[Jas15](#), Thm. 1.3] where the self-injective cluster tilted algebras of canonical type are classified.

Remark 6.3.4. In relation to [Theorem 6.3.1](#), we note that Keller and Liu have recently announced [[KL23](#)] a proof of the following (modified) conjecture of Amiot [[Ami08](#)]: Suppose that the ground field \mathbf{k} is algebraically closed and of characteristic 0. Given a Karoubian pre-triangulated DG category \mathcal{A} (enriched in cochain complexes of pseudo-compact vector spaces) equipped with a right 2-Calabi–Yau structure in the sense of [[KS06](#)] which admits a basic 2-cluster tilting object $c \in H^0(\mathcal{A})$ with finite-dimensional endomorphism algebra, there exists a quiver with potential (Q, W) and an equivalence of triangulated categories

$$\mathcal{C}(Q, W) \xrightarrow{\simeq} H^0(\mathcal{A}), \quad \Gamma(Q, W) \mapsto c.$$

In particular, $H^0(\mathcal{A}(c, c)) \cong H^0(\Gamma(Q, W)) = J(Q, W)$. Notice, however, the explicit assumption on the enhancement. On the other hand, Keller’s theorem is more general than [Theorem 6.1.5](#) in that the object c need not be $2\mathbb{Z}$ -cluster tilting and its endomorphism algebra is known to be isomorphic to a completed Jacobian algebra only *a posteriori*, and this is one place where the existence of a right 2-Calabi–Yau structure plays an important role in Keller’s proof (compare with [[VdB15](#)]).

6.4. Cohen–Macaulay modules for one-dimensional hypersurface singularities. Let (R, \mathfrak{m}) be a local complete d -dimensional commutative noetherian Gorenstein isolated singularity. We are interested in the category

$$\text{CM}(R) := \{M \in \text{mod}(R) \mid \text{depth } M = \dim R\}$$

of *maximal Cohen–Macaulay R -modules*; this is a Frobenius exact category whose projective objects are the projective R -modules. Moreover, it is well known that there is an equivalence of triangulated categories [[Buc21](#)]

$$\underline{\text{CM}}(R) \xrightarrow{\simeq} \text{D}_{\text{sing}}(R) := \text{D}^b(R) / \text{D}^c(R)$$

between the stable category of maximal Cohen–Macaulay R -modules and the singularity category of R . The following result of Burban, Iyama, Keller and Reiten gives a large class of one-dimensional hypersurface singularities whose stable category of maximal Cohen–Macaulay modules admits a $2\mathbb{Z}$ -cluster tilting object.

Theorem 6.4.1 ([[BIKR08](#), Thm. 1.5]). *Let \mathbf{k} be an algebraically closed field of characteristic zero. Let $R = \mathbf{k}[[x, y]]/(f)$ be a one-dimensional reduced hypersurface singularity. The following statements are equivalent:*

- (1) The stable category $\underline{\mathbf{CM}}(R)$ of Cohen–Macaulay R -modules admits a $2\mathbb{Z}$ -cluster tilting object.
- (2) f is a product $f = f_1 f_2 \cdots f_n$ with $f_i \notin (x, y)^2$.

Remark 6.4.2. In [BIKR08], the authors only discuss 2-cluster tilting objects. However, for hypersurface singularities the stable category of Cohen–Macaulay modules is 2-periodic, that is the shift functor squares to the identity up to natural isomorphism [Eis80] (see also [BIKR08, Sec. 1]). In particular, all 2-cluster tilting objects are also $2\mathbb{Z}$ -cluster tilting.

Theorem 6.4.3. *Let \mathbf{k} be an algebraically closed field of characteristic zero. Let $R = \mathbf{k}\llbracket x, y \rrbracket / (f)$ be a one-dimensional reduced hypersurface singularity such that f is a product $f = f_1 f_2 \cdots f_n$ with $f_i \notin (x, y)^2$. Then, the stable category $\underline{\mathbf{CM}}(R)$ of Cohen–Macaulay R -modules admits a unique enhancement.*

Proof. In view of [Theorem 6.4.1](#), the claim follows immediately from [Theorem A](#). \square

6.5. Stable categories of self-injective higher Nakayama algebras. Given integers $n \geq 1$ and $\ell \geq 2$, using results of Darpö and Iyama [DI20], Külshammer and the first-named author introduced in [JK19] a family of finite-dimensional algebras

$$\tilde{A}_{n-1, \ell}^{(d)}, \quad n, \ell \geq 1,$$

called the *self-injective d -Nakayama algebras*. For example, if $d = 1$, then $\tilde{A}_{n-1, \ell}^{(1)}$ is the classical (connected) Nakayama algebra with n -simples and indecomposable projective modules of Loewy length ℓ . On the other hand, if $d = 2$ and $n = 1$, then $\tilde{A}_{0, \ell}^{(2)}$ is isomorphic to the Gelfand–Ponomarev preprojective algebra of Dynkin type A_ℓ . More generally, if $d \geq 2$ and $n = 1$, then $\tilde{A}_{0, \ell}^{(d)}$ is isomorphic to the *d -preprojective algebra of Dynkin type A_ℓ* first investigated by Iyama and Oppermann in [IO11, Sec. 5] (see also [GKO13, 6.3 and 6.5]).

Theorem 6.5.1 ([JK19, Thm. 4.10]). *Let $n \geq 1$ and $\ell \geq 2$ and $A = \tilde{A}_{n-1, \ell}^{(d)}$ the corresponding self-injective d -Nakayama algebra. The following statements hold:*

- (1) The algebra A is indeed a finite-dimensional self-injective algebra.
- (2) There exists a distinguished $d\mathbb{Z}$ -cluster tilting A -module $M = M_{n-1, \ell}^{(d)}$.
- (3) The triangulated category $\underline{\mathbf{mod}}(A)$ admits a $d\mathbb{Z}$ -cluster tilting object (namely, the image of M).

We obtain the following recognition theorem for the stable module category of a self-injective d -Nakayama algebra.

Theorem 6.5.2. *Let \mathbf{k} be an arbitrary field. Let $n \geq 1$ and $\ell \geq 2$ and $A = \tilde{A}_{n-1, \ell}^{(d)}$ the corresponding self-injective d -Nakayama algebra. The following statements hold:*

- (1) The stable module category $\underline{\mathbf{mod}}(A)$ admits a unique enhancement.
- (2) Let \mathcal{T} be an algebraic triangulated category with finite-dimensional morphism spaces and split idempotents. If there exists a $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$ such that the algebra $\mathcal{T}(c, c)$ is isomorphic to the stable endomorphism algebra $\underline{\mathbf{End}}_A(M)$ of the distinguished $d\mathbb{Z}$ -cluster tilting A -module M . Then,

$$\mathcal{T} \simeq \underline{\mathbf{mod}}(A)$$

as triangulated categories.

Proof. In view of [Theorem 6.5.1](#), both claims are immediate from [Theorem A](#). Indeed, using the explicit description of the endomorphism algebra of the distinguished $d\mathbb{Z}$ -cluster tilting A -module M given in [JK19, Def. 4.9], it is easy to verify

that there is an isomorphism of algebras

$$\underline{\text{End}}_A(M) \cong A_{n-1, \ell-1}^{(d+1)},$$

that is the length parameter ℓ is reduced by 1 while the dimension parameter d increases by 1. In particular, $\underline{\text{End}}_A(M)$ is isomorphic to the bounded path algebra of a finite quiver. \square

6.6. Singularity categories of Iwanaga–Gorenstein algebras with a $d\mathbb{Z}$ -cluster tilting module. Recall that a finite-dimensional algebra A is *Iwanaga–Gorenstein* if

$$\text{inj. dim } A_A < \infty \quad \text{and} \quad \text{inj. dim } {}_A A < \infty.$$

We are interested in the category

$$\text{Gproj}(A) := \{M \in \text{mod}(A) \mid \forall i > 0, \text{Ext}_A^i(M, A) = 0\}$$

of (finite-dimensional) *Gorenstein projective A -modules*; this is a Frobenius exact category whose projective objects are the projective A -modules. Moreover, it is well known that there is an equivalence of triangulated categories [Buc21]

$$\underline{\text{Gproj}}(A) \xrightarrow{\sim} \text{D}_{\text{sing}}(A) := \text{D}^b(A) / \text{D}^c(A)$$

between the stable category of Gorenstein projective A -modules and the singularity category of A . We recall the following theorem of Kvamme.

Theorem 6.6.1 ([Kva21, Cor. 1.3]). *Let A be a finite-dimensional Iwanaga–Gorenstein algebra. Suppose that there exists $d\mathbb{Z}$ -cluster tilting A -module $M \in \text{mod}(A)$ and let \overline{M} be the largest direct summand of M that is Gorenstein projective. Then, \overline{M} is a $d\mathbb{Z}$ -cluster tilting object in $\underline{\text{Gproj}}(A) \simeq \text{D}_{\text{sing}}(A)$.*

We obtain the following recognition theorem for the singularity category of an Iwanaga–Gorenstein algebra with a $d\mathbb{Z}$ -cluster tilting module.

Theorem 6.6.2. *Let \mathbf{k} be an arbitrary field. Let A be a finite-dimensional Iwanaga–Gorenstein algebra. Suppose that there exists a $d\mathbb{Z}$ -cluster tilting A -module $M \in \text{mod}(A)$ and let \overline{M} be the largest direct summand of M that is Gorenstein projective. The following statements hold:*

- (1) *The singularity category $\text{D}_{\text{sing}}(A)$ admits a unique enhancement.*
- (2) *Suppose, moreover, that the algebra $\underline{\text{End}}_A(\overline{M})$ is connected and non-semisimple. Let \mathcal{T} be an algebraic triangulated category with finite-dimensional morphism spaces and split idempotents. If there exists a $d\mathbb{Z}$ -cluster tilting object $c \in \mathcal{T}$ such that the algebra $\mathcal{T}(c, c)$ is isomorphic to the stable endomorphism algebra $\underline{\text{End}}_A(\overline{M})$, then*

$$\mathcal{T} \simeq \text{D}_{\text{sing}}(A)$$

as triangulated categories.

Proof. In view of [Theorem 6.6.1](#), both claims are immediate from [Theorem A](#). \square

For instances where [Theorem 6.6.2](#) can be applied see [McM20, Xin23] for example.

APPENDIX A. THE DONOVAN–WEMYSS CONJECTURE, BY BERNHARD KELLER

We work over the field of complex numbers $\mathbf{k} = \mathbb{C}$. A *compound Du Val (=cDV) singularity* is a singularity of the form

$$R = \mathbb{C}[[u, v, x, y]] / (f(u, v, x) + yg(u, v, x, y))$$

such that $\mathbb{C}[[u, v, x]]/(f(u, v, x))$ is a Kleinian surface singularity, cf. Definition 4.2 in [Aug20]. These singularities were introduced by Miles Reid [Rei83] at the beginning of the eighties and play an important role in the minimal model program in birational geometry. We refer to the introduction of [Aug19] as well as to [Wem23] for excellent introductory surveys on this subject.

From now on, we make the blanket assumption that all the cDV singularities we consider are *isolated* and admit a (*smooth*) *resolution* because the conjecture only applies to these.

Let us fix a cDV singularity R . Before stating the conjecture, let us list the most important properties of its singularity category

$$\mathrm{sg}(R) = \mathrm{D}_{\mathrm{sing}}(R) = \mathrm{D}^{\mathrm{b}}(R)/\mathrm{D}^{\mathrm{c}}(R).$$

Notice first that since R is a hypersurface, it is Gorenstein and the canonical functor

$$\mathrm{CM}(R) \rightarrow \mathrm{mod} R \rightarrow \mathrm{D}^{\mathrm{b}}(R)$$

induces an equivalence from the stable category of Cohen-Macaulay modules $\underline{\mathrm{CM}}(R)$ to $\mathrm{sg}(R)$, cf. [Buc21]. Since the singularity is isolated, the category $\mathrm{sg}(R)$ is Hom-finite. Moreover, it is Krull–Schmidt with split idempotents since R is complete. By definition, it is algebraic. Since R is a hypersurface, we can describe $\mathrm{sg}(R)$ using matrix factorisations and thus, the square Σ^2 of the suspension functor is isomorphic to the identity functor, i.e. the category $\mathrm{sg}(R)$ is 2-periodic. Finally, it is 2-Calabi–Yau since it is the stable category of Cohen–Macaulay modules over a Gorenstein ring of Krull dimension 3, cf. Prop. 1.3 in Ch. III of [Aus78].

A *contraction algebra* for R is [DW16, Wem18] the stable endomorphism algebra of a 2-cluster-tilting object T of the singularity category $\mathrm{sg}(R)$. Now let R_1 and R_2 be two cDV singularities and let A_i be a contraction algebra for R_i , $i = 1, 2$. In its original form the Donovan–Wemyss conjecture states

Conjecture A.1 ([DW16]). *Suppose that A_1 and A_2 are local. Then A_1 is isomorphic to A_2 if and only if R_1 is isomorphic to R_2 .*

The sufficiency was shown in [DW16]. In type A, the conjecture follows from the work of Reid [Rei83]. In type D, evidence is given in [BW18, vG, Kaw]. Further evidence comes from [Hua18, HT18, HK, Boo21], where the authors use enhancements of the contraction algebra.

Later Donovan–Wemyss generalized their conjecture as follows for not necessarily local contraction algebras:

Conjecture A.2. *The contraction algebras A_1 and A_2 are derived equivalent if and only if the singularities R_1 and R_2 are isomorphic.*

In this form, the conjecture appears as Conjecture 1.3 in [Aug20]. The sufficiency follows by combining results of [Wem18] with [Dug15]. We will now deduce the necessity from **Theorem A**. Let us emphasize that our strategy of proof has been known to the experts since the appearance of [HK] on the arXiv in October 2018. The missing puzzle piece was precisely (a special case of) **Theorem A**.

Let R_i be cDV singularities with contraction algebra A_i , $i = 1, 2$, and suppose that A_1 is derived equivalent to A_2 . Then, by Theorem 1.6 of [Aug20], there is a contraction algebra A'_2 for R_2 which is isomorphic to A_1 . Thus, we may and will assume that A_1 and A_2 are isomorphic. By definition, the algebras A_i are endomorphism algebras of 2-cluster-tilting objects T_i in $\mathrm{sg}(R_i)$. By 2-periodicity, these are in fact $2\mathbb{Z}$ -cluster-tilting objects. The 2-periodicity also yields the commutative

square

$$\begin{array}{ccc} \mathrm{sg}(R_i) & \longleftarrow \mathrm{add}(T_i) \xrightarrow{\mathrm{Hom}(T_i, ?)} & \mathrm{proj}(A_i) \\ \downarrow \Sigma^2 & & \downarrow ? \otimes_{A_i} \sigma_i A_i \\ \mathrm{sg}(R_i) & \longleftarrow \mathrm{add}(T_i) \xrightarrow{\mathrm{Hom}(T_i, ?)} & \mathrm{proj}(A_i) \end{array}$$

where σ_i is the identity automorphism of A_i . Now clearly the given isomorphism $A_1 \xrightarrow{\sim} A_2$ yields an equivalence between the pairs (A_1, σ_1) and (A_2, σ_2) . Thus, by [Theorem A](#) applied in dimension $d = 2$, we obtain a triangle equivalence

$$\mathrm{sg}(R_1) \xrightarrow{\sim} \mathrm{sg}(R_2).$$

Moreover, by the uniqueness of the dg enhancement in [Theorem A](#), we even obtain an isomorphism in the homotopy category of dg categories

$$\mathrm{sg}_{dg}(R_1) \xrightarrow{\sim} \mathrm{sg}_{dg}(R_2),$$

where $\mathrm{sg}_{dg}(R_i)$ denotes the canonically dg enhanced singularity category. Using [Theorem 5.9](#) of [[HK](#)], we deduce that there is an isomorphism $R_1 \cong R_2$.

REFERENCES

- [AI12] Takuma Aihara and Osamu Iyama. Silting mutation in triangulated categories. *J. Lond. Math. Soc. (2)*, 85(3):633–668, 2012.
- [AIR15] Claire Amiot, Osamu Iyama, and Idun Reiten. Stable categories of Cohen-Macaulay modules and cluster categories. *Amer. J. Math.*, 137(3):813–857, 2015.
- [Ami07] Claire Amiot. On the structure of triangulated categories with finitely many indecomposables. *Bull. Soc. Math. France*, 135(3):435–474, 2007.
- [Ami08] Claire Amiot. *Sur les petites catégories triangulées*. PhD thesis, Université Paris 7, July 2008.
- [Ami09] Claire Amiot. Cluster categories for algebras of global dimension 2 and quivers with potential. *Ann. Inst. Fourier (Grenoble)*, 59(6):2525–2590, 2009.
- [Ang11] Vignleik Angeltveit. Uniqueness of Morava K -theory. *Compos. Math.*, 147(2):633–648, 2011.
- [Ant] Benjamin Antieau. On the uniqueness of infinity-categorical enhancements of triangulated categories, arXiv:1812.01526 [math.AG].
- [AR75] Maurice Auslander and Idun Reiten. Representation theory of Artin algebras. III. Almost split sequences. *Comm. Algebra*, 3:239–294, 1975.
- [AR78] Maurice Auslander and Idun Reiten. Representation theory of Artin algebras. VI. A functorial approach to almost split sequences. *Comm. Algebra*, 6(3):257–300, 1978.
- [AR91] Maurice Auslander and Idun Reiten. On a theorem of E. Green on the dual of the transpose. In *Representations of finite-dimensional algebras (Tsukuba, 1990)*, volume 11 of *CMS Conf. Proc.*, pages 53–65. Amer. Math. Soc., Providence, RI, 1991.
- [AR96] Maurice Auslander and Idun Reiten. D tr-periodic modules and functors. In *Representation theory of algebras (Cocoyoc, 1994)*, volume 18 of *CMS Conf. Proc.*, pages 39–50. Amer. Math. Soc., Providence, RI, 1996.
- [AS80] M. Auslander and Sverre O. Smalø. Preprojective modules over Artin algebras. *J. Algebra*, 66(1):61–122, 1980.
- [AS81] M. Auslander and Sverre O. Smalø. Almost split sequences in subcategories. *J. Algebra*, 69(2):426–454, 1981.
- [AS93] M. Auslander and Ø. Solberg. Gorenstein algebras and algebras with dominant dimension at least 2. *Comm. Algebra*, 21(11):3897–3934, 1993.
- [Aug19] Jenny August. The tilting theory of contraction algebras. Ph. D. thesis, University of Edinburgh, 2019.
- [Aug20] Jenny August. On the finiteness of the derived equivalence classes of some stable endomorphism rings. *Math. Z.*, 296(3-4):1157–1183, 2020.
- [Aus71] Maurice Auslander. *Representation dimension of Artin algebras*. Lecture Notes. Queen Mary College, 1971.
- [Aus78] Maurice Auslander. Functors and morphisms determined by objects. In *Representation theory of algebras (Proc. Conf., Temple Univ., Philadelphia, Pa., 1976)*, Lecture Notes in Pure Appl. Math., Vol. 37, pages 1–244. Dekker, New York, 1978.

- [Bal02] P. Balmer. Triangulated categories with several triangulations. Available online at the author's website: <https://www.math.ucla.edu/~balmer/Pubfile/TriangulationS.pdf>, 2002.
- [BD89] Hans Joachim Baues and Winfried Dreckmann. The cohomology of homotopy categories and the general linear group. *K-Theory*, 3(4):307–338, 1989.
- [Bel15] Apostolos Beligiannis. Relative homology, higher cluster-tilting theory and categorified Auslander-Iyama correspondence. *J. Algebra*, 444:367–503, 2015.
- [BGL87] Dagmar Baer, Werner Geigle, and Helmut Lenzing. The preprojective algebra of a tame hereditary Artin algebra. *Comm. Algebra*, 15(1-2):425–457, 1987.
- [BIKR08] Igor Burban, Osamu Iyama, Bernhard Keller, and Idun Reiten. Cluster tilting for one-dimensional hypersurface singularities. *Adv. Math.*, 217(6):2443–2484, 2008.
- [BIRS09] A. B. Buan, O. Iyama, I. Reiten, and J. Scott. Cluster structures for 2-Calabi-Yau categories and unipotent groups. *Compos. Math.*, 145(4):1035–1079, 2009.
- [BK72] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin-New York, 1972.
- [BK89] A. I. Bondal and M. M. Kapranov. Representable functors, Serre functors, and reconstructions. *Izv. Akad. Nauk SSSR Ser. Mat.*, 53(6):1183–1205, 1337, 1989.
- [BK90] A. I. Bondal and M. M. Kapranov. Framed triangulated categories. *Mat. Sb.*, 181(5):669–683, 1990.
- [BKS04] David Benson, Henning Krause, and Stefan Schwede. Realizability of modules over Tate cohomology. *Trans. Amer. Math. Soc.*, 356(9):3621–3668, 2004.
- [BLM08] Yu. Bernalov, V. Lyubashenko, and O. Manzyuk. *Pretriangulated A_∞ -categories*, volume 76 of *Proceedings of Institute of Mathematics of NAS of Ukraine. Mathematics and its Applications*. Institute of Mathematics of NAS of Ukraine, Kiev, 2008.
- [BMFM20] Urtzi Buijs, José M. Moreno-Fernández, and Aniceto Murillo. A_∞ structures and Massey products. *Mediterr. J. Math.*, 17(1):Paper No. 31, 15, 2020.
- [BMR⁺06] Aslak Bakke Buan, Robert Marsh, Markus Reineke, Idun Reiten, and Gordana Todorov. Tilting theory and cluster combinatorics. *Adv. Math.*, 204(2):572–618, 2006.
- [BO01] Alexei Bondal and Dmitri Orlov. Reconstruction of a variety from the derived category and groups of autoequivalences. *Compositio Math.*, 125(3):327–344, 2001.
- [Bod14] Agnieszka Bodzenta-Skibińska. *DG Categories and Derived Categories of Coherent Sheaves*. PhD thesis, University of Warsaw, 2014.
- [Bol84] Michael L. Bolla. Isomorphisms between endomorphism rings of progenerators. *J. Algebra*, 87(1):261–281, 1984.
- [Boo21] Matt Booth. Singularity categories via the derived quotient. *Adv. Math.*, 381:Paper No. 107631, 56, 2021.
- [BS01] Paul Balmer and Marco Schlichting. Idempotent completion of triangulated categories. *J. Algebra*, 236(2):819–834, 2001.
- [BT13] Petter Andreas Bergh and Marius Thauale. The axioms for n -angulated categories. *Algebr. Geom. Topol.*, 13(4):2405–2428, 2013.
- [Buc21] Ragnar-Olaf Buchweitz. *Maximal Cohen-Macaulay modules and Tate cohomology*, volume 262 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, [2021] ©2021. With appendices and an introduction by Luchezar L. Avramov, Benjamin Briggs, Srikanth B. Iyengar and Janina C. Letz.
- [BW18] Gavin Brown and Michael Wemyss. Gopakumar-Vafa invariants do not determine flops. *Comm. Math. Phys.*, 361(1):143–154, 2018.
- [CDIM] Aaron Chan, Erik Darpö, Osamu Iyama, and René Marczinzik. Periodic trivial extension algebras and fractionally Calabi–Yau algebras, arXiv:2012.11927 [math.RT].
- [CF17] J. Daniel Christensen and Martin Frankland. Higher Toda brackets and the Adams spectral sequence in triangulated categories. *Algebr. Geom. Topol.*, 17(5):2687–2735, 2017.
- [Che21] Justin Chen. Surjections of unit groups and semi-inverses. *J. Commut. Algebra*, 13(3):323–331, 2021.
- [CNS22] Alberto Canonaco, Amnon Neeman, and Paolo Stellari. Uniqueness of enhancements for derived and geometric categories. *Forum Math. Sigma*, 10:Paper No. e92, 65, 2022.
- [Coh68] Joel M. Cohen. The decomposition of stable homotopy. *Ann. of Math. (2)*, 87:305–320, 1968.
- [CS18] Alberto Canonaco and Paolo Stellari. Uniqueness of dg enhancements for the derived category of a Grothendieck category. *J. Eur. Math. Soc. (JEMS)*, 20(11):2607–2641, 2018.
- [DI20] Erik Darpö and Osamu Iyama. d -representation-finite self-injective algebras. *Adv. Math.*, 362:106932, 50, 2020.

- [DJL21] Tobias Dyckerhoff, Gustavo Jasso, and Yankı Lekili. The symplectic geometry of higher Auslander algebras: Symmetric products of disks. *Forum of Mathematics, Sigma*, 9:e10, 2021.
- [DK08] Raika Dehy and Bernhard Keller. On the combinatorics of rigid objects in 2-Calabi-Yau categories. *Int. Math. Res. Not. IMRN*, (11):Art. ID rnn029, 17, 2008.
- [DS07] Daniel Dugger and Brooke Shipley. Topological equivalences for differential graded algebras. *Adv. Math.*, 212(1):37–61, 2007.
- [Dug10] Alex S. Dugas. Periodic resolutions and self-injective algebras of finite type. *J. Pure Appl. Algebra*, 214(6):990–1000, 2010.
- [Dug12] Alex Dugas. Periodicity of d -cluster-tilted algebras. *J. Algebra*, 368:40–52, 2012.
- [Dug15] Alex Dugas. A construction of derived equivalent pairs of symmetric algebras. *Proc. Amer. Math. Soc.*, 143(6):2281–2300, 2015.
- [DW16] Will Donovan and Michael Wemyss. Noncommutative deformations and flops. *Duke Math. J.*, 165(8):1397–1474, 2016.
- [DWZ08] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky. Quivers with potentials and their representations. I. Mutations. *Selecta Math. (N.S.)*, 14(1):59–119, 2008.
- [Eis80] David Eisenbud. Homological algebra on a complete intersection, with an application to group representations. *Trans. Amer. Math. Soc.*, 260(1):35–64, 1980.
- [EKMM07] A. Elmendorf, I. Kriz, M. Mandell, and J. May. *Rings, Modules, and Algebras in Stable Homotopy Theory*, volume 47 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, Rhode Island, April 2007.
- [Eno81] Edgar E. Enochs. Injective and flat covers, envelopes and resolvents. *Israel J. Math.*, 39(3):189–209, 1981.
- [ERZ57] Samuel Eilenberg, Alex Rosenberg, and Daniel Zelinsky. On the dimension of modules and algebras. VIII. Dimension of tensor products. *Nagoya Math. J.*, 12:71–93, 1957.
- [ES06] Karin Erdmann and Andrzej Skowroński. The stable Calabi-Yau dimension of tame symmetric algebras. *J. Math. Soc. Japan*, 58(1):97–128, 2006.
- [ES08] Karin Erdmann and Andrzej Skowroński. Periodic algebras. In *Trends in representation theory of algebras and related topics*, EMS Ser. Congr. Rep., pages 201–251. Eur. Math. Soc., Zürich, 2008.
- [Fed19] Francesca Fedele. Auslander-Reiten $(d + 2)$ -angles in subcategories and a $(d + 2)$ -angulated generalisation of a theorem by Brüning. *J. Pure Appl. Algebra*, 223(8):3554–3580, 2019.
- [Fre66] Peter Freyd. Stable homotopy. In *Proc. Conf. Categorical Algebra (La Jolla, Calif., 1965)*, pages 121–172. Springer, New York, 1966.
- [Fre09] Benoit Fresse. *Operadic cobar constructions, cylinder objects and homotopy morphisms of algebras over operads*, volume 504 of *Contemp. Math.*, page 125–188. Amer. Math. Soc., Providence, RI, 2009.
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [Ger63] Murray Gerstenhaber. The cohomology structure of an associative ring. *Ann. of Math. (2)*, 78:267–288, 1963.
- [GHW21] Eugene Gorsky, Matthew Hogancamp, and Paul Wedrich. Derived Traces of Soergel Categories. *International Mathematics Research Notices*, 04 2021. rnab019.
- [Gin] Victor Ginzburg. Calabi–Yau algebras, math/0612139.
- [GKO13] Christof Geiss, Bernhard Keller, and Steffen Oppermann. n -angulated categories. *J. Reine Angew. Math.*, 675:101–120, 2013.
- [GLS06] Christof Geiss, Bernard Leclerc, and Jan Schröer. Rigid modules over preprojective algebras. *Invent. Math.*, 165(3):589–632, 2006.
- [GP79] I. M. Gel’fand and V. A. Ponomarev. Model algebras and representations of graphs. *Funktsional. Anal. i Prilozhen.*, 13(3):1–12, 1979.
- [GSS03] Edward L. Green, Nicole Snashall, and Øyvind Solberg. The Hochschild cohomology ring of a selfinjective algebra of finite representation type. *Proc. Amer. Math. Soc.*, 131(11):3387–3393, 2003.
- [Guo11] Lingyan Guo. Cluster tilting objects in generalized higher cluster categories. *J. Pure Appl. Algebra*, 215(9):2055–2071, 2011.
- [Han20] Norihiro Hanihara. Auslander correspondence for triangulated categories. *Algebra Number Theory*, 14(8):2037–2058, 2020.
- [Han22] Norihiro Hanihara. Morita theorem for hereditary Calabi-Yau categories. *Adv. Math.*, 395:Paper No. 108092, 60, 2022.
- [Hap88] Dieter Happel. *Triangulated categories in the representation theory of finite-dimensional algebras*, volume 119 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1988.

- [Hel68] Alex Heller. Stable homotopy categories. *Bull. Amer. Math. Soc.*, 74:28–63, 1968.
- [Her16] Stephen Hermes. Minimal model of Ginzburg algebras. *J. Algebra*, 459:389–436, 2016.
- [HI11] Martin Herschend and Osamu Iyama. Selfinjective quivers with potential and 2-representation-finite algebras. *Compos. Math.*, 147(6):1885–1920, 2011.
- [HIMO] Martin Herschend, Osamu Iyama, Hiroyuki Minamoto, and Steffen Oppermann. Representation theory of Geigle–Lenzing complete intersections, 1409.0668. To appear in *Mem. Amer. Math. Soc.*
- [HIO14] Martin Herschend, Osamu Iyama, and Steffen Oppermann. n -representation infinite algebras. *Adv. Math.*, 252:292–342, 2014.
- [HJR10] Thorsten Holm, Peter Jørgensen, and Raphaël Rouquier, editors. *Triangulated categories*, volume 375 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2010.
- [HK] Zheng Hua and Bernhard Keller. Cluster categories and rational curves, arXiv:1810.00749 [math.AG]. Accepted for publication in *Geom. Topol.*
- [HKK17] F. Haiden, L. Katzarkov, and M. Kontsevich. Flat surfaces and stability structures. *Publ. Math. Inst. Hautes Études Sci.*, 126(1):247–318, 2017.
- [Hov99] Mark Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [HPS97] Mark Hovey, John H. Palmieri, and Neil P. Strickland. Axiomatic stable homotopy theory. *Mem. Amer. Math. Soc.*, 128(610):x+114, 1997.
- [HT18] Zheng Hua and Yukinobu Toda. Contraction algebra and invariants of singularities. *Int. Math. Res. Not. IMRN*, (10):3173–3198, 2018.
- [Hua18] Zheng Hua. Contraction algebra and singularity of three-dimensional flopping contraction. *Math. Z.*, 290(1-2):431–443, 2018.
- [IJ17] Osamu Iyama and Gustavo Jasso. Higher Auslander correspondence for dualizing R -varieties. *Algebr. Represent. Theory*, 20(2):335–354, 2017.
- [IO11] Osamu Iyama and Steffen Oppermann. n -representation-finite algebras and n -APR tilting. *Trans. Amer. Math. Soc.*, 363(12):6575–6614, 2011.
- [IO13] Osamu Iyama and Steffen Oppermann. Stable categories of higher preprojective algebras. *Adv. Math.*, 244:23–68, 2013.
- [IS18] Osamu Iyama and Øyvind Solberg. Auslander-Gorenstein algebras and precluster tilting. *Adv. Math.*, 326:200–240, 2018.
- [IT13] Osamu Iyama and Ryo Takahashi. Tilting and cluster tilting for quotient singularities. *Math. Ann.*, 356(3):1065–1105, 2013.
- [IV14] S. O. Ivanov and Y. V. Volkov. Stable Calabi-Yau dimension of self-injective algebras of finite type. *J. Algebra*, 413:72–99, 2014.
- [IW14] Osamu Iyama and Michael Wemyss. Maximal modifications and Auslander-Reiten duality for non-isolated singularities. *Invent. Math.*, 197(3):521–586, 2014.
- [IY08] Osamu Iyama and Yuji Yoshino. Mutation in triangulated categories and rigid Cohen-Macaulay modules. *Invent. Math.*, 172(1):117–168, 2008.
- [IY18] Osamu Iyama and Dong Yang. Silting reduction and Calabi-Yau reduction of triangulated categories. *Trans. Amer. Math. Soc.*, 370(11):7861–7898, 2018.
- [Iya07a] Osamu Iyama. Auslander correspondence. *Adv. Math.*, 210(1):51–82, 2007.
- [Iya07b] Osamu Iyama. Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories. *Adv. Math.*, 210(1):22–50, 2007.
- [Iya11] Osamu Iyama. Cluster tilting for higher Auslander algebras. *Adv. Math.*, 226(1):1–61, 2011.
- [Jas15] Gustavo Jasso. τ^2 -stable tilting complexes over weighted projective lines. *Adv. Math.*, 273:1–31, 2015.
- [Jas16] Gustavo Jasso. n -abelian and n -exact categories. *Math. Z.*, 283(3-4):703–759, 2016.
- [JK19] Gustavo Jasso and Julian Külshammer. Higher Nakayama algebras I: Construction. *Adv. Math.*, 351:1139–1200, 2019.
- [Kad80] T. V. Kadeishvili. On the theory of homology of fiber spaces. *Uspekhi Mat. Nauk*, 35(3(213)):183–188, 1980. International Topology Conference (Moscow State Univ., Moscow, 1979).
- [Kad82] T. V. Kadeishvili. The algebraic structure in the homology of an $A(\infty)$ -algebra. *Soobshch. Akad. Nauk Gruzin. SSR*, 108(2):249–252 (1983), 1982.
- [Kad88] T. V. Kadeishvili. The structure of the $A(\infty)$ -algebra, and the Hochschild and Harrison cohomologies. *Trudy Tbiliss. Mat. Inst. Razmadze Akad. Nauk Gruzin. SSR*, 91:19–27, 1988.
- [Kaj13] Hiroshige Kajiura. On A_∞ -enhancements for triangulated categories. *J. Pure Appl. Algebra*, 217(8):1476–1503, 2013.

- [Kaw] Yujiro Kawamata. Non-commutative deformations of perverse coherent sheaves and rational curves, arXiv:2006.09547 [math.AG].
- [Kel65] G. M. Kelly. Chain maps inducing zero homology maps. *Proc. Cambridge Philos. Soc.*, 61:847–854, 1965.
- [Kel94] Bernhard Keller. Deriving DG categories. *Ann. Sci. École Norm. Sup. (4)*, 27(1):63–102, 1994.
- [Kel01] Bernhard Keller. Introduction to A -infinity algebras and modules. *Homology Homotopy Appl.*, 3(1):1–35, 2001.
- [Kel05a] Bernhard Keller. On triangulated orbit categories. *Doc. Math.*, 10:551–581, 2005.
- [Kel05b] G. M. Kelly. Basic concepts of enriched category theory. *Repr. Theory Appl. Categ.*, (10):vi+137, 2005.
- [Kel06] Bernhard Keller. On differential graded categories. In *International Congress of Mathematicians. Vol. II*, pages 151–190. Eur. Math. Soc., Zürich, 2006.
- [Kel08] Bernhard Keller. Calabi-Yau triangulated categories, 2008.
- [Kel11] Bernhard Keller. Deformed Calabi-Yau completions. *J. Reine Angew. Math.*, 654:125–180, 2011. With an appendix by Michel Van den Bergh.
- [Kel13] Bernhard Keller. The periodicity conjecture for pairs of Dynkin diagrams. *Ann. of Math. (2)*, 177(1):111–170, 2013.
- [KL23] Bernhard Keller and Junyang Liu. Calabi-yau structures on drinfeld quotients and amiot’s conjecture. February 2023, 2302.03681.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [KR08] Bernhard Keller and Idun Reiten. Acyclic Calabi-Yau categories. *Compos. Math.*, 144(5):1332–1348, 2008. With an appendix by Michel Van den Bergh.
- [Kra15] Henning Krause. Krull-Schmidt categories and projective covers. *Expo. Math.*, 33(4):535–549, 2015.
- [KS06] M. Kontsevich and Y. Soibelman. Notes on A_∞ -algebras, A_∞ -categories and non-commutative geometry. In *Homological mirror symmetry*, volume 757 of *Lecture Notes in Phys.*, pages 153–219. Springer, Berlin, 2006.
- [KV87] Bernhard Keller and Dieter Vossieck. Sous les catégories dérivées. *C. R. Acad. Sci. Paris Sér. I Math.*, 305(6):225–228, 1987.
- [Kva21] Sondre Kvanne. $d\mathbb{Z}$ -cluster tilting subcategories of singularity categories. *Math. Z.*, 297(1-2):803–825, 2021.
- [KY] Martin Kalck and Dong Yang. Relative singularity categories III: Cluster resolutions, arXiv:2006.09733 [math.RT].
- [KY11] Bernhard Keller and Dong Yang. Derived equivalences from mutations of quivers with potential. *Adv. Math.*, 226(3):2118–2168, 2011.
- [Lad] Sefi Ladkani. Finite-dimensional algebras are $(m > 2)$ -Calabi-Yau-tilted, arXiv:1603.09709 [math.RT].
- [LH] Kenji Lefèvre-Hasegawa. Sur les A -infini catégories, math/0310337.
- [Lin19] Zengqiang Lin. A general construction of n -angulated categories using periodic injective resolutions. *J. Pure Appl. Algebra*, 223(7):3129–3149, 2019.
- [Lin21] Zengqiang Lin. Idempotent completion of n -angulated categories. *Appl. Categ. Structures*, 29(6):1063–1071, 2021.
- [LO10] Valery A. Lunts and Dmitri O. Orlov. Uniqueness of enhancement for triangulated categories. *J. Amer. Math. Soc.*, 23(3):853–908, 2010.
- [Lor] Antonio Lorenzin. Formality and strongly unique enhancements, arXiv:2204.09527 [math.KT].
- [LP11] Yankı Lekili and Timothy Perutz. Fukaya categories of the torus and Dehn surgery. *Proc. Natl. Acad. Sci. USA*, 108(20):8106–8113, 2011.
- [Lur17] Jacob Lurie. Higher algebra. Available online at the author’s webpage: <https://www.math.ias.edu/~lurie/papers/HA.pdf>, May 2017.
- [LV12] Jean-Louis Loday and Bruno Vallette. *Algebraic operads*, volume 346 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, 2012.
- [Lyu11] Volodymyr Lyubashenko. Homotopy unital A_∞ -algebras. *J. Algebra*, 329:190–212, 2011.
- [May69] J. Peter May. Matric Massey products. *J. Algebra*, 12:533–568, 1969.
- [McM20] Jordan McMahon. Fabric idempotents and higher Auslander-Reiten theory. *J. Pure Appl. Algebra*, 224(8):106343, 19, 2020.
- [Mit72] Barry Mitchell. Rings with several objects. *Adv. Math.*, 8:1–161, 1972.

- [MSS07] Fernando Muro, Stefan Schwede, and Neil Strickland. Triangulated categories without models. *Invent. Math.*, 170(2):231–241, 2007.
- [Mur06] Fernando Muro. On the functoriality of cohomology of categories. *J. Pure Appl. Algebra*, 204(3):455–472, 2006.
- [Mur11] Fernando Muro. Homotopy theory of nonsymmetric operads. *Algebr. Geom. Topol.*, 11(3):1541–1599, 2011.
- [Mur14] Fernando Muro. Moduli spaces of algebras over nonsymmetric operads. *Algebr. Geom. Topol.*, 14(3):1489–1539, 2014.
- [Mur16] Fernando Muro. Cylinders for non-symmetric dg-operads via homological perturbation theory. *Journal of Pure and Applied Algebra*, 220(9):3248–3281, 2016.
- [Mur20a] Fernando Muro. Enhanced A_∞ -obstruction theory. *J. Homotopy Relat. Struct.*, 15(1):61–112, 2020.
- [Mur20b] Fernando Muro. The first obstructions to enhancing a triangulated category. *Math. Z.*, 296(1-2):719–759, 2020.
- [Mur22] Fernando Muro. Enhanced Finite Triangulated Categories. *J. Inst. Math. Jussieu*, 21(3):741–783, 2022.
- [Nee01] Amnon Neeman. *Triangulated categories*, volume 148 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001.
- [Pal08] Yann Palu. Cluster characters for 2-Calabi-Yau triangulated categories. *Ann. Inst. Fourier (Grenoble)*, 58(6):2221–2248, 2008.
- [Pas20] Andrea Pasquali. Self-injective Jacobian algebras from Postnikov diagrams. *Algebr. Represent. Theory*, 23(3):1197–1235, 2020.
- [Pup62] Dieter Puppe. On the formal structure of stable homotopy theory. In *Colloquium on algebraic topology*, pages 65–71. Aarhus Universitet, 1962.
- [Rei83] Miles Reid. Minimal models of canonical 3-folds. In *Algebraic varieties and analytic varieties (Tokyo, 1981)*, volume 1 of *Adv. Stud. Pure Math.*, pages 131–180. North-Holland, Amsterdam, 1983.
- [Rou08] Raphaël Rouquier. Dimensions of triangulated categories. *J. K-Theory*, 1(2):193–256, 2008.
- [RVdB02] I. Reiten and M. Van den Bergh. Noetherian hereditary abelian categories satisfying Serre duality. *J. Amer. Math. Soc.*, 15(2):295–366, 2002.
- [RVdB19] Alice Rizzardo and Michel Van den Bergh. A note on non-unique enhancements. *Proc. Amer. Math. Soc.*, 147(2):451–453, 2019.
- [RVdB20] Alice Rizzardo and Michel Van den Bergh. A k -linear triangulated category without a model. *Ann. of Math. (2)*, 191(2):393–437, 2020.
- [Sag08] Steffen Sagave. Universal Toda brackets of ring spectra. *Trans. Amer. Math. Soc.*, 360(5):2767–2808, 2008.
- [Sai20] Shunya Saito. Tilting objects in periodic triangulated categories, 2020, arXiv:2011.14096 [math.RT].
- [Sch01] Stefan Schwede. The stable homotopy category has a unique model at the prime 2. *Adv. Math.*, 164(1):24–40, 2001.
- [Sch02] Marco Schlichting. A note on K -theory and triangulated categories. *Invent. Math.*, 150(1):111–116, 2002.
- [Sch07] Stefan Schwede. The stable homotopy category is rigid. *Ann. of Math. (2)*, 166(3):837–863, 2007.
- [Sch10] Stefan Schwede. Algebraic versus topological triangulated categories. In *Triangulated categories*, volume 375 of *London Math. Soc. Lecture Note Ser.*, pages 389–407. Cambridge Univ. Press, Cambridge, 2010.
- [Sei08] Paul Seidel. *Fukaya categories and Picard-Lefschetz theory*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [Sei15] Paul Seidel. Homological mirror symmetry for the quartic surface. *Mem. Amer. Math. Soc.*, 236(1116):vi+129, 2015.
- [Shi02] Brooke Shipley. An algebraic model for rational S^1 -equivariant stable homotopy theory. *Q. J. Math.*, 53(1):87–110, 2002.
- [Shi07] Brooke Shipley. $H\mathbb{Z}$ -algebra spectra are differential graded algebras. *Amer. J. Math.*, 129(2):351–379, 2007.
- [SS02] Stefan Schwede and Brooke Shipley. A uniqueness theorem for stable homotopy theory. *Math. Z.*, 239(4):803–828, 2002.
- [SS03] Stefan Schwede and Brooke Shipley. Stable model categories are categories of modules. *Topology*, 42(1):103–153, 2003.
- [ST01] Paul Seidel and Richard Thomas. Braid group actions on derived categories of coherent sheaves. *Duke Math. J.*, 108(1):37–108, 2001.

- [Tab05] Gonalo Tabuada. Invariants additifs de DG-cat6gories. *Int. Math. Res. Not.*, (53):3309–3339, 2005.
- [Tab06] Gonalo Tabuada. Addendum  Invariants Additifs de dg-Cat6gories. *Int. Math. Res. Not.*, 2006:Art. ID 75853, 3, 2006.
- [Tab07a] Gonalo Tabuada. Corrections  Invariants Additifs de DG-cat6gories. *Int. Math. Res. Not. IMRN*, 2007(24):Art. ID rnm149, 17, 2007.
- [Tab07b] Gonalo Tabuada. On the structure of Calabi-Yau categories with a cluster tilting subcategory. *Doc. Math.*, 12:193–213, 2007.
- [Tac64] Hiroyuki Tachikawa. On dominant dimensions of QF-3 algebras. *Trans. Amer. Math. Soc.*, 112:249–266, 1964.
- [To611] Bertrand To6en. Lectures on dg-categories. In *Topics in algebraic and topological K-theory*, volume 2008 of *Lecture Notes in Math.*, pages 243–302. Springer, Berlin, 2011.
- [TV08] Bertrand To6en and Gabriele Vezzosi. Homotopical algebraic geometry. II. Geometric stacks and applications. *Mem. Amer. Math. Soc.*, 193(902):x+224, 2008.
- [Val14] Bruno Vallette. Algebra + homotopy = operad. In *Symplectic, Poisson, and non-commutative geometry*, volume 62 of *Math. Sci. Res. Inst. Publ.*, pages 229–290. Cambridge Univ. Press, New York, 2014.
- [VdB15] Michel Van den Bergh. Calabi-Yau algebras and superpotentials. *Selecta Math. (N.S.)*, 21(2):555–603, 2015.
- [Ver96] Jean-Louis Verdier. Des cat6gories d6riv6es des cat6gories ab6eliennes. *Ast6risque*, (239):xii+253 pp. (1997), 1996. With a preface by Luc Illusie, Edited and with a note by Georges Maltsiniotis.
- [vG] Okke van Garderen. Donaldson–Thomas invariants of length 2 flops, arXiv:2008.02591 [math.AG].
- [Wei94] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.
- [Wem18] Michael Wemyss. Flops and clusters in the homological minimal model programme. *Invent. Math.*, 211(2):435–521, 2018.
- [Wem23] Michael Wemyss. A lockdown survey on cDV singularities. In Yukari Ito, Akira Ishii, and Osamu Iyama, editors, *McKay Correspondence, Mutation and Related Topics*, volume 88 of *Adv. Stud. Pure Math.*, pages 47–93, 2023.
- [Xin23] Wei Xing. Singularity categories of higher nakayama algebras, 2023, arXiv:2306.07006.
- [Zho22] Panyue Zhou. Higher-dimensional Auslander-Reiten theory on $(d + 2)$ -angulated categories. *Glasg. Math. J.*, 64(3):527–547, 2022.
- [Zim14] Alexander Zimmermann. *Representation theory*, volume 19 of *Algebra and Applications*. Springer, Cham, 2014. A homological algebra point of view.
- [ZZ] Yu Zhou and Bin Zhu. Cotorsion pairs and t-structures in a 2–Calabi-Yau triangulated category, arXiv:1210.6424 [math.RT].

(G. Jasso) LUND UNIVERSITY, CENTRE FOR MATHEMATICAL SCIENCES, SLVEGATAN 18A, 221 00 LUND, SWEDEN

Email address: gustavo.jasso@math.lu.se

URL: <https://gustavo.jasso.info>

(B. Keller) UNIVERSIT6 PARIS CIT6, UFR DE MATH6MATIQUES, CASE 7012, BTIMENT SOPHIE GERMAIN, 8 PLACE AUR6LIE NEMOURS, 75013 PARIS CEDEX 13, FRANCE

Email address: bernhard.keller@imj-prg.fr

URL: <https://webusers.imj-prg.fr/~bernhard.keller/>

(F. Muro) UNIVERSIDAD DE SEVILLA, FACULTAD DE MATEMATICAS, DEPARTAMENTO DE LGBRA, CALLE TAFIA S/N, 41012 SEVILLA, SPAIN

Email address: fmuro@us.es

URL: <https://personal.us.es/fmuro/>