

Realistic compact stars in conformal teleparallel gravity

G.G.L. Nashed^{1,*} and Kazuharu Bamba^{2,†}

¹*Centre for Theoretical Physics, The British University in Egypt,
P.O. Box 43, El Sherouk City, Cairo 11837, Egypt*

²*Faculty of Symbiotic Systems Science, Fukushima University, Fukushima 960-1296, Japan*

(Dated: September 1, 2022)

We explore an interior solution of a physically symmetric vierbein with two unknown functions in conformal teleparallel gravity. The field equations can be described in a closed system for a particular form of the metric potentials and an appropriate anisotropic function. As a result, we find a new set of configurations consistent with observed pulsars. In particular, the boundary conditions for the interior spacetime are used for the Schwarzschild spacetime to constrain the conformal field that with a unit value through the surface of a compact object. Furthermore, we apply the present model to the pulsar $4U1608-52$ with approximated radius $R = 9.52 \pm 0.15$ km and mass $M = 1.74 \pm 0.14 M_{\odot}$. To analyze the stability, we also study the causality conditions and the adiabatic index by assuming the Tolman-Oppenheimer-Volkov equation, adiabatic index and the static state. Moreover, the consistency of the model under consideration with other pulsars is investigated.

PACS numbers: 11.30.-j; 04.50.Kd; 97.60.Jd.

Keywords: Conformal teleparallel, compact stars; stability using TOV.

* nashed@bue.edu.eg

† bamba@sss.fukushima-u.ac.jp

I. INTRODUCTION

It is one of the most important challenges to explore gravity theories with conformal invariance [1–3]. This was because conformal invariant theories are predicted to be renormalizable and thus a conceivable model of quantum gravity, and the conceivable way to reach the issues of dark energy and dark matter [4]. It is an important defect that in such a formulation, the Lagrangian includes higher derivative expressions, yielding a fourth-order theory with the Gauss-Ostrogradsky ghost. However, under specific situations, the ghost will vanish and the theories are expected to be consistent ones [5, 6].

Another method to overcome the problem of ghosts is to modify gravitational theories and consider the construction of torsional gravity, in which instead of the Riemannian geometry, the teleparallelism is used [7]. In the teleparallel spacetime, the connection is written based on the teleparallel constraints in which the curvature vanishes. In this regard, two different classes exist.

The first class is the one known as the symmetric teleparallel gravity theories constructed using symmetric teleparallel geometry. In this geometry, the connection is symmetric where the torsion is zero. Consequently, the geometry is prescribed by the spacetime non-metricity. This geometry was used first in [8] to construct the symmetric teleparallel equivalent of general relativity (STEGR). Further, the geometry of STEGR was extended to different modified structures [9–13].

In the second class, the connection is completely fixed by extra constraints of the metric compatibility. Thus, this geometry is depicted by the spacetime torsion which allows the construction of completely torsion gravitational theories. A specific model of this class is the so-called “teleparallel equivalent of general relativity (TEGR)”, that is completely equivalent to general relativity [14, 15]. This theory, TEGR, was first used by Einstein attempting to unify electromagnetic and gravitational fields [16, 17]. Moreover, TEGR theory was used as an alternative construction of Einstein’s GR in [18–25]. The TEGR theory was also employed in the frame of a metric-affine gravitational theory [26, 27]. Furthermore, an alternative TEGR theory have been constructed, namely, $f(T)$ gravity [28–45], which became a familiar theory to discuss the cosmic accelerated expansions and investigate gravitational waves [46–50]. Moreover, a non-metricity conformal symmetric teleparallel gravity have been studied in [51]. Also, new anisotropic star solutions in mimetic gravity by imposing the Tolman-Finch-Skea metric have been discussed in [52].

In this work, we concentrate on TEGR constructed in [53, 54]. There are many advantages for using TEGR theory. For instance, energy, momentum and angular momentum can be defined in a consistent manner [55–57]. Also in the extend TEGR theory, the conformal invariance should be taken into account. The main feature of physical properties is that it is scale invariance, therefore, conformal invariance can be regarded as an extension of this invariance. The conformal term will have a the coordinate dependence. We here explain the meaning of a conformal transformation by consider the spacetime (\mathcal{M}, g_{ab}) , with \mathcal{M} a smooth D -dimensional manifold, g_{ab} the metric in \mathcal{M} . We use the following conformal transformation:

$$\tilde{g}_{\mu\nu} = e^{2\xi(x)} g_{\mu\nu}(x), \quad (1)$$

with $e^{2\xi(x)}$ being a smooth function of the coordinates. Therefore, the expression $e^{2\xi(x)}$ is the conformal factor. By this transformations, angles are kept and distances between two points prescribed by the same coordinate are alerted. The transformation (1) becomes trivial when $\xi(x) = \text{const}$. Any theory that respect transformation given by Eq. (1) is called a conformal field theory in which the nature of physics has a scale invariance.

There are two explanations of the transformation (1). One is for a constant background metric and the other is for a dynamical metric. For a constant metric, the transformation can be regarded as a physical symmetry of a dynamical system. On the other hand, for a dynamical metric, the transformation has a diffeomorphism, in other words, a gauge symmetry appears. For a review on the topic of conformal field theory one can refer to [58–61]. Recently, there are many papers including conformal transformations have been studies [62–67], in the area of cosmology [68, 69]; in higher-order curvature terms [70], and in quantum gravity [71]. Our main purpose is to find a novel spherically symmetric interior solution in a conformal TEGR theory and study its relevant physics.

The organization of this paper is as follows. In Sec. II, we explain the conformal TEGR theory including mathematical definitions of tensors and field equations of gravitation. In Sec. III, the field equations are applied to a four dimensional spherically symmetric vierbein field with non-diagonal components. We study the system of differential equations with specific forms of the unknowns of the ansate and the anisotropic function and succeed to derive the remaining unknowns with three constants in Sec. III. In Sec. IV, we present the necessary conditions that must be satisfied by real compact stellar objects. In Sec. V we show that the model under consideration is non-singular in the center of a star and derive the gradient of its energy-momentum components as well as its speed of sound. In Sec. VI we apply the junction conditions and derive the form of the constants in terms of the physical quantities numerically for the stellar $4U1608 - 52$, with approximated radius $R = 9.52 \pm 0.15$ km and mass $M = 1.74 \pm 0.14 M_\odot$. Moreover, in Sec. VI we show that the model bypass a real compact stellar graphically. In Sec. VII we show that our model

is stable using different methods, Tolman-Oppenheimer-Volkoff (TOV) equation, Adriatic index, and the static state. Also in Sec. VI, we show the consistency of our model with other pulsars. Finally, the section is devoted to the arguments of the astrophysical consequences.

II. THE CONFORMAL TEGR

The TEGR geometry is described by ¹ $\{M, e_i\}$, where M is a four-dimensional manifold and e_i ($i = 1, 2, 3, 4$) is the parallel vector on M . The contra-variant derivative of the vierbein field reads

$$D_\mu e_i^\nu := \partial_\mu e_i^\nu + \Gamma^\nu_{\lambda\mu} e_i^\lambda = 0, \quad (2)$$

where the Weitzenböck connection is given by

$$\Gamma^\lambda_{\mu\nu} := e_i^\lambda \partial_\nu e^i_\mu, \quad (3)$$

with $\partial_\rho = \frac{\partial}{\partial x^\rho}$. In the TEGR theory, we can defined the metric as:

$$g_{\mu\nu} := \eta_{ij} e^i_\mu e^j_\nu, \quad (4)$$

with the Minkowski metric $\eta_{ij} = (+, -, -, -)$. The torsion $T^\alpha_{\mu\nu}$ and the contortion tensors $K^{\mu\nu}_\alpha$ are defined as

$$T^\alpha_{\mu\nu} := \Gamma^\alpha_{\nu\mu} - \Gamma^\alpha_{\mu\nu} = e_i^\alpha (\partial_\mu e^i_\nu - \partial_\nu e^i_\mu), \quad K^{\mu\nu}_\alpha := -\frac{1}{2} (T^{\mu\nu}_\alpha - T^{\nu\mu}_\alpha - T_\alpha^{\mu\nu}).$$

The contraction of the torsion yields

$$T_\nu := T^\mu_{\mu\nu}. \quad (5)$$

The torsion scalar is

$$T := T^\alpha_{\mu\nu} S_\alpha^{\mu\nu}, \quad (6)$$

where $S_\alpha^{\mu\nu}$ is the superpotential tensor whose first pair are skew symmetry. The superpotential tensor is described by:

$$S_\alpha^{\mu\nu} := \frac{1}{2} (K^{\mu\nu}_\alpha + \delta_\alpha^\mu T^{\beta\nu}_\beta - \delta_\alpha^\nu T^{\beta\mu}_\beta). \quad (7)$$

The Lagrangian is given by

$$\mathcal{L}(e^i_\mu)_g = \frac{|e|T}{2\kappa}, \quad \text{where} \quad e = \sqrt{-g}. \quad (8)$$

In this study, we use the relativistic units in which $c = G = 1$, where c is the speed of light and G is the Newtonian constant².

In the conformal formalism, the Lagrangian of Eq. (8) changes through the conformal transformation

$$\bar{e}_{a\mu} = e^{\zeta(x)} e_{a\mu}, \quad (9)$$

where $\xi(x)$ is an arbitrary function of the coordinate x [54]. Thus, Eq. (8) must be altered, so that Eq. (8) will not be changed by the conformal transformation (9). Under the conformal transformation, the Lagrangian becomes [54]

$$\mathcal{L}(e^i_\mu, \xi)_G = 2\kappa|e| [6g^{\mu\nu}\partial_\mu\xi\partial_\nu\xi - \xi^2 T - 4g^{\mu\nu}\xi(\partial_\nu\xi)T_\mu] + \mathcal{L}_m, \quad (10)$$

where ξ is a scalar field and \mathcal{L}_m is the Lagrangian of the matter. The Lagrangian (10) is invariant through

$$\xi \rightarrow \bar{\xi} = e^{-\xi(x)} \xi.$$

¹ The Latin labels i, j, \dots refer to the coordinates of the tangent spacetime and the Greek indices α, β, \dots denote the components of the (co-)frame.

² In the following, we put the coupling of the gravitational constant as $\kappa = 1$.

Making the variation of the Lagrangian (10) in terms of ξ , we find [54]

$$I \equiv \partial_\mu (e g^{\mu\nu} \partial_\nu \xi) \xi - \frac{e}{6} \xi^2 R - \frac{e \mathcal{T}}{6} = 0, \quad (11)$$

with R the scalar curvature given by $eR = 2\partial_\nu(eT^\nu) - eT$, where $|e| = e = \sqrt{-g} = \det(e^\alpha_\mu)$ the determinant of the metric and T being the scalar torsion defined from Eq. (6). Here \mathcal{T} is the trace of the energy momentum tensor defined as $\mathcal{T} = g_{\mu\nu} \mathcal{T}^{\mu\nu}$.

The variation of the Lagrangian (10) in terms of the vierbein $e_{a\mu}$ yields [54]

$$\begin{aligned} I^{a\nu} \equiv & \partial_\alpha (e \xi^2 S^{a\nu\alpha}) - e \xi^2 (S^{b\alpha\nu} T_{b\alpha}{}^a - \frac{1}{4} e^{a\nu} T) - \frac{3}{2} e e^{a\nu} g^{\beta\mu} \partial_\beta \xi \partial_\mu \xi + 3 e e^{a\mu} g^{\beta\nu} \partial_\beta \xi \partial_\mu \xi + e e^{a\nu} g^{\beta\mu} T_\mu \xi \partial_\beta \xi \\ & - e \xi e^{a\beta} g^{\nu\mu} (T_\mu \partial_\beta \xi + T_\beta \partial_\mu \xi) - e g^{\beta\mu} \xi T^{\nu a}{}_\mu \partial_\beta \xi - \partial_\mu [e g^{\beta\nu} e^{a\mu} \xi \partial_\beta \xi] + \partial_\rho [e g^{\beta\rho} e^{a\nu} \xi \partial_\beta \xi] - e^a{}_\beta \mathcal{T}^{\beta\nu} = 0. \end{aligned} \quad (12)$$

Here, $\mathcal{T}_\alpha{}^\beta = \mathcal{T}_{G\alpha}{}^\beta + \mathcal{T}_{S\alpha}{}^\beta$ with $\mathcal{T}_{G\alpha}{}^\beta$ the energy-momentum tensor of anisotropic fluid, given by

$$\mathcal{T}_{G\nu}^\mu = (p_\perp + \rho) u^\mu u_\nu + p_\perp \delta_\nu^\mu + (p_r - p_\perp) \varepsilon_a \varepsilon^\mu, \quad (13)$$

where $u^\mu = [1, 0, 0, 0]$ is a time-like vector and $\varepsilon^\mu = [0, 1, 0, 0]$ is the unit space-like vector along the radial direction, so that $u^\mu u_\mu = -1$ and $\varepsilon^\mu \varepsilon_\mu = 1$. Moreover, ρ denotes the energy density, p_r and p_\perp are the pressures of the radial and tangential directions, respectively. Furthermore, $\mathcal{T}_{S\alpha}{}^\beta$ is the energy-momentum tensor of the scalar field, represented as

$$\mathcal{T}_S^\beta{}_\nu = \partial^\beta \xi \partial_\nu \xi - \delta_\nu^\beta \left[g^{\alpha\alpha_1} \partial_\alpha \xi \partial_{\alpha_1} \xi - \frac{V(\xi)}{2} \right]. \quad (14)$$

The above equations determine the conformal TEGR gravitational theory in 4-dimension space.

III. INTERIOR SPHERICALLY SYMMETRIC SOLUTION

We explore the metric with spherical symmetry metric

$$ds^2 = -a^2(r) dt^2 + \frac{dr^2}{b^2(r)} + r^2 d\Omega^2, \quad \text{where} \quad d\Omega^2 = (d\theta^2 + \sin^2 \theta d\phi^2), \quad (15)$$

where $a(r)$ and $b(r)$ are functions of the radial coordinate r . Equation (15) can be generated using the following covariant vierbein field [72]

$$e^i{}_\mu = \begin{pmatrix} a(r) & 0 & 0 & 0 \\ 0 & \frac{\cos(\phi) \sin(\theta)}{b(r)} & r \cos(\phi) \cos(\theta) & -r \sin(\phi) \sin(\theta) \\ 0 & \frac{\sin(\phi) \sin(\theta)}{b(r)} & r \sin(\phi) \cos(\theta) & r \cos(\phi) \sin(\theta) \\ 0 & \frac{\cos(\theta)}{b(r)} & -r \sin(\theta) & 0 \end{pmatrix}. \quad (16)$$

Moreover, Eq. (16) can be rewritten in a diagonal vierbein product and a local Lorentz transformation [20].

Substituting Eq. (16) into Eq. (6), we find

$$T = \frac{2(b-1)(2a'b'r - a + ab)}{ar^2}. \quad (17)$$

Employing Eq. (17) in the field equations (11) and (12), we get

$$-1/2 \kappa^2 r^2 (2\rho - V + \xi^2 b^2) = \xi^2 b^2 - \xi^2 + 2r^2 b \xi \xi' b' + 4\xi r b^2 \xi' - r^2 b^2 \xi'^2 + 2\xi^2 r b b' + 2r^2 b^2 \xi \xi'',$$

$$1/2 \kappa^2 a r^2 (2p_r + \xi'^2 b^2 + V) = 2b^2 r^2 \xi \xi' a' + \xi^2 a b^2 + 4a r b^2 \xi \xi' + 3a r^2 b^2 \xi'^2 + 2\xi^2 b^2 r a' - \xi^2 a,$$

$$\begin{aligned} \frac{\kappa^2 ar \left(2 p_t + V - \xi'^2 b^2 \right)}{2b} &= 2 \xi \xi' r a' b + 2 \xi \xi' a r b' + \xi^2 r a' b' + 2 \xi \xi' a b + \xi^2 r b a'' - a b r \xi'^2 + \xi^2 a b' + b \xi'^2 a' + 2 a b r \xi \xi'', \\ \kappa^2 r^2 a(p_t - p_r - b^2 \xi'^2) &\equiv r^2 a(\Delta(r) - \kappa^2 b^2 \xi'^2) = 2 a r^2 b \xi \xi' b' + \xi^2 b r^2 a' b' - \xi'^2 a b^2 - 2 a r b^2 \xi \xi' + \xi^2 b^2 r^2 a'' - 4 a r^2 b^2 \xi'^2 \\ &+ \xi^2 b a b' r - \xi^2 b^2 r a' + 2 a r^2 b^2 \xi \xi'' + \xi^2 a, \end{aligned} \quad (18)$$

and the field equation of the scalar field takes the form:

$$\frac{1}{2} \kappa^2 a r^2 \left(2 V - \xi'^2 b^2 \right) = \xi \left(\xi b [2 r b a' + 2 a r b' + a b + r^2 a' b' + b r^2 a'' - a/b] + 3 \xi' r b [b r a' + a r^2 b' + 2 a b + a r b \xi''/\xi'] \right), \quad (19)$$

where the prime ' means derivatives in terms of r . The expression of the energy-momentum tensor \mathcal{T}_ν^μ is given by

$$\text{diag } \mathcal{T}_\nu^\mu = \left(-\rho + \frac{V(r)}{2} - \frac{\xi'^2 b^2}{2}, p_r + \frac{V(r)}{2} + \frac{\xi'^2 b^2}{2}, p_t + \frac{V(r)}{2} + \frac{\xi'^2 b^2}{2}, p_t + \frac{V(r)}{2} + \frac{\xi'^2 b^2}{2} \right), \quad (20)$$

where $V(r)$ is the potential field and ξ is the conformal field. $\Delta(r)$ in (18) is a parameter to express the anisotropy of compact objects. It is confirmed that Eqs. (18) are equivalent to those shown in [20, 73, 74] for $\xi(r) = V(r) = 0$.

The above system of differential equations consist of five independent equations in eight unknown functions, $a(r)$, $b(r)$, $\rho(r)$, $p_r(r)$, $p_t(r)$, $V(r)$, $\xi(r)$ and $\Delta(r)$. Therefore, three more conditions are necessary to make this system a closed system to be solved. As two conditions among them, the forms of $a(r)$ and $b(r)$ are supposed to be:

$$a(r) = a_1 - a_0 r^2, \quad b(r) = 1 + a_0 r^2 \quad (21)$$

with a_1 a dimensionless constant and a_0 a one with the dimension of length. These constants will be determined from the junction condition to match the interior solution with the exterior one. Equation (21) indicates that when $r = 0$, $a(r) = a_1$ and $b(r) = 1$, which are finite in the center of a star. Moreover, the derivative of the two functions a and b are finite in the center. The third condition is acquired by substituting Eq. (21) into the last equation of (18), i.e., $\Delta(r)$ which yields the form³

$$\Delta(r) = (b^2 r^2 a'' - b^2 r a' + b r^2 a' b' + b a b' r + a - a b^2) \xi^2 + (2 a b r^2 \xi' b' + 2 a b^2 r^2 \xi'' - 2 a b^2 r \xi') \xi - 4 a b^2 r^2 \xi'^2 + a b^2 r^2 \xi'^2, \quad (22)$$

that is a second order differential equation in terms of ξ if

$$0 = (2 a b r^2 \xi' b' + 2 a b^2 r^2 \xi'' - 2 a b^2 r \xi') \xi - 4 a b^2 r^2 \xi'^2 + a b^2 r^2 \xi'^2, \quad (23)$$

whose solution reads

$$\begin{aligned} \xi &= \frac{16 a_0^2}{(c_1 S + 2 a_0^2)^2}, \\ \implies \Delta(r) &= r^4 a_0^2 a_1 - 4 r^4 a_0^2 - 5 r^6 a_0^3, \end{aligned} \quad (24)$$

where $S = \ln(1 + a_0 r^2)$. Using Eq. (21) as well as Eq. (24) in Eqs. (18), we get the unknowns $V(r)$, $\rho(r)$, $p_r(r)$ and $p_t(r)$ in the forms

³ From now on we are going to use the relativistic units in which $G = c = 1$, namely, $\kappa = 1$.

$$\begin{aligned}
\rho &= 128a_0^5 \left\{ 6a_1 c_1 S + 5a_0^2 r^4 c_1 S + 6c_1 S + 5r^2 a_1 a_0 c_1 S + 10c_1 S r^2 a_0 - 12c_1 r^4 a_0^2 - 12c_1 a_0 r^2 a_1 - 12c_1 a_1 - 12c_1 r^2 a_0 \right. \\
&\quad \left. + 20a_0^3 r^2 + 12a_1 a_0^2 + 10a_0^4 r^4 + 10r^2 a_1 a_0^3 + 12a_0^2 \right\} \left\{ S^5 c_1^5 r^2 a_0 - S^5 c_1^5 a_1 - 10S^4 c_1^4 a_1 a_0^2 + 10S^4 c_1^4 r^2 a_0^3 + 32r^2 a_0^{11} \right. \\
&\quad \left. + 40S^3 c_1^3 r^2 a_0^5 - 40S^3 c_1^3 a_1 a_0^4 + 80S^2 c_1^2 r^2 a_0^7 - 80S^2 c_1^2 a_1 a_0^6 + 80Sc_1 r^2 a_0^9 - 80Sc_1 a_0^8 a_1 - 32a_1 a_0^{10} \right\}^{-1}, \\
p_r &= -128a_0^5 \left\{ 2c_1 S r^2 a_0 + 5a_0^2 r^4 c_1 S - 2c_1 S - 3r^2 a_1 a_0 c_1 S + 4c_1 a_0 r^2 a_1 - 2a_1 c_1 S + 4c_1 a_1 + 4c_1 r^2 a_0 + 4c_1 r^4 a_0^2 \right. \\
&\quad \left. + 4a_0^3 r^2 - 6r^2 a_1 a_0^3 - 4a_0^2 + 10a_0^4 r^4 - 4a_1 a_0^2 \right\} \left\{ S^5 c_1^5 r^2 a_0 - S^5 c_1^5 a_1 - 10S^4 c_1^4 a_1 a_0^2 + 10S^4 c_1^4 r^2 a_0^3 - 32a_1 a_0^{10} \right. \\
&\quad \left. + 40S^3 c_1^3 r^2 a_0^5 - 40S^3 c_1^3 a_1 a_0^4 + 80S^2 c_1^2 r^2 a_0^7 - 80S^2 c_1^2 a_1 a_0^6 + 80Sc_1 r^2 a_0^9 - 80Sc_1 a_0^8 a_1 + 32r^2 a_0^{11} \right\}^{-1}, \\
p_t &= 128a_0^5 \left\{ 6c_1 S r^2 a_0 + 5a_0^2 r^4 c_1 S + 2c_1 S + r^2 a_1 a_0 c_1 S - 4c_1 a_0 r^2 a_1 + 2a_1 c_1 S - 4c_1 a_1 - 4c_1 r^2 a_0 - 4c_1 r^4 a_0^2 \right. \\
&\quad \left. + 12a_0^3 r^2 + 2r^2 a_1 a_0^3 + 4a_0^2 + 10a_0^4 r^4 + 4a_1 a_0^2 \right\} \left\{ S^5 c_1^5 r^2 a_0 - S^5 c_1^5 a_1 - 10S^4 c_1^4 a_1 a_0^2 + 10S^4 c_1^4 r^2 a_0^3 + 32r^2 a_0^{11} \right. \\
&\quad \left. + 40S^3 c_1^3 r^2 a_0^5 - 40S^3 c_1^3 a_1 a_0^4 + 80S^2 c_1^2 r^2 a_0^7 - 32a_1 a_0^{10} - 80S^2 c_1^2 a_1 a_0^6 + 80Sc_1 r^2 a_0^9 - 80Sc_1 a_0^8 a_1 \right\}^{-1}, \\
V(r) &= \frac{256a_0^5}{S} \left\{ 24a_0^4 - 6S^2 c_1^2 a_1 + 6S^2 c_1^2 - 20r^2 a_1 a_0^3 c_1 S - 20r^2 a_1 a_0^5 + 36r^2 a_0 c_1^2 a_1 S - 5r^2 a_1 a_0 c_1^2 S^2 + 80a_0^2 r^4 c_1^2 \right. \\
&\quad \left. + 36c_1^2 a_1 S + 72a_0^2 c_1 a_1 - 120r^4 a_0^4 c_1 + 24c_1 S a_0^2 - 120r^2 a_0^3 c_1 + 88r^2 a_0^5 + 60a_0^6 r^4 - 24a_1 a_0^4 + 60a_0^4 r^4 c_1 S \right. \\
&\quad \left. - 24a_1 c_1 S a_0^2 - 60r^4 a_0^2 c_1^2 S + 22r^2 a_0 c_1^2 S^2 + 88r^2 a_0^3 c_1 S - 60r^2 a_0 c_1^2 S - 80a_1 c_1^2 r^2 a_0 + 15a_0^2 r^4 c_1^2 S^2 + 72r^2 a_0^3 c_1 a_1 \right\} \\
&\quad \times \left\{ c_1^6 S^6 + 12c_1^5 S^5 a_0^2 + 60c_1^4 S^4 a_0^4 + 160c_1^3 S^3 a_0^6 + 240c_1^2 S^2 a_0^8 + 192c_1 S a_0^{10} + 64a_0^{12} \right\}^{-1}. \tag{25}
\end{aligned}$$

It is important to note that the anisotropic force is given by $\frac{2\Delta}{r}$, which is repulsive if $p_r - p_t < 0$, while it is attractive when $p_r - p_t > 0$. The mass of a sphere with its radius r is expressed by [75, 76]

$$M(r) = \int_0^r \left[4\pi\rho(\zeta) - \frac{\xi'(\zeta)^2 b(\zeta)}{2} + \frac{V(\zeta)}{2} \right] \zeta^2 d\zeta. \tag{26}$$

Combining Eq. (25) with Eq. (26), we have a very length form of $M(r)$ which has the following asymptotic form⁴:

$$M(r) \approx -6.358950030 \times 10^{11} r + 3.712536047 \times 10^{12} r^3 - 2.148986652 \times 10^{13} r^5, \tag{27}$$

With Eq. (27), we find a parameter describing the compactness of a spherically symmetric star [73]

$$u(r) \approx \frac{2M(r)}{r} = -1.271790006 \times 10^{12} + 7.425072094 \times 10^{12} r^2 - 4.297973304 \times 10^{13} r^4. \tag{28}$$

In the next section, we investigate the conditions, with which a realistic star must satisfy and test whether Eq. (25) can be met.

⁴ The numerical values of the constants a_0 , a_1 , and c_1 that will be fixed from the junction condition in Sec. IV using the pulsar 4U1608–52, whose approximated radius $R = 9.52 \pm 0.15$ km and mass $M = 1.74 \pm 0.14 M_\odot$, are used in Eq. (26).

IV. NECESSARY CONDITIONS OF A REALISTIC STAR

Necessary conditions that a realistic star must satisfy are summarized as follows.

- ★ The metric potentials $a(r)$ and $b(r)$ and the matter components $\rho(r)$, $p_r(r)$, $p_t(r)$ have finite values in the center of a star and are regular and free from singularity.
- ★ The matter density ρ should be positive in all regions of a star, $0 \leq \rho(r)$. Moreover, the value of ρ should be positive in the center and finite and it decreases across the boundary of the interior of a star, namely, $0 \geq \rho'(r) = \frac{d\rho}{dr}$.
- ★ The radial pressure p_r as well as the tangential one p_t should be positive within a star, $p_r \geq 0$, $p_t \geq 0$. Moreover, the derivative of the pressure have to be negative within a star, $\frac{dp_r}{dr} < 0$, $\frac{dp_t}{dr} < 0$. In the center of a star, the anisotropy vanish, namely, $\Delta(r \rightarrow 0) = 0$.
- ★ The requirement of the energy conditions is necessary for anisotropic fluid sphere. These conditions are described as the following forms, that is satisfied in all points of a star
 - (i) The dominant energy conditions (DEC): $\rho \geq |p_r|$ and $\rho \geq |p_t|$.
 - (ii) The strong energy condition (SEC): $p_r + \rho > 0$, $p_t + \rho > 0$, $\rho - p_r - 2p_t > 0$.
 - (iii) The weak energy condition (WEC): $p_r + \rho > 0$, $\rho > 0$.
 - (iv) The null energy condition (NEC): $p_t + \rho > 0$, $\rho > 0$.
- ★ The causality condition for a realistic star, that is, the sound speed must be smaller than unity (in the relativistic units of $c = G = 1$) in the interior of the star, that is, $1 \geq \frac{dp_t}{dr} \geq 0$ and $1 \geq \frac{dp_r}{dr} \geq 0$.
- ★ The interior solution have to match with the exterior Schwarzschild one in the star surface for $\xi = 1$ [64, 67].
- ★ The adiabatic index have to be larger than $\frac{4}{3}$ to have a stable model.
- ★ For the stability analysis of an anisotropic star, it has been suggested [77] that the relation $0 < v_t^2 - v_r^2 < 1$ must be met. Here, v_t and v_r are the transverse and radial speeds, respectively.

By using the physical conditions, we study whether the present model can satisfy the relations mentioned above.

V. THE PHYSICAL CONDITIONS OF THE PRESENT MODEL

We examine Eq. (25) and confirm that it is appropriate for a realistic star.

A. A free singular model

First, the following conditions are satisfied by the metric potentials

$$a(r \rightarrow 0) = a_1 \quad \text{and} \quad b(r \rightarrow 0) = 1. \quad (29)$$

If this condition is met, the metric potentials have finite value in the center of a star. Additionally, the gradients of the metric potentials have finite value in the center such as $a'_{r \rightarrow 0} = b'_{r \rightarrow 0} = 0$. This means that the metric potentials are not singular in the center.

Second, the components of the fluid given by Eq. (25) in the center of a star are express as

$$\rho(r \rightarrow 0) = \frac{48a_1c_1 - 48a_0^2 - 48a_1a_0^2}{a_1a_0^5}, \quad p_r(r \rightarrow 0) = p_t(r \rightarrow 0) = \frac{16a_1c_1 - 16a_0^2 - 16a_1a_0^2}{a_1a_0^5}. \quad (30)$$

The above equations show that the energy density is positive if $\frac{48a_1c_1 - 48a_0^2 - 48a_1a_0^2}{a_1a_0^5} > 0$, and that the space-time becomes isotropic in the center. Moreover, the radial and tangential pressures become positive value when $\frac{16a_1c_1 - 16a_0^2 - 16a_1a_0^2}{a_1a_0^5} > 0$, otherwise they become negative. Furthermore, in Ref. [78], radial pressure should equal or to or weaker than density in the center, $\frac{p_r(0)}{\rho(0)} \leq 1$. With the constrains $\frac{p_r(0)}{\rho(0)}$ given by Eq. (30), it is seen that this condition can be satisfied.

Third, we describe the gradient of energy density, radial and tangential pressures in Appendix (A). In the next section, we consider these quantities by fixing the numerical values of a_0 , a_1 , and c_1 .

Fourth, the radial and transverse sound speeds ($c = 1$) are presented in Appendix (B). We show these components for the concrete values of a_0 , a_1 , and c_1 .

B. Junction conditions

It is supposed that the exterior region of a non-rotating star is vacuum and is expressed by the Schwarzschild metric, given by⁵:

$$ds^2 = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2, \quad (31)$$

with M the total mass. We have to match

The interior metric (15) must be matched with the exterior Schwarzschild one (31) in the star boundary $r = l$. By using the continuity of the metric at $r = l$, we find

$$a(r \rightarrow l) = \left(1 - \frac{2M}{l}\right), \quad b(r \rightarrow l) = \left(1 - \frac{2M}{l}\right), \quad \xi(r \rightarrow l) = 1. \quad (32)$$

With these conditions, the constants a_0 , a_1 and c_1 are determined as

$$a_0 = -\frac{2M}{l^3}, \quad a_1 = -\frac{4M - l}{l}, \quad c_1 = -\frac{8M(M \pm l^3)}{l^6 \ln\left(\frac{l-2M}{l}\right)}. \quad (33)$$

We use the numerical values of these constants when we deal with observations.

VI. COMPARISON OF THEORETICAL MODEL WITH THE OBSERVATIONS OF REAL STARS

Now, we convert back to the standard physical units to accurately determine the values of the model parameters under consideration numerically. Through the use of the masses and the radii of observed pulsars with the physical conditions we referred above, one can fix the constant parameters a_0 and a_1 and check its viability. Now, we employ the observational limits of the pulsar 4U 1608-52, whose mass $M = 1.74 \pm 0.14M_{\odot}$ and $R = 9.52 \pm 0.15$ km, where $M (= 1.9891030 kg)$ denotes the solar mass. Then, the boundary conditions given by Eq. (33) are able to fix the dimensionless constants.

We adopt the conditions listed previously to compare the present model to observational data. We study the pulsar 4U 1608-52 whose mass and radius are $M = 1.74 \pm 0.14M_{\odot}$ and $R = 9.52 \pm 0.15$ km, respectively [79], and therefore we take $M = 1.78M_{\odot}$ and $R = 9.67$ km. We use the boundary conditions to examine the values of the constants $a_0 = -0.005826829242$, $a_1 = -0.08972078594$, and $c_1 = -0.02952344694$. With these constants, we depict physical quantities. We examine whether the present model can fit the observational data of realistic stars.

Figure 1 (a) shows the metric potentials for 4U 1608-52. From this figure, we see that $a(r \rightarrow 0) = -0.08972078594$, and $b(r \rightarrow 0) = 1$, both of which are finite in the center of a star.

Figures 1 (b), 1 (c) and 1 (d) plot the density, radial and tangential pressures, respectively. We see that the density, radial and tangential pressures are positive and they become large in the center and become smaller as the radii becomes larger. These behaviors are consistent with realistic stars. From figure 2 (a), we understand that the spacetime becomes isotropic in the center of a star and the anisotropy is smaller in the surface of a star. Specifically, we find the negative anisotropic force $\frac{\Delta}{r}$, that is, t is an attractive one due to the fact that $p_t - p_r < 0$. Figure 2 (b) means that the gradients of the energy-momentum components, density, radial and tangential pressures presented in Appendix (A), are negative. This ensures the decreasing of all of them as a whole of a star. In Figure 2 (c), we depict the radial and tangential speeds of sound, presented in Appendix (B). These values are positive and less than unity, so that the causality condition can be met within a star. Figure 2 (d) shows the difference between the tangential and radial speed of sounds that has a positive value as required for any realistic stellar.

Figure 3 describes the behavior of the energy conditions. Particularly, Figures. 3 (a), (b), (c) and (d) means that all of the energy conditions, WEC, NEC, SEC and DEC ones, are satisfied. In Figure 4, we have depicted the radial and

⁵ It is shown that the general solution of a non-charged spherically symmetric spacetime in conformal teleparallel theory is just the Schwarzschild solution [67].




Figure 1. Schematic plots of (a) the metric potentials; (b) the energy-density; (c) the radial pressure and (d) the tangential pressure. All of these plots are depicted vs. the radial coordinate r .

the tangential equation of state (EoS). As Fig. 4 (a) shows, the EoS is not linear contrary to what discussed in Ref. [80]. From Fig. 4 (a), it can be found that the radial and tangential EoS are non-linear, and that the mass function in Eq. (27) becomes $M(r \rightarrow 0) = 0$. Additionally, Fig. 4 (b) suggests that the compactness parameter becomes larger. Moreover, the radial variation of the surface red-shift is depicted in Fig. 4 (c). The present result of ≈ -1 for 4U 1608-52 is compatible with the constraints of the surface red-shift $Z \leq 5$ [81].

VII. STUDY OF THE STABILITY OF THE MODEL

It is the aim of this section, to study the stability problem through three different methods; the Tolman-Oppenheimer-Volkoff equations (TOV), the adiabatic index and the static state.

A. Equilibrium process using the TOV equation

We explore a hydrostatic equilibrium with the TOV equation [82–84]

$$\frac{2[p_t - p_r]}{r} - \frac{M_g(r)[\rho(r) + p_r]\sqrt{ab}}{r} - \frac{dp_r}{r} + \frac{\xi[p_t - p_r]}{r} = 0, \quad (34)$$




Figure 2. Schematic plots of (a) the anisotropy and anisotropic force; (b) the gradient of the energy-momentum components; (c) the speed of sound; and (d) the difference between the tangential and radial speed of sounds. All of the figures are depicted vs. the radial coordinate r .

with $M_g(r)$ the gravitational mass at radius r , given by

$$M_g(r) = 4\pi \int_0^r \left(T_t^t - T_r^r - T_\theta^\theta - T_\phi^\phi \right) r^2 \sqrt{\frac{a}{b}} dr = \frac{ra'}{2a\sqrt{b}}, \quad (35)$$

Inserting Eq. (35) into (34), we find

$$\frac{2(p_t - p_r)}{r} - \frac{dp_r}{dr} - \frac{a'[\rho(r) + p_r]}{2\sqrt{a}} + \frac{\xi[p_t - p_r]}{r} = F_g + F_a + F_h + F_h + F_s = 0. \quad (36)$$

Here, $F_g = -\frac{a'[\rho(r) + p_r]}{2\sqrt{a}}$, $F_a = \frac{2(p_t - p_r)}{r}$, $F_h = -\frac{dp_r}{dr}$ and $F_s = \frac{\xi[p_t - p_r]}{r}$ means the gravitational, anisotropic, hydrostatic and scalar field forces, respectively. The TOV equation for Eq. (25) is found in Fig. 5 (a), which shows the four different forces. It can be understood that F_g, F_a, F_h are positive and F_s is dominant, so that the equilibrium of a star can be maintained.




Figure 3. Plot of the (a) weak, (b) null, (c) strong and (d) dominant energy conditions given by Eq. (25). All these plots are depicted vs. the radial coordinate r .




Figure 4. Plot of (a) EoS, (b) mass and compactness, and (c) the red shift. All of the figures are depicted vs. the radial coordinate r .

B. Adiabatic index

For examine the stability of a star, we investigate the equilibrium state for a spherically symmetric spacetime. We analyze the adiabatic index Γ , given by [85–87]

$$\Gamma = \left(\frac{\rho + p}{p} \right) \left(\frac{dp}{d\rho} \right), \quad (37)$$

where ρ and p denote the density and pressure of isotropic system. The equilibrium state is realized for $\Gamma > \frac{4}{3}$ [88]. If $\Gamma = \frac{4}{3}$, the isotropic spacetime is in equilibrium.

For an anisotropic system, the stability of a relativistic sphere is realized for $\Gamma > \gamma$ [87]. Here,

$$\gamma = \frac{4}{3} - \left\{ \frac{4(p_r - p_t)}{3|p'_r|} \right\}_{max}. \quad (38)$$

From Eqs. (37) and (38), we get the components of the adiabatic index that are listed in Appendix (C).

In Fig. 5 (b), we depict γ , Γ_r and Γ_t . It is seen that Γ_r and Γ_t are larger than γ within a star, and thus the stability conditions are met.

C. Stability in the static state

Another test to ensure the stability of the model under consideration is the one of the static state given by Harrison, Zeldovich and Novikov [78, 89, 90]. They showed that for a stable star the derivative of the mass in terms of the central density $\rho(\rightarrow 0)$, must give a positive increasing value, i.e., $\frac{\partial M}{\partial \rho_0} > 0$. Using this condition in our model we obtain the form of the central density as

$$\rho_0(r \rightarrow 0) = \frac{48(a_1 c_1 - a_0^2 - a_1 a_0^2)}{a_1 a_0^5} \implies c_1 = \frac{a_0^2(\rho_0 a_1 a_0^3 + 48[1 + a_1])}{48 a_1}. \quad (39)$$

Using Eq. (39) in Eq. (26) we find the mass of the central density, represented as

$$M(\rho_0) \approx \frac{663552 a_1^3}{a_0^7 (\rho_0 a_1 a_0^3 + 48 + 48 a_1)^3 l^5} + \frac{1}{a_0^6 (\rho_0 a_1 a_0^3 + 48 + 48 a_1)^3 l^3} \left[\frac{27648(\rho_0 a_1 a_0^3 + 48 + 48 a_1)}{a_1} - 9216 \left(\frac{288}{a_1} \right. \right. \\ \left. \left. - \frac{a_0 (1344 a_0^7 a_1^3 - 7 a_0^7 (\rho_0 a_1 a_0^3 + 48 + 48 a_1) a_1^3 + 192 a_0^7 a_1^4) - 288 a_1^4 a_0^8}{a_0^8 a_1^4} \right) \right] a_1^3. \quad (40)$$

The pattern of the derivative of the mass in terms of the central density is given by the following asymptotic form

$$\frac{\partial M(\rho_0)}{\partial \rho_0} \approx \frac{1990656 a_1^4}{a_0^4 (\rho_0 a_1 a_0^3 + 48 + 48 a_1)^4 l^5} + \frac{11520 a_1^3}{a_0^3 (\rho_0 a_1 a_0^3 + 48 + 48 a_1)^3 l^3} - \frac{a_1^4}{a_0^3 (\rho_0 a_1 a_0^3 + 48 + 48 a_1)^4 l^3} \left[\frac{82944}{a_1} \left(48 \right. \right. \\ \left. \left. - \frac{a_0 (1344 a_0^7 a_1^3 - 7 a_0^7 (\rho_0 a_1 a_0^3 + 48 + 48 a_1) a_1^3 + 192 a_0^7 a_1^4) - 288 a_1^4 a_0^8}{a_0^8 a_1^4} \right) \right]. \quad (41)$$

Equations (40) and (41) are depicted in Figs. 5 (c) and Fig. 5 (d) which ensure the stability of the model.




Figure 5. Plot of (a) the different forces; (b) the radial and tangential adiabatic indices; (c) and (d) the mass and its derivative in terms of the central density. All of the figures are depicted vs. the radial coordinate r .

Table I. Model parameters

Pulsar	Mass (M_\odot)	Radius (km)	a_0	a_1	c_1
Her X-1	0.85 ± 0.15	8.1 ± 0.41	-0.004802894320	0.3043478261	-0.04483731442
Cen X-3	1.49 ± 0.08	9.178 ± 0.13	-0.005762649954	0.001461108724	-0.03322916139
RX J 1856 - 37	0.9 ± 0.2	$\simeq 6$	-0.01507407407	-0.085333333333	-0.07649056652
4U1608 - 52	1.74 ± 0.14	9.52 ± 0.15	-0.0061541792	-0.1509410548	-0.02864346182
EXO 1785 - 248	1.3 ± 0.2	8.849 ± 0.4	-0.00561175409	0.039896205	-0.0342249748
4U1820 - 30	1.58	9.1	-0.006206183624	-0.02786813187	-0.03430457972

In addition to PSR J 1614-2230, the same procedure can be performed for other pulsars. The results for other stellar objects are shown in Table I and II

VIII. DISCUSSIONS AND CONCLUSIONS

A TEGR theory is known to be identical with general relativity [91–93]. However, conformal TEGR is not equivalent to the conformal GR theory. In this research, we studied conformal TEGR field equations. In the present paper, we have explored the role of the conformal field on the anisotropic compact objects. Applying the field equations of

Table II. Values of physical quantities

Pulsar	$\rho _0$	$\rho _l$	$\frac{dp_r}{d\rho} _0$	$\frac{dp_r}{d\rho} _l$	$\frac{dp_t}{d\rho} _0$	$\frac{dp_t}{d\rho} _l$	$(\rho - p_r - 2p_t) _0$	$(\rho - p_r - 2p_t) _l$	$z _l$
Her X-1	$\approx 8.44 \times 10^{11}$	≈ 0.0012	≈ 0.3	≈ -2.1	≈ 0.3	≈ 1.2	≈ 2508.8	$\approx 2.3 \times 10^{-14}$	-1
Cen X-3	$\approx 4.23 \times 10^{11}$	≈ -0.04	≈ 0.3	≈ 0.7	≈ 0.3	≈ 0.16	$\approx 1.41 \times 10^8$	$\approx 1.49 \times 10^{-13}$	-1
RX J 1856 - 37	4.57×10^9	-0.052	.33	3.76	.33	55.7	1.5×10^6	-0.25×10^{-4}	-1
4U1608 - 52	1.55×10^{11}	-0.02051145188	0.33	-2	0.33	0.36	5.2×10^7	-2.34×10^{-13}	-1
EXO 1785 - 248	3×10^{11}	0.0105	.3	1.57	.3	.1	1.48×10^{11}	.041	-1
4U1820 - 30	3×10^{11}	-0.007	0.3	-0.3	0.3	1.36	1.48×10^{11}	.06	-1

conformal TEGR theory to a non-diagonal vierbein field. For exterior solution, the scalar torsion is vanishing when the two unknown functions approach Minkowski spacetime which ensures that the vierbein field we begin is a physical one. The output differential equations consist of five non-linear ones in eight unknowns, $a(r)$, $b(r)$, $\rho(r)$, $p_r(r)$, $p_t(r)$, $V(r)$, $\xi(r)$ and $\Delta(r)$.

As a result, we have obtained an analytic solution for realistic compact star configurations. It has been demonstrated that the model proposed in this work is regular at the center as well as at the surface according to the stellar structure. In particular, the properties of the density as well as the radial and tangential pressures are regular for the case of a pulsar 4U1608-52 [94].

Moreover, we have shown that the anisotropy and anisotropic forces have negative values which can be considered as attractive forces. This result originates from the fact that the radial pressure is stronger than the tangential pressure, that is, $p_r > p_t$ [95]. The present model is stable owing to the balance of the hydrostatic, gravitational, anisotropic and scalar forces. Also with the procedures of the adiabatic index for a static star, the model is shown to be stable. Furthermore, we have evaluated the sound speed and confirmed that it is consistent with the observations. According to this result, the metric potentials (21) and the anisotropy expression (24) are considered to be realistic for astrophysical compact objects. In addition, we have analyzed the present model by using the observational data of pulsars (Tables I and II) and shown that the present theoretical model fit can fit the observational date well.

The consequences obtained in this work are summarized as follows. We have employed a non-diagonal vierbein field yielding the vanishing torsion scalar as the metric potentials are close to unity. Such a behavior is important for the vierbein field as shown in Refs. [96–102]. It has also been demonstrated that the theoretical predictions of the present model are compatible with the real observational data of pulsars. It is expected that the present study will be able to be tested for a wide range of the metric and anisotropy.

ACKNOWLEDGEMENT

The work of KB was supported in part by the JSPS KAKENHI Grant Number JP21K03547.

Appendix A: The gradients of the components of Eqs. (25)

The explicite form of the gradients of the components of the energy-momentum components given by Eq. (25) have the following form:

$$\begin{aligned}
\rho' = 256 \left\{ & \left[20 a_0^6 r^4 - 24 a_0^4 - 24 r^2 a_0^5 - 64 a_1 a_0^4 - 6 c_1^2 S^2 - 64 a_1 c_1 S a_0^2 - 52 r^4 a_0^2 c_1^2 S - 6 r^2 a_0 c_1^2 S^2 - 24 r^2 a_0^3 c_1 S \right. \right. \\
& - 24 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 + 128 r^2 a_0^3 c_1 a_1 + 20 a_0^4 r^4 c_1 S - 104 r^2 a_1 a_0^3 c_1 S + 64 r^2 a_0 c_1^2 a_1 S - 26 r^2 a_1 a_0 c_1^2 S^2 \\
& + 72 c_1 a_1^2 a_0^2 + 24 S c_1^2 r^4 a_0^2 a_1 + 32 S c_1^2 a_0 r^2 a_1^2 - 40 S c_1 r^4 a_0^4 a_1 - 20 S c_1 a_0^3 r^2 a_1^2 - 10 S^2 c_1^2 r^4 a_0^2 a_1 - 5 S^2 c_1^2 a_0 r^2 a_1^2 \\
& - 60 c_1^2 a_1^2 - 20 a_1^2 a_0^4 + 20 r^6 a_0^7 + 48 c_1 r^4 a_0^4 a_1 + 64 c_1 a_0^3 r^2 a_1^2 - 60 c_1^2 a_0 r^2 a_1^2 - 20 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 \\
& - 32 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 + 60 c_1^2 r^6 a_0^3 - 64 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 - 20 a_0^5 r^2 a_1^2 + 36 S c_1^2 a_1^2 - 5 S^2 c_1^2 a_1^2 \\
& \left. \left. + 96 a_0^2 c_1 a_1 - 104 r^2 a_1 a_0^5 - 16 a_1 c_1^2 S^2 - 24 c_1 S a_0^2 - 104 r^4 a_0^4 c_1 + 60 a_0^2 r^4 c_1^2 + 48 c_1^2 a_1 S - 48 r^2 a_0^3 c_1 \right] a_0^6 r \right\} \\
& \left\{ S \left[c_1 S r^2 a_0 - a_1 c_1 S - 2 a_1 a_0^2 + 2 a_0^3 r^2 \right] \left(S^5 c_1^5 r^2 a_0 - S^5 c_1^5 a_1 - 10 S^4 c_1^4 a_1 a_0^2 + 10 S^4 c_1^4 r^2 a_0^3 + 40 S^3 c_1^3 r^2 a_0^5 \right. \right. \\
& \left. \left. \right. \right. \right.
\end{aligned}$$

$$\begin{aligned}
& -40 S^3 c_1^3 a_1 a_0^4 + 80 S^2 c_1^2 r^2 a_0^7 - 80 S^2 c_1^2 a_1 a_0^6 + 80 S c_1 r^2 a_0^9 - 80 S c_1 a_0^8 a_1 - 32 a_1 a_0^{10} + 32 r^2 a_0^{11} \Bigg) \Bigg\}^{-1}, \\
p'_r &= -256 \left\{ \left[12 a_0^5 r^2 a_1^2 - 4 r^4 a_0^2 c_1^2 S + 2 r^2 a_0 c_1^2 S^2 + 8 r^2 a_0^3 c_1 S + 8 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 + 20 a_0^4 r^4 c_1 S \right. \right. \\
&+ 48 c_1 r^4 a_0^4 a_1 - 32 c_1 a_0^3 r^2 a_1^2 + 20 c_1^2 a_0 r^2 a_1^2 + 12 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 - 16 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 \\
&- 40 r^2 a_1 a_0^3 c_1 S - 10 r^2 a_1 a_0 c_1^2 S^2 + 24 S c_1^2 r^4 a_0^2 a_1 - 16 S c_1^2 a_0 r^2 a_1^2 - 40 S c_1 r^4 a_0^4 a_1 \\
&+ 12 S c_1 a_0^3 r^2 a_1^2 - 10 S^2 c_1^2 r^4 a_0^2 a_1 + 3 S^2 c_1^2 a_0 r^2 a_1^2 + 8 r^2 a_0^5 + 20 a_0^6 r^4 + 2 c_1^2 S^2 + 20 c_1^2 a_1^2 + 12 a_1^2 a_0^4 + 20 r^6 a_0^7 \\
&+ 3 S^2 c_1^2 a_1^2 - 40 r^4 a_0^6 a_1 - 12 S c_1^2 a_1^2 - 32 a_0^2 c_1 a_1 - 32 c_1 r^6 a_0^5 - 20 c_1^2 r^6 a_0^3 - 24 c_1 a_1^2 a_0^2 - 40 r^2 a_1 a_0^5 + 8 c_1 S a_0^2 \\
&\left. \left. - 8 r^4 a_0^4 c_1 - 20 a_0^2 r^4 c_1^2 - 16 c_1^2 a_1 S + 16 r^2 a_0^3 c_1 + 8 a_0^4 \right] a_0^6 r \right\} \left\{ (1 + r^2 a_0) (c_1 S r^2 a_0 - a_1 c_1 S - 2 a_1 a_0^2 + 2 a_0^3 r^2) \right. \\
&\left. \left[S^5 c_1^5 r^2 a_0 - S^5 c_1^5 a_1 - 10 S^4 c_1^4 a_1 a_0^2 + 10 S^4 c_1^4 r^2 a_0^3 + 40 S^3 c_1^3 r^2 a_0^5 - 40 S^3 c_1^3 a_1 a_0^4 + 80 S^2 c_1^2 r^2 a_0^7 - 80 S^2 c_1^2 a_1 a_0^6 \right. \right. \\
&\left. \left. + 80 S c_1 r^2 a_0^9 - 80 S c_1 a_0^8 a_1 - 32 a_1 a_0^{10} + 32 r^2 a_0^{11} \right] \right\}^{-1} \\
p'_t &= -256 \left\{ \left[32 r^2 a_0 c_1^2 a_1 S - 4 a_0^5 r^2 a_1^2 - 28 r^4 a_0^2 c_1^2 S - 2 r^2 a_0 c_1^2 S^2 - 8 r^2 a_0^3 c_1 S - 8 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 \right. \right. \\
&+ 20 a_0^4 r^4 c_1 S + 48 c_1 r^4 a_0^4 a_1 + 16 c_1 a_0^3 r^2 a_1^2 - 20 c_1^2 a_0 r^2 a_1^2 - 4 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 - 24 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 \\
&- 32 a_1 c_1 S a_0^2 - 72 r^2 a_1 a_0^3 c_1 S - 18 r^2 a_1 a_0 c_1^2 S^2 + 24 S c_1^2 r^4 a_0^2 a_1 + 8 S c_1^2 a_0 r^2 a_1^2 - 40 S c_1 r^4 a_0^4 a_1 - 4 S c_1 a_0^3 r^2 a_1^2 \\
&- 10 S^2 c_1^2 r^4 a_0^2 a_1 - S^2 c_1^2 a_0 r^2 a_1^2 - 8 r^2 a_0^5 + 20 a_0^6 r^4 - 2 c_1^2 S^2 - 20 c_1^2 a_1^2 - 4 a_1^2 a_0^4 + 20 r^6 a_0^7 - S^2 c_1^2 a_1^2 - 40 r^4 a_0^6 a_1 \\
&+ 12 S c_1^2 a_1^2 + 32 a_0^2 c_1 a_1 - 48 c_1 r^6 a_0^5 + 20 c_1^2 r^6 a_0^3 + 24 c_1 a_1^2 a_0^2 - 72 r^2 a_1 a_0^5 - 8 c_1 S a_0^2 - 56 r^4 a_0^4 c_1 + 20 a_0^2 r^4 c_1^2 \\
&\left. \left. + 16 c_1^2 a_1 S - 16 r^2 a_0^3 c_1 - 8 a_0^4 + 64 r^2 a_1 a_0^3 c_1 - 8 c_1^2 S^2 a_1 - 32 a_1 a_0^4 \right] a_0^6 r \right\} \left\{ (1 + r^2 a_0) \left[c_1 S r^2 a_0 - a_1 c_1 S - 2 a_1 a_0^2 \right. \right. \\
&\left. \left. + 2 a_0^3 r^2 \right] \left(S^5 c_1^5 r^2 a_0 - S^5 c_1^5 a_1 - 10 S^4 c_1^4 a_1 a_0^2 + 10 S^4 c_1^4 r^2 a_0^3 + 40 S^3 c_1^3 r^2 a_0^5 - 40 S^3 c_1^3 a_1 a_0^4 + 80 S^2 c_1^2 r^2 a_0^7 \right. \right. \\
&\left. \left. - 80 S^2 c_1^2 a_1 a_0^6 + 80 S c_1 r^2 a_0^9 - 80 S c_1 a_0^8 a_1 - 32 a_1 a_0^{10} + 32 r^2 a_0^{11} \right] \right\}^{-1}, \tag{A1}
\end{aligned}$$

where $\rho' = \frac{d\rho}{dr}$, $p'_r = \frac{dp_r}{dr}$ and $p'_t = \frac{dp_t}{dr}$. Eqs. (A1) show that the gradients of density, radial and tangential pressures are negative as we will show when we plot them.

Appendix B: The components of the speed of sound

Using Eq. (25) we evaluate the value of the components of the sound and get:

$$\begin{aligned}
v_r^2 &= - \left\{ 20 c_1^2 a_1^2 - 24 c_1 a_1^2 a_0^2 + 12 a_1^2 a_0^4 + 20 r^6 a_0^7 + 8 c_1 S a_0^2 + 16 r^2 a_0^3 c_1 - 32 a_0^2 c_1 a_1 - 8 r^4 a_0^4 c_1 - 16 c_1^2 a_1 S \right. \\
&- 20 a_0^2 r^4 c_1^2 - 40 r^2 a_1 a_0^5 + 8 r^2 a_0^5 + 20 a_0^6 r^4 + 2 c_1^2 S^2 - 12 S c_1^2 a_1^2 + 20 c_1^2 a_0 r^2 a_1^2 - 32 c_1 a_0^3 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 \\
&+ 12 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 - 16 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 - 32 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 + 12 a_0^5 r^2 a_1^2 - 20 c_1^2 r^6 a_0^3 \\
&+ 3 S^2 c_1^2 a_1^2 + 3 S^2 c_1^2 a_0 r^2 a_1^2 + 24 S c_1^2 r^4 a_0^2 a_1 - 16 S c_1^2 a_0 r^2 a_1^2 + 12 S c_1 a_0^3 r^2 a_1^2 - 40 S c_1 r^4 a_0^4 a_1 - 10 S^2 c_1^2 r^4 a_0^2 a_1 \\
&+ 8 a_0^4 - 40 r^2 a_1 a_0^3 c_1 S - 10 r^2 a_1 a_0 c_1^2 S^2 + 20 a_0^4 r^4 c_1 S - 4 r^4 a_0^2 c_1^2 S + 2 r^2 a_0 c_1^2 S^2 + 8 r^2 a_0^3 c_1 S + 8 r^2 a_0 c_1^2 S \\
&\left. + 5 a_0^2 r^4 c_1^2 S^2 \right\} \times \left\{ 72 c_1 a_1^2 a_0^2 - 60 c_1^2 a_1^2 - 20 a_1^2 a_0^4 + 20 r^6 a_0^7 - 24 c_1 S a_0^2 - 48 r^2 a_0^3 c_1 + 96 a_0^2 c_1 a_1 - 104 r^4 a_0^4 c_1 \right. \\
&\left. + 4 c_1^2 [12 a_1 S + 15 a_0^2 r^4 - 4 a_1 S^2] - 104 r^2 a_1 a_0^5 - 24 r^2 a_0^5 + 20 a_0^6 r^4 - 64 a_1 a_0^4 - 6 c_1^2 S^2 + 36 S c_1^2 a_1^2 - 60 c_1^2 a_0 r^2 a_1^2 \right. \\
&\left. + 64 c_1 a_0^3 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 - 20 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 - 32 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 - 64 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 \right\}
\end{aligned}$$

$$\begin{aligned}
& -20 a_0^5 r^2 a_1^2 + 60 c_1^2 r^6 a_0^3 - 5 S^2 c_1^2 a_1^2 - 5 S^2 c_1^2 a_0 r^2 a_1^2 + 24 S c_1^2 r^4 a_0^2 a_1 + 32 S c_1^2 a_0 r^2 a_1^2 - 20 S c_1 a_0^3 r^2 a_1^2 \\
& - 40 S c_1 r^4 a_0^4 a_1 - 10 S^2 c_1^2 r^4 a_0^2 a_1 - 24 a_0^4 - 104 r^2 a_1 a_0^3 c_1 S + 64 r^2 a_0 c_1^2 a_1 S - 26 r^2 a_1 a_0 c_1^2 S^2 + 20 a_0^4 r^4 c_1 c_1 S \\
& - 64 a_1 c_1 S a_0^2 - 52 r^4 a_0^2 c_1^2 S - 6 r^2 a_0 c_1^2 S^2 - 24 r^2 a_0^3 c_1 S - 24 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 + 128 r^2 a_0^3 c_1 a_1 \Big\}^{-1}, \\
v_r^2 = & \left\{ 24 c_1 a_1^2 a_0^2 - 20 c_1^2 a_1^2 - 4 a_1^2 a_0^4 + 20 r^6 a_0^7 - 8 c_1 S a_0^2 - 16 r^2 a_0^3 c_1 + 32 a_0^2 c_1 a_1 - 56 r^4 a_0^4 c_1 + 16 c_1^2 a_1 S \right. \\
& + 20 a_0^2 r^4 c_1^2 - 8 a_1 c_1^2 S^2 - 72 r^2 a_1 a_0^5 - 8 r^2 a_0^5 + 20 a_0^6 r^4 - 32 a_1 a_0^4 - 2 c_1^2 S^2 + 12 S c_1^2 a_1^2 - 20 c_1^2 a_0 r^2 a_1^2 \\
& + 16 c_1 a_0^3 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 - 4 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 - 24 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 - 48 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 \\
& - 4 a_0^5 r^2 a_1^2 + 20 c_1^2 r^6 a_0^3 - S^2 c_1^2 a_1^2 - S^2 c_1^2 a_0 r^2 a_1^2 + 24 S c_1^2 r^4 a_0^2 a_1 + 8 S c_1^2 a_0 r^2 a_1^2 - 4 S c_1 a_0^3 r^2 a_1^2 - 40 S c_1 r^4 a_0^4 a_1 \\
& - 10 S^2 c_1^2 r^4 a_0^2 a_1 - 8 a_0^4 - 72 r^2 a_1 a_0^3 c_1 S + 32 r^2 a_0 c_1^2 a_1 S - 18 r^2 a_1 a_0 c_1^2 S^2 + 20 a_0^4 r^4 c_1 S - 32 a_1 c_1 S a_0^2 - 28 r^4 a_0^2 c_1^2 S \\
& \left. - 2 r^2 a_0 c_1^2 S^2 - 8 r^2 a_0^3 c_1 S - 8 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 + 64 r^2 a_0^3 c_1 a_1 \right\} \left\{ 72 c_1 a_1^2 a_0^2 - 60 c_1^2 a_1^2 - 20 a_1^2 a_0^4 + 20 r^6 a_0^7 \right. \\
& - 24 c_1 S a_0^2 - 48 r^2 a_0^3 c_1 + 96 a_0^2 c_1 a_1 - 104 r^4 a_0^4 c_1 + 48 c_1^2 a_1 S + 60 a_0^2 r^4 c_1^2 - 16 a_1 c_1^2 S^2 - 104 r^2 a_1 a_0^5 - 24 r^2 a_0^5 \\
& + 20 a_0^6 r^4 - 64 a_1 a_0^4 - 6 c_1^2 S^2 + 36 S c_1^2 a_1^2 - 60 c_1^2 a_0 r^2 a_1^2 + 64 c_1 a_0^3 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 - 20 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 \\
& - 32 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 - 64 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 - 20 a_0^5 r^2 a_1^2 + 60 c_1^2 r^6 a_0^3 - 5 S^2 c_1^2 a_1^2 - 5 S^2 c_1^2 a_0 r^2 a_1^2 \\
& + 24 S c_1^2 r^4 a_0^2 a_1 + 32 S c_1^2 a_0 r^2 a_1^2 - 20 S c_1 a_0^3 r^2 a_1^2 - 40 S c_1 r^4 a_0^4 a_1 - 10 S^2 c_1^2 r^4 a_0^2 a_1 - 24 a_0^4 - 104 r^2 a_1 a_0^3 c_1 S \\
& + 64 r^2 a_0 c_1^2 a_1 S - 26 r^2 a_1 a_0 c_1^2 S^2 + 20 a_0^4 r^4 c_1 S - 64 a_1 c_1 S a_0^2 - 52 r^4 a_0^2 c_1^2 S - 6 r^2 a_0 c_1^2 S^2 - 24 r^2 a_0^3 c_1 S - 24 r^2 a_0 c_1^2 S \\
& \left. + 5 a_0^2 r^4 c_1^2 S^2 + 128 r^2 a_0^3 c_1 a_1 \right\}. \tag{B1}
\end{aligned}$$

Appendix C: The components of the adiabatic index

The explicite form of the components of the adiabatic index given by Eq. (25) take the following form:

$$\begin{aligned}
\Gamma = & -8 \left\{ 16 c_1 a_0^3 r^2 a_1^2 - 4 S c_1 a_1^2 a_0^2 + 8 S c_1^2 r^6 a_0^3 - 8 a_1 c_1 S a_0^2 - 4 c_1 S a_0^2 - 8 r^2 a_0^3 c_1 + 16 a_0^2 c_1 a_1 + 4 r^4 a_0^4 c_1 \right. \\
& + 8 c_1^2 a_1 S + 10 a_0^2 r^4 c_1^2 + 12 c_1 a_1^2 a_0^2 + 6 S c_1^2 a_1^2 + 10 c_1^2 r^6 a_0^3 + 16 c_1 r^6 a_0^5 + 8 r^4 a_0^6 a_1 - 4 a_0^5 r^2 a_1^2 - S^2 c_1^2 a_1^2 \\
& - 2 a_1 c_1^2 S^2 + 4 r^2 a_0^5 + 8 a_0^6 r^4 - c_1^2 S^2 - 10 c_1^2 a_1^2 - 4 a_1^2 a_0^4 - 8 a_1 a_0^4 - 4 a_0^4 + 8 S c_1 r^4 a_0^4 a_1 - 4 S c_1 a_0^3 r^2 a_1^2 \\
& + 2 S^2 c_1^2 r^4 a_0^2 a_1 - 12 S c_1^2 r^4 a_0^2 a_1 + 8 S c_1^2 a_0 r^2 a_1^2 - S^2 c_1^2 a_0 r^2 a_1^2 + 8 a_0^4 r^4 c_1 S + 2 r^4 a_0^2 c_1^2 S + r^2 a_0 c_1^2 S^2 + 4 r^2 a_0^3 c_1 S \\
& \left. - 4 r^2 a_0 c_1^2 S + 2 a_0^2 r^4 c_1^2 S^2 - 10 c_1^2 a_0 r^2 a_1^2 - 24 c_1 r^4 a_0^4 a_1 \right\} \left\{ 3 \left(-32 c_1 a_0^3 r^2 a_1^2 + 12 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 \right. \right. \\
& - 16 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 + 8 c_1 S a_0^2 + 16 r^2 a_0^3 c_1 - 32 a_0^2 c_1 a_1 - 8 r^4 a_0^4 c_1 - 16 c_1^2 a_1 S - 20 a_0^2 r^4 c_1^2 - 40 r^2 a_1 a_0^5 \\
& - 24 c_1 a_1^2 a_0^2 - 12 S c_1^2 a_1^2 - 20 c_1^2 r^6 a_0^3 - 32 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 + 12 a_0^5 r^2 a_1^2 + 3 S^2 c_1^2 a_1^2 + 8 r^2 a_0^5 + 20 a_0^6 r^4 \\
& + 2 c_1^2 S^2 + 20 c_1^2 a_1^2 + 12 a_1^2 a_0^4 + 20 r^6 a_0^7 + 8 a_0^4 - 40 r^2 a_1 a_0^3 c_1 S - 10 r^2 a_1 a_0 c_1^2 S^2 - 40 S c_1 r^4 a_0^4 a_1 + 12 S c_1 a_0^3 r^2 a_1^2 \\
& - 10 S^2 c_1^2 r^4 a_0^2 a_1 + 24 S c_1^2 r^4 a_0^2 a_1 - 16 S c_1^2 a_0 r^2 a_1^2 + 3 S^2 c_1^2 a_0 r^2 a_1^2 + 20 a_0^4 r^4 c_1 S - 4 r^4 a_0^2 c_1^2 S + 2 r^2 a_0 c_1^2 S^2 \\
& \left. \left. + 8 r^2 a_0^3 c_1 S + 8 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 + 20 c_1^2 a_0 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 \right) \right\}. \tag{C1}
\end{aligned}$$

From Eq. (37), we obtain the adiabatic index of solution (25) in the form

$$\begin{aligned}
\Gamma_r = & 8 \left\{ \left[-2 c_1 r^4 a_0^2 - 2 c_1 r^2 a_0 + 2 r^2 a_1 a_0^3 - 2 c_1 a_1 + 2 a_0^3 r^2 + 2 a_1 a_0^2 + 2 a_0^2 - 2 c_1 a_0 r^2 a_1 + a_1 c_1 S + c_1 S r^2 a_0 + c_1 S \right. \right. \\
& + r^2 a_1 a_0 c_1 S \left. \right] \times \left(-32 c_1 a_0^3 r^2 a_1^2 + 12 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 - 16 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 + 8 c_1 S a_0^2 + 16 r^2 a_0^3 c_1 \right. \\
& - 32 a_0^2 c_1 a_1 - 8 r^4 a_0^4 c_1 - 16 C1^2 a_1 S - 20 a_0^2 r^4 c_1^2 - 40 r^2 a_1 a_0^5 - 24 c_1 a_1^2 a_0^2 - 12 S c_1^2 a_1^2 - 20 c_1^2 r^6 a_0^3 - 32 c_1 r^6 a_0^5 \\
& - 40 r^4 a_0^6 a_1 + 12 a_0^5 r^2 a_1^2 + 3 S^2 c_1^2 a_1^2 + 8 r^2 a_0^5 + 20 a_0^6 r^4 + 2 c_1^2 (S)^2 + 20 c_1^2 a_1^2 + 12 a_1^2 a_0^4 + 20 r^6 a_0^7 + 8 a_0^4 \\
& - 40 r^2 a_1 a_0^3 c_1 S - 10 r^2 a_1 a_0 c_1^2 S^2 - 40 S c_1 r^4 a_0^4 a_1 + 12 S c_1 a_0^3 r^2 a_1^2 - 10 S^2 c_1^2 r^4 a_0^2 a_1 + 24 S c_1^2 r^4 a_0^2 a_1 - 16 S c_1^2 a_0 r^2 a_1^2 \\
& \left. + 3 S^2 c_1^2 a_0 r^2 a_1^2 + 20 a_0^4 r^4 c_1 S - 4 r^4 a_0^2 c_1^2 \ln(1+r^2 a_0) + 2 r^2 a_0 c_1^2 S^2 + 8 r^2 a_0^3 c_1 S + 8 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 \right)
\end{aligned}$$

$$\begin{aligned}
& +20 c_1^2 a_0 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 \Big) \Big\} \times \left\{ \left[2 c_1 S r^2 a_0 + 5 a_0^2 r^4 c_1 S - 2 c_1 S - 3 r^2 a_1 a_0 c_1 S - 2 a_1 c_1 S + 4 c_1 a_0 r^2 a_1 + 4 c_1 a_1 \right. \right. \\
& +4 c_1 r^2 a_0 + 4 c_1 r^4 a_0^2 + 4 a_0^3 r^2 - 6 r^2 a_1 a_0^3 - 4 a_0^2 + 10 a_0^4 r^4 - 4 a_1 a_0^2 \Big] \times \left(64 c_1 a_0^3 r^2 a_1^2 - 20 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 \right. \\
& -32 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 - 64 a_1 c_1 S a_0^2 + 128 r^2 a_0^3 c_1 a_1 - 24 c_1 S a_0^2 - 48 r^2 a_0^3 c_1 + 96 a_0^2 c_1 a_1 - 104 r^4 a_0^4 c_1 \\
& +48 c_1^2 a_1 S + 60 a_0^2 r^4 c_1^2 - 104 r^2 a_1 a_0^5 + 72 c_1 a_1^2 a_0^2 + 36 S c_1^2 a_1^2 + 60 c_1^2 r^6 a_0^3 - 64 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 - 20 a_0^5 r^2 a_1^2 \\
& -5 S^2 c_1^2 a_1^2 - 16 a_1 c_1^2 (S)^2 - 24 r^2 a_0^5 + 20 a_0^6 r^4 - 6 c_1^2 S^2 - 60 c_1^2 a_1^2 - 20 a_1^2 a_0^4 + 20 r^6 a_0^7 - 64 a_1 a_0^4 - 24 a_0^4 \\
& -104 r^2 a_1 a_0^3 c_1 S - 26 r^2 a_1 a_0 c_1^2 S^2 - 40 S c_1 r^4 a_0^4 a_1 - 20 S c_1 a_0^3 r^2 a_1^2 - 10 S^2 c_1^2 r^4 a_0^2 a_1 + 24 S c_1^2 r^4 a_0^2 a_1 \\
& +32 S c_1^2 a_0 r^2 a_1^2 - 5 S^2 c_1^2 a_0 r^2 a_1^2 + 64 r^2 a_0 c_1^2 a_1 S + 20 a_0^4 r^4 c_1 S - 52 r^4 a_0^2 c_1^2 S - 6 r^2 a_0 c_1^2 S^2 - 24 r^2 a_0^3 c_1 S \\
& \left. \left. -24 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 - 60 c_1^2 a_0 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 \right) \right\}^{-1}, \\
\Gamma_t = & 2 \left\{ \left(6 r^2 a_1 a_0^3 - 8 c_1 r^4 a_0^2 - 8 c_1 r^2 a_0 - 8 c_1 a_1 + 16 a_0^3 r^2 + 8 a_1 a_0^2 + 10 a_0^4 r^4 + 8 a_0^2 - 8 c_1 a_0 r^2 a_1 + 8 c_1 S r^2 a_0 \right. \right. \\
& +5 a_0^2 r^4 c_1 S + 3 r^2 a_1 a_0 c_1 S + 4 c_1 S + 4 a_1 c_1 S \Big) \left[16 c_1 a_0^3 r^2 a_1^2 - 4 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 - 24 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 \right. \\
& -32 a_1 c_1 S a_0^2 + 64 r^2 a_0^3 c_1 a_1 - 8 c_1 S a_0^2 - 16 r^2 a_0^3 c_1 + 32 a_0^2 c_1 a_1 - 56 r^4 a_0^4 c_1 + 16 c_1^2 a_1 S + 20 a_0^2 r^4 c_1^2 - 72 r^2 a_1 a_0^5 \\
& +24 c_1 a_1^2 a_0^2 + 12 S c_1^2 a_1^2 + 20 c_1^2 r^6 a_0^3 - 48 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 - 4 a_0^5 r^2 a_1^2 - S^2 c_1^2 a_1^2 - 8 a_1 c_1^2 S^2 - 8 r^2 a_0^5 \\
& +20 a_0^6 r^4 - 2 c_1^2 S^2 - 20 c_1^2 a_1^2 - 4 a_1^2 a_0^4 + 20 r^6 a_0^7 - 32 a_1 a_0^4 - 8 a_0^4 - 72 r^2 a_1 a_0^3 c_1 S - 18 r^2 a_1 a_0 c_1^2 S^2 \\
& -40 S c_1 r^4 a_0^4 a_1 - 4 S c_1 a_0^3 r^2 a_1^2 - 10 S^2 c_1^2 r^4 a_0^2 a_1 + 24 S c_1^2 r^4 a_0^2 a_1 + 8 S c_1^2 a_0 r^2 a_1^2 - S^2 c_1^2 a_0 r^2 a_1^2 + 32 r^2 a_0 c_1^2 a_1 S \\
& \left. \left. +20 a_0^4 r^4 c_1 S - 28 r^4 a_0^2 c_1^2 S - 2 r^2 a_0 c_1^2 (S)^2 - 8 r^2 a_0^3 c_1 S - 8 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 - 20 c_1^2 a_0 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 \right] \right\} \\
& \times \left\{ \left[6 c_1 S r^2 a_0 + 5 a_0^2 r^4 c_1 S + 2 c_1 S + r^2 a_1 a_0 c_1 S + 2 a_1 c_1 S - 4 c_1 a_0 r^2 a_1 - 4 c_1 a_1 - 4 c_1 r^2 a_0 - 4 c_1 r^4 a_0^2 + 12 a_0^3 r^2 \right. \right. \\
& +2 r^2 a_1 a_0^3 + 4 a_0^2 + 10 a_0^4 r^4 + 4 a_1 a_0^2 \Big) \left(64 c_1 a_0^3 r^2 a_1^2 - 20 S c_1 a_1^2 a_0^2 + 5 S^2 c_1^2 r^6 a_0^3 - 32 S c_1^2 r^6 a_0^3 + 20 S c_1 r^6 a_0^5 \right. \\
& -64 a_1 c_1 S a_0^2 + 128 r^2 a_0^3 c_1 a_1 - 24 c_1 S a_0^2 - 48 r^2 a_0^3 c_1 + 96 a_0^2 c_1 a_1 - 104 r^4 a_0^4 c_1 + 48 c_1^2 a_1 S + 60 a_0^2 r^4 c_1^2 \\
& -104 r^2 a_1 a_0^5 + 72 c_1 a_1^2 a_0^2 + 36 S c_1^2 a_1^2 + 60 c_1^2 r^6 a_0^3 - 64 c_1 r^6 a_0^5 - 40 r^4 a_0^6 a_1 - 20 a_0^5 r^2 a_1^2 - 5 S^2 c_1^2 a_1^2 - 16 a_1 c_1^2 S^2 \\
& -24 r^2 a_0^5 + 20 a_0^6 r^4 - 6 c_1^2 S^2 - 60 c_1^2 a_1^2 - 20 a_1^2 a_0^4 + 20 r^6 a_0^7 - 64 a_1 a_0^4 - 24 a_0^4 - 104 r^2 a_1 a_0^3 c_1 S - 26 r^2 a_1 a_0 c_1^2 S^2 \\
& -40 S c_1 r^4 a_0^4 a_1 - 20 S c_1 a_0^3 r^2 a_1^2 - 10 S^2 c_1^2 r^4 a_0^2 a_1 + 24 S c_1^2 r^4 a_0^2 a_1 + 32 S c_1^2 a_0 r^2 a_1^2 - 5 S^2 c_1^2 a_0 r^2 a_1^2 \\
& \left. \left. +64 r^2 a_0 c_1^2 a_1 S + 20 a_0^4 r^4 c_1 S - 52 r^4 a_0^2 c_1^2 S - 6 r^2 a_0 c_1^2 S^2 - 24 r^2 a_0^3 c_1 S - 24 r^2 a_0 c_1^2 S + 5 a_0^2 r^4 c_1^2 S^2 \right. \right. \\
& \left. \left. -60 c_1^2 a_0 r^2 a_1^2 + 48 c_1 r^4 a_0^4 a_1 \right) \right\}^{-1}. \tag{C2}
\end{aligned}$$

-
- [1] G. 't Hooft, (2010), arXiv:1009.0669 [gr-qc].
- [2] P. D. Mannheim, Found. Phys. **42**, 388 (2012), arXiv:1101.2186 [hep-th].
- [3] G. 't Hooft, (2014), arXiv:1410.6675 [gr-qc].
- [4] P. D. Mannheim, Prog. Part. Nucl. Phys. **56**, 340 (2006), arXiv:astro-ph/0505266.
- [5] J. Maldacena, (2011), arXiv:1105.5632 [hep-th].
- [6] G. Anastasiou and R. Olea, Phys. Rev. D **94**, 086008 (2016), arXiv:1608.07826 [hep-th].
- [7] Y.-F. Cai, S. Capozziello, M. De Laurentis, and E. N. Saridakis, Rept. Prog. Phys. **79**, 106901 (2016), arXiv:1511.07586 [gr-qc].
- [8] J. M. Nester and H.-J. Yo, Chin. J. Phys. **37**, 113 (1999), arXiv:gr-qc/9809049.
- [9] M. Adak and O. Sert, Turk. J. Phys. **29**, 1 (2005), arXiv:gr-qc/0412007.
- [10] M. Adak, M. Kalay, and O. Sert, Int. J. Mod. Phys. D **15**, 619 (2006), arXiv:gr-qc/0505025.
- [11] J. Beltrán Jiménez, L. Heisenberg, and T. Koivisto, Phys. Rev. D **98**, 044048 (2018), arXiv:1710.03116 [gr-qc].
- [12] J. Beltrán Jiménez, L. Heisenberg, and T. S. Koivisto, JCAP **08**, 039 (2018), arXiv:1803.10185 [gr-qc].
- [13] L. Järv, M. Rünklä, M. Saal, and O. Vilson, Phys. Rev. D **97**, 124025 (2018), arXiv:1802.00492 [gr-qc].
- [14] R. Aldrovandi and J. G. Pereira, *Teleparallel Gravity: An Introduction* (Springer, 2013).
- [15] M. Krssak, R. J. van den Hoogen, J. G. Pereira, C. G. Böhmer, and A. A. Coley, Class. Quant. Grav. **36**, 183001 (2019), arXiv:1810.12932 [gr-qc].

- [16] A. Einstein, Sitz. Preuss. Akad. Wiss **217** (1928).
- [17] T. Sauer, *Historia Math.* **33**, 399 (2006), arXiv:physics/0405142.
- [18] K. Hayashi and T. Nakano, Progress of Theoretical Physics **38**, 491 (1967).
- [19] G. G. L. Nashed, *Annalen Phys.* **523**, 450 (2011), arXiv:1105.0328 [gr-qc].
- [20] G. G. L. Nashed and S. Capozziello, *Eur. Phys. J. C* **80**, 969 (2020), arXiv:2010.06355 [gr-qc].
- [21] G. G. L. Nashed, *Astrophys. Space Sci.* **330**, 173 (2010), arXiv:1503.01379 [gr-qc].
- [22] T. Shirafuji, G. G. L. Nashed, and Y. Kobayashi, *Prog. Theor. Phys.* **96**, 933 (1996), arXiv:gr-qc/9609060.
- [23] G. G. L. Nashed, *Eur. Phys. J. C* **49**, 851 (2007), arXiv:0706.0260 [gr-qc].
- [24] Y. Cho, Physical Review D **14**, 2521 (1976).
- [25] K. Hayashi, Physics Letters B **69**, 441 (1977).
- [26] F. W. Hehl, J. D. McCrea, E. W. Mielke, and Y. Ne'eman, *Phys. Rept.* **258**, 1 (1995), arXiv:gr-qc/9402012.
- [27] Y. N. Obukhov and J. G. Pereira, *Phys. Rev. D* **67**, 044008 (2003), arXiv:gr-qc/0212078.
- [28] R. Ferraro and F. Fiorini, *Phys. Rev. D* **75**, 084031 (2007), arXiv:gr-qc/0610067.
- [29] R. Ferraro and F. Fiorini, *Phys. Rev. D* **78**, 124019 (2008), arXiv:0812.1981 [gr-qc].
- [30] G. R. Bengochea and R. Ferraro, *Phys. Rev. D* **79**, 124019 (2009), arXiv:0812.1205 [astro-ph].
- [31] E. V. Linder, *Phys. Rev. D* **81**, 127301 (2010), [Erratum: Phys.Rev.D 82, 109902 (2010)], arXiv:1005.3039 [astro-ph.CO].
- [32] S.-H. Chen, J. B. Dent, S. Dutta, and E. N. Saridakis, *Phys. Rev. D* **83**, 023508 (2011), arXiv:1008.1250 [astro-ph.CO].
- [33] K. Hayashi and T. Shirafuji, Physical Review D **19**, 3524 (1979).
- [34] G. G. L. Nashed, *Int. J. Mod. Phys. D* **27**, 1850074 (2018).
- [35] G. G. L. Nashed and E. N. Saridakis, *Class. Quant. Grav.* **36**, 135005 (2019), arXiv:1811.03658 [gr-qc].
- [36] K. Hayashi and T. Shirafuji, *Phys. Rev. D* **24**, 3312 (1981).
- [37] G. Kofinas and E. N. Saridakis, *Phys. Rev. D* **90**, 084044 (2014), arXiv:1404.2249 [gr-qc].
- [38] G. Kofinas and E. N. Saridakis, *Phys. Rev. D* **90**, 084045 (2014), arXiv:1408.0107 [gr-qc].
- [39] S. Bahamonde and C. G. Böhmer, *Eur. Phys. J. C* **76**, 578 (2016), arXiv:1606.05557 [gr-qc].
- [40] T. Harko, F. S. N. Lobo, G. Otalora, and E. N. Saridakis, *JCAP* **12**, 021 (2014), arXiv:1405.0519 [gr-qc].
- [41] S. Bahamonde, C. G. Böhmer, and M. Krššák, *Phys. Lett. B* **775**, 37 (2017), arXiv:1706.04920 [gr-qc].
- [42] Y. Itin, F. W. Hehl, and Y. N. Obukhov, *Phys. Rev. D* **95**, 084020 (2017), arXiv:1611.05759 [gr-qc].
- [43] G. G. L. Nashed and W. El Hanafy, *Eur. Phys. J. C* **77**, 90 (2017), arXiv:1612.05106 [gr-qc].
- [44] M. Hohmann, L. Järv, M. Krššák, and C. Pfeifer, *Phys. Rev. D* **97**, 104042 (2018), arXiv:1711.09930 [gr-qc].
- [45] Z. Chen, W. Luo, Y.-F. Cai, and E. N. Saridakis, *Phys. Rev. D* **102**, 104044 (2020), arXiv:1907.12225 [astro-ph.CO].
- [46] K. Bamba, S. Capozziello, M. De Laurentis, S. Nojiri, and D. Sáez-Gómez, *Phys. Lett. B* **727**, 194 (2013), arXiv:1309.2698 [gr-qc].
- [47] Y.-F. Cai, C. Li, E. N. Saridakis, and L. Xue, *Phys. Rev. D* **97**, 103513 (2018), arXiv:1801.05827 [gr-qc].
- [48] G. Farrugia, J. Levi Said, V. Gakis, and E. N. Saridakis, *Phys. Rev. D* **97**, 124064 (2018), arXiv:1804.07365 [gr-qc].
- [49] M. Hohmann, M. Krššák, C. Pfeifer, and U. Ualikhanova, *Phys. Rev. D* **98**, 124004 (2018), arXiv:1807.04580 [gr-qc].
- [50] R. C. Nunes, S. Pan, and E. N. Saridakis, *Phys. Rev. D* **98**, 104055 (2018), arXiv:1810.03942 [gr-qc].
- [51] V. Gakis, M. Krššák, J. Levi Said, and E. N. Saridakis, *Phys. Rev. D* **101**, 064024 (2020), arXiv:1908.05741 [gr-qc].
- [52] G. G. L. Nashed and E. N. Saridakis, (2022), arXiv:2206.12256 [gr-qc].
- [53] J. W. Maluf and F. F. Faria, *Phys. Rev. D* **85**, 027502 (2012), arXiv:1110.3095 [gr-qc].
- [54] J. W. Maluf and F. F. Faria, *Annalen Phys.* **524**, 366 (2012), arXiv:1203.0040 [gr-qc].
- [55] J. W. Maluf, *Journal of Mathematical Physics* **35**, 335 (1994).
- [56] J. W. Maluf, S. C. Ulhoa, F. F. Faria, and J. F. da Rocha-Neto, *Class. Quant. Grav.* **23**, 6245 (2006), arXiv:gr-qc/0609018.
- [57] J. W. Maluf, *Annalen Phys.* **14**, 723 (2005), arXiv:gr-qc/0504077.
- [58] M. R. Gaberdiel, *Rept. Prog. Phys.* **63**, 607 (2000), arXiv:hep-th/9910156.
- [59] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2011).
- [60] Y. Fujii and K. Maeda, *The scalar-tensor theory of gravitation*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2007).
- [61] S. M. Carroll, *Spacetime and geometry. An introduction to general relativity* (2004).
- [62] A. H. Chamseddine and V. Mukhanov, *JHEP* **11**, 135 (2013), arXiv:1308.5410 [astro-ph.CO].
- [63] A. H. Chamseddine, V. Mukhanov, and A. Vikman, *JCAP* **06**, 017 (2014), arXiv:1403.3961 [astro-ph.CO].
- [64] G. G. L. Nashed and K. Bamba, *Phys. Rev. D* **101**, 044029 (2020), arXiv:1905.07229 [gr-qc].
- [65] S. W. Hawking, T. Hertog, and H. S. Reall, *Phys. Rev. D* **63**, 083504 (2001), arXiv:hep-th/0010232.
- [66] R. Armillis, A. Monin, and M. Shaposhnikov, *JHEP* **10**, 030 (2013), arXiv:1302.5619 [hep-th].
- [67] G. G. L. Nashed and K. Bamba, *JCAP* **09**, 020 (2018), arXiv:1805.12593 [gr-qc].
- [68] I. L. Buchbinder, S. D. Odintsov, and I. L. Shapiro, *Effective action in quantum gravity* (1992).
- [69] I. L. Buchbinder and S. D. Odintsov, Sov. J. Nucl. Phys. **40**, 848 (1984).
- [70] T. Moon, P. Oh, and J. Sohn, *JCAP* **11**, 005 (2010), arXiv:1002.2549 [hep-th].
- [71] G. 't Hooft, *Found. Phys.* **41**, 1829 (2011), arXiv:1104.4543 [gr-qc].
- [72] S. Bahamonde, K. Flathmann, and C. Pfeifer, (2019), arXiv:1907.10858 [gr-qc].
- [73] K. N. Singh, F. Rahaman, and A. Banerjee, (2019), arXiv:1909.10882 [gr-qc].
- [74] G. G. L. Nashed, A. Abebe, and K. Bamba, *Eur. Phys. J. C* **80**, 1109 (2020).
- [75] P. D. Lasky, H. Sotani, and D. Giannios, *Phys. Rev. D* **78**, 104019 (2008), arXiv:0811.2006 [astro-ph].

- [76] F. J. Fattoyev, *Arab. J. Math.* **8**, 293 (2019), arXiv:1905.10767 [gr-qc].
- [77] L. Herrera, *Physics Letters A* **165**, 206 (1992).
- [78] Y. B. Zeldovich and I. D. Novikov, *Relativistic astrophysics. Vol.1: Stars and relativity* (1971).
- [79] T. Gangopadhyay, S. Ray, X.-D. Li, J. Dey, and M. Dey, *Mon. Not. Roy. Astron. Soc.* **431**, 3216 (2013), arXiv:1303.1956 [astro-ph.HE].
- [80] S. Das, F. Rahaman, and L. Baskey, *Eur. Phys. J.* **C79**, 853 (2019).
- [81] C. G. Böhmer and T. Harko, *Classical and Quantum Gravity* **23**, 6479 (2006).
- [82] R. C. Tolman, *Phys. Rev.* **55**, 364 (1939).
- [83] J. R. Oppenheimer and G. M. Volkoff, *Phys. Rev.* **55**, 374 (1939).
- [84] J. Ponce de Leon, *General Relativity and Gravitation* **25**, 1123 (1993).
- [85] S. Chandrasekhar, *Astrophys. J.* **140**, 417 (1964).
- [86] M. Merafina and R. Ruffini, *aap* **221**, 4 (1989).
- [87] R. Chan, L. Herrera, and N. O. Santos, *Monthly Notices of the Royal Astronomical Society* **265**, 533 (1993), <http://oup.prod.sis.lan/mnras/article-pdf/265/3/533/3807712/mnras265-0533.pdf>.
- [88] H. Heintzmann and W. Hillebrandt, *aap* **38**, 51 (1975).
- [89] B. K. Harrison, K. S. Thorne, M. Wakano, and J. A. Wheeler, *Gravitation Theory and Gravitational Collapse* (1965).
- [90] I. B. Zeldovich and I. D. Novikov, *Relativistic astrophysics. Vol.2: The structure and evolution of the universe* (1983).
- [91] A. Golovnev, T. Koivisto, and M. Sandstad, *Classical and Quantum Gravity* **34**, 145013 (2017).
- [92] M. Kršák, R. J. van den Hoogen, J. G. Pereira, C. G. Bohmer, and A. A. Coley, *Classical and Quantum Gravity* **36**, 183001 (2019).
- [93] C. Bejarano, R. Ferraro, F. Fiorini, and M. J. Guzmán, *Universe* **5** (2019), 10.3390/universe5060158.
- [94] L. du Buisson, S. E. Motta, and R. P. Fender, *Mon. Not. Roy. Astron. Soc.* **486**, 4485 (2019), arXiv:1905.00366 [astro-ph.HE].
- [95] J. Sunzu, A. Mathias, and S. Maharaj, *Journal of Astrophysics and Astronomy* **40** (2019), 10.1007/s12036-019-9575-4.
- [96] S. c. v. Ilijic and M. Sossich, *Phys. Rev. D* **98**, 064047 (2018).
- [97] G. Abbas, A. Kanwal, and M. Zubair, *Astrophys. Space Sci.* **357**, 109 (2015), arXiv:1501.05829 [physics.gen-ph].
- [98] D. Momeni, G. Abbas, S. Qaisar, Z. Zaz, and R. Myrzakulov, *Can. J. Phys.* **96**, 1295 (2018), arXiv:1611.03727 [gr-qc].
- [99] G. Abbas, S. Qaisar, and A. Jawad, *apss* **359**, 17 (2015), arXiv:1509.06711 [physics.gen-ph].
- [100] A. Chanda, S. Dey, and B. C. Paul, *Eur. Phys. J.* **C79**, 502 (2019).
- [101] U. Debnath, *Eur. Phys. J.* **C79**, 499 (2019), arXiv:1901.04303 [gr-qc].
- [102] S. Ilijic and M. Sossich, *Phys. Rev. D* **98**, 064047 (2018), arXiv:1807.03068 [gr-qc].

