# Associative submanifolds in twisted connected sum $G_2$ -manifolds

Gorapada Bera

2023-08-07

#### Abstract

We introduce a new method to construct closed rigid (unobstructed) associative submanifolds in the twisted connected sum  $G_2$ -manifolds [Kovo3; KL11; CHNP15]. More precisely, we prove a gluing theorem of asymptotically cylindrical (ACyl) associative submanifolds in ACyl  $G_2$ -manifolds under a hypothesis that can be interpreted as a transverse Lagrangian intersection condition. This is analogous to the gluing theorem for  $G_2$ -instantons introduced in [SW15]. We rephrase the hypothesis in the special cases where the ACyl associative submanifolds are obtained from holomorphic curves or special Lagrangians in ACyl Calabi-Yau 3-folds. In this way we find many new associative submanifolds which are diffeomorphic to  $S^3$ ,  $RP^3$  or  $RP^3$ # $RP^3$ . To lay the foundation for the gluing theorem we also study the moduli space of ACyl associative submanifolds with a natural topology. We prove that the moduli space is locally homeomorphic to the zero set of a smooth map between two finite dimensional spaces whose index depends only on the asymptotic cross section.

#### 1 Introduction: Main results

A  $G_2$ -manifold is a Riemannian 7-manifold whose holonomy group is contained in the exceptional Lie group  $G_2$  or, equivalently a smooth 7-manifold with a torsion free  $G_2$ -structure [Joy96b, Introduction]. The group  $G_2$  is one of the two exceptional holonomy groups in the Berger's classification [Ber55, Theorem 3] of Riemannian manifolds. Simple examples of G<sub>2</sub>-manifolds include product Riemannian manifolds  $S^1 \times Z$ , where Z is a Calabi-Yau 3-fold. However, the holonomy group of these examples is included in the proper subgroup  $SU(3) \subset G_2$ . The most effective method to date of constructing compact manifolds whose holonomy is exactly  $G_2$ , is the twisted connected sum (TCS) construction, which glues a suitable matching pair of asymptotically cylindrical (ACyl)  $G_2$ -manifolds. Kovalev [Kovo3] has pioneered this construction and later was improved by Kovalev-Lee [KL11]. They produce  $G_2$ -manifolds from matching pairs of ACyl Calabi-Yau 3-folds which are constructed from Fano 3-folds or K3-surfaces with non-symplectic involutions. Corti, Haskins, Nordström, and Pacini [CHNP15] have produced hundreds of thousands of  $G_2$ -manifolds by extending this construction to a larger class of ACyl Calabi-Yau 3-folds which are obtained from semi-Fano 3-folds. There has been a considerable enthusiasm in understanding the geometry of  $G_2$ -manifolds in the last decade. This surge in interest can be attributed, in part, to the role that  $G_2$ -manifolds play in M-theory of theoretical high energy physics, which is analogous to the role that Calabi-Yau 3-folds play in string theory. To define an enumerative invariant of  $G_2$ -manifolds which is unchanged under the deformation of  $G_2$ -metrics, Joyce [Joy18], Doan and Walpuski [DW19] have outlined proposals which are based on counting closed associative submanifolds. An associative submanifold in a  $G_2$ -manifold is a 3-dimensional calibrated submanifold calibrated by the 3-form defining the  $G_2$ -structure [HL82, IV.2.A; Joyo7, Chapter 12]. These are volume minimizing in their homology class and hence are minimal submanifolds. They can be regarded as analogous to holomorphic curves in Calabi-Yau 3-folds. Donaldson and Thomas [DT98, Section 3] made a different proposal to define an invariant of  $G_2$ -manifolds by counting  $G_2$ -instantons. But the  $G_2$ -instantons can degenerate by bubbling off along associative submanifolds [Tiaoo] which could cause serious problems to this definition [DS11]. The twisted connected sum (TCS)  $G_2$ -manifolds and associative submanifolds inside them can be a testing ground for these enumerative theories.

The holomorphic curves in Calabi-Yau 3-folds can be constructed directly using algebraic geometry, but the process becomes significantly more challenging when attempting to construct associative submanifolds. Nevertheless, the ACyl Calabi-Yau 3-folds can be obtained from building blocks, that is, smooth projective 3-folds Z together with a projective morphism  $\pi: Z \to \mathbb{CP}^1$  such that  $X := \pi^{-1}(\infty)$  is a smooth anti-canonical K3 surface together with some additional data. A natural idea constructing associative submanifolds in TCS G2-manifolds is therefore to find closed rigid holomorphic curves C in one of the building blocks Z avoiding X, and then deform the product  $S^1 \times C$ . Unfortunately, it is not easy to find such holomorphic curves C, at least not when the building blocks are obtained from Fano 3-folds W, because  $-K_W$  is ample. Corti, Haskins, Nordström, and Pacini [CHNP15] overcome this situation by considering building blocks obtained from semi-Fano 3 folds. They produce some closed rigid associative submanifolds diffeomorphic to  $S^1 \times S^2$  by finding rigid holomorphic lines in the semi-Fano 3-folds. In this article, we improve significantly our abilities of constructing associative submanifolds using gluing technique. This technique now can be applied to holomorphic curves in the building blocks Z which intersects X. More generally, we prove a gluing theorem for a pair of ACyl associative submanifolds in a matching pair of ACyl G2-manifolds, which is similar to the gluing theorem for  $G_2$ -instantons introduced in [SW15, Theorem 2.3.4]. Clearly, the pregluing construction yields approximate associative submanifolds. These can be deformed to genuine associative submanifolds if they are unobstructed in their deformation theory. We impose a set of conditions (mostly algebraic geometric data in the special cases) that guarantee the unobstructedness. In this way we obtain associative 3-spheres in many TCS  $G_2$ -manifolds arising from Fano 3-folds. Our theorem also applies to ACyl special Lagrangian 3-folds, which in turn can be obtained from anti-holomorphic involutions on building blocks. In particular, we construct new rigid associative submanifolds which are diffeomorphic to RP3 and RP3#RP3.

We wish to emphasize that the examples presented in this article only offer a glimpse of the vast potential. Among other things, we expect that this gluing theorem will be important for investigating the interesting conjectures by the physicists Braun, Del Zotto, Halverson, Larfors, Morrison, and Schäfer-Nameki [BDHLMS18] about producing infinitely many closed associative submanifolds in a concrete TCS  $G_2$ -manifold.

The proof of the gluing theorem and the inputs required for the gluing hypothesis motivates us to study the deformation theory and moduli space of ACyl associative submanifolds. In this article, we define the moduli space of ACyl associative submanifolds with a natural topology

and prove that the moduli space is locally homeomorphic to the zero set of a smooth map between two finite-dimensional spaces. We also compute the virtual dimension (which need not be zero). In the best scenario, the moduli space of ACyl associative submanifolds is a Lagrangian submanifold in the moduli space of holomorphic curves in the limiting Calabi-Yau 3-fold and the gluing hypothesis is a transverse Lagrangian intersection condition. We would like to emphasize that the work here can have stand-alone value for various other future purposes. For instance, one could aim to formulate a significant conjecture similar to the Atiyah-Floer conjecture (Remark 1.15). We would like to acknowledge that there exist related studies addressing deformation problems for ACyl special Lagrangian submanifolds [ST10], coassociative submanifolds [JS05] or Caley submanifolds [Ohs15].

The remainder of this section summarizes the main results obtained in this article.

# Gluing of asymptotically cylindrical associative submanifolds

Let  $(Y_{\pm},\phi_{\pm})$  be a matching pair of ACyl  $G_2$ -manifolds with a matching map  $f:Z_{\pm}\to Z_{\pm}$  and let  $\{(Y_T,\phi_T):T\geqslant T_0\}$  be the family of generalized connected sum  $G_2$ -manifolds (see Theorem 2.32). Let  $P_{\pm}$  be a pair of ACyl associative submanifolds in  $Y_{\pm}$  with asymptotic cross sections  $\Sigma_{\pm}$  (see Definition 3.20). Here  $\Sigma_{\pm}$  refer to holomorphic curves in the Calabi-Yau 3-folds  $Z_{\pm}$ . We can preglue  $P_{\pm}$  to construct approximate associative submanifolds  $P_T$ , but these can be deformed to genuine associative submanifolds if they are unobstructed in their deformation theory. Therefore we need to impose a hypothesis on  $P_{\pm}$  that guarantee the unobstructedness. Merely assuming that  $P_{\pm}$  are rigid would have sufficed for our purpose, but this is not the case in the primary application involving twisted connected sum  $G_2$ -manifolds. Instead, we impose a hypothesis that essentially states that the images of the moduli spaces of ACyl associative submanifolds intersect transversely in the moduli space of possible cross sections, so that there is only a discrete set of matching pairs  $P_{\pm}$  that can be glued.




Figure 1: Gluing of a pair of ACyl associative submanifolds  $P_{\pm}$ .

**Hypothesis 1.1.** The ACyl associative submanifolds  $P_{\pm}$  satisfy the following:

• The asymptotic cross sections  $\Sigma_{\pm}$  are matched by f, that is,  $f(\Sigma_{+}) = \Sigma_{-}$ ,

- There are no infinitesimal deformations of  $P_{\pm}$  fixing the asymptotic cross sections  $\Sigma_{\pm}$ , that is, the asymptotic limit maps  $\iota_{\infty}^{\pm}$ :  $\ker \mathbf{D}_{P_{\pm},0} \to \ker \mathbf{D}_{\Sigma_{\pm}}$  are injective (where  $\mathbf{D}_{\Sigma_{\pm}}$  are defined in (3.7) and  $\mathbf{D}_{P_{\pm},0}$  are defined in Definition 3.58 and  $\iota_{\infty}^{\pm}$  are defined in Definition 3.59),
- The images of  $f_*\iota_\infty^+$  and  $\iota_\infty^-$  intersect trivially, that is,  $\operatorname{im}(f_*\iota_\infty^+) \cap \operatorname{im}\iota_\infty^- = \{0\}$ .

Remark 1.2. The first condition in the Hypothesis 1.1 is obviously about the matching of the asymptotic cross sections of  $P_{\pm}$ . The second condition is equivalent to  $P_{\pm}$  being unobstructed, in particular, the moduli spaces of ACyl associative submanifolds near  $P_{\pm}$  are smooth. The last condition concerns the transverse intersection at  $\Sigma \pm$  of these moduli spaces within the moduli space of possible cross sections. Of course, the Hypothesis 1.1 is sufficient to carry out the gluing but not necessary. In fact, it is plausible to relax the requirements of the Hypothesis 1.1 by some transversality conditions on a family of matching pair of ACyl associative submanifolds in a family of matching pair of ACyl  $G_2$ -manifolds. Nevertheless, this aspect will not be addressed in this particular article.

By imposing the Hypothesis 1.1 we prove the following gluing theorem.

**Theorem 1.3.** Let  $(Y_{\pm}, \phi_{\pm})$  be a matching pair of ACyl  $G_2$ -manifolds with a matching map  $f: Z_+ \to Z_-$  and let  $\{(Y_T, \phi_T): T \geq T_0\}$  be the family of generalized connected sum  $G_2$ -manifolds. Let  $P_{\pm}$  be a pair of ACyl associative submanifolds in  $Y_{\pm}$  with asymptotic cross sections  $\Sigma_{\pm}$  satisfying Hypothesis 1.1. Then there exist a constant  $T_1 \geq T_0$  and a family of smooth unobstructed (rigid) associative submanifolds  $\tilde{P}_T$  in  $(Y_T, \phi_T)$  for all  $T \geq T_1$ , which are all diffeomorphic to the connected sum  $P_+ \#_f P_-$ .

The idea of the proof of this theorem is to realize the ends of  $P_{\pm}$  as graphs of sections of the normal bundles of the asymptotic associative cylinders and then glue these graphs together using cut-off functions. Depending on the parameter T we obtain a family of approximate associative submanifolds  $P_T$ , that is, the error of being associative is very small when T is sufficiently large. Under the Hypothesis 1.1 we can then deform  $P_T$  for sufficiently large T using an application of the Banach contraction principle (see Lemma 4.16) on suitable Banach spaces to construct a family of rigid associative submanifolds  $\tilde{P}_T$ . This is to say that we solve a nonlinear PDE:  $\mathfrak{F}_T u = 0$  (see Definition 4.4) whose solutions are associative submanifolds in a tubular neighbourhood of  $P_T$ . To apply Lemma 4.16 we decompose the operator  $\mathfrak{F}_T$  as

$$\mathfrak{F}_T = \mathfrak{L}_T + Q_T + e_T$$

where  $\mathfrak{L}_T$  is the linearization of  $\mathfrak{F}_T$  (see Definition 4.5),  $e_T$  is the error of  $P_T$  to be an associative (see Definition 4.8) and prove that the error  $e_T$  (see Proposition 4.11) is very small compared to the lower bound of  $\mathfrak{L}_T$  (see Proposition 4.14) when T is sufficiently large. The Hypothesis 1.1 has been only used to show that the index zero operator  $\mathfrak{L}_T$  has zero kernel (see Proposition 4.14) and hence is invertible.

It seems very difficult to produce examples of associative submanifolds using Theorem 1.3, since testing the Hypothesis 1.1 in the way it has been stated is a hard task. Therefore we restrict ourselves to the twisted connected sum  $G_2$ -manifolds that are obtained from ACyl Calabi-Yau 3-folds and rewrite the Hypothesis 1.1 for pairs of ACyl associative submanifolds that are either coming from ACyl holomorphic curves or ACyl special Lagrangian 3-folds.

Haskins, Hein, and Nordström [HHN15] have proved (see Theorem 2.22) that ACyl Calabi-Yau 3-folds are essentially equivalent to framed building blocks (see Definition 2.21). As an application to the Theorem 1.3 we prove the following gluing theorem which constructs closed associative submanifolds from holomorphic curves in the building blocks.

**Theorem 1.4.** Let  $(Z_{\pm}, X_{\pm}, \omega_{\pm})$  be a pair of framed building blocks with a hyperkähler rotation  $\mathfrak{r}: X_{+} \to X_{-}$ . Let  $V_{\pm} := Z_{\pm} \backslash X_{\pm}$  be a pair of corresponding ACyl Calabi-Yau 3-folds and let  $\{(Y_{T}, \phi_{T})\}$  be the family of  $G_{2}$ -manifolds obtained from the twisted connected sum construction. Let  $C_{\pm}$  be a pair of holomorphic curves in  $Z_{\pm}$  intersecting  $X_{\pm}$  transversely at  $\bar{x}_{\pm} := \{x_{1,\pm}, x_{2,\pm}, ..., x_{m,\pm}\}$ . Then  $C_{\pm} \backslash \bar{x}_{\pm}$  are ACyl holomorphic curves in  $V_{\pm}$  with asymptotic cross sections  $\coprod_{j=1}^{m} S^{1} \times \{x_{j,\pm}\} \subset S^{1} \times X_{\pm}$ . Assume that

- $\mathfrak{r}(\bar{x}_+) = \bar{x}_-$
- $H^0(C_{\pm}, NC_{\pm}(-\bar{x}_{\pm})) = 0$ , that is, there are no holomorphic normal vector fields of  $C_{\pm}$  vanishing at  $\bar{x}_{\pm}$ ,
- $\operatorname{im}(\mathfrak{r}_* \operatorname{ev}_+) \cap \operatorname{im}(\operatorname{ev}_-) = \{0\}, where$  $\operatorname{ev}_{\pm} := \bigoplus_{j=1}^m \operatorname{ev}_{X_{j,\pm}} : H^0(C_{\pm}, NC_{\pm}) \to \bigoplus_{j=1}^m N_{X_{j,\pm}} C_{\pm} \cong \bigoplus_{j=1}^m T_{X_{j,\pm}} X_{\pm}$

are the evaluations maps.

Then the pair of ACyl associative submanifolds  $S^1 \times (C_{\pm} \setminus \bar{x}_{\pm})$  satisfies Hypothesis 1.1 and therefore there is a family of closed rigid associative submanifolds  $\tilde{P}_T$  for sufficiently large T. These are diffeomorphic to the connected sum  $(S^1 \times (C_{+} \setminus \bar{x}_{+})) \#_{\tau}(S^1 \times (C_{-} \setminus \bar{x}_{-}))$  along the m Clifford tori, where  $\tau: T^2 \to T^2$  is given by  $\tau(\theta, s) = (s, \theta)$ .

As an application to the Theorem 1.4 we prove the next proposition that will help us to produce examples of associative submanifolds in the twisted connected sum  $G_2$ -manifolds. In particular, we construct associative 3-spheres in a lot of twisted connected sum  $G_2$ -manifolds arising from Fano 3-folds (see Example 4.20).

**Proposition 1.5.** Let  $(Z_+, X_+)$  be a building block and let  $\ell_+$  be a unobstructed holomorphic line in  $Z_+$  (that is,  $H^1(\ell_+, N\ell_+) = 0$ ) intersecting  $X_+$  transversely at a point. Let  $W_-$  be a semi-Fano 3-fold and let  $X_- \in |-K_{W_-}|$  be a smooth K3-surface in  $W_-$  such that  $-K_{W_-|_{X_-}}$  is very ample and there exists a hyperkähler rotation  $\mathfrak{r}: X_+ \to X_-$ . Then there is a building block  $(Z_-, X_-)$  constructed by blowing up a base locus of an anti-canonical pencil in  $W_-$  (see Theorem 2.25) such that the corresponding family of twisted connected sum  $G_2$ -manifolds  $(Y_T, \phi_T)$  contain always closed rigid associative 3-spheres  $\tilde{P}_T$  for sufficiently large T.

As another application to the Theorem 1.3 we prove the following theorem which produces associative submanifolds from ACyl special Lagrangian 3-folds.

**Theorem 1.6.** Let  $(V_{\pm}, \omega_{\pm}, \Omega_{\pm})$  be a pair of ACyl Calabi-Yau 3-folds with asymptotic cross sections  $(X_{\pm}, \omega_{I,\pm}, \omega_{J,\pm}, \omega_{K,\pm})$  with  $\mathfrak{r}: X_{+} \to X_{-}$  a hyperkähler rotation. Let  $\{(Y_{T}, \phi_{T})\}$  be the family of  $G_{2}$ -manifolds obtained from the twisted connected sum construction. Let  $L_{\pm}$  be a matching pair of ACyl special Lagrangian 3-folds with cross sections  $\Sigma_{s_{\pm}} := \{e^{is_{\pm}}\} \times \Sigma_{\pm}$  in  $V_{\pm}$ , where  $\Sigma_{\pm}$  are  $K_{\pm}$ -holomorphic curves in  $X_{\pm}$  (see Example 3.25). Let  $i_{\pm}^{*}: H^{1}(L_{\pm}, \mathbb{R}) \to H^{1}(\Sigma_{\pm}, \mathbb{R})$  be the map induced by the inclusion  $\Upsilon_{\pm} \circ \iota_{\pm} : \Sigma_{\pm} \cong \{T\} \times \Sigma_{\pm} \to L_{\pm}$  for any large T. Assume that

- $\mathfrak{r}(\Sigma_+) = \Sigma_-$ ,
- $b^2(L_{\pm}) b^0(\Sigma_{\pm}) + b^0(L_{\pm}) = 0$ ,
- $\operatorname{im}(K_{+} \circ i_{+}^{*}) \cap \operatorname{im}(\mathfrak{r}^{*} \circ i_{-}^{*}) = \{0\}.$

Then  $L_{s_{\pm}} := \{e^{is_{\pm}}\} \times L_{\pm}$  satisfies Hypothesis 1.1 and hence we obtain a family of closed rigid associative submanifolds  $\tilde{P}_T$  for sufficiently large T which are diffeomorphic to the connected sum  $L_{+}\#_r L_{-}$ .

The last two conditions in the assumption of Theorem 1.6 hold immediately if  $b^1(L_\pm) = 0$ . An easy way to find ACyl special Lagrangian 3-folds is to find anti-holomorphic involutions on building blocks. The following proposition explains how to use anti-holomorphic involutions on building blocks to construct closed associative submanifolds in the twisted connected sum  $G_2$ -manifolds. In particular, we construct associative submanifolds which are diffeomorphic to  $RP^3$  (see Example 4.22) or  $RP^3$ # $RP^3$  (see Example 4.23, Example 4.24). We impose the following hypothesis on building blocks to satisfy the conditions in Theorem 1.6.

**Hypothesis 1.7.** The building block  $(Z_+, X_+)$  satisfies the following:

- $Z_+$  admits an anti-holomorphic involution  $\sigma_+:Z_+\to Z_+$  which takes  $X_+$  to itself,
- $X_+$  admits a non-symplectic involution  $\rho_+$  which commutes with  $\sigma_{+|X_+}$ ,
- $\operatorname{Fix}_{\sigma_+}(X_+) \cap \operatorname{Fix}_{\rho_+}(X_+) = \emptyset$  and  $b^1(\Sigma_+/\langle \rho_+ \rangle) = 0$  where  $\Sigma_+ := \operatorname{Fix}_{\sigma_+}(X_+)$ ,
- Fix<sub> $\sigma_+$ </sub>(Z<sub>+</sub>)\X<sub>+</sub> has a connected, non-compact component L<sub>+</sub> with  $b^1(L_+) = 0$ .

**Proposition 1.8.** Let  $(Z_+, X_+)$  be a building block satisfying Hypothesis 1.7. Then  $V_+ := Z_+ \setminus X_+$  admits an ACyl Calabi-Yau structure  $(\omega_+, \Omega_+)$  such that  $\sigma_+$  is an anti-holomorphic involutive isometry on  $V_+$ . If the hyperkähler structure of  $X_+$  is  $(\omega_{I,+}, \omega_{J,+}, \omega_{K,+})$ , then take  $X_-$  to be same as  $X_+$  but with hyperkähler structure  $(\omega_{J,+}, \omega_{I,+}, -\omega_{K,+})$  so that the identity map  $\mathbf{1}: X_+ \to X_-$  becomes a hyperkähler rotation. Then  $\rho_- := \rho_+ \circ \sigma_+$  is a non-symplectic involution on  $X_-$ . Define  $W_- := \frac{\mathrm{CP}^1 \times X_-}{\langle \iota \times \rho_- \rangle}$  and  $Z_-$  to be the blow-up of  $W_-$  along the fixed point locus (see Theorem 2.28) and  $\tilde{L}_-$  to be the proper transform of  $\frac{\mathrm{RP}^1 \times \Sigma_+}{\langle \iota \times \rho_+ \rangle}$ . Denote  $L_- := \tilde{L}_- \setminus X_-$ . Then the pair of ACyl special Lagrangians  $L_+$  satisfies Theorem 1.6 and hence we obtain a family of closed rigid associative submanifolds  $\tilde{P}_T$  with sufficiently large T.

#### Moduli space of asymptotically cylindrical associative submanifolds

We will consider first the moduli space of closed holomorphic curves in Calabi-Yau 3-folds, since they appear as the asymptotic cross sections of ACyl associative submanifolds. We express the moduli space of holomorphic curves in a Calabi-Yau 3-fold locally as the zero set of a nonlinear map (see Definition 3.9) and use Fredholm theory [DK90, Section 4.2.4] to prove the following theorem.

**Theorem 1.9.** Let  $\Sigma$  be a closed holomorphic curve in a Calabi-Yau 3-fold (Z, J). Denote by  $\mathfrak{M}_Z^{hol}$  the moduli space of closed holomorphic curves in Z (see Definition 3.2). Then  $\mathfrak{M}_Z^{hol}$  near  $\Sigma$  is homeomorphic to the zero set of a smooth map (obstruction map/Kuranishi map)

$$ob_{\Sigma}: \mathcal{I}_{\Sigma} \to coker \mathbf{D}_{\Sigma},$$

where the operator  $\mathbf{D}_{\Sigma}$  is a Dirac operator defined in (3.7) and  $\mathfrak{I}_{\Sigma}$  is an open neighbourhood of 0 in  $\ker \mathbf{D}_{\Sigma}$ . Moreover the index of the operator  $\mathbf{D}_{\Sigma}$  is zero and there is a J-anti-linear isomorphism  $\boldsymbol{\gamma}_{\Sigma}$ :  $\overline{\operatorname{Hom}}_{\mathbb{C}}(T\Sigma, N\Sigma) \to N\Sigma$  such that the normal Cauchy-Riemann operator  $\bar{\partial}_{\nabla,J}^N$  (see Definition 3.3) satisfies

$$\boldsymbol{\gamma}_{\Sigma} \circ \bar{\partial}_{\nabla,I}^{N} = \mathbf{D}_{\Sigma}.$$

**Definition 1.10.** A closed holomorphic curve  $\Sigma$  in a Calabi-Yau 3-fold Z is called **Morse-Bott** if the obstruction map  $ob_{\Sigma}$  vanishes near 0 in  $\mathfrak{I}_{\Sigma}$ .

*Remark* 1.11. If a closed holomorphic curve  $\Sigma$  in a Calabi-Yau 3-fold Z is Morse-Bott (see Example 3.11, Example 3.12) then in Proposition 3.65 we see that the moduli space of holomorphic curves near  $\Sigma$  is a smooth symplectic manifold whose dimension is dim ker  $\mathbf{D}_{\Sigma}$ .

The following theorem is about the moduli space of ACyl associative submanifolds with a fixed rate and a fixed asymptotic cross section.

**Theorem 1.12.** Let P be an ACyl associative submanifold in an ACyl  $G_2$ -manifold Y with asymptotic cross section  $\Sigma := \coprod_{i=1}^m \Sigma_i$  and rate  $\mu := (\mu_1, \ldots, \mu_m) < 0$  (see Definition 3.20). Assume that  $\mu$  lies outside the wall of critical rates (see Definition 3.31). Then the moduli space  $\mathcal{M}_{ACyl,\Sigma}^{\mu}$  of ACyl associative submanifolds with the fixed rate  $\mu$  and the fixed asymptotic cross section  $\Sigma$  near P is homeomorphic (with  $C_{\mu}^{\infty}$ -topology, see Definition 3.46) to the zero set of a smooth map (obstruction map/Kuranishi map)

$$ob_{P,\mu}: \mathcal{I}_{P,\mu} \to coker \mathbf{D}_{P,\mu}$$

where  $\mathfrak{I}_{P,\mu}$  is a neighbourhood of 0 in  $\ker \mathbf{D}_{P,\mu}$  and  $\mathbf{D}_{P,\mu}$  is the Fueter operator defined in Definition 3.58. Moreover,

index 
$$D_{P,\mu} = -\sum_{i=1}^{m} \frac{d_{0,i}}{2} - \sum_{i=1}^{m} \sum_{\lambda_i \in (\mu_i,0)} d_{\lambda_i}$$

where  $d_{\lambda_i}$  is the dimension of the  $\lambda_i$ -eigenspace of  $JD_{\Sigma_i}$  and  $d_{0,i} := \dim \ker JD_{\Sigma_i}$  (see Definition 3.14). Moreover  $\mathcal{M}^{\mu}_{ACyl,\Sigma} = \mathcal{M}^{\mu'}_{ACyl,\Sigma}$ , if  $\mu$  and  $\mu'$  are lying in the same connected component of the complement of the wall of critical rates.

The following theorem is about the moduli space of ACyl associative submanifolds where the asymptotic cross sections are allowed to vary.

**Theorem 1.13.** Let P be an ACyl associative submanifold in an ACyl  $G_2$ -manifold Y with asymptotic cross section  $\Sigma := \coprod_{i=1}^m \Sigma_i$ . Then the moduli space  $\mathcal{M}_{ACyl}$  of ACyl associative submanifolds near P is homeomorphic (with weighted  $C^{\infty}$ -topology, see Definition 3.47) to the zero set of a smooth map (obstruction map/Kuranishi map)

$$\widetilde{\operatorname{ob}}_{P,u}:\widetilde{\mathbb{I}}_{P,u}\to\operatorname{coker}\widetilde{\mathbf{D}}_{P,u}\cong\ker\mathbf{D}_{P,u}$$

for some rate  $\mu < 0$ . Here  $\tilde{\mathbb{J}}_{P,\mu}$  is a neighbourhood of (0,0) in  $\ker \widetilde{\mathbf{D}}_{P,\mu}$  and  $\widetilde{\mathbf{D}}_{P,\mu}$  is defined in Definition 3.64. Moreover,

$$\operatorname{index} \widetilde{\mathbf{D}}_{P,\mu} = \frac{1}{2} \sum_{i=1}^{m} \dim \ker \mathbf{D}_{\Sigma_{i}} = \sum_{i=1}^{m} \frac{d_{0,i}}{2}.$$

*Remark* 1.14. Suppose P is an ACyl associative submanifold with Morse-Bott asymptotic cross section  $\Sigma$  (see Definition 1.10) and ker  $\mathbf{D}_{P,\mu} = 0$ . In Proposition 3.65 we see that the moduli space of ACyl associative submanifolds near P is a Lagrangian submanifold of the moduli space of holomorphic curves near  $\Sigma$ .

Remark 1.15. If a pair of ACyl associative submanifolds  $P_{\pm}$  satisfies Hypothesis 1.1 and the asymptotic cross sections  $\Sigma_{\pm}$  of  $P_{\pm}$  are Morse-Bott, then the moduli spaces of ACyl associative submanifolds around  $P_{\pm}$  are Lagrangian submanifolds in the moduli space holomorphic curves which intersect transversally at  $\Sigma_{\pm}$ . This Lagrangian intersection yields a unique closed rigid associative submanifold in the twisted connected sum  $G_2$ -mannifold by Theorem 1.3. This possibly gives a hint for formulating an Atiyah-Floer type conjecture [Ati88] by considering twisted connected sum  $G_2$ -manifolds as an analog of Heegaard splitting of 3-manifolds. The possible analog of Atiyah-Floer conjecture would be some relationship between yet-to-be defined Floer homology of closed associative submanifolds say, HA(Y) and Lagrangian Floer homology say,  $HF(\mathcal{M}_{ACyl}(Y_+), \mathcal{M}_{ACyl}(Y_-))$  where  $Y = Y_+ \bigcup_Z Y_-$  is the twisted connected sum  $G_2$ -manifold. A similar analog with  $G_2$ -instantons has been pointed out in [SW15, Remark 1.7].

Structure of the article. In Section 2 we review the basic definitions and facts about  $G_2$ -manifolds, associative submanifolds and twisted connected sum construction. Section 3 focuses on ACyl associative submanifolds. In particular, we prove Theorem 1.9 and study the Fueter operator on associative cylinders in Section 3.1. The definition and examples of ACyl associative submanifolds have been discussed in Section 3.2. The Fredholm theory of asymptotically translation invariant elliptic operators appears in Section 3.3. The moduli space of ACyl associative submanifolds, and the proof of Theorem 1.12 and Theorem 1.13 are presented in Section 3.4. The focus of Section 4 is on gluing ACyl associative submanifolds in ACyl  $G_2$ -manifolds, where the proof of the main gluing theorem, Theorem 1.3 has been completed. In addition, we prove Theorem 1.4 and Proposition 1.5 in Section 4.3 that are based on the groundwork in Section 3.6 about ACyl holomorphic curves. Finally, the proof of Theorem 1.6 and Proposition 1.8 is presented in Section 4.3 that are based on the groundwork in Section 3.5 about ACyl special Lagrangian 3-folds.

Acknowledgements. I am grateful to my PhD supervisor Thomas Walpuski for suggesting the problem solved in this article. I am also indebted to him for his constant encouragement and advice during this work. Additionally, I extend my thanks to Thomas Walpuski, Johannes Nordström, and Jason Lotay for their meticulous review of the draft and valuable feedback. I also thank to Dominik Gutwein and Viktor Majewski for their careful reading and helpful comments on the draft. This material is based upon work supported by the Simons Collaboration on Special

Holonomy in Geometry, Analysis, and Physics and the National Science Foundation under Grant No. DMS-1928930, while the author was in residence at the Simons Laufer Mathematical Institute (previously known as MSRI), Berkeley, during the Fall 2022 semester.

#### 2 Preliminaries

In this section we review definitions and basic facts about  $G_2$ -manifolds and associative submanifolds, which are important to understand this article. To delve further into these topics, we refer the reader to [SW17; Joyo7; KLL20; Har90] and other relevant sources mentioned throughout the discourse.

#### 2.1 $G_2$ -manifolds and associative submanifolds

The group  $G_2$  is the automorphism group of the normed division algebra of octonions O, i.e.  $G_2 := \operatorname{Aut}(O) \subset \operatorname{SO}(7)$ . This is a simple, compact, connected, simply connected Lie group of dimension 14. Writing  $O = \operatorname{Re} O \oplus \operatorname{Im} O \cong \mathbf{R} \oplus \mathbf{R}^7$ , we define the **cross-product**  $\times : \Lambda^2 \mathbf{R}^7 \to \mathbf{R}^7$  by  $(u, v) \mapsto u \times v := \operatorname{Im}(uv)$ . The 3-form  $\phi_e \in \Lambda^3(\mathbf{R}^7)^*$  is defined by  $\phi_e(u, v, w) := g_e(u \times v, w)$ , where  $g_e : S^2(\mathbf{R}^7) \to \mathbf{R}$  is  $g_e(u, v) = -\operatorname{Re}(uv)$ , the standard Euclidean metric on  $\mathbf{R}^7$ . These are related by an important identity

(2.1) 
$$\iota_{u}\phi_{e} \wedge \iota_{v}\phi_{e} \wedge \phi_{e} = 6q_{e}(u, v)\operatorname{vol}_{a_{e}}.$$

The Lie group  $G_2$  can also be expressed as  $G_2 := \{A \in GL(\mathbf{R}^7) : A^*\phi_e = \phi_e\}$ . There is also a 4-form  $\psi_e := *_{g_e}\phi_e \in \Lambda^4(\mathbf{R}^7)^*$ , which also can be defined by  $\psi_e(u, v, w, z) := g_e([u, v, w], z)$ , where  $[\cdot, \cdot, \cdot] : \Lambda^3(\mathbf{R}^7) \to \mathbf{R}^7$  is the associator, defined as follows:

$$[u, v, w] := (u \times v) \times w + \langle v, w \rangle u - \langle u, w \rangle v.$$

**Definition 2.2.** A  $G_2$ -manifold is a 7-dimensional manifold Y equipped with a torsion-free  $G_2$ -structure, that is, equipped with a definite  $^1$  3-form  $\phi \in \Omega^3(Y)$  such that  $\nabla_{g_\phi} \phi = 0$ , or equivalently  $^2$ 

$$d\phi = 0$$
 and  $d\psi = 0$ .

The definite 3-form  $\phi$  on Y defines uniquely a Riemannian metric  $g_{\phi}$  and a volume form  $\operatorname{vol}_{g_{\phi}}$  on Y satisfying the identity (2.1). Moreover it defines

- a cross product  $\times : \Lambda^2(TY) \to TY$ ,
- an associator  $[\cdot,\cdot,\cdot]:\Lambda^3(TY)\to TY$ ,
- a 4-form  $\psi := *_{a_{\phi}} \phi \in \Omega^{4}(Y)$ .

Every  $G_2$ -manifold  $(Y, \phi)$  is a spin manifold (see [Gra69, Theorem 3.1-3.2]) and admits a nowhere vanishing parallel spinor, and therefore the metric  $g_{\phi}$  is Ricci-flat [LM89, pg. 321; Hit74, Theorem 1.2]. A compact  $G_2$ -manifold Y has holonomy exactly equal to  $G_2$  if and only if  $\pi_1(Y)$  is finite [Joyo7, Proposition 11.2.1].

<sup>&</sup>lt;sup>1</sup>i.e. the bilinear form  $G_{\phi}: S^2TY \to \Lambda^7(T^*Y)$  defined by  $G_{\phi}(u,v) := \iota_u \phi \wedge \iota_v \phi \wedge \phi$  is definite.

<sup>&</sup>lt;sup>2</sup>The equivalence was established by Fernández and Gray [FG82, Theorem 5.2]

**Example 2.3.** ( $\mathbb{R}^7$ ,  $\phi_e$ ) is a  $G_2$ -manifold.

**Example 2.4.** Let  $(X, \omega_I, \omega_J, \omega_K)$  be a hyperkähler 4-manifold. The manifolds  $\mathbb{R}^3 \times X$  and  $T^3 \times X$  are  $G_2$ -manifolds with  $G_2$ -structure

$$\phi := dt^1 \wedge dt^2 \wedge dt^3 + dt^1 \wedge \omega_I + dt^2 \wedge \omega_I + dt^3 \wedge \omega_K,$$

where  $(t^1, t^2, t^3)$  are the coordinates of  $\mathbb{R}^3$ .

**Example 2.5.** Let  $(Z, \omega, \Omega)$  be a Calabi-Yau 3-fold, where  $\omega$  is a Kähler form and  $\Omega$  is a holomorphic volume form on Z satisfying  $\frac{\omega^3}{3!} = \frac{i}{8}\Omega \wedge \bar{\Omega}$ . The product with the unit circle,  $Y := S^1 \times Z$  is a  $G_2$ -manifold with the  $G_2$ -structure

$$\phi := dt \wedge \omega + \operatorname{Re} \Omega, \ \psi = \frac{1}{2}\omega \wedge \omega + dt \wedge \operatorname{Im} \Omega,$$

where t denotes the coordinate on  $S^1$ . In this case the holonomy group  $\operatorname{Hol}(Y, g_{\phi}) \subset \operatorname{SU}(3)$ .

**Example 2.6.** Bryant [Bry87], Bryant and Salamon [BS89] first constructed local and complete manifolds with holonomy equal to  $G_2$  respectively. Joyce [Joy96c] first constructed compact manifolds with holonomy equal to  $G_2$  by smoothing flat  $T^7/\Gamma$  where  $\Gamma$  is a finite group of isometries of  $T^7$ . This has been generalized later by Joyce and Karigiannis [JK17]. Kovalev [Kovo3] introduced the twisted connected sum construction of  $G_2$ -manifolds which glues a suitable matching pair of asymptotically cylindrical  $G_2$ -manifolds. This construction was later improved by Kovalev and Lee [KL11] and Corti, Haskins, Nordström, and Pacini [CHNP15] to produce hundreds of thousands examples of compact manifolds with holonomy equal to  $G_2$ .

In a  $G_2$ -manifold Harvey and Lawson [HL82] considered a special class of 3-dimensional calibrated submanifolds, called associative submanifolds.

**Definition 2.7.** Let  $(Y, \phi)$  be a  $G_2$ -manifold. A 3-dimensional oriented submanifold P of Y is called an **associative submanifold** if it is calibrated by the 3-form  $\phi$ , that is,  $\phi_{|P}$  is the volume form  $\operatorname{vol}_{P,g_{\phi}}$  on P, or equivalently  $^3[u,v,w] = 0$ , for all  $x \in P$  and  $u,v,w \in T_xP$ .

Any closed associative submanifold is volume minimizing in its homology class and hence is a minimal submanifold [HL82, Theorem 4.2].

**Example 2.8.** In Example 2.4,  $\mathbb{R}^3$  and  $T^3$  are associative submanifolds.

**Example 2.9.** Let Z be a Calabi-Yau 3-fold and  $S^1 \times Z$  be the  $G_2$ -manifold as in Example 2.5. For any holomorphic curve  $\Sigma \subset Z$  and special Lagrangian (i.e. calibrated by Re  $\Omega$ )  $L \subset Z$  the 3-dimensional submanifolds  $S^1 \times \Sigma$  and  $\{e^{i\theta}\} \times L$  are associative submanifolds of  $S^1 \times Z$  for all  $e^{i\theta} \in S^1$ .

**Example 2.10.** Joyce [Joy96a, Section 4.2] has produced examples of closed associative submanifolds which are the fixed point loci of  $G_2$ -involutions in his generalized Kummer constructions. Recently new examples of associative submanifolds in generalized Kummer constructions have been constructed by Dwivedi, Platt, and Walpuski [DPW23].

<sup>&</sup>lt;sup>3</sup>The equivalence follows from the identity  $|u \wedge v \wedge w|^2 = \phi(u, v, w)^2 + |[u, v, w]|^2$ , see [HL82, Chapter IV, Theorem 1.6]

**Example 2.11.** Examples of closed associative submanifolds in the twisted connected sum (TCS)  $G_2$ -manifolds first constructed by Corti, Haskins, Nordström, and Pacini [CHNP15, Section 5, Section 7.2.2] from closed holomorphic curves and closed special Lagrangians in asymptotically cylindrical (ACyl) Calabi-Yau submanifolds.

There is Dirac operator  $D_P$  which controls the deformation theory of associative submanifolds (for more details, see [McL98, Theorem 5.2; Joy18, Theorem 2.12; DW19, Theorem 2.20, Proposition 2.23, Section 2.7]).

**Definition 2.12.** Let P be an associative submanifold (compact or noncompact) of a  $G_2$ -manifold  $(Y, \phi)$ . The **Fueter operator**  $\mathbf{D}_P : C^{\infty}(NP) \to C^{\infty}(NP)$  is defined by

$$\mathbf{D}_P v := \sum_{i=1}^3 e_i \times \nabla_{P,e_i}^{\perp} v$$

Here NP is the normal bundle of P and  $\nabla_P^{\perp}$  is the normal connection and  $\{e_1, e_2, e_3\}$  is an oriented local orthonormal frame for TP with respect to the metric  $g_{\phi}$ .

A closed associative submanifold P of a  $G_2$ -manifold is called **rigid** or equivalently **unobstructed** if ker  $\mathbf{D}_P = \{0\}$ .

The following definitions about normal bundles and tubular neighbourhood maps are useful to set up the deformation theory for associative submanifolds in this article.

**Definition 2.13 (Normal bundles and tubular neighbourhood maps).** Let M be any submanifold of a manifold Y. The **normal bundle**  $\pi: NM \to M$  is characterised by the exact sequence

$$(2.14) 0 \to TM \to TY_{|M} \to NM \to 0.$$

The tangent bundle TNM fits into the exact sequence  $0 \to \pi^* NM \xrightarrow{i} TNM \xrightarrow{d\pi} \pi^* TM \to 0$ . This induces an inclusion of the space normal vector fields of M into the space vector fields of NM,

$$\tilde{\cdot}: C^{\infty}(NM) \hookrightarrow \text{Vect}(NM), \quad \tilde{u}:=i(\pi^*u).$$

Here  $\pi^*u \in C^{\infty}(\pi^*NM)$  is the pull back section. Observe that,  $[\tilde{u}, \tilde{v}] = 0$  for all  $u, v \in C^{\infty}(NM)$ . For any section  $u \in C^{\infty}(NM)$  we define the **graph of** u by

$$\Gamma_u := \{ (x, u(x)) \in NM : x \in M \}.$$

For any section  $u \in C^{\infty}(NM)$ , the graph,  $\Gamma_u$  is a submanifold of NM and the bundle  $\pi^*NM_{|\Gamma_u}$  fits into the split exact sequence

$$(2.15) 0 \longrightarrow T\Gamma_{u} \xrightarrow{d(u \circ \pi)} TNM_{|\Gamma_{u}} \xrightarrow{1-d(u \circ \pi)} \pi^{*}NM_{|\Gamma_{u}} \longrightarrow 0.$$

In particular, this induces an isomorphism  $N\Gamma_u \cong \pi^* NM_{|\Gamma_u}$ . Moreover, the composition  $T\Gamma_u \to TNM_{|\Gamma_u} \xrightarrow{d\pi} \pi^* TM_{|\Gamma_u}$  is an isomorphism.

The choice of a Riemannian metric g on Y induces a splitting of the exact sequence (2.14), that is,  $TY_{|M} = TM \perp NM$ . Denote by  $\cdot^{\parallel}$  and  $\cdot^{\perp}$  the projections onto the first and second summands respectively. The Levi-Civita connection  $\nabla$  on  $TY_{|M}$  decomposes as

$$\nabla = \begin{bmatrix} \nabla^{\parallel} & -\mathrm{II}^* \\ \mathrm{II} & \nabla^{\perp} \end{bmatrix}.$$

Here II  $\in \text{Hom}(S^2TM, NM)$  is the **second fundamental form** of M,  $\nabla^{\parallel}$  is the Levi-Civita connection on M and  $\nabla^{\perp}$  is the **normal connection** on NM.

A tubular neighbourhood map of M is a diffeomorphism between an open neighbouhood  $V_M$  of the zero section of the normal bundle NM of M which is convex in each fiber and an open neighbourhood  $U_M$  (tubular neighbourhood) of M in Y,

$$\Upsilon_M:V_M\to U_M$$

that takes the zero section 0 to M and the composition  $NM \to 0^*TNM \xrightarrow{dY_M} TY_{|M} \to NM$  is the identity.

Let  $\Upsilon_M: V_M \to U_M$  be a tubular neighbourhood map of M. We define

$$C^{\infty}(V_M) := \{ u \in C^{\infty}(NM) : \Gamma_u \subset V_M \}.$$

Let  $u \in C^{\infty}(V_M)$ . Denote by  $M_u$  the submanifold  $\Upsilon_M(\Gamma_u)$  of Y. Then there is a **canonical** bundle isomorphism

$$NM \xrightarrow{\Theta_u^M} NM_u$$

$$\downarrow \qquad \qquad \downarrow$$

$$M \xrightarrow{\Upsilon_M \circ u} M_u.$$

induced by the following commutative diagram of bundle isomorphisms.

$$NM \xrightarrow{\Theta_u^M} NM_u$$

$$\pi^* \left( \downarrow u^* \qquad d\Upsilon_M \right)$$

$$\pi^* NM_{|\Gamma_u} \xleftarrow{1 - d(u \circ \pi)} N\Gamma_u.$$

**Notation 2.16.** By abusing notation we will always denote the above  $\tilde{u}$  again by u.

# 2.2 The twisted connected sum (TCS) construction

In this section we briefly discuss the twisted connected sum construction following [CHNP15].

**Definition 2.17.** Let  $(Z, \omega, \Omega)$  be a compact Calabi-Yau 3-fold, where  $\omega$  is the Kähler form and  $\Omega$  is the holomorphic volume form. A  $G_2$ -manifold  $(Y, \phi)$  is called an **asymptotically cylindrical** (ACyl)  $G_2$ -manifold with asymptotic cross section  $(Z, \omega, \Omega)$  and rate  $\nu < 0$  if there exist

• a compact submanifold  $K_Y$  with boundary and a diffeomorphism

$$\Upsilon: \mathbf{R}^+ \times Z \to Y \backslash K_Y$$

• a 2-form  $\varrho$  on  $\mathbb{R}^+ \times Z$  such that  $\Upsilon^* \varphi = dt \wedge \omega + \operatorname{Re} \Omega + d\varrho$  with

$$|\nabla^k \varrho| = O(e^{\nu t}) \text{ as } t \to \infty, \forall k \in \mathbb{N} \cup \{0\}.$$

Here t denotes the coordinate on  $\mathbb{R}^+$ ,  $|\cdot|$  and Levi-Civita connection  $\nabla$  are induced by the product metric on  $\mathbb{R}^+ \times Z$ .

Remark 2.18. Since ACyl  $G_2$ -manifolds are Ricci flat and complete, by the Cheeger-Gromoll splitting theorem [CG71; Salo6, Theorem 1.1] they can have at most two ends, if they are connected. If there are two ends, then the holonomy group reduces. It was shown in [Noro8a, Theorem 3.8] that  $Hol(Y) = G_2$  if and only if the fundamental group  $\pi_1(Y)$  is finite and neither Y nor any double cover of Y is homeomorphic to a cylinder. Examples of ACyl  $G_2$ -manifolds with holonomy exactly  $G_2$  are hard to find and the first examples of this kind were found by Kovalev-Nordström [KN10, Section 4-5].

**Definition 2.19**. Let  $(X, \omega_I, \omega_J, \omega_K)$  be a hyperkähler 4-manifold. A Calabi-Yau 3-fold  $(V, \omega, \Omega)$  is called an **asymptotically cylindrical (ACyl) Calabi-Yau** 3-fold with asymptotic cross section  $(X, \omega_I, \omega_J, \omega_K)$  and rate v < 0 if there exist

• a compact submanifold  $K_V$  with boundary, a diffeomorphism

$$\Upsilon_V : \mathbf{R}^+ \times S^1 \times X \to V \backslash K_V$$

• a 1-form  $\varrho_1$  on  $\mathbb{R}^+ \times S^1 \times X$  such that  $\Upsilon_V^* \omega = dt \wedge ds + \omega_I + d\varrho_1$  with

$$|\nabla^k \rho_1| = O(e^{vt})$$
 as  $t \to \infty, \forall k \in \mathbb{N} \cup \{0\}$ .

• a 2-form  $\varrho_2$  on  $\mathbb{R}^+ \times S^1 \times X$  such that  $\Upsilon_V^* \Omega = (ds - idt) \wedge (\omega_I + i\omega_K) + d\varrho_2$  with

$$|\nabla^k \varrho_2| = O(e^{\nu t}) \text{ as } t \to \infty, \forall k \in \mathbb{N} \cup \{0\}.$$

Here t and s denote the coordinates on  $\mathbb{R}^+$  and  $S^1$  respectively,  $|\cdot|$  and Levi-Civita connection  $\nabla$  are induced by the product metric on  $\mathbb{R}^+ \times S^1 \times X$ .

*Remark* 2.20. Let  $(V, \omega, \Omega)$  be an ACyl Calabi-Yau 3-fold with asymptotic cross section  $(X, \omega_I, \omega_J, \omega_K)$ . Then

$$(Y := S^1 \times V, \phi := d\theta \wedge \omega + \operatorname{Re} \Omega)$$

is an ACyl G2-manifold with asymptotic cross section

$$(S^1 \times S^1 \times X, ds \wedge d\theta + \omega_K, (d\theta - ids) \wedge (\omega_I + i\omega_I)).$$

Again by the Cheeger-Gromoll splitting theorem [CG71] connected ACyl Calabi-Yau 3-folds whose holonomy is exactly SU(3) can have at most one end.

**Definition 2.21.** A **building block** is a pair (Z, X) in which Z is a non-singular complex projective 3-fold with primitive anti-canonical class  $-K_Z$  in  $H^2(Z)$ , and  $X \in |-K_Z|$  is a smooth K3 surface divisor having trivial holomorphic normal bundle or equivalently there exists a projective morphism  $\mathfrak{f}: Z \to \mathbb{CP}^1$  with  $\mathfrak{f}^{-1}(\infty) = X \in |-K_Z|$  is a smooth K3 surface.

A framing of a building block (Z, X) is a hyperkähler structure  $\boldsymbol{\omega} = (\omega_I, \omega_J, \omega_K)$  on X such that  $\omega_J + i\omega_K$  is of type (2, 0) and  $[\omega_I] \in H^{1,1}(X)$  is the restriction of a Kähler class on Z. A framed building block<sup>4</sup> is such a triple  $(Z, X, \boldsymbol{\omega})$ .

The following theorem summarizes the relationship between simply connected irreducible ACyl Calabi-Yau 3-folds and building blocks.

**Theorem 2.22** (Haskins, Hein, and Nordström [HHN15, Theorem C, D]). Let  $(Z, X, \omega)$  be a framed building block. Then  $V := Z \setminus X$  is simply connected and there is a irreducible ACyl Calabi-Yau structure  $(\omega, \Omega)$  on V with asymptotic cross section  $(X, \omega)$ . Conversely, let  $(V, \omega, \Omega)$  be a simply connected irreducible ACyl Calabi-Yau 3-fold with asymptotic cross section  $(X, \omega)$ . Then there is a complex projective 3-fold Z with  $X \in |-K_Z|$  and  $V = Z \setminus X$  such that  $(Z, X, \omega)$  is a framed building block.

Remark 2.23. Definition 2.19 of ACyl Calabi-Yau 3-fold can be generalized by letting the asymptotic cross section becomes the quotient  $\frac{S^1 \times X}{\langle \tau \rangle}$ , where  $\tau(z,x) = (e^{\frac{2\pi i}{m}}z, \varsigma(x))$  and  $\varsigma$  is a Kähler isometry of X of order m. There is a generalization of Theorem 2.22 with orbifold building blocks. Actually the theorems in [HHN15, Theorem C, D] have been formulated in this general form.

**Definition 2.24.** A **Fano 3-fold** is a smooth projective 3-fold W such that the anti-canonical line bundle  $-K_W$  is ample.

A weak Fano 3-fold is a smooth projective 3-fold W such that the anti-canonical line bundle  $-K_W$  is nef and big, that is,  $-K_W \cdot C \ge 0$  for all compact algebraic curves C in Z and  $-K_W^3 > 0$  respectively.

A weak Fano 3-fold W is called **semi-Fano** if the anticanonical morphism

$$W \longrightarrow R(W, -K_W) := \bigoplus_{l \ge 0} H^0(W, -lK_W)$$

is semi-small, that is, it does not contract any divisor to a point.

**Theorem 2.25** (Corti, Haskins, Nordström, and Pacini [CHNP13, Proposition 4.25]). Let W be a weak Fano 3-fold, and suppose that  $|X_0, X_\infty| \subset |-K_W|$  is an anti-canonical pencil with smooth (reduced) base locus B, and that  $X \in |X_0, X_\infty|$  is a smooth divisor. Let Z be the blow-up of W along the base locus B. Denote the proper transform of X by X again. Then (Z, X) is a building block and  $V := Z \setminus X$  admits an ACyl Calabi-Yau structure.

Remark 2.26. There are precisely 105 deformation families of Fano 3-folds [IP99, Chapter 12] and all but two have a choice of pencils as in Theorem 2.25 [CHNP15, Proposition 3.15]. The deformation families of weak Fano 3-folds are also finite but there exist at least hundreds of thousands. Again, all but a few have a choice of pencils as in Theorem 2.25(see [CHNP13, Section 4, Theorem 4.13]).

<sup>&</sup>lt;sup>4</sup>By Yau's proof of the Calabi conjecture, one can see that each building block admits a framing.

**Definition 2.27.** A holomorphic involution  $\rho$  of a K3 surface X is called **non-symplectic** if  $\rho^*(\alpha) = -\alpha$  for all  $\alpha \in H^{2,0}(X)$ .

**Theorem 2.28** (Kovalev and Lee [KL11, Proposition 5.1]). Let X be a K3 surface with a non-symplectic involution  $\rho$ . Suppose that the fixed point loci of  $\rho$ ,  $C := \operatorname{Fix}_{\rho}(X)$  is nonempty. Denote  $W := \frac{\operatorname{CP}^1 \times X}{\iota \times \rho}$ , where  $\iota : \operatorname{CP}^1 \to \operatorname{CP}^1$  is defined by  $\iota(z) = \frac{1}{z}$ . Let Z be the blow-up of W along the singular set  $\{\pm 1\} \times C$ . Denote the proper transform of the equivalence class of  $\{\infty\} \times X$  by X again. Then (Z,X) is a building block and  $V := Z \setminus X$  admits an ACyl Calabi-Yau structure.

Remark 2.29. There are exactly 75 deformation families of K3 surfaces with non-symplectic involutions and all but one satisfy the nonempty assumption of the fixed point loci in Theorem 2.28 (see [KL11, Proposition 3.2]).

We can glue a pair of ACyl  $G_2$ -manifolds if we can identify the asymptotic cross sections such that the asymptotic  $G_2$ -structures are preserved.

**Definition 2.30.** A pair of ACyl  $G_2$ -manifolds  $(Y_{\pm}, \phi_{\pm})$  with asymptotic cross sections  $(Z_{\pm}, \omega_{\pm}, \Omega_{\pm})$  is said to be a **matching pair** if there exists a diffeomorphism  $f: Z_{+} \to Z_{-}$  such that

$$f^*\omega^- = -\omega^+, \quad f^* \operatorname{Re} \Omega^- = \operatorname{Re} \Omega^+.$$

Let  $(Y_{\pm}, \phi_{\pm})$  be a matching pair of ACyl  $G_2$ -manifolds. Let  $\Upsilon_{\pm} : \mathbb{R}^+ \times Z_{\pm} \to Y_{\pm} \setminus K_{Y_{\pm}}$  be the diffeomorphisms in Definition 2.17. For  $T \ge 1$ , the compact 7-manifold  $Y_T$  is defined by

$$Y_T := Y_{T,+} \cup_F Y_{T,-}$$

where  $Y_{T,\pm} := K_{Y_{\pm}} \cup \Upsilon_{\pm}((0, T+1] \times Z_{\pm})$  and  $F : [T, T+1] \times Z_{+} \to [T, T+1] \times Z_{-}$  is given by

$$F(t,z) = (2T - t + 1, f(z))$$

Here  $Y_T$  is obtained by gluing  $Y_{T,\pm}$  through the identification map F. Let  $\chi: \mathbf{R} \to [0,1]$  (**cut-off function**) be a smooth function such that

(2.31) 
$$\chi(t) = \begin{cases} 0, & t \le 0 \\ 1, & t \ge 1. \end{cases}$$

Define,  $\chi_T(t) := \chi(t-T)$ . The 3-form  $\hat{\phi}_T$  on  $Y_T$  defined by

$$\hat{\phi}_T := \phi_{\pm} - d((\Upsilon_{\pm}^{-1})^* \chi_{T-1} \varrho_{\pm}) \text{ on } Y_{T,\pm},$$

is a closed  $G_2$ -structure. This  $G_2$ -structure might not be torsion free on the gluing region, but for sufficiently large T (as the error is small enough) we may deform it to a torsion free  $G_2$ -structure.

**Theorem 2.32** (Kovalev [Kovo3, Theorem 5.34]). Let  $(Y_{\pm}, \phi_{\pm})$  be a matching pair of ACyl  $G_2$ -manifolds. Then there are constants  $T_0 > 1$ ,  $\delta > 0$  and torsion free  $G_2$  structures  $\phi_T$  on  $Y_T$  for all  $T \ge T_0$  such that

(2.33) 
$$\|\phi_T - \hat{\phi}_T\|_{C^{0,\gamma}} = O(e^{-\delta T})$$




Figure 2: Gluing of a matching pair of ACyl  $G_2$ -manifolds  $Y_{\pm}$ .

**Definition 2.34.** The  $G_2$ -manifold  $(Y_T, \phi_T)$  in Theorem 2.32 is called a **generalized connected** sum  $G_2$ -manifold.

**Definition 2.35.** Let  $(X_{\pm}, \omega_{I,\pm}, \omega_{J,\pm}, \omega_{K,\pm})$  be a pair of hyperkähler 4-manifolds. A diffeomorphism  $\mathfrak{r}: X_+ \to X_-$  is said to be a **hyperkähler rotation** if

$$\mathbf{r}^* \omega_{I,-} = \omega_{I,+}, \quad \mathbf{r}^* \omega_{I,-} = \omega_{I,+} \text{ and } \mathbf{r}^* \omega_{K,-} = -\omega_{K,+}.$$

**Definition 2.36.** Let  $(V_{\pm}, \omega_{\pm}, \Omega_{\pm})$  be a pair of ACyl Calabi-Yau 3-folds with hyperkähler asymptotic cross sections  $(X_{\pm}, \omega_{I,\pm}, \omega_{J,\pm}, \omega_{K,\pm})$  (see Definition 2.19) and  $\mathbf{r}: X_{+} \to X_{-}$  be a hyperkähler rotation. Then  $Y_{\pm} := S^{1} \times V_{\pm}$  is a matching pair of ACyl  $G_{2}$ -manifolds matched by the diffeomorphism

$$f: S^1 \times S^1 \times X_+ \to S^1 \times S^1 \times X_-,$$

defined by  $f(\theta, s, x) = (s, \theta, \mathfrak{r}(x))$ . Therefore by Theorem 2.32 we obtain a family of compact generalized connected sum  $G_2$ -manifolds  $(Y_T, \phi_T)$ . These  $G_2$ -manifolds will be called **twisted connected sum** (TCS)  $G_2$ -manifolds.




Figure 3: TCS construction for a pair of ACyl Calabi-Yau 3-folds  $V_{\pm}$ .

Remark 2.37. If the matching pair of ACyl Calabi-Yau 3-folds  $V_{\pm}$  in Definition 2.36 have holonomy exactly SU(3) (indeed this is true if obtained from building blocks) then the TCS  $G_2$ -manifolds  $Y_T$  have holonomy exactly  $G_2$  [CHNP15, paragraph after Theorem 3.13].

Remark 2.38. Given a pair of building blocks  $(Z_{\pm}, X_{\pm})$  there are no systematic ways to find hyperkähler rotations  $\mathbf{r}: X_{+} \to X_{-}$ . Corti, Haskins, Nordström, and Pacini [CHNP15, Proposition 6.18, Proposition 6.2, Remark 6.19] have rather shown that for any pair of  $N_{\pm}$ -marked full

deformation types of semi-Fano 3-folds (see Definition 2.24) whose  $N_{\pm}$ -polarized K3 lattices have an orthogonal push out (in particular this exists if the sum of the ranks of these lattices is less than equal to 11), there exists always a pair of  $N_{\pm}$ -marked semi-Fano 3-folds  $W_{\pm}$  in that pair of deformation types such that the building blocks  $(Z_{\pm}, X_{\pm})$  arising from them (by blowing up along the base locus of a general anti-canonical pencil) admits a hyperkähler rotation  $\mathfrak{r}: X_{\pm} \to X_{-}$ .

# 3 Moduli space of asymptotically cylindrical associative submanifolds

This section focuses on ACyl associative submanifolds and their moduli spaces. In particular, we prove Theorem 1.9 about moduli space of holomorphic curves and study the Fueter operator on associative cylinders. The definition and examples of ACyl associative submanifolds and the Fredholm theory of asymptotically translation invariant elliptic operators have been also discussed here. The definition of moduli space of ACyl associative submanifolds, and the proof of Theorem 1.12 and Theorem 1.13 are presented in this section. ACyl special Lagrangian 3-folds and ACyl holomorphic curves have been considered at the end because of their importance in the special cases of the gluing theorem in Section 4.

#### 3.1 Associative cylinder

Since ACyl associative submanifolds are asymptotic to associative cylinders, we choose first to study the associative cylinders. In fact, the analysis on them will be useful for the deformation theory of ACyl associative submanifolds.

**Definition 3.1.** Let  $(Z, J, \omega, \Omega)$  be a compact Calabi-Yau 3-fold in which J is the complex structure,  $\omega$  is the Kähler form and  $\Omega$  is the holomorphic volume form. Then

$$(Y_0 := \mathbf{R} \times Z, \phi_0 := dt \wedge \omega + \operatorname{Re} \Omega)$$

is a  $G_2$ -manifold. Let  $\Sigma$  be a 2-dimensional closed submanifold of Z and let C be the cylinder  $\mathbf{R} \times \Sigma$ . The Calabi-Yau metric on Z induces a metric  $g_\Sigma$  on  $\Sigma$  and a metric  $g_C = dt^2 + g_\Sigma$  on C. Moreover, it induces a normal metric  $g_{NC}$  and a normal connection  $\nabla_C^{\perp}$  on the normal bundle NC of the cylinder C. Let  $\pi: \mathbf{R} \times Z \to Z$  be the standard projection. Then  $NC = \pi^*N\Sigma$ , the pullback of the normal bundle  $N\Sigma$  of  $\Sigma$  in Z,  $g_{NC} = \pi^*g_{N\Sigma}$  and  $\nabla_C^{\perp} = \pi^*\nabla_{\Sigma}^{\perp}$ . Let  $\Upsilon_{\Sigma}: V_{\Sigma} \subset N\Sigma \to U_{\Sigma} \subset Z$  be a tubular neighbourhood map of  $\Sigma$  (see Definition 2.13). Define

$$V_C := \pi^* V_{\Sigma} \subset NC, \quad U_C := \mathbf{R} \times U_{\Sigma} \subset Y_0.$$

The diffeomorphism  $\Upsilon_C: V_C \to U_C$  defined by

$$\Upsilon_C((t,\sigma),v) := (t,\Upsilon_{\Sigma}(\sigma,v)), t \in \mathbf{R}, \sigma \in \Sigma, v \in N\Sigma.$$

is a tubular neighbourhood map of the cylinder C. Actually, all the translation invariant tubular neighbourhood maps are of this form.

The cylinder  $C = \mathbf{R} \times \Sigma$  is called an **associative cylinder** in  $Y_0 = \mathbf{R} \times Z$  if it is an associative submanifold, or equivalently,  $\Sigma$  is holomorphic in (Z, J).

#### 3.1.1 Moduli space of holomorphic curves in Calabi-Yau 3-folds

**Definition 3.2.** Let Z be a Calabi-Yau 3-fold and S be the set of all 2-dimensional oriented closed smooth embedded submanifolds. We define the  $C^k$ -topology on the set S by specifying a basis which is a collection of all sets of the form  $\{\Upsilon_{\Sigma}(\Gamma_u): u \in \mathcal{V}_{\Sigma}^k\}$ , where  $\Sigma \in S$ ,  $\Upsilon_{\Sigma}$  is a tubular neighbourhood map of  $\Sigma$  and  $\mathcal{V}_{\Sigma}^k$  is an open set in  $C^{\infty}(V_{\Sigma})$ . Here the topology on  $C^{\infty}(V_{\Sigma})$  is induced by the  $C^k$ -norm on  $C^{\infty}(N_{\Sigma})$ .

The  $C^{\infty}$ -topology on the set S is the inverse limit topology of  $C^k$ -topologies on it, that is, a set is open with  $C^{\infty}$ -topology if and only if it is open in every  $C^k$ -topology.

The **moduli space**  $\mathcal{M}_Z^{\text{hol}}$  of embedded closed holomorphic curves in Z is the subset of all submanifolds  $\Sigma$  in S which are J-holomorphic in Z. The topologies on  $\mathcal{M}_Z^{\text{hol}}$  are the subspace topologies of the above.

For  $\Sigma \in \mathcal{M}_Z^{\text{hol}}$  we denote the complex structure on  $\Sigma$  by j, which is just the restriction of J. Let  $\Upsilon_\Sigma : V_\Sigma \subset N\Sigma \to U_\Sigma \subset Z$  be a tubular neighbourhood map of  $\Sigma$ . For  $u \in C^\infty(V_\Sigma)$  we denote by  $\Sigma_u$  the submanifold  $\Upsilon_\Sigma(\Gamma_u)$ . Then  $\Sigma_u$  is J-holomorphic if and only if u satisfies the following non-linear Cauchy-Riemann equation

$$0 = \bar{\partial}_J u := \frac{1}{2} (du + \Upsilon_{\Sigma}^* J(u) \circ du \circ j) \in C^{\infty}(\overline{\operatorname{Hom}}_{\mathbb{C}}(T\Sigma, u^* T V_{\Sigma})).$$

The linearization of the nonlinear map  $\bar{\partial}_J: C^{\infty}(V_{\Sigma}) \to C^{\infty}(\overline{\operatorname{Hom}}_{\mathbb{C}}(T\Sigma, u^*TV_{\Sigma}))$  at the zero section, is described in [MS12, Proposition 3.1.1]. This is the linear map  $\mathfrak{d}_{\Sigma,J}: C^{\infty}(N\Sigma) \to C^{\infty}(\overline{\operatorname{Hom}}_{\mathbb{C}}(T\Sigma, TV_{\Sigma|\Sigma}))$ , defined by

$$\mathfrak{d}_{\Sigma,J}\xi:=\frac{1}{2}(\nabla_{\Sigma}\xi+J\circ(\nabla_{\Sigma}\xi)\circ j),\ \xi\in C^{\infty}(N\Sigma).$$

The tangential component of  $\mathfrak{d}_{\Sigma,J}$  can be discarded for the deformation theory. The following normal component actually controls the deformation theory.

**Definition 3.3.** For  $\Sigma \in \mathcal{M}_Z^{\mathrm{hol}}$  the **normal Cauchy-Riemann operator**  $\bar{\partial}_{\nabla,J}^N : C^{\infty}(N\Sigma) \to C^{\infty}(\overline{\mathrm{Hom}}_{\mathbb{C}}(T\Sigma,N\Sigma)) \cong \Omega^{0,1}(\Sigma,N\Sigma)$  is defined by

$$\bar{\partial}^N_{\nabla,J}\xi:=\frac{1}{2}(\nabla^\perp_\Sigma\xi+J\circ(\nabla^\perp_\Sigma\xi)\circ j).$$

We now study the moduli space of holomorphic curves in a set up which will be useful later.

**Definition 3.4.** The *J*-anti-linear multiplication map,  $\times_Z : TZ \times TZ \to TZ$  is defined by the orthogonal projection of the cross product in  $Y_0$  onto TZ or equivalently, for all vector fields x, y, w on Z

$$q(x \times_Z y, w) = \text{Re } \Omega(x, y, w).$$

Remark 3.5. Let  $Y_0$  be as in Definition 3.1. Let  $\Sigma$  be an oriented smooth surface in Z. The following are equivalent: (i)  $\Sigma$  is J-holomorphic in Z, (ii)  $C = \mathbb{R} \times \Sigma$  is associative in  $Y_0$ , (iii) for all  $x, y \in T\Sigma$  and  $w \in TZ$ ,  $\mathbb{R} \in \Omega(x, y, w) = 0$ , (iv) for all  $x, y \in T\Sigma$ ,  $x \times_Z y = 0$ , (v) for all  $x \in T\Sigma$  and  $x \in T\Sigma$  and  $x \in T\Sigma$  and  $x \in T\Sigma$ .

**Definition 3.6.** Let Σ be an embedded closed *J*-holomorphic curve in *Z*. The map  $\gamma_{\Sigma} : T\Sigma \to \overline{\operatorname{End}}_{\mathbb{C}}(N\Sigma)$  given by

$$\gamma_{\Sigma}(f_{\Sigma})(v_{\Sigma}) := f_{\Sigma} \times v_{\Sigma}, \quad \forall \ v_{\Sigma} \in N\Sigma, \ f_{\Sigma} \in T\Sigma,$$

is a skew symmetric J-anti-linear Clifford multiplication. Moreover the normal bundle  $N\Sigma$  together with the metric  $g_{N\Sigma}$ , Clifford multiplication  $\gamma_{\Sigma}$  and the metric connection  $\nabla^{\perp} := \nabla^{\perp}_{\Sigma}$  is a Dirac bundle, that is,  $\nabla^{\perp}\gamma_{\Sigma} = 0$ . The Dirac operator is given by

(3.7) 
$$\mathbf{D}_{\Sigma} := \gamma_{\Sigma} \circ \nabla^{\perp} = \sum_{i=1}^{2} f_{i} \times \nabla_{f_{i}}^{\perp}$$

where  $\{f_i\}$  is a local orthonormal oriented frame on  $\Sigma$ . Note that  $\mathbf{D}_{\Sigma}$  is J-anti-linear. The map  $\gamma_{\Sigma}$  induces a J-anti-linear isomorphism  $\gamma_{\Sigma} : \overline{\mathrm{Hom}}_{\mathbb{C}}(T\Sigma, N\Sigma) \to N\Sigma$  which is defined by

$$\boldsymbol{\gamma}_{\Sigma}(f_{\Sigma}^{*}\otimes v_{\Sigma})=\boldsymbol{\gamma}_{\Sigma}(f_{\Sigma})(v_{\Sigma}).$$

Remark 3.8.  $\gamma_{\Sigma}$  satisfies the following commutative diagram:

$$C^{\infty}(N\Sigma) \xrightarrow{\bar{\partial}_{\nabla J}^{N}} C^{\infty}(\overline{\operatorname{Hom}}_{\mathbb{C}}(T\Sigma, N\Sigma))$$

$$\downarrow^{\gamma_{\Sigma}}$$

$$C^{\infty}(N\Sigma).$$

**Definition 3.9.** Let  $\Sigma \in S$  and let  $\Upsilon_{\Sigma} : V_{\Sigma} \subset N\Sigma \to U_{\Sigma} \subset Z$  be a tubular neighbourhood map of  $\Sigma$ . We define  $\mathcal{F} : C^{\infty}(V_{\Sigma}) \to C^{\infty}(N\Sigma)$  as follows. If  $u \in C^{\infty}(V_{\Sigma})$  and  $v \in C^{\infty}(N\Sigma)$ ,

$$\langle \mathfrak{F}(u), v \rangle_{L^2} := \int_{\Gamma_{\cdot}} \iota_v(\Upsilon_{\Sigma}^* \operatorname{Re} \Omega).$$

The notation v in the integrand is the extension vector field of  $v \in C^{\infty}(N\Sigma)$  in the tubular neighbourhood as in Notation 2.16.

**Proposition 3.10.** For  $u \in C^{\infty}(V_{\Sigma})$ , we have  $\mathcal{F}(u) = 0$  if and only if  $\Sigma_u := \Upsilon_{\Sigma}(\Gamma_u)$  is J-holomorphic. If  $\Sigma \in \mathcal{M}_Z^{hol}$ , that is, J-holomorphic then the linearization of  $\mathcal{F}$  at zero,  $d\mathcal{F}_0 : C^{\infty}(N\Sigma) \to C^{\infty}(N\Sigma)$  is given by

$$d\mathcal{F}_0 = J\mathbf{D}_{\Sigma}$$
.

This is a first order self adjoint elliptic operator and thus the index of  $d\mathcal{F}_0$  is 0.

*Proof.* The first part follows from the Remark 3.5. Let  $\{f_1, f_2 = Jf_1\}$  be a local oriented orthonormal frame for  $T\Sigma$ . Then for  $u, v \in C^{\infty}(N\Sigma)$ ,

$$\frac{d}{dt}\big|_{t=0}\langle \mathfrak{F}(tu),v\rangle_{L^2} = \frac{d}{dt}\big|_{t=0} \int_{\Gamma_{tu}} \iota_v(\Upsilon_{\Sigma}^*\operatorname{Re}\Omega) = \int_{\Sigma} \mathcal{L}_u \iota_v(\Upsilon_{\Sigma}^*\operatorname{Re}\Omega).$$

This is same as  $\int_{\Sigma} \iota_v \mathcal{L}_u(\Upsilon_{\Sigma}^* \operatorname{Re} \Omega) + \iota_{[u,v]}(\Upsilon_{\Sigma}^* \operatorname{Re} \Omega)$ . Following Notation 2.16 we have [u,v] = 0 and therefore this is equal to

$$\int_{\Sigma} \iota_{v}(\Upsilon_{\Sigma}^{*} \operatorname{Re} \Omega)(\nabla_{f_{1}} u, f_{2}) + \iota_{v}(\Upsilon_{\Sigma}^{*} \operatorname{Re} \Omega)(f_{1}, \nabla_{f_{2}} u) + \iota_{\nabla_{v} u}(\Upsilon_{\Sigma}^{*} \operatorname{Re} \Omega) + \iota_{v} \nabla_{u}(\Upsilon_{\Sigma}^{*} \operatorname{Re} \Omega).$$

As,  $\nabla \Omega = 0$  and  $\iota_v$  Re  $\Omega_{|\Sigma} = 0$ , the whole expression above becomes

$$\int_{\Sigma} -\langle f_2 \times \nabla_{f_1}^{\perp} u, v \rangle + \langle f_1 \times \nabla_{f_2}^{\perp} u, v \rangle = \int_{\Sigma} \langle J(f_1 \times \nabla_{f_1}^{\perp} u), v \rangle + \langle J(f_2 \times \nabla_{f_2}^{\perp} u), v \rangle.$$

The right hand side is  $\int_{\Sigma} \langle J\mathbf{D}_{\Sigma}u, v \rangle$ , which proves the second part of the proposition. Since  $\mathbf{D}_{\Sigma}$  is a *J*-anti-linear and self adjoint Dirac operator, we obtain the last part of the proposition.

**Proof of Theorem 1.9.** We observe the fact  $\gamma_{\Sigma} \circ \bar{\partial}_{\nabla,J}^{N} = \mathbf{D}_{\Sigma}$ . in Remark 3.8. Extending the nonlinear map  $\mathcal{F}$  to Hölder spaces we get a smooth map

$$\mathcal{F}: C^{2,\gamma}(V_{\Sigma}) \to C^{1,\gamma}(N\Sigma).$$

Proposition 3.10 implies that the linearization of  $\mathcal{F}$  at zero is an elliptic operator and hence Fredholm. By implicit function theorem applied to  $\mathcal{F}$  we obtain the map  $ob_{\Sigma}$  as stated in the theorem (see [DK90, Proposition 4.2.19]). We need to only prove that, if  $u \in C^{2,\gamma}(V_{\Sigma})$  with  $\mathcal{F}(u) = 0$ , then  $u \in C^{\infty}(V_{\Sigma})$ . To prove this, we observe

$$0 = \mathbf{D}_{\Sigma}(\mathcal{F}(u)) = a(u, \nabla_{\Sigma}^{\perp} u)(\nabla_{\Sigma}^{\perp})^{2} u + b(u, \nabla_{\Sigma}^{\perp} u)$$

Since  $a(u, \nabla_{\Sigma}^{\perp} u) \in C^{1,\gamma}$  and  $b(u, \nabla_{\Sigma}^{\perp} u) \in C^{1,\gamma}$ , by Schauder elliptic regularity (see [Joyo7, Theorem 1.4.2]) we obtain  $u \in C^{3,\gamma}$ . By repeating this argument we get higher regularity, which completes the proof of the theorem.

**Example 3.11.** We consider a special case where the Calabi-Yau 3-fold  $(Z, J, \omega, \Omega)$  is  $T^2 \times X$  with  $(X, \omega_I, \omega_J, \omega_K)$  a hyperkähler 4-manifold and

$$\omega = ds \wedge d\theta + \omega_K$$
,  $\Omega = (d\theta - ids) \wedge (\omega_I + i\omega_I)$ ,

where  $(\theta, s)$  denotes the coordinate on  $T^2$ . For each point  $x \in X$ , consider the J-holomorphic tori  $T_x^2 := T^2 \times \{x\} \subset Z$ . Then

$$NT_x^2 = T^2 \times T_x X$$
,  $(K(x)\mathbf{D}_{T_x^2})^2 = \mathbf{D}_{T_x^2}^2 = \Delta_{T_x^2}$  and  $\ker \mathbf{D}_{T_x^2} \cong T_x X$ .

Therefore  $T_x^2$  is Morse-Bott and the moduli space  $\mathcal{M}_Z^{\text{hol}}$  near  $T_x^2$  is  $\{T_y^2:y\in U_x\}$  for some open neighbourhood  $U_x$  of x in X.

**Example 3.12.** We again consider the Calabi-Yau 3-fold  $T^2 \times X$  as in Example 3.11. For each point  $z \in T^2$  and K-holomorphic curve  $\Sigma$  in X consider the holomorphic curve  $\Sigma_z := \{z\} \times \Sigma$  in Z. Then  $\Sigma$  is a special Lagrangian in  $(X, \omega_I, \omega_I + i\omega_K)$  with phase i. The moduli space of special

Lagrangians  $\mathcal{M}_{SL}$  near  $\Sigma$  in X is homeomorphic to an open neighbourhood of 0 in  $\mathcal{H}^1(\Sigma)$ , space of harmonic 1-forms on  $\Sigma$ . We have an isomorphism

$$N\Sigma_z = T_z T^2 \oplus N_X \Sigma \cong \mathbf{R}^2 \oplus T^* \Sigma$$

where the last isomorphism  $N_X\Sigma \to T^*\Sigma$  is given by  $\{v \to \iota_v \omega_I\}$ . A simple computation shows under this isomorphism  $D_{\Sigma_z}$ , J,  $JD_{\Sigma_z}$  are congruent to

$$\hat{D}, \hat{J}, \hat{J}\hat{D}: \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^1(\Sigma, \mathbf{R}) \to \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^0(\Sigma, \mathbf{R}) \oplus \Omega^1(\Sigma, \mathbf{R})$$

respectively, where

$$\hat{D} := \begin{bmatrix} 0 & 0 & d^* \\ 0 & 0 & *d \\ d & -*d & 0 \end{bmatrix}, \quad \hat{J} := \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -* \end{bmatrix}, \quad \hat{J}\hat{D} = \begin{bmatrix} 0 & 0 & *d \\ 0 & 0 & -d^* \\ -*d & -d & 0 \end{bmatrix}.$$

Thus

$$(\hat{J}\hat{D})^2=\hat{D}^2=\Delta,\ \ker \mathbf{D}_{\Sigma_z}\cong\mathcal{H}^0(\Sigma)\oplus\mathcal{H}^0(\Sigma)\oplus\mathcal{H}^1(\Sigma).$$

Hence by [McL98, Theorem 3.6]  $\Sigma_z$  is Morse-Bott and the moduli space  $\mathcal{M}_Z^{\text{hol}}$  near  $\Sigma_z$  is  $\{\Sigma_{z'}':=\{z'\}\times\Sigma':(z',\Sigma')\in U_z\times U_\Sigma\}$ , for some open neighbourhoods  $U_z$  of z in  $T^2$  and  $U_\Sigma$  of  $\Sigma$  in  $\mathcal{M}_{SL}$ 

#### 3.1.2 Fueter operator on associative cylinder

We will see later that the deformation operator controlling the deformations of an ACyl associative submanifold is an asymptotically translation invariant self-adjoint elliptic operator. It will be asymptotic to a translation invariant elliptic operator, namely the Fueter operator on the asymptotic associative cylinder. The Fredholm theory of asymptotically translation invariant elliptic operators has been studied by Lockhart and McOwen [LM85]. For this we need to study indicial roots and homogeneous kernels of this Fueter operator.

Let  $C = \mathbf{R} \times \Sigma$  be a connected associative cylinder in a  $G_2$ -manifold  $Y_0 = \mathbf{R} \times Z$ . Recall that  $\Sigma$  is J-holomorphic in Z if and only if C is associative in  $Y_0$ . Denote the coordinate on  $\mathbf{R}$  by t.

**Definition 3.13.** The **Fueter operator** on the associative cylinder *C* is denoted by  $\mathbf{D}_C : C^{\infty}(NC) \to C^{\infty}(NC)$  and defined by

$$\mathbf{D}_C := \sum_{i=1}^3 e_i \times \nabla_{C,e_i}^{\perp},$$

where  $\{e_i\}$  is a local oriented orthonormal frame on C and  $\nabla_C^{\perp}$  is the normal connection induced by the Levi-Civita connection on  $Y_0$ .

**Definition 3.14.** For  $\lambda \in \mathbb{R}$ , we define the **homogeneous kernel** of rate  $\lambda$  by

$$V_{\lambda} := \{ e^{\lambda t} v_{\Sigma} : v_{\Sigma} \in C^{\infty}(N\Sigma), \ \mathbf{D}_{C}(e^{\lambda t} v_{\Sigma}) = 0 \}, \ d_{\lambda} := \dim V_{\lambda}.$$

The set of **indicial roots** is defined by

$$\mathfrak{D} := \{ \lambda \in \mathbf{R} : d_{\lambda} \neq 0 \}.$$

**Proposition 3.15.** Let  $v_{\Sigma} \in C^{\infty}(N\Sigma)$ . Then the following holds

(i) 
$$\mathbf{D}_C = J\partial_t + \mathbf{D}_{\Sigma}$$
.

(ii) 
$$\mathbf{D}_C(e^{\lambda t}v_{\Sigma}) = e^{\lambda t}(\mathbf{D}_{\Sigma}v_{\Sigma} + \lambda Jv_{\Sigma}).$$

(iii) 
$$\mathbf{D}_C(e^{\lambda t}t^j\nu_{\Sigma}) = e^{\lambda t}t^j(\lambda J\nu_{\Sigma} + \mathbf{D}_{\Sigma}\nu_{\Sigma}) + je^{\lambda t}t^{j-1}J\nu_{\Sigma}.$$

(iv) 
$$\mathbf{D}_{\Sigma}(J\nu_{\Sigma}) = -J\mathbf{D}_{\Sigma}\nu_{\Sigma}$$
.

(v) 
$$V_{\lambda} = \{e^{\lambda t} v_{\Sigma} : \mathbf{D}_{\Sigma} v_{\Sigma} = -\lambda J v_{\Sigma}\} = \{e^{\lambda t} v_{\Sigma} : (J\mathbf{D}_{\Sigma}) v_{\Sigma} = \lambda v_{\Sigma}\}$$

(vi) 
$$JV_{\lambda} = V_{-\lambda}$$
 and  $d_{\lambda} = d_{-\lambda}$  for all  $\lambda \in \mathbb{R}$ .

*Proof.* Observe that  $\nabla_C^{\perp} = dt \otimes \partial_t + \nabla_{\Sigma}^{\perp}$ . Now for a local oriented orthonormal frame  $\{f_i\}$  on  $\Sigma$  and  $v \in C^{\infty}(NC) = C^{\infty}(\mathbb{R}, N\Sigma)$  we have

$$\mathbf{D}_{C}v = J\partial_{t}v + \sum_{i=1}^{2} f_{i} \times \nabla_{\Sigma, f_{i}}^{\perp}v = J\partial_{t}v + \mathbf{D}_{\Sigma}v$$

This proves (i). And (ii) and (iii) follows from (i), indeed

$$\mathbf{D}_{C}(e^{\lambda t}t^{j}\nu_{\Sigma}) = e^{\lambda t}t^{j}(\lambda J\nu_{\Sigma} + \mathbf{D}_{\Sigma}\nu_{\Sigma}) + je^{\lambda t}t^{j-1}J\nu_{\Sigma}.$$

To see (iv), we compute

$$\mathbf{D}_{\Sigma}(J\nu_{\Sigma}) = \sum_{i=1}^{2} f_{i} \times \left(\nabla_{\Sigma,f_{i}}^{\perp}(\partial_{t} \times \nu_{\Sigma})\right) = \sum_{i=1}^{2} f_{i} \times (\partial_{t} \times \nabla_{\Sigma,f_{i}}^{\perp} \nu_{\Sigma}) + \sum_{i=1}^{2} f_{i} \times (\partial_{t} f_{i} \times \nu_{\Sigma})$$
$$= -\sum_{i=1}^{2} \partial_{t} \times (f_{i} \times \nabla_{\Sigma,f_{i}}^{\perp} \nu_{\Sigma}) = -J\mathbf{D}_{\Sigma}(\nu_{\Sigma})$$

Finally, (v) follows from (ii), and (vi) follows from (iv).

**Proposition 3.16.** Let  $m \in \mathbb{N}_0$  and  $v(t, \sigma) = \sum_{j=0}^m e^{\lambda t} t^j v_{\Sigma, j} \in C^{\infty}(NC)$  with  $v_{\Sigma, j} \in C^{\infty}(N\Sigma)$  and  $v_{\Sigma, m} \neq 0$ . If  $\mathbf{D}_C v(t, \sigma) = 0$ , then m = 0 and thus  $v(t, \sigma) = e^{\lambda t} v_{\Sigma, 0}$ .

*Proof.* If  $\mathbf{D}_C v(t, \sigma) = 0$  then by Proposition 3.15(iii) and comparing the coefficients of  $e^{\lambda t} t^{j-1}$ ,  $j \ge 1$  we can see that

$$\mathbf{D}_{\Sigma} v_{\Sigma,m} + \lambda J v_{\Sigma,m} = 0$$
 and  $j J v_{\Sigma,j} + \mathbf{D}_{\Sigma} v_{\Sigma,j-1} + \lambda J v_{\Sigma,j-1} = 0$ .

Thus we have

$$\begin{split} m\|v_{\Sigma,m}\|_{L^{2}(\Sigma)}^{2} &= m\|Jv_{\Sigma,m}\|_{L^{2}(\Sigma)}^{2} = -\langle Jv_{\Sigma,m}, \mathbf{D}_{\Sigma}v_{\Sigma,m-1} + \lambda Jv_{\Sigma,m-1}\rangle_{L^{2}(\Sigma)} \\ &= -\langle \mathbf{D}_{\Sigma}Jv_{\Sigma,m} + \lambda v_{\Sigma,m}, v_{\Sigma,m-1}\rangle_{L^{2}(\Sigma)} \\ &= \langle J(\mathbf{D}_{\Sigma}v_{\Sigma,m} + \lambda Jv_{\Sigma,m}), v_{\Sigma,m-1}\rangle_{L^{2}(\Sigma)} = 0. \end{split}$$

The last inequality uses the fact from Proposition 3.15(iv). The conclusion now follows by backwards induction starting with j = m.

Remark 3.17. Although it follows from [LM85, Equation 1.11] that a general element in the homogeneous kernel is of the form  $\sum_{j=0}^{m} e^{\lambda t} t^{j} v_{\Sigma,j}$ , Proposition 3.16 implies that it must be of the form we have defined in Definition 3.14. We also see that the indicial roots are in a canonical one to one correspondence with the eigenvalues of a self adjoint elliptic operator  $JD_{\Sigma}$ . In particular they are countable and discrete subset of R.

**Example 3.18.** Consider the special case where Z is  $T^2 \times X$  as in Example 3.11. For the holomorphic tori  $T_x^2$  we have seen that

$$(K(x)\mathbf{D}_{T_x^2})^2 = \mathbf{D}_{T_x^2}^2 = \Delta_{T^2}^0 \oplus \Delta_{T^2}^0 \oplus \Delta_{T^2}^0 \oplus \Delta_{T^2}^0.$$

Therefore the set of indicial roots is  $\mathcal{D} = \{\pm \sqrt{\lambda} : \lambda \text{ is an eigenvalue of } \Delta_{T^2}^0\}$ . If  $\lambda > 0$ , the direct sum of homogeneous kernels,

$$V_{\sqrt{\lambda}} \oplus V_{-\sqrt{\lambda}} = E_{\Delta^0_{T^2}}^{\lambda} \oplus E_{\Delta^0_{T^2}}^{\lambda} \oplus E_{\Delta^0_{T^2}}^{\lambda} \oplus E_{\Delta^0_{T^2}}^{\lambda},$$

where  $E^{\lambda}_{\Delta^0_{T^2}}$  is the  $\lambda$ -eigenspace of  $\Delta^0_{T^2}$ . Thus  $d_{\sqrt{\lambda}}=d_{-\sqrt{\lambda}}=2\dim E^{\lambda}_{\Delta^0_{T^2}}$ . It follows from [Maro2, Section 6.3.4, p. 132] that  $\mathcal{D}\cap[-\sqrt{2},\sqrt{2}]=\{0,\pm\sqrt{2}\}$  and  $d_{\sqrt{2}}=12$ . If  $\lambda=0$ ,  $V_0\cong T_xX$ . Thus  $d_0=4$ .

**Example 3.19.** Consider the special case where Z is  $T^2 \times X$  as in Example 3.12. For the holomorphic curve  $\Sigma_z$  we have seen that

$$(\hat{J}\hat{D})^2 = \hat{D}^2 = \Delta_{\Sigma}^0 \oplus \Delta_{\Sigma}^0 \oplus \Delta_{\Sigma}^1.$$

Therefore the set of indicial roots is  $\mathcal{D} = \{\pm \sqrt{\lambda} : \lambda \text{ is an eigenvalue of } \Delta_{\Sigma}^0 \oplus \Delta_{\Sigma}^1 \}$ . If  $\lambda > 0$ , the direct sum of homogeneous kernels,

$$V_{\sqrt{\lambda}} \oplus V_{-\sqrt{\lambda}} = E_{\Delta^0_{\Sigma}}^{\lambda} \oplus E_{\Delta^0_{\Sigma}}^{\lambda} \oplus E_{\Delta^1_{\Sigma}}^{\lambda},$$

where  $E_{\Delta_{\Sigma}^{i}}^{\lambda}$  is the  $\lambda$ -eigenspace of  $\Delta_{\Sigma}^{i}$ . If  $\lambda = 0$ ,

$$V_0 \cong H^0(\Sigma, \mathbf{R}) \oplus H^0(\Sigma, \mathbf{R}) \oplus H^1(\Sigma, \mathbf{R}).$$

Thus  $d_0 = 2b^0(\Sigma) + b^1(\Sigma)$ .

#### 3.2 Asymptotically cylindrical (ACyl) associative submanifolds

Let  $(Y, \phi)$  be an ACyl  $G_2$ -manifold with asymptotic cross section  $(Z, \omega, \Omega)$  and rate  $\nu < 0$  as in Definition 2.17.

**Definition 3.20.** Let  $C = \mathbb{R} \times \Sigma$  be a cylinder (possibly disconnected) in  $Y_0 = \mathbb{R} \times Z$ . Let  $\Sigma = \coprod_{i=1}^m \Sigma_i$  be the decomposition of  $\Sigma$  into connected components, and subsequently  $C = \coprod_{i=1}^m C_i$ , where  $C_i = \mathbb{R} \times \Sigma_i$ . Let  $\Upsilon_C : V_C \to U_C \subset \mathbb{R} \times Z$  be a translation invariant tubular neighbourhood map of C as in Definition 3.1.

A smooth three dimensional oriented submanifold P of Y is said to be **asymptotically** cylindrical (ACyl) submanifold with asymptotic cross section  $\Sigma$  and rate  $\mu = (\mu_1, \mu_2, ..., \mu_m)$  with  $\nu \leq \mu_i < 0$  for all i = 1, 2..., m if there exist

- a constant  $T_0 > 0$ , a compact submanifold with boundary  $K_P$  of P,
- a smooth embedding  $\Psi_P: (T_0, \infty) \times \Sigma \to U_C \subset \mathbb{R}^+ \times Z$  such that  $\Upsilon \circ \Psi_P: (T_0, \infty) \times \Sigma \to Y$  is a diffeomorphism onto  $P \setminus K_P$ ,
- on  $(T_0, \infty) \times \Sigma$ ,  $\Psi_P = \Upsilon_C \circ \alpha$  for some smooth section  $\alpha$  of the normal bundle NC of C which lies in  $V_C$ , and

(3.21) 
$$|(\nabla_{C_i}^{\perp})^k \alpha| = O(e^{\mu_i t}) \text{ as } t \to \infty, i = 1, 2..., m, \ \forall k \in \mathbb{N} \cup \{0\}.$$

Here  $\nabla_C^{\perp}$  is the normal connection on NC induced from the Levi-Civita connection on  $\mathbb{R}^+ \times Z$  and  $|\cdot|$  is taken with respect to the normal metric on NC and cylindrical metric on C.

An asymptotically cylindrical submanifold is said to be **end cylindrical (ECyl)** if  $\alpha = 0$  in (3.21).

P is said to be an **asymptotically cylindrical (ACyl) associative submanifold** with asymptotic cross section  $\Sigma$  if it is associative and asymptotically cylindrical submanifold as above. lacktriangle

Remark 3.22. We make a few remarks about the above definition.

- (1) An ACyl submanifold with rate  $\mu$  is also an ACyl submanifold with any rate  $\mu' \geqslant \mu$ . Moreover, it is also an ACyl Riemannian manifold with rate  $\mu$  with the metric induced by  $g_{\phi}$  corresponding to the  $G_2$ -structure  $\phi$ .
- (2) If P is an ACyl associative submanifold, then the asymptotic cylinder C is an associative cylinder in  $Y_0$ . Indeed, the associator on  $C_i$

$$|[\cdot,\cdot,\cdot]_{|C_i}| = O(e^{\mu_i T})$$
, as  $T \to \infty$ .

But  $|[\cdot, \cdot, \cdot]|_{C_i}$  is translation invariant and therefore  $[\cdot, \cdot, \cdot]_{C_i} = 0$ .

**Example 3.23**. Let  $(V, \omega, \Omega)$  be an ACyl Calabi-Yau 3-fold with asymptotic cross section  $(X, \omega_I, \omega_I, \omega_K)$  and let  $Y := S^1 \times V$  be the ACyl  $G_2$ -manifold as in Remark 2.20.

- (1) Let C be an ACyl holomorphic curve in V with asymptotic cross section  $\coprod_{j=1}^m \left(S^1 \times \{x_j\}\right) \subset S^1 \times X$ . Then  $P := S^1 \times C$  is an ACyl associative submanifold in Y with asymptotic cross section  $\Sigma := \coprod_{j=1}^m \left(T^2 \times \{x_j\}\right)$ .
- (2) Let (Z,X) be a building block and let  $V := Z \setminus X$  be the corresponding ACyl Calabi-Yau 3-fold. Let C be a closed holomorphic curve in Z intersecting X transversally at  $\bar{x} := \{x_1, \ldots, x_m\}$ . Then  $C \setminus X$  is an ACyl holomorphic curve in V with asymptotic cross section  $\Sigma := \coprod_{i=1}^m (T^2 \times \{x_j\})$  and rate -1. This has been proved in Proposition 3.72.
- (3) Let W be a weak-Fano 3-fold and let B be the base locus of an anti-canonical pencil. Let  $\pi: Z \to W$  be the blow up of W along B as given in Theorem 2.25. Let C be a holomorphic curve in W avoiding B and satisfying  $-K_W \cdot C > 0$ . Then for general  $X \in |-K_W|$  in the pencil, the proper transform of C in Z is an example of a closed holomorphic curve in part (2), where the building block is (Z, X). Moreover, for each  $b \in B$ ,  $\ell_b := \pi^{-1}(b)$  is a rational curve in Z which intersects X transversally at one point and

$$N\ell_b \cong \mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-1).$$

(4) Let X be a K3 surface with a non-symplectic involution  $\rho$  and let  $\pi: Z \to W$  be the blow-up given in Theorem 2.28 so that (Z,X) is a building block. Denote by  $\operatorname{Fix}_{\rho}(X)$  the fixed point loci of  $\rho$ . Let  $x \notin \operatorname{Fix}_{\rho}(X)$ . Then the proper transform of  $[\mathbf{P}^1 \times \{x\}] \subset W$  in Z, call it  $\ell_x$  is an example of C in part (2). In this case,

$$N\ell_{x} \cong \mathcal{O}_{\mathbf{P}^{1}} \oplus \mathcal{O}_{\mathbf{P}^{1}}.$$

Let  $y \in \text{Fix}_{\rho}(X)$ . Set,  $\ell_y^{\pm} := \pi^{-1}(\pm 1, y)$ . Then  $\ell_y^{\pm}$  are lines in the exceptional divisor of Z and are examples of C in part (2). In this case,

$$N\ell_u^{\pm} \cong \mathcal{O}_{\mathbf{P}^1} \oplus \mathcal{O}_{\mathbf{P}^1}(-1).$$

**Example 3.24.** Let  $(Y, \phi)$  be an ACyl  $G_2$ -manifold with asymptotic cross section  $(Z, \omega, \Omega)$ . A nontrivial  $G_2$ -involution  $\sigma: Y \to Y$  (i.e.  $\sigma^2 = \mathbf{1}, \sigma^* \phi = \phi$ ) is always an ACyl  $G_2$ -involution [Noro8b, Proposition 2.3.7], that is, there exist constants T > 0,  $\varepsilon > 0$ , a non trivial SU(3)-involution  $\tau: Z \to Z$  (i.e.  $\tau^2 = \mathbf{1}, \tau^* \omega = \omega$  and  $\tau^* \Omega = \Omega$ ) and a vector field N on  $\mathbf{R}^+ \times Z$  such that over  $(T, \infty) \times Z$ ,  $\sigma \circ \Upsilon = \Upsilon \circ \exp(N \circ (\mathbf{1} \times \tau))$  with

$$|\nabla^k N| = O(e^{-\varepsilon t})$$
 as  $t \to \infty, \forall k \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$ .

Here t denotes the coordinate on  $\mathbb{R}^+$ ,  $|\cdot|$  and Levi-Civita connection  $\nabla$  are induced by the product metric on  $\mathbb{R}^+ \times Z$ . Let P be a connected non-compact component of  $\mathrm{Fix}_\sigma(Y)$ . Then it is an ACyl associative [Joyo7, Proposition 12.3.7] with cross section  $\Sigma := \mathrm{Fix}_\tau(Z)$ .

**Example 3.25**. Let  $(V, \omega, \Omega)$  be an ACyl Calabi-Yau 3-fold with asymptotic cross section  $(X, \omega_I, \omega_I, \omega_K)$  and  $Y := S^1 \times V$  be the ACyl  $G_2$ -manifold as in Remark 2.20.

- (1) Let L be an ACyl Special Lagrangian 3-fold in V with asymptotic cross section  $\{e^{is}\} \times \Sigma$ , where  $\Sigma$  is a K-holomorphic curve in X. Then for each  $\theta \in [0, 2\pi)$ ,  $\{e^{i\theta}\} \times L$  is an ACyl associative in  $Y = S^1 \times V$  with cross section  $\{e^{i\theta}\} \times \{e^{is}\} \times \Sigma$ . By abusing notation we will denote them by L and  $\Sigma$  respectively.
- (2) Let  $\sigma_V: V \to V$  be an anti-holomorphic involutive isometry (i.e.  $\sigma_V^2 = 1, \sigma_V^* \omega = -\omega, \sigma_V^* \Omega = \bar{\Omega}$ ). Let L be a connected non-compact component of the fixed point loci  $\operatorname{Fix}_{\sigma_V}(Z)$ . Then it is an ACyl special Lagrangian 3-fold (see [Tal17, Proposition 3.11]). Thus we can apply part (1) to it.
- (3) Let  $(Z, J, X, \omega)$  be a framed building block and let  $\sigma_Z : Z \to Z$  be an anti-holomorphic involution (i.e.  $\sigma_Z^2 = 1$ ,  $\sigma_Z^* J = -J$ ) such that it is an anti-holomorphic involutive isometry on  $(X, \omega)$ . Then  $V := Z \setminus X$  admits an ACyl Calabi-Yau structure  $(\omega, \Omega)$  such that  $\sigma_Z$  is an anti-holomorphic involutive isometry on V [Kov13, Proposition 5.2]. If a building block (Z, X) admits an anti-holomorphic involution  $\sigma_Z$  which takes X to X, then there is a framing  $\omega$  such that it is an anti-holomorphic involutive isometry on  $(X, \omega)$ . Thus we can apply part (2) to it.
- (4) Building blocks obtained from weak Fano 3-folds with anti-holomorphic involution which acts on the anti-canonical divisor and base locus, admit a lifting of the anti-holomorphic involution [Kov13, pg. 19] so that part (3) applies.

(5) Building blocks obtained from *K*3 surfaces with commuting non-symplectic involution and anti-holomorphic involution admit a lifting of the anti-holomorphic involution. This type of *K*3 surfaces have been studied in [NSo5; NSo7].

**Example 3.26.** Let  $(Y, \phi)$  be an ACyl  $G_2$ -manifold with asymptotic cross section  $(Z, \omega, \Omega)$ . Let  $p_Y: Y \to M$  be an ACyl coassociative fibration over an ACyl 3-manifold M with asymptotic cross section  $\Sigma$  and asymptotic holomorphic fibration  $p_Z: Z \to \Sigma$ . Let  $u_\Sigma$  be a holomorphic section of  $p_Z$  and u be a section of  $p_Y$  such that u is asymptotic to  $u_\Sigma$ . If the graph of u is associative then it is an ACyl associative in Y whose asymptotic cross section is the graph of  $u_\Sigma$ . Let (Z,X) be a building block with projective morphism  $f:Z \to \mathbf{P}^1$  and let  $V:=Z\setminus X$  be a corresponding ACyl Calabi-Yau 3-fold. Then  $f:V\to C$  is a holomorphic K3 fibration (possibly some of the fibers are singular) and  $\tilde{f}:=\mathbf{1}\times f:S^1\times V\to S^1\times C$  is an ACyl coassociative fibration. Moreover the graph of any holomorphic section of  $f:Z\to \mathbf{P}^1$  is an example of holomorphic curve C considered in Example 3.23 (1).

One can also expect to construct ACyl associative submanifolds from gradient graphs in adiabatic limits of ACyl co-associative fibrations following [Don17] and [DS22].

### 3.3 Linear analysis on ACyl associative submanifolds

By introducing appropriate weighted function spaces of sections of vector bundles over ACyl submanifolds (where the weights represent decay rates of the sections at infinity) we obtain a Fredholm theory for asymptotically translation invariant elliptic operators provided the weights avoid wall of critical weights or rates. The index of the operator changes when the weights lie on various connected components of the complement of the wall of the critical rates. This theory originally appeared in Lockhart-McOwen [LM85]. A very good exposition can be found in [Maro2, Chapter 4].

#### 3.3.1 Weighted function spaces

Let P be an ACyl submanifold in an ACyl  $G_2$ -manifold  $(Y, \phi)$  with asymptotic cylinder  $C = \mathbb{R} \times \Sigma$  as in Definition 3.20. Let  $\Sigma = \coprod_{i=1}^m \Sigma_i$  be the decomposition of  $\Sigma$  into connected components. Then  $C = \coprod_{i=1}^m C_i$ , where  $C_i = \mathbb{R} \times \Sigma_i$ . We would like to define the weighted Sobolev and Hölder spaces with rate  $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_m) \in \mathbb{R}^m$ .

**Notation 3.27.** Let  $\lambda, \lambda' \in \mathbf{R}^m$ . The notation  $\lambda < \lambda'$  means for all i = 1, 2, ..., m we have  $\lambda_i < \lambda_i'$ . Similarly we denote  $\lambda > \lambda'$  and  $\lambda = \lambda'$  by replacing < with > and = respectively. For any  $s \in \mathbf{R}$ ,  $\lambda + s := (\lambda_1 + s, \lambda_2 + s, ..., \lambda_m + s)$ . Set,  $|\lambda| := \sum_{i=1}^m |\lambda_i|$ .

**Definition 3.28.** For each  $\lambda = (\lambda_1, \lambda_2, ..., \lambda_m) \in \mathbb{R}^m$ , a weight function  $w_{P,\lambda} : P \to (0, \infty)$  is a smooth function on P such that for all  $x = \Upsilon \circ \Psi_P(t, \sigma)$  with  $(t, \sigma)$  in  $(T_0, \infty) \times \Sigma_i$ ,

$$w_{P\lambda}(x) = e^{-\lambda_i t}$$
.

Let  $\lambda \in \mathbb{R}^m$  and  $k \ge 0$ ,  $p \ge 1$ ,  $\gamma \in (0,1)$ . For a continuous section u of NP we define the weighted  $L^{\infty}$  norm and the weighted Hölder semi-norm by

$$||u||_{L_{P,\lambda}^{\infty}} := ||w_{P,\lambda}u||_{L^{\infty}(NP)}, \quad [u]_{C_{P,\lambda}^{0,\gamma}} := [w_{P,\lambda}u]_{C^{0,\gamma}(NP)}.$$

For a continuous section u of NP with k continuous derivatives we define the **weighted**  $C^k$  **norm** and the **weighted Hölder norm** by

$$\|u\|_{C^k_{P,\lambda}}:=\sum_{j=0}^k\|(\nabla^{\perp}_P)^ju\|_{L^{\infty}_{P,\lambda}},\ \, \|u\|_{C^{k,\gamma}_{P,\lambda}}:=\sum_{j=0}^k\|(\nabla^{\perp}_P)^ju\|_{L^{\infty}_{P,\lambda}}+\big[(\nabla^{\perp}_P)^ku\big]_{C^{0,\gamma}_{P,\lambda}},$$

and the weighted Sobolev norm by

$$||u||_{W^{k,p}_{P,\lambda}} := \Big(\sum_{j=0}^k \int_P |w_{P,\lambda}(\nabla_P^\perp)^j u|^p dV_P\Big)^{\frac{1}{p}}.$$

Here the connection  $\nabla_P^{\perp}$  on NP is the projection of the Levi-Civita connection induced by  $g_{\phi}$  for the decomposition  $TY_{|P} = TP \oplus NP$  and the  $|\cdot|$  is with respect to the the metric  $g_{\phi}$ . We denote always  $W_{P,\lambda}^{0,p}$  by  $L_{P,\lambda}^p$ .

We define the **weighted Hölder space**  $C_{P,\lambda}^{k,\gamma}$  to be the space of continuous sections of NP with k continuous derivatives and finite weighted Hölder norm  $\|\cdot\|_{C_{P,\lambda}^{k,\gamma}}$ . This is a Banach space but not separable.

We define the weighted Sobolev space  $W_{P,\lambda}^{k,p}$ , the weighted  $C^k$ -space  $C_{P,\lambda}^k$ , the weighted  $L^p$ -space  $L_{P,\lambda}^p$ , the weighted  $L^\infty$ -space  $L_{P,\lambda}^\infty$  to be the completion of the space of compactly supported smooth sections of NP, namely  $C_c^\infty(NP)$  with respect to the weighted Hölder norm  $\|\cdot\|_{C_{P,\lambda}^{k,\gamma}}$ , the weighted Sobolev norm  $\|\cdot\|_{W_{P,\lambda}^{k,p}}$ , the weighted  $C^k$ -norm  $\|\cdot\|_{C_{P,\lambda}^k}$ , the weighted  $L^p$ -norm  $\|\cdot\|_{L_{P,\lambda}^p}$ .

Furthermore, we define the weighted  $C^{\infty}$ -space  $C_{P\lambda}^{\infty}$  by

$$C_{P,\lambda}^{\infty} := \bigcap_{k=0}^{\infty} C_{P,\lambda}^{k}.$$

**Proposition 3.29.** Let  $k, l \ge 0$ , p, q > 1 with  $\frac{1}{p} + \frac{1}{q} = 1$  and  $\lambda, \lambda_1, \lambda_2 \in \mathbb{R}^m$ . Then the  $L^2$ -inner product map

$$W_{\lambda_1,P}^{k,p} \times W_{\lambda_2,P}^{l,q} \xrightarrow{\langle \cdot, \cdot \rangle_{L^2}} \mathbf{R}$$

is continuous if  $\lambda_1 + \lambda_2 \leq 0$ . Moreover, this  $L^2$ -inner product yields a Banach space isomorphism

$$(L^p_{\lambda,P})^*\cong L^q_{-\lambda,P}.$$

*Proof.* For  $u \in L^p_{\lambda_1,P}$  and  $v \in L^q_{\lambda_2,P}$  we obtain

$$\langle u, v \rangle_{L^{2}} = \int_{P} (w_{P,\lambda_{1}} u)(w_{P,\lambda_{2}} v) w_{P,-\lambda_{1}-\lambda_{2}} dV_{P}$$

$$\lesssim ||u||_{L_{P,\lambda_{1}}^{P}} ||v||_{L_{P,\lambda_{2}}^{q}}, \text{ provided } \lambda_{1} + \lambda_{2} \leqslant 0.$$

This proves that the  $L^2$ -inner product map is continuous. The isomorphism stated in the proposition follows from the standard fact that  $\langle \cdot, \cdot \rangle_{L^2} : L^p \times L^q \to \mathbb{R}$  is a dual pairing.

The following is a weighted embedding and compactness theorem [Maro2, Theorem 4.2, Theorem 4.3].

**Proposition 3.30.** Let  $\lambda, \lambda_1, \lambda_2 \in \mathbb{R}^m$ ,  $k, l \in \mathbb{N} \cup \{0\}$ ,  $p, q \ge 1$ ,  $\alpha, \beta \in (0, 1)$ . Then

- i) If  $k \ge l$  and  $k \frac{n}{p} \ge l \frac{n}{q}$ , the inclusion  $W_{P,\lambda_1}^{k,p} \longleftrightarrow W_{P,\lambda_2}^{l,q}$  is a continuous embedding, provided either  $p \le q, \lambda_1 \le \lambda_2$  or,  $p > q, \lambda_1 < \lambda_2$ .
- ii) If  $k + \alpha \geqslant l + \beta$ , the inclusion  $C_{P,\lambda_1}^{k,\alpha} \hookrightarrow C_{P,\lambda_2}^{l,\beta}$  is a continuous embedding, provided  $\lambda_1 \leqslant \lambda_2$ .
- iii) If  $k \frac{n}{p} \ge l + \alpha$ , the inclusion  $W_{P,\lambda}^{k,p} \hookrightarrow C_{P,\lambda}^{l,\alpha}$  is a continuous embedding.
- iv) Each of the above embeddings are compact provided the inequalities in the corresponding hypotheses are strict inequalities.

# 3.3.2 Asymptotically translation invariant elliptic operator

Let P be an ACyl associative submanifold with asymptotic cylinder C and rate  $\mu < 0$  in an ACyl  $G_2$  manifold  $(Y, \phi)$  with asymptotic cross section Z and rate  $\nu$ . Then C is an associative cylinder in  $Y_0 = \mathbf{R} \times Z$  and we have the translation invariant (cylindrical) Fueter operator

$$\mathbf{D}_C = J\partial_t + \mathbf{D}_{\Sigma}.$$

**Definition 3.31.** Let  $C = \coprod_{i=1}^{m} C_i$ , where each  $C_i = \mathbf{R} \times \Sigma_i$  is connected. Then the set of critical rates  $\mathcal{D}_C$  is defined by

$$\mathcal{D}_C := \{(\lambda_1, \lambda_2, ..., \lambda_m) \in \mathbf{R}^m : \lambda_i \in \mathcal{D}_{C_i} \text{ for some } i\},\$$

where  $\mathcal{D}_{C_i}$  is the set of all indicial roots of  $\mathbf{D}_{C_i}$  (see Definition 3.14). We call  $\mathcal{D}_C$  as wall of critical rates in  $\mathbf{R}^m$ . We also define

$$V_{\lambda} := \bigoplus_{i=1}^{m} V_{\lambda_i}, \ d_{\lambda_i} := \dim V_{\lambda_i} \text{ and } d_{\lambda} := \sum_{i=1}^{m} d_{\lambda_i}.$$

**Definition 3.32.** Let P be an ACyl associative submanifold with asymptotic cylinder C. The translation invariant tubular neighbourhood map  $\Upsilon_C$  induces canonical bundle isomorphisms between NP, TP and NC, TC respectively over the end  $C_\infty := (T_0, \infty) \times \Sigma$  (see Definition 2.13). Let  $\mathbf{D}_P : C_c^\infty(NP) \to C_c^\infty(NP)$  be a first order, formally self-adjoint elliptic operator. It is called asymptotically translation invariant uniformly elliptic operator asymptotic to  $\mathbf{D}_C$  if over  $C_\infty$  with the canonical bundle isomorphisms we have

$$\mathbf{D}_P = \sigma(\mathbf{D}_P) \circ \nabla_C^{\perp} + A_P$$

such that

$$|\nabla_C^k(\sigma(\mathbf{D}_P) - \sigma(\mathbf{D}_C))| = O(e^{\mu t})$$
 and  $|(\nabla_C^\perp)^k(A_P)| = O(e^{\mu t})$ 

as  $t \to \infty$  for all  $k \in \mathbb{N}_0$ . Here t denotes the coordinate on  $\mathbb{R}^+$ , and  $|\cdot|$  and the normal connection  $\nabla_C^{\perp}$  on NC are induced by the product metric on  $\mathbb{R}^+ \times Z$ . Here  $\sigma$  is the symbol of a differential operator.

The operator  $\mathbf{D}_P$  has canonical extensions to weighted function spaces and we denote them as follows:

$$\mathbf{D}^{k,p}_{P,\lambda}:W^{k+1,p}_{P,\lambda}\to W^{k,p}_{P,\lambda},\quad \mathbf{D}^{k,\gamma}_{P,\lambda}:C^{k+1,\gamma}_{P,\lambda}\to C^{k,\gamma}_{P,\lambda}.$$

The following proposition is about elliptic regularity and estimates for asymptotically translation invariant uniformly elliptic operator [Maro2, Theorem 4.6].

**Proposition 3.33.** Suppose  $\lambda \in \mathbb{R}^m$ ,  $k \geq 0$ , p > 1,  $\gamma \in (0,1)$ . Let  $u, v \in L^1_{loc}$  be two locally integrable sections of NP such that u is a weak solution of the equation  $\mathbf{D}_P u = v$ .

i) If  $v \in C_{P,\lambda}^{k,\gamma}$ , then  $u \in C_{P,\lambda}^{k+1,\gamma}$  is a strong solution and there exists a constant C > 0 such that

$$||u||_{C^{k+1,\gamma}_{P,\lambda}} \le C\Big(||\mathbf{D}^{k,\gamma}_{P,\lambda}u||_{C^{k,\gamma}_{P,\lambda}} + ||u||_{L^{\infty}_{P,\lambda}}\Big).$$

ii) If  $v \in W_{P,\lambda}^{k,p}$ , then  $u \in W_{P,\lambda}^{k+1,p}$  is a strong solution and there exists a constant C > 0 such that

$$||u||_{W^{k+1,p}_{P,\lambda}} \leq C\Big(||\mathbf{D}^{k,p}_{P,\lambda}u||_{W^{k,p}_{P,\lambda}} + ||u||_{L^{\infty}_{P,\lambda}}\Big).$$

**Proposition 3.34.** Suppose  $k \ge 0$ , p, q > 1 with  $\frac{1}{p} + \frac{1}{q} = 1$  and  $\lambda_1, \lambda_2 \in \mathbb{R}^m$ . For all  $u \in W^{k+1,p}_{P,\lambda_1}, v \in W^{k+1,q}_{P,\lambda_2}$  we have  $\langle \mathbf{D}^{k,p}_{P,\lambda_1} u, v \rangle_{L^2} = \langle u, \mathbf{D}^{k,q}_{P,\lambda_2} v \rangle_{L^2}$  if  $\lambda_1 + \lambda_2 \le 0$ .

*Proof.* The result is true for  $u, v \in C_c^{\infty}(NP)$  and therefore the general statement follows from Proposition 3.29.

**Definition 3.35.** Let  $C = \mathbb{R} \times \Sigma$  be an associative cylinder. We can define the Banach spaces  $W_{C,\lambda}^{k,p}$ ,  $C_{C,\lambda}^{k,\gamma}$  and all others over C similar to Section 3.3.1 replacing P by C, NP by NC and weight function  $w_{P,\lambda}$  by  $w_{C,\lambda}: C = \coprod_i C_i \to \mathbb{R}$  where  $w_{C,\lambda}(t,\sigma_i) = e^{-\lambda_i t}$ .

The following lemma is explained in [Dono2, Lemma 3.1, Proposition 3.4, Section 3.3.1] and [MP78, Theorem 5.1].

**Lemma 3.36.** Let  $C = \mathbf{R} \times \Sigma$  be a connected associative cylinder. Then the translation invariant Fueter operator  $\mathbf{D}_C = J\partial_t + \mathbf{D}_\Sigma : C_{C,\lambda}^{k+1,\gamma} \to C_{C,\lambda}^{k,\gamma}$  is invertible if and only if  $\lambda \in \mathbf{R}^m \setminus \mathcal{D}_C$ . Moreover, any element  $u \in \ker \mathbf{D}_C$  has a  $L^2$ -decomposition

$$u=\sum_{\lambda\in\mathcal{D}_C}e^{\lambda t}u_{\Sigma,\lambda}.$$

where  $u_{\Sigma,\lambda}$  are  $\lambda$ -eigensections of  $J\mathbf{D}_{\Sigma}$ .

**Proposition 3.37.** Let  $\lambda \in \mathbb{R}^m$ ,  $k \ge 0$ , p > 1,  $\gamma \in (0,1)$ . Then  $\mathbf{D}_{P,\lambda}^{k,p}$  and  $\mathbf{D}_{P,\lambda}^{k,\gamma}$  are Fredholm for all  $\lambda \in \mathbb{R}^m \setminus \mathcal{D}_C$ . Moreover, for all  $\lambda \in \mathbb{R}^m$ , Ker  $\mathbf{D}_{P,\lambda}^{k,p} = \operatorname{Ker} \mathbf{D}_{P,\lambda}^{k,\gamma}$  is finite dimensional, independent of k, p and  $\gamma$ .

*Proof.* The operators  $\mathbf{D}_{P,\lambda}^{k,p}$  and  $\mathbf{D}_{P,\lambda}^{k,\gamma}$  are Fredholm with weight  $\lambda \in \mathbf{R}^m \setminus \mathcal{D}_C$  follows from [Dono2, Proposition 3.6, Section 3.3.1] or [LM85, Theorem 6.2] . Independence of  $k, p, \gamma$  follows from Proposition 3.33 and Proposition 3.30.

**Lemma 3.38.** Let  $\lambda \in \mathbb{R}^m$  and  $\lambda_1, \lambda_2$  be two elements in  $\mathbb{R}^m \setminus \mathcal{D}_C$  with  $\lambda_1 < \lambda < \lambda_2, |\lambda_2 - \lambda_1| \leq |\mu|$  and there are no critical rates between  $\lambda_1, \lambda_2$  except possibly  $\lambda$ . We define the set

$$S_{\lambda_2} := \{ u \in C_{P,\lambda_2}^{k+1,\gamma} : \mathbf{D}_{P,\lambda_2}^{k,\gamma} u \in C_{P,\lambda_1}^{k,\gamma} \}.$$

Define a linear map  $e_{P,\lambda}: V_{\lambda} \to C_{P,\lambda_2}^{k+1,\gamma}$  by  $e_{P,\lambda}(w) := \chi_{T_0} w$  (under the identification of the canonical bundle isomorphisms  $NP \cong NC$  over the end  $(T_0, \infty) \times \Sigma$ ). Then there exists a unique linear map  $\widetilde{i}_{P,\lambda}: S_{\lambda_2} \to V_{\lambda}$  such that for any  $u \in S_{\lambda_2}$  we have

$$u - e_{P,\lambda} \circ \widetilde{i}_{P,\lambda} u \in C_{P,\lambda_1}^{k+1,\gamma}$$

Moreover, the following statements hold.

(i)  $S_{\lambda_2} \subset C_{P,\lambda}^{k+1,\gamma}$ ,  $\operatorname{Ker} \mathbf{D}_{P,\lambda_2}^{k,\gamma} = \operatorname{Ker} \mathbf{D}_{P,\lambda}^{k,\gamma}$ , and  $\operatorname{Ker} \widetilde{i}_{P,\lambda} = C_{P,\lambda_1}^{k+1,\gamma}$ . The restriction map

(3.39) 
$$i_{P,\lambda} := \widetilde{i}_{P,\lambda} : \operatorname{Ker} \mathbf{D}_{P,\lambda_2}^{k,\gamma} \to V_{\lambda}$$

satisfies the following: Ker  $i_{P,\lambda} = \text{Ker } \mathbf{D}_{P,\lambda}^{k,\gamma}$ .

(ii)  $S_{\lambda_2} = C_{P,\lambda_1}^{k+1,\gamma} + \text{im } e_{P,\lambda}$  and the restriction of  $D_{M,\lambda_2}^{k,\gamma}$ , denoted by

(3.40) 
$$\widetilde{\mathbf{D}}_{P,\lambda_1}^{k,\gamma}: C_{P,\lambda_1}^{k+1,\gamma} + \operatorname{im} e_{P,\lambda} \to C_{P,\lambda_1}^{k,\gamma}$$

has the property that  $\operatorname{Ker} \mathbf{D}_{P,\lambda_2}^{k,\gamma} = \operatorname{Ker} \widetilde{\mathbf{D}}_{P,\lambda_1}^{k,\gamma}$ ,  $\operatorname{Coker} \mathbf{D}_{P,\lambda_2}^{k,\gamma} \cong \operatorname{Coker} \widetilde{\mathbf{D}}_{P,\lambda_1}^{k,\gamma}$ .

(iii) (Wall crossing formula)  $\operatorname{Ind}(\mathbf{D}_{P,\lambda_2}) = \operatorname{Ind}(\mathbf{D}_{P,\lambda_1}) + d_{\lambda}.$ 

*Proof.* Let u be an element of  $\mathcal{S}_{\lambda_2}$ . Define,  $\tilde{u}:=\chi_{T_0}u\in C^\infty(NC)$ . Now  $\mathbf{D}_C\tilde{u}\in C^\infty_{C,\lambda_1}$ . By Lemma 3.36 there exists a unique  $v\in C^\infty_{C,\lambda_1}$  such that  $\mathbf{D}_C(\tilde{u}-v)=0$ . Define

$$\widetilde{i}_{P,\lambda}(u) := e^{\lambda t} (\widetilde{u} - v)_{\Sigma,\lambda}$$

following the decomposition in Lemma 3.36. Since  $\tilde{u} - v - e^{\lambda t} (\tilde{u} - v)_{\Sigma,\lambda} \in C^{\infty}_{C,\lambda_1}$ , this also proves (i) and  $\operatorname{Ker} \mathbf{D}^{k,\gamma}_{P,\lambda_2} = \operatorname{Ker} \widetilde{\mathbf{D}}^{k,\gamma}_{P,\lambda_1}$ . Note that  $\widetilde{\mathbf{D}}^{k,\gamma}_{P,\lambda_1}$  is well-defined as  $|\lambda - \lambda_1| \leq |\mu|$ .

Denote W to be the closure of im  $\widetilde{\mathbf{D}}_{P,\lambda_1}^{k,\gamma}$  in  $C_{P,\lambda_2}^{k,\gamma}$ . Then we have bijective linear maps,  $\operatorname{Coker} \widetilde{\mathbf{D}}_{P,\lambda_1}^{k,\gamma} \to C_{P,\lambda_2}^{k,\gamma}/W$  and  $C_{P,\lambda_2}^{k,\gamma}/W \to \operatorname{Coker} \mathbf{D}_{P,\lambda_2}^{k,\gamma}$ . Indeed, the first map is bijective as  $C_{P,\lambda_1}^{k,\gamma} \hookrightarrow C_{P,\lambda_2}^{k,\gamma}$  is continuous with dense image and the  $\operatorname{Coker} \widetilde{\mathbf{D}}_{P,\lambda_1}^{k,\gamma}$  is finite dimensional. Therefore every  $[v] \in C_{P,\lambda_2}^{k,\gamma}/W$  has representative  $v \in C_{P,\lambda_1}^{k,\gamma}$ . Moreover, if  $v = \mathbf{D}_{P,\lambda_2}^{k,\gamma}u$ , then  $u \in S_{\lambda_2}$  and hence, [v] = 0. This proves the last map is injective whose surjectivity is obvious. Thus, the obvious map  $\operatorname{Coker} \widetilde{\mathbf{D}}_{P,\lambda_1}^{k,\gamma} \to \operatorname{Coker} \mathbf{D}_{P,\lambda_2}^{k,\gamma}$  is an isomorphism and this finishes the proof of (ii). Since  $\operatorname{Ind}(\widetilde{\mathbf{D}}_{P,\lambda_1}) = \operatorname{Ind}(\mathbf{D}_{P,\lambda_1}) + d_{\lambda}$ , (iii) follows from (ii).

**Proposition 3.41.** Ker  $D_{P,\lambda}^{k,\gamma}$  is independent of  $\lambda$  in each connected component of  $\lambda \in \mathbb{R}^m \backslash \mathcal{D}_C$ . Moreover, for all  $\lambda \in \mathbb{R}^m \backslash \mathcal{D}_C$ , we have

- (i) Coker  $D_{P,\lambda} \cong \text{Ker } D_{P,-\lambda}$ . If  $\lambda \geqslant 0$ , then  $\text{Ker } D_{P,-\lambda} = \text{Coker } D_{P,\lambda}$ .
- (ii) If s > 0 such that 0 is the possibly only critical rates in between -s and s, then

$$\dim \operatorname{Ker} \mathbf{D}_{P,0} = \dim \operatorname{Ker} \mathbf{D}_{P,s}, \quad and \quad \operatorname{Ind}(\mathbf{D}_{P,s}) = \dim \operatorname{Ker} \mathbf{D}_{P,s} - \dim \operatorname{Ker} \mathbf{D}_{P,-s} = \frac{d_0}{2}$$

$$(iii) \operatorname{Ind}(\mathbf{D}_{P,\lambda}) = \sum_{\lambda_i \geqslant 0} \left( \frac{d_{0,i}}{2} + \sum_{\zeta_i \in \mathcal{D}_{C_i} \cap (0,\lambda_i)} d_{\zeta_i} \right) - \sum_{\lambda_i < 0} \left( \frac{d_{0,i}}{2} + \sum_{\zeta_i \in \mathcal{D}_{C_i} \cap (\lambda_i,0)} d_{\zeta_i} \right).$$

*Proof.* The statement  $\operatorname{Ker} \mathbf{D}_{P,\lambda}^{k,\gamma}$  is independent of  $\lambda$  in each connected component of  $\lambda \in \mathbf{R}^m \backslash \mathcal{D}_C$  is a direct consequence of part (i) of Lemma 3.38. Now, (i) follows from Proposition 3.29 and the fact that  $\mathbf{D}_P$  is formally self adjoint, and if  $\lambda \geq 0$  then  $C_{P,-\lambda}^{k,\gamma} \subset C_{P,\lambda}^{k,\gamma}$ . To see (ii), observe from Lemma 3.38 that,  $\operatorname{Ker} \mathbf{D}_{P,s} = \operatorname{Ker} \mathbf{D}_{P,0}$  and

$$\dim \operatorname{Ker} \mathbf{D}_{P,s} - \dim \operatorname{Ker} \mathbf{D}_{P,-s} = \dim \operatorname{Ker} \mathbf{D}_{P,-s} - \dim \operatorname{Ker} \mathbf{D}_{P,s} + d_0.$$

Finally, (iii) is an immediate consequences of (ii) and the wall crossing formula in the Lemma 3.38.

# 3.4 Moduli space of ACyl associative submanifolds

In this subsection we will consider the moduli space of ACyl associative submanifolds (possibly with multiple ends) in an ACyl  $G_2$ -manifold and prove Theorem 1.12 and Theorem 1.13. We will see that this moduli space is locally homeomorphic to the zero set of a smooth nonlinear Fredholm map between suitable Banach manifolds. Therefore by the implicit function theorem we can conclude that this moduli space is locally homeomorphic to the zero set of a smooth nonlinear map, called Kuranishi map or obstruction map between two finite dimensional vector spaces.

Let  $(Y, \phi)$  be an ACyl  $G_2$ -manifold with asymptotic cross section  $(Z, \omega, \Omega)$  with rate  $\nu < 0$  as in Definition 2.17. Recall that we have denoted in Section 3.1.1 the set of all oriented two dimensional connected, closed submanifolds of Z by S and the moduli space of holomorphic curves in S by  $\mathcal{M}_Z^{\text{hol}}$ .

**Definition 3.42.** Let  $S_{ACyl}$  be the set of all ACyl submanifolds in Y as in Definition 3.20. Denote by  $S_{ACyl,\Sigma}^{\mu}$  the space of all ACyl submanifolds with fixed asymptotic cross section  $\Sigma$  and rate  $\mu$ . Note that if  $\Sigma = \coprod_{i=1}^{m} \Sigma_i$  and  $\mu \in \mathbb{R}^m$  then any  $P \in S_{ACyl,\Sigma}^{\mu}$  has m ends. The **moduli space**  $\mathcal{M}_{ACyl}$  of ACyl associative submanifolds is defined by

$$\mathfrak{M}_{ACyl} := \{P \in \mathbb{S}_{ACyl} : P \text{ is an ACyl associative submanifold}\}.$$

The **moduli space**  $\mathcal{M}^{\mu}_{ACyl,\Sigma}$  of all ACyl associative submanifolds with fixed asymptotic cross section  $\Sigma$  and rate  $\mu$  is defined by

$$\mathcal{M}^{\mu}_{ACyl,\Sigma} := \mathcal{S}^{\mu}_{ACyl,\Sigma} \cap \mathcal{M}_{ACyl}.$$

We will now define topologies on  $\mathcal{M}_{ACyl,\Sigma}^{\mu}$  and  $\mathcal{M}_{ACyl}$ . They will be called  $C_{\mu}^{\infty}$ -topology and weighted  $C^{\infty}$ -topology respectively. The topology on each of these moduli spaces will be defined by specifying a basis. The open sets in the basis will be constructed using end cylindrical (ECyl) tubular neighbourhood maps of only ECyl submanifolds (see Definition 3.43). Indeed given an ACyl submanifold we will chose an ECyl submanifold with an ECyl tubular neighbourhood containing the ACyl submanifold (see Definition 3.44). This will suffice to define the  $C_{\mu}^{\infty}$ -topology of  $\mathcal{M}_{ACyl,\Sigma}^{\mu}$ . For the weighted  $C^{\infty}$ -topology of  $\mathcal{M}_{ACyl}$  we aditionally need to vary the ECyl submanifolds and the ECyl tubular neighbourhood maps. This will be done by choosing suitable ECyl diffeomorphisms on Y.

**Definition 3.43.** Let P be an end cylindrical (ECyl) submanifold of Y with end cylinder C as in Definition 3.20. Let  $\Upsilon_C: V_C \to U_C$  be a translation invariant tubular neighbourhood map of C. A tubular neighbourhood map  $\Upsilon_P: V_P \to U_P$  of P is called **end cylindrical (ECyl)** if  $V_P$  and  $\Upsilon_P$  agree with  $\Upsilon_*(V_C)$  and  $\Upsilon \circ \Upsilon_C \circ \Upsilon_*^{-1}$  on  $\Upsilon((T_0 + 1, \infty) \times \Sigma)$  respectively.

**Definition 3.44.** Given an ACyl submanifold P with asymptotic cross section  $\Sigma$  and  $\alpha$  as in Definition 3.20, we define an ECyl submanifold  $P_C$  which is diffeomorphic to P,

$$P_C := K_P \cup (\Upsilon \circ \Upsilon_C)((1 - \chi_{T_0})\alpha)$$

We define  $K_{P_C} := P_C \setminus \Upsilon((T_0 + 1, \infty) \times Z)$ .




Figure 4: Construction of ECyl submanifold  $P_C$  from an ACyl submanifold P.

**Definition 3.45.** Given an ACyl submanifold P in Y with asymptotic cylinder C as in Definition 3.20, we choose an ECyl submanifold  $P_C$  as in Definition 3.44. Let

$$\Upsilon_{PC}: V_{PC} \to U_{PC}$$

be an end cylindrical (ECyl) tubular neighbourhood map of  $P_C$  as in Definition 3.43. There is a section  $\beta = (\beta_1, \beta_2, ..., \beta_m)$  in  $V_{P_C} \subset NP_C$  which is zero on  $K_P$  and  $\Upsilon_* \circ \alpha = \beta \circ \Upsilon$  on  $\Upsilon((T_0 + 1, \infty) \times \Sigma)$  such that  $\Upsilon_{P_C}(\Gamma_\beta)$  is P.

Denote

$$C_{\infty}:=(T_0+1,\infty)\times\Sigma,\ \ V_{C_{\infty}}:=V_{C_{|C_{\infty}}},\ \ U_{C_{\infty}}:=U_{C_{|C_{\infty}}},$$

$$P_{C_{\infty}} := P_{C_{|\Upsilon(C_{\infty})}}, \quad V_{P_{C_{\infty}}} := V_{P_{C_{|\Upsilon(C_{\infty})}}}.$$

The following commutative diagram helps us to keep track of the definitions above.

$$C_{\infty} \xleftarrow{\alpha} V_{C_{\infty}} \xrightarrow{\Upsilon_{C}} U_{C_{\infty}}$$

$$\downarrow^{\Upsilon} \qquad \qquad \downarrow^{\Upsilon} \qquad \qquad \downarrow^{\Upsilon}$$

$$P_{C_{\infty}} \xleftarrow{\beta} V_{P_{C_{\infty}}} = \Upsilon_{*}(V_{C_{\infty}}) \xrightarrow{\Upsilon_{P_{C}}} Y \backslash K_{Y}.$$

**Definition 3.46** ( $C_{\mu}^{\infty}$ -topology). We define the  $C_{\mu}^{k}$ -topology on the set  $\mathcal{S}_{ACyl,\Sigma}^{\mu}$  by specifying a basis which is a collection of all sets of the form  $\{\Upsilon_{P_{C}}(\Gamma_{u}): u \in \mathcal{V}_{P_{C},\mu}^{k}\}$ , where  $P \in \mathcal{S}_{ACyl,\Sigma}^{\mu}$ ,  $\Upsilon_{P_{C}}$  is an ECyl tubular neighbourhood map of  $P_{C}$  and  $\mathcal{V}_{P_{C},\mu}^{k}$  is an open set in  $C_{\mu}^{\infty}(V_{P_{C}}):=C_{P_{C},\mu}^{\infty}\cap C^{\infty}(V_{P_{C}})$ . Here the topology on  $C_{\mu}^{\infty}(V_{P_{C}})$  is induced by the  $C_{\mu}^{k}$ -norm on  $C^{\infty}(N_{P_{C}})$ .

The  $C_{\mu}^{\infty}$ -topology on the set  $\mathcal{S}_{ACyl,\Sigma}^{\mu}$  is the inverse limit topology of  $C_{\mu}^{k}$ -topologies on it, that is, a set is open with  $C_{\mu}^{\infty}$ -topology if and only if it is open in every  $C_{\mu}^{k}$ -topology.

The  $C^{\infty}_{\mu}$ -topology on the moduli space  $\mathfrak{M}^{\mu}_{ACyl,\Sigma}$  is then the subspace topology of  $C^{\infty}_{\mu}$ -topology of  $\mathcal{S}^{\mu}_{ACyl,\Sigma}$ .

**Definition 3.47** (Weighted  $C^{\infty}$ -topology). We now define weighted  $C^{\infty}$ -topology on the moduli space  $\mathfrak{M}_{ACyl}$  by specifying a basis. Let  $P \in \mathfrak{M}_{ACyl}$  be an ACyl assocciative submanifold with an asymptotic cone  $C = \mathbb{R} \times \Sigma$ . Let  $P_C$  be an ECyl submanifold as in Definition 3.44 and  $\Upsilon_{P_C}$  be an ECyl tubular neighbourhood map as in Definition 3.43. Since C is allowed to vary, we need to also vary the  $P_C$  and  $\Upsilon_{P_C}$ . This can be done as follows.

By Theorem 1.9, there is a tubular neighbourhood map  $\Upsilon_{\Sigma}: V_{\Sigma} \to U_{\Sigma}$  of  $\Sigma$  and an obstruction map  $ob_{\Sigma}: \mathcal{I}_{\Sigma} \to \mathcal{O}_{\Sigma}$  where  $\mathcal{I}_{\Sigma}:= C^{\infty}(V_{\Sigma}) \cap \ker \mathbf{D}_{\Sigma}$  and  $\mathcal{O}_{\Sigma}:= \operatorname{coker} \mathbf{D}_{\Sigma}$ . The inclusion map of normal vector fields in Definition 2.13 induces a smooth map

$$J_{\Sigma} \xrightarrow{:} \operatorname{Vect}(V_{\Sigma}) \xrightarrow{d\Upsilon_{\Sigma}} \operatorname{Vect}(U_{\Sigma}) \hookrightarrow \operatorname{Vect}(Z) \xrightarrow{\chi_{T_0}} \operatorname{Vect}(\mathbf{R} \times Z).$$

The time-1 flows of this family of vector fields defines a smooth family of diffeomorphisms,

$$\Phi: \mathcal{I}_{\Sigma} \to \mathrm{Diff}(\mathbf{R} \times Z), \ \xi \mapsto \Phi_{\xi}.$$

In particular, the map  $\Phi$  satisfies the following:

- (i)  $\Phi_0$  is identity and  $\Phi_{\xi}$  is identity on  $(-\infty, T_0) \times Z$  for all  $\xi \in \mathcal{I}_{\Sigma}$ .
- (ii)  $\Phi_{\xi}$  is independent of  $t \in [T_0 + 1, \infty)$  and  $\Phi_{\xi}(\{T_0 + 1\} \times \Sigma) = \{T_0 + 1\} \times \Upsilon_{\Sigma}(\xi)$ .

For any  $\xi$  in  $\mathcal{I}_{\Sigma}$  we define

$$\Upsilon_{\xi} := \Upsilon \circ \Phi_{\xi}, \quad P_C^{\xi} := (\Upsilon_{\xi} \circ \Upsilon^{-1}(P_C)) \cup K_P$$

and

$$\Upsilon_{P_C^{\xi}} := \left( \Upsilon_{\xi} \circ \Upsilon^{-1} \circ \Upsilon_{P_C} \right) \cup \Upsilon_{P_C|K_P} : V_{P_C} \to Y.$$

Finally, the basis of the **weighted**  $C^{\infty}$ -topology is a collection of all sets of the form

$$\{\Upsilon_{P_C^{\xi}}(\Gamma_u): u \in \mathcal{V}_{P_C,\mu}, \xi \in \mathcal{V}_{\Sigma}\}$$

where  $\mathcal{V}_{\Sigma}$  is an open set in  $\mathcal{I}_{\Sigma}$  and  $\mathcal{V}_{PC,\mu}$  is an open set in  $C^{\infty}_{\mu}(V_{PC})$ . Here  $\mathcal{I}_{\Sigma}$  is always chosen small enough so that we can consider any rate  $\mu < 0$  satisfying  $(\mu, 0) \cap \mathcal{D}_{C_{\xi}} = \emptyset$  for all associative cones  $C_{\xi} := \mathbf{R} \times \Upsilon_{\Sigma}(\xi)$  where  $\xi \in \mathcal{I}_{\Sigma}$ . Anyway this definition will be independent of such choices of  $\mu$  (see the last statement of Theorem 1.12).




Figure 5: Construction of  $P_C^{\xi}$ , a small ECyl deformation of  $P_C$ .

The following definition identifies two normal bundles  $NP_C$  and NP.

**Definition 3.48.** Let  $P \in \mathcal{M}_{ACyl}$  be an ACyl assocciative submanifold with an asymptotic cone C. Let  $P_C$  be an ECyl submanifold as in Definition 3.44 and  $\Upsilon_{P_C}$  be an ECyl tubular neighbourhood map as in Definition 3.43. Then there is a **canonical** bundle isomorphism as in Definition 2.13,

$$\Theta_P^C: NP_C \to NP.$$

**Definition 3.49.** Let  $P \in \mathcal{M}_{ACyl,\Sigma}^{\mu}$  and let  $\Upsilon_{PC}: V_{PC} \to U_{PC}$  be an end cylindrical (ECyl) tubular neighbourhood map of  $P_C$  as in Definition 3.43. We define  $\mathfrak{F}: C_{\mu}^{\infty}(V_{PC}) \times \mathcal{I}_{\Sigma} \to C^{\infty}(NP_C)$  as follows: for all  $u \in C_{\mu}^{\infty}(V_{PC})$ ,  $\xi \in \mathcal{I}_{\Sigma}$  and  $w \in C_{c}^{\infty}(NP_{C})$ ,

$$\langle \mathfrak{F}(u,\xi), w \rangle_{L^2} := \int_{\Gamma_u} \iota_w \Upsilon_{P_C^{\xi}}^* \psi.$$

The notation w in the integrand is actually  $\tilde{w}$ , the extension vector field of  $w \in C^{\infty}(NP_C)$  in the tubular neighbourhood as in Notation 2.16. The  $L^2$  inner product we choose here is coming from the canonical bundle isomorphism  $\Theta_P^C: NP_C \to NP$  as in Definition 3.48 and the induced metric on NP of  $g_{\phi}$ .

Remark 3.50. Let  $\beta \in C^{\infty}_{\mu}(V_{P_C})$  be the section corresponding to the ACyl associative P as in Definition 3.45. Then  $\mathfrak{F}(\beta,0)=0$ . Moreover, for  $u\in C^{\infty}_{\mu}(V_{P_C})$  and  $\xi\in \mathbb{J}_{\Sigma}$ , we have  $\Upsilon_{P_C^{\Sigma}}(\Gamma_u)\in \mathbb{M}_{ACyl}$  if and only if  $\mathfrak{F}(u,\xi)=0$ . Hence  $\mathbb{M}_{ACyl}$  is locally homeomorphic to  $\mathfrak{F}^{-1}(0)$ , provided  $\mathbb{J}_{\Sigma}$  and  $\mu$  are chosen so that  $(\mu,0)\cap \mathbb{D}_{C_{\xi}}=\emptyset$  for all associative cones  $C_{\xi}:=\mathbf{R}\times\Upsilon_{\Sigma}(\xi)$  where  $\xi\in \mathbb{J}_{\Sigma}$ .

Remark 3.51. Let  $u \in C^{\infty}_{\mu}(V_{P_C})$ ,  $\xi \in \mathcal{I}_{\Sigma}$  and  $w \in C^{\infty}_{c}(NP_C)$ . The restriction of the associator  $\Upsilon^*_{P^{\xi}_{\Sigma}}[\cdot,\cdot,\cdot]_{|\Gamma_{u}} \in C^{\infty}(N\Gamma_{u})$ . The map  $\mathfrak{F}$  can be rewritten as follows.

$$\langle \mathfrak{F}(u,\xi), w \rangle_{L^2} = \int_{\Gamma_u} \langle \Upsilon_{P_C^{\xi}}^* [\cdot, \cdot, \cdot]_{|\Gamma_u}, w \rangle.$$

This implies that  $\mathfrak{F}_{\xi} := \mathfrak{F}(\cdot, \xi)$  is of the form

$$\mathfrak{F}_{\xi}(u) = A_{\xi}(u, \nabla^{\perp}_{P_C} u) \nabla^{\perp}_{P_C} u + B_{\xi}(u, \nabla^{\perp}_{P_C} u).$$

Hence  $\mathfrak{F}_{\xi}: C_c^{\infty}(V_{P_C}) \to C_c^{\infty}(NP_C)$  is a non-linear first order differential operator.

**Proposition 3.52.** For any  $\xi \in \mathcal{I}_{\Sigma}$  the linearization of  $\mathfrak{F}_{\xi} := \mathfrak{F}(\cdot, \xi)$  at  $u \in C^{\infty}_{\mu}(V_{P_C})$ ,

$$d\mathfrak{F}_{\xi|_{\mathcal{U}}}: C^{\infty}_{P_{C},\mathcal{U}} \to C^{\infty}_{P_{C},\mathcal{U}}$$

is given by the following: for any  $v \in C^{\infty}_{P_C,\mu}$  and  $w \in C^{\infty}_c(NP_C)$ ,

$$\langle d\mathfrak{F}_{\xi|u}(v), w \rangle_{L^{2}} = \int_{\Gamma_{u}} \iota_{w} \mathcal{L}_{v}(\Upsilon_{P_{C}^{\xi}}^{*} \psi) = \int_{\Gamma_{u}} \left\langle \sum_{\substack{cyclic \\ permutations}} [e_{2}, e_{3}, \nabla_{e_{1}} v], w \right\rangle + \int_{\Gamma_{u}} \iota_{\nabla_{w} v}(\Upsilon_{P_{C}^{\xi}}^{*} \psi).$$

where  $\{e_1,e_2,e_3\}$  is a local oriented orthonormal frame for  $T\Gamma_u$  and by abusing notation we are denoting  $\Upsilon_{P_C^\xi}^* \nabla$  by  $\nabla$  and the associator  $\Upsilon_{P_C^\xi}^* [\cdot,\cdot,\cdot]$  by  $[\cdot,\cdot,\cdot]$  which are induced from the 3-form  $\Upsilon_{P_C^\xi}^* \phi$ .

*Proof.* For the family of sections  $\{u + tv : |t| \ll 1\}$  we have

$$\langle d\mathfrak{F}_{\xi|u}(v), w \rangle_{L^2} = \langle \frac{d}{dt} \Big|_{t=0} \mathfrak{F}_{\xi}(u+tv), w \rangle_{L^2} = \frac{d}{dt} \Big|_{t=0} \int_{\Gamma_{u+tv}} \iota_w(\Upsilon_{P_C^{\xi}}^* \psi).$$

This is equal to  $\int_{\Gamma_u} \mathcal{L}_v \iota_w (\Upsilon_{P_C^{\xi}}^* \psi) = \int_{\Gamma_u} \iota_w \mathcal{L}_v (\Upsilon_{P_C^{\xi}}^* \psi) + \iota_{[v,w]} (\Upsilon_{P_C^{\xi}}^* \psi)$ . As [v,w] = 0, using the definition of Lie derivative this is further equal to

$$\int_{\Gamma_{u}} \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \iota_{w}(\Upsilon_{p_{C}^{\sharp}}^{*}\psi)(\nabla_{e_{1}}v, e_{2}, e_{3}) + \int_{\Gamma_{u}} \iota_{\nabla_{w}v}(\Upsilon_{p_{C}^{\sharp}}^{*}\psi) + \iota_{w}\nabla_{v}(\Upsilon_{p_{C}^{\sharp}}^{*}\psi).$$

Corollary 3.53. Let  $u \in C^\infty_\mu(V_{P_C})$  and  $\xi \in \mathcal{I}_\Sigma$ . If  $\Upsilon_{P^\xi_C}(\Gamma_u) \in \mathcal{M}_{ACyl}$ , an ACyl associative submanifold then  $d\mathfrak{F}_{\xi|u}$  is given by the following: for all  $v \in C^\infty_{P_C,\mu}$  and  $w \in C^\infty_c(NP_C)$ ,

$$\langle d\mathfrak{F}_{\xi_{|u}}(v), w \rangle_{L^{2}} = \int_{\Gamma_{u}} \left\langle \sum_{\substack{cyclic \\ permutations}} [e_{2}, e_{3}, \nabla_{e_{1}}v], w \right\rangle.$$

**Proposition 3.54.** For any  $\xi \in \mathcal{I}_{\Sigma}$  the linearization of  $\mathfrak{F}_{\xi}$  at  $u \in C^{\infty}_{P,\mu}(V_{PC})$ ,  $d\mathfrak{F}_{\xi|u}$  is a formally self adjoint first order differential operator.

*Proof.* For all  $v, w \in C_c^{\infty}(NP_C)$ , the difference  $d\mathcal{F}_{\xi|u}(v)(w) - d\mathcal{F}_{\xi|u}(w)(v)$  is

$$\int_{\Gamma_u} \mathcal{L}_v \iota_w (\Upsilon_{P_C^\xi}^* \psi) - \mathcal{L}_w \iota_v (\Upsilon_{P_C^\xi}^* \psi) = \int_{\Gamma_u} \iota_w \mathcal{L}_v (\Upsilon_{P_C^\xi}^* \psi) + \iota_{[v,w]} (\Upsilon_{P_C^\xi}^* \psi) - \mathcal{L}_w \iota_v (\Upsilon_{P_C^\xi}^* \psi).$$

As [v, w] = 0, this is equal to  $\int_{\Gamma_u} \iota_w \iota_v (\Upsilon_{P_C^{\xi}}^* d\psi) - d(\iota_w \iota_v (\Upsilon_{P_C^{\xi}}^* \psi)) = 0$ .

**Definition 3.55.** Let  $P \in \mathcal{M}_{ACyl,\Sigma}^{\mu}$  and  $\beta \in C_{\mu}^{\infty}(V_{PC})$  representing P as in Definition 3.45. We define  $\mathfrak{L}_P : C_c^{\infty}(NP_C) \to C_c^{\infty}(NP_C)$  by

$$\mathfrak{L}_P u := d\mathfrak{F}_{0|B}(u).$$

**Proposition 3.56.** Let  $P \in \mathcal{M}_{ACyl}$  be an ACyl assocciative submanifold and  $\beta \in C^{\infty}_{\mu}(V_{PC})$  representing P as in Definition 3.45. Then

$$\mathfrak{L}_P = (\Theta_P^C)^{-1} \circ \mathbf{D}_P \circ \Theta_P^C,$$

where  $\mathbf{D}_P: C_c^{\infty}(NP) \to C_c^{\infty}(NP)$  is the Fueter operator defined in Definition 2.12.

*Proof.* For all  $v, w \in C_c^{\infty}(NP_C)$  we have by definition  $\langle \mathfrak{L}_P v, w \rangle_{L^2} = \langle \Theta_P^C \mathfrak{L}_P v, \Theta_P^C w \rangle_{L^2(NP)}$ . This is further equal to

$$\int_{\Gamma_{\beta}} \left\langle \sum_{\substack{\text{cyclic} \\ \text{permutations}}} [e_2, e_3, \nabla_{e_1}^{\perp} v], w \right\rangle = \int_{P} \left\langle \sum_{\substack{\text{cyclic} \\ \text{permutations}}} [e_2, e_3, \nabla_{e_1}^{\perp} (\Theta_P^C v)], \Theta_P^C w \right\rangle = \langle \mathbf{D}_P \Theta_P^C v, \Theta_P^C w \rangle_{L^2}.$$

The equality before the last equality holds, because  $w - \Theta_P^C w \in TP$  and  $v - \Theta_P^C v \in TP$ , and  $[\cdot, \cdot, \cdot]_{P} = 0$ .

**Proposition 3.57.** The operator  $\mathfrak{L}_P$  and  $\mathbf{D}_P$  are asymptotically translation invariant uniformly elliptic operators and asymptotic to the translation invariant Fueter operator  $\mathbf{D}_C$ .

*Proof.* Since  $\mathbf{D}_P$  is elliptic and by Proposition 3.56  $\mathfrak{L}_P = (\Theta_P^C)^{-1} \circ \mathbf{D}_P \circ \Theta_P^C$ , we obtain that  $\mathfrak{L}_P$  is elliptic.

It remains to prove that  $\mathfrak{L}_P$  is asymptotic to  $\mathbf{D}_C$ . For all  $v, w \in C_c^{\infty}(NC_t)$  where  $C_t := (T_0 + 1, t) \times \Sigma$  we have

$$\langle d\mathfrak{F}_{0|0}\Upsilon_* v, \Upsilon_* w \rangle_{L^2} = \int_{C_t} \iota_w \mathcal{L}_v(\Upsilon_C^* \Upsilon^* \psi) = \int_{C_t} \iota_w \mathcal{L}_v(\Upsilon_C^* \psi_0) + O(e^{\nu t}) \|v\|_{W^{1,2}(NC)} \|w\|_{L^2(NC)}.$$

By Proposition 3.61 (i) we obtain

$$\langle \Upsilon^* \mathfrak{Q}_P v, w \rangle_{L^2(NC)} = \langle \mathbf{D}_C v, w \rangle_{L^2(NC)} + O(e^{\mu t}) \|v\|_{W^{1,2}(NC)} \|w\|_{L^2(NC)}.$$

This completes the proof of the proposition for  $\mathfrak{L}_P$  and hence for  $D_P$  by Proposition 3.56.

**Definition 3.58.** Let P be an ACyl associative submanifold and  $\mathbf{D}_P: C_c^\infty(NP) \to C_c^\infty(NP)$  be the Fueter operator defined in Definition 2.12. Proposition 3.57 implies that it is an asymptotically translation invariant uniformly elliptic operator. Therefore it has the following canonical extensions:

$$\mathbf{D}^{k,p}_{P,\lambda}:W^{k+1,p}_{P,\lambda}\to W^{k,p}_{P,\lambda},\quad \mathbf{D}^{k,\gamma}_{P,\lambda}:C^{k+1,\gamma}_{P,\lambda}\to C^{k,\gamma}_{P,\lambda}.$$

Similarly we have canonical extensions

$$\mathfrak{Q}^{k,p}_{P,\lambda}:W^{k+1,p}_{P_C,\lambda}\to W^{k,p}_{P_C,\lambda},\quad \mathfrak{Q}^{k,\gamma}_{P,\lambda}:C^{k+1,\gamma}_{P_C,\lambda}\to C^{k,\gamma}_{P_C,\lambda}.$$

**Definition 3.59.** Proposition 3.57 implies that  $\mathbf{D}_P$  is an asymptotically translation invariant uniformly elliptic operator and asymptotic to the translation invariant Fueter operator  $\mathbf{D}_C$ . We define the **asymptotic limit map** 

$$i_{\infty}: \operatorname{Ker} \mathbf{D}_{P,0} \to \operatorname{Ker} \mathbf{D}_{\Sigma}$$

to be the map  $i_{P,0}$ , defined in (3.39) of Lemma 3.38.

**Definition 3.60.** For all  $\xi \in \mathcal{I}_{\Sigma}$  the nonlinear map  $Q_{\xi} : C_{\mu}^{\infty}(V_{P_C}) \to C^{\infty}(NP_C)$  is defined by

$$Q_{\xi} := \mathfrak{F}_{\xi} - d\mathfrak{F}_{\xi_{|0}} - \mathfrak{F}_{\xi}(0).$$

**Proposition 3.61.** For all  $\xi \in \mathcal{I}_{\Sigma}$ ,  $u, v \in C^{\infty}_{\mu}(V_{P_C})$  and  $\eta \in C^{\infty}_{P_{C,\mu}}$  we have

(i) 
$$|d\mathfrak{F}_{\xi_{|u}}(\eta) - d\mathfrak{F}_{\xi_{|v}}(\eta)| \leq (|u-v| + |\nabla^{\perp}(u-v)|)(|\eta| + |\nabla^{\perp}\eta|).$$

(ii) 
$$|Q_{\xi}(u) - Q_{\xi}(v)| \leq (|u| + |\nabla^{\perp}u| + |v| + |\nabla^{\perp}v|)(|u - v| + |\nabla^{\perp}(u - v)|)$$

(iii) 
$$\|Q_{\xi}(u)\|_{C^0_{P_C,2\mu}} \lesssim \|u\|_{C^1_{P_C,\mu}}^2$$
.

*Proof.* For all  $w \in C_c^{\infty}(NP_C)$  and  $\eta \in C_{P_C,u}^{\infty}$  we write

$$\langle d\mathfrak{F}_{\xi|u}(\eta) - d\mathfrak{F}_{\xi|v}(\eta), w \rangle_{L^2} = \int_0^1 \frac{d}{dt} \Big( d\mathfrak{F}_{\xi|tu+(1-t)v}(\eta)(w) \Big) dt$$

and using Proposition 3.52 this becomes

$$\int_{0}^{1} \left( \frac{d}{dt} \int_{\Gamma_{tu+(1-t)v}} L_{\eta} \iota_{w} (\Upsilon_{P_{C}^{\xi}}^{*} \psi) \right) dt = \int_{0}^{1} \int_{\Gamma_{tu+(1-t)v}} L_{(u-v)} L_{\eta} \iota_{w} (\Upsilon_{P_{C}^{\xi}}^{*} \psi) dt.$$

As, [u - v, w] = 0 and  $[\eta, w] = 0$  the last expression is same as

$$\int_0^1 \int_{\Gamma_{tu+(1-t)v}} \iota_w L_{(u-v)} L_{\eta}(\Upsilon_{P_C^{\xi}}^* \psi) dt.$$

The required estimate in (i) now follows from Lemma A.1. The estimate in (ii) follows from (i) after writing

$$Q_{\xi}(u) - Q_{\xi}(v) = \int_{0}^{1} dQ_{\xi|tu+(1-t)v}(u-v)dt = \int_{0}^{1} \left(d\mathfrak{F}_{\xi|tu+(1-t)v}(u-v) - d\mathfrak{F}_{\xi|0}(u-v)\right)dt.$$

Finally (iii) follows from (ii). Indeed, substituting v = 0 we have

$$w_{P_C,2\mu}|Q_{\xi}(u)| \lesssim w_{P_C,2\mu}(|u| + |\nabla^{\perp}u|)^2 \lesssim w_{P_C,\mu}^2(|u| + |\nabla^{\perp}u|)^2.$$

**Proposition 3.62.** For all  $\xi \in \mathcal{I}_{\Sigma}$  and  $u, v \in C^{k+1,\gamma}_{\mu}(V_{P_C})$  we have

$$\|Q_{\xi}(u) - Q_{\xi}(v)\|_{C^{k,\gamma}_{P_{C},2\mu}} \lesssim \|u - v\|_{C^{k+1,\gamma}_{P_{C},\mu}} (\|u\|_{C^{k+1,\gamma}_{P_{C},\mu}} + \|v\|_{C^{k+1,\gamma}_{P_{C},\mu}}).$$

*Proof.* With the above notation and appropriate product operation '·', one can express  $\mathcal{L}_u \mathcal{L}_v \psi$  formally as a quadratic polynomial

$$\mathcal{L}_{u}\mathcal{L}_{v}\psi = O(f_{1}) \cdot u \cdot v + O(f_{2}) \cdot (u \cdot \nabla^{\perp}v + v \cdot \nabla^{\perp}u) + \psi \cdot \nabla^{\perp}u \cdot \nabla^{\perp}v$$

where  $O(f_1) := \psi \cdot \nabla B + \psi \cdot B \cdot B + B \cdot \nabla \psi + \nabla^2 \psi + R \cdot \psi$  and  $O(f_2) := \nabla \psi + B \cdot \psi$ . With this observation and a similar computations as in Lemma A.1 we can prove this Proposition.

**Proof of Theorem 1.12.** Extending the nonlinear map  $\mathcal{F}_0$  to weighted Hölder spaces and applying Proposition 3.61, Proposition 3.62 and Proposition 3.52 we obtain that

$$\mathfrak{F}_0: C^{2,\gamma}_{\mu}(V_{P_C}) \to C^{1,\gamma}_{P_C,\mu}$$

since,  $\mathfrak{F}_0=\mathfrak{F}_0(0)+d\mathfrak{F}_{0|0}+Q_0$  and  $\mathfrak{F}_0(0)\in C^{1,\gamma}_{P_C,\nu}$ . As  $\mu\in(\nu,0)\backslash\mathcal{D}_C$ , by Proposition 3.37 we have that the linearization of  $\mathfrak{F}_0$  at  $\beta$ ,  $\mathfrak{L}_{P,\mu}$  is a Fredholm operator. Applying the implicit function theorem to  $\mathfrak{F}_0$  and Proposition 3.33 of elliptic regularity to  $\mathfrak{L}_{P,\mu}$  we obtain the map  $\mathrm{ob}_{P,\mu}$  as stated in the theorem (see [DK90, Proposition 4.2.19]). We need to only prove that, if  $u\in C^{2,\gamma}_{\mu}(V_{P_C})$  with  $\mathfrak{F}_0(u)=0$  then  $u\in C^{\infty}_{\mu}(V_{P_C})$ . To prove this, we observe

$$0=\mathfrak{L}_P(\mathcal{F}_0(u))=a(u,\nabla^\perp_{P_C}u)(\nabla^\perp_{P_C})^2u+b(u,\nabla^\perp_{P_C}u)$$

Since  $a(u, \nabla^{\perp}_{P_C} u) \in C^{1,\gamma}_{P_C,\mu}$  and  $b(u, \nabla^{\perp}_{P_C} u) \in C^{1,\gamma}_{P_C,\mu}$ , by Schauder elliptic regularity (see [Joyo7, Theorem 1.4.2]) and Proposition 3.33 we obtain  $u \in C^{3,\gamma}_{P_C,\mu}$ . By repeating this argument we can get higher regularity. The index formula follows from Proposition 3.41.

Finally we prove the last statement of the theorem. If  $\mathfrak{F}_0(u)=0$ , then  $d\mathfrak{F}_{0|0}(u)=-Q_0(u)-\mathfrak{F}_0(0)$ . By Proposition 3.61 we see that  $d\mathfrak{F}_{0|0}$  is also an ACyl uniformly elliptic operator asymptotic to  $\mathbf{D}_C$ . Also we have  $Q_0(u)\in C_{P_C,2\mu}^{k,\gamma}$  and  $\mathfrak{F}_0(0)\in C_{P_C,\nu}^{k,\gamma}$ . Therefore by an inductive procedure with the help of Lemma 3.38 we can conclude that  $u\in C_{\mu'}^{\infty}(V_{P_C})$ .

**Lemma 3.63.** The linearization of  $\mathfrak{F}(\beta,\cdot)$  at  $0 \in \mathcal{I}_{\Sigma}$ ,  $L : \ker \mathbf{D}_{\Sigma} \to C^{\infty}_{P_{C,U}}$  is given by

$$L(\xi) = \mathfrak{L}_P(\chi_{T_0} \Upsilon_* \xi).$$

*Proof.* For all  $w \in C_c^{\infty}(NP_C)$  and  $\xi \in \ker \mathbf{D}_{\Sigma}$ ,

$$\langle L(\xi), w \rangle_{L^2} = \int_{\Gamma_{\beta}} \frac{d}{dt} \Big|_{t=0} \iota_w(\Upsilon_{P_C^{t\xi}}^* \psi) = \int_{\Gamma_{\beta}} \iota_w \mathcal{L}_{(\chi_{T_0} \Upsilon_* \xi)}(\Upsilon_{P_C}^* \psi)$$

Therefore the lemma now follows from Proposition 3.52.

**Definition 3.64.** Let P be an ACyl associative and  $\Theta_P^C: NP_C \to NP$  be the isomorphism in Definition 3.48. We define

$$\widetilde{\mathfrak{L}}_{P,\mu}: C^{\infty}_{P_C,\mu} \oplus \ker D_{\Sigma} \to C^{\infty}_{P_C,\mu}$$

by  $\widetilde{\mathfrak{L}}_{P,\mu}(u,\xi) := \mathfrak{L}_P u + L\xi$ , where  $\mathfrak{L}_P$  and L are as in Definition 3.55 and Lemma 3.63. The operator

$$\widetilde{\mathbf{D}}_{P,\mu}: C_{P,\mu}^{\infty} \oplus \ker D_{\Sigma} \to C_{P,\mu}^{\infty}$$

is the operator  $\widetilde{\mathfrak{Q}}_{P,\mu}$  under the identification map  $\Theta_P^C$ , that is,

$$\widetilde{\mathbf{D}}_{P,\mu}(u,\xi) := \Theta_P^C \widetilde{\mathfrak{D}}_{P,\mu} \big( (\Theta_P^C)^{-1} u, \xi \big) = \mathbf{D}_P u + \Theta_P^C L \xi.$$

**Proof of Theorem 1.13.** Using Proposition 3.61 and Proposition 3.62 we obtain that the non-linear map

$$\mathcal{F}: C^{2,\gamma}_{\mu}(V_{P_C}) \times \mathcal{I}_{\Sigma} \to C^{1,\gamma}_{P_C,\mu}$$

since,  $\mathcal{F}(0,\xi) \in C_{P,\mu}^{1,\gamma}$ . By Lemma 3.63, the linearization of  $\mathcal{F}$  at  $(\beta,0)$ ,

$$d\mathfrak{F}_{|(\beta,0)}(u,\xi) = \mathfrak{L}_{P,\mu}u + L\xi.$$

Thus  $d\mathfrak{F}_{|(\beta,0)} = \widetilde{\mathfrak{L}}_{P,\mu}$  which is defined in Definition 3.64. Now the rest of the proof is similar to the proof of Theorem 1.12. The index formula follows from Proposition 3.41.

**Proposition 3.65.** Let P be an ACyl associative submanifold asymptotic to  $\Sigma$ . Assume that the asymptotic limit map  $\iota_{\infty}$ :  $\ker \mathbf{D}_{P,0} \to \ker \mathbf{D}_{\Sigma}$  is injective. Then  $\operatorname{im} \iota_{\infty}$  is Lagrangian subspace of  $\ker \mathbf{D}_{\Sigma}$  with respect to the symplectic form  $\omega$  which is defined as follows: for  $\xi_1, \xi_2 \in \ker \mathbf{D}_{\Sigma}$ ,

$$\boldsymbol{\omega}(\xi_1,\xi_2) \coloneqq \int_{\Sigma} \omega(\xi_1,\xi_2).$$

*Proof.* By the proof of Proposition 3.54, if  $\xi_k = i_{P,0}(u_k)$  with  $u_k \in \ker \mathcal{L}_{P,0}$ , k = 1, 2, then

$$\boldsymbol{\omega}(\xi_1,\xi_2) = \int_{\Sigma} \omega(\xi_1,\xi_2) = \lim_{T \to \infty} \int_{\{T\} \times \Sigma} \iota_{u_1} \iota_{u_2} \Upsilon_{P_C}^* \psi = \int_{P} \langle \mathcal{L}_P u_1, u_2 \rangle - \langle u_1, \mathcal{L}_P u_2 \rangle = 0.$$

Therefore im  $\iota_{\infty}$  is an isotropic immersion by Proposition 3.56. By Proposition 3.41 we see that dim ker  $D_{P,0} = \frac{1}{2} \dim \ker D_{\Sigma}$  and that proves im  $\iota_{\infty}$  is a Lagrangian subspace.

### 3.5 ACyl associative submanifolds from ACyl special Lagrangian 3-folds

Let  $(V, \omega, \Omega)$  be an ACyl Calabi-Yau 3-fold with asymptotic cross section  $(X, \omega_I, \omega_J, \omega_K)$  and  $(Y := S^1 \times V, \phi := d\theta \wedge \omega + \operatorname{Re}\Omega)$  be an ACyl  $G_2$ -manifold with asymptotic cross section  $(T^2 \times X, ds \wedge d\theta + \omega_K, (d\theta - ids) \wedge (\omega_I + i\omega_J))$  as in Remark 2.20. Let L be an ACyl Special Lagrangian 3-fold in V with cross section  $\Sigma_s := \{e^{is}\} \times \Sigma$ , where  $\Sigma$  is a K-holomorphic curve in X. Then for each  $\theta \in [0, 2\pi)$ ,  $\{e^{i\theta}\} \times L$  is an ACyl associative in  $Y = S^1 \times V$  with cross section  $\{e^{i\theta}\} \times \{e^{is}\} \times \Sigma$ . By abusing notation we denote them always by L and  $\Sigma$  respectively. This already has been mentioned in Example 3.25.

**Definition 3.66.** We define an isometry  $\Phi_L : C^{\infty}(NL) \to \Omega^0(L, \mathbf{R}) \oplus \Omega^1(L, \mathbf{R})$  by

$$\Phi_L(u) = (\langle \partial_{\theta}, u \rangle, -(\partial_{\theta} \times u)^{\flat}).$$

Let  $D_L$  be the Fueter operator defined in Definition 2.12. Define

$$\check{\mathbf{D}}_L:\Omega^0(L,\mathbf{R})\oplus\Omega^1(L,\mathbf{R})\to\Omega^0(L,\mathbf{R})\oplus\Omega^1(L,\mathbf{R})$$

by 
$$\check{\mathbf{D}}_L := \Phi_L \mathbf{D}_L \Phi_I^{-1}$$
.

**Proposition 3.67.** We have  $\Phi_L^{-1}(f, \sigma) = \partial_\theta \times \sigma_\sharp + f \partial_\theta$  and

$$\check{\mathbf{D}}_L = \begin{bmatrix} 0 & d^* \\ d & *d \end{bmatrix}, \quad \check{\mathbf{D}}_L^2 = \begin{bmatrix} \Delta_L^0 & 0 \\ 0 & \Delta_L^1 \end{bmatrix}$$

*Proof.* The first part is a simple direct computation. Now given a 1-form  $\sigma \in \Omega^1(L, \mathbf{R})$ , we observe that  $\nabla_L^{\perp}(\partial_{\theta} \times \sigma_{\sharp}) = \partial_{\theta} \times \nabla_L^{\parallel} \sigma^{\sharp}$ . Therefore,

$$\Phi_{L*}\nabla_L^{\perp}(f,\sigma) = \Phi_L\nabla_L^{\perp}\Phi_L^{-1}(f,\sigma) = (\nabla f,\nabla_L^{\parallel}\sigma).$$

Denote the Clifford multiplication for  $D_L$  by  $\gamma$ . We see that

$$\Phi_{L*}\gamma(v)(f,\sigma) = (\langle v, \sigma_{\mathsf{tl}} \rangle, fv^{\mathsf{b}} + (v \times \sigma_{\mathsf{tl}})^{\mathsf{b}}) = (\iota_v \sigma, fv^{\mathsf{b}} + *(v^{\mathsf{b}} \wedge \sigma)).$$

Hence,  $\check{\mathbf{D}}_L = \Phi_{L*} \gamma(\Phi_{L*} \nabla_L^{\perp})$  has to be the one in the statement of the proposition.

Definition 3.68. We define,

$$\Phi_{\Sigma}: \Omega^{0}(\Sigma, \mathbf{R}) \oplus \Omega^{0}(\Sigma, \mathbf{R}) \oplus C^{\infty}(N_{X}\Sigma) \to \Omega^{0}(\Sigma, \mathbf{R}) \oplus \Omega^{0}(\Sigma, \mathbf{R}) \oplus \Omega^{1}(\Sigma, \mathbf{R})$$

by  $\Phi_{\Sigma}(f_1, f_2, u) = (-f_2, -f_1, -\iota_u \omega_I)$ . Let  $\mathbf{D}_{\Sigma}$  be the Dirac operator defined in Definition 3.6. Define

$$\check{\mathbf{D}}_{\Sigma} := \Phi_{\Sigma} \mathbf{D}_{\Sigma} \Phi_{\Sigma}^{-1} = \begin{bmatrix} 0 & 0 & d_{\Sigma}^* \\ 0 & 0 & *d_{\Sigma} \\ d_{\Sigma} & -*d_{\Sigma} & 0 \end{bmatrix}.$$

**Proposition 3.69**. Assume the rate  $\mu < 0$  satisfies  $[\mu, 0) \cap \mathcal{D} = \emptyset$ . Then the map  $\Phi_L$  induces the following isomorphisms

$$\Phi_L : \ker \mathbf{D}_{L,u} \xrightarrow{\sim} \ker \{ H^1(L, \mathbf{R}) \to H^1(\Sigma, \mathbf{R}) \}$$

and

$$\Phi_L : \ker \mathbf{D}_{L,0} \xrightarrow{\sim} H^0(L,\mathbf{R}) \oplus H^1(L,\mathbf{R}).$$

Moreover, the following diagram commutes:

$$\begin{split} \ker \mathfrak{Q}_{L,0} & \xrightarrow{\Theta_L^C} & \ker \mathbf{D}_{L,0} & \xrightarrow{\Phi_L} & \ker \check{\mathbf{D}}_{L,0} \cong H^0(L,\mathbf{R}) \oplus H^1(L,\mathbf{R}) \\ \downarrow \iota_{L,0} & \iota_{L,0} \downarrow & & \downarrow i^* \oplus (0 \oplus i^*) \\ \ker \mathbf{D}_{\Sigma} & & \ker \mathbf{D}_{\Sigma} & \xrightarrow{\Phi_{\Sigma}} & \ker \check{\mathbf{D}}_{\Sigma} \cong H^0(\Sigma,\mathbf{R}) \oplus \big(H^0(\Sigma,\mathbf{R}) \oplus H^1(\Sigma,\mathbf{R})\big). \end{split}$$

Here  $\Theta^C_L: NL_C \to NL$  is the canonical bundle isomorphism as in Definition 3.48. The maps  $\iota_{L,0}$  are the asymptotic limit maps as in (3.39). The operators  $\mathbf{D}_{\lambda}$  are defined in Definition 3.58.

*Proof.* Set  $\Omega_{\lambda}^k(L, \mathbf{R}) := \{ \sigma \in \Omega^k(L, \mathbf{R}) : |\nabla^l \sigma| = O(e^{\lambda t}) \text{ as } t \to \infty, \forall l \in \mathbf{N} \cup \{0\} \}.$  First we prove that, the linear map

$$\Pi: \mathcal{H}_0^1 := \{ \sigma \in \Omega_0^1(L, \mathbf{R}) : d\sigma = 0, d^*\sigma = 0 \} \to H^1(L, \mathbf{R})$$

given by  $\sigma \mapsto [\sigma]$ , is an isomorphism. Let  $[\sigma] = 0$  for some  $\sigma \in \mathcal{H}_0^1$ . Then  $\sigma = dh$  for some smooth function  $h \in \Omega^0(L, \mathbf{R})$ . Since  $d^*\sigma = 0$  therefore  $\int_L |dh|^2 = 0$ . Hence  $\sigma = dh = 0$ , which shows  $\Pi$  is injective. Given  $[\sigma] \in H^1(L, \mathbf{R})$ , we denote the image of  $[\sigma]$  under the map  $H^1(L, \mathbf{R}) \to H^1(\Sigma, \mathbf{R})$  by  $[\sigma_{\Sigma}]$  with  $\sigma_{\Sigma}$  is harmonic. There exists  $\eta \in \Omega^1(L, \mathbf{R})$  such that  $d\eta = 0, d^*\eta \in \Omega^0_\lambda(L, \mathbf{R})$  with  $\lambda < 0$  sufficiently small,  $\Psi_L^* \Upsilon^* \eta = \pi^* \sigma_{\Sigma}$  and  $[\eta] = [\sigma]$ . Integrating by parts we get  $\int_L d^*\eta = 0$  and hence [Joyo4, Proposition 2.18] there exist  $\tilde{h} \in C^\infty_\lambda(L, \mathbf{R})$  and  $d^*\eta = \Delta_L h$ . Define  $\tilde{\eta} := \eta - d\tilde{h} \in \mathcal{H}_0^1$ . Then,  $[\tilde{\eta}] = [\sigma]$  and hence,  $\Pi$  is surjective. Moreover,  $\Psi_L^* \Upsilon^* \tilde{\eta} = \pi^* (\sigma_{\Sigma}) - \pi^* (dh)$ .

Now,  $(f, \sigma) \in \ker \check{\mathbf{D}}_{L,0}$  if and only if df = 0,  $d\sigma = 0$ ,  $d^*\sigma = 0$ . Therefore,

$$\ker \check{\mathbf{D}}_{L,0} \cong H^0(L,\mathbf{R}) \oplus H^1(L,\mathbf{R}).$$

The commutativity of the above diagram now follows from the following. Replacing L by the associative cylinder  $C = \mathbb{R} \times \Sigma$  we see  $NC = \pi^*(N_Z\Sigma)$  and  $N_Z\Sigma = \mathbb{R}^2 \oplus N_X\Sigma$  and

$$\Phi_C: \Omega^0(C, \mathbf{R}) \oplus \Omega^0(C, \mathbf{R}) \oplus C^{\infty}(\mathbf{R}, C^{\infty}(N_X\Sigma)) \to \Omega^0(C, \mathbf{R}) \oplus \Omega^0(C, \mathbf{R}) \oplus C^{\infty}(\mathbf{R}, \Omega^1(\Sigma, \mathbf{R}))$$

is given by  $\Phi_C(f_1, f_2, u) = (-f_2, -f_1, -\iota_u \omega_I)$ . Also,

$$\check{\mathbf{D}}_C := \Phi_C \mathbf{D}_C \Phi_C^{-1} = \begin{bmatrix} 0 & -\partial_t & d_{\Sigma}^* \\ \partial_t & 0 & *d_{\Sigma} \\ d_{\Sigma} & -*d_{\Sigma} & -*\partial_t \end{bmatrix} = J\partial_t + \check{\mathbf{D}}_{\Sigma}.$$

Finally, by Lemma 3.38 and the above commutative diagram we have

$$\ker \mathbf{D}_{L,u} = \ker \iota_{L,0} \cong \{0\} \oplus \ker \{H^1(L,\mathbf{R}) \to H^1(\Sigma,\mathbf{R})\}.$$

The deformation theory of ACyl special Lagrangian 3-folds was studied by Salur and Todd [ST10]. Combining results proved there with the Proposition 3.69 and Proposition 3.65 we get the following.

**Corollary 3.70.** In the above  $ACyl G_2$ -manifold  $Y = S^1 \times V$ , the following hold.

- i) The moduli space of ACyl associative submanifolds near a special Lagrangian 3-fold L with fixed asymptotic cross section  $\Sigma$  contains only ACyl special Lagrangians and is a smooth manifold of dimension  $b^2(L) + b^0(L) b^0(\Sigma)$ .
- ii) The moduli space of ACyl associative submanifolds near a special Lagrangian 3-fold L contains only ACyl special Lagrangians and is a smooth manifold of dimension  $b^1(L) + b^0(L) = \frac{1}{2}b^1(\Sigma) + b^0(L)$ . Moreover, if  $b^2(L) + b^0(L) b^0(\Sigma) = 0$ , then the asymptotic limit map  $\iota_{\infty}$  near L is a Lagrangian immersion.

# 3.6 ACyl associative submanifolds from ACyl holomorphic curves

Let  $(V, J, \omega, \Omega)$  be an ACyl Calabi-Yau 3-fold with asymptotic cross section  $(X, \omega_I, \omega_J, \omega_K)$  and  $Y := S^1 \times V$  be the ACyl  $G_2$ -manifold with asymptotic cross section  $T^2 \times X$  as in Remark 2.20. Let C be an ACyl holomorphic curve in V with asymptotic cross section  $\coprod_{j=1}^m S^1 \times \{x_j\} \subset S^1 \times X$ . Then  $P := S^1 \times C$  is an ACyl associative submanifold in  $Y = S^1 \times V$  with asymptotic cross section  $\Sigma := \coprod_{j=1}^m T^2 \times \{x_j\}$ . Moreover,  $C^\infty(NP) = C^\infty(S^1, C^\infty(NC))$  and by Proposition 3.15 (i),

$$\mathbf{D}_P = J\partial_\theta + \mathbf{D}_C.$$

We have also seen that, the isomorphism  $\gamma_C: \Omega^{0,1}(NC) \cong \overline{\mathrm{Hom}}(TC,NC) \to \Omega^0(NC)$  satisfies  $\gamma_C \circ \bar{\partial}_J^N = \mathbf{D}_C$ .

**Proposition 3.71.** Assume that the rate  $\mu < 0$  satisfies  $[\mu, 0) \cap \mathcal{D} = \emptyset$ . Then for all  $\lambda \leq 0$ ,  $\ker \mathbf{D}_{P,\lambda} = \ker \mathbf{D}_{C,\lambda}$ . Moreover, we have the following commutative diagram:

$$\ker \mathbf{D}_{P,\mu} \longrightarrow \ker \mathbf{D}_{P,0} \xrightarrow{\iota_{P,0}} \ker \mathbf{D}_{\Sigma}$$

$$\parallel \qquad \qquad \parallel \qquad \qquad \parallel$$

$$H^0_{\mu}(C,NC) \longrightarrow H^0_0(C,NC) \xrightarrow{\oplus_{j=1}^m \iota_{\infty}(x_j)} \oplus_{j=1}^m N_{x_j}C \cong \oplus_{j=1}^m T_{x_j}X.$$

Here,  $H^0_{\lambda}(C, NC) := \{ u \in C^{\infty}_{\lambda}(C, NC) : \bar{\partial}_I^N u = 0 \}.$ 

*Proof.* Since  $\mathbf{D}_C$  is *J*-antilinear, for all  $u \in C^{\infty}(S^1, C^{\infty}(NC))$  we have

$$\langle \mathbf{D}_C u, J \partial_{\theta} u \rangle_{L^2(NC)} = 0.$$

Let  $\lambda \leq 0$ . If  $u \in \ker \mathbf{D}_{P,\lambda}$ , then  $\|\mathbf{D}_C u\| = \|\partial_\theta u\| = 0$ . Hence,  $\ker \mathbf{D}_{P,\lambda} = \ker \bar{\partial}_I^N$ .

It remains to see the asymptotic limit map is  $\bigoplus_{j=1}^m \iota_\infty(x_j): H_0^0(C, NC) \to \bigoplus_{j=1}^m T_{x_j} X$ . Given,  $u \in \ker \mathbf{D}_{P,0}$ , by Lemma 3.38  $\iota_{P,0}(u) = (\chi_{T_0} \Upsilon^* u - v)_0$  for some  $v \in C_\lambda^\infty(S^1, C^\infty(NC))$  with  $\lambda < 0$ . Now,  $(\chi_{T_0} \Upsilon^* u - v)_0 = \bigoplus_{j=1}^m \iota_\infty(x_j) (\chi_{T_0} \Upsilon^* u - v)$  and hence this is equal to  $\bigoplus_{j=1}^m \iota_\infty(x_j)(u)$ .

**Proposition 3.72.** Let  $(Z, X, \omega)$  be a framed building block and  $V := Z \setminus X$  be the corresponding ACyl Calabi-Yau 3-fold. Let C be a holomorphic curve in Z intersecting X transversely at  $\bar{x} := \{x_1, x_2, ..., x_m\}$ . Then  $\hat{C} := C \setminus \bar{x}$  is an ACyl holomorphic curve in V with any rate  $\mu \ge -1$ . If  $\mu > -1$  then

$$H^0_{\mu}(\hat{C}, N\hat{C}) = H^0_{-1}(\hat{C}, N\hat{C}).$$

Moreover, there are isomorphisms

$$\Lambda_{\mu}: H^0_{\mu}(\hat{C}, N\hat{C}) \to H^0(C, NC(-\bar{x}))$$

and

$$\Lambda_0: H^0_0(\hat{C}, N\hat{C}) \to H^0(C, NC)$$

which satisfy the following commutative diagram:

$$\begin{split} H^0_{\mu}(\hat{C},N\hat{C}) & \longrightarrow H^0_0(\hat{C},N\hat{C}) & \stackrel{\oplus_{j=1}^m \iota_\infty(x_j)}{\longrightarrow} \oplus_{j=1}^m N_{x_j}C \cong \oplus_{j=1}^m T_{x_j}X \\ & \downarrow^{\Lambda_{\mu}} & \downarrow^{\Lambda_0} & \parallel \\ & H^0(C,NC(-\bar{x})) & \longrightarrow H^0(C,NC) & \stackrel{\oplus_{j=1}^m \operatorname{ev}_{x_j}}{\longrightarrow} \oplus_{j=1}^m N_{x_j}C \cong \oplus_{j=1}^m T_{x_j}X. \end{split}$$

Here,  $H^0(C, NC(-\bar{x}))$  is the space of holomorphic sections of the normal bundle NC which vanishes at each  $x_i$ .

*Proof.* In [HHN15], the diffeomorphism  $\Upsilon: (T, \infty) \times S^1 \times X \to V \backslash K_V$  for some compact submanifold with boundary  $K_V \subset V$  in the definition of ACyl  $G_2$ -manifold is given, by restricting a smooth embedding  $\tilde{\Upsilon}: \Delta \times X \to Z$ , (where  $\Delta$  is an open disc in C around 0) pre-composed by the biholomorphism  $(T, \infty) \times S^1 \times X \to \Delta \times X$  taking  $(t, z, x) \to (\frac{e^{-t}}{z}, x)$ . Denote the coordinate for  $\Delta$  by w. As C intersects X transversely at  $\bar{x} := \{x_1, x_2, ..., x_m\}$ , near each  $x_j$  it can be written as  $\tilde{\Upsilon}_*(\Gamma_{\tilde{\alpha}_j})$ , where  $\Gamma_{\tilde{\alpha}_j}$  is the graph of a map  $\tilde{\alpha}_j: \Delta \times \{x_j\} \to T_{x_j}X$  (by choosing  $\Delta$  sufficiently small disc) satisfying

$$|(\nabla^{\perp})^k \tilde{\alpha}_i| = O(|w|^{1-k})$$

for all  $j = 1, 2..., m, k \in \mathbb{N}_0$  as  $w \to 0$ . Therefore  $\hat{C} := C \setminus \bar{x}$  is an ACyl holomorphic curve in V with rate -1.

We will now prove that for each  $-1 \le \lambda \le 0$ , there is a canonical injective linear map

$$\Lambda_{\lambda}: H_{\lambda}^{0}(\hat{C}, N\hat{C}) \to H^{0}(C, NC(\lfloor \lambda \rfloor \bar{x})).$$

As  $\tilde{\Upsilon}^*J - J_0 = 0$  along  $\Delta \times \{x\}$  for all  $x \in X$  (see [HHN15, Appendix A]),  $\tilde{w} := \tilde{\Upsilon}_* w$  is a holomorphic function around  $\bar{x}$  in C. Now, given  $u \in H^0_\lambda(\hat{C}, N\hat{C})$ , we have  $\tilde{w}^{-\lfloor \lambda \rfloor}u$  is a bounded weak holomorphic section around  $\bar{x}$  in C and therefore it can be extended as a holomorphic section. Hence, u can be extended uniquely to get an element  $\Lambda_\lambda(u)$  in  $H^0(C, NC(\lfloor \lambda \rfloor \bar{x}))$ .

Moreover,  $\Lambda_{\lambda}$  is surjective if  $\lambda \in \mathbb{Z}^m$  or,  $(\lfloor \lambda \rfloor, \lambda] \cap \mathcal{D} = \emptyset$ . In our case, as the asymptotic cross section is the  $T_{\tilde{x}}^2$ , we have  $(-1, \mu] \cap \mathcal{D} = \emptyset$  (see Example 3.18). This completes the proof of the proposition.

**Example 3.73.** Let (Z, X) be a building block and  $\ell \cong \mathbf{P}^1$  be a rational curve in Z intersecting X transversely at  $\bar{x} := \{x_1, x_2, ..., x_m\}, m \ge 1$ . By Grothendieck's lemma

$$N\ell \cong \mathcal{O}_{\mathbf{P}^1}(k_1) \oplus \mathcal{O}_{\mathbf{P}^1}(k_2).$$

As,  $\ell \cdot c_1(Z) = \ell \cdot [X] = m$ , we have  $k_1 + k_2 = \ell \cdot c_1(N\ell) = \ell \cdot ([X] - c_1(T\ell)) = m - 2$ . Thus for some  $k \in \mathbb{Z}$ ,

$$N\ell \cong \mathcal{O}_{\mathbf{P}^1}(k) \oplus \mathcal{O}_{\mathbf{P}^1}(m-k-2).$$

Moreover,

$$\begin{split} H^{1}(\ell,N\ell) &= H^{1}(\mathbf{P}^{1}, \mathcal{O}_{\mathbf{P}^{1}}(k)) \oplus H^{1}(\mathbf{P}^{1}, \mathcal{O}_{\mathbf{P}^{1}}(m-k-2)) \\ &= H^{0}(\mathbf{P}^{1}, \mathcal{O}_{\mathbf{P}^{1}}(-k-2)) \oplus H^{0}(\mathbf{P}^{1}, \mathcal{O}_{\mathbf{P}^{1}}(k-m)) \ \ \text{(by Serre duality)}, \end{split}$$

and

$$H^{0}(\ell, N\ell(-\bar{x})) = H^{0}(\ell, N\ell \otimes \mathcal{O}_{\mathbf{P}^{1}}(-m)) = H^{0}(\mathbf{P}^{1}, \mathcal{O}_{\mathbf{P}^{1}}(k-m)) \oplus H^{0}(\mathbf{P}^{1}, \mathcal{O}_{\mathbf{P}^{1}}(-k-2)).$$

Thus we obtain the following:

a)  $H^0(\ell, N\ell(-\bar{x})) \cong H^1(\ell, N\ell) = 0$  if and only if  $-1 \le k \le m-1$ . In this case, the moduli space of ACyl associative submanifolds around  $S^1 \times (\ell \setminus \underline{x})$  contains only product of  $S^1$  with ACyl holomorphic curves. Furthermore, it is a smooth manifold of dimension 2m and the asymptotic limit map  $\iota_{\infty}$  is a Lagrangian immersion.

b) 
$$H^0(\ell, N\ell) \neq 0$$
, as  $m \ge 1$ .

*Remark* 3.74. In the above example if we force  $H^0(\ell, N\ell) = 0$ , then  $\ell$  does not intersect X and  $S^1 \times \ell$  will be closed rigid associative in  $S^1 \times V$  and  $N\ell \cong \mathcal{O}_{\mathbf{P}^1}(-1) \oplus \mathcal{O}_{\mathbf{P}^1}(-1)$ .

# 4 Gluing ACyl associative submanifolds in ACyl $G_2$ -manifolds

This section is about gluing ACyl associative submanifolds in ACyl  $G_2$ -manifolds. Here we prove the main gluing theorem, Theorem 1.3. As an application of Theorem 1.3, we prove Theorem 1.4 and Proposition 1.5 about associative submanifolds by gluing ACyl holomorphic curves with the help of Section 3.6. As another application of Theorem 1.3, we prove Theorem 1.6 and Proposition 1.8 about associative submanifolds by gluing ACyl special Lagrangian 3-folds with the help of Section 3.5.

#### 4.1 Pregluing construction and estimates

Let  $(Y_{\pm}, \phi_{\pm})$  be a matching pair of ACyl  $G_2$ -manifolds of rates  $\nu_{\pm} < 0$  with a matching map  $f: Z_{+} \to Z_{-}$  and let  $\{(Y_{T}, \phi_{T}): T \geq T_{0}\}$  be the family of generalized connected sum  $G_2$ -manifolds. Let  $P_{\pm}$  be a pair of ACyl associative submanifolds in  $Y_{\pm}$  with asymptotic cross sections  $\Sigma_{\pm}$  and rates  $\mu_{\pm}$  (see Definition 3.20) satisfying Hypothesis 1.1.

Assumption 4.1. With out loss of generality we will always assume in the following that  $P_{\pm}$  have rates  $\mu_{\pm} \in (\nu_{\pm}, 0)$  such that

$$[\mu_{\pm},0)\cap \mathcal{D}_{C_{+}}=\varnothing.$$

Notation 4.2. We denote

$$\mu := \max\{\mu_+, \mu_-\} < 0, \quad \nu := \max\{\nu_+, \nu_-\} < 0.$$

For all  $T \ge T_0$ , we construct approximate associative submanifolds  $P_T$  in  $Y_T$  as follows. We will carry over the notations  $\Upsilon_{\pm}$  from Definition 2.17, and  $P_{C,\pm}$ ,  $K_{P_{C,\pm}}$  from Definition 3.44.

**Definition 4.3.** We define a closed 3-dimensional submanifold  $P_{T,C}$  of  $Y_T$  by

$$P_{T,C} := P_{T,C,+} \bigcup_F P_{T,C,-}$$

where  $P_{T,C,\pm} := K_{P_{C,\pm}} \cup \Upsilon_{\pm}((T_0, T+1] \times \Sigma_{\pm})$ . Here the identification map  $F : [T, T+1] \times Z_+ \to [T, T+1] \times Z_-$  is given by F(t, z) = (2T - t + 1, f(z)). The normal bundle of  $P_{T,C}$  is

$$NP_{T,C} = NP_{T,C,+} \bigcup_F NP_{T,C,-}.$$

The tubular neighbourhood map is defined by

$$\Upsilon_{P_{T,C}} := \Upsilon_{P_{T,C,+}} \bigcup_{F} \Upsilon_{P_{T,C,-}} : V_{P_{T,C}} \to U_{P_{T,C}}$$

where  $\Upsilon_{P_{T,C,\pm}}$  is the restriction of the ECyl tubular neighbourhood maps  $\Upsilon_{P_{C,\pm}}$  (see Definition 3.43) on  $P_{T,C,\pm}$ . Similarly,

$$\Upsilon_T := \Upsilon_{T,+} \bigcup_F \Upsilon_{T,-}$$

where  $\Upsilon_{T,\pm}$  is the restriction of  $\Upsilon_{\pm}$  on  $Y_{T,\pm}$ .

Let  $\beta_{\pm} \in C^{\infty}(NP_{C,\pm})$  represent  $P_{\pm}$  as in Definition 3.45 such that  $\Upsilon \circ \Upsilon_{P_{C,\pm}}(\beta_{\pm})$  is  $P_{\pm}$ . We define the approximate associative  $P_T$  by

$$P_T := \Upsilon_T \circ \Upsilon_{P_{T,C}}(\beta_T),$$

where  $\beta_T := (1 - \chi_{T-1})\beta_{\pm} \in C^{\infty}(NP_{T,C,\pm})$ . Here  $\chi$  is the cut-off function as in (2.31). Finally there is also a canonical bundle isomorphism

$$\Theta^{C}_{P_{T}}:NP_{T,C}\rightarrow NP_{T},\ \Theta^{C}_{P_{T}}:=\Theta^{C}_{P_{+}}\bigcup_{F}\Theta^{C}_{P_{-}},$$

where  $\Theta_{P_{+}}^{C}$  are as in Definition 3.48.




Figure 6: Pregluing construction: approximate associative  $P_T$ .

Note that all  $P_T$  are diffeomorphic to the connected sum  $P_+\#_f P_-$  along the cross sections  $\Sigma_\pm$  on the ends of  $P_\pm$ . From the definition of  $P_T$  we see that it may not be associative in  $(Y_T, \phi_T)$ , so we deform this to construct an associative  $\tilde{P}_T$ .

**Definition 4.4.** Define  $\mathfrak{F}_T: C^{\infty}(V_{P_{T,C}}) \to C^{\infty}(NP_{T,C})$  as follows: for all  $u \in C^{\infty}(V_{P_{T,C}})$  and  $w \in C^{\infty}(NP_{T,C})$ ,

$$\langle \mathfrak{F}_T(u), w \rangle_{L^2} := \int_{\Gamma_{\mathcal{U}}} \iota_w \Upsilon_{P_{T,C}}^* \psi_T.$$

The notation w in the integrand is the extension vector field of  $w \in C^{\infty}(NP_{T,C})$  in the tubular neighbourhood as in Notation 2.16. The  $L^2$  inner product we choose here is coming from the canonical bundle isomorphism  $\Theta_{P_T}^C: NP_{T,C} \to NP_T$  as in Definition 4.3 and the induced metric on  $NP_T$  of  $g_{\phi_T}$ .

**Definition 4.5.** The linearization of  $\mathfrak{F}_T$  at  $\beta_T \in C^{\infty} \in (V_{P_{T,C}})$  is denoted by  $\mathfrak{L}_T$ , that is,

$$\mathfrak{L}_T := d\mathfrak{F}_{T|_{\beta_T}} : C^{\infty}(NP_{T,C}) \to C^{\infty}(NP_{T,C}).$$

**Proposition 4.6.** The linearization  $\mathfrak{L}_T$  is given by the following: for all  $v \in C^{\infty}(NP_{T,C})$  and  $w \in C^{\infty}(NP_{T,C})$ ,

$$\langle \mathfrak{L}_{T}(v), w \rangle = \int_{\Gamma_{\beta_{T}}} \iota_{w} \mathcal{L}_{v}(\Upsilon_{P_{T,C}}^{*} \psi_{T}) = \int_{\Gamma_{\beta_{T}}} \langle \sum_{\substack{cyclic \\ permutations}} [e_{2}, e_{3}, \nabla_{e_{1}} v], w \rangle + \iota_{\nabla_{w} v}(\Upsilon_{P_{T,C}}^{*} \psi_{T}).$$

where  $\{e_1, e_2, e_3\}$  is a local oriented orthonormal frame for  $T\Gamma_{\beta_T}$  and by abusing notation we are denoting  $\Upsilon_{P_{T,C}}^* \nabla$  by  $\nabla$  and the associator  $\Upsilon_{P_{T,C}}^* [\cdot, \cdot, \cdot]$  by  $[\cdot, \cdot, \cdot]$  which are induced from the 3-form  $\Upsilon_{P_{T,C}}^* \phi_T$ .

Moreover,  $\mathfrak{L}_T$  is a formally self adjoint elliptic operator for all sufficiently large  $T\geqslant T_0.$ 

*Proof.* The first part is exactly same as Proposition 3.52. It remains to prove that  $\mathfrak{L}_T$  is a formally self adjoint elliptic operator for all sufficiently large  $T \geq T_0$ . We denote the restrictions of  $\mathfrak{L}_T$  and  $\beta_T$  over  $P_{T,C,\pm}$  by  $\mathfrak{L}_{T,\pm}$  and  $\beta_{T,\pm}$  respectively. As  $|\phi_T - \phi_{\pm}| = O(e^{-\delta T})$  with  $\delta > 0$  by Theorem 2.32 we have,

$$\mathfrak{L}_{T,\pm}=d\mathfrak{F}_{0|\beta_{T,\pm}}+O(e^{-\delta T}).$$

By Definition 3.45, we have  $|\nabla^k(\beta_T - \beta_{\pm})| = O(e^{\mu T})$ ,  $k \in \mathbb{N}_0$  and therefore using Proposition 3.61 we get

$$d\mathfrak{F}_{0|\beta_{T,\pm}}=\mathfrak{L}_{P_{\pm}}+O(e^{\mu T}).$$

Thus

(4.7) 
$$\mathfrak{L}_{T,\pm} = \mathfrak{L}_{P_{\pm}} + O(e^{\mu T}) + O(e^{-\delta T})$$

Therefore for all sufficiently large  $T \ge T_0$ ,  $\mathfrak{L}_T$  is an elliptic operator.

Similar to Proposition 3.54 we obtain that for all  $v \in C^{\infty}(NP_{T,C})$  and  $w \in C^{\infty}(NP_{T,C})$ ,

$$\langle \mathfrak{L}_T v, w \rangle_{L^2} - \langle w, \mathfrak{L}_T v \rangle_{L^2} = \int_{\Gamma_{\beta_T}} \iota_w \iota_v (\Upsilon_{P_{T,C}}^* d\psi_T) = 0$$

and hence  $\mathfrak{L}_T$  is formally self adjoint.

**Definition 4.8.** The nonlinear map  $\mathfrak{F}_T: C^\infty(V_{P_{T,C}}) \to C^\infty(NP_{T,C})$  can be written as

$$\mathfrak{F}_T := \mathfrak{L}_T + Q_T + e_T,$$

where 
$$e_T := \mathfrak{F}_T(\beta_T) \in C^{\infty}(NP_{T,C})$$
. Note that  $Q_T(\beta_T) = -\mathfrak{L}_T(\beta_T)$ .

**Notation 4.9.** We will always write the Hölder spaces  $C^{k,\gamma}(NP_{T,C})$  by  $C^{k,\gamma}$  only with out writing the T-dependency but it will be clear from the context that which T we are using. Similarly we will also write for other Banach spaces, for example  $L^2$  spaces or Sobolev spaces without T.

**Proposition 4.10 (Schauder estimate).** There exists a constant C > 0 independent of T such that for all  $u \in C^{k+1,\gamma}$  we have

$$||u||_{C^{k+1,\gamma}} \leq C(||\Omega_T u||_{C^{k,\gamma}} + ||u||_{L^{\infty}}).$$

*Proof.* For  $u \in C^{k+1,\gamma}$ , define  $u_{\pm} \in C^{k+1,\gamma}(NP_{T,C,\pm})$  by restricting u over  $P_{T,C,\pm}$ . Using interior Schauder estimates for  $\mathfrak{L}_{P_{+}}$  applied to  $u_{\pm}$  we get,

$$||u||_{C^{k+1,\gamma}} \lesssim ||\Omega_{P_+}u_+||_{C^{k,\gamma}} + ||\Omega_{P_-}u_-||_{C^{k,\gamma}} + ||u||_{L^{\infty}}.$$

Similar to (4.7) we obtain

$$\|\mathfrak{Q}_{T}u_{\pm} - \mathfrak{Q}_{P_{\pm}}u_{\pm}\|_{C^{k,\gamma}} \lesssim \|u_{\pm}\|_{C^{k+1,\gamma}} \|\beta_{T} - \beta_{\pm}\|_{C^{k+1,\gamma}} + O(e^{-\delta T}) \|u\|_{C^{k+1,\gamma}} \lesssim (O(e^{-\delta T}) + O(e^{\mu T})) \|u\|_{C^{k+1,\gamma}}.$$

Hence, we get the required estimate. Here  $\delta$  is the constant from Theorem 2.32.

**Proposition 4.11** (Error estimate). For all  $T \ge T_0$  we have,

$$\|\mathfrak{F}_T(\beta_T)\|_{C^{k,\gamma}} = \|e_T\|_{C^{k,\gamma}} \lesssim O(e^{-\delta_e T}),$$

where  $\delta_e := \min\{\delta, -\mu\}$  and  $\delta$  is the constant from Theorem 2.32.

*Proof.* As  $P_{\pm}$  are associative submanifolds,  $\mathfrak{F}_T(\beta_T) = 0$  over  $P_{T-1,C,\pm} \subset P_{T,C}$ . Denote,  $A_{T,\pm} := P_{T,C,\pm} \setminus P_{T-1,C,\pm}$ . By Theorem 2.32, we have

$$|\Upsilon_{P_{T,C,+}}^* \psi_T - \Upsilon_* \psi_0| = O(e^{-\delta T}) + O(e^{\nu_{\pm} T})$$

where  $\delta > 0$ . Therefore for all  $w \in C^{\infty}(NP_{T,C})$ , we have

$$\langle \mathfrak{F}_{T}(\beta_{T}), w \rangle_{L^{2}} \leq (O(e^{-\delta T}) + O(e^{\nu T})) \|w\|_{L^{2}} + \int_{A_{T}} \iota_{w} \Upsilon_{*} \psi_{0} + \int_{A_{T}} \iota_{w} \Upsilon_{*} \psi_{0}.$$

Thus,

$$|\mathfrak{F}_T(\beta_T)| \lesssim O(e^{-\delta T}) + O(e^{\nu T}) + \max \|\beta_T - \beta_{\pm}\|_{C^1(A_{T+})} \lesssim O(e^{-\delta_e T}).$$

Estimates for higher derivatives and Hölder norms of  $\mathfrak{F}_T(\beta_T)$  also hold similarlly.

**Proposition 4.12 (Quadratic estimate).** For all sufficiently large  $T \ge T_0$ ,  $u, v \in C^{\infty}(V_{P_{T,C}})$  and  $\eta \in C^{\infty}(NP_{T,C})$  we have,

(i) 
$$|d\mathfrak{F}_{T|_{\mathcal{U}}}(\eta) - d\mathfrak{F}_{T|_{\mathcal{U}}}(\eta)| \lesssim (|u-v| + |\nabla^{\perp}(u-v)|)(|\eta| + |\nabla^{\perp}\eta|),$$

(ii) 
$$||Q_T(u) - Q_T(v)||_{C^{k,\gamma}} \lesssim ||u - v||_{C^{k+1,\gamma}} (||u - \beta_T||_{C^{k+1,\gamma}} + ||v - \beta_T||_{C^{k+1,\gamma}}).$$

*Proof.* The proofs are exactly like in Proposition 3.61, Proposition 3.62. An important observation here is that one can express  $\mathcal{L}_u \mathcal{L}_v \psi_T$  formally as a quadratic polynomial

$$\mathcal{L}_{u}\mathcal{L}_{v}\psi_{T} = O(f_{1,T}) \cdot u \cdot v + O(f_{2,T}) \cdot (u \cdot \nabla^{\perp}v + v \cdot \nabla^{\perp}u) + \psi_{T} \cdot \nabla^{\perp}u \cdot \nabla^{\perp}v$$

where  $O(f_{1,T}) = O(1)$  and  $O(f_{2,T}) = O(1)$ . Alternatively, we can prove (i) just like in the proof of (4.7) with the estimates from Proposition 3.61.

Finally to see (ii) we write  $Q_T(u) - Q_T(v)$  as

$$\int_0^1 dQ_{T_{|tu+(1-t)v}}(u-v)dt = \int_0^1 \left( d\mathfrak{F}_{T_{|tu+(1-t)v}}(u-v) - d\mathfrak{F}_{T_{|\beta_T}}(u-v) \right) dt$$

and (ii) follows from (i).

**Definition 4.13.** Denote the asymptotic limit maps  $\iota_{P_{\pm},0}$  of Lemma 3.38 by

$$\iota_{\infty}^{\pm}: \ker \mathfrak{L}_{P_{\pm},0} \to \ker \mathbf{D}_{\Sigma_{\pm}}.$$

It follows from Lemma 3.38 that  $\ker \iota_{\infty}^{\pm} = \ker \mathfrak{L}_{P_{\pm},\mu_{\pm}}$ . We define the following spaces which are useful to formulate Proposition 4.14:

(i) the matching kernel  $K^{\mathfrak{m}}$  by

$$\mathcal{K}^{\mathfrak{m}} := \{(u_+, u_-) \in \ker \mathfrak{L}_{P_+,0} \times \ker \mathfrak{L}_{P_-,0} : f_* \iota_{\infty}^+ u_+ = \iota_{\infty}^- u_-\},$$

(ii) the approximate kernel of  $\mathfrak{L}_T$  by

$$\mathcal{K}_{T}^{\mathfrak{m}} := \{ u_{+} \#_{T} u_{-} \in C^{\infty}(NP_{T,C}) : (u_{+}, u_{-}) \in \mathcal{K}^{\mathfrak{m}} \}$$

where  $u_+\#_T u_-$  over  $P_{T,C,\pm}$  is

$$u_+ \#_T u_- := u_\pm - \chi_{T-1} (u_\pm - u_\mp),$$

(iii) 
$$\mathcal{X}_{T}^{k+1,\gamma} := \{ u \in C^{k+1,\gamma} : \langle u, \xi \rangle_{L^{2}(K_{P_{\pm}})} = 0, \forall \xi \in \mathcal{K}_{T}^{\mathfrak{m}} \}.$$

**Proposition 4.14 (Linear estimate).** For all  $\Lambda > 0$  there exists  $T_{\Lambda} \ge T_0$  such that for all  $T \ge T_{\Lambda}$  and  $u \in \mathcal{X}_T^{k+1,\gamma}$ , we have

$$||u||_{C^{k+1,\gamma}} \lesssim e^{\Lambda T} ||\mathfrak{L}_T u||_{C^{k,\gamma}}.$$

*Proof.* By the Schauder estimate, Proposition 4.10 we see it is enough to prove that for all sufficiently large  $T \ge T_0$  and for all  $u \in \mathcal{X}_T^{k+1,\gamma}$ , we have

$$||u||_{L^{\infty}} \lesssim e^{\Lambda T} ||\Omega_T u||_{C^{k,\gamma}}$$

We will prove this by contradiction. Suppose this is not true, then there exists an increasing sequence  $T_n \ge T_0$  which tends to  $\infty$  as  $n \to \infty$  and  $u_n$  in  $\mathcal{X}_{T_n}^{k+1,\gamma}$  such that

$$||u_n||_{L^\infty} = 1$$
,  $e^{\Lambda T_n} ||\mathfrak{Q}_{T_n} u_n||_{C^{k,\gamma}} \to 0$  as  $n \to \infty$ .

Define  $u_{n,\pm} \in C^{k+1,\gamma}(NP_{T_n,C,\pm}) =: C^{k+1,\gamma}_{P_{T_n,C,\pm}}$  by restricting  $u_n$  over  $P_{T_n,C,\pm}$ . By (4.7) and Proposition 3.33 we see that  $\|u_n\|_{C^{k+1,\gamma}}$  is bounded and hence  $\|u_{n,\pm}\|_{C^{k+1,\gamma}}$  is also bounded. Therefore the Arzelà-Ascoli theorem implies that there exists a subsequence which we call again  $u_n$ , and there exists  $u_\pm$  in  $C^{k+1,\frac{\gamma}{2}}_{P_\pm}$  such that  $\mathfrak{L}_{P_\pm}u_\pm=0$  and  $u_{n,\pm}\to u_\pm$  in  $C^{k+1,\frac{\gamma}{2}}_{P_\pm,\mathrm{loc}}$ . Moreover, by elliptic regularity of Proposition 3.33 we get,  $u_\pm\in C^{k+1,\gamma}_{P_\pm}$ . By taking further subsequences if necessary we claim that as  $n\to\infty$ ,

$$(4.15) ||u_{n,\pm} - u_{\pm}||_{L^{\infty}(NP_{T_n,C,\pm})} \leq ||u_{n,\pm} - u_{\pm}||_{C^{k+1,\gamma}_{P_{T_n,C,\pm}}} \to 0.$$

To prove this claim we argue as follows. Define,  $\tilde{u}_{n,\pm} := (1 - \chi_{\frac{3}{2}T_n})u_n \in C_{P_{\pm}}^{k+1,\gamma}$ . Again by (4.7), we obtain that for any sufficiently small  $0 < s < \Lambda$  we have

$$\begin{split} \|\mathfrak{Q}_{P_{\pm}}\tilde{u}_{n,\pm}\|_{C^{k,\gamma}_{P_{\pm},s}} & \lesssim e^{-(\delta+s)T_n} \|u_n\|_{C^{k+1,\gamma}} + e^{(\mu-s)T_n} \|u_n\|_{C^{k+1,\gamma}} + \|\mathfrak{Q}_{T_n}\tilde{u}_{n,\pm}\|_{C^{k,\gamma}} \\ & \lesssim (e^{-(\delta+s)T_n} + e^{(\mu-s)T_n} + e^{-\frac{3}{2}sT_n}) \|u_n\|_{C^{k+1,\gamma}} + \|\mathfrak{Q}_{T_n}u_n\|_{C^{k,\gamma}} \\ & \lesssim e^{-(\delta+s)T_n} + e^{(\mu-s)T_n} + e^{-\frac{3}{2}sT_n} + e^{-\Lambda T_n}. \end{split}$$

Then there exists  $v_{n,\pm} \in \ker \mathfrak{L}_{P_{\pm},s} = \ker \mathfrak{L}_{P_{\pm},0}$  such that

$$\|\tilde{u}_{n,\pm} - v_{n,\pm}\|_{C^{1,\gamma}_{p_{+,s}}} \lesssim e^{-(\delta+s)T_n} + e^{(\mu-s)T_n} + e^{-\frac{3}{2}sT_n} + e^{-\Lambda T_n}.$$

and hence,  $\|u_{n,\pm}-v_{n,\pm}\|_{C^{k+1,\gamma}_{P_{T_n,C,\pm}}}\lesssim e^{-\delta T_n}+e^{\mu T_n}+e^{-\frac{s}{2}T_n}+e^{(s-\Lambda)T_n}\to 0 \quad n\to\infty.$  As,  $\ker\mathfrak{L}_{P_\pm,0}$  is finite dimensional, the norms  $\|\cdot\|_{C^{k+1,\frac{\gamma}{2}}_{K_{P_\pm}}}$  and  $\|\cdot\|_{C^{k+1,\gamma}_{P_\pm}}$  are equivalent on it. Taking further subsequence yields  $\|v_{n,\pm}-u_\pm\|_{C^{k+1,\frac{\gamma}{2}}_{K_{P_\pm}}}\to 0$  and hence as  $n\to\infty$ ,  $\|v_{n,\pm}-u_\pm\|_{C^{k+1,\gamma}_{P_\pm}}\to 0$ . This proves (4.15).

This will also further imply that  $f_*\iota_\infty^+ u_+ = \iota_\infty^- u_-$ , that is  $(u_+, u_-) \in \mathcal{K}^{\mathfrak{m}}$ . Indeed, for all  $z \in \Sigma_+$ ,

$$f_*\iota_{\infty}^+ u_+(z) = \lim_{n \to \infty} F_* u_{n,+}(T_n, z) = \lim_{n \to \infty} u_{n,-}(T_n, f(z)) = \iota_{\infty}^- u_-(f(z)).$$

As,  $u_n \in \mathfrak{X}_{T_n}^{k+1,\gamma}$ , we have

$$||u_{\pm}||_{L^{2}(K_{P_{\pm}})} = \langle u_{\pm} - u_{n_{\pm}}, u_{\pm} \rangle_{L^{2}(K_{P_{\pm}})} \leq ||u_{n_{\pm}} - u_{\pm}||_{L^{\infty}(K_{P_{\pm}})} ||u_{\pm}||_{L^{1}(K_{P_{\pm}})} \to 0.$$

Therefore  $u_{\pm} = 0$ , which is a contradiction because as  $n \to \infty$ ,

$$1 = ||u_n||_{L^{\infty}} \le ||u_{n,+}||_{L^{\infty}(NP_{T_n,C,+})} + ||u_{n,-}||_{L^{\infty}(NP_{T_n,C,-})} \to 0.$$

### 4.2 Proof of the gluing theorem

Let  $P_{\pm}$  be a pair of ACyl associative submanifolds in  $Y_{\pm}$  with asymptotic cross sections  $\Sigma_{\pm}$  and rates  $\mu_{\pm}$ . We also have Assumption 4.1 about the rates. To prove the gluing theorem we solve the nonlinear PDE,  $\mathfrak{F}_T u = 0$  for u very close to  $\beta_T$  by using the following Lemma 4.16, which is an application of Banach contraction principle.

**Lemma 4.16.** Let X, Y be two Banach spaces and let  $V \subset X$  be a neighbourhood of  $0 \in X$ . Let  $x_0 \in V$ . Let  $F: V \to Y$  be a smooth map of the form

$$F(x) = L(x) + Q(x) + F(x_0)$$

such that:

- $L: \mathcal{X} \to \mathcal{Y}$  is a linear bounded invertible operator and there exists a constant  $c_L > 0$  such that for all  $x \in \mathcal{X}$ ,  $||x||_{\mathcal{X}} \le c_L ||Lx||_{\mathcal{Y}}$ .
- $Q: V \to \mathcal{Y}$  is a smooth map and there exists a constant  $c_O > 0$  such that for all  $x_1, x_2 \in V$ ,

$$||Q(x_1) - Q(x_2)||_{\mathcal{Y}} \le c_O ||x_1 - x_2||_{\mathcal{X}} (||x_1 - x_0||_{\mathcal{X}} + ||x_2 - x_0||_{\mathcal{X}}).$$

If  $||F(x_0)||_{\mathcal{Y}} \leq \frac{1}{10c_L^2c_Q}$  and  $B(x_0, \frac{1}{5c_Lc_Q}) \subset V$ , then there exists an unique  $x \in \mathcal{X}$  with  $||x-x_0||_{\mathcal{X}} \leq \frac{1}{5c_Lc_Q}$  solving F(x) = 0.

*Proof.* Denote  $\varepsilon:=\frac{1}{5c_Lc_O}$  and define,  $\varphi:B(0,\varepsilon)\subset \mathfrak{X}\to B(0,\varepsilon)$  by

$$\varphi(x) = -L^{-1}(F(x_0) + Q(x_0 + x)) - x_0.$$

Therefore,  $\varphi(x) = x$  if and only if  $L(x_0 + x) + Q(x_0 + x) + F(x_0) = 0$ . It is a simple computation which shows that, for all  $x, x_1, x_2 \in B(0, \varepsilon)$  we have

$$\|\varphi(x)\|_{\mathcal{X}} \leqslant \frac{7}{10}\varepsilon < \varepsilon$$
, and  $\|\varphi(x_1) - \varphi(x_2)\|_{\mathcal{X}} \leqslant \frac{1}{2}\|x_1 - x_2\|_{\mathcal{X}}$ .

Apply Banach's contraction principle to get the required conclusion.

**Proof of Theorem 1.3.** The nonlinear map  $\mathfrak{F}_T: C^{k+1,\gamma}(V_{P_{T,C}}) \to C^{k,\gamma}(NP_{T,C})$  can be written as

$$\mathfrak{F}_T = \mathfrak{Q}_T + O_T + \mathfrak{F}_T(\beta_T).$$

The Hypothesis 1.1 implies that the matching kernel  $\mathfrak{K}^{\mathfrak{m}}=0$ . Therefore by Proposition 4.14, there exists  $T'_0\geqslant T_0$  such that for all  $T\geqslant T'_0$ , we have  $\mathfrak{L}_T:C^{k+1,\gamma}(NP_{T,C})\to C^{k,\gamma}(NP_{T,C})$  is an invertible operator and there exists a constant  $c_{\mathfrak{L}_T}=O(e^{\frac{\delta_e}{4}T})>0$  such that for all  $u\in C^{k+1,\gamma}$ , we have

$$||u||_{C^{k+1,\gamma}} \leq c_{\mathfrak{L}_T} ||\mathfrak{L}_T u||_{C^{k,\gamma}}.$$

By Proposition 4.12 there exists a constant  $c_{Q_T} = O(1) > 0$  such that for all  $u, v \in C^{k+1,\gamma}(V_{P_{T,C}})$  we have,

$$||Q_T(u) - Q_T(v)||_{C^{k,\gamma}} \le c_{O_T} ||u - v||_{C^{k+1,\gamma}} (||u - \beta_T||_{C^{k+1,\gamma}} + ||v - \beta_T||_{C^{k+1,\gamma}}).$$

Therefore by Proposition 4.11, there exists  $T_0'' \ge T_0'$  such that for all  $T \ge T_0''$ , we have

$$\|\mathfrak{F}_T(\beta_T)\|_{C^{k,\gamma}} = O(e^{-\delta_e T}) \leqslant \frac{1}{10c_{\mathfrak{D}_T}^2 c_{Q_T}}$$

and hence by Lemma 4.16, there exists a unique family  $\{\tilde{\beta}_T \in C^{k+1,\gamma}(V_{P_{T,C}}) : T \geqslant T_0''\}$  such that  $\mathfrak{F}_T(\tilde{\beta}_T) = 0$  and

(4.17) 
$$\|\tilde{\beta}_T - \beta_T\|_{C^{k+1,\gamma}} = O(e^{-\frac{\delta_e}{4}T}).$$

Thus we obtain a family of closed associative submanifolds  $\tilde{P}_T := \Upsilon_{P_{T,C}}(\tilde{\beta}_T)$ .

It remains to prove that  $\tilde{P}_T$  is unobstructed, that is  $d\mathfrak{F}_{T|\tilde{P}_T}$  is invertible. We choose  $\Lambda := \frac{\delta_e}{8}$ . Then by Proposition 4.14, there exists  $T_{\Lambda} > 0$  such that for all  $T \ge T_{\Lambda}$ ,  $\|\mathfrak{L}_T^{-1}\| = O(e^{\Lambda T})$ . Now by Proposition 4.12 and (4.17),

$$\|\mathfrak{L}_{T}^{-1}d\mathfrak{F}_{T_{|\tilde{\beta}_{T}}}-\mathbf{1}\| \leq \|\mathfrak{L}_{T}^{-1}\|\|d\mathfrak{F}_{T_{|\tilde{\beta}_{T}}}-\mathfrak{L}_{T}\| = O(e^{(\Lambda-\frac{\delta_{e}}{4})T}) = O(e^{-\frac{\delta_{e}}{8}T}).$$

Therefore by defining  $T_1 := \max\{T_{\Lambda}, T_0^{\prime\prime}\}$ , we obtain the required theorem.

*Remark* 4.18. If the approximate associative  $P_T$  is  $\Upsilon_{P_{T,C}}(\beta_T)$ , then  $\tilde{P}_T$  can be taken as  $\Upsilon_{P_{T,C}}(\tilde{\beta}_T)$  such that

$$\|\tilde{\beta}_T - \beta_T\|_{C^{k+1,\gamma}} = O(e^{-\frac{\delta_e}{4}T}).$$

## 4.3 Associative submanifolds by gluing ACyl holomorphic curves

**Proof of Theorem 1.4.** The theorem follows from Theorem 1.3 with the help of Proposition 3.71 and Proposition 3.72.

To prove Proposition 1.5 we need the following lemma (for a proof, see [MNE21, Lemma 2.5]).

**Lemma 4.19.** Let W be a semi-Fano 3-fold and  $X = X_{\infty} \in |-K_W|$  be a smooth anti-canonical K3-surface such that  $-K_{W|_X}$  is very ample. Let  $x \in X$  and  $(y,v) \in P(TX)$  be such that x and y are distinct. Then there is an anti-canonical K3-surface  $X_0$  such that the base locus B of the pencil  $|X_0:X_{\infty}|$  is smooth and

$$x \notin B$$
,  $y \in B$  and  $T_uB = v$ .

**Proof of Proposition 1.5.** Suppose  $\ell_+ \cap X_+ = \{x_+\}$ . Define  $x_- := \mathfrak{r}(x_+) \in X_-$ . Choose  $v \in \mathbf{P}(T_{x_-}X_-)$  such that  $v \cap \operatorname{im} \mathfrak{r}_* \operatorname{ev}_+ = \{0\}$ . By Lemma 4.19 we can choose an anti-canonical pencil in  $W_-$  so that the base locus  $B_-$  is smooth,  $x_- \in B_-$  and  $T_{x_-}B_- = v$ . The building block  $Z_-$  is defined now by blowing up  $W_-$  along  $B_-$ . Consider the unobstructed line  $\ell_- := \{x_-\} \times \mathbf{P}^1 \subset Z_-$ . Then im  $\operatorname{ev}_- = v \in \mathbf{P}(T_{x_-}X_-)$ . Hence by Theorem 1.4 we obtain the required proposition.

**Example 4.20.** Proposition 1.5 produces lot of rigid associative 3-spheres as follows.

- (1) We apply Proposition 1.5 to all very ample Fano 3-folds  $W_{\pm}$  that have been used in [CHNP15] to obtain twisted connected sum  $G_2$  manifolds. Let  $Z_{\pm}$  be a building block that comes from  $W_{\pm}$ . More explicitly,  $\pi: Z_{\pm} \to W_{\pm}$  is a blow-up of  $W_{\pm}$  along a base locus  $B_{\pm}$  as in Theorem 2.25. Then for every  $x \in B_{\pm}$  we can use the unobstructed holomorphic line  $\pi^{-1}(x)$  in Proposition 1.5 to obtain rigid associative 3-sphere.
- (2) Šokurov [Šok79] proved that every Fano 3-fold W of index 1 except  $\mathbf{P}^1 \times \mathbf{P}^2$  contains a line  $\ell$  with  $-K_W \cdot \ell = 1$ . Any line in a general Fano 3-fold of index 1 and Picard rank 1 is unobstructed [IP99, Proposition 4.2.2 and Theorem 4.2.7]. In the Mori-Mukai list there are 97 deformation types of Fano 3-folds with very ample anti-canonical bundle and 8 of them are of index 1 and Picard rank 1 namely,

$$\#_3^1, \#_4^1, \#_5^1, \#_6^1, \#_7^1, \#_8^1, \#_9^1, \#_{10}^1$$

where  $\#_n^{\rho}$  is the *n*-th Fano 3-fold with Picard rank  $\rho$  in the Mori-Mukai list (see [IP99, Appendix 12.2]). The Fano 3-folds with not very ample anti-canonical bundle [IP99, Theorem 2.4.5, Theorem 2.1.16] are

$$\#_1^1, \#_2^1, \#_{12}^1, \#_1^2, \#_2^2, \#_3^2, \#_1^7, \#_1^8$$

Therefore we are able to take 8 and 97 Fano 3-folds as  $W_+$  and  $W_-$  respectively to apply Proposition 1.5 (as,  $\operatorname{rk}\operatorname{Pic}(W_+) + \operatorname{rk}\operatorname{Pic}(W_-) \le 11$ ). In this way we obtain  $8 \times 97 = 776$  twisted connected sum  $G_2$  manifolds containing a rigid associative 3-sphere.

Remark 4.21. This list of examples is not exhaustive. The reader may find more examples by applying Proposition 1.5 to many other Fano 3-folds or more generally to semi-Fano 3-folds.

## 4.4 Associative submanifolds by gluing ACyl special Lagrangian 3-folds

**Proof of Theorem 1.6.** Since  $\mathfrak{r}(\Sigma_+) = \Sigma_-$ , therefore by definition of f we have  $f(\Sigma_{s_+}) = \Sigma_{s_-}$ . By Proposition 3.69 we see that  $\iota_{\infty}^{\pm}$  is injective if and only if  $i_{\pm}^* : H^1(L_{\pm}, \mathbb{R}) \to H^1(\Sigma_{\pm}, \mathbb{R})$  is injective which is equivalent to  $b^2(L_{\pm}) - b^0(\Sigma_{\pm}) + b^0(L_{\pm}) = 0$ . Again by Proposition 3.69 we have,

$$\Phi_{\Sigma_-}\mathfrak{r}_*\Phi_{\Sigma_+}^{-1}=(\mathfrak{r}^{-1})^*K_+.$$

Therefore,  $\operatorname{im}(f_*\iota_{\infty}^+) \cap \operatorname{im} \iota_{\infty}^- = \{0\}$  if and only if  $\operatorname{im}(K_+ \circ i_+^*) \cap \operatorname{im} (\mathfrak{r}^* \circ i_-^*) = \{0\}.$ 

**Proof of Proposition 1.8.** By [Kov13, Proposition 5.2],  $V_+ := Z_+ \setminus X_+$  admits an ACyl Calabi-Yau structure  $(\omega_+, \Omega_+)$  such that  $\sigma_+$  is an anti-holomorphic involutive isometry on  $V_+$ . Now by definition we have  $\rho_+^*(\omega_{I,+}) = \omega_{I,+}$  and  $\rho_+^*(\omega_{J,+} + i\omega_{K,+}) = -\omega_{J,+} - i\omega_{K,+}$ . Also  $\sigma_+^*(\omega_{I,+}) = -\omega_{I,+}$  and  $\sigma_+^*(\omega_{J,+} + i\omega_{K,+}) = -\omega_{J,+} + i\omega_{K,+}$ . We can now easily check that  $\rho_-$  is a non-symplectic involution on  $X_-$ . Since  $\frac{\Sigma_+}{\langle \rho_+ \rangle}$  is a deformation retract of  $L_-$ , therefore  $b^1(L_-) = 0$ . Thus the pair of ACyl special Lagrangian  $L_\pm$  satisfies the hypotheses of Theorem 1.6.

**Example 4.22** (Nordström [Nor13]). Let  $W_+$  be the Fano 3-fold in  $\mathbb{CP}^4$  defined by the quartic polynomial

$$Q(z_0, z_1, z_2, z_3, z_4) = -z_0^4 + z_1^4 + z_2^4 + z_3^4 + z_4^4 = 0.$$

Then  $X_{\infty} := \{z_4 = 0\}$  and  $X_0 := \{z_0 = 0\}$  are anti-canonical divisors in  $W_+$ . The base locus B of the anti-canonical pencil  $|X_0, X_{\infty}|$  is  $\{z_0 = 0, z_4 = 0\}$ . Let  $Z_+$  be the blow-up of  $W_+$  along B and let  $X_+$  be the proper transform of  $X_{\infty}$  as in Theorem 2.25. Then  $(Z_+, X_+)$  is a building block. As the complex conjugation in  $\mathbb{CP}^4$  acts on  $X_{\infty}$  and B, it induces an anti-holomorphic involution  $\sigma_+$  on  $Z_+$  which acts also on  $X_+$  by [Kov13, pg. 19]. The involution  $\rho_{\infty} : X_{\infty} \to X_{\infty}$  defined by

$$\rho_{\infty}(z_0, z_1, z_2, z_3, 0) = (-z_0, z_1, z_2, z_3, 0)$$

induces a non-symplectic involution  $\rho_+$  on  $X_+$  which commutes with  $\sigma_{+|X_+}$  acts freely on  $\Sigma_+ := \operatorname{Fix}_{\sigma_+}(X_+)$ . Since  $\operatorname{Fix}_{\sigma_+}(W_+)$  disjoint from B therefore  $\operatorname{Fix}_{\sigma_+}(Z_+)$  is homeomorphic to  $W_+ \cap \mathbf{RP}^4 \cong S^3$  and  $\Sigma_+ := \operatorname{Fix}_{\sigma_+}(X_+)$  is homeomorphic to  $X_{\infty} \cap \mathbf{RP}^4 \cong S^2$ . The involution  $\rho_{\infty}$  acts on  $S^2$  as an antipodal map. Thus  $\operatorname{Fix}_{\sigma_+}(Z_+) \setminus \Sigma_+$  is a disjoint union of two 3-balls. Let  $L_+$  be any one of these 3-balls. Applying Proposition 1.8 we obtain a family of closed associative submanifolds  $P_T$  in  $Y_T$ , each of which is diffeomorphic to  $\mathbf{RP}^3$ .

Example 4.23. Let  $X_{\infty}$  be the K3-surface with non-symplectic involution  $\rho_{\infty}$  as in Example 4.22. Let  $W_+ := \frac{\operatorname{CP}^1 \times X_{\infty}}{\langle \iota \times \rho_{\infty} \rangle}$  and let  $Z_+$  be the blow-up of  $W_+$  as in Theorem 2.28. Let  $X_+$  be the proper transform of  $\{\infty\} \times X_{\infty}$  as in Theorem 2.28. Then  $(Z_+, X_+)$  is a building block. As the complex conjugation in  $\operatorname{CP}^3$  acts on  $X_{\infty}$  and  $\operatorname{Fix}_{\rho_{\infty}}(X_{\infty})$ , it induces an anti-holomorphic involution  $\sigma_+$  on  $Z_+$  which acts also on  $X_+$  by  $[\operatorname{Kov13}, \operatorname{pg}. \operatorname{19}]$ . The non-symplectic involution  $\rho_{\infty}$  on  $X_+$  commutes with  $\sigma_{+|X_+}$  and acts freely on  $\Sigma_+ := \operatorname{Fix}_{\sigma_+}(X_+) \cong S^2$ . The involution  $\rho_{\infty}$  acts on  $S^2$  as an antipodal map. Let  $\tilde{L}_+$  be the proper transform of  $\frac{\operatorname{RP}^1 \times \Sigma_{\infty}}{\langle \iota \times \rho_{\infty} \rangle}$ , where  $\Sigma_{\infty} := \operatorname{Fix}_{\sigma_{\infty}}(X_{\infty})$ . Applying Proposition 1.8 we obtain a family of closed associative submanifolds  $P_T$  in  $Y_T$ , each of which is diffeomorphic to  $\operatorname{RP}^3 \# \operatorname{RP}^3$ .

**Example 4.24**. Let  $X_{\infty} \subset \mathbf{CP}(1,1,1,3)$  be the *K*3-surface defined by the polynomial

$$P(z_0, z_1, z_2, z_3) = z_3^2 - z_0^6 - z_1^6 - z_2^6 = 0$$

which is a double cover of  $\mathbb{CP}^2$  branched along the curve  $\{z_0^6 + z_1^6 + z_2^6 = 0\}$ . The non-symplectic involution on  $X_\infty$  taking  $(z_0, z_1, z_2, z_3) \to (z_0, z_1, z_2, -z_3)$ . Let  $W_+ := \frac{\mathbb{CP}^1 \times X_\infty}{\langle \iota \times \rho_\infty \rangle}$  and let  $Z_+$  be the blow-up of  $W_+$  as in Theorem 2.28. Let  $X_+$  be the proper transform of  $\{\infty\} \times X_\infty$  as in Theorem 2.28. Then  $(Z_+, X_+)$  is a building block. The complex conjugation in  $\mathbb{CP}(1, 1, 1, 3)$  induces an anti-holomorphic involution  $\sigma_+$  on  $Z_+$  which acts also on  $X_+$  by [Kov13, pg. 19]. The non-symplectic involution  $\rho_\infty$  on  $X_+$  commutes with  $\sigma_+|_{X_+}$  and acts freely on  $\Sigma_+ := \mathrm{Fix}_{\sigma_+}(X_+)$ . Let  $\tilde{L}_+$  be the proper transform of  $\frac{\mathrm{RP}^1 \times \Sigma_\infty}{\langle \iota \times \rho_\infty \rangle}$ , where  $\Sigma_\infty := \mathrm{Fix}_{\sigma_\infty}(X_\infty)$ . Under the homeomorphism  $\mathbb{RP}(1,1,1,3) \to \mathbb{RP}^4$  taking  $(x_0,x_1,x_2,x_3) \to (x_0^3,x_1^3,x_2^3,x_3)$  we see that  $\Sigma_+ \cong S^2$  and the involution  $\rho_\infty$  acts on  $S^2$  as an antipodal map. Applying Proposition 1.8 we obtain a family of closed associative submanifolds  $P_T$  in  $Y_T$ , each of which is diffeomorphic to  $\mathbb{RP}^3 \# \mathbb{RP}^3$ .

Remark 4.25. This list of examples is not exhaustive. The reader may find more examples by applying Proposition 1.8 to *K*3-surfaces having commuting non-symplectic involutions and anti-holomorphic involutions, studied by Nikulin and Saito [NSo5; NSo7].

# A Lemma for Quadratic estimate

**Lemma A.1.** Let M be a 3-dimensional oriented submanifold of a  $G_2$ -manifold  $(Y, \phi)$ . Let  $\Upsilon_M : V_M \subset NM \to Y$  be a tubular neighbourhood map. There exists a constant C > 0 such that if  $w \in C^{\infty}(NM)$  and  $u, v, s \in C^{\infty}(V_M)$  then over  $\Gamma_s := \operatorname{graph} s \subset V_M$  we have<sup>5</sup>

$$|\iota_w \mathcal{L}_u \mathcal{L}_v \psi| \lesssim |w| \Big( f_1 |u| |v| + f_2 \big( |u| |\nabla v| + |v| |\nabla u| \big) + |\nabla u| |\nabla v| |\psi| \Big),$$

where  $f_1 := |\psi| |\nabla B| + |\psi| |B|^2 + |R| |\psi|$  and  $f_2 := |B| |\psi|$ .

Here R is the Riemann curvature tensor and  $B: C^{\infty}(NM) \times C^{\infty}(NM) \to C^{\infty}(TV_M)$  is defined by  $B(u,v) := \nabla_u v$ 

*Proof.* Let  $\{e_1, e_2, e_3\}$  be a local oriented orthonormal frame for  $T\Gamma_s$ . Then

$$\mathcal{L}_v \psi(w, e_1, e_2, e_3) = \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \psi(w, \nabla_{e_1} v, e_2, e_3) + \psi(\nabla_w v, e_1, e_2, e_3),$$

and

$$\mathcal{L}_{u}\mathcal{L}_{v}\psi(w,e_{1},e_{2},e_{3})=u\big(\mathcal{L}_{v}\psi(w,e_{1},e_{2},e_{3})\big)-\sum_{\substack{\text{cyclic}\\ \text{permutations}}}\mathcal{L}_{v}\psi(w,[u,e_{1}],e_{2},e_{3}).$$

Now,

$$u(\psi(w, \nabla_{e_1} v, e_2, e_3)) = \psi(w, \nabla_u \nabla_{e_1} v, e_2, e_3) + \psi(w, \nabla_{e_1} v, \nabla_u e_2, e_3) + \psi(w, \nabla_{e_1} v, e_2, \nabla_u e_3) + \psi(\nabla_w u, \nabla_{e_1} v, e_2, e_3).$$

Furthermore,

$$u(\psi(\nabla_w v, e_1, e_2, e_3)) = \sum_{\substack{\text{cyclic} \\ \text{permutations}}} \psi(\nabla_w v, \nabla_u e_1, e_2, e_3) + \psi(\nabla_w \nabla_u v + R(u, w)v, e_1, e_2, e_3),$$

and

$$\begin{split} \mathcal{L}_v \psi(w, [u, e_1], e_2, e_3) &= \psi(w, \nabla_{[u, e_1]} v, e_2, e_3) + \psi(\nabla_w v, [u, e_1], e_2, e_3) \\ &+ \psi(w, [u, e_1], \nabla_{e_2} v, e_3) + \psi(w, [u, e_1], e_2, \nabla_{e_3} v). \end{split}$$

Finally, putting together after some cancellations we obtain

$$\begin{split} |\mathcal{L}_{u}\mathcal{L}_{v}\psi(w,e_{1},e_{2},e_{3})| &\lesssim |\psi| \big( |\nabla v| |\nabla_{w}u| + |\nabla_{w}v| |\nabla u| + |\nabla_{w}\nabla_{u}v| \big) \\ &+ |w| |\psi| \Big( \sum_{i=1}^{3} |\nabla_{e_{i}}(\nabla_{u}v)| + |R||u||v| + |\nabla u||\nabla v| \Big). \end{split}$$

<sup>&</sup>lt;sup>5</sup>By abusing notation we denote again by u, v, w the extension of the vector field u, v, w over NM as in Notation 2.16. We also denote  $\Upsilon_M^* \psi$  by  $\psi$ , where  $\psi = *\phi$ . The notation  $\mathfrak L$  stands for the Lie derivative.

Observe that,

$$|\nabla_{e_i}(\nabla_u v)| \lesssim |\nabla B||u||v| + |B||u||\nabla v| + |B||v||\nabla u|,$$
  
$$|\nabla_w(\nabla_u v)| \lesssim |w||u||v|(|\nabla B| + |B|^2),$$

Therefore,

$$|\iota_{w}\mathcal{L}_{u}\mathcal{L}_{v}\psi| \lesssim f_{1}|w||u||v| + f_{2}|w|(|u||\nabla^{\perp}v| + |v||\nabla^{\perp}u|) + |w||\nabla^{\perp}u||\nabla^{\perp}v||\psi|.$$

# References

- [Ati88] M. Atiyah. New invariants of 3– and 4–dimensional manifolds. The mathematical heritage of Hermann Weyl (Durham, NC, 1987). Proceedings of Symposia in Pure Mathematics 48. Providence, RI, 1988. MR: 974342 (cit. on p. 8)
- [Ber55] M. Berger. Sur les groupes d'holonomie homogène des variétés à connexion affine et des variétés riemanniennes. Bulletin de la Société Mathématique de France 83 (1955), pp. 279–330. MR: 0079806. Zbl: 0068.36002 (cit. on p. 1)
- [BDHLMS18] A. P. Braun, M. Del Zotto, J. Halverson, M. Larfors, D. R. Morrison, and S. Schäfer-Nameki. *Infinitely many M2-instanton corrections to M-theory on G2-manifolds. Journal of High Energy Physics* 2018.9 (2018). DOI: 10.1007/jhepo9(2018)077 (cit. on p. 2)
- [BS89] R. L. Bryant and S. M. Salamon. On the construction of some complete metrics with exceptional holonomy. Duke Mathematical Journal 58.3 (1989), pp. 829–850. DOI: 10.1215/S0012-7094-89-05839-0. MR: 1016448 (cit. on p. 10)
- [Bry87] R. L. Bryant. Metrics with exceptional holonomy. Annals of Mathematics 126.3 (1987), pp. 525–576. DOI: 10.2307/1971360. MR: 916718. Zbl: 0637.53042 (cit. on p. 10)
- [CG71] J. Cheeger and D. Gromoll. The splitting theorem for manifolds of nonnegative Ricci curvature. Journal of Differential Geometry 6 (1971), pp. 119–128. MR: 0303460. Zbl: 0223.53033. (a) (cit. on p. 13)
- [CHNP13] A. Corti, M. Haskins, J. Nordström, and T. Pacini. Asymptotically cylindrical Calabi–Yau 3–folds from weak Fano 3–folds. Geometry and Topology 17.4 (2013), pp. 1955–2059. DOI: 10.2140/gt.2013.17.1955. MR: 3109862. Zbl: 1273.14081 (cit. on p. 14)
- [CHNP15] A. Corti, M. Haskins, J. Nordström, and T. Pacini.  $G_2$ -manifolds and associative submanifolds via semi-Fano 3-folds. Duke Mathematical Journal 164.10 (2015), pp. 1971–2092. DOI: 10.1215/00127094-3120743. MR: 3369307. Zbl: 06486366 (cit. on pp. 1, 2, 10–12, 14, 16, 52)
- [DW19] A. Doan and T. Walpuski. On counting associative submanifolds and Seiberg-Witten monopoles. Pure and Applied Mathematics Quarterly 15.4 (2019), pp. 1047–1133. DOI: 10.4310/PAMQ.2019.V15.n4.a4. arXiv: 1712.08383. (cit. on pp. 2, 11)

- [Dono2] S. K. Donaldson. Floer homology groups in Yang-Mills theory. Cambridge Tracts in Mathematics 147. With the assistance of M. Furuta and D. Kotschick. Cambridge: Cambridge University Press, 2002, pp. viii+236. DOI: 10.1017/CBO9780511543098. MR: 1883043. Zbl: 0998.53057 (cit. on pp. 29, 30)
- [DK90] S. K. Donaldson and P. B. Kronheimer. *The geometry of four-manifolds*. Oxford Mathematical Monographs. New York, 1990. MR: MR1079726. Zbl: 0904.57001 (cit. on pp. 6, 20, 38)
- [DS11] S. K. Donaldson and E. P. Segal. *Gauge theory in higher dimensions, II. Surveys in differential geometry.* Vol. 16. Geometry of special holonomy and related topics. International Press, 2011, pp. 1–41. arXiv: 0902.3239. MR: 2893675. Zbl: 1256.53038 (cit. on p. 2)
- [DT98] S. K. Donaldson and R. P. Thomas. Gauge theory in higher dimensions. The geometric universe. Oxford University Press, 1998, pp. 31–47. MR: 1634503. Zbl: 0926.58003. (cit. on p. 2)
- [Don17] Simon Donaldson. Adiabatic Limits of Co-associative Kovalev-Lefschetz Fibrations. Algebra, Geometry, and Physics in the 21st Century. Springer International Publishing, 2017, pp. 1–29. DOI: 10.1007/978-3-319-59939-7\_1. (cit. on p. 26)
- [DS22] Simon Donaldson and Christopher Scaduto. Associative submanifolds and gradient cycles. Surveys in differential geometry 2019. Differential geometry, Calabi-Yau theory, and general relativity. Part 2. Vol. 24. Surv. Differ. Geom. Int. Press, Boston, MA, [2022] ©2022, pp. 39–65. (cit. on p. 26)
- [DPW23] Shubham Dwivedi, Daniel Platt, and Thomas Walpuski. Associative Submanifolds in Joyce's Generalised Kummer Constructions. Communications in Mathematical Physics 401.3 (Aug. 2023), pp. 2327–2353. DOI: 10.1007/s00220-023-04716-7. (cit. on p. 10)
- [FG82] M. Fernández and A. Gray. *Riemannian manifolds with structure group G*<sub>2</sub>. *Ann. Mat. Pura Appl.* (4) 132 (1982), 19–45 (1983). DOI: 10.1007/BF01760975. MR: 696037. Zbl: 0524.53023 (cit. on p. 9)
- [Gra69] A. Gray. Vector cross products on manifolds. Transactions of the American Mathematical Society 141 (1969), pp. 465–504. MR: 0243469. Zbl: 0182.24603 (cit. on p. 9)
- [Har90] F. R. Harvey. *Spinors and calibrations*. Vol. 9. Perspectives in Mathematics. Boston, MA: Academic Press Inc., 1990, pp. xiv+323 (cit. on p. 9)
- [HL82] R. Harvey and H. B. Lawson Jr. *Calibrated geometries. Acta Math.* 148 (1982), pp. 47–157. DOI: 10.1007/BF02392726. MR: MR6666108. Zbl: 0584.53021 (cit. on pp. 2, 10)
- [HHN15] M. Haskins, H.-J. Hein, and J. Nordström. Asymptotically cylindrical Calabi–Yau manifolds. Journal of Differential Geometry 101.2 (2015), pp. 213–265. DOI: 10.4310/jdg/1442364651. MR: 3399097. Zbl: 1332.32028 (cit. on pp. 5, 14, 43)

- [Hit74] N. Hitchin. *Harmonic spinors. Advances in Math.* 14 (1974), pp. 1–55. MR: 0358873 (cit. on p. 9)
- [IP99] V. A. Iskovskih and Yu. G. Prokhorov. Fano varieties. Algebraic geometry, V.
   Vol. 47. Encyclopaedia Math. Sci. Springer, 1999, pp. 1–247. MR: 1668579
   (cit. on pp. 14, 52)
- [Joy96a] D. D. Joyce. *Compact* 8-manifolds with holonomy Spin(7). *Invent. Math.* 123.3 (1996), pp. 507–552. DOI: 10.1007/s002220050039. MR: MR1383960. Zbl: 0858.53037 (cit. on p. 10)
- [Joy96b] D. D. Joyce. Compact Riemannian 7-manifolds with holonomy  $G_2$ . I. Journal of Differential Geometry 43.2 (1996), pp. 291-328. DOI: 10.4310/jdg/1214458109. MR: MR1424428. Zbl: 0861.53022 (cit. on p. 1)
- [Joy96c] D. D. Joyce. Compact Riemannian 7-manifolds with holonomy  $G_2$ . II. Journal of Differential Geometry 43.2 (1996), pp. 329-375. DOI: 10.4310/jdg/1214458110. MR: MR1424428. Zbl: 0861.53023 (cit. on p. 10)
- [Joyo4] D. D. Joyce. Special Lagrangian submanifolds with isolated conical singularities. I. Regularity. Annals of Global Analysis and Geometry 25.3 (2004), pp. 201–251. DOI: 10.1023/B:AGAG.0000023229.72953.57. MR: 2053761 (cit. on p. 41)
- [Joy18] D. D. Joyce. Conjectures on counting associative 3–folds in  $G_2$ –manifolds. Modern geometry: a celebration of the work of Simon Donaldson. Proceedings of Symposia in Pure Mathematics 99. American Mathematical Society, 2018, pp. 97–160. DOI: 10.1090/pspum/099/01739. arXiv: 1610.09836. MR: 3838881. Zbl: 1448.53061 (cit. on pp. 2, 11)
- [JK17] D.D. Joyce and S. Karigiannis. A new construction of compact torsion-free  $G_2$ -manifolds by gluing families of Eguchi-Hanson spaces. Journal of Differential Geometry 117.2 (2017), pp. 255-343. DOI: 10.4310/jdg/1612975017. arXiv: 1707.09325. Zbl: 1464.53067. (cit. on p. 10)
- [JSo5] Dominic Joyce and Sema Salur. Deformations of asymptotically cylindrical coassociative submanifolds with fixed boundary. Geom. Topol. 9 (2005), pp. 1115–1146. DOI: 10.2140/gt.2005.9.1115. arXiv: math/0408137. (cit. on p. 3)
- [Joy07] Dominic D. Joyce. Riemannian holonomy groups and calibrated geometry. Vol. 12. Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, 2007, pp. x+303. (cit. on pp. 2, 9, 20, 25, 38)
- [KLL20] Spiro Karigiannis, Naichung Conan Leung, and Jason D. Lotay, eds. *Lectures and Surveys on G2-Manifolds and Related Topics*. Springer US, 2020. DOI: 10.1007/978-1-0716-0577-6. (cit. on p. 9)
- [Kovo3] A. Kovalev. Twisted connected sums and special Riemannian holonomy. Journal für die Reine und Angewandte Mathematik 565 (2003), pp. 125–160. DOI: 10.1515/crll.2003.097. MR: MR2024648. Zbl: 1043.53041 (cit. on pp. 1, 10, 15)

- [KL11] A. Kovalev and N.-H. Lee. K3 surfaces with non-symplectic involution and compact irreducible  $G_2$ -manifolds. Mathematical Proceedings of the Cambridge Philosophical Society 151.2 (2011), pp. 193–218. DOI: 10.1017/S030500411100003X. MR: 2823130. Zbl: 1228.53064 (cit. on pp. 1, 10, 15)
- [KN10] A. Kovalev and J. Nordström. Asymptotically cylindrical 7-manifolds of holonomy  $G_2$  with applications to compact irreducible  $G_2$ -manifolds. Annals of Global Analysis and Geometry 38.3 (2010), pp. 221–257. DOI: 10.1007/s10455-010-9210-8. MR: 2721660 (cit. on p. 13)
- [Kov13] Alexei Kovalev. *Asymptotically cylindrical manifolds with holonomy Spin(7). I* (Sept. 2013). arXiv: 1309.5027. (cit. on pp. 25, 52, 53)
- [LM89] H. B. Lawson Jr. and M.-L. Michelsohn. *Spin geometry*. Vol. 38. Princeton Mathematical Series. Princeton, NJ: Princeton University Press, 1989, pp. xii+427. MR: 1031992. Zbl: 0688.57001 (cit. on p. 9)
- [LM85] R. B. Lockhart and R. C. McOwen. Elliptic differential operators on noncompact manifolds. Annali della Scuola Normale Superiore di Pisa 12.3 (1985), pp. 409–447. MR: 837256. Zbl: 0615.58048. (cit. on pp. 21, 23, 26, 30)
- [Maro2] S. Marshall. Deformations of special Lagrangian submanifolds. DPhil. Thesis. University of Oxford, 2002. (cit. on pp. 23, 26, 28, 29)
- [MP78] V. G. Maz'ya and B. A. Plamenevskiĭ. Estimates in  $L_p$  and in Hölder classes, and the Miranda–Agmon maximum principle for the solutions of elliptic boundary value problems in domains with singular points on the boundary. Mathematische Nachrichten 81 (1978), pp. 25–82. MR: 0492821 (cit. on p. 29)
- [MS12] D. McDuff and D. Salamon. *J-holomorphic curves and symplectic topology*. Second. Vol. 52. American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 2012, pp. xiv+726. MR: 2954391. Zbl: 1272.53002 (cit. on p. 18)
- [McL98] R. C. McLean. Deformations of calibrated submanifolds. Communications in Analysis and Geometry 6.4 (1998), pp. 705–747. DOI: 10.4310/CAG.1998.v6.n4.a4. MR: 1664890. Zbl: 0929.53027 (cit. on pp. 11, 21)
- [MNE21] G. Menet, J. Nordström, and H. N. Sá Earp. Construction of  $G_2$ -instantons via twisted connected sums. Mathematical Research Letters 28.2 (2021), pp. 471–509. DOI: 10.4310/mrl.2021.v28.n2.a6. (cit. on p. 51)
- [NSo5] Viacheslav V. Nikulin and Sachiko Saito. Real K3 surfaces with non-symplectic involution and applications. Proc. London Math. Soc. (3) 90.3 (2005), pp. 591–654. DOI: 10.1112/S0024611505015212. (cit. on pp. 26, 53)
- [NSo7] Viacheslav V. Nikulin and Sachiko Saito. Real K3 surfaces with non-symplectic involution and applications. II. Proc. Lond. Math. Soc. (3) 95.1 (2007), pp. 20–48. DOI: 10.1112/plms/pdlo23. (cit. on pp. 26, 53)

- [Noro8a] J. Nordström. Deformations of asymptotically cylindrical  $G_2$ -manifolds. Mathematical Proceedings of the Cambridge Philosophical Society 145.2 (2008), pp. 311–348. DOI: 10.1017/S0305004108001333. MR: MR2442130 (cit. on p. 13)
- [Nor13] J. Nordström. *Desingularising intersecting associatives.* unpublished. 2013 (cit. on p. 52)
- [Noro8b] Johannes Nordström. Deformations and gluing of asymptotically cylindrical manifolds with exceptional holonomy. en (2008). DOI: 10.17863/CAM.16204. (cit. on p. 25)
- [Ohs15] Matthias Ohst. Deformations of Asymptotically Cylindrical Cayley Submanifolds (June 2015). arXiv: 1506.00110. (cit. on p. 3)
- [SW15] H. N. Sá Earp and T. Walpuski.  $G_2$ -instantons over twisted connected sums. Geometry and Topology 19.3 (2015), pp. 1263–1285. DOI: 10.2140/gt.2015.19.1263. arXiv: 1310.7933. MR: 3352236. Zbl: 06441803. ⓐ (cit. on pp. 1, 2, 8)
- [SW17] D. A. Salamon and T. Walpuski. Notes on the octonions. Proceedings of the 23rd Gökova Geometry-Topology Conference. 2017, pp. 1-85. arXiv: 1005.2820. MR: 3676083. Zbl: 06810387. (cit. on p. 9)
- [Salo6] Sema Salur. Asymptotically cylindrical Ricci-flat manifolds. Proc. Amer. Math. Soc. 134.10 (2006), pp. 3049–3056. DOI: 10.1090/S0002-9939-06-08313-4. (cit. on p. 13)
- [ST10] Sema Salur and A. J. Todd. Deformations of asymptotically cylindrical special Lagrangian submanifolds. Proceedings of the Gökova Geometry-Topology Conference 2009. Int. Press, Somerville, MA, 2010, pp. 99–123. (cit. on pp. 3, 41)
- [Šok79] V. V. Šokurov. *The existence of a line on Fano varieties. Izv. Akad. Nauk SSSR Ser. Mat.* 43.4 (1979), pp. 922–964, 968. MR: 548510 (cit. on p. 52)
- [Tal17] Tim Talbot. Gluing and deformations of asymptotically cylindrical special Lagrangians (Sept. 2017). arXiv: 1709.09564. (cit. on p. 25)
- [Tiaoo] G. Tian. Gauge theory and calibrated geometry. I. Annals of Mathematics 151.1 (2000), pp. 193–268. DOI: 10.2307/121116. arXiv: math/0010015. MR: MR1745014. Zbl: 0957.58013 (cit. on p. 2)