

Analysis and Numerical Approximation of Stationary Second-Order Mean Field Game Partial Differential Inclusions*

Yohance A. P. Osborne and Iain Smears

September 5, 2022

Abstract

The formulation of Mean Field Games (MFG) typically requires continuous differentiability of the Hamiltonian in order to determine the advective term in the Kolmogorov–Fokker–Planck equation for the density of players. However in many cases of practical interest, the underlying optimal control problem may exhibit bang-bang controls, which typically lead to nondifferentiable Hamiltonians. We develop the analysis and numerical analysis of stationary MFG for the general case of convex, Lipschitz, but possibly nondifferentiable Hamiltonians. In particular, we propose a generalization of the MFG system as a Partial Differential Inclusion (PDI) based on interpreting the derivative of the Hamiltonian in terms of sub-differentials of convex functions. We establish existence of a weak solution to the MFG PDI system, and we further prove uniqueness under a similar monotonicity condition to the one considered by Lasry and Lions. We then propose a monotone finite element discretization of the problem, and we prove strong H^1 -norm convergence of the approximations to the value function and strong L^q -norm convergence of the approximations of the density function. We illustrate the performance of the numerical method in numerical experiments featuring nonsmooth solutions.

Key words: Mean Field Games, Hamilton–Jacobi–Bellman equations, Nondifferentiable Hamiltonians, Partial Differential Inclusions, Monotone finite element method, Convergence analysis

AMS subject classifications: 65N15, 65N30

1 Introduction

Mean Field Games (MFG), as introduced by Lasry and Lions [12, 13, 14] and independently by Huang, Caines & Malhamé [11], consider the asymptotic be-

*University College London, Department of Mathematics, 25 Gordon Street, London, WC1H 0AY, UK; yohance.osborne.16@ucl.ac.uk, i.smears@ucl.ac.uk

behaviour of rational stochastic differential games as the number of players approaches infinity. Under suitable assumptions, the system of equations consists of a Hamilton–Jacobi–Bellman (HJB) equation for the value function associated to the underlying stochastic optimal control problem faced by the players, coupled with a Kolmogorov–Fokker–Planck (KFP) for the density of players within the state space of the game. MFG systems find applications in a broad range of areas, such as economics, population dynamics, and mass transport [21, 18, 22]. We refer the reader to the surveys in [31, 22, 25] for extensive reviews of the literature on the theory and applications for a variety of MFG problems.

The numerical solution of MFG systems is an active area of research and has led to various approaches. Monotone finite difference methods on Cartesian grids are considered in [15, 19, 24]. In particular, under the assumption that the continuous problem admits a unique classical solution, [19] shows the convergence of the approximation of the value function in some first-order Sobolev space for the stationary case, and in some Bochner–Sobolev space for the time-dependent case, along with convergence of the approximations of the density function in some Lebesgue spaces. The assumption of the existence of a classical solution was then removed in [24], which showed convergence of the approximations to a weak solution of the system. There is also an alternative approach to the solution of the problem when the couplings of the system are local. In this case, the MFG system can at least formally be related to the first-order optimality conditions of convex optimisation problems, which leads to other methods based on optimization, see for example [27, 29].

We now outline the motivation for the present paper. Recall that MFG PDE systems are derived from models of large numbers of players solving stochastic optimal control problems. It is well-known from stochastic optimal control that in many applications of practical interest, the underlying controls may be of *bang-bang* type, which typically leads to discontinuities in the optimal control policies. In turn, this generally leads to nondifferentiable Hamiltonians, which pose special challenges for the analysis and numerical analysis of MFG systems.

To illustrate these challenges, we consider as a model problem a stationary MFG system of the form

$$-\nu\Delta u + H(x, \nabla u) + \kappa u = F[m] \quad \text{in } \Omega, \quad (1a)$$

$$-\nu\Delta m - \operatorname{div} \left(m \frac{\partial H}{\partial p}(x, \nabla u) \right) + \kappa m = G(x) \quad \text{in } \Omega, \quad (1b)$$

along with homogeneous Dirichlet boundary conditions $u = 0$ and $m = 0$ on $\partial\Omega$. The unknowns u and m denote, respectively, the value function and the player distribution function of the game. Here, the domain $\Omega \subset \mathbb{R}^n$ is a bounded connected open set in \mathbb{R}^n , $n \geq 2$, and $\nu > 0$ and $\kappa \geq 0$ are constants. Precise assumptions on the data H , F , and G are given below in Section 2. The system (1) includes as special cases the stationary MFG model considered in [12] (in which case κ and G vanish) and some models of discounted MFGs [32]. However note that in contrast to the periodic boundary conditions considered in [12], we consider (1) along with Dirichlet boundary conditions, which arise in models

where players may enter or exit the game, and thus m is not a probability density function in general. This explains why there is no Lagrange multiplier term in the first equation (1a). The source term G and the term involving κ in (1) are also relevant in the context of temporal semi-discretizations of time-dependent MFG systems.

The Hamiltonian in (1) is given in terms of components of the underlying stochastic optimal control problem; we therefore consider Hamiltonians H of the form

$$H(x, p) := \sup_{\alpha \in \mathcal{A}} (b(x, \alpha) \cdot p - f(x, \alpha)) \quad \forall (x, p) \in \bar{\Omega} \times \mathbb{R}^n, \quad (2)$$

where \mathcal{A} denotes the set of controls, and where b is the controlled drift and f is a control-dependent running cost component set by the underlying stochastic optimal control problem. For simplicity, we assume that \mathcal{A} is a compact metric space, and $b : \bar{\Omega} \times \mathcal{A} \rightarrow \mathbb{R}^n$, $f : \bar{\Omega} \times \mathcal{A} \rightarrow \mathbb{R}$ are uniformly continuous, so that the supremum in (2) is achieved. In many applications, the controls that achieve the supremum (2) may be non-unique for some $(x, p) \in \Omega \times \mathbb{R}^n$, which often leads to discontinuous optimal controls of *bang-bang* type. In these cases, the Hamiltonian H is then typically Lipschitz continuous but not differentiable everywhere. However, most works so far on MFGs require differentiable or even C^1 Hamiltonians, which can be quite restrictive in practice.

Nondifferentiable Hamiltonians pose an immediate and obvious challenge for analysis since the advective term in (1b) is then no longer well-defined in a classical sense. This leads to the problem of finding a suitable relaxed meaning for the equation in these situations. To the best of our knowledge, the analysis of MFG with nondifferentiable Hamiltonians seems to only have been considered in [33] for the special case of Hamiltonians of the form $\bar{H}(m, p) = \bar{\kappa}(m)|p|$, for some given function $\bar{\kappa}$; see Remark 1 for further comments. Otherwise, the analysis and numerical analysis of MFG problems with nondifferentiable Hamiltonians remains largely untouched.

Our first main contribution in this work is to provide a suitable generalized meaning for the system (1) when H is nondifferentiable, and to prove results on the existence and uniqueness of solutions under conditions where they are expected to hold. Using the fact that the Hamiltonian H is convex with respect to its second argument, our approach is based on relaxing (1b) as the following *Partial Differential Inclusion* (PDI):

$$-\nu \Delta m + \kappa m - G(x) \in \operatorname{div} (m \partial_p H(x, \nabla u)) \quad \text{in } \Omega, \quad (3)$$

where $\partial_p H$ denotes the Moreau–Rockafellar pointwise partial subdifferential of H with respect to p , and where the equation is understood in a suitable weak sense. The resulting MFG PDI is then

$$\begin{aligned} -\nu \Delta u + H(x, \nabla u) + \kappa u &= F[m] && \text{in } \Omega, \\ -\nu \Delta m + \kappa m - G(x) &\in \operatorname{div} (m \partial_p H(x, \nabla u)) && \text{in } \Omega, \\ u = 0, \quad m = 0, &&& \text{on } \partial\Omega. \end{aligned} \quad (4)$$

We first prove existence of weak solutions of (4) for rather general problem data. Then, crucially, we show uniqueness of solutions for (4) for monotone couplings, thus extending the important uniqueness result from [12, 14] to the case of nondifferentiable Hamiltonians.

Our second main contribution is to propose and study a monotone Finite Element Method (FEM) for approximating weak solutions to the MFG PDI (4). In this context, monotonicity of the FEM refers to the presence of a discrete maximum principle. There is a wide range of approaches to constructing monotone FEMs; see for instance [4, 5, 6, 8, 10]. The discretisation considered here is based on the one from [20] for degenerate fully nonlinear HJB equations, where convergence to the unique viscosity solution was shown; see also [28, 34]. To keep the analysis as simple as possible, we concentrate on a monotone FEM where the discrete maximum principle is achieved via artificial diffusion on strictly acute meshes.

The main result on the analysis of the numerical approximations is given in Theorem 5, which shows convergence of the numerical approximations in the small mesh limit for uniquely solvable MFG PDI systems. In particular, we prove strong convergence in the H^1 -norm of the approximations to the value function u . We also show that the approximations of the density function m converge strongly in L^q -norms for $q \in [1, 2^*)$, $2^* = \frac{2n}{n-2}$, as well as weak convergence in H^1 . For general nondifferentiable Hamiltonians, a proof of strong convergence in H^1 of the density approximations is not currently available owing to a lack of continuity in the advective terms of the KFP equation. However, if some additional continuity is assumed (which holds for instance when the Hamiltonian is C^1), then the density approximations also converge strongly in H^1 , see Corollary 1. We then complement the convergence analysis with two numerical experiments that illustrate the performance of the method.

This paper is organised as follows. We outline basic notation in Section 2, after which in Section 3 we formulate the notion of weak solution for the MFG PDI (4) and state the main results on the continuous problem. These results are then proved in Section 4. In Section 5 we introduce a monotone finite element scheme along with main results on well-posedness and convergence. This is followed by the proofs of these results in Section 6. Section 7 concludes the paper with a discussion of two numerical experiments.

2 Notation

We denote $\mathbb{N} := \{1, 2, 3, \dots\}$, and let $n \in \mathbb{N}$, $n \geq 2$. For a Lebesgue measurable set $\omega \subset \mathbb{R}^n$, let $\|\cdot\|_\omega$ denote the standard L^2 -norm for scalar- and vector-valued functions on ω . Let Ω be a bounded, open connected subset of \mathbb{R}^n with Lipschitz boundary $\partial\Omega$. The n -dimensional open ball of radius r and centre $x_0 \in \mathbb{R}^n$ is denoted by $B_r(x_0)$. For a set $\mathcal{C} \subset \mathbb{R}^n$, we denote its closed convex hull by $\overline{\text{conv}} \mathcal{C}$.

We make the following assumptions on the data appearing in (4). Let $\nu > 0$ and $\kappa \geq 0$ be constants, and let $G \in H^{-1}(\Omega)$. Next, let $F : L^2(\Omega) \rightarrow H^{-1}(\Omega)$

be a possibly nonlocal operator that satisfies

$$\|F[z]\|_{H^{-1}(\Omega)} \leq c_1 (\|z\|_{\Omega} + 1) \quad \forall z \in L^2(\Omega), \quad (5a)$$

$$\|F[m_1] - F[m_2]\|_{H^{-1}(\Omega)} \leq c_2 \|m_1 - m_2\|_{\Omega} \quad \forall m_1, m_2 \in L^2(\Omega) \quad (5b)$$

where $c_1, c_2 \geq 0$ are constants. We will say that F is strictly monotone if

$$\langle F[m_1] - F[m_2], m_1 - m_2 \rangle_{H^{-1} \times H_0^1} \leq 0 \implies m_1 = m_2 \quad (6)$$

whenever $m_1, m_2 \in H_0^1(\Omega)$. Note that although the domain of F is $L^2(\Omega)$, the monotonicity condition (6) is needed only for arguments in the smaller space $H_0^1(\Omega)$. In addition, we will say that $G \in H^{-1}(\Omega)$ is non-negative in the sense of distributions if $\langle G, \phi \rangle_{H^{-1} \times H_0^1} \geq 0$ for all functions $\phi \in H_0^1(\Omega)$ that are non-negative a.e. in Ω . Recall that the Hamiltonian $H: \bar{\Omega} \times \mathbb{R}^n \rightarrow \mathbb{R}$ is defined by (2), and that b and f are uniformly continuous on $\bar{\Omega} \times \mathcal{A}$ with \mathcal{A} a compact metric space. The Hamiltonian H then satisfies the following bounds

$$|H(x, p)| \leq c_3 (|p| + 1) \quad \forall (x, p) \in \Omega \times \mathbb{R}^n \quad (7a)$$

$$|H(x, p) - H(x, q)| \leq c_4 |p - q| \quad \forall (x, p, q) \in \Omega \times \mathbb{R}^n \times \mathbb{R}^n \quad (7b)$$

with $c_3 := \max \left\{ \|b\|_{C(\bar{\Omega} \times \mathcal{A})}, \|f\|_{C(\bar{\Omega} \times \mathcal{A})} \right\}$ and $c_4 := \|b\|_{C(\bar{\Omega} \times \mathcal{A})}$. It is then clear that the mapping $v \mapsto H(\cdot, \nabla v)$ is Lipschitz continuous from $H^1(\Omega)$ into $L^2(\Omega)$.

Given arbitrary sets A and B , an operator \mathcal{M} that maps each point $x \in A$ to a subset of B is called a *set-valued map from A to B* , and we write $\mathcal{M}: A \rightrightarrows B$. For the Hamiltonian given by (2) its pointwise Moreau–Rockafellar partial subdifferential with respect to p is the set-valued map $\partial_p H: \Omega \times \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ defined by

$$\partial_p H(x, p) := \left\{ \tilde{b} \in \mathbb{R}^n : H(x, q) \geq H(x, p) + \tilde{b} \cdot (q - p) \quad \forall q \in \mathbb{R}^n \right\}. \quad (8)$$

Note that $\partial_p H(x, p)$ is non-empty for all $x \in \Omega$ and $p \in \mathbb{R}^n$ because H is real-valued and convex in p for each fixed $x \in \Omega$. Note also that for the special case of a differentiable convex function, the (partial) subdifferential at a point is simply the singleton set containing the value of the (partial) derivative at the point. Furthermore, the subdifferential $\partial_p H$ is uniformly bounded since (7b) implies that for all $(x, p) \in \Omega \times \mathbb{R}^n$, the set $\partial_p H(x, p)$ is contained in the closed ball of radius $c_4 = \|b\|_{C(\bar{\Omega} \times \mathcal{A})}$ centred at the origin.

Given a function $v \in W^{1,1}(\Omega)$, we say that a real-valued vector field $\tilde{b}: \Omega \rightarrow \mathbb{R}^n$ is a *measurable selection of $\partial_p H(\cdot, \nabla v)$* if \tilde{b} is Lebesgue measurable and $\tilde{b}(x) \in \partial_p H(x, \nabla v(x))$ for a.e. $x \in \Omega$. The uniform boundedness of the subdifferential sets implies that any measurable selection \tilde{b} of $\partial_p H(\cdot, \nabla v)$ must belong to $L^\infty(\Omega; \mathbb{R}^n)$. Thus the correspondence between a function $v \in W^{1,1}(\Omega)$ and the set of all measurable selections of $\partial_p H(\cdot, \nabla v)$ defines a set-valued map between $W^{1,1}(\Omega)$ and $L^\infty(\Omega; \mathbb{R}^n)$.

Definition 1. Let H be the function given by (2). We define the set-valued map $D_p H: W^{1,1}(\Omega) \rightrightarrows L^\infty(\Omega; \mathbb{R}^n)$ by

$$D_p H[v] := \left\{ \tilde{b} \in L^\infty(\Omega; \mathbb{R}^n) : \tilde{b}(x) \in \partial_p H(x, \nabla v(x)) \text{ for a.e. } x \in \Omega \right\}.$$

We show in Lemma 2 below that $D_p H[v]$ is non-empty for all v in $W^{1,1}(\Omega)$.

3 Continuous Problem and Main Results

3.1 Problem Statement

We now introduce the notion of weak solution for the MFG PDI (4).

Definition 2 (Weak Solution of (4)). We say that a pair $(u, m) \in H_0^1(\Omega) \times H_0^1(\Omega)$ is a weak solution of (4) if there exists a vector field $\tilde{b}_* \in D_p H[u]$ such that, for all $\psi, \phi \in H_0^1(\Omega)$, there hold

$$\int_{\Omega} \nu \nabla u \cdot \nabla \psi + H(x, \nabla u) \psi + \kappa u \psi \, dx = \langle F[m], \psi \rangle_{H^{-1} \times H_0^1}, \quad (9a)$$

$$\int_{\Omega} \nu \nabla m \cdot \nabla \phi + \tilde{b}_* \cdot \nabla \phi + \kappa m \phi \, dx = \langle G, \phi \rangle_{H^{-1} \times H_0^1}. \quad (9b)$$

The weak formulation of the problem given in Definition 2 can be reformulated in terms of a PDI. In particular, recalling the definition of the set-valued map $D_p H$ in Definition 1 above, for given $m, u \in H_0^1(\Omega)$, let

$$\begin{aligned} & \text{div}(m D_p H[u]) := \\ & \left\{ g \in H^{-1}(\Omega) : \exists \tilde{b} \in D_p H[u] \text{ s.t. } \langle g, \phi \rangle_{H^{-1} \times H_0^1} = - \int_{\Omega} m \tilde{b} \cdot \nabla \phi \, dx \, \forall \phi \in H_0^1(\Omega) \right\}. \end{aligned} \quad (10)$$

In other words, the set $\text{div}(m D_p H[u])$ is the set of all distributions in $H^{-1}(\Omega)$ of the form $\text{div}(m \tilde{b})$ where $\tilde{b} \in D_p H[u]$. Then, the definition of a weak solution in Definition 2 is equivalent to requiring that $(u, m) \in H_0^1(\Omega) \times H_0^1(\Omega)$ solves the following pair of conditions which hold in the sense of distributions in $H^{-1}(\Omega)$:

$$- \nu \Delta u + H(x, \nabla u) + \kappa u = F[m], \quad (11a)$$

$$- \nu \Delta m + \kappa m - G(x) \in \text{div}(m D_p H[u]). \quad (11b)$$

Therefore, the PDI system (11) is the weak formulation of (4).

Remark 1. In [33, Definition 3.1], Ducasse *et al.* propose a definition of weak solutions for problems with Hamiltonians of the form $\bar{H}(\bar{m}, p) := \bar{\kappa}(\bar{m})|p|$. In particular, their definition for a weak solution (\bar{u}, \bar{m}) involves an advective velocity term V in the KFP equation replaced by a possibly non-unique vector field V that satisfies the conditions (in the present notation)

$$\|V\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq \bar{\kappa}(\bar{m}), \quad V(x) \cdot \nabla \bar{u}(x) = \bar{\kappa}(\bar{m}) |\nabla \bar{u}(x)| \text{ for a.e. } x \in \Omega. \quad (12)$$

Although it is not stated therein, it is straightforward to check that the conditions in (12) are equivalent to requiring that V belongs to the partial subdifferential $\partial_p \overline{H}(\overline{m}, \nabla \overline{u})$. Thus, modulo the dependence of the Hamiltonian on the density of players, our approach significantly generalizes that of [33] to more general nondifferentiable Hamiltonians.

3.2 Main Results

The first main result for the continuous problem (9) is the following.

Theorem 1 (Existence of Weak Solutions). *There exists a pair $(u, m) \in H_0^1(\Omega) \times H_0^1(\Omega)$ that is a weak solution of (4) in the sense of Definition 2 satisfying*

$$\|m\|_{H^1(\Omega)} \leq C^* \|G\|_{H^{-1}(\Omega)}, \quad (13)$$

$$\|u\|_{H^1(\Omega)} \leq C^{**} \left(\|G\|_{H^{-1}(\Omega)} + \|f\|_{C(\overline{\Omega} \times \mathcal{A})} + 1 \right), \quad (14)$$

for some constants $C^*, C^{**} \geq 0$ depending only on n , Ω , ν , $\|b\|_{C(\overline{\Omega} \times \mathcal{A})}$, κ and c_1 .

The second main result ensures uniqueness of weak solutions to (4) under a monotonicity condition on F that is similar to the one that was used by Lasry and Lions in [14]. Since we also consider problems with source terms, we shall further require non-negativity of G in the sense of distributions.

Theorem 2 (Uniqueness of Weak Solutions). *If F is strictly monotone and G is non-negative in the sense of distributions in $H^{-1}(\Omega)$, then there exists a unique weak solution pair $(u, m) \in H_0^1(\Omega) \times H_0^1(\Omega)$ to (4) in the sense of Definition 2.*

Remark 2. To avoid any confusion, we stress that Theorem 2 guarantees the uniqueness of the weak solution pair (u, m) under the relevant hypotheses, although the advective vector field $\tilde{b} \in D_p H[u]$ that appears in Definition 2 may be non-unique. Note also that the monotonicity condition on F is similar to the monotonicity condition on the coupling term used by Lasry & Lions in [12] for classical solutions to ergodic mean field game systems with C^1 Hamiltonians.

4 Analysis of Continuous Problem

4.1 Preliminary Results

To begin, we introduce the pointwise maximising set corresponding to the Hamiltonian (2). Define the set-valued map $\Lambda: \Omega \times \mathbb{R}^n \rightrightarrows \mathcal{A}$ via

$$\Lambda(x, p) := \operatorname{argmax}_{\alpha \in \mathcal{A}} \{b(x, \alpha) \cdot p - f(x, \alpha)\}.$$

Note that $\Lambda(x, p)$ is non-empty for all $x \in \Omega$ and all $p \in \mathbb{R}^n$ since \mathcal{A} is compact and the functions b and f are continuous. The following Lemma, which is a consequence of [7, Proposition 4.4], shows the link between the sets of maximising controls and the subdifferentials of the Hamiltonian.

Lemma 1. *Let H be given by (2). Then*

$$\partial_p H(x, p) = \overline{\text{conv}} \{b(x, \alpha) : \alpha \in \Lambda(x, p)\} \quad \forall (x, p) \in \bar{\Omega} \times \mathbb{R}^n. \quad (15)$$

To show that $D_p H$ possess non-empty images, we introduce an auxiliary set-valued map. For a given $v \in W^{1,1}(\Omega)$, let $\Lambda[v]$ denote the set of all Lebesgue measurable functions $\alpha^* : \Omega \rightarrow \mathcal{A}$ that satisfy $\alpha^*(x) \in \Lambda(x, \nabla v(x))$ for a.e. $x \in \Omega$. We will refer to each element of $\Lambda[v]$ as a *measurable selection* of $\Lambda(\cdot, \nabla v(\cdot))$. It is known that $\Lambda[v]$ is non-empty for each $v \in W^{1,1}(\Omega)$, see e.g. [23, Theorem 10], where the proof the existence of measurable selections ultimately rests upon the Kuratowski and Ryll–Nardzewski Selection Theorem [3].

We now show that the set-valued map $D_p H$ has non-empty images.

Lemma 2. *For each $v \in W^{1,1}(\Omega)$, the set $D_p H[v]$ is a non-empty subset of $L^\infty(\Omega; \mathbb{R}^n)$, and we have the uniform bound*

$$\sup_{v \in W^{1,1}(\Omega)} \left[\sup_{\tilde{b} \in D_p H[v]} \|\tilde{b}\|_{L^\infty(\Omega; \mathbb{R}^n)} \right] \leq \|b\|_{C(\bar{\Omega} \times \mathcal{A})}. \quad (16)$$

Proof. Let $v \in W^{1,1}(\Omega)$ be given. We need to show $D_p H[v]$ is non-empty. Non-emptiness of $\Lambda[v]$ implies that there exists a Lebesgue measurable map $\alpha^* : \Omega \rightarrow \mathcal{A}$ such that $\alpha^*(x) \in \Lambda(x, \nabla v(x))$ for a.e. $x \in \Omega$, and thus $H(x, \nabla v(x)) = b(x, \alpha^*(x)) \cdot \nabla v(x) - f(x, \alpha^*(x))$ for a.e. $x \in \Omega$. Now, suppose $q \in \mathbb{R}^n$ is arbitrary. We find by definition of $H(x, q)$ that, for a.e. $x \in \Omega$,

$$\begin{aligned} H(x, q) &\geq b(x, \alpha^*(x)) \cdot q - f(x, \alpha^*(x)) \\ &= b(x, \alpha^*(x)) \cdot q + H(x, \nabla v(x)) - b(x, \alpha^*(x)) \cdot \nabla v(x) \\ &= H(x, \nabla v(x)) + b(x, \alpha^*(x)) \cdot (q - \nabla v(x)). \end{aligned}$$

It follows then that $b(x, \alpha^*(x)) \in \partial_p H(x, \nabla v(x))$ for a.e. $x \in \Omega$. Furthermore, we have $b(\cdot, \alpha^*(\cdot)) \in L^\infty(\Omega; \mathbb{R}^n)$ since $\|b(\cdot, \alpha^*(\cdot))\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq \|b\|_{C(\bar{\Omega} \times \mathcal{A})}$. Hence, $D_p H[v]$ is non-empty, as claimed. Finally, the bound (16) follows immediately from the fact that, for all $(x, p) \in \Omega \times \mathbb{R}^n$, the subdifferential $\partial_p H(x, p)$ is contained in the closed ball of radius $\|b\|_{C(\bar{\Omega} \times \mathcal{A})}$ centred at the origin. \square

The following Lemma shows that $D_p H$ has a certain closure property with respect to convergent sequences of its arguments and their measurable selections.

Lemma 3. *Suppose $\{v_j\}_{j \in \mathbb{N}} \subset H^1(\Omega)$, $\{\tilde{b}_j\}_{j \in \mathbb{N}} \subset L^\infty(\Omega; \mathbb{R}^n)$ are sequences such that $\tilde{b}_j \in D_p H[v_j]$ for all $j \in \mathbb{N}$. If $v_j \rightarrow v$ in $H^1(\Omega)$ and $\tilde{b}_j \rightharpoonup^* \tilde{b}$ in $L^\infty(\Omega; \mathbb{R}^n)$ as $j \rightarrow \infty$, then $\tilde{b} \in D_p H[v]$.*

Proof. Introduce the set $Y := \{v \in L^2(\Omega) : v(x) \geq 0 \text{ for a.e. } x \in \Omega\}$ of nonnegative a.e. functions in $L^2(\Omega)$, and note that Mazur's Theorem implies that Y is weakly closed in $L^2(\Omega)$ since it is convex and strongly closed. Let $q \in \mathbb{R}^n$ be a fixed but arbitrary vector. Define the sequence of real-valued functions $\{\omega_j\}_{j=1}^\infty \subset L^2(\Omega)$ by

$$\omega_j(x) := H(x, q) - H(x, \nabla v_j(x)) - \tilde{b}_j(x) \cdot (q - \nabla v_j(x)) \quad (17)$$

for each $j \in \mathbb{N}$ and a.e. $x \in \Omega$. It follows from the definitions of the subdifferential sets (8) and Definition 1 that $\omega_j \in Y$ for each $j \geq 1$. The hypothesis of strong convergence of $\{v_j\}_{j \in \mathbb{N}}$ and weak-* convergence of $\{\tilde{b}_j\}_{j \in \mathbb{N}}$ implies the weak convergence $\omega_j \rightharpoonup \omega$ in $L^2(\Omega)$ where

$$\omega(x) := H(x, q) - H(x, \nabla v(x)) - \tilde{b}(x) \cdot (q - \nabla v(x)) \quad (18)$$

for a.e. $x \in \Omega$. Since Y is weakly closed, it follows that $\omega \in Y$. Since $q \in \mathbb{R}^n$ is arbitrary and since \mathbb{R}^n is separable, we conclude that $\tilde{b} \in D_p H[v]$. \square

4.2 Existence of weak solutions

In this section, we prove Theorem 1. To begin, we introduce notation describing a collection of linear differential operators in weak form. Given $C_0 \geq 0$, let $\mathcal{G}(C_0)$ denote the set of all operators $L : H_0^1(\Omega) \rightarrow H^{-1}(\Omega)$ of the form

$$\langle Lu, v \rangle_{H^{-1} \times H_0^1} = \int_{\Omega} \nu \nabla u \cdot \nabla v + \tilde{b} \cdot \nabla uv + cuv \, dx, \quad (19)$$

where the coefficients satisfy

$$\|\tilde{b}\|_{L^\infty(\Omega; \mathbb{R}^n)} + \|c\|_{L^\infty(\Omega)} \leq C_0, \text{ and } c \geq 0 \text{ a.e. in } \Omega. \quad (20)$$

Moreover, given an operator $L \in \mathcal{G}(C_0)$ for some $C_0 \geq 0$, we define $L^* : H_0^1(\Omega) \rightarrow H^{-1}(\Omega)$, the formal adjoint of L , by $\langle L^* w, v \rangle_{H^{-1} \times H_0^1} := \langle Lv, w \rangle_{H^{-1} \times H_0^1}$ for all $w, v \in H_0^1(\Omega)$.

The invertibility of operators L and their adjoints L^* from the class $\mathcal{G}(C_0)$ is well-known, and follows from the the Fredholm Alternative together with the Weak Maximum Principle and the Comparison Principle (see [9, Ch. 8, Ch. 10]). In the analysis below, we will use the following stronger result, which shows that for fixed C_0 , there is a uniform bound on the norm of the inverses of all operators and their adjoints from the class $\mathcal{G}(C_0)$.

Lemma 4. *Let $C_0 \geq 0$ be given. For every operator $L \in \mathcal{G}(C_0)$, both L and L^* are boundedly invertible as mappings from $H_0^1(\Omega)$ to $H^{-1}(\Omega)$, and there exists a constant $C_1 > 0$ depending on only Ω , n , ν , and C_0 such that*

$$\sup_{L \in \mathcal{G}(C_0)} \max \left\{ \|L^{-1}\|_{\mathcal{L}(H^{-1}(\Omega), H_0^1(\Omega))}, \|L^{*-1}\|_{\mathcal{L}(H^{-1}(\Omega), H_0^1(\Omega))} \right\} \leq C_1. \quad (21)$$

The proof of Lemma 4 is obtained by a contradiction argument and using the well-known compactness theorems of Rellich–Kondrachov and Banach–Alaoglu. We leave the details as an exercise to the reader.

With the help of Schaefer’s fixed point theorem (see [16]), the existence of measurable selections guaranteed by Lemma 2, and the uniform inverse operator estimate given in Lemma 4, we obtain well-posedness for a class of HJB equations along with a continuity result.

Lemma 5 (Well-posedness of the HJB equation). *Let $m \in L^2(\Omega)$ be given. Then, there exists a unique $u \in H_0^1(\Omega)$ such that*

$$\int_{\Omega} \nu \nabla u \cdot \nabla \psi + H(x, \nabla u) \psi + \kappa u \psi \, dx = \langle F[m], \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in H_0^1(\Omega). \quad (22)$$

There exists a constant C_1 depending only on Ω , n , ν , κ , $\|b\|_{C(\bar{\Omega} \times \mathcal{A}; \mathbb{R}^n)}$ and c_1 such that

$$\|u\|_{H^1(\Omega)} \leq C_1 \left(\|m\|_{\Omega} + \|f\|_{C(\bar{\Omega} \times \mathcal{A})} + 1 \right). \quad (23)$$

Moreover, the solution u depends continuously on m , i.e., if $\{m_j\}_{j \in \mathbb{N}} \subset L^2(\Omega)$ is such that $m_j \rightarrow m$ in $L^2(\Omega)$ as $j \rightarrow \infty$, then the corresponding sequence of solutions $\{u_j\}_{j \in \mathbb{N}} \subset H_0^1(\Omega)$ to the problem (22) converges in $H_0^1(\Omega)$ to the unique solution u of (22).

We can now show existence of weak solutions to (4) in the sense of Definition 2 by a fixed point argument.

Proof of Theorem 1. We start by developing an iterative sequence of function pairs $\{(u_j, m_j)\}_{j \in \mathbb{N}}$ in $H_0^1(\Omega) \times H_0^1(\Omega)$ as follows: Let $m_1 \in H_0^1(\Omega)$ be given. Then, for each $j \geq 1$, we define inductively $u_j, m_{j+1} \in H_0^1(\Omega)$ as the unique solutions of

$$\int_{\Omega} \nu \nabla u_j \cdot \nabla \psi + H(x, \nabla u_j) \psi + \kappa u_j \psi \, dx = \langle F[m_j], \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in H_0^1(\Omega), \quad (24)$$

$$\int_{\Omega} \nu \nabla m_{j+1} \cdot \nabla \phi + m_{j+1} \tilde{b}_j \cdot \nabla \phi + \kappa m_{j+1} \phi \, dx = \langle G, \phi \rangle_{H^{-1} \times H_0^1} \quad \forall \phi \in H_0^1(\Omega), \quad (25)$$

where \tilde{b}_j is chosen from $D_p H[u_j]$. Note that the unique solvability of (24) is given by Lemma 5. Then, the existence of a choice $\tilde{b}_j \in D_p H[u_j]$ is assured by Lemma 2, and the unique solvability of (25) follows from Lemma 4. We consequently obtain sequences $\{m_j\}_{j \in \mathbb{N}} \subset H_0^1(\Omega)$, $\{u_j\}_{j \in \mathbb{N}} \subset H_0^1(\Omega)$ and $\{\tilde{b}_j\}_{j \in \mathbb{N}} \subset L^\infty(\Omega; \mathbb{R}^n)$. Note that the sequence $\{\tilde{b}_j\}_{j \in \mathbb{N}}$ is uniformly bounded as a result of (16), and thus by Banach–Alaoglu’s Theorem we may pass to a subsequence, without change of notation, which satisfies $\tilde{b}_j \rightharpoonup^* \tilde{b}_*$ in $L^\infty(\Omega; \mathbb{R}^n)$ for some $\tilde{b}_* \in L^\infty(\Omega; \mathbb{R}^n)$ as $j \rightarrow \infty$. Furthermore, Lemma 4 implies that there exists a constant C_1 such that

$$\|m_{j+1}\|_{H^1(\Omega)} \leq C_1 \|G\|_{H^{-1}(\Omega)} \quad \forall j \in \mathbb{N}, \quad (26)$$

and thus $\{m_j\}_{j \in \mathbb{N}}$ is uniformly bounded in $H_0^1(\Omega)$. By Rellich–Kondrachov’s Compactness Theorem we may pass to a further subsequence (without change of notation) that satisfies $m_j \rightarrow m$ in $H_0^1(\Omega)$ and $m_j \rightarrow m$ in $L^2(\Omega)$ for some

$m \in H_0^1(\Omega)$ as $j \rightarrow \infty$. Hence, from (26) and the weak convergence of $\{m_j\}_{j \in \mathbb{N}}$ to m in $H_0^1(\Omega)$ we find

$$\|m\|_{H^1(\Omega)} \leq C_1 \|G\|_{H^{-1}(\Omega)}. \quad (27)$$

Moreover, since $\tilde{b}_j \rightharpoonup^* \tilde{b}$ in $L^\infty(\Omega; \mathbb{R}^n)$ and since $m_j \rightarrow m$ in $L^2(\Omega)$, we have $m_j \tilde{b}_j \rightharpoonup m \tilde{b}$ in $L^2(\Omega)$ as $j \rightarrow \infty$. Therefore, we may pass to the limit in (25) to find that

$$\int_{\Omega} \nu \nabla m \cdot \nabla \phi + m \tilde{b}_* \cdot \nabla \phi + \kappa m \phi \, dx = \langle G, \phi \rangle_{H^{-1} \times H_0^1} \quad \forall \phi \in H_0^1(\Omega). \quad (28)$$

Due to the property of continuous dependence of solutions of the HJB equation on the data, as stated in Lemma 5, the sequence u_j converges to $u \in H_0^1(\Omega)$ that solves

$$\int_{\Omega} \nu \nabla u \cdot \nabla \psi + H(x, \nabla u) \psi + \kappa u \psi \, dx = \langle F[m], \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in H_0^1(\Omega). \quad (29)$$

The fact that $\tilde{b}_* \in D_p H[u]$ follows directly from Lemma 3. Thus it is seen from (28) and (29) that (u, m) is a weak solution pair of the MFG PDI system in the sense of Definition 2. The bounds (13) and (14) then follow directly from (27) and (23). \square

4.3 Uniqueness of weak solutions

Using Definition 1, in addition to a strict monotonicity condition on F and the non-negativity of G in the sense of distributions in $H^{-1}(\Omega)$, we obtain uniqueness of weak solutions in the sense of Definition 2.

Proof of Theorem 2. Suppose there are two solutions to (9), i.e. there exist (u_i, m_i) , $i \in \{1, 2\}$, that each satisfy (9) with

$$\int_{\Omega} \nu \nabla u_i \cdot \nabla \psi + H(x, \nabla u_i) \psi + \kappa u_i \psi \, dx = \langle F[m_i], \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in H_0^1(\Omega), \quad (30a)$$

$$\int_{\Omega} \nu \nabla m_i \cdot \nabla \phi + m_i \tilde{b}_i \cdot \nabla \phi + \kappa m_i \phi \, dx = \langle G, \phi \rangle_{H^{-1} \times H_0^1} \quad \forall \phi \in H_0^1(\Omega), \quad (30b)$$

for some $\tilde{b}_i \in D_p H[u_i]$. Since G is non-negative in the sense of distributions in $H^{-1}(\Omega)$, the Comparison Principle (see [9, Theorem 10.7]) applied to (30b) implies that $m_i \geq 0$ a.e. in Ω for each $i \in \{1, 2\}$. After choosing as test functions $\psi = m_1 - m_2$ and $\phi = u_1 - u_2$ in (30), and subtracting the equations, we eventually find that

$$\int_{\Omega} m_1 \lambda_{12} + m_2 \lambda_{21} \, dx = \langle F[m_1] - F[m_2], m_1 - m_2 \rangle_{H^{-1} \times H_0^1}, \quad (31)$$

where the functions λ_{ij} are defined by

$$\lambda_{ij} := H(\cdot, \nabla u_i) - H(\cdot, \nabla u_j) + \tilde{b}_i \cdot \nabla(u_j - u_i), \quad i, j \in \{1, 2\}. \quad (32)$$

By definition of $D_p H[u_i]$, in particular that $\tilde{b}_i(x) \in \partial_p H(x, \nabla u_i(x))$ for a.e. $x \in \Omega$, we see that $\lambda_{ij} \leq 0$ a.e. in Ω for $i, j \in \{1, 2\}$. Therefore, (31) implies that $\langle F[m_1] - F[m_2], m_1 - m_2 \rangle_{H^{-1} \times H_0^1} \leq 0$, and thus the strict monotonicity condition (6) on F implies that $m_1 = m_2$. Consequently u_1 and u_2 satisfy (30a) with identical right-hand side $F[m_1] = F[m_2]$. Therefore, Lemma 5 implies that $u_1 = u_2$. This shows that there is at most one weak solution to (4) in the sense of Definition 2. \square

Remark 3. In cases where the mean field game PDI (9) admits a unique solution (u, m) , the collection of transport vectors $\tilde{b}_* \in D_p H[u]$ for which (9) holds constitutes an equivalence class of vector fields under the following equivalence relation: for vector fields $\tilde{b}_1, \tilde{b}_2 \in D_p H[u]$,

$$\tilde{b}_1 \sim \tilde{b}_2 \quad \text{if and only if} \quad \int_{\Omega} m \tilde{b}_1 \cdot \nabla \phi \, dx = \int_{\Omega} m \tilde{b}_2 \cdot \nabla \phi \, dx \quad \forall \phi \in H_0^1(\Omega).$$

Therefore, whenever \tilde{b}_1, \tilde{b}_2 are in the above equivalence class, $\operatorname{div}(m \tilde{b}_1 - m \tilde{b}_2) = 0$ in the sense of distributions in $H^{-1}(\Omega)$, and so the vector $m \tilde{b}_1 - m \tilde{b}_2$ is the curl of a vector potential in a distributional sense.

5 Monotone Continuous Galerkin Finite Element Scheme

In this section we introduce a monotone finite element scheme for approximating solutions to the weak formulation (9). In the sequel, we shall further assume that Ω is a polyhedron, in addition to the earlier assumption that it is a bounded connected open set with Lipschitz boundary.

5.1 Notation

A mesh \mathcal{T} is a collection of closed n -dimensional simplices, called elements, K with non-overlapping interiors that satisfy $\bar{\Omega} = \bigcup_{K \in \mathcal{T}} K$. Each vertex of an element K of a given mesh is called a node of the element. A face of an element $K \in \mathcal{T}$ is the convex hull of a collection of n nodes of K which has positive $(n-1)$ -dimensional Hausdorff measure (c.f. [17]). For instance, when the space dimension $n = 2$, each face of an element $K \in \mathcal{T}$ is one of its three edges. We will always assume that a given mesh \mathcal{T} is conforming (or often called matching) [17], i.e. for any element $K \in \mathcal{T}$ with nodes $\{x_0, \dots, x_n\}$, the set $\partial K \cap \partial K'$ for each element $K' \in \mathcal{T}$, $K' \neq K$, is the convex hull of a (possibly empty) subset of $\{x_0, \dots, x_n\}$. Let $\{\mathcal{T}_k\}_{k \in \mathbb{N}}$ be a given sequence of conforming meshes. For each $k \in \mathbb{N}$, let the mesh-size of a given mesh \mathcal{T}_k be defined by $h_k := \max_{K \in \mathcal{T}_k} \operatorname{diam}(K)$. We assume that $h_k \rightarrow 0$ as $k \rightarrow \infty$. We assume that

$\{\mathcal{T}_k\}_{k \in \mathbb{N}}$ is shape-regular, i.e. there exists a real-number $\delta > 1$, independent of $k \in \mathbb{N}$, such that $\forall k \in \mathbb{N}, \forall K \in \mathcal{T}_k, \text{diam}(K) \leq \delta \rho_K$, where ρ_K denotes the radius of the largest inscribed ball in the element K . We assume in addition that the family of meshes $\{\mathcal{T}_k\}_{k \in \mathbb{N}}$ is nested, i.e. for each $k \in \mathbb{N}$ the mesh \mathcal{T}_{k+1} is obtained from \mathcal{T}_k via an admissible subdivision of each element of \mathcal{T}_k into simplices.

Given an element K , we let $\mathcal{P}_1(K)$ denote the vector space of n -variate real-valued polynomials of total degree 1 that are defined on K . The discretization of the continuous problem (9) is based on the following finite element spaces:

$$V_k := \{v \in H_0^1(\Omega) : v|_K \in \mathcal{P}_1(K) \ \forall K \in \mathcal{T}_k\} \quad \forall k \in \mathbb{N}.$$

Given $k \in \mathbb{N}$, the space V_k admits a unique nodal basis of hat functions that we denote by $\{\xi_1, \dots, \xi_{N_k}\}$, which corresponds to a maximal collection of nodes $\{x_1, \dots, x_{N_k}\}$ of the mesh \mathcal{T}_k , such that $\xi_i(x_j) = \delta_{ij}$ for $i, j \in \{1, \dots, N_k\}$, where δ_{ij} is the Kronecker delta. Moreover, V_k inherits the standard norm on $H_0^1(\Omega)$ and we denote this norm by $\|\phi\|_{V_k} := \|\phi\|_{H^1(\Omega)}$ for $\phi \in V_k$. Note that, due to nestedness of the sequence of meshes $\{\mathcal{T}_k\}_{k \in \mathbb{N}}$, V_k is a closed subspace of V_{k+1} for each $k \in \mathbb{N}$. In addition, the union $\bigcup_{k \in \mathbb{N}} V_k$ is dense in $H_0^1(\Omega)$. We let V_k^* denote the space of continuous linear functionals on V_k with standard norm denoted by $\|\cdot\|_{V_k^*}$. For any operator $\mathcal{Z} : V_k \rightarrow V_k^*$ we define the adjoint operator $\mathcal{Z}^* : V_k \rightarrow V_k^*$ by $\langle \mathcal{Z}^* w, v \rangle_{V_k^* \times V_k} := \langle \mathcal{Z} v, w \rangle_{V_k^* \times V_k}$ for all $w, v \in V_k$.

Let $k \in \mathbb{N}$ be given. For $K \in \mathcal{T}_k$, we denote by $\{\psi_{k,0}^K, \dots, \psi_{k,n}^K\} \subset V_k$ the set of nodal basis functions associated with the $n+1$ nodes of K and let

$$\sigma_K^k := \text{diam}(K) \min_{0 \leq i \leq n} |\nabla \psi_{k,i}^K|, \quad \sigma^k := \min_{K \in \mathcal{T}_k} \sigma_K^k. \quad (33)$$

We note that, owing to the shape regularity of the family of meshes $\{\mathcal{T}_k\}_{k \in \mathbb{N}}$, there exist constants $\underline{\sigma}, \bar{\sigma} > 0$, independent of $k \in \mathbb{N}$, such that

$$\underline{\sigma} \leq \sigma^k \leq \bar{\sigma} \quad \forall k \in \mathbb{N}.$$

We assume in addition, for Subsection 5.2, Subsection 5.3 and Section 6, that the family of meshes $\{\mathcal{T}_k\}_{k \in \mathbb{N}}$ is *strictly acute* [10] in the following sense: there exists $\theta \in (0, \pi/2)$, independent of $k \in \mathbb{N}$, such that, for each $k \in \mathbb{N}$, the nodal basis $\{\xi_1, \dots, \xi_{N_k}\}$ of V_k satisfies

$$\nabla \xi_i \cdot \nabla \xi_j|_K \leq -\sin(\theta) |\nabla \xi_i|_K |\nabla \xi_j|_K \quad \forall 1 \leq i, j \leq N_k, i \neq j, \forall K \in \mathcal{T}. \quad (34)$$

The condition (34) can be interpreted geometrically. For instance, in two space dimensions the strict acuteness condition (34) indicates that the largest angle of a given triangle $K \in \mathcal{T}_k$ is at most $\frac{\pi}{2} - \theta$, while in three space dimensions (34) indicates that each angle formed by the six pairs of faces of any tetrahedron $K \in \mathcal{T}_k$ is at most $\frac{\pi}{2} - \theta$ (see [10]).

5.2 A Monotone Finite Element Method

The proof of uniqueness of weak solutions of (9) uses the Comparison Principle of elliptic operators (see Section 4). In order to preserve this approach on the

discrete level, we consider here approximations by a monotone finite element method that satisfies a discrete maximum principle (see, e.g., [4]). As such, we will consider a finite element discretisation of (9) that employs the method of artificial diffusion on strictly acute meshes [10, 20] to ensure non-negativity of the approximations for the density.

We introduce a family of artificial diffusion coefficients that will be used in the finite element discretisation of (9). Let $\mu > 1$ be a fixed constant. Then, for each $k \in \mathbb{N}$, we define the artificial diffusion coefficient $\gamma_k : \Omega \rightarrow \mathbb{R}$ element-wise over \mathcal{T}_k by

$$\gamma_k|_K := \max \left(\mu \frac{\|b\|_{C(\bar{\Omega} \times \mathcal{A})} \text{diam}(K) + \kappa \text{diam}(K)^2}{\sigma^k \sin(\theta)} - \nu, 0 \right) \quad \text{on } K \in \mathcal{T}_k. \quad (35)$$

With the artificial diffusion coefficients $\{\gamma_k\}_{k \in \mathbb{N}}$ given by (35), the P1-continuous Galerkin finite element discretisation of (9) that we consider is the following: *Given $k \in \mathbb{N}$, find $(u_k, m_k) \in V_k \times V_k$ such that there exists $\tilde{b}_k \in D_p H[u_k]$ satisfying*

$$\int_{\Omega} (\nu + \gamma_k) \nabla u_k \cdot \nabla \psi + H(x, \nabla u_k) \psi + \kappa u_k \psi \, dx = \langle F[m_k], \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in V_k, \quad (36a)$$

$$\int_{\Omega} (\nu + \gamma_k) \nabla m_k \cdot \nabla \phi + m_k \tilde{b}_k \cdot \nabla \phi + \kappa m_k \phi \, dx = \langle G, \phi \rangle_{H^{-1} \times H_0^1} \quad \forall \phi \in V_k. \quad (36b)$$

Remark 4 (Basic Properties of Artificial Diffusion Coefficients). We observe some key properties of the artificial diffusion coefficients given by (35). Firstly, for each $k \in \mathbb{N}$, γ_k is in $L^\infty(\Omega)$, constant element-wise, and non-negative a.e. in Ω . Moreover, there holds $\sup_{k \in \mathbb{N}} \|\gamma_k\|_{L^\infty(\Omega)} < \infty$ due to shape regularity of the family $\{\mathcal{T}_k\}_{k \in \mathbb{N}}$. Secondly, since the sequence of mesh sizes $\{h_k\}_{k \in \mathbb{N}}$ satisfies $h_k \rightarrow 0$ as $k \rightarrow \infty$ by assumption, there exists $k_* \in \mathbb{N}$ that depends on ν, μ, θ, κ and $\|b\|_{C(\bar{\Omega} \times \mathcal{A})}$ such that, for all $k \geq k_*$, we have $\gamma_k = 0$ a.e. in Ω . Hence,

$$\|\gamma_k\|_{L^\infty(\Omega)} = 0 \quad \forall k \geq k_*, \quad (37)$$

and thus we recover consistency of the discrete problems (36) with the continuous problem (9) in the limit as the mesh-size vanishes.

5.3 Main Results

We can now state the main results concerning the finite element scheme (36). The first result concerns the existence of solutions to (36).

Theorem 3 (Existence). *For each $k \in \mathbb{N}$ there exists a discrete solution pair $(u_k, m_k) \in V_k \times V_k$ that solves (36). Moreover, there exist constants $C_1^*, C_2^* \geq 0$,*

independent of $k \in \mathbb{N}$, such that

$$\sup_{k \in \mathbb{N}} \|m_k\|_{H^1(\Omega)} \leq C_1^* \|G\|_{H^{-1}(\Omega)}, \quad (38a)$$

$$\sup_{k \in \mathbb{N}} \|u_k\|_{H^1(\Omega)} \leq C_2^* \left(\|G\|_{H^{-1}(\Omega)} + \|f\|_{C(\bar{\Omega} \times \mathcal{A})} + 1 \right). \quad (38b)$$

Next, uniqueness of solutions to (36) holds under the same monotonicity and non-negativity assumptions of Theorem 2.

Theorem 4 (Uniqueness). *Suppose the coupling term F is strictly monotone and G is non-negative in the sense of distributions in $H^{-1}(\Omega)$. Then, for each $k \in \mathbb{N}$, there exists a unique discrete solution pair $(u_k, m_k) \in V_k \times V_k$ to (36).*

The proofs of Theorem 3 and Theorem 4 are given in Section 6 below.

We now state the first main result on the convergence of the scheme (36).

Theorem 5 (Convergence). *Assume the coupling term F is strictly monotone and G is non-negative in the sense of distributions in $H^{-1}(\Omega)$. Let (u, m) denote the unique pair that solves (4) in the sense of Definition 2 and let $\{(u_k, m_k)\}_{k \in \mathbb{N}}$ denote the sequence of solutions generated by (36). Then, as $k \rightarrow \infty$,*

$$u_k \rightarrow u \quad \text{in } H_0^1(\Omega), \quad m_k \rightarrow m \quad \text{in } L^q(\Omega), \quad m_k \rightharpoonup m \quad \text{in } H_0^1(\Omega) \quad (39)$$

for any $q \in [1, 2^*)$ where $2^* = \infty$ if $n = 2$ and $2^* = \frac{2n}{n-2}$ if $n \geq 3$.

In general, the strong convergence of ∇m_k to ∇m in $L^2(\Omega; \mathbb{R}^n)$ is not known. The difficulty lies in the fact that it appears possible that the sequence $\{\tilde{b}_k\}_{k \in \mathbb{N}}$ might be such that there is no subsequence that converges in a sufficiently strong sense. However, under additional conditions, the weak convergence of the density approximations in $H_0^1(\Omega)$ can be improved to strong convergence. This is the case, for instance, if the Hamiltonian H given by (2) is such that partial derivative $\frac{\partial H}{\partial p}$ exists and is continuous in $\Omega \times \mathbb{R}^n$. In fact, a weaker hypothesis than this can be formulated that ensures strong convergence of the density approximations in $H_0^1(\Omega)$.

Corollary 1 (Strong H^1 -Convergence for Density Approximations). *In addition to the hypotheses of Theorem 5, suppose that the sequence of transport vector fields $\{\tilde{b}_k\}_{k \in \mathbb{N}}$ from (36) is pre-compact in $L^1(\Omega; \mathbb{R}^n)$. Then, m_k converges to m strongly in $H_0^1(\Omega)$ as $k \rightarrow \infty$.*

Remark 5. The compactness hypothesis on the sequence of transport vector fields introduced above is satisfied when the partial derivative $\frac{\partial H}{\partial p}$ exists and is continuous in $\Omega \times \mathbb{R}^n$. Indeed, in this case $\tilde{b}_k = \frac{\partial H}{\partial p}(x, \nabla u_k)$ in $L^\infty(\Omega; \mathbb{R}^n)$ for all $k \in \mathbb{N}$. With the strong convergence of the value function approximations in $H_0^1(\Omega)$ guaranteed by Theorem 5, it is easy to see that the entire sequence $\{\tilde{b}_k\}_{k \in \mathbb{N}}$ converges strongly to $\frac{\partial H}{\partial p}(x, \nabla u)$ in $L^s(\Omega; \mathbb{R}^n)$ for any $s \in [1, \infty)$. Hence, the sequence is pre-compact in $L^1(\Omega; \mathbb{R}^n)$.

6 Analysis of Monotone Finite Element Scheme

6.1 Stabilisation of Linear Differential Operators

We will say that a linear operator $L : V_k \rightarrow V_k^*$ exhibits the *discrete maximum principle* provided that the following condition holds: if $w \in V_k$ and $\langle Lw, \xi_i \rangle_{V_k^* \times V_k} \geq 0$ for all $i \in \{1, \dots, N_k\}$, then $w \geq 0$ in Ω . For the analysis of (36), we introduce a collection of linear operators that are perturbations of discrete relatives to the operators considered in the continuous setting of Lemma 4. Recall the definition of the sequence $\{\gamma_k\}_{k \in \mathbb{N}}$ given in (35). For each $k \in \mathbb{N}$, let W_k denote the collection of linear operators $L : V_k \rightarrow V_k^*$ of the form

$$\langle Lw, v \rangle_{V_k^* \times V_k} := \int_{\Omega} (\nu + \gamma_k) \nabla w \cdot \nabla v + \tilde{b} \cdot \nabla w v + \kappa w v \, dx \quad \forall w, v \in V_k,$$

with $\tilde{b} : \Omega \rightarrow \mathbb{R}^n$ denoting a Lebesgue measurable vector field satisfying the uniform bound $\|\tilde{b}\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq \|b\|_{C(\bar{\Omega} \times \mathcal{A})}$.

By adaptating the proof of [10, Theorem 4.2] and [20, Section 8], we obtain the following result that will assist with ensuring non-negativity of finite element approximations $\{m_k\}_{k \in \mathbb{N}}$ of the density function and proving convergence of the scheme (36).

Lemma 6 (Stabilisation via Artificial Diffusion). *Given $k \in \mathbb{N}$, each operator $A \in W_k$, and its adjoint A^* , exhibits the discrete maximum principle.*

6.2 Well-posedness

First, we establish that the numerical scheme (36) is well-posed, i.e. it admits a unique numerical solution for each $k \in \mathbb{N}$. Suppose $C_0 \geq 0$ is given. Fundamental to the conclusion of Lemma 4 is the fact that operators from $\mathcal{G}(C_0)$, and their adjoints, are invertible as maps from $H_0^1(\Omega)$ into $H^{-1}(\Omega)$. We will employ a discrete version of Lemma 4 in the analysis of (36).

Lemma 7. *There exists a constant $C_1^* > 0$, independent of $k \in \mathbb{N}$, such that*

$$\sup_{k \in \mathbb{N}} \sup_{L \in W_k} \max \left\{ \|L^{-1}\|_{\mathcal{L}(V_k^*, V_k)}, \|L^{*-1}\|_{\mathcal{L}(V_k^*, V_k)} \right\} \leq C_1^*. \quad (40)$$

Proof. It suffices to show that, for any $k \in \mathbb{N}$ and any operator $L \in W_k$ we have

$$\|L^{*-1}\|_{\mathcal{L}(V_k^*, V_k)} \leq C_1^*, \quad (41)$$

for some constant $C_1^* > 0$ independent of $k \in \mathbb{N}$. Once proved, an application of the Hanh–Banach theorem allows us to deduce that for any $k \in \mathbb{N}$ and any operator $L \in W_k$ we also have $\|L^{-1}\|_{\mathcal{L}(V_k^*, V_k)} \leq C_1^*$.

Let $k \in \mathbb{N}$ and $L \in W_k$ be given. We know by Lemma 6 that the operator L and their adjoint L^* exhibit the discrete maximum principle. Since V_k is a

finite dimensional vector space and $L : V_k \rightarrow V_k^*$ is a linear map, the discrete maximum principle ensures the invertibility of both L and L^* as maps from V_k to V_k^* . Moreover, for each $f \in V_k^*$ the unique solution $\bar{u}_k \in V_k$ to the equation $L^* \bar{u}_k = f$ in V_k^* satisfies the following Garding inequality for some constant $C^* > 0$ that is independent of k :

$$\|\bar{u}_k\|_{H^1(\Omega)} \leq C^* (\|f\|_{V_k^*} + \|\bar{u}_k\|_{\Omega}) \quad \forall f \in V_k^*. \quad (42)$$

Suppose for contradiction that (41) does not hold. Then for every integer $j \in \mathbb{N}$ there exists an integer $k_j \in \mathbb{N}$ and an operator $L_j \in W_{k_j}$ such that

$$\|L_j^{*-1}\|_{\mathcal{L}(V_{k_j}^*, V_{k_j})} > j,$$

with the sequence $\{k_j\}_{j \in \mathbb{N}}$ strictly increasing. This implies, together with (42), that there exist sequences $\{\bar{u}_j\}_{j \in \mathbb{N}} \subset H_0^1(\Omega)$, and $\{f_j\}_{j \in \mathbb{N}} \subset V_1^*$ such that

$$\bar{u}_j := L_j^{*-1} f_j \in V_{k_j}, \quad f_j \in V_{k_j}^*, \quad \|\bar{u}_j\|_{\Omega} = 1 \quad \forall j \in \mathbb{N}, \quad (43a)$$

$$\|f_j\|_{V_{k_j}^*} < \frac{C^*}{j - C^*} \quad \forall j > C^*. \quad (43b)$$

In particular, for all $j \in \mathbb{N}$, there holds

$$\langle L_j^* \bar{u}_j, v \rangle_{V_{k_j}^* \times V_{k_j}} = \int_{\Omega} (\nu + \gamma_{k_j}) \nabla \bar{u}_j \cdot \nabla v + \bar{u}_j \tilde{b}_j \cdot \nabla v + \kappa \bar{u}_j v \, dx = \langle f_j, v \rangle_{V_{k_j}^* \times V_{k_j}} \quad (44)$$

for all $v \in V_{k_j}$, with some \tilde{b}_j satisfying the uniform bound $\|\tilde{b}_j\|_{L^\infty(\Omega; \mathbb{R}^n)} \leq \|b\|_{C(\bar{\Omega} \times \mathcal{A})}$ by definition of the inclusion $L_j \in W_{k_j}$. Garding's inequality (42) and (43b) imply there exists a constant $C > 0$, independent of j , such that $\|\bar{u}_j\|_{H^1(\Omega)} \leq C$ for all $j \in \mathbb{N}$. Since the sequence $\{\tilde{b}_j\}_{j \in \mathbb{N}} \subset L^\infty(\Omega; \mathbb{R}^n)$ is uniformly bounded, we may pass to a subsequence without change of notation to get, as $j \rightarrow \infty$,

$$\bar{u}_j \rightharpoonup \bar{u} \text{ in } H_0^1(\Omega), \quad \bar{u}_j \rightarrow \bar{u} \text{ in } L^2(\Omega), \quad (45a)$$

$$\tilde{b}_j \rightharpoonup^* \tilde{b}_* \text{ in } L^\infty(\Omega; \mathbb{R}^n). \quad (45b)$$

Let $\phi \in V_l$ be given for some fixed $l \in \mathbb{N}$. By nestedness subspaces $\{V_k\}_{k \in \mathbb{N}}$ in $H_0^1(\Omega)$, we get by (44) that

$$\int_{\Omega} \nu \nabla \bar{u}_j \cdot \nabla v_l + \bar{u}_j \tilde{b}_j \cdot \nabla v_l + \kappa \bar{u}_j v_l \, dx + \int_{\Omega} \gamma_{k_j} \nabla \bar{u}_j \cdot \nabla v_l \, dx = \langle f_j, v_l \rangle_{V_{k_j}^* \times V_{k_j}} \quad (46)$$

for all $j \in \mathbb{N}$ such that $k_j \geq l$. We have for all $l \in \mathbb{N}$ that $\langle f_j, v_l \rangle_{V_{k_j}^* \times V_{k_j}} = 0$ as $j \rightarrow \infty$ by (43b). Moreover, uniform boundedness of the sequence $\{\bar{u}_j\}_{j \in \mathbb{N}}$ in $H_0^1(\Omega)$ and (37) imply that

$$\lim_{j \rightarrow \infty} \left| \int_{\Omega} \gamma_{k_j} \nabla \bar{u}_j \cdot \nabla v_l \, dx \right| = 0 \quad \forall l \in \mathbb{N}.$$

Recalling (45a), (45b), we send $j \rightarrow \infty$ in (46) to then obtain

$$\int_{\Omega} \nu \nabla \bar{u} \cdot \nabla v_l + \bar{u} \tilde{b}_* \cdot \nabla v_l + \kappa \bar{u} v_l \, dx = 0 \quad \forall l \in \mathbb{N}.$$

Since l was arbitrary, density of the union $\bigcup_{l \in \mathbb{N}} V_l$ in $H_0^1(\Omega)$ allows us to conclude

$$\int_{\Omega} \nu \nabla \bar{u} \cdot \nabla \phi + \bar{u} \tilde{b}_* \cdot \nabla \phi + \kappa \bar{u} \phi \, dx = 0 \quad \forall \phi \in H_0^1(\Omega).$$

Thus, \bar{u} solves an elliptic equation with an operator that is the adjoint of an operator from the class $\mathcal{G}(C_0)$ for some constant C_0 . Lemma 4 then implies that $\bar{u} = 0$ in $H_0^1(\Omega)$ necessarily, contradicting the fact that $\|\bar{u}\|_{\Omega} = 1$ which follows from (43a) and (45a). Hence, (40) is proved for some constant C_1^* independent of $k \in \mathbb{N}$, as required. \square

By using Lemma 7 and Schaefer's fixed point theorem, in a fashion similar to the proof of Lemma 5, we obtain a well-posedness result for the discrete HJB equation in (36).

Lemma 8. *Let $k \in \mathbb{N}$ be given. Then, for each $\bar{m} \in L^2(\Omega)$ there exists unique $u_k \in V_k$ such that*

$$\int_{\Omega} (\nu + \gamma_k) \nabla u_k \cdot \nabla \psi + H(x, \nabla u_k) \psi + \kappa u_k \psi \, dx = \langle F[\bar{m}], \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in V_k. \quad (47)$$

There exists a constant C_1^* independent of $k \in \mathbb{N}$ such that

$$\|u_k\|_{H^1(\Omega)} \leq C_1^* \left(\|\bar{m}\|_{\Omega} + \|f\|_{C(\bar{\Omega} \times \mathcal{A})} + 1 \right). \quad (48)$$

Moreover, the solution u_k depends continuously on \bar{m} , i.e., if $\{\bar{m}_j\}_{j \in \mathbb{N}} \subset L^2(\Omega)$ is such that $\bar{m}_j \rightarrow \bar{m}$ in $L^2(\Omega)$ as $j \rightarrow \infty$, then the corresponding sequence of solutions $\{\bar{u}_j\}_{j \in \mathbb{N}} \subset V_k$ to the equation (47) converges in V_k to the solution u_k of (47).

Proof of Theorem 3 and Theorem 4. Let $k \in \mathbb{N}$ be given. We deduce the existence of discrete solutions to (36) using Lemma 7 and Lemma 8 in an adaptation of the proof of Theorem 1. The uniform estimate (38a) follows immediately from the discrete KFP equation (36b) and the uniform bound (40). The bound (38b) then follows directly from (38a) and (48).

Uniqueness of solutions to (36) follows when F is strictly monotone and G is non-negative in the sense of distributions in $H^{-1}(\Omega)$, with the details being similar to the proof of Theorem 2 since the space V_k is a subspace of $H_0^1(\Omega)$. Indeed, this follows through since the linear differential operator featuring in the discrete KFP equation of (36) is the adjoint of an operator in the class W_k . Therefore, we have access to the discrete maximum principle via Lemma 6 that ensures non-negativity everywhere in Ω for the density approximation m_k . \square

6.3 Convergence

In this section we employ a compactness argument to prove the convergence result Theorem 5 when unique solutions to (9) are ensured under the hypotheses of Theorem 2. Note that, in addition, the following proof yields an alternative method of showing the existence of weak solutions to (4) in the sense of Definition 2.

Proof of Theorem 5. Let $\{(u_k, m_k)\}_{k \in \mathbb{N}}$ denote the sequence of solutions given by Theorem 4 with associated vector fields $\tilde{b}_k \in D_p H[u_k]$ ($k \in \mathbb{N}$). Since Theorem 3 indicates that the sequences $\{m_k\}_{k \in \mathbb{N}}$, $\{u_k\}_{k \in \mathbb{N}}$ are uniformly bounded in $H_0^1(\Omega)$, while $\{\tilde{b}_k\}_{k \in \mathbb{N}}$ is uniformly bounded in $L^\infty(\Omega; \mathbb{R}^n)$, by the Rellich–Kondrachov compactness theorem and Banach–Alaoglu theorem we may pass to subsequences, without change of notation, that satisfy as $k \rightarrow \infty$

$$m_k \rightharpoonup m \quad \text{in } H_0^1(\Omega), \quad m_k \rightarrow m \quad \text{in } L^q(\Omega), \quad (49)$$

$$u_k \rightharpoonup u \quad \text{in } H_0^1(\Omega), \quad u_k \rightarrow u \quad \text{in } L^q(\Omega), \quad (50)$$

$$\tilde{b}_k \rightharpoonup^* \tilde{b}_* \quad \text{in } L^\infty(\Omega; \mathbb{R}^n), \quad (51)$$

for some $m, u \in H_0^1(\Omega)$, some $\tilde{b}_* \in L^\infty(\Omega; \mathbb{R}^n)$, and for any $q \in [1, 2^*)$ where the critical exponent $2^* = \infty$ if $n = 2$ and $2^* = \frac{2n}{n-2}$ if $n \geq 3$. Notice in particular that $m_k \tilde{b}_k$ converges weakly to $m \tilde{b}_*$ in $L^2(\Omega; \mathbb{R}^n)$ as $k \rightarrow \infty$.

Let $v \in V_j$ be given, for some fixed $j \in \mathbb{N}$. Since the sequence $\{m_k\}_{k \in \mathbb{N}}$ satisfies

$$\int_{\Omega} (\nu + \gamma_k) \nabla m_k \cdot \nabla v + m_k \tilde{b}_k \cdot \nabla v + \kappa m_k v \, dx = \langle G, v \rangle_{H^{-1} \times H_0^1} \quad \forall k \geq j,$$

and we have the convergence given by (49) and (51), along with the uniform boundedness of the sequence $\{\nabla m_k\}_{k \in \mathbb{N}}$ in $L^2(\Omega; \mathbb{R}^n)$, and the vanishing of the artificial diffusion coefficients given by (37), we obtain in the limit as $k \rightarrow \infty$

$$\int_{\Omega} \nu \nabla m \cdot \nabla v + m \tilde{b}_* \cdot \nabla v + \kappa m v \, dx = \langle G, v \rangle_{H^{-1} \times H_0^1} \quad \forall v \in V_j.$$

Since j was arbitrary, density of the union $\bigcup_{j \in \mathbb{N}} V_j$ in $H_0^1(\Omega)$ allows us to conclude

$$\int_{\Omega} \nu \nabla m \cdot \nabla \phi + m \tilde{b}_* \cdot \nabla \phi + \kappa m \phi \, dx = \langle G, \phi \rangle_{H^{-1} \times H_0^1} \quad \forall \phi \in H_0^1(\Omega). \quad (52)$$

Next, observe that the boundedness of $\{u_k\}_{k \in \mathbb{N}}$ in $H_0^1(\Omega)$ and the linear p -growth of the Hamiltonian $H = H(x, p)$ imply that the sequence $\{H(\cdot, \nabla u_k)\}_{k \in \mathbb{N}}$ is bounded in $L^2(\Omega)$. Therefore, there exists $g \in L^2(\Omega)$ such that, by passing to a subsequence without change of notation, we have as $k \rightarrow \infty$

$$H(\cdot, \nabla u_k) \rightharpoonup g \quad \text{in } L^2(\Omega). \quad (53)$$

For fixed $j \in \mathbb{N}$, the definition of the sequence $\{u_k\}_{k \in \mathbb{N}}$ implies that for all $k \geq j$,

$$\int_{\Omega} (\nu + \gamma_k) \nabla u_k \cdot \nabla v + H(x, \nabla u_k) v + \kappa u_k v \, dx = \langle F[m_k], v \rangle_{H^{-1} \times H_0^1} \quad \forall v \in V_j. \quad (54)$$

Therefore, the convergence given by (49), (50), and (53), together with the vanishing of the artificial diffusion coefficients given by (37), and the uniform boundedness of the sequence $\{\nabla u_k\}_{k \in \mathbb{N}}$ in $L^2(\Omega; \mathbb{R}^n)$, imply that we obtain

$$\int_{\Omega} \nu \nabla u \cdot \nabla v + gv + \kappa uv \, dx = \langle F[m], v \rangle_{H^{-1} \times H_0^1} \quad \forall v \in V_j,$$

after sending $k \rightarrow \infty$ in (54). As j was arbitrary, we conclude from above that

$$\int_{\Omega} \nu \nabla u \cdot \nabla \psi + g\psi + \kappa u \psi \, dx = \langle F[m], \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in H_0^1(\Omega), \quad (55)$$

and in particular $\|\nabla u\|_{\Omega}^2 = \nu^{-1} \left(\langle F[m], u \rangle_{H^{-1} \times H_0^1} - \int_{\Omega} gu \, dx - \kappa \|u\|_{\Omega}^2 \right)$.

On the other hand, the definition of $\{u_k\}_{k \in \mathbb{N}}$ gives

$$\nu \|\nabla u_k\|_{\Omega}^2 + \int_{\Omega} H(x, \nabla u_k) u_k \, dx + \kappa \|u_k\|_{\Omega}^2 + \int_{\Omega} \gamma_k |\nabla u_k|^2 \, dx = \langle F[m_k], u_k \rangle_{H^{-1} \times H_0^1}$$

for each $k \in \mathbb{N}$. In view of the convergence given by (49), (50), and (53), along with the uniform boundedness of the sequence $\{\nabla u_k\}_{k \in \mathbb{N}}$ in $L^2(\Omega; \mathbb{R}^n)$, the vanishing of the artificial diffusion coefficients given by (37), and the Lipschitz continuity of the coupling term F , we find that

$$\lim_{k \rightarrow \infty} \|\nabla u_k\|_{\Omega}^2 = \nu^{-1} \left(\langle F[m], u \rangle_{H^{-1} \times H_0^1} - \int_{\Omega} gu \, dx - \kappa \|u\|_{\Omega}^2 \right) = \|\nabla u\|_{\Omega}^2. \quad (56)$$

Because (50) holds, we deduce via (56) that

$$\lim_{k \rightarrow \infty} \|u_k\|_{H^1(\Omega)} = \|u\|_{H^1(\Omega)}. \quad (57)$$

Since u_k converges weakly to u in $H_0^1(\Omega)$ by (50) and we have convergence of norms by (57), we deduce strong convergence in $H_0^1(\Omega)$: $u_k \rightarrow u$ as $k \rightarrow \infty$. Because the mapping $v \mapsto H(\cdot, \nabla v)$ is Lipschitz continuous from $H^1(\Omega)$ into $L^2(\Omega)$, it then follows that (53) is in fact strong convergence in $L^2(\Omega)$ with $g = H(\cdot, \nabla u)$. Hence, (55) gives the weak HJB equation:

$$\int_{\Omega} \nu \nabla u \cdot \nabla \psi + H(x, \nabla u) \psi + \kappa u \psi \, dx = \langle F[m], \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in H_0^1(\Omega). \quad (58)$$

To deduce convergence to a weak solution of (4), we need to show that $\tilde{b}_* \in D_p H[u]$ in addition to the established equations (52), (58). But this follows by applying Lemma 3 to $\{(\tilde{b}_k, u_k)\}_{k \in \mathbb{N}}$ after passing to an appropriate subsequence.

In summary, we have shown that a subsequence of the finite element approximations $\{(u_k, m_k)\}_{k \in \mathbb{N}}$ converges to a weak solution (u, m) of (4) in the sense that

$$u_k \rightarrow u \quad \text{in } H_0^1(\Omega), \quad m_k \rightarrow m \quad \text{in } L^q(\Omega), \quad m_k \rightharpoonup m \quad \text{in } H_0^1(\Omega), \quad (59)$$

as $k \rightarrow \infty$, for any $q \in [1, 2^*)$. But uniqueness of solutions to (9) then implies that the whole sequence $\{(u_k, m_k)\}_{k \in \mathbb{N}}$ exhibits this convergence towards the unique solution of (9), as required. \square

Under the additional hypothesis on the transport vector fields given in Corollary 1 we obtain strong convergence of the density approximations in $H_0^1(\Omega)$.

Proof of Corollary 1. To prove strong convergence of $\{m_k\}_{k \in \mathbb{N}}$ in $H_0^1(\Omega)$, let us consider an arbitrary subsequence $\{m_{k_j}\}_{j \in \mathbb{N}}$ with corresponding subsequence of transport vector fields $\{\tilde{b}_{k_j}\}_{j \in \mathbb{N}}$. By the stated hypotheses in the Corollary 1, together with Hölder's inequality and the fact that the sequence $\{\tilde{b}_{k_j}\}_{j \in \mathbb{N}}$ is uniformly bounded in $L^\infty(\Omega; \mathbb{R}^n)$, we deduce that there exists $s > n$ such that the sequence $\{\tilde{b}_{k_j}\}_{j \in \mathbb{N}}$ is pre-compact in $L^s(\Omega; \mathbb{R}^n)$. Therefore, there exists a subsequence of $\{\tilde{b}_{k_j}\}_{j \in \mathbb{N}}$, to which we pass without change of notation, that converges in $L^s(\Omega; \mathbb{R}^n)$, $s > n$, to a Lebesgue measurable vector field $\bar{b} : \Omega \rightarrow \mathbb{R}^n$ that is in $L^\infty(\Omega; \mathbb{R}^n)$. From Theorem 5, we know that $m_{k_j} \rightarrow m$ in $L^r(\Omega)$ as $j \rightarrow \infty$ for any $r \in [1, 2^*)$ (where we recall $2^* = \frac{2n}{n-2}$ when $n \geq 3$ and $2^* = \infty$ when $n = 2$). Hence, by Hölder's inequality and a suitable choice of $r \in [1, 2^*)$, we deduce $m_{k_j} \tilde{b}_{k_j} \rightarrow m \bar{b}$ strongly in $L^2(\Omega; \mathbb{R}^n)$ as $j \rightarrow \infty$. Consequently, the weak convergence of $\{m_{k_j}\}_{j \in \mathbb{N}}$ to m in $H_0^1(\Omega)$ and hence weak convergence of the gradients $\{\nabla m_{k_j}\}_{j \in \mathbb{N}}$ to ∇m in $L^2(\Omega)$ implies that

$$\lim_{j \rightarrow \infty} \int_{\Omega} m_{k_j} \tilde{b}_{k_j} \cdot \nabla m_{k_j} \, dx = \int_{\Omega} m \bar{b} \cdot \nabla m \, dx. \quad (60)$$

From the discrete KFP equation (36b), we obtain

$$\int_{\Omega} (\nu + \gamma_{k_j}) |\nabla m_{k_j}|^2 \, dx + \int_{\Omega} m_{k_j} \tilde{b}_{k_j} \cdot \nabla m_{k_j} \, dx + \kappa \|m_{k_j}\|_{\Omega}^2 = \langle G, m_{k_j} \rangle_{H^{-1} \times H_0^1}. \quad (61)$$

We deduce from (49), (38a), (37), and (61) that

$$\lim_{j \rightarrow \infty} \|\nabla m_{k_j}\|_{\Omega}^2 = \nu^{-1} \left(\langle G, m \rangle_{H^{-1} \times H_0^1} - \kappa \|m\|_{\Omega}^2 - \int_{\Omega} m \bar{b} \cdot \nabla m \, dx \right). \quad (62)$$

We also deduce from the discrete KFP equation (36b) that \bar{b} and m satisfy

$$\int_{\Omega} \nu \nabla m \cdot \nabla \phi + m \bar{b} \cdot \nabla \phi + \kappa m \phi \, dx = \langle G, \phi \rangle_{H^{-1} \times H_0^1} \quad \forall \phi \in H_0^1(\Omega).$$

Hence,

$$\|\nabla m\|_{\Omega}^2 = \nu^{-1} \left(\langle G, m \rangle_{H^{-1} \times H_0^1} - \kappa \|m\|_{\Omega}^2 - \int_{\Omega} m \bar{b} \cdot \nabla m \, dx \right). \quad (63)$$

We thus obtain from (62) and (63), together with the fact that $m_{k_j} \rightarrow m$ in $L^2(\Omega)$ as $j \rightarrow \infty$, that

$$\lim_{j \rightarrow \infty} \|m_{k_j}\|_{H^1(\Omega)} = \|m\|_{H^1(\Omega)}. \quad (64)$$

Since m_{k_j} converges weakly to m in $H_0^1(\Omega)$ by Theorem 5 and we have convergence of norms by (64), we deduce that $m_{k_j} \rightarrow m$ as $j \rightarrow \infty$ strongly in $H_0^1(\Omega)$. The argument above shows that any subsequence of $\{m_k\}_{k \in \mathbb{N}}$ has a further subsequence that converges to m in $H_0^1(\Omega)$, and thus the whole sequence is convergent. This completes the proof. \square

Remark 6 (Convergence in Hölder Norms). When the space dimension $n = 2$ and the domain Ω is convex, one can derive a uniform Hölder-norm bound for the approximating sequence $\{m_k\}_{k \in \mathbb{N}}$ (see [30, Theorem 3.20]). It follows that $\{m_k\}_{k \in \mathbb{N}}$ converges strongly to m in some Hölder space. This likewise holds for the corresponding sequence of value function approximations $\{u_k\}_{k \in \mathbb{N}}$.

7 Numerical Experiments

The numerical experiments shown below are designed to study the performance of the method for problems with nonsmooth solutions, and when relaxing the condition on the meshes to being weakly acute rather than strictly acute.

7.1 Set-up

For each numerical experiment we take the domain Ω of the continuous problem (9) to be the unit square. The computations are performed on a sequence of uniform, conforming, shape-regular, weakly acute meshes on Ω . The artificial diffusion coefficient we use in each experiment is identically zero. The choice of Hamiltonian $H : \mathbb{R}^2 \rightarrow \mathbb{R}$ for each experiment is set to be

$$H(x, p) := \max_{\alpha \in \overline{B_1(0)}} (\alpha \cdot p) = |p| \quad \forall (x, p) \in \overline{\Omega} \times \mathbb{R}^2. \quad (65)$$

It is clear from (65) that $b(x, \alpha) := \alpha$ for $(x, \alpha) \in \overline{\Omega} \times \overline{B_1(0)} \subset \mathbb{R}^2$ and hence $\|b\|_{C(\overline{\Omega} \times \mathcal{A})} = 1$. Moreover, the subdifferential $\partial_p H : \mathbb{R}^2 \rightrightarrows \mathbb{R}^2$ is given by

$$\partial_p H(p) = \begin{cases} \left\{ \frac{1}{|p|} p \right\} & \text{if } p \neq 0, \\ \overline{B_1(0)} & \text{if } p = 0. \end{cases} \quad (66)$$

In each experiment we consider the continuous problem (9) where we assume the diffusion to be $\nu = 1$ and the reaction coefficient to be $\kappa = 0$. The choices of the coupling term F and source $G \in H^{-1}(\Omega)$ differ between both experiments.

We employ a fixed-point approximation of (36) that follows the fixed point process described in the proof of Theorem 3. In each iteration we approximate the discrete HJB equation (36a) via a policy iteration method and we resolve the linear system resulting from the discrete KFP equation (36b) via LU factorisation. For an introduction to policy iteration in general we refer the reader to [1, 2]. We used the open-source finite element software Firedrake [26] to perform the computations.

7.2 First Experiment

We consider the approximation of a known solution pair that uniquely satisfies (9) with suitable coupling term F and source term $G \in H^{-1}(\Omega)$. For this experiment we take the coupling term $F : L^2(\Omega) \rightarrow H^{-1}(\Omega)$ defined via

$$\langle F[v], \psi \rangle_{H^{-1} \times H_0^1} := \int_{\Omega} \tanh(v) \psi \, dx + \langle J, \psi \rangle_{H^{-1} \times H_0^1} \quad \forall \psi \in H_0^1(\Omega), \forall v \in L^2(\Omega),$$

where the functional $J \in H^{-1}(\Omega)$ is given by

$$\langle J, \psi \rangle_{H^{-1} \times H_0^1} := \int_{\Omega} h \psi + \tilde{t} \cdot \nabla \psi \, dx \quad \forall \psi \in H_0^1(\Omega),$$

with $\tilde{t} : \Omega \rightarrow \mathbb{R}^2$ and $h : \Omega \rightarrow \mathbb{R}$ defined by

$$\tilde{t}(x, y) := \begin{pmatrix} (1 + \log(x))y \log(y) \\ (1 + \log(y))x \log(x) \end{pmatrix}, \quad h(x, y) := |\tilde{t}(x, y)| - \tanh(xy(1-x)(1-y)).$$

Note that $\tilde{t} \in L^2(\Omega; \mathbb{R}^2)$ and $h \in L^2(\Omega)$ so indeed $J \in H^{-1}(\Omega)$. Next, we take the source term $G \in H^{-1}(\Omega)$ to be

$$\langle G, \phi \rangle_{H^{-1} \times H_0^1} := \int_{\Omega} r \phi + \tilde{g} \cdot \nabla \phi \, dx \quad \forall \phi \in H_0^1(\Omega),$$

where $r : \Omega \rightarrow \mathbb{R}$ is given by $r(x, y) := 2(x(1-x) + y(1-y))$ and $\tilde{g} : \Omega \rightarrow \mathbb{R}^2$ is the vector field whose image is $\frac{xy(1-x)(1-y)}{|\tilde{t}|} \tilde{t}$ whenever $\tilde{t}(x, y) \neq (0, 0)^T$ and is $(0, 0)^T$ whenever $\tilde{t}(x, y) = (0, 0)^T$. It can be shown that F is strictly monotone and that $\langle G, \phi \rangle_{H^{-1} \times H_0^1} \geq 0$ if $\phi \in H_0^1(\Omega)$ is non-negative a.e. in Ω . Therefore, (9) is indeed well-posed according to Theorem 2. Moreover, the unique solution (u, m) to (9) in this case is given by

$$u(x, y) := xy \log(x) \log(y), \quad m(x, y) := xy(1-x)(1-y), \quad \forall (x, y) \in \Omega. \quad (67)$$

In Figure 1 we plot the relative errors in various norms versus the mesh-size h for a sequence of finite element approximations obtained from (36). Observe that strong convergence is seen in each convergence plot. In particular, we see that in the H^1 -norm the approximations of the value function converge at a slower rate than the approximations of the density function. This is due to the fact that u has lower regularity compared to m : $m \in C^\infty(\Omega) \cap C(\bar{\Omega})$ but




Figure 1: First Experiment – Convergence plots for approximations of the value function, density function, and transport vector. Almost optimal rates of convergence are observed for the value function and density function errors in the H^1 -norm.

$u \notin H^2(\Omega)$. In fact, u is in the Besov space $H^{\frac{3}{2}-\epsilon}(\Omega)$ for all arbitrary small $0 < \epsilon < \frac{1}{2}$. We thus see that the convergence rate of the H^1 -norm error of the value function approximation is close to the optimal value of $\frac{1}{2}$. The convergence rate of the H^1 -norm error of the density approximations is close to the maximal rate allowed by continuous P1 finite element approximations. Moreover, the transport vector field approximations converge strongly in the L^2 -norm. This is likely due to the transport vector field approximations converging a.e. to the vector field $\frac{1}{m}\tilde{g}$, along with the fact that the gradient $\nabla u \neq 0$ a.e. in Ω .

7.3 Second Experiment

For this experiment we choose the coupling term F and source $G \in H^{-1}(\Omega)$ in such a way that the conditions of Theorem 2 hold so that the MFG PDI (9) admits a unique solution. In this case, we set the coupling term $F : L^2(\Omega) \rightarrow H^{-1}(\Omega)$ via

$$\langle F[v], \psi \rangle_{H^{-1} \times H_0^1} := \int_{\Omega} (\arctan(v) + 2\operatorname{sgn}((x - 0.5) \cos(8\pi y))) \psi \, dx$$

for all $\psi \in H_0^1(\Omega)$ and $v \in L^2(\Omega)$, and we let $G \in H^{-1}(\Omega)$ be given by

$$\langle G, \phi \rangle_{H^{-1} \times H_0^1} := \int_{\Omega} \frac{1}{2} (\operatorname{sgn}(\sin(4\pi x) \sin(4\pi y)) + 1) \phi + \tilde{c} \cdot \nabla \phi \, dx \quad \forall \phi \in H_0^1(\Omega),$$

where $\tilde{c} : \Omega \rightarrow \mathbb{R}^2$ is defined by

$$\tilde{c}(x, y) := \begin{cases} y(1, 0)^T & \text{if } 0 < x < 2/3, \\ y^2(-1, 0)^T & \text{otherwise.} \end{cases}$$

It is easy to show that F is strictly monotone and that $\langle G, \phi \rangle_{H^{-1} \times H_0^1} \geq 0$ for all $\phi \in H_0^1(\Omega)$ that is non-negative a.e. in Ω . The exact solution is not known, so we measure the errors using a computed reference solution on a very fine mesh. To illustrate the discrete reference solution (u, m) that we will consider, in Figure 2 we display contour plots of a discrete approximation for (u, m) that we computed on a fine mesh. We note that the displayed approximation for m is non-negative everywhere in Ω .

In Figure 3 relative errors versus mesh size h are plotted. These results were obtained with respect to a discrete reference solution (u, m) that was computed on a much finer mesh than the approximation displayed in Figure 2. The convergence rate of the value function approximations is now of optimal rate 1 since u is smooth. The convergence rate of H^1 -norm errors of the approximation of the density is now close to $\frac{1}{2}$, which is again optimal given the limited regularity of the density function m as a result of the line singularity in the right-hand side source term G .




Figure 2: Second Experiment – Approximate Contour plot of u (Left), m (Right) computed on a fine mesh




Figure 3: Second Experiment – Convergence plots for approximations of the value function, density function, and transport vector. The optimal rate of convergence is observed for the value function errors in the H^1 -norm, with an approximate rate of $\frac{1}{2}$ seen in the H^1 -norm errors of the density approximations.

8 Conclusions

We have introduced the notion of Mean Field Game Partial Differential Inclusions (MFG PDIs) as a generalization of Mean Field Game systems with possibly nondifferentiable Hamiltonians. We established the existence and uniqueness of weak solutions of the MFG PDI system (4) under appropriate hypotheses. We proved the well-posedness and convergence of numerical approximations of the system by a monotone FEM. The numerical experiments suggest that the convergence rates of the errors for the different functions u and m seem to be optimal, even in the case of non-smooth solution pairs. Interestingly, the experiments suggest that the lower regularity in one of the two functions only seems to affect the convergence rate for the error of the approximations of that function, but not in the other. This is of significant interest for further work on the analysis of convergence rates of the method.

Acknowledgments

We would like to acknowledge the assistance of Jack Betteridge in enabling the use of Firedrake on the UCL Myriad High Performance Computing Facility (Myriad@UCL). Moreover, the authors acknowledge the use of the UCL Myriad High Performance Computing Facility (Myriad@UCL), and associated support services, in the completion of this work.

References

- [1] Richard Bellman. “The theory of dynamic programming”. In: *Bulletin of the American Mathematical Society* 60.6 (1954), pp. 503–515.
- [2] Ronald A Howard. *Dynamic Programming and Markov Processes*. Cambridge, Massachusetts: The M.I.T. Press, 1960.
- [3] Kazimierz Kuratowski and Czesław Ryll-Nardzewski. “A general theorem on selectors”. In: *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys* 13.6 (1965), pp. 397–403.
- [4] Philippe G Ciarlet and P-A Raviart. “Maximum principle and uniform convergence for the finite element method”. In: *Computer methods in applied mechanics and engineering* 2.1 (1973), pp. 17–31. DOI: 10.1016/0045-7825(73)90019-4.
- [5] Kinji Baba and Masahisa Tabata. “On a conservative upwind finite element scheme for convective diffusion equations”. en. In: *ESAIM: Mathematical Modelling and Numerical Analysis - Modélisation Mathématique et Analyse Numérique* 15.1 (1981), pp. 3–25. URL: http://www.numdam.org/item/M2AN_1981__15_1_3_0/.

- [6] Akira Mizukami and Thomas J.R. Hughes. “A Petrov-Galerkin finite element method for convection-dominated flows: An accurate upwinding technique for satisfying the maximum principle”. In: *Computer Methods in Applied Mechanics and Engineering* 50.2 (1985), pp. 181–193. ISSN: 0045-7825. DOI: 10.1016/0045-7825(85)90089-1.
- [7] Jean-Pierre Aubin. *Optima and Equilibria: An Introduction to Nonlinear Analysis*. Vol. 140. Springer-Verlag, 1993.
- [8] Jinchao Xu and Ludmil Zikatanov. “A monotone finite element scheme for convection-diffusion equations”. In: *Mathematics of Computation* 68.228 (1999), pp. 1429–1446. DOI: 10.1090/S0025-5718-99-01148-5.
- [9] David Gilbarg and Neil S Trudinger. *Elliptic Partial Differential Equations of Second Order*. Springer, 2001. DOI: 10.1007/978-3-642-61798-0.
- [10] Erik Burman and Alexandre Ern. “Nonlinear diffusion and discrete maximum principle for stabilized Galerkin approximations of the convection-diffusion-reaction equation”. In: *Computer Methods in Applied Mechanics and Engineering* 191.35 (2002), pp. 3833–3855. DOI: 10.1016/S0045-7825(02)00318-3.
- [11] Minyi Huang, Roland P Malhamé, and Peter E Caines. “Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle”. In: *Communications in Information & Systems* 6.3 (2006), pp. 221–252.
- [12] Jean-Michel Lasry and Pierre-Louis Lions. “Jeux à champ moyen. I—Le cas stationnaire”. In: *Comptes Rendus Mathématique* 343.9 (2006), pp. 619–625. DOI: 10.1016/j.crma.2006.09.019.
- [13] Jean-Michel Lasry and Pierre-Louis Lions. “Jeux à champ moyen. II—Horizon fini et contrôle optimal”. In: *Comptes Rendus Mathématique* 343.10 (2006), pp. 679–684. DOI: 10.1016/j.crma.2006.09.018.
- [14] Jean-Michel Lasry and Pierre-Louis Lions. “Mean field games”. In: *Japanese journal of mathematics* 2.1 (2007), pp. 229–260. DOI: 10.1007/s11537-007-0657-8.
- [15] Yves Achdou and Italo Capuzzo-Dolcetta. “Mean Field Games: Numerical Methods”. In: *SIAM Journal on Numerical Analysis* 48.3 (2010), pp. 1136–1162. DOI: 10.1137/090758477.
- [16] Lawrence C Evans. *Partial Differential Equations*. Vol. 19 of Graduate Studies in Mathematics. American Mathematical Society, 2010.
- [17] Daniele Antonio Di Pietro and Alexandre Ern. *Mathematical aspects of discontinuous Galerkin methods*. Vol. 69. Springer Science & Business Media, 2011. DOI: 10.1007/978-3-642-22980-0.
- [18] Olivier Guéant, Jean-Michel Lasry, and Pierre-Louis Lions. “Mean field games and applications”. In: *Paris-Princeton Lectures on Mathematical Finance 2010*. Vol. 2003. Lecture Notes in Math. Springer, Berlin, 2011, pp. 205–266. DOI: 10.1007/978-3-642-14660-2_3. URL: https://doi.org/10.1007/978-3-642-14660-2_3.

- [19] Yves Achdou, Fabio Camilli, and Italo Capuzzo-Dolcetta. “Mean Field Games: Convergence of a Finite Difference Method”. In: *SIAM Journal on Numerical Analysis* 51.5 (2013), pp. 2585–2612. DOI: 10.1137/120882421.
- [20] Max Jensen and Iain Smears. “On the convergence of finite element methods for Hamilton-Jacobi-Bellman equations”. In: *SIAM J. Numer. Anal.* 51.1 (2013), pp. 137–162. ISSN: 0036-1429. DOI: 10.1137/110856198. URL: <https://doi.org/10.1137/110856198>.
- [21] Yves Achdou et al. “Partial differential equation models in macroeconomics”. In: *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* 372.2028 (2014), pp. 20130397, 19. ISSN: 1364-503X. DOI: 10.1098/rsta.2013.0397. URL: <https://doi.org/10.1098/rsta.2013.0397>.
- [22] Diogo A. Gomes and João Saúde. “Mean field games models—a brief survey”. In: *Dyn. Games Appl.* 4.2 (2014), pp. 110–154. ISSN: 2153-0785. DOI: 10.1007/s13235-013-0099-2. URL: <https://doi.org/10.1007/s13235-013-0099-2>.
- [23] Iain Smears and Endre Suli. “Discontinuous Galerkin finite element approximation of Hamilton–Jacobi–Bellman equations with Cordes coefficients”. In: *SIAM Journal on Numerical Analysis* 52.2 (2014), pp. 993–1016. DOI: 10.1137/130909536.
- [24] Yves Achdou and Alessio Porretta. “Convergence of a finite difference scheme to weak solutions of the system of partial differential equations arising in mean field games”. In: *SIAM Journal on Numerical Analysis* 54.1 (2016), pp. 161–186. DOI: 10.1137/15M1015455.
- [25] Diogo A Gomes, Edgard A Pimentel, and Vardan Voskanyan. *Regularity theory for mean-field game systems*. Springer, 2016. DOI: 10.1007/978-3-319-38934-9.
- [26] Florian Rathgeber et al. “Firedrake: automating the finite element method by composing abstractions”. In: *ACM Transactions on Mathematical Software (TOMS)* 43.3 (2016), pp. 1–27. DOI: 10.1145/2998441.
- [27] Roman Andreev. “Preconditioning the augmented Lagrangian method for instationary mean field games with diffusion”. In: *SIAM Journal on Scientific Computing* 39.6 (2017), A2763–A2783. DOI: 10.1137/16M1072346.
- [28] Max Jensen. “ $L^2(H_\gamma^1)$ finite element convergence for degenerate isotropic Hamilton-Jacobi-Bellman equations”. In: *IMA J. Numer. Anal.* 37.3 (2017), pp. 1300–1316. ISSN: 0272-4979. DOI: 10.1093/imanum/drw055. URL: <https://doi.org/10.1093/imanum/drw055>.
- [29] L. M. Briceño-Arias, D. Kalise, and F. J. Silva. “Proximal methods for stationary mean field games with local couplings”. In: *SIAM J. Control Optim.* 56.2 (2018), pp. 801–836. ISSN: 0363-0129. DOI: 10.1137/16M1095615. URL: <https://doi.org/10.1137/16M1095615>.

- [30] Seungchan Ko, Petra Pustějovská, and Endre Süli. “Finite element approximation of an incompressible chemically reacting non-Newtonian fluid”. In: *ESAIM: Mathematical Modelling and Numerical Analysis* 52.2 (2018), pp. 509–541. DOI: 10.1051/m2an/2017043.
- [31] Yves Achdou et al. *Mean Field Games*. Vol. 2281. Springer, 2019. DOI: 10.1007/978-3-030-59837-2.
- [32] Rita Ferreira, Diogo Gomes, and Teruo Tada. “Existence of weak solutions to first-order stationary mean-field games with Dirichlet conditions”. In: *Proceedings of the American Mathematical Society* 147.11 (2019), pp. 4713–4731. DOI: 10.1090/proc/14475.
- [33] Romain Ducasse, Guilherme Mazanti, and Filippo Santambrogio. “Second order local minimal-time Mean Field Games”. In: *Nonlinear Differ. Equ. Appl.* 29.37 (2022). DOI: 10.1007/s00030-022-00767-2.
- [34] Bartosz Jaroszkowski and Max Jensen. “Finite element methods for isotropic Isaacs equations with viscosity and strong Dirichlet boundary conditions”. In: *Appl. Math. Optim.* 85.2 (2022), Paper No. 8, 32. ISSN: 0095-4616. DOI: 10.1007/s00245-022-09860-5. URL: <https://doi.org/10.1007/s00245-022-09860-5>.