

# SEVERAL INTEGRAL ESTIMATES AND SOME APPLICATIONS

XUEJUN ZHANG \*, HONGXIN CHEN, MIN ZHOU, YUTING GUO, PENGCHENG TANG

**ABSTRACT.** In this paper, the authors first consider the bidirectional estimates of several typical integrals. As some applications of these integral estimates, the authors investigate the pointwise multipliers from the normal weight general function space  $F(p, \mu, s)$  to the normal weight Bloch type space  $\mathcal{B}_\nu(B_n)$  on the unit ball  $B_n$  of  $\mathbb{C}^n$ , where  $\mu$  and  $\nu$  are two normal functions on  $[0, 1)$ . For the special normal function  $\mu(r) = (1 - r^2)^\alpha \log^\beta \frac{e}{1 - r^2}$  ( $\alpha > 0$ ,  $-\infty < \beta < \infty$ ), the authors give the necessary and sufficient conditions of pointwise multipliers from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  for all cases.

## 1. Introduction

In this paper, we call “ $P \asymp Q$ ” if there exist two positive constants  $A_1$  and  $A_2$  such that  $A_1 P \leq Q \leq A_2 P$ . We call “ $Q \lesssim P$ ” (or “ $Q \gtrsim P$ ”) if there exists a positive constant  $A$  such that  $Q \leq AP$  (or  $Q \geq AP$ ).

Let  $\mathbb{C}$  denote the set of complex numbers. Fixed a positive integer  $n$ , let  $\mathbb{C}^n = \mathbb{C} \times \cdots \times \mathbb{C}$  denote the Euclidean space of complex dimension  $n$ . For  $z = (z_1, \dots, z_n)$  and  $w = (w_1, \dots, w_n)$  in  $\mathbb{C}^n$ , the inner product of  $z$  and  $w$  is defined by  $\langle z, w \rangle = z_1 \overline{w_1} + \cdots + z_n \overline{w_n}$ . The open unit ball in  $\mathbb{C}^n$  is the set  $B_n = \{z \in \mathbb{C}^n : |z| = \sqrt{\langle z, z \rangle} < 1\}$  (we write as  $D$  when  $n = 1$ ), and the boundary of  $B_n$  is the set  $S_n = \{z \in \mathbb{C}^n : |z| = 1\}$ . The class of holomorphic functions and the space of bounded holomorphic functions on  $B_n$  are denoted by  $H(B_n)$  and  $H^\infty(B_n)$ , respectively. For  $a \in B_n$ , let  $\varphi_a$  be the holomorphic automorphism of  $B_n$  satisfying with  $\varphi_a(0) = a$ ,  $\varphi_a(a) = 0$  and  $\varphi_a^{-1} = \varphi_a$ . It follows from Lemma 1.3 in [1] that

$$1 - \langle \varphi_a(z), \varphi_a(w) \rangle = \frac{(1 - |a|^2)(1 - \langle z, w \rangle)}{(1 - \langle z, a \rangle)(1 - \langle a, w \rangle)} \quad (z, w \in B_n). \quad (1.1)$$

In particular, if  $w = z$  or  $w = 0$ , then there are

$$1 - |\varphi_a(z)|^2 = \frac{(1 - |a|^2)(1 - |z|^2)}{|1 - \langle z, a \rangle|^2}, \quad 1 - \langle \varphi_a(z), a \rangle = \frac{1 - |a|^2}{1 - \langle z, a \rangle}. \quad (1.2)$$

---

2010 *Mathematics Subject Classification.* 32A37; 47B38.

*Key words and phrases.* Integral estimate;  $F(p, \mu, s)$  space; normal weight Bloch type space; pointwise multiplier; unit ball.

\* Corresponding author.

The research is supported by the National Natural Science Foundation of China (No. 11942109) and the Natural Science Foundation of Hunan Province (No. 2022JJ30369).

Given  $r > 0$ , the Bergman ball with  $a$  as the center and  $r$  as the radius is the set

$$D(a, r) = \{z \in B_n : \beta(z, a) < r\}, \quad \text{where } \beta(z, a) = \frac{1}{2} \log \frac{1 + |\varphi_a(z)|}{1 - |\varphi_a(z)|}.$$

1996, R. H. Zhao first introduced the general function space  $F(p, q, s)$  on  $D$  (see [2]). Soon after, the space  $F(p, q, s)$  was extended to  $B_n$ . If  $f \in F(p, q, s)$ , then

$$\begin{aligned} & \sup_{w \in B_n} \int_{B_n} |\nabla f(z)|^p (1 - |z|^2)^q \log^s \frac{1}{|\varphi_w(z)|} dv(z) \\ & \asymp \sup_{w \in B_n} (1 - |w|^2)^s \int_{B_n} \frac{(1 - |z|^2)^{q+s} |\nabla f(z)|^p}{|1 - \langle z, w \rangle|^{2s}} dv(z), \end{aligned}$$

where  $\nabla f(z) = \left( \frac{\partial f}{\partial z_1}(z), \dots, \frac{\partial f}{\partial z_n}(z) \right)$  (see [3]).

In recent years, there have been a lot of results related to  $F(p, q, s)$  space, such as [2]-[18] etc. In order to consider the general function space in a broader field of vision, we first give the definition of normal function.

**Definition 1.1** A positive continuous function on  $[0, 1)$  is called normal if there exist constants  $0 < a \leq b < \infty$  and  $0 \leq r_0 < 1$  such that

$$(i) \quad \frac{\mu(r)}{(1 - r^2)^a} \text{ is decreasing on } [r_0, 1); \quad (ii) \quad \frac{\mu(r)}{(1 - r^2)^b} \text{ is increasing on } [r_0, 1).$$

For example,  $\mu(r) = (1 - r^2)^\alpha \log^\beta \frac{e}{1 - r^2} \left\{ \log \log \frac{e^2}{1 - r^2} \right\}^\gamma$  ( $\alpha > 0, -\infty < \beta, \gamma < \infty$ ) is this kind of normal function.

Without losing generality, let  $r_0 = 0$  in this paper.

Recently, S. L. Li ([4]) extended  $F(p, q, s)$  to a kind of abstract form as follows:

**Definition 1.2** Let  $\mu$  be a normal function on  $[0, 1)$ . For  $p > 0$  and  $s \geq 0$ , if  $f \in H(B_n)$

$$\text{and } \|f\|_{p, \mu, s} = |f(0)| + \left\{ \sup_{w \in B_n} \int_{B_n} \frac{(1 - |w|^2)^s |\nabla f(z)|^p \mu^p(|z|)}{|1 - \langle z, w \rangle|^{2s}} \frac{1}{1 - |z|^2} dv(z) \right\}^{\frac{1}{p}} < \infty,$$

then  $f$  is said to belong to the normal weight general function space  $F(p, \mu, s)$ .

It is easy to prove that  $F(p, \mu, s)$  is a Banach space with the norm  $\|\cdot\|_{p, \mu, s}$  when  $p \geq 1$ , and  $F(p, \mu, s)$  is a complete distance space with  $\rho(f, g) = \|f - g\|_{p, \mu, s}^p$  when  $0 < p < 1$ .

This paper also involves the normal weight Bloch type space and the weighted Bergman space. We give their definitions, respectively.

**Definition 1.3** Let  $\mu$  be a normal function on  $[0, 1)$ . The normal weight Bloch type space  $\mathcal{B}_\mu(B_n)$  consists of holomorphic function  $f$  in  $B_n$  such that

$$\|f\|_{\mathcal{B}_\mu} = |f(0)| + \sup_{z \in B_n} \mu(|z|) |\nabla f(z)| < \infty.$$

The space  $\mathcal{B}_\mu(B_n)$  is a Banach space under the norm  $||\cdot||_{\mathcal{B}_\mu}$ . In particular,  $\mathcal{B}_\mu(B_n)$  is just the  $\alpha$ -Bloch space  $\mathcal{B}^\alpha(B_n)$  when  $\mu(r) = (1 - r^2)^\alpha$  ( $\alpha > 0$ ).

**Definition 1.4** For  $\alpha > -1$  and  $p > 0$ , the weighted Bergman space  $A_\alpha^p(B_n)$  consists of holomorphic function  $f$  in  $B_n$  such that  $\int_{B_n} |f(z)|^p dv_\alpha(z) < \infty$ , where  $dv_\alpha(z) = c_\alpha(1 - |z|^2)^\alpha dv(z)$ ,  $c_\alpha = \Gamma(n + 1 + \alpha)/n!\Gamma(\alpha + 1)$  and  $v(B_n) = 1$ .

It is known that the application of integral estimate is a basic tool for the study of function space theory and operator theory. Next, we introduce several integral estimates.

For a point in  $B_n$ , W. Rudin gave the following Proposition in [19]:

**Proposition A** Let  $t > -1$  and  $c$  be real. Then the integrals

$$I(z) = \int_{S_n} \frac{d\sigma(\xi)}{|1 - \langle \xi, z \rangle|^{n+c}} \quad \text{and} \quad J(z) = \int_{B_n} \frac{(1 - |w|^2)^t}{|1 - \langle z, w \rangle|^{n+1+t+c}} dv(w) \quad (z \in B_n)$$

have the following asymptotic properties:

- (1)  $I(z) \asymp J(z) \asymp 1$  when  $c < 0$ . (2)  $I(z) \asymp J(z) \asymp \log \frac{e}{1 - |z|^2}$  when  $c = 0$ .
- (3)  $I(z) \asymp J(z) \asymp \frac{1}{(1 - |z|^2)^c}$  when  $c > 0$ .

For two points in  $B_n$ , the following Proposition B comes from Lemma 2.2 in [10] (case  $k = 0$ ) and Theorem 3.1 in [12] (case  $k > 0$ ), and the following Proposition C comes from Proposition 3.1 in [36].

**Proposition B** Let  $\delta > -1$ ,  $r \geq 0$ ,  $t \geq 0$  and  $k \geq 0$ . Then the integral

$$J_{w,a} = \int_B \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^t |1 - \langle z, a \rangle|^r} \log^k \frac{e}{1 - |z|^2} dv(z) \quad (w, a \in B_n)$$

has the following bidirectional estimates:

- (1)  $J_{w,a} \asymp 1$  when  $t + r - \delta < n + 1$ .
- (2)  $J_{w,a} \asymp \log^{k+1} \frac{e}{|1 - \langle w, a \rangle|}$  when  $t + r - \delta = n + 1$ ,  $t > 0$  and  $r > 0$ .

$$(3) \quad J_{w,a} \asymp \frac{1}{|1 - \langle w, a \rangle|^r} \log^k \frac{e}{1 - |w|^2} \log \frac{e}{|1 - \langle w, \varphi_w(a) \rangle|}$$

when  $t - \delta = n + 1 > r - \delta$  and  $r > 0$ .

$$(4) \quad J_{w,a} \asymp \frac{1}{|1 - \langle w, a \rangle|^{t+r-\delta-n-1}} \log^k \frac{e}{|1 - \langle w, a \rangle|}$$

when  $t + r - \delta > n + 1 > \max\{r - \delta, t - \delta\}$ .

$$(5) \quad J_{w,a} \asymp \frac{1}{|1 - \langle w, a \rangle|^{\delta+n+1}} \log^k \frac{e}{1 - |w|^2} \log \frac{e}{|1 - \langle w, \varphi_w(a) \rangle|} \\ + \frac{1}{|1 - \langle w, a \rangle|^{\delta+n+1}} \log^k \frac{e}{1 - |a|^2} \log \frac{e}{|1 - \langle a, \varphi_a(w) \rangle|} \quad \text{when } t - \delta = n + 1 = r - \delta.$$

$$(6) \quad I_{w,a} \asymp \frac{1}{(1 - |w|^2)^{t-\delta-n-1} |1 - \langle w, a \rangle|^r} \log^k \frac{e}{1 - |w|^2} \quad \text{when } t - \delta > n + 1 > r - \delta.$$

$$(7) \quad I_{w,a} \asymp \frac{(1-|w|^2)^{n+1+\delta-t}}{|1-\langle w,a \rangle|^r} \log^k \frac{e}{1-|w|^2} + \frac{(1-|a|^2)^{n+1+\delta-r}}{|1-\langle w,a \rangle|^t} \log^k \frac{e}{1-|a|^2}$$

when  $t - \delta > n + 1$ ,  $r - \delta > n + 1$ .

$$(8) \quad I_{w,a} \asymp \frac{1}{(1-|w|^2)^{t-\delta-n-1} |1-\langle w,a \rangle|^{\delta+n+1}} \log^k \frac{e}{1-|w|^2} \\ + \frac{1}{|1-\langle w,a \rangle|^t} \log^k \frac{e}{1-|a|^2} \log \frac{e}{|1-\langle a, \varphi_a(w) \rangle|} \quad \text{when } t - \delta > n + 1 = r - \delta.$$

$$(9) \quad I_{w,a} \asymp \log^{k+1} \frac{e}{1-|w|^2} \quad \text{when } t - \delta = n + 1 \quad \text{and } r = 0.$$

**Proposition C** Let  $r + n > 0$  and  $t + n > 0$ . Then the integral

$$I_{w,a} = \int_{\partial B} \frac{d\sigma(\xi)}{|1-\langle \xi, w \rangle|^{n+t} |1-\langle \xi, a \rangle|^{n+r}} \quad (w, a \in B_n)$$

has the following bidirectional estimates:

- (1)  $I_{w,a} \asymp 1$  when  $t + r + n < 0$ .
- (2)  $I_{w,a} \asymp \log \frac{e}{|1-\langle w,a \rangle|}$  when  $t + r + n = 0$ .
- (3)  $I_{w,a} \asymp \frac{1}{|1-\langle w,a \rangle|^{n+r}} \log \frac{e}{|1-\langle w, \varphi_w(a) \rangle|}$  when  $t = 0 > r$ .
- (4)  $I_{w,a} \asymp \frac{1}{|1-\langle w,a \rangle|^{t+r+n}}$  when  $t + r + n > 0 > \max\{r, t\}$ .
- (5)  $I_{w,a} \asymp \frac{1}{|1-\langle w,a \rangle|^n} \log \frac{e}{1-|\varphi_w(a)|^2}$  when  $t = 0 = r$ .
- (6)  $I_{w,a} \asymp \frac{1}{(1-|w|^2)^t |1-\langle w,a \rangle|^{n+r}}$  when  $t > 0 > r$ .
- (7)  $I_{w,a} \asymp \frac{(1-|w|^2)^{-t}}{|1-\langle w,a \rangle|^{n+r}} + \frac{(1-|a|^2)^{-r}}{|1-\langle w,a \rangle|^{n+t}}$  when  $t > 0, r > 0$ .
- (8)  $I_{w,a} \asymp \frac{(1-|w|^2)^{-t}}{|1-\langle w,a \rangle|^n} + \frac{1}{|1-\langle w,a \rangle|^{n+t}} \log \frac{e}{1-|\varphi_a(w)|^2}$  when  $t > 0 = r$ .

**Note 1** When  $t - \delta > n + 1 = r - \delta$ , it follows from Lemma 2.2 in [10] that

$$I_{w,a} \asymp \frac{(1-|w|^2)^{\delta+n+1-t}}{|1-\langle w,a \rangle|^{\delta+n+1}} + \frac{1}{|1-\langle w,a \rangle|^t} \log \frac{e}{1-|\varphi_a(w)|^2} = L_1.$$

However, it follows from Theorem 3.1 (case  $k = 0$ ) in [12] that

$$J_{w,a} \asymp \frac{(1-|w|^2)^{\delta+n+1-t}}{|1-\langle w,a \rangle|^{\delta+n+1}} + \frac{1}{|1-\langle w,a \rangle|^t} \log \frac{e}{|1-\langle \varphi_a(w), a \rangle|} = L_2.$$

Although the expressions of  $L_1$  and  $L_2$  are different,  $L_1$  and  $L_2$  are equivalent.

In fact,  $L_2 \lesssim L_1$  is obvious. Conversely, let  $\sup_{0 < x < 2} x^{t-\delta-n-1} \log \frac{e}{x} = M$ . Then

$$\frac{\frac{1}{|1-\langle w, a \rangle|^t} \log \frac{e|1-\langle w, a \rangle|}{1-|w|^2}}{\frac{(1-|w|^2)^{\delta+n+1-t}}{|1-\langle w, a \rangle|^{\delta+n+1}}} = \left( \frac{1-|w|^2}{|1-\langle w, a \rangle|} \right)^{t-\delta-n-1} \log \frac{e|1-\langle w, a \rangle|}{1-|w|^2} \leq M \Rightarrow$$

$$L_1 \leq L_2 + \frac{1}{|1-\langle w, a \rangle|^t} \log \frac{e|1-\langle w, a \rangle|}{1-|w|^2} \leq L_2 + \frac{M(1-|w|^2)^{\delta+n+1-t}}{|1-\langle w, a \rangle|^{\delta+n+1}} \leq (M+1)L_2.$$

In practical applications, the following integrals are often encountered (for example, Zhou and Chen took advantage of the case  $k = 2$  in [8]):

$$G(w) = \int_{S_n} \frac{1}{|1-\langle \xi, w \rangle|^{n+c}} \left| \log \frac{e}{1-\langle \xi, w \rangle} \right|^k d\sigma(\xi) \quad \text{and}$$

$$F(w) = \int_{B_n} \frac{(1-|z|^2)^\delta}{|1-\langle z, w \rangle|^{n+1+\delta+c}} \left| \log \frac{e}{1-\langle z, w \rangle} \right|^k dv(z) \quad (w \in B_n),$$

where  $\delta > -1$ ,  $c$  and  $k$  are real numbers.

There is a natural problem. Do  $G(w)$  and  $F(w)$  have bidirectional estimates similar to Proposition A? In this paper, we first discuss this problem. Since  $k$  is an abstract real number, the original proof method of Proposition A is no longer suitable for  $F(w)$  and  $G(w)$ . Therefore, we need to deal with the two integrals in a completely different way. Otherwise, for two points in  $B_n$ , we also consider the bidirectional estimates of the following integral in some cases:

$$J_{w,\eta} = \int_{B_n} \frac{(1-|z|^2)^\delta}{|1-\langle z, w \rangle|^t |1-\langle z, \eta \rangle|^r} \log^{-k} \frac{e}{|1-\langle z, \eta \rangle|} \log^k \frac{e}{1-|z|^2} dv(z) \quad (w, \eta \in B_n),$$

where  $\delta > -1$ ,  $r > 0$ ,  $t > 0$ ,  $k > 0$ .

In the study of function space theory, we often encounter the following problems:

(1) Let  $g$  be a given function. Suppose that  $X$  and  $Y$  are two function spaces. Does  $fg$  belong to  $Y$  if  $f$  belongs to  $X$ ?

(2) What conditions do the function  $g$  need if we want to have  $fg \in Y$  for all  $f \in X$ ? In particular, does  $fg$  still belong to  $X$  if  $f$  and  $g$  belong to  $X$ ?

This is the pointwise multiplier problem between function spaces. In general, a function in space  $X$  multiplied by another function may not be in space  $Y$ . For example, we take  $\psi(w_1, w_2) = w_2$  and  $X = Y = \mathcal{B}(U^2)$ , the Bloch space with 2-dimensional unit cylinder as support domain. It is easy to prove that

$$f(w_1, w_2) = \log \frac{1}{1-w_1} + \log \frac{1}{1-w_2} \in \mathcal{B}(U^2).$$

But  $\psi f$  does not belong to  $\mathcal{B}(U^2)$ . This shows that it is very meaningful for us to consider pointwise multipliers between function spaces.

**Definition 1.5** Let  $X$  and  $Y$  be two function spaces on domain  $\Omega$ . If  $\psi f$  belongs to  $Y$  for all  $f \in X$ , then  $\psi$  is called a pointwise multiplier from  $X$  to  $Y$ .

The study of multiplier theory in function spaces has a long history, and there have been a lot of research results, which are closely related to this paper, such as [5]-[6], [20]-[30] etc. As some applications of these integral estimates, we discuss pointwise multipliers from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  in this paper.

## 2. Some Lemmas

In order to prove the main results, we first give several lemmas.

**Lemma 2.1** Let  $c \geq 0$ ,  $\delta > -1$ ,  $k$  be real. Then the integrals

$$I_1(\rho) = \int_0^1 \frac{(1-r)^\delta}{(1-r\rho)^{\delta+1+c}} \log^k \frac{e}{1-\rho r} dr \quad \text{and}$$

$$I_2(\rho) = \int_0^1 \frac{(1-r)^\delta}{(1-r\rho)^{\delta+1+c}} \log^k \frac{e(1-\rho r)}{1-\rho} dr \quad (0 \leq \rho < 1)$$

have the following bidirectional estimates:

- (1)  $I_1(\rho) \asymp I_2(\rho) \asymp 1$  when  $c = 0$  and  $k < -1$ .
- (2)  $I_1(\rho) \asymp \frac{1}{(1-\rho)^c} \log^k \frac{e}{1-\rho}$  when  $c > 0$ .
- (3)  $I_1(\rho) \asymp I_2(\rho) \asymp \log^{k+1} \frac{e}{1-\rho}$  when  $c = 0$  and  $k > -1$ .
- (4)  $I_1(\rho) \asymp I_2(\rho) \asymp \log \log \frac{e^2}{1-\rho}$  when  $c = 0$  and  $k = -1$ .

**Proof** If there exists a constant  $0 < \rho_0 < 1$  such that  $0 \leq \rho \leq \rho_0$ , then it is obvious that these quantities are equivalent. We only need to consider  $\rho$  that is sufficiently close to 1. Therefore, we may let  $\rho > 1/2$ .

By a change of variables  $x = \frac{(1-r)\rho}{1-r\rho}$ , we have that

$$I_1(\rho) = \frac{1}{(1-\rho)^c \rho^{\delta+1}} \int_0^\rho x^\delta (1-x)^{c-1} \log^k \frac{e(1-x)}{1-\rho} dx.$$

It is clear that

$$\int_0^{\frac{1}{2}} x^\delta (1-x)^{c-1} \log^k \frac{e(1-x)}{1-\rho} dx \asymp \log^k \frac{e}{1-\rho}.$$

Otherwise, let  $y = \frac{1-x}{1-\rho}$ . We have that

$$\begin{aligned} & \int_{\frac{1}{2}}^\rho x^\delta (1-x)^{c-1} \log^k \frac{e(1-x)}{1-\rho} dx \asymp \int_{\frac{1}{2}}^\rho (1-x)^{c-1} \log^k \frac{e(1-x)}{1-\rho} dx \\ &= (1-\rho)^c \int_1^{\frac{1}{2(1-\rho)}} y^{c-1} \log^k ey dy. \end{aligned}$$

This means that

$$I_1(\rho) \asymp \frac{1}{(1-\rho)^c} \log^k \frac{e}{1-\rho} + \int_1^{\frac{1}{2(1-\rho)}} y^{c-1} \log^k ey \, dy.$$

By changes of variables  $x = \frac{(1-r)\rho}{1-\rho}$  and  $y = 1+x$ , we have that

$$\begin{aligned} I_2(\rho) &= \frac{1}{(1-\rho)^c \rho^{\delta+1}} \int_0^{\frac{\rho}{1-\rho}} \frac{x^\delta}{(1+x)^{\delta+1+c}} \log^k e(1+x) \, dx \\ &\asymp \frac{1}{(1-\rho)^c} \left\{ 1 + \int_2^{\frac{1}{1-\rho}} \frac{1}{y^{1+c}} \log^k ey \, dy \right\}. \end{aligned}$$

If  $\rho \rightarrow 1^-$ , then there are the following results:

$$\begin{aligned} \int_1^{\frac{1}{2(1-\rho)}} y^{c-1} \log^k ey \, dy &\asymp \frac{1}{(1-\rho)^c} \log^k \frac{e}{1-\rho} \quad \text{when } c > 0, \\ \int_1^{\frac{1}{2(1-\rho)}} y^{-1} \log^k ey \, dy &\asymp \int_2^{\frac{1}{1-\rho}} \frac{1}{y} \log^k ey \, dy \asymp \log^{k+1} \frac{e}{1-\rho} \quad \text{when } k > -1, \\ \int_1^{\frac{1}{2(1-\rho)}} y^{-1} \log^{-1} ey \, dy &\asymp \int_2^{\frac{1}{1-\rho}} \frac{1}{y} \log^{-1} ey \, dy \asymp \log \log \frac{e^2}{1-\rho}, \\ \int_1^{\frac{1}{2(1-\rho)}} y^{-1} \log^k ey \, dy &\asymp \int_2^{\frac{1}{1-\rho}} \frac{1}{y} \log^k ey \, dy \asymp 1 \quad \text{when } k < -1. \end{aligned}$$

This proof is complete.  $\square$

**Lemma 2.2** For  $t > n > r > 0$  and  $k < 0$ , let

$$L_{w,\eta} = \int_{S_n} \frac{1}{|1 - \langle \xi, w \rangle|^t |1 - \langle \xi, \eta \rangle|^r} \log^k \frac{e}{|1 - \langle \xi, \eta \rangle|} \, d\sigma(\xi) \quad (w, \eta \in B_n). \quad \text{Then}$$

$$L_{w,\eta} \gtrsim \frac{1}{(1 - |w|^2)^{t-n} |1 - \langle w, \eta \rangle|^r} \log^k \frac{e}{|1 - \langle w, \eta \rangle|}.$$

**Proof** For  $\varepsilon > 0$  and real number  $y$ , it is easy to obtain that

$$1 \leq \sup_{0 < x < 2} x^\varepsilon \log^y \frac{e}{x} \leq \max \left\{ e^{\varepsilon-|y|} \left( \frac{|y|+1}{\varepsilon} \right)^{|y|}, 2^\varepsilon \log^y \frac{e}{2} \right\}. \quad (2.1)$$

For any  $u \in S_n$  and  $k < 0$ , it follows from (2.1) that

$$\begin{aligned} &|1 - \langle u, \varphi_w(\eta) \rangle|^r \log^{-k} \frac{e|1 - \langle u, w \rangle|}{|1 - \langle w, \eta \rangle| |1 - \langle u, \varphi_w(\eta) \rangle|} \\ &\lesssim |1 - \langle u, \varphi_w(\eta) \rangle|^r \left\{ \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} + \log^{-k} \frac{e}{|1 - \langle u, \varphi_w(\eta) \rangle|} \right\} \\ &\lesssim \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} + 1 \asymp \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|}. \end{aligned}$$

By a change of variables  $\xi = \varphi_w(u)$ , (1.1)-(1.2), (4.7) in [1], the increasing property of integral mean of holomorphic function, we obtain that

$$\begin{aligned} L_{w,\eta} &= \frac{(1-|w|^2)^{n-t}}{|1-\langle w, \eta \rangle|^r} \int_{S_n} \frac{|1-\langle u, w \rangle|^{t+r-2n}}{|1-\langle u, \varphi_w(\eta) \rangle|^r} \log^k \frac{e|1-\langle u, w \rangle|}{|1-\langle w, \eta \rangle||1-\langle u, \varphi_w(\eta) \rangle|} d\sigma(u) \\ &\gtrsim \frac{1}{(1-|w|^2)^{t-n}|1-\langle w, \eta \rangle|^r} \log^k \frac{e}{|1-\langle w, \eta \rangle|} \int_{S_n} |1-\langle u, w \rangle|^{t+r-2n} d\sigma(u) \\ &\geq \frac{1}{(1-|w|^2)^{t-n}|1-\langle w, \eta \rangle|^r} \log^k \frac{e}{|1-\langle w, \eta \rangle|}. \end{aligned}$$

This proof is complete.  $\square$

**Lemma 2.3 ([31])** Let  $\mu$  be a normal function on  $[0, 1)$ . Suppose that  $a$  and  $b$  are the two parameters in the definition of  $\mu$ . Then there are the following results.

(1) For all  $z, w \in B_n$ ,

$$\frac{\mu(|z|)}{\mu(|w|)} \leq \left( \frac{1-|z|^2}{1-|w|^2} \right)^a + \left( \frac{1-|z|^2}{1-|w|^2} \right)^b.$$

(2) If  $z \in B_n$ , then  $\mu(|w|) \asymp \mu(|z|)$  for all  $w \in D(z, 1)$ .

This Lemma comes from Lemma 2.2 in [31].

**Lemma 2.4** Let  $\mu$  be a normal function on  $[0, 1)$ , and let  $\mu_1(r) = (1-r^2)^{\frac{n-s}{p}} \mu(r)$  be also a normal function on  $[0, 1)$ . If  $f \in F(p, \mu, s)$ , then

$$\begin{aligned} |\nabla f(w)| &\lesssim \frac{\|f\|_{p,\mu,s}}{(1-|w|^2)^{\frac{n-s}{p}} \mu(|w|)} \quad \text{and} \\ |f(w)| &\lesssim \left\{ 1 + \int_0^{|w|} \frac{d\rho}{(1-\rho^2)^{\frac{n-s}{p}} \mu(\rho)} \right\} \|f\|_{p,\mu,s} \quad \text{for all } w \in B_n. \end{aligned}$$

In particular,  $F(p, \mu, s) = \mathcal{B}_{\mu_1}(B_n)$  when  $s > n$ .

**Proof** For any  $l \in \{1, 2, \dots, n\}$  and  $w \in B_n$ , by Lemma 2.20 and Lemma 2.24 in [1], Lemma 2.3, we have that

$$\begin{aligned} \left| \frac{\partial f}{\partial w_l}(w) \right|^p &\lesssim \frac{1}{(1-|w|^2)^{n+1}} \int_{D(w,1)} \left| \frac{\partial f}{\partial z_l}(z) \right|^p dv(z) \\ &\lesssim \frac{(1-|w|^2)^{s-n}}{\mu^p(|w|)} \int_{D(w,1)} \frac{(1-|w|^2)^s |\nabla f(z)|^p \mu^p(|z|)}{|1-\langle z, w \rangle|^{2s} (1-|z|^2)} dv(z) \\ &\leq \frac{(1-|w|^2)^{s-n}}{\mu^p(|w|)} \|f\|_{p,\mu,s}^p. \end{aligned}$$

This shows that

$$|\nabla f(w)| \asymp \sum_{l=1}^n \left| \frac{\partial f}{\partial w_l}(w) \right| \asymp \left\{ \sum_{l=1}^n \left| \frac{\partial f}{\partial w_l}(w) \right|^p \right\}^{\frac{1}{p}} \lesssim \frac{\|f\|_{p,\mu,s}}{(1-|w|^2)^{\frac{n-s}{p}} \mu(|w|)}.$$

Therefore,  $F(p, \mu, s) \subseteq \mathcal{B}_{\mu_1}(B_n)$ . Moreover, we have that

$$\begin{aligned} |f(w)| &= \left| f(0, \dots, 0) + \int_0^1 \langle \nabla f(\rho w), \bar{w} \rangle d\rho \right| \\ &\lesssim \left\{ 1 + \int_0^{|w|} \frac{d\rho}{(1 - \rho^2)^{\frac{n-s}{p}} \mu(\rho)} \right\} \|f\|_{p, \mu, s}. \end{aligned}$$

When  $s > n$ , for any  $f \in \mathcal{B}_{\mu_1}(B_n)$  and  $w \in B_n$ , it follows from Proposition A that

$$\begin{aligned} &\int_{B_n} \frac{(1 - |w|^2)^s |\nabla f(z)|^p \mu^p(|z|)}{|1 - \langle z, w \rangle|^{2s} (1 - |z|^2)} dv(z) \\ &\lesssim \|f\|_{\mathcal{B}_{\mu_1}}^p \int_{B_n} \frac{(1 - |w|^2)^s (1 - |z|^2)^{s-n-1}}{|1 - \langle z, w \rangle|^{2s}} dv(z) \\ &\asymp \|f\|_{\mathcal{B}_{\mu_1}}^p \Rightarrow f \in F(p, \mu, s) \Rightarrow \mathcal{B}_{\mu_1}(B_n) \subseteq F(p, \mu, s). \end{aligned}$$

This proof is complete.  $\square$

**Lemma 2.5** ([32, 33]) Let  $\mu$  be a normal function on  $[0, 1)$  and

$$g(u) = 1 + \sum_{j=1}^{\infty} 2^j u^{n_j} \quad (u \in D),$$

where  $n_j$  is the integral part of  $(1 - r_j)^{-1}$ ,  $\mu(r_j) = 2^{-j}$  ( $j = 1, 2, \dots$ ). Then there are the following results:

(1)  $g(r)$  is increasing on  $[0, 1)$ , and there exist  $M_0 > 0$  and  $M_1 < \infty$  such that

$$\inf_{r \in [0, 1)} \mu(r) g(r) \geq M_0, \quad \sup_{u \in D} \mu(|u|) |g(u)| \leq M_1$$

(2) There exists a constant  $M_2 < \infty$  such that

$$\sup_{r \in [0, 1)} (1 - r) \mu(r) g'(r) \leq M_2.$$

The first part of this Lemma comes from Theorem 1 in [32], and the second part comes from Lemma 2.4 in [33].

**Lemma 2.6** ([1]) For  $\alpha > -1$ , if  $f \in A_{\alpha}^1(B_n)$ , then

$$f(z) = \int_B \frac{f(w)}{(1 - \langle z, w \rangle)^{n+1+\alpha}} dv_{\alpha}(w) \quad (z \in B_n).$$

This Lemma comes from Theorem 2.2 in [1].

### 3. Main Results

We first prove three propositions.

**Proposition 3.1** Let  $c$  and  $k$  be real,  $\delta > -1$ . Then the integrals

$$G(w) = \int_{S_n} \frac{1}{|1 - \langle \xi, w \rangle|^{n+c}} \left| \log \frac{e}{1 - \langle \xi, w \rangle} \right|^k d\sigma(\xi) \quad \text{and}$$

$$F(w) = \int_{B_n} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^{n+1+\delta+c}} \left| \log \frac{e}{1 - \langle z, w \rangle} \right|^k dv(z) \quad (w \in B_n)$$

have the following bidirectional estimates:

- (1)  $G(w) \asymp F(w) \asymp 1$  when  $c < 0$ , or  $c = 0$  and  $k < -1$ .
- (2)  $G(w) \asymp F(w) \asymp \frac{1}{(1 - |w|^2)^c} \log^k \frac{e}{1 - |w|^2}$  when  $c > 0$ .
- (3)  $G(w) \asymp F(w) \asymp \log^{k+1} \frac{e}{1 - |w|^2}$  when  $c = 0$  and  $k > -1$ .
- (4)  $G(w) \asymp F(w) \asymp \log \log \frac{e^2}{1 - |w|^2}$  when  $c = 0$  and  $k = -1$ .

**Proof** If there exists a constant  $0 < \rho_0 < 1$  such that  $1 - |w|^2 \geq \rho_0$ , then these bidirectional estimates are obvious. Therefore, we let  $1 - |w|^2$  be sufficiently close to 0. When  $|z| \geq 1$ , it is clear that

$$\log |z| \leq |\log z| = \sqrt{(\log |z|)^2 + (\arg z)^2} \leq \log e^\pi |z|. \quad (3.1)$$

This shows that

$$G(w) \asymp \int_{S_n} \frac{1}{|1 - \langle \xi, w \rangle|^{n+c}} \log^k \frac{e}{|1 - \langle \xi, w \rangle|} d\sigma(\xi).$$

When  $c < 0$ , it follows from (2.1) that we may take  $c < c' < 0$  such that

$$\frac{1}{|1 - \langle \xi, w \rangle|^{n+c}} \log^k \frac{e}{|1 - \langle \xi, w \rangle|} \lesssim \frac{1}{|1 - \langle \xi, w \rangle|^{n+c'}}$$

for all  $\xi \in S_n$  and  $w \in B_n$ . It follows from the increasing property of integral mean of holomorphic function and Proposition A that

$$1 \leq G(w) \lesssim \int_{S_n} \frac{d\sigma(\xi)}{|1 - \langle \xi, w \rangle|^{n+c'}} \asymp 1.$$

By a change of variables  $\xi = \varphi_w(\eta)$ , (4.7) in [1], (1.1)-(1.2), we have that

$$G(w) \asymp \frac{1}{(1 - |w|^2)^c} \int_{S_n} \frac{1}{|1 - \langle \eta, w \rangle|^{n-c}} \log^k \frac{e|1 - \langle \eta, w \rangle|}{1 - |w|^2} d\sigma(\eta) = \frac{J(w)}{(1 - |w|^2)^c}.$$

Next, we consider  $J(w)$  when  $c \geq 0$ .

When  $n = 1$ , it follows from the rotation invariance of the integral that

$$\begin{aligned}
J(w) &= \int_{-\pi}^{\pi} \frac{1}{|1 - |w|e^{i\theta}|^{1-c}} \log^k \frac{e|1 - |w|e^{i\theta}|}{1 - |w|^2} \frac{d\theta}{2\pi} \\
&= \int_0^{\pi} \frac{1}{(1 + |w|^2 - 2|w|\cos\theta)^{\frac{1-c}{2}}} \log^k \frac{e^2(1 + |w|^2 - 2|w|\cos\theta)}{(1 - |w|^2)^2} \frac{d\theta}{2^k\pi} \\
&= \frac{1}{2^k\pi} \int_{-1}^1 \frac{(1 - x^2)^{-\frac{1}{2}}}{(1 + |w|^2 - 2|w|x)^{\frac{1-c}{2}}} \log^k \frac{e^2(1 + |w|^2 - 2|w|x)}{(1 - |w|^2)^2} dx \\
&\asymp \log^k \frac{e}{1 - |w|^2} + \int_0^1 \frac{(1 - x)^{-\frac{1}{2}}}{(1 + |w|^2 - 2|w|x)^{\frac{1-c}{2}}} \log^k \frac{e^2(1 + |w|^2 - 2|w|x)}{(1 - |w|^2)^2} dx.
\end{aligned}$$

Without losing generality, we let  $|w| > 1/2$ . By a change of variables

$$\rho = \frac{(1 + |w|^2)(1 - x)}{1 + |w|^2 - 2|w|x}, \quad \text{we have that}$$

$$\begin{aligned}
J(w) &\asymp \log^k \frac{e}{1 - |w|^2} + (1 - |w|)^c \int_0^1 \frac{\log^k \frac{e}{1 - \frac{2|w|\rho}{1 + |w|^2}}}{\rho^{\frac{1}{2}} \left(1 - \frac{2|w|\rho}{1 + |w|^2}\right)^{1 + \frac{c}{2}}} d\rho \\
&\asymp \log^k \frac{e}{1 - |w|^2} + (1 - |w|)^c \left\{ 1 + \int_{\frac{1}{2}}^1 \frac{\log^k \frac{e}{1 - \frac{2|w|\rho}{1 + |w|^2}}}{\left(1 - \frac{2|w|\rho}{1 + |w|^2}\right)^{1 + \frac{c}{2}}} d\rho \right\} \\
&\asymp \log^k \frac{e}{1 - |w|^2} + (1 - |w|)^c \int_0^1 \frac{\log^k \frac{e}{1 - \frac{2|w|\rho}{1 + |w|^2}}}{\left(1 - \frac{2|w|\rho}{1 + |w|^2}\right)^{1 + \frac{c}{2}}} d\rho.
\end{aligned}$$

It follows from Lemma 2.1 that we can get the estimates of  $J(w)$  by different cases.

When  $n > 1$ , it follows from (1.13) in [1] that

$$J(w) = (n - 1) \int_D \frac{(1 - |z|^2)^{n-2}}{|1 - |w|z|^{n-c}} \log^k \frac{e|1 - |w|z|}{1 - |w|^2} dA(z).$$

If  $c > 0$  and  $k \geq 0$ , then it follows from Proposition A that

$$J(w) \lesssim \log^k \frac{e}{1 - |w|^2} \int_D \frac{(1 - |z|^2)^{n-2}}{|1 - |w|z|^{n-c}} dA(z) \asymp \log^k \frac{e}{1 - |w|^2}.$$

If  $c > 0$  and  $k < 0$ , then we take  $0 < \varepsilon < c$ . It is easy to obtain that

$$\sup_{1 - |w| \leq x \leq 1 + |w|} x^\varepsilon \log^k \frac{ex}{1 - |w|^2} \asymp \max \left\{ (1 - |w|^2)^\varepsilon, \log^k \frac{e}{1 - |w|} \right\} \lesssim \log^k \frac{e}{1 - |w|^2}.$$

Therefore, it follows from Proposition A that

$$J(w) \lesssim \log^k \frac{e}{1 - |w|^2} \int_D \frac{(1 - |z|^2)^{n-2}}{|1 - |w|z|^{n-c+\varepsilon}} dA(z) \asymp \log^k \frac{e}{1 - |w|^2}.$$

On the other hand, we have that

$$J(w) \gtrsim \int_{|z| \leq \frac{1}{2}} \frac{(1 - |z|^2)^{n-2}}{|1 - |w|z|^{n-c}} \log^k \frac{e|1 - |w|z|}{1 - |w|^2} dA(z) \asymp \log^k \frac{e}{1 - |w|^2}.$$

This means that  $J(w) \asymp \log^k \frac{e}{1 - |w|^2}$  when  $c > 0$ . Therefore,

$$G(w) \asymp \frac{1}{(1 - |w|^2)^c} \log^k \frac{e}{1 - |w|^2} \quad \text{when } c > 0.$$

For any  $1/2 < \rho < 1$  and any real number  $k$ , let  $x = \frac{2\rho|w|(1-r)}{(1-\rho|w|)^2}$ . Similar to the previous calculation, we can obtain that

$$\begin{aligned} & \int_{-\pi}^{\pi} \frac{1}{|1 - \rho|w|e^{i\theta}|^n} \log^k \frac{e|1 - \rho|w|e^{i\theta}|}{1 - |w|^2} \frac{d\theta}{2\pi} \asymp \log^k \frac{e}{1 - |w|^2} \\ & + \int_0^1 \frac{1}{(1 + \rho^2|w|^2 - 2\rho|w|r)^{\frac{n}{2}}} \log^k \frac{e^2(1 + \rho^2|w|^2 - 2\rho|w|r)}{(1 - |w|^2)^2} \frac{dr}{\sqrt{1-r}} \\ & \asymp \log^k \frac{e}{1 - |w|^2} + \frac{1}{(1 - \rho|w|)^{n-1}} \int_0^{\frac{2\rho|w|}{(1-\rho|w|)^2}} \frac{\log^k \frac{e^2(1-\rho|w|)^2}{(1-|w|^2)^2} (1+x)}{x^{\frac{1}{2}}(1+x)^{\frac{n}{2}}} dx. \end{aligned}$$

It is clear that

$$\int_0^{\frac{8}{9}} \frac{1}{x^{\frac{1}{2}}(1+x)^{\frac{n}{2}}} \log^k \frac{e^2(1-\rho|w|)^2}{(1-|w|^2)^2} (1+x) dx \asymp \log^k \frac{e(1-\rho|w|)}{1-|w|}.$$

When  $k \geq 0$ , we have that

$$\log^k \frac{e^2(1-\rho|w|)^2}{(1-|w|^2)^2} (1+x) \asymp \log^k \frac{e^2(1-\rho|w|)^2}{(1-|w|^2)^2} + \log^k(1+x). \quad \text{Therefore,}$$

$$\begin{aligned} & \int_{\frac{8}{9}}^{\frac{2\rho|w|}{(1-\rho|w|)^2}} \frac{1}{x^{\frac{1}{2}}(1+x)^{\frac{n}{2}}} \log^k \frac{e^2(1-\rho|w|)^2}{(1-|w|^2)^2} (1+x) dx \\ & \lesssim \int_{\frac{8}{9}}^{\infty} \frac{1}{x^{\frac{n+1}{2}}} \left\{ \log^k \frac{e(1-\rho|w|)}{1-|w|} + \log^k(x+1) \right\} dx \\ & \asymp \log^k \frac{e(1-\rho|w|)}{1-|w|} + 1 \asymp \log^k \frac{e(1-\rho|w|)}{1-|w|}. \end{aligned}$$

When  $k < 0$ , we have that

$$\log^k \frac{e^2(1-\rho|w|)^2}{(1-|w|^2)^2} (1+x) \leq \log^k \frac{e^2(1-\rho|w|)^2}{(1-|w|^2)^2}. \quad \text{Therefore,}$$

$$\begin{aligned} & \int_{\frac{8}{9}}^{\frac{2\rho|w|}{(1-\rho|w|)^2}} \frac{1}{x^{\frac{1}{2}}(1+x)^{\frac{n}{2}}} \log^k \frac{e^2(1-\rho|w|)^2}{(1-|w|^2)^2} (1+x) dx \\ & \lesssim \log^k \frac{e(1-\rho|w|)}{1-|w|} \int_{\frac{8}{9}}^{\infty} \frac{1}{x^{\frac{n+1}{2}}} dx \asymp \log^k \frac{e(1-\rho|w|)}{1-|w|}. \end{aligned}$$

This means that

$$\begin{aligned} & \int_{-\pi}^{\pi} \frac{1}{|1 - \rho|w|e^{i\theta}|^n} \log^k \frac{e|1 - \rho|w|e^{i\theta}|}{1 - |w|} \frac{d\theta}{2\pi} \\ & \asymp \log^k \frac{e}{1 - |w|^2} + \frac{1}{(1 - \rho|w|)^{n-1}} \log^k \frac{e(1 - \rho|w|)}{1 - |w|}. \end{aligned}$$

If  $c = 0$ , then it follows from the polar coordinate and the above result that

$$J(w) \asymp \log^k \frac{e}{1 - |w|^2} \int_0^1 (1 - \rho)^{n-2} d\rho + \int_0^1 \frac{(1 - \rho)^{n-2}}{(1 - \rho|w|)^{n-1}} \log^k \frac{e(1 - \rho|w|)}{1 - |w|} d\rho.$$

It follows from Lemma 2.1 that we can get the estimates of  $J(w)$  for all cases.

Finally, we consider  $F(w)$ .

When  $c < 0$ , let  $c < c' < 0$ . By (3.1), (2.1) and Proposition A, we have that

$$\begin{aligned} 1 & \lesssim F(w) \asymp \int_{B_n} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^{n+1+\delta+c}} \log^k \frac{e}{|1 - \langle z, w \rangle|} dv(z) \\ & \lesssim \int_{B_n} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^{n+1+\delta+c'}} dv(z) \lesssim 1. \end{aligned}$$

When  $c \geq 0$ , by (3.1), Lemma 1.8 in [1] and the estimate of  $G(w)$ , we have that

$$F(w) \asymp \int_0^1 \frac{(1 - \rho)^\delta}{(1 - \rho|w|)^{\delta+1+c}} \log^k \frac{e}{1 - \rho|w|} d\rho,$$

By Lemma 2.1, we can get the estimates of  $F(w)$  in different cases.

The proof is complete.  $\square$

**Proposition 3.2** For  $\delta > -1$ ,  $r > 0$ ,  $t > 0$ ,  $k > 0$ , the integral

$$J_{w,\eta} = \int_{B_n} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^t |1 - \langle z, \eta \rangle|^r} \left| \log \frac{e}{1 - \langle z, \eta \rangle} \right|^{-k} \log^k \frac{e}{1 - |z|^2} dv(z)$$

( $w, \eta \in B_n$ ) has the following bidirectional estimates:

(1) When  $r + t - \delta > n + 1 > \max\{t - \delta, r - \delta\}$ ,

$$J_{w,\eta} \asymp \frac{1}{|1 - \langle w, \eta \rangle|^{r+t-\delta-n-1}}.$$

(2) When  $t - \delta = n + 1 > r - \delta$ ,

$$J_{w,\eta} \asymp \frac{1}{|1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \log^k \frac{e}{1 - |w|^2} \log \frac{e}{|1 - \langle w, \varphi_w(\eta) \rangle|}.$$

(3) When  $t - \delta > n + 1 > r - \delta$ ,

$$J_{w,\eta} \asymp \frac{1}{(1 - |w|^2)^{t-\delta-n-1} |1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \log^k \frac{e}{1 - |w|^2}.$$

**Proof** Without losing generality, let  $1 - |w|^2$  and  $|1 - \langle w, \eta \rangle|$  be sufficiently close to 0. It follows from (3.1) that

$$J_{w,\eta} \asymp \int_{B_n} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^t |1 - \langle z, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle z, \eta \rangle|} \log^k \frac{e}{1 - |z|^2} dv(z).$$

By a partition of  $B_n$ , there exist  $\Omega_1, \Omega_2, \Omega_3, \Omega_4$  such that  $B = \Omega_1 \cup \Omega_2 \cup \Omega_3 \cup \Omega_4$ , where  $\Omega_j$  and  $\Omega_k$  ( $j \neq k$ ) are mutually disjoint. By Lemma 3.3 in [34], we have

$$\begin{aligned} |1 - \langle z, w \rangle| &\lesssim |1 - \langle w, \eta \rangle| \lesssim |1 - \langle z, \eta \rangle| \quad (z \in \Omega_1); \\ |1 - \langle z, \eta \rangle| &\lesssim |1 - \langle w, \eta \rangle| \lesssim |1 - \langle z, w \rangle| \quad (z \in \Omega_2); \\ |1 - \langle w, \eta \rangle| &\lesssim |1 - \langle z, w \rangle| \lesssim |1 - \langle z, \eta \rangle| \quad (z \in \Omega_3); \\ |1 - \langle w, \eta \rangle| &\lesssim |1 - \langle z, \eta \rangle| \lesssim |1 - \langle z, w \rangle| \quad (z \in \Omega_4). \end{aligned}$$

When  $z \in \Omega_1 \cup \Omega_3$ , we have that  $|1 - \langle z, \eta \rangle| \gtrsim |1 - \langle w, \eta \rangle|$ . If  $t - \delta > n + 1$ , then it follows from Proposition B(6) that

$$\begin{aligned} J_1 &= \int_{\Omega_1 \cup \Omega_3} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^t |1 - \langle z, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle z, \eta \rangle|} \log^k \frac{e}{1 - |z|^2} dv(z) \\ &\lesssim \frac{1}{|1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \int_{B_n} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^t} \log^k \frac{e}{1 - |z|^2} dv(z) \\ &\asymp \frac{1}{(1 - |w|^2)^{t-\delta-n-1} |1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \log^k \frac{e}{1 - |w|^2}. \end{aligned} \quad (3.2)$$

When  $t - \delta = n + 1$ , we may take  $0 < \varepsilon < r$ . This means that  $(r - \varepsilon) - \delta < n + 1$  and  $r - \varepsilon > 0$ . By a change of variables  $z = \varphi_w(u)$ , (1.1)-(1.2), Proposition B(2), we have that

$$\begin{aligned} J_1 &\lesssim \frac{1}{|1 - \langle w, \eta \rangle|^\varepsilon} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \int_{B_n} \frac{(1 - |z|^2)^\delta \log^k \frac{e}{1 - |z|^2} dv(z)}{|1 - \langle z, w \rangle|^{\delta+n+1} |1 - \langle z, \eta \rangle|^{r-\varepsilon}} \\ &= \frac{1}{|1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \int_{B_n} \frac{(1 - |u|^2)^\delta \log^k \frac{e}{1 - |\varphi_w(u)|^2} dv(u)}{|1 - \langle u, w \rangle|^{\delta+n+1-(r-\varepsilon)} |1 - \langle u, \varphi_w(\eta) \rangle|^{r-\varepsilon}} \\ &\lesssim \frac{1}{|1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \int_{B_n} \frac{(1 - |u|^2)^\delta \left\{ \log^k \frac{e}{1 - |w|^2} + \log^k \frac{e}{1 - |u|^2} \right\} dv(u)}{|1 - \langle u, w \rangle|^{\delta+n+1-(r-\varepsilon)} |1 - \langle u, \varphi_w(\eta) \rangle|^{r-\varepsilon}} \\ &\asymp \frac{\log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|}}{|1 - \langle w, \eta \rangle|^r} \left\{ \log^k \frac{e}{1 - |w|^2} \log \frac{e}{|1 - \langle w, \varphi_w(\eta) \rangle|} + \log^{k+1} \frac{e}{|1 - \langle w, \varphi_w(\eta) \rangle|} \right\} \\ &\asymp \frac{1}{|1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \log^k \frac{e}{1 - |w|^2} \log \frac{e}{|1 - \langle w, \varphi_w(\eta) \rangle|}. \end{aligned} \quad (3.3)$$

Let  $m = |1 - \langle w, \eta \rangle|$ . When  $z \in \Omega_2 \cup \Omega_4$ , it is clear that

$$|1 - \langle z, w \rangle| \gtrsim (m + |1 - \langle z, \eta \rangle|) \geq |m + 1 - \langle z, \eta \rangle|.$$

When  $n > 1$ , it is similar to the proof of (3.17) in [12] (we omit some processes). If  $r - \delta < n + 1 < t + r - \delta$ , then it follows from Lemma 1.8 and (1.13) in [1] that

$$\begin{aligned} J_2 &= \int_{\Omega_2 \cup \Omega_4} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^t |1 - \langle z, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle z, \eta \rangle|} \log^k \frac{e}{1 - |z|^2} dv(z) \\ &\lesssim \int_0^1 (1 - \rho^2)^\delta \log^k \frac{e}{1 - \rho^2} \left\{ \int_D \frac{\rho^{2n-1} (1 - |\xi|^2)^{n-2}}{|m + 1 - \rho \xi|^t |1 - \rho \xi|^r} \log^{-k} \frac{e}{|1 - \rho \xi|} dA(\xi) \right\} d\rho \\ &\lesssim \int_0^1 (1 - \rho)^\delta \log^k \frac{e}{1 - \rho} \left\{ \int_{|u-1| < \rho} \frac{(\rho - |u - 1|)^{n-2}}{|m + u|^t |u|^r} \log^{-k} \frac{e}{|u|} dA(u) \right\} d\rho \end{aligned}$$

$$\begin{aligned}
&\lesssim \int_0^1 (1-\rho)^\delta \log^k \frac{e}{1-\rho} \left\{ \int_{1-\rho}^1 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\{R-(1-\rho)\}^{n-2}}{(m+R)^t R^{r-1}} \log^{-k} \frac{e}{R} dR d\theta \right\} d\rho \\
&+ \int_0^1 (1-\rho)^\delta \log^k \frac{e}{1-\rho} \left\{ \int_1^{1+\rho} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\rho^{n-2}}{(m+R)^t R^{r-1}} \log^{-k} \frac{e}{R} dR d\theta \right\} d\rho \\
&\lesssim \int_0^1 (1-\rho)^\delta \log^k \frac{e}{1-\rho} \left\{ \int_{1-\rho}^1 \frac{R^{n-1-r}}{(m+R)^t} \log^{-k} \frac{e}{R} dR \right\} d\rho + 1 \\
&\lesssim \int_0^1 \frac{(1-\rho)^\delta}{m^{r+t-n}} \log^k \frac{e}{1-\rho} \left\{ \int_{\frac{1-\rho}{m}}^{\frac{1}{m}} \frac{x^{n-1-r}}{(1+x)^t} \log^{-k} \frac{e}{mx} dx \right\} d\rho + 1 \\
&= \frac{1}{m^{r+t-n}} \int_0^{\frac{1}{m}} \frac{x^{n-1-r}}{(1+x)^t} \log^{-k} \frac{e}{mx} \left\{ \int_{1-mx}^1 (1-\rho)^\delta \log^k \frac{e}{1-\rho} d\rho \right\} dx + 1 \\
&\asymp \frac{1}{m^{r+t-\delta-n-1}} \int_0^{\frac{1}{m}} \frac{x^{n+\delta-r}}{(1+x)^t} dx + 1 \\
&\leq \frac{1}{m^{r+t-\delta-n-1}} \int_0^1 y^{n+\delta-r} (1-y)^{r+t-\delta-n-2} dy \asymp \frac{1}{|1-\langle w, \eta \rangle|^{r+t-\delta-n-1}}. \quad (3.4)
\end{aligned}$$

When  $n > 1$ , it is similar to the proof of (3.4). If  $t - \delta < n + 1 < r + t - \delta$ , then we can also get that

$$\begin{aligned}
J_1 &\lesssim \int_{\Omega_1 \cup \Omega_3} \frac{(1-|z|^2)^\delta \log^k \frac{e}{1-|z|^2}}{|1-\langle z, w \rangle|^t |m+1-\langle z, w \rangle|^r} \log^{-k} \frac{e}{|m+1-\langle z, w \rangle|} dv(z) \\
&\lesssim \frac{1}{m^{r+t-\delta-n-1}} \int_0^{\frac{1}{m}} \frac{x^{n+\delta-t}}{(1+x)^r \log^k \frac{e}{m(1+x)}} \log^k \frac{e}{mx} dx + 1.
\end{aligned}$$

It is clear that

$$\begin{aligned}
\int_0^{\frac{1}{2}} \frac{x^{n+\delta-t}}{(1+x)^r \log^k \frac{e}{m(1+x)}} \log^k \frac{e}{mx} dx &\lesssim \int_0^{\frac{1}{2}} \frac{x^{n+\delta-t}}{\log^k \frac{e}{m}} \log^k \frac{e}{mx} dx \\
&\lesssim \int_0^{\frac{1}{2}} x^{n+\delta-t} dx + \log^{-k} \frac{e}{m} \int_0^{\frac{1}{2}} x^{n+\delta-t} \log^k \frac{1}{x} dx \asymp 1.
\end{aligned}$$

Otherwise, let  $y = x/(x+1)$ . We also have that

$$\begin{aligned}
&\int_{\frac{1}{2}}^{\frac{1}{m}} \frac{x^{n+\delta-t}}{(1+x)^r \log^k \frac{e}{m(1+x)}} \log^k \frac{e}{mx} dx \\
&= \int_{\frac{1}{3}}^{\frac{1}{m+1}} y^{n+\delta-t} (1-y)^{r+t-\delta-n-2} \log^k \frac{e(1-y)}{my} \log^{-k} \frac{e(1-y)}{m} dy \\
&\lesssim \int_{\frac{1}{3}}^1 y^{n+\delta-t} (1-y)^{r+t-\delta-n-2} dy \asymp 1.
\end{aligned}$$

This shows that

$$J_1 \lesssim \frac{1}{|1-\langle w, \eta \rangle|^{r+t-\delta-n-1}} \text{ when } t - \delta < n + 1 < r + t - \delta \text{ and } n > 1. \quad (3.5)$$

The proof of case  $n = 1$  is similar to the proof of Lemma 1 in [35].

It follows from (3.4) and (3.5) that the “ $\lesssim$ ” part of (1) is true. It follows from (3.3) and (3.4) that the “ $\lesssim$ ” part of (2) is true. It follows from (3.2) and (3.4) that the “ $\lesssim$ ” part of (3) is true. Next, we consider “ $\gtrsim$ ” parts.

When  $t - \delta < n + 1$  and  $r - \delta < n + 1$ , it follows from Proposition B(4) that

$$J_{w,\eta} \gtrsim \int_{B_n} \frac{(1 - |z|^2)^\delta}{|1 - \langle z, w \rangle|^t |1 - \langle z, \eta \rangle|^r} dv(z) \asymp \frac{1}{|1 - \langle w, \eta \rangle|^{r+t-\delta-n-1}}.$$

This shows that the “ $\gtrsim$ ” part of (1) is true.

When  $t - \delta > n + 1$ , by a change of variables  $z = \varphi_w(u)$  and (1.1), we have that

$$\begin{aligned} J_{w,\eta} &= \frac{(1 - |w|^2)^{\delta+n+1-t}}{|1 - \langle w, \eta \rangle|^r} \int_{B_n} \frac{|1 - \langle u, \varphi_w(\eta) \rangle|^{-r} (1 - |u|^2)^\delta \log^k \frac{e|1 - \langle u, w \rangle|^2}{(1 - |u|^2)(1 - |w|^2)} dv(u)}{|1 - \langle u, w \rangle|^{2\delta+2n+2-r-t} \log^k \frac{e|1 - \langle u, \eta \rangle|}{|1 - \langle w, \eta \rangle| |1 - \langle u, \varphi_w(\eta) \rangle|}} \\ &\geq \frac{(1 - |w|^2)^{\delta+n+1-t}}{|1 - \langle w, \eta \rangle|^r} \int_{|u| \leq \frac{1}{9}} \frac{|1 - \langle u, \varphi_w(\eta) \rangle|^{-r} (1 - |u|^2)^\delta \log^k \frac{e|1 - \langle u, w \rangle|^2}{(1 - |u|^2)(1 - |w|^2)} dv(u)}{|1 - \langle u, w \rangle|^{2\delta+2n+2-r-t} \log^k \frac{e|1 - \langle u, \eta \rangle|}{|1 - \langle w, \eta \rangle| |1 - \langle u, \varphi_w(\eta) \rangle|}} \\ &\asymp \frac{1}{(1 - |w|^2)^{t-\delta-n-1} |1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \log^k \frac{e}{1 - |w|^2}. \end{aligned}$$

This shows that the “ $\gtrsim$ ” part of (3) is true.

When  $t - \delta = n + 1 > r - \delta$ , by Lemma 1.8 in [1], Lemma 2.2 and  $(1 - \rho|w|^2)/2 < 1 - \rho|w| \leq 1 - \rho|w|^2$  for any  $0 \leq \rho < 1$  and  $w \in B_n$ , we have that

$$\begin{aligned} J_{w,\eta} &\gtrsim \int_0^1 \frac{(1 - \rho^2)^\delta}{(1 - \rho^2|w|^2)^{\delta+1} |1 - \rho\langle w, \eta \rangle|^r} \log^k \frac{e}{1 - \rho^2} \log^{-k} \frac{e}{|1 - \rho\langle w, \eta \rangle|} d\rho \\ &\gtrsim \int_0^1 \frac{(1 - \rho)^\delta}{(1 - \rho|w|^2)^{\delta+1} |1 - \rho\langle w, \eta \rangle|^r} \log^k \frac{e}{1 - \rho} \log^{-k} \frac{e}{|1 - \rho\langle w, \eta \rangle|} d\rho. \end{aligned} \quad (3.6)$$

When  $|w| > 1/\sqrt{2}$ , it follows from a change of variables  $x = \frac{|w|^2(1 - \rho)}{1 - \rho|w|^2}$  that

$$\begin{aligned} &\int_0^1 \frac{(1 - \rho)^\delta}{(1 - \rho|w|^2)^{\delta+1} |1 - \rho\langle w, \eta \rangle|^r} \log^k \frac{e}{1 - \rho} \log^{-k} \frac{e}{|1 - \rho\langle w, \eta \rangle|} d\rho \\ &= \frac{1}{|w|^{2(\delta+1)}} \int_0^{|w|^2} \frac{x^\delta}{1 - x} \left| 1 - \langle w, \eta \rangle + \frac{\langle w, \eta \rangle(1 - |w|^2)x}{|w|^2(1 - x)} \right|^{-r} \log^k \frac{e|w|^2(1 - x)}{(1 - |w|^2)x} \\ &\quad \times \log^{-k} \frac{e}{\left| 1 - \langle w, \eta \rangle + \frac{\langle w, \eta \rangle(1 - |w|^2)x}{|w|^2(1 - x)} \right|} dx \\ &\gtrsim \int_{\frac{1}{2}}^{|w|^2} \frac{1}{1 - x} \left( m + \frac{1 - |w|^2}{1 - x} \right)^{-r} \log^k \frac{e(1 - x)}{1 - |w|^2} \log^{-k} \frac{e}{m + \frac{1 - |w|^2}{1 - x}} dx \\ &\gtrsim \log^{-k} \frac{e}{m} \int_{\frac{1}{2}}^{|w|^2} \frac{1}{1 - x} \left( m + \frac{1 - |w|^2}{1 - x} \right)^{-r} \log^k \frac{e(1 - x)}{1 - |w|^2} dx. \end{aligned}$$

It follows from (3.12) and (3.13) in [12] that

$$\int_0^1 \frac{(1 - \rho)^\delta}{(1 - \rho|w|^2)^{\delta+1} |1 - \rho\langle w, \eta \rangle|^r} \log^k \frac{e}{1 - \rho} \log^{-k} \frac{e}{|1 - \rho\langle w, \eta \rangle|} d\rho$$

$$\gtrsim \frac{1}{m^r} \log^{-k} \frac{e}{m} \log \frac{em}{1-|w|^2} \log^k \frac{e}{1-|w|^2}. \quad (3.7)$$

When  $t - \delta = n + 1 > r - \delta$ , it follows from (3.6) and (3.7) that

$$J_{w,\eta} \gtrsim \frac{1}{|1 - \langle w, \eta \rangle|^r} \log^{-k} \frac{e}{|1 - \langle w, \eta \rangle|} \log^k \frac{e}{1 - |w|^2} \log \frac{e}{|1 - \langle w, \varphi_w(\eta) \rangle|}.$$

This shows that the “ $\gtrsim$ ” part of (2) is true. This proof is complete.  $\square$

**Proposition 3.3** Let  $\mu_1, \mu_2$  be two normal functions on  $[0, 1)$ ,  $h \in H(B_n)$ . If

$$\sup_{z \in B_n} \frac{\mu_1(|z|)|h(z)|}{\mu_2(|z|)} = M < \infty, \quad \text{then}$$

$$\sup_{z \in B_n} \frac{(1 - |z|^2)\mu_1(|z|)|\nabla h(z)|}{\mu_2(|z|)} \lesssim M.$$

**Proof** Let  $g$  be this function in Lemma 2.5, matching  $\mu_2$ . For any  $w \in B_n$ , let  $F_w(z) = h(z)g(\langle z, w \rangle)$ . It follows from Lemma 2.5 that

$$\mu_1(|z|)|F_w(z)| = \frac{\mu_1(|z|)|h(z)|}{\mu_2(|z|)} \mu_2(|z|)|g(\langle z, w \rangle)| \leq M \mu_2(|\langle z, w \rangle|)|g(\langle z, w \rangle)| \leq M_1 M$$

for all  $z \in B_n$ . Moreover, we have that  $|F_w(z)| \leq M_1 M / \mu_1(0)(1 - |z|^2)^{b_1}$ , where  $b_1$  the larger parameter in the definition of  $\mu_1$ . When  $\alpha > b_1 - 1$ , it is clear that  $F_w \in A_\alpha^1(B_n)$ . It follows from Lemma 2.6 that

$$F_w(z) = \int_{B_n} \frac{F_w(u) dv_\alpha(u)}{(1 - \langle z, u \rangle)^{n+1+\alpha}} \quad (z \in B_n).$$

Let  $a_1$  be the smaller parameter in the definition of  $\mu_1$ . By Lemma 2.3 and Proposition A, we have that

$$\begin{aligned} \mu_1(|z|)|(\nabla F_w)(z)| &\leq \int_{B_n} \frac{(n+1+\alpha)M_1 M c_\alpha \mu_1(|z|)(1 - |u|^2)^\alpha dv(u)}{\mu_1(|u|)|1 - \langle z, u \rangle|^{n+2+\alpha}} \\ &\lesssim \int_{B_n} \frac{M(1 - |z|^2)^{a_1}(1 - |u|^2)^{\alpha-a_1} dv(u)}{|1 - \langle z, u \rangle|^{n+2+\alpha}} + \int_{B_n} \frac{M(1 - |z|^2)^{b_1}(1 - |u|^2)^{\alpha-b_1} dv(u)}{|1 - \langle z, u \rangle|^{n+2+\alpha}} \\ &\asymp \frac{M}{1 - |z|^2}. \end{aligned}$$

In particular, if we take  $z = w$ , then it is clear that

$$(1 - |w|^2)\mu_1(|w|)|g(|w|^2)\nabla h(w) + h(w)g'(|w|^2)\overline{w}| \lesssim M.$$

Suppose that  $b_2$  is the larger parameter in the definition of  $\mu_2$ . By Lemma 2.5 and the definition of normal function, it is clear that

$$\begin{aligned} \frac{(1 - |w|^2)\mu_1(|w|)|\nabla h(w)|}{\mu_2(|w|)} &\leq \frac{2^{b_2}}{M_0}(1 - |w|^2)\mu_1(|w|)|\nabla h(w)|g(|w|^2) \\ &\lesssim M + \frac{\mu_1(|w|)|h(w)|}{\mu_2(|w|)}(1 - |w|^2)\mu_2(|w|)g'(|w|^2) \leq M(1 + M_2). \end{aligned}$$

This proof is complete.  $\square$

**Theorem 3.4** Let  $\mu$  and  $\nu$  be two normal functions on  $[0, 1)$ . Suppose that  $\psi \in H(B_n)$  and  $0 \leq s \leq n$ . Then there are the following results:

(1) If  $\int_0^1 \mu^{-1}(\rho)(1-\rho^2)^{\frac{s-n}{p}} d\rho < \infty$ , then  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  if and only if  $\psi \in \mathcal{B}_\nu(B_n)$  and

$$\sup_{z \in B_n} \frac{\nu(|z|)|\psi(z)|}{\mu(|z|)(1-|z|^2)^{\frac{n-s}{p}}} < \infty. \quad (3.8)$$

(2) If  $\int_0^1 \mu^{-1}(\rho)(1-\rho^2)^{\frac{s-n}{p}} d\rho = \infty$  and  $1 + \int_0^r \mu^{-1}(\rho)(1-\rho^2)^{\frac{s-n}{p}} d\rho \asymp \mu^{-1}(r)(1-r^2)^{1+\frac{s-n}{p}}$ , then  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  if and only if (3.8) holds.

**Proof** Suppose that  $\psi \in \mathcal{B}_\nu(B_n)$  and (3.8) holds. For any  $f \in F(p, \mu, s)$  and  $z \in B_n$  with  $|z| > 1/2$ , it follows from Lemma 2.4 and Proposition 3.3 that

$$\begin{aligned} \nu(|z|)|\nabla(\psi f)(z)| &\leq \nu(|z|)|\psi(z)||\nabla f(z)| + \nu(|z|)|\nabla\psi(z)||f(z)| \\ &\lesssim \frac{\nu(|z|)|\psi(z)|}{(1-|z|^2)^{\frac{n-s}{p}}\mu(|z|)} \|f\|_{p,\mu,s} \\ &\quad + \nu(|z|)|\nabla\psi(z)| \left\{ 1 + \int_0^{|z|} \frac{d\rho}{\mu(\rho)(1-\rho^2)^{\frac{n-s}{p}}} \right\} \|f\|_{p,\mu,s} \\ &\lesssim \|f\|_{p,\mu,s} \Rightarrow \|\psi f\|_{p,\mu,s} < \infty. \end{aligned}$$

This means that  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$ .

Conversely, let  $\psi$  be a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$ . It follows from the closed graph theorem that  $M_\psi : f \rightarrow \psi f$  is a bounded operator from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$ . First, it is clear that  $\psi \in \mathcal{B}_\nu(B_n)$  by taking  $f(z) = 1$ . Suppose that  $b$  is the larger parameter in the definition of  $\mu$ . For any  $w \in B_n$  with  $|w| > 1/2$ , we take that

$$f_w(z) = \frac{(1-|w|^2)^{1+b+\frac{s}{p}}}{\mu(|w|)} \left\{ \frac{1-|w|^2}{(1-\langle z, w \rangle)^{b+1+\frac{n}{p}}} - \frac{1}{(1-\langle z, w \rangle)^{b+\frac{n}{p}}} \right\} \quad (z \in B_n).$$

Then  $f_w(w) = 0$  and it follows from Lemma 2.3 that

$$\begin{aligned} &\int_{B_n} \frac{(1-|u|^2)^s |\nabla f_w(z)|^p \mu^p(|z|)}{|1-\langle z, u \rangle|^{2s}(1-|z|^2)} dv(z) \\ &\lesssim \int_{B_n} \frac{(1-|u|^2)^s (1-|w|^2)^{p+pb+s} \mu^p(|z|)}{\mu^p(|w|)|1-\langle z, u \rangle|^{2s}|1-\langle z, w \rangle|^{p+pb+n}(1-|z|^2)} dv(z) \\ &\lesssim \int_{B_n} \frac{(1-|u|^2)^s (1-|w|^2)^{p+s}(1-|z|^2)^{pb-1}}{|1-\langle z, u \rangle|^{2s}|1-\langle z, w \rangle|^{p+pb+n}} dv(z) \\ &\quad + \int_{B_n} \frac{(1-|u|^2)^s (1-|w|^2)^{p+s+pb-pa}(1-|z|^2)^{pa-1}}{|1-\langle z, u \rangle|^{2s}|1-\langle z, w \rangle|^{p+pb+n}} dv(z) = J_1 + J_2. \end{aligned}$$

When  $0 \leq 2s < pb + n$ , it follows from Proposition B(6) or Proposition A that

$$J_1 \asymp \frac{(1-|u|^2)^s (1-|w|^2)^{p+s}}{(1-|w|^2)^p |1-\langle u, w \rangle|^{2s}} \lesssim 1.$$

When  $2s > pb + n$ , it follows from Proposition B(7) and  $0 \leq s \leq n$  that

$$J_1 \asymp \frac{(1 - |u|^2)^s(1 - |w|^2)^{p+s}}{(1 - |w|^2)^p|1 - \langle u, w \rangle|^{2s}} + \frac{(1 - |u|^2)^s(1 - |w|^2)^{p+s}}{(1 - |u|^2)^{2s-pb-n}|1 - \langle u, w \rangle|^{p+pb+n}} \lesssim 1.$$

When  $2s = pb + n$ , we choose  $0 < \varepsilon < s$ . By Proposition B(8) and (2.1), we have

$$\begin{aligned} J_1 &\asymp \frac{(1 - |u|^2)^s(1 - |w|^2)^{p+s}}{(1 - |w|^2)^p|1 - \langle u, w \rangle|^{2s}} + \frac{(1 - |u|^2)^s(1 - |w|^2)^{p+s}}{|1 - \langle u, w \rangle|^{p+pb+n}} \log \frac{e}{|1 - \langle \varphi_u(w), u \rangle|} \\ &\lesssim \frac{(1 - |u|^2)^s(1 - |w|^2)^{p+s}}{(1 - |w|^2)^p|1 - \langle u, w \rangle|^{2s}} + \frac{(1 - |u|^2)^{s-\varepsilon}(1 - |w|^2)^{p+s}}{|1 - \langle u, w \rangle|^{p+pb+n-\varepsilon}} \lesssim 1. \end{aligned}$$

Similarly, we may prove that  $J_2 \lesssim 1$ . This shows that  $\|f_w\|_{p,\mu,s} \lesssim 1$ .

If  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$ , then we have that

$$\|M_\psi\| \gtrsim \|M_\psi\| \cdot \|f_w\|_{p,\mu,s} \geq \nu(|w|)|\nabla(\psi f_w)(w)| \asymp \frac{\nu(|w|)|\psi(w)|}{(1 - |w|^2)^{\frac{n-s}{p}} \mu(|w|)}.$$

This shows that (3.8) holds. This proof is complete.  $\square$

**Corollary 3.5** Let  $\mu$  and  $\nu$  be two normal functions on  $[0, 1)$ . Suppose that  $\psi \in H(B_n)$ ,  $s > n$  and  $\gamma(\rho) = (1 - \rho^2)^{\frac{n-s}{p}} \mu(\rho)$  is a normal function on  $[0, 1)$ . Then  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  if and only if (3.8) holds and

$$\sup_{z \in B_n} \nu(|z|)|\nabla\psi(z)| \int_0^{|z|} \frac{d\rho}{\mu(\rho)(1 - \rho^2)^{\frac{n-s}{p}}} < \infty.$$

**Proof** It follows from Lemma 2.4 that  $F(p, \mu, s) = \mathcal{B}_\gamma(B_n)$  when  $s > n$ . By Theorem A in [6], we may obtain this result.

In Theorem 3.4, if  $\int_0^1 \mu^{-1}(\rho)(1 - \rho^2)^{\frac{s-n}{p}} d\rho = \infty$  and  $1 + \int_0^r \mu^{-1}(\rho)(1 - \rho^2)^{\frac{s-n}{p}} d\rho$  is not equivalent to  $\mu^{-1}(r)(1 - r^2)^{1+\frac{s-n}{p}}$  for  $0 \leq r < 1$ , then we do not give the necessary and sufficient conditions. However, there are the following results for the normal weight  $\mu(r) = (1 - r^2)^\alpha \log^\beta \frac{e}{1-r^2}$  ( $\alpha > 0$ ,  $-\infty < \beta < \infty$ ).

**Corollary 3.6** Let  $\nu$  be a normal function on  $[0, 1)$ . Suppose that  $\mu(r) = (1 - r^2)^\alpha \log^\beta \frac{e}{1-r^2}$  ( $\alpha > 0$ ,  $-\infty < \beta < \infty$ ),  $\psi \in H(B_n)$  and  $0 \leq s \leq n$ . Then there are the following results:

(1) If  $p\alpha + n - p < s$ , or  $p\alpha + n - p = s > 0$  and  $\beta > 1$ , then  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  if and only if  $\psi \in \mathcal{B}_\nu(B_n)$  and (3.8) holds.

(2) If  $p\alpha + n - p > s$ , then  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  if and only if (3.8) holds.

(3) If  $p\alpha + n - p = s > 0$  and  $\beta = 1$ , then  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  if and only if (3.8) holds and

$$\sup_{z \in B_n} \nu(|z|)|\nabla\psi(z)| \log \log \frac{e}{1 - |z|^2} < \infty. \quad (3.9)$$

(4) If  $p\alpha + n - p = s > 0$  and  $\beta < 1$ , then  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  if and only if (3.8) holds and

$$\sup_{z \in B_n} \nu(|z|) |\nabla \psi(z)| \log^{1-\beta} \frac{e}{1-|z|^2} < \infty. \quad (3.10)$$

**Proof** The necessary and sufficient conditions of (1)-(2), and the sufficient conditions of (3)-(4) come from Theorem 3.4.

For any  $w \in B_n$  with  $|w| > 1/2$ , we take the test function

$$G_w(z) = \left\{ \int_0^{\langle z, w \rangle} \frac{1}{1-t} \log^{-\beta} \frac{e}{1-t} dt \right\}^2 / \int_0^{|w|^2} \frac{1}{1-t} \log^{-\beta} \frac{e}{1-t} dt \\ - 2 \int_0^{\langle z, w \rangle} \frac{1}{1-t} \log^{-\beta} \frac{e}{1-t} dt \quad (z \in B_n).$$

We know that  $p+s-n-1 = p\alpha-1 > -1$ ,  $2s+p-(p+s-n-1) = s+n+1 > n+1$  and  $p-(p+s-n-1) = -s+n+1 < n+1$ . When  $\beta > 0$  and  $0 \leq s < p$ , for any  $u \in B_n$ , it follows from Proposition 3.2(1) that

$$\int_{B_n} \frac{(1-|u|^2)^s |\nabla G_w(z)|^p \mu^p(|z|) dv(z)}{(1-|z|^2)|1-\langle z, u \rangle|^{2s}} \\ \lesssim \int_{B_n} \frac{(1-|u|^2)^s (1-|z|^2)^{p+s-n-1} \log^{p\beta} \frac{e}{1-|z|^2} dv(z)}{|1-\langle z, u \rangle|^{2s} |1-\langle z, w \rangle|^p \log^{p\beta} \frac{e}{|1-\langle z, w \rangle|}} \\ \asymp \frac{(1-|u|^2)^s}{|1-\langle u, w \rangle|^{2s+p-(p+s-n-1)-n-1}} = \frac{(1-|u|^2)^s}{|1-\langle u, w \rangle|^s} \lesssim 1.$$

When  $\beta > 0$  and  $s = p$ , we choose  $0 < \varepsilon < p$ . For any  $u \in B_n$ , it follows from Proposition 3.2(2) and (2.1) that

$$\int_{B_n} \frac{(1-|u|^2)^s |\nabla G_w(z)|^p \mu^p(|z|) dv(z)}{(1-|z|^2)|1-\langle z, u \rangle|^{2s}} \\ \lesssim \frac{(1-|u|^2)^s}{|1-\langle u, w \rangle|^s} \log^{-p\beta} \frac{e}{|1-\langle u, w \rangle|} \log^{p\beta} \frac{e}{1-|u|^2} \log \frac{e}{|1-\langle u, \varphi_u(w) \rangle|} \\ = \frac{(1-|u|^2)^{s-\varepsilon}}{|1-\langle u, w \rangle|^{s-\varepsilon}} |1-\langle u, \varphi_u(w) \rangle|^{\frac{\varepsilon}{2}} \log \frac{e}{|1-\langle u, \varphi_u(w) \rangle|} \\ \times (1-|u|^2)^{\frac{\varepsilon}{2}} \log^{p\beta} \frac{e}{1-|u|^2} |1-\langle u, w \rangle|^{-\frac{\varepsilon}{2}} \log^{-p\beta} \frac{e}{|1-\langle u, w \rangle|} \lesssim 1.$$

When  $\beta > 0$  and  $s > p$ , it follows from Proposition 3.2(3) and (2.1) that

$$\int_{B_n} \frac{(1-|u|^2)^s |\nabla G_w(z)|^p \mu^p(|z|) dv(z)}{(1-|z|^2)|1-\langle z, u \rangle|^{2s}} \\ \lesssim \frac{(1-|u|^2)^s}{(1-|u|)^{s-p} |1-\langle u, w \rangle|^p} \log^{-p\beta} \frac{e}{|1-\langle u, w \rangle|} \log^{p\beta} \frac{e}{1-|u|^2} \lesssim 1.$$

When  $\beta \leq 0$ , it follows from Proposition B(6)-(8) that

$$\begin{aligned} & \int_{B_n} \frac{(1 - |u|^2)^s |\nabla G_w(z)|^p \mu^p(|z|) dv(z)}{(1 - |z|^2) |1 - \langle z, u \rangle|^{2s}} \\ & \lesssim \int_{B_n} \frac{(1 - |u|^2)^s (1 - |z|^2)^{p+s-n-1} dv(z)}{|1 - \langle z, u \rangle|^{2s} |1 - \langle z, w \rangle|^p} \lesssim 1. \end{aligned}$$

This shows that  $\|G_w\|_{p,\mu,s} \lesssim 1$ . If  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$ , then we may obtain that

$$\nu(|w|) |\nabla \psi(w)| \int_0^{|w|^2} \frac{1}{1-t} \log^{-\beta} \frac{e}{1-t} dt \lesssim \|M_\psi\|,$$

According to the arbitrariness of  $w$ , we can get (3.9) and (3.10) by different cases.

This proof is complete.  $\square$

In particular, when  $\nu(r) = (1 - r^2)^{\frac{n-s}{p}} \mu(r)$  in Corollary 3.6, it follows from Lemma 2.4 that  $F(p, \mu, s) \subseteq \mathcal{B}_\nu(B_n)$ . Moreover, there are the following results:

**Corollary 3.7** Let  $\mu(r) = (1 - r^2)^\alpha \log^\beta \frac{e}{1-r^2}$ . If  $\nu(r) = (1 - r^2)^{\alpha + \frac{n-s}{p}} \log^\beta \frac{e}{1-r^2}$  ( $\alpha > 0$ ,  $-\infty < \beta < \infty$ ) is a normal function on  $[0, 1)$  and  $\psi \in H(B_n)$ , then  $\psi$  is a pointwise multiplier from  $F(p, \mu, s)$  to  $\mathcal{B}_\nu(B_n)$  if and only if

- (1)  $\psi \in \mathcal{B}_\nu(B_n)$  when  $p\alpha + n - p < s$ , or  $p\alpha + n - p = s > 0$  and  $\beta > 1$ .
- (2)  $\psi \in H^\infty(B_n)$  when  $p\alpha + n - p > s$ .
- (3)  $\psi \in H^\infty(B_n)$  and

$$\sup_{z \in B_n} (1 - |z|^2) |\nabla \psi(z)| \left( \log \frac{e}{1 - |z|^2} \right) \log \log \frac{e}{1 - |z|^2} < \infty \quad (3.11)$$

when  $p\alpha + n - p = s > 0$  and  $\beta = 1$ .

- (4)  $\psi \in H^\infty(B_n)$  and

$$\sup_{z \in B_n} (1 - |z|^2) |\nabla \psi(z)| \log \frac{e}{1 - |z|^2} < \infty \quad (3.12)$$

when  $p\alpha + n - p = s > 0$  and  $\beta < 1$ .

**Proof** It follows from Lemma 2.4 that  $\mathcal{B}_\nu(B_n) \subseteq H^\infty(B_n)$  when  $\int_0^1 \mu^{-1}(\rho)(1 - \rho^2)^{\frac{s-n}{p}} < \infty$ . By Corollary 3.5 and Corollary 3.6, we may obtain these results.

**Note 2** In Corollary 3.7, no condition can be removed. For example, if  $\psi(z) = e^{\frac{z_1+1}{z_1-1}}$ , then  $\psi \in H^\infty(B_n)$ . But

$$\begin{aligned} & \sup_{z \in B_n} (1 - |z|^2) |\nabla \psi(z)| \log \frac{e}{1 - |z|^2} \geq \sup_{z_1 \in D} (1 - |z_1|^2) |\nabla \psi(z_1)| \log \frac{e}{1 - |z_1|^2} \\ & = \sup_{re^{i\theta} \in D} \frac{2(1 - r^2)}{1 + r^2 - 2r \cos \theta} e^{\frac{r^2-1}{1+r^2-2r \cos \theta}} \log \frac{e}{1 - r^2} \geq \sup_{0 < \theta < \frac{\pi}{2}} \frac{2}{e} \log \frac{e}{\sin^2 \theta} = \infty. \end{aligned}$$

This means that  $\psi$  does not meet (3.11) and (3.12). If  $\psi(z) = \log \log \frac{e^2}{1-z_1}$ , then it meets (3.12). But  $\psi$  does not belong to  $H^\infty(B_n)$ . If  $\psi(z) = \log \log \log \frac{e^4}{1-z_1}$ , then it meets (3.11). But  $\psi$  does not belong to  $H^\infty(B_n)$ .

**Statement:** No potential conflict of interest.

## References

- [1] K. H. Zhu, Spaces of Holomorphic Functions in the Unit Ball, Springer-Verlag (GTM 226), New York, 2005.
- [2] R. H. Zhao, On a general family of function spaces, Ann. Acad. Sci. Fenn. Math. Diss. 105(1996) 1-56.
- [3] X. J. Zhang, C. Z. He, F. F. Cao, The equivalent norms of  $F(p,q,s)$  space in  $\mathbb{C}^n$ , J. Math. Anal. Appl. 401(2013) 601-610.
- [4] S. L. Li, Bergman type operator on spaces of holomorphic functions in the unit ball of  $\mathbb{C}^n$ , J. Math. Anal. Appl. 514(2022) 126088.
- [5] R. H. Zhao, On  $F(p,q,s)$  spaces, Acta Math. Sci. 41B (2021) 1985-2020.
- [6] X. J. Zhang, The multipliers on several holomorphic function spaces, Chin. J. Conte. Math. 26(2005): 249-258.
- [7] L. J. Jiang, Y. Z. He, Composition operators from  $\beta^\alpha$  to  $F(p,q,s)$ , Acta Math. Sci. 23B(2003) 252-260.
- [8] Z. H. Zhou, R. Y. Chen, Weighted composition operator from  $F(p,q,s)$  to Bloch type spaces on the unit ball, Int. J. Math. 19(2008) 899-926.
- [9] S. L. Ye, Weighted composition operators from  $F(p,q,s)$  into logarithmic Bloch space, J. Kore. Math. Soc. 45(2008) 977-991.
- [10] S. L. Li, X. J. Zhang, S. Xu, The Bergman type operators on the  $F(p,q,s)$  type spaces in  $\mathbb{C}^n$ , Chin. J. Conte. Math. 38(2017) 303-316.
- [11] W. F. Yang, Volterra composition operators from  $F(p,q,s)$  spaces to Bloch-type spaces, Bull. Malay. Math. Sci. 34(2011) 267-277.
- [12] X. J. Zhang, S. L. Li, Q. L. Shang, Y. T. Guo, An integral estimate and the equivalent norms on  $F(p,q,s,k)$  spaces in the unit ball, Acta Math. Sci. 38B(2018) 1861-1880.
- [13] Y. X. Liang, On an integral-type operator from a weighted-type space to  $F(p,q,s)$  on the unit ball, Complex Var. Ellip. Equ. 60(2015) 282-291.
- [14] X. J. Zhang, Y. T. Guo, Q. L. Shang, S. L. Li, The Gleason's problem on  $F(p,q,s)$  type spaces in the unit ball of  $\mathbb{C}^n$ , Complex Anal. Oper. Theory. 12(2018) 1251-1265.
- [15] S. L. Li, X. J. Zhang, S. Xu, The equivalent characterization of  $F(p,q,s)$  space on bounded symmetric domains of  $\mathbb{C}^n$ , Acta Math. Sci. 37B(2017) 1791-1802.
- [16] J. Xiao,  $Q_\alpha$  Analysis on Euclidean Spaces, De Gruyter, 2019.
- [17] J. Xiao, Geometric  $Q_p$  Functions, Frontiers in Math. Birkhauser-Verlag, 2006.
- [18] H. S. Wulan, K. H. Zhu, Möbius Invariant  $Q_K$  Spaces, Springer-Nature, Switzerland, 2017.
- [19] W. Rudin, Function Theory in the Unit Ball of  $\mathbb{C}^n$ , Springer, New York, 1980.
- [20] G. D. Taylor, Multipliers on  $D_\alpha$ , Trans. Amer. Math. Soc. 123(1966) 229-240.
- [21] D. Stegenga, Multipliers of the Dirichlet space, Illinois J. Math. 24(1980) 113-139.
- [22] P. Y. Hu, J. H. Shi, Multipliers on Dirichlet type spaces, Acta Math. Sin. 17(2001) 263-272.
- [23] K. H. Zhu, Multipliers of BMO in the Bergman metric with applications to Toeplitz operators, J. Funct. Anal. 87(1989) 31-50.
- [24] S. Axler, A. Shields, Univalent multipliers of the Dieichlet space, Michigan Math. J. 32(1985) 65-80.
- [25] J. Ortega, J. Fàbrega, Pointwise multipliers and corona type decomposition in BMOA, Ann. Inst. Fourier (Grenoble). 46(1996) 111-137.

- [26] J. Pau, J. Peláez, Multipliers of Möbius invariant  $Q_s$  spaces, *Math. Z.* 261(2009) 545-555.
- [27] D. Stegenga, Bounded Toeplitz operators on  $H^1$  and applications of the duality between  $H^1$  and the functions of bounded mean oscillation, *Amer. J. Math.* 98(1976) 573-589.
- [28] R. H. Zhao, On logarithmic Carleson measures, *Acta Sci. Math (Szeged)*. 69(2003) 605-618.
- [29] S. Xu, X. J. Zhang, Multiplier and composition operator between several holomorphic function spaces in  $\mathbb{C}^n$ , *Complex Anal. Oper. Theory*. 15(2021) 36.
- [30] M. J. Tan, X. J. Zhang, Equivalent characterizations and pointwise multipliers of normal weight Dirichlet space on the unit ball, *Complex Anal. Oper. Theory*. 15(2021) 24.
- [31] X. J. Zhang, L. H. Xi, H. X. Fan, J. F. Li, Atomic decomposition of  $\mu$ -Bergman space in  $\mathbb{C}^n$ , *Acta Math. Sci.* 34B(2014) 779-789.
- [32] Z. J. Hu, Composition operators between Bloch-type spaces in the polydisc, *Science in China, Ser. A.* 48 Supp(2005) 1-15.
- [33] X. J. Zhang, S. Xu, Weighted differentiation composition operators between normal weight Zygmund spaces and Bloch spaces in the unit ball of  $\mathbb{C}^n$  for  $n > 1$ , *Complex Anal. Oper. Theory*. 13(2019) 859-878.
- [34] J. Ortega, J. Fabrega, Corona type decomposition in some Besov spaces, *Math. Scand.* 78(1996) 93-111.
- [35] R. H. Zhao, Distances from Bloch functions to some Möbius invariant spaces, *Annl. Acad. Sci. Fen. Math.* 33(2008) 303-313.
- [36] X. J. Zhang, R. X. Lv, P. C. Tang, Several equivalent characterizations of general Hardy type spaces on the unit ball in  $\mathbb{C}^n$ , *Chin. J. Conte. Math.* 40 (2019) 101-114.

COLLEGE OF MATHEMATICS AND STATISTICS, HUNAN NORMAL UNIVERSITY, CHANGSHA,  
HUNAN 410006, CHINA  
*Email address:* xuejunttt@263.net