Quantum Back-action Limits in Dispersively Measured Bose-Einstein Condensates

Emine Altuntaş^{*} and I. B. Spielman[†]

National Institute of Standards and Technology, Gaithersburg, MD, 20899, USA. and

Joint Quantum Institute, University of Maryland, College Park, MD, 20742, USA.

(Dated: April 13, 2023)

A fundamental tenet of quantum mechanics is that measurements change a system's wavefunction to that most consistent with the measurement outcome, even if no observer is present. Weak measurements produce only limited information about the system, and as a result only minimally change the system's state. Here, we theoretically and experimentally characterize quantum back-action in atomic Bose-Einstein condensates interacting with a far-from resonant laser beam. We theoretically describe this process using a quantum trajectories approach where the environment measures the scattered light and present a measurement model based on an ideal photodetection mechanism. We experimentally quantify the resulting wavefunction change in terms of the contrast of a Ramsey interferometer and control parasitic effects associated with the measurement process. The observed back-action is in good agreement with our measurement model; this result is a necessary precursor for achieving true quantum back-action limited measurements of quantum gases.

Back-action limited weak measurements are essential for advancing quantum technologies, enable new probes of quantum systems, and offer new ways to understand the measurement process. Most quantum technologies simultaneously require quantum limited measurements and feedback control to establish and maintain quantum coherence and entanglement, with applications ranging from quantum state preparation [1, 2] to quantum error correction[3]. Even without feedback, system dynamics combined with weak measurements can lead to entangled states in the thermodynamic limit [4–7]. Large-scale applications of these capabilities hinge on understanding system-reservoir dynamics of many-body quantum systems, whose Hilbert space grows exponentially with system size. Ultracold atoms, a workhorse for quantum simulation[8, 9], are an ideal platform for studying the system-reservoir dynamics of large-scale many-body systems.

Weakly measured quantum systems can be understood using the robust framework of quantum trajectories[10, 11]. In these descriptions, the system and a larger reservoir interact and become weakly entangled, at which point the reservoir is projectively measured. This destroys the system-reservoir (SR) entanglement and leads to a change in the system's wavefunction. We develop such a measurement model to study the interplay between the system-reservoir interaction, the scattered light, and the post-measurement system state.

Very far from atomic resonance light Rayleigh-scatters from atomic ensembles, changing the incident light's wavevector in proportion to the Fourier transform of the atomic density distribution. The straightforward interpretation of back-action resulting from scattered photons makes quantum trajectories an ideal tool for both intuitively and quantitatively understanding the systemreservoir interaction. When the reservoir-measurement outcomes are rejected, quantum trajectories methods form a specific physically motivated "unraveling" of the master equation[11]. In the quantum problem, light scattering gives information both about the expectation value of the density—essentially classical scattering—as well as quantum fluctuations, which contribute to spontaneous emission. Quite recently a trio of papers observed the predicted suppression of light-scattering from deeply degenerate Fermi gases[12–14] as well as amplification from ultracold Bose gases[15]; these effects result from scattering atoms into occupied quantum states.

Ultracold atoms have multiple well-established "nondestructive" measurement techniques[16–21]. While backaction-induced heating of a single motional degree of freedom of a BEC was observed in a single mode optical cavity[22], previous demonstrations of such methods with spatial resolution did not quantify quantum backaction.

Here we characterize measurement back-action in atomic Bose-Einstein condensates (BECs), weakly interacting with a far-from resonant laser beam. The information extracted by light-scattering can be treated as a quantum measurement process where the scattered light is detected by the environment [Fig. 1a-b], and we-the observer-detect only the resulting back-action on the system. The wavefunction change is quantified by the phase shift and contrast of a Ramsey interferometer. In our Ramsey interferometer [Fig. 1c], spontaneously scattered light measures atoms to be in the detected spin state, thereby breaking its coherence and reducing the interferometer contrast. We further distinguish between non-destructive measurements (where the system is apparently undisturbed) and back-action limited measurements (where observed quantum projection noise dominates the change in the post-measurement state). We systematically control for two stray effects that otherwise lead to excess excitation or loss: inhomogeneities in the probe beam, and a weak optical lattice from weak backreflections of the probe beam. We explore a third systematic effect: light induced collisions—intrinsic atomic

 $[\]ast$ altuntas@umd.edu

[†] ian.spielman@nist.gov

processes—that were found to have limited impact on our Ramsey data. We demonstrate that these technical artifacts can be eliminated, bringing the observed backaction into agreement with our measurement model.

RESULTS

Quantum trajectories model

We consider a weakly interacting atomic BEC (the system) dispersively coupled to the optical electric field $\hat{\mathbf{E}}(\mathbf{x},t)$ (the reservoir) by the ac Stark shift with interaction picture Hamiltonian

$$\hat{H}_{\rm SR}(t) = \int \frac{d^3 \mathbf{x}}{\hbar \Delta} \hat{n}_{\rm g}(\mathbf{x}) \otimes [\hat{\mathbf{E}}(\mathbf{x},t) \cdot \mathbf{d}_{\rm ge}] [\mathbf{d}_{\rm ge}^* \cdot \hat{\mathbf{E}}^{\dagger}(\mathbf{x},t)].$$
(1)

Here $\hat{n}_{g}(\mathbf{x}) = \hat{b}_{g}^{\dagger}(\mathbf{x})\hat{b}_{g}(\mathbf{x})$ is the atomic density operator in terms of the bosonic field operators $\hat{b}_{g}(\mathbf{x})$ for ground state atoms at position \mathbf{x} ; \mathbf{d}_{ge} is the dipole matrix element for transitions between ground and excited state atoms with energy difference $\hbar\omega_{ge}$; lastly, $\Delta = \omega_{0} - \omega_{ge}$ is the detuning from atomic resonance of a probe laser with frequency ω_{0} .

For $|\Delta| \ll \omega_{\rm ge}$, the optical electric field operator is

$$\hat{\mathbf{E}}(\mathbf{x},t) = i\sqrt{\frac{\hbar\omega_{\text{ge}}}{2\epsilon_0}} \sum_{\sigma} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \hat{a}_{\sigma}(\mathbf{k}) \boldsymbol{\epsilon}_{\sigma}(\mathbf{k}) e^{i(\mathbf{k}\cdot\mathbf{x}-c|\mathbf{k}|t)},$$
(2)

expressed in terms of field operators $\hat{a}_{\sigma}(\mathbf{k})$ describing states with wavevector \mathbf{k} and polarization σ . Here, cis the speed of light; ϵ_0 is the electric constant; and $\epsilon_{\sigma}(\mathbf{k})$ are a pair orthogonal polarization vectors transverse to \mathbf{k} , labeled by $\sigma = \pm$. Figure 1a depicts the full systemreservoir coupling scheme with the BEC interacting with outgoing transverse modes and a probe laser in mode (\mathbf{k}_0, σ_0) for a duration $t_{\rm m}$.

During this time the atomic ensemble scatters monochromatic light into outgoing modes of wavevector \mathbf{k}_{\perp} with coupling strength

$$g_{\sigma}(\mathbf{k}_{\perp}) \equiv -i \left(\frac{\omega_{\rm ge}}{2\hbar\epsilon_0}\right)^{1/2} \left[\mathbf{d}_{\rm ge} \cdot \boldsymbol{\epsilon}_{\sigma}(\mathbf{k}_{\perp})\right]. \tag{3}$$

Since each outgoing mode is in a specific polarization state $\epsilon(\mathbf{k}_{\perp})$ the polarization subscript is redundant.

Assuming that the probe laser of wavelength λ occupies a single optical mode (\mathbf{k}_0, σ_0) with $k_0 \equiv |\mathbf{k}_0| = 2\pi/\lambda$, we make the replacement $\hat{a}_{\sigma}(\mathbf{k}) \rightarrow \delta(\mathbf{k} - \mathbf{k}_0)\delta_{\sigma,\sigma_0}\alpha_0 + \hat{a}_{\sigma}(\mathbf{k})$, which describes a coherent driving field with amplitude α_0 . In this expression the modes $\hat{a}_{\sigma}(\mathbf{k})$ are initially empty. This replacement allows us to expand Eq. (1) in decreasing powers of the large parameter α_0 . The leading term describes the ac Stark shift, and the next term

$$\hat{H}_{\rm eff} = \frac{\hbar P_{\rm e}^{1/2}}{(ct_{\rm m})^{1/2}} \oint_{k_0} \frac{d^2 \mathbf{k}_{\perp}}{(2\pi)^2} g^*(\mathbf{k}_{\perp}) \hat{n}_{\mathcal{F}}(\mathbf{k}_{\perp} - \mathbf{k}_0) \hat{a}^{\dagger}(\mathbf{k}_{\perp}) + \text{H.c},$$

describes scattering from the probe field into outgoing modes by any structure in the atomic density, with Fourier components

$$\hat{n}_{\mathcal{F}}(\mathbf{k}_{\perp} - \mathbf{k}_0) = \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \hat{b}^{\dagger} [\mathbf{k} - (\mathbf{k}_{\perp} - \mathbf{k}_0)] \hat{b}(\mathbf{k}).$$

Here $P_{\rm e} = |\alpha_0 g_{\sigma_0}(\mathbf{k}_0)|^2 / \Delta^2$ is the excited state occupation probability. In the far-detuned limit, the outgoing wavenumber is fixed at k_0 leading to the surface integral over the sphere of radius k_0 .

We model the larger environment as performing measurements on the outgoing light in the far-field with an ideal photo detection process, a strong measurement of the photon density $\hat{a}^{\dagger}(\mathbf{k}_{\perp})\hat{a}(\mathbf{k}_{\perp})$ [Fig. 1b]. In the abstract, this process begins with the combined system reservoir state $|0\rangle \otimes |\Psi_{\rm S}\rangle$, describing a reservoir with no photons but with the system in an arbitrary state. This state evolves briefly for a time $t_{\rm m}$ via the time evolution operator $\hat{U}_{SR}(t_{\rm m}) = \mathcal{T} \exp \left[-i \int_{-t_{\rm m}/2}^{t_{\rm m}/2} \hat{H}_{\rm eff}(t) dt/\hbar\right]$. This entangles the system and reservoir; as depicted in Fig. 1a amplitude can be present in every reservoir mode prior to measurement by the environment.

Photodetection

We turn to the photodetection model shown in Fig. 1b. In this case, the measurement of the reservoir collapses the superposition by measuring either no photons or a single photon in final state $|\mathbf{k}_{\perp}\rangle$. The back-action of this measurement is described by a conditional change in the system wavefunction $|\Psi'_{S}\rangle = \hat{M}(\mathbf{k}_{\perp}) |\Psi_{S}\rangle$, an operation described by Kraus operator $\hat{M}(\mathbf{k}_{\perp}) = \langle \mathbf{k}_{\perp} | \hat{U}_{SR}(t_m) | 0 \rangle$. Taken together this schema is a generalized measurement of the system effected by projective measurements on the reservoir.

In the limit of small $t_{\rm m}$, such that at most one photon is scattered, we obtain the Kraus operator

$$\hat{M}(\mathbf{k}_{\perp}) = -iP_{\rm e}^{1/2} \left(\frac{t_{\rm m}}{c}\right)^{1/2} g^*(\mathbf{k}_{\perp}) \hat{n}_{\mathcal{F}}(\mathbf{k}_{\perp} - \mathbf{k}_0) \quad (4)$$

describing the recoil of the system from momentumconserving scattering out of every occupied state.

The Kraus operator contains information both about the change in the system as well as the probability density

$$P(\mathbf{k}_{\perp}) \equiv \langle \Psi_S | \, \hat{M}^{\dagger}(\mathbf{k}_{\perp}) \hat{M}(\mathbf{k}_{\perp}) | \Psi_S \rangle$$

$$= \frac{t_{\rm m} P_{\rm e}}{c} |g(\mathbf{k}_{\perp})|^2 \langle \Psi_S | \left| \hat{n}_{\mathcal{F}}(\mathbf{k}_0 - \mathbf{k}_{\perp}) \right|^2 | \Psi_S \rangle$$
(5)

that this change occurred. Bringing $|\hat{n}_{\mathcal{F}}|^2$ into a normalordered form shows that the scattering probability has two contributions. For a BEC with condensate mode $\tilde{\psi}(\mathbf{k})$ the scattering probability is

$$P(\mathbf{k}_{\perp}) = \frac{t_{\rm m} P_{\rm e}}{c} |g(\mathbf{k}_{\perp})|^2 N \left[(N-1) \left| n_{\mathcal{F}}(\mathbf{k}_0 - \mathbf{k}_{\perp}) \right|^2 + 1 \right],$$



FIG. 1. Photodetection measurement model and Ramsey interferometry. a Interaction. A Bose-Einstein condensate (BEC) illustrated in blue, is illuminated with far-detuned laser light (red) for a time $t_{\rm m}$ and scatters light (wiggly lines) into both occupied (red) and reservoir (orange) modes. b Measurement. The reservoir modes are projectively measured by an array of photo-detectors encompassing 4π steradians yielding the outgoing wavevector and polarization. c Level diagram. d Time sequence for Ramsey interferometry. An initial $\pi/2$ microwave pulse (light blue) is followed by a 15 μ s evolution period; then the $t_{\rm m} = 20 \ \mu$ s off-resonant light pulse (red) is applied, and after a 5 μ s delay (giving a total $T = 40 \ \mu$ s free evolution time) the Ramsey sequence is completed with a second $\pi/2$ pulse (light blue). The optical dipole trap (ODT), denoted by the orange dashed line, is extinguished immediately following the Ramsey sequence. A Stern-Gerlach (SG) gradient (grey) is applied during time-of-flight (TOF) and the final density is detected using absorption imaging (purple). e Bloch sphere depiction of Ramsey interferometry. The dark blue arrows depict the axes of rotation for each microwave pulse and the light blue arrows mark the associated trajectories. The green circles show the coherent evolution during each step of our sequence. Red arrows depict evolution associated with the measurement pulse with the solid curve resulting from the Stark shift and dashed curves resulting from measurement back-action. The red circles are the states that were measured to be in $|g_2\rangle$. Translucent (solid) symbols indicate the initial (final) state.

where in analogy with the operator expression, $n_{\mathcal{F}}(\mathbf{k}_0 - \mathbf{k}_\perp)$ describes the Fourier components of the probabilitydensity. The first term describes collective scattering from the overall density profile (including thermal fluctuations), a.k.a. classical scattering[23], while the second results from scattering from quantum fluctuations, here giving rise to spontaneous emission. For extended systems such as our BEC, the collective term is dominated by small angle forward scattering while the spontaneous term is nominally isotropic. Notably, this result illustrates that the ratio between collective and spontaneous scattering depends on N but not the measurement parameters.

Integrating over the final \mathbf{k}_{\perp} states gives $P_{\text{tot}} = P_{\text{col}} + P_{\text{sp}}$ with the spontaneous scattering probability $P_{\text{sp}} = \Gamma t_{\text{m}} P_{\text{e}} = g^2/8$. We introduced an overall measurement strength $g = \sqrt{\bar{t}_{\text{m}} \bar{I}}/\bar{\delta}$ in terms of dimensionless: time $\bar{t}_{\text{m}} = \Gamma t_{\text{m}}$ scaled by the natural linewidth Γ ; detuning $\bar{\delta} = \Delta/\Gamma$ in units of Γ ; and laser intensity $\bar{I} = I/I_{\text{sat}}$ in units of the saturation intensity I_{sat} . Thus when $g = \sqrt{8}$ each atom will have on average spontaneously scattered a single photon (The relation between g and the signal to noise ratio of a measurement outcome is briefly discussed in Supplementary Note 1.).

In experiment, a single measurement pulse can lead to thousands of photodetection events, each described by a Kraus operator. The concatenation of many such Kraus operators—one for each scattering event—describes the evolution of our system. By contrast with master equation methods that trace out the environment, quantum trajectories approaches predict individual measurement outcomes and the associated back-action, drawn from a suitable statistical distribution. Thus, the final postmeasurement state can be predicted given an experimentally observed measurement record. For ensemble averaged predictions, our technique and standard methods such as those used in Appel et al.[24] give the same results. We compare the predictions of this theoretical description with an observable, contrast in a Ramsey interferometer, that does not rely on knowledge of the specific quantum trajectory that the system followed.

Experimental system

Our experiments started with highly elongated ⁸⁷Rb BECs prepared in a crossed opdipole tical trap (ODT)with frequencies $(\omega_x, \omega_y, \omega_z) = 2\pi \times [9.61(3), 113.9(3), 163.2(3)]$ Hz in the $|g_1\rangle \equiv |F=1, m_F=1\rangle$ electronic ground state (All uncertainties herein reflect the uncorrelated combination of single-sigma statistical and systematic This trap configuration yielded conuncertainties). densates with $N_{\rm c} = 0.70(15) \times 10^5$ atoms[25, 26], condensate fraction $R_c = 78(3)\%$, and chemical potential $\mu = h \times 0.76(6)$ kHz. We drove transitions between $|q_1\rangle$ and $|q_2\rangle \equiv |F=2, m_F=2\rangle$ using an ≈ 6.8 GHz microwave magnetic field with Rabi frequency ≈ 7.5 kHz.

In our experiments we illuminated the BEC in situ with an off-resonant probe laser beam that drove the $|g_2\rangle$ to $|e\rangle \equiv |F' = 3, m'_F = 3\rangle$ ground to excited state transition. This probe laser was blue detuned by $0 < \bar{\delta} < 317$, and had intensity $\bar{I} \leq 10$. We theoretically describe the light scattered at large angle as being subsequently projectively measured by the environment, as described above. We then detected the post-measurement density distribution using absorption imaging after a longer 20 ms TOF during which a Stern-Gerlach gradient spatially separated the $|g_1\rangle$ and $|g_2\rangle$ components.

Detecting Wavefunction Change via Ramsey Interferometry

We characterize the light matter interaction, as well as back-action, predicted by our quantum trajectories model using Ramsey interferometry (RI). Our Ramsey interferometer [Fig. 1d-e] commenced with a resonant microwave pulse driving a $\pi/2$ rotation about \mathbf{e}_{u} , taking the atoms from $-\mathbf{e}_z$ (in $|g_1\rangle$) to \mathbf{e}_x . Then during the free evolution time we applied the probe laser detuned by δ from the $|g_2\rangle$ to $|e\rangle$ transition for a time $t_{\rm m}$; the resulting ac Stark shift drove a rotation about \mathbf{e}_z by ϕ (solid red arc). A second microwave pulse drove a $\pi/2$ rotation about an axis rotated by $\delta \phi_{\rm P}$ at which time we measured the final populations N_1 and N_2 in $|g_1\rangle$ and $|g_2\rangle$ respectively in TOF, giving the fraction in $|g_2\rangle$ as $R_2 = N_2/(N_1 + N_2)$. The black data (squares) in Fig. 2a, taken with the probe laser off, shows that the resulting fractional population R_2 is cosinusoidal, and the red data (circles), with the probe on, is phase shifted (from the ac Stark shift on $|g_2\rangle$). We obtain the phase shift ϕ , contrast A, and center shift b with fits to $R_2 = [1 + A\cos(\delta\phi_P + \phi)]/2 + b.$

The RI phase shift is a direct measure of the differential phase acquired during free evolution, here $-V_{\rm ac}t_{\rm m}/\hbar$ from the ac Stark shift of $|g_2\rangle$ due to the probe beam, with $V_{\rm ac} = \Gamma \bar{I}/(8\bar{\delta})$. The Stark shift of $|g_1\rangle$ is a small contribution that we do not include in our fits. The data in Fig. 2b was taken at $\overline{\delta} = 63.4$ and 116.2 (circles and squares respectively). As expected the slope is larger for smaller δ , but in both cases the acquired phase can exceed 2π at which point it wraps back to zero. The intensity of the probe laser is difficult to obtain in-vacuo[27, 28]; however, fitting $t_{\rm m}V_{\rm ac}$ to these data gives a direct calibration of the laser intensity, providing a conversion between our camera signal and I_{sat} with < 5 % fractional uncertainty. We imaged the *in situ* probe beam (with no atoms present) on a charge coupled device camera to obtain the local probe intensity (in arbitrary camera units) at the location of the BEC. Further details are described in Altuntas et al. [29]. The solid lines in Fig. 2b are the result of this fitting process.

Figure 2a shows a second effect of increasing measurement strength (blue data): the Ramsey contrast decreases with increasing measurement strength, implying that the post-measurement many-body wave function is



FIG. 2. Ramsey interferometry data. a Ramsey oscillation without (black squares) and with the light pulse at $\bar{\delta} = 63.4$ and $\bar{I} \approx 2$ (red circles) and $\bar{I} \approx 7$ (blue triangles), taken *in situ* as the phase jump $\delta\phi_{\rm P}$ between the two pulses is varied. Solid curves are fits to the equation given in the text. Blue Ramsey data $\delta\phi_P$ values are deliberately shifted by 2π to visualize the reduction in contrast and the larger phase shift. **b** Optically induced phase shift ϕ as a function \bar{I} at $\bar{\delta} = 63.4$ (red circles) and $\bar{\delta} = 116.2$ (green squares). The same-color lines are fits to $-V_{\rm ac}t_{\rm m}/\hbar \mod 2\pi$. Shaded regions indicate the $\pm 1\sigma$ statistical uncertainty range.

not described by a coherent superposition of $|g_1\rangle$ and $|g_2\rangle$.

Our measurement model predicts this effect: as illustrated in the middle Bloch sphere in Fig. 1e, each time a photon is spontaneously scattered and detected by the environment, the wavefunction of a single atom collapses into $|q_2\rangle$ (along \mathbf{e}_z), losing any coherence with $|q_1\rangle$ (red dashed arrows). The second $\pi/2$ pulse always returns that atom to the equator of the Bloch sphere, reducing the contrast by 1/N. In this situation, the per-atom probability of scattering a single photon at large angle is $g^2/16$ (see Supplementary Note 2 for the complete calculation). By contrast for collective scattering (generally at small-angle), a detected photon scattered off of the global density distribution yields Mössbauer-like collective back-action and no reduction in contrast. As a result, the change in contrast measures the number of spontaneously scattered photons [24].

The ideal Ramsey interferometry scheme presented in Fig. 1d is sensitive to additional systematic effects leading to contrast reduction. In the following sections we identify such factors, and develop an enhanced RI scheme that detects the post-measurement wavefunction change in agreement with the theoretical prediction.



FIG. 3. Improved Ramsey interferometers. a, b Pulse sequences for spin-echo (SE) and pulse-evolve-pulse with spinecho (PEP-SE) Ramsey interferometers. The red blocks denote the dispersive measurement pulses, and blue bars indicate the microwave pulses. c Interferometer signal measured using: a conventional Ramsey sequence (blue squares), a spinecho Ramsey sequence (green circles), and a pulse-evolvepulse with spin-echo sequence (magenta diamonds). All measurements were performed *in situ* at $\bar{\delta} = 63.4$ with $\bar{I} \approx 7$ yielding $g \approx 1$.

Spin-echo Ramsey Interferometer

Spatial inhomogeneities in the probe beam as well as near-dc magnetic field noise can reduce the RI contrast. In the first case, the resulting position-dependent ac Stark shift imprints spatial structure to the RI phase ϕ , thereby reducing the spatially averaged contrast. Second, because the $|g_1\rangle$ - $|g_2\rangle$ transition is first-order sensitive to the external magnetic field, the RI contrast is reduced when field noise randomly shifts the resonance condition between different repetitions of the experiment.

We added a spin-echo pulse to our interferometer [see Fig. 3a] to compensate for both of these parasitic effects. As Fig. 3c shows, the noise in the spin-echo signal (circles) is reduced compared to the standard RI measurement (squares). Although the measurement noise is reduced, the contrast with spin echo is unchanged (Fig. 4a-b), leaving the substantial disagreement with our theory prediction (black curve) due to the systematic factor we report next.

Ramsey Interferometer with Split Measurement Pulses

Contrary to our predictions, Fig. 4a-b show that the contrast depends on probe detuning (green squares versus red circles). This difference signifies the presence of the second parasitic effect: a weak optical lattice generated by the probe beam interfering with its retroreflections off subsequent optical elements. The probe beam is nearly perfectly concentric with our imaging system and intersects each optical element at normal incidence. While it is common practice in optical setups to slightly tilt optical elements to eliminate back-reflections, in the high-resolution imaging context optimized alignment is a necessary condition for minimizing optical aberrations.

As a result, each probe pulse corresponds to the sudden application of a lattice potential. Weak lattices create populations in matterwave diffraction orders with momentum $\pm 2\hbar k_0$. In principle a suitable spin-echo sequence could remedy this, nonetheless, the rapidly moving diffracted atoms experience different lattice potentials during our first and second pulses precluding effective cancellation.

Instead we extended the ideas in Wu et al. [30] and Herold et al. [31] by splitting each probe pulse into two pulses of duration $t_{\rm p} = 8.2 \ \mu s$ spaced in time by a carefully chosen $t_{\rm d} = 25.6 \ \mu {\rm s}$ of free evolution, essentially unwinding the phase imprinted by the lattice (see Supplementary Note 3). Fig. 3b shows such a pulse-evolvepulse with spin-echo (PEP-SE) sequence. The near-full contrast magenta Ramsey fringe in Fig. 3c results from this PEP-SE sequence applied *in-situ* for $g \approx 1$. As seen in Fig. 4c, there is negligible difference in the extracted contrast between measurements at the same qvalue but with different probe detunings (squares and circles) further confirming control over systematic effects. The PEP-SE Ramsey contrast is in good agreement with our theoretical model (black curve) and provides a mechanism for identifying the regime of backaction limited measurements of ultracold gases. In order to obtain a quantitative metric for comparison with theoretical prediction, we fit the $\bar{\delta} = 63.4$ data in $g \leq 1$ to $A = A_0 - \alpha g^2$, where A_0 describes a small overall reduction in contrast. PEP-SE scheme measurements yield $\alpha = 0.083(10)$, which is in good agreement with the theortical prediction $\alpha_{\rm th} = 1/16 \approx 0.063$.

Light induced collisions

We used the post-measurement atom number as an auxiliary probe of measurement back-action and found that, although photoassociation (PA) is suppressed at blue detuning, at our high *in situ* atomic densities of 1×10^{14} cm⁻³, light induced collisions lead to rampant atom loss[15]. We quantify the importance of these losses by preparing BECs with N_0 total atoms in $|g_2\rangle$ and measuring fractional change in total atom number N_t/N_0 and in uncondensed number $N_{\rm nc}/N_0$. $N_{\rm nc}/N_0$ counts both thermal atoms as well as atoms that have undergone large-angle light scattering.

Fig. 5a confirms that this is a 2-body process by reducing the atomic density with a short TOF. We find that the losses rapidly drop starting at $t_{\rm TOF} \approx 0.5$ ms



FIG. 4. Ramsey interferometer contrast A dependence on the measurement strength g for different schemes. a In situ Ramsey interferometry measurements. b In situ Ramsey interferometry with spin-echo measurements. c In situ Pulse-evolve-pulse with spin-echo Ramsey interferometry measurements. d Pulse-evolve-pulse with spin-echo sequence after 2 ms TOF measurements. The horizontal blue lines show the RI contrast observed without the measurement pulse with the shaded regions indicating the $\pm 1\sigma$ statistical uncertainty range. The black curves plot the prediction of our photodetection model. The pink curves depict a fit of the $\bar{\delta} = 63.4$ data to $A = A_0 - \alpha g^2$ with each best-fit alpha value quoted on the respective figure.

(when mean-field driven expansion becomes significant) and vanish after 3 ms (at which time the density has dropped by a factor of nearly 20). We also investigated another potential loss mechanism due to two-color PA resulting from the combination of the intense dipole trapping beam and the probe beam. Data taken just before (star symbols at negative time for clarity) and just after the ODT turn-off have no difference in loss, confirming the absence of any two-color PA effects.

Panel b, taken in situ, shows that the fractional number is independent of $\bar{\delta}$. These data were taken at constant $P_{\rm e}$ (achieved by tuning \bar{I}) and demonstrate that there are no PA resonances. Figure 5c shows that in situ the total number drops rapidly with increasing g^2 while the number outside the BEC remains constant. This verifies that the high-density BEC experiences light induced collisions while the low density thermal cloud is left mostly unchanged. Lastly Fig. 5d plots these quantities following a 2 ms TOF, confirming the same reduced losses found in Fig. 5a. Furthermore $N_{\rm nc}$ increases linearly with slope $g^2/8$ (red dashed curve) as expected from photon scattering.

All of these data are well described by a 2-body loss model (solid curves), however, these simulations require a 2-body coefficient that is about $20 \times$ in excess of the PA loss coefficient found in Fuhrmanek et al. [32]. In fact these observations reflect different processes: in the blue-detuned case light induced collision leads to rapidly accelerated atom pairs rather than PA[33].

Lastly, we note that light traversing the BEC acquires a phase shift causing the atomic cloud to act as a lens. When the phase shift is in excess of about 1 radian the scattering is no longer described by our model and atomic cloud experiences excess compression, potentially enhancing 3-body loss. The absence of $\bar{\delta}$ dependence in Fig. 5b affirms that effects such as this arising from the ac Stark shift do not contribute to loss.

As light assisted collisions precipitate atom loss, we added a short TOF to the spin-echo pulse-evolve-pulse Ramsey sequence to study the impact of light assisted collisions on RI contrast. As shown in Fig. 4d, the contrast is modestly reduced, and as with Fig. 4a-b, data taken at larger detuning are impacted more significantly. We attribute this reduction to the changing optical intensity profile that the falling BEC experiences as it traverses different regions of the probe beam during the pulse sequence; this compromises the PEP-SE sequence.

DISCUSSION

Even though RI contrast is a direct measure of the overall wavefunction change, our light assisted collision data show that RI contrast alone is insufficient to identify back-action dominated measurement regimes. For our *in situ* results—with rampant light induced losses photon scattering from the measurement process does not fully explain the change of the system's state. Consequently such measurements are not back-action limited, even in principle. An interesting question that we did not touch on, is how light induced collisions are able to remove atoms while leaving the Ramsey contrast largely unchanged.

For the modest range of detuning explored here, the two-body loss rate scales as the excited state probability $P_{\rm e} \propto g^2$; this implies that for a target measurement strength, light induced collisions are not reduced until vastly larger detuning when this scaling breaks down[34]. In our experiment, data taken with $g \lesssim 0.3$ (with peratom spontaneous scattering probability $P_{\rm sp} \lesssim 0.01$) had no discernible loss in Ramsey contrast or reduction in atom number: functionally non-destructive[16].



FIG. 5. Light induced collisions. Fractional total number of atoms (green squares) and number of atoms outside the BEC but within a 1 recoil momentum circle (blue circles) following measurement. The symbols mark experimental data while the curves are the result of our 2-body model. In all cases, the opacity of the points reflects the detuning. a Dispersive measurements at two different probe detunings ($\bar{\delta} = 84.5$ and $\bar{\delta} = 126.7$) after a variable short time-of-flight (TOF). All measurements were at measurement strength $g \approx 1$, which was attained by adjusting intensity to $\bar{I} \approx 9.5$ and $\bar{I} \approx 21$ respectively. Star symbols mark in situ measurements with the optical dipole trap on (plotted at negative TOF for display purposes). **b** In situ measurements at different $\bar{\delta}$ all with $g \approx 1$. Measurement time was $t_{\rm m} = 25 \ \mu {\rm s}$ in (a) and (b). c and d Loss as a function of q^2 for measurements made *in situ* in (c) and with a 2 ms TOF in (d). The red dashed line in (d) plots the expected $q^2/8$ light-scattering behavior. Both cases were at $\bar{\delta} = 84.5$ and $\bar{I} \approx 9.5$ with $t_{\rm m}$ varied from 4 μ s to 36 μ s.

ever, our results demonstrate that such functionally nondestructive measurements can be far from quantum backaction limited. As a consequence, back-action limited measurements of BECs can be achieved either by managing the atom density, or by careful control of molecular resonances[35]. In degenerate Fermi gases the Pauli pressure leads to much lower densities[36], typically diluted by an order of magnitude or more compared to BECs, making two-body losses less significant.

Employing the strategies identified here is necessary to achieve back-action limited measurements, and as a next step the scattered light must actually be detected. There are multiple imaging techniques for quantum gases based on the dispersive light-matter interaction[16, 20, 37, 38] that in principle can give back-action limited measurement outcomes. Implementing these requires an imaging system with minimal losses and large numerical aperture in conjunction with a high efficiency detector, as any scattered light that is not detected is effectively measured by the environment and its information lost. Furthermore, the captured signal must lead to a faithful representation of the atomic ensemble, necessitating an imaging system with minimal or well-calibrated aberrations as we demonstrated previously[21]. Lastly, the initial optical field must be well known, for which techniques such as outlined here and described in more detail in Altuntas et al. [29], are essential. These physical considerations do not touch on technical matters such as calibrating the response and hardware specific noise properties of the physical detector, i.e., a charge coupled device (CCD) or complementary metal oxide semiconductor (CMOS) camera. Future work needs to account for these sources of technical noise.

Looking forward, back-action limited weak measurements coupled with real-time control are enabling tools for quantum technology. Feedback cooling is one application of closed loop quantum control, and the interplay between measurement back-action and the actual information extracted from the system limits the achievable temperature[39–41]. In addition to simply cooling into established quantum states (both weakly and strongly correlated), closed-loop feedback enables the engineering of artificial, non-local, and non-Markovian, reservoirs. Existing proposals with engineered reservoirs show that suitable quantum jumps lead to equilibration into strongly correlated states [42]; and schemes using feedback can generate new Mott insulating phases [43] and squeezed states [44, 45]. In the latter case a single measurement locally creates conditional squeezing that requires a second spatially resolved control pulse conditioned on the measurement outcome-to obtain useful unconditional squeezing. Metrological implementations would also require atoms individually confined in the sites of an optical lattice to prevent spatial diffusion and clock shifts.

In addition, weak measurements offer new ways to explore fundamental concepts in quantum mechanics. For example, a weak measurement of strength g can be decomposed into a series of N sub-measurements[46, 47] each with strength g/\sqrt{N} . In this configuration, the total outcome of these measurements recovers an individual measurement of strength g, but the quantum backaction of earlier sub-measurements, giving information that is erased in a single stronger measurement. For example, correlating the outcome of two sub-measurements can isolate the measurement back-action of the first measurement.

METHODS

A. Magnetic field lock

Our interferometry measurements operate on the magnetic field sensitive $|F = 1, m_F = 1\rangle$ to $|F = 2, m_F = 2\rangle$ transition, and as a result are negatively impacted by magnetic field noise. To minimize any effect on contrast, we monitored the field shifts using a microwave based monitoring scheme first implemented in LeBlanc et al.[48].

Our two level system is well described by the Hamiltonian

$$\hat{H}_{\mu} = \frac{\hbar}{2} \left(\begin{array}{cc} \Delta_{\mu} + \delta_{\mu} & \Omega_{\mu} \\ \Omega_{\mu} & -(\Delta_{\mu} + \delta_{\mu}) \end{array} \right),$$

where Δ_{μ} describes an unknown detuning from resonance, δ_{μ} is an adjustable detuning, and Ω_{μ} is the microwave Rabi frequency.

Our protocol began with optically trapped atoms just above T_c in the $|F = 1, m_F = 1\rangle$ hyperfine state. We applied a microwave pulse of duration $t_{\mu} = 100 \ \mu s$ and Rabi frequency $\Omega_{\mu}/(2\pi) \approx 0.1/t_{\mu}$ detuned by $\delta_{\mu}/(2\pi) =$ $1/(2t_{\mu}) = 5$ kHz from resonance and absorption-imaged the atoms transferred to $|F = 2, m_F = 2\rangle$ in-situ ($\approx 10\%$ fractional transfer) leaving $|F = 1, m_F = 1\rangle$ state atoms undisturbed. We used these data to obtain the transferred atom number N_+ . Then after a ≈ 34 ms delay, we repeated the processes with $\delta \rightarrow -\delta$, giving N_- . The delay between the transfer pulses was selected to be an integer multiple of the $T_{\text{line}} = (60 \text{ Hz})^{-1} \approx 17 \text{ ms line}$ period.

The fractional imbalance between the transferred numbers

$$\varepsilon = \frac{N_+ - N_-}{N_+ + N_-} \approx -4t_\mu \frac{\Delta_\mu}{2\pi},\tag{6}$$

provides an error signal that can be related to any overall shift in detuning Δ_{μ} (see Supplementary Figure 2 in Supplementary Note 4). For example $\epsilon = 0.5$ corresponds to a detuning of just $\Delta_{\mu}/2\pi \approx 1.25$ kHz.

We employed a two step procedure to minimize the impact of field noise during interferometry experiments. First, prior to any measurement sequence we optimized the bias field to minimize δB . Second, we post-selected data to exclude cases with $|\epsilon| > 0.5$; this value was determined empirically to retain most of the data while notably removing outliers in measured contrast.

B. Lattice pulse sequence

An intuitive picture of our scheme for mitigating the effect of the optical lattice begins with a three-state truncation[30, 31] of the full lattice Hamiltonian

$$\frac{\hat{H}(k)}{E_0} = \begin{pmatrix} (k+2k_0)^2 & s/4 & 0\\ s/4 & k^2 & s/4\\ 0 & s/4 & (k-2k_0)^2 \end{pmatrix}, \quad (7)$$

describing a lattice of depth sE_0 , with single photon recoil momentum $\hbar k_0 = 2\pi\hbar/\lambda$, energy $E_0 = \hbar^2 k_0^2/(2m)$, and time $T_0 = 2\pi\hbar/E_0 \approx 265 \ \mu$ s. For atoms initially at rest, i.e. k = 0, this is a resonant lambda coupling scheme with bright state subspace spanned by $|b_0\rangle = |k = 0\rangle$ and $|b_1\rangle = (|k = +2k_0\rangle + |k = -2k_0\rangle)/\sqrt{2}$ and an uncoupled dark state $|d\rangle = (|k = +2k_0\rangle - |k = -2k_0\rangle)/\sqrt{2}$. Since our initial state $|k=0\rangle$ is in the bright state manifold, we focus on the bright state Hamiltonian

$$\frac{\hat{H}_{\rm b}(0)}{E_0} = \begin{pmatrix} 0 & s/(2\sqrt{2}) \\ s/(2\sqrt{2}) & 4 \end{pmatrix} \\ = 2\hat{I} + \frac{1}{2} \left[4\hat{\sigma}_z + \frac{s}{\sqrt{2}}\hat{\sigma}_x \right].$$
(8)

When the lattice is off, this Hamiltonian describes Larmour precession around \mathbf{e}_z with Rabi frequency $4E_0/\hbar$ and when the lattice is on it describes precession about $4\mathbf{e}_z + [s/\sqrt{2}]\mathbf{e}_x$ with Rabi frequency $\sqrt{16 + s^2/2}E_0/\hbar$. In the limit $s \ll 4\sqrt{2}$, the axis of rotation is tipped by $\theta = 4s/\sqrt{2}$, and the Rabi frequency is nearly unchanged from $4E_0/\hbar$. In Supplementary Note 3 Supplementary Figure 1a plots the top of the Bloch sphere with two example orbits in this limit (dashed lines), both for zero (red) and non-zero s (blue).

The solid curves in Supplementary Figure 1a show the trajectory for a two pulse sequence that also returns to the origin. In the small s limit, the condition to return to the initial state is $t_d/T_0 = 1/8 - t_p/T_0$, where t_d is the delay time between pulses and t_p is the pulse duration. Supplementary Figure 1b plots the probability that the final state returns to k = 0 for a shallow lattice with s = 1 (computed using 7 momentum states). The red line indicates the predicted minimum which is in good agreement with the numerically evaluated optimum configuration.

Supplementary Figure 1c plots the same quantity, now with s = 10, showing the narrow range of parameters for which our scheme is expected to be successful. For most parameters, the large s simulation is qualitatively different from the small s results, with the exception of very short pulse times and the region following our scheme. In practice we selected $t_{\rm d} = T_0/10 = 26.5 \ \mu {\rm s}$ and $t_{\rm p} = T_0/32 = 8.2 \ \mu {\rm s}$, marked by the red star in Supplementary Figure 1c.

C. Conventional parameters

Here we outline the relationships between conventional experimental parameters and the relatively abstract quantities employed in deriving the coupling strength $g_{\sigma}(\mathbf{k}_{\perp})$ in Eq. (3).

We start with the coherent state amplitude α_0 and relate it to the optical intensity

$$I = \frac{1}{2} \epsilon_0 c |E|^2 = \hbar \omega_{\rm ge} c |\alpha_0|^2.$$
(9)

In the second statement we inserted the expression

$$|E|^2 = \frac{2\hbar\omega_{\rm ge}|\alpha_0|^2}{\epsilon_0} \tag{10}$$

for the magnitude of the electric field. The saturation intensity is a key metric of the light-matter interaction; for arbitrary light polarization

$$I_{\text{sat}} = \frac{\epsilon_0 c \Gamma^2 \hbar^2}{4 |\boldsymbol{\epsilon}_{\sigma_0}(\mathbf{k}_0) \cdot \mathbf{d}_{\text{ge}}|^2}.$$
 (11)

As detailed in the main text, \mathbf{d}_{ge} is the dipole matrix element for transitions between the ground and excited state with energy difference $\hbar \omega_{ge}$ and where $\epsilon_{\sigma}(\mathbf{k}_0)$ are the polarization vectors as pairs of orthogonal vectors transverse to \mathbf{k}_0 . In terms of these parameters the transition linewidth is

$$\Gamma = \left(\frac{|\mathbf{k}_0|^3}{3\pi\hbar\epsilon_0}\right)|\mathbf{d}_{ge}|^2.$$
 (12)

We recall the standard definition for saturation intensity

$$\frac{I}{I_{\text{sat}}} = 2 \left| \frac{\Omega}{\Gamma} \right|^2, \tag{13}$$

acquired from a more traditional treatment, where

$$\Omega = |\boldsymbol{\epsilon}_{\sigma_0}(\mathbf{k}_0) \cdot \mathbf{d}_{ge}| \frac{E_0}{\hbar}$$
(14)

is the Rabi frequency. We next express Ω in terms of the coupling strength and the optical field amplitude α_0 giving

$$\Omega = 2|g_{\sigma_0}(\mathbf{k}_0)\alpha_0|. \tag{15}$$

These relations allow us to bridge between conventional laboratory parameters and those employed in our model. For example, we combine Eq. (9) and Eq. (11), to obtain

$$\bar{I} = \frac{I}{I_{\text{sat}}} = \frac{8|\alpha_{\sigma_0}g_{\sigma_0}(\mathbf{k}_0)|^2}{\Gamma^2}$$
(16)

- C. L. Degen, F. Reinhard, and P. Cappellaro, Quantum sensing, Rev. Mod. Phys. 89, 035002 (2017).
- [2] K. Hammerer, A. S. Sørensen, and E. S. Polzik, Quantum interface between light and atomic ensembles, Rev. Mod. Phys. 82, 1041 (2010).
- [3] B. M. Terhal, Quantum error correction for quantum memories, Rev. Mod. Phys. 87, 307 (2015).
- [4] H. Krauter, C. A. Muschik, K. Jensen, W. Wasilewski, J. M. Petersen, J. I. Cirac, and E. S. Polzik, Entanglement generated by dissipation and steady state entanglement of two macroscopic objects, Phys. Rev. Lett. 107, 080503 (2011).
- [5] M. J. Gullans and D. A. Huse, Dynamical purification phase transition induced by quantum measurements, Phys. Rev. X 10, 041020 (2020).
- [6] C. Noel, P. Niroula, D. Zhu, A. Risinger, L. Egan, D. Biswas, M. Cetina, A. V. Gorshkov, M. J. Gullans, D. A. Huse, and C. Monroe, Measurement-induced quantum phases realized in a trapped-ion quantum computer, Nature Physics 18, 760 (2022).

in agreement with Eq. (13) and Eq. (15).

We now turn to the scattering probability $P_{\rm sp} = \Gamma t_{\rm m} P_{\rm e}$, which contains the excited state probability

$$P_{\rm e} = \frac{|\alpha_0 g_\sigma(\mathbf{k}_0)|^2}{\Delta^2}.$$
 (17)

This, along with Eq. (16), allows us to rewrite the scattering probability as

$$P_{\rm sp} = |\alpha_0 g_{\sigma_0}(\mathbf{k}_0)|^2 \frac{\Gamma t_{\rm m}}{\Delta^2} = \frac{\Gamma t_{\rm m}}{8} \frac{I}{I_{\rm sat}} \frac{\Gamma^2}{\Delta^2}.$$
 (18)

This expression is organized into the physically relevant dimensionless quantities $\bar{t}_{\rm m}$, \bar{I} and $\bar{\delta}$ introduced in the main text. In the main text, we defined the overall measurement strength using these parameters and made the choice to not include the factor of 8 so

$$g^2 = \frac{I}{I_{\text{sat}}} \frac{\Gamma t_{\text{m}}}{(\Delta/\Gamma)^2} = 8P_{\text{sp}}.$$
 (19)

As such, a measurement strength of $g^2 = 1$ signifies a probability of 1/8 for an atom to scatter a single photon at large angle.

ACKNOWLEDGMENTS

The authors thank J. V. Porto and M. Gullans for carefully reading the manuscript. This work was partially supported by the National Institute of Standards and Technology, and the National Science Foundation through the Physics Frontier Center at the Joint Quantum Institute (PHY-1430094) and the Quantum Leap Challenge Institute for Robust Quantum Simulation (OMA-2120757).

- [7] M. Block, Y. Bao, S. Choi, E. Altman, and N. Y. Yao, Measurement-induced transition in long-range interacting quantum circuits, Phys. Rev. Lett. **128**, 010604 (2022).
- [8] I. Bloch, J. Dalibard, and W. Zwerger, Many-body physics with ultracold gases, Rev. Mod. Phys. 80, 885 (2008).
- [9] M. Müller, S. Diehl, G. Pupillo, and P. Zoller, Engineered open systems and quantum simulations with atoms and ions, in *Advances in Atomic, Molecular, and Optical Physics*, Vol. 61, edited by P. Berman, E. Arimondo, and C. Lin (Academic Press, 2012) pp. 1–80.
- [10] H. J. Carmichael, Quantum trajectory theory for cascaded open systems, Phys. Rev. Lett. 70, 2273 (1993).
- [11] K. Mølmer, Y. Castin, and J. Dalibard, Monte carlo wave-function method in quantum optics, J. Opt. Soc. Am. B 10, 524 (1993).
- [12] A. B. Deb and N. Kjærgaard, Observation of pauli blocking in light scattering from quantum degenerate fermions, Science 374, 972 (2021).

- [13] C. Sanner, L. Sonderhouse, R. B. Hutson, L. Yan, W. R. Milner, and J. Ye, Pauli blocking of atom-light scattering, Science **374**, 979 (2021).
- [14] Y. Margalit, Y.-K. Lu, and W. Top, Furkan Çağrı and Ketterle, Pauli blocking of light scattering in degenerate fermions, Science **374**, 976 (2021).
- [15] Y.-K. Lu, Y. Margalit, and W. Ketterle, Bosonic stimulation of atom–light scattering in an ultracold gas, Nature Physics 19, 210 (2023).
- [16] M. R. Andrews, M.-O. Mewes, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, Direct, nondestructive observation of a bose condensate, Science 273, 84 (1996).
- [17] J. M. Higbie, L. E. Sadler, S. Inouye, A. P. Chikkatur, S. R. Leslie, K. L. Moore, V. Savalli, and D. M. Stamper-Kurn, Direct nondestructive imaging of magnetization in a spin-1 bose-einstein gas, Phys. Rev. Lett. **95**, 050401 (2005).
- [18] A. Ramanathan, S. R. Muniz, K. C. Wright, R. P. Anderson, W. D. Phillips, K. Helmerson, and G. K. Campbell, Partial-transfer absorption imaging: A versatile technique for optimal imaging of ultracold gases, Review of Scientific Instruments 83, 083119 (2012).
- [19] D. V. Freilich, D. M. Bianchi, A. M. Kaufman, T. K. Langin, and D. S. Hall, Real-time dynamics of single vortex lines and vortex dipoles in a bose-einstein condensate, Science **329**, 1182 (2010).
- [20] M. Gajdacz, P. L. Pedersen, T. Mørch, A. J. Hilliard, J. Arlt, and J. F. Sherson, Non-destructive Faraday imaging of dynamically controlled ultracold atoms, Review of Scientific Instruments 84, 83105 (2013).
- [21] E. Altuntaş and I. B. Spielman, Self-bayesian aberration removal via constraints for ultracold atom microscopy, Phys. Rev. Research 3, 043087 (2021).
- [22] K. W. Murch, K. L. Moore, S. Gupta, and D. M. Stamper-Kurn, Observation of quantum-measurement backaction with an ultracold atomic gas, Nature Physics 4, 561 (2008).
- [23] M. Born and E. Wolf, Principles of Optics: Electromagnetic Theory of Propagation, Interference and Diffraction of Light (7th Edition), 7th ed. (Cambridge University Press, 1999).
- [24] J. Appel, P. J. Windpassinger, D. Oblak, U. B. Hoff, N. Kjærgaard, and E. S. Polzik, Mesoscopic atomic entanglement for precision measurements beyond the standard quantum limit, Proceedings of the National Academy of Sciences **106**, 10960 (2009).
- [25] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, Theory of bose-einstein condensation in trapped gases, Rev. Mod. Phys. **71**, 463 (1999).
- [26] Y. Castin and R. Dum, Bose-einstein condensates in time dependent traps, Phys. Rev. Lett. 77, 5315 (1996).
- [27] G. Reinaudi, T. Lahaye, Z. Wang, and D. Guéry-Odelin, Strong saturation absorption imaging of dense clouds of ultracold atoms, Opt. Lett. **32**, 3143 (2007).
- [28] K. Hueck, N. Luick, L. Sobirey, J. Siegl, T. Lompe, H. Moritz, L. W. Clark, and C. Chin, Calibrating high intensity absorption imaging of ultracold atoms, Opt. Express 25, 8670 (2017).
- [29] E. Altuntaş and I. B. Spielman, Direct calibration of laser intensity via ramsey interferometry for cold atom imaging, arXiv preprint arXiv:2304.00656 (2023).
- [30] S. Wu, Y.-J. Wang, Q. Diot, and M. Prentiss, Splitting matter waves using an optimized standing-wave light-

pulse sequence, Phys. Rev. A 71, 43602 (2005).

- [31] C. D. Herold, V. D. Vaidya, X. Li, S. L. Rolston, J. V. Porto, and M. S. Safronova, Precision measurement of transition matrix elements via light shift cancellation, Phys. Rev. Lett. **109**, 243003 (2012).
- [32] A. Fuhrmanek, R. Bourgain, Y. R. P. Sortais, and A. Browaeys, Light-assisted collisions between a few cold atoms in a microscopic dipole trap, Phys. Rev. A 85, 062708 (2012).
- [33] Y. H. Fung, A. V. Carpentier, P. Sompet, and M. Andersen, Two-atom collisions and the loading of atoms in microtraps, Entropy 16, 582 (2014).
- [34] N. S. Kampel, A. Griesmaier, M. P. H. Steenstrup, F. Kaminski, E. S. Polzik, and J. H. Müller, Effect of light assisted collisions on matter wave coherence in superradiant bose-einstein condensates, Phys. Rev. Lett. 108, 090401 (2012).
- [35] A. Urvoy, Z. Vendeiro, J. Ramette, A. Adiyatullin, and V. Vuletić, Direct laser cooling to bose-einstein condensation in a dipole trap, Phys. Rev. Lett. **122**, 203202 (2019).
- [36] W. Ketterle and M. W. Zwierlein, Making, probing and understanding ultracold fermi gases, Rivista del Nuovo Cimento **31**, 247 (2008).
- [37] M. Inguscio, S. Stringari, and C. E. Wieman, eds., Bose-Einstein condensation in atomic gases, proceedings of the international school of physics "Enrico Fermi", course CXL (IOS Press, 1999) Chap. Making, probing and understanding Bose-Einstein condensates, pp. 67–176.
- [38] B. P. Anderson, P. C. Haljan, C. A. Regal, D. L. Feder, L. A. Collins, C. W. Clark, and E. A. Cornell, Watching dark solitons decay into vortex rings in a bose-einstein condensate, Phys. Rev. Lett. 86, 2926 (2001).
- [39] T. Y. Ivanova and D. A. Ivanov, Quantum limits of feedback cooling in optical lattices, Journal of Experimental and Theoretical Physics Letters 82, 482 (2005).
- [40] M. Koch, C. Sames, A. Kubanek, M. Apel, M. Balbach, A. Ourjoumtsev, P. W. H. Pinkse, and G. Rempe, Feedback cooling of a single neutral atom, Phys. Rev. Lett. 105, 173003 (2010).
- [41] N. Behbood, G. Colangelo, F. Martin Ciurana, M. Napolitano, R. J. Sewell, and M. W. Mitchell, Feedback cooling of an atomic spin ensemble, Phys. Rev. Lett. 111, 103601 (2013).
- [42] S. Diehl, A. Tomadin, A. Micheli, R. Fazio, and P. Zoller, Dynamical phase transitions and instabilities in open atomic many-body systems, Phys. Rev. Lett. 105, 015702 (2010).
- [43] J. T. Young, A. V. Gorshkov, and I. B. Spielman, Feedback-stabilized dynamical steady states in the bosehubbard model, Phys. Rev. Research 3, 043075 (2021).
- [44] L. Walker, Measurement and control of atomic and nanomechanical systems for quantum technologies, Ph.D. thesis, University of Strathclyde (2020).
- [45] A. C. J. Wade, J. F. Sherson, and K. Mølmer, Squeezing and entanglement of density oscillations in a boseeinstein condensate, Phys. Rev. Lett. 115, 060401 (2015).
- [46] C. M. Caves and G. J. Milburn, Quantum-mechanical model for continuous position measurements, Phys. Rev. A 36, 5543 (1987).
- [47] T. A. Brun, A simple model of quantum trajectories, American Journal of Physics 70, 719 (2002).
- [48] L. J. LeBlanc, M. C. Beeler, K. Jiménez-García, A. R. Perry, S. Sugawa, R. A. Williams, and I. B. Spielman,

Supplementary Information Quantum Back-action Limits in Dispersively Measured Bose-Einstein Condensates

Supplementary Note 1. Measurement strength and signal to noise

In this section we briefly consider the relation of the measurement strength g to signal to noise ratio (SNR) for the determination of atom number using either forward-directed collectively scattered light or spontaneously scattered light. For standard imaging, directly detecting the collectively scattered light would typically be performed using dark-field imaging (which in principle has the same SNR as phase-contrast imaging), and detecting the spontaneously scattered light would be realized via fluorescence imaging. In principle there is no hard distinction between these techniques since large numerical aperture imaging systems can capture significant contributions from both. For practical experiments with degenerate gases the forward scattered light dominates.

For a Bose-Einstein condensate (BEC), and for simplicity assuming an isotropic (rather than dipole) scattering distribution, the scattering probabilities are

$$P_{\rm tot} = P_{\rm col} + P_{\rm sp} \propto N^2 G + \frac{k_0^2 N}{\pi},\tag{S1}$$

where

$$G = \oint \frac{d^2 \mathbf{k}_{\perp}}{(2\pi)^2} |n_{\mathcal{F}}(\mathbf{k}_0 - \mathbf{k}_{\perp})|^2 \sim \frac{1}{A}$$
(S2)

is a geometric factor and A is the cross-sectional area of the system normal to the imaging axis (in detail this depends on the exact density distribution). Thus the ratio of collective to spontaneous scattering is $\sim \pi N/(k_0^2 A)$ and the total number of spontaneously and collectively scattered photons is

$$N_{\rm sp} = N \frac{g^2}{8},$$
 and $N_{\rm col} = \frac{\pi N^2 G}{k_0^2} \frac{g^2}{8}.$ (S3)

For our system with $N \approx 7 \times 10^4$ and transverse Thomas-Fermi radii $(R_x, R_y) \approx (43, 4) \ \mu m$ the ratio of these is ≈ 6 .

The integrated number of spontaneously scattered photons gives an SNR of $g\sqrt{N/8}$ for the determination of the atom number N. For example, for our BEC and g = 0.3 about $N_{\rm sp} = 800$ photons would be spontaneously scattered. This implies that the forward scattered light would contain about $N_{\rm col} = 5 \times 10^3$ photons.

In both cases the uncertainty in the photon number is simply the square root of the photon number. The SNR in the atom number derived from spontaneously scattered light is therefore ≈ 30 since the atom number is proportional to $N_{\rm sp}$. For the forward scattered light, the atom number is proportional to $N_{\rm col}^{1/2}$ giving a SNR of ≈ 140 . In both cases the SNR is proportional to g.

Supplementary Note 2. Ramsey interferometry with measurement induced decoherence

We consider two-states $\{|g_1\rangle \equiv |\downarrow\rangle, |g_2\rangle \equiv |\uparrow\rangle\}$ that are microwave coupled with the rotating wave Hamiltonian

$$\hat{H}'(\phi_P) = \frac{1}{2} \left[\Delta \hat{\sigma}_z + \Omega_\mu \sin(\phi_P) \hat{\sigma}_x - \Omega_\mu \cos(\phi_P) \hat{\sigma}_y \right],$$

in terms of the detuning Δ (in this case $\Delta > 0$ corresponds to red detuning and $\Delta < 0$ yields blue detuning, this is reversed with respect to the spectroscopy convention), coupling strength Ω_{μ} and microwave oscillator phase ϕ_{P} .

A. Standard Ramsey interferometer

We first outline the basic framework describing Ramsey interferometry (RI) for an arbitrary many-body state. Consider an atomic ensemble in an initial state $|\Psi_i\rangle$ along $-\mathbf{e}_z$ on the Bloch sphere (with no population in $|\uparrow\rangle$), i.e., $\hat{b}_{\uparrow} |\Psi_i\rangle = 0$. In the second quantized notation, the operator

$$\hat{R}_z(\theta) \equiv \exp\left[i\frac{\theta}{2}\hat{b}_i^{\dagger}\sigma_{z,ij}\hat{b}_j\right] = \exp\left[i\frac{\theta}{2}\left(\hat{b}_{\uparrow}^{\dagger}\hat{b}_{\uparrow} - \hat{b}_{\downarrow}^{\dagger}\hat{b}_{\downarrow}\right)\right],$$

implements a spin-rotation about \mathbf{e}_z by the angle θ , and similarly for the other two axes (note that we used an implied summation convention in this expression). Here $\sigma_{x,y,z}$ is the Pauli matrix for the specified axis. Our experimental sequence began with a $\pi/2$ pulse to rotate the system into the equal superposition state $(|\uparrow\rangle + |\downarrow\rangle)/\sqrt{2}$, aligned along \mathbf{e}_x on the Bloch sphere.

The RI completed with a second $\pi/2$ pulse, that drove rotations about the $\mathbf{e}_x \sin \delta \phi_{\rm P} - \mathbf{e}_y \cos \delta \phi_{\rm P}$ axis (implemented by phase shifting the microwave oscillator by $\delta \phi_{\rm P}$). The final state is

$$|\Psi_f\rangle = \left[\hat{R}_z^{\dagger}(\delta\phi_{\rm P})\hat{R}_y(\pi/2)\hat{R}_z(\delta\phi_{\rm P})\right]\hat{R}_y(\pi/2)\left|\Psi_i\right\rangle,\tag{S4}$$

where the quantity in square brackets implements the rotation about the new axis. In this standard RI scheme, for the initial state with all atoms in $|\downarrow\rangle$ the number of atoms in state $|\uparrow\rangle$ in the final state is

$$\langle \hat{n}_{\uparrow} \rangle = \langle \Psi_f | \, \hat{b}_{\uparrow}^{\dagger} \hat{b}_{\uparrow} \, | \Psi_f \rangle = N \cos^2(\delta \phi_{\rm P}/2). \tag{S5}$$

Here, the total number of atoms N in the initial state is conserved in the RI process.

B. Ramsey interferometer with a weak measurement

In our experimental sequence with the dispersive-measurement pulse, the above classic RI is augmented with further evolution from the Kraus operator $\hat{M}(\mathbf{k}_{f})$ [see Eq. (4) in the main text] describing the measurement. The final state is therefore

$$|\Psi_f\rangle = \hat{R}_z^{\dagger}(\delta\phi_{\rm P})\hat{R}_y(\pi/2)\hat{R}_z(\delta\phi_{\rm P})\hat{M}(\mathbf{k}_{\rm f})\hat{R}_y(\pi/2)|\Psi_i\rangle$$

Our discussion for conventional RI was completely agnostic regarding the initial state. This is not the case with the addition of a measurement pulse. We describe our BEC as containing N atoms in the same spatial mode $\psi(\mathbf{x})$ [with Fourier transform $\tilde{\psi}(\mathbf{k})$], and adopt the notation where

$$\hat{b}^{\dagger} = \int d^3 \mathbf{x} \psi(\mathbf{x}) \hat{b}^{\dagger}(\mathbf{x}) = \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \tilde{\psi}(\mathbf{k}) \hat{b}^{\dagger}(\mathbf{k})$$
(S6)

describes the creation of a single particle in that mode (we will add \uparrow / \downarrow subscripts as needed in what follows). Accordingly our initial state is $|N\rangle = (\hat{b}^{\dagger})^{N} |0\rangle / \sqrt{N!}$ and we will make frequent use of the relation $\hat{b}(\mathbf{k}) |N\rangle = \sqrt{N}\tilde{\psi}(\mathbf{k}) |N-1\rangle$.

An alternate description in terms of coherent states is possible but makes non-physical predictions. In this description the initial state is an eigenstate of the annihilation operator with $\hat{b} |\beta\rangle = \beta |\beta\rangle$ where $N \rightarrow |\beta|^2$ is the average atom number. In the measurement problem, this has the implication that each scattering processes creates a recoiling atom, *but* the number of particles in the condensate mode does not decrease since the action of \hat{b} leaves $|\beta\rangle$ unchanged.

We are interested in both the probability of detecting scattered photons and the change in RI contrast. As before, we assume all atoms are initially in $|\downarrow\rangle$ number state. We begin with the scattering probability $P(\mathbf{k}_{f})$ giving

$$P(\mathbf{k}_{\perp}) = \langle N | \hat{R}_{y}^{\dagger}(\pi/2) \hat{R}_{z}^{\dagger}(\delta\phi_{\mathrm{P}}) \hat{M}^{\dagger}(\mathbf{k}_{\perp}) R_{y}^{\dagger}(\pi/2) \hat{R}_{y}(\pi/2) \hat{M}(\mathbf{k}_{\perp}) \hat{R}_{z}(\delta\phi_{\mathrm{P}}) \hat{R}_{y}(\pi/2) | N \rangle$$

$$= \frac{t_{\mathrm{m}} P_{\mathrm{e}}}{c} |g(\mathbf{k}_{\perp})|^{2} \frac{N}{2} \left[\frac{N-1}{2} \left| \int \frac{d^{3} \mathbf{k}_{1}}{(2\pi)^{3}} \tilde{\psi}_{\downarrow}^{*}(\mathbf{k}_{1}) \tilde{\psi}_{\downarrow} [\mathbf{k}_{1} - (\mathbf{k}_{\perp} - \mathbf{k}_{0})] \right|^{2} + 1 \right].$$
(S7)

This reproduces Eq. (5) in the main manuscript but with atom number N essentially reduced by 1/2 (owing to the $\pi/2$ pulse prior to applying the measurement pulse). As elaborated on in the main text, the first term describes collective (stimulated) scattering with integrated probability $P_{\rm col} \propto N^2$ whereas the second term stems from spontaneous emission with probability $P_{\rm sp} \propto N$.

We are interested in the mean number of atoms in momentum state \mathbf{k} and internal state $|\uparrow\rangle$, conditioned on detecting a photon in state \mathbf{k}_{\perp} :

$$\langle \hat{n}_{\uparrow}(\mathbf{k}) \rangle_{|\mathbf{k}_{\perp}} = \frac{1}{P(\mathbf{k}_{\perp})} \langle N | \hat{R}_{y}^{\dagger}(\pi/2) \hat{R}_{z}^{\dagger}(\delta\phi_{\mathrm{P}}) \hat{M}^{\dagger}(\mathbf{k}_{\perp}) R_{y}^{\dagger}(\pi/2) \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}) \hat{b}_{\uparrow}(\mathbf{k}) \hat{R}_{y}(\pi/2) \hat{M}(\mathbf{k}_{\perp}) \hat{R}_{z}(\delta\phi_{\mathrm{P}}) \hat{R}_{y}(\pi/2) | N \rangle$$

$$= \frac{1}{2P(\mathbf{k}_{\perp})} \langle N | \hat{R}_{y}^{\dagger}(\pi/2) \hat{R}_{z}^{\dagger}(\delta\phi_{\mathrm{P}}) \hat{M}^{\dagger}(\mathbf{k}_{\perp}) \left[\hat{b}_{\uparrow}^{\dagger}(\mathbf{k}) + \hat{b}_{\downarrow}^{\dagger}(\mathbf{k}) \right] \left[\hat{b}_{\uparrow}(\mathbf{k}) + \hat{b}_{\downarrow}(\mathbf{k}) \right] \hat{M}(\mathbf{k}_{\perp}) \hat{R}_{z}(\delta\phi_{\mathrm{P}}) \hat{R}_{y}(\pi/2) | N \rangle$$

$$\propto \frac{1}{2} \int \frac{d^{3}\mathbf{k}_{1}}{(2\pi)^{3}} \frac{d^{3}\mathbf{k}_{2}}{(2\pi)^{3}} \langle N | \hat{R}_{y}^{\dagger}(\pi/2) \left\{ \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}_{1}) \hat{b}_{\uparrow}[\mathbf{k}_{1} - (\mathbf{k}_{\perp} - \mathbf{k}_{0})] \left[e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}) + e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\downarrow}^{\dagger}(\mathbf{k}) \right]$$

$$\times \left[e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}(\mathbf{k}) + e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\downarrow}(\mathbf{k}) \right] \hat{b}_{\uparrow}^{\dagger}[\mathbf{k}_{2} - (\mathbf{k}_{\perp} - \mathbf{k}_{0})] \hat{b}_{\uparrow}(\mathbf{k}_{2}) \right\} \hat{R}_{y}(\pi/2) | N \rangle .$$

$$(S8)$$

In the last line, we omitted the numerical prefactor for brevity, but will reinstate it at the end of our computation. In the next step we normal order the field operators (with all creation operators moved to the left and the annihilation operators to the right), by performing commutators term-by-term on the chain of field operators inside the curly braces in Eq. (S8) (without the leading and trailing rotation operators). This procedure yields an expression

$$\begin{split} &= \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}_{1}) \left[e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}) + e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\downarrow}^{\dagger}(\mathbf{k}) \right] \hat{b}_{\uparrow}^{\dagger}[\mathbf{k}_{2} - (\mathbf{k}_{\perp} - \mathbf{k}_{0})] \\ &\times \hat{b}_{\uparrow}[\mathbf{k}_{1} - (\mathbf{k}_{\perp} - \mathbf{k}_{0})] \left[e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}(\mathbf{k}) + e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\downarrow}(\mathbf{k}) \right] \hat{b}_{\uparrow}(\mathbf{k}_{2}) \\ &+ \delta(\mathbf{k}_{1} - \mathbf{k}_{2}) \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}_{1}) \left[e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}) + e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\downarrow}^{\dagger}(\mathbf{k}) \right] \left[e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}(\mathbf{k}) + e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\downarrow}(\mathbf{k}) \right] \hat{b}_{\uparrow}(\mathbf{k}_{2}) \\ &+ \delta(\mathbf{k}_{1} - (\mathbf{k}_{\perp} - \mathbf{k}_{0}) - \mathbf{k}) e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}_{1}) \hat{b}_{\uparrow}^{\dagger}[\mathbf{k}_{2} - (\mathbf{k}_{\perp} - \mathbf{k}_{0})] \left[e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}(\mathbf{k}) + e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\downarrow}(\mathbf{k}) \right] \hat{b}_{\uparrow}(\mathbf{k}_{2}) \\ &+ \delta(\mathbf{k}_{2} - (\mathbf{k}_{\perp} - \mathbf{k}_{0}) - \mathbf{k}) e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}_{1}) \left[e^{-i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}) + e^{i\delta\phi_{\mathrm{P}}/2} \hat{b}_{\downarrow}^{\dagger}(\mathbf{k}) \right] \hat{b}_{\uparrow}[\mathbf{k}_{1} - (\mathbf{k}_{\perp} - \mathbf{k}_{0})] \hat{b}_{\uparrow}(\mathbf{k}_{2}) \\ &+ \delta(\mathbf{k}_{1} - (\mathbf{k}_{\perp} - \mathbf{k}_{0}) - \mathbf{k}) \delta(\mathbf{k}_{2} - (\mathbf{k}_{\perp} - \mathbf{k}_{0}) - \mathbf{k}) \hat{b}_{\uparrow}^{\dagger}(\mathbf{k}_{1}) \hat{b}_{\uparrow}(\mathbf{k}_{2}), \end{split}$$

with five terms that we label (I) to (V) from top to bottom. We evaluate each term (now accounting for the rotation operators) by keeping only the operators that act on the initial $|\downarrow\rangle$ state. Integrating over all BEC momentum states **k**, we find

$$(I) = \frac{N(N-1)(N-2)}{4} \cos^2\left(\frac{\delta\phi_{\rm P}}{2}\right) \left| \int \frac{d^3\mathbf{k}_1}{(2\pi)^3} \tilde{\psi}^*_{\downarrow}(\mathbf{k}_1) \tilde{\psi}_{\downarrow}[\mathbf{k}_1 - (\mathbf{k}_{\perp} - \mathbf{k}_0)] \right|^2,$$
(S9)

$$(\mathrm{II}) = \frac{N(N-1)}{2} \cos^2\left(\frac{\delta\phi_{\mathrm{P}}}{2}\right),\tag{S10}$$

$$(\text{III}) + (\text{IV}) = \frac{N(N-1)}{2} \cos^2\left(\frac{\delta\phi_{\text{P}}}{2}\right) \left| \int \frac{d^3\mathbf{k}_2}{(2\pi)^3} \tilde{\psi}^*_{\downarrow} [\mathbf{k}_2 - (\mathbf{k}_{\perp} - \mathbf{k}_0)] \tilde{\psi}_{\downarrow}(\mathbf{k}_2) \right|^2, \tag{S11}$$

$$(\mathbf{V}) = \frac{N}{4}.\tag{S12}$$

Together terms (I), (III) and (IV) describe collective scattering (the usual Born and Wolf [S1] forward scattering from the overall density distribution, as previously noted in the main manuscript) and combine to give

$$(I) + (III) + (IV) = \frac{N(N-1)N}{4} \cos^2\left(\frac{\delta\phi_{\rm P}}{2}\right) \left| \int \frac{d^3\mathbf{k}_1}{(2\pi)^3} \tilde{\psi}^*_{\downarrow}(\mathbf{k}_1) \tilde{\psi}_{\downarrow}[\mathbf{k}_1 - (\mathbf{k}_{\perp} - \mathbf{k}_0)] \right|^2.$$
(S13)

On the other hand, terms (II) and (V) give

(II) + (V) =
$$\frac{N}{2} \left[(N-1)\cos^2\left(\frac{\delta\phi_{\rm P}}{2}\right) + \frac{1}{2} \right],$$
 (S14)

describing independent scattering, i.e., spontaneous emission. These two terms convey that the contrast will be reduced by the fractional count of an atom, i.e. 1/N, for each spontaneous scattering event.

Finally, we combine all five terms, re-insert the prefactor, integrate over \mathbf{k}_{\perp} and organize into contributions from single atom scattering $P_{\rm sp}$ and collective scattering $P_{\rm col}$. This elucidates the distinction between large angle and small

$$\langle \hat{N}_{\uparrow}(\delta\phi_{\rm P}) \rangle = \frac{P_{\rm col}}{P_{\rm tot}} N \cos^2\left(\frac{\delta\phi_{\rm P}}{2}\right) + \frac{P_{\rm sp}}{P_{\rm tot}} \left[(N-1)\cos^2\left(\frac{\delta\phi_{\rm P}}{2}\right) + \frac{1}{2} \right],\tag{S15}$$

with total scattering probability P_{tot} . In the first term, small angle collective scattering returns atoms into their initial spatial mode (Mössbauer scattering); thereby leaving the RI contrast unchanged. In the second term, larger angle scattering sends the recoiling atom into a previously empty mode. In this process the environment counts the scattered atom to be in $|\uparrow\rangle$ and removes its contribution to the contrast, as motivated in the main manuscript.

Supplementary Note 3. Lattice pulse sequence



Supplementary Figure 1. Lattice pulse sequence. **a** Top of Bloch sphere for the spherical polar angles ϕ and θ showing trajectories for zero (red) and non-zero (blue) s. The dashed curve shows complete orbits while the solid curve with arrows results from a pulsed sequence that combine to form trajectory that returns to the initial state. **b** and **c** Return probability computed including 7 momentum states for s = 1 and s = 10. The star marks the parameters used in our experiment.

This section shows the diagrams elucidating the lattice pulse sequence used to remedy the effect of the stray weak optical lattice. See the second subsection of the Method section in the main manuscript for a more detailed analysis.

Supplementary Note 4. Magnetic field lock



Supplementary Figure 2. Microwave field lock. **a** Transfer fraction for individual pulses with parameters as described in subsection A of the Method section in the main manuscript. **b** Error function (solid curve) and linear approximation (dashed line).

This section presents the diagrams for the microwave field lock employed to minimize the impact of magnetic field shifts in Ramsey contrast measurements. See the first subsection of the main manuscript Method section for a more detailed description of this procedure. [S1] M. Born and E. Wolf, *Principles of Optics: Electromagnetic Theory of Propagation, Interference and Diffraction of Light* (7th Edition), 7th ed. (Cambridge University Press, 1999).