

Non-SUSY Lepton Flavor Model with 3HDM

Yukimura Izawa ^{1*}, Yusuke Shimizu ^{1,2†}, Hironori Takei ^{1‡}

¹*Physics Program, Graduate School of Advanced Science
and Engineering, Hiroshima University,
Higashi-Hiroshima 739-8526, Japan*

²*Core of Research for the Energetic Universe, Hiroshima University,
Higashi-Hiroshima 739-8526, Japan*

Abstract

We propose a simple non-supersymmetric lepton flavor model with A_4 symmetry. The A_4 group is a minimal one which includes triplet irreducible representation. We introduce three Higgs doublets which are assigned as triplet of the A_4 symmetry. It is natural that there are three generations of the Higgs fields as same as the standard model fermions. We study the potential analysis and we get the vacuum expectation values for the local minimum. In the vacuum expectation values, we obtain the charged lepton, Dirac neutrino, and right-handed Majorana neutrino mass matrices. By using type-I seesaw mechanism, we get the left-handed Majorana neutrino mass matrix. In the NuFIT 5.1 data, we predict the Dirac CP phase and the Majorana phases for the only inverted neutrino mass hierarchy. Especially, the Dirac CP phase and lepton mixing angle θ_{23} are strongly correlated. If the θ_{23} is more precise measured, the Dirac CP phase is more precise predicted, and vice versa. We also predict the effective mass for neutrino-less double beta decay $m_{ee} \simeq 47.1$ [meV] and the lightest neutrino mass $m_3 \simeq 0.789 - 1.43$ [meV]. It is testable for our model in the near future neutrino experiments.

*izawa-yukimura@hiroshima-u.ac.jp

†yu-shimizu@hiroshima-u.ac.jp

‡t-hironori@hiroshima-u.ac.jp

1 Introduction

The standard model (SM) is the successful one with the discovery of the Higgs boson. In the particle physics, the gauge theory is applied and tested by the electroweak precision measurements for the SM. However there are still mysterious puzzle, e.g. the origin of the generations which are differences of the mixing angles and masses for quark and lepton sectors. Actually, the Yukawa couplings are completely free parameters so that the mixing angles and masses cannot be predicted in the SM. In addition, the neutrinos are massless for renormalizable operators. One of the attractive phenomena to solve the puzzle is the neutrino oscillation which provides us the useful information such as three lepton mixing angles and two neutrino mass squared differences which means that the neutrinos have non-zero masses. The T2K and NO ν A experiments have confirmed the neutrino oscillation in the $\nu_\mu \rightarrow \nu_e$ appearance events [1–3], which are one of the clues of the new physics beyond the SM such as the Dirac CP violating phase for the lepton sector by combining the data of the reactor neutrino experiments [4, 5]. The KamLAND-Zen [6, 7], GERDA [8, 9], and CUORE [10, 11] experiments also provide us the significant information which are whether the neutrinos are Dirac or Majorana particles, the lepton number violation, and Majorana phases if the neutrinos are Majorana particles. Thus the neutrino oscillation experiments go into a new phase of the precise determinations of the lepton mixing angles, the neutrino mass squared differences, and the CP violating phases.

The SM particles obey the gauge theory. After spontaneous symmetry breaking (SSB), the gauge bosons and fermions get the masses through the Higgs mechanism. However the Yukawa couplings cannot be controlled by the gauge symmetry, the Yukawa couplings are completely free parameters in the SM. The flavor symmetry can apply to the generations. The Froggatt-Nielsen mechanism which was proposed by C. D. Froggatt and H. B. Nielsen are introduced global $U(1)_{\text{FN}}$ symmetry [12]. Thanks for the $U(1)_{\text{FN}}$ symmetry, it is natural to explain the fermion mass hierarchies. On the other hand, the non-Abelian discrete symmetry (See for the review [13]–[17].) can naturally explain the lepton mixing angles so-called “tri-bimaximal mixing (TBM)” [18, 19] before the reactor experiments reported the non-zero reactor angle θ_{13} [4, 5]. Actually, many authors have studied the breaking or deviation from the TBM [20]–[42] or other patterns of the lepton mixing angles, e.g. tri-bimaximal-Cabibbo mixing [43, 44]. One of the successful flavor models was proposed by G. Altarelli and F. Feruglio [45, 46]. The Altarelli and Feruglio (AF) model show the TBM by using the non-Abelian discrete symmetry A_4 . They introduced two SM gauge singlet scalar fields so-called “flavons” and taking the vacuum expectation value (VEV) alignments of the A_4 triplets as $(1, 0, 0)$ and $(1, 1, 1)$, which naturally explained that the charged lepton is diagonal and neutrino mixing is TBM, respectively. However the corrected VEV alignments cannot be driven from the potential analysis. Then, they applied to the supersymmetry (SUSY) and introduced the driving fields. There are so many scalar fields in addition to the no evidence of the SUSY particles for the accelerator experiments, e.g. Large Hadron Collider experiment.

In this paper, we propose a simple non-SUSY lepton flavor model with A_4 symmetry. The A_4 group is a minimal one which includes triplet irreducible representation. We introduce three Higgs doublets which is assigned as triplet of the A_4 symmetry. It is natural that there are three generations as same as the SM fermions. We study the potential analysis and we get the VEV for the local minimum in the three Higgs doublet model (3HDM) with A_4 symmetry. The left-handed lepton doublets are assigned to triplet and the right-handed charged leptons are assigned

to different singlets of the A_4 symmetry, respectively. We introduce the right-handed Majorana neutrinos which are assigned to triplet of the A_4 symmetry. In our model, the right-handed Majorana neutrino mass matrix has a simple flavor structure. On the other hand, the Dirac neutrino mass matrix has symmetric and anti-symmetric Yukawa couplings for A_4 symmetry. By using the type-I seesaw mechanism [47–51], we obtain the left-handed Majorana neutrino mass matrix. After diagonalizing the charged lepton and left-handed Majorana neutrino mass matrices, we get the lepton mixing matrix which is Pontecorvo-Maki-Nakagawa-Sakata (PMNS) one [52,53]. In the numerical analysis, we use the NuFIT 5.1 data [54,55]. We find that only inverted ordering is acceptable and we cannot find the solutions for normal ordering in the neutrino mass hierarchy. We obtain relevant relations for mixing angles and the effective mass for the neutrino-less double beta ($0\nu\beta\beta$) decay as a function of the lightest neutrino mass. Especially, the Dirac CP phase and lepton mixing angle θ_{23} are strongly correlated. If the neutrinos are Majorana particles, the effective mass for the neutrino-less double beta decay and the lightest neutrino mass are also predicted in our model.

This paper is organized as follows. In Section 2, we briefly explain the potential analysis of the three Higgs doublet model and we apply to the A_4 flavor model. In Section 3, we present the A_4 flavor model and study mass matrices. In Section 4, we show the numerical analysis of our model. Section 5 is devoted to a summary and discussions. In Appendix A, we show the Higgs mass matrices in the general 3HDM. We also show the multiplication rule of the A_4 group in Appendix B.

2 Three Higgs doublet model

In this section, we discuss the 3HDM. First, we briefly introduce the general 3HDM and discuss the potential analysis. Next, we discuss the A_4 potential analysis in the 3HDM, where we assign the three Higgs doublets as the triplet of the A_4 symmetry. It is natural that the Higgs fields are three generations as same as the SM fermions.

2.1 The general 3HDM

At first, we briefly introduce the general 3HDM. The general 3HDM means that we consider the SM Higgs sector extend to the three Higgs doublets, denoted as

$$\phi_1 = \begin{pmatrix} \phi_1^+ \\ \phi_1^0 \end{pmatrix}, \quad \phi_2 = \begin{pmatrix} \phi_2^+ \\ \phi_2^0 \end{pmatrix}, \quad \phi_3 = \begin{pmatrix} \phi_3^+ \\ \phi_3^0 \end{pmatrix}, \quad (1)$$

where the Higgs doublets ϕ_i (with $i = 1, 2, 3$) have the same $U(1)_Y$ hypercharge. One Higgs doublet consists of four real scalar fields. Thus, there are twelve real scalar fields in the 3HDM.

The general 3HDM potential which is invariant under $SU(2)_L \times U(1)_Y$ is written as follows [56]:

$$\begin{aligned}
V = & - \sum_{i,j=1}^3 m_{ij}^2 (\phi_i^\dagger \phi_j) + \frac{1}{2} \sum_{i,j,k,l=1}^3 \lambda_{ijkl} (\phi_i^\dagger \phi_j) (\phi_k^\dagger \phi_l) \quad (2) \\
= & - m_{11}^2 (\phi_1^\dagger \phi_1) - m_{22}^2 (\phi_2^\dagger \phi_2) - m_{33}^2 (\phi_3^\dagger \phi_3) \\
& - [m_{12}^2 (\phi_1^\dagger \phi_2) + m_{13}^2 (\phi_1^\dagger \phi_3) + m_{23}^2 (\phi_2^\dagger \phi_3) + h.c.] \\
& + \lambda_{11} (\phi_1^\dagger \phi_1)^2 + \lambda_{22} (\phi_2^\dagger \phi_2)^2 + \lambda_{33} (\phi_3^\dagger \phi_3)^2 \\
& + \lambda_{1122} (\phi_1^\dagger \phi_1) (\phi_2^\dagger \phi_2) + \lambda_{1133} (\phi_1^\dagger \phi_1) (\phi_3^\dagger \phi_3) + \lambda_{2233} (\phi_2^\dagger \phi_2) (\phi_3^\dagger \phi_3) \\
& + \lambda_{1221} (\phi_1^\dagger \phi_2) (\phi_2^\dagger \phi_1) + \lambda_{1331} (\phi_1^\dagger \phi_3) (\phi_3^\dagger \phi_1) + \lambda_{2332} (\phi_2^\dagger \phi_3) (\phi_3^\dagger \phi_2) \\
& + \left[\frac{\lambda_{1212}}{2} (\phi_1^\dagger \phi_2)^2 + \frac{\lambda_{1313}}{2} (\phi_1^\dagger \phi_3)^2 + \frac{\lambda_{2323}}{2} (\phi_2^\dagger \phi_3)^2 \right. \quad (3) \\
& + \lambda_{1213} (\phi_1^\dagger \phi_2) (\phi_1^\dagger \phi_3) + \lambda_{2113} (\phi_2^\dagger \phi_1) (\phi_1^\dagger \phi_3) + \lambda_{1323} (\phi_1^\dagger \phi_3) (\phi_2^\dagger \phi_3) \\
& + \lambda_{1332} (\phi_1^\dagger \phi_3) (\phi_3^\dagger \phi_2) + \lambda_{2123} (\phi_2^\dagger \phi_1) (\phi_2^\dagger \phi_3) + \lambda_{1223} (\phi_1^\dagger \phi_2) (\phi_2^\dagger \phi_3) \\
& + \lambda_{1112} (\phi_1^\dagger \phi_1) (\phi_1^\dagger \phi_2) + \lambda_{2212} (\phi_2^\dagger \phi_2) (\phi_1^\dagger \phi_2) + \lambda_{1113} (\phi_1^\dagger \phi_1) (\phi_1^\dagger \phi_3) \\
& + \lambda_{1123} (\phi_1^\dagger \phi_1) (\phi_2^\dagger \phi_3) + \lambda_{2213} (\phi_2^\dagger \phi_2) (\phi_1^\dagger \phi_3) + \lambda_{2223} (\phi_2^\dagger \phi_2) (\phi_2^\dagger \phi_3) \\
& \left. + \lambda_{3312} (\phi_3^\dagger \phi_3) (\phi_1^\dagger \phi_2) + \lambda_{3313} (\phi_3^\dagger \phi_3) (\phi_1^\dagger \phi_3) + \lambda_{3323} (\phi_3^\dagger \phi_3) (\phi_2^\dagger \phi_3) + h.c. \right],
\end{aligned}$$

where $\lambda_{ii} = \lambda_{iiii}/2$. As you can see, the $SU(2)_L \times U(1)_Y$ invariant potential has 9 real mass term, m_{ij}^2 and 45 real quartic couplings, λ_{ijkl} . The scalar field VEVs are generally written as follows

$$\langle \phi_i \rangle = \left(\frac{1}{\sqrt{2}} (v_i + \rho_i + i\chi_i) \right), \quad i = 1, 2, 3. \quad (4)$$

Furthermore, the potential satisfies following three minimization conditions:

$$\begin{aligned}
0 = & - 2v_1 m_{11}^2 - 2m_{12}^2 v_2 - 2m_{13}^2 v_3 + 2\lambda_{11} v_1^3 + (\lambda_{1133} + \lambda_{1313} + \lambda_{1331}) v_1 v_3^2 \quad (5) \\
& + (\lambda_{1122} + \lambda_{1212} + \lambda_{1221}) v_1 v_2^2 + 2(\lambda_{1123} + \lambda_{1213} + \lambda_{2113}) v_1 v_2 v_3 \\
& + (\lambda_{1223} + \lambda_{2123} + \lambda_{2213}) v_2^2 v_3 + (\lambda_{1323} + \lambda_{1332} + \lambda_{3312}) v_2 v_3^2 \\
& + \lambda_{2212} v_2^3 + 3\lambda_{1112} v_1^2 v_2 + 3\lambda_{1113} v_1^2 v_3 + \lambda_{3313} v_3^3,
\end{aligned}$$

$$\begin{aligned}
0 = & - 2v_2 m_{22}^2 - 2m_{12}^2 v_1 - 2m_{23}^2 v_3 + 2\lambda_{22} v_2^3 + (\lambda_{1122} + \lambda_{1212} + \lambda_{1221}) v_1^2 v_2 \quad (6) \\
& + (\lambda_{2233} + \lambda_{2323} + \lambda_{2332}) v_2 v_3^2 + 2(\lambda_{2213} + \lambda_{1223} + \lambda_{2123}) v_1 v_2 v_3 \\
& + (\lambda_{1323} + \lambda_{1332} + \lambda_{3312}) v_1 v_3^2 + (\lambda_{1123} + \lambda_{1213} + \lambda_{2113}) v_1^2 v_3 \\
& + \lambda_{1112} v_1^3 + 3\lambda_{2212} v_1 v_2^2 + 3\lambda_{2223} v_2^2 v_3 + \lambda_{3323} v_3^3,
\end{aligned}$$

$$\begin{aligned}
0 = & - 2v_3 m_{33}^2 - 2m_{13}^2 v_1 - 2m_{23}^2 v_2 + 2\lambda_{33} v_3^3 + (\lambda_{1133} + \lambda_{1313} + \lambda_{1331}) v_1^2 v_3 \quad (7) \\
& + (\lambda_{2233} + \lambda_{2323} + \lambda_{2332}) v_2^2 v_3 + 2(\lambda_{1223} + \lambda_{2123} + \lambda_{2213}) v_1^2 v_3 \\
& + (\lambda_{1123} + \lambda_{1213} + \lambda_{2113}) v_1^2 v_2 + 2(\lambda_{1323} + \lambda_{1332} + \lambda_{3312}) v_1 v_2 v_3 \\
& + \lambda_{1113} v_1^3 + \lambda_{2223} v_2^3 + 3\lambda_{3313} v_1 v_3^2 + 3\lambda_{3323} v_2 v_3^2.
\end{aligned}$$

The VEVs v_1, v_2, v_3 of the three Higgs doublets above are given by

$$v_1 = v \cos \beta_1 \cos \beta_2, \quad v_2 = v \sin \beta_1 \cos \beta_2, \quad v_3 = v \sin \beta_2, \quad (8)$$

where $\tan \beta_1 = v_2/v_1$, $\tan \beta_2 = v_3/\sqrt{v_1^2 + v_2^2}$, $v \equiv \sqrt{v_1^2 + v_2^2 + v_3^2}$ and the v is the SM Higgs doublet VEV.

After the SSB, the Z and W^\pm gauge bosons get masses from three Goldstone bosons. Consequently, 3HDM has nine scalar mass eigenstates, three CP-even scalars: $H, h_{1,2}$, two CP-odd scalars: $a_{1,2}$, and four charged scalars: $h_{1,2}^\pm$. See Appendix A for Higgs mass matrices.

2.2 Potential analysis in the 3HDM with A_4 symmetry

Let us discuss the potential analysis in the 3HDM with A_4 symmetry. The A_4 group which is a minimal including triplet irreducible representation is the symmetry group of a tetrahedron or even permutation of four elements. There are twelve elements and four irreducible representations such as three different singlets $\mathbf{1}, \mathbf{1}', \mathbf{1}''$ and triplet $\mathbf{3}$ in the A_4 group, respectively. Also the A_4 can be defined as the group generated by two elements S and T which satisfy the following algebraic relations as

$$S^2 = T^3 = (ST)^3 = \mathbf{1}. \quad (9)$$

These generators are represented by

$$\begin{aligned} \mathbf{1} : \quad & S = 1, \quad T = 1, \\ \mathbf{1}' : \quad & S = 1, \quad T = e^{\frac{4\pi i}{3}} \equiv \omega^2, \\ \mathbf{1}'' : \quad & S = 1, \quad T = e^{\frac{2\pi i}{3}} \equiv \omega, \end{aligned} \quad (10)$$

on the one-dimensional representations. These generators are also represented by

$$\mathbf{3} : \quad S = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}, \quad (11)$$

on the three-dimensional representation. In these bases Eqs. (10) and (11), we can make the character table and obtain the multiplication of the A_4 group. The relevant multiplication rule is shown in Appendix B.

We introduce three Higgs doublets ϕ_1, ϕ_2, ϕ_3 which are assigned as triplet ϕ :

$$\phi = (\phi_1, \phi_2, \phi_3), \quad (12)$$

of the A_4 symmetry. On the other hand, the complex conjugate of the ϕ is considered by the conjugate of the generator T of Eq. (11) as

$$T^* = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}. \quad (13)$$

Then, the complex conjugate of the ϕ is given by

$$\phi^* = (\phi_1^*, \phi_3^*, \phi_2^*), \quad (14)$$

and the multiplication rule of A_4 is kept as Eq. (81) in Appendix B. In this case, the Higgs potential is written as

$$V = -\mu^2\phi^\dagger\phi + \lambda(\phi^\dagger\phi)^2. \quad (15)$$

According to the multiplication rule of A_4 in Appendix B, we obtain

$$\begin{aligned} V = & -\mu^2(|\phi_1|^2 + |\phi_2|^2 + |\phi_3|^2) \\ & + \lambda_1|\phi_1^2 + 2\phi_2\phi_3|^2 + \lambda_2|\phi_2^2 + 2\phi_3\phi_1|^2 + \lambda_3|\phi_3^2 + 2\phi_1\phi_2|^2 \\ & + \lambda_4\left[|\phi_1^2 - \phi_2\phi_3|^2 + |\phi_2^2 - \phi_3\phi_1|^2 + |\phi_3^2 - \phi_1\phi_2|^2\right]. \end{aligned} \quad (16)$$

For the sake of simplicity in the potential analysis, ϕ_1, ϕ_2, ϕ_3 are considered as three Higgs fields. The VEVs of these Higgs fields are as follows:

$$\langle\phi_1\rangle = v_1, \quad \langle\phi_2\rangle = v_2, \quad \langle\phi_3\rangle = v_3, \quad (17)$$

where v_1, v_2, v_3 are real. When the Higgs fields have these values, the potential takes the minimum value, so the following holds:

$$\left(\frac{\partial V}{\partial\phi_1}\right)_{\phi_1=\langle\phi_1\rangle, \phi_2=\langle\phi_2\rangle, \phi_3=\langle\phi_3\rangle} = 0, \quad (18)$$

$$\left(\frac{\partial V}{\partial\phi_2}\right)_{\phi_1=\langle\phi_1\rangle, \phi_2=\langle\phi_2\rangle, \phi_3=\langle\phi_3\rangle} = 0, \quad (19)$$

$$\left(\frac{\partial V}{\partial\phi_3}\right)_{\phi_1=\langle\phi_1\rangle, \phi_2=\langle\phi_2\rangle, \phi_3=\langle\phi_3\rangle} = 0. \quad (20)$$

From the above equations we obtain the following conditions:

$$\begin{aligned} 0 = & -2\mu^2v_1 + 4\lambda_1v_1(v_1^2 + 2v_2v_3) + 4\lambda_2v_3(v_2^2 + 2v_3v_1) + 4\lambda_3v_2(v_3^2 + 2v_1v_2) \\ & + 4\lambda_4\left[v_1(v_1^2 - v_2v_3) - \frac{1}{2}v_2(v_3^2 - v_1v_2) - \frac{1}{2}v_3(v_2^2 - v_3v_1)\right], \end{aligned} \quad (21)$$

$$\begin{aligned} 0 = & -2\mu^2v_2 + 4\lambda_1v_3(v_1^2 + 2v_2v_3) + 4\lambda_2v_2(v_2^2 + 2v_3v_1) + 4\lambda_3v_1(v_3^2 + 2v_1v_2) \\ & + 4\lambda_4\left[v_2(v_2^2 - v_3v_1) - \frac{1}{2}v_1(v_3^2 - v_1v_2) - \frac{1}{2}v_3(v_1^2 - v_2v_3)\right], \end{aligned} \quad (22)$$

$$\begin{aligned} 0 = & -2\mu^2v_3 + 4\lambda_1v_2(v_1^2 + 2v_2v_3) + 4\lambda_2v_1(v_2^2 + 2v_3v_1) + 4\lambda_3v_3(v_3^2 + 2v_1v_2) \\ & + 4\lambda_4\left[v_3(v_3^2 - v_1v_2) - \frac{1}{2}v_1(v_2^2 - v_3v_1) - \frac{1}{2}v_2(v_1^2 - v_2v_3)\right]. \end{aligned} \quad (23)$$

For these conditions, when $\lambda_1 \neq \lambda_4$ and $2\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 \neq 0$ hold, we obtain the following

local solution¹:

$$\langle \phi_1 \rangle = v_1, \tag{24}$$

$$\langle \phi_2 \rangle = v_2 \tag{25}$$

$$\begin{aligned} &= -\frac{\lambda_2 + \lambda_3 - \lambda_4}{2\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4} v_1 \\ &\pm \frac{\sqrt{\{-2\lambda_1^2 + \lambda_2^2 + \lambda_3^2 - (\lambda_1 - 3\lambda_4)(\lambda_2 + \lambda_3) - 3\lambda_1\lambda_4\}v_1^2 + \frac{1}{2}(2\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4)\mu^2}}{2\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4}, \\ \langle \phi_3 \rangle &= v_3 \tag{26} \end{aligned}$$

$$\begin{aligned} &= -\frac{\lambda_2 + \lambda_3 - \lambda_4}{2\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4} v_1 \\ &\pm \frac{\sqrt{\{-2\lambda_1^2 + \lambda_2^2 + \lambda_3^2 - (\lambda_1 - 3\lambda_4)(\lambda_2 + \lambda_3) - 3\lambda_1\lambda_4\}v_1^2 + \frac{1}{2}(2\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4)\mu^2}}{2\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4}. \end{aligned}$$

Expressed in the form of the general 3HDM VEVs, these local VEVs can be written as

$$v_1 = v \cos \beta, \quad v_2 = \frac{v}{\sqrt{2}} \sin \beta, \quad v_3 = \frac{v}{\sqrt{2}} \sin \beta. \tag{27}$$

In the next section, we present our model by using the VEVs in Eq. (27).

3 Lepton flavor model in the A_4 symmetry

In this section, we present a non-SUSY lepton flavor model in the A_4 symmetry. The left-handed lepton doublets are assigned to triplet and the right-handed charged leptons are assigned to different singlets as $\mathbf{1}$, $\mathbf{1}''$, and $\mathbf{1}'$ of the A_4 symmetry, respectively. We introduce the right-handed Majorana neutrinos which are assigned to triplet of the A_4 symmetry. We also introduce three Higgs doublets which are assigned as triplet of the A_4 symmetry as discussed in Section 2. In Table 1, we summarize the particle assignments of $SU(2)_L$ and A_4 symmetry².

	$\bar{\ell} = (\bar{\ell}_e, \bar{\ell}_\mu, \bar{\ell}_\tau)$	e_R	μ_R	τ_R	$\nu_R = (\nu_{R1}, \nu_{R2}, \nu_{R3})$	$\phi = (\phi_1, \phi_2, \phi_3)$
$SU(2)_L$	$\mathbf{2}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{2}$
A_4	$\mathbf{3}$	$\mathbf{1}$	$\mathbf{1}''$	$\mathbf{1}'$	$\mathbf{3}$	$\mathbf{3}$

Table 1: The charge assignments of $SU(2)_L \times A_4$ symmetry in our model.

We can now write down the Lagrangian for Yukawa interactions and Majorana mass term in our model. The $SU(2)_L \times A_4$ invariant Lagrangian is written as

$$\mathcal{L}_Y = \mathcal{L}_\ell + \mathcal{L}_D + \mathcal{L}_M + h.c., \tag{28}$$

¹We also find the trivial solution $(0, 0, 0)$ and general solution (v_1, v_2, v_3) , $v_1 \neq v_2 \neq v_3$. Of course, the trivial solution cannot be realized the current experimental data. In the general solution, we cannot predict the physical quantity because there are many free parameters. Then, we discuss one of the solutions which can be predicted the Dirac CP phase, Majorana phases, and effective mass for the $0\nu\beta\beta$ decay.

²In the AF model [45, 46], they introduce the Z_3 symmetry in order to obtain the relevant couplings. Thanks for the $SU(2)_L$ gauge symmetry, we do not need to add the extra symmetry in our model.

where

$$\mathcal{L}_\ell = y_e \bar{\ell} \phi e_R + y_\mu \bar{\ell} \phi \mu_R + y_\tau \bar{\ell} \phi \tau_R, \quad (29)$$

$$\mathcal{L}_D = y_D \bar{\ell} \tilde{\phi} \nu_R, \quad (30)$$

$$\mathcal{L}_M = \frac{1}{2} M \bar{\nu}_R^c \nu_R. \quad (31)$$

Note that y_e, y_μ, y_τ , and y_D are Yukawa couplings and M is the right-handed Majorana neutrino mass. After the SSB, three Higgs doublets have VEVs as $\langle \phi \rangle = (v_1, v_2, v_3)$. In the charged lepton sector Eq. (29), the Yukawa interactions are rewritten as

$$\begin{aligned} y_e \bar{\ell} \phi e_R &= y_e (\bar{\ell}_e \phi_1 + \bar{\ell}_\mu \phi_3 + \bar{\ell}_\tau \phi_2) e_R \\ &= \frac{1}{\sqrt{2}} y_e (\bar{e}_L v_1 + \bar{\mu}_L v_3 + \bar{\tau}_L v_2) e_R, \end{aligned} \quad (32)$$

$$\begin{aligned} y_\mu \bar{\ell} \phi \mu_R &= y_\mu (\bar{\ell}_\tau \phi_3 + \bar{\ell}_e \phi_2 + \bar{\ell}_\mu \phi_1) \mu_R \\ &= \frac{1}{\sqrt{2}} y_\mu (\bar{\tau}_L v_3 + \bar{e}_L v_2 + \bar{\mu}_L v_1) \mu_R, \end{aligned} \quad (33)$$

$$\begin{aligned} y_\tau \bar{\ell} \phi \tau_R &= y_\tau (\bar{\ell}_\mu \phi_2 + \bar{\ell}_e \phi_3 + \bar{\ell}_\tau \phi_1) \tau_R \\ &= \frac{1}{\sqrt{2}} y_\tau (\bar{\mu}_L v_2 + \bar{e}_L v_3 + \bar{\tau}_L v_1) \tau_R. \end{aligned} \quad (34)$$

Then, the charged lepton mass matrix M_ℓ is obtained as

$$M_\ell = \frac{1}{\sqrt{2}} \begin{pmatrix} y_e v_1 & y_\mu v_2 & y_\tau v_3 \\ y_e v_3 & y_\mu v_1 & y_\tau v_2 \\ y_e v_2 & y_\mu v_3 & y_\tau v_1 \end{pmatrix}_{LR}. \quad (35)$$

Here and Hereafter we take the left-right basis in the mass matrices. In order to obtain the left-side unitary mixing matrix U_ℓ , we consider $M_\ell M_\ell^\dagger$ as

$$\begin{aligned} M_\ell M_\ell^\dagger &= \frac{1}{2} \times \\ &\begin{pmatrix} |y_e|^2 v_1^2 + |y_\mu|^2 v_2^2 + |y_\tau|^2 v_3^2 & |y_e|^2 v_1 v_3 + |y_\mu|^2 v_1 v_2 + |y_\tau|^2 v_2 v_3 & |y_e|^2 v_1 v_2 + |y_\mu|^2 v_2 v_3 + |y_\tau|^2 v_1 v_3 \\ \dots & |y_e|^2 v_3^2 + |y_\mu|^2 v_1^2 + |y_\tau|^2 v_2^2 & |y_e|^2 v_2 v_3 + |y_\mu|^2 v_1 v_3 + |y_\tau|^2 v_1 v_2 \\ \dots & \dots & |y_e|^2 v_2^2 + |y_\mu|^2 v_3^2 + |y_\tau|^2 v_1^2 \end{pmatrix}. \end{aligned} \quad (36)$$

Then, the elements of the mass matrix Eq. (36) are real and we can numerically calculate the charged lepton masses in Section 4. Next, we derive the Dirac neutrino mass matrix from Eq. (30). By A_4 product rule (See appendix B.), we can make the singlet term. When we take the complex conjugate, we need to take care $\tilde{\phi} = (\tilde{\phi}_1, \tilde{\phi}_3, \tilde{\phi}_2)$ because of the complex conjugate for the generator T in Eq. (13). Since Eq. (30) is $3 \otimes 3 \otimes 3$ form in A_4 symmetry, we first make $3_S \oplus 3_A \in 3 \otimes 3$ as

$$y_D \bar{\ell} \tilde{\phi} = \frac{y_{DS}}{3} \begin{pmatrix} 2\bar{\ell}_e \tilde{\phi}_1 - \bar{\ell}_\mu \tilde{\phi}_2 - \bar{\ell}_\tau \tilde{\phi}_3 \\ 2\bar{\ell}_\tau \tilde{\phi}_2 - \bar{\ell}_e \tilde{\phi}_3 - \bar{\ell}_\mu \tilde{\phi}_1 \\ 2\bar{\ell}_\mu \tilde{\phi}_3 - \bar{\ell}_\tau \tilde{\phi}_1 - \bar{\ell}_e \tilde{\phi}_2 \end{pmatrix}_{3_S} + \frac{y_{DA}}{2} \begin{pmatrix} \bar{\ell}_\mu \tilde{\phi}_2 - \bar{\ell}_\tau \tilde{\phi}_3 \\ \bar{\ell}_e \tilde{\phi}_3 - \bar{\ell}_\mu \tilde{\phi}_1 \\ \bar{\ell}_\tau \tilde{\phi}_1 - \bar{\ell}_e \tilde{\phi}_2 \end{pmatrix}_{3_A}, \quad (37)$$

where y_{DS} and y_{DA} are the symmetric and anti-symmetric Dirac Yukawa couplings, respectively. Then, the Dirac neutrino Yukawa interaction in Eq. (30) can be written as follows:

$$y_D \bar{\ell} \tilde{\phi} \nu_R = \frac{y_{DS}}{3} \begin{pmatrix} 2\bar{\ell}_e \tilde{\phi}_1 - \bar{\ell}_\mu \tilde{\phi}_2 - \bar{\ell}_\tau \tilde{\phi}_3 \\ 2\bar{\ell}_\tau \tilde{\phi}_2 - \bar{\ell}_e \tilde{\phi}_3 - \bar{\ell}_\mu \tilde{\phi}_1 \\ 2\bar{\ell}_\mu \tilde{\phi}_3 - \bar{\ell}_\tau \tilde{\phi}_1 - \bar{\ell}_e \tilde{\phi}_2 \end{pmatrix}_{3_S} \otimes \begin{pmatrix} \nu_{R1} \\ \nu_{R2} \\ \nu_{R3} \end{pmatrix} + \frac{y_{DA}}{2} \begin{pmatrix} \bar{\ell}_\mu \tilde{\phi}_2 - \bar{\ell}_\tau \tilde{\phi}_3 \\ \bar{\ell}_e \tilde{\phi}_3 - \bar{\ell}_\mu \tilde{\phi}_1 \\ \bar{\ell}_\tau \tilde{\phi}_1 - \bar{\ell}_e \tilde{\phi}_2 \end{pmatrix}_{3_A} \otimes \begin{pmatrix} \nu_{R1} \\ \nu_{R2} \\ \nu_{R3} \end{pmatrix} \quad (38)$$

$$= \frac{y_{DS}}{3} \left[(2\bar{\ell}_e \tilde{\phi}_1 - \bar{\ell}_\mu \tilde{\phi}_2 - \bar{\ell}_\tau \tilde{\phi}_3) \nu_{R1} + (2\bar{\ell}_\tau \tilde{\phi}_2 - \bar{\ell}_e \tilde{\phi}_3 - \bar{\ell}_\mu \tilde{\phi}_1) \nu_{R3} \right. \\ \left. + (2\bar{\ell}_\mu \tilde{\phi}_3 - \bar{\ell}_\tau \tilde{\phi}_1 - \bar{\ell}_e \tilde{\phi}_2) \nu_{R2} \right] \\ + \frac{y_{DA}}{2} \left[(\bar{\ell}_\mu \tilde{\phi}_2 - \bar{\ell}_\tau \tilde{\phi}_3) \nu_{R1} + (\bar{\ell}_e \tilde{\phi}_3 - \bar{\ell}_\mu \tilde{\phi}_1) \nu_{R3} + (\bar{\ell}_\tau \tilde{\phi}_1 - \bar{\ell}_e \tilde{\phi}_2) \nu_{R2} \right], \quad (39)$$

where $\tilde{\phi} = -i\sigma_2 \phi^*$ and the VEVs of $\tilde{\phi}_i$ are $\langle \tilde{\phi}_i \rangle = (v_i, 0)$. Therefore, the Dirac neutrino mass matrix M_D is obtained as

$$M_D = \frac{y_{DS}}{3\sqrt{2}} \begin{pmatrix} 2v_1 & -v_2 & -v_3 \\ -v_2 & 2v_3 & -v_1 \\ -v_3 & -v_1 & 2v_2 \end{pmatrix}_{LR} + \frac{y_{DA}}{2\sqrt{2}} \begin{pmatrix} 0 & -v_2 & v_3 \\ v_2 & 0 & -v_1 \\ -v_3 & v_1 & 0 \end{pmatrix}_{LR}. \quad (40)$$

Next we discuss the right-handed Majorana neutrino mass matrix. In Eq. (31), the right-handed Majorana neutrino mass term is decomposed as follows:

$$M \bar{\nu}_R^c \nu_R = M (\bar{\nu}_{R1}^c \nu_{R1} + \bar{\nu}_{R2}^c \nu_{R3} + \bar{\nu}_{R3}^c \nu_{R2}). \quad (41)$$

Then, the right-handed Majorana neutrino mass matrix M_R is obtained as

$$M_R = M \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (42)$$

In order to obtain the left-handed Majorana neutrino mass matrix, we consider the type-I seesaw mechanism [47–51]. By using the type-I seesaw mechanism, the left-handed Majorana neutrino mass matrix m_ν is written as

$$m_\nu = -M_D M_R^{-1} M_D^T \quad (43)$$

$$= \frac{1}{M} \begin{pmatrix} 2v_2 v_3 (y_{DA}^2 - y_{DS}^2) & v_1 v_2 (y_{DS}^2 - 4y_{DS} y_{DA} - y_{DA}^2) & v_1 v_3 (y_{DS}^2 + 4y_{DS} y_{DA} - y_{DA}^2) \\ v_1 v_2 (y_{DS}^2 - 4y_{DS} y_{DA} - y_{DA}^2) & 4v_1 v_3 y_{DS} (y_{DS} + y_{DA}) & -y_{DS}^2 (v_1^2 + 5v_2 v_3) \\ v_1 v_3 (y_{DS}^2 + 4y_{DS} y_{DA} - y_{DA}^2) & -y_{DS}^2 (v_1^2 + 5v_2 v_3) & 4v_1 v_2 y_{DS} (y_{DS} - y_{DA}) \end{pmatrix} \\ + \frac{1}{M} \begin{pmatrix} -4v_1^2 y_{DS}^2 & 2v_3^2 y_{DS} (y_{DS} - y_{DA}) & 2v_2^2 y_{DS} (y_{DS} + y_{DA}) \\ 2v_3^2 y_{DS} (y_{DS} - y_{DA}) & -v_2^2 (y_{DA} - y_{DS})^2 & y_{DA}^2 (v_1^2 + v_2 v_3) \\ 2v_2^2 y_{DS} (y_{DS} + y_{DA}) & y_{DA}^2 (v_1^2 + v_2 v_3) & -v_3^2 (y_{DA} - y_{DS})^2 \end{pmatrix}, \quad (44)$$

where we redefine the Dirac Yukawa couplings $y_{DS}/3\sqrt{2} \rightarrow y_{DS}$ and $y_{DA}/2\sqrt{2} \rightarrow y_{DA}$ for simplicity. In our model, the Dirac neutrino mass matrix has symmetric and anti-symmetric Yukawa couplings for the A_4 symmetry in Eq. (40). On the other hand, in Eq. (42), the right-handed Majorana neutrino mass matrix has a simple flavor structure, c.f., in the AF Model the Dirac neutrino mass matrix is simple. On the other hand, the right-handed Majorana neutrino mass matrix is the structure which derives the TBM in their model. In the next section, we show the numerical analysis.

Inverted Ordering	bfp $\pm 1\sigma$	3σ range
$\sin^2 \theta_{12}$	$0.304^{+0.013}_{-0.012}$	$0.269 \rightarrow 0.343$
$\theta_{12}/^\circ$	$33.45^{+0.78}_{-0.75}$	$31.27 \rightarrow 35.87$
$\sin^2 \theta_{23}$	$0.570^{+0.016}_{-0.022}$	$0.410 \rightarrow 0.613$
$\theta_{23}/^\circ$	$49.0^{+0.9}_{-1.3}$	$39.8 \rightarrow 51.6$
$\sin^2 \theta_{13}$	$0.02241^{+0.00074}_{-0.00062}$	$0.02055 \rightarrow 0.02457$
$\theta_{13}/^\circ$	$8.61^{+0.14}_{-0.12}$	$8.24 \rightarrow 9.02$
$\delta_{\text{CP}}/^\circ$	278^{+22}_{-30}	$194 \rightarrow 345$
$\frac{\Delta m_{21}^2}{10^{-5}[\text{eV}^2]}$	$7.42^{+0.21}_{-0.20}$	$6.82 \rightarrow 8.04$
$\frac{\Delta m_{32}^2}{10^{-3}[\text{eV}^2]}$	$-2.490^{+0.026}_{-0.028}$	$-2.574 \rightarrow -2.410$

Table 2: NuFIT 5.1 data [54, 55] in the inverted neutrino mass ordering, where Δm_{ij}^2 is mass squared difference between m_i and m_j .

4 Numerical analysis

In this section, we show the numerical analysis such as the lepton flavor mixing angles, Dirac CP phase, Majorana phases, and the effective mass for the $0\nu\beta\beta$ decay. We discuss what is our model verifiable in the near future experiments.

In section 2.2, we have analyzed the Higgs potential and found the following solution which is derived from Higgs potential minimization as

$$(v_1, v_2, v_3) = (v \cos \beta, \frac{v}{\sqrt{2}} \sin \beta, \frac{v}{\sqrt{2}} \sin \beta). \quad (45)$$

Since charged lepton mass matrix Eq. (36) is only depend on three Yukawa couplings and Higgs VEV. Then once we fix the Higgs VEV, we can obtain charged lepton Yukawa couplings by solving following equations and unitary matrix which diagonalizes charged lepton mass matrix.

$$\begin{aligned} \text{Tr}(M_\ell M_\ell^\dagger) &= m_e^2 + m_\mu^2 + m_\tau^2, \\ \text{Det}(M_\ell M_\ell^\dagger) &= m_e^2 m_\mu^2 m_\tau^2, \\ \left(\text{Tr} M_\ell M_\ell^\dagger\right)^2 - \text{Tr} \left[(M_\ell M_\ell^\dagger)^2 \right] &= 2(m_e^2 m_\mu^2 + m_\mu^2 m_\tau^2 + m_\tau^2 m_e^2), \end{aligned} \quad (46)$$

where m_e, m_μ, m_τ are charged lepton masses. The Dirac neutrino Yukawa couplings similarly obtained by solving equation which is substituted Eq. (43) into Eq. (46), where we use the left-handed Majorana neutrino masses instead of the charged lepton masses in the right-side of Eq. (46). However neutrino masses are only known mass squared differences Δm_{21}^2 and $\Delta m_{32(31)}^2$ in the inverted (normal) ordering of the neutrino mass hierarchy. Then we need to decide the Dirac neutrino Yukawa couplings y_{DS} and y_{DA} to satisfy experimental data in Table 2. In our numerical analysis, we simulate by assigning different numerical values to the lightest neutrino mass at the range $m_3 \in [0, 15.9]$ [meV], where the upper limit of the lightest neutrino mass is estimated in Ref. [57]. We analyze our model in normal and inverted neutrino mass orderings.




Figure 1: (a) The relation between the Dirac neutrino Yukawa couplings y_{DS} and y_{DA} . We simulate by assigning different numerical values to the Dirac neutrino Yukawa couplings at $\mathcal{O}(1)$. (b) Higgs VEV ratio $\tan\beta$ and complex phase of the Dirac Yukawa coupling ϕ_{DA} .

When we assume the normal ordering, there are no realistic parameters which satisfy the current experimental data in Table 2. We then assume the inverted ordering for the rest of our discussion. In Fig. 1a, we show the symmetric and anti-symmetric Dirac Yukawa couplings for Eq. (40) which are satisfied the current experimental data in Table 2. These Yukawa couplings look like proportional each other and $\mathcal{O}(1)$. The Majorana mass M in Eq. (42) is allowed at $\mathcal{O}(10^{15})$ [GeV]. We also show the numerical results of the Higgs VEV ratio $\tan\beta$ and the complex phase of the Dirac Yukawa coupling in Fig. 1b. The $\tan\beta$ is localized around $\tan\beta \in [0.4, 0.6], [0.8, 1.2]$. The complex phase of the Dirac Yukawa coupling ϕ_{DA} only appear in $70[^\circ] - 100[^\circ]$. In our model, the complex phase of the Dirac Yukawa coupling which contribute to the mixing matrix is only ϕ_{DA} , then this result has a strong influence to the CP phases.

In our calculation, we can obtain two unitary matrices U_ℓ , U_ν which diagonalize mass matrices Eq. (35) and Eq. (43) and also obtain the PMNS matrix $U_{\text{PMNS}} = U_\ell^\dagger U_\nu$,

$$\begin{aligned}
 U_{\text{PMNS}} &= \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu1} & U_{\mu2} & U_{\mu3} \\ U_{\tau1} & U_{\tau2} & U_{\tau3} \end{pmatrix} \\
 &= \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{\text{CP}}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{\text{CP}}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{\text{CP}}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{\text{CP}}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{\text{CP}}} & c_{23}c_{13} \end{pmatrix},
 \end{aligned} \tag{47}$$

where $U_{\alpha i}$ ($\alpha = e, \mu, \tau$, $i = 1, 2, 3$) are the PMNS matrix elements, c_{ij} and s_{ij} denote $\cos\theta_{ij}$ and $\sin\theta_{ij}$, and δ_{CP} is the Dirac CP phase, respectively [58]. The lepton mixing angles are obtained as follows:

$$\sin\theta_{13} = |U_{e3}|, \quad \tan\theta_{12} = \left| \frac{U_{e2}}{U_{e1}} \right|, \quad \tan\theta_{23} = \left| \frac{U_{\mu3}}{U_{\tau3}} \right|. \tag{48}$$

In addition, the Dirac CP phase is determined by one of the Jarlskog Invariants [59] as

$$J_{\text{CP}} = \text{Im} [U_{e1}U_{e2}^*U_{\mu2}U_{\mu1}^*], \tag{49}$$

and

$$J_{\text{CP}} = \sin\theta_{12} \cos\theta_{12} \sin\theta_{23} \cos\theta_{23} \sin\theta_{13} \cos^2\theta_{13} \sin\delta_{\text{CP}}, \tag{50}$$

in the PDG parametrization in Ref. [58]. The δ_{CP} is also determined by one of the absolute values for PMNS mixing matrix element:

$$|U_{\tau 1}|^2 = \sin^2 \theta_{12} \sin^2 \theta_{23} + \cos^2 \theta_{12} \cos^2 \theta_{23} \sin^2 \theta_{13} - 2 \sin \theta_{12} \sin \theta_{23} \cos \theta_{12} \cos \theta_{23} \sin \theta_{13} \cos \delta_{\text{CP}}. \quad (51)$$

Then, we can determine the δ_{CP} . In Fig. 2, we show the allowed region for the $\sin^2 \theta_{23}$ and δ_{CP} with 3σ range of the Table 2. The gray area is without 3σ range of δ_{CP} for NuFIT 5.1 data in Ref. [54, 55]. The relation between $\sin^2 \theta_{23}$ and Dirac CP phase δ_{CP} has strong correlation. Especially, in $\sin^2 \theta_{23} \in [0.41, 0.52]$, this relation has one to one correspondence because the complex phase comes from one Yukawa coupling phase ϕ_{DA} . Then if the θ_{23} is more precise measured by the future neutrino oscillation experiments, the Dirac CP phase is more precise predicted, and vice versa.




Figure 2: The allowed region for the $\sin^2 \theta_{23}$ and δ_{CP} with 3σ range of the Table 2. The gray area is without 3σ range in NuFIT 5.1 data [54, 55].

The parametrization in Eq. (47) is the case of the Dirac neutrino. If neutrinos are Majorana particles, there are three CP phases such as one Dirac CP phase δ_{CP} and two Majorana phases η_1, η_2 when we consider three generations. The above unitary matrix U_ν which we obtained from analysis has complex phases of mass eigenstates:

$$U_\nu^\dagger m_\nu U_\nu^* = \text{diag}(m_1 e^{i\phi_1}, m_2 e^{i\phi_2}, m_3 e^{i\phi_3}). \quad (52)$$

To remove these phases from mass diagonal matrix, we need to multiply phase diagonal matrix $P_\nu = \text{diag}(e^{i\phi_1/2}, e^{i\phi_2/2}, e^{i\phi_3/2})$ on both side of Eq (52),

$$P_\nu^\dagger (U_\nu^\dagger m_\nu U_\nu^*) P_\nu^* = \text{diag}(m_1, m_2, m_3). \quad (53)$$

The unitary matrix which makes real mass diagonal matrix is $U_\nu P_\nu$. Therefore the full parametrization of PMNS matrix when there are three generation Majorana neutrinos is written as [58]

$$U_{\text{PMNS}} = U_\ell^\dagger U_\nu P_\nu \quad (54)$$

$$= \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{\text{CP}}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{\text{CP}}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{\text{CP}}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{\text{CP}}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{\text{CP}}} & c_{23}c_{13} \end{pmatrix} \begin{pmatrix} e^{i\eta_1} & & \\ & e^{i\eta_2} & \\ & & 1 \end{pmatrix},$$




Figure 3: (a) The relation between the effective mass for $0\nu\beta\beta$ decay m_{ee} and lightest neutrino mass. The red area and blue area are model independent analyses for the inverted ordering and normal ordering of the neutrino mass hierarchies, respectively. The gray area is upper limit on the effective mass of 36 - 156 [meV] at 90% C.L. in Ref. [7]. The yellow area is upper limit on the lightest neutrino mass $m_3 \simeq 15.9$ [meV] which is estimated in Ref. [57]. (b) The relation among Majorana phases η_1 and η_2 .

where $\eta_1 = \frac{1}{2}(\phi_1 - \phi_3)$ and $\eta_2 = \frac{1}{2}(\phi_2 - \phi_3)$. In our numerical calculation, these Majorana phases η_1 and η_2 are determined as follows:

$$\eta_1 = \arg \left[\frac{U_{e1}U_{e3}^*}{\cos \theta_{12} \cos \theta_{13} \sin \theta_{13} e^{i\delta_{CP}}} \right], \quad \eta_2 = \arg \left[\frac{U_{e2}U_{e3}^*}{\sin \theta_{12} \cos \theta_{13} \sin \theta_{13} e^{i\delta_{CP}}} \right]. \quad (55)$$

The $0\nu\beta\beta$ decay is one of the attractive phenomena which is determined the magnitude of the lightest neutrino mass and neutrino mass ordering. In Fig. 3a, we show the prediction of the the effective mass m_{ee} for the $0\nu\beta\beta$ decay as

$$m_{ee} = \left| \sum_{i=1}^3 m_i U_{ei}^2 \right| = |m_1 U_{e1}^2 + m_2 U_{e2}^2 + m_3 U_{e3}^2|. \quad (56)$$

Note that only inverted ordering is acceptable then the lightest neutrino mass is m_3 . The lightest neutrino mass m_3 appears in 0.789 - 1.43 [meV] which is the within the Planck data [57] and the effective mass take very restricted region $m_{ee} \simeq 47.1$ [meV]. This value is in upper limit on the effective mass of 36 - 156 [meV] at 90% C.L. in Ref. [7]. Then this model can be verified in the near future neutrino experiments, e.g. the KamLAND-Zen [6, 7], GERDA [8, 9], and CUORE [10, 11] experiments. In Fig. 3b, we show the relation among Majorana phases η_1 and η_2 . Since the phase of our model parameter is only ϕ_{DA} , then the Majorana phases are strongly correlated.

5 Summary and Discussions

We have proposed the simple non-SUSY lepton flavor model with A_4 symmetry. The A_4 group is a minimal one which includes triplet irreducible representation. We have introduced three Higgs doublets which are assigned as triplet of the A_4 symmetry. It is natural that there are

three generations as same as the SM fermions. First, we have studied the potential analysis and we have got the VEV for the local minimum. Next, we have presented our A_4 model. The left-handed lepton doublets are assigned to triplet and the right-handed leptons are assigned to different singlets of the A_4 symmetry, respectively. We have introduced the right-handed Majorana neutrinos which are assigned to triplet of the A_4 symmetry. In our model, the right-handed Majorana neutrino mass matrix has a simple flavor structure. On the other hand, the Dirac neutrino mass matrix has symmetric and anti-symmetric Yukawa couplings for A_4 symmetry. By using the type-I seesaw mechanism, we have obtained the left-handed Majorana neutrino mass matrix. After diagonalizing the charged lepton and left-handed Majorana neutrino mass matrices, we have got the PMNS mixing matrix. In our numerical analyses, we have used the NuFIT 5.1 data. We found that only inverted ordering is acceptable and we could not find the solutions for normal ordering in the neutrino mass hierarchy. We have obtained relevant relations for mixing angles and neutrino effective mass m_{ee} as a function of the lightest neutrino mass. Especially, the Dirac CP phase and lepton mixing angle θ_{23} are strongly correlated. If the θ_{23} is more precise measured, the Dirac CP phase is more precise predicted, and vice versa. In this model, the effective mass for the $0\nu\beta\beta$ decay can be predicted as $m_{ee} \simeq 47.1$ [meV] and the lightest neutrino mass can be also predicted as $m_3 \simeq 0.789 - 1.43$ [meV]. It is testable for near future neutrino experiments.

The flavor symmetry also apply to the quark sector. In the same assignments as the charged leptons, the elements of the mass matrices are real. Then, we take quarks different assignments, e.g. left-handed quark doublets are assigned to the different singlets and right-handed up and down-type quarks are assigned to the triplets of the A_4 symmetry, respectively. The more details are in the future work. In our model, the right-handed Majorana mass matrix is very simple and masses are degenerated. Then, we cannot apply to the leptogenesis. Fortunately, there are three Higgs doublets, we can discuss the electroweak baryogenesis which are also in the future work.

Acknowledgement

We thank Y. Kawamura, Y. Matsuo, H. Shimoji, S. Takahashi, and A. Yuu for useful discussions.

Appendix

A Higgs mass matrices in the general 3HDM

The general 3HDM has nine scalar mass eigenstates, three CP-even scalars: $H, h_{1,2}$, two CP-odd scalars: $a_{1,2}$, and four charged scalars: $h_{1,2}^\pm$. More details are shown in Ref. [56].

A.1 CP-even scalar mass matrix

The mass matrix of CP-even scalars in the general 3HDM is written by

$$\mathcal{M}_s^2 = \begin{pmatrix} A & C_1 & C_2 \\ C_1 & B_1 & C_3 \\ C_2 & C_3 & B_2 \end{pmatrix}, \quad (57)$$

with

$$\begin{aligned}
A = & m_{12}^2 t_{\beta_1} + m_{13}^2 c_{\beta_1}^{-1} t_{\beta_2} + 2\lambda_{11} v^2 c_{\beta_1}^2 c_{\beta_2}^2 - \frac{1}{2} v^2 s_{\beta_1} c_{\beta_2}^2 (\lambda_{2212} t_{\beta_1}^2 - 3\lambda_{1112}) \\
& - \frac{1}{2} v^2 c_{\beta_1}^2 s_{\beta_2} c_{\beta_2} (t_{\beta_1}^2 (\lambda_{2213} + \lambda_{1223} + \lambda_{2123}) - 3\lambda_{1113}) \\
& - \frac{1}{2} v^2 s_{\beta_2}^2 (\lambda_{3313} c_{\beta_1}^{-1} t_{\beta_2} + t_{\beta_1} (\lambda_{1323} + \lambda_{1332} + \lambda_{3312})),
\end{aligned} \tag{58}$$

$$\begin{aligned}
B_1 = & m_{12}^2 t_{\beta_1}^{-1} + m_{23}^2 s_{\beta_1}^{-1} t_{\beta_2} + 2\lambda_{22} v^2 s_{\beta_1}^2 c_{\beta_2}^2 - \frac{1}{2} v^2 s_{\beta_1} c_{\beta_2}^2 (\lambda_{1112} t_{\beta_1}^2 - 3\lambda_{2212}) \\
& - \frac{1}{2} v^2 s_{\beta_1} s_{\beta_2} c_{\beta_2} (t_{\beta_1}^2 (\lambda_{1123} + \lambda_{1213} + \lambda_{2113}) - 3\lambda_{2223}) \\
& - \frac{1}{2} v^2 s_{\beta_2}^2 (\lambda_{3323} s_{\beta_1}^{-1} t_{\beta_2} + t_{\beta_1}^{-1} (\lambda_{1323} + \lambda_{1332} + \lambda_{3312})),
\end{aligned} \tag{59}$$

$$\begin{aligned}
B_2 = & 2m_{13}^2 c_{\beta_1} t_{\beta_2}^{-1} + 2m_{23}^2 s_{\beta_1} t_{\beta_2}^{-1} + 4\lambda_{33} v^2 s_{\beta_2}^2 + 3v^2 s_{\beta_2} c_{\beta_2} (\lambda_{3313} c_{\beta_1} + \lambda_{3323} s_{\beta_1}) \\
& v^2 c_{\beta_2}^2 t_{\beta_2}^{-1} (\lambda_{1113} c_{\beta_1}^3 + s_{\beta_1} c_{\beta_1}^2 (\lambda_{1123} + \lambda_{1213} + \lambda_{2113}) \\
& + s_{\beta_1}^2 c_{\beta_1} (\lambda_{1223} + \lambda_{2123} + \lambda_{2213}) + \lambda_{2223} s_{\beta_1}^3),
\end{aligned} \tag{60}$$

$$\begin{aligned}
C_1 = & -m_{12}^2 + \frac{1}{2} v^2 s_{2\beta_1} c_{\beta_2}^2 (\lambda_{1122} + \lambda_{1221}) + \frac{1}{2} v^2 c_{\beta_2}^2 (3\lambda_{1122} c_{\beta_1}^2 + \lambda_{1212} s_{2\beta_1} + 3\lambda_{2212} s_{\beta_1}^2) \\
& + \frac{1}{2} v^2 s_{2\beta_2} (c_{\beta_1} (\lambda_{1123} + \lambda_{1213} + \lambda_{2113}) + s_{\beta_1} (\lambda_{1223} + \lambda_{2123} + \lambda_{2213})) \\
& + \frac{1}{2} v^2 s_{\beta_2}^2 (\lambda_{1323} + \lambda_{1332} + \lambda_{3312}),
\end{aligned} \tag{61}$$

$$\begin{aligned}
C_2 = & -m_{13}^2 + \frac{1}{2} v^2 c_{\beta_1} s_{2\beta_2} (\lambda_{1133} + \lambda_{1331}) \\
& + \frac{1}{2} v^2 c_{\beta_2}^2 (3\lambda_{1113} c_{\beta_1}^2 + s_{2\beta_1} (\lambda_{1123} + \lambda_{1213} + \lambda_{2113}) + s_{\beta_1}^2 (\lambda_{1223} + \lambda_{2123} + \lambda_{2213})) \\
& + \frac{1}{2} v^2 s_{2\beta_2} (\lambda_{1313} c_{\beta_1} + s_{\beta_1} (\lambda_{1323} + \lambda_{1332} + \lambda_{3312})) + \frac{1}{2} v^2 3\lambda_{3313} s_{\beta_2}^2,
\end{aligned} \tag{62}$$

$$\begin{aligned}
C_3 = & -m_{23}^2 + \frac{1}{2} v^2 s_{\beta_1} s_{2\beta_2} (\lambda_{2233} + \lambda_{2332}) \\
& + \frac{1}{2} v^2 c_{\beta_2}^2 (c_{\beta_1}^2 \lambda_{1123} + \lambda_{1213} + \lambda_{2113}) + s_{2\beta_1} (\lambda_{1223} + \lambda_{2123}) + 3\lambda_{2223} s_{\beta_1}^2 \\
& + \frac{1}{2} v^2 s_{2\beta_2} (c_{\beta_1} (\lambda_{1323} + \lambda_{1332} + \lambda_{3312}) + \lambda_{2323} s_{\beta_1}) + 3\lambda_{3323} s_{\beta_2}^2.
\end{aligned} \tag{63}$$

A.2 CP-odd pseudo-scalar mass matrix

The mass matrix of the CP-odd scalars in the general 3HDM is written by

$$\mathcal{M}_P^2 = \begin{pmatrix} M_{P,22}^2 & M_{P,23}^2 \\ M_{P,32}^2 & M_{P,33}^2 \end{pmatrix}, \tag{64}$$

with

$$\begin{aligned}
M_{P,22}^2 = & m_{12}^2 s_{2\beta_1} + m_{12}^2 c_{\beta_1}^2 t_{\beta_1}^{-1} + s_{\beta_1} t_{\beta_1} (m_{12}^2 s_{\beta_1} + m_{13}^2 t_{\beta_2}) + m_{23}^2 c_{\beta_1} t_{\beta_1}^{-1} t_{\beta_2} \\
& + \frac{1}{4} t_{\beta_1}^{-1} (-s_{\beta_1} (v^2 s_{\beta_1} t_{\beta_1}^2 c_{\beta_2}^2 (c_{2\beta_1} (\lambda_{1112} + \lambda_{2212}) + 2\lambda_{1212} s_{2\beta_1} + \lambda_{1112} + 3\lambda_{2212}) \\
& + \frac{1}{2} v^2 t_{\beta_1}^2 (s_{2\beta_2} (c_{2\beta_1} (\lambda_{1113} - \lambda_{1223} + 3\lambda_{2123} - \lambda_{2213}) + 4\lambda_{1213} s_{2\beta_1} + \lambda_{1113} + \lambda_{1223} + 5\lambda_{2123} \\
& + \lambda_{2213}) + 4\lambda_{3313} s_{\beta_2}^2 t_{\beta_2}) + v^2 t_{\beta_1} c_{\beta_1}^{-1} s_{\beta_2}^2 (2\lambda_{1313} s_{2\beta_1} + c_{2\beta_1} (-3\lambda_{1323} + \lambda_{1332} + \lambda_{3312}) \\
& - \lambda_{1323} + 3(\lambda_{1332} + \lambda_{3312})) - \lambda_{2223} v^2 s_{2\beta_1} s_{2\beta_2}) + v^2 c_{\beta_1}^2 (c_{\beta_2}^2 (c_{2\beta_1} (\lambda_{1112} + \lambda_{2212}) - 3\lambda_{1112} \\
& - \lambda_{2212}) - 8\lambda_{2123} s_{\beta_1} s_{\beta_2} c_{\beta_2} - 2s_{\beta_2}^2 (\lambda_{1323} + \lambda_{1332} + \lambda_{3312})) - 2v^2 c_{3\beta_1}^3 c_{\beta_2} (2\lambda_{1212} s_{\beta_1} c_{\beta_2} \\
& + s_{\beta_2} (\lambda_{1123} + \lambda_{1213} + \lambda_{2113})) - 2v^2 c_{\beta_1} (4\lambda_{1212} s_{3\beta_1}^3 c_{\beta_2}^2 + s_{\beta_1}^2 s_{\beta_2} c_{\beta_2} (4\lambda_{1213} + 3\lambda_{2223}) \\
& + s_{\beta_2}^2 (2\lambda_{2223} s_{\beta_1} + \lambda_{3323} t_{\beta_2}))),
\end{aligned} \tag{65}$$

$$\begin{aligned}
M_{P,33}^2 = & s_{\beta_2}^{-1} c_{\beta_2}^{-1} (m_{13}^2 c_{\beta_1} + m_{23}^2 s_{\beta_1}) + \frac{1}{16} (-c_{\beta_1} s_{\beta_2}^{-1} c_{\beta_2}^{-1} (v^2 c_{2\beta_2} (3\lambda_{1113} + \lambda_{1223} + \lambda_{2123} + \lambda_{2213} \\
& - 4\lambda_{3313}) + v^2 (3\lambda_{1223} + \lambda_{2123} + 4\lambda_{3313})) + 2v^2 c_{3\beta_1} t_{\beta_2}^{-1} (-\lambda_{1113} + \lambda_{1223} \\
& + \lambda_{2123} + \lambda_{2213}) - s_{\beta_1} s_{\beta_2}^{-1} c_{\beta_2}^{-1} (v^2 c_{2\beta_2} (\lambda_{1123} + \lambda_{1213} + \lambda_{2113} + 3\lambda_{2223} - 4\lambda_{3323}) \\
& + v^2 (\lambda_{1123} + \lambda_{1213} + \lambda_{2113} + 3\lambda_{2223} + 4\lambda_{3323})) - 2v^2 s_{3\beta_1} t_{\beta_2}^{-1} (\lambda_{1123} + \lambda_{1213} + \lambda_{2113} \\
& - \lambda_{2223}) + 8v^2 c_{2\beta_1} (\lambda_{2323} - \lambda_{1313}) - 16\lambda_{1323} v^2 s_{2\beta_1} - 8v^2 (\lambda_{1313} + \lambda_{2323})),
\end{aligned} \tag{66}$$

$$\begin{aligned}
M_{P,23}^2 = & c_{\beta_2}^{-1} (m_{13}^2 s_{\beta_1} - m_{23}^2 c_{\beta_1} + \frac{1}{8} (v^2 c_{\beta_2} (-s_{\beta_1} (\lambda_{1113} + 3\lambda_{1223} - 5\lambda_{2123} + 3\lambda_{2213} - 4\lambda_{3313}) \\
& + s_{3\beta_1} (-\lambda_{1113} + \lambda_{1223} + \lambda_{2123} + \lambda_{2213}) + c_{\beta_1} (3\lambda_{1123} - 5\lambda_{1213} + 3\lambda_{2113} + \lambda_{2223} - 4\lambda_{3323}) \\
& + c_{3\beta_1} (\lambda_{1123} + \lambda_{1213} + \lambda_{2113} + -\lambda_{2223})) + 4v^2 s_{\beta_2} (s_{2\beta_1} (\lambda_{2323} - \lambda_{1313}) + 2\lambda_{1323} c_{2\beta_1}) \\
& + 4c_{\beta_2}^{-1} (\lambda_{3323} v^2 c_{\beta_1} - \lambda_{3313} v^2) s_{\beta_1})).
\end{aligned} \tag{67}$$

The above mass matrix \mathcal{M}_P^2 can be diagonalized by using a rotational matrix $R(\rho)$ in the form

$$\bar{\mathcal{M}}_P^2 = R(\sigma) \mathcal{M}_P^2 R^\top(\sigma) = \begin{pmatrix} M_{a1}^2 & 0 \\ 0 & M_{a2}^2 \end{pmatrix}, \tag{68}$$

with

$$M_{a1}^2 = M_{P,22}^2 c_\sigma^2 + M_{P,23}^2 s_{2\rho} + M_{P,33}^2 s_\sigma^2, \tag{69}$$

$$M_{a2}^2 = \frac{1}{2} (M_{P,33}^2 - M_{P,22}^2) s_{2\sigma} + M_{P,23}^2 c_{2\sigma}, \tag{70}$$

where

$$\tan 2\sigma = \frac{2M_{P,23}^2}{M_{P,22}^2 - M_{P,33}^2}. \tag{71}$$

The rotational matrix $R(\sigma)$ is

$$R(\sigma) = \begin{pmatrix} \cos \sigma & \sin \sigma \\ -\sin \sigma & \cos \sigma \end{pmatrix}. \tag{72}$$

A.3 Charged scalar mass matrix

The mass matrix of the charged scalars in the general 3HDM is written by

$$\mathcal{M}_{\pm}^2 = \begin{pmatrix} M_{\pm,22}^2 & M_{\pm,23}^2 \\ M_{\pm,23}^2 & M_{\pm,33}^2 \end{pmatrix}, \quad (73)$$

with

$$\begin{aligned} M_{\pm,22}^2 = & c_{\beta_1}^{-1} (m_{12}^2 c_{\beta_1}^{-3} + t_{\beta_2} (m_{13}^2 t_{\beta_1}^3 + m_{23}^2)) - \frac{1}{2} c_{\beta_1}^3 t_{\beta_1}^{-1} \left(\frac{1}{2} v^2 c_{\beta_1}^{-5} c_{\beta_2}^2 (c_{2\beta_1} (\lambda_{1112} - \lambda_{2212}) \right. \\ & + s_{2\beta_1} (\lambda_{1212} + \lambda_{1221}) + \lambda_{1112} + \lambda_{2212}) + v^2 s_{\beta_2} c_{\beta_2} (\lambda_{1113} t_{\beta_1}^3 + t_{\beta_1}^5 (\lambda_{1223} + \lambda_{2123} + \lambda_{2213}) \\ & + \lambda_{2223} t_{\beta_1}^2 + \lambda_{1123} + \lambda_{1213} + \lambda_{2113}) + v^2 t_{\beta_1} c_{\beta_1}^{-2} s_{2\beta_2} (t_{\beta_1} (\lambda_{1213} + \lambda_{2113}); \lambda_{1223} + \lambda_{2123}) \\ & + \frac{1}{8} v^2 c_{\beta_1}^{-5} s_{\beta_2}^2 (2s_{2\beta_1} (\lambda_{1313} + \lambda_{1331} + \lambda_{2323} + \lambda_{2332} + 2c_{4\beta_1} (\lambda_{1323} + \lambda_{1332}) + s_{4\beta_2} (\lambda_{2323} \\ & + \lambda_{2332} - \lambda_{1313} - \lambda_{1331}) + 6(\lambda_{1323} + \lambda_{1332}) + 8\lambda_{3312}) + v^2 c_{\beta_1}^{-2} s_{\beta_2}^2 t_{\beta_2} (\lambda_{3313} t_{\beta_1}^3 + \lambda_{3323}), \end{aligned} \quad (74)$$

$$\begin{aligned} M_{\pm,33}^2 = & s_{\beta_2}^{-1} c_{\beta_2}^{-1} (m_{13}^2 c_{\beta_1} + m_{23}^2 s_{\beta_1}) - \frac{1}{32} s_{\beta_2}^{-1} c_{\beta_2}^{-1} (16\lambda_{1113} v^2 c_{\beta_1}^3 c_{\beta_2}^2 + \lambda_{1123} v^2 (s_{\beta_1-2\beta_2} + s_{3\beta_1-2\beta_2}) \\ & + s_{\beta_1+2\beta_2} + s_{3\beta_1+2\beta_2}) + \lambda_{1213} v^2 (s_{\beta_1-2\beta_2} + s_{3\beta_1-2\beta_2} + s_{\beta_1+2\beta_2} + s_{3\beta_1+2\beta_2}) \\ & + \lambda_{1223} v^2 (c_{\beta_1-2\beta_2} - c_{3\beta_1-2\beta_2} + c_{\beta_1+2\beta_2} - c_{3\beta_1+2\beta_2}) - 2\lambda_{1313} v^2 (s_{2\beta_1-2\beta_2} + s_{2\beta_1+2\beta_2}) \\ & 4\lambda_{1323} v^2 (c_{2\beta_1-2\beta_2} - c_{2\beta_1+2\beta_2}) - 2\lambda_{1331} v^2 (s_{2\beta_1-2\beta_2} + s_{2\beta_1+2\beta_2}) + 4\lambda_{1332} v^2 (c_{2\beta_1-2\beta_2} \\ & + c_{2\beta_1+2\beta_2}) + \lambda_{2113} v^2 (s_{\beta_1-2\beta_2} + s_{3\beta_1-2\beta_2} + s_{\beta_1+2\beta_2} + s_{3\beta_1+2\beta_2}) + \lambda_{2123} v^2 (c_{\beta_1-2\beta_2} \\ & - c_{3\beta_1-2\beta_2} + c_{\beta_1+2\beta_2} - c_{3\beta_1+2\beta_2}) + \lambda_{2213} v^2 (c_{\beta_1-2\beta_2} - c_{3\beta_1-2\beta_2} + c_{\beta_1+2\beta_2} - c_{3\beta_1+2\beta_2}) \\ & + \lambda_{2223} v^2 (3s_{\beta_1-2\beta_2} - s_{3\beta_1-2\beta_2} + 3s_{\beta_1+2\beta_2} - s_{3\beta_1+2\beta_2}) + 2\lambda_{2323} v^2 (s_{2\beta_1-2\beta_2} - s_{2\beta_1+2\beta_2}) \\ & + 2\lambda_{2332} v^2 (s_{2\beta_1-2\beta_2} - s_{2\beta_1+2\beta_2}) - 4\lambda_{3313} v^2 (c_{\beta_1-2\beta_2} - c_{\beta_1+2\beta_2}) - 4\lambda_{3323} v^2 (s_{\beta_1-2\beta_2} \\ & - s_{\beta_1+2\beta_2}) + 2\lambda_{1123} v^2 (s_{\beta_1} + s_{3\beta_1}) + 2\lambda_{1213} v^2 (s_{\beta_1} + s_{3\beta_1}) + 2v^2 c_{\beta_1} (\lambda_{1223} + \lambda_{2123} + \lambda_{2213} \\ & + 4\lambda_{3313}) - 2v^2 c_{\beta_1} (\lambda_{1223} + \lambda_{2123} + \lambda_{2213}) + 2v^2 s_{\beta_1} (\lambda_{2123} + 3\lambda_{2223} + 4\lambda_{3323}) \\ & + 2v^2 s_{3\beta_1} (\lambda_{2113} - \lambda_{2223}) + 4v^2 s_{2\beta_2} (\lambda_{1313} + \lambda_{1331} + \lambda_{2323} + \lambda_{2332}), \end{aligned} \quad (75)$$

$$\begin{aligned} M_{\pm,23}^2 = & c_{\beta_2}^{-1} (m_{13}^2 s_{\beta_1} - m_{23}^2 c_{\beta_1}) + \frac{1}{16} c_{\beta_1} (8v^2 c_{\beta_1}^2 (s_{\beta_1} (-\lambda_{1113} + \lambda_{1223} + \lambda_{2123}) + t_{\beta_2} (\lambda_{1323} + \lambda_{1332})) \\ & - 4v^2 s_{\beta_1} (2s_{\beta_1} t_{\beta_2} (\lambda_{1323} + \lambda_{1332}) + s_{2\beta_1} (\lambda_{1213} + \lambda_{2113} - \lambda_{2223}) + 2\lambda_{2213} s_{\beta_1}^2 + 2\lambda_{3313} t_{\beta_2}^2) \\ & - v^2 s_{2\beta_1} s_{3\beta_2} c_{\beta_2}^{-3} (\lambda_{1313} + \lambda_{1331} - \lambda_{2323} - \lambda_{2332}) + v^2 s_{2\beta_1} t_{\beta_2} c_{\beta_2}^{-2} (-\lambda_{1313} - \lambda_{1331} + \lambda_{2323} \\ & + \lambda_{2332}) + 8\lambda_{3323} v^2 c_{\beta_1} t_{\beta_2}^2 + 8\lambda_{1123} v^2 c_{\beta_1}^3). \end{aligned} \quad (76)$$

The above mass matrix \mathcal{M}_{\pm}^2 can be diagonalised by using a rotational matrix $R(\tau)$ in the form

$$\bar{\mathcal{M}}_{\pm}^2 = R(\tau) \mathcal{M}_{\pm}^2 R(\tau)^{-1} = \begin{pmatrix} M_{h_1^{\pm}}^2 & 0 \\ 0 & M_{h_2^{\pm}}^2 \end{pmatrix}, \quad (77)$$

with

$$M_{h_1^{\pm}}^2 = M_{\pm,22}^2 c_{\tau}^2 + M_{\pm,23}^2 s_{2\tau} + M_{\pm,33}^2 s_{\tau}^2, \quad (78)$$

$$M_{h_2^{\pm}}^2 = \frac{1}{2} (M_{\pm,33}^2 - M_{\pm,22}^2) s_{2\tau} + M_{\pm,23}^2 c_{2\tau}, \quad (79)$$

where

$$\tan 2\tau = \frac{2M_{\pm,23}^2}{M_{\pm,22}^2 - M_{\pm,33}^2}. \quad (80)$$

B Multiplication rule of A_4 group

We use the multiplication rule of the A_4 triplet as follows:

$$\begin{aligned} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}_{\mathbf{3}} \otimes \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}_{\mathbf{3}} &= (a_1b_1 + a_2b_3 + a_3b_2)_{\mathbf{1}} \oplus (a_3b_3 + a_1b_2 + a_2b_1)_{\mathbf{1}'} \\ &\oplus (a_2b_2 + a_1b_3 + a_3b_1)_{\mathbf{1}''} \\ &\oplus \frac{1}{3} \begin{pmatrix} 2a_1b_1 - a_2b_3 - a_3b_2 \\ 2a_3b_3 - a_1b_2 - a_2b_1 \\ 2a_2b_2 - a_3b_1 - a_1b_3 \end{pmatrix}_{\mathbf{3}} \oplus \frac{1}{2} \begin{pmatrix} a_2b_3 - a_3b_2 \\ a_1b_2 - a_2b_1 \\ a_3b_1 - a_1b_3 \end{pmatrix}_{\mathbf{3}}. \end{aligned} \quad (81)$$

More details are shown in the review [14, 15].

References

- [1] K. Abe *et al.* [T2K], Phys. Rev. D **88** (2013) no.3, 032002 [arXiv:1304.0841 [hep-ex]].
- [2] K. Abe *et al.* [T2K], Phys. Rev. Lett. **112** (2014), 061802 [arXiv:1311.4750 [hep-ex]].
- [3] P. Adamson *et al.* [NOvA], Phys. Rev. Lett. **116** (2016) no.15, 151806 [arXiv:1601.05022 [hep-ex]].
- [4] F. P. An *et al.* [Daya Bay], Phys. Rev. Lett. **108** (2012), 171803 [arXiv:1203.1669 [hep-ex]].
- [5] J. K. Ahn *et al.* [RENO], Phys. Rev. Lett. **108** (2012), 191802 [arXiv:1204.0626 [hep-ex]].
- [6] A. Gando *et al.* [KamLAND-Zen], Phys. Rev. Lett. **117** (2016) no.8, 082503 [arXiv:1605.02889 [hep-ex]].
- [7] S. Abe *et al.* [KamLAND-Zen], [arXiv:2203.02139 [hep-ex]].
- [8] M. Agostini *et al.* [GERDA], Phys. Rev. Lett. **111** (2013) no.12, 122503 [arXiv:1307.4720 [nucl-ex]].
- [9] M. Agostini *et al.* [GERDA], Phys. Rev. Lett. **120** (2018) no.13, 132503 [arXiv:1803.11100 [nucl-ex]].
- [10] C. Alduino *et al.* [CUORE], Phys. Rev. Lett. **120** (2018) no.13, 132501 [arXiv:1710.07988 [nucl-ex]].
- [11] D. Q. Adams *et al.* [CUORE], Phys. Rev. Lett. **124** (2020) no.12, 122501 [arXiv:1912.10966 [nucl-ex]].

- [12] C. D. Froggatt and H. B. Nielsen, Nucl. Phys. B **147** (1979), 277-298 doi:10.1016/0550-3213(79)90316-X
- [13] G. Altarelli and F. Feruglio, Rev. Mod. Phys. **82** (2010), 2701-2729 [arXiv:1002.0211 [hep-ph]].
- [14] H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada and M. Tanimoto, Prog. Theor. Phys. Suppl. **183** (2010), 1-163 [arXiv:1003.3552 [hep-th]].
- [15] H. Ishimori, T. Kobayashi, H. Ohki, H. Okada, Y. Shimizu and M. Tanimoto, Lect. Notes Phys. **858** (2012), 1-227, Springer.
- [16] S. F. King, A. Merle, S. Morisi, Y. Shimizu and M. Tanimoto, New J. Phys. **16** (2014), 045018 [arXiv:1402.4271 [hep-ph]].
- [17] T. Kobayashi, H. Ohki, H. Okada, Y. Shimizu and M. Tanimoto, Lect. Notes Phys. **995** (2022), 1-353, Springer.
- [18] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530** (2002), 167 [arXiv:hep-ph/0202074 [hep-ph]].
- [19] P. F. Harrison and W. G. Scott, Phys. Lett. B **535** (2002), 163-169 [arXiv:hep-ph/0203209 [hep-ph]].
- [20] Z. z. Xing, Phys. Lett. B **533** (2002), 85-93 [arXiv:hep-ph/0204049 [hep-ph]].
- [21] Z. z. Xing and S. Zhou, Phys. Lett. B **653** (2007), 278-287 [arXiv:hep-ph/0607302 [hep-ph]].
- [22] B. Adhikary and A. Ghosal, Phys. Rev. D **75** (2007), 073020 [arXiv:hep-ph/0609193 [hep-ph]].
- [23] S. F. King, Phys. Lett. B **659** (2008), 244-251 [arXiv:0710.0530 [hep-ph]].
- [24] M. Honda and M. Tanimoto, Prog. Theor. Phys. **119** (2008), 583-598 [arXiv:0801.0181 [hep-ph]].
- [25] B. Brahmachari, S. Choubey and M. Mitra, Phys. Rev. D **77** (2008), 073008 [erratum: Phys. Rev. D **77** (2008), 119901] [arXiv:0801.3554 [hep-ph]].
- [26] B. Adhikary and A. Ghosal, Phys. Rev. D **78** (2008), 073007 [arXiv:0803.3582 [hep-ph]].
- [27] M. Hirsch, S. Morisi and J. W. F. Valle, Phys. Rev. D **79** (2009), 016001 [arXiv:0810.0121 [hep-ph]].
- [28] S. Morisi, Phys. Rev. D **79** (2009), 033008 [arXiv:0901.1080 [hep-ph]].
- [29] A. Hayakawa, H. Ishimori, Y. Shimizu and M. Tanimoto, Phys. Lett. B **680** (2009), 334-342 [arXiv:0904.3820 [hep-ph]].
- [30] S. Goswami, S. T. Petcov, S. Ray and W. Rodejohann, Phys. Rev. D **80** (2009), 053013 [arXiv:0907.2869 [hep-ph]].

- [31] J. Barry and W. Rodejohann, Phys. Rev. D **81** (2010), 093002 [erratum: Phys. Rev. D **81** (2010), 119901] [arXiv:1003.2385 [hep-ph]].
- [32] C. H. Albright, A. Dueck and W. Rodejohann, Eur. Phys. J. C **70** (2010), 1099-1110 [arXiv:1004.2798 [hep-ph]].
- [33] H. Ishimori, Y. Shimizu, M. Tanimoto and A. Watanabe, Phys. Rev. D **83** (2011), 033004 [arXiv:1010.3805 [hep-ph]].
- [34] S. F. King and C. Luhn, JHEP **09** (2011), 042 [arXiv:1107.5332 [hep-ph]].
- [35] S. F. King and C. Luhn, JHEP **03** (2012), 036 [arXiv:1112.1959 [hep-ph]].
- [36] Y. Shimizu, M. Tanimoto and A. Watanabe, Prog. Theor. Phys. **126** (2011), 81-90 [arXiv:1105.2929 [hep-ph]].
- [37] S. Antusch and V. Maurer, Phys. Rev. D **84** (2011), 117301 [arXiv:1107.3728 [hep-ph]].
- [38] Y. H. Ahn and S. K. Kang, Phys. Rev. D **86** (2012), 093003 [arXiv:1203.4185 [hep-ph]].
- [39] H. Ishimori and E. Ma, Phys. Rev. D **86** (2012), 045030 [arXiv:1205.0075 [hep-ph]].
- [40] W. Rodejohann and H. Zhang, Phys. Rev. D **86** (2012), 093008 [arXiv:1207.1225 [hep-ph]].
- [41] C. Hagedorn, S. F. King and C. Luhn, Phys. Lett. B **717** (2012), 207-213 [arXiv:1205.3114 [hep-ph]].
- [42] S. F. King and C. Luhn, Rept. Prog. Phys. **76** (2013), 056201 [arXiv:1301.1340 [hep-ph]].
- [43] S. F. King, Phys. Lett. B **718** (2012), 136-142 [arXiv:1205.0506 [hep-ph]].
- [44] Y. Shimizu, R. Takahashi and M. Tanimoto, PTEP **2013** (2013) no.6, 063B02 [arXiv:1212.5913 [hep-ph]].
- [45] G. Altarelli and F. Feruglio, Nucl. Phys. B **720** (2005), 64-88 [arXiv:hep-ph/0504165 [hep-ph]].
- [46] G. Altarelli and F. Feruglio, Nucl. Phys. B **741** (2006), 215-235 [arXiv:hep-ph/0512103 [hep-ph]].
- [47] P. Minkowski, Phys. Lett. B **67** (1977), 421-428.
- [48] T. Yanagida, in Proceedings of the Workshop on Unified Theories and Baryon Number in the Universe, eds. O. Sawada and A. Sugamoto (KEK report 79-18, 1979).
- [49] M. Gell-Mann, P. Ramond and R. Slansky, Conf. Proc. C **790927** (1979), 315-321 [arXiv:1306.4669 [hep-th]].
- [50] R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44** (1980), 912.
- [51] J. Schechter and J. W. F. Valle, Phys. Rev. D **22** (1980), 2227.

- [52] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962), 870-880.
- [53] B. Pontecorvo, Zh. Eksp. Teor. Fiz. **53** (1967), 1717-1725.
- [54] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz and A. Zhou, JHEP **09** (2020), 178 [arXiv:2007.14792 [hep-ph]].
- [55] M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, Universe **7** (2021) no.12, 459 [arXiv:2111.03086 [hep-ph]].
- [56] N. Darvishi, M. R. Masouminia and A. Pilaftsis, Phys. Rev. D **104** (2021) no.11, 115017 [arXiv:2106.03159 [hep-ph]].
- [57] N. Aghanim *et al.* [Planck], Astron. Astrophys. **641** (2020), A6 [erratum: Astron. Astrophys. **652** (2021), C4] [arXiv:1807.06209 [astro-ph.CO]].
- [58] R. L. Workman [Particle Data Group], PTEP **2022** (2022), 083C01.
- [59] C. Jarlskog, Phys. Rev. Lett. **55** (1985), 1039.