

GENERALIZING LIQUID DEMOCRACY TO MULTI-AGENT DELEGATION: A VOTING POWER MEASURE AND EQUILIBRIUM ANALYSIS

FRANCISCO M. BERSETCHE

ABSTRACT. In this study, we propose a generalization of the classic model of liquid democracy that allows fractional delegation of voting weight, while simultaneously allowing for the existence of equilibrium states. Our approach empowers agents to partition and delegate their votes to multiple representatives, all while retaining a fraction of the voting power for themselves. We introduce a penalty mechanism for the length of delegation chains. We discuss the desirable properties of a reasonable generalization of the classic model, and prove that smaller penalty factors bring the model closer to satisfying these properties. In the subsequent section, we explore the presence of equilibrium states in a general delegation game utilizing the proposed voting measure. In contrast to the classical model, we demonstrate that this game exhibits pure strategy Nash equilibria, contingent upon the imposition of a penalty on the length of delegation chains.

1. INTRODUCTION

Liquid democracy (LD) is a collective decision making method that lies between representative democracy and direct democracy. In LD, an agent can choose to either vote directly or to select another agent as their proxy. Unlike classic proxy voting, the delegation is transitive; this means that the proxy may delegate their voting weight further, creating delegation paths along which the voting weight flows and accumulates.

Over the past decade, several practical implementations of LD have been conducted [23]. Notable examples include Zupa [22], which was employed by the Student Union of the Faculty of Information Studies in Novo mesto, Slovenia; LiquidFeedback [2], which was used by the German Pirate Party; and Sovereign [24][Section 2], a blockchain-based platform developed and utilized by the Democracy Earth Foundation in Argentina. Of these implementations, the use of LiquidFeedback by the German Pirate Party is the best documented in literature and will be considered as a benchmark for LD practical applications throughout this work

Proxy voting models have been studied since Carroll's work in 1884 [13]. In the last decade, these models have been investigated by political science [18] and artificial intelligence communities [9]. Subsequent research has focused on potential improvements and modifications, as well as possible weaknesses of these models [5, 11, 12, 19, 20], among others. For a comprehensive review of this topic see [23] and the references therein.

Recently, there has been an increasing interest in examining the existence of equilibrium states in these systems. In [16], the authors investigate this matter from a game theory perspective. In this context, they establish that the classical model of liquid democracy generally lacks equilibrium states. Furthermore, determining the existence of an equilibrium state is proven to be NP-complete.

The author was partially supported by ANID through FONDECYT Project 3220254. Part of this work was performed in Centro Atómico Bariloche under a postdoctoral fellowship funded by CONICET.

To address this problem, one may consider a model that allows delegation to multiple agents through the use of mixed strategies. However, while this model ensures the existence of equilibrium states, it becomes evident that it does not preserve desirable properties for a liquid democracy model.

The aim of this work is to provide a model of liquid democracy that ensures the existence of equilibrium states while simultaneously preserving desirable properties in a liquid democracy system. The proposed model also generalizes the single delegation paradigm by allowing delegation to multiple agents.

Contribution. The prevalent approach in liquid democracy implementations and existing literature typically adopts a simple delegation paradigm, where each agent has the option to delegate their entire voting weight to a single agent or cast a direct vote ¹. This paper generalizes this model by allowing agents to divide their voting weight arbitrarily and delegate it to multiple agents, while also retaining part of it for themselves, and enabling the existence of equilibrium states. This generalization is achieved through a novel voting weight measure that preserves the main properties of classic delegation. The introduced measure incorporates a penalty factor based on the length of delegation chains, allowing the existence of equilibrium states. As in most platforms and available works, we consider the voting weight trapped in delegation cycles as null votes (see [1]).

Throughout this work, we posit that a reasonable generalization of the classical LD model should fulfill, at least, the following properties:

- (P1) It generalizes the concept of simple delegation, whereby classic delegation is recovered when agents choose to delegate their voting weight without dividing it.
- (P2) Delegation makes sense. If agent a delegates her voting weight to agent b , the distribution of voting weight is equivalent to what would be obtained if agent a replicated the delegation preference of agent b . This implies that delegating to an agent is equivalent to replicating their actions. Although we are being advisedly vague here, this property is formalized and proven in Theorem 3.7.
- (P3) If an agent chooses to retain a fraction $q \in (0, 1]$ of the vote for herself, then her voting weight should be no less than q . Conversely, if an agent opts to delegate all the power, her voting weight should be rendered null. This property is referred to as *self-selection*.

Note that property (P2) implies that, for an agent, delegating is equivalent to allowing their proxy to make decisions on their behalf. This intuitively corresponds with the expected outcome of delegation. We refer to (P2) as the *delegation property*. We emphasize that the properties (P1)-(P3) are desirable, but do not necessarily provide a comprehensive characterization of a reasonable generalization of the classic LD model. In this work we establish the fulfillment of properties (P1)-(P3) by the proposed voting measure within a defined margin of error determined by the penalty factor.

The temporal variable is a crucial factor to consider in our analysis. As reported in [3], the experience with LiquidFeedback reveals that the behavior of agents changes over time due to their observations of other agents' actions, which is essential for the development of decision-making processes. Therefore, we assume that an agent's delegation preference can be represented as a function that evolves over time and depends on real-time feedback received by the agent. This leads us to ask whether this process can reach an equilibrium state, then in the second part of

¹Sovereign [24] introduces a multi-agent delegation system, granting users a finite number of independently delegatable votes; however, each received vote must be independently dispensed by each proxy.

this work, we introduce a game-theoretic model where agents serve as players and their delegation preferences represent their strategies. Each agent i is endowed with a utility function that measures the amount of their own vote attributed to a specific player j , weighted by a satisfaction factor which accounts for i 's contentment with the proportion of their voting power ceded to j . We demonstrate the existence of Nash equilibria in this game under the condition of non-zero penalties imposed on extended delegation chains. Furthermore, we present the main result of this work, where we establish the possibility of defining a voting power measure such that the fundamental properties of classic delegation are upheld within an acceptable margin of error, while simultaneously attaining pure Nash equilibria for the examined game. To the best of our knowledge, this study offers the first formal proof establishing the feasibility of a liquid democracy system that guarantees the existence of equilibrium states.

Related work. Regarding models that allow multi-agent delegation, we can mention the model proposed in [15], which is similar to the model presented here when the penalty factor is not considered. Spectral type centrality measures have been proposed for measuring voting weight; [25] explores the use of PageRank as a voting measure, while [7] propose another voting measure based on a PageRank-type centrality which seeks to penalize long delegation paths. Although simple delegation is used here for theoretical analysis purposes, these techniques can easily be generalized to multi-agent delegation. On the other hand, following a different approach, in [14], the authors propose and analyze a model based on multi-agent ranked delegation, in which agents provide a ranking over multiple delegations to be used as backups when solving possible delegation cycles. Similarly, [10] and [11] study the delegation of vote through a ranking of preferences, while [17] proposes a technique based on fluid dynamics that attempts to minimize concentration of votes by choosing an appropriate delegation graph according to the agents' preferences.

In Section 4, we define a game in which the delegation preference of each agent is considered as a strategy, and an utility function based on the preferences of each agent over the others. This type of game has been studied in the literature with regard to simple delegation, e.g., [16] which analyzes this kind of games with three different utility functions. It can be seen that the problem addressed in Section 4 is a generalization of the **MINDIS** problem defined in [16, Section 3.2]. Other works dealing with similar topics include [4, 21, 26].

Organization of the paper. The rest of the paper is organized as follows. Section 2 provides an intuitive description of the proposed voting power measure, beginning with its probabilistic interpretation. These ideas are formalized in Section 3, where we define the measure and prove its main properties, as well as provide an efficient algorithm for computation. In Section 4, game theory is employed to investigate the existence of equilibrium states in the context of agents' ability to modify their delegation preference according to system outcomes. Ultimately, our main result is demonstrated, which establishes the feasibility of constructing a voting power measure through imposition of penalties on long delegation chains, that both preserves classic delegation properties and allows for pure strategy Nash equilibria.

2. PRELIMINARIES AND MAIN IDEAS

2.1. Preliminaries. Let $N = \{1, \dots, n\}$ be the set of agents. Each agent i expresses to whom they want to delegate their voting weight, and the fraction they want to delegate, through the delegation preference $x_i \in X$, with $X := \{x \in \mathbb{R}_{\geq 0}^n : \sum_k x_k = 1\}$. Here x_{ij} represents the fraction of voting weight that the agent i wants to delegate to agent j . We define the matrix

$P := (x_1 | \dots | x_n) \in \mathbb{R}^{n \times n}$ with the delegation preferences as columns, and we call it delegation matrix. Clearly P is a (left)stochastic matrix (since $\sum_i P_{ij} = 1$), with the delegation preference of agent j stored in its j -th column. We denote by $\mathcal{P}^n \subset \mathbb{R}^{n \times n}$ the set of all possible delegation matrices. Note that while the delegation preferences from set X and delegation matrices from \mathcal{P}^n , are strongly related, it is convenient to treat these two concepts separately, especially in Section 4, where delegation preferences represent agents' strategies.

Given i and $j \in N$, we say that there is a delegation path between i and j if there is a strictly positive sequence $(x_{ik_1}, x_{k_1 k_2}, \dots, x_{k_{q-1} k_q}, x_{k_q j})$ with $k_1, \dots, k_q \in N$. Given a vector $v \in \mathbb{R}^n$, we define the matrix $diag(v) \in \mathbb{R}^{n \times n}$ as $diag_{ii}(v) := v_i$ for all $1 \leq i \leq n$, and $diag_{ij}(v) := 0$ for all $i \neq j$. As is common in packages designed for numerical methods, if $A \in \mathbb{R}^{n \times n}$, we define $diag(A) \in \mathbb{R}^n$ as $diag_i(A) := A_{ii}$ for all $1 \leq i \leq n$. Unless specified otherwise, throughout this paper vectors will always be regarded as column vectors.

2.2. Description of the problem. Let $N = \{1, \dots, n\}$ denote the set of agents in the classic model of liquid democracy (LD), where agents can delegate their voting power to a single agent, including themselves, and we refer to agents who preserve all their voting power as *candidates*. The delegation matrices obtained under this paradigm correspond to adjacency matrices associated with delegation graphs defined by agent preferences. Defining $\mathcal{A}^n := \{A \in \mathcal{P}^n : A_{ij} \in \{0, 1\} \text{ for all } 1 \leq i, j \leq n\}$, the classic measure of voting power or voting weight is a function $V^c : \mathcal{A}^n \rightarrow \mathbb{R}_{\geq 0}^n$, where $V_i^c(A)$ represents the voting power of agent i under the delegation configuration given by the matrix $A \in \mathcal{A}^n$. In the classic model of LD, the number of votes received by a candidate is determined as the sum of votes received by all agents who delegate their power to the candidate, plus the votes of all agents who delegate to these agents, and so on. Although this procedure can be easily described recursively, we provide an equivalent vectorial definition, which is more manageable in the context of this work. To this end, for a given $A \in \mathcal{A}^n$, we can provide a vectorial definition of $V^c : \mathcal{A}^n \rightarrow \mathbb{R}_{\geq 0}^n$ as follows:

$$(2.1) \quad V^c(A) := \lim_{k \rightarrow \infty} (A^k \mathbf{1}) \odot diag(A),$$

where $\mathbf{1} = (1, \dots, 1) \in \mathbb{R}^n$ and \odot denotes the element wise product. Here c stands for "classic". Given the known properties of adjacency matrices, it can be easily observed that successive products $A^k \mathbf{1}$ flow and accumulate votes from agents to candidates through delegation chains. Note that $diag_i(A) \in \{0, 1\}$ for all i , and $diag_i(A) = 1$ if and only if i is a candidate. Thus, the element-wise product with $diag(A)$ does not affect the entries of $A^k \mathbf{1}$ corresponding to candidates and nullifies the remaining entries. This ensures that the limit in (2.1) is well-defined as the votes trapped in delegation cycles are multiplied by zero. We now consider multi-agent delegation, that is, we aim to extend the function V^c to the entire set \mathcal{P}^n in such a way that this function constitutes a reasonable generalization of the classic LD model. As a first idea one can use the definition (2.1) by considering $A \in \mathcal{P}^n$, since the limit in this expression remains well-defined. This generalization is, in fact, closely related to the Eigenvector Centrality Measure [8], and related in general to spectral centrality measures, as the PageRank used as voting measure in [6, 7, 25]. However, it is possible to see that this generalization does not satisfy the properties stated in the previous section. However, it should be mentioned that if we prohibit agents from preserving a fraction of votes for themselves that is different from 1 or 0, the aforementioned generalization preserves the properties stated in the previous section.

In [16], it is shown that the classic LD model typically lacks equilibrium states in the context of delegation games. However, Nash equilibria can be achieved by incorporating mixed strategies,

suggesting a potential multi-agent delegation model. Unfortunately, this model does not possess the delegation property (P2) (see A.2), making it unsuitable as a generalization of the classic model. Our goal is to establish a measure of multi-agent voting power to ensure equilibrium states while maintaining properties (P1)-(P3). To achieve this, we propose a generalization of the classic model by introducing penalties on the delegation chain length. Specifically, we demonstrate that as the penalty approaches zero, a voting power measure that satisfies properties (P1)-(P3) is attained. However, like the classic model, the acceptance of equilibrium states by this limiting case voting measure is not guaranteed in general (see A.3). Nonetheless, we show that with a positive penalty factor, the model achieves at least one Nash equilibrium in the delegation game. Moreover, properties (P1)-(P3) are upheld within a defined margin of error depending on the penalty factor. These findings are formally presented and summarized in Theorem 4.2, which is our main result.

2.3. The voting weight measure. Our goal is to construct a function $V : \mathcal{P}^n \rightarrow \mathbb{R}^n$ that represents the voting weight of each agent i , which is a reasonable generalization of V^c . In Section 3, we provide a formal definition of the proposed function V , but in this section, we first introduce its probabilistic interpretation. Specifically, we construct V using a particle system that depends on the agents and their delegation preferences, governed by the following rules:

Given a delegation matrix $P \in \mathcal{D}$, within a time interval of length $\delta t > 0$:

- Each agent j receives 1 particle with probability δt .
- A particle in position j has a probability $\delta t P_{ij}$ to jump to position i , with $j \neq i$.
- A particle in position j leaves the system with probability $\delta t P_{jj}$. In such a case we shall say that the particle was consumed by agent j .

We consider the limit as $\delta t \rightarrow 0$, and define $V_i(P)$ as the mean rate of particles consumed by agent i when the system reaches equilibrium.

To compute $V_i(P)$, we introduce u_i as the average number of particles occupying position i in the equilibrium state. We can then estimate $V_i(P)$ as the expected number of particles consumed by agent i per unit time, given by $P_{ii}u_i$. To determine u_i , we set up a set of balance equations for the particle system described in the previous section. Using the rules governing the system, the evolution of the number of particles in position i is given by:

$$u_i(t + \delta t) = u_i(t) + \delta t \left(\sum_{j \neq i} u_j(t) P_{ij} + 1 - u_i(t) \right),$$

$$\frac{u_i(t + \delta t) - u_i(t)}{\delta t} = \sum_{j \neq i} u_j(t) P_{ij} + 1 - u_i(t).$$

Since we look for u_i at the equilibrium state, we have

$$(2.2) \quad 0 = \sum_{j \neq i} u_j P_{ij} + 1 - u_i.$$

The above expression can be written in matrix form:

$$(2.3) \quad \begin{aligned} 0 &= \tilde{P}u - u + \mathbf{1}, \\ u &= (I - \tilde{P})^{-1}\mathbf{1}, \end{aligned}$$

with $\tilde{P} = P - \text{diag}((P_{11}, \dots, P_{nn}))$.

At this point, we can verify that $V(P)$ satisfies some of the properties enlisted in the preceding section. In particular, part of the property (P3) can be readily established, as when an agent i delegates all their voting weight, we get $P_{ii} = 0$, and thus $V_i(P) = P_{ii}u_i = 0$. Furthermore, we can observe that to determine which agents consume a part of the voting weight of a certain agent i , we may use the same particle system defined above, with the constraint that the influx of particles is only through the agent i . That is, we calculate $V(P)$ from $u = (I - \tilde{P})^{-1}\delta_i$, where $\delta_i \in \mathbb{R}^n$, $\delta_{ii} = 1$ and $\delta_{ij} = 0$ if $i \neq j$. Consequently, $V_j(P)$ will quantify the amount of voting weight consumed by agent j , coming from agent i .

2.4. Solving delegation cycles. It is clear that equation (2.3) is not always solvable. This is due to the possible existence of delegation cycles. We formally define a delegation cycle as follows:

Definition 2.1 (Delegation cycle). *Given a set of agents $N = \{1, \dots, n\}$ and their delegation preferences x_1, \dots, x_n , we say that $C \subseteq N$ is a delegation cycle if $x_{ii} = 0$ for all $i \in C$, and given $i \in C$, $x_{ij} = 0$ for all $j \in N \setminus C$.*

Based on this definition, it follows that for a cycle C , any particle that reaches an agent $i \in C$ cannot exit the system. To address this issue, we propose the introduction of an artificial agent $n+1$ into the system, where each particle has a probability $\delta t \varepsilon$ of transitioning to agent $n+1$ from any position within a time interval δt , where $0 < \varepsilon < 1$ and $P_{n+1n+1} = 1$. Specifically, we add agent $n+1$ with delegation preference $x_{n+1} = (0, \dots, 0, 1) \in \mathbb{R}^{n+1}$, and redefine the preferences of the remaining agents as $x_i := ((1 - \varepsilon)x_i, \varepsilon)$, where $i \in N$. It is worth noting that every particle that reaches agent $n+1$ is absorbed. We denote the voting weight measure obtained through this method as V^ε .

Following the modification introduced, delegation cycles no longer pose a challenge, as any particle trapped in a cycle is consumed by agent $n+1$. However, this modification leads to the penalization of extended delegation paths. To illustrate, consider a classic delegation chain of length k between agent i and agent j . For a particle originating from agent i , the probability of exiting the system through agent $n+1$ before reaching agent j is given by $p_\varepsilon^k = 1 - (1 - \varepsilon)^k > 0$. Consequently, the voting weight of agent i dissipates through agent $n+1$, which results in the penalization of long delegation paths². Moreover, V_{n+1}^ε denotes the aggregate of all the voting weight dissipated through cycles and extended delegation paths. Note that $p_\varepsilon^k \rightarrow 0$ as $\varepsilon \rightarrow 0$, yet the voting weight dissipated by cycles remains invariant. To illustrate this, classic delegation cycles can be considered infinite-length delegation chains, and $p_\varepsilon^\infty := \lim_{k \rightarrow \infty} p_\varepsilon^k = 1$, which establishes the independence of ε . In order to eliminate all penalties in long but finite chains, we may define

$$(2.4) \quad V(P) := \lim_{\varepsilon \rightarrow 0} V^\varepsilon(P),$$

where $V_{n+1}(P)$ accumulates the total amount of voting weight lost in delegation cycles.

Remark 1. It is worth mentioning that V can be constructed in a much simpler manner. For instance, one can use a straightforward generalization of the model proposed in [15], or alternatively, a construction based on Markov chains. Starting from these simpler constructions, it can be easily shown that the measure V satisfies properties (P1)-(P3). However, it can be proven that using this measure, in the context of the delegation game defined in Section 4, it is not true that an equilibrium state always exists, similar to what occurs in the classic delegation model (see Section

²This damping factor is similar to the one employed in PageRank computation, and is examined in [6, 7]

A.3). As mentioned earlier, the main contribution of this work focuses on the definition and analysis of the measure V^ε . We will see below that this measure possesses the following properties:

- $V^\varepsilon \rightarrow V$ when $\varepsilon \rightarrow 0$, meaning the limiting case defines a reasonable voting measure,
- V^ε has at least one equilibrium state whenever $\varepsilon > 0$,
- V^ε satisfies properties (P1)-(P3) within a certain margin of error depending on ε .

3. FORMAL FRAMEWORK

Let $N = \{1, \dots, n\}$ be a set of agents with delegation preferences x_1, \dots, x_n , and P their delegation matrix. We define

$$(3.1) \quad P^\varepsilon := \begin{pmatrix} (1-\varepsilon)P & \mathbf{0} \\ \varepsilon & 1 \end{pmatrix},$$

$$u^\varepsilon := (I - \tilde{P}^\varepsilon)^{-1} \mathbf{1}_0,$$

and

$$(3.2) \quad V^\varepsilon(P) := (P_{11}^\varepsilon u_1^\varepsilon, \dots, P_{nn}^\varepsilon u_n^\varepsilon, u_{n+1}^\varepsilon),$$

where $\varepsilon = (\varepsilon, \dots, \varepsilon) \in \mathbb{R}^{1 \times n}$, $\tilde{P}^\varepsilon = P^\varepsilon - \text{diag}((P_{11}^\varepsilon, \dots, P_{nn}^\varepsilon, P_{n+1n+1}^\varepsilon))$ and $\mathbf{1}_0 = (1, \dots, 1, 0) \in \mathbb{R}^{n+1}$. Note that, since $\varepsilon > 0$ we have $\|\tilde{P}^\varepsilon\| < 1$, as a consequence $(I - \tilde{P}^\varepsilon)^{-1}$ exists and $V^\varepsilon(P)$ is well defined.

We define the voting power (or voting weight) $V(P)$ associated to a delegation matrix P as

$$(3.3) \quad V(P) := \lim_{\varepsilon \rightarrow 0} V^\varepsilon(P).$$

3.1. Well-definedness. To establish the existence of the limit in (3.3), it is necessary to introduce the following auxiliary result.

Lemma 3.1. *Let $\{a_k\}_{k \in \mathbb{N}_0}$ and $\{b_k\}_{k \in \mathbb{N}_0} \in \ell^\infty$ be two uniformly bounded sequences, where a_k is periodic with period $q \in \mathbb{N}_0$, and $\sum_k |b_k| < \infty$. Then:*

- (i) $\lim_{\varepsilon \rightarrow 0} \sum_{k \geq 0} \varepsilon (1-\varepsilon)^k b_k = 0$,
- (ii) $\lim_{\varepsilon \rightarrow 0} \sum_{k \geq 0} \varepsilon (1-\varepsilon)^k a_k = \ell$, for some ℓ depending on $\{a_k\}_{k \in \mathbb{N}_0}$.

Proof. Since $\sum_{k \geq 0} |b_k| = c$ for some constant c , we proceed

$$\left| \sum_{k \geq 0} \varepsilon (1-\varepsilon)^k b_k \right| \leq \varepsilon \sum_{k \geq 0} |b_k| = \varepsilon c \rightarrow 0$$

when $\varepsilon \rightarrow 0$. And then (i) is proved.

On the other hand, since $\{a_k\}$ is periodic with period q , there are constants c_j , with $j = 0, \dots, q-1$ such that $a_{qk+j} = c_j$ for all $k \geq 0$. Then, for every fixed j we have

$$\sum_{k \geq 0} \varepsilon (1-\varepsilon)^{qk+j} a_{qk+j} = c_j (1-\varepsilon)^j \sum_{k \geq 0} \varepsilon (1-\varepsilon)^{qk} = \frac{c_j (1-\varepsilon)^j \varepsilon}{1 - (1-\varepsilon)^q}.$$

Observe that the denominator in the right-hand side corresponds to a polynomial $S(\varepsilon)$ of degree q . Additionally, one can readily confirm that $S(0) = 0$ and $S'(0) \neq 0$. Therefore, it follows that $S(\varepsilon) = \varepsilon R(\varepsilon)$, where R denotes a polynomial of degree $q - 1$, and satisfies $R(0) \neq 0$. Then

$$\frac{c_j(1-\varepsilon)^j\varepsilon}{1-(1-\varepsilon)^q} = \frac{c_j(1-\varepsilon)^j}{R(\varepsilon)} \rightarrow \frac{c_j}{R(0)}$$

when $\varepsilon \rightarrow 0$. Therefore, we can infer that $\lim_{\varepsilon \rightarrow 0} \sum_{k \geq 0} \varepsilon(1-\varepsilon)^k a_k = \sum_{j=0}^{q-1} \frac{c_j}{R(0)}$, establishing the proof of (ii). \square

Using the previous lemma, we can prove the main result of this section.

Theorem 3.1. *Let $N = \{1, \dots, n\}$ be a set of agents with delegation preferences x_1, \dots, x_n , and P their delegation matrix. Then the limit in (3.3) is well defined.*

Proof. We have

$$I - \tilde{P}^\varepsilon = \begin{pmatrix} I - (1-\varepsilon)\tilde{P} & \mathbf{0} \\ -\varepsilon & 1 \end{pmatrix},$$

with $\tilde{P} = P - \text{diag}((P_{11}, \dots, P_{nn}))$. For the sake of simplicity we define $A_\varepsilon := I - (1-\varepsilon)\tilde{P}$. Since $\varepsilon > 0$ and P is a stochastic matrix, we have $\|(1-\varepsilon)\tilde{P}\| < 1$. Thus A_ε^{-1} exists, and can be computed using the identity

$$(3.4) \quad A_\varepsilon^{-1} = \sum_{k \geq 0} (1-\varepsilon)^k \tilde{P}^k.$$

Utilizing the blockwise inversion formula yields:

$$(I - \tilde{P}^\varepsilon)^{-1} = \begin{pmatrix} A_\varepsilon & \mathbf{0} \\ -\varepsilon & 1 \end{pmatrix}^{-1} = \begin{pmatrix} A_\varepsilon^{-1} & \mathbf{0} \\ \varepsilon A_\varepsilon^{-1} & 1 \end{pmatrix}.$$

from we can derive

$$u^\varepsilon = \begin{pmatrix} A_\varepsilon^{-1} & \mathbf{0} \\ \varepsilon A_\varepsilon^{-1} & 1 \end{pmatrix} \mathbf{1}_0 = \begin{pmatrix} A_\varepsilon^{-1} \mathbf{1} \\ \varepsilon A_\varepsilon^{-1} \mathbf{1} \end{pmatrix},$$

where $\mathbf{1} = (1, \dots, 1) \in \mathbb{R}^n$.

Now we focus on the computation of $A_\varepsilon^{-1} \mathbf{1}$. Using (3.4) we obtain $A_\varepsilon^{-1} \mathbf{1} = \sum_{k \geq 0} (1-\varepsilon)^k \tilde{P}^k \mathbf{1}$. In order to analyze the powers of \tilde{P} we define:

$$(3.5) \quad N_1 := \{i \in N \text{ such that } P_{ii} \neq 0\},$$

$$(3.6) \quad N_2 := \{i \in N \setminus N_1 \text{ such that exists a delegation path from } i \text{ to } j \in N_1\},$$

$$(3.7) \quad N_3 := N \setminus (N_1 \cup N_2).$$

Given that $N = N_1 \overset{d}{\cup} N_2 \overset{d}{\cup} N_3$, we adopt, without loss of generality, an indexing of the agents such that those belonging to N_1 appear first, those belonging to N_2 appear second, and those belonging to N_3 appear last. From this, we observe that \tilde{P} can be written as

$$(3.8) \quad \tilde{P} = \begin{pmatrix} P_1 & \mathbf{0} \\ P_2 & P_3 \end{pmatrix},$$

with P_1 being a sub-stochastic matrix, and P_3 a stochastic matrix. It is well known that the powers of \tilde{P} can be computed as

$$(3.9) \quad \tilde{P}^k = \begin{pmatrix} P_1^k & \mathbf{0} \\ B_k & P_3^k \end{pmatrix},$$

with $k \in \mathbb{N}$, and $B_k = \sum_{j=0}^{k-1} P_3^j P_2 P_1^{k-j-1}$. Then we have

$$\tilde{P}^k \mathbf{1} = \begin{pmatrix} P_1^k & \mathbf{0} \\ B_k & P_3^k \end{pmatrix} \mathbf{1} = \begin{pmatrix} P_1^k \mathbf{1} \\ B_k \mathbf{1} + P_3^k \mathbf{1} \end{pmatrix},$$

as a consequence

$$(3.10) \quad A_\varepsilon^{-1} \mathbf{1} = \sum_{k \geq 0} (1 - \varepsilon)^k \tilde{P}^k \mathbf{1} = \begin{pmatrix} \sum_{k \geq 0} (1 - \varepsilon)^k P_1^k \mathbf{1} \\ \sum_{k \geq 0} (1 - \varepsilon)^k (B_k \mathbf{1} + P_3^k \mathbf{1}) \end{pmatrix}.$$

Since P_1 is a sub-stochastic matrix, its spectral radius $\rho(P_1) < 1$. Then, each entry of $P_1^k \mathbf{1}$ decays exponentially, and $\lim_{\varepsilon \rightarrow 0} \sum_{k \geq 0} (1 - \varepsilon)^k P_1^k \mathbf{1}$ exists. Note that we do not need to compute $\sum_{k \geq 0} (1 - \varepsilon)^k (B_k \mathbf{1} + P_3^k \mathbf{1})$ since these entries correspond to agents $i \in N_3$ with $P_{ii} = 0$, and then $V_i^\varepsilon(\tilde{P}) = 0$ for all $\varepsilon > 0$.

Now our aim is to compute the limit of $\lim_{\varepsilon \rightarrow 0} \varepsilon A^{-1} \varepsilon \mathbf{1}$. By exploiting the fact that $\varepsilon A^{-1} \varepsilon \mathbf{1} = \mathbf{1} \varepsilon A^{-1} \varepsilon \mathbf{1}$, our focus is solely on studying the behavior of $\varepsilon A^{-1} \varepsilon \mathbf{1}$. Specifically, we obtain

$$\varepsilon A_\varepsilon^{-1} \mathbf{1} = \begin{pmatrix} \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k P_1^k \mathbf{1} \\ \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k (B_k \mathbf{1} + P_3^k \mathbf{1}) \end{pmatrix}.$$

As previously established, the limit of $\lim_{\varepsilon \rightarrow 0} \sum_{k \geq 0} (1 - \varepsilon)^k P_1^k \mathbf{1}$ exists. In this regard, we proceed to examine $\sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k (B_k \mathbf{1} + P_3^k \mathbf{1})$. Notably, the definition of B_k , coupled with the sub-stochasticity of P_1 , enables us to confirm that each element of $B_k \mathbf{1}$ is an exponentially convergent sequence. We define $\lim_{k \rightarrow \infty} [B_k \mathbf{1}]_i =: \ell_i$, where $i \in \{1, \dots, N+1\}$, and further introduce the vector $L \in \mathbb{R}^N$ such that $L_i = \ell_i$. Consequently, each element of the sequence $B_k \mathbf{1} - L$ converges exponentially to zero. This leads to the expression:

$$\begin{aligned} \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k B_k \mathbf{1} &= \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k (B_k \mathbf{1} - L) + \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k L, \\ &= \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k (B_k \mathbf{1} - L) + L, \end{aligned}$$

where in the last step we computed the geometric summation. Since every entry of $B_k \mathbf{1}$ is a positive summable sequence, from Lemma 3.1 we conclude that $\lim_{\varepsilon \rightarrow 0} \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k (B_k \mathbf{1} - L) = 0$, and then

$$\lim_{\varepsilon \rightarrow 0} \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k B_k \mathbf{1} = L.$$

We proceed to compute the expression $\lim_{\varepsilon \rightarrow 0} \sum_{k \geq 0} \varepsilon (1 - \varepsilon)^k P_3^k \mathbf{1}$. It is pertinent to note that the only information available regarding P_3 is that it is a stochastic matrix. It is a well-established fact that the powers of P_3 can be represented as:

$$P_3^k = C_1 + C_2^k + C_3^k,$$

where $C_i = X J_i X^{-1}$, being J the Jordan normal form $P_3 = X J X^{-1}$, with $J = J_1 + J_2 + J_3$. Here, J_1 consists solely of the Jordan block corresponding to the eigenvalue $\lambda = 1$, J_2 encompasses the

Jordan blocks of the eigenvalues λ satisfying $|\lambda| = 1$ and $\lambda \neq 1$, and J_3 comprises the remaining blocks. Since P_3 is a stochastic matrix, the geometric multiplicity of every eigenvalue λ with $|\lambda| = 1$ is 1. Consequently, the sequence $C_2^k \mathbf{1}$ is periodic. Further, as per the definition of J_3 , it follows that the sequence $C_3^k \mathbf{1}$ converges exponentially to $\mathbf{0}$.

From this we have

$$\begin{aligned} \sum_{k \geq 0} \varepsilon(1 - \varepsilon)^k P_3^k \mathbf{1} &= \sum_{k \geq 0} \varepsilon(1 - \varepsilon)^k C_1 \mathbf{1} + \sum_{k \geq 0} \varepsilon(1 - \varepsilon)^k C_2^k \mathbf{1} + \sum_{k \geq 0} \varepsilon(1 - \varepsilon)^k C_3^k \mathbf{1}, \\ &= C_1 \mathbf{1} + \sum_{k \geq 0} \varepsilon(1 - \varepsilon)^k C_2^k \mathbf{1} + \sum_{k \geq 0} \varepsilon(1 - \varepsilon)^k C_3^k \mathbf{1}. \end{aligned}$$

Based on the findings in Lemma 3.1, it is concluded that the limit as ε tends to zero exists for both the second and third terms. Thus, the theorem is proven. \square

3.2. Generalization. Up to this point, we have made the assumption that each agent holds an equal role in the decision-making process. In the context of the particle system introduced in Section 2.3, this assumption is reflected in the distribution of one particle to each agent per unit time. This is formalized in the definition of V , where the vector $\mathbf{1}_0$ is used to construct u^ε in (3.1). To modify the inherent voting weight of each agent, we need only adjust the definition of u^ε by replacing the vector $\mathbf{1}_0$ with an alternative vector $f \in \mathbb{R}^{n+1}$, where f_i represents the inherent voting weight of agent i . We can then generalize the definitions of u^ε , $V^\varepsilon(P)$, and $V^\varepsilon(P)$ as follows: Let $N = \{1, \dots, n\}$ denote a set of agents with delegation preferences x_1, \dots, x_n , P their delegation matrix, and $f \in \mathbb{R}^{n+1}$. We define

$$(3.11) \quad u^\varepsilon := (I - \tilde{P}^\varepsilon)^{-1} f,$$

$$(3.12) \quad V^\varepsilon(P, f) := (P_{11}^\varepsilon u_1^\varepsilon, \dots, P_{nn}^\varepsilon u_n^\varepsilon, u_{n+1}^\varepsilon),$$

and

$$(3.13) \quad V(P, f) := \lim_{\varepsilon \rightarrow 0} V^\varepsilon(P, f).$$

It is noteworthy that the proof of Theorem (3.1) can be straightforwardly adapted to encompass the extended definitions. Therefore, we obtain:

Theorem 3.2. *Let $N = \{1, \dots, n\}$ be a set of agents with delegation preferences x_1, \dots, x_n , P their delegation matrix, and $f \in \mathbb{R}^{n+1}$. Then the limit in definition (3.13) is well defined.*

3.3. Self-selection. The following lemma asserts that a positive value of the term f implies a positive voting weight for each agent.

Lemma 3.2. *Let $N = \{1, \dots, n\}$ be a set of agents with delegation preferences $x_1, \dots, x_n \in X$, P their delegation matrix, and $f \in \mathbb{R}^{n+1}$ with $f_i \geq 0$ for all $i \in N$. Then $V_i^\varepsilon(P, f) \geq 0$ for all $i \in N$, for all $\varepsilon \in [0, 1)$.*

Proof. Given $\varepsilon \in (0, 1)$, consider u^ε as in (3.11). Define $u^m = \min_{1 \leq i \leq n} u_i^\varepsilon$, and observe that from (3.11) we can estimate

$$\begin{aligned} u^m &= f_k + \tilde{P}_{k1}^\varepsilon u_1 + \dots + \tilde{P}_{kk-1}^\varepsilon u_{k-1} + \tilde{P}_{kk+1}^\varepsilon u_{k+1} + \dots + \tilde{P}_{kn}^\varepsilon u_n \\ &\geq f_k + \tilde{P}_{k1}^\varepsilon u^m + \dots + \tilde{P}_{kk-1}^\varepsilon u^m + \tilde{P}_{kk+1}^\varepsilon u^m + \dots + \tilde{P}_{kn}^\varepsilon u^m, \\ u^m &\geq \frac{f_k}{1 - \tilde{P}_{k1}^\varepsilon - \dots - \tilde{P}_{kk-1}^\varepsilon - \tilde{P}_{kk+1}^\varepsilon - \dots - \tilde{P}_{kn}^\varepsilon} \geq 0, \end{aligned}$$

for some $k \in N$ such that $u_k = u^m$. Henceforth, it is established that $u_i^\varepsilon \geq 0$ for all $i \in N$. As a consequence, it can be deduced that $V_i^\varepsilon(P, f) \geq 0$ holds true for all $i \in N$ and for all $\varepsilon \in (0, 1)$. Furthermore, the scenario $\varepsilon = 0$ can be trivially derived by taking the limit as ε approaches 0. \square

The application of the aforementioned result allows us to establish the generalized version of the self-selection property (P3). The subsequent theorem states that an agent's voting weight is zero if they choose to delegate their entire voting power, whereas if they retain a fraction of their voting power for themselves, then their voting weight is no less than their inherent voting weight scaled by this fraction. The theorem is stated as follows.

Theorem 3.3 (Self-selection). *Let $N = \{1, \dots, n\}$ be a set of agents with delegation preferences $x_1, \dots, x_n \in X$, P their delegation matrix, and $f \in \mathbb{R}^{n+1}$ with $f_i \geq 0$ for all $i \in N$. Then we have that $P_{ii} = 0$ implies $V_i^\varepsilon(P, f) = 0$, and $V_i^\varepsilon(P, f) \geq (1 - \varepsilon)P_{ii}f_i$ for all $i \in N$ and $\varepsilon \in [0, 1)$.*

Proof. Given that $V_i^\varepsilon(P, f) = P_{ii}u_i^\varepsilon$, it can be readily verified that $P_{ii} = 0$ implies $V_i^\varepsilon(P, f) = 0$ for all $1 \leq i \leq n$ and all $\varepsilon \in (0, 1)$. Moreover, from Lemma 3.2, which establishes that $u_i^\varepsilon \geq 0$ for all $i \in N$, since $f_i \geq 0$ for all $i \in N$, we can estimate using (3.11) that:

$$u_i^\varepsilon = f_i + \tilde{P}_{i1}^\varepsilon u_1 + \dots + \tilde{P}_{in}^\varepsilon u_n \geq f_i,$$

As a result, we obtain $V_i^\varepsilon(P, f) = P_{ii}u_i^\varepsilon \geq (1 - \varepsilon)P_{ii}f_i$ for all $i \in N$, from which we can deduce the desired outcome for all $\varepsilon \in (0, 1)$. The scenario $\varepsilon = 0$ can be easily handled by means of the aforementioned derivation, as we take the limit $\varepsilon \rightarrow 0$. \square

3.4. Conservation of voting power. The preservation of voting power is a desired property in the generalization of the classic LD model, which requires that the sum of the voting power of each agent and the lost voting power in delegation cycles equals the number of agents. Although this property is usually achieved through the normalization of the voting weight vector, it is demonstrated in our case that this step is unnecessary and, therefore, preserves the interpretation of the proposed voting measure discussed in Section 2. The proof of this property is presented in the following result. This property is formalized in the following result, and its proof is given in Appendix A.

Theorem 3.4. *Let $N = \{1, \dots, n\}$ be a set of agents with delegation preferences $x_1, \dots, x_n \in X$, P their delegation matrix, and $f \in \mathbb{R}^{n+1}$. Then $\sum_{i=1}^{n+1} V_i^\varepsilon(P, f) = \sum_{i=1}^{n+1} f_i$ for all $\varepsilon \in [0, 1)$.*

3.5. Numerical computation. The computation of V^ε requires the inverse of a badly conditioned matrix when ε is small. However, from the proof of Theorem 3.1, it is possible to obtain a procedure for computing V that avoids the need for computing V^ε . Specifically, (3.8) and (3.10) imply that only the inverse of $(I - P_1)$ needs to be computed, where P_1 is a sub-stochastic matrix. This is due to the fact that the voting weight of agents in N_3 (and N_2) is zero, and therefore, it suffices to obtain the voting weight of agents in N_1 . This observation is summarized in Algorithm 1.

Algorithm 1 Computation of $V(P, f)$

Input: A set of agents $N = \{1, \dots, n\}$, their delegation preferences x_1, \dots, x_n , and a vector $f \in \mathbb{R}^{n+1}$;

1: Define the delegation matrix P using the preferences x_1, \dots, x_n , and N_1 and N_2 as in (3.5) and (3.6) respectively. Define I_i as the i -th element of $N_1 \cup N_2$, with $1 \leq i \leq \#(N_1 \cup N_2)$;

2: Define $\tilde{P} := P - \text{diag}((P_{11}, \dots, P_{nn}))$, and \tilde{P}_r as the restriction of \tilde{P} to $N_1 \cup N_2$. That is, remove from \tilde{P} the i -th row and the i -th column, for all $i \notin N_1 \cup N_2$. Analogously, define f_r removing the i -th entry of f for all $i \notin N_1 \cup N_2$.

3: If $\tilde{P}_r \neq \emptyset$, compute $u_r = (I - \tilde{P}_r)^{-1} f_r$;

4: Compute $V_{I_i}(P, f) = P_{I_i I_i} [u_r]_i$ for all $1 \leq i \leq \#(N_1 \cup N_2)$, and $V_i(P, f) = 0$ for all $i \notin N_1 \cup N_2 \cup \{n+1\}$. Finally, using Theorem 3.4, compute $V_{n+1}(P, f) = \sum_{i=1}^{n+1} f_i - \sum_{i=1}^n V_i(P, f)$;

3.6. An alternative definition of V . As we mentioned before, V can be defined directly without relying on the functions V^ε . Algorithm 1 provides a way to compute V directly. It is worth noting that this procedure removes agents belonging to delegation cycles from the system to ensure that the matrix $I - P_1$ is invertible. However, considering that entries of V that correspond to agents belonging to cycles are multiplied by zero in the last step, V can be defined in terms of powers of the delegation matrix. Specifically, let $P \in \mathcal{P}^n$ and $f \in \mathbb{R}^n$, then $V : \mathcal{P}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ can be defined as:

$$(3.14) \quad V(P, f) := \lim_{k \rightarrow \infty} \left(\sum_{\ell=0}^k \tilde{P}^\ell f \right) \odot \text{diag}(P),$$

where $\tilde{P} = P - \text{diag}((P_{11}, \dots, P_{nn}))$. It should be noted that the value associated with node $n+1$ may be computed utilizing Theorem 3.4, as stipulated in step 4 of Algorithm 1.

3.7. The generalization property. Expanding upon the previous section's alternative definition of V , we establish that this voting measure is a generalization of the classic LD model, as it satisfies the property (P1) outlined in Section 1. Moreover, the following result demonstrates that the proposed measure V and the standard generalization of the classic model defined in (2.1) are equivalent, provided that agents are unable to retain a portion $q \in (0, 1)$ of their vote. Its proof is given in Appendix A.

Theorem 3.5. *Let $\mathcal{B}^n := \{B \in \mathcal{P}^n : B_{ii} \in \{0, 1\} \text{ for all } 1 \leq i \leq n\}$, and let V and V^c the voting measures defined in (3.3), and (2.1) respectively. Then $V_i(P) = V_i^c(P)$ for all $P \in \mathcal{B}^n$, and $i \in N$.*

Analogous results hold for V^ε , which establish that this voting power measure can be interpreted as a generalization of the classic LD model with a specific margin of error, subject to the penalty parameter ε . As before, its proof is provided in Appendix A.

Theorem 3.6. *Let $\mathcal{A}^n := \{A \in \mathcal{P}^n : A_{ij} \in \{0, 1\} \text{ for all } 1 \leq i, j \leq n\}$, and let V^ε and V^c the voting measures defined in (3.2), and (2.1) respectively. Given $\delta > 0$ there exist $\varepsilon_\delta > 0$ such that if $\varepsilon < \varepsilon_\delta$ then $\max_{1 \leq i \leq n} |V_i^\varepsilon(P) - V_i^c(P)| < \delta$ for all $P \in \mathcal{A}^n$.*

3.8. The delegation property. This section presents the formalization and proof of the delegation property (P2), which establishes the legitimacy of delegation. Specifically, the property posits that if agent i delegates her voting weight to agent j , who subsequently delegates all of her voting

weight to other agents, the resulting weight distribution would be equivalent to that obtained if agent i had utilized delegation preference x_j instead of x_i . Furthermore, when agent i assigns, let say, $\frac{1}{4}$ and $\frac{3}{4}$ of her voting weight to agents j and k , respectively, and these proxies also delegate all their voting weight, the corresponding voting weight distribution would match that produced by the delegation preference $\frac{1}{4}x_j + \frac{3}{4}x_k$ instead of x_i . Our examples assume that agents j and k do not delegate a portion of their voting weight back to agent i . However, in such cases, the fraction of the voting weight is distributed proportionally among the proxies. We formally state this observation in the following result:

Theorem 3.7 (Delegation property). *Consider a set $N = \{1, \dots, n\}$ of agents with delegation preferences $x_1, \dots, x_n \in X$, their delegation matrix P , and $f \in \mathbb{R}^{n+1}$. Let $D \subset N$, such that $x_{ii} = 0$ for all $i \in D$, and suppose that there exists an agent $k \in N \setminus D$ such that $x_{kj} > 0$ for all $j \in D$, and $x_{kj} = 0$ otherwise. Moreover, assume that agent k is not involved in a delegation cycle. For each $i \in D$, we define $x_i^* \in \mathbb{R}^n$ as $x_{ij}^* := x_{ij}$ for all $j \neq k$ and $x_{ik}^* := 0$, and $x_k^* \in X$ as*

$$(3.15) \quad x_k^* := \frac{\sum_{i \in D} x_{ki} x_i^*}{1 - \sum_{i \in D} x_{ki} x_{ik}};$$

We also define $P^* \in \mathbb{R}^{n \times n}$ as $P^* := (x_1 | \dots | x_{k-1} | x_k^* | \dots | x_n)$, which is the delegation matrix obtained by replacing x_k with x_k^* in P . Then, we have $V(P, f) = V(P^*, f)$.

Proof. We begin by verifying the well-definedness of (3.15), that is, $1 - \sum_{i \in D} x_{ki} x_{ik} > 0$. To establish this, we assume the contrary, i.e., $\sum_{i \in D} x_{ki} x_{ik} = 1$. Since $0 \leq x_{ik}, x_{ki} \leq 1$ for all $i \in N$, and $\sum_{i \in D} x_{ki} = 1$, it follows that $x_{ik} = 1$ for all $i \in D$. Subsequently, since $x_{kj} = 0$ for all $j \notin D$, it implies that $D \cup k$ forms a delegation cycle, which contradicts our assumption that agent k is not part of a cycle.

To apply the procedure detailed in Section 3.5, we introduce the sets N_1 , N_2 , and N_3 as specified in (3.5), (3.6), and (3.7), respectively. Similarly, we define N_1^* , N_2^* , and N_3^* using the delegation preference x_k^* instead of x_k . We assume that the agents in N are arranged in a way that agents in N_1 appear first, followed by agents in N_2 , then agents in N_3 , and finally $k = \#(N_1 \cup N_2)$, for simplicity. Note that an agent $i \in N$ belongs to a cycle if and only if $i > k$. Let us define $D' := D \cap (N_1 \cup N_2)$, and observe that D' cannot be empty, otherwise $D \cup k$ would form a cycle, which contradicts our assumption.

We now establish that $N_3^* = N_3$. Firstly, since agent k is not part of any cycle, changing her delegation preference by x_k^* cannot resolve any cycle. Therefore, $N_3 \subseteq N_3^*$. On the other hand, since $D' \neq \emptyset$, there exists an agent $q \in D'$ such that $x_{qs} > 0$, where $s \in (N_1 \cup N_2)$. By definition (3.15), it follows that $x_{ks}^* > 0$. Thus, agent k does not belong to N_3^* . Consequently, no new delegation cycle is formed when we replace x_k with x_k^* , and we can deduce that $N_3^* \subseteq N_3$.

The equality $N_3 = N_3^*$ implies that $N_1 = N_1^*$ and $N_2 = N_2^*$. Consequently, Algorithm 1 can be applied to P and P^* , and it can be observed that their restricted matrices, denoted by \tilde{P}_r and \tilde{P}_r^* respectively, share the same dimension. Specifically, we have $\tilde{P}_r, \tilde{P}_r^* \in \mathbb{R}^{k \times k}$. To determine $V(P, f)$ and $V(P^*, f)$, we need to solve the linear systems:

$$(I - \tilde{P}_r)u = f_r,$$

and

$$(I - \tilde{P}_r^*)u^* = f_r$$

respectively, where f_r is the restricted source term defined in Algorithm 1. It suffices to show that $u_i = u_i^*$ holds for all $i \in N_1$, given that $V_i(P, f) = V_i(P^*, f) = 0$ for all $i \in N_2 \cup N_3$. To this end, we introduce the matrix $A := (I - \tilde{P}_r)$, and manipulate its columns. By changing variables according to these operations, we define the variable u' as follows:

$$(3.16) \quad \begin{cases} u'_i = u_i - x_{ki}u_k & \text{for all } i \in D', \\ u'_k = u_k(1 - \sum_{i \in D'} x_{ki}x_{ik}), \\ u'_i = u_i & \text{for all } i \notin D' \cup \{k\}. \end{cases}$$

From this, it can be seen that $u' \in \mathbb{R}^k$ solves the linear system

$$\left(A_1 \mid \dots \mid A_{k-1} \mid \frac{A_k + \sum_{i \in D'} x_{ki}A_i}{(1 - \sum_{i \in D'} x_{ki}x_{ik})} \right) u' = f_r,$$

where A_i denotes the i -th column of A . Now, we observe that $A_i = I_i - x_i$, then

$$(3.17) \quad A_{ki} - x_{ki}A_{ii} = x_{ki} - x_{ki} - x_{ki}x_{ii} = -x_{ki}x_{ii},$$

and

$$(3.18) \quad A_{kk} - x_{ki}A_{kk} = 1 - x_{ki}x_{ik}.$$

Then we have

$$(3.19) \quad A_k + \sum_{i \in D'} x_{ki}A_i = (1 - \sum_{i \in D'} x_{ki}x_{ik})I_k - \sum_{i \in D} x_{ki}x_i^*,$$

Based on the dependence of \tilde{P}_r^* construction solely on x_{ki}^* values for $i \leq k$, that are determined by x_i for $i \in D' \cup k$, we can infer that

$$(I - \tilde{P}_r^*) = \left(A_1 \mid \dots \mid A_{k-1} \mid \frac{A_k + \sum_{i \in D'} x_{ki}A_i}{(1 - \sum_{i \in D'} x_{ki}x_{ik})} \right).$$

In conclusion, we establish the equality $u' = u^*$ and, by virtue of the expression given in (3.16), deduce that $u_i = u'_i = u_i^*$ holds for every $i \in N_1$. Thus, the theorem is demonstrated. \square

The definition of the voting measure V^ε includes a penalty on the length of delegation chains, which causes these measures to violate the delegation property. We observe that replicating an agent's delegation preference results in a configuration with shorter delegation chains, which in turn applies less penalty and produces a different distribution of voting weights. Nevertheless, we demonstrate that V^ε conforms to the delegation property within a certain margin of error that relies on the penalty ε . Specifically, a reduction in the penalty leads to a decrease in the error. We establish and verify this property, denoted as the δ -Delegation property, in the following theorem.

Theorem 3.8 (δ -Delegation property). *Assuming the identical hypotheses as stipulated in Theorem 3.7, and $f \in \mathbb{R}_{\geq 0}^{n+1}$. Given $\delta > 0$, there exists a positive ε_δ such that if $\varepsilon < \varepsilon_\delta$ then $\|V^\varepsilon(P, f) - V^\varepsilon(P^*, f)\|_\infty < \delta$.*

Proof. Define

$$\begin{aligned} A &:= (I - \tilde{P}^\varepsilon), \\ A^* &:= (I - \tilde{P}^{*\varepsilon}), \\ u &:= A^{-1}f, \end{aligned}$$

and

$$u^* := A^{*-1}f.$$

Operating with the columns of the matrix A and making the appropriate change of variables as in the proof of Theorem 3.7, we introduce a new variable, denoted by $u' \in \mathbb{R}^{n+1}$, which is defined as follows:

$$(3.20) \quad \begin{cases} u'_i = u_i - (1 - \varepsilon)x_{ki}u_k & \text{for all } i \in D, \\ u'_k = u_k(1 - \sum_{i \in D} (1 - \varepsilon)^2 x_{ki}x_{ik}), \\ u'_i = u_i & \text{for all } i \notin D \cup \{k\}, \end{cases}$$

and fulfills the equation $A'u' = f$, with

$$A' := \left(A_1 \mid \dots \mid A_{k-1} \mid \frac{A_k + \sum_{i \in D'} (1 - \varepsilon)x_{ki}A_i}{(1 - \sum_{i \in D'} (1 - \varepsilon)^2 x_{ki}x_{ik})} \mid A_{k+1} \mid \dots \mid A_{n+1} \right).$$

Let us define $\Delta := A' - A^*$. From this definition, it follows that $A^*u' = f - \Delta u'$. By combining this expression with the definition of u^* , we arrive at:

$$(3.21) \quad A^*(u^* - u') = \Delta u'.$$

By invoking the definition of A^* and following a similar approach as carried out in equations (3.17), (3.18), and (3.19), and denoting Δ_i as the i -th column of Δ , it can be established that $\Delta_i = \mathbf{0}$ for all $i \neq k$ and

$$(3.22) \quad \Delta_k = \begin{pmatrix} 0 \\ (1 - \varepsilon) \frac{\sum_{i \in D} x_{ki}x_{i2}^*}{1 - \sum_{i \in D} x_{ki}x_{ik}} - (1 - \varepsilon)^2 \frac{\sum_{i \in D} x_{ki}x_{i2}^*}{1 - (1 - \varepsilon)^2 \sum_{i \in D} x_{ki}x_{ik}} \\ \vdots \\ (1 - \varepsilon) \frac{\sum_{i \in D} x_{ki}x_{in}^*}{1 - \sum_{i \in D} x_{ki}x_{ik}} - (1 - \varepsilon)^2 \frac{\sum_{i \in D} x_{ki}x_{in}^*}{1 - (1 - \varepsilon)^2 \sum_{i \in D} x_{ki}x_{ik}} \\ (1 - \varepsilon)\varepsilon \end{pmatrix},$$

from we have $\Delta u' = u'_k \Delta_k$. Upon utilizing Theorem 3.4 and considering the constraint that $f \in \mathbb{R}_{\geq 0}$, a direct deduction leads to the following estimate:

$$(3.23) \quad 0 \leq u'_k \leq \sum_i f_i.$$

On the other hand, it is possible to verify that $\Delta_k \in \mathbb{R}_{\geq 0}^n$ and $\Delta_k \rightarrow \mathbf{0}$ as $\varepsilon \rightarrow 0$. This is evident from (3.22), where it can be observed that $\Delta_{k1} = 0$, $\Delta_{kn+1} \geq 0$, and $\Delta_{kn+1} \rightarrow 0$ as $\varepsilon \rightarrow 0$. It can also be easily established that $\Delta_{kj} \geq 0$ for all $2 \leq j \leq n+1$, by making use of (3.22) and the facts that $\sum_{i \in D} x_{ki}x_{ij}^* \geq 0$ and $\sum_{i \in D} x_{ki}x_{ik} \geq 0$. Consequently, we can derive an estimate as follows:

$$(3.24) \quad 0 \leq \Delta_{kj} = (1 - \varepsilon) \frac{\sum_{i \in D} x_{ki}x_{ij}^*}{1 - \sum_{i \in D} x_{ki}x_{ik}} - (1 - \varepsilon)^2 \frac{\sum_{i \in D} x_{ki}x_{ij}^*}{1 - (1 - \varepsilon)^2 \sum_{i \in D} x_{ki}x_{ik}} \leq \\ (1 - \varepsilon) \frac{\sum_{i \in D} x_{ki}x_{ij}^*}{1 - \sum_{i \in D} x_{ki}x_{ik}} - (1 - \varepsilon)^2 \frac{\sum_{i \in D} x_{ki}x_{ij}^*}{1 - \sum_{i \in D} x_{ki}x_{ik}} = \varepsilon(1 - \varepsilon) \frac{\sum_{i \in D} x_{ki}x_{ij}^*}{1 - \sum_{i \in D} x_{ki}x_{ik}} \leq \\ \frac{\varepsilon}{1 - \sum_{i \in D} x_{ki}x_{ik}}.$$

Now, by applying Theorem 3.4 to (3.21) and using the estimates (3.23) and (3.24), we obtain:

$$0 \leq (u^* - u')_i \leq \sum_{1 \leq j \leq n+1} u'_k \Delta_{kj} \leq \left(\sum_{1 \leq j \leq n+1} f_j \right) \left(\sum_{1 \leq j \leq n+1} \Delta_{kj} \right) \leq \varepsilon \frac{(n+1) \sum_{1 \leq j \leq n+1} f_j}{1 - \sum_{j \in D} x_{kj} x_{jk}}.$$

Defining the constant

$$(3.25) \quad C(P, f, n, k) := \frac{(n+1) \sum_{1 \leq j \leq n+1} f_j}{1 - \sum_{j \in D} x_{kj} x_{jk}},$$

and upon observing that (3.20) entails $u'_i = u_i$ for all $i \notin D \cup \{k\}$, one can establish that $\|V^\varepsilon(P, f) - V^\varepsilon(P^*, f)\|_\infty < C(P, f, n, k)\varepsilon$. Taking $\varepsilon_\delta = \delta/C$ we conclude the proof. \square

Remark 2. Assuming fixed parameters n and $f \in \mathbb{R}^{n+1}$, the dependence of ε_δ on P and k is apparent from (3.25). However, the dependence on k can be eliminated as follows: given $P \in \mathcal{P}^n$, let $K_P \subset N$ be the set of agents satisfying the conditions required for agent k as stated in Theorem 3.7. We define

$$C'(P) := \max_{k \in K_P} C(P, f, n, k) = \frac{(n+1) \sum_{1 \leq j \leq n+1} f_j}{\min_{k \in K_P} 1 - \sum_{j \in D} x_{kj} x_{jk}},$$

and set $\varepsilon_\delta = \delta/C'$. On the other hand, the dependence on P cannot be easily disregarded since the constant C' deteriorates as $\min_{k \in K_P} 1 - \sum_{j \in D} x_{kj} x_{jk}$ approaches zero. However, it can be deduced that, for a fixed value of δ , taking $\varepsilon < \varepsilon_\delta$ implies that $\|V^\varepsilon(P, f) - V^\varepsilon(P^*, f)\|_\infty < \delta$ holds for all $P \in \{P \in \mathcal{P}^n \text{ such that } C'(P) > \varepsilon/\delta\}$.

4. EQUILIBRIUM ANALYSIS

4.1. The Delegation Game. In this paper, the decision-making process is examined as a continuous dynamic over time, rather than as a single event, as previously stated. The actions taken by agents, whether it be delegating voting weight or making use of it, have an impact on other agents. As a result, feedback is received by agents, who then adjust their delegation preferences and actions accordingly. This raises the question of whether or not equilibrium states exist for this type of mechanics.

To partially model this dynamic, we introduce the game $\mathcal{G}(N, W, \varepsilon)$, wherein $N = \{1, \dots, n\}$ represents the set of agents, and $W = \{w_1, \dots, w_n\}$, with $w_i \in \mathbb{R}^{n+1}$ denotes the set of preferences. Here, $w_i \in W$ assigns to each agent a weight, which signifies the preferences of agent i . Specifically, w_{ij} reflects the level of satisfaction of agent i with the consumption of her voting weight by j (note that w_i may contain negative entries). The strategy of each agent i is given by its delegation preference $x_i \in X$. The goal of each agent is to maximize its utility function $U_i : X \rightarrow \mathbb{R}$, given by

$$U_i(x, x_{-i}) := \sum_{j=1}^{n+1} w_{ij} V_j^\varepsilon((x|x_{-i}), \delta_i) = w_i \cdot V^\varepsilon((x|x_{-i}), \delta_i).$$

Here $\delta_i \in \mathbb{R}^{n+1}$ is such that $\delta_{ii} = 1$ and $\delta_{ik} = 0$ if $i \neq k$, and $(x|x_{-i})$ denotes the matrix $(x_1, \dots, x, \dots, x_n)$ with column x in the i -th place, where x_k is the delegation preference of agent k with $k \neq i$. We take $0 < \varepsilon < 1$. As illustrated in Section 2.3, the value function $V_j^\varepsilon(P, \delta_i) \in \mathbb{R}$ quantifies the amount of voting weight consumed by agent j that comes from agent i . Hence, U_i can be interpreted as the sum of the voting weight delegated by agent i to other agents, weighted in accordance with her preferences.

Through conventional arguments, one can establish that the game $\mathcal{G}(N, W, \varepsilon)$ always possesses at least one Nash equilibrium. To achieve this, we begin by proving the following lemma.

Lemma 4.1. *Let $N = \{1, \dots, n\}$ be a set of agents with delegation preferences $x_1, \dots, x_n \in X$, P their delegation matrix, and $f \in \mathbb{R}_{\geq 0}^{n+1}$ in such a way that $f_j > 0$ for some $j \in N$. Then $u_j^\varepsilon > 0$ for all $\varepsilon > 0$.*

Proof. From the definition of u^ε we have:

$$(4.1) \quad \begin{pmatrix} I - (1 - \varepsilon)\tilde{P} & \mathbf{0} \\ -\varepsilon & 1 \end{pmatrix} u^\varepsilon = f,$$

Since $f_i \geq 0$ for all i , we know from Lemma 3.2 that $u^\varepsilon \geq 0$. From (4.1) we have

$$u_j^\varepsilon - (1 - \varepsilon) \sum_{i \neq j} P_{ji} u_i^\varepsilon = f_j > 0, \quad \text{if } j < n + 1.$$

It follows that u_j^ε is greater than zero for all $j < n + 1$. The scenario where $j = n + 1$ can be dealt with in a similar manner. \square

The subsequent result demonstrates that $\mathcal{G}(N, W, \varepsilon)$ possesses at least one Nash equilibrium for all $0 < \varepsilon < 1$.

Theorem 4.1. *Let $N = \{1, \dots, n\}$ be a set of agents with preferences $W = \{w_1, \dots, w_n\} \subset \mathbb{R}^{n+1}$, and $f \in \mathbb{R}^{n+1}$. The game $\mathcal{G}(N, W, \varepsilon)$ has a Nash equilibrium for all $0 < \varepsilon < 1$.*

Proof. For the sake of convenience, we shall treat the elements of W as row vectors, that is to say $W \subset \mathbb{R}^{1 \times n+1}$. Let $r_i(x_{-i})$ denote the optimal responses of agent i to the strategies employed by all other agents:

$$r_i(x_{-i}) := \arg \max_{x_i} U_i(x_i, x_{-i}).$$

Let x be an element of $\mathbf{X} := X^n = X \times \dots \times X$, and let the set valued function $r : \mathbf{X} \rightarrow 2^{\mathbf{X}}$ be defined as $r(x) := r_1(x_{-1}) \times \dots \times r_n(x_{-n})$. It is noteworthy that any stationary point of r constitutes a Nash equilibrium. Our aim is to demonstrate the existence of a stationary point of r . To this end, we will employ Kakutani's fixed point theorem, which ensures the existence of a fixed point given the fulfillment of the following conditions:

- (1) \mathbf{X} is compact, convex, and nonempty,
- (2) $r(x)$ is nonempty,
- (3) $r(x)$ is upper hemicontinuous,
- (4) $r(x)$ is convex.

Condition 1 is easily fulfilled. Conditions 2 and 3 can be verified by exploiting the continuous nature of the function U_i for all i , and for all $0 < \varepsilon < 1$, combined with Berge's maximum theorem.

In order to establish 4, it suffices to demonstrate that $r_i(x_{-i})$ is convex for all i , based on the definition of $r(x)$. Let x and $x' \in r_i(x_{-i})$ be two strategies. Let $\eta := x' - x$ and $x(t) := x + t\eta$. Our objective is to show that $x(t) \in r_i(x_{-i})$ for all $t \in [0, 1]$. Without loss of generality and for the sake of simplicity, we assume that $i = 1$. Referring to the definition given in (3.2), if we define $P(t) := (x(t)|_{x_{-1}})$, the utility function U_1 can be expressed as:

$$U_1(x(t), x_{-1}) = w_1 D_t u(t),$$

where $D_t := \text{diag}((P_{11}^\varepsilon(t), \dots, P_{nn}^\varepsilon(t), P_{n+1n+1}^\varepsilon(t)))$, and $u(t) := (I - \tilde{P}^\varepsilon(t))^{-1} \delta_1$. We observe that $u(t)$ satisfies the equation

$$\begin{pmatrix} I - (1 - \varepsilon)\tilde{P}(t) & \mathbf{0} \\ -\varepsilon & 1 \end{pmatrix} u(t) = \delta_1,$$

with $\tilde{P}(t) = P(t) - \text{diag}((P_{11}(t), \dots, P_{nn}(t)))$. Using the definition of $x(t)$, we can rewrite the former expression as

$$\left[\begin{pmatrix} I - (1 - \varepsilon)\tilde{P}(0) & \mathbf{0} \\ -\varepsilon & 1 \end{pmatrix} - \begin{pmatrix} (1 - \varepsilon)t\tilde{\eta} & \mathbf{0} \\ \mathbf{0} & 0 \end{pmatrix} \right] u(t) = \delta_1,$$

where $\tilde{\eta} \in \mathbb{R}^n$, with $\tilde{\eta}_i := \eta_i$ for all $i > 1$ and $\tilde{\eta}_1 := 0$. Defining

$$(4.2) \quad A_0 := \begin{pmatrix} I - (1 - \varepsilon)\tilde{P}(0) & \mathbf{0} \\ -\varepsilon & 1 \end{pmatrix} \text{ and } \Delta := \begin{pmatrix} (1 - \varepsilon)\tilde{\eta} & \mathbf{0} \\ \mathbf{0} & 0 \end{pmatrix},$$

we have $(A_0 - t\Delta)u(t) = \delta_1$, and then

$$(4.3) \quad u(t) = A_0^{-1} \delta_1 + tA_0^{-1} \Delta u(t).$$

Also, we observe that $D_t = D_0 + tD_\Delta$, with

$$(4.4) \quad D_\Delta := \begin{pmatrix} x'_1 - x_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix}.$$

On the other hand, let us consider a constant vector $c \in \mathbb{R}^{1 \times n+1}$, with $c_j = C \in \mathbb{R}$ for all j . If we define $w^* = w_1 + c$, then it can be observed that maximizing (or minimizing) the utility function $w_1 D_t u(t)$ is equivalent to maximizing (or minimizing) the utility function $U(x(t)|x_{-1}) := w_t^D u(t)$. In fact, we have $w^* D_t u(t) = w_1 D_t u(t) + c D_t u(t)$, and

$$c D_t u(t) = c \cdot V^\varepsilon((x(t)|x_{-1}), \delta_1) = C \sum_{j=1}^{n+1} V_j^\varepsilon((x(t)|x_{-1}), \delta_1) = C,$$

where in the last step we use Theorem 3.4. Then $U^*(x(t)|x_{-1}) = U(x(t)|x_{-1}) + C$, and therefore

$$(4.5) \quad \arg \max_x U_1(x, x_{-1}) = \arg \max_x U^*(x, x_{-1}).$$

Now, we choose $c \in \mathbb{R}^{1 \times n+1}$ with $c_j = -w_{11}$ for all j , so that $w_1^* = 0$. As a consequence, from the definition of w^* and (4.4), we have $w^* D_\Delta = \mathbf{0}$, thus

$$U^*(x(t)|x_{-1}) = w^* D_t u(t) = w^*(D_0 + D_\Delta)u(t) = w^* D_0 u(t),$$

and using (4.3) we obtain

$$(4.6) \quad U^*(x(t)|x_{-1}) = w^* D_0 A_0^{-1} \delta_1 + t w^* D_0 A_0^{-1} \Delta u(t).$$

From the definition of Δ (4.2) we have

$$\Delta u(t) = u_1(t) \begin{pmatrix} \tilde{\eta} \\ 0 \end{pmatrix},$$

and we can rewrite (4.6) as follow

$$(4.7) \quad U^*(x(t)|x_{-1}) = w^* D_0 A_0^{-1} \delta_1 + t u_1(t) w^* D_0 A_0^{-1} \begin{pmatrix} \tilde{\eta} \\ 0 \end{pmatrix}.$$

Since x and x' are maximizers of U_1 , from (4.5) we know that they are also maximizers of U^* , then

$$\begin{aligned} U^*(x(0)|x_{-1}) &= U^*(x(1)|x_{-1}), \\ w^* D_0 A_0^{-1} \delta_1 &= w^* D_0 A_0^{-1} \delta_1 + u_1(1) w^* D_0 A_0^{-1} \begin{pmatrix} \tilde{\eta} \\ 0 \end{pmatrix}, \\ 0 &= u_1(1) w^* D_0 A_0^{-1} \begin{pmatrix} \tilde{\eta} \\ 0 \end{pmatrix}. \end{aligned}$$

From Lemma 4.1, we know that $u_1(1) > 0$. Consequently, we have

$$w^* D_0 A_0^{-1} \begin{pmatrix} \tilde{\eta} \\ 0 \end{pmatrix} = 0,$$

which, by using (4.7), leads to $U^*(x(t)|x_{-1}) = U^*(x(0)|x_{-1})$ for all $t \in [0, 1]$. Therefore, $x(t)$ maximizes U^* , and from (4.5), we conclude that $x(t)$ maximizes U_1 for all $t \in [0, 1]$. Hence, $r_1(x_{-1})$ is a convex set. Similarly, $r_i(x_{-i})$ is a convex set for all $1 \leq i \leq n + 1$. This completes the proof of the theorem. \square

Remark 3. We note that if we restrict agents' strategies to convex, compact, and non-empty spaces, Theorem 4.1 can be trivially reproduced. For example, we can consider a social network in which $\mathcal{N}_i \subseteq N$ denotes the neighbors of the agent i , and restrict the strategies of each agent to its neighborhood, that is $x_i \in X_i := \{x \in \mathbb{R}_{\geq 0}^n : x_k = 0 \ \forall k \notin \mathcal{N}_i, \text{ and } \sum_k x_k = 1\}$. For this case the proof of the Theorem 4.1 can be easily replicated using $\mathbf{X} := X_1 \times \dots \times X_n$. This setting can be seen as a generalization of problem **MINDIS**, form [16][Section 3.2]. In that work, the authors establish that a pure strategy Nash equilibrium may not exist in the context of simple delegation, and it is even NP-complete to determine whether one exists or not. The aforementioned problem is also discussed in [21]. In that work, the equilibrium states with mixed strategies are examined, and the authors establish the bicriteria approximation guarantees of the game with respect to both rationality and social welfare. However, mixed strategies do not serve as a good generalization of the classical model when deployed to assess the voting weight of agents (see Appendix A.2).

4.2. A stable Liquid Democracy model. As we mentioned before, we can demonstrate that employing the measure V^ε in a liquid democracy system guarantees the existence of equilibria, while ensuring that desirable properties are maintained within a certain error margin that depends on the penalty parameter $\varepsilon > 0$. This observation is formalized in the following result.

Theorem 4.2. *Consider a system consisting of agents $N = \{1, 2, \dots, n\}$, where each agent $i \in N$ delegates their vote by using a delegation preference $x_i \in X$. Let $P \in \mathcal{P}^n$ represent the delegation matrix formed from these preferences. Let $f \in \mathbb{R}_{> 0}^{n+1}$, where f_i represents the inherent voting weight of agent i . Let $W = \{w_1, w_2, \dots, w_n\} \subset \mathbb{R}^{n+1}$ denote the preferences of these agents over others, as specified in Theorem 4.1. Given $0 < \varepsilon < 1$, using the voting power measure V^ε defined in (3.13) the system has the following properties:*

- (I) *δ -Generalization property. Given $\delta > 0$ and $f = \mathbf{1}$, there exists a positive ε_δ such that for any $\varepsilon < \varepsilon_\delta$, $\max_{1 \leq i \leq n} |V_i^\varepsilon(P) - V_i^c(P)| < \delta$ for all $P \in \mathcal{A}^n$, where V^c denotes the classic LD measure as defined in (2.1).*
- (II) *δ -Delegation property, as introduced in Theorem 3.8, which asserts the existence of delegation property (P2) subject to a degree of error that is contingent upon the penalty ε .*

- (III) *δ -Self-selection.* For all $i \in N$ such that $P_{ii} = 0$, it holds that $V_i^\varepsilon(P, f) = 0$. Moreover, given $\delta > 0$, there exists $\varepsilon_\delta > 0$ such that for all $\varepsilon < \varepsilon_\delta$ and for all $P \in \mathcal{P}^n$, we have $(1 - \delta)f_i P_{ii} \leq V_i^\varepsilon(P, f)$.
- (IV) *Conservation of voting weight.* Namely, $\sum_{i=1}^{n+1} V_i^\varepsilon(P, f) = \sum_{i=1}^{n+1} f_i$ for all $P \in \mathcal{P}^n$.
- (V) *Existence of equilibrium states.* The game $\mathcal{G}(N, W, \varepsilon)$ has pure strategy Nash equilibria.

Proof. The property (I) may be straightforwardly deduced from Theorem 3.6. On the other hand, the properties (II), (III), and (IV) are readily derivable from theorems 3.8, 3.3, and 3.4, respectively. Finally, property (V) can be inferred from Theorem 4.1. \square

5. CONCLUSIONS

This paper presents a novel method for computing the voting power (or voting weight) of agents in the liquid democracy paradigm. The proposed method extends the classical model by letting agents to fractionate and delegate their voting power to multiple agents, allowing for the imposition of a penalty factor on the length of the delegation chains. This generalization is achieved while preserving the most reasonable properties of the classical model.

We explore the existence of equilibrium states in the proposed liquid democracy system. Specifically, we define a game in which each agent's delegation preference constitutes their strategy, and the goal of each agent is to maximize a utility function that quantifies their satisfaction regarding their voting power. By leveraging the voting power measure outlined in the first part of this paper and imposing a non-zero penalty factor on the length of delegation chains, we prove that the game admits pure strategy Nash equilibria.

Finally, we present Theorem 4.2, which summarizes the obtained results of this work. Our results demonstrate that by introducing a penalty on the length of delegation chains that is suitably small, it is possible to construct a liquid democracy system that maintains the main features of the classical model within a prescribed error margin. Furthermore, such a system allows for the existence of pure strategy Nash equilibria. To the best of our knowledge, this study provides the first formal proof that establishes the feasibility of a liquid democracy system that guarantees the existence of equilibrium states.

APPENDIX A.

Proof. (of Theorem 3.4). We consider $V^\varepsilon(P, f) \in \mathbb{R}^{n+1}$, defined as in (3.12), for some $\varepsilon > 0$. Firstly, our aim is to demonstrate that the equality $\sum_{i=1}^{n+1} V_i^\varepsilon(P, f) = \sum_{i=1}^{n+1} f_i$ holds for all $\varepsilon > 0$. To this end, using the definitions provided in equations (3.12) and (3.11), we can write

$$(A.1) \quad (I - \tilde{P}^\varepsilon)u^\varepsilon = f.$$

Let $D \in \mathbb{R}^{(n+1) \times (n+1)}$ be the diagonal matrix $D := \text{diag}((P_{11}^\varepsilon, \dots, P_{nn}^\varepsilon, P_{n+1n+1}^\varepsilon))$. Since $\tilde{P}^\varepsilon = P^\varepsilon - D$, from (A.1) we obtain

$$(I - \tilde{P}^\varepsilon)u^\varepsilon = u^\varepsilon - P^\varepsilon u^\varepsilon + Du^\varepsilon = f.$$

By summing all the equations of the preceding linear system, we obtain

$$\sum_{i=1}^{n+1} u_i^\varepsilon - \sum_{i=1}^{n+1} \sum_{j=1}^{n+1} P_{ij}^\varepsilon u_j^\varepsilon + \sum_{i=1}^{n+1} V_i^\varepsilon(P, f) = \sum_{i=1}^{n+1} f_i.$$

Since P^ε is a stochastic matrix, $\sum_{i=1}^{n+1} P_{ij}^\varepsilon = 1$ for all j , and then

$$\sum_{i=1}^{n+1} \sum_{j=1}^{n+1} P_{ij}^\varepsilon u_j^\varepsilon = \sum_{j=1}^{n+1} u_j^\varepsilon \sum_{i=1}^{n+1} P_{ij}^\varepsilon = \sum_{j=1}^{n+1} u_j^\varepsilon.$$

By combining this with the previous equality, we obtain:

$$\sum_{i=1}^{n+1} V_i^\varepsilon(P, f) = \sum_{i=1}^{n+1} f_i.$$

Given that $W := \{w \in \mathbb{R}^{n+1} : \sum_{i=1}^{n+1} w_i = \sum_{i=1}^{n+1} f_i\}$ a closed set and $V^\varepsilon(P, f) \in W$ for all $\varepsilon > 0$, it follows that $V(P, f) = \lim_{\varepsilon \rightarrow 0} V^\varepsilon(P, f)$, and therefore, $V(P, f) \in W$. This concludes the proof. \square

Proof. (of Theorem 3.5). Let $P \in \mathcal{B}^n$ and $N = \{1, \dots, n\}$ be the set of agents associated with this delegation matrix. From the definitions of V and V^c , we observe that $V_i(P) = V_i^c(P) = 0$ for every agent i that belongs to a delegation cycle (according to Definition 2.1). Therefore, without loss of generality, we can assume that the delegation matrix P does not contain cycles. This allows us to omit the elementwise product in the definition of V^c (2.1). Starting from the alternative definition of V given in (3.14), it suffices to prove that

$$(A.2) \quad \lim_{k \rightarrow \infty} P^k \mathbf{1} = \lim_{k \rightarrow \infty} \left(\sum_{\ell=0}^k \tilde{P}^\ell \mathbf{1} \right) \odot \text{diag}(P), \quad \forall P \in \mathcal{B}^n.$$

To this end, without loss of generality, we can assume that the candidate agents are listed last. In this way, the matrix P can be written as

$$P = \begin{pmatrix} P_1 & \mathbf{0} \\ P_2 & I \end{pmatrix},$$

with P_1 being a sub-stochastic matrix. As in (3.9), the powers of P can be computed as

$$P^k = \begin{pmatrix} P_1^k & \mathbf{0} \\ B_k & I \end{pmatrix},$$

with $k \in \mathbb{N}$, and $B_k = \sum_{\ell=0}^{k-1} I^\ell P_2 P_1^{k-\ell-1} = \sum_{\ell=0}^{k-1} P_2 P_1^\ell$. Applying the same idea, we can see that

$$\tilde{P} = \begin{pmatrix} P_1 & \mathbf{0} \\ P_2 & \mathbf{0} \end{pmatrix},$$

and the powers of this matrix can be computed as

$$\tilde{P}^\ell = \begin{pmatrix} P_1^\ell & \mathbf{0} \\ P_2 P_1^{\ell-1} & \mathbf{0} \end{pmatrix},$$

with $\ell \in \mathbb{N}$.

Now, we have

$$(A.3) \quad P^k \mathbf{1} = \begin{pmatrix} P_1^k & \mathbf{0} \\ \sum_{\ell=0}^{k-1} P_2 P_1^\ell & I \end{pmatrix} \mathbf{1} = \begin{pmatrix} P_1^k \mathbf{1} \\ (\sum_{\ell=0}^{k-1} P_2 P_1^\ell) \mathbf{1} + \mathbf{1} \end{pmatrix}.$$

Also we have

$$(A.4) \quad \sum_{\ell=0}^k \tilde{P}^\ell \mathbf{1} = \mathbf{1} + \sum_{\ell=1}^k \tilde{P}^\ell \mathbf{1} = \mathbf{1} + \begin{pmatrix} \sum_{\ell=1}^k P_1^\ell & \mathbf{0} \\ \sum_{\ell=1}^k P_2 P_1^{\ell-1} & \mathbf{0} \end{pmatrix} \mathbf{1} = \begin{pmatrix} (\sum_{\ell=0}^k P_1^\ell) \mathbf{1} \\ (\sum_{\ell=0}^{k-1} P_2 P_1^\ell) \mathbf{1} + \mathbf{1} \end{pmatrix}.$$

Note that $\lim_{k \rightarrow \infty} P_1^k \mathbf{1} = 0$ since P_1 is a sub-stochastic matrix. This, together with (A.3), yields

$$(A.5) \quad \lim_{k \rightarrow \infty} P^k \mathbf{1} = \begin{pmatrix} \lim_{k \rightarrow \infty} P_1^k \mathbf{1} \\ \lim_{k \rightarrow \infty} (\sum_{\ell=0}^{k-1} P_2 P_1^\ell) \mathbf{1} + \mathbf{1} \end{pmatrix} = \begin{pmatrix} \mathbf{0} \\ \lim_{k \rightarrow \infty} (\sum_{\ell=0}^{k-1} P_2 P_1^\ell) \mathbf{1} + \mathbf{1} \end{pmatrix}.$$

We now observe that by listing the candidates last, we have $\text{diag}(P) = \begin{pmatrix} \mathbf{0} \\ \mathbf{1} \end{pmatrix}$, combining this with

(A.4) gives us

$$(A.6) \quad \lim_{k \rightarrow \infty} \left(\sum_{\ell=0}^k \tilde{P}^\ell \mathbf{1} \right) \odot \text{diag}(P) = \begin{pmatrix} \mathbf{0} \\ \lim_{k \rightarrow \infty} (\sum_{\ell=0}^{k-1} P_2 P_1^\ell) \mathbf{1} + \mathbf{1} \end{pmatrix}.$$

Finally, from (A.5) and (A.6), we can deduce (A.2). \square

Proof. (of Theorem 3.6). We posit the assumption that not all the agents within the system are part of a cycle, otherwise, there would be nothing to demonstrate. By applying Theorem 3.5, we obtain $|V_i^\varepsilon(P) - V_i^c(P)| = |V_i^\varepsilon(P) - V_i(P)|$ for all $P \in \mathcal{A}^n$. Therefore, it suffices to establish that the convergence addressed in Theorem 3.1 uniformly holds in \mathcal{A}^n . The proof of this result reveals that the convergence of V_i^ε , where $1 \leq i \leq n$, relies on the powers of the matrix $P_1 \in \mathbb{R}^{r \times r}$, $1 \leq r \leq n$, defined in (3.8). It is noteworthy that this matrix can be represented as:

$$P_1 = \begin{pmatrix} \mathbf{0} & B \end{pmatrix},$$

with $B_{ij} \in \{0, 1\}$, and $\mathbf{0} \in \mathbb{R}^{s \times r}$ with $s \geq 1$, since we are considering $P \in \mathcal{A}^n$. Furthermore, in view of the fact that P_1 accounts for interactions among agents who not belong to any delegation cycle, it can be deduced based on the construction of P_1 that this matrix is nilpotent. Given that $P_1 \in \mathbb{R}^{r \times r}$ with $r \leq n$, it follows that $\sum_{k \geq 0} (1 - \varepsilon)^k P_1^k \mathbf{1} = \sum_{0 \leq k \leq n+1} (1 - \varepsilon)^k P_1^k \mathbf{1}$. Thus, the uniform convergence of $V_i^\varepsilon(P)$ in \mathcal{A}^n can be readily inferred. \square

A.1. Spectral LD generalization. We shall denote the generalized Spectral LD measure as the natural extension of the function V^c defined in (2.1) to the set \mathcal{P}^n . Specifically, we consider $V^c : \mathcal{P}^n \rightarrow \mathbb{R}_{\geq 0}$,

$$(A.7) \quad V^c(P) := \lim_{k \rightarrow \infty} (P^k \mathbf{1}) \odot \text{diag}(P).$$

As discussed in Section 2.2, the function under consideration does not preserve self-selection (property (P3)). Additionally, it is apparent that this generalization does not fulfill the delegation property (P2). This can be demonstrated by examining the example presented in Figure A.1. Considering this, we have

$$P_l = \begin{pmatrix} 1/2 & 1/2 & 0 \\ 1/2 & 0 & 1 \\ 0 & 1/2 & 0 \end{pmatrix}, \text{ and } P_r = \begin{pmatrix} 1/2 & 1/2 & 1 \\ 1/2 & 0 & 0 \\ 0 & 1/2 & 0 \end{pmatrix}$$

where the delegation matrix P_l corresponds to the left setting, whereas P_r corresponds to the right setting. By applying (A.7), it is readily apparent that $V^c(P_l) = (\frac{9}{16}, 0, 0)$ and $V^c(P_r) = (\frac{13}{16}, 0, 0)$.




FIGURE A.1. We consider two settings with $N = \{1, 2, 3\}$, where the delegation property establishes that the voting weight distribution should be the same for both settings. However, it can be verified that this distribution differs when evaluated in each of the aforementioned settings using the measure V^c .

Therefore, it can be deduced that V^c fails to satisfy the delegation property. It is worth noting, however, that $V(P_l) = V(P_r) = (3, 0, 0, 0)$.

A.2. Mixed strategies as multi-agent delegation. As mentioned earlier, in [21], the authors examine a game analogous to the one defined in Section 4 within the framework of the classic model of Liquid Democracy. There, they establish that the game has bicriteria approximation guarantees concerning both rationality and social welfare when mixed strategies are taken into consideration. However, mixed strategies are not a suitable generalization of the classical model when employed to assess the voting power of agents.

To see this, we consider a system consisting of agents $N = \{1, \dots, n\}$, where $x_1, \dots, x_n \in X$ represent their respective delegation preferences, and $P \in \mathcal{P}^n$ is the delegation matrix derived from these preferences. It is noteworthy that $\mathcal{A}^n = \{A \in \mathcal{P}^n : A_{ij} \in \{0, 1\} \text{ for all } 1 \leq i, j \leq n\}$. We define the function $p : \mathcal{A}^n \times \mathcal{P}^n \rightarrow [0, 1]$.

$$p(A, P) := \prod_{i=1}^n A_i^t \cdot P_i,$$

where, A_i and P_i correspond to the i -th column of A and P , respectively. It is easy to see that $p(\cdot, P)$ defines a point probability function on the set \mathcal{A}^n . We can therefore define the mixed strategy-based voting power measure $V^{MS} : \mathcal{P}^n \rightarrow \mathbb{R}^n$ as follows:


$$(A.8) \quad V^{MS}(P) := \sum_{A \in \mathcal{A}^n} p(A, P) V^C(A).$$

Here, V^C represents the voting power measure of the classic model defined in (2.1).

We can easily observe that the voting power measure V^{MS} does not satisfy the delegation property (P2). To demonstrate this, we can consider the example illustrated in Figure A.2, where the delegation matrices are given by:

$$P_l = \begin{pmatrix} 1 & 1/2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1/4 & 0 & 0 \\ 0 & 1/4 & 0 & 1 \end{pmatrix}, \text{ and } P_r = \begin{pmatrix} 1 & 1/2 & 2/3 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1/4 & 0 & 0 \\ 0 & 1/4 & 1/3 & 1 \end{pmatrix}$$

for the left and right settings, respectively. Formula (A.8) yields $V^{MS}(P_l) = (2, 0, 0, \frac{3}{2})$ and $V^{MS}(P_r) = (\frac{7}{3}, 0, 0, \frac{5}{3})$, leading to the conclusion that V^{MS} violates the delegation property. It is noteworthy that even normalizing the output vectors we obtain nonidentical outcomes.



A.3. V is not stable. The game $\mathcal{G}(N, W, 0)$ does not necessarily have a Nash equilibrium. This is illustrated by the example: $N = \{1, 2\}$, $w_1 = (0, -1, 1)$, and $w_2 = (1, 0, -1)$. Indeed, it can be easily verified that no classic delegation strategy pair constitutes an equilibrium state. On the other hand, considering fractional strategies $x_1 = (1 - \eta_1, \eta_1)$ and $x_2 = (1 - \eta_2, \eta_2)$, if $0 < \eta_1 < 1$ and $0 < \eta_2 < 1$, it cannot be an equilibrium as $\eta_2 = 0$ dominates for agent 2. Similarly, for $0 < \eta_1 < 1$ and $\eta_2 = 0$, $\eta_1 = 1$ dominates for agent 1. Likewise, if $\eta_2 = 1$, $\eta_1 = 0$ dominates for agent 1. Therefore, no equilibrium exists for $0 < \eta_1 < 1$. Now, let us consider the cases $\eta_1 = 0$ and $\eta_1 = 1$. For $\eta_1 = 0$, agent 2's dominant strategy is $\eta_2 = 0$, yet agent 1's dominant strategy is $\eta_1 = 1$. Finally, with $\eta_1 = 1$, any $\eta_2 > 0$ strategy for agent 2 is optimal. However, with $\eta_2 > 0$, agent 1's dominant strategy is $\eta_1 = 0$, indicating it cannot be an equilibrium state.

REFERENCES

- [1] J. Behrens. Circular delegations – myth or disaster? *The Liquid Democracy Journal*, 3, 2015.
- [2] J. Behrens, A. Kistner, A. Nitsche, and B. Swierczek. *The principles of LiquidFeedback*. Interaktive Demokratie, 2014.
- [3] J. Behrens, A. Kistner, A. Nitsche, and B. Swierczek. The temporal dimension in the analysis of liquid democracy delegation graphs. *Interaktive Demokratie*, 2021.
- [4] D. Bloembergen, D. Grossi, and M. Lackner. On rational delegations in liquid democracy. In *Proc. AAAI Conf. Artif. Intell.*, volume 33, pages 1796–1803, 2019.
- [5] C. Blum and C. I. Zuber. Liquid democracy: Potentials, problems, and perspectives. *J. Polit. Philos.*, 24(2):162–182, 2016.
- [6] P. Boldi, F. Bonchi, C. Castillo, and S. Vigna. Voting in social networks. In *CIKM '09: Proc. 18th ACM Conf. Inf. Knowl. Manag.*, pages 777–786, 2009.
- [7] P. Boldi, F. Bonchi, C. Castillo, and S. Vigna. Viscous democracy for social networks. *Commun. ACM*, 54(6):129–137, 2011.
- [8] P. Bonacich. Some unique properties of eigenvector centrality. *Soc. Netw.*, 29(4):555–564, 2007.
- [9] F. Brandt, V. Conitzer, U. Endriss, J. Lang, and A. D. Procaccia. *Handbook of computational social choice*. CUP, 2016.
- [10] M. Brill, T. Delemazure, A.-M. George, M. Lackner, and U. Schmidt-Kraepelin. Liquid democracy with ranked delegations. In *Proc. AAAI Conf. Artif. Intell.*, volume 36, pages 4884–4891, 2022.
- [11] M. Brill and N. Talmon. Pairwise liquid democracy. In *IJCAI*, volume 18, pages 137–143, 2018.
- [12] I. Caragiannis and E. Micha. A contribution to the critique of liquid democracy. In *IJCAI*, pages 116–122, 2019.
- [13] L. Carroll. *The principles of parliamentary representation*. Harrison and Sons, 1884.
- [14] R. Colley, U. Grandi, and A. Novaro. Unravelling multi-agent ranked delegations. *Auton. Agents Multi-Agent Syst.*, 36(1):1–35, 2022.
- [15] J. Degraeve. Resolving multi-proxy transitive vote delegation. *arXiv preprint arXiv:1412.4039*, 2014.
- [16] B. Escoffier, H. Gilbert, and A. Pass-Lanneau. The convergence of iterative delegations in liquid democracy in a social network. In *SAGT*, pages 284–297. Springer, 2019.

- [17] P. Gözl, A. Kahng, S. Mackenzie, and A. D. Procaccia. The fluid mechanics of liquid democracy. *ACM Trans. Econ. Comput.*, 9(4):1–39, 2021.
- [18] J. Green-Armytage. Direct voting and proxy voting. *Constitut. Political Econ.*, 26(2):190–220, 2015.
- [19] D. Halpern, J. Y. Halpern, A. Jadbabaie, E. Mossel, A. D. Procaccia, and M. Revel. In defense of liquid democracy. *arXiv preprint arXiv:2107.11868*, 2021.
- [20] A. Kahng, S. Mackenzie, and A. Procaccia. Liquid democracy: An algorithmic perspective. *J. Artif. Intell. Res.*, 70:1223–1252, 2021.
- [21] J. A. Noel, M. Suzuki, and A. Vetta. Pirates in wonderland: Liquid democracy has bicriteria guarantees. In *SAGT*, pages 391–405. Springer, 2021.
- [22] A. Paulin. Župa-grassroots e-democracy revolution on the web. In *E-Democracy*, pages 113–123, 2010.
- [23] A. Paulin. An overview of ten years of liquid democracy research. In *dg.o '20: Proc. 21st Annu. Int. Conf. Digit. Gov. Res.*, pages 116–121, 2020.
- [24] S. Siri, V. Deville, P. Berman, E. Medina, H. Stephens, S. Miller, A. Bacelar, D. Wilson, M. Williams, L. Margot-Duclot, et al. The social smart contract. *The Democracy Earth Foundation*, 2018.
- [25] H. Yamakawa, M. Yoshida, and M. Tsuchiya. Toward delegated democracy: Vote by yourself, or trust your network. *IJHSS*, 1(2):146–150, 2007.
- [26] Y. Zhang and D. Grossi. Power in liquid democracy. In *Proc. AAAI Conf. Artif. Intell.*, volume 35, pages 5822–5830, 2021.

(F.M. Bersetche) DEPARTAMENTO DE MATEMÁTICA, FCEYN, UNIVERSIDAD DE BUENOS AIRES, CIUDAD UNIVERSITARIA 1428, BUENOS AIRES, ARGENTINA.

(F.M. Bersetche) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA, AV. ESPAÑA 1680, VALPARAÍSO, CHILE.

Email address: bersetche@gmail.com