

Chiral current induced by torsional Weyl anomaly

Chong-Sun Chu ^{2,3,4*}, Rong-Xin Miao ^{1†}

¹*School of Physics and Astronomy, Sun Yat-Sen University, 2 Daxue Road, Zhuhai 519082, China¹*

²*Department of Physics, National Tsing-Hua University, Hsinchu 30013, Taiwan*

³*Center of Theory and Computation, National Tsing-Hua University, Hsinchu 30013, Taiwan*

⁴*National Center for Theoretical Sciences, Taipei 10617, Taiwan*

Abstract

Torsion can be realized as dislocation in the crystal lattice of material. It is particularly interesting if the material has fermions in the spectrum, such as graphene, topological insulators, Dirac and Weyl semimetals, as its transport properties can be affected by the torsion. In this letter, we find that, due to Weyl anomaly, torsion in Dirac and Weyl semimetals can induce novel chiral currents, either near a boundary or in a “conformally flat space”. We briefly discuss how to measure this interesting effect in experiment. It is remarkable that these experiments can help to clarify the theoretical controversy of whether an imaginary Pontryagin density could appear in the Weyl anomaly.

*Email: cschu@phys.nthu.edu.tw

†Email: miaorx@mail.sysu.edu.cn

¹All the Institutes of authors contribute equally to this work, the order of Institutes is adjusted for the assessment policy of SYSU.

Contents

1	Introduction	1
2	Torsion and Weyl Anomaly	4
3	Chiral Current	5
3.1	Boundary Theory	5
3.2	Conformally Flat Spacetime	8
4	Weyl Fermions	9

1 Introduction

The study of anomaly induced transport is an interesting subject (see [1] for a recent review). Although anomaly was originally discovered in particle physics, due to its universal nature, anomaly has non-trivial implications to a large number of physical phenomena ranging over vastly different scales. For example, the chiral anomaly of nonabelian gauge theory imposes nontrivial constraints on the fundamental interaction of chiral fermions in the standard model [2]. Chiral anomaly also affects the transport dynamics of systems with chiral fermions [3–12] due to the well-known chiral magnetic and chiral vortical effects (see [13–15] for review). Interestingly, this kind of anomalous transport occurs only in a material system since nonvanishing chemical potentials are required. As anomaly itself is intrinsic to the quantum vacuum, it is an interesting question to ask if anomaly induced transport can occur independent of the chemical potentials.

Recently, chiral response associated with chiral anomaly in a torsional background has been a subject of intensive study. In general, a curved spacetime is equipped with a metric which fixes the causal structure and metric relations, and a connection which defines the parallel transport of tensors on the manifold. While the path dependence of parallel transport is measured by the curvature, the non-closure of parallelism of parallel transport is measured by the torsion. For the parallel transport

$$\nabla_{\mu} V^{\rho} = \partial_{\mu} V^{\rho} + \Gamma^{\rho}_{\mu\nu} V^{\nu}, \quad (1)$$

the torsion

$$T^{\rho}_{\mu\nu} := \Gamma^{\rho}_{\mu\nu} - \Gamma^{\rho}_{\nu\mu}, \quad (2)$$

is given by the antisymmetric part of the connection. In Einstein general relativity (GR), the geometry of spacetime is taken to be torsion free since there is seemingly no observational

evidence for torsion in the spacetime of our universe [16]. However, perhaps unexpectedly, torsion finds a legitimate position in condensed matter physics since torsion appears to be naturally suited for the geometrical description of dislocation defects in crystals [17–20]. Torsion has been realized and studied in diverse material systems such as graphene [21, 22], topological insulators [23–25], Dirac and Weyl semimetal [26–31]. As a result of the specific manner fermion is coupled to torsion, chiral anomaly could emerge and give rise to novel chiral response in torsional material systems [32–40].

Just as a system may possess a chiral anomaly which characterizes the quantum chiral dependence of the system, generally a system may also possess a Weyl anomaly which characterizes the quantum scale dependence of the system. In general, the Weyl anomaly is defined as a difference between the trace of renormalized stress tensor and the renormalized trace of stress tensor [41, 42]

$$\mathcal{A} = \int_M \sqrt{-g} \left[g^{\mu\nu} \langle T_{\mu\nu} \rangle - \langle g^{\mu\nu} T_{\mu\nu} \rangle \right]. \quad (3)$$

In the presence of a background gauge field, the Weyl anomaly receives a contribution

$$\mathcal{A} = \int_M \sqrt{-g} b_1 F_{\mu\nu} F^{\mu\nu} \quad (4)$$

whose form is universal and is entirely determined by the coefficient b_1 , a bulk central charge of the theory. For the normalization of the gauge field kinetic term $S = -1/4 \int F^2$, b_1 is given by the beta function of the theory as $b_1 = -\beta/2$. As Weyl anomaly is independent of the chiral anomaly, it is interesting to ask if and how it give rises to any transport phenomena in a system. The answer is positive. Recently, a new kind of induced transport was discovered for boundary vacuum system as a result of the Weyl anomaly. It was found that [43, 44] for any renormalizable quantum field theory with a current coupled to an external electromagnetic (EM) field

$$S_A = \int_M \sqrt{-g} J^\mu A_\mu, \quad (5)$$

the Weyl anomaly give rises to an induced magnetization current in the vicinity of the boundary of the vacuum system

$$\langle J_\mu \rangle = \frac{-2\beta F_{\mu\nu} n^\nu}{x} + \dots, \quad x \sim 0. \quad (6)$$

Here x is the proper distance to the boundary, n_μ is the inner normal vector, ... denote higher order terms in $O(x)$ and β is the beta function. Hereafter we will drop the symbol $\langle \rangle$ for the expectation value. It is instructive to review the derivation of this result to appreciate how it could be derived from the Weyl anomaly. In general, for a boundary quantum field theory, the renormalized current is generally singular near the boundary and the expectation value takes the asymptotic form near $x \sim 0$:

$$J_\mu = \frac{1}{x^3} J_\mu^{(3)} + \frac{1}{x^2} J_\mu^{(2)} + \frac{1}{x} J_\mu^{(1)} + J^{(0)} \log x + \dots, \quad (7)$$

where \dots denotes terms regular at $x = 0$, and $J_\mu^{(n)}$ depend only the background geometry, the background vector field strength and the type of fields under consideration. For current that is conserved

$$D_\mu J^\mu = 0 \quad (8)$$

up to possibly an anomaly term, one can easily obtain the gauge invariant solution

$$\begin{aligned} J_\mu^{(3)} &= 0, & J_\mu^{(2)} &= 0, \\ J_\mu^{(1)} &= \alpha_1 F_{\mu\nu} n^\nu + \alpha_2 \mathcal{D}_\mu k + \alpha_3 \mathcal{D}_\nu k_\mu^\nu + \alpha_4 \star F_{\mu\nu} n^\nu \end{aligned} \quad (9)$$

where $F_{\mu\nu}$, $\star F_{\mu\nu}$, n_μ , \mathcal{D}_m , $k_{\mu\nu}$ and $h_{\mu\nu}$ are respectively the background field strength, Hodge dual of field strength, the normal vector, induced covariant derivative, extrinsic curvature and induced metric of the boundary. Now the Weyl anomaly \mathcal{A} is a function of the background gauge field. Since it is related to the Logarithmic UV divergent term of effective action, one can establish the following ‘‘integrability’’ relation [43, 54]

$$(\delta\mathcal{A})_{\partial M_\epsilon} = \left(\int_{M_\epsilon} dx^4 \sqrt{g} J^\mu \delta A_\mu \right)_{\log \frac{1}{\epsilon}}, \quad (10)$$

where here a regulator $x \geq \epsilon$ to the boundary is introduced for the integral on the right hand side (RHS) of (10) and J^μ is the renormalized current. The relation (10) identifies the boundary contribution of the variation of the integrated anomaly \mathcal{A} under an arbitrary variation of the gauge field δA_μ with the UV logarithmic divergent part of the integral involving the expectation value J^μ of the renormalized $U(1)$ current. The power of the relation (10) lies in the fact that the left hand side of (10) is a total variation and impose constraints on the RHS of (10) that are powerful enough to to fix completely the asymptotic behavior of the current in terms of the Weyl anomaly of the theory. Using (10), one obtain the result (6) for the renormalized current immediately. We note that the result (6) is universal in two remarkable ways. First, it works for any quantum field theory, and not just conformal field theory. Moreover, it is independent of the choices of boundary conditions since only the bulk central charge, instead of boundary central charge, appears. It should be mentioned that the induced magnetization current for free theories in the vicinity of the the boundary was first obtained by Osborn and McAvity in [45], the universal result (6) as well as its intimate relation with the Weyl anomaly were originally established in [43]. Weyl anomaly also has interesting effect in cosmological spacetime. It was found that in a conformally flat spacetime $ds^2 = e^{2\sigma} \eta_{\mu\nu} dx^\mu dx^\nu$ without boundaries, the anomalous current is given by [46, 47]

$$\langle J^\mu \rangle = -2\beta F^{\mu\nu} \partial_\nu \sigma + O(\sigma^2), \quad (11)$$

to the leading order of small σ . Generalization of the result (6) to higher dimensions and the result (11) for arbitrary finite σ can be found in [48, 49] and [50] respectively. In addition, there is also a novel effect of vacuum spin transport induced by electromagnetic field [51].

Central to these results is the fact that the anomalous currents (6), (11) emerge as direct response of the Weyl anomaly arising from background. Motivated by this observation, it is

natural to expect that similar induced phenomena may occur if the fermions are allowed to couple to other external backgrounds. Now apart from EM background or spacetime curvature, spacetime torsion is another interesting background to consider. In this paper, we will study the quantum transport phenomena induced by Weyl anomaly in a torsional material. In the following, we first derive in section 2 the Weyl anomaly for Dirac fermions coupled to torsion. Using this result, we derive the Weyl anomaly induced chiral and vector currents in section 3. The discussion is extended to Weyl fermions in section 4. We propose that measurements of the induced currents in Weyl semimetal could help to clarify the theoretical controversy of whether the Pontryagin density appears in the Weyl anomaly.

2 Torsion and Weyl Anomaly

Generally a spacetime is equipped with a metric and a connection. In Einstein general relativity (GR), the metricity condition and a symmetric connection are adopted so that connection is not independent but given by the metric. In general, departure from GR is characterized [16] by the non-metricity tensor $Q_{\mu\nu\rho} := \nabla_\mu g_{\nu\rho}$ and the torsion tensor (2) defined as the antisymmetric part of the connection. For simplicity, we will focus in this paper the particular interesting generalization of GR called the Einstein-Cartan theory where $Q = 0$ and the connection is independently characterized by the torsion tensor. In this case, the gravitational coupling in 4-dimensions takes the general form

$$S = \int_M d^4x \sqrt{-g} \bar{\psi} i \gamma^\mu (\nabla_\mu - iV_\mu - i\gamma_5 S_\mu) \psi, \quad (12)$$

where ∇_μ is the covariant derivative defined with the standard Levi-Civita metric connection and the components of torsion

$$V_\mu := T^\rho_{\rho\mu}, \quad S_\mu := \epsilon_{\mu\nu\rho\sigma} T^{\nu\rho\sigma}, \quad (13)$$

behave effectively as vectors and axial vectors [21]. For 3-dimensions, there is no S_μ . Instead

$$\hat{S} := \epsilon_{\nu\rho\sigma} T^{\nu\rho\sigma} \quad (14)$$

behaves as pseudo-scalar and the action in 3-dimensions becomes [21, 22]

$$S = \int_M d^3x \sqrt{-g} \bar{\psi} i \gamma^\mu (\nabla_\mu - iV_\mu - i\gamma_5 \hat{S}) \psi. \quad (15)$$

We use the mostly negative convention for the signature of the metric and the torsion will be taken as a background.

The action (12) is classically Weyl invariant under the local scaling transformation: $\psi \rightarrow e^{-3\sigma/2} \psi$, $g_{\mu\nu} \rightarrow e^{2\sigma} g_{\mu\nu}$, $V_\mu \rightarrow V_\mu$ and $S_\mu \rightarrow S_\mu$. However, quantum mechanically there is an anomaly. For a manifold with a boundary, boundary conditions should be imposed on half of

the spinor components. It can be shown that [52] Hermiticity of the Dirac operator selects out of the general chiral bag boundary conditions the following specific ones:

$$(1 \pm i\gamma^n \gamma^5)\psi|_{\partial M} = 0 \quad (16)$$

where n denote the normal direction. The one-loop Weyl anomaly can be obtained by applying the heat kernel expansion [52]. Let us focus on 4-dimensions, the Weyl anomaly reads

$$\begin{aligned} \mathcal{A} = & \frac{1}{24\pi^2} \int_M d^4x \sqrt{-g} [F_{\mu\nu} F^{\mu\nu} + H_{\mu\nu} H^{\mu\nu}] \\ & + \frac{1}{12\pi^2} \int_{\partial M} d^3x \sqrt{-h} [B_1(S) - B_2(S) - B_3(S)], \end{aligned} \quad (17)$$

where $F = dV$, $H = dS$, $h_{\mu\nu}$ is the induced metric on the boundary ∂M , $k_{\mu\nu} = h_\mu^\rho h_\nu^\sigma \nabla_\rho n_\sigma$ is the extrinsic curvature, $\bar{k}_{\mu\nu}$ and k denote the traceless part and the trace of extrinsic curvatures respectively, $B_1(S) := \frac{2}{3}k(h^{\mu\nu} + n^\mu n^\nu)S_\mu S_\nu + S_n \nabla_\mu S^\mu + 2S_\mu h^{\mu\nu} \nabla_n S_\nu$, $B_2(S) := \frac{1}{3}k S_\mu S^\mu + n^\mu S^\nu \nabla_\nu S_\mu$ and $B_3(S) := \frac{1}{5}\bar{k}_{\mu\nu} S^\mu S^\nu$. Here we choose the normal vector so that $n^\mu = -n_\mu = (0, -1, 0, 0)$ in a flat half space. The Weyl anomaly (17) is Weyl invariant and satisfies the Wess-Zumino consistency condition [53]. Note that the bulk contribution to the torsional Weyl anomaly is discussed in [55–57]. To the best of our knowledge, the boundary contribution to the torsional Weyl anomaly (17) is new.

We are interested in the expectation value of the chiral current and vector current in the theory. In 4-dimensions, the renormalized vacuum expectation value of the chiral current derived by the variation of effective action with respect to the background ‘axial vector’

$$J_S^\mu = \langle \bar{\psi} \gamma^\mu \gamma_5 \psi \rangle = \frac{1}{\sqrt{-g}} \frac{\delta I_{\text{eff}}}{\delta S_\mu}. \quad (18)$$

In the following, we show that the knowledge of the Weyl anomaly (17) allows one immediately to determine (18) in closed analytic form.

3 Chiral Current

3.1 Boundary Theory

Let us first study the chiral current in 4-dimensional spacetime with a boundary, say, at $x = 0$ of the coordinate system. We follow the methods of [43, 54], where we have studied the expectation value of current and stress tensor in boundary quantum field theories [58]. To start with, we note that since the mass dimension of chiral current is 3, it takes the asymptotic form [59]

$$J_S^\mu = \frac{J_0^\mu}{x^3} + \frac{J_1^\mu}{x^2} + \frac{J_2^\mu}{x} + O(\ln x) \quad (19)$$

near the boundary. Here x is the proper distance from the boundary, J_n^μ have mass dimension n and depend on only the background geometry and the background torsion. J_n^μ can be solved by imposing the conservation law $\nabla_\mu J_S^\mu = O(1)$ [43], where $O(1)$ denotes the finite part of the chiral anomaly which is irrelevant to the divergent part of renormalized current. We obtain

$$J_0^\mu = 0, \quad J_1^\mu = \lambda h^{\mu\nu} S_\nu, \quad (20)$$

where λ is some constant. The key point in the above derivations is that the leading term of chiral current cannot be proportional to the normal vector n^μ , otherwise it cannot satisfy the conservation law $\nabla_\mu (n^\mu/x^3) \sim 1/x^4 \neq O(1)$. We note that unlike the case of gauge field [43], the transformation $\delta S_\mu = \partial_\mu \alpha$ changes the torsion and is not required to be a symmetry of the theory, therefore a non-vanishing J_1^μ term as in (20) is allowed.

Now let us use the ‘‘integrability’’ relation (10) for the renormalized chiral current J_S^μ . To proceed, let us employ the Gauss normal coordinates to express the metric $ds^2 = -dx^2 + (h_{ij} - 2xk_{ij} + \dots)dy^i dy^j$ and expand $S_\mu(x) = S_{0\mu} + xS_{1\mu} + O(x^2)$, where $x \in [0, +\infty)$ and S_i give the i^{th} derivatives of S at $x = 0$. Substituting (17),(19), (20) into (10), after some calculations we obtain the chiral current near the boundary:

$$\begin{aligned} J_S^a &= \frac{S_0^a}{6\pi^2 x^2} + \frac{2k^a_b S_0^b + k S_0^a}{10\pi^2 x} + O(\ln x), \\ J_S^n &= \frac{(D_a S^a)_0}{6\pi^2 x} + O(\ln x), \quad x \sim 0, \end{aligned} \quad (21)$$

where n and a denote respectively the normal and tangential directions, D_a is the covariant derivative on the boundary, and we denote for any function $F(x)$ that $F_0 = F(x=0)$. A couple of remarks are in order. We note that as in the discussion [43, 54], the total chiral current is finite since there are boundary contributions to the chiral current which cancel the divergence from the bulk contribution (21). We note that the result (21) applies not only to conformal field theory (CFT) but also the general quantum field theory (QFT) since the Weyl anomaly is well-defined for general quantum field theories [41, 42]. We remark that (21) can be verified by the Green’s function method [61, 62]. In a flat half space with $k_{ab} = 0$, the correction of Green function due to torsion is given by

$$G_c(x, x') = - \int_M d^4 y \sqrt{|g|} G_0(x, y) \gamma^\mu \gamma_5 S_\mu(y) G_0(y, x') + O(S^2) \quad (22)$$

where G_0 is Green’s function without torsion. From (22), we can obtain the chiral current by

$$J_S^\mu(x) = -i \lim_{x' \rightarrow x} \text{Tr}_{\text{reg}} \left[\gamma^\mu \gamma_5 G_c(x, x') \right], \quad (23)$$

which agrees with (21). Here Tr_{reg} means we have subtracted the reference current without boundaries. Another interesting remark is about the universal nature of the boundary current (21). In [38] it was shown that torsion does not lead to new chiral transport effects in the bulk since the response to torsion can be viewed as a manifestation of the chiral vortical effect. To




Figure 1: Chiral current $\vec{J}_S \sim \vec{b}/x^2$ induced by Screw dislocation, where \vec{b} is the Burgers vector.

see this, it was noted that in the presence of torsion, the chiral anomaly receives in addition to the electromagnetic contribution a torsion contribution

$$\partial_\mu J_S^\mu - T_{\lambda\mu}^\lambda J_S^\mu = c_F \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} + c_T \epsilon^{\mu\nu\rho\sigma} \eta_{ab} T_{\mu\nu}^a T_{\rho\sigma}^b, \quad (24)$$

where the last term denotes the famous Nieh-Yan term (here we have focused on the space without curvatures for simplicity) and the coefficients c_F and c_T have mass dimensions zero and two respectively. As a result, unlike c_F , c_T depends on regularization and can be removed by a suitable local counterterm in the background fields [1]. Thus the Nieh-Yan anomaly is not independent, and can be considered a manifestation of the chiral anomaly [38]. In fact the form of current non-conservation will depend on the precise definition of the current [38]. Moreover, a choice of current that is based on physical symmetries was suggested and it was shown that the Nieh-Yan anomaly does not appear [38].

Remarkably, the boundary chiral current induced by the Weyl anomaly (17) is different as, unlike the effect of torsion on the bulk current in (24), the effect of torsion on the boundary current (17) is non-removable. Note that all the coefficients of Weyl anomaly (17) are dimensionless. As a result, the Weyl-anomaly induced chiral current is universal near the boundary, and cannot be removed by local counterterms. The universality of the current (21) near the boundary does not contradict the non-universality of torsion-induced chiral current in the bulk. It arises from novel boundary effect that is independent of renormalization scheme.

Let us briefly comment on how the chiral current (21) may be measured in Dirac semimetals. As shown in Fig. 1, we perform the Screw dislocation of lattices so that the red parallelogram does not close, and the missing part is defined by the blue Burgers vector. The density of the Burgers vector \vec{b} behaves as the axial vector in Dirac and Weyl semimetals [21]. From (21)

together with $\vec{S} \sim \vec{b}$, we draw conclusions that the Screw dislocation induces an anomalous chiral current near the boundary in Dirac and Weyl semimetals

$$\vec{J}_S \sim \frac{\vec{b}}{x^2}, \quad x \gtrsim a \quad (25)$$

where a denotes the lattice length, and we mainly focus on spatial \vec{b} in this paper. Note that our result (21) for the continuum is UV finite and is independent of regularization. However, as we go from the continuum to a lattice, the details of the lattice will enter in general, such as a in (25). In principle, other parameters of the system such as mass, temperature, hydrodynamic velocity etc may also appear in dimensionless combination and correct the overall coefficient of the induced current. Note that, to obey the bag boundary condition (16), we should place an insulator on the boundary of the materials so that no current can flow out of the boundary $x = 0$.

Finally we make a remark for 3-dimensions. Following the same analysis as for the derivation of (19), (20), the renormalized expectation value of the vector current $J_V^\mu = \langle \bar{\psi} \gamma^\mu \psi \rangle$ takes the form

$$J_V^\mu = F^{\eta\mu}(\alpha_1 + \alpha_2 \ln x), \quad (26)$$

where α_1, α_2 are constant parameters which are sensitive to the boundary condition. We note that in dimensions $d < 4$, the current is not related to the Weyl anomaly. Hence the parameters α_1, α_2 are not determined just by the central charges, but by further specific details of theory.

3.2 Conformally Flat Spacetime

There are also novel chiral current in 4-dimensional conformally flat spacetime without boundaries. To demonstrate this, let us start by deriving the anomalous transformation rule for the chiral current. Consider the theory (12) with metric and chiral vector field given by $(g_{\mu\nu}, S_\mu)$. Due to the anomaly, the renormalized effective action I_{eff} is not invariant under the Weyl transformation. Generally, we have [63]

$$\frac{\delta}{\delta\sigma} I_{\text{eff}}(e^{-2\sigma} g_{\mu\nu}) = \mathcal{A}(e^{-2\sigma} g_{\mu\nu}) \quad (27)$$

for arbitrary finite $\sigma(x)$. This can be integrated to give the effective action [53, 64, 65]. Using the fact that the anomaly (17) is Weyl invariant, we obtain immediately the transformation rule for the effective action:

$$I_{\text{eff}}(e^{-2\sigma} g_{\mu\nu}) = I_{\text{eff}}(g_{\mu\nu}) + \frac{1}{24\pi^2} \int_M d^4x \sqrt{-g} H_{\mu\nu} H^{\mu\nu} \sigma, \quad (28)$$

plus a boundary term $\frac{1}{12\pi^2} \int_{\partial M} \sqrt{-h} [\frac{-1}{5} \bar{k}_{\mu\nu} S^\mu S^\nu + B_1 - B_2] \sigma$, which we drop in spacetime without boundaries. One can check that the dilaton effective action satisfies Wess-Zumino consistency

$[\delta_{\sigma_1}, \delta_{\sigma_2}]I_{\text{eff}} = 0$. Using (28), we obtain finally the transformation rule for the chiral current (18) under Weyl transformation $g_{\mu\nu} \rightarrow g'_{\mu\nu} = e^{-2\sigma}g_{\mu\nu}$, $S_\mu \rightarrow S'_\mu = S_\mu$

$$J_S^\mu = \frac{1}{6\pi^2} \nabla_\nu (H^{\nu\mu} \sigma), \quad (29)$$

plus a trivial term $e^{-4\sigma} J_S'^\mu$. Here J_S^μ (resp. $J_S'^\mu$) denotes the vev of the chiral current of the theory (12) in the background spacetime $g_{\mu\nu}$ (resp. $g'_{\mu\nu}$). Taking $g'_{\mu\nu}$ to be the flat spacetime metric and assuming that the chiral current vanishes in some region of the flat spacetime, we finally obtain (29) as the chiral current in conformally flat spacetime

$$ds^2 = e^{2\sigma} \eta_{\mu\nu} dx^\mu dx^\nu. \quad (30)$$

Note that the conformal factor σ in (29) is arbitrary and needs not to be small. Therefore we can use (29) to calculate the current in general conformally flat spacetimes such as Anti-de-Sitter space, de-Sitter space and Robertson-Walker universe. For Robertson-Walker universe $ds^2 = dt^2 - a(t)^2(dx^2 + dy^2 + dz^2)$, we have at time $t = t_*$

$$J_S^\mu = \frac{1}{6\pi^2} H^{0\mu} H \quad (31)$$

where $H = \dot{a}/a$ is the Hubble parameter. For simplicity we have chosen $a(t_*) = 1$. In materials, curvature and torsion can be mimicked by disclinations and dislocations, respectively. Thus, one may measure the effect (29) in Dirac semimetals with suitable disclinations and dislocations.

4 Weyl Fermions

So far we have focused on Dirac fermions. The discussions can be generalized to Weyl fermions straightforwardly. The real part of Weyl anomaly for Weyl fermions is half of that of Dirac fermions (17). As a result, the anomalous chiral current is also half of that Dirac fermions (21), (29). The imaginary part of Weyl anomaly is parity odd and it is controversial whether such term exists [66, 67]. This imaginary part implies that the theory is non-unitary or there is absorption and dissipation in materials. For simplicity, let us take the vector parts of torsion V_μ as an example. The discussion for axial vector parts of torsion S_μ is similar. The Weyl anomaly of Weyl fermions related to V_μ is

$$\mathcal{A} = \frac{1}{48\pi^2} \int_M d^4x \sqrt{-g} [F_{\mu\nu} F^{\mu\nu} + i\frac{3}{2}\eta F_{\mu\nu} {}^*F^{\mu\nu}] \quad (32)$$

where ${}^*F^{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\alpha\beta} F^{\alpha\beta}$, $\eta = 0$ or 1 denote the controversy. Following the above approach, we derive the currents

$$J_V^\mu = \frac{n_\nu F^{\nu\mu} + i\frac{3}{2}\eta n_\nu {}^*F^{\nu\mu}}{12\pi^2 x} + O(\ln x), \quad (33)$$

near a boundary and

$$J_V^\mu = e^{-4\sigma} J_V^{\prime\mu} + \frac{1}{12\pi^2} \nabla_\nu \left(F^{\nu\mu} \sigma + i \frac{3}{2} \eta^* F^{\nu\mu} \sigma \right), \quad (34)$$

in a conformally flat space without boundaries. Recall that the edge dislocations and screw dislocations can induce effective vectors and axial vectors coupled with fermions in materials, respectively. Thus, the vector V_μ can be realized by either an electromagnetic field or suitable edge dislocations in Weyl semimetals. It is interesting to measure the predicted current (33), (34) in Weyl semimetals, which can help to clarify the theoretical controversy that if an imaginary Pontryagin density could appear in the Weyl anomaly [66, 67].

Summarizing, we have shown in this paper that, due to Weyl anomaly, torsion can lead to novel currents and chiral currents for Dirac and Weyl fermions. We propose to measure these interesting effects in Dirac and Weyl semimetals with suitable dislocations. These experiments can help to clarify the theoretical controversy that if an imaginary Pontryagin density could appear in the Weyl anomaly.

Acknowledgments

C.S.Chu acknowledge support of this work by NCTS and the grant 110-2112-M-007-015-MY3 of the National Science and Technology Council of Taiwan. R.X.Miao acknowledges the supports from Guangdong Basic and Applied Basic Research Foundation (No.2020A1515010900) and National Natural Science Foundation of China (No. 11905297).

References

- [1] M. N. Chernodub, Y. Ferreiros, A. G. Grushin, K. Landsteiner and M. A. H. Vozmediano, “Thermal transport, geometry, and anomalies,” Phys. Rept. **977**, 1-58 (2022) [arXiv:2110.05471 [cond-mat.mes-hall]].
- [2] M. E. Peskin and D. V. Schroeder, “An Introduction to quantum field theory,” Addison-Wesley, 1995, ISBN 978-0-201-50397-5
- [3] A. Vilenkin, “Parity nonconservation and neutrino transport in magnetic fields,” Astrophys. J. **451** (1995) 700.
- [4] A. Vilenkin, “Equilibrium Parity Violating Current In A Magnetic Field,” Phys. Rev. D **22** (1980) 3080.

- [5] M. Giovannini and M. E. Shaposhnikov, “Primordial hypermagnetic fields and triangle anomaly,” *Phys. Rev. D* **57** (1998) 2186 [hep-ph/9710234].
- [6] A.Y. Alekseev, V. V. Cheianov, and J. Frohlich, “Universality of transport properties in equilibrium, Goldstone theorem and chiral anomaly,” *Phys. Rev. Lett.* **81** (1998) 3503 [cond-mat/9803346].
- [7] K. Fukushima, “Views of the Chiral Magnetic Effect,” *Lect. Notes Phys.* **871** (2013) 241 [arXiv:1209.5064 [hep-ph]].
- [8] D. Kharzeev and A. Zhitnitsky, “Charge separation induced by P-odd bubbles in QCD matter,” *Nucl. Phys. A* **797**, 67 (2007) [arXiv:0706.1026 [hep-ph]].
- [9] J. Erdmenger, M. Haack, M. Kaminski and A. Yarom, “Fluid dynamics of R-charged black holes,” *JHEP* **0901** (2009) 055 [arXiv:0809.2488 [hep-th]].
- [10] N. Banerjee, J. Bhattacharya, S. Bhattacharyya, S. Dutta, R. Loganayagam and P. Surowka, “Hydrodynamics from charged black branes,” *JHEP* **1101** (2011) 094 [arXiv:0809.2596 [hep-th]].
- [11] D. T. Son and P. Surowka, “Hydrodynamics with Triangle Anomalies,” *Phys. Rev. Lett.* **103** (2009) 191601 [arXiv:0906.5044 [hep-th]].
- [12] K. Landsteiner, E. Megias and F. Pena-Benitez, “Gravitational Anomaly and Transport,” *Phys. Rev. Lett.* **107** (2011) 021601 [arXiv:1103.5006 [hep-ph]].
- [13] K. Landsteiner, E. Megías, F. Pena-Benitez, in: *Strongly Interacting Matter in Magnetic Fields*, Springer, 2013, pp. 433–468.
- [14] D.E. Kharzeev, “The Chiral Magnetic Effect and Anomaly-Induced Transport,” *Prog. Part. Nucl. Phys.* **75** (2014) 133, arXiv:1312.3348 [hep-ph].
- [15] D.E. Kharzeev, J. Liao, S.A. Voloshin, G. Wang, “Chiral Magnetic and Vortical Effects in High-Energy Nuclear Collisions — A Status Report,” *Prog. Part. Nucl. Phys.* **88** (2016) 1, arXiv:1511.04050 [hep-ph].
- [16] R. T. Hammond, “Torsion gravity,” *Rept. Prog. Phys.* **65** (2002), 599-649
- [17] K. Kondo. On the geometrical and physical foundations of the theory of yielding. In *Proc. 2nd Japan Nat. Congr. Applied Mechanics*, pages 41–47, Tokyo, 1952.
- [18] H. Kleinert, *Gauge Fields in Condensed Matter, Vol II: Stresses and Defects* (World Scientific, Singapore, 1989).
- [19] M.O Katanaev and I.V Volovich, “Theory of defects in solids and three-dimensional gravity,” *Annals of Physics* **216**, 1–28 (1992).
- [20] D. R. Nelson, *Defects and geometry in condensed matter physics* (Cambridge University Pres, 2002).

- [21] F. de Juan, A. Cortijo and M. A. Vozmediano, “Dislocations and torsion in graphene and related systems,” Nucl. Phys. B **828**, 625 (2010) [arXiv:0909.4068 [cond-mat.mes-hall]].
- [22] A. Mesaros, D. Sadri and J. Zaanen, “Parallel Transport of Electrons in Graphene Parallels Gravity,” Phys. Rev. B **82** (2010), 073405
- [23] T. L. Hughes, R. G. Leigh and E. Fradkin, “Torsional Response and Dissipationless Viscosity in Topological Insulators,” Phys. Rev. Lett. **107**, 075502 (2011) [arXiv:1101.3541 [cond-mat.mes-hall]].
- [24] T. L. Hughes, R. G. Leigh and O. Parrikar, “Torsional anomalies, Hall viscosity, and bulk-boundary correspondence in topological states,” Phys. Rev. D **88**, 025040 (2013).
- [25] C. Hoyos, “Hall viscosity, topological states and effective theories,” Int. J. Mod. Phys. B **28**, 1430007 (2014).
- [26] M. N. Chernodub and M. Zubkov, “Chiral anomaly in Dirac semimetals due to dislocations,” Phys. Rev. B **95**, no.11, 115410 (2017) [arXiv:1508.03114 [cond-mat.mes-hall]].
- [27] H. Sumiyoshi and S. Fujimoto, “Torsional Chiral Magnetic Effect in a Weyl Semimetal with a Topological Defect,” Phys. Rev. Lett. **116**, no.16, 166601 (2016) [arXiv:1509.03981 [cond-mat.mes-hall]].
- [28] Y. You, G. Y. Cho and T. L. Hughes, “Response Properties of Axion Insulators and Weyl Semimetals Driven by screw dislocations and dynamical axion strings,” Phys. Rev. B **94** (2016) no.8, 085102 doi:10.1103/PhysRevB.94.085102
- [29] Z. M. Huang, L. Li, J. Zhou and H. H. Zhang, “Torsional response and Liouville anomaly in Weyl semimetals with dislocations,” Phys. Rev. B **99** (2019) no.15, 155152 doi:10.1103/PhysRevB.99.155152
- [30] Y. Ferreira, Y. Kedem, E. J. Bergholtz and J. H. Bardarson, “Mixed axial-torsional anomaly in Weyl semimetals,” Phys. Rev. Lett. **122** (2019) no.5, 056601
- [31] J. Nissinen, “Emergent spacetime and gravitational Nieh-Yan anomaly in chiral $p + ip$ Weyl superfluids and superconductors,” Phys. Rev. Lett. **124**, no.11, 117002 (2020) [arXiv:1909.05846 [cond-mat.supr-con]].
- [32] Z. Khaidukov and M. Zubkov, “Chiral torsional effect,” JETP Lett. **108**, no.10, 670-674 (2018) [arXiv:1812.00970 [cond-mat.mes-hall]].
- [33] S. Imaki and Z. Qiu, “Chiral torsional effect with finite temperature, density and curvature,” [arXiv:2004.11899 [hep-th]].
- [34] T. Amitani and Y. Nishida, “Torsion-induced chiral magnetic current in equilibrium,” Annals Phys. **448**, 169181 (2022) [arXiv:2204.13415 [hep-th]].
- [35] Z. M. Huang, B. Han and M. Stone, “Nieh-Yan anomaly: Torsional Landau levels, central charge, and anomalous thermal Hall effect,” Phys. Rev. B **101**, no.12, 125201 (2020) doi:10.1103/PhysRevB.101.125201 [arXiv:1911.00174 [cond-mat.mes-hall]].

- [36] J. Nissinen and G. E. Volovik, “Thermal Nieh-Yan anomaly in Weyl superfluids,” *Phys. Rev. Res.* **2**, no.3, 033269 (2020) [arXiv:1909.08936 [cond-mat.str-el]].
- [37] S. Laurila and J. Nissinen, “Torsional Landau levels and geometric anomalies in condensed matter Weyl systems,” *Phys. Rev. B* **102**, no.23, 235163 (2020) [arXiv:2007.10682 [cond-mat.str-el]].
- [38] Y. Ferreira and K. Landsteiner, “On chiral responses to geometric torsion,” *Phys. Lett. B* **819**, 136419 (2021) [arXiv:2011.10535 [cond-mat.mes-hall]].
- [39] J. Nissinen and G. E. Volovik, “Anomalous chiral transport with vorticity and torsion: Cancellation of two mixed gravitational anomaly currents in rotating chiral $p + ip$ Weyl condensates,” *Phys. Rev. D* **106**, no.4, 045022 (2022) [arXiv:2111.08639 [cond-mat.supr-con]].
- [40] M. Valle and M. A. Vazquez-Mozo, *JHEP* **03**, 177 (2022) [arXiv:2112.02003 [hep-th]].
- [41] M. J. Duff, “Twenty years of the Weyl anomaly,” *Class. Quant. Grav.* **11**, 1387 (1994) [hep-th/9308075].
- [42] L. S. Brown, “Stress Tensor Trace Anomaly in a Gravitational Metric: Scalar Fields,” *Phys. Rev. D* **15**, 1469 (1977).
- [43] C. S. Chu and R. X. Miao, “Weyl Anomaly Induced Current in Boundary Quantum Field Theories,” *Phys. Rev. Lett.* **121**, no. 25, 251602 (2018) [arXiv:1803.03068 [hep-th]].
- [44] C. S. Chu and R. X. Miao, “Anomalous Transport in Holographic Boundary Conformal Field Theories,” *JHEP* **1807**, 005 (2018).
- [45] D. M. McAvity and H. Osborn, “A DeWitt expansion of the heat kernel for manifolds with a boundary,” *Class. Quant. Grav.* **8**, 603-638 (1991)
- [46] M. N. Chernodub, “Anomalous Transport Due to the Conformal Anomaly,” *Phys. Rev. Lett.* **117** (2016) no.14, 141601 [arXiv:1603.07993 [hep-th]].
- [47] M. N. Chernodub, A. Cortijo and M. A. H. Vozmediano, “Generation of a Nernst Current from the Conformal Anomaly in Dirac and Weyl Semimetals,” *Phys. Rev. Lett.* **120**, no.20, 206601 (2018).
- [48] C. Chu and R. Miao, “Boundary String Current & Weyl Anomaly in Six-dimensional Conformal Field Theory,” *JHEP* **07** (2019), 151 [arXiv:1812.10273 [hep-th]].
- [49] C. Chu, “Weyl Anomaly and Vacuum Magnetization Current of M5-brane in Background Flux,” *Fortsch. Phys.* **67** (2019) no.8-9, 1910005 [arXiv:1903.02817 [hep-th]].
- [50] J. J. Zheng, D. Li, Y. Q. Zeng and R. X. Miao, “Anomalous Current Due to Weyl Anomaly for Conformal Field Theory,” *Phys. Lett. B* **797**, 134844 (2019) [arXiv:1904.07017 [hep-th]].
- [51] C. S. Chu and C. H. Leung, “Induced Quantized Spin Current in Vacuum,” *Phys. Rev. Lett.* **127** (2021) no.11, 111601

- [52] D. V. Vassilevich, “Heat kernel expansion: User’s manual,” Phys. Rept. **388**, 279 (2003) [hep-th/0306138].
- [53] J. Wess and B. Zumino, “Consequences of anomalous Ward identities,” Phys. Lett. **37B**, 95 (1971).
- [54] R. X. Miao and C. S. Chu, “Universality for Shape Dependence of Casimir Effects from Weyl Anomaly,” JHEP **1803**, 046 (2018) [arXiv:1706.09652 [hep-th]].
- [55] Y. n. Obukhov, “SPECTRAL GEOMETRY OF THE RIEMANN-CARTAN SPACE-TIME,” Nucl. Phys. B **212**, 237-254 (1983)
- [56] I. L. Buchbinder, S. D. Odintsov and I. L. Shapiro, “NONSINGULAR COSMOLOGICAL MODEL WITH TORSION INDUCED BY VACUUM QUANTUM EFFECTS,” Phys. Lett. B **162**, 92-96 (1985)
- [57] G. H. S. Camargo and I. L. Shapiro, “Anomaly-induced vacuum effective action with torsion: Covariant solution and ambiguities,” Phys. Rev. D **106**, no.4, 045004 (2022) [arXiv:2206.02839 [hep-th]].
- [58] J. L. Cardy, “Boundary conformal field theory,” hep-th/0411189.
- [59] D. Deutsch and P. Candelas, “Boundary Effects in Quantum Field Theory,” Phys. Rev. D **20**, 3063 (1979).
- [60] Note that $\log \epsilon$ appears on the right hand side of the relation (10) since we adopt here the signature convention $(1, -1, -1, -1)$. In contrast, the signature convention $(-1, +1, +1, +1)$ was adopted in [43] and $\log 1/\epsilon$ appears.
- [61] P. J. Hu, Q. L. Hu and R. X. Miao, “Note on anomalous currents for a free theory,” Phys. Rev. D **101**, no.12, 125010 (2020) [arXiv:2004.06924 [hep-th]].
- [62] C. S. Chu and R. X. Miao, “Weyl Anomaly induced Fermi Condensation and Holography,” JHEP **08**, 134 (2020) [arXiv:2005.12975 [hep-th]].
- [63] We have find it conveniently to consider the Weyl transformation $g_{\mu\nu} \rightarrow e^{-2\sigma} g_{\mu\nu}$ here.
- [64] A. Cappelli and A. Coste, “On the Stress Tensor of Conformal Field Theories in Higher Dimensions,” Nucl. Phys. B **314**, 707 (1989).
- [65] A. Schwimmer and S. Theisen, “Spontaneous Breaking of Conformal Invariance and Trace Anomaly Matching,” Nucl. Phys. B **847**, 590 (2011) [arXiv:1011.0696 [hep-th]].
- [66] F. Bastianelli and R. Martelli, “On the trace anomaly of a Weyl fermion,” JHEP **11**, 178 (2016) [arXiv:1610.02304 [hep-th]].
- [67] L. Bonora, S. Giaccari and B. Lima de Souza, “Trace anomalies in chiral theories revisited,” JHEP **07**, 117 (2014) [arXiv:1403.2606 [hep-th]].