

Mutually unbiased maximally entangled bases from difference matrices

Yajuan Zang,¹ Zihong Tian,² Hui-Juan Zuo,² and Shao-Ming Fei^{1,*}

¹*School of Mathematical Sciences, Capital Normal University, Beijing, 100048, China*

²*School of Mathematical Sciences, Hebei Normal University, Shijiazhuang, 050024, China*

Abstract

Based on maximally entangled states, we explore the constructions of mutually unbiased bases in bipartite quantum systems. We present a new way to construct mutually unbiased bases by difference matrices in the theory of combinatorial designs. In particular, we establish q mutually unbiased bases with $q - 1$ maximally entangled bases and one product basis in $\mathbb{C}^q \otimes \mathbb{C}^q$ for arbitrary prime power q . In addition, we construct maximally entangled bases for dimension of composite numbers of non-prime power, such as five maximally entangled bases in $\mathbb{C}^{12} \otimes \mathbb{C}^{12}$ and $\mathbb{C}^{21} \otimes \mathbb{C}^{21}$, which improve the known lower bounds for $d = 3m$, with $(3, m) = 1$ in $\mathbb{C}^d \otimes \mathbb{C}^d$. Furthermore, we construct $p + 1$ mutually unbiased bases with p maximally entangled bases and one product basis in $\mathbb{C}^p \otimes \mathbb{C}^{p^2}$ for arbitrary prime number p .

Keywords: mutually unbiased bases, maximally entangled states, difference matrices, Latin squares

* feishm@cnu.edu.cn

I. INTRODUCTION

The mutually unbiased bases (MUBs) are tightly related to the complementarity of measurements on complementary quantum mechanical observables [34]. They are widely utilized in quantum state tomography [20, 40], quantum kinematics [14] and quantum error correction codes [31]. Moreover, MUBs also play an important role in various tasks in quantum information processing such as quantum key distribution [7], cryptographic protocols [5, 7], mean king problem [2], quantum teleportation and superdense coding [13, 32, 33]. Two orthonormal bases $\mathcal{B}_1 = \{|\phi_i\rangle\}_{i=0}^{d-1}$ and $\mathcal{B}_2 = \{|\psi_j\rangle\}_{j=0}^{d-1}$ of a d -dimensional vector space \mathbb{C}^d are called *mutually unbiased* if

$$|\langle\phi_i|\psi_j\rangle| = \frac{1}{\sqrt{d}}, \quad (i, j \in [d]), \quad (1)$$

where $[d] = \{0, 1, \dots, d - 1\}$. It implies that if a system is in an eigenstate of a certain basis, the measurement outcome with respect to the corresponding MUBs is completely uncertain.

It has been shown that there are no more than $d + 1$ MUBs in \mathbb{C}^d , which can be attained when d is a single prime power [20, 40]. For instance, for any prime dimension $d = p$, the eigenstates of the $p + 1$ operators, $Z, X, XZ, XZ^2, \dots, XZ^{p-1}$, form a full set of $(p + 1)$ Heisenberg-Weyl MUBs, where Z and X are the generalized Pauli operators such that $X|j\rangle = |j + 1 \bmod d\rangle$ and $Z|j\rangle = \omega_d^j|j\rangle$ for the given d -dimensional basis $|j\rangle$, $j \in [d]$. For $d = p^n$, the unitary operators $X^{k_1}Z^{l_1} \otimes \dots \otimes X^{k_n}Z^{l_n}$, $k_j, l_j \in [p]$, acting on the Hilbert space $\mathbb{C}^p \otimes \dots \otimes \mathbb{C}^p$, generate the complete set of MUBs. By partitioning the operators into $d + 1$ commuting classes, the common eigenvectors of the operators in each class just constitute one of the bases. Moreover, the $d + 1$ classes yield $d + 1$ unbiased bases [19]. Nevertheless, when d is a composite number of non-prime power, the maximum number of MUBs of \mathbb{C}^d is still unknown.

Quantum entanglement is considered to be one of the most striking features of quantum mechanics. Especially the maximally entangled states such that whose reduced states are maximally mixed play an important role in quantum information processing. Recent years, the study on mutually unbiased maximally entangled bases (MUMEBs) has attracted much attention [8, 24, 35, 37, 39, 42]. Let $M(d, d')$ be the maximal cardinality of any set of MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$. Liu et al. constructed $p_1^{\alpha_1} - 1$ MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^d$ for $d = p_1^{\alpha_1} \cdots p_s^{\alpha_s}$ with $p_1^{\alpha_1} \leq \dots \leq p_s^{\alpha_s}$ [24]. Xu constructed $2(d-1)$ MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^d$ for $d = p^\alpha$, and $\min\{q, M(d, d)\}$ MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^{qd}$ for $d = p^\alpha$ and $q = p'^{\alpha'}$ [42]. Cheng et al. constructed $2(p_1^{\alpha_1} - 1)$ MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^d$ for odd $d = p_1^{\alpha_1} \cdots p_s^{\alpha_s}$ with $p_1^{\alpha_1} \leq \dots \leq p_s^{\alpha_s}$, and $\min\{(p'_1)^{\alpha'_1} + 1, M(d, d)\}$ MUMEBs in $\mathbb{C}^d \otimes \mathbb{C}^{kd}$ for

odd d and $k = (p'_1)^{\alpha'_1} \cdots (p'_l)^{\alpha'_l}$ with $(p'_1)^{\alpha'_1} \leq \cdots \leq (p'_l)^{\alpha'_l}$, which improved Liu's and Xu's results [8].

Difference matrix was first introduced by Bose and Bush in combinatorial designs [4]. It has been studied primarily as a consequence of their uses in the constructions of orthogonal arrays and Latin squares (see, for example, [3, 11]). Difference matrix has also been found useful in the constructions of authentication codes without secrecy [38], data compression [22], software testing [9, 10] and general Steiner triple systems related to constant weight codes [41].

In this paper, by using difference matrices we investigate the mutually unbiased bases for maximally entangled bases in bipartite systems. We introduce some basic definitions and facts needed in Sec. II. In Sec. III, we present q MUBs with $q - 1$ MEBs and one product basis (PB) in $\mathbb{C}^q \otimes \mathbb{C}^q$ for arbitrary prime power q . In addition, we present MUMEBs for the dimension of composite numbers of non-prime power such as 5 MUMEBs in $\mathbb{C}^{12} \otimes \mathbb{C}^{12}$, 5 MUMEBs in $\mathbb{C}^{21} \otimes \mathbb{C}^{21}$. In Sec. IV, we directly establish a connection between a difference matrix and maximally entangled bases. Finally, we provide $p + 1$ MUBs with p MEBs and one PB in $\mathbb{C}^p \otimes \mathbb{C}^{p^2}$ for arbitrary prime number p . By the way, we also give the corresponding examples in $\mathbb{C}^{12} \otimes \mathbb{C}^{12}$, $\mathbb{C}^3 \otimes \mathbb{C}^6$ and $\mathbb{C}^3 \otimes \mathbb{C}^9$. Conclusions and discussions are given in Sec. V.

II. DIFFERENCE MATRICES AND LATIN SQUARES

In this section, we introduce the concepts of difference matrix and Latin square. As well as we give the connection between a difference matrix and a Latin square. Meanwhile we also prove some preliminary results needed for the rest sections.

Let (G, \odot) be a group of order d . A (d, N, λ) -difference matrix denoted by (d, N, λ) -DM is an $N \times \lambda d$ matrix $M = (m_{il})$ with entries from G , so that for different $i, j \in [N]$, the multiset $\{m_{il} \odot m_{jl}^{-1} : l \in [\lambda d]\}$ contains every element of G λ times. When G is abelian, typically additive notation is used, so that difference $m_{il} - m_{jl}$ is employed. In this paper, we default that the index of the row of a DM begins at 0.

A difference matrix is *normalized* if all entries in its first row are all the identity element of the group G . Obviously, any difference matrix can be written as normalized by $m_{0,l}^{-1}$ action on any element $m_{i,l}$ in i th-row in M for each $i \in [N]$ and $l \in [\lambda d]$. Note that if a difference matrix is normalized, then each element occurs exactly λ times in arbitrary row except for the 0th-row.

Otherwise, if a difference matrix only consists of rows with each element occurring exactly λ times, then it can always be written as normalized by adding a row with all identity.

Example II.1 Below is a normalized $(12, 6, 1)$ -DM in $\mathbb{Z}_2 \times \mathbb{Z}_6$ which will be used to construct MUMEBs and PB in $\mathbb{C}^{12} \otimes \mathbb{C}^{12}$.

$$M = \begin{pmatrix} 00 & 00 & 00 & 00 & 00 & 00 & 00 & 00 & 00 & 00 & 00 & 00 \\ 00 & 01 & 02 & 03 & 04 & 05 & 10 & 11 & 12 & 13 & 14 & 15 \\ 00 & 03 & 10 & 01 & 13 & 15 & 02 & 12 & 05 & 04 & 11 & 14 \\ 00 & 12 & 01 & 15 & 05 & 13 & 03 & 14 & 02 & 11 & 10 & 04 \\ 00 & 04 & 15 & 14 & 02 & 11 & 12 & 10 & 13 & 01 & 03 & 05 \\ 00 & 10 & 12 & 02 & 11 & 01 & 13 & 15 & 04 & 14 & 05 & 03 \end{pmatrix}. \quad (2)$$

A *Latin square* of order d , denoted by $LS(d)$, is a $d \times d$ array in which each cell contains a single symbol from a d -set S , such that each symbol occurs exactly once in each row and exactly once in each column. Let $M = (m_{il})$ be a (d, N, λ) -DM in a group (G, \odot) , and m_i be the multiset formed by the elements of the i th-row of M , $i \in [N]$. For any $g \in G$, define $m_i \odot g = \{m_{il} \odot g : l \in [\lambda d]\}$. Any multiset $m_i \odot g$ is called a *translate* of the i th-row of M . Then, define $Dev(M)$ to be the collection of all d translates of all rows of M , i.e. $Dev(M) = \{m_i \odot g : g \in G, i \in [N]\}$. $Dev(M)$ is called the *development* of M . Actually, when M is normalized and $\lambda = 1$, the development of the i th-row of M forms a Latin square of order d for $1 \leq i \leq N - 1$.

Concerning the development, let us consider a non-normalized $(q, q+1, q)$ -DM for detail which will be needed for rest sections. Let q be a prime power and \mathbb{F}_q be a finite field of order q . Denote $\mathbb{F}_q = \{\alpha_0 = 0, \alpha_1 = 1, \alpha_2, \dots, \alpha_{q-1}\}$ and $\mathbb{F}_q^* = \{\alpha_1 = 1, \alpha_2, \dots, \alpha_{q-1}\}$ the multiplicative group of \mathbb{F}_q . Then M is a $(q, q+1, q)$ -DM as follows [11]:

$$M = \begin{pmatrix} 0 & 0 & \cdots & 0 & \cdots \alpha_i \cdots & \alpha_i & \cdots & \alpha_i & \cdots \alpha_{q-1} \cdots & \alpha_{q-1} \\ 0 & 1 & \cdots & \alpha_{q-1} & \cdots 0 \cdots & \alpha_j & \cdots & \alpha_{q-1} & \cdots 0 & \cdots & \alpha_{q-1} \\ 0 & 1 & \cdots & \alpha_{q-1} & \cdots \alpha_i \cdots & \alpha_i + \alpha_j & \cdots & \alpha_i + \alpha_{q-1} & \cdots \alpha_{q-1} \cdots & \alpha_{q-1} + \alpha_{q-1} \\ & & & & & \cdots & & & & & \\ 0 & \alpha_g & \cdots \alpha_g \alpha_{q-1} \cdots \alpha_i \cdots & \alpha_i + \alpha_g \alpha_j & \cdots \alpha_i + \alpha_g \alpha_{q-1} \cdots \alpha_{q-1} \cdots \alpha_{q-1} + \alpha_g \alpha_{q-1} \\ & & & & & \cdots & & & & & \\ 0 & \alpha_{q-1} \cdots & \alpha_{q-1}^2 & \cdots \alpha_i \cdots \alpha_i + \alpha_{q-1} \alpha_j \cdots & \alpha_i + \alpha_{q-1}^2 & \cdots \alpha_{q-1} \cdots & \alpha_{q-1} + \alpha_{q-1}^2 \end{pmatrix},$$

$$\underbrace{\phantom{\alpha_{q-1} \cdots \alpha_{q-1}^2}}_{P_0} \quad \cdots \quad \underbrace{\phantom{\alpha_i + \alpha_{q-1} \cdots \alpha_{q-1} + \alpha_g \alpha_{q-1}}}_{P_i} \quad \cdots \quad \underbrace{\phantom{\alpha_{q-1} + \alpha_{q-1}^2}}_{P_{q-1}}$$

(3)

with the partitions P_0, P_1, \dots, P_{q-1} , where P_i is a $(q+1) \times q$ submatrix for $i \in [q]$. Actually, if $r > 1$, each translate of the r th-row of M forms a Latin square. Namely, for any fixed $\alpha_t \in \mathbb{F}_q$, we can define a Latin square of order q with $L(i, j) = \alpha_i + \alpha_{r-1}\alpha_j + \alpha_t$, $i, j \in [q]$, where i (j) is the index of row (column) in Latin square. On the other hand, there contain exactly q intersections in any two translates respectively from the r th-row and the r' th-row of M for $r \neq r'$. Moreover each symbol in \mathbb{F}_q appears exactly once in these intersections. Especially, when q is a prime number, we have the following results, see Appendix A for the proof.

Lemma II.1 *When p is a prime number, for positive $r \neq r'$ the p intersection points of any two translates respectively from the r th-row and the r' th-row in the difference matrix M (3) are different and have the properties below:*

1. *if $r = 1$ and $r' = 2$, the p intersection points just appear at the development of submatrix P_i for some $i \in [p]$;*
2. *if $r = 1$ and $r' > 2$, the p intersection points are respectively from the development of submatrix P_i for each $i \in [p]$. Moreover, the difference of two adjoining intersections is a fixed number;*
3. *if $r, r' > 1$, the p intersection points are respectively from the development of submatrix P_i with the same column index j for each $i \in [p]$. Moreover, the intersections just constitute a shift of $(0, 1, \dots, p-1)$.*

Musto introduced the concept of weak orthogonal Latin squares (WOLS) in order to construct MUMEBs [25]. Given a pair of Latin squares L and K with entries l_{ij} and k_{ij} respectively, they are *weak orthogonal* when for all different $i, j \in [d]$, if there exists a unique $s \in [d]$ such that $l_{is} = k_{js}$. A set of $t \geq 2$ Latin squares of order d , say L^1, L^2, \dots, L^t , is said to be *mutually weak orthogonal*, denoted by $t\text{-MWOLS}(d)$, if L^i and L^j are weak orthogonal for all $1 \leq i < j \leq t$.

Actually, difference matrix has a close relationship with weak orthogonal Latin squares.

Lemma II.2 *Corresponding to every normalized $(d, N, 1)$ -DM, there is an $(N - 1)$ -MWOLS(d).*

Proof. Assume $M = (m_{ij})$ is a normalized $(d, N, 1)$ -DM based on group G . Let L^i be the

development of the i th-row of M , $1 \leq i \leq N - 1$, i.e.,

$$L^i = \begin{pmatrix} m_{i1} \odot g_0 & m_{i2} \odot g_0 & \cdots & m_{id} \odot g_0 \\ m_{i1} \odot g_1 & m_{i2} \odot g_1 & \cdots & m_{id} \odot g_1 \\ \vdots & \vdots & \ddots & \vdots \\ m_{i1} \odot g_{d-1} & m_{i2} \odot g_{d-1} & \cdots & m_{id} \odot g_{d-1} \end{pmatrix}, \quad (4)$$

where $g_s \in G, s \in [d]$. Obviously, L^1, L^2, \dots and L^{N-1} are LS(d)s. For any $i \neq j, s, s' \in [d]$, if $m_{ix} \odot g_s = m_{jx} \odot g_{s'}$, then $m_{jx}^{-1} \odot m_{ix} = g_{s'} \odot g_s^{-1}$. Since M is a difference matrix, so there exactly exists one solution of $x \in [d]$. Therefore, L^i and L^j are weak orthogonal for any $1 \leq i \neq j \leq N - 1$. \square

Corollary II.3 *For a normalized $(d, N, 1)$ -DM, any translate of the r th-row and any translate of the r' th-row of the DM just intersect at one point for arbitrary $1 \leq r \neq r' \leq N - 1$.*

III. CONSTRUCTIONS FOR MUMEBs IN $\mathbb{C}^d \otimes \mathbb{C}^d$

In this section, we construct mutually unbiased maximally entangled bases and product basis from a $(d, N, 1)$ -DM in $\mathbb{C}^d \otimes \mathbb{C}^d$. A bipartite pure state $|\Psi\rangle$ in systems A and B is said to be maximally entangled in $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ ($d \leq d'$) if and only if for an arbitrary given orthonormal complete basis $\{|\phi_i\rangle\}_{i=0}^{d-1}$ of the subsystem A , there exists an orthonormal basis $\{|\psi_j\rangle\}_{j=0}^{d'-1}$ of the subsystem B such that

$$|\Psi\rangle = \frac{1}{\sqrt{d}} \sum_{i \in [d]} |\phi_i\rangle \otimes |\psi_i\rangle.$$

A basis of $\mathbb{C}^d \otimes \mathbb{C}^{d'}$ ($d \leq d'$) constituted by maximally entangled states is called a maximally entangled basis (MEB).

A complex Hadamard matrix of order d is a $d \times d$ matrix H with entries H_{ij} such that

$$|H_{ij}| = 1, \quad H_{ij}H_{ij}^* = 1, \quad (5)$$

$$HH^\dagger = dI_d, \quad \sum_{p \in [d]} H_{ip}H_{jp}^* = d\delta_{ij}, \quad (6)$$

$$H^\dagger H = dI_d, \quad \sum_{p \in [d]} H_{pi}H_{pj}^* = d\delta_{ij}, \quad (7)$$

where I_d is the identity matrix of order d . Obviously, $\hat{F}_d \equiv \sqrt{d}F_d$ is a complex Hadamard matrix with F_d corresponding with the $d \times d$ Fourier matrix.

Lemma III.1 Given N indexed families of complex Hadamard matrices H_0, H_1, \dots, H_{N-1} of order d . With respect to a normalized $(d, N, 1)$ -DM, there exist N MUBs with $N - 1$ MEBs and a PB in $\mathbb{C}^d \otimes \mathbb{C}^d$.

Proof. Assume that M is a normalized $(d, N, 1)$ -DM, and $L^r = (l_{ij}^r)$ is the development of the r th-row of M for $1 \leq r \leq N - 1$. Define

$$A_{i,j}^0 = \frac{1}{\sqrt{d}} |i\rangle \otimes \sum_{k \in [d]} |k\rangle \langle k| H_0 |j\rangle, i, j \in [d], \quad (8)$$

$$A_{i,j}^r = \frac{1}{\sqrt{d}} \sum_{k \in [d]} |l_{ik}^r\rangle \otimes |k\rangle \langle k| H_r |j\rangle, i, j \in [d], 1 \leq r \leq N - 1. \quad (9)$$

It is easy to check that $\{A_{i,j}^0 : i, j \in [d]\}$ is a set of PB and $\{A_{i,j}^r : i, j \in [d]\}$ is a set of MEB for each $1 \leq r \leq N - 1$. Next we show the mutual unbiasedness.

For any $1 \leq r \leq N - 1$, we have

$$\begin{aligned} |(A_{i,j}^0, A_{i',j'}^r)| &= \frac{1}{d} |\langle k| H_r |j\rangle^* \langle k| H_0 |j'\rangle| \\ &= \frac{1}{d}. \end{aligned}$$

For any $1 \leq r \neq r' \leq N - 1$, by Corollary II.3 we have

$$\begin{aligned} |(A_{i,j}^r, A_{i',j'}^{r'})| &= \frac{1}{d} \left| \sum_{k, k' \in [d]} \langle l_{ik}^r | l_{i'k'}^{r'} \rangle \langle k| k' \rangle \langle k| H_r |j\rangle^* \langle k'| H_{r'} |j'\rangle \right| \\ &= \frac{1}{d} \left| \sum_{k \in [d]} \langle l_{ik}^r | l_{i'k}^{r'} \rangle \langle k| H_r |j\rangle^* \langle k| H_{r'} |j'\rangle \right| \\ &= \frac{1}{d} |\langle k| H_r |j\rangle^* \langle k| H_{r'} |j'\rangle| \\ &= \frac{1}{d}, \end{aligned}$$

which completes the proof. \square

Example III.1 As an example of the Lemma III.1, we have that there exist 5 MUMEBs and a PB from a normalized $(12, 6, 1)$ -DM in $\mathbb{C}^{12} \otimes \mathbb{C}^{12}$, see Appendix B for the detailed expressions.

There have been fruitful results on the researches of difference matrices, not only for different orders, but also for abelian and non-abelian groups [6, 12, 15, 16, 27–29]. Below are some useful properties of difference matrices.

Lemma III.2 ([6, 12, 15, 16, 27])

1. For any prime power q , there exists a $(q, q, 1)$ -DM and a $(q, q + 1, q)$ -DM.
2. Suppose G is any group of order d , then there exists a $(d, p, 1)$ -DM over G where p is the smallest prime dividing d .
3. A $(d, 4, 1)$ -DM exists if and only if $d \geq 4$ and $d \not\equiv 2 \pmod{4}$.
4. A $(d, 5, 1)$ -DM exists over \mathbb{Z}_d if and only if d is odd and $d \notin \{3, 9\}$, except possibly when $d = 9p$ and p is an odd prime other than 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, or 109.
5. If $m \neq 3, 9$ is odd and not of the form $9n$, n not divisible by 3, 5, 7, 11, 13, 17, 23, 29, 31, or 109, then there exists a $(16m, 5, 1)$ -DM over Dihedral group $D(8m)$.

Lemma III.3 ([1, 11, 21, 23, 36]) There exists a $(12, 6, 1)$ -DM, $(21, 6, 1)$ -DM, $(24, 8, 1)$ -DM, $(33, 6, 1)$ -DM, $(39, 6, 1)$ -DM, $(48, 9, 1)$ -DM, $(51, 6, 1)$ -DM, $(57, 7, 1)$ -DM, $(75, 8, 1)$ -DM and $(273, 16, 1)$ -DM .

Combing Lemma III.1-III.3, we have the main results of this section.

Theorem III.4

1. For any prime power q , $M(q, q) \geq q - 1$.
2. For any group G of order d , $M(d, d) \geq p - 1$, where p is the smallest prime dividing d .
3. If $d \geq 4$ and $d \not\equiv 2 \pmod{4}$, then $M(d, d) \geq 3$.
4. If d is odd and $d \notin \{3, 9\}$, then $M(d, d) \geq 4$ except possibly when $d = 9p$ and p is an odd prime other than 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, or 109.
5. If $m \neq 3, 9$ is odd and not of the form $9n$, n not divisible by 3, 5, 7, 11, 13, 17, 23, 29, 31, or 109, then $M(16m, 16m) \geq 4$.

Theorem III.5 $M(12, 12), M(21, 21), M(33, 33), M(39, 39), M(51, 51) \geq 5$; $M(57, 57) \geq 6$; $M(24, 24), M(75, 75) \geq 7$; $M(48, 48) \geq 8$; and $M(273, 273) \geq 15$.

Particularly, we compare the $M(d, d)$ in the above theorem with other known results for some small d with $d = 3m$ in Table 1.

TABLE I. Lower bound on $M(d, d)$ of MUMEBs for some small d with $d = 3m$, $(3, m) = 1$. a is the lower bound value $2(p^\alpha - 1)$ when d is odd [8] and $p^\alpha - 1$ when d is even [24], where p^α is the minimal prime power factor of d ; b is the lower bound value $(p^{2\alpha} - 1)/2$ when d is odd and $\min\{3(2^t - 1), (p^{2\alpha} - 1)/2\}$ when d is even [35], where p^α is the minimal odd prime power factor of d and t is the maximal positive integer such that $2^t | d$; c is the lower bound in Theorem III.5 in this paper.

d	12	21	24	33	39	48	51	57	75	273
$M(d, d)^a$	2	4	2	4	4	2	4	4	4	4
$M(d, d)^b$	4	4	4	4	4	4	4	4	4	4
$M(d, d)^c$	5	5	7	5	5	8	5	6	7	15

IV. CONSTRUCTIONS FOR MUMEBs IN $\mathbb{C}^p \otimes \mathbb{C}^{p^2}$

Similar to the case of $\lambda = 1$, for $\lambda \geq 2$ the MEBs in $\mathbb{C}^d \otimes \mathbb{C}^{\lambda d}$ also can be obtained from a normalized (d, N, λ) -DM. In this section, we introduce a method for constructing MEBs from a normalized (d, N, λ) -DM and a family of complex Hadamard matrices. Furthermore, we construct mutually unbiased bases with MEBs and one PB in $\mathbb{C}^p \otimes \mathbb{C}^{p^2}$ for arbitrary prime number p .

Lemma IV.1 *Given a normalized $(d, N+1, \lambda)$ -DM with $\lambda \geq 2$ and a family of complex Hadamard matrices H_i^1 of order d for $1 \leq i \leq N - 1$ and H_j^2 of order λ for $j \in [N]$, then $N - 1$ maximally entangled bases and one product basis can be obtained in $\mathbb{C}^d \otimes \mathbb{C}^{\lambda d}$.*

Proof. Assume $M = (m_{ij})$ to be the (d, N, λ) -DM after removing the 0th-row of all identity in a normalized $(d, N + 1, \lambda)$ -DM. Without loss of generality, suppose that the 0th-row in M is $(0, \dots, 0, \dots, d - 1, \dots, d - 1)$ with every element repeated λ times.

Let $S^r = (s_{ij}^r)$ be the development of the r th-row of M for $r \in [N]$. Denote

$$A_{i,j,l}^0 = \frac{1}{\sqrt{\lambda}}|j\rangle \otimes \sum_{\{x \in [\lambda d] : s_{ix}^0 = j\}} |x\rangle \langle j_x| H_0^2 |l\rangle, \quad (10)$$

$$A_{i,j,l}^r = \frac{1}{\sqrt{\lambda d}} \sum_{k \in [d]} \langle k | H_r^1 | j \rangle |k\rangle \otimes \sum_{\{x \in [\lambda d] : s_{ix}^r = k\}} |x\rangle \langle k_x| H_r^2 |l\rangle, \quad (11)$$

where $i, j \in [d]$, $l \in [\lambda]$, $1 \leq r \leq N - 1$, j_x or k_x is the rank of element x in the set $\{x \in [\lambda d] : s_{ix}^r = j\}$ and here we default the rank starts from zero. Obviously $A_{i,j,l}^0$ is a product state, and $A_{i,j,l}^r$ is a maximally entangled state under LU operation in $\mathbb{C}^d \otimes \mathbb{C}^{\lambda d}$ for each $1 \leq r \leq N - 1$, $i, j \in [d]$, $l \in [\lambda]$.

Next we show the orthogonality. Firstly, for any $i, j, i', j' \in [d]$ and $l, l' \in [\lambda]$, we have

$$\begin{aligned}
(A_{i,j,l}^0, A_{i',j',l'}^0) &= \frac{1}{\lambda} \langle j | j' \rangle \sum_{\{x' \in [\lambda d] : s_{i'x}^0 = j'\}} \sum_{\{x \in [\lambda d] : s_{ix}^0 = j\}} \langle x | x' \rangle \langle j_x | H_0^2 | l \rangle^* \langle j'_{x'} | H_0^2 | l' \rangle \\
&= \frac{1}{\lambda} \sum_{\{x \in [\lambda d] : s_{ix}^0 = j\}} \langle j_x | H_0^2 | l \rangle^* \langle j_x | H_0^2 | l' \rangle \delta_{ii'} \delta_{jj'} \\
&= \delta_{ii'} \delta_{jj'} \delta_{ll'}.
\end{aligned} \tag{12}$$

Secondly, for any $i, j, i', j' \in [d]$, $l, l' \in [\lambda]$ and $1 \leq r \leq N - 1$, we have

$$\begin{aligned}
(A_{i,j,l}^r, A_{i',j',l'}^r) &= \frac{1}{\lambda d} \sum_{k \in [d]} \langle k | H_r^1 | j \rangle^* \langle k | H_r^1 | j' \rangle \sum_{\{x \in [\lambda d] : s_{ix}^r = s_{i'x}^r = k\}} \langle k_x | H_r^2 | l \rangle^* \langle k_x | H_r^2 | l' \rangle \\
&= \frac{1}{\lambda d} \sum_{k \in [d]} \langle k | H_r^1 | j \rangle^* \langle k | H_r^1 | j' \rangle \sum_{h \in [\lambda]} \langle h | H_r^2 | l \rangle^* \langle h | H_r^2 | l' \rangle \delta_{ii'} \\
&= \delta_{ii'} \delta_{jj'} \delta_{ll'}.
\end{aligned} \tag{13}$$

□

Example IV.1 There exist 4 MEBs and one PB from a non-normalized $(3, 5, 2)$ -DM in $\mathbb{C}^3 \otimes \mathbb{C}^6$. See Appendix C for the details.

We now use the Heisenberg-Weyl MUBs as the complex Hadamard matrices. In fact, when p is prime, the Heisenberg-Weyl MUBs can be written in matrix form, with $M_0 = I$, $M_1 = F_p$ and $M_{i+1} = D^i F_p$ for $i = 1, \dots, p-1$, where I is the identity matrix; F_p is the $p \times p$ Fourier matrix and D is a diagonal matrix [19]. For instance, when $p = 3$ or 5 , the complete sets are generated by using $D_3 = \text{diag}(1, \omega_3, \omega_3)$ and $D_5 = \text{diag}(1, \omega_5, \omega_5^4, \omega_5^4, \omega_5)$. We show some properties on these Heisenberg-Weyl MUBs below.

Lemma IV.2

1. $\sum_{i \in [p]} |\langle i | D^m \hat{F}_p | j \rangle^* \langle i | D^{m'} \hat{F}_p | j' \rangle| = \sqrt{p}$. Moreover
 $\sum_{i \in [p]} |\langle i | I | j \rangle^* \langle i | D^m \hat{F}_p | j' \rangle| = \sqrt{p}$ for any $m \neq m' \in [p]$ and $j, j' \in [p]$.
2. $\langle i | D^m \hat{F}_p | j \rangle^* \langle i | D^{m'} \hat{F}_p | j' \rangle = \langle i | D^{m'-m} \hat{F}_p | j' - j \rangle$ for any $i, j, m, m' \in [p]$.

Proof. (i) is obvious since $I, D F_p, \dots, D^{p-1} F_p$ are constituted by $p + 1$ MUBs. For the case (ii), suppose $D = \text{diag}(\omega_p^{n_0} = 1, \omega_p^{n_1}, \dots, \omega_p^{n_{p-1}})$. Then we have

$$\begin{aligned}
\langle i | D^m \hat{F}_p | j \rangle^* \langle i | D^{m'} \hat{F}_p | j' \rangle &= ((\omega_p^{n_i})^m \omega_p^{ij})^* ((\omega_p^{n_i})^{m'} \omega_p^{ij'}) \\
&= (\omega_p^{n_i})^{m'-m} \omega_p^{i(j'-j)} \\
&= \langle i | D^{m'-m} \hat{F}_p | j' - j \rangle.
\end{aligned}$$

□

Theorem IV.3 For prime number p , there exist $p+1$ MUBs with p MEBs and one PB in $\mathbb{C}^p \otimes \mathbb{C}^{p^2}$.

Proof. Suppose $H_i^1 = D^i \hat{F}_p$ and $H_j^2 = \hat{F}_p$ for $1 \leq i \leq p$, $j \in [p]$. Assume that M is a non-normalized $(p, p+1, p)$ -DM defined in Eq. (3) and $S^r = (s_{ij}^r)$ is the development of the r th-row of M for $1 \leq r \leq p$. Let $\{A_{i,j,l}^0 : i, j, l \in [p]\}$ and $\{A_{i,j,l}^r : i, j, l \in [p]\}$, $1 \leq r \leq p$, be the PB and MEBs defined in Lemma IV.1, where $d = p$, $N = p+1$ and $\lambda = p$. Namely,

$$A_{i,j,l}^0 = \frac{1}{\sqrt{p}} |j\rangle \otimes \sum_{\{x \in [p^2] : s_{ix}^0 = j\}} |x\rangle \langle j_x | \hat{F}_p |l\rangle, \quad (14)$$

$$A_{i,j,l}^r = \frac{1}{p} \sum_{k \in [p]} \langle k | D^r \hat{F}_p | j \rangle |k\rangle \otimes \sum_{\{x \in [p^2] : s_{ix}^r = k\}} |x\rangle \langle k_x | \hat{F}_p | l \rangle. \quad (15)$$

Next we show the unbiasedness. Firstly for any $1 \leq r \leq p$, we have

$$\begin{aligned} |(A_{i,j,l}^0, A_{i',j',l'}^r)| &= \frac{1}{p\sqrt{p}} |\langle j | D^r \hat{F}_p | j' \rangle| \left| \sum_{\{x \in [p^2] : s_{ix}^0 = s_{i'x}^r = j\}} \langle j_x | \hat{F}_p | l \rangle^* \langle j_x | \hat{F}_p | l' \rangle \right| \\ &= \frac{1}{p\sqrt{p}} |\langle j | D^r \hat{F}_p | j' \rangle| |\langle j_x | \hat{F}_p | l \rangle^* \langle j_x | \hat{F}_p | l' \rangle| \\ &= \frac{1}{p\sqrt{p}}. \end{aligned}$$

Secondly for any $1 \leq r \neq r' \leq p$, by Lemma II.1 and Lemma IV.2 we have

$$\begin{aligned} |(A_{i,j,l}^r, A_{i',j',l'}^{r'})| &= \frac{1}{p^2} \left| \sum_{k \in [p]} \langle k | D^r \hat{F}_p | j \rangle^* \langle k | D^{r'} \hat{F}_p | j' \rangle \sum_{\{x \in [p^2] : s_{ix}^r = s_{i'x}^{r'} = k\}} \langle k_x | \hat{F}_p | l \rangle^* \langle k_x | \hat{F}_p | l' \rangle \right| \\ &= \frac{1}{p^2} \left| \sum_{k \in [p]} \langle k | D^r \hat{F}_p | j \rangle^* \langle k | D^{r'} \hat{F}_p | j' \rangle \langle k_x | \hat{F}_p | l \rangle^* \langle k_x | \hat{F}_p | l' \rangle \right| \\ &= \frac{1}{p^2} \left| \sum_{k \in [p]} \langle k | D^{r'-r} \hat{F}_p | j' - j \rangle \langle k_x | \hat{F}_p | l' - l \rangle \right|. \end{aligned} \quad (16)$$

If $r = 1$ and $r' = 2$, since the i th-row and the i' th-row in S^0 and S^1 just intersect at p points in the development of P_h for a certain h by Lemma II.1, one has $k_x = h$ for any $k \in [p]$. Thus $(\langle 0_x | \hat{F}_p | l' - l \rangle, \langle 1_x | \hat{F}_p | l' - l \rangle, \dots, \langle p-1_x | \hat{F}_p | l' - l \rangle) = \omega^{h(l'-l)}(1, 1, \dots, 1)$. Therefore, Eq. (16) can be further written as

$$\begin{aligned}
& \frac{1}{p^2} \left| \sum_{k \in [p]} \langle k | D \hat{F}_p | j' - j \rangle \langle k_x | \hat{F}_p | l' - l \rangle \right| \\
&= \frac{1}{p^2} \left| \sum_{k \in [p]} \omega^h \langle k | D \hat{F}_p | j' - j \rangle \langle 0 | \hat{F}_p | l' - l \rangle \right| \\
&= \frac{1}{p^2} \left| \sum_{k \in [p]} \langle k | D \hat{F}_p | j' - j \rangle \langle 0 | \hat{F}_p | l' - l \rangle \right| \\
&= \frac{1}{p\sqrt{p}}.
\end{aligned}$$

If $r = 1$ and $r' > 2$ with $g > 1$, by Lemma II.1 the difference of the two adjoining intersections of the i th-row and the i' th-row is $(g-1)^{-1}$, i.e., if $k_x = h$ and $k'_x = h+1$, then $k - k' = (g-1)^{-1}$. Hence, we have $0_x - 1_x = g-1$ and $k_x - (k+1)_x = g-1$, which implies that $(\langle 0_x | \hat{F}_p | l' - l \rangle, \langle 1_x | \hat{F}_p | l' - l \rangle, \dots, \langle p-1_x | \hat{F}_p | l' - l \rangle) = \omega^{h(l'-l)}(1, w^{-(g-1)(l'-l)}, \dots, w^{-(g-1)(p-1)(l'-l)})$ with $0_x = h$. Therefore, Eq. (16) can be written as

$$\begin{aligned}
& \frac{1}{p^2} \left| \sum_{k \in [p]} \langle k | D^{r'-r} \hat{F}_p | j' - j \rangle \langle k_x | \hat{F}_p | l' - l \rangle \right| \\
&= \frac{1}{p^2} \left| \sum_{k \in [p]} \omega^{h(l'-l)} \langle k | D^{r'-r} \hat{F}_p | j' - j \rangle \langle k | \hat{F}_p | - (g-1)(l' - l) \rangle \right| \\
&= \frac{1}{p\sqrt{p}}.
\end{aligned}$$

If $r, r' > 1$, with $g \neq g' \in \{1, \dots, p-1\}$, since the intersections of the i th-row and the i' th-row just constitute a shift of $(0, 1, 2, \dots, p-1)$ by Lemma II.1, we have $(\langle 0_x | \hat{F}_p | l' - l \rangle, \langle 1_x | \hat{F}_p | l' - l \rangle, \dots, \langle p-1_x | \hat{F}_p | l' - l \rangle) = \omega^{h(l'-l)}(1, w^{(l'-l)}, \dots, w^{(p-1)(l'-l)})$ with $0_x = h$. Therefore, Eq. (16) can also be written as

$$\begin{aligned}
& \frac{1}{p^2} \left| \sum_{k \in [p]} \langle k | D^{r'-r} \hat{F}_p | j' - j \rangle \langle k_x | \hat{F}_p | l' - l \rangle \right| \\
&= \frac{1}{p^2} \left| \sum_{k \in [p]} \omega^{h(l'-l)} \langle k | D^{r'-r} \hat{F}_p | j' - j \rangle \langle k | \hat{F}_p | l' - l \rangle \right| \\
&= \frac{1}{p\sqrt{p}}.
\end{aligned}$$

□

Example IV.2 There exist 3 MUMEBs and one PB from a non-normalized $(3, 4, 3)$ -DM in $\mathbb{C}^3 \otimes \mathbb{C}^9$, see Appendix D for the details.

V. CONCLUSIONS

The Latin square has been used as a quite facilitating tool in the research of MUBs. After Hayashi et al. first presented the proof of the existence of the solutions of the mean king's problem with maximal MUBs in prime power dimensions in terms of the mutually orthogonal Latin squares [18], the mutually orthogonal Latin squares have been employed to study MUBs for single systems as well as bipartite systems [25, 30, 37]. While the difference matrix can not only give rise to mutually orthogonal Latin squares [1, 3, 11, 15, 21, 36], but also to the MUMEBs in a more direct way, as shown in this paper. We conjecture that the difference matrix may be also applied to construct MUBs in multipartite systems. Moreover, the difference matrix takes up less storage space than Latin squares, since one row (except for the 0th row) in a normalized difference matrix brings one or λ Latin square(s).

We have introduced a new method for constructing MUMEBs via difference matrices in the theory of combinatorial designs. By using difference matrices, we constructed q mutually unbiased bases with $q - 1$ MEBs and one product basis in $\mathbb{C}^q \otimes \mathbb{C}^q$ for arbitrary prime power q . Furthermore, we constructed MUMEBs for some dimension $d = 3m$, where $(3, m) = 1$ in $\mathbb{C}^d \otimes \mathbb{C}^d$ such as 5 MUMEBs in $\mathbb{C}^{12} \otimes \mathbb{C}^{12}$ and 5 MUMEBs in $\mathbb{C}^{21} \otimes \mathbb{C}^{21}$ etc. (see Table 1), which implied that the bounds of MUMEBs has large probability to be improved especially for bigger dimensions. In addition, we constructed $p + 1$ mutually unbiased bases with p MEBs and one product basis in $\mathbb{C}^p \otimes \mathbb{C}^{p^2}$ for arbitrary prime number p . Concerning the existence of MUMEBs there are still many open problems, for example, about the improvement of the lower bound of $M(3m, 3m) \geq 4$ for any $d = 3m$ with $(3, m) = 1$, and that of $M(p, p^2) \geq p$ for prime number.

Recently, many concepts related to the combinatorial designs have been generalized to the field of quantum information, such as quantum Latin squares, quantum orthogonal arrays and quantum Sudoku etc., which have a close relationship with the absolutely maximally entangled (AME) states, k -uniform states, orthogonal quantum measurements and MUBs [17, 25, 26, 43]. Concerning the relationship among difference matrices, Latin squares and orthogonal arrays, it would be also interesting to consider quantum Latin squares, quantum orthogonal arrays or quantum Sudoku from the view of difference matrices. Moreover, a quantum version of difference matrix would also shed new light on the investigation of AME states, k -uniform states, orthogonal quantum measurements and the related applications in quantum information processing.

ACKNOWLEDGEMENTS

This work is supported by Beijing Postdoctoral Research Foundation (2022ZZ071), Natural Science Foundation of Hebei Province (F2021205001), NSFC (Grant Nos. 11871019, 12075159, 12171044, 62272208), Beijing Natural Science Foundation (Z190005), Academy for Multidisciplinary Studies, Capital Normal University, the Academician Innovation Platform of Hainan Province, and Shenzhen Institute for Quantum Science and Engineering, Southern University of Science and Technology (No. SIQSE202001).

REFERENCES

-
- [1] R. J. R. Abel, C. J. Colbourn, M. Wojtas. Concerning seven and eight mutually orthogonal latin squares. *J. Combin. Des.*, **12**, 123-131 (2004).
 - [2] Y. Aharonov, B.-G. Englert. The mean king's problem: Spin. *Z. Naturforsch. A*, **56**, 16-19 (2001).
 - [3] T. Beth, D. Jungnickel, H. Lenz. *Design Theory*. Bibliographisches Institut, Zurich, 1985.
 - [4] R. C. Bose, K. A. Bush. Orthogonal arrays of strength two and three. *Ann. Math. Statist.*, **23**, 508-524 (1952).
 - [5] S. Brierley. Quantum Key Distribution Highly Sensitive to Eavesdropping. [arXiv:0910.2578](https://arxiv.org/abs/0910.2578) (2009).
 - [6] M. Buratti. Recursive constructions for difference matrices and relative difference families. *J. Combin. Des.*, **6**, 165-182 (1998).
 - [7] N. J. Cerf, M. Bourennane, A. Karlsson, N. Gisin. Security of quantum key distribution using d -level systems. *Phys. Rev. Lett.*, **88**, 127902 (2002).
 - [8] X. Cheng, Y. Shang. New bounds of mutually unbiased maximally entangled bases in $\mathbb{C}^d \otimes \mathbb{C}^{kd}$. *Quantum Inf. Comput.*, **18**, 1152-1164 (2018).
 - [9] D. M. Cohen, S. R. Dalal, M. L. Fredman, G. C. Patton. The AETG system: an approach to testing based on combinatorial design. *IEEE Trans. Software Eng.*, **23**, 437-444 (1997).
 - [10] D. M. Cohen, S. R. Dalal, J. Parelus, G. C. Patton. The combinatorial design approach to automatic test generation. *IEEE Trans. Software*, **13**, 83-88 (1996).
 - [11] C. J. Colbourn, J. H. Dinitz. *The CRC Handbook of Combinatorial Designs*. Chapman and Hall/CRC Press, 2007.
 - [12] D. A. Drake. Partial λ -geometries and generalized Hadamard matrices over groups. *Canad. J. Math.*, **31**, 617-627 (1979).


- [13] T. Durt. If $1 = 2 + 3$, then $1 = 2 \times 3$: Bell states, finite groups, and mutually unbiased bases, a unifying approach. arXiv: [quant-ph/0401046](https://arxiv.org/abs/quant-ph/0401046) (2004).
- [14] T. Durt, B.-G. Englert, I. Bengtsson, K. Życzkowski. On mutually unbiased bases. *Int. J. Quant. Inf.* **8**, 535-640 (2010).
- [15] A. B. Evans. On orthogonal orthomorphisms of cyclic and non-abelian groups II. *J. Combin. Des.*, **15**, 346-412 (2007).
- [16] G. N. Ge. On $(g, 4; 1)$ -difference matrices. *Discrete Math.*, **301**, 164-174 (2005).
- [17] D. Goyeneche, Z. Raissi, S. Di Martino, K. Życzkowski. Entanglement and quantum combinatorial designs. *Phys. Rev. A*, **97**, 062326 (2018) .
- [18] A. Hayashi, M. Horibe, T. Hashimoto. Mean king's problem with mutually unbiased bases and orthogonal Latin squares. *Phys. Rev. A*, **71**, 052331 (2005).
- [19] B. C. Hiesmayr, D. McNulty, S. Baek, S. S. Roy, J. Bae, D. Chruscinski. Detecting entanglement can be more effective with inequivalent mutually unbiased bases. [arXiv:2011.15078](https://arxiv.org/abs/2011.15078) (2021).
- [20] I. D. Ivanović. Geometrical description of quantal state determination. *J. Phys. A*, **14**, 3241-3245 (1981).
- [21] D. M. Johnson, A. L. Dulmage, N. S. Mendelsohn. Orthomorphisms of groups and orthogonal latin squares I. *Canad. J. Math.* **13**, 356-372 (1961).
- [22] J. Korner, M. Lucertini. Compressing inconsistent data. *IEEE Trans. Inform. Theory*, **40**, 706-715 (1994).
- [23] W. F. Kufeld. Table of difference matrices.
http://support.sas.com/techsup/technote/ts723_DifferenceSchemes.txt
(10 January 2006).
- [24] J. Liu, M. Yang, K. Feng. Mutually unbiased maximally entangled bases in $\mathbb{C}^d \otimes \mathbb{C}^d$. *Quantum Inf. Process.*, **16**, 159 (2017).
- [25] B. Musto. Constructing mutually unbiased bases from quantum Latin squares. *Electron. Proc. Theor. Comput. Sci.*, **236**, 108-126 (2017).
- [26] J. Paczos, M. Wierzbniński, G. Rajchel-Mieldzioć, A. Burchardt, K. Życzkowski. Genuinely quantum solutions of the game Sudoku and their cardinality. *Phys. Rev. A*, **104**, 042423 (2021).
- [27] R. Pan, R. J. R. Abel, Y. A. Benjamin, T. Feng, T. J. Tsang Ung, X. M. Wang. Difference matrices with five rows over finite abelian groups. *Des. Codes Cryptogr.*, **90**, 367-386 (2022).

- [28] R. Pan, R. J. R. Abel, Y. A. Bunjamin, T. Feng, X. M. Wang, M. L. Zhang. Difference matrices with four rows over generalized dihedral groups. *J. Combin. Des.*, **30**, 298-314 (2022).
- [29] R. Pan, Y. Chang, A note on difference matrices over noncyclic finite abelian groups, *Discrete Math.*, **339**, 822-830 (2016).
- [30] T. Paterek, B. Daki, Č. Brukner. Mutually unbiased bases, orthogonal Latin squares, and hidden-variable models. *Phys. Rev. A*, **79**, 012109 (2009).
- [31] M. Pawłowski, M. Żukowski. Entanglement-assisted random access codes, *Phys. Rev. A*, **81**, 042326 (2010).
- [32] M. Revzen. Maximally entangled states via mutual unbiased collective bases. *Phys. Rev. A*, **81**, 012113 (2010).
- [33] D. Sangare, P. M. Adler. Continuum percolation of isotropically oriented circular cylinders. *Phys. Rev. E*, **79**, 052101 (2009).
- [34] M. O. Scully, M. S. Zubairy. *Quantum Optics*. Cambridge University Press, Cambridge, 1997.
- [35] F. Shi, Y. Shen, L. Chen, X. D. Zhang. Bounds on the number of mutually unbiased entangled bases. *Quantum Inf. Process.*, **19**, 383 (2020).
- [36] H. Shen. *Theory of Combinatorial Designs*. ShangHai Jiao Tong University Press, ShangHai, 2008.
- [37] Y. Y. Song, G. J. Zhang, L. S Xu, Y. H. Tao. Construction of mutually unbiased bases using mutually orthogonal Latin squares. *Internat. J. Theoret. Phys.*, **59**, 1777-1787 (2020).
- [38] D. R. Stinson. Combinatorial characterizations of authentication codes. *Des. Codes Cryptogr.*, **2**, 175-187 (1992).
- [39] Y. H. Tao, H. Nan, J. Zhang, S. M. Fei. Mutually unbiased maximally entangled bases in $\mathbb{C}^d \otimes \mathbb{C}^{kd}$. *Quantum Inf. Process.*, **14**, 2291-2300 (2015).
- [40] W. K. Wootters, B. D. Fields. Optimal state-determination by mutually unbiased measurements. *Ann. Phys.*, **191**, 363-381 (1989).
- [41] D. Wu, G. Ge, L. Zhu. Generalized Steiner triple systems with group size $g= 7, 8$. *Ars Combin.* **57** 175-192 (2000).
- [42] D. Xu. Construction of mutually unbiased maximally entangled bases through permutations of hadamard matrices. *Quantum Inf. Process.*, **16**, 1-11 (2017).
- [43] Y. J. Zang, P. Facchi, Z. H. Tian. Quantum combinatorial designs and k -uniform states. *J. Phys. A*, **54**, 505204 (2021).

Appendix A: Proof of Lemma II.1

When p is a prime number, M can be written as

$$M = \begin{pmatrix} 0 & 0 & \cdots & 0 & \cdots i \cdots & i & \cdots & i & \cdots p-1 \cdots & p-1 \\ 0 & 1 & \cdots & p-1 & \cdots 0 \cdots & j & \cdots & p-1 & \cdots 0 & \cdots & p-1 \\ 0 & 1 & \cdots & p-1 & \cdots i \cdots & i+j & \cdots & i+p-1 & \cdots p-1 \cdots & p-2 \\ & & & & & & \cdots & & & & \\ 0 & g & \cdots g(p-1) \cdots i \cdots & & i+gj & \cdots i+g(p-1) \cdots p-1 \cdots (g+1)(p-1) \\ & & & & & & \cdots & & & & \\ 0 & p-1 \cdots (p-1)^2 \cdots i \cdots i+(p-1)j \cdots i+(p-1)^2 \cdots p-1 \cdots & & & & & & & & 0 \end{pmatrix}.$$


...

...


 P_0 P_i P_{p-1}

Case $r = 1$ and $r' = 2$. For any $s, s' \in [p]$, the equation $j + s = i + j + s'$ has p solutions given by $i = s - s'$ and $j \in [p]$. Thus the p intersections of the s th translate and the s' th translate of the 1st-row and the 2nd-row are different and just located at the development of the submatrix $P_{s-s'}$.

Case $r = 1$ and $r' > 2$ with $g > 1$. For any $s, s' \in [p]$, the equation $j + s = i + gj + s'$ has p solutions given by $i = s - s' + (1 - g)j$ and $j \in [p]$. Thus the p intersections are different and respectively from the development of the submatrix P_i , $i \in [p]$. Suppose $j_0 + s = i_0 + gj_0 + s'$ and $j_1 + s = (i_0 + 1) + gj_1 + s'$. Then we have $j_0 = (g - 1)^{-1}(-i_0 + s - s')$ and $j_1 = (g - 1)^{-1}(-i_0 - 1 + s - s')$. Therefore, the difference of arbitrary two adjoining intersections is $\pm(g - 1)^{-1}$.

Case $r, r' > 1$, with $g \neq g' \in \{1, \dots, p-1\}$. For any $s, s' \in [p]$, the equation $i + gj + s = i + g'j + s'$ has p solutions given by $j = (g' - g)^{-1}(s - s')$ and $i \in [p]$. Thus the p intersection points are different and respectively located at the same column index j of the development of submatrix P_i for each $i \in [p]$. Moreover they just constitute a shift of $(0, 1, \dots, p-1)$.

Appendix B: Example III.1: MUMEBs and PB in $\mathbb{C}^{12} \otimes \mathbb{C}^{12}$

Let L^i , $i = 0, 1, \dots, 5$, be the development of the i th-row of difference matrix (2) in Example II.1 as follows (Here we map $\mathbb{Z}_2 \times \mathbb{Z}_6$ onto \mathbb{Z}_{12} with bijection map $f : (i, j) \rightarrow 6i + j$).

0	0	0	0	0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9	9	9	9	9
10	10	10	10	10	10	10	10	10	10	10	10
11	11	11	11	11	11	11	11	11	11	11	11

0	1	2	3	4	5	6	7	8	9	10	11
1	2	3	4	5	0	7	8	9	10	11	6
2	3	4	5	0	1	8	9	10	11	6	7
3	4	5	0	1	2	9	10	11	6	7	8
4	5	0	1	2	3	10	11	6	7	8	9
5	0	1	2	3	4	11	6	7	8	9	10
6	7	8	9	10	11	0	1	2	3	4	5
7	8	9	10	11	6	1	2	3	4	5	0
8	9	10	11	6	7	2	3	4	5	0	1
9	10	11	6	7	8	3	4	5	0	1	2
10	11	6	7	8	9	4	5	0	1	2	3
11	6	7	8	9	10	5	0	1	2	3	4

0	3	6	1	9	11	2	8	5	4	7	10
1	4	7	2	10	6	3	9	0	5	8	11
2	5	8	3	11	7	4	10	1	0	9	6
3	0	9	4	6	8	5	11	2	1	10	7
4	1	10	5	7	9	0	6	3	2	11	8
5	2	11	0	8	10	1	7	4	3	6	9
6	9	0	7	3	5	8	2	11	10	1	4
7	10	1	8	4	0	9	3	6	11	2	5
8	11	2	9	5	1	10	4	7	6	3	0
9	6	3	10	0	2	11	5	8	7	4	1
10	7	4	11	1	3	6	0	9	8	5	2
11	8	5	6	2	4	7	1	10	9	0	3

L^0											
0	8	1	11	5	9	3	10	2	7	6	4
1	9	2	6	0	10	4	11	3	8	7	5
2	10	3	7	1	11	5	6	4	9	8	0
3	11	4	8	2	6	0	7	5	10	9	1
4	6	5	9	3	7	1	8	6	11	10	2
5	7	0	10	4	8	2	9	1	6	11	3
6	2	7	5	11	3	9	4	8	1	0	10
7	3	8	0	6	4	10	5	9	2	1	11
8	4	9	1	7	5	11	0	10	3	2	6
9	5	10	2	8	0	6	1	11	4	3	7
10	0	11	3	9	1	7	2	6	5	4	8
11	1	6	4	10	2	8	3	7	0	5	9

L^1											
0	4	11	10	2	7	8	6	9	1	3	5
1	5	6	11	3	8	9	7	10	2	4	0
2	0	7	6	4	9	10	8	11	3	5	1
3	1	8	7	5	10	11	9	6	4	0	2
4	2	9	8	0	11	6	10	7	5	1	3
5	3	10	9	1	6	7	11	8	0	2	4
6	10	5	4	8	1	2	0	3	7	9	11
7	11	0	5	9	2	3	1	4	8	10	6
8	6	1	0	10	3	4	2	5	9	11	7
9	7	2	1	11	4	5	3	0	10	6	8
10	8	3	2	6	5	0	4	1	11	7	9
11	9	4	3	7	0	1	5	2	6	8	10

L^2											
0	6	8	2	7	1	9	11	4	10	5	3
1	7	9	3	8	2	10	6	5	11	0	4
2	8	10	4	9	3	11	7	0	6	1	5
3	9	11	5	10	4	6	8	1	7	2	0
4	10	6	0	11	5	7	9	2	8	3	1
5	11	7	1	6	0	8	10	3	9	4	2
6	0	2	8	1	7	3	5	10	4	11	9
7	1	3	9	2	8	4	0	11	5	6	10
8	2	4	10	3	9	5	1	6	0	7	11
9	3	5	11	4	10	0	2	7	1	8	6
10	4	0	6	5	11	1	3	8	2	9	7
11	5	1	7	0	6	2	4	9	3	10	8

Actually, L^i is a Latin square of order 12 on \mathbb{Z}_{12} for $1 \leq i \leq 5$. Set the complex Hadamard matrix $H_0 = H_1 = \dots = H_5 = \hat{F}_{12}$ with

$$\hat{F}_{12} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \omega & \omega^2 & \omega^3 & \omega^4 & \omega^5 & \omega^6 & \omega^7 & \omega^8 & \omega^9 & \omega^{10} & \omega^{11} \\ 1 & \omega^2 & \omega^4 & \omega^6 & \omega^8 & \omega^{10} & 1 & \omega^2 & \omega^4 & \omega^6 & \omega^8 & \omega^{10} \\ 1 & \omega^3 & \omega^6 & \omega^9 & 1 & \omega^3 & \omega^6 & \omega^9 & 1 & \omega^3 & \omega^6 & \omega^9 \\ 1 & \omega^4 & \omega^8 & 1 & \omega^4 & \omega^8 & 1 & \omega^4 & \omega^8 & 1 & \omega^4 & \omega^8 \\ 1 & \omega^5 & \omega^{10} & \omega^3 & \omega^8 & \omega & \omega^6 & \omega^{11} & \omega^4 & \omega^9 & \omega^2 & \omega^7 \\ 1 & \omega^6 & 1 & \omega^6 \\ 1 & \omega^7 & \omega^2 & \omega^9 & \omega^4 & \omega^{11} & \omega^6 & \omega^1 & \omega^8 & \omega^3 & \omega^{10} & \omega^5 \\ 1 & \omega^8 & \omega^4 & 1 & \omega^8 & \omega^4 & 1 & \omega^8 & \omega^4 & 1 & \omega^8 & \omega^4 \\ 1 & \omega^9 & \omega^6 & \omega^3 & 1 & \omega^9 & \omega^6 & \omega^3 & 1 & \omega^9 & \omega^6 & \omega^3 \\ 1 & \omega^{10} & \omega^8 & \omega^6 & \omega^4 & \omega^2 & 1 & \omega^{10} & \omega^8 & \omega^6 & \omega^4 & \omega^2 \\ 1 & \omega^{11} & \omega^{10} & \omega^9 & \omega^8 & \omega^7 & \omega^6 & \omega^5 & \omega^4 & \omega^3 & \omega^2 & \omega \end{pmatrix},$$

where $\omega = e^{\frac{\pi\sqrt{-1}}{6}}$. Then the six bases can be written as follows:

$$\left\{ \begin{array}{l} A_{0,j}^0 = \frac{1}{2\sqrt{3}}|0\rangle(|0\rangle + \omega^j|1\rangle + \omega^{2j}|2\rangle + \omega^{3j}|3\rangle + \omega^{4j}|4\rangle + \omega^{5j}|5\rangle + \omega^{6j}|6\rangle + \omega^{7j}|7\rangle + \omega^{8j}|8\rangle \\ \quad + \omega^{9j}|9\rangle + \omega^{10j}|10\rangle + \omega^{11j}|11\rangle) \\ A_{1,j}^0 = \frac{1}{2\sqrt{3}}|1\rangle(|0\rangle + \omega^j|1\rangle + \omega^{2j}|2\rangle + \omega^{3j}|3\rangle + \omega^{4j}|4\rangle + \omega^{5j}|5\rangle + \omega^{6j}|6\rangle + \omega^{7j}|7\rangle + \omega^{8j}|8\rangle \\ \quad + \omega^{9j}|9\rangle + \omega^{10j}|10\rangle + \omega^{11j}|11\rangle) \\ \dots \\ A_{11,j}^0 = \frac{1}{2\sqrt{3}}|11\rangle(|0\rangle + \omega^j|1\rangle + \omega^{2j}|2\rangle + \omega^{3j}|3\rangle + \omega^{4j}|4\rangle + \omega^{5j}|5\rangle + \omega^{6j}|6\rangle + \omega^{7j}|7\rangle + \omega^{8j}|8\rangle \\ \quad + \omega^{9j}|9\rangle + \omega^{10j}|10\rangle + \omega^{11j}|11\rangle) \end{array} \right.$$

$$\left\{
\begin{aligned}
A_{0,j}^5 &= \frac{1}{2\sqrt{3}}(|0\rangle|0\rangle + \omega^j|6\rangle|1\rangle + \omega^{2j}|8\rangle|2\rangle + \omega^{3j}|2\rangle|3\rangle + \omega^{4j}|7\rangle|4\rangle + \omega^{5j}|1\rangle|5\rangle + \omega^{6j}|9\rangle|6\rangle \\
&\quad + \omega^{7j}|11\rangle|7\rangle + \omega^{8j}|4\rangle|8\rangle + \omega^{9j}|10\rangle|9\rangle + \omega^{10j}|5\rangle|10\rangle + \omega^{11j}|3\rangle|11\rangle) \\
A_{1,j}^5 &= \frac{1}{2\sqrt{3}}(|1\rangle|1\rangle + \omega^j|7\rangle|1\rangle + \omega^{2j}|9\rangle|2\rangle + \omega^{3j}|3\rangle|3\rangle + \omega^{4j}|8\rangle|4\rangle + \omega^{5j}|2\rangle|5\rangle + \omega^{6j}|10\rangle|6\rangle \\
&\quad + \omega^{7j}|6\rangle|7\rangle + \omega^{8j}|5\rangle|8\rangle + \omega^{9j}|11\rangle|9\rangle + \omega^{10j}|0\rangle|10\rangle + \omega^{11j}|4\rangle|11\rangle) \\
A_{2,j}^5 &= \frac{1}{2\sqrt{3}}(|2\rangle|0\rangle + \omega^j|8\rangle|1\rangle + \omega^{2j}|10\rangle|2\rangle + \omega^{3j}|4\rangle|3\rangle + \omega^{4j}|9\rangle|4\rangle + \omega^{5j}|3\rangle|5\rangle + \omega^{6j}|11\rangle|6\rangle \\
&\quad + \omega^{7j}|7\rangle|7\rangle + \omega^{8j}|0\rangle|8\rangle + \omega^{9j}|6\rangle|9\rangle + \omega^{10j}|1\rangle|10\rangle + \omega^{11j}|5\rangle|11\rangle) \\
A_{3,j}^5 &= \frac{1}{2\sqrt{3}}(|3\rangle|0\rangle + \omega^j|9\rangle|1\rangle + \omega^{2j}|11\rangle|2\rangle + \omega^{3j}|5\rangle|3\rangle + \omega^{4j}|10\rangle|4\rangle + \omega^{5j}|4\rangle|5\rangle + \omega^{6j}|6\rangle|6\rangle \\
&\quad + \omega^{7j}|8\rangle|1\rangle + \omega^{8j}|1\rangle|8\rangle + \omega^{9j}|7\rangle|9\rangle + \omega^{10j}|2\rangle|10\rangle + \omega^{11j}|0\rangle|11\rangle) \\
A_{4,j}^5 &= \frac{1}{2\sqrt{3}}(|4\rangle|0\rangle + \omega^j|10\rangle|1\rangle + \omega^{2j}|6\rangle|2\rangle + \omega^{3j}|0\rangle|3\rangle + \omega^{4j}|11\rangle|4\rangle + \omega^{5j}|5\rangle|5\rangle + \omega^{6j}|7\rangle|6\rangle \\
&\quad + \omega^{7j}|9\rangle|7\rangle + \omega^{8j}|2\rangle|8\rangle + \omega^{9j}|8\rangle|9\rangle + \omega^{10j}|3\rangle|10\rangle + \omega^{11j}|1\rangle|11\rangle) \\
A_{5,j}^5 &= \frac{1}{2\sqrt{3}}(|5\rangle|0\rangle + \omega^j|11\rangle|1\rangle + \omega^{2j}|7\rangle|2\rangle + \omega^{3j}|1\rangle|3\rangle + \omega^{4j}|6\rangle|4\rangle + \omega^{5j}|0\rangle|5\rangle + \omega^{6j}|8\rangle|6\rangle \\
&\quad + \omega^{7j}|10\rangle|7\rangle + \omega^{8j}|3\rangle|8\rangle + \omega^{9j}|9\rangle|9\rangle + \omega^{10j}|4\rangle|10\rangle + \omega^{11j}|2\rangle|11\rangle) \\
A_{6,j}^5 &= \frac{1}{2\sqrt{3}}(|6\rangle|0\rangle + \omega^j|0\rangle|1\rangle + \omega^{2j}|2\rangle|2\rangle + \omega^{3j}|8\rangle|3\rangle + \omega^{4j}|1\rangle|4\rangle + \omega^{5j}|7\rangle|5\rangle + \omega^{6j}|3\rangle|6\rangle \\
&\quad + \omega^{7j}|5\rangle|7\rangle + \omega^{8j}|10\rangle|8\rangle + \omega^{9j}|4\rangle|9\rangle + \omega^{10j}|11\rangle|10\rangle + \omega^{11j}|9\rangle|11\rangle) \\
A_{7,j}^5 &= \frac{1}{2\sqrt{3}}(|7\rangle|0\rangle + \omega^j|1\rangle|1\rangle + \omega^{2j}|3\rangle|2\rangle + \omega^{3j}|9\rangle|3\rangle + \omega^{4j}|2\rangle|4\rangle + \omega^{5j}|8\rangle|5\rangle + \omega^{6j}|4\rangle|6\rangle \\
&\quad + \omega^{7j}|0\rangle|7\rangle + \omega^{8j}|11\rangle|8\rangle + \omega^{9j}|5\rangle|9\rangle + \omega^{10j}|6\rangle|10\rangle + \omega^{11j}|10\rangle|11\rangle) \\
A_{8,j}^5 &= \frac{1}{2\sqrt{3}}(|8\rangle|0\rangle + \omega^j|2\rangle|1\rangle + \omega^{2j}|4\rangle|2\rangle + \omega^{3j}|10\rangle|3\rangle + \omega^{4j}|3\rangle|4\rangle + \omega^{5j}|9\rangle|5\rangle + \omega^{6j}|5\rangle|6\rangle \\
&\quad + \omega^{7j}|1\rangle|7\rangle + \omega^{8j}|6\rangle|8\rangle + \omega^{9j}|0\rangle|9\rangle + \omega^{10j}|7\rangle|10\rangle + \omega^{11j}|11\rangle|11\rangle) \\
A_{9,j}^5 &= \frac{1}{2\sqrt{3}}(|9\rangle|0\rangle + \omega^j|3\rangle|1\rangle + \omega^{2j}|5\rangle|2\rangle + \omega^{3j}|11\rangle|3\rangle + \omega^{4j}|4\rangle|4\rangle + \omega^{5j}|10\rangle|5\rangle + \omega^{6j}|0\rangle|6\rangle \\
&\quad + \omega^{7j}|2\rangle|7\rangle + \omega^{8j}|7\rangle|8\rangle + \omega^{9j}|1\rangle|9\rangle + \omega^{10j}|8\rangle|10\rangle + \omega^{11j}|6\rangle|11\rangle) \\
A_{10,j}^5 &= \frac{1}{2\sqrt{3}}(|10\rangle|0\rangle + \omega^j|4\rangle|1\rangle + \omega^{2j}|0\rangle|2\rangle + \omega^{3j}|6\rangle|3\rangle + \omega^{4j}|5\rangle|4\rangle + \omega^{5j}|11\rangle|5\rangle + \omega^{6j}|1\rangle|6\rangle \\
&\quad + \omega^{7j}|3\rangle|7\rangle + \omega^{8j}|8\rangle|8\rangle + \omega^{9j}|2\rangle|9\rangle + \omega^{10j}|9\rangle|10\rangle + \omega^{11j}|7\rangle|11\rangle) \\
A_{11,j}^5 &= \frac{1}{2\sqrt{3}}(|11\rangle|0\rangle + \omega^j|5\rangle|1\rangle + \omega^{2j}|1\rangle|2\rangle + \omega^{3j}|7\rangle|3\rangle + \omega^{4j}|0\rangle|4\rangle + \omega^{5j}|6\rangle|5\rangle + \omega^{6j}|2\rangle|6\rangle \\
&\quad + \omega^{7j}|4\rangle|7\rangle + \omega^{8j}|9\rangle|8\rangle + \omega^{9j}|3\rangle|9\rangle + \omega^{10j}|10\rangle|10\rangle + \omega^{11j}|8\rangle|11\rangle)
\end{aligned}
\right.$$

where $0 \leq j \leq 11$. It is easy to check that the above six bases are mutually unbiased.

Appendix C: Example IV.1: MEBs and PB in $\mathbb{C}^3 \otimes \mathbb{C}^6$

A (3, 5, 2)-DM without one row all being 0s on \mathbb{Z}_3 exists as follows.

$$M = \begin{pmatrix} 0 & 0 & 1 & 1 & 2 & 2 \\ 0 & 1 & 0 & 2 & 2 & 1 \\ 0 & 1 & 2 & 0 & 1 & 2 \\ 0 & 2 & 1 & 2 & 1 & 0 \\ 0 & 2 & 2 & 1 & 0 & 1 \end{pmatrix}. \quad (\text{C1})$$

Let $H_i^1 = \hat{F}_3$, $H_j^2 = \hat{F}_2$, $1 \leq i \leq 4$, $0 \leq j \leq 4$. Denote S^r the development of the r th-row of M for $0 \leq r \leq 4$:

	0 1 2 3 4 5	0 1 2 3 4 5	0 1 2 3 4 5	0 1 2 3 4 5	0 1 2 3 4 5
0	0 0 1 1 2 2	0 0 1 0 2 2 1	0 0 1 2 0 1 2	0 0 2 1 2 1 0	0 0 2 2 1 0 1
1	1 1 2 2 0 0	1 1 2 1 0 0 2	1 1 2 0 1 2 0	1 1 0 2 0 2 1	1 1 0 0 2 1 2
2	2 2 0 0 1 1	2 2 0 2 1 1 0	2 2 0 1 2 0 1	2 2 1 0 1 0 2	2 2 1 1 0 2 0

$S^0 \quad S^1 \quad S^2 \quad S^3 \quad S^4$

Then the five bases can be written as,

$$\begin{aligned}
& \left\{ \begin{array}{l} A_{0,0,l}^0 = \frac{1}{\sqrt{2}}|0\rangle(|0\rangle \pm |1\rangle) \\ A_{0,1,l}^0 = \frac{1}{\sqrt{2}}|1\rangle(|2\rangle \pm |3\rangle) \\ A_{0,2,l}^0 = \frac{1}{\sqrt{2}}|2\rangle(|4\rangle \pm |5\rangle) \\ A_{1,0,l}^0 = \frac{1}{\sqrt{2}}|0\rangle(|4\rangle \pm |5\rangle) \\ A_{1,1,l}^0 = \frac{1}{\sqrt{2}}|1\rangle(|0\rangle \pm |1\rangle) \\ A_{1,2,l}^0 = \frac{1}{\sqrt{2}}|2\rangle(|2\rangle \pm |3\rangle) \\ A_{2,0,l}^0 = \frac{1}{\sqrt{2}}|0\rangle(|2\rangle \pm |3\rangle) \\ A_{2,1,l}^0 = \frac{1}{\sqrt{2}}|1\rangle(|4\rangle \pm |5\rangle) \\ A_{2,2,l}^0 = \frac{1}{\sqrt{2}}|2\rangle(|0\rangle \pm |1\rangle) \end{array} \right. \\
& \left\{ \begin{array}{l} A_{0,j,l}^1 = \frac{1}{\sqrt{6}}|0\rangle(|0\rangle \pm |2\rangle) + \omega^j|1\rangle(|1\rangle \pm |5\rangle) + \omega^{2j}|2\rangle(|3\rangle \pm |4\rangle) \\ A_{1,j,l}^1 = \frac{1}{\sqrt{6}}|0\rangle(|3\rangle \pm |4\rangle) + \omega^j|1\rangle(|0\rangle \pm |2\rangle) + \omega^{2j}|2\rangle(|1\rangle \pm |5\rangle) \\ A_{2,j,l}^1 = \frac{1}{\sqrt{6}}|0\rangle(|1\rangle \pm |5\rangle) + \omega^j|1\rangle(|3\rangle \pm |4\rangle) + \omega^{2j}|2\rangle(|0\rangle \pm |2\rangle) \end{array} \right. \\
& \left\{ \begin{array}{l} A_{0,j,l}^2 = \frac{1}{\sqrt{6}}|0\rangle(|0\rangle \pm |3\rangle) + \omega^j|1\rangle(|1\rangle \pm |4\rangle) + \omega^{2j}|2\rangle(|2\rangle \pm |5\rangle) \\ A_{1,j,l}^2 = \frac{1}{\sqrt{6}}|0\rangle(|2\rangle \pm |5\rangle) + \omega^j|1\rangle(|0\rangle \pm |3\rangle) + \omega^{2j}|2\rangle(|1\rangle \pm |4\rangle) \\ A_{2,j,l}^2 = \frac{1}{\sqrt{6}}|0\rangle(|1\rangle \pm |4\rangle) + \omega^j|1\rangle(|2\rangle \pm |5\rangle) + \omega^{2j}|2\rangle(|0\rangle \pm |3\rangle) \end{array} \right. \\
& \left\{ \begin{array}{l} A_{0,j,l}^3 = \frac{1}{\sqrt{6}}|0\rangle(|0\rangle \pm |5\rangle) + \omega^j|1\rangle(|2\rangle \pm |4\rangle) + \omega^{2j}|2\rangle(|1\rangle \pm |3\rangle) \\ A_{1,j,l}^3 = \frac{1}{\sqrt{6}}|0\rangle(|1\rangle \pm |3\rangle) + \omega^j|1\rangle(|0\rangle \pm |5\rangle) + \omega^{2j}|2\rangle(|2\rangle \pm |4\rangle) \\ A_{2,j,l}^3 = \frac{1}{\sqrt{6}}|0\rangle(|2\rangle \pm |4\rangle) + \omega^j|1\rangle(|1\rangle \pm |3\rangle) + \omega^{2j}|2\rangle(|0\rangle \pm |5\rangle) \end{array} \right. \\
& \left\{ \begin{array}{l} A_{0,j,l}^4 = \frac{1}{\sqrt{6}}|0\rangle(|0\rangle \pm |4\rangle) + \omega^j|1\rangle(|3\rangle \pm |5\rangle) + \omega^{2j}|2\rangle(|1\rangle \pm |2\rangle) \\ A_{1,j,l}^4 = \frac{1}{\sqrt{6}}|0\rangle(|1\rangle \pm |2\rangle) + \omega^j|1\rangle(|0\rangle \pm |4\rangle) + \omega^{2j}|2\rangle(|3\rangle \pm |5\rangle) \\ A_{2,j,l}^4 = \frac{1}{\sqrt{6}}|0\rangle(|3\rangle \pm |5\rangle) + \omega^j|1\rangle(|1\rangle \pm |2\rangle) + \omega^{2j}|2\rangle(|0\rangle \pm |4\rangle) \end{array} \right.
\end{aligned}$$

where $0 \leq j \leq 2$, $0 \leq l \leq 1$ and $\omega = e^{\frac{2\pi\sqrt{-1}}{3}}$.

Appendix D: Example IV.2: MUMEBs and PB in $\mathbb{C}^3 \otimes \mathbb{C}^9$

A (3, 4, 3)-DM exists on \mathbb{Z}_3 as follows:

$$M = \begin{pmatrix} 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 2 \\ 0 & 1 & 2 & 0 & 1 & 2 & 0 & 1 & 2 \\ 0 & 1 & 2 & 1 & 2 & 0 & 2 & 0 & 1 \\ 0 & 2 & 1 & 1 & 0 & 2 & 2 & 1 & 0 \end{pmatrix}. \quad (\text{D1})$$

Let H_i^1, H_j^2 be the following complex Hadamard matrices for $1 \leq i \leq 3, 0 \leq j \leq 3$,

$$H_1^1 = D\hat{F}_3 = \begin{pmatrix} 1 & 1 & 1 \\ \omega & \omega^2 & 1 \\ \omega & 1 & \omega^2 \end{pmatrix}, \quad H_2^1 = D^2\hat{F}_3 = \begin{pmatrix} 1 & 1 & 1 \\ \omega^2 & 1 & \omega \\ \omega^2 & \omega & 1 \end{pmatrix},$$

$$H_j^2 = H_3^1 = \hat{F}_3 = \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix},$$

where $D = \text{diag}(1, \omega, \omega)$, $\omega = e^{\frac{2\pi\sqrt{-1}}{3}}$.

Denote the development of the r th-row as S^r for $0 \leq r \leq 3$,

0 1 2 3 4 5 6 7 8	0 1 2 3 4 5 6 7 8	0 1 2 3 4 5 6 7 8	0 1 2 3 4 5 6 7 8
0 0 0 1 1 1 2 2 2	0 0 1 2 0 1 2 0 1 2	0 0 1 2 1 2 0 2 0 1	0 0 2 1 1 0 2 2 1 0
1 1 1 2 2 2 0 0 0	1 1 2 0 1 2 0 1 2 0	1 1 2 0 2 0 1 0 1 2	1 1 0 2 2 1 0 0 2 1
2 2 2 0 0 0 1 1 1	2 2 0 1 2 0 1 2 0 1	2 2 0 1 0 1 2 1 2 0	2 2 1 0 0 2 1 1 0 2

$S^0 \qquad S^1 \qquad S^2 \qquad S^3$

The four bases can be written as

$$\left\{ \begin{array}{l} A_{0,0,l}^0 = \frac{1}{\sqrt{3}}|0\rangle(|0\rangle + \omega^l|1\rangle + \omega^{2l}|2\rangle) \\ A_{0,1,l}^0 = \frac{1}{\sqrt{3}}|1\rangle(|3\rangle + \omega^l|4\rangle + \omega^{2l}|5\rangle) \\ A_{0,2,l}^0 = \frac{1}{\sqrt{3}}|2\rangle(|6\rangle + \omega^l|7\rangle + \omega^{2l}|8\rangle) \\ A_{1,0,l}^0 = \frac{1}{\sqrt{3}}|0\rangle(|6\rangle + \omega^l|7\rangle + \omega^{2l}|8\rangle) \\ A_{1,1,l}^0 = \frac{1}{\sqrt{3}}|1\rangle(|0\rangle + \omega^l|1\rangle + \omega^{2l}|2\rangle) \\ A_{1,2,l}^0 = \frac{1}{\sqrt{3}}|2\rangle(|3\rangle + \omega^l|4\rangle + \omega^{2l}|5\rangle) \\ A_{2,0,l}^0 = \frac{1}{\sqrt{3}}|0\rangle(|3\rangle + \omega^l|4\rangle + \omega^{2l}|5\rangle) \\ A_{2,1,l}^0 = \frac{1}{\sqrt{3}}|1\rangle(|6\rangle + \omega^l|7\rangle + \omega^{2l}|8\rangle) \\ A_{2,2,l}^0 = \frac{1}{\sqrt{3}}|2\rangle(|0\rangle + \omega^l|1\rangle + \omega^{2l}|2\rangle) \end{array} \right.$$

$$\left\{ \begin{array}{l} A_{0,j,l}^1 = \frac{1}{3}[|0\rangle(|0\rangle + \omega^l|3\rangle + \omega^{2l}|6\rangle) + \omega^{j+1}|1\rangle(|1\rangle + \omega^l|4\rangle + \omega^{2l}|7\rangle) \\ \quad + \omega^{2j+1}|2\rangle(|2\rangle + \omega^l|5\rangle + \omega^{2l}|8\rangle)] \\ A_{1,j,l}^1 = \frac{1}{3}[|0\rangle(|2\rangle + \omega^l|5\rangle + \omega^{2l}|8\rangle) + \omega^{j+1}|1\rangle(|0\rangle + \omega^l|3\rangle + \omega^{2l}|6\rangle) \\ \quad + \omega^{2j+1}|2\rangle(|1\rangle + \omega^l|4\rangle + \omega^{2l}|7\rangle)] \\ A_{2,j,l}^1 = \frac{1}{3}[|0\rangle(|1\rangle + \omega^l|4\rangle + \omega^{2l}|7\rangle) + \omega^{j+1}|1\rangle(|2\rangle + \omega^l|5\rangle + \omega^{2l}|8\rangle) \\ \quad + \omega^{2j+1}|2\rangle(|0\rangle + \omega^l|3\rangle + \omega^{2l}|6\rangle)] \end{array} \right.$$

$$\left\{ \begin{array}{l} A_{0,j,l}^2 = \frac{1}{3}[|0\rangle(|0\rangle + \omega^l|5\rangle + \omega^{2l}|7\rangle) + \omega^{j+2}|1\rangle(|1\rangle + \omega^l|3\rangle + \omega^{2l}|8\rangle) \\ \quad + \omega^{2j+2}|2\rangle(|2\rangle + \omega^l|4\rangle + \omega^{2l}|6\rangle)] \\ A_{1,j,l}^2 = \frac{1}{3}[|0\rangle(|2\rangle + \omega^l|4\rangle + \omega^{2l}|6\rangle) + \omega^{j+2}|1\rangle(|0\rangle + \omega^l|5\rangle + \omega^{2l}|7\rangle) \\ \quad + \omega^{2j+2}|2\rangle(|1\rangle + \omega^l|3\rangle + \omega^{2l}|8\rangle)] \\ A_{2,j,l}^2 = \frac{1}{3}[|0\rangle(|1\rangle + \omega^l|3\rangle + \omega^{2l}|8\rangle) + \omega^{j+2}|1\rangle(|2\rangle + \omega^l|4\rangle + \omega^{2l}|6\rangle) \\ \quad + \omega^{2j+2}|2\rangle(|0\rangle + \omega^l|5\rangle + \omega^{2l}|7\rangle)] \\ \\ A_{0,j,l}^3 = \frac{1}{3}[|0\rangle(|0\rangle + \omega^l|4\rangle + \omega^{2l}|8\rangle) + \omega^j|1\rangle(|2\rangle + \omega^l|3\rangle + \omega^{2l}|7\rangle) \\ \quad + \omega^{2j}|2\rangle(|1\rangle + \omega^l|5\rangle + \omega^{2l}|6\rangle)] \\ A_{1,j,l}^3 = \frac{1}{3}[|0\rangle(|1\rangle + \omega^l|5\rangle + \omega^{2l}|6\rangle) + \omega^j|1\rangle(|0\rangle + \omega^l|4\rangle + \omega^{2l}|8\rangle) \\ \quad + \omega^{2j}|2\rangle(|2\rangle + \omega^l|3\rangle + \omega^{2l}|7\rangle)] \\ A_{2,j,l}^3 = \frac{1}{3}[|0\rangle(|2\rangle + \omega^l|3\rangle + \omega^{2l}|7\rangle) + \omega^j|1\rangle(|1\rangle + \omega^l|5\rangle + \omega^{2l}|6\rangle) \\ \quad + \omega^{2j}|2\rangle(|0\rangle + \omega^l|4\rangle + \omega^{2l}|8\rangle)] \end{array} \right.$$

where $0 \leq j, l \leq 2$. It is easy to check that the above four bases are mutually unbiased.