Probability of Causation with Sample Selection: A Reanalysis of the Impacts of *Jóvenes en Acción* on Formality[∗]

Vitor Possebom[†] Flavio Riva[‡]

Sao Paulo School of Economics - FGV Sao Paulo School of Business Administration - FGV

First Draft: October 2022; This Draft: November 2022

[Please click here for the most recent version](https://www.sites.google.com/site/vitorapossebom/working-papers)

Abstract

This paper identifies the probability of causation when there is sample selection. We show that the probability of causation is partially identified for individuals who are always observed regardless of treatment status and derive sharp bounds under three increasingly restrictive sets of assumptions. The first set imposes an exogenous treatment and a monotone sample selection mechanism. To tighten these bounds, the second set also imposes the monotone treatment response assumption, while the third set additionally imposes a stochastic dominance assumption. Finally, we use experimental data from the Colombian job training program *J*_{ovenes} en Acción to empirically illustrate our approach's usefulness. We find that, among women who are always employed regardless of treatment, at least 12% and at most 19% transition to the formal labor market because of this training program.

Keywords: Probability of Causation, Sample Selection, Partial Identification, Job Training Programs.

JEL Codes: C31, C35, J24.

[∗]We thank Xiaohong Chen, Bruno Ferman, John Eric Humphries, Helena Laneuville, Yusuke Narita, Cormac O'Dea, Giovanni Di Pietra, Edward Vytlacil, Siu Yuat Wong, and seminar participants at Yale University, EPGE Brazilian School of Economics and Finance, and Sao Paulo School of Economics for helpful suggestions.

† vitor.possebom@fgv.br

[‡]flaviorussoriva@gmail.com

1 Introduction

Many policy evaluation questions involve two simultaneous identification challenges: the causal parameter of interest depends on the joint distribution of potential outcomes and sample selection is present.^{[1](#page-1-0)} For example, when evaluating the effects of job training programs, the researcher may be interested in learning to what extent the transition from informal to formal employment can be attributed to the policy but she only observes formality status among those who are employed. Moreover, when analyzing the effects of a political campaign, the researcher may be interested in identifying the share of the population who supports policy A when treated, given that they would support policy B if untreated. In this case, the researcher only observes the agents' opinions if they reply to a survey. This double identification challenge also arises when researchers consider the effects of health interventions on health quality if agents may pass away, or the effects of educational interventions on learning if there is selection into test-taking.[2](#page-1-1)

In this paper, we derive novel sharp bounds around the probability of causation parameter [\(Pearl,](#page-24-0) [1999;](#page-24-0) [Tian and Pearl,](#page-24-1) [2000;](#page-24-1) [Jun and Lee,](#page-23-0) [2019;](#page-23-0) [Cinelli and Pearl,](#page-23-1) [2021\)](#page-23-1) for individuals who would self-select into the sample regardless of their treatment assignment. The probability of causation parameter summarizes one crucial aspect of the effects of treatments on binary outcomes: the proportion of individuals who benefit from being treated within the subgroup who would, counterfactually, experience a negative untreated outcome. Thus, our target parameter helps researchers gauge to what extent the transition from one state to another can be attributed to the treatment in a relevant latent sub-population.^{[3](#page-1-2)}

¹For a detailed discussion about causal parameters that depend on the joint distribution of potential outcomes, see [Heckman, Smith, and Clements](#page-23-2) [\(1997\)](#page-23-2); [Pearl](#page-24-0) [\(1999\)](#page-24-0); [Tian and Pearl](#page-24-1) [\(2000\)](#page-24-1); [Jun and Lee](#page-23-0) [\(2019\)](#page-23-0); [Cinelli and Pearl](#page-23-1) [\(2021\)](#page-23-1). For a recent discussion on sample selection, see [Lee](#page-24-2) [\(2009\)](#page-24-2); [Chen and Flores](#page-22-0) [\(2015\)](#page-22-0); [Bartalotti, Kedagni, and Possebom](#page-22-1) [\(2022\)](#page-22-1).

²Training programs are studied by [Heckman, LaLonde, and Smith](#page-23-3) [\(1999\)](#page-23-3); [Attanasio, Kugler, and Meghir](#page-22-2) [\(2011\)](#page-22-2); [Attanasio et al.](#page-22-3) [\(2017\)](#page-22-3); [Blanco and Flores-Lagunes](#page-22-4) [\(2018\)](#page-22-4). Persuasion effects of political campaigns are analyzed by [DellaVigna and Kaplan](#page-23-4) [\(2007\)](#page-23-4); [DellaVigna and Gentzkow](#page-23-5) [\(2010\)](#page-23-5). Education interventions are investigated by [Krueger and Whitmore](#page-24-3) [\(2001\)](#page-24-3); [Angrist, Bettinger, and Kremer](#page-22-5) [\(2006\)](#page-22-5); [Angrist, Lang, and](#page-22-6) [Oreopoulos](#page-22-6) [\(2009\)](#page-22-6); [Chetty et al.](#page-22-7) [\(2011\)](#page-22-7); [Dobbie and Jr.](#page-23-6) [\(2015\)](#page-23-6). Medical treatments are studied by [CASS](#page-22-8) [\(1984\)](#page-22-8); [Sexton and Hebel](#page-24-4) [\(1984\)](#page-24-4); [U.S. Department of Health and Human Services](#page-24-5) [\(2004\)](#page-24-5).

 3 The probability of causation parameter is also called persuasion effect when treatments are predominantly informational [\(Jun and Lee,](#page-23-7) [2020\)](#page-23-7).

Our partial identification strategies are based on three increasingly restrictive sets of assumptions. They extend the identification of probabilities of causation to scenarios with endogenous sample selection. In our model, treatment effects can be related to the sample selection mechanism even though treatment take-up is exogenous.

Our first identification result relies on a monotone sample selection mechanism. This condition imposes that treatment has a non-negative effect on the sample selection indicator for all individuals. In the job training example, this restriction implies that the treatment can move workers into employment but never out of employment.

To tighten the identified bounds, our second result further assumes a monotone treatment response. This condition imposes that treatment has a non-negative effect on the potential outcomes for all individuals. In the job training example, this restriction implies that the treatment can move workers into formal jobs but never into informal jobs.

To further reduce the identified set, our final result additionally relies on a stochastic dominance assumption. This condition imposes that the sub-population that self-selects into the sample regardless of the treatment status has higher treated potential outcomes than the sub-population that self-selects into the sample only when treated. In the job training example, this restriction implies that the agents who are always employed are more likely to have a formal job if treated than the agents who are employed only when treated.

To empirically illustrate the usefulness of our approach, we provide bounds for the probability of causation of an intensive training program: *Jóvenes en Acción*. This program aimed to improve the labor market prospects and, in particular, the quality of jobs held by disadvantaged youths in seven large cities in Colombia. It offered in-classroom intensive training in occupational skills to qualify unemployed individuals for locally demanded jobs. Additionally, it focused on socioemotional development and offered on-the-job internships with formal employers.

Previous research [\(Attanasio, Kugler, and Meghir,](#page-22-2) [2011;](#page-22-2) [Attanasio et al.,](#page-22-3) [2017\)](#page-22-3) finds that this program had positive effects on employment and unconditional formality. However, less is known about whether the program achieves its goal of improving job quality conditioning on having a job. We study effects on the quality margin of jobs by considering the share of women that transitioned to the formal labor market *because* they participated in the training program.[4](#page-3-0) We find that incorporating selection and bounding the probability of causation leads to a less optimistic view of the program's impacts. Despite its efforts to connect youths with formal enterprises as trainees, we find that at most 19% of the women switched their status because they were assigned to the *Jovenes en Acción* program.

Concerning its theoretical contribution, our work is inserted in two research areas: identification of probabilities of causation and identification in the presence of sample selection.

[Heckman, Smith, and Clements](#page-23-2) [\(1997\)](#page-23-2) motivate the focus on a parameter closely connected to the probability of causation based on the political economy of policy evaluation. They argue that a program would only be adopted in a democracy if it benefited most people in the population. They make either strong probabilistic assumptions or parametric assumptions to point-identify this parameter, while we focus entirely on partial identification strategies that are based on a menu of easily interpretable assumptions.

[Pearl](#page-24-0) [\(1999\)](#page-24-0) and [Tian and Pearl](#page-24-1) [\(2000\)](#page-24-1) discuss how to interpret and partially identify probabilities of causation in a single population where agents are always observed. [Cinelli](#page-23-1) [and Pearl](#page-23-1) [\(2021\)](#page-23-1) extend their work by combining experimental results from multiple trials to extrapolate probabilities of causation from one population to a different population. Moreover, [Jun and Lee](#page-23-0) [\(2019\)](#page-23-0) extend their work by considering endogenous selection into treatment.

We extend the work by [Pearl](#page-24-0) [\(1999\)](#page-24-0) and [Tian and Pearl](#page-24-1) [\(2000\)](#page-24-1) to a different direction. We identify probabilities of causation when the agents' realized outcomes may not be observed due to endogenous sample selection. To do so, we combine the tools developed in the literature about probabilities of causation with the trimming bounds developed in the sample selection literature [\(Horowitz and Manski,](#page-23-8) [1995;](#page-23-8) [Lee,](#page-24-2) [2009;](#page-24-2) [Chen and Flores,](#page-22-0) [2015;](#page-22-0) [Bartalotti, Kedagni,](#page-22-1) [and Possebom,](#page-22-1) [2022\)](#page-22-1).

Concerning its empirical contribution, our work is inserted in the literature about job training programs. [Attanasio, Kugler, and Meghir](#page-22-2) [\(2011\)](#page-22-2) and [Attanasio et al.](#page-22-3) [\(2017\)](#page-22-3) analyze

⁴In Colombia, as in many developing countries, formal jobs are usually linked to benefits, such as health insurance, which may greatly extend individual welfare.

the average treatment effect (ATE) of *J*óvenes en Acción on short and long-term outcomes associated with labor force attachment. We extend their work by analyzing a treatment effect parameter that focuses on job quality instead of labor force attachment. Importantly, [Blanco](#page-22-4) [and Flores-Lagunes](#page-22-4) [\(2018\)](#page-22-4) also analyzes the impact of a job training program on job quality using partial identification strategies. However, we focus on different contexts (Job Corps v. Jóvenes en $Acci\acute{o}n$) and on different target parameters (Quantile Treatment Effects v. Probabilities of Causation).

This paper is organized as follows. Section [2](#page-4-0) presents our structural model, sample selection mechanism, and identifying assumptions. It also discusses the testable restrictions imposed by our model. Section [3](#page-9-0) describes our main identification results, while Section [4](#page-12-0) intuitively explains them using a numerical example. Moreover, Section [5](#page-16-0) discusses how to estimate our identified bounds and the results of our empirical application. At the end, Section [6](#page-21-0) concludes. We also have an online appendix with the proofs of all our results.

2 Analytical Framework

We aim to identify the probability of causation [\(Pearl,](#page-24-0) [1999;](#page-24-0) [Tian and Pearl,](#page-24-1) [2000;](#page-24-1) [Jun](#page-23-0) [and Lee,](#page-23-0) [2019;](#page-23-0) [Cinelli and Pearl,](#page-23-1) [2021\)](#page-23-1) within the always-observed subsample. To do so, we consider the generalized sample selection model [\(Lee,](#page-24-2) [2009\)](#page-24-2), described in the potential outcomes framework:

$$
\begin{cases}\nY^* &= Y_1^* \cdot D + Y_0^* \cdot (1 - D) \\
S &= S_1 \cdot D + S_0 \cdot (1 - D) \\
Y &= Y^* \cdot S\n\end{cases}
$$
\n(1)

where D is the treatment status indicator (in our application, being selected to enroll in the *Jóvenes in Acción* training program). The variable Y^* is the possibly censored realized outcome variable (indicator for whether the agent has a formal or informal job) with support $\mathcal{Y} = \{0, 1\}$, while Y_0^* and Y_1^* are the possibly censored potential outcomes when the person is untreated and treated, respectively. Similarly, S is the realized sample selection indicator (indicator for whether the agent holds a job), and S_0 and S_1 are potential sample selection indicators when individuals are untreated and treated. Finally, Y is the uncensored observed outcome. The researcher observes only the vector (Y, D, S) , while Y_1^*, Y_0^*, S_1 and S_0 are latent variables.^{[5](#page-5-0)}

In the setting analyzed here, the task of learning about the probability of causation [\(Pearl,](#page-24-0) [1999;](#page-24-0) [Tian and Pearl,](#page-24-1) [2000;](#page-24-1) [Jun and Lee,](#page-23-0) [2019;](#page-23-0) [Cinelli and Pearl,](#page-23-1) [2021\)](#page-23-1) is further complicated by the potential for nonrandom sample selection. As pointed out by [Lee](#page-24-2) [\(2009\)](#page-24-2), even in the simpler case of the average treatment effect (ATE), point identification is no longer possible, leading him to derive bounds for the ATE.

This paper combines the insights of these literatures to develop sharp bounds for the probability of causation under sample selection. To do so, we define four latent groups based on the potential sample selection indicators. The sub-populations are defined as: alwaysobserved $(S_0 = 1, S_1 = 1)$, observed-only-when-treated $(S_0 = 0, S_1 = 1)$, observed-only-whenuntreated $(S_0 = 1, S_1 = 0)$, and never-observed $(S_0 = 0, S_1 = 0)$.^{[6](#page-5-1)} They are denoted by OO, NO, ON and NN respectively.

Following [Zhang, Rubin, and Mealli](#page-24-6) [\(2008\)](#page-24-6) and [Lee](#page-24-2) [\(2009\)](#page-24-2), we focus on the alwaysobserved sub-population $(S_0 = 1, S_1 = 1)$. Importantly, this sub-population is the only group with censored potential outcomes observed in both treatment arms. For the other three subpopulations, treatment effect parameters are not point-identified or bounded in a non-trivial way without further parametric assumptions since at least one of the potential outcomes (Y_0^*) or Y_1^*) is never observed.^{[7](#page-5-2)} Since our focus is on a fully non-parametric identification strategy, we do not discuss parametric identification of unconditional treatment effect parameters or treatment effect parameters associated with the latent groups ON , NO and NN .

Our target parameter is the probability of causation within the sub-population that is

 5 For simplicity, we drop exogenous covariates from the model. All results derived in the paper hold conditionally on covariates.

 6 Since the conditioning sub-population is determined by post-treatment outcomes, our work is also connected to the statistical literature known as principal stratification [\(Frangakis and Rubin,](#page-23-9) [2002\)](#page-23-9), in which the four latent groups would be called strata.

⁷In some applications (e.g., analyzing the impact of a medical treatment on a health quality measure where selection is given by whether the patient is alive), the potential censored outcome Y_d^* is not even properly defined when $S_d = 0$ for $d \in \{0, 1\}.$

always observed:

$$
\theta^{OO} = \mathbb{P}\left[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1\right].\tag{2}
$$

The unconditional probability of causation $(\mathbb{P}[Y_1^* = 1 | Y_0^* = 0])$ captures, within the subpopulation whose untreated potential outcome is equal to zero, the share whose treated potential outcome is equal to one. Intuitively, it measures the share of agents who benefited from the treatment within the subgroup with a negative untreated outcome. In our empirical application, the unconditional probability of causation captures, within the population with an informal job if untreated, the share of workers with a formal job if treated. Our target parameter in Equation [\(2\)](#page-6-0) focuses on the probability of causation for the always-observed latent group. In our empirical application, our target parameter captures, within the population who is employed regardless of treatment status and has an informal job if untreated, the share of workers with a formal job if treated. Intuitively, we focus on the population who is always-employed and found a job of higher observable quality because they were assigned to the *J*_{ovenes} in *Accion* training program.

Analogously to [Heckman, Smith, and Clements](#page-23-2) [\(1997\)](#page-23-2), [Jun and Lee](#page-23-0) [\(2019\)](#page-23-0) and [Cinelli and](#page-23-1) [Pearl](#page-23-1) [\(2021\)](#page-23-1), identification of θ^{OO} is complicated because it depends on the joint distribution of the potential outcomes (Y_0^*, Y_1^*) while, even in a randomized controlled trial, we can only identify the marginal distributions of the potential outcomes. Analogously to [Lee](#page-24-2) [\(2009\)](#page-24-2), identification of θ^{OO} is complex because sample selection is nonrandom and possibly impacted by the treatment.

To simultaneously address these issues, we consider three sets of assumptions to partially identify our target parameter. The identified set shrinks when stronger assumptions are used.^{[8](#page-6-1)} Assumptions [1-](#page-6-2)[3](#page-7-0) are sufficient to partially identify θ^{OO} .

Assumption 1 (Random Assignment) Treatment D is randomly assigned, i.e., D \perp $(Y_0^*, Y_1^*, S_0, S_1).$

⁸According to [Tamer](#page-24-7) [\(2010,](#page-24-7) p. 167), this approach to identification "characterizes the informational content of various assumptions by providing a menu of estimates, each based on different sets of assumptions, some of which are plausible and some of which are not." This approach was called a layered policy analysis by [Manski](#page-24-8) [\(2011\)](#page-24-8) and is illustrated empirically by [Manski and Molinari](#page-24-9) [\(2020\)](#page-24-9).

Assumption [1](#page-6-2) is a modification of the standard independence assumption [\(Imbens and](#page-23-10) [Wooldridge,](#page-23-10) [2009\)](#page-23-10) to account for sample selection. Instead of assuming that the treatment variable is independent of the potential outcomes only, we also assume independence between the treatment variable and the potential sample selection indicators similarly to [Lee](#page-24-2) [\(2009\)](#page-24-2). In our empirical application, it holds conditionally on course indicators because the possibility of enrolling in the *J*óvenes in Acción training program was randomly allocated within oversubscribed courses.

Assumption 2 (Positive Mass) Both treatment groups and the always-observed sub-population who chooses $Y_0^* = 0$ exist, i.e., $0 < \mathbb{P}[D = 1] < 1$ and $\mathbb{P}[Y_0^* = 0, S_0 = 1, S_1 = 1] > 0$.

Assumption [2](#page-7-1) is crucial for the identification results because it ensures that our subpopulation of interest exists. In our empirical application, it requires that there are alwaysemployed individuals who have an informal jobs when untreated.

Assumption 3 (Monotone Sample Selection) Treatment has a non-negative effect on the sample selection indicator for all individuals, i.e., $S_1 \geq S_0$.

Assumption [3](#page-7-0) is a monotonicity restriction that rules out the existence of the observedonly-when-untreated sub-population and is commonly used in the literature about sample selection [\(Lee,](#page-24-2) [2009;](#page-24-2) [Chen and Flores,](#page-22-0) [2015;](#page-22-0) [Bartalotti, Kedagni, and Possebom,](#page-22-1) [2022\)](#page-22-1).^{[9](#page-7-2)} In our empirical application, it imposes that the *Jóvenes in Acción* training program can only move agents into employment.

Assumptions [1-](#page-6-2)[3](#page-7-0) form our first set of assumptions required to partially identify the probability of causation within the always-observed individuals. Importantly, this set of assumptions has a testable implication, as discussed in Lemma [1.](#page-9-1) Even though these assumptions are sufficient to partially identify θ^{OO} , the identified set can be substantially tightened by additionally imposing that the treatment can only increase the possibly censored potential outcome.

⁹As in [Lee](#page-24-2) [\(2009\)](#page-24-2), this assumption can be stated as $S_1 \geq S_0$ with probability 1. For the sake of simplicity, we assume it holds for all individuals. All results can be stated with some straightforward changes if the inequality in Assumption [3](#page-7-0) holds in the opposite direction.

Assumption 4 (Monotone Treatment Response) Treatment has a non-negative effect on the censored outcome variable for all individuals, i.e., $Y_1^* \ge Y_0^*$.

Assumption [4](#page-7-3) is a monotonicity restriction that is common in the partial identification literature [\(Manski,](#page-24-10) [1997;](#page-24-10) [Manski and Pepper,](#page-24-11) [2000;](#page-24-11) [Jun and Lee,](#page-23-0) [2019\)](#page-23-0).^{[10](#page-8-0)} In our empirical application, it imposes that the *J*_{ovenes} in Acci^{on} training program can only move agents from informal jobs to formal ones.

Assumptions [1](#page-6-2)[-4](#page-7-3) form our second set of assumptions required to partially identify the probability of causation within the always-observed individuals. Importantly, this set of assumptions has an extra testable implication, as discussed in Proposition [1.](#page-9-2)

We can further shrink the identified set around θ^{OO} by adding Assumption [5](#page-8-1) and completing our final set of identifying assumptions.

Assumption 5 (Stochastic Dominance) The treated counterfactual for the always-observed group stochastically dominates the treated counterfactual for the observed-only-when-treated, *i.e.*, $\mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] \ge \mathbb{P}[Y_1^* = 1 | S_0 = 0, S_1 = 1].$

Assumption [5](#page-8-1) is a stochastic dominance restriction that imposes that the always-observed sub-population has higher potential treated outcomes than the observed-only-when-treated group. This type of assumption is common in the literature [\(Imai,](#page-23-11) [2008;](#page-23-11) [Blanco, Flores, and](#page-22-9) [Flores-Lagunes,](#page-22-9) [2013;](#page-22-9) [Huber and Mellace,](#page-23-12) [2015;](#page-23-12) [Huber, Laffers, and Mellace,](#page-23-13) [2017;](#page-23-13) [Bartalotti,](#page-22-1) [Kedagni, and Possebom,](#page-22-1) [2022\)](#page-22-1) and is intuitively based on the argument that some sub-groups have more favorable underlying characteristics than others.^{[11](#page-8-2)} In our empirical application, it imposes that the always-employed sub-population has higher potential formality when treated than the employed-only-when-treated sub-population.

¹⁰This assumption can be stated as $Y_1^* \geq Y_0^*$ with probability 1. For the sake of simplicity, we assume it holds for all individuals. All results can be stated with straightforward changes if the inequality in Assumption [4](#page-7-3) holds in the opposite direction.

 11 All of our results can be stated if the inequality in Assumption 5 holds in the opposite direction.

2.1 Testable Restrictions

This subsection discusses testable restrictions implied by the assumptions described in Section [2.](#page-4-0)

First, the testable restriction implied by Assumptions [1-](#page-6-2)[3](#page-7-0) was already derived by [Lee](#page-24-2) [\(2009\)](#page-24-2). We state it here for completeness.

Lemma 1 Under Assumptions [1-](#page-6-2)[3,](#page-7-0) the following inequality holds:

$$
\mathbb{P}[S=1|D=1]-\mathbb{P}[S=1|D=0]\geq 0.
$$

Second, we derive a set of testable restrictions implied by Assumptions [1-](#page-6-2)[4](#page-7-3) as detailed in Proposition [1.](#page-9-2) Its proof is in Appendix [A.1.](#page-25-0)

Proposition 1 Under Assumptions [1-](#page-6-2)[4,](#page-7-3) the following inequalities hold:

$$
\mathbb{P}\left[S=1|D=1\right] - \mathbb{P}\left[S=1|D=0\right] \ge 0,\tag{3}
$$

$$
\mathbb{P}\left[Y=1|D=1\right] - \mathbb{P}\left[Y=1|D=0\right] \ge 0. \tag{4}
$$

Intuitively, the monotonicity of the sample selection indicator and the censored potential outcome implies that treatment positively affects the uncensored potential outcome.

These restrictions can be easily tested using two one-sided tests of mean differences. To be conservative, we recommend testing them as separate null hypotheses without correcting for multiple hypotheses testing.

3 Identification Results

In this section, we partially identify the probability of causation within the always-observed sub-population (Equation (2)). In the next section, we illustrate these results with a numerical example that captures the intuition behind the formal results presented here.

Combining Assumptions [1-](#page-6-2)[3,](#page-7-0) we derive sharp bounds around θ^{OO} as detailed in Proposition [2.](#page-10-0) Its proof is in Appendix [A.2.](#page-25-1)

Proposition 2 Under Assumptions [1-](#page-6-2)[3,](#page-7-0) the probability of causation is partially identified for the always-observed subgroup, i.e.,

$$
LB_1 \leq \theta^{OO} \leq UB_1,
$$

where

$$
LB_{1} := \max \left\{ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)}{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]} + \mathbb{P}\left[Y=0|S=1,D=0\right] - 1}{\mathbb{P}\left[Y=0|S=1,D=0\right]}, 0 \right\}
$$

and

$$
UB_1 := \min\left\{\frac{\mathbb{P}\left[Y=1|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]}}{\mathbb{P}\left[Y=0|S=1,D=0\right]},1\right\}.
$$

In practice, these bounds may be uninformative even though they are sharp. To derive tighter bounds, researchers can add increasingly stronger assumptions. Even though the credibility of these assumptions depends on their empirical contexts, applied researchers frequently have some prior about the direction of the treatment effect. Using this prior, the researcher can impose the monotone treatment response condition.

Formally, combining Assumptions [1-](#page-6-2)[4,](#page-7-3) we derive sharp bounds around the probability of causation within the always-observed sub-population as detailed in Proposition [3.](#page-10-1) Its proof is in Appendix [A.4.](#page-44-0)

Proposition 3 Under Assumptions [1-](#page-6-2)[4,](#page-7-3) the probability of causation is partially identified for

the always-observed subgroup, i.e.,

$$
LB_2 \leq \theta^{OO} \leq UB_2,
$$

where

$$
LB_{2} := \max \left\{ \frac{\frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)}{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}} + \mathbb{P}\left[Y=0|S=1,D=0\right] - 1}{\mathbb{P}\left[Y=0|S=1,D=0\right]}, 0 \right\}
$$

and

$$
UB_2 := \min\left\{\frac{\mathbb{P}\left[Y=1|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]} + \mathbb{P}\left[Y=0|S=1,D=0\right] - 1}{\mathbb{P}\left[Y=0|S=1,D=0\right]},1\right\}
$$

.

Moreover, these bounds are sharp.

Note that the identifying power of Assumption [4](#page-7-3) is illustrated by a smaller upper bound in Proposition [3](#page-10-1) in comparison with Proposition [2.](#page-10-0)

To achieve even tighter bounds, researchers can impose the stochastic dominance condi-tion. Formally, combining Assumptions [1-](#page-6-2)[5,](#page-8-1) we derive sharp bounds around θ^{OO} as detailed in Proposition [4.](#page-11-0) Its proof is in Appendix [A.6.](#page-55-0)

Proposition 4 Under Assumptions [1-](#page-6-2)[5,](#page-8-1) the probability of causation is partially identified for the always-observed subgroup, i.e.,

$$
LB_3 \leq \theta^{OO} \leq UB_3,
$$

where

$$
LB_3 := \max\left\{\frac{\mathbb{P}\left[Y=1|S=1,D=1\right]+\mathbb{P}\left[Y=0|S=1,D=0\right]-1}{\mathbb{P}\left[Y=0|S=1,D=0\right]},0\right\}
$$

and

$$
UB_3 := \min\left\{\frac{\mathbb{P}\left[Y=1|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]}+\mathbb{P}\left[Y=0|S=1,D=0\right]-1}{\mathbb{P}\left[Y=0|S=1,D=0\right]},1\right\}.
$$

Moreover, these bounds are sharp.

Note that the identifying power of Assumption [5](#page-8-1) is illustrated by a larger lower bound in Proposition [4](#page-11-0) in comparison with Proposition [3.](#page-10-1)

4 Numerical Example

In this section, we use a numerical example to intuitively explain our partial identification results from Section [3.](#page-9-0) We focus on understanding the factors that determine the length of our bounds in each proposition and the reason why each additional assumption tightens our bounds.

Let our data-generating process be given by

$$
\mathbb{P}[Y_0^* = 1, Y_1^* = 1, S_0 = 1, S_1 = 1 | D = d] = \frac{2}{16}
$$

$$
\mathbb{P}[Y_0^* = 0, Y_1^* = 1, S_0 = 1, S_1 = 1 | D = d] = \frac{4}{16}
$$

$$
\mathbb{P}[Y_0^* = 1, Y_1^* = 0, S_0 = 1, S_1 = 1 | D = d] = 0
$$

$$
\mathbb{P}[Y_0^* = 0, Y_1^* = 0, S_0 = 1, S_1 = 1 | D = d] = \frac{3}{16}
$$

$$
\mathbb{P}[Y_0^* = 1, Y_1^* = 1, S_0 = 0, S_1 = 1 | D = d] = \frac{1}{16}
$$

$$
\mathbb{P}[Y_0^* = 0, Y_1^* = 1, S_0 = 0, S_1 = 1 | D = d] = \frac{1}{16}
$$

$$
\mathbb{P}[Y_0^* = 1, Y_1^* = 0, S_0 = 0, S_1 = 1 | D = d] = 0
$$

$$
\mathbb{P}[Y_0^* = 0, Y_1^* = 0, S_0 = 0, S_1 = 1 | D = d] = \frac{2}{16}
$$

$$
\mathbb{P}[Y_0^* = 1, Y_1^* = 1, S_0 = 1, S_1 = 0 | D = d] = 0
$$

$$
\mathbb{P}[Y_0^* = 0, Y_1^* = 1, S_0 = 1, S_1 = 0 | D = d] = 0
$$

$$
\mathbb{P}[Y_0^* = 1, Y_1^* = 0, S_0 = 1, S_1 = 0 | D = d] = 0
$$

$$
\mathbb{P}[Y_0^* = 1, Y_1^* = 0, S_0 = 1, S_1 = 0 | D = d] = 0
$$

$$
\mathbb{P}\left[Y_0^* = 0, Y_1^* = 0, S_0 = 1, S_1 = 0 | D = d\right] = 0
$$

$$
\mathbb{P}\left[Y_0^* = 1, Y_1^* = 1, S_0 = 0, S_1 = 0 | D = d\right] = \frac{1}{16}
$$

$$
\mathbb{P}\left[Y_0^* = 0, Y_1^* = 1, S_0 = 0, S_1 = 0 | D = d\right] = \frac{1}{16}
$$

$$
\mathbb{P}\left[Y_0^* = 1, Y_1^* = 0, S_0 = 0, S_1 = 0 | D = d\right] = 0
$$

$$
\mathbb{P}\left[Y_0^* = 0, Y_1^* = 0, S_0 = 0, S_1 = 0 | D = d\right] = \frac{1}{16}
$$

for any $d \in \{0, 1\}$ and $\mathbb{P}\left[D = 1\right] = \frac{1}{2}$.

Note that this data-generating process satisfies Assumptions [1](#page-6-2)[-4](#page-7-3) by construction. Observe also that $\mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] = \frac{2}{3}$ and $\mathbb{P}[Y_1^* = 1 | S_0 = 0, S_1 = 1] = \frac{1}{2}$, implying that Assumption [5](#page-8-1) is valid too.

Finally, notice that our target parameter — the probability of causation for the alwaysemployed — is given by

$$
\theta^{OO} = \mathbb{P}\left[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1\right] \approx 0.571.
$$

Now, we carefully derive our bounds to understand the factors that determine the length of our bounds in each proposition and the reason why each additional assumption tightens our bounds.

To understand the intuition behind Proposition [2,](#page-10-0) note that

$$
\theta^{OO} = \mathbb{P}\left[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1\right]
$$

$$
= \frac{\mathbb{P}\left[Y_0^* = 0, Y_1^* = 1 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y_0^* = 0 | , S_0 = 1, S_1 = 1\right]}.
$$

Since the denominator is point-identified by $\mathbb{P}[Y=0|S=1,D=0]$ (Lemma [A.4\)](#page-26-0), we have that

$$
\theta^{OO} = \frac{\mathbb{P}\left[Y_0^* = 0, Y_1^* = 1 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y = 0 | S = 1, D = 0\right]}.
$$
\n(5)

We want to bound the numerator in Equation [\(5\)](#page-13-0) using information from the marginal distributions of $Y_0^* | (S_0 = 1, S_1 = 1)$ and $Y_1^* | (S_0 = 1, S_1 = 1)$. To do so, we use the BooleFrechet inequalities (Lemma [A.1\)](#page-9-1) and find that

$$
\theta^{OO} \le \frac{\min \{ \mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1], \mathbb{P}[Y_0^* = 0 | S_0 = 1, S_1 = 1] \}}{\mathbb{P}[Y = 0 | S = 1, D = 0]}
$$

and that

$$
\theta^{OO} \ge \frac{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1}{\mathbb{P}\left[Y = 0 | S = 1, D = 0\right]}.\tag{6}
$$

Note, once more, that $\mathbb{P}[Y_0^* = 0 | S_0 = 1, S_1 = 1]$ is point-identified by $\mathbb{P}[Y = 0 | S = 1, D = 0]$ (Lemma [A.4\)](#page-26-0), implying that

$$
\theta^{OO} \le \min\left\{ \frac{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y = 0 | S = 1, D = 0\right]}, 1 \right\}
$$

and that

$$
\theta^{OO} \ge \frac{\mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] + \mathbb{P}[Y = 0 | S = 1, D = 0] - 1}{\mathbb{P}[Y = 0 | S = 1, D = 0]}.
$$
\n(7)

Now, we address the sample selection issue that is present in the term $\mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1]$. To do so, we use the trimming bounds proposed by [Horowitz and Manski](#page-23-8) [\(1995\)](#page-23-8) and [Lee](#page-24-2) [\(2009\)](#page-24-2) (Lemma [A.2\)](#page-26-1) and find that

$$
\theta^{OO} \le \min \left\{ \frac{\frac{\mathbb{P}[Y=1|S=1,D=1]}{\mathbb{P}[S_0=1,S_1=1|S_1=1]}}{\frac{\mathbb{P}[Y=0|S=1,D=0]}{\mathbb{P}[Y=0|S=1,D=0]}}, 1 \right\}
$$

and that

$$
\theta^{OO} \ge \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - (1-\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right)]}{\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]} + \mathbb{P}\left[Y=0|S=1,D=0\right] - 1}{\mathbb{P}\left[Y=0|S=1,D=0\right]}.
$$
\n(8)

The last two inequalities illustrate the first factor that intuitively explains the length of our bounds. Observe that the upper bound is smaller and the lower bound is greater if the share of the always-employed among the ones who are employed when treated ($\mathbb{P}\left[S_0 = 1, S_1 = 1 | S_1 = 1\right]$) is large.

Finally, to derive the last expression of the bounds in Proposition [2,](#page-10-0) we use Assumption [3](#page-7-0) to pointy identify $\mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]$ (Lemma [A.3\)](#page-26-2). Applying the analytic expressions from Proposition [2,](#page-10-0) our data generating process implies that $LB_1 \approx 0.286$ and $UB_1 = 1$.

Now, we focus on the bounds in Proposition [3.](#page-10-1) Since $LB_2 = LB_1$ and $UB_2 \le UB_1$, we want to understand why Assumption [4](#page-7-3) is able to reduce the upper bound around the target parameter. Using the Monotone Treatment Response Assumption, the joint probability $\mathbb{P}[Y_0^* = 0, Y_1^* = 1 | S_0 = 1, S_1 = 1]$ is equal to $\mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] + \mathbb{P}[Y_0^* = 0 | S_0 = 1, S_1 = 1] -$ 1 (Lemma [A.5\)](#page-44-1). Combining this result with Equation [5,](#page-13-0) we find that

$$
\theta^{OO} = \frac{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1}{\mathbb{P}\left[Y = 0 | S = 1, D = 0\right]}.
$$
\n(9)

Since the right-hand side term in Equation (9) is equal to the lower bound in Inequality (6) , we can conclude that the upper bound in Proposition [3](#page-10-1) is less than or equal to the upper bound in Proposition [2.](#page-10-0) This result intuitively explains the identifying power of Assumption [4.](#page-7-3)

Now, to derive the last expression of the bounds in Proposition [3,](#page-10-1) we follow the same steps used to derive the bounds in Proposition [2.](#page-10-0) Finally, applying the analytic expressions from Proposition [3,](#page-10-1) our data generating process implies that $LB_2 \approx 0.286$ and $UB_2 \approx 0.857$, numerically illustrating that Assumption [4](#page-7-3) reduces the upper bound substantially.

To end this section, we focus on the bounds in Proposition [4.](#page-11-0) Since $UB_3 = UB_2$ and $LB_3 \geq$ $LB₂$, we want to understand why Assumption 5 is able to increase the lower bound around the target parameter. To do so, we go back to Inequality [\(7\)](#page-14-1). Since $\mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] \ge$ $\mathbb{P}[Y=1|S=1,D=1]$ due to the stochastic dominance assumption (Lemma [A.6\)](#page-55-1), there is no need to use the trimming bounds in Inequality [\(8\)](#page-14-2). Consequently, we have that

$$
\theta^{OO} \ge \frac{\mathbb{P}\left[Y=1|S=1,D=1\right]+\mathbb{P}\left[Y=0|S=1,D=0\right]-1}{\mathbb{P}\left[Y=0|S=1,D=0\right]},
$$

which is greater than the expression in Inequality (8) and the lower bound in Proposition [3.](#page-10-1) This result intuitively explains the identifying power of Assumption [5.](#page-8-1)

Finally, applying the analytic expressions from Proposition [4,](#page-11-0) our data generating process implies that $LB_3 \approx 0.505$ $LB_3 \approx 0.505$ $LB_3 \approx 0.505$ and $UB_3 \approx 0.857$, numerically illustrating that Assumption 5 increases the lower bound substantially. Importantly, our shortest identified interval contains the target parameter and is informative.

We can also compare our identified bounds against an estimand that would identify the probability of causation if Assumptions [1](#page-6-2)[-4](#page-7-3) were valid and all agents were observed $(\mathbb{P}[S_0 = 1, S_1 = 1] = 1)$. In this case, the probability of causation would be point-identified by the lower bound LB_3 in Proposition [4.](#page-11-0) If we ignored sample selection and used this estimand, we would underestimate the true probability of causation for the always-employed in this numerical example.

5 Empirical Application: Trasition into Formality in the Jóvenes in Acción Training Program

Our application relates to the large job training program *Jóvenes en Acción*. The program's main goals were to increase labor market attachment and job quality among disadvantaged youth aged between 18 and 28 living in Colombia's largest cities. The intervention lasted approximately six months and was based on three main components: (i) three months in private institutions, during which trainees received occupational-specific training and developed soft skills, such as proactive behavior, resourcefulness, openness to feedback and teamwork; (ii) three months of on-the-job training provided by legally registered companies in the form of an unpaid internship; (iii) elaboration of a project of life, orienting youth towards a positive visualization of their abilities and work perspectives.[12](#page-16-1)

We are interested in learning about the share of the population who became formal because they were selected for the *J*óvenes en Acción training program. The short term experimental effects of the program have been described extensively in [Attanasio, Kugler, and Meghir](#page-22-2) [\(2011\)](#page-22-2). In what follows, we focus on women since the results for this sub-sample are starker

¹²Training providers were paid according to students' course progression and received a bonus if the trainees were hired by the firms that employed them during the internship. One interesting feature of the program was its high take-up: 97% of the individuals accepted it, which minimizes concerns about compliance problems and their consequences for the identification of policy-relevant parameters.

and, as described in [Attanasio, Kugler, and Meghir](#page-22-2) [\(2011\)](#page-22-2), there was significant differential sample selection into employment among women in the short-run. Women were 8% more likely to be employed, earned 20% higher salaries and were 35% more likely to have a formal job one year after the end of the program.^{[13](#page-17-0)}

As explained in Section [2,](#page-4-0) our target parameter is the probability of causation for the latent group that would be employed regardless of treatment assignment. We compute bounds around this parameter by considering the selection indicator as employment (either in the formal or the informal sector) and the variable of interest as an indicator that equals one if the person has a formal job and zero if the person has an informal job.

First, we report the share of the female population who is employed regardless of being assigned to the *J*_{ovenes} en Acci_{on} training program and, within this group, the probability of having an informal job when assigned to the control group. Since both objects are pointidentified under Assumptions [1-](#page-6-2)[3,](#page-7-0) we focus on our first set of assumptions when estimating them. In our sample, we find that 69.0% of the women are always-employed. Within this subgroup, we also estimate the probability of having an informal job when untreated as equal to 61.8% ^{[14](#page-17-1)}

Our main results are presented in Figure [1](#page-18-0) and Table [1.](#page-19-0) In Figure [1,](#page-18-0) the leftmost bars represent bounds for the target parameter without considering the stratified nature of the assignment mechanism, while the rightmost bars condition on a full set of course fixed ef-fects.^{[15,](#page-17-2)[16](#page-17-3)} Each bar represents the bounds under each group of assumptions [1-](#page-6-2)[3,](#page-7-0) 1-[4](#page-7-3) and

¹³These results suggest that our identifying assumptions are valid according to the testable restrictions described in Subsection [2.1.](#page-9-3)

¹⁴The percentile 90%-confidence intervals around these probabilities are $[65.5, 72.5]$ and $[57.3, 66.8]$, respectively.

 $15T₀$ estimate the unconditional bounds on the left side, we use sample means to estimate the probabilities in Propositions [2-](#page-10-0)[4](#page-11-0)

¹⁶When conditioning on covariates X in the right side, we identify bounds around the conditional probabilities of causation for the always-employed: $\theta^{OO}(x) := \mathbb{P}[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1, X = x]$ for every course x . These bounds can be expressed similarly to the ones derived in Propositions [2-](#page-10-0)[4](#page-11-0) by conditioning on X all probabilities that appear in these propositions. To estimate these conditional probabilities, we use fixed-effect models. After estimating bounds for each course, we average these bounds across the sample distribution of courses. By doing so, we estimate bounds around a summary measure of the conditional probabilities of causation for the always-employed: $SM = \int \theta^{OO}(x) dF_X(x)$. If $X \perp (Y_0^*, S_0, S_1)$, then $SM = \theta^{OO}$, implying that the bounds around the summary measure partially identify the unconditional probability of causation for the always-employed. Alternatively, we can analyze $\theta^{OO}(x)$ for pre-specified values of the covariates. These results are available upon request.

[1](#page-6-2)[-5,](#page-8-1) as presented in the graph's legend. Table [1](#page-19-0) presents, apart from the plotted estimates, 90%-confidence intervals around the target parameter [\(Imbens and Manski,](#page-23-14) [2004\)](#page-23-14). Since the bounds with course fixed effects and without them are very similar, we focus our discussion on the results that take the stratified randomization into account.

Figure 1: Bounds on the Probability of Causation in the *Jóvenes in Acción*

Notes: This figure presents lower and upper bounds on the probability of causation for the always-employed women using data from the job training program $Jóvenes en Acción$. The outcome of interest is formal employment one year after the training program, the selection indicator is employment, and the treatment is a randomized assignment indicator. The leftmost bars depict sample analogs of the expressions in Propositions [2-](#page-10-0)[4](#page-11-0) without considering the stratified nature of the assignment mechanism. The rightmost bars do the same considering the stratified nature of the assignment mechanism.

	Without Incorporating Strata			Incorporating Strata		
Assumptions	$1-3$	$1-4$	$1-5$	$1-3$	$1-4$	$1-5$
	$\left(1\right)$	$\left(2\right)$	$\left(3\right)$	(4)	(5)	(6)
LВ	0.014	0.014	0.106	0.057	0.057	0.121
UB	0.609	0.163	0.163	0.696	0.188	0.188
IM 90%-CI	[0, 0.664]	[0, 0.218]	[0.058, 0.219]	[0, 0.741]	[0, 0.318]	[0.010, 0.327]
Number of Observations	1,769	1,769	1,769	741	741	741

Table 1: Bounds on the Probability of Causation for Always Employed in the *Jóvenes in Acción* Training Program)

Notes: This table depicts bounds on the probability of causation for the always-employed women using data from the job training program *J*óvenes en Acción. The outcome of interest is formal employment one year after the training program, the selection indicator is employment, and the treatment is a randomized assignment indicator. Rows LB and UB report the lower and upper bounds around the target parameter. Row "IM 90%-CI" report 90%-confidence intervals around the target parameters as proposed by Imbens and [Manski](#page-23-15) [\(2004\)](#page-23-15). To estimate the standard errors that enter the formula proposed by Imbens and [Manski](#page-23-15) [\(2004\)](#page-23-15), we use the empirical bootstrap with 1,000 replications. Columns (1) , (2) and (3) do not take into consideration the stratified nature of the assignment mechanism. Columns (4) , (5) and (6) incorporate the stratified nature of the assignment mechanism by conditioning on course fixed effects. The sample size in columns (4), (5) and (6) is smaller due to restrictions on strata for which we can estimate meaningful probabilities of causation: we drop strata where (i) there was only one woman; (ii) there was no treated women; (iii) all treated or all control women were unemployed; (iv) all employed women were either in the formal or in the informal labor market. The numbers above the column numbers denote which assumptions areused to estimate the bounds around the target parameter.

We, now, consider the probability of causation for the always-employed women under Assumptions [1-](#page-6-2)[3.](#page-7-0) In this case, we only impose, beyond the usual independence assumption, that participation in the program does not deter employment (monotone sample selection). We find that the bounds are uninformative. In particular, the upper bound in Column (4) in Table [1](#page-19-0) is large, implying that our estimates are consistent with a wide range of values for the probability of causation $([0.057, 0.696])$.

We now discuss the bounds obtained by additionally imposing Assumption [4.](#page-7-3) In this case, we assume that participation in the program can only move agents from informal jobs to formal ones. By imposing a monotone treatment response, we decrease the upper bound substantially. Column (5) in Table [1](#page-19-0) suggests that the *J*óvenes en Acción Training Program formalized at most 19% of the population that is always employed and would have an informal job if untreated.

Finally, we discuss the bounds obtained by additionally imposing Assumption [5.](#page-8-1) In this case, we assume that the always-employed sub-population has higher potential formality when treated than the employed-only-when-treated sub-population. By imposing this stochastic dominance assumption, we increase the lower bound from 0.057 to 0.121. Consequently, Column (6) in Table [1](#page-18-0) suggests that the *J*óvenes en Acción Training Program formalized at least 12% of the population that is always employed and would have an informal job if untreated.

To incorporate sample uncertainty around our estimates, we use the confidence intervals proposed by [Imbens and Manski](#page-23-14) [\(2004\)](#page-23-14). These confidence intervals contain the probability of causation for the always-employed women with a probability of at least 90% asymptotically. We estimate them as equal to $[0, 0.741]$ $[0, 0.741]$ $[0, 0.741]$, $[0, 0.318]$ and $[0.010, 0.327]$ under Assumptions 1[-3,](#page-7-0) [1](#page-6-2)[-4](#page-7-3) and [1](#page-6-2)[-5,](#page-8-1) respectively. Consequently, using our most informative bounds, we can reject the null that no one benefits from the *J*óvenes en Acción Training Program.

6 Conclusion

This paper partially identifies the probability of causation for the always-observed subgroup when there is sample selection. This parameter is relevant for researchers aiming to describe policy-relevant treatment effects. Intuitively, it describes the share of the population that was induced by the treatment to switch from a negative to a positive state. We derive sharp bounds around this parameter under three increasingly restrictive sets of assumptions. The first set imposes an exogenous treatment and a monotone sample selection mechanism. To find a smaller upper bound, the second set additionally imposes the monotone treatment response assumption. As our final set of assumptions, we also impose a stochastic dominance assumption to find a larger lower bound.

To illustrate the usefulness of our partial identification strategy, we use experimental data from the Colombian job training program *Jóvenes en Acción*. This policy aimed to improve the labor market prospects and, in particular, the quality of jobs held by disadvantaged youths in seven large cities in Colombia.

Focusing on the sub-sample of women, we study the effects on the quality margin of jobs by considering the share of women that transitioned to the formal labor market because they were assigned to the program. Despite the positive effects on the share of women employed in the formal labor market [\(Attanasio, Kugler, and Meghir,](#page-22-2) [2011\)](#page-22-2), we find that incorporating selection and bounding the probability of causation leads to a less optimistic view of the program's impacts. More precisely, we find that at most 1 in every 5 women switched their formality status because they were assigned to the *J*_{ovenes} en Acción training program.

Beyond the analysis of job training programs, our partial identification strategy can be useful for researchers interested in assessing the impacts of interventions in the presence of sample selection. For example, when analyzing the effects of a political campaign in the presence of survey non-response, the researcher may be interested in identifying the share of the population who supports policy A when treated, given that they would support policy B if untreated. This double identification challenge also arises when researchers consider the effects of health interventions on health quality if agents may pass away, or the effects of

educational interventions on learning if there is selection into test-taking.

References

- Angrist, Joshua, Eric Bettinger, and Michael Kremer. 2006. "Long-Term Educational Consequences of Secondary School Vouchers: Evidence from Administrative Records in Colombia." The American Economic Review 96 (3):847–862. URL [http://www.jstor.org/](http://www.jstor.org/stable/30034075) [stable/30034075](http://www.jstor.org/stable/30034075). (Cited on page [2.](#page-1-3))
- Angrist, Joshua, Daniel Lang, and Philip Oreopoulos. 2009. "Incentives and Services for College Achievement: Evidence from a Randomized Trial." American Economic Journal: Applied Economics 1 (1):pp. 1–28. (Cited on page [2.](#page-1-3))
- Attanasio, Orazio, Arlen Guarin, Carlos Medina, and Costas Meghir. 2017. "Vocational Training for Disadvantaged Youth in Colombia: A Long-Term Follow-Up." American Economic Journal: Applied Economics 9 (2):pp. 131–143. (Cited on pages [2,](#page-1-3) [3,](#page-2-0) and [4.](#page-3-1))
- Attanasio, Orazio, Adriana Kugler, and Costas Meghir. 2011. "Subsidizing Vocational Training for Disadvantaged Youth in Colombia: Evidence from a Randomized Trial." American Economic Journal: Applied Economics 3 (3):pp. 188–220. (Cited on pages [2,](#page-1-3) [3,](#page-2-0) [4,](#page-3-1) [17,](#page-16-2) [18,](#page-17-4) and [22.](#page-21-1))
- Bartalotti, Otavio, Desire Kedagni, and Vitor Possebom. 2022. "Identifying Marginal Treatment Effects in the Presence of Sample Selection." Journal of Econometrics Available at <https://doi.org/10.1016/j.jeconom.2021.11.011>. (Cited on pages [2,](#page-1-3) [4,](#page-3-1) [8,](#page-7-6) and [9.](#page-8-4))
- Blanco, German, Carlos A. Flores, and Alfonso Flores-Lagunes. 2013. "Bounds on Average and Quantile Treatment Effects of Job Corps Training on Wages." Journal of Human Resources 48 (3):pp. 659–701. (Cited on page [9.](#page-8-4))
- Blanco, German and Alfonso Flores-Lagunes. 2018. "Does Youth Training Lead to Better Job Quality: Evidence from Job Corps." Available at [https://drive.google.com/file/](https://drive.google.com/file/d/1gkkvK_gupfEyYGgDr3b-K8-n9pDpLBfe/view) [d/1gkkvK_gupfEyYGgDr3b-K8-n9pDpLBfe/view](https://drive.google.com/file/d/1gkkvK_gupfEyYGgDr3b-K8-n9pDpLBfe/view). (Cited on pages [2](#page-1-3) and [5.](#page-4-1))
- CASS. 1984. "Myocardial Infarction and Mortality in the Coronary Artery Surgery Study (CASS) Randomized Trial." The New England Journal of Medicine 310 (12):pp. 750–758. (Cited on page [2.](#page-1-3))
- Chen, Xuan and Carlos A. Flores. 2015. "Bounds on Treatment Effects in the Presence of Sample Selection and Noncompliance: The Wage Effects of Job Corps." Journal of Business and Economic Statistics 33 (4):pp. 523–540. (Cited on pages [2,](#page-1-3) [4,](#page-3-1) [8,](#page-7-6) and [56.](#page-55-2))
- Chetty, Raj, John N. Friedman, Nathaniel Hilger, Emmanuel Saez, Diane Whitmore Schanzenbach, and Danny Yagan. 2011. "How Does Your Kindergarten Classroom Affect Your Earnings? Evidence from Project Star." The Quarterly Journal of Economics 126 (4):1593–1660. URL <http://dx.doi.org/10.1093/qje/qjr041>. (Cited on page [2.](#page-1-3))
- Cinelli, Carlos and Judea Pearl. 2021. "Generalizing Experimental Results by Leveraging Knowledge of Mechanisms." European Journal of Epidemiology 36:pp. 149–164. (Cited on pages [2,](#page-1-3) [4,](#page-3-1) [5,](#page-4-1) [6,](#page-5-3) and [7.](#page-6-4))
- DellaVigna, Stefano and Matthew Gentzkow. 2010. "Persuasion: Empirical Evidence." Annual Review of Economics 2 (1):pp. 643–669. (Cited on page [2.](#page-1-3))
- DellaVigna, Stefano and Ethan Kaplan. 2007. "The Fox News Effect: Media Bias and Voting." The Quarterly Journal of Economics 122 (3):pp. 1187–1234. (Cited on page [2.](#page-1-3))
- Dobbie, Will and Roland G. Fryer Jr. 2015. "The Medium-Term Impacts of High-Achieving Charter Schools." Journal of Political Economy 123 (5):pp. 985–1037. (Cited on page [2.](#page-1-3))
- Frangakis, Constantine and Donald Rubin. 2002. "Principal Stratification in Causal Inference." Biometrics 58 (1):pp. 21–29. (Cited on page [6.](#page-5-3))
- Heckman, James, Robert LaLonde, and Jeff Smith. 1999. "The Economics and Econometrics of Active Labor Market Programs." In Handbook of Labor Economics, vol. 3A, edited by Orley Ashenfelter and David Card. Elsevier, pp. 1865–2097. (Cited on page [2.](#page-1-3))
- Heckman, James J., Jeffrey Smith, and Nancy Clements. 1997. "Making the Most Out of Programme Evaluations and Social Experiments: Accounting for Heterogeneity in Programme Impacts." Review of Economic Studies 64:pp. 487–535. (Cited on pages [2,](#page-1-3) [4,](#page-3-1) and [7.](#page-6-4))
- Horowitz, Joel L. and Charles F. Manski. 1995. "Identification and Robustness with Contaminated and Corrupted Data." Econometrica 63 (2):pp. 281–302. (Cited on pages [4,](#page-3-1) [15,](#page-14-3) and [27.](#page-26-3))
- Huber, Martin, Lukas Laffers, and Giovanni Mellace. 2017. "Sharp IV Bounds on Average Treatment Effects on the Treated and Other Populations under Endogeneity and Noncompliance." Journal of Applied Econometrics 32:pp. 56–79. (Cited on page [9.](#page-8-4))
- Huber, Martin and Giovanni Mellace. 2015. "Sharp Bounds on Causal Effects under Sample Selection." Oxford Bulletin of Economics and Statistics 77 (1):pp. 129–151. (Cited on page [9.](#page-8-4))
- Imai, Kosuke. 2008. "Sharp Bounds on the Causal Effects in Randomized Experiments with Truncation- by- Death." Statistics and Probability Letters 78 (2):pp. 144–149. (Cited on pages [9](#page-8-4) and [27.](#page-26-3))
- Imbens, Guido and Charles Manski. 2004. "Confidence Intervals for Partially Identified Parameters." Econometrica 72 (6):pp. 1845–1857. (Cited on pages [19,](#page-18-1) [20,](#page-19-1) and [21.](#page-20-0))
- Imbens, Guido W. and Jeffrey M. Wooldridge. 2009. "Recent Developments in the Econometrics of Program Evaluation." Journal of Economic Literature 47 (1):pp. 5–86. (Cited on page [8.](#page-7-6))
- Jun, Sung Jae and Sokbae Lee. 2019. "Identifying the Effect of Persuasion." Available at <https://arxiv.org/abs/1812.02276>. (Cited on pages [2,](#page-1-3) [4,](#page-3-1) [5,](#page-4-1) [6,](#page-5-3) [7,](#page-6-4) [9,](#page-8-4) and [45.](#page-44-2))
	- ———. 2020. "Causal Inference in Case-Control Studies." Working Paper. (Cited on page [2.](#page-1-3))
- Krueger, Alan B. and Diane M. Whitmore. 2001. "The Effect of Attending a Small Class in the Early Grades on College-Test Taking and Middle School Test Results: Evidence from Project STAR." The Economic Journal 111 (468):pp. 1–28. (Cited on page [2.](#page-1-3))
- Lee, David S. 2009. "Training, Wages, and Sample Selection: Estimating Sharp Bounds on Treatment Effects." The Review of Economic Studies 76:pp. 1071–1102. (Cited on pages [2,](#page-1-3) [4,](#page-3-1) [5,](#page-4-1) [6,](#page-5-3) [7,](#page-6-4) [8,](#page-7-6) [10,](#page-9-4) [15,](#page-14-3) and [27.](#page-26-3))
- Manski, Charles F. 1997. "Monotone Treatment Response." Econometrica 65 (6):pp. 1311– 1334. (Cited on page [9.](#page-8-4))
- ———. 2011. "Policy Analysis with Incredible Certitude." The Economic Journal 121 (554):pp. F261–F289. (Cited on page [7.](#page-6-4))
- Manski, Charles F. and Francesca Molinari. 2020. "Estimating the COVID-19 infection rate: Anatomy of an Inference Problem." Journal of Econometrics 220 (1):181 – 192. Themed Issue: Pandemic Econometrics / Covid Pandemics. (Cited on page [7.](#page-6-4))
- Manski, Charles F. and John V. Pepper. 2000. "Monotone Instrumental Variables: With an Application to the Returns to Schooling." Econometrica 68 (4):pp. 997–1010. (Cited on page [9.](#page-8-4))
- Pearl, Judea. 1999. "Probabilities of Causation: Three Counterfactual Interpretations and their Identification." Synthese 121 (1-2):pp. 93–149. (Cited on pages [2,](#page-1-3) [4,](#page-3-1) [5,](#page-4-1) and [6.](#page-5-3))
- Sexton, Mary and Richard Hebel. 1984. "A Clinical Trial of Change in Maternal Smoking and its Effects on Birth Weight." Journal of the American Medical Association 251 (7):pp. 911–915. (Cited on page [2.](#page-1-3))
- Tamer, Elie. 2010. "Partial Identification in Econometrics." The Annual Review of Economics 2:pp. 167–195. (Cited on page [7.](#page-6-4))
- Tian, Jin and Judea Pearl. 2000. "Probabilities of Causation: Bounds and Identification." Annals of Mathematics and Artificial Intelligence 28:pp. 287–313. (Cited on pages [2,](#page-1-3) [4,](#page-3-1) [5,](#page-4-1) and [6.](#page-5-3))
- U.S. Department of Health and Human Services. 2004. The Health Consequences of Smoking: A Report of the Surgeon General. U.S. Department of Health and Human Services, Public Health Service, Office on Smoking and Health. Available at: https://www.cdc.gov/tobacco/data_statistics/sgr/2004/index.htm. (Cited on page [2.](#page-1-3))
- Zhang, Junni L., Donald B. Rubin, and Fabrizia Mealli. 2008. "Evaluating the Effects of Job Training Programs on Wages through Principal Stratification." In Modelling and Evaluating Treatment Effects in Econometrics. Emerald Group Publishing Limited, $117-145$. (Cited on page [6.](#page-5-3))

Supporting Information (Online Appendix)

A Proofs

A.1 Proof of Proposition [1](#page-9-2)

To prove Proposition [1,](#page-9-2) we must prove that Inequalities [\(3\)](#page-9-5) and [\(4\)](#page-9-6) hold. Since the validity of Inequality [\(3\)](#page-9-5) is a direct consequence of Lemma [1,](#page-9-1) we focus on proving Inequality [\(4\)](#page-9-6). Note that

$$
\mathbb{P}[Y = 1 | D = 1] - \mathbb{P}[Y = 1 | D = 0]
$$

\n
$$
= \mathbb{P}[Y_1^* \cdot S_1 = 1 | D = 1] - \mathbb{P}[Y_0^* \cdot S_0 = 1 | D = 0]
$$

\nby Equation (1)
\n
$$
= \mathbb{P}[Y_1^* \cdot S_1 = 1] - \mathbb{P}[Y_0^* \cdot S_0 = 1]
$$

\nby Assumption 1
\n
$$
= \mathbb{P}[Y_1^* = 1, S_1 = 1] - \mathbb{P}[Y_0^* = 1, S_0 = 1]
$$

\n
$$
\ge \mathbb{P}[Y_1^* = 1, S_0 = 1] - \mathbb{P}[Y_0^* = 1, S_0 = 1]
$$

\nby Assumption 3
\n
$$
\ge \mathbb{P}[Y_0^* = 1, S_0 = 1] - \mathbb{P}[Y_0^* = 1, S_0 = 1]
$$

\nby Assumption 4
\n
$$
= 0.
$$

A.2 Proof of Proposition [2](#page-10-0)

To prove Proposition [2,](#page-10-0) we first show that $LB_1 \n\t\leq \theta^{OO}$ and $\theta^{OO} \leq UB_1$. Then, we show that LB_1 and UB_1 are sharp bounds. For completeness, we state four lemmas that were previously derived in the literature and are used in our proofs. We prove then in Appendix [A.3.](#page-42-0)

Lemma A.1 Boole-Frechet Bounds [\(Imai,](#page-23-11) [2008\)](#page-23-11): We have that

$$
\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1
$$

\n
$$
\leq \mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]
$$

\n
$$
\leq \min \left\{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right], \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]\right\}.
$$

Lemma A.2 [Horowitz and Manski](#page-23-8) [\(1995,](#page-23-8) Corollary 1.2): Under Assumptions [1](#page-6-2) and [2,](#page-7-1) we have that

$$
\frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]\right)}{\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]} \le \mathbb{P}\left[Y_1^* = 1|S_0=1,S_1=1\right] \le \frac{\mathbb{P}\left[Y=1|S=1,D=1\right]}{\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]}.
$$

Lemma A.3 [Lee](#page-24-2) (2009) : Under Assumptions [1-](#page-6-2)[3,](#page-7-0) we have that

$$
\mathbb{P}\left[S_0 = 1, S_1 = 1 | S_1 = 1\right] = \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}.
$$

Lemma A.4 [Lee](#page-24-2) (2009) : Under Assumptions [1-](#page-6-2)[3,](#page-7-0) we have that

$$
\mathbb{P}[Y_0^* = 0 | S_0 = 1, S_1 = 1] = \mathbb{P}[Y = 0 | S = 1, D = 0].
$$

A.2.1 Lower Bound: $LB_1 \n\leq \theta^{OO}$

Note that

$$
\theta^{OO} := \mathbb{P}\left[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1\right]
$$

=
$$
\frac{\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

$$
\ge \frac{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

by Lemma [A.1](#page-9-1)

$$
\geq \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]\right)}{\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]} + \mathbb{P}\left[Y_0^*=0|S_0=1,S_1=1\right] - 1}{\mathbb{P}\left[Y_0^*=0|S_0=1,S_1=1\right]}
$$

by Lemma [A.2](#page-26-1)

$$
\frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)}{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]} + \mathbb{P}\left[Y_0^* = 0|S_0=1,S_1=1\right] - 1}
$$
\n
$$
= \frac{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{\mathbb{P}\left[Y_0^* = 0|S_0=1,S_1=1\right]}
$$

by Lemma [A.3](#page-26-2)

$$
\frac{\mathbb{P}[Y=1|S=1,D=1] - \left(1 - \frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]}\right)}{\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=0]}} + \mathbb{P}[Y=0|S=1,D=0] - 1
$$
\n
$$
= \frac{\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[Y=0|S=1,D=0]}}{\mathbb{P}[Y=0|S=1,D=0]}
$$

by Lemma [A.4.](#page-26-0)

Moreover, $\theta^{OO} \geq 0$ by definition.

A.2.2 Upper Bound: $\theta^{OO} \leq UB_1$

Note that

$$
\theta^{OO} := \mathbb{P}\left[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1\right]
$$

=
$$
\frac{\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

$$
\leq \frac{\min\left\{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right], \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]\right\}}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

by Lemma [A.1](#page-9-1) $=\min\left\{\frac{\mathbb{P}[Y_1^*=1|S_0=1,S_1=1]}{\mathbb{P}[Y_1^*=0|S_0=1,S_1=1]} \right\}$ $\frac{\mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1]}{\mathbb{P}[Y_0^* = 0 | S_0 = 1, S_1 = 1]}, 1$ \leq min $\sqrt{ }$ \int $\overline{\mathcal{L}}$ $\mathbb{P}[Y = 1 | S = 1, D = 1]$ $\overline{\mathbb{P}\left[S_0 = 1, S_1 = 1 | S_1 = 1 \right]}$ $\frac{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}, 1$ \mathcal{L} \downarrow \int

by Lemma [A.2](#page-26-1)

$$
= \min \left\{ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] \cdot \frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]}}{\mathbb{P}\left[Y_0^*=0|S_0=1,S_1=1\right]}, 1 \right\}
$$

by Lemma A.3

$$
= \min \left\{ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] \cdot \frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]}}{\mathbb{P}\left[Y=0|S=1,D=0\right]}, 1 \right\}
$$

by Lemma [A.4.](#page-26-0)

A.2.3 LB_1 and UB_1 are sharp bounds

To show that LB_1 and UB_1 are sharp bounds, we have to show that, for any $\tilde{\theta} \in$ $[LB_1, UB_1]$, there exist candidate random variables $(\tilde{Y}_0^*, \tilde{Y}_1^*, \tilde{S}_0, \tilde{S}_1, \tilde{D})$ that satisfy the fol-lowing conditions:^{[17](#page-28-0)}

- (A) The model restrictions hold, i.e., $(\tilde{Y}_0^*, \tilde{Y}_1^*, \tilde{S}_0, \tilde{S}_1, \tilde{D})$ satisfy Assumptions [1-](#page-6-2)[3.](#page-7-0)
- (B) The data restrictions hold, i.e., $\mathbb{P}\left[\tilde{Y} = 1\right]$ $\tilde{S} = 1, \tilde{D} = d$ = $\mathbb{P}[Y = 1 | S = 1, D = d]$, $\mathbb{P}\left[\tilde{S}=1 \middle| \tilde{D}=d\right] = \mathbb{P}\left[S=1 | D=d\right] \text{ for any } d \in \{0,1\} \text{ and } \mathbb{P}\left[\tilde{D}=1 \right] = \mathbb{P}\left[D=1 \right],$ \vert where $\tilde{Y}^* = \tilde{Y}_1^* \cdot \tilde{D} + \tilde{Y}_0^* \cdot (1 - \tilde{D}), \, \tilde{S} = \tilde{S}_1 \cdot \tilde{D} + \tilde{S}_0 \cdot (1 - \tilde{D})$ and $\tilde{Y} = \tilde{Y}^* \cdot \tilde{S}^{18}$ $\tilde{Y} = \tilde{Y}^* \cdot \tilde{S}^{18}$ $\tilde{Y} = \tilde{Y}^* \cdot \tilde{S}^{18}$.
- (C) $\tilde{\theta}$ is attained, i.e., $\mathbb{P}\left[\tilde{Y}_1^* = 1\right]$ $\tilde{Y}_0^* = 0, \tilde{S}_0 = 1, \tilde{S}_1 = 1 = \tilde{\theta}.$

To do so, we construct random variables $(\tilde{Y}_0^*, \tilde{Y}_1^*, \tilde{S}_0, \tilde{S}_1, \tilde{D})$ by:

¹⁸From the observable data, one can estimate:

- (a) The joint distribution of (S, D) , which is equivalent to estimating $P[S = 1 | D = d]$ for all $d \in \{0, 1\}$ and $P[D = 1]$ given that S and D are binary;
- (b) The joint distribution of $(Y, D)|S = 1$, which is equivalent to estimating $P[Y = 1 | S = 1, D = d]$ for all $d \in \{0,1\}$ and $P[D=1]$ because Y and D are binary.

Hence, the data restrictions guarantee that the proposed latent variables are indistinguishable from the real latent variables in the data.

¹⁷Intuitively, the definition of sharpness says that there exist candidate random variables $(\tilde{Y}_0^*, \tilde{Y}_1^*, \tilde{S}_0, \tilde{S}_1, \tilde{D})$ that attain the candidate target parameter $\tilde{\theta}$, satisfy the model restrictions and are indistinguishable from the true latent variables $(Y_0^*, Y_1^*, S_0, S_1, D)$ in the sense that they generate the same distribution of the observable data $(\tilde{Y}, \tilde{S}, \tilde{D})$ as the distribution of the data that is actually observed, i.e., (Y, S, D) .

- Part 1. imposing a joint distribution that satisfies Assumptions [1-](#page-6-2)[2](#page-7-1) and ensures that the marginal distribution of \tilde{D} is the same as the marginal distribution of D;
- Part 2. imposing a joint distribution of $(\tilde{S}_0, \tilde{S}_1)$ that satisfies Assumptions [2-](#page-7-1)[3](#page-7-0) and ensures that the conditional distribution of \tilde{S} ^{\vert} \tilde{D} is the same as the conditional distribution of $S|D;$
- Part 3. constructing a conditional distribution $(\tilde{Y}_0^*, \tilde{Y}_1^*)$ $(\tilde{S}_0^*, \tilde{S}_1^*)$ that is a probability distribution, satisfies data restrictions, and generates a probability of causation parameter $\hat{\theta}$ respectively equal to:
	- (3.a) the lower bound;
	- (3.b) the upper bound;
	- (3.c) any value in the interval (LB_1, UB_1) .

Part 1: The distribution of \tilde{D} and Assumptions [1-](#page-6-2)[2](#page-7-1)

Fix $(y_0, y_1, s_0, s_1, d) \in \{0, 1\}^5$ arbitrarily.

To ensure that Assumption [1](#page-6-2) holds, we impose that $\mathbb{P}\left[\tilde{Y}^*_0 = y_0, \tilde{Y}^*_1 = y_1, \tilde{S}_0 = s_0, \tilde{S}_1 = s_1, \tilde{D} = d\right] =$ $\mathbb{P}\left[\tilde{Y}^*_0 = y_0, \tilde{Y}^*_1 = y_1, \tilde{S}_0 = s_0, \tilde{S}_1 = s_1\right] \cdot \mathbb{P}\left[\tilde{D} = d\right].$

We set

$$
\mathbb{P}\left[\tilde{D}=1\right]=\mathbb{P}\left[D=1\right].\tag{A.1}
$$

Note that Assumption [2](#page-7-1) holds because $\mathbb{P}[D = 1] \in (0, 1)$ according to Assumption 2 for the true variable D.

We also impose that

$$
\mathbb{P}\left[\tilde{D}=0\right] = 1 - \mathbb{P}\left[\tilde{D}=1\right],\tag{A.2}
$$

so that \tilde{D} has a probability distribution.

Part [2](#page-7-1): The distribution of $(\tilde{S}_0, \tilde{S}_1)$ and Assumptions 2[-3](#page-7-0)

Since we have defined $\mathbb{P}\left[\tilde{D} = d\right]$ in Part 1, it remains to define

$$
\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1, \tilde{S}_0 = s_0, \tilde{S}_1 = s_1\right].
$$

Since $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1, \tilde{S}_0 = s_0, \tilde{S}_1 = s_1\right] = \mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = s_0, \tilde{S}_1 = s_1\Big] . \mathbb{P}\left[\tilde{S}_0 = s_0, \tilde{S}_1 = s_1\right],$ we define $\mathbb{P}\left[\tilde{S}_0 = s_0, \tilde{S}_1 = s_1\right]$ here and $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = s_0, \tilde{S}_1 = s_1$ in Part 3.

We set

$$
\mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = \mathbb{P}\left[S = 1 | D = 0\right],\tag{A.3}
$$

implying that Assumption [2](#page-7-1) holds because $\mathbb{P}\left[S=1|\ D=0\right]=\mathbb{P}\left[S_0=1\right]=\mathbb{P}\left[S_0=1,S_1=1\right] >$ 0 according to Assumption [1](#page-6-2)[-3](#page-7-0) for the true latent variables.

To ensure that Assumption [3](#page-7-0) holds, we set $\mathbb{P} \left[\tilde{S}_0 = 1, \tilde{S}_1 = 0 \right] = 0.$ To finish defining the distribution of $(\tilde{S}_0, \tilde{S}_1)$, let

$$
\mathbb{P}\left[\tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = \mathbb{P}\left[S = 1 | D = 1\right] - \mathbb{P}\left[S = 1 | D = 0\right]
$$
\n(A.4)

and

$$
\mathbb{P}\left[\tilde{S}_0 = 0, \tilde{S}_1 = 0\right] = 1 - \mathbb{P}\left[S = 1 | D = 1\right].\tag{A.5}
$$

To see that what we have indeed defined a probability distribution for $(\tilde{S}_0, \tilde{S}_1)$, note that

$$
\mathbb{P}\left[\tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = \mathbb{P}\left[S_1 = 1\right] - \mathbb{P}\left[S_0 = 1\right] = \mathbb{P}\left[S_0 = 0, S_1 = 1\right] \ge 0
$$

by Assumptions [1](#page-6-2) and [3](#page-7-0) for the true latent variables, and

$$
\mathbb{P}\left[\tilde{S}_0=0,\tilde{S}_1=0\right]+\mathbb{P}\left[\tilde{S}_0=1,\tilde{S}_1=0\right]+\mathbb{P}\left[\tilde{S}_0=0,\tilde{S}_1=1\right]+\mathbb{P}\left[\tilde{S}_0=1,\tilde{S}_1=1\right]=1
$$

by construction.

We conclude this part by showing that the distribution of $\tilde{S}|\tilde{D}$ is the same as that of $S|D$.

Note that

$$
\mathbb{P}\left[\tilde{S}=1 \middle| \tilde{D}=0\right] = \mathbb{P}\left[\tilde{S}_0=1\right] = \mathbb{P}\left[\tilde{S}_0=1, \tilde{S}_1=1\right] = \mathbb{P}\left[S=1 \middle| D=0\right]
$$

and that

$$
\mathbb{P}\left[\tilde{S}=1\middle| \tilde{D}=1\right] = \mathbb{P}\left[\tilde{S}_1=1\right] = \mathbb{P}\left[\tilde{S}_0=1, \tilde{S}_1=1\right] + \mathbb{P}\left[\tilde{S}_0=0, \tilde{S}_1=1\right]
$$

$$
= \mathbb{P}\left[S=1\middle| D=0\right] + \mathbb{P}\left[S=1\middle| D=1\right] - \mathbb{P}\left[S=1\middle| D=0\right]
$$

$$
= \mathbb{P}\left[S=1\middle| D=1\right].
$$

Part 3: The distribution of $(\tilde{\mathbf{Y}}^*_1, \tilde{\mathbf{Y}}^*_0)|(\tilde{\mathbf{S}}_1, \tilde{\mathbf{S}}_0)$

Since we have defined $\mathbb{P}\left[\tilde{D} = d\right]$ in Part 1 and $\mathbb{P}\left[\tilde{S}_0 = s_0, \tilde{S}_1 = s_1\right]$ in Part 2, it remains to define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = s_0, \tilde{S}_1 = s_1.$

We will define $(\tilde{Y}_1^*, \tilde{Y}_0^*) | (\tilde{S}_1, \tilde{S}_0)$ in three different ways so that $\tilde{\theta}$ attains each value in the identified interval $[LB_1, UB_1]$ and \tilde{Y} $\tilde{S} = 1, \tilde{D}$ has the same distribution as $Y | S = 1, D$.

(Part 3.a) Constructing a conditional distribution such that $\tilde{\theta} = LB_1$

Since $\mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 0\right] = 0$, we do not need to define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 0.$ We define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $[\tilde{S}_0 = 0, \tilde{S}_1 = 0] = \frac{1}{4}$ for any $(y_0, y_1) \in \{0, 1\}^2$. We also define the constant

$$
\blacklozenge = \max \left\{ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)}{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}, 0 \right\}
$$

,

and the conditional probabilities

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = \max\{\blacklozenge + \mathbb{P}\left[Y = 0 | S = 1, D = 0 \right] - 1, 0\} \tag{A.6}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = \min\{1 - \mathbb{P}\left[Y = 0 | S = 1, D = 0\right], \blacklozenge\}
$$
\n(A.7)

$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=0|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]
$$
\n
$$
=\mathbb{P}\left[Y=0|S=1,D=0\right]-\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right],
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=0|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]
$$
\n
$$
=\mathbb{P}\left[Y=1|S=1,D=0\right]-\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=1|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right],
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]
$$
\n
$$
=\frac{\mathbb{P}\left[Y=1,|S=1,D=1\right]-\mathbb{P}\left[\tilde{Y}_{1}^{*}=1|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]},
$$
\n
$$
1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=1|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]=0,
$$
\n(A.11)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right],\tag{A.12}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0. \tag{A.13}
$$

(Part 3.a.1) The candidate conditional distribution is a probability distribution

Now, we want to show that the functions described by equations $(A.6)-(A.13)$ $(A.6)-(A.13)$ $(A.6)-(A.13)$ are a probability mass function. First, note that:

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 1
$$

and

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 1.
$$

We must show that all values in $(A.6)-(A.13)$ $(A.6)-(A.13)$ $(A.6)-(A.13)$ are in the interval $[0, 1]$. Note that

$$
\blacklozenge \in [0,1]
$$

because $\blacklozenge \geq 0$ by construction, and, using Lemma [A.3,](#page-26-2) the expression in the definition of \blacklozenge becomes the expression on the left hand side of Lemma [A.2](#page-26-1) and, therefore, \blacklozenge \le $\mathbb{P}[Y_1^*|S_0 = 1, S_1 = 1] \leq 1.$

Furthermore, by construction, we have that:

$$
\max\{0, \blacklozenge -1 + \mathbb{P}[Y=0|S=1, D=0]\} = \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1|\tilde{S}_0=1, \tilde{S}_1=1\right] \le \blacklozenge \quad \text{(A.14)}
$$

$$
0 \le \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \le 1 - \mathbb{P}\left[Y = 0 | S = 1, D = 0\right] \le 1
$$
\n(A.15)

Given Equation [\(A.15\)](#page-33-0) and the fact that $1-\mathbb{P}[Y=0|S=1,D=0] = \mathbb{P}[Y=1|S=1,D=0]$, Equation [\(A.9\)](#page-32-1) implies that

$$
0 \le \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \le 1.
$$
\n(A.16)

Given Equations $(A.7)$ and $(A.14)$, Equation $(A.8)$ implies that

$$
1 - \Phi \le \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \le \mathbb{P}\left[Y = 0 | S = 1, D = 0\right] \le 1. \tag{A.17}
$$

In order to bound $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 0, \tilde{S}_1 = 1$, consider three cases:

Case 1) $\blacklozenge = 0$:

In this case, using Equations $(A.6)$ and $(A.7)$, we get that:

$$
\mathbb{P}\left[\tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$

= $\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]$
= 0

Also, by the definition of \blacklozenge , it is the case that:

$$
\mathbb{P}[Y=1|S=1,D=1] \le 1 - \frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]},
$$

implying, by Equation [\(A.10\)](#page-32-3), that

$$
0\leq\frac{\mathbb{P}\left[Y=1\right|S=1,D=1]}{1-\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]}}=\mathbb{P}\left[\left.\tilde{Y}^*_0=0,\tilde{Y}^*_1=1\right|\tilde{S}_0=0,\tilde{S}_1=1\right]\leq 1.
$$

Case 2) \blacklozenge > 1 – $\mathbb{P}\left[Y = 0 | S = 1, D = 0 \right]$.

In this case, Equations $(A.6)$ and $(A.7)$ imply that

$$
\mathbb{P}\left[\tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$

= $\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]$
= $1 - \mathbb{P}\left[Y = 0 | S = 1, D = 0\right] + \blacklozenge - (1 - \mathbb{P}\left[Y = 0 | S = 1, D = 0\right])$
= \blacklozenge .

Case 3) $\blacklozenge \in (0, 1 - \mathbb{P}[Y = 0 | S = 1, D = 0]].$

In this case, we have that $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $[\tilde{S}_0 = 1, \tilde{S}_1 = 1] = 0$ by Equation [\(A.6\)](#page-31-0) and $\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1\right]$ $[\tilde{S}_0 = 1, \tilde{S}_1 = 1] = \bigotimes$ by Equation [\(A.7\)](#page-31-1), implying that

$$
\mathbb{P}\left[\tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = \blacklozenge.
$$

In Cases 2 and 3, we can use Equation $(A.10)$ to see that

$$
\mathbb{P}\left[\tilde{Y}^*_0=0,\tilde{Y}^*_1=1|\tilde{S}_0=0,\tilde{S}_1=1\right]=\frac{\mathbb{P}\left[Y=1|S=1,D=1\right]-\blacklozenge\cdot\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]}}{1-\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]}} ,
$$

implying, by the definition of $\blacklozenge,$ that

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$
\n(A.18)\n
$$
= \frac{\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] - \left(\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] - \left(1 - \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}\right)\right)}{1 - \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}}
$$
\n(A.18)

Since $\mathbb{P} \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right] \in [0, 1]$, Equation [\(A.12\)](#page-32-4) ensures that

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 | \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] \in [0, 1].
$$

(Part 3.a.2) The candidate conditional distribution satisfies its data restrictions The data restrictions for \tilde{Y} $\tilde{S} = 1, \tilde{D}$ are satisfied because:

•
$$
\mathbb{P}\left[\tilde{Y} = 1 | \tilde{S} = 1, \tilde{D} = 0\right] = \mathbb{P}\left[Y = 1 | S = 1, D = 0\right];
$$

To see that, use Equations [\(A.7\)](#page-31-1) and [\(A.9\)](#page-32-1) and the fact that $\mathbb{P} \left[\tilde{S}_0 = 1, \tilde{S}_1 = 0 \right] = 0$ to write:

$$
\begin{aligned} \mathbb{P}\left[\tilde{Y}=1 \middle| \tilde{S}=1, \tilde{D}=0\right] \\ &= \mathbb{P}\left[\tilde{Y}_0^* = 1 \middle| \tilde{S}_0=1\right] \\ &= \mathbb{P}\left[\tilde{Y}_0^* = 1 \middle| \tilde{S}_0=1, \tilde{S}_1=1\right] \\ &= \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0=1, \tilde{S}_1=1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0=1, \tilde{S}_1=1\right] \\ &= \mathbb{P}\left[Y=1 \middle| S=1, D=0\right]. \end{aligned}
$$

•
$$
\mathbb{P}\left[\tilde{Y} \middle| \tilde{S} = 1, \tilde{D} = 1\right] = \mathbb{P}\left[Y | S = 1, D = 1\right].
$$

To see that, note that we can write:

$$
\mathbb{P}\left[\tilde{Y} = 1 | \tilde{S} = 1, \tilde{D} = 1\right]
$$

= $\mathbb{P}\left[\tilde{Y}_1^* = 1 | \tilde{S}_1 = 1\right]$
= $\mathbb{P}\left[\tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \cdot \mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 1 | \tilde{S}_1 = 1\right]$
+ $\mathbb{P}\left[\tilde{Y}_1^* = 1 | \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] \cdot \left(1 - \mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 1 | \tilde{S}_1 = 1\right]\right)$

Now, note that we can sum Equations $(A.10)$ and $(A.11)$ and find that

$$
\mathbb{P}\left[\tilde{Y}_1^* = 1 | \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = \frac{\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] - \mathbb{P}\left[\tilde{Y}_1^* | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \cdot \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}}{1 - \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}}\tag{A.19}
$$

Using Equations $(A.4)$ and $(A.3)$ from Part 1, we get:

$$
\mathbb{P}\left[\tilde{S}_1 = 1, \tilde{S}_0 = 1 | \tilde{S}_1 = 1\right] = \frac{\mathbb{P}\left[\tilde{S}_1 = 1, \tilde{S}_0 = 1\right]}{\mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{S}_0 = 0, \tilde{S}_1 = 1\right]} = \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}
$$
\n(A.20)

Plugging $(A.19)$ and $(A.20)$ in the expression above, we get:

$$
\begin{aligned}\n\mathbb{P}\left[\tilde{Y}=1\Big|\tilde{S}=1,\tilde{D}=1\right] \\
&= \mathbb{P}\left[\tilde{Y}_{1}^{*}=1|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right] \cdot \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]} \\
&+ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \mathbb{P}\left[\tilde{Y}_{1}^{*}|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right] \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{\mathbb{1} - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}} \cdot \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right) \\
&= \mathbb{P}\left[Y=1|S=1,D=1\right]\n\end{aligned}
$$

(Part 3.a.3) The probability of causation $\tilde{\theta}$ reaches the lower bound LB_1

Finally, note that the lower bound ${\it LB}_1$ is attained because

$$
\begin{split} &\mathbb{P}\left[\left.\tilde{Y}_{1}^{*}=1\right|\tilde{Y}_{0}^{*}=0,\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\\ &=\frac{\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}\\ &=\frac{\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=0\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}\\ &=\frac{\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\left.Y=0\right|S=1,D=0]-\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}\\ &\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\left.Y=0\right|S=1,D=0]-\mathbb{P}\left[\left.\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\right|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\end{split}
$$

$$
\max \left\{ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)}{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}} + \mathbb{P}\left[Y=0|S=1,D=0\right] - 1, 0 \right\}
$$
\n
$$
= \frac{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{\mathbb{P}\left[Y=0|S=1,D=0\right]}
$$

 $=$ $LB₁$.

(Part 3.b) Constructing a conditional distribution such that $\tilde{\theta} = \text{UB}_1$

Since $\mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 0\right] = 0$, we do not need to define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 0.$ We define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = 0, \tilde{S}_1 = 0$ = 1/4 for any $(y_0, y_1) \in \{0, 1\}^2$. We also define:

$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]
$$
\n
$$
=\min\left\{\mathbb{P}\left[Y=1|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]},\mathbb{P}\left[Y=0|S=1,D=0\right]\right\},
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]
$$
\n
$$
=\max\left\{\min\left\{\mathbb{P}\left[Y=1|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]},1\right\}-\mathbb{P}\left[Y=0|S=1,D=0\right],0\right\}
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]
$$
\n(A.23)

,

$$
= \mathbb{P}\left[Y = 0| S = 1, D = 0\right] - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right],
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$

$$
- \mathbb{P}\left[V - 1 | S = 1, D = 0\right] - \mathbb{P}\left[\tilde{Y}^* = 1 | \tilde{Y}_0 = 1 | \tilde{S}_0 = 1 | \tilde{S}_1 = 1\right]
$$

(A.24)

$$
= \mathbb{P}\left[Y = 1| S = 1, D = 0\right] - \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right],
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$

$$
= \max\left\{\frac{\mathbb{P}\left[Y = 1| S = 1, D = 1\right] - \frac{\mathbb{P}\left[S = 1| D = 0\right]}{\mathbb{P}\left[S = 1| D = 1\right]}{0}, 0\right\},
$$

(A.25)

$$
= \max \left\{ \frac{\mathbb{P}\left[S=1|D=0\right]}{1 - \frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=1\right]}}, 0 \right\},
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right] = 0,
$$
\n(A.26)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right],\tag{A.27}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0. \tag{A.28}
$$

Observe that

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$

\n
$$
\ge \mathbb{P}\left[Y = 0 | S = 1, D = 0\right] - \mathbb{P}\left[Y = 0 | S = 1, D = 0\right]
$$

\n
$$
\ge 0,
$$

and

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$

\n
$$
\ge \mathbb{P}\left[Y = 1 | S = 1, D = 0 \middle| -1 + \mathbb{P}\left[Y = 0 | S = 1, D = 0\right]\right]
$$

\n= 0

and

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 1.
$$

Moreover, note that $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 0, \tilde{S}_1 = 1 \vert \in [0, 1)$ by construction.

Notice also that the data restrictions are satisfied because

$$
\mathbb{P}\left[\tilde{Y} = 1 | \tilde{S} = 1, \tilde{D} = 0\right]
$$

= $\mathbb{P}\left[\tilde{Y}_0^* = 1 | \tilde{S}_0 = 1\right]$
= $\mathbb{P}\left[\tilde{Y}_0^* = 1 | \tilde{S}_0 = 1, S_1 = 1\right]$
= $\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, S_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 | \tilde{S}_0 = 1, S_1 = 1\right]$
= $\mathbb{P}\left[Y = 1 | S = 1, D = 0\right]$

and

$$
\mathbb{P}\left[\tilde{Y}=1\middle|\tilde{S}=1,\tilde{D}=1\right]
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{1}=1\right]
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\cdot\mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\middle|\tilde{S}_{1}=1\right]
$$
\n
$$
+ \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]\cdot\left(1-\mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\middle|\tilde{S}_{1}=1\right]\right),
$$
\n
$$
= \left(\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\right)
$$
\n
$$
\cdot \mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\middle|\tilde{S}_{1}=1\right]
$$
\n
$$
+ \left(\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]\right)
$$
\n
$$
\cdot \left(1-\mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\middle|\tilde{S}_{1}=1\right]\right)
$$
\n
$$
= \min\left\{\mathbb{P}\left[Y=1|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]},1\right\}\cdot\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}
$$
\n
$$
+ \max\left\{\mathbb{P}\left[Y=1|S=1,D=1\
$$

Finally, note that

$$
\begin{split} &\mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{Y}_{0}^{*}=0,\tilde{S}_{0}=1,\tilde{S}_{1}=1\right] \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \end{split}
$$

$$
= \frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[Y=0|S=1,D=0\right]-\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}\\=\frac{\min\left\{\mathbb{P}\left[Y=1|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]},\mathbb{P}\left[Y=0|S=1,D=0\right]\right\}}{\mathbb{P}\left[Y=0|S=1,D=0\right]}
$$

 $= UB_1.$

(Part 3.c) Constructing a conditional distribution that attains any $\tilde{\theta}\in (\mathbf{LB_1}, \mathbf{UB_1})$ Since $\tilde{\theta} \in (LB_1, UB_1)$, there exists $\omega \in (0, 1)$ such that $\tilde{\theta} = \omega \cdot LB_1 + (1 - \omega)UB_1$. Since $\mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 0\right] = 0$, we do not need to define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 0.$ We define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $[\tilde{S}_0 = 0, \tilde{S}_1 = 0] = \frac{1}{4}$ for any $(y_0, y_1) \in \{0, 1\}^2$. We also define

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$
\n
$$
= \omega \cdot \mathbb{P}_L\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + (1 - \omega) \cdot \mathbb{P}_U\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right],
$$
\n(A.29)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$
\n(A.30)\n
$$
\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}
$$
\n(A.31)

$$
= \omega \cdot \mathbb{P}_L \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right] + (1 - \omega) \cdot \mathbb{P}_U \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right],
$$

$$
\mathbb{P} \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right]
$$
(A.31)

$$
= \omega \cdot \mathbb{P}_L \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right] + (1 - \omega) \cdot \mathbb{P}_U \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right],
$$

$$
\mathbb{P} \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right]
$$
(A.32)

$$
= \omega \cdot \mathbb{P}_L \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right] + (1 - \omega) \cdot \mathbb{P}_U \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right],
$$

$$
\mathbb{P} \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right]
$$
(A.33)

$$
= \omega \cdot \mathbb{P}_L \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right] + (1 - \omega) \cdot \mathbb{P}_U \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right],
$$

$$
\mathbb{P} \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right]
$$
(A.34)

$$
= \omega \cdot \mathbb{P}_L \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right] + (1 - \omega) \cdot \mathbb{P}_U \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right],
$$

$$
\mathbb{P} \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right]
$$
(A.35)

$$
\begin{aligned} \left[\begin{array}{c} 1 \end{array} \right] \left[\begin{array} \end{array} \right
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$
\n(A.36)

$$
= \omega \cdot \mathbb{P}_L \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right] + (1 - \omega) \cdot \mathbb{P}_U \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right].
$$
\n(A.37)

where the sub-index L denotes the conditional probabilities defined for the lower bound (Equations $(A.6)-(A.13)$ $(A.6)-(A.13)$ $(A.6)-(A.13)$ and the sub-index U denotes the conditional probabilities defined for the upper bound (Equations $(A.21)-(A.28)$ $(A.21)-(A.28)$ $(A.21)-(A.28)$).

Notice that the data restrictions are satisfied because

$$
\mathbb{P}\left[\tilde{Y} = 1 | \tilde{S} = 1, \tilde{D} = d\right]
$$

= $\omega \cdot \mathbb{P}_L \left[\tilde{Y} = 1 | \tilde{S} = 1, \tilde{D} = d\right] + (1 - \omega) \cdot \mathbb{P}_U \left[\tilde{Y} = 1 | \tilde{S} = 1, \tilde{D} = d\right]$
= $\mathbb{P}\left[Y = 1 | S = 1, D = d\right].$

Finally, note that

$$
\begin{split} \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{Y}_{0}^{*}=0,\tilde{S}_{0}=1,\tilde{S}_{1}=1\right] \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &=\frac{\omega\cdot\mathbb{P}_{L}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+(1-\omega)\cdot\mathbb{P}_{U}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[Y=0\right|S=1,D=0]} \\ &=\omega\cdot LB_{1}+(1-\omega)\cdot UB_{1} \\ &=\tilde{\theta}. \end{split}
$$

A.3 Proofs of Lemmas [A.1](#page-9-1)[-A.4](#page-26-0)

A.3.1 Lemma [A.1](#page-9-1)

For the upper bound, note that

$$
\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]
$$

\n
$$
\leq \mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_1^* = 1, Y_0^* = 1 | S_0 = 1, S_1 = 1\right]
$$

\n
$$
= \mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right]
$$

and

$$
\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]
$$

\n
$$
\leq \mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_1^* = 0, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]
$$

\n
$$
= \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right].
$$

For the lower bound, observe that

$$
\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]
$$

= $\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - \mathbb{P}\left[Y_1^* = 1 \text{ or } Y_0^* = 0 | S_0 = 1, S_1 = 1\right]$
 $\ge \mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1.$

A.3.2 Lemma [A.2](#page-26-1)

Note that

$$
\mathbb{P}[Y=1|S=1, D=1] = \mathbb{P}[Y_1^* = 1|S_1 = 1, D=1]
$$

=
$$
\frac{\mathbb{P}[Y_1^* = 1, S_1 = 1|D=1]}{\mathbb{P}[S_1 = 1|D=1]}
$$

=
$$
\frac{\mathbb{P}[Y_1^* = 1, S_1 = 1]}{\mathbb{P}[S_1 = 1]}
$$
 by Assumption 1
=
$$
\mathbb{P}[Y_1^* = 1|S_1 = 1]
$$

$$
= \mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

$$
+ \mathbb{P}[Y_1^* = 1 | S_0 = 0, S_1 = 1] \cdot (1 - \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]),
$$

implying that

$$
\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right]
$$

=
$$
\frac{\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] - \mathbb{P}\left[Y_1^* = 1 | S_0 = 0, S_1 = 1\right] \cdot (1 - \mathbb{P}\left[S_0 = 1, S_1 = 1 | S_1 = 1\right])}{\mathbb{P}\left[S_0 = 1, S_1 = 1 | S_1 = 1\right]}.
$$

Since $\mathbb{P}[Y_1^* = 1 | S_0 = 0, S_1 = 1] \in [0, 1]$, we can conclude that the bounds above hold.

A.3.3 Lemma [A.3](#page-26-2)

Note that

$$
\mathbb{P}\left[S_0 = 1, S_1 = 1 | S_1 = 1\right] = \frac{\mathbb{P}\left[S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[S_1 = 1\right]}
$$

$$
= \frac{\mathbb{P}\left[S_0 = 1\right]}{\mathbb{P}\left[S_1 = 1\right]} \text{ by Assumption 3}
$$

$$
= \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]} \text{ by Assumption 1.}
$$

A.3.4 Lemma [A.4](#page-26-0)

Note that

$$
\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] = \frac{\mathbb{P}\left[Y_0^* = 0, S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[S_0 = 1, S_1 = 1\right]}
$$

\n
$$
= \frac{\mathbb{P}\left[Y_0^* = 0, S_0 = 1\right]}{\mathbb{P}\left[S_0 = 1\right]} \text{ by Assumption 3}
$$

\n
$$
= \frac{\mathbb{P}\left[Y = 0, S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 0\right]} \text{ by Assumption 1}
$$

\n
$$
= \mathbb{P}\left[Y = 0 | S = 1, D = 0\right].
$$

A.4 Proof of Proposition [3](#page-10-1)

To prove Proposition [3,](#page-10-1) we first show that $LB_2 \leq \theta^{OO}$ and $\theta^{OO} \leq UB_2$. Then, we show that LB_2 and UB_2 are sharp bounds. For completeness, we state one lemma that was previously derived in the literature and is used in our proofs. We prove then in Appendix [A.5.](#page-54-0)

Lemma A.5 [Jun and Lee](#page-23-0) [\(2019\)](#page-23-0): Under Assumption [4,](#page-7-3) we have that

 $\mathbb{P}[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1] = \mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] + \mathbb{P}[Y_0^* = 0 | S_0 = 1, S_1 = 1] - 1.$

A.4.1 Lower Bound: $LB_2 \leq \theta^{OO}$

Note that

$$
\theta^{OO} := \mathbb{P}\left[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1\right]
$$

=
$$
\frac{\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

=
$$
\frac{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

by Lemma [A.5](#page-44-1)

$$
\geq \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]\right)}{\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]} + \mathbb{P}\left[Y_0^* = 0|S_0=1,S_1=1\right] - 1}{\mathbb{P}\left[Y_0^* = 0|S_0=1,S_1=1\right]}
$$

by Lemma [A.2](#page-26-1)

$$
\frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)}{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}} + \mathbb{P}\left[Y_0^* = 0|S_0=1, S_1=1\right] - 1
$$
\n
$$
= \frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[Y_0^* = 0|S_0=1, S_1=1\right]}
$$

by Lemma [A.3](#page-26-2)

$$
\frac{\mathbb{P}[Y=1|S=1,D=1] - \left(1 - \frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]}\right)}{\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]}} + \mathbb{P}[Y=0|S=1,D=0] - 1
$$
\n
$$
= \frac{\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]}}{\mathbb{P}[Y=0|S=1,D=0]}
$$

by Lemma [A.4.](#page-26-0)

Moreover, $\theta^{OO} \geq 0$ by definition.

A.4.2 Upper Bound: $\theta^{OO} \leq UB_2$

Note that

$$
\theta^{OO} := \mathbb{P}\left[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1\right]
$$

=
$$
\frac{\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

=
$$
\frac{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

by Lemma [A.5](#page-44-1)

$$
\leq \frac{\mathbb{P}\left[Y=1|S=1,D=1\right]}{\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]} + \mathbb{P}\left[Y_0^*=0|S_0=1,S_1=1\right] - 1
$$

$$
\mathbb{P}\left[Y_0^*=0|S_0=1,S_1=1\right]
$$

by Lemma [A.2](#page-26-1)

$$
= \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] \cdot \frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]} + \mathbb{P}\left[Y_0^* = 0|S_0=1,S_1=1\right] - 1}{\mathbb{P}\left[Y_0^* = 0|S_0=1,S_1=1\right]}
$$

by Lemma [A.3](#page-26-2)

$$
= \frac{\mathbb{P}[Y=1|S=1,D=1] \cdot \frac{\mathbb{P}[S=1|D=1]}{\mathbb{P}[S=1|D=0]} + \mathbb{P}[Y=0|S=1,D=0] - 1}{\mathbb{P}[Y=0|S=1,D=0]}
$$

by Lemma [A.4.](#page-26-0)

Moreover, $\theta^{OO} \leq 1$ by definition.

A.4.3 LB_2 and UB_2 are sharp bounds

The only difference between this proof and the proof in Appendix [A.2](#page-25-1) is the definition of $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = s_0, \tilde{S}_1 = s_1$ for any $(y_0, y_1, s_0, s_1) \in \{0, 1\}^4$. For this reason, we will only construct a conditional distribution $(\tilde{Y}_0^*, \tilde{Y}_1^*)$ $(\tilde{S}_0^*, \tilde{S}_1^*)$ that is a probability distribution,

satisfies Assumption [4,](#page-7-3) satisfies the data restrictions, and generates a probability of causation $\tilde{\theta}$ respectively equal to:

- (a) the lower bound LB_2 ;
- (b) the upper bound UB_2 ;
- (c) any value in the interval (LB_2, UB_2) .

(Part a) Constructing a conditional distribution such that $\tilde{\theta} = \mathbf{LB_2}$ Since $\mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 0\right] = 0$, we do not need to define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 0.$ We define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $[\tilde{S}_0 = 0, \tilde{S}_1 = 0] = \frac{1}{4}$ for any $(y_0, y_1) \in \{0, 1\}^2$. We also define the constant

$$
\blacklozenge = \max \left\{ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)}{\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}, 1 - \mathbb{P}\left[Y=0|S=1,D=0\right] \right\},\right.
$$

and the conditional probabilities

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = \bigoplus \mathbb{P}\left[Y = 0 | S = 1, D = 0\right] - 1\tag{A.38}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[Y = 0 | S = 1, D = 0\right]
$$
\n(A.39)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$
\n(A.40)

$$
= \mathbb{P}\left[Y = 0| S = 1, D = 0\right] - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right],
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 0
$$
 (A.41)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$
\n(A.42)

$$
\mathbb{P}\left[Y=1, | S=1, D=1\right] - \mathbb{P}\left[\tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \cdot \frac{\mathbb{P}\left[S=1 | D=0\right]}{\mathbb{P}\left[S=1 | D=1\right]} \\
1 - \frac{\mathbb{P}\left[S=1 | D=0\right]}{\mathbb{P}\left[S=1 | D=1\right]} \\
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0,\n\tag{A.43}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right],\tag{A.44}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0. \tag{A.45}
$$

Note that Equations $(A.41)$ and $(A.45)$ ensure that Assumption [4](#page-7-3) holds.

(Part a.1) The candidate conditional distribution is a probability distribution

Now, we want to show that the functions described by equations $(A.38)-(A.45)$ $(A.38)-(A.45)$ $(A.38)-(A.45)$ are a probability mass function. First, note that:

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 1
$$

and

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 1.
$$

We must show that all values in $(A.38)-(A.45)$ $(A.38)-(A.45)$ $(A.38)-(A.45)$ are in the interval $[0, 1]$.

Note that $\blacklozenge \in [0,1]$ for the same reasons explained in Appendix [A.2,](#page-25-1) implying that $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 1$ $\in [0, 1]$. Moreover, observe that Equation [\(A.40\)](#page-46-2) implies that

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 1 - \blacklozenge \ge 0.
$$

In order to bound, $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 0, \tilde{S}_1 = 1$, note that Equations [\(A.38\)](#page-46-1) and [\(A.39\)](#page-46-3) imply that $\mathbb{P}\left[\tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 1$ = \bullet . Consequently, Equation [\(A.42\)](#page-46-4) imply that

$$
\mathbb{P}\left[\tilde{Y}^*_0=0, \tilde{Y}^*_1=1 | \tilde{S}_0=0, \tilde{S}_1=1 \right] = \frac{\mathbb{P}\left[Y=1 | S=1, D=1 \right] - \blacklozenge \cdot \frac{\mathbb{P}\left[S=1 | D=0 \right]}{\mathbb{P}\left[S=1 | D=1 \right]}}{1 - \frac{\mathbb{P}\left[S=1 | D=0 \right]}{\mathbb{P}\left[S=1 | D=1 \right]}}
$$

.

Now, consider two cases:

Case 1) \blacklozenge > 1 – $\mathbb{P}\left[Y = 0 | S = 1, D = 0 \right]$.

In this case, we have that

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$
\n(A.46)\n
$$
= \frac{\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] - \left(\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] - \left(1 - \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}\right)\right)}{1 - \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}}
$$
\n(A.46)

Case 2) $\blacklozenge = 1 - \mathbb{P}[Y = 0 | S = 1, D = 0].$

In this case, we have that

$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]
$$
\n
$$
=\frac{\mathbb{P}\left[Y=1|S=1,D=1\right]-(1-\mathbb{P}\left[Y=0|S=1,D=0\right])\cdot\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}\\=\frac{\mathbb{P}\left[Y_{1}^{*}=1|S_{1}=1\right]-\mathbb{P}\left[Y_{0}^{*}=1|S_{0}=1,S_{1}=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}
$$
\n(A.47)

by Lemma [A.4](#page-26-0)

$$
\alpha \mathbb{P}[Y_1^* = 1 | S_1 = 1] - \mathbb{P}[Y_0^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

by Lemma A.3

$$
= \mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

$$
+ \mathbb{P}[Y_1^* = 1 | S_0 = 0, S_1 = 1] \cdot \mathbb{P}[S_0 = 0, S_1 = 1 | S_1 = 1]
$$

$$
- \mathbb{P}[Y_0^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

$$
= \mathbb{P}[Y_0^* = 0, Y_1^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

$$
+ \mathbb{P}[Y_0^* = 1, Y_1^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

$$
+ \mathbb{P}[Y_0^* = 0, Y_1^* = 1 | S_0 = 0, S_1 = 1] \cdot \mathbb{P}[S_0 = 0, S_1 = 1 | S_1 = 1]
$$

$$
+ \mathbb{P}[Y_0^* = 0, Y_1^* = 1 | S_0 = 0, S_1 = 1] \cdot \mathbb{P}[S_0 = 0, S_1 = 1 | S_1 = 1]
$$

$$
+ \mathbb{P}[Y_0^* = 1, Y_1^* = 1 | S_0 = 0, S_1 = 1] \cdot \mathbb{P}[S_0 = 0, S_1 = 1 | S_1 = 1]
$$

$$
-\mathbb{P}[Y_0^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

\n
$$
=\mathbb{P}[Y_0^* = 0, Y_1^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

\n
$$
+\mathbb{P}[Y_0^* = 1, Y_1^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

\n
$$
+\mathbb{P}[Y_0^* = 0, Y_1^* = 1 | S_0 = 0, S_1 = 1] \cdot \mathbb{P}[S_0 = 0, S_1 = 1 | S_1 = 1]
$$

\n
$$
+\mathbb{P}[Y_0^* = 1, Y_1^* = 1 | S_0 = 0, S_1 = 1] \cdot \mathbb{P}[S_0 = 0, S_1 = 1 | S_1 = 1]
$$

\n
$$
-\mathbb{P}[Y_0^* = 1, Y_1^* = 1 | S_0 = 1, S_1 = 1] \cdot \mathbb{P}[S_0 = 1, S_1 = 1 | S_1 = 1]
$$

by Assumption [4](#page-7-3) for the true latent variables

$$
= \mathbb{P}\left[Y_0^* = 0, Y_1^* = 1 | S_0 = 1, S_1 = 1\right] \cdot \mathbb{P}\left[S_0 = 1, S_1 = 1 | S_1 = 1\right]
$$

+ $\mathbb{P}\left[Y_0^* = 0, Y_1^* = 1 | S_0 = 0, S_1 = 1\right] \cdot \mathbb{P}\left[S_0 = 0, S_1 = 1 | S_1 = 1\right]$
+ $\mathbb{P}\left[Y_0^* = 1, Y_1^* = 1 | S_0 = 0, S_1 = 1\right] \cdot \mathbb{P}\left[S_0 = 0, S_1 = 1 | S_1 = 1\right]$
 ≥ 0 (A.48)

by the definition of a probability.

Moreover, we have that

$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]
$$
\n
$$
=\frac{\mathbb{P}\left[Y=1|S=1,D=1\right]-(1-\mathbb{P}\left[Y=0|S=1,D=0\right])\cdot\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}\n\leq \frac{\mathbb{P}\left[Y=1|S=1,D=1\right]-\left(\mathbb{P}\left[Y=1|S=1,D=1\right]-\left(1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)\right)}{1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}
$$
\n(A.49)

by the definition of \blacklozenge

$$
= 1.
$$

Since $\mathbb{P} \left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 0, \tilde{S}_1 = 1 \right] \in [0, 1]$, Equation [\(A.44\)](#page-47-1) ensures that $\mathbb{P}\left[\tilde{Y}^*_0 = 0, \tilde{Y}^*_1 = 0 | \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] \in [0, 1].$

(Part 3.a.2) The candidate conditional distribution satisfies its data restrictions The data restrictions for \tilde{Y} ^{\vert} $\tilde{S} = 1, \tilde{D}$ are satisfied because:

• $\mathbb{P}\left[\tilde{Y}=1\right]$ $\tilde{S} = 1, \tilde{D} = 0$ = $\mathbb{P}[Y = 1 | S = 1, D = 0];$

To see that, use Equations [\(A.39\)](#page-46-3) and [\(A.41\)](#page-46-0) and the fact that $\mathbb{P} \left[\tilde{S}_0 = 1, \tilde{S}_1 = 0 \right] = 0$ to write:

$$
\mathbb{P}\left[\tilde{Y} = 1 \middle| \tilde{S} = 1, \tilde{D} = 0\right]
$$

= $\mathbb{P}\left[\tilde{Y}_0^* = 1 \middle| \tilde{S}_0 = 1\right]$
= $\mathbb{P}\left[\tilde{Y}_0^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]$
= $\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]$
= $\mathbb{P}\left[Y = 1 | S = 1, D = 0\right].$

•
$$
\mathbb{P}\left[\tilde{Y} = 1 | \tilde{S} = 1, \tilde{D} = 1\right] = \mathbb{P}\left[Y = 1 | S = 1, D = 1\right].
$$

To see that, note that we can write:

$$
\mathbb{P}\left[\tilde{Y}=1\Big|\tilde{S}=1,\tilde{D}=1\right]
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{1}=1\right]
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\cdot\mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\Big|\tilde{S}_{1}=1\right]
$$
\n
$$
+ \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]\cdot\left(1-\mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\Big|\tilde{S}_{1}=1\right]\right)
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{1}^{*}=1|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}
$$
\n
$$
+ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right]-\mathbb{P}\left[\tilde{Y}_{1}^{*}|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}} \cdot \left(1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right)
$$
\n
$$
= \mathbb{P}\left[Y=1|S=1,D=1\right].
$$

(Part a.3) The probability of causation $\tilde{\theta}$ reaches the lower bound LB_1

Finally, note that the lower bound $LB₁$ is attained because

$$
\begin{split} &\mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{Y}_{0}^{*}=0,\tilde{S}_{0}=1,\tilde{S}_{1}=1\right] \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\Big]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[Y=0|S=1,D=0\right]-\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[Y=0|S=1,D=0\right]-\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &\max\left\{\frac{\mathbb{P}\left[Y=1|S=1,D=1\right]-\left(1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]},1-\mathbb{P}\left[Y=0|S=1,D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}\right.\end{split}
$$

$$
+\mathbb{P}[Y=0|S=1,D=0]-1
$$
\n
$$
=\frac{\mathbb{P}[Y=1|S=1,D=1]-\left(1-\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[S=1|D=1]}\right)}{\max\left\{\frac{\mathbb{P}[Y=1|S=1|D=0]}{\mathbb{P}[S=1|D=0]}\right\}}+\mathbb{P}[Y=0|S=1,D=0]-1,0
$$
\n
$$
=\frac{\mathbb{P}[S=1|D=0]}{\mathbb{P}[Y=0|S=1,D=0]}
$$

 $=$ $LB₂$.

(Part b) Constructing a conditional distribution such that $\tilde{\theta} = \text{UB}_2$

Since $\mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 0\right] = 0$, we do not need to define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 0.$ We define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = 0, \tilde{S}_1 = 0$ = 1/4 for any $(y_0, y_1) \in \{0, 1\}^2$. We also define:

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$
\n
$$
= \min\left\{\mathbb{P}\left[Y = 1 \middle| S = 1, D = 1\right] \cdot \frac{\mathbb{P}\left[S = 1 \middle| D = 1\right]}{\mathbb{P}\left[S = 1 \middle| D = 0\right]}, 1\right\} + \mathbb{P}\left[Y = 0 \middle| S = 1, D = 0\right] - 1,
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[Y = 0 \middle| S = 1, D = 0\right]
$$
\n(A.51)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$
\n
$$
= \mathbb{P}\left[Y = 0 \middle| S = 1, D = 0\right] - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right],
$$
\n(A.52)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 0
$$
\n(A.53)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$
\n(A.54)

$$
= \max \left\{ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]} }{1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}, 0 \right\},
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0,
$$
\n(A.55)\n
$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0.
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right],\tag{A.56}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0. \tag{A.57}
$$

Note that Equations [\(A.53\)](#page-52-0) and [\(A.57\)](#page-52-1) ensure that Assumption [4](#page-7-3) hold. Moreover, observe that

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$

\n
$$
= \min\left\{\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] \cdot \frac{\mathbb{P}\left[S = 1 | D = 1\right]}{\mathbb{P}\left[S = 1 | D = 0\right]}, 1\right\} + \mathbb{P}\left[Y = 0 | S = 1, D = 0\right] - 1
$$

\n
$$
\ge \mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y = 0 | S = 1, D = 0\right] - 1
$$

\nby Lemmas A.2 and A.3
\n
$$
= \mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1
$$

\nby Lemma A.4
\n
$$
= \mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]
$$

\nby Lemma A.5

 ≥ 0 ,

and

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$

$$
\geq \mathbb{P}[Y=0|S=1,D=0] - \mathbb{P}[Y=0|S=1,D=0]
$$

$$
\geq 0,
$$

and

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 1.
$$

Moreover, note that $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 0, \tilde{S}_1 = 1 \in [0, 1]$ by construction.

Notice also that the data restrictions are satisfied because

$$
\mathbb{P}\left[\tilde{Y} = 1 | \tilde{S} = 1, \tilde{D} = 0\right]
$$

= $\mathbb{P}\left[\tilde{Y}_0^* = 1 | \tilde{S}_0 = 1\right]$
= $\mathbb{P}\left[\tilde{Y}_0^* = 1 | \tilde{S}_0 = 1, S_1 = 1\right]$
= $\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, S_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 | \tilde{S}_0 = 1, S_1 = 1\right]$
= $\mathbb{P}\left[Y = 1 | S = 1, D = 0\right]$

and

$$
\begin{aligned}\n\mathbb{P}\left[\tilde{Y}=1\Big|\tilde{S}=1,\tilde{D}=1\right] \\
&= \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{1}=1\right] \\
&= \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\cdot\mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\Big|\tilde{S}_{1}=1\right] \\
&+ \mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]\cdot\left(1-\mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\Big|\tilde{S}_{1}=1\right]\right), \\
&= \left(\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]\right) \\
&\cdot \mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\Big|\tilde{S}_{1}=1\right] \\
&\quad + \left(\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]\right) \\
&\cdot\left(1-\mathbb{P}\left[\tilde{S}_{0}=1,\tilde{S}_{1}=1\Big|\tilde{S}_{1}=1\right]\right)\n\end{aligned}
$$

$$
= \min \left\{ \mathbb{P}\left[Y=1|S=1,D=1\right] \cdot \frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]}, 1 \right\} \cdot \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]} + \max \left\{ \frac{\mathbb{P}\left[Y=1|S=1,D=1\right] - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}, 0 \right\} \cdot \left(1 - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]} \right) = \min \left\{ \mathbb{P}\left[Y=1|S=1,D=1\right], \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]} \right\} + \max \left\{ \mathbb{P}\left[Y=1|S=1,D=1\right] - \frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}, 0 \right\} = \mathbb{P}\left\{ Y=1|S=1,D=1 \right].
$$

Finally, note that

$$
\begin{split} &\mathbb{P}\left[\tilde{Y}_{1}^{*}=1\Big|\tilde{Y}_{0}^{*}=0,\tilde{S}_{0}=1,\tilde{S}_{1}=1\right] \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=0\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &=\frac{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]}{\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]+\mathbb{P}\left[Y=0|S=1,D=0\right]-\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=1,\tilde{S}_{1}=1\right]} \\ &=\frac{\min\left\{\mathbb{P}\left[Y=1|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=1\right]}{\mathbb{P}\left[S=1|D=0\right]},1\right\}+\mathbb{P}\left[Y=0|S=1,D=0\right]-1}{\mathbb{P}\left[Y=0|S=1,D=0\right]} \end{split}
$$

 $= U B_2.$

(Part c) Constructing a conditional distribution that attains any $\tilde{\theta}\in (\mathbf{LB_2}, \mathbf{UB_2})$ This part of the proof is identical to the proof explained in Appendix [A.2.](#page-25-1)

A.5 Proof of Lemma [A.5](#page-44-1)

Observe that

$$
\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]
$$

$$
= \mathbb{P}[Y_1^* = 1, Y_0^* = 1 | S_0 = 1, S_1 = 1] + \mathbb{P}[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1]
$$

\n
$$
- \mathbb{P}[Y_1^* = 1, Y_0^* = 1 | S_0 = 1, S_1 = 1]
$$

\n
$$
= \mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] - \mathbb{P}[Y_1^* = 1, Y_0^* = 1 | S_0 = 1, S_1 = 1]
$$

\n
$$
= \mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] - \mathbb{P}[Y_0^* = 1 | S_0 = 1, S_1 = 1]
$$

\nby Assumption 4
\n
$$
= \mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] + \mathbb{P}[Y_0^* = 0 | S_0 = 1, S_1 = 1] - 1.
$$

A.6 Proof of Proposition [4](#page-11-0)

To prove Proposition [4,](#page-11-0) we first show that $LB_3 \leq \theta^{OO}$ and $\theta^{OO} \leq UB_3$. Then, we show that LB_3 and UB_3 are sharp bounds. For completeness, we state one lemma that was previously derived in the literature and is used in our proofs. We prove it in Appendix [A.7.](#page-61-0)

Lemma A.6 [Chen and Flores](#page-22-0) [\(2015\)](#page-22-0): Under Assumptions [1,](#page-6-2) [2](#page-7-1) and [5,](#page-8-1) we have that

$$
\mathbb{P}[Y_1^* = 1 | S_0 = 1, S_1 = 1] \ge \mathbb{P}[Y = 1 | S = 1, D = 1].
$$

A.6.1 Lower Bound: $LB_3 \n\leq \theta^{OO}$

Note that

$$
\theta^{OO} := \mathbb{P}\left[Y_1^* = 1 | Y_0^* = 0, S_0 = 1, S_1 = 1\right]
$$

=
$$
\frac{\mathbb{P}\left[Y_1^* = 1, Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

=
$$
\frac{\mathbb{P}\left[Y_1^* = 1 | S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right] - 1}{\mathbb{P}\left[Y_0^* = 0 | S_0 = 1, S_1 = 1\right]}
$$

by Lemma [A.5](#page-44-1) $\geq \frac{\mathbb{P}[Y=1|S=1,D=1]+\mathbb{P}[Y_0^*=0|S_0=1,S_1=1]-1}{\sum_{i=1}^{m}Y_i^*=[0|S_0=1,S_1=1]}$ $\mathbb{P}[Y_0^* = 0 | S_0 = 1, S_1 = 1]$

by Lemma A.6
=
$$
\frac{\mathbb{P}[Y=1|S=1,D=1]+\mathbb{P}[Y=0|S=1,D=0]-1}{\mathbb{P}[Y=0|S=1,D=0]}
$$

by Lemma [A.4.](#page-26-0)

Moreover, $\theta^{OO} \geq 0$ by definition.

A.6.2 Upper Bound: $\theta^{OO} \leq UB_2$

The proof is identical to the proof explained in Appendix [A.4.](#page-44-0)

A.6.3 LB_2 and UB_2 are sharp bounds

The only difference between this proof and the proof in Appendix [A.2](#page-25-1) is the definition of $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = s_0, \tilde{S}_1 = s_1$ for any $(y_0, y_1, s_0, s_1) \in \{0, 1\}^4$. For this reason, we will only construct a conditional distribution $(\tilde{Y}_0^*, \tilde{Y}_1^*)$ $(\tilde{S}_0^*, \tilde{S}_1^*)$ that is a probability distribution, satisfies Assumption [5,](#page-8-1) satisfies the data restrictions, and generates a probability of causation θ respectively equal to:

- (a) the lower bound LB_3 ;
- (b) the upper bound UB_3 ;
- (c) any value in the interval (LB_3, UB_3) .

(Part a) Constructing a conditional distribution such that $\tilde{\theta} = \text{LB}_3$ Since $\mathbb{P}\left[\tilde{S}_0 = 1, \tilde{S}_1 = 0\right] = 0$, we do not need to define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 0.$ We define $\mathbb{P}\left[\tilde{Y}_0^* = y_0, \tilde{Y}_1^* = y_1\right]$ $[\tilde{S}_0 = 0, \tilde{S}_1 = 0] = \frac{1}{4}$ for any $(y_0, y_1) \in \{0, 1\}^2$. We also define

the conditional probabilities

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$
\n
$$
= \max \left\{\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] + \mathbb{P}\left[Y = 0 | S = 1, D = 0\right] - 1, 0\right\}
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[Y = 0 | S = 1, D = 0\right]
$$
\n(A.59)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right]
$$
\n(A.60)

$$
= \min \left\{ 1 - \mathbb{P} \left[Y = 1 | S = 1, D = 1 \right], \mathbb{P} \left[Y = 0 | S = 1, D = 0 \right], 0 \right\},\
$$

$$
\mathbb{P} \left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1 \right] = 0
$$
 (A.61)

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_0^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$
\n
$$
\mathbb{P}\left[\tilde{Y}_0^* = 1 \middle| \tilde{S}_0 = 1\right] = \mathbb{P}\left[\tilde{S}_0^* = 1 \middle| \tilde{S}_0 = 1\right] = \mathbb{P}\left[S = 1 \middle| D = 0\right]
$$
\n(A.62)

$$
= \frac{\mathbb{P}[Y=1, | S=1, D=1] - \mathbb{P}\left[\tilde{Y}_1^* = 1 | \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \cdot \frac{\mathbb{P}[S=1 | D=0]}{\mathbb{P}[S=1 | D=1]}}{1 - \frac{\mathbb{P}[S=1 | D=0]}{\mathbb{P}[S=1 | D=1]}},
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0,\tag{A.63}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 1 - \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right],\tag{A.64}
$$

$$
\mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 0 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = 0. \tag{A.65}
$$

To check that Assumption [5](#page-8-1) holds, we have to analyze two cases.

Case 1) $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0=1, \tilde{S}_1=1\big]>0$

In this case, we have that

$$
\mathbb{P}\left[\tilde{Y}_{1}^{*} = 1 | \tilde{S}_{0} = 1, \tilde{S}_{1} = 1\right]
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{0}^{*} = 0, \tilde{Y}_{1}^{*} = 1 | \tilde{S}_{0} = 1, \tilde{S}_{1} = 1\right] + \mathbb{P}\left[\tilde{Y}_{0}^{*} = 1, \tilde{Y}_{1}^{*} = 1 | \tilde{S}_{0} = 1, \tilde{S}_{1} = 1\right]
$$
\n
$$
= \mathbb{P}\left[Y = 1 | S = 1, D = 1\right]
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{0}^{*} = 0, \tilde{Y}_{1}^{*} = 1 | \tilde{S}_{0} = 0, \tilde{S}_{1} = 1\right] + \mathbb{P}\left[\tilde{Y}_{0}^{*} = 1, \tilde{Y}_{1}^{*} = 1 | \tilde{S}_{0} = 0, \tilde{S}_{1} = 1\right]
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{1}^{*} = 1 | \tilde{S}_{0} = 0, \tilde{S}_{1} = 1\right].
$$

Case 2) $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 1 = 0$

In this case, we have that

$$
\begin{aligned} \mathbb{P}\left[\tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \\ &= \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \\ &= \mathbb{P}\left[Y = 1 | S = 1, D = 0\right] \end{aligned}
$$

$$
\mathbb{P}\left[\tilde{Y}_{1}^{*} = 1 \middle| \tilde{S}_{0} = 0, \tilde{S}_{1} = 1\right]
$$
\n
$$
= \mathbb{P}\left[\tilde{Y}_{0}^{*} = 0, \tilde{Y}_{1}^{*} = 1 \middle| \tilde{S}_{0} = 0, \tilde{S}_{1} = 1\right] + \mathbb{P}\left[\tilde{Y}_{0}^{*} = 1, \tilde{Y}_{1}^{*} = 1 \middle| \tilde{S}_{0} = 0, \tilde{S}_{1} = 1\right]
$$
\n
$$
= \frac{\mathbb{P}\left[Y = 1, | S = 1, D = 1\right] - \mathbb{P}\left[Y = 1 | S = 1, D = 0\right] \cdot \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}
$$
\n
$$
1 - \frac{\mathbb{P}\left[S = 1 | D = 0\right]}{\mathbb{P}\left[S = 1 | D = 1\right]}
$$

by Equation [\(A.62\)](#page-57-0) and the last result,

implying that

$$
\mathbb{P}\left[\tilde{Y}_{1}^{*} = 1 \middle| \tilde{S}_{0} = 1, \tilde{S}_{1} = 1\right] - \mathbb{P}\left[\tilde{Y}_{1}^{*} = 1 \middle| \tilde{S}_{0} = 0, \tilde{S}_{1} = 1\right] \\
= \frac{\left(1 - \frac{\mathbb{P}\left[S = 1 \middle| D = 0\right]}{\mathbb{P}\left[S = 1 \middle| D = 1\right]}\right) \cdot \mathbb{P}\left[Y = 1 \middle| S = 1, D = 0\right]}{1 - \frac{\mathbb{P}\left[S = 1 \middle| D = 0\right]}{\mathbb{P}\left[S = 1 \middle| D = 1\right]}} \\
- \frac{\mathbb{P}\left[Y = 1, | S = 1, D = 1] - \mathbb{P}\left[Y = 1 | S = 1, D = 0\right] \cdot \frac{\mathbb{P}\left[S = 1 \middle| D = 0\right]}{\mathbb{P}\left[S = 1 \middle| D = 1\right]}}{1 - \frac{\mathbb{P}\left[S = 1 \middle| D = 0\right]}{\mathbb{P}\left[S = 1 \middle| D = 1\right]}} \\
= \frac{\mathbb{P}\left[Y = 1 | S = 1, D = 0] - \mathbb{P}\left[Y = 1, | S = 1, D = 1\right]}{1 - \frac{\mathbb{P}\left[S = 1 \middle| D = 0\right]}{\mathbb{P}\left[S = 1 \middle| D = 1\right]}} \\
\geq 0
$$

by Equation [\(A.58\)](#page-56-0) and the assumption that $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 1 = 0.$

(Part a.1) The candidate conditional distribution is a probability distribution Now, we only have to show that $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $(\tilde{S}_0 = 0, \tilde{S}_1 = 1) \in [0, 1].$ We have to analyze two cases.

Case 1) $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0=1, \tilde{S}_1=1\big]>0$

and

In this case, we have that

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right] = \mathbb{P}\left[Y = 1 | S = 1, D = 1\right] \in [0, 1]
$$

according to Equations [\(A.58\)](#page-56-0), [\(A.59\)](#page-56-1) and [\(A.62\)](#page-57-0).

Case 2) $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0=1, \tilde{S}_1=1 \Bigr]=0$

In this case, we have that

$$
\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 0, \tilde{S}_1 = 1\right]
$$
\n
$$
= \frac{\mathbb{P}\left[Y = 1 \middle| S = 1, D = 1\right] - \mathbb{P}\left[Y = 1 \middle| S = 1, D = 0\right] \cdot \frac{\mathbb{P}\left[S = 1 \middle| D = 0\right]}{\mathbb{P}\left[S = 1 \middle| D = 1\right]}}{1 - \frac{\mathbb{P}\left[S = 1 \middle| D = 0\right]}{\mathbb{P}\left[S = 1 \middle| D = 1\right]}}
$$
\n
$$
\propto \mathbb{P}\left[Y = 1 \middle| S = 1, D = 1\right] - \mathbb{P}\left[Y = 1 \middle| S = 1, D = 0\right] \cdot \frac{\mathbb{P}\left[S = 1 \middle| D = 0\right]}{\mathbb{P}\left[S = 1 \middle| D = 1\right]}
$$
\nby Lemma A.3

$$
\propto \mathbb{P}[Y=1|S=1, D=1] \cdot \mathbb{P}[S=1|D=1] - \mathbb{P}[Y=1|S=1, D=0] \cdot \mathbb{P}[S=1|D=0]
$$

= $\mathbb{P}[Y_1^* = 1, S_1 = 1] - \mathbb{P}[Y_0^* = 1, S_0 = 1]$

by Assumption [1](#page-6-2)

$$
= \mathbb{P}\left[Y_1^* = 0, Y_1^* = 1, S_0 = 0, S_1 = 1\right] + \mathbb{P}\left[Y_1^* = 1, Y_1^* = 1, S_0 = 0, S_1 = 1\right]
$$

+ $\mathbb{P}\left[Y_1^* = 0, Y_1^* = 1, S_0 = 1, S_1 = 1\right] + \mathbb{P}\left[Y_1^* = 1, Y_1^* = 1, S_0 = 1, S_1 = 1\right]$
- $\mathbb{P}\left[Y_0^* = 1, Y_1^* = 1, S_0 = 1, S_1 = 1\right]$

by Assumptions [3](#page-7-0) and [4](#page-7-3)

$$
= \mathbb{P}\left[Y_1^* = 0, Y_1^* = 1, S_0 = 0, S_1 = 1\right] + \mathbb{P}\left[Y_1^* = 1, Y_1^* = 1, S_0 = 0, S_1 = 1\right]
$$

+ $\mathbb{P}\left[Y_1^* = 0, Y_1^* = 1, S_0 = 1, S_1 = 1\right]$
 $\geq 0.$

We also have that

$$
\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\Big|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]
$$
\n
$$
=\frac{\mathbb{P}\left[Y=1,|S=1,D=1\right]-\mathbb{P}\left[Y=1|S=1,D=0\right]\cdot\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}
$$
\n
$$
\leq\frac{\mathbb{P}\left[Y=1,|S=1,D=1\right]-\mathbb{P}\left[Y=1,|S=1,D=1\right]\cdot\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}{1-\frac{\mathbb{P}\left[S=1|D=0\right]}{\mathbb{P}\left[S=1|D=1\right]}}
$$
\n
$$
=\mathbb{P}\left[Y=1,|S=1,D=1\right]
$$
\n
$$
\leq 1
$$

by Equation [\(A.58\)](#page-56-0) and the assumption that $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 1 = 0.$

(Part a.2) The candidate conditional distribution satisfies its data restrictions This part of the proof is identical to the proof explained in Appendix [A.4.](#page-44-0)

(Part a.3) The probability of causation $\tilde{\theta}$ reaches the lower bound LB_1

This part of the proof is identical to the proof explained in Appendix [A.4.](#page-44-0)

(Part b) Constructing a conditional distribution such that $\tilde{\theta} = \text{UB}_3$

Here, we use the same distribution that attains the upper bound UB_2 in Appendix [A.4.](#page-44-0) For this reason, we only have to show that the distribution in Appendix [A.4](#page-44-0) also satisfies Assumption [5.](#page-8-1) Note that Equations $(A.50)-(A.57)$ $(A.50)-(A.57)$ $(A.50)-(A.57)$ imply that

$$
\begin{aligned} \mathbb{P}\left[\tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \\ &= \mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] + \mathbb{P}\left[\tilde{Y}_0^* = 1, \tilde{Y}_1^* = 1 \middle| \tilde{S}_0 = 1, \tilde{S}_1 = 1\right] \\ &= \min\left\{\mathbb{P}\left[Y = 1 | S = 1, D = 1\right] \cdot \frac{\mathbb{P}\left[S = 1 | D = 1\right]}{\mathbb{P}\left[S = 1 | D = 0\right]}, 1\right\} \end{aligned}
$$

$$
\mathbb{P}\left[\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]
$$
\n
$$
=\mathbb{P}\left[\tilde{Y}_{0}^{*}=0,\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]+\mathbb{P}\left[\tilde{Y}_{0}^{*}=1,\tilde{Y}_{1}^{*}=1\middle|\tilde{S}_{0}=0,\tilde{S}_{1}=1\right]
$$
\n
$$
=\max\left\{\frac{\mathbb{P}\left[Y=1\middle|S=1,D=1\right]-\frac{\mathbb{P}\left[S=1\middle|D=0\right]}{\mathbb{P}\left[S=1\middle|D=1\right]}}{1-\frac{\mathbb{P}\left[S=1\middle|D=0\right]}{\mathbb{P}\left[S=1\middle|D=1\right]}},0\right\}.
$$

Consequently, we have to analyze two cases. If $\mathbb{P}\left[\tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 1 \Big] < 1$, then $\mathbb{P}\left[\tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 0, \tilde{S}_1 = 1\Big]$ = 0 and Assumption [5](#page-8-1) holds. If $\mathbb{P}\left[\tilde{Y}_1^* = 1\right]$ $\tilde{S}_0 = 1, \tilde{S}_1 = 1$] = 1, then $\mathbb{P}\left[\tilde{Y}_1^* = 1\right]$ $\tilde{S}_0=0,\tilde{S}_1=1\Big]=$ $\mathbb{P}\left[\tilde{Y}_0^* = 0, \tilde{Y}_1^* = 1\right]$ $[\tilde{S}_0 = 0, \tilde{S}_1 = 1] \leq 1$ according to Appendix [A.4,](#page-44-0) implying that Assumption [5](#page-8-1) holds.

(Part c) Constructing a conditional distribution that attains any $\tilde{\theta}\in (\mathbf{LB_2}, \mathbf{UB_2})$

This part of the proof is identical to the proof explained in Appendix [A.2.](#page-25-1)

A.7 Proof of Lemma [A.6](#page-55-1)

Observe that

$$
\mathbb{P}\left[Y=1|S=1,D=1\right]
$$
\n
$$
= \mathbb{P}\left[Y_1^*=1|S_1=1\right]
$$
\nby Assumption 1\n
$$
= \mathbb{P}\left[Y_1^*=1|S_0=1,S_1=1\right] \cdot \mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]
$$
\n
$$
+ \mathbb{P}\left[Y_1^*=1|S_0=0,S_1=1\right] \cdot \left(1-\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]\right)
$$
\n
$$
\leq \mathbb{P}\left[Y_1^*=1|S_0=1,S_1=1\right] \cdot \mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]
$$
\n
$$
+ \mathbb{P}\left[Y_1^*=1|S_0=1,S_1=1\right] \cdot \left(1-\mathbb{P}\left[S_0=1,S_1=1|S_1=1\right]\right)
$$
\nby Assumption 5\n
$$
= \mathbb{P}\left[Y_1^*=1|S_0=1,S_1=1\right].
$$

and