# Doubly Robust Proximal Synthetic Controls

Hongxiang Qiu<sup>1</sup>, Xu Shi<sup>2</sup>, Wang Miao<sup>3</sup>, Edgar Dobriban<sup>4</sup>, and Eric Tchetgen Tchetgen<sup>∗</sup><sup>4</sup>

<sup>1</sup>Department of Epidemiology and Biostatistics, Michigan State University

<sup>2</sup>Department of Biostatistics, University of Michigan

<sup>3</sup>Department of Probability and Statistics, Peking University

<sup>4</sup>Department of Statistics and Data Science, University of Pennsylvania

#### Abstract

To infer the treatment effect for a single treated unit using panel data, synthetic control methods construct a linear combination of control units' outcomes that mimics the treated unit's pre-treatment outcome trajectory. This linear combination is subsequently used to impute the counterfactual outcomes of the treated unit had it not been treated in the post-treatment period, and used to estimate the treatment effect. Existing synthetic control methods rely on correctly modeling certain aspects of the counterfactual outcome generating mechanism and may require near-perfect matching of the pre-treatment trajectory. Inspired by proximal causal inference, we obtain two novel nonparametric identifying formulas for the average treatment effect for the treated unit: one is based on weighting, and the other combines models for the counterfactual outcome and the weighting function. We introduce the concept of covariate shift to synthetic controls to obtain these identification results conditional on the treatment assignment. We also develop two treatment effect estimators based on these two formulas and the generalized method of moments. One new estimator is doubly robust: it is consistent and asymptotically normal if at least one of the outcome and weighting models is correctly specified. We demonstrate the performance of the methods via simulations and apply them to evaluate the effectiveness of a Pneumococcal conjugate vaccine on the risk of all-cause pneumonia in Brazil.

<sup>∗</sup>Author e-mail addresses: qiuhongx@msu.edu, shixu@umich.edu, mwfy@pku.edu.cn, dobriban@wharton.upenn.edu, ett@wharton.upenn.edu

# 1 Introduction

#### 1.1 Background

Interventions such as policies are often implemented in a single unit such as a state, a city, or a school. Causal inference in these cases is challenging due to the small number of treated units, and due to the lack of randomization and independence. In various fields including economics, public health, and biometry, synthetic control (SC) methods [\[Abadie and Gardeazabal, 2003,](#page-58-0) [Abadie et al., 2010,](#page-58-1) [2015,](#page-58-2) [Doudchenko and Imbens, 2016\]](#page-60-0) are a common tool to estimate the intervention (or treatment) effect for the treated unit in time series from a single treated unit and multiple untreated units in both pre- and post-treatment periods. For example, SC methods have been used to estimate the effects of terrorist conflicts on GDP [\[Abadie and Gardeazabal, 2003\]](#page-58-0), tobacco control program on tobacco consumption [\[Abadie et al., 2010\]](#page-58-1), Kansas's tax cut on GDP [\[Ben-Michael](#page-59-0) [et al., 2021b,](#page-59-0) [Rickman and Wang, 2018\]](#page-62-0), Florida's "stand your ground" law on homicide rates [\[Bonander et al., 2021\]](#page-59-1), and pneumococcal conjugate vaccines on pneumonia [\[Bruhn](#page-59-2) [et al., 2017\]](#page-59-2).

Classical SCs are linear combinations of control units that mimic the treated unit before the treatment. Outcome differences between the treated unit and the SC in the post-treatment period are used to make inferences about the treatment effect for the treated unit. In [Abadie and Gardeazabal](#page-58-0) [\[2003\]](#page-58-0) and [Abadie et al.](#page-58-1) [\[2010\]](#page-58-1), a SC is a weighted average of a pool of control units, called the donors. The weights are obtained by minimizing a distance between the SC and the treated unit in the pre-treatment period, under the constraint that the weights are non-negative and sum to unity. Many extensions have been proposed. For example, [Abadie and L'Hour](#page-58-3) [\[2021\]](#page-58-3) proposed methods for multiple treated units, and [Ben-Michael et al.](#page-59-3) [\[2021a\]](#page-59-3) further considered the case where these treated units initiate treatment at different time points; [Doudchenko and Imbens](#page-60-0) [\[2016\]](#page-60-0) and [Ben-Michael et al.](#page-59-3) [\[2021a,](#page-59-3)[b\]](#page-59-0) introduced penalization to improve performance; [Athey et al.](#page-59-4) [\[2021\]](#page-59-4) and [Bai and Ng](#page-59-5) [\[2021\]](#page-59-5) used techniques from matrix completion; [Li](#page-62-1) [\[2020\]](#page-62-1) studied statistical inference for SC methods; [Chernozhukov et al.](#page-60-1) [\[2021\]](#page-60-1) and [Cattaneo et al.](#page-59-6) [\[2021\]](#page-59-6) considered prediction intervals for treatment effects. Among these extensions, some also incorporate the idea that, similarly to the control units' outcomes in the post-treatment period, the treated unit's outcomes in the pre-treatment period can be used to impute the counterfactual outcome had it not been treated [\[Ben-Michael](#page-59-0) [et al., 2021b,](#page-59-0) [Arkhangelsky et al., 2021\]](#page-58-4).

Existing methods often rely on assuming linear models and on the existence of nearperfectly matching weights in the observed data. Under such assumptions, valid SCs are linear combinations, often weighted averages, of donors. However, if such assumptions do not hold, these methods may not produce a valid SC. This may happen if the outcomes in the donors have a different measuring scale from the treated unit, or if the treated unit's and the donors' outcomes have a nonlinear relationship.

To relax these assumptions, [Shi et al.](#page-63-0) [\[2023\]](#page-63-0) viewed SCs from the proximal causal inference perspective. For independent and identically distributed (i.i.d.) observations, [Miao et al.](#page-62-2) [\[2018\]](#page-62-2), [Deaner](#page-60-2) [\[2018,](#page-60-2) [2021\]](#page-60-3), [Cui et al.](#page-60-4) [\[2020\]](#page-60-4), [Tchetgen Tchetgen et al.](#page-64-0) [\[2020\]](#page-64-0) derived nonparametric identification using proxies, variables capturing the effect of the unmeasured confounders. [Shi et al.](#page-63-0) [\[2023\]](#page-63-0) viewed control units' outcomes as proxies and obtained nonparametric identification results for the potential outcome of the treated unit had it not been treated as well as the treatment effect in a general setting, beyond the common linear factor model [e.g., [Abadie et al., 2010\]](#page-58-1). They assumed the existence of a function of these proxies, termed confounding bridge function, that captures the (possibly nonlinear) effects of unobserved confounders. With this function, they imputed the expected counterfactual outcomes for the treated unit. Estimation of, and inference about, the average treatment effect for the treated unit (ATT) followed from this identification result. Instrumental variables have also been used. For example, [Holtz-Eakin](#page-61-0) [et al.](#page-61-0) [\[1988\]](#page-61-0) considered a linear model with interactive fixed effects and showed how to identify it using appropriate instruments. The solution to this problem relies on a particular differencing strategy, which may be viewed as an application of a confounding bridge function. [Cunha et al.](#page-60-5) [\[2010\]](#page-60-5), [Freyberger](#page-60-6) [\[2018\]](#page-60-6) and references therein considered general nonparametric models with interactive effects, showing how to identify it using appropriate instruments. While treatment confounding proxies in proximal causal inference are sometimes described as instruments, it is crucial to note that they are more general than instrumental variables (IVs), in the sense that valid IVs are valid treatment confounding proxies, but invalid IVs dependent on hidden confounders are also valid treatment confounding proxies [\[Tchetgen Tchetgen et al., 2020\]](#page-64-0). In addition, while IVs require a form of homogeneity condition for nonparametric identification (e.g., separable errors or monotonicity), proxies do not require such a condition.

#### 1.2 Our contribution

Existing methods rely on correctly specifying an outcome model, based on which one can impute the counterfactual outcome trajectory of the treated unit, had it not been treated, after treatment. This outcome bridge function model may be difficult to specify correctly, or may not exist. In this paper, we relax this requirement by leveraging the proximal causal inference framework as in [Shi et al.](#page-63-0) [\[2023\]](#page-63-0). We develop two novel methods to estimate the ATT. One method relies on weighting and is a building block to a second method which we rigorously prove is doubly robust [\[Bang and Robins, 2005,](#page-59-7) [Scharfstein](#page-63-1) [et al., 1999\]](#page-63-1). It is consistent and asymptotically normal if either the outcome model or the weighting function is correctly specified, without requiring that both are. An advantage of the doubly robust method compared to existing methods is that it allows for misspecifing one of the two models, without the user necessarily knowing which might be misspecified.

We observed that our estimand of interest, the ATT, is closely related to the average

treatment effect on the treated for i.i.d. data [e.g., [Hahn, 1998,](#page-61-1) [Imbens, 2004,](#page-61-2) [Chen](#page-60-7) [et al., 2008,](#page-60-7) [Shu and Tan, 2018\]](#page-64-1). The method in [Shi et al.](#page-63-0) [\[2023\]](#page-63-0) corresponds to using an outcome confounding bridge function [\[Miao et al., 2018\]](#page-62-2), which is the proximal causal inference counterpart of G-computation, or an outcome regression-based approach in causal inference under unconfoundedness [\[Robins, 1986\]](#page-62-3). Our proposed methods are motivated by the existing identification results in proximal causal inference in the i.i.d. setting [\[Cui et al., 2020\]](#page-60-4): one result is based on weighting and the other is based on the influence function.

Despite these similarities, it remains challenging to adapt these ideas from the i.i.d. setting to panel data. Since treatment assignment is often viewed as fixed in SC problems, a key concept from the i.i.d. setting, the propensity score [\[Rosenbaum and Rubin,](#page-63-2) [1983\]](#page-63-2), is undefined. Thus, existing results for the i.i.d. setting cannot be directly applied to SC problems. We leverage the notion of *covariate shift* [e.g., Quiñonero-Candela et al., [2009\]](#page-62-4) to circumvent this issue. We also find a relaxed version of the i.i.d. assumption to allow for serial correlation, while still obtaining identification via weighting. We illustrate our proposed methods in simulations and three empirical examples: two examples concern public health outcomes, one studying the effect of the PCV10 vaccine in Brazil on pneumonia [\[Bruhn et al., 2017\]](#page-59-2), and the other studying the effect of Florida's "stand your ground" law on homicide rates [\[Bonander et al., 2021\]](#page-59-1); the third example concerns economic outcomes, studying the effect of Kansas's tax cut on GDP [\[Rickman and Wang,](#page-62-0) [2018\]](#page-62-0).

Both our doubly robust method and the method in [Ben-Michael et al.](#page-59-0) [\[2021b\]](#page-59-0) take the form of augmented weighted moment equations, but the known robustness properties of these methods differ. In [Ben-Michael et al.](#page-59-0) [\[2021b\]](#page-59-0), the treated unit's counterfactual outcome had it not been treated after treatment can be imputed with two approaches. One is a weighted average of control units' outcomes, identical to classical SC methods [\[Abadie et al., 2010\]](#page-58-1); the other is a prediction model obtained using the treated unit's outcomes before treatment. By combining these two approaches, [Ben-Michael et al.](#page-59-0) [\[2021b\]](#page-59-0) developed a SC method with improved performance. [Arkhangelsky et al.](#page-58-4) [\[2021\]](#page-58-4) proposed a method combining two imputation approaches based on similar ideas. Nevertheless, to date, neither method has formally been shown to be doubly robust. In contrast, we formally establish double robustness, inherited from the influence function of the ATT in i.i.d cases.

### 2 Problem setup

We observe data over T time periods. The first  $T_0$  time periods are the pre-treatment periods, and the last  $T - T_0$  time periods are the post-treatment periods. For the treated unit, at each time period  $t = 1, \ldots, T$ , let  $Y_t(0), Y_t(1) \in \mathbb{R}$  be the counterfactual outcome corresponding to no treatment and treatment, respectively, and  $Y_t = \mathbb{1}(t \leq T_0)Y_t(0)$  +  $\mathbb{1}(t > T_0)Y_t(1)$  be the observed outcome. At each time period t, other variables such as other control units' outcomes are observed. We provide more details about these variables below. We treat  $T$ ,  $T_0$  and the treated unit as deterministic, and treat other variables such as the treated unit's potential outcomes  $Y_t(1)$  and  $Y_t(0)$  as random. In other words, our proposed methods are conditional on the study design. We study the ATT causal estimand, that is,

$$
\phi^*(t) = E\{Y_t(1) - Y_t(0)\}
$$

in a post-treatment period  $t > T_0$ . We treat times, namely T and  $T_0$ , and all units as deterministic, and treat the potential outcomes as stochastic [\[Greenland, 1987,](#page-61-3) [Robins](#page-63-3) [and Greenland, 1989,](#page-63-3) [2000,](#page-63-4) [VanderWeele and Robins, 2012\]](#page-64-2); that is,  $Y_t(0)$  and  $Y_t(1)$  are both stochastic processes indexed by t that are randomly generated over time, rather than fixed unknown scalar sequences. In the frequentist interpretation, under repeated sampling, the times and units are all fixed and hence identical for all samples, but the outcomes are randomly generated from a fixed unknown joint distribution and hence may differ across samples. Stochastic counterfactuals are commonly assumed in the SC literature, implicitly in the random noise or residuals of a linear latent factor model or an autoregressive model [e.g., [Abadie et al., 2010,](#page-58-1) [Abadie and L'Hour, 2021,](#page-58-3) [Ben-Michael](#page-59-0) [et al., 2021b,](#page-59-0)[a,](#page-59-3) Athey et al., 2021. This notion of stochastic counterfactuals  $Y_t(0)$  and  $Y_t(1)$  as time series is required in our paper because the expectation in  $\phi^*(t)$  is taken over the joint distribution of  $(Y_t(0), Y_t(1))$ .

In the main text, we focus on the case without covariates, and discuss using covariates in Web Appendix [S5.](#page-32-0) We assume that all unmeasured confounding is captured by a latent factor  $U_t$ , with assumptions stated in later sections. We use  $t_-\,$  and  $t_+$  to denote generic times before and after treatment, respectively; that is,  $t_-\leq T_0$  and  $t_+ > T_0$ . When stating asymptotic results, we consider the asymptotic regime where  $T \to \infty$ with  $T_0/T \to \gamma \in (0,1)$ . This asymptotic regime may be interpreted as the number of observations in both pre- and post-treatment periods growing to infinity, collecting more data before and after the treatment time. Beyond this asymptotic regime, finite-sample results concerning the error in pre-treatment fitting or treatment effect estimation have been established in previous works [e.g., [Abadie and L'Hour, 2021,](#page-58-3) [Athey et al., 2021,](#page-59-4) [Ben-Michael et al., 2021a,](#page-59-3)[b\]](#page-59-0).

### 3 Review of identification via outcome modeling

In classical SC methods, a weighted average of a pool of control units called donors forms the SC [\[Abadie et al., 2010\]](#page-58-1). The motivation for using control units' outcomes to learn about  $Y_{t+}(0)$ , despite the presence of potential unmeasured confounder  $U_t$ , is that these control units may be affected by, and thus contains information about,  $U_t$ . This characteristic resembles that of proxies in proximal causal inference. In an i.i.d. setting, proxies

<span id="page-5-1"></span>

Figure [1](#page-5-0): Causal graphs satisfying Condition 1 at each time period t. The variable  $U_t$  is the unobserved confounder. In Figure [1a,](#page-5-1) additional unmeasured confounding between proxy  $W_t$  and the treated unit's outcome  $Y_t$  may be present. In Figure [1b,](#page-5-1)  $Z_t$ ,  $W_t$  and  $Y_t$  are mutually independent conditional on  $U_t$ , which is often sensible when  $(W_t, Z_t)$  are control units' outcomes

.

capture the effect of the unmeasured confounder on the outcome or treatment assignment. They may also be viewed as noisy observations of the unmeasured confounder. In a panel data setting, for each time t, we use  $W_t \in \mathcal{W}$  to denote a proxy, the vector of the donors' outcomes in classical SC settings. Under commonly assumed data-generating assumptions such as the linear factor model [\[Abadie et al., 2010\]](#page-58-1), donors' outcomes are ideal proxies of  $U_t$ : any variation in  $U_t$  induces some variation in  $W_t$ . We use  $Z_t \in \mathcal{Z}$  to denote a general supplemental proxy. We next state the causal conditions required and discuss the role of and the choice of  $Z_t$ .

<span id="page-5-0"></span>Condition 1. For all pre-treatment time points  $t_-,$ , the supplemental proxies are independent of the outcomes and the proxies, conditional on the confounders:  $Z_{t-} \perp (Y_{t-}, W_{t-})$  $U_{t-}.$ 

Conditions similar to Condition [1](#page-5-0) are common in proximal causal inference literature [e.g., [Miao and Tchetgen Tchetgen, 2016,](#page-62-5) [Miao et al., 2018,](#page-62-2) [Cui et al., 2020,](#page-60-4) [Tchet](#page-64-0)[gen Tchetgen et al., 2020\]](#page-64-0). As we will show later,  $W_t$  is used to model the outcome  $Y_t$ with  $Z_t$  supplementing for identification, while  $Z_t$  is used to model the weighting process introduced in the following section with  $W_t$  supplementing for identification. The conditional independence in Condition [1](#page-5-0) is implied by the factor model in classical SC [e.g., [Abadie et al., 2010\]](#page-58-1), so such assumptions are commonly made implicitly. Causal graphs of Condition [1](#page-5-0) are in Figure [1.](#page-5-1)

<span id="page-5-2"></span>**Condition 2.** There exists a function  $h^*: W \to \mathbb{R}$ ,  $h^*(W_t)$  that captures the conditional mean of  $Y_t(0)$  given  $U_t$ :  $E\{h^*(W_t) | U_t\} = E\{Y_t(0) | U_t\}$ , for all time points t.

The function  $h^*$  is called an *outcome confounding bridge function* in proximal causal inference, a terminology we adopt. Condition [2](#page-5-2) states that the transformation  $h^*(W_t)$ of the observable proxy  $W_t$  matches the unobservable  $Y_t(0)$  in expectation conditional

on  $U_t$ . Sufficient conditions for the existence of  $h^*$  and examples of  $h^*$  can be found in [Shi et al.](#page-63-0) [\[2023\]](#page-63-0). The most popular model for  $h^*$  is the linear model in classical SC methods [\[Abadie et al., 2010\]](#page-58-1)—that is, the average of  $Y_t(0)$  may be imputed by a linear combination of the donors' outcomes  $W_t$ —but  $h^*$  may also be nonlinear. This assumption substantially generalizes the form of SCs by allowing more flexible models. In this condition, we implicitly assume that  $W_t$  is not causally impacted by the treatment because of the constant relationship over time.

Throughout, proxy  $W_t$  will consist of donors' outcomes a priori known not to be causally impacted by the treatment, to make Conditions [1](#page-5-0)[–2](#page-5-2) plausible. When many donors are viable, a subset may be selected based on, for example, the similarity of their outcome trajectories to the treated unit's trajectory before treatment. Typical choices of  $Z_t$  that may satisfy Condition [1](#page-5-0) include (i) outcomes of control units that are not valid donors, (ii) outcomes of donors excluded from the model  $h^*$  to impute  $Y_t(0)$ , and (iii) covariates of donors that are contemporaneous with  $(Y_t, W_t)$ . The proxy  $Z_t$  may be impacted by the treatment. [Shi et al.](#page-63-0) [\[2023\]](#page-63-0) Section 2.2. contains more discussion about how to choose proxies  $W_t$  and  $Z_t$ .

[Shi et al.](#page-63-0) [\[2023\]](#page-63-0) showed that, under Condition [2,](#page-5-2) for all post-treatment time points  $t_{+} > T_{0},$ 

<span id="page-6-2"></span>
$$
E\{Y_{t+}(0)\} = E\{h^*(W_{t+})\} \tag{1}
$$

and thus the ATT  $\phi^*(t_+) = E\{Y_{t_+} - h^*(W_{t_+})\}\$ . Additionally under Condition [1,](#page-5-0)  $h^*$  solves

<span id="page-6-0"></span> $E{Y_{t-} - h(W_{t-}) | Z_{t-}} = 0$  for all pre-treatment time points  $t- \leq T_0$  (2)

in  $h : \mathcal{W} \to \mathbb{R}$ . Further, under Condition [S1](#page-19-0) in Web Appendix [S1,](#page-19-1) [\(2\)](#page-6-0) has a unique solution.

**Remark 1.** In principle, it may be possible to allow  $h^*$  to depend on t, but it may be challenging to obtain a consistent estimator because only one observation is available at each time point; assumptions such as smoothness may be necessary. Our subsequent confounding bridge functions can also depend on t, but we do not pursue this direction.

# <span id="page-6-1"></span>4 Weighted and doubly robust identification of ATT

The method reviewed above is solely based on the treated unit's outcome process model, similar to G-computation under unconfoundedness [\[Robins, 1986\]](#page-62-3) and proximal G-computation [\[Tchetgen Tchetgen et al., 2020\]](#page-64-0). Now we introduce our first novel SC method, which is based on weighting and similar to inverse probability weighting under unconfoundedness [\[Robins et al., 1994\]](#page-63-5) and proximal inverse probability weighting [\[Cui et al., 2020\]](#page-60-4).

To illustrate the idea, for the moment, suppose that the observations are i.i.d. across time. Then, the estimand  $\phi^*(t_+) = E\{Y_{t+}(1) - Y_{t+}(0)\}\)$  corresponds to an average treatment effect on the treated, for which several identification formulas exist [e.g., [Hahn,](#page-61-1)

[1998,](#page-61-1) [Imbens, 2004,](#page-61-2) [Chen et al., 2008\]](#page-60-7), including those based on outcome regression and weighting. Therefore, one may identify  $E\{Y_{t+}(1) - Y_{t+}(0)\}\$  in a SC setting via weighting. To avoid the issue that propensity scores are undefined conditional on the design, we use the concept of covariate shift  $[e.g.,\ Qui\~{nonero-}\$ Candela et al., 2009 and likelihood ratio weighting. We next describe our needed causal conditions to identify the ATT via weighting for panel data.

<span id="page-7-0"></span>**Condition 3.** The joint conditional distribution of the counterfactual outcome  $Y_t(0)$  and proxy  $W_t$  given  $U_t$  is identical for all time points t.

Intuitively, this condition states that once the process  $U_t$  is generated,  $(Y_t(0), W_t)$  are then generated in the same way for all  $t$ . The invariance of the conditional distribution of  $W_t$  given  $U_t$  rules out a causal effect of the treatment on the proxies  $W_t$ , analogously to the exclusion restriction property of negative control outcomes [\[Lipsitch et al., 2010,](#page-62-6) [Miao et al., 2018\]](#page-62-2).

The next condition states that the marginal distribution of the confounder  $U_{t+}$  is identical for all  $t_{+}$ , which is implied by stationarity of  $U_{t_{+}}$ . Moreover, this condition ensures that only one approach is needed to impute post-treatment counterfactual outcomes  $Y_{t+}(0)$ .

<span id="page-7-1"></span>**Condition 4.** The distribution of the unobserved confounder  $U_{t_+}$  is identical for all post-treatment time points  $t_{+} > T_{0}$ .

Conditions [3](#page-7-0) and [4](#page-7-1) together imply that the distribution of  $(Y_{t+}(0), W_{t+}, U_{t+})$  is identical for all  $t_{+}$ . This implication holds under stationarity after treatment, without necessarily requiring stationarity before treatment. Even if stationarity also holds before treatment, these two conditions hold if the distributions in these two periods differ. Stationarity or similar are often used in time series analysis and other areas, including SCs [e.g., [Hsiao et al., 2012,](#page-61-4) [Hahn and Shi, 2017,](#page-61-5) [Li, 2020,](#page-62-1) [Cattaneo et al., 2021,](#page-59-6) [Chernozhukov](#page-60-1) [et al., 2021,](#page-60-1) [Ferman and Pinto, 2021\]](#page-60-8). Conditions [3](#page-7-0)[–4](#page-7-1) may thus be plausible in certain applications. Stationarity in Condition [4](#page-7-1) might be implausible when the number  $T - T_0$ of post-treatment time periods is large. We assume Condition [4](#page-7-1) to facilitate the presentation and present an identification approach applicable to non-stationary cases in Web Appendix [S3.1.](#page-23-0) These two conditions allow an instantaneous distributional shift in the unobserved confounder  $U_t$  at treatment  $T_0$ . Such an instantaneous shift is a source of confounding. Therefore, in general, pre-treatment outcomes  $Y_{t-}$  cannot be directly used to impute post-treatment counterfactual outcomes  $Y_{t+}(0)$ . We also note that, although these two conditions appear to require instantaneous distributional shift, in practice one may specify a window around the treatment in which the transition may not be instantaneous, and restrict the pre- and post-treatment periods before and after the specified window, respectively, so that Conditions [3](#page-7-0) and [4](#page-7-1) are plausible.

The next condition states the existence of a treatment confounding bridge function [\[Cui et al., 2020\]](#page-60-4), which models the likelihood ratio, namely a Radon–Nikodym derivative,  $dP_{U_{t-}}$  /  $dP_{U_{t-}}$  via a regression of the supplemental proxy  $Z_{t-}$  on the unobserved confounder  $U_t$ .

<span id="page-8-1"></span>**Condition 5.** For all times  $t_{-} \leq T_0$  and  $t_{+} > T_0$ , the distribution  $P_{U_{t_{+}}}$  of the unobserved confounder  $U_{t+}$  after treatment is dominated by that before treatment, namely  $P_{U_{t-}}$ , and there exists a function  $q^* : \mathcal{Z} \to \mathbb{R}$  capturing the distributional shift in  $U_t$ ; that is,

<span id="page-8-0"></span>
$$
E\{q^*(Z_{t-}) \mid U_{t-} = u\} = \frac{\mathrm{d}P_{U_{t+}}}{\mathrm{d}P_{U_{t-}}} (u). \tag{3}
$$

The left-hand side of [\(3\)](#page-8-0) does not depend on  $t_{+}$ , due to Condition [4.](#page-7-1) We call  $q^*$ the treatment confounding bridge function, encoding information on how pre-treatment outcomes  $Y_{t-}$  can be used to impute post-treatment counterfactual outcomes  $Y_{t+}(0)$ . The function  $q^*$  is closely related to that for estimating the ATT in i.i.d settings, such as in Theorem 5.1 of [Cui et al.](#page-60-4) [\[2020\]](#page-60-4). One consequence of Condition [5](#page-8-1) is that the distribution of  $U_{t_+}$  must be dominated by that of  $U_{t_-}$ , for all  $t_-\leq T_0$  and  $t_+ > T_0$ . As shown in Theorem [1](#page-9-0) Equation [6](#page-9-1) below, under Conditions [1](#page-5-0) and [3–](#page-7-0)[5,](#page-8-1) this further implies that the distribution of  $W_{t_+}$  is dominated by that of  $W_{t_-}$  for all  $t_-\leq T_0$  and  $t_+ > T_0$ , which is a testable condition. Thus, for time series data subject to a significant secular trend, especially a monotone trend, we recommend a pre-processing step to remove, to the extent possible, any significant secular trends to make Condition [5](#page-8-1) as plausible as possible. We provide concrete approaches for removing time trends in Web Appendix [S8](#page-43-0) and the corresponding simulation results in Web Appendix [S7.6.](#page-40-0) Although detrending with our approach may lead to slightly to moderately anti-conservative inference, failing to correct for such a trend will likely compromise one's ability to implement the proposed weighted approach successfully. It is still an open question how to account for time trends appropriately to obtain asymptotically valid inference in our proposed methods. Sufficient conditions for the existence of  $q^*$  can be found in Web Appendix [S9.](#page-49-0) We list a few examples of treatment confounding bridge functions below.

<span id="page-8-2"></span>**Example 1.** Suppose that, for all  $t_{-} \leq T_0$ ,  $t_{+} > T_0$  and  $t_{+} \sim N(0, \sigma_{-}^2)$ ,  $U_{t_{+}} \sim$  $N(0, \sigma_+^2), Z_t \mid U_t \sim N(aU_t, \sigma^2)$  for some  $a \neq 0$ . If  $\sigma^{-2}a^2 - \sigma_+^{-2} + \sigma_-^{-2} > 0$ , then  $q^*$ :  $z \mapsto \exp(\alpha + \beta z^2)$  for  $\beta = \sigma^{-2}(\sigma_{-}^{-2} - \sigma_{+}^{-2}) / \{2(\sigma^{-2}a^2 - \sigma_{+}^{-2} + \sigma_{-}^{-2})\}$  and some  $\alpha \in \mathbb{R}$ satisfies Condition [5.](#page-8-1) In this example, we only specify the marginal distributions of  $(U_t, Z_t)$  for each t and allow for serial correlation. For instance,  $U_t$  could be generated from an autoregressive model that is stationary before and after treatment, respectively. This might occur if the treatment is implemented at a random time point  $T_0$  due to an abrupt change in  $U_t$  in a time window around  $T_0$ ; with the observations in this window excluded from analysis,  $U_t$  can be stationary before and after treatment with possibly different distributions. Moreover, if  $U_t$  is stationary, Condition [5](#page-8-1) holds with the constant

bridge function  $q^* : z \mapsto 1$  even if  $T_0$  is random. Similar results hold for data marginally distributed as multivariate normal at each time t.

<span id="page-9-3"></span>**Example 2.** Suppose that  $U_t = (U_{t,1}, \ldots, U_{t,K})$  has all coordinates mutually independent at any time  $t$ , while there may be serial correlations for each element over time. Let  $U_{t_-,k} \sim \text{Exponential}(\lambda_-)$  and  $U_{t_+,k} \sim \text{Exponential}(\lambda_+),$  for  $k = 1, ..., K$ , with  $\lambda_+ \geq \lambda_-$ . Suppose that  $Z_t = (Z_{t,1},..., Z_{t,K})$  with mutually independent entries distributed as, conditional on  $U_t$ ,  $Z_{t,k} \mid U_t \sim \text{Poisson}(\alpha + \beta U_{t,k})$  for some scalars  $\alpha \geq 0$  and  $\beta > 0$ . Then, direct calculation shows that  $q^* : z \mapsto \prod_{k=1}^K \exp(a + bz_k)$  satisfies Condition [5](#page-8-1) for some scalars a and b.

**Remark 2.** Condition [5](#page-8-1) may fail when  $T_0$  is random and depends on the unobserved confounder  $U_t$ , even when we condition on  $T_0$  to mimic a fixed treatment time design. Consider the following counterexample. Suppose that  $U_t, t \geq 0$  are i.i.d. across t, with support being R, and  $T_0 = \max\{t : U_t < a\}$  for an unknown fixed number a. This corresponds to the case where the treatment initiates immediately when  $U_t$  crosses the threshold a. We assume i.i.d. and unbounded  $U_t$ s only for simplicity. Conditional on  $T_0$ , since  $pr(U_{t-} < a) = 1$  but  $pr(U_{t+} \ge a) > 0$ , the Radon-Nikodym derivative  $dP_{U_{t+}} / dP_{U_{t-}}$ in Condition [5](#page-8-1) does not exist. Thus, this condition fails. Therefore, a fixed treatment time is crucial to the weighted approach to ATT, and conditioning on a random  $T_0$  might not suffice.

The above conditions lead to our first formal identification result.

<span id="page-9-0"></span>**Theorem 1** (Identification of ATT with  $q^*$ ). Let  $f : \mathbb{R} \to \mathbb{R}$  be any square-integrable function. Under Conditions [1](#page-5-0) and [3–](#page-7-0)[5,](#page-8-1) for all  $t_-\leq T_0$  and  $t_+ > T_0$ ,

<span id="page-9-4"></span>
$$
E[f\{Y_{t+}(0)\}] = E\{q^*(Z_{t-})f(Y_{t-})\}.
$$
\n(4)

In particular, taking f to be the identity function, it holds that

<span id="page-9-2"></span>
$$
E\{Y_{t+}(0)\} = E\{q^*(Z_{t-})Y_{t-}\}\tag{5}
$$

and thus the ATT is identified as  $\phi^*(t_+) = E{Y_{t_+} - q^*(Z_{t_-})Y_{t_-}}$  for all  $t_- \leq T_0$  and  $t_+ > T_0$ . In addition,  $P_{W_{t_+}}$  is dominated by  $P_{W_{t_-}}$ ; the treatment confounding bridge function  $q^*$  solves

<span id="page-9-1"></span>
$$
E\{q(Z_{t-}) \mid W_{t-} = w\} = \frac{\mathrm{d}P_{W_{t+}}}{\mathrm{d}P_{W_{t-}}} (w) \quad \text{for all } t_- \le T_0 \tag{6}
$$

in  $q : \mathcal{Z} \to \mathbb{R}$ . Further, under Condition [S2](#page-19-2) in Web Appendix [S1,](#page-19-1) [\(6\)](#page-9-1) has a unique solution.

In contrast to SC methods based on outcome modeling, the identifying expression [\(5\)](#page-9-2) cannot be directly interpreted as an outcome trajectory. Indeed, given a treatment confounding bridge function  $q^*$ , the right-hand side of  $(5)$  only depends on observations before treatment but not after treatment. We also obtain a novel doubly robust identification result, which is motivated by doubly robust estimation of the average treatment effect on the treated [\[Cui et al., 2020\]](#page-60-4) and is the basis of doubly robust inference.

<span id="page-10-0"></span>**Theorem 2** (Doubly robust identification). Let  $h : \mathcal{W} \to \mathbb{R}$  and  $q : \mathcal{Z} \to \mathbb{R}$  be square-integrable. Under Conditions [1,](#page-5-0) [3](#page-7-0) and [4,](#page-7-1) for all  $t_− ≤ T_0$  and  $t_+ > T_0$ , we have

<span id="page-10-2"></span>
$$
\phi^*(t_+) = E\left[Y_{t_+} - q(Z_{t_-})\{Y_{t_-} - h(W_{t_-})\} - h(W_{t_+})\right],\tag{7}
$$

if (i) Condition [2](#page-5-2) holds and  $h = h^*$ , or (ii) Condition [5](#page-8-1) holds and  $q = q^*$ .

Theorem [2](#page-10-0) states that the ATT  $\phi^*(t_+)$  is identified by a single formula if at least one of the two nuisance functions  $h^*$  or  $q^*$  is known, and thus doubly robust estimation [\[Bang](#page-59-7) [and Robins, 2005,](#page-59-7) [Robins and Rotnitzky, 1995,](#page-63-6) [Robins et al., 1995,](#page-63-7) [Scharfstein et al.,](#page-63-1) [1999\]](#page-63-1) is possible. This result holds even if one of  $h^*$  and  $q^*$  exists but the other does not; that is, Conditions [2](#page-5-2) and [5](#page-8-1) need not hold simultaneously.

### <span id="page-10-1"></span>5 Doubly robust inference about ATT

We assume that one specifies parametric models for the confounding bridge functions h<sup>\*</sup> and q<sup>\*</sup>. We use  $h_{\alpha}$  and  $q_{\beta}$  to denote the models for confounding bridge functions parameterized by  $\alpha \in \mathcal{A} \subseteq \mathbb{R}^{d_{\alpha}}$  and  $\beta \in \mathcal{B} \subseteq \mathbb{R}^{d_{\beta}}$ , respectively. For example, in classical SC methods, it is typically assumed that  $h^*(w) = w^\top \alpha_0$  for a vector  $\alpha_0$  of non-negative numbers that sum up to unity [\[Abadie et al., 2010\]](#page-58-1). In this case, we may take  $h_{\alpha}$  to be  $w \mapsto w^{\top} \alpha$ . In some cases, e.g., in Example [2,](#page-9-3) we may take  $q_{\beta}(z) = \exp(\beta_0 + z^{\top} \beta_1)$  where  $\beta = (\beta_0, \beta_1^{\top})^{\top}.$ 

We assume that the function  $t_+ \mapsto \phi^*(t_+)$  encoding the potentially time-varying ATT is correctly parameterized by  $\lambda \in \Lambda \subseteq \mathbb{R}^{d_{\lambda}}$ . We use  $\phi_{\lambda}$  to denote this model. For example, the ATT is commonly assumed to be constant overtime, which holds under stationarity of  $\{Y_{t+}\}_{t+>T_0}$  and Conditions [3–](#page-7-0)[4.](#page-7-1) In this case, we may set  $\phi_{\lambda}$  to be constant  $\lambda \in \mathbb{R}$ .

By [\(2\)](#page-6-0) and [\(6\)](#page-9-1) respectively, we have that

<span id="page-10-3"></span>
$$
E[{Y_{t-} - h^*(W_{t-})}g_Z(Z_{t-})] = 0 \text{ and } E{q^*(Z_{t-})g_W(W_{t-}) - g_W(W_{t+})} = 0 \quad (8)
$$

for any functions  $q_Z : \mathcal{Z} \to \mathbb{R}$  and  $q_W : \mathcal{W} \to \mathbb{R}$ . We propose to use the generalized method of moments (GMM) to estimate  $h^*$ ,  $q^*$ , and the ATT [e.g., [Hansen,](#page-61-6) [1982,](#page-61-6) [Wooldridge, 1994,](#page-64-3) Hall, 2007. This method involves a parameter vector  $\theta =$  $(\alpha, \beta, \lambda, \psi, \psi) \in \Theta$  including nuisance parameters  $\psi$  and  $\psi$ , a moment equation  $G_t$ for each time t, and a weight matrix  $\Omega_T$  that may depend on sample size T. Details of this method are presented in Web Appendix [S2.](#page-20-0) We need some additional conditions to obtain valid inferences about  $\phi^*$ .

<span id="page-11-0"></span>**Condition 6.** (i) There is a unique parameter value  $\theta_{\infty} = (\alpha_{\infty}, \beta_{\infty}, \lambda_{\infty}, \psi_{\infty}, \psi_{-\infty}) \in \Theta$ such that  $E\{G_t(\theta_\infty)\}=0$  for all time points t. (ii) The function  $h^* = h_{\alpha_\infty}$  is a valid outcome confounding bridge function satisfying Condition [2,](#page-5-2) or the function  $q^* = q_{\beta_{\infty}}$  is a valid treatment confounding bridge function satisfying Condition [5.](#page-8-1)

Part (i) of Condition [6](#page-11-0) is an identifying condition to ensure a unique solution to the population moment equation, which is standard for GMM. If both  $h^*$  and  $q^*$  are correctly specified, part (i) would hold if they are both unique. Part (ii) requires correct parametric specification of at least one of, but not necessarily both of,  $h^*$  or  $q^*$ , without necessarily knowing a priori which model might be incorrect. Under Condition [6,](#page-11-0) Theorem [2](#page-10-0) implies that  $\phi_{\lambda_{\infty}}$  equals the target estimand, ATT  $\phi^*$ . By standard asymptotic theory for GMM [e.g., Theorems 7.1 and 7.2 in [Wooldridge, 1994\]](#page-64-3) along with Theorem [2,](#page-10-0) we have that  $\theta_T$ is consistent for  $\theta_{\infty}$  and asymptotically normal under conditions, as stated in Theorem [3](#page-11-1) below. If the data are i.i.d., as we argued in Section [4,](#page-6-1) the estimand reduces to the average treatment effect on the treated and our estimator is locally efficient [\[Cui et al.,](#page-60-4) [2020\]](#page-60-4).

<span id="page-11-1"></span>Theorem 3. Under Conditions [1,](#page-5-0) [3,](#page-7-0) [4,](#page-7-1) [6,](#page-11-0) as well as [S3–](#page-20-1)[S5](#page-21-0) and [S7](#page-21-1) in Web Appendix [S2,](#page-20-0) with the estimator  $\theta_T$  from [\(S1\)](#page-20-2) and  $\theta_{\infty}$  in Condition [6,](#page-11-0) it holds that, as  $T \to \infty$ ,  $\hat{\theta}_T$  is consistent for  $\theta_{\infty}$ . Additionally, under Conditions [S6,](#page-21-2) [S8](#page-22-0) and [S9,](#page-22-1) as  $T \to \infty$ ,  $\overline{T}(\hat{\theta}_T - \theta_\infty) \stackrel{d}{\rightarrow} \text{N}(0, A^{-1}BA^{-1}),$  where  $A = R^{\top} \Omega R$ ,  $\Omega$  is from Condition [S3,](#page-20-1)

$$
R = \lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^{T} E\{\nabla_{\theta} G_t(\theta) |_{\theta = \theta_{\infty}}\}, \ B = R^{\top} \Omega \left[ \lim_{T \to \infty} \text{var}\left\{ T^{-1/2} \sum_{t=1}^{T} G_t(\theta_{\infty}) \right\} \right] \Omega R.
$$

Suppose that the ATT function  $\phi_{\lambda}$  is differentiable with respect to  $\lambda$  and let  $\dot{\phi}_{\lambda}$  denote this partial derivative. Thus, with  $\Pi := (0_{d_{\lambda} \times (d_{\alpha} + d_{\beta})}, I_{d_{\lambda}}, 0_{d_{\lambda} \times (d_{\beta'} + 1}))$  being the matrix consisting of zeros and ones with dimensions denoted in the subscript, for every  $t_+ > T_0$ , it holds that  $\phi_{\lambda_{\infty}}$  is the ATT at time  $t_+$  and  $\sqrt{T}(\phi_{\hat{\lambda}_T}(t_+) - \phi_{\lambda_{\infty}}(t_+)) \stackrel{d}{\rightarrow}$  $N(0, \dot{\phi}_{\lambda_{\infty}}(t_{+})^{\top} \Pi^{\top} A^{-1} B A^{-1} \Pi \dot{\phi}_{\lambda_{\infty}}(t_{+})).$ 

One can in principle use any GMM implementation, and we use the standard R package gmm for our simulation and data analyses. Confidence intervals of the ATT follows from standard outputs of gmm. In particular, when  $\phi_{\lambda}$  is a constant function, since  $\lambda_{\infty}$  is a component of  $\theta_{\infty}$ , a Wald test or confidence interval about the ATT  $\phi_{\lambda_{\infty}}(t_{+}) = \lambda_{\infty}$  follows immediately from inference about  $\theta_{\infty}$ . We have noted some numerical instabilities with this implementation in our simulations, and provide our empirical suggestions to alleviate numerical issues in Web Appendix [S6.](#page-33-0) We expect these issues to be alleviated by using an improved GMM software implementation with, for example, more numerically stable

optimization algorithms that are more capable to handle nonconvex problems, potentially under constraints.

Our asymptotic results in Theorem [3](#page-11-1) rely on the total number  $T$  of time periods tending to infinity. Thus, our proposed doubly robust method is applicable to cases with a large number of time periods and a relatively small number of model parameters. With many control units, expertise might be required to reduce the number of parameters before analysis using our method. Methods with different theoretical results that allow for a large number of control units include [Athey et al.](#page-59-4) [\[2021\]](#page-59-4), [Ben-Michael et al.](#page-59-3) [\[2021a,](#page-59-3)[b\]](#page-59-0), among others.

The method and results corresponding to the identification results in Theorem [1](#page-9-0) based on weighting alone are similar and can be found in Web Appendix [S4.](#page-30-0) This is largely based on the outcome modeling-based approach developed by [Shi et al.](#page-63-0) [\[2023\]](#page-63-0). Compared to these two methods, the doubly robust method has the advantage that it only requires correct specification of one of  $h^*$  and  $q^*$  in a parametric model, but not necessarily both. One potential drawback of the doubly robust method, however, is that more parameters need to be estimated in GMMs compared to the other two methods. This issue of dimensionality might limit the usage of complicated models for  $h^*, q^*$  and  $\phi^*$  when the time series is short. In particular, if the number of parameters is comparable to the total number  $T$  of time periods, the doubly robust method might lead to numerical instability due to too many parameters being estimated and might be impractical. Our methods have no guarantee in such scenarios. However, the number of parameters is much smaller than  $T$  in many applications, for example, in our data analyses in Section [7](#page-13-0) and Web Appendix [S8.](#page-43-0) The number of control units is often highly related to the number of parameters, and the number of control units is commonly small compared to the number of time periods in SC applications, including the motivating ones [e.g., [Abadie and Gardeazabal, 2003,](#page-58-0) [Abadie et al., 2010\]](#page-58-1). Models with many variables might possibly be used by utilizing methods in [Deaner](#page-60-3) [\[2021\]](#page-60-3) under linearity and sparsity. [Chamberlain](#page-60-9) [\[1992\]](#page-60-9), [Hansen](#page-61-6) [\[1982,](#page-61-6) [1985\]](#page-61-8) showed that GMM attains the efficiency bound under conditional moment restrictions with a suitably chosen weighting matrix  $\Omega_T$  in i.i.d settings as well as panel data settings where asymptotics are in the number of units. We do not pursue high-dimensional models or efficiency further in this work and restrict ourselves to the setting of a bounded number of units (a single treated unit and a finite number of control units) while relying on large T asymptotics; to the best of our knowledge, none of these prior results can directly apply to our more challenging setting and existing semiparametric efficiency theory does not appear to directly apply without additional restrictions (e.g., Markov restrictions).

# <span id="page-13-1"></span>6 Simulations

We investigate the performance of our methods to estimate the constant ATT  $\phi^* \equiv \phi^*(t_+)$ in several simulations. Here we present the first simulation where the moment equations in the GMM are just identified. We compare the following methods: correct.DR, the doubly robust method with correctly specified parametric  $h^*$  and  $q^*$ ; correct.h, the outcome confounding bridge method from [Shi et al.](#page-63-0) [\[2023\]](#page-63-0) with correctly specified  $h^*$ ; correct.q, the treatment confounding bridge method described in Web Appendix [S4](#page-30-0) with correctly specified  $q^*$ ; mis.h.DR, the doubly robust method with misspecified  $h^*$  and correctly specified  $q^*$ ; mis.q.DR, the doubly robust method with misspecified  $q^*$  and correctly specified  $h^*$ ; mis.h, the outcome confounding bridge method with misspecified  $h^*$ ; mis.q, the treatment confounding bridge method with misspecified  $q^*$ ; OLS, the method based on unconstrained ordinary least squares and similar to the method from [Abadie et al.](#page-58-1) [\[2010\]](#page-58-1). OLS finds the linear combination of donors' outcomes that best fits the treated unit's outcome before treatment, and uses this combination as the SC. All methods except OLS are based on the proximal causal inference perspective. We let the number of latent confounders, the number of donors and of the other control units, all equal to K, range in {2,3,4,5}. Details of the data-generating mechanism are presented in Wed Appendix [S7.1.](#page-36-0)

We next present the simulation results. We have run 16,000 GMM involving weighting via a treatment confounding bridge function. Among them, only one run had numerical errors. The sampling distributions of the estimated ATT is presented in Figure [2.](#page-14-0) In all settings, the OLS estimator is biased. When at least one of  $h^*$  or  $q^*$  is correctly specified, our proposed doubly robust method appears consistent and asymptotically normal, aligning with Theorem [3.](#page-11-1) The other two methods, based on only one of  $h^*, q^*,$ are not doubly robust. They are consistent and asymptotically normal only when the bridge function they rely on is correctly specified, but are biased otherwise. The 95%- Wald confidence interval coverage of all above methods is presented in Figure [3.](#page-15-0) In large samples, confidence intervals based on consistent and asymptotically normal estimators have coverage close to the nominal level; otherwise, the confidence interval coverage is much lower than the nominal level.

# <span id="page-13-0"></span>7 Analysis of Brazil all-cause pneumonia hospitalization

We study the effect of the introduction of the PCV10 vaccine in Brazil on the number of hospitalizations due to all-cause pneumonia [\[Bruhn et al., 2017\]](#page-59-2). Monthly hospitalizations and their causes were collected from 2003 to 2013. We focus on the subpopulation of children less than 12 months old in this analysis. PCV10 was introduced to Brazil in January 2010. Following the analysis in [Bradley and Tone](#page-59-8) [\[2017\]](#page-59-8), we allow two years for the introduction of the vaccine to take effect and set the evaluation period to be

<span id="page-14-0"></span>

Figure 2: Sampling distribution of estimated average treatment effect. The horizontal dotted line is the true average treatment effect for the treated unit.

2012–2013.

We view each group of causes of hospitalization as a unit and dismiss units with missing data. The time series data does not have a clear monotone trend; thus, Condition [5](#page-8-1) may be plausible. To alleviate numerical issues due to non-linearity in GMM and to reduce differences in scaling between units, we scale the numbers of hospitalizations due to each group of causes to the unit interval before analysis (see Web Appendix [S6](#page-33-0) for more details). We model the outcome bridge function as a linear function  $h_{\alpha}: w \mapsto (1, w^{\top})^{\top} \alpha$ , and select the number of hospitalizations due to the following three groups of causes (units) as the proxies  $W_t$ : (i) bronchitis, bronchiolitis and unspecified acute lower respiratory infection, (ii) endocrine, nutritional, metabolic disorders, and (iii) malnutrition. This choice is motivated by [Bradley and Tone](#page-59-8) [\[2017\]](#page-59-8) and the prior knowledge about their relation to pneumonia.

We use all other groups of causes as control units in the supplemental proxies  $Z_t$ . We consider three parametrizations of the treatment confounding bridge function with increasing sets of units included in the model:  $q_\beta(Z_t) = \exp(\beta_0 + \sum_{j=1}^J \beta_j Z_j)$  for  $J =$ 1, 2, 3, where  $Z_1$ ,  $Z_2$  and  $Z_3$ , respectively, are the numbers of hospitalizations due to (i) certain infectious and parasitic diseases, except intestinal, (ii) diseases of blood and blood-forming organs and certain disorders involving the immune mechanism, and (iii) premature delivery and low birth weight. The motivation for choosing these supplemental proxies  $Z_j$  is to capture the effect of unmeasured confounders  $U_t$  on general health issues

<span id="page-15-0"></span>

Figure 3: 95%-Wald confidence interval coverage of the average treatment effect for the treated unit for all methods in the simulation under the just-identified setting. The error bars represent 95%-Wilson score intervals for the confidence interval coverage. The horizontal dotted line is the nominal coverage 95%. We set the confidence interval to be the empty set when numerical errors occur to obtain a conservative Monte Carlo estimate of confidence interval coverage.

related to infections and the immune system, which are both associated with the outcome of interest, all-cause pneumonia hospitalizations, among children less than 12 months old. We refer to the corresponding doubly robust (resp., weighted) estimators as DR, DR2 and DR3 (resp., treatment bridge, treatment bridge2 and treatment bridge3). As shown in the simulation in Web Appendix [S7.7,](#page-41-0) choosing  $W$  and  $Z$  completely based on the data might lead to estimates with increased uncertainty due to the possibility of model misspecification. Such estimates may not be informative. When selecting units empirically, we recommend, when possible, using prior knowledge and not solely relying on the data, in which case this issue might be avoided. Ideally, proxies should be selected a priori to avoid potential post-selection inference issues.

The GMM estimator outlined in Section [5](#page-10-1) is implemented with the user-specified functions being  $g_h : z \mapsto (1, z^{\top})^{\top}$ , where we recall that z is the collection of all supplemental proxies, namely hospitalizations due to non-donor causes, and  $g_q : w \mapsto (1, w^{\top})^{\top}$ . We set the weight matrix to equal the identity matrix.

Besides proximal SC estimators, we also report results for the standard OLS estimator described in Section [6.](#page-13-1) We also consider the regression-based SC method (Abadie's SC)

<span id="page-16-0"></span>

Method	PCV <sub>10</sub>	placebo
Abadie's SC	409	3092
OLS	$-3533$ $(-4137, -2930)$	253 (-287, 794)
DR.	$-2745$ $(-3559, -1931)$	1192 (501, 1884)
DR2	$-3527$ $(-4663, -2392)$	$317$ (-407, 1042)
DR <sub>3</sub>	$-3548$ $(-6036, -1061)$	$260$ (-246, 767)
Outcome bridge	$-3646$ $(-4693, -2598)$	$565$ $(-224, 1355)$
Treatment bridge	$-3989$ $(-4373, -3605)$	$-532$ $(-1638, 574)$
Treatment bridge2	$-3814$ $(-4941, -2688)$	$-205$ $(-1542, 1133)$
Treatment bridge3	$-3895$ $(-6401, -1388)$	$97$ ( $-502, 695$ )

Table 1: Estimate of the average treatment effect of PCV10 and placebo treatment on the number of hospitalizations due to all-cause pneumonia among children less than 12 months old in Brazil for various methods with 95%-Wald confidence intervals. Abadie's SC [\[Abadie et al., 2010\]](#page-58-1) does not readily provide confidence intervals.

proposed by [Abadie et al.](#page-58-1) [\[2010\]](#page-58-1) with the same three donors forming the proxy  $W_t$  as above. The point estimates and 95% confidence intervals for the ATT are presented in Table [1.](#page-16-0) The trajectories of SCs and the actual number of hospitalizations are presented in Figure [4a.](#page-17-0)

Although we have scaled all outcomes to fall in the unit interval, Abadie's SC does not output a good pre-treatment fit and its ATT estimate appears unreliable. Because the original scales of the outcomes across units differ substantially, the constraints in Abadie's SC (i) that the intercept vanishes, and (ii) that the weights are non-negative and sum to one, might not be appropriate in this application [\[Doudchenko and Imbens,](#page-60-0) [2016\]](#page-60-0); scaling to unit interval might still fail to justify the adequacy of these constraints. All other methods conclude a significant decrease in hospitalizations due to all-cause pneumonia after the introduction of PCV10, as expected. The estimate from DR is somewhat different from other proximal methods involving a treatment confounding bridge function with at least two units, suggesting a model misspecification affecting DR in finite samples. Though the theory suggests that DR should be consistent, when at least one nuisance function is correctly specified, a larger sample size  $T$  than this data set might be needed for DR to be close to the truth.

We also conduct a falsification analysis. We consider a hypothetical placebo treatment in January 2009 and estimate its effect in the year 2009. The analysis results are presented in Table [1](#page-16-0) and Figure [4b.](#page-17-0) Similarly to the main analysis, the pre-treatment fit from Abadie's SC appears unreliable in this setting; therefore confirming the recommendation that SCs might perform poorly when the pre-treatment fit is suboptimal [\[Abadie](#page-58-1) [et al., 2010\]](#page-58-1). Our proposed methods perform better and offer alternative justification for SC methods in such settings. In fact, 95%-confidence intervals from most proximal SC

<span id="page-17-0"></span>

Figure 4: Trajectories of synthetic controls (green dashed) and the number of hospitalizations due to all-cause pneumonia (red solid) among children less than 12 months old in Brazil for various methods. The vertical line is the last time point (month) before implementing the treatment. Methods based on weighting only (treatment bridge, treatment bridge2 and treatment bridge3) do not have synthetic control trajectories.

methods cover zero, correctly indicating a non-significant effect due to the placebo treatment; so does OLS in this case. The only exception DR echoes the poor performance from the main analysis. We conclude that, in this application, when the nuisance functions are approximately correctly specified, the proximal methods are among the best.

We also study the effect of Florida's "stand your ground" law on homicide rates and the effect of a tax cut in Kansas on economic outcomes (see Web Appendix [S8\)](#page-43-0). In these analyses, our proposed doubly robust method outperforms Abadie's SC or OLS in several cases.

### 8 Discussion

To estimate the ATT in panel data settings, classical SC methods often require correct specification of nuisance functions. Our proposed doubly robust methods involve estimating two nuisance functions but allow for misspecification of one, without knowledge about which is misspecified. Our identification results may enable the development of new SC methods, especially nonparametric or semiparametric ones that rely on weaker conditions. Such methods can extend the idea of SCs and proximal causal inference to more general applications.

# Acknowledgement

This work was supported in part by the U.S. National Science Foundation and National Institutes of Health, and by Analytics at Wharton. We are grateful to an anonymous reviewer for a counterexample of Condition [5.](#page-8-1)

# Supplementary Materials for "Doubly Robust Proximal Synthetic Controls" by Hongxiang Qiu, Xu Shi, Wang Miao, Edgar Dobriban, and Eric Tchetgen Tchetgen

### <span id="page-19-1"></span>S1 Completeness conditions

The following completeness conditions are sufficient to ensure the uniqueness of the so-lutions to the equations [\(2\)](#page-6-0) and [\(6\)](#page-9-1) used to estimate confounding bridge functions  $h^*$ and  $q^*$ . We emphasize that these conditions are not necessary for valid estimation of the average treatment effect for the treated unit, as shown in the simulation in Section [S7.2.](#page-37-0)

<span id="page-19-0"></span>**Condition S1.** Let  $g : \mathcal{W} \to \mathbb{R}$  be any square-integrable function. For all time points  $t_-\leq T_0$  before treatment, the following two statements are equivalent: (i)  $g(W_{t-})=0$ , (ii)  $E\{g(W_{t-}) \mid Z_{t-}\}=0.$ 

<span id="page-19-2"></span>**Condition S2.** Let  $g : \mathcal{Z} \to \mathbb{R}$  be any square-integrable function. For all time points  $t_-\leq T_0$  before treatment, the following two statements are equivalent: (i)  $g(Z_{t-})=0$ , (ii)  $E\{g(Z_{t-}) \mid W_{t-}\}=0.$ 

We next discuss Condition [S1](#page-19-0) in more detail; Condition [S2](#page-19-2) is similar. Intuitively, Condition [S1](#page-19-0) requires that  $Z_t$  captures all variation in  $W_t$ , because any transformation  $g(W_t)$  of  $W_t$  whose regression on  $Z_t$  is zero must be zero. Equivalent conditions to Condition [S1](#page-19-0) have been found. Lemma 2.1 in [Severini and Tripathi](#page-63-8) [\[2006\]](#page-63-8) and Proposition 1 in [Andrews](#page-58-5) [\[2017\]](#page-58-5) imply that Condition [S1](#page-19-0) is equivalent to the following: for every nonconstant random variable  $a(W_{t-})$ , there exists a random variable  $b(Z_{t-})$  that is correlated with  $a(W_t)$ . According to this equivalent condition, it is evident that the completeness condition [S1](#page-19-0) can be interpreted as that any variation in  $W_{t-}$  can be captured by a variation in  $Z_{t-}$ . We refer readers to Supplemental Section D in [Ying et al.](#page-64-4) [\[2021\]](#page-64-4) for more details on completeness conditions.

[Shi et al.](#page-63-0) [\[2023\]](#page-63-0) recommended measuring a rich set of proxies  $Z_t$  to make this condition plausible. Similarly, to make Condition [S2](#page-19-2) plausible, we recommend measuring a rich set of proxies  $W_t$ . In the common setting where both  $W_t$  and  $Z_t$  are control units' outcomes, we thus recommend measuring contemporary outcomes in many control units and using many of them as proxies, either in  $W_t$  or  $Z_t$ , to make Conditions [S1](#page-19-0) and [S2](#page-19-2) plausible.

# <span id="page-20-0"></span>S2 Technical details and formal results for generalized method of moments

We first describe our estimation method based on GMM. Let  $g_h : \mathcal{Z} \to \mathbb{R}^{d'_\alpha}$ , with  $d'_\alpha \geq d_\alpha$ , and  $g_q: \mathcal{W} \to \mathbb{R}^{d'_{\beta}}$ , with  $d'_{\beta} \geq d_{\beta}$ , be two user-specified functions. In practice, these functions can be chosen from known classes of basis functions used in non-parametric estimation. For instance, they can be a truncated polynomial basis [\[Hastie et al., 2009\]](#page-61-9): with the support  $\mathcal Z$  of  $Z_t$  being  $\mathbb R^{dz}$ ,  $g_h(z)$  may consist of monomials up to order K, namely  $\left(\prod_{k=1}^K z_{i_k}^{j_k}\right)$  $i_k^{j_k}$ ) with  $i_k \in \{1, ..., d_Z\}$ ,  $j_k$  being non-negative integers, and  $\sum_{k=1}^{\alpha} j_k \leq K$ , such that the dimension of the output of  $g_h$  is at least  $\alpha$ . Let  $\alpha \in \mathcal{A}, \beta \in \mathcal{B}, \lambda \in \Lambda$ ,  $\psi \in \Psi \subseteq \mathbb{R}^{d_{\beta'}}$ , and  $\psi_- \in \Psi_- \subseteq \mathbb{R}$  be GMM parameters,  $\theta = (\alpha, \beta, \lambda, \psi, \psi_-)$  be the collection of parameters, and  $\Theta = \mathcal{A} \times \mathcal{B} \times \Lambda \times \Psi \times \Psi$  be the parameter space. Define the moment equation  $G_t : \Theta \to \mathbb{R}^{2d'_{\beta}+3}$  as the vertical stacking of

 $G_{t1} : \theta \mapsto \mathbb{1}(t \leq T_0) \left[ \{ Y_t - h_\alpha(W_t) \} g_h(Z_t) \right], \quad G_{t2} : \theta \mapsto \mathbb{1}(t > T_0) \left\{ \psi - g_\alpha(W_t) \right\}$  $G_{t3} : \theta \mapsto \mathbb{1}(t \leq T_0) \{q_\beta(Z_t)g_q(W_t) - \psi\}, \quad G_{t4} : \theta \mapsto \mathbb{1}(t > T_0) \left[\phi_\lambda(t) - \{Y_t - h_\alpha(W_t)\} + \psi_-\right]$ and  $G_{t5} : \theta \mapsto \mathbb{1}(t \leq T_0) [\psi_- - q_\beta(Z_t) \{Y_t - h_\alpha(W_t)\}]$ .

Let  $\Omega_T$  be a user-specified symmetric positive semi-definite  $(d'_\alpha + 2d'_\beta + 2) \times (d'_\alpha + 2d'_\beta + 2)$ matrix, for example, the identity. Consider the GMM estimator

<span id="page-20-2"></span>
$$
\hat{\theta}_T = \left(\hat{\alpha}_T, \hat{\beta}_T, \hat{\lambda}_T, \hat{\psi}_T, , \hat{\psi}_{-,T}\right) = \underset{\theta \in \Theta}{\text{argmin}} \left\{ \frac{1}{T} \sum_{t=1}^T G_t(\theta) \right\}^\top \Omega_T \left\{ \frac{1}{T} \sum_{t=1}^T G_t(\theta) \right\}. \tag{S1}
$$

We use  $\phi_{\hat{\lambda}_T}$  as the doubly robust estimator of the ATT  $\phi^*.$ 

Remark 3. Since [\(7\)](#page-10-2) and [\(8\)](#page-10-3) involve expectations over different time periods, directly using these equations for estimation leads to moment equations that might not have mean zero for each time period  $t$ , but only when averaged over all time periods. This may invalidate the corresponding GMM procedure. To circumvent this, we introduce two centering parameters  $\psi$  and  $\psi$  to decouple different time periods in [\(7\)](#page-10-2) and [\(8\)](#page-10-3). The second moment equation in [\(8\)](#page-10-3) is decoupled via the introduction of  $\psi$  and represented by the second and third elements in  $G_t$ . Further, [\(7\)](#page-10-2) is decoupled via the introduction of  $\psi$ <sub>-</sub> and split into the fourth and fifth elements in  $G_t$ .

We next present the additional technical conditions we rely on for the consistency and asymptotic normality results for the generalized method of moments estimators. We put conditions and their explanation under stationarity of confounders (Condition [4\)](#page-7-1) in the text and those without stationarity in square brackets.

<span id="page-20-1"></span>**Condition S3.** For the doubly robust method,  $\Omega_T$  [ $\mathcal{Q}_T$ ] converges in probability to  $\Omega$  [ $\mathcal{Q}$ ] for some fixed symmetric positive-definite matrix  $\Omega$  [Q]; for the weighting method,  $\Omega_q^q$ 

 $[\mathcal{Q}_7^q$  $\Omega^{q}$  [ $\mathcal{Q}^{q}$ ].  $T_T^q$  converges in probability to  $\Omega^q$  [ $\Omega^q$ ] for some fixed symmetric positive-definite matrix

Condition [S3](#page-20-1) can be easily satisfied by taking  $\Omega_T$  or  $\Omega_T^q$  [ $\Omega_T$  or  $\Omega_T^q$  $T$  as a fixed positivedefinite matrix.

<span id="page-21-3"></span>Condition S4. For the doubly robust method,  $\Theta$   $[Q]$  is compact; for the weighting method,  $\Theta^q$  [ $\Theta^q$ ] is compact.

Condition [S4](#page-21-3) is a regularity compactness condition to establish theoretical results. In computations, the parameter space is often taken to be unrestricted.

<span id="page-21-0"></span>**Condition S5.** For the doubly robust method,  $E\{G_t(\theta)\}\left[\lim_{T\to\infty}\sum_{t=1}^T E\{G_t(\theta)\}\right]$ e  $_{T,t}(\theta$  $\alpha$  $)\}/T]$ exists and is finite for all  $\theta \in \Theta$  [ $\underline{\theta}$  $\overline{f}$ ∈ Θ  $\vec{r}_t$ ];  $G_t$  [ $\theta$  $\lfloor$  $\mapsto \lim_{T \to \infty} \sum_{t=1}^{T} E\{Q_t\}$  $\left(\frac{t}{2}\right)$  $_{T,t}(\theta$  $\bigg/$  $\langle$  \equal uous; for the weighting method,  $E\{G_t^q\}$  $_{t}^{q}(\theta^{q})\}$  [lim<sub>T→∞</sub>  $\sum_{t=1}^{T} E\{Q_t\}$  $\overline{\varrho}$  ${}_{T,t}^q(\theta$  $\sqrt{2}$  $\binom{q}{\ell}$   $\binom{T}{\ell}$  exists and is finite for all  $\theta^q \in \Theta^q$  [ $\underline{\theta}^q \in \Theta^q$ ];  $G_t^q$  $t^q \left[ \theta^q \mapsto \lim_{T \to \infty} \sum_{t=1}^T E \{ \mathcal{G}_{T,t}^q(\theta^q) \} / T \right]$  is continuous.

Condition [S5](#page-21-0) is also a regularity condition, which ensures that the population expectation of the moment equation exists.

<span id="page-21-2"></span>**Condition S6.** For the doubly robust method, rank $(R) = d_{\alpha} + d_{\beta} + 1$  [rank $(R) =$  $d_{\alpha} + d_{\beta} + 1$ ; for the weighting method,  $rank(R_q) = d_{\beta} + 1$  [rank $(R_q) = d_{\beta} + 1$ ].

Condition [S6](#page-21-2) is a full-rank condition to ensure that the parameter in the is locally identified.

We next list uniform weak laws of large numbers and central limit theorems as Conditions [S7,](#page-21-1) [S8](#page-22-0) and [S9.](#page-22-1) In turn, these hold under various sets of assumptions.

<span id="page-21-1"></span>**Condition S7.** For the doubly robust method, with  $\bar{V}_{T}(\theta) = \sum_{t=1}^{T'} E\{G_{T',t}(\theta)\}/T',$ 

$$
\sup_{\theta \in \Theta} \left\| \frac{1}{T} \sum_{t=1}^T G_t(\theta) - \lim_{T' \to \infty} \frac{1}{T'} \sum_{t=1}^{T'} E\{G_t(\theta)\} \right\|, \qquad \left[ \sup_{\underline{\theta} \in \underline{\Theta}} \left\| \frac{1}{T} \sum_{t=1}^T G_t(\theta) - \lim_{T' \to \infty} \bar{V}_{T'}(\underline{\theta}) \right\| \right]
$$

converge to zero in probability. For the weighting method,

$$
\sup_{\theta^q \in \Theta^q} \left\| \frac{1}{T} \sum_{t=1}^T G_t^q(\theta^q) - \lim_{T' \to \infty} \frac{1}{T'} \sum_{t=1}^{T'} E\{G_t^q(\theta^q)\} \right\|,
$$
  

$$
\left[ \sup_{\theta^q \in \Theta^q} \left\| \frac{1}{T} \sum_{t=1}^T \mathcal{G}_{T,t}^q(\theta^q) - \lim_{T' \to \infty} \frac{1}{T'} \sum_{t=1}^{T'} E\{G_{T',t}^q(\theta^q)\} \right\| \right]
$$

e e converge to zero in probability.

<span id="page-22-0"></span>**Condition S8.** For the doubly robust method,  $\theta_{\infty}$  [ $\theta_{\infty}$ ] is in the interior of  $\Theta$  [ $\Theta$ ],  $G_t$ is continuously differentiable  $[G_{T,t}]$  is continuously differentiable and the derivative of  $\sum_{t=1}^{T} G_{T,t}/T$  is uniformly bounded over all  $T \in \{1, 2, \ldots\}$ , and

$$
\sup_{\theta \in \Theta} \left\| \frac{1}{T} \sum_{t=1}^T \nabla_{\theta} G_t(\theta) - \lim_{T' \to \infty} \frac{1}{T'} \sum_{t=1}^{T'} E \left\{ \nabla_{\theta} G_t(\theta) \right\} \right\|
$$
  

$$
\left[ \sup_{\underline{\theta} \in \Theta} \left\| \frac{1}{T} \sum_{t=1}^T \nabla_{\underline{\theta}} G_{T,t}(\underline{\theta}) - \lim_{T' \to \infty} \frac{1}{T'} \sum_{t=1}^{T'} E \left\{ \nabla_{\underline{\theta}} G_{T',t}(\underline{\theta}) \right\} \right\| \right]
$$

converges to zero in probability; for the weighting method,  $\theta_{\infty}^{q}$  [ $\theta_{\infty}^{q}$ ] lies in the interior of  $\Theta^q$  is continuously differentiable  $[\mathcal{G}^q_{T,t}]$  is continuously differentiable and the derivative of  $\sum_{t=1}^{T} \mathcal{G}_{T,t}^{q}/T$  is uniformly bounded over all  $T \in \{1, 2, \ldots\}$ , and

$$
\sup_{\theta^q \in \Theta^q} \left\| \frac{1}{T} \sum_{t=1}^T \nabla_{\theta^q} G_t^q(\theta^q) - \lim_{T' \to \infty} \frac{1}{T'} \sum_{t=1}^{T'} E \left\{ \nabla_{\theta^q} G_t^q(\theta^q) \right\} \right\|
$$
  

$$
\left[ \sup_{\theta^q \in \Theta^q} \left\| \frac{1}{T} \sum_{t=1}^T \nabla_{\theta^q} G_{T,t}^q(\theta^q) - \lim_{T' \to \infty} \frac{1}{T'} \sum_{t=1}^{T'} E \left\{ \nabla_{\theta^q} G_{T',t}^q(\theta^q) \right\} \right\| \right].
$$

converges to zero in probability.

<span id="page-22-1"></span>Condition S9. For the doubly robust method,

$$
T^{-1/2} \sum_{t=1}^{T} R^{\top} \Omega G_t(\theta_{\infty}) \qquad \left[ T^{-1/2} \sum_{t=1}^{T} R^{\top} \Omega G_{T,t}(\theta_{\infty}) \right]
$$

is asymptotically  $N(0, B)$  [N $(0, B)$ ]; for the weighting method,

$$
T^{-1/2} \sum_{t=1}^T R_q^{\top} \Omega^q G_t^q(\theta_\infty^q) \qquad \left[ T^{-1/2} \sum_{t=1}^T R_q^{\top} \Omega^q G_{T,t}^q(\theta_\infty^q) \right]
$$

is asymptotically  $N(0, B_q)$  [N $(0, \underline{B}_q)$ ].

In Condition [S8,](#page-22-0) the assumption that the probability limit of the estimator lies in the interior of the parameter space and the differentiability of the moment equation are usually satisfied under stationarity (Condition [4\)](#page-7-1). Without stationarity, we require a slightly stronger version of differentiability, namely differentiability of the average moment equation with uniformly bounded derivative. This difference appears a necessary (and low) price for non-stationairty to establish asymptotic normality in the generalized method of moments.

The uniform weak laws of large numbers, Conditions [S7](#page-21-1) and [S8,](#page-22-0) can be verified by using, for example, Theorem 5.1 and 5.2 in Pötscher and Prucha [\[1997\]](#page-62-7), Theorems 1 and 2 in [Andrews](#page-58-6) [\[1988\]](#page-58-6), or Corollary 1.4 in [van Handel](#page-64-5) [\[2013\]](#page-64-5), among others. For example, if the underlying process is strictly stationary and the involved function class is uniformly bounded with a finite bracketing number for every size of the brackets, then Corollary 1.4 in [van Handel](#page-64-5) [\[2013\]](#page-64-5) implies Conditions [S7](#page-21-1) and [S8.](#page-22-0) Uniform laws of large numbers can also be obtained without stationarity assumptions for strongly mixing (e.g.,  $\alpha$ -mixing) processes, or  $\phi$ -mixing processes [see, e.g., Chapter 5 in Pötscher and Prucha, 1997].

The central limit theorem, Condition [S9,](#page-22-1) can be verified by using, for example, results in [Rosenblatt](#page-63-9) [\[1956\]](#page-63-9), Theorems 5 and 6 in [Philipp](#page-62-8) [\[1969\]](#page-62-8), Corollary 2.11 in [McLeish](#page-62-9) [\[1977\]](#page-62-9), Corollary 1 in [Herrndorf](#page-61-10) [\[1984\]](#page-61-10), Theorem 3.6 in [Davidson](#page-60-10) [\[1992\]](#page-60-10), Theorem 2.1 and Corollary 2.1 in [Arcones and Yu](#page-58-7) [\[1994\]](#page-58-7), Theorems  $10.1$  and  $10.2$  in Pötscher and [Prucha](#page-62-7) [\[1997\]](#page-62-7), or Theorem 3.23 in [Dehling and Philipp](#page-60-11) [\[2002\]](#page-60-11), Theorem 1.1 and 6.1 in [Bradley and Tone](#page-59-8) [\[2017\]](#page-59-8), among others. For example, define  $||A||_q = (E|A|^q)^{1/q}$  for any random variable A. For the doubly robust method, suppose that the underlying process is strongly mixing with coefficients  $\alpha(k)$ ,  $k \geq 1$  [\[Rosenblatt, 1956\]](#page-63-9). If, for some  $s \in (2,\infty]$ and with  $r = 2/s$ ,  $\sum_{k=1}^{\infty} \alpha(k)^{1-r} < \infty$  and  $\limsup_{t \to \infty} ||R^{\top} \Omega \nabla_{\theta} G_t(\theta) ||_{\theta = \theta_{\infty}} ||_s < \infty$ , then Condition [S9](#page-22-1) holds by Corollary 1 in [Herrndorf](#page-61-10) [\[1984\]](#page-61-10).

# <span id="page-23-1"></span>S3 Relaxing stationarity of confounders

#### <span id="page-23-0"></span>S3.1 Estimand and identification

As mentioned in Section [4,](#page-6-1) stationarity of confounders, namely Condition [4,](#page-7-1) may be too strong in certain conditions. In this section, we drop this condition and present more general identification results. We will generally add a tilde symbol under various symbols for objects similar to those from Section [4.](#page-6-1)

Without Condition [4,](#page-7-1) it is challenging to identify the average treatment effect for the treated unit  $\phi_{t_+}^*$  at each post-treatment time period  $t_+$ . We therefore consider another causal estimand, the time-averaged average treatment effect for the treated unit. We first consider the case where T and  $T_0$  are fixed. Let  $\ell_T : t_+ \mapsto \ell_T(t_+) \geq 0$  be a user-specified importance weighting function that satisfies  $\sum_{t_+ = T_0+1}^{T} \ell_T(t_+) = 1$ , and define  $\phi$ ะ<br>. \* to be  $\sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+})\phi^{*}(t_{+}) = \sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+})E\{Y_{t_{+}}(1)-Y_{t_{+}}(0)\}$ . Thus,  $\phi$ 1<sup>p</sup> ∗ T is a weighted average of time-specific ATTs  $\phi_{t_+}^*$  over post-treatment time periods with averaging weights given by  $\ell_T(\cdot)$ . We refer to  $\ell_T$  as an importance weight because it encodes the importance of each post-treatment time period in the average treatment effect for the treated unit  $\phi_T^*$  averaged over time. In the notation, we suppress the dependence of  $\phi_T^*$  on  $\ell_T$  and  $T_0$  for conciseness.

Our approach to this problem via weighting is similar to the case with stationarity in Section [5.](#page-10-1) We again use a treatment bridge function to capture the covariate shift of confounders between pre- and post-treatment periods and subsequently use this function to weight pre-treatment observations to impute post-treatment outcomes. A key difference is that the treatment bridge function in this case also needs to incorporate the importance weight  $\ell_T(\cdot)$  and therefore is a proxy of a Radon-Nikodym derivative between two mixtures of distributions. Specifically, we rely on the following condition on the existence of a treatment confounding bridge function  $q^*$ .

<span id="page-24-0"></span>**Condition S10.** There exists a function  $q_T^* : \mathcal{Z} \to \mathbb{R}$  such that, for any square-integrable function  $g: \mathcal{U} \to \mathbb{R}$ :

<span id="page-24-1"></span>
$$
\frac{1}{T_0} \sum_{t=1}^{T_0} E\{q_T^*(Z_{t-})g(U_{t-})\} = \sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E\{g(U_{t+})\}.
$$
\n(S2)

We have dropped the dependence of the treatment confounding bridge function  $q_T^*$  on the importance weighting function  $\ell_T$  from the notation for conciseness. Condition [S10,](#page-24-0) namely [\(S2\)](#page-24-1), might appear non-intuitive. We next rewrite [\(S2\)](#page-24-1) in an integral form and argue that  $q^*$  can be interpreted as a proxy of a Radon-Nikodym derivative, similarly to  $q^*$  in Condition [5.](#page-8-1) Let  $\mathscr{U}_T$  be the uniform law over pre-treatment time periods  $\{1, \ldots, T_0\}$ and  $\mathscr{L}_T$  be the law over post-treatment time periods  $\{T_0 + 1, T_0 + 2, \ldots, T\}$  defined by  $\mathscr{L}_T(\{t_+\}) = \ell_T(t_+)$  for  $t_+ > T_0$ . With these definitions, [\(S2\)](#page-24-1) is equivalent to

$$
\int_{\{1,\ldots,T\}} E\{q_T^*(Z_{t-})g(U_{t-})\} \mathscr{U}_T(\mathrm{d}t_-) = \int_{\{T_0+1,\ldots,T\}} E\{g(U_{t+})\} \mathscr{L}_T(\mathrm{d}t_+),
$$

which is also equivalent to

$$
\int_{\{1,\ldots,T\}} E[E\{q_T^*(Z_{t-}) \mid U_{t-}\}g(U_{t-})] \mathscr{U}_T(\mathrm{d}t_-) = \int_{\{T_0+1,\ldots,T\}} E\{g(U_{t+})\} \mathscr{L}_T(\mathrm{d}t_+).
$$

Since g is an arbitrary square-integrable function, clearly  $u \mapsto E\{q_T^*(Z_{t-}) \mid U_{t-} = u\}$  may be viewed as a Radon-Nikodym derivative, of the law of  $U_{t_{+}}$  ( $t_{+} \sim \mathscr{L}_{T}$ ), with respect to the law of  $U_{t-}$  ( $t-\sim \mathscr{U}_T$ ), namely

$$
u \mapsto \frac{\int_{\{T_0+1,\ldots,T\}} f_{U_{t_+}}(u) \mathscr{L}_T(\mathrm{d} t_+)}{\int_{\{1,\ldots,T_0\}} f_{U_{t_-}}(u) \mathscr{U}_T(\mathrm{d} t_-)},
$$

where we have informally used  $f_A$  to denote the density of a random variable A.

The next condition is a completeness condition that is similar to Condition [S2.](#page-19-2)

<span id="page-24-2"></span>**Condition S11.** Let  $g: \mathcal{Z} \to \mathbb{R}$  be any square-integrable function. The following two statements are equivalent: (i)  $g(Z_{t-}) = 0$  for all  $t_- \leq T_0$ , (ii)  $\sum_{t=1}^{T_0} E\{g(Z_{t-}) f(W_{t-})\}/T_0$  $= 0$  for any square-integrable function f.

The second statement of Condition [S11](#page-24-2) is a generalization of the second statement of Condition [S2.](#page-19-2) Indeed, the latter is equivalent to the following:  $E\{g(Z_{t-})f(W_{t-})\}=0$  for any square-integrable function f.

With the above two conditions, we have the following identification result for  $\phi^*_7$ without stationarity for  $U_{t_+}$  in Condition [4.](#page-7-1)

<span id="page-25-1"></span>**Theorem S1** (Identification of average treatment effect with  $q_T^*$ ). Let  $f : \mathbb{R} \to \mathbb{R}$  be any square-integrable function. Under Conditions [1,](#page-5-0) [3,](#page-7-0) and [S10,](#page-24-0) it holds that

<span id="page-25-3"></span>
$$
\sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+}) E\{f(Y_{t_{+}}(0))\} = \frac{1}{T_{0}} \sum_{t_{-}=1}^{T_{0}} E\{\underline{q}_{T}^{*}(Z_{t_{-}})f(Y_{t_{-}})\}.
$$
\n(S3)

In particular, taking f to be the identity function, it holds that

<span id="page-25-4"></span>
$$
\sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+}) E\{Y_{t_{+}}(0)\} = \frac{1}{T_{0}} \sum_{t_{-}=1}^{T_{0}} E\{\underline{q}_{T}^{*}(Z_{t_{-}})Y_{t_{-}}\}.
$$
\n(S4)

and thus  $\oint_T^*$  is identified as  $\oint_T^* = \sum_{t_+ = T_0 + 1}^T E\{Y_{t_+}\} - \sum_{t_- = 1}^{T_0} E\{q^*(Z_{t_-})Y_{t_-}\}/T_0$ . In addition, the treatment confounding bridge function  $q_T^*$  is a solution to

<span id="page-25-0"></span>
$$
\frac{1}{T_0} \sum_{t_0=1}^{T_0} E\{q(Z_{t_0})g(W_{t_0})\} = \sum_{t_0=T_0+1}^{T} \ell_T(t_+) E\{g(W_{t_0})\} \quad \text{for all square-integrable } g \tag{S5}
$$

in  $q: \mathcal{Z} \to \mathbb{R}$ . Further, under Condition [S11,](#page-24-2) [\(S5\)](#page-25-0) has a unique solution almost surely.

We also have the following doubly robust identification result similar to Theorem [2.](#page-10-0)

<span id="page-25-5"></span>**Theorem S2** (Doubly robust identification with  $h^*$  and  $q_T^*$ ). Let  $h : \mathcal{W} \to \mathbb{R}$  and  $q: \mathcal{Z} \to \mathbb{R}$  be any square-integrable functions. Under Conditions [1](#page-5-0) and [3,](#page-7-0) if either (i) Condition [2](#page-5-2) holds and  $h = h^*$ , or (ii) Condition [S10](#page-24-0) holds and  $q = q_T^*$ , then

$$
\sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+}) E[Y_{t_{+}}(0)] = \frac{1}{T_{0}} \sum_{t_{-}=1}^{T_{0}} E\left[g(Z_{t_{-}}) \{Y_{t_{-}} - h(W_{t_{-}})\}\right] + \sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+}) E\{h(W_{t_{+}})\},
$$
\n
$$
\phi_{T}^{*} = \sum_{t_{-}}^{T} \ell_{T}(t_{+}) E\left\{Y_{t_{+}} - h(W_{t_{+}})\right\} - \frac{1}{T_{0}} \sum_{t_{-}}^{T_{0}} E\left[q(Z_{t_{-}}) \{Y_{t_{-}} - h(W_{t_{-}})\}\right].
$$
\n(S6)

<span id="page-25-2"></span>
$$
\phi_T^* = \sum_{t_+ = T_0 + 1} \ell_T(t_+) E\left\{ Y_{t_+} - h(W_{t_+}) \right\} - \frac{1}{T_0} \sum_{t_- = 1} E\left[ q(Z_{t_-}) \{ Y_{t_-} - h(W_{t_-}) \} \right].
$$
 (S6)  
Since our estimators and associated theoretical results are asymptotic as  $T \to \infty$ , we

also present identification results under the same asymptotic regime. The causal estimand of interest is  $\phi^*$  defined as  $\lim_{T\to\infty} \phi^*_T = \lim_{T\to\infty} \sum_{t_+ = T_0 + 1}^T \ell_T(t_+) E\{Y_{t_+}(1) - Y_{t_+}(0)\}.$  The associated identification results are similar to Theorem [S1](#page-25-1) with the key difference being the limit as  $T \to \infty$ , so we abbreviate our presentation. Recall that we assume that  $T_0/T \to \gamma \in (0,1)$  as  $T \to \infty$ .

<span id="page-26-0"></span>**Condition S12.** There exists a function  $q^* : \mathcal{Z} \to \mathbb{R}$  such that, for any square-integrable function  $g: \mathcal{U} \to \mathbb{R}$ , it holds that

$$
\lim_{T \to \infty} \frac{1}{T_0} \sum_{t=1}^{T_0} E\{g^*(Z_{t-})g(U_{t-})\} = \lim_{T \to \infty} \sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E\{g(U_{t+})\}.
$$
 (S7)

<span id="page-26-1"></span>**Condition S13.** Let  $g : \mathcal{Z} \to \mathbb{R}$  be any square-integrable function. The following two statements are equivalent: (i)  $g(Z_t) = 0$  for all  $t \in \{1, 2, \ldots\}$ , (ii)  $\lim_{T \to \infty} \sum_{t=1}^{T_0} E\{g(Z_{t-})\}$  $f(W_{t-})/T_0 = 0$  for any square-integrable function f.

<span id="page-26-5"></span>**Theorem S3** (Identification of average treatment effect with  $q^*$ ). Let  $f : \mathbb{R} \to \mathbb{R}$  be any square-integrable function. Under Conditions [1,](#page-5-0) [3,](#page-7-0) and [S12,](#page-26-0) it holds that

<span id="page-26-3"></span>
$$
\lim_{T \to \infty} \sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E\{f(Y_{t_+}(0))\} = \lim_{T \to \infty} \frac{1}{T_0} \sum_{t_- = 1}^{T_0} E\{\underline{q}^*(Z_{t_-})f(Y_{t_-})\}.
$$
 (S8)

In particular, taking f to be the identity function, it holds that

$$
\lim_{T \to \infty} \sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E\{Y_{t_+}(0)\} = \lim_{T \to \infty} \frac{1}{T_0} \sum_{t_- = 1}^{T_0} E\{\underline{q}^*(Z_{t_-})Y_{t_-}\}.
$$
 (S9)

and thus  $\phi^* = \lim_{T \to \infty} \left[ \sum_{t_+ = T_0 + 1}^T E\{Y_{t_+}\} - \sum_{t_- = 1}^{T_0} E\{q^*(Z_{t_-})Y_{t_-}\}/T_0 \right]$ . In addition, the treatment confounding  $\tilde b$ ridge function  $q_T^*$  is a solution to

<span id="page-26-2"></span>
$$
\lim_{T \to \infty} \frac{1}{T_0} \sum_{t=1}^{T_0} E\{q(Z_{t-})g(W_{t-})\} = \lim_{T \to \infty} \sum_{t_{+}=T_0+1}^{T} \ell_T(t_{+}) E\{g(W_{t+})\}
$$
\n(S10)

\nfor all square-integrable function  $g$ 

in  $q: \mathcal{Z} \to \mathbb{R}$ . Further, under Condition [S13,](#page-26-1) [\(S10\)](#page-26-2) has a unique solution almost surely.

<span id="page-26-4"></span>**Theorem S4** (Doubly robust identification with  $h^*$  and  $q^*$ ). Let  $h : \mathcal{W} \to \mathbb{R}$  and  $q : \mathcal{Z} \to$  $\mathbb R$  be any square-integrable functions. Under Conditions  $\widetilde{I}$  and [3,](#page-7-0) if either (i) Condition [2](#page-5-2) holds and  $h = h^*$ , or (ii) Condition [S12](#page-26-0) holds and  $q = q^*$ , then

$$
\lim_{T \to \infty} \sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E\{Y_{t_+}(0)\}
$$
\n
$$
= \lim_{T \to \infty} \left( \frac{1}{T_0} \sum_{t_- = 1}^{T_0} E\left[ q(Z_{t_-}) \{Y_{t_-} - h(W_{t_-})\} \right] + \sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E\{h(W_{t_+})\} \right),
$$

<span id="page-27-0"></span>
$$
\phi_T^* = \lim_{T \to \infty} \left( \sum_{t_+ = T_0 + 1}^T \ell_T(t_+) E\left\{Y_{t_+} - h(W_{t_+})\right\} - \frac{1}{T_0} \sum_{t_- = 1}^{T_0} E\left[q(Z_{t_-}) \{Y_{t_-} - h(W_{t_-})\}\right]\right). \tag{S11}
$$

Remark 4. We use causal conditions on the treatment confounding bridge that connect to the unobserved confounder  $U_t$ , namely Conditions [5,](#page-8-1) [S10](#page-24-0) and [S12.](#page-26-0) With such conditions, we do not rely on the existence of the outcome confounding bridge function in Condition [2](#page-5-2) to obtain the doubly robust identification formulas [\(7\)](#page-10-2), [\(S6\)](#page-25-2) and [\(S11\)](#page-27-0). An alternative set of conditions that leads to the same identification formulas is the existence of the treatment confounding bridge function in terms of  $W_t$  (namely [\(6\)](#page-9-1), [\(S5\)](#page-25-0) and [\(S10\)](#page-26-2)) as well as the existence of the outcome confounding bridge function in Condition [2.](#page-5-2) Thus, these two sets of causal conditions do not imply each other. One potential drawback of the alternative set of conditions is that, for example, to identify  $E[f\{Y_{t+}(0)\}]$  for some given function f under stationarity of  $U_{t_{+}}$  (Condition [4\)](#page-7-1), a different outcome confounding bridge function that can identify the same causal estimand is required to exist, but the approach we present in this paper does not require this (see [\(4\)](#page-9-4), [\(S3\)](#page-25-3) and [\(S8\)](#page-26-3)). Identification of  $E[f\{Y_{t+}(0)\}]$  can be useful, for example, in constructing prediction intervals for the actual treatment effect for the treated unit. Since our main focus is the average treatment effect for the treated unit and both approaches lead to the same nonparametric identification formulas and therefore the same estimation procedures, we do not present this alternative approach in detail.

#### <span id="page-27-1"></span>S3.2 Doubly robust estimation

In this section, we describe a doubly robust method to estimate the average treatment effect for the treated unit based on Theorem [S4,](#page-26-4) along with its theoretical properties. Due to the similarity between Theorems [2](#page-10-0) and [S4,](#page-26-4) this method is also similar to that described in Section [5.](#page-10-1) We focus on the estimation of  $\phi^*$ , the limit (as the number of time periods tends to infinity) of an average of time-specific average treatment effect for the treated unit  $\phi_t^*$  over post-treatment time periods. We propose to use a generalized method of moments method that is similar to that described in Section [5.](#page-10-1) We therefore abbreviate our presentation with emphasis on key differences.

We still use  $\alpha$  and  $\beta$  to parameterize  $h^*$  and  $q^*$ , and will use  $h_{\alpha}$  and  $q_{\beta}$  to denote the models of confounding bridge functions with parameter  $\alpha$  and  $\beta$ , respectively. Key changes of the method without stationarity of confounders are (i) that we no longer parameterize the average treatment effect for the treated unit with  $\lambda$  since the estimand  $\phi^*$  is a scalar, and (ii) that we need to incorporate the user-specified importance weighting function  $\ell_T$ . We still let  $g_h: \mathcal{Z} \to \mathbb{R}^{d'_{\alpha}}$ , with  $d'_{\alpha} \geq d_{\alpha}$ , and  $g_q: \mathcal{W} \to \mathbb{R}^{d'_{\beta}}$ , with  $d'_{\beta} \geq d_{\beta}$ , be two user-specified functions. Let  $\alpha \in \mathcal{A}, \beta \in \mathcal{B}, \phi \in \Phi \subseteq \mathbb{R}, \psi \in \Psi \subseteq \mathbb{R}^{d_{\beta'}},$ and  $\psi_-\in\Psi_-\subseteq\mathbb{R}$  be parameters in the generalized method of moments procedure,  $\theta = (\alpha, \beta, \phi, \psi, \psi)$  be the collection of parameters, and  $\Theta = \mathcal{A} \times \mathcal{B} \times \Phi \times \Psi \times \Psi$  be the parameter space. For each  $t = 1, ..., T$ , define  $\mathcal{G}_{T,t} : \mathcal{Q} \mapsto \mathbb{R}^{2d'_{\beta}+3}$  as

<span id="page-28-2"></span><span id="page-28-1"></span>
$$
G_{T,t}: \varrho \mapsto \begin{pmatrix} \mathbb{1}(t \leq T_0) \left[ \{Y_t - h_{\alpha}(W_t)\} g_h(Z_t) \right] \\ \mathbb{1}(t > T_0) \left\{ \psi - g_q(W_t) \right\} \\ \mathbb{1}(t \leq T_0) \left\{ g_{\beta}(Z_t) g_q(W_t) - \psi \right\} \\ \mathbb{1}(t > T_0) \left[ \phi - (T - T_0) \ell_T(t) \{ Y_t - h_{\alpha}(W_t) \} + \psi_- \right] \\ \mathbb{1}(t \leq T_0) \left[ \psi_- - g_{\beta}(Z_t) \{ Y_t - h_{\alpha}(W_t) \} \right] \end{pmatrix} . \tag{S12}
$$

Let  $\Omega_T$  be a user-specified symmetric positive semi-definite  $(d'_\alpha + 2d'_\beta + 2) \times (d'_\alpha + 2d'_\beta + 2)$ matrix, for example, the identity matrix. Consider the generalized method of moments estimator

$$
\hat{\mathcal{Q}}_T = \left( \hat{\mathcal{Q}}_T, \hat{\mathcal{Q}}_T, \hat{\mathcal{Q}}_T, \hat{\mathcal{Q}}_T, \hat{\mathcal{W}}_{-T} \right) = \underset{\mathcal{Q} \in \mathcal{Q}}{\operatorname{argmin}} \left\{ \frac{1}{T} \sum_{t=1}^T \mathcal{G}_{T,t}(\mathcal{Q}) \right\}^\top \mathcal{Q}_T \left\{ \frac{1}{T} \sum_{t=1}^T \mathcal{G}_{T,t}(\mathcal{Q}) \right\}. \tag{S13}
$$

The entry  $\hat{\phi}_T$  in the generalized method of moments estimator is the estimator of the average treatment effect for the treated unit  $\phi^*$  averaged over the post-treatment time e periods. This generalized method of moments estimator coincides with the estimator  $\phi_{\hat{\lambda}_T}(t)$  in Section [5](#page-10-1) if the time-varying model  $\phi_{\lambda}$  for the average treatment effect for the treated unit is a constant function  $t \mapsto \lambda$  and the user-specified importance weight  $\ell_T(t) = 1/(T - T_0)$ . This is the case in all our simulations in Sections [6](#page-13-1) and [S7.2,](#page-37-0) as well as in our analysis of Kansas GDP in Section [S8.2.](#page-47-0)

Similarly to Condition [6,](#page-11-0) we make the following regularity conditions to obtain consistency and valid inference about  $\phi^*$ .

<span id="page-28-0"></span>**Condition S14.** (i) There exists a unique  $\ell_{\infty} = (\alpha_{\infty}, \beta_{\infty}, \phi_{\infty}, \psi_{\infty}, \psi_{-\infty}) \in \mathcal{Q}$  such that  $\lim_{T\to\infty}\sum_{t=1}^T E\{\mathcal{G}_{T,t}(\underline{\theta}_{\infty})\}/T = 0$ , (ii)  $h^* = h_{\underline{\alpha}_{\infty}}$  satisfies Condition [2,](#page-5-2) or  $\underline{q}^* = \underline{q}_{\beta_{\infty}}$ satisfies Condition [S10.](#page-24-0)

Though similar to Condition [6,](#page-11-0) Condition [S14](#page-28-0) is different. In part (i) of Condition [6,](#page-11-0) we require that the moment equation  $G_t$  at the true parameter has mean zero for all time periods; that is,  $E\{G_t(\theta_\infty)\}=0$  for all  $t=1,\ldots,T$ . This is a standard assumption for the generalized method of moments, and therefore Theorem [3](#page-11-1) follows immediately from standard theory for the generalized method of moments. In contrast, part (i) of Condition [S14](#page-28-0) only requires that the average of the moment equation means over all time periods has approximately mean zero, namely  $\lim_{T\to\infty}\sum_{t=1}^T E\{\mathcal{G}_{T,t}(\mathcal{Q}_\infty)\}/T=0$ . However, the moment equation might not have mean zero for all time periods; that is, we do not require that  $E\{G_{T,t}(\ell_\infty)\}=0$  for all  $t=1,\ldots,T$ . This relaxed condition deviates from the standard generalized method of moments assumptions, and therefore the standard theory for the generalized method of moments does not apply.

Another deviation from standard assumptions for the generalized method of moments is that the moment equation  $G_{t,T}$  depends on the sample size T. Therefore, when deriving the asymptotic normality of the estimator  $\theta_T$ , in order to apply an argument based on Taylor series, we require a slightly different differentiability condition (see Condition [S8](#page-22-0) in Web Appendix [S2\)](#page-20-0) on the moment equation from standard conditions for the generalized method of moments. We obtain the following theoretical result for the above generalized method of moments method after carefully modifying the proof for the standard theory of the generalized method of moments.

<span id="page-29-0"></span>**Theorem S5.** Under Conditions [1,](#page-5-0) [3,](#page-7-0) [S14,](#page-28-0) [S3](#page-20-1)[–S5](#page-21-0) and [S7,](#page-21-1) with the estimator  $\hat{\theta}_T$  from [\(S13\)](#page-28-1) and  $\ell_{\infty}$  in Condition [S14,](#page-28-0) it holds that  $\phi_{\infty} = \phi^*$  and, as  $T \to \infty$ ,  $\hat{\ell}_{T}$  is consistent for  $\theta_{\infty}$  and  $\hat{\phi}_{T}$  is consistent for  $\phi^{*}$ . Additionally, under Conditions [S6,](#page-21-2) [S8](#page-22-0) and [S9,](#page-22-1) it holds that, as  $T \to \infty$ ,  $\sqrt{T}(\hat{\theta}_T - \theta_\infty)$  is asymptotically distributed as  $N(0, \mathcal{A}^{-1} \mathcal{B} \mathcal{A}^{-1})$ , √ where, with  $G_{T,t}$  in [\(S12\)](#page-28-2) and  $\Omega$  being the probability limit of  $\Omega_T$  in Condition [S3,](#page-20-1) we use the following quantities, whose existence follows based on our assumptions:

$$
R = \lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^{T} E\{\nabla_{\underline{\theta}} \underline{G}_{T,t}(\underline{\theta})|_{\underline{\theta} = \underline{\theta}_{\infty}}\}, \qquad A = R^{\top} \Omega R,
$$
  

$$
B = R^{\top} \Omega \left[ \lim_{T \to \infty} \text{var} \left\{ T^{-1/2} \sum_{t=1}^{T} \underline{G}_{T,t}(\underline{\theta}_{\infty}) \right\} \right] \Omega R.
$$

Theorem [S5](#page-29-0) appears similar to Theorem [3,](#page-11-1) but there is a key difference in their practical implications in data analysis. Under stationarity of confounders, the moment equation for  $G_t$  has mean zero at the limiting parameter value  $\theta_{\infty}$  at each time period (see Condition [6\)](#page-11-0). In this case, the matrix  $B$  in Theorem [3](#page-11-1) can be consistently estimated with heteroskedasticity and autocorrelation consistent estimators. In contrast, without this stationarity, the moment equation  $\mathcal{G}_{T,t}$  might not have mean zero at any parameter e value at all time periods; only the average of the moment equation over all time periods is zero at the true parameter value (see Condition [S14\)](#page-28-0).

Therefore, it is challenging, if possible at all, to consistently estimate  $\underline{B}$  in Theorem [S5,](#page-29-0) which is a key component of obtaining a consistent variance estimator. Indeed, standard generalized method of moments software outputs standard errors under the assumption that the moment equation has mean zero at all time periods, and the limit of the variance in the "meat" of  $\underline{B}$  is typically estimated as a weighted average of squares of  $\underline{G}_{T,t}$ . This standard error is conservative, because  $E(X^2) \geq \text{var}(X)$  for any random variable X. Therefore, implementing the generalized method of moments without assuming the stationarity would lead to conservative statistical inference.

# <span id="page-30-0"></span>S4 Estimation of average treatment effect based on weighting

In this section, we describe the generalized method of moments estimators of the average treatment effect for the treated unit  $\phi^*(t)$  under stationarity of confounders and  $\phi^*$ without stationarity. Since the estimators are similar to the doubly robust generalized method of moments estimators, we abbreviate our presentation.

#### S4.1 Estimation under stationarity

Let  $g_q: \mathcal{W} \to \mathbb{R}^{d'_\beta}$ , with  $d'_\beta \geq d_\beta$ , be a user-specified function. For  $\beta \in \mathcal{B}$ ,  $\lambda \in \Lambda$ ,  $\psi \in \Psi$ , and  $\psi_-\in\Psi_-,$  define  $\theta^q=(\beta,\lambda,\psi,\psi_-),\,\Theta^q=\mathcal{B}\times\Lambda\times\Psi\times\Psi_-,$  and

$$
G_t^q : \theta^q \mapsto \begin{pmatrix} \mathbb{1}(t > T_0) \{ \psi - g_q(W_t) \} \\ \mathbb{1}(t \le T_0) \{ q_\beta(Z_t) g_q(W_t) - \psi \} \\ \mathbb{1}(t > T_0) \{ \phi_\lambda(t) - Y_t + \psi_- \} \\ \mathbb{1}(t \le T_0) \{ \psi_- - q_\beta(Z_t) Y_t \} \end{pmatrix}.
$$

In case of non-convexity, the interpretation is as for  $\hat{\theta}_T$  from [\(S1\)](#page-20-2). Let  $\Omega_T^q$  be a userspecified symmetric positive semi-definite  $(2d'_{\beta} + 2) \times (2d'_{\beta} + 2)$  matrix. Consider the generalized method of moments estimator

$$
\hat{\theta}_T^q = \left(\hat{\beta}_T^q, \hat{\lambda}_T^q, \hat{\psi}_T^q, \hat{\psi}_{-,T}^q\right) = \underset{\theta \in \Theta^q}{\text{argmin}} \left\{ \frac{1}{T} \sum_{t=1}^T G_t^q(\theta) \right\}^{\top} \Omega_T^q \left\{ \frac{1}{T} \sum_{t=1}^T G_t^q(\theta) \right\}.
$$

We propose to use  $\phi_{\hat{\lambda}^q_T}(t_+)$  as the estimator of the average treatment effect for the treated unit  $\phi^*(t_+)$  at time period  $t_+ > T_0$  based on the treatment confounding bridge function. We require the following assumption, similarly to condition [6.](#page-11-0)

<span id="page-30-1"></span>**Condition S15.** (i) There exists a unique  $\theta_{\infty}^q = (\beta_{\infty}^q, \lambda_{\infty}^q, \psi_{\infty}^q, \psi_{-\infty}^q, \phi_{\infty}^q) \in \Theta$  such that  $E\{G_t^q$  $\{f_t^q(\theta_{\infty}^q)\}=0$  for all  $t=1,\ldots,T$ . (ii)  $q^*=q_{\beta_{\infty}^q}$  satisfies Condition [5.](#page-8-1)

Under Condition [S15,](#page-30-1) Theorem [1](#page-9-0) implies that  $\phi_{\lambda^q_\infty}(t_+)$  equals the average treatment effect for the treated unit  $\phi^*(t_+)$  at time period  $t_+ > T_0$ . Similarly to Theorem [3,](#page-11-1) we have the following asymptotic result about the generalized method of moments estimator  $\hat{\theta}^q_7$  $T<sub>T</sub>$ , under the additional assumptions [S3–](#page-20-1)[S7.](#page-21-1)

<span id="page-30-2"></span>**Theorem S6.** Under Conditions [1,](#page-5-0) [3,](#page-7-0) [4,](#page-7-1) [S15,](#page-30-1) [S3](#page-20-1)[–S5](#page-21-0) and [S7,](#page-21-1) as  $T \rightarrow \infty$ , we have that  $\hat{\theta}_7^q$ that  $\hat{\theta}_T^q$  is consistent for  $\theta_\infty^q$ . Additionally under Conditions [S6,](#page-21-2) [S8](#page-22-0) and [S9,](#page-22-1) as  $T \to \infty$ ,  $\overline{T}(\hat{\theta}_{T}^{\bar{q}}-\theta_{\infty}^{q})$  is asymptotically  $\mathrm{N}(0,A_{q}^{-1}B_{q}A_{q}^{-1}),$  where we define

$$
R_q = \lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^T E\{\nabla_{\theta^q} G_t^q(\theta^q) |_{\theta^q = \theta^q_\infty}\}, \qquad A_q = R_q^\top \Omega^q R_q,
$$

$$
B_q = R_q^{\top} \Omega^q \left[ \lim_{T \to \infty} \text{var} \left\{ T^{-1/2} \sum_{t=1}^T G_t^q(\theta^q_{\infty}) \right\} \right] \Omega^q R_q.
$$

#### S4.2 Estimation without stationarity

Let  $g_q: \mathcal{W} \to \mathbb{R}^{d'_\beta}$ , with  $d'_\beta \geq d_\beta$ , be a user-specified function. For  $\beta \in \mathcal{B}$ ,  $\phi \in \Phi$ ,  $\psi \in \Psi$ , and  $\psi_-\in\Psi_-,$  define  $\underline{\theta}^q=(\beta,\phi,\psi,\psi_-),$   $\ominus^q=\mathcal{B}\times\Phi\times\Psi\times\Psi_-,$  and

$$
G_{T,t}^{q}: \theta^{q} \mapsto \begin{pmatrix} \mathbb{1}(t > T_{0}) \left\{ \psi - g_{q}(W_{t}) \right\} \\ \mathbb{1}(t \leq T_{0}) \left\{ g_{\beta}(Z_{t}) g_{q}(W_{t}) - \psi \right\} \\ \mathbb{1}(t > T_{0}) \left\{ \phi - (T - T_{0}) \ell_{T}(t) Y_{t} + \psi_{-} \right\} \\ \mathbb{1}(t \leq T_{0}) \left\{ \psi_{-} - g_{\beta}(Z_{t}) Y_{t} \right\} \end{pmatrix}.
$$

Let  $\Omega^q_T$  be a user-specified symmetric positive semi-definite  $(2d'_{\beta} + 2) \times (2d'_{\beta} + 2)$  matrix. Consider the generalized method of moments estimator

$$
\hat{\mathcal{Q}}_T^q = \left(\hat{\beta}_T^q, \hat{\phi}_T^q, \hat{\psi}_T^q, \hat{\psi}_T^q\right) = \underset{\underline{\theta} \in \Theta^q}{\text{argmin}} \left\{ \frac{1}{T} \sum_{t=1}^T \mathcal{G}_{T,t}^q(\theta) \right\}^{\top} \Omega_T^q \left\{ \frac{1}{T} \sum_{t=1}^T \mathcal{G}_t^q(\underline{\theta}) \right\}.
$$

We propose to use  $\hat{\phi}_T$  as the estimator of the average treatment effect for the treated unit  $\phi^*$  based on the treatment confounding bridge function. We require the following e assumption, similarly to condition [S14.](#page-28-0)

<span id="page-31-0"></span>**Condition S16.** (i) There exists a unique  $\mathcal{Q}_{\infty}^q = (\mathcal{Q}_{\infty}^q, \phi_{\infty}^q, \psi_{\infty}^q, \psi_{\infty}^q, \psi_{\infty}^q, \phi_{\infty}^q) \in \mathcal{Q}$  such that  $\lim_{T\to\infty}\sum_{t=1}^T E\{\mathcal{G}^q_{T,t}(\mathcal{Q}^q_{\infty})\}=0.$  (ii)  $\underline{q}^*=\underline{q}_{\beta^q_{\infty}}$  satisfies Condition [5.](#page-8-1)

Under Condition [S16,](#page-31-0) Theorem [S3](#page-26-5) implies that  $\phi^q_{\infty}$  equals the average treatment effect for the treated unit  $\phi^*$  averaged over post-treatment periods. Similarly to Theorem [S5,](#page-29-0) we have the following asymptotic result about the generalized method of moments estimator  $\hat{\theta}^q_{\tau}$  $T_T^q$ , under the additional assumptions [S3–](#page-20-1)[S7.](#page-21-1)

<span id="page-31-1"></span>**Theorem S7.** Under Conditions [1,](#page-5-0) [3,](#page-7-0) [S16,](#page-31-0) [S3–](#page-20-1)[S5](#page-21-0) and [S7,](#page-21-1) it holds that  $\phi^q_{\infty} = \phi^*$ , and as  $T\to\infty$ , we have that  $\hat{\theta}^q_{T}$  is consistent for  $\theta^q_{\infty}$ , which implies that  $\hat{\phi}^q_{T}$  is consister Additionally under Conditions [S6,](#page-21-2) [S8](#page-22-0) and [S9,](#page-22-1) as  $T \to \infty$ ,  $\sqrt{T}(\hat{\theta}_T^q - \theta_\infty^q)$  is asymptotically  $\frac{q}{T}$  is consistent for  $\theta_{\infty}^{q}$ , which implies that  $\hat{\theta}_{T}^{q}$  $\int_T^q$  is consistent for  $\phi^*$ . √  $N(0, A_q^{-1}B_qA_q^{-1}),$  where we define

$$
R_q = \lim_{T \to \infty} \frac{1}{T} \sum_{t=1}^T E\{\nabla_{\theta^q} G_{T,t}^q(\theta^q) |_{\theta^q = \theta^q_{\infty}}\}, \qquad A_q = R_q^\top \Omega^q R_q,
$$
  

$$
B_q = R_q^\top \Omega^q \left[ \lim_{T \to \infty} \text{var}\left\{ T^{-1/2} \sum_{t=1}^T G_{T,t}^q(\theta^q_{\infty}) \right\} \right] \Omega^q R_q.
$$

### <span id="page-32-0"></span>S5 Incorporating covariates

For some set X, let  $X_t \in \mathcal{X}$  be the observed covariates in set at time period t. A natural way to incorporate covariate  $X_t$  is to condition on  $X_t$ , treating  $X_t$  similarly to  $U_t$ , but as observed. Below we list the causal assumptions and identification results with a one-to-one correspondence to those in the main text under stationarity of covariates in the post-treatment periods (similar to Condition [4\)](#page-7-1). In fact, the identification result Theorem [2](#page-10-0) in the main text is a special case of Theorem [S9](#page-33-1) below with  $X_t = \emptyset$ . The proof of Theorem [S9](#page-33-1) below is also similar with the only modification being to condition on  $X_t$  throughout. We therefore omit their proofs. The results without stationarity are strikingly similar and thus omitted. The generalized method of moments estimation methods corresponding to the results below can also be derived similarly to those without covariates.

An issue with the results below is that the covariates  $X_t$  must be used in the posttreatment period in the bridge functions. We therefore need to assume that  $X_t$  is exogenous. However, this assumption may well be violated, since distributional shift from  $X_{t-}$  to  $X_{t+}$  may be a consequence of the treatment, thus being endogenous. This is in contrast to the classical SC method from [Abadie et al.](#page-58-1) [\[2010\]](#page-58-1), where only covariates in the pre-treatment period are used.

Another possible way to use covariates is to view them as proxies and concatenate them to  $W_t$  or  $Z_t$ . The user can decide which set of proxies each covariate belongs to based on Conditions [1–](#page-5-0)[S2.](#page-19-2) For example, covariates of donors and the treated unit may be part of  $W_t$ , while covariates of the other control units may be part of  $Z_t$ .

Our assumptions are as follows:

<span id="page-32-1"></span>**Condition S17.** For all  $t_-\leq T_0$ ,  $Z_{t_-}\perp (Y_{t_-}, W_{t_-})$  |  $(X_{t_-}, U_{t_-})$ .

<span id="page-32-4"></span>**Condition S18.** There exists a function  $h^* : \mathcal{W} \times \mathcal{X} \to \mathbb{R}$  such that  $E\{h^*(W_t, X_t) \mid$  $X_t, U_t$  =  $E\{Y_t(0) | X_t, U_t\}$  for all t.

Condition S19. Let  $g : W \to \mathbb{R}$  be any square-integrable function. For all  $t_-\leq T_0$ and  $P_{X_{t-}}$ -a.e.  $x \in \mathcal{X}$ , the following two statements are equivalent: (i)  $g(W_{t-}) = 0$ ,  $P_{W_{t-}|X_{t-}=x}$ -almost surely, (ii)  $E\{g(W_{t-}) | Z_{t-}, X_{t-}=x\} = 0, P_{Z_{t-}|X_{t-}=x}$ -almost surely.

<span id="page-32-2"></span>**Condition S20.** The conditional distribution  $(Y_t(0), W_t) | (X_t, U_t)$  is identical for all t.

<span id="page-32-5"></span>**Condition S21.** The distribution of  $(X_{t_+}, U_{t_+})$  is identical for all  $t_+ > T_0$ .

<span id="page-32-3"></span>**Condition S22.** Suppose that  $P_{X_{t_+}, U_{t_+}}$  is dominated by  $P_{X_{t_-}, U_{t_-}}$  and there exists a function  $q^*$ :  $\mathcal{Z} \times \mathcal{X} \to \mathbb{R}$  such that, for  $P_{X_{t-},U_{t-}}$ -a.e.  $(x, u)$  and for all  $t- \leq T_0$  and  $t_{+} > T_{0}$ ,

$$
E\{q^*(Z_{t-},x) \mid X_{t-}=x, U_{t-}=u\} = \frac{\mathrm{d}P_{X_{t+},U_{t+}}}{\mathrm{d}P_{X_{t-},U_{t-}}}(x,u).
$$

<span id="page-33-2"></span>Condition S23. Let  $g: \mathcal{Z} \to \mathbb{R}$  be any square-integrable function. For all  $t_-\leq T_0$ and  $P_{X_{t-}}$ -a.e.  $x \in \mathcal{X}$ , the following two statements are equivalent: (i)  $g(Z_{t-}) = 0$ ,  $P_{Z_{t-}|X_{t-}=x}$ -almost surely, (ii)  $E\{g(Z_{t-}) | W_{t-}, X_{t-}=x\} = 0$ ,  $P_{W_{t-}|X_{t-}=x}$ -almost surely.

The resulting theorems are analogous to those from the main text.

**Theorem S8.** Let  $f : \mathbb{R} \to \mathbb{R}$  be any square-integrable function. Under Conditions [S17](#page-32-1) and [S20](#page-32-2)[–S22,](#page-32-3) it holds that

$$
E[f\{Y_{t+}(0)\}] = E\{q^*(Z_{t-}, X_{t-})f(Y_{t-})\}.
$$
\n(S14)

In particular, taking f to be the identity function, it holds that

$$
E\{Y_{t+}(0)\} = E\{q^*(Z_{t-}, X_{t-})Y_{t-}\}\tag{S15}
$$

and thus  $\phi^*(t_+) = E\{Y_{t_+} - q^*(Z_{t_-}, X_{t_-})Y_{t_-}\}\$  for all  $t_- \leq T_0$  and  $t_+ > T_0$ . In addition,  $P_{W_{t_+},X_{t_+}}$  is dominated by  $P_{W_{t_-},X_{t_-}}$  and the treatment confounding bridge function  $q^*$  is a solution to

<span id="page-33-3"></span>
$$
E\{q(Z_{t-},x) \mid W_{t-}=w, X_{t-}=x\} = \frac{\mathrm{d}P_{W_{t+},X_{t+}}}{\mathrm{d}P_{W_{t-},X_{t-}}}(w,x) \quad \text{for } P_{W_{t-},X_{t-}}\text{-a.e. } (w,x) \in \mathcal{W} \times \mathcal{X}
$$
\n(S16)

in  $q : \mathcal{Z} \times \mathcal{X} \to \mathbb{R}$ . Also assuming Condition [S23,](#page-33-2) [\(S16\)](#page-33-3) has a unique solution up to probability zero sets.

<span id="page-33-1"></span>**Theorem S9.** Let  $h : \mathcal{W} \times \mathcal{X} \to \mathbb{R}$  and  $q : \mathcal{Z} \times \mathcal{X} \to \mathbb{R}$  be any square-integrable functions. Under Conditions [S17,](#page-32-1) [S18,](#page-32-4) [S20,](#page-32-2) [S21](#page-32-5) and [S22,](#page-32-3) it holds that

$$
E\{Y_{t+}(0)\} = E\left[q(Z_{t-}, X_{t-})\{Y_{t-} - h(W_{t-}, X_{t-})\} + h(W_{t+}, X_{t+})\right],
$$
  
\n
$$
\phi^*(t_+) = E\left[Y_{t+} - q(Z_{t-}, X_{t-})\{Y_{t-} - h(W_{t-}, X_{t-})\} - h(W_{t+}, X_{t+})\right]
$$
\n(S17)

for all  $t_-\leq T_0$  and  $t_+ > T_0$ , if (i) Condition [S18](#page-32-4) holds and  $h = h^*$ , or (ii) Condition [S22](#page-32-3) holds and  $q = q^*$ .

# <span id="page-33-0"></span>S6 Generalized method of moments implementation in simulations & data analysis

In this appendix, we describe some implementation details for the generalized method of moments used in our simulations and data analysis with the R package gmm. One challenge with the generalized method of moments in our methods is that the treatment confounding bridge function  $q^*$  is often parameterized as an exponential function, and thus the value of the moment equation is sensitive to small changes in the coefficients.

This seems to have led to numerical instability in our experience. We expect better software implementation of the generalized method of moments to resolve these issues by default, but with the current software implementation, we describe some non-default options that we have found to alleviate numerical issues for our methods.

We recommend providing the analytic gradient function to gmm function. Compared to the default numerical gradients, in our experiments, the analytic gradient functions could improve stability and speed in numerical optimization in the generalized method of moments.

Because the generalized method of moments can be highly nonlinear, the associated optimization problem can have several local optima. Thus, proper initialization can be crucial to obtaining an estimator that is close to the truth. When feasible, we recommend fitting a GLM assuming no latent confounders and taking the fitted coefficients as the initial values. For example, when  $h^*$  is specified as a linear function of  $W_t$ , we can fit an ordinary least squares regression with outcome  $Y_t$  and covariates  $W_t$  for  $t \leq T_0$ , and take the fitted coefficients as the initial coefficient for  $h^*$ .

When  $q^*$  is specified as a log-linear model, namely  $q^* : z \mapsto \exp\left(\beta_0 + \sum_{k=1}^K \beta_k z_k\right)$ , we may fit a GLM as follows. If we treat time period  $t$  as a random variable uniform over the observed time points, by Bayes' theorem, we find that

$$
\frac{\mathrm{d}P_{Z_{t+}}}{\mathrm{d}P_{Z_{t-}}}(z) = \frac{\mathrm{d}P_{Z_{t|t>T_0}}}{\mathrm{d}P_{Z_t|t\leq T_0}}(z) = \frac{\mathrm{pr}(t>T_0 \mid Z_t=z)}{\mathrm{pr}(t \leq T_0 \mid Z_t=z)} \frac{\mathrm{pr}(t \leq T_0)}{\mathrm{pr}(t > T_0)},
$$

where  $pr(t \leq T_0)$  and  $pr(t \leq T_0)$  may be interpreted as the proportions of pre- and post-treatment periods. We set the above to be equal to  $\exp\left(\beta_0 + \sum_{k=1}^K \beta_k z_k\right)$  for initialization. Then, with  $A_t$  defined to be  $\mathbb{1}(t > T_0)$ , we have that

<span id="page-34-0"></span>
$$
\log \text{OR}(A_t \mid Z_t = z) = \log \frac{\text{pr}(A_t = 1)}{\text{pr}(A_t = 0)} + \beta_0 + \sum_{k=1}^{K} \beta_k z_k \tag{S18}
$$

where OR stands for odds ratio, and  $pr(A_t = 1)/pr(A_t = 0) = (T - T_0)/T_0$ . Therefore, we can run a logistic regression with outcome  $A_t$  and covariate  $Z_t$  for all t, and derive an initial value for the coefficient  $\beta = (\beta_0, \ldots, \beta_K)$  with an adjustment in the intercept  $\beta_0$  according to [\(S18\)](#page-34-0). In simulations, we have found this initialization approach effective in removing a potentially large proportion of estimates that are far from the truth, thus achieving consistency and approximately correct confidence interval coverage. In practice, a sensitivity analysis can also be conducted by, for example, randomly initializing parameter values or optimizing the objective function via stochastic methods like stochastic gradient descent.

Many choices of the weight matrix  $\Omega_T$  lead to asymptotically normal generalized method of moments estimators since the only requirement is that the probability limit  $\Omega$  of  $\Omega_T$  is positive definite (see Condition [S3\)](#page-20-1). In addition to setting  $\Omega_T$  to be a fixed weight matrix, many choices of  $\Omega_T$  have been proposed, including the continuously updated efficient generalized method of moments, the two-step generalized method of moments estimator, and the iteratively updated generalized method of moments estimator. These three data-adaptive choices all lead to asymptotically efficient generalized method of moments estimators [\[Hansen et al., 1996\]](#page-61-11), while a fixed weight matrix is generally asymptotically inefficient. In finite samples, the continuously updated efficient generalized method of moments estimator has been found to perform better than the two-step generalized method of moments estimator and the iteratively updated estimator [\[Hansen](#page-61-11) [et al., 1996\]](#page-61-11), which are preferable to using a fixed weight matrix [\[Hall, 2007\]](#page-61-7). Nevertheless, we have found more numerical issues with these more complicated methods for our nonlinear generalized method of moments in finite samples. We thus recommend using a fixed weight matrix because it still leads to asymptotically valid inference despite its asymptotic inefficiency, and it has led to fewer numerical issues in our experience. In particular, we chose the weight matrix to be the identity matrix throughout; that is, we set wmatrix="'ident'' in the gmm function.

In addition, the option  $\mathbf{v}\circ\mathbf{v} =$  'HAC'' in the gmm function in the R package gmm has led to numerical errors in our experiments. The issue seems to be caused by prewhitening. Since our theory does not require prewhitening, we recommend not prewhitening and instead setting vcov=''iid'' in the gmm function, and then using vcovHAC with the default option prewhite=FALSE from the sandwich package to estimate the variance for dependent time series data. This has led to significantly fewer numerical issues and appears to have correct asymptotic behavior in our simulations. It is also feasible to use the NeweyWest function with the option prewhite=FALSE from the sandwich package to estimate the variance.

In some applications, the magnitude of the variables can be large. As an example, the number of hospitalizations in Section [7](#page-13-0) is of order  $10^2-10^3$ . Directly using these values in the GMM involving estimation of the treatment confounding bridge function  $q^*$  can lead to severe numerical instability. The reason seems that, when  $Z_t$  has large magnitudes and the treatment confounding bridge function is parameterized by a log-linear model as in our simulation and data analysis, even a tiny change in the parameter value will lead to a drastic change in the confounding bridge function value. The above issue with the magnitude of  $Z_t$  appears to be another challenge caused by strong nonlinearity and large gradients. Scaling all outcomes to be of a unit order before feeding the data into GMMs appears to significantly improve the numerical stability.

#### S7 Additional simulation details and results

#### <span id="page-36-0"></span>S7.1 Data-generating mechanism for just-identified seeting

We let  $T \in \{500, 1000, 2000, 4000\}$  and  $T_0 = T/2$ . For each T, we run 200 Monte Carlo simulations.

We let the number of latent confounders, the number of donors and of the other control units all be K. We generate random vectors  $U_t = (U_{t,1}, \ldots, U_{t,K})^\top$ ,  $W_t =$  $(W_{t,1},...,W_{t,K})^{\top}$  and  $Z_t = (Z_{t,1},...,Z_{t,K})^{\top}$ . We consider  $K \in \{2,3,4,5\}$ . We generate serially correlated latent confounders  $U_t$  via a Gaussian copula [e.g., [Jaworski](#page-62-10) [et al., 2010\]](#page-62-10) as follows:

$$
\epsilon_t \stackrel{i.i.d.}{\sim} \mathcal{N}(0, I_K), \qquad U_1 = \epsilon_1, \qquad U_t = 0.1 U_{t-1} + 0.9 \epsilon_t, \quad t \ge 2,
$$
  

$$
U_{t-,k} = F_1^{-1} \{ \Phi(U_{t-,k}) \}, \qquad U_{t+,k} = F_2^{-1} \{ \Phi(U_{t+,k}) \}, \quad t_- \le T_0, \ t_+ > T_0, \ k = 1, \dots, K,
$$

where  $I_K$  is the  $K \times K$  identity matrix,  $\Phi$  is the cumulative distribution function of the standard normal distribution, and  $F_{\lambda}$  is the cumulative distribution function of the Exponential( $\lambda$ ) distribution with rate  $\lambda > 0$ . Subsequently, for each  $t \geq 1$ , the observed outcomes  $(Y_t, W_t, Z_t)$  are generated independently conditional on  $U_t$  as follows:

$$
Y_t | U_t \sim \text{Unif}\left(2\mathbb{1}(t > T_0) + 2\sum_{k=1}^K U_{t,k} - 1, 2\mathbb{1}(t > T_0) + 2\sum_{k=1}^K U_{t,k} + 1\right)
$$
  

$$
W_{t,k} | U_t \sim \text{Unif}\left(2U_{t,k} - 1, 2U_{t,k} + 1\right), \qquad Z_{t,k} | U_t \sim \text{Unif}\left(2U_{t,k} - 1, 2U_{t,k} + 1\right)
$$

for  $k = 1, ..., K$ . Since  $U_t$  contains serial correlation, so do the observed variables  $(W_t, Z_t, Y_t)$ . As an example,  $(Y_t, W_t, Z_t)$  may be numbers of hospitalizations due to various causes up to shifting and scaling, while  $U_t$  may be the unobserved confounding factors, such as the overall infection level and the overall immune status. From the above formula for  $Y_t | U_t$ , we see that the true ATT equals two.

Therefore, the outcomes are generated from a factor model [\[Abadie et al., 2010\]](#page-58-1) and  $h^*: w \mapsto \alpha_0 + \sum_{k=1}^K \alpha_k w_k$  satisfies Condition [2](#page-5-2) for some coefficients  $\alpha_0, \ldots, \alpha_K \in \mathbb{R}$ . By the properties of the copula generating  $U_t$ , the likelihood ratio in Condition [5](#page-8-1) is  $C \exp(-\sum_{k=1}^K u_k)$  for some constant  $C > 0$ . By a simple calculation, we can check that  $q^*: z \mapsto \exp\left(\beta_0 + \sum_{k=1}^K \beta_k z_k\right)$  satisfies Condition [5](#page-8-1) for some coefficients  $\beta_0, \ldots, \beta_K \in \mathbb{R}$ . For methods that involve a correctly specified outcome confounding bridge function  $h^*$ , we parameterize  $h^*$  as above and choose  $g_h: z \mapsto (1, z^{\top})^{\top}$ . When  $q^*$  is correctly specified, we parameterize  $q^*$  as above and choose  $g_q: w \mapsto (1, w^{\top})^{\top}$ . When  $h^*$  is misspecified, we omit the last several proxies: we parameterize  $h^*$  as  $h: w \mapsto \alpha_0 + \alpha_1w_1$  with coefficients  $\alpha_0, \alpha_1 \in$ R, and choose  $g_h : z \mapsto (1, z_1)^\top$ . Similarly, when  $q^*$  is misspecified, we parameterize  $q^*$  as  $q: z \mapsto \exp(\beta_0 + \beta_1 z_1)$  with coefficients  $\beta_0, \beta_1 \in \mathbb{R}$ , and choose  $g_q: w \mapsto (1, w_1)^\top$ .

#### <span id="page-37-0"></span>S7.2 Over-identified setting

The data generating mechanism is similar to the just identified setting, and we abbreviate the description. There are three latent confounders  $U_t \in \mathbb{R}^3$ , five donors  $W_t \in \mathbb{R}^5$  and ten other control units  $Z_t \in \mathbb{R}^{10}$ . The latent confounders are generated as follows:

$$
\epsilon_t \stackrel{iid}{\sim} \mathcal{N}(0, I_3), \qquad U_1 = \epsilon_1, \qquad U_t = 0.1 U_{t-1} + 0.9 \epsilon_t \quad (t \ge 2),
$$
  
\n
$$
U_{t-,k} = F_1^{-1} \{ \Phi(U_{t-,k}) \}, \qquad U_{t+,k} = F_2^{-1} \{ \Phi(U_{t+,k}) \}, \quad t_- \le T_0, \ t_+ > T_0, \ k = 1, \ldots, 3.
$$

The observed outcomes are generated as follows:

$$
Y_t | U_t \sim \text{Unif}\left(2\mathbb{1}(t > T_0) + 2\sum_{k=1}^3 U_{t,k} - 1, 2\mathbb{1}(t > T_0) + 2\sum_{k=1}^3 U_{t,k} + 1\right),
$$
  
\n
$$
W_{t,k} | U_t \sim \text{Unif}(a_k^\top U_t - 1, a_k^\top U_t + 1), \quad k = 1, ..., 5,
$$
  
\n
$$
Z_{t,k} | U_t \sim \text{Unif}(b_k^\top U_t - 1, b_k^\top U_t + 1), \quad k = 1, ..., 10,
$$

where

$$
a_1 = (1, 0, 0)^{\top}, \quad a_2 = (0, 1, 0)^{\top}, \quad a_3 = (0, 0, 1)^{\top}, \quad a_4 = (1, 1, 0)^{\top}, \quad a_5 = (1, 0, 1)^{\top}, b_1 = (2, 0, 0)^{\top}, \quad b_2 = (0, 2, 0)^{\top}, \quad b_3 = (0, 0, 2)^{\top}, \quad b_4 = (-3, 0, 0)^{\top}, \quad b_5 = (0, -3, 0)^{\top}, b_6 = (0, 0, -3)^{\top}, \quad b_7 = (1, -1, 0)^{\top}, \quad b_8 = (1, 0, -1)^{\top}, \quad b_9 = (0, 1, -1)^{\top}, \quad b_{10} = (2, -0.5, -0.5)^{\top}.
$$

From the form of  $Y_t \mid U_t$ , the true average treatment effect for the treated unit equals two.

One can verify that there is a valid outcome confounding bridge function takes of form  $h^*: w \mapsto \alpha_0 + \sum_{k=1}^5 \alpha_k w_k$  for some coefficients  $\alpha_0, \alpha_1, \ldots, \alpha_5 \in \mathbb{R}$ . Further, there is a valid treatment confounding bridge function of the form  $q^* : z \mapsto \exp(\beta_0 + \sum_{k=1}^{10} \alpha_k z_k)$  for some coefficients  $\beta_0, \beta_1, \ldots, \beta_{10} \in \mathbb{R}$ . When  $h^*$  is correctly specified, we parameterize  $h^*$ as above and take  $g_h : z \mapsto (1, z^\top)^\top$ . When  $q^*$  is correctly specified, we also parameterize  $q^*$  as above and take  $g_q(w)$  to be the vector in  $\mathbb{R}^{21}$  consisting of  $w_{k_1}^{\gamma_{k_1}}$  $\chi_{k_1}^{\gamma_{k_1}} w_{k_2}^{\gamma_{k_2}}$  $k_2^{k_2}$  with  $k_1 \neq k_2$ ,  $\gamma_{k_1} + \gamma_{k_2} \leq 2$ , and both of  $\gamma_{k_1}$  and  $\gamma_{k_2}$  are non-negative integers. This can be concisely expressed as cbind(1,poly(w,degree=2,raw=TRUE)) in R.

We did not encounter numerical issues in this simulation. The sampling distributions of the estimated average treatment effect for the treated unit and the 95%-Wald confidence interval coverage are presented in Figure [S1](#page-38-0) and [S2,](#page-39-0) respectively. As shown in these figures, the performance of the methods is similar to the just identified setting from Section [6.](#page-13-1)

#### S7.3 Smaller sample size T

We investigate the performance of our methods when the sample size  $T$  is relatively small. In this simulation, we consider a similar setting as in Section [6](#page-13-1) except that  $T \in \{80, 100\}$ ,

<span id="page-38-0"></span>

Figure S1: Figures similar to Figure [2](#page-14-0) for the over-identified setting.

 $T - T_0 = 20$ , and we restrict to  $K = 2$ . The choice of T and  $T - T_0$  mimics the values in the analysis of Brazil all-cause pneumonia hospitalization in Section [7.](#page-13-0) The simulation results are presented in Figures [S3](#page-39-1) and [S4.](#page-40-1) We observe similar phenomena as in Section [6.](#page-13-1) However, due to smaller sample sizes  $T$ , the estimators' distributions are further from Gaussian than in Section [6,](#page-13-1) suggesting that the asymptotic results (Theorems [3,](#page-11-1) [S5,](#page-29-0) [S6,](#page-30-2) and [S7\)](#page-31-1) might not approximate the sampling distributions well at such small sample sizes. Thus, the somewhat outlying estimate from DR in Section [7,](#page-13-0) which uses a very simple model for the treatment confounding bridge function under a relatively small sample size  $(T = 108, T_0 = 84)$ , might not be surprising.

#### S7.4 Short post-treatment period

We also investigate the performance of our methods when the post-treatment period is relatively short. The simulation setting is also similar to Section [6,](#page-13-1) except that  $T_0 =$ relatively short. The simulation setting is also similar to Section 6, except that  $I_0 = T - \lfloor \sqrt{T} \rfloor$  and we restrict to  $K = 2$ , where  $\lfloor \cdot \rfloor$  denotes the floor function. Although the number of post-treatment time periods  $T - T_0$  still grows to infinity as  $T \to \infty$ , the growth has a slower rate and thus  $T_0/T \rightarrow 1$ . The results are presented in Figures [S5](#page-41-1) and [S6.](#page-42-0) Unlike Section [6](#page-13-1) where  $T_0/T = 1/2$ , proximal causal inference methods whose validity relies on the correct weighting function (corect.q and mis.h.DR) perform poorly with large bias, even if the treatment confounding bridge function is correctly specified. In contrast, methods whose validity can be justified by correct outcome bridge functions

<span id="page-39-0"></span>

Figure S2: Figures similar to Figure [3](#page-15-0) for the overidentified setting.

(correct.DR, correct.h and mis.q.DR) still appear to perform well as in Section [6.](#page-13-1) Thus, when  $T_0/T \rightarrow 1$ , our proposed method is not robust against misspecification of the outcome bridge function, even if the treatment confounding bridge function is correctly specified. Having a sufficient proportion of post-treatment periods appears necessary for our proposed method to be doubly robust. One explanation for this phenomenon is that, to obtain a stable estimator of the treatment confounding bridge function capturing the likelihood ratio in [\(3\)](#page-8-0), a comparable amount of data from pre- and post-treatment periods is needed. Having too little data from either period could lead to substantially larger variances for the treatment confounding bridge function estimator.

<span id="page-39-1"></span>

Figure S3: Figures similar to Figure [2](#page-14-0) for small sample sizes T.

<span id="page-40-1"></span>

Figure S4: Figures similar to Figure [3](#page-15-0) for small sample sizes T.

#### <span id="page-40-2"></span>S7.5 Autoregressive data-generating model

We consider the following autoregressive data-generating model:

$$
\epsilon_t \stackrel{i.i.d.}{\sim} \mathcal{N}(0, I_K), \qquad U_1 = \epsilon_1, \qquad U_t = 0.1U_{t-1} + 0.9\epsilon_t, \quad t \ge 2,
$$
  
\n
$$
Y_t \mid U_t \sim \mathcal{N}\left(2\mathbb{1}(t > T_0) + 2\sum_{k=1}^K U_{t,k}, 1\right),
$$
  
\n
$$
W_{t,k} \mid U_t \sim \mathcal{N}(2U_{t,k}, 1), \qquad Z_{t,k} \mid U_t \sim \mathcal{N}(2U_{t,k}, 1).
$$

We also restrict to  $K = 2$  and consider the seven methods as in Section [6.](#page-13-1) The only difference is the forms of the confounding bridge functions because  $U_t$  is marginally Gaussian, rather than exponentially, distributed (see Examples [1](#page-8-2) and [2\)](#page-9-3). In particular, when the outcome confounding bridge function is correctly specified, we parameterize  $h^*$  as  $w \mapsto \alpha_0 + \alpha_1 w_1 + \alpha_2 w_2 + \alpha_3 w_1^2 + \alpha_4 w_1 w_2 + \alpha_5 w_2^2$  and choose  $g_h : z \mapsto$  $(1, z_1, z_2, z_1^2, z_1 z_2, z_2^2)^\top$ ; when the treatment confounding bridge function is correctly specified, we parameterize  $q^*$  as  $z \mapsto \exp(\beta_0 + \beta_1 z_1 + \beta_2 z_2 + \beta_3 z_1^2 + \beta_4 z_1 z_2 + \beta_5 z_2^2)$  and choose  $g_q: w \mapsto (1, w_1, w_2, w_1^2, w_1w_2, w_2^2)^\top$ . We also consider misspecified confounding bridge functions similar to Section [6.](#page-13-1)

The simulation results are shown in Figures [S7](#page-43-1) and [S8.](#page-44-0) We overall performance of all methods are similar to Section [6.](#page-13-1) The only exception is that mis.h and OLS happen to also perform reasonably well. However, as we show in Section [6,](#page-13-1) these two methods are not guaranteed to produce consistent and asymptotically normal estimators.

#### <span id="page-40-0"></span>S7.6 Detrending before analysis

We investigate the effect of detrending the panel data before applying our methods. We consider a scenario similar to that in Section [S7.5,](#page-40-2) except that we add a time trend  $t \mapsto 10t/T + 10(t/T)^2$  when generating the observed data  $(W_t, Z_t, Y_t)$ . In all methods, we detrend the data as described in the following Section [S8](#page-43-0) before applying all methods.

<span id="page-41-1"></span>

Figure S5: Figures similar to Figure [2](#page-14-0) for short post-treatment periods.

We focus on the methods with correctly specified confounding bridge functions and the case  $K = 2$ . We run 1,000 Monte Carlo simulations for each T to obtain more accurate inference about the CI coverage.

The results are shown in Figures [S9](#page-45-0) and [S10.](#page-46-0) Although the estimators still appear normally distributed, the CI coverage is lower than the nominal level 95% by around 2–5%. Such anti-conservativeness might be caused by the fact that the uncertainty in estimating the trend is not accounted for in this procedure.

#### <span id="page-41-0"></span>S7.7 Post-selection

We investigate the effect of post-selecting  $W$  and  $Z$  before applying our methods. We consider a scenario similar to that in Section [S7.5,](#page-40-2) except that we first use Abadie's synthetic control method on all control units, and then select those with top  $K$  weights to be donors W and the others to be supplemental proxies  $Z$ . We also focus on the methods with correctly specified confounding bridge functions and the case  $K = 2$ .

The results are shown in Figures [S9](#page-45-0) and [S10.](#page-46-0) Among the 2,400 estimates, 202 estimates (8.4%) are outside the range (1, 3) shown in Figure [S9.](#page-45-0) The estimators' distributions appear further from normal distributions compared to the case without postselection, but the CI coverage still appears close to the nominal level 95%. With postselection, proximal methods appear to produce estimates with extremely large magnitudes and standard errors more often. This is likely due to the fact that post-selection

<span id="page-42-0"></span>

Figure S6: Figures similar to Figure [3](#page-15-0) for short post-treatment periods.

sometimes selects wrong units for proxies  $W$  and  $Z$ , and thus wrong models for both confounding bridge functions. However, in general, we do not expect the CI coverage to be close to the nominal level.

#### S7.8 Nonstationarity and time-varying treatment effect

We also investigate the performance of our proposed method with nonstationarity and time-varying treatment effects. We consider a scenario similar to that in Section [S7.5,](#page-40-2) except that (i)  $(W_t, Z_t, Y_t(0))$  has a seasonal trend  $1.5 \sin(20\pi t/T)$ , and (ii) the true treatment effect at time t is  $(8t)/(3T)$ , and thus the treatment effect averaged over posttreatment periods approaches the value two as  $T \to \infty$ . This scenario corresponds to Section [S3.](#page-23-1) We focus on the proximal methods based on the generalized method of moments involving weighting, and we expect them to be consistent, with conservative inference.

The results are shown in Figures [S13](#page-49-1)[–S15.](#page-51-0) As discussed in Section [S3.2,](#page-27-1) the estimates appear consistent and asymptotically normal, but the CI coverage is above the nominal level. Although the CI coverage is as high as  $100\%$  for large sample sizes ( $T > 2000$ ), the standard error overestimates the estimator's standard deviation by around 50–60%, and thus the CI may still be meaningful.

<span id="page-43-1"></span>

Figure S7: Figures similar to Figure [2](#page-14-0) for an autoregressive data-generating model.

# <span id="page-43-0"></span>S8 Additional examples

#### <span id="page-43-2"></span>S8.1 Analysis of Florida homicide rates

We apply our methods to study the effect of Florida's "stand your ground" law implemented in October 2005 on homicide rates [\[Bonander et al., 2021\]](#page-59-1). In these data, monthly public safety measures are collected in 16 states including Florida from January 1999 to December 2014.

Condition [5](#page-8-1) requires that the distribution of  $U_{t_+}$  is dominated by  $U_{t_-}$ , and [\(6\)](#page-9-1) can only be solved if the distribution of  $W_{t_+}$  is dominated by  $W_{t_-}$ . In practice, however, there might be a time trend in the time series  $(W_t, Z_t, Y_t)$ , in which case the treatment confounding bridge function would fail to exist. Therefore, we recommend preprocessing the data by detrending to account for the time trend, so that Condition [5](#page-8-1) may be plausible. One may fit a regression model  $f : \{1, \ldots, T\} \to \mathbb{R}$  with covariate being time t and outcome being the observed outcomes  $W_t$  and  $Z_t$  in all control units, use this model to predict an outcome  $f(t)$  for each time period t, and finally consider the residuals  $(\tilde{W}_t, \tilde{Z}_t, \tilde{Y}_t) = (W_t - f(t), Z_t - f(t), Y_t - f(t))$  as the new detrended outcomes.

We then use these detrended outcomes for all subsequent analyses. This does not impact the classical synthetic control method of [Abadie et al.](#page-58-1) [\[2010\]](#page-58-1), because, if we find  $Y_t(0) \approx \sum_i \alpha_i W_{ti}$  where  $W_t$  is the collection of  $W_{ti}$  and  $\sum_i \alpha_i = 1$ , then the same holds for detrended outcomes:  $\tilde{Y}_t(0) = Y_t(0) - f(t) \approx \sum_i \alpha_i \tilde{W}_{ti}^i$  where  $\tilde{W}_{ti} = W_{ti} - f(t)$ . If

<span id="page-44-0"></span>

Figure S8: Figures similar to Figure [3](#page-15-0) for an autoregressive data-generating model.

 $\sum_i \alpha_i \neq 1$ , we still have  $\tilde{Y}_t(0) \approx \sum_i \alpha_i \tilde{W}_{ti} + (\sum_i \alpha_i - 1) f(t)$ , that is, one may include an additional time-varying intercept to explicitly capture the time trend. Under the linear factor model of [Abadie et al.](#page-58-1) [\[2010\]](#page-58-1), this intercept vanishes, and thus the model remains correctly specified. Therefore, the treatment effect on the residual can still be interpreted as the treatment effect on the original outcome. The above discussion is meant to illustrate the interpretation of the detrended outcomes, but we do not make this assumption throughout the analysis. However, as shown in Section [S7.6,](#page-40-0) detrending might lead to slightly to moderately anti-conservative inference.

We detrend the Florida data by fitting a quadratic function of time to homicide rates. We model the time-varying average treatment effect  $\phi_{\lambda}(t)$  for the treated unit as a constant. We also model the outcome bridge function as a linear function  $h_{\alpha}: w \mapsto$  $(1, w^{\top})^{\top} \alpha$ , and select the homicide rates in the states of Arkansas, Maryland, New Jersey, and New York as the proxy  $W_t$ . This choice is motivated by Abadie's synthetic control method [\[Abadie et al., 2010\]](#page-58-1), for which weights are negligible for all but these four states. Our inference might be non-informative due to increased uncertainty associated with empirical selection of the proxy  $W_t$ . Proxy selection remains an open problem we are actively working on. Similar challenges are known to impact IV analysis with potentially invalid IVs. We use all other states as control units in  $Z_t$ . We consider the following two parametrizations of the treatment confounding bridge function, where the second has a

<span id="page-45-0"></span>

Figure S9: Figures similar to Figure [2](#page-14-0) with detrending.

larger set of states included in the model:

$$
q_{\beta}(Z_t) = \exp \{ \beta_0 + \beta_{\text{Delaware}} Z_{t, \text{Delaware}} \};
$$
  

$$
q_{\beta}(Z_t) = \exp \{ \beta_0 + \beta_{\text{Delaware}} Z_{t, \text{Delaware}} + \beta_{\text{Ohio}} Z_{t, \text{Ohio}} \}.
$$

We considered these two states in the treatment confounding bridge model due to their relative proximity and similarities to Florida. Thus, we hypothesize that these two states capture variation in the unmeasured confounder  $U_t$ . More states similar to Florida could have been included, had data from more control units been available.

The point estimates and 95% confidence intervals for the average treatment effect for the treated unit produced by the above methods are presented in Table [S1.](#page-46-1) The trajectories of synthetic controls and the actual residuals of log GDP per capita in Kansas are presented in Figure [S16a.](#page-52-0) We also conduct a falsification analysis of the placebo effect in October 2002. The results are presented in Table [S1](#page-46-1) and Figure [S16.](#page-52-0)

Because this time series is quite noisy, compared to typical data from econometric applications (e.g., data from Section [S8.2\)](#page-47-0), OLS does not yield a good pre-treatment fit. Abadie's SC appears to yield a reasonable pre-treatment fit because of the regularization, but the high noise level might still be a concern. The pre-treatment fits for proximal causal inference methods also appear reasonable and qualitatively similar to that of Abadie's SC. The point estimates of the effect of "stand your ground" law on homicide rates from all methods are positive. However, methods based on proximal causal inference involving the

<span id="page-46-0"></span>

Figure S10: Figures similar to Figure [3](#page-15-0) with detrending.

outcome confounding bridge function report an insignificant effect and much smaller point estimates than Abadie's SC, while other methods yield similar conclusions to Abadie's SC. Because of the concern about the noise level, the results from DR, DR2 and Outcome bridge might be more reliable. The methods based on weighting might not be reliable due to a lack of data from control states that are similar to Florida. These results suggest that, with a high noise level present, doubly robust methods may outperform OLS and Abadie's SC because doubly robust methods do not rely on a near-perfect pre-treatment fit.

<span id="page-46-1"></span>

Method	Florida's "stand your ground" law	placebo
Abadie's SC	0.083	$-0.025$
<b>OLS</b>	$0.066$ $(0.045, 0.086)$	$-0.032$ $(-0.062, -0.002)$
DR.	$0.024$ ( $-0.061$ , $0.108$ )	$-0.075$ $(-0.239, 0.088)$
DR <sub>2</sub>	$0.006$ ( $-0.099$ , $0.110$ )	$-0.042$ ( $-0.086$ , $0.002$ )
Outcome bridge	$0.012$ ( $-0.087, 0.112$ )	$-0.094$ $(-0.262, 0.075)$
Treatment bridge	$0.058$ $(0.038, 0.078)$	$-0.033$ $(-0.061, -0.005)$
Treatment bridge2	$0.057$ $(0.036, 0.078)$	$-0.039$ $(-0.092, 0.013)$

Table S1: Table similar to Table [1](#page-16-0) for Florida's "stand your ground" law.



Figure S11: Figures similar to Figure [2](#page-14-0) with post-selection.

#### <span id="page-47-0"></span>S8.2 Analysis of Kansas GDP

We also illustrate our methods by studying the effect of an aggressive tax cut in Kansas in the first quarter of 2012 on the logarithm of GDP per capita. This question has been studied in [Rickman and Wang](#page-62-0) [\[2018\]](#page-62-0) using the classical synthetic control method proposed by [Abadie et al.](#page-58-1) [\[2010,](#page-58-1) [2015\]](#page-58-2). In these data, economic outcomes are measured every quarter from 1990 to 2016 for all US states. GDP was measured in millions of U.S. dollars.

We preprocess the data, choose proxies  $(W_t, Z_t)$  and specify the confounding bridge functions  $(h^*, q^*)$  similarly to Section [S8.1.](#page-43-2) Specifically, the donors forming the proxy  $W_t$  are the states of North Dakota, South Carolina, Texas, and Washington as donors in  $W_t$ ; the quarterly log GDP per capita of all other control states form the proxy  $Z_t$ . We consider the following three parametrizations of the treatment confounding bridge function with increasing sets of states included in the model:

$$
q_{\beta}(Z_{t}) = \exp \{ \beta_{0} + \beta_{\text{Iowa}} Z_{t,\text{Iowa}} \};
$$
  
\n
$$
q_{\beta}(Z_{t}) = \exp \{ \beta_{0} + \beta_{\text{Iowa}} Z_{t,\text{Iowa}} + \beta_{\text{South Dakota}} Z_{t,\text{South Dakota}} \};
$$
  
\n
$$
q_{\beta}(Z_{t}) = \exp \{ \beta_{0} + \beta_{\text{Iowa}} Z_{t,\text{Iowa}} + \beta_{\text{South Dakota}} Z_{t,\text{South Dakota}} + \beta_{\text{Oklahoma}} Z_{t,\text{Oklahoma}} \}.
$$

These states are chosen because they are similar to Kansas and likely to capture the effect of the unmeasured confounder  $U_t$ .



Figure S12: Figures similar to Figure [3](#page-15-0) with post-selection.

The analysis results are presented in Table [S2](#page-52-1) and Figure [S17.](#page-52-2) All methods support a decrease in log GDP per capita due to the tax cut; the decrease is significant for all methods providing 95% confidence intervals (namely all methods except Abadie's SC); the only exception is the weighted method based on a single state included in the treatment confounding bridge model, treatment bridge. This result may be due to model misspecification, and the doubly robust method with the same treatment confounding bridge model is still significant. Point estimates of the average treatment effect for the treated unit from doubly robust methods are almost twice those of Abadie's SC and weighted methods. Doubly robust estimates are well within sampling variability based on the proximal outcome bridge estimates, suggesting that the outcome confounding bridge function may be correctly specified. The synthetic controls projected potential outcome trajectories of all methods have similar fits in the pre-treatment period; however, Abadie's SC and OLS appear to lead to a lower synthetic control trajectory in the post-treatment period.

We further conduct a falsification analysis of the placebo effect in the first quarter of 2008. The analysis results are presented in Table [S2](#page-52-1) and Figure [S17b.](#page-52-2) The 95% confidence intervals from all methods cover zero, correctly indicating a non-significant effect in the placebo period, with the exception of OLS and the weighted method with two states only included in the treatment confounding bridge model. The point estimates of doubly robust methods are also closer to zero. These results suggest a potentially superior performance of doubly robust methods and a potential bias of OLS.

<span id="page-49-1"></span>

Figure S13: Figures similar to Figure [2](#page-14-0) with nonstationarity and time-varying treatment effects.

# <span id="page-49-0"></span>S9 Existence of treatment confounding bridge function

In this appendix, we briefly state examples of sufficient conditions for Condition [5.](#page-8-1) Sufficient conditions for Conditions [S10](#page-24-0) and [S12](#page-26-0) are similar. Sufficient conditions for existence of various confounding bridge functions have been considered in [Miao et al.](#page-62-2) [\[2018\]](#page-62-2), [Cui](#page-60-4) [et al.](#page-60-4) [\[2020\]](#page-60-4) and [Shi et al.](#page-63-0) [\[2023\]](#page-63-0). In particular, sufficient conditions for Condition [2](#page-5-2) are presented in [Shi et al.](#page-63-0) [\[2023\]](#page-63-0). We refer readers to these works for more details. These conditions are only sufficient and not necessary.

#### S9.1 Discrete random variables

Fix any  $t_-\leq T_0$  and  $t_+ > T_0$ . Suppose that,  $(Z_{t_-}, U_{t_-})$  is a discrete random vector with finite support. Suppose that the supports of  $U_{t-}$  and  $Z_{t-}$  are  $\{u_1, \ldots, u_I\}$  and  $\{z_1, \ldots, z_J\}$ , respectively. Let  $P_{Z|U}$  be a  $J \times I$  matrix with the  $(j, i)$ -th entry being  $pr(Z_{t-} = z_j \mid U_{t-} = z_j)$ u<sub>i</sub>). Let  $R_U$  be the I-dimensional vector of likelihood ratios  $pr(U_{t_+} = u_i)/pr(U_{t_-} = u_i)$ . For any function  $q : \mathcal{Z} \to \mathbb{R}$ , we may equivalently represent  $z \mapsto q(z)$  by a J-dimensional vector Q with entries  $q(z_i)$ ,  $1 \leq j \leq J$ . Condition [5](#page-8-1) is then equivalent to the existence of



Figure S14: Figures similar to Figure [3](#page-15-0) with nonstationarity and time-varying treatment effects.

a solution in Q to

<span id="page-50-0"></span>
$$
P_{Z|U}^{\top} Q = R_U. \tag{S19}
$$

Therefore, a sufficient condition for Condition [5](#page-8-1) is that  $P_{Z|U}$  has full column rank, namely rank( $P_{Z|U}$ ) = I, and the solution to [\(S19\)](#page-50-0) is identical for all  $t_-\leq T_0$ , which is implied by stationarity of  $(Z_{t-}, U_{t-})$   $(t_-\leq T_0)$ .

#### S9.2 Continuous random variables

We first fix any  $t_-\leq T_0$  and  $t_+ > T_0$  in this appendix. Suppose that  $(Z_{t_-}, U_{t_-})$  is a continuous random vector. For any distribution P, recall that  $L^2(P)$  denotes the space of all square-integrable functions with respect to  $P$ , which is a Hilbert space equipped with inner product  $\langle f, g \rangle = \int fg \ dP$ . Let  $K_{t-} : L^2(P_{Z_{t-}}) \to L^2(P_{U_{t-}})$  be the operator defined pointwise by  $K_{t-}q: u \mapsto E\{q(Z_{t-}) \mid U_{t-} = u\}$  for  $q \in L^2(P_{Z_{t-}})$ . We assume the following regularity conditions:

<span id="page-50-1"></span>**Condition S24.** For all  $t_{-} \leq T_0$ ,  $\iint p_{Z_{t-}|U_{t-}}(z \mid u)p_{U_{t-}|Z_{t-}}(u \mid z) dz du < \infty$ .

By Example 2.3 in [Carrasco et al.](#page-59-9) [\[2007\]](#page-59-9) (pages 5656 and 5659), under Condi-tion [S24,](#page-50-1) the operator  $K_{t-}$  is compact [see, e.g., Definition 2.17, [Kress, 2014\]](#page-62-11). Let  $(\lambda_{t_{-},m}, \varphi_{t_{-},m}, \psi_{t_{-},m})_{m=1}$  be a singular system of  $K_{t_{-}}$  [see, e.g., Theorem 15.16 of [Kress,](#page-62-11) [2014,](#page-62-11) for more about singular systems].

<span id="page-51-0"></span>

Figure S15: Ratio of the standard error to the standard deviation of the estimator's sampling distribution with nonstationarity and time-varying treatment effects.

<span id="page-51-2"></span>Condition S25. For all  $t_-\leq T_0$ ,

$$
\int \left\{ \frac{\mathrm{d}P_{U_{t_+}}}{\mathrm{d}P_{U_{t_-}}}(u) \right\}^2 p_{U_{t_-}}(u) \, \mathrm{d}u < \infty.
$$

Condition S26. For all  $t_-\leq T_0$ ,

$$
\sum_{m=1}^{\infty}\lambda_{t_{-},m}^{-2}\left|\left\langle \frac{\mathrm{d}P_{U_{t_{+}}}}{\mathrm{d}P_{U_{t_{-}}}},\psi_{t_{-},m}\right\rangle \right|^{2}<\infty.
$$

The next assumption is a completeness condition, which is similar to Conditions [S1](#page-19-0) and [S2.](#page-19-2)

<span id="page-51-1"></span>**Condition S27.** Let  $g \in L^2(P_{U_{t-}})$ . For all  $t_{-} \leq T_0$ , the following two statements are equivalent: (i)  $g(U_{t-}) = 0$ , (ii)  $E\{g(U_{t-}) | Z_{t-}\} = 0$ .

Under Conditions [S24–](#page-50-1)[S27,](#page-51-1) Condition [5](#page-8-1) holds for the fixed  $t_$ ; that is, a solution to [\(3\)](#page-8-0) exists for the fixed t\_. This claim can be proved by Picard's Theorem [Theorem 15.18 of [Kress, 2014\]](#page-62-11). We next sketch the proof. The orthogonal complement  $N(K_{t-}^*)^{\perp}$  of the nullspace of the adjoint of  $K_{t-}$  equals  $L^2(P_{Z_{t-}})$  by Condition [S27.](#page-51-1) By Condition [S25,](#page-51-2)  $dP_{U_{t_+}}/dP_{U_{t_-}}$  lies in  $N(K_{t_-}^*)^{\perp}$ . The desired existence result then follows by directly applying Picard's Theorem. Condition [5](#page-8-1) then holds if the solution to [\(3\)](#page-8-0) is identical for all  $t_-\leq T_0$ .

<span id="page-52-0"></span>



(b) Placebo

<span id="page-52-1"></span>

Method	Kansas tax cut	placebo
Abadie's SC	$-0.048$	0.029
OLS	$-0.069$ $(-0.087, -0.050)$	$0.026~(2.6\times10^{-6}, 0.052)$
DR.	$-0.077$ $(-0.126, -0.028)$	$0.004$ ( $-0.068$ , $0.077$ )
DR <sub>2</sub>	$-0.095$ $(-0.147, -0.043)$	$-0.005$ $(-0.039, 0.030)$
DR <sub>3</sub>	$-0.103$ $(-0.228, -0.021)$	$-0.007$ $(-0.059, 0.046)$
Outcome bridge	$-0.104$ $(-0.150, -0.058)$	$0.012$ ( $-0.069$ , $0.093$ )
Treatment bridge	$-0.031$ $(-0.087, 0.024)$	$-0.028$ $(-0.063, 0.008)$
Treatment bridge2	$-0.017$ $(-0.032, -0.002)$	$-0.042$ $(-0.056, -0.0027)$
Treatment bridge3	$-0.016$ $(-0.029, -0.003)$	$-0.048$ $(-0.097, 0.001)$

Table S2: Table similar to Table [1](#page-16-0) for tax cut in Kansas.

<span id="page-52-2"></span>

Figure S17: Figures similar to Figure [4](#page-17-0) for tax cut in Kansas.

# S10 Proofs

### S10.1 Proof of Theorem [1](#page-9-0)

We first prove [\(4\)](#page-9-4). Let  $f : \mathbb{R} \to \mathbb{R}$  be any square-integrable function. Then,

$$
E[q^*(Z_{t-})f\{Y_{t-}(0)\}] = E(E[q^*(Z_{t-})f\{Y_{t-}(0)\} | U_{t-}])
$$
  
\n
$$
= E(E\{q^*(Z_{t-}) | U_{t-}\}E[f\{Y_{t-}(0)\} | U_{t-}])
$$
(Condition 1)  
\n
$$
= E\left(\frac{dP_{U_{t+}}}{dP_{U_{t-}}}(U_{t-})E[f\{Y_{t-}(0)\} | U_{t-}]\right)
$$
(Condition 4 & 5)  
\n
$$
= \int_{\mathcal{U}} \frac{dP_{U_{t+}}}{dP_{U_{t-}}}(u)E[f\{Y_{t-}(0)\} | U_{t-} = u]dP_{U_{t-}}(u)
$$
  
\n
$$
= \int_{\mathcal{U}} E[f\{Y_{t+}(0)\} | U_{t+} = u]dP_{U_{t+}}(u)
$$
(Condition 3)  
\n
$$
= E[f\{Y_{t+}(0)\}].
$$

Thus, [\(4\)](#page-9-4) has been proved and [\(5\)](#page-9-2) follows immediately. We next prove [\(6\)](#page-9-1). Let  $f : \mathcal{W} \to$ R be any square-integrable function.

$$
E\{q^*(Z_{t-})f(W_{t-})\} = E[E\{q^*(Z_{t-})f(W_{t-}) | U_{t-}\}]
$$
  
\n
$$
= E[E\{q^*(Z_{t-}) | U_{t-}\}E\{f(W_{t-}) | U_{t-}\}]
$$
 (Condition 1)  
\n
$$
= E\left[\frac{dP_{U_{t+}}}{dP_{U_{t-}}}(U_{t-})E\{f(W_{t-}) | U_{t-}\}\right]
$$
 (Conditions 4 & 5)  
\n
$$
= \int_{\mathcal{U}} \frac{dP_{U_{t+}}}{dP_{U_{t-}}}(u)E\{f(W_{t-}) | U_{t-} = u\} dP_{U_{t-}}(u)
$$
  
\n
$$
= \int_{\mathcal{U}} E\{f(W_{t+}) | U_{t+} = u\} dP_{U_{t+}}(u) = E\{f(W_{t+})\}.
$$
 (Condition 3)

For a Borel set  $B \subseteq U$ , take  $f(w) = \mathbb{1}(w \in B)$ . If  $P_{W_{t-}}(B) = 0$ , the above equality implies that  $P_{W_{t+}}(B) = E\{q^*(Z_{t-})\mathbb{1}(W_{t-} \in B)\} = 0$ . Therefore,  $P_{W_{t+}}$  is dominated by  $P_{W_{t-}}$  and the Radon-Nikodym derivative  $dP_{W_{t+}}/dP_{W_{t-}}$  is well defined. Thus, for any integrable function  $f$ ,

$$
E\{q^*(Z_{t-})f(W_{t-})\} = E\left\{\frac{\mathrm{d}P_{W_{t+}}}{\mathrm{d}P_{W_{t-}}}(W_{t-})f(W_{t-})\right\},\,
$$

that is,

$$
E\left[\left\{q^*(Z_{t-}) - \frac{\mathrm{d}P_{W_{t+}}}{\mathrm{d}P_{W_{t-}}}(W_{t-})\right\}f(W_{t-})\right] = 0.
$$

Since  $f$  is arbitrary, we have that

$$
E\left\{q^*(Z_{t-}) - \frac{\mathrm{d}P_{W_{t+}}}{\mathrm{d}P_{W_{t-}}}(w) \mid W_{t-} = w\right\} = 0
$$

for  $P_{W_{t-}}$ -a.e.  $w \in \mathcal{W}$ . Equation [6](#page-9-1) follows.

We finally prove the uniqueness of  $q^*$  under Condition [S2.](#page-19-2) Suppose that two functions  $q_1$  and  $q_2$  both solve [\(6\)](#page-9-1). Then,  $E\{q_1(Z_{t-}) - q_2(Z_{t-}) | W_{t-}\} = 0$ , and thus  $q_1(Z_{t-}) =$  $q_2(Z_{t-})$  by Condition [S2.](#page-19-2)

#### S10.2 Proof of Theorem [2](#page-10-0)

We study the two cases where  $h = h^*$  and  $q = q^*$  separately. If Condition [2](#page-5-2) holds and  $h = h^*$ , then, by [\(1\)](#page-6-2) and [\(2\)](#page-6-0),

<span id="page-54-0"></span>
$$
E[q(Z_{t-})\{Y_{t-} - h(W_{t-})\} + h(W_{t+})] = E\{Y_{t+}(0)\}
$$
\n(S20)

and

<span id="page-54-1"></span>
$$
E[Y_{t_{+}} - q(Z_{t_{-}})\{Y_{t_{-}} - h(W_{t_{-}})\} - h(W_{t_{+}})] = E\{Y_{t_{+}}(1) - Y_{t_{+}}(0)\} = \phi^{*}(t_{+}),
$$
(S21)

as desired.

If Condition [5](#page-8-1) holds and  $q = q^*$ , then by a similar argument as in the proof of Theorem [1,](#page-9-0)

$$
E[q^*(Z_{t-})\{Y_{t-}(0) - h(W_{t-})\}] = E(E[q^*(Z_{t-})\{Y_{t-}(0) - h(W_{t-})\} | U_{t-}])
$$
  
\n
$$
= E[E\{q^*(Z_{t-}) | U_{t-}\} E\{Y_{t-}(0) - h(W_{t-}) | U_{t-}\}]
$$
 (Condition 1)  
\n
$$
= \int_{\mathcal{U}} \frac{dP_{U_{t+}}}{dP_{U_{t-}}} (u) E\{Y_{t-}(0) - h(W_{t-}) | U_{t-} = u\} P_{U_{t-}} (du)
$$
 (Conditions 4 & 5)  
\n
$$
= \int_{\mathcal{U}} E\{Y_{t+}(0) - h(W_{t+}) | U_{t+} = u\} P_{U_{t+}} (du) = E\{Y_{t+}(0) - h(W_{t+})\}.
$$
 (Condition 3)

Therefore, [\(S20\)](#page-54-0) and [\(S21\)](#page-54-1) hold, as desired. We have proved Theorem [2.](#page-10-0)

#### S10.3 Proof of Theorems [S1–](#page-25-1)[S4](#page-26-4)

The proofs of these theorems are similar to those of Theorems [1](#page-9-0) and [2](#page-10-0) and thus we abbreviate the presentation.

*Proof of Theorem [S1.](#page-25-1)* By Condition [3,](#page-7-0) for any square-integrable functions f and g,  $E[f\{Y_t(0)\}]$  $U_t = u$  and  $E[g(W_t) | U_t = u]$  do not depend on t. We first prove [\(S3\)](#page-25-3):

$$
\sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+}) E[f\{Y_{t_{+}}(0)\}] = \sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+}) E(E[f\{Y_{t_{+}}(0)\} | U_{t_{+}}])
$$

$$
= \frac{1}{T_0} \sum_{t=1}^{T_0} E[q_T^*(Z_{t-})E\{f(Y_{t-}) | U_{t-}\}] = \frac{1}{T_0} \sum_{t=1}^{T_0} E\{q_T^*(Z_{t-})f(Y_{t-})\}. \quad \text{(Conditions S10 and 1)}
$$

Then, [\(S4\)](#page-25-4) follows by taking f to be the identity function in [\(S3\)](#page-25-3). We next show that  $q_T^*$ is a solution to  $(S5)$ . For any square-integrable function  $g$ ,

$$
\frac{1}{T_0} \sum_{t_{-1}}^{T_0} E\{q(Z_{t_{-}})g(W_{t_{-}})\} = \frac{1}{T_0} \sum_{t_{-1}}^{T_0} E[q(Z_{t_{-}})E\{g(W_{t_{-}}) | U_{t_{-}}\}] \text{ (Condition 1)}
$$
\n
$$
= \sum_{t_{+}=T_0+1}^{T} \ell_T(t_{+}) E[E\{g(W_{t_{+}}) | U_{t_{+}}\}] = \sum_{t_{+}=T_0+1}^{T} \ell_T(t_{+}) E\{g(W_{t_{+}})\}.
$$

Hence,  $q_T^*$  is a solution to [\(S5\)](#page-25-0). We finally prove the uniqueness of the solution to (S5). Suppose that two functions  $q_{T,1}$  and  $q_{T,2}$  are solutions to [\(S5\)](#page-25-0). Then, for any squareintegrable function g,

$$
\frac{1}{T_0} \sum_{t_{-}=1}^{T_0} E[\{q_{T,1}(Z_{t_{-}}) - q_{T,2}(Z_{t_{-}})\} g(W_{t_{-}})] = 0.
$$

By Condition [S11,](#page-24-2)  $q_{T,1}(Z_{t-}) - q_{T,2}(Z_{t-}) = 0$  for all  $t_{-} \leq T_0$  and thus the solution is unique almost surely.  $\Box$ 

*Proof of Theorem [S2.](#page-25-5)* First consider the case where Condition [2](#page-5-2) holds and  $h = h^*$ . In this case, using the identification result in [Shi et al.](#page-63-0) [\[2023\]](#page-63-0) (Theorem 4), we have that

$$
\frac{1}{T_0} \sum_{t_{-1}}^{T_0} E \left[ \underbrace{q(Z_{t_{-}})} \{Y_{t_{-}} - h(W_{t_{-}})\} \right] + \sum_{t_{+} = T_0 + 1}^{T} \ell_T(t_{+}) E \{h(W_{t_{+}})\}
$$
\n
$$
= \frac{1}{T_0} \sum_{t_{-} = 1}^{T_0} E \left[ E \{ \underbrace{q(Z_{t_{-}})} | U_{t_{-}} \} E \{Y_{t_{-}} - h(W_{t_{-}}) | U_{t_{-}} \} \right] + \sum_{t_{+} = T_0 + 1}^{T} \ell_T(t_{+}) E \{h(W_{t_{+}})\}
$$
\n
$$
= 0 + \sum_{t_{+} = T_0 + 1}^{T} \ell_T(t_{+}) E \{Y_{t_{+}}(0)\}.
$$

The second line follows by Condition [1.](#page-5-0) Therefore,  $\sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+}) E\{Y_{t_{+}}(0)\}$  is identified as in [\(S6\)](#page-25-2). Next suppose that Condition [S10](#page-24-0) holds and  $q = q_T^*$ . Using Theorem [S2,](#page-25-5) we have that

$$
\frac{1}{T_0} \sum_{t=1}^{T_0} E\left[q(Z_{t-})\{Y_{t-} - h(W_{t-})\}\right] + \sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E[h(W_{t+})]
$$

$$
= \frac{1}{T_0} \sum_{t=1}^{T_0} E\left\{ \underline{q}(Z_{t-})Y_{t-} \right\} - \frac{1}{T_0} \sum_{t=1}^{T_0} E\left\{ \underline{q}(Z_{t-})h(W_{t-}) \right\} + \sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E\{h(W_{t+}) \}
$$
  
= 
$$
\sum_{t_+ = T_0 + 1}^{T} \ell_T(t_+) E\{Y_{t+}(0)\} + 0.
$$

Therefore,  $\sum_{t_{+}=T_{0}+1}^{T} \ell_{T}(t_{+}) E\{Y_{t_{+}}(0)\}\)$  is identified as in [\(S6\)](#page-25-2). Finally, the identification formula for  $\phi_T^*$  in [\(S6\)](#page-25-2) follows by noting that

$$
\phi_T^* = \sum_{t_+ = T_0 + 1}^T \ell_T(t_+) E\{Y_{t_+} - Y_{t_+}(0)\}.
$$

<span id="page-56-0"></span> $\Box$ 

The proof of Theorems [S3](#page-26-5) and [S4](#page-26-4) is almost identical and thus omitted.

#### S10.4 Proof of Theorems [3](#page-11-1) & [S6](#page-30-2)

Theorem [3](#page-11-1) follows immediately from standard estimation theory of the generalized method of moments, for example, Theorem 7.1 and 7.2 in [Wooldridge](#page-64-3) [\[1994\]](#page-64-3), along with Theorem [2.](#page-10-0) The proof of Theorem [S6](#page-30-2) is similar.

#### S10.5 Proof of Theorems [S5](#page-29-0) & [S7](#page-31-1)

The proofs of Theorems [S5](#page-29-0) & [S7](#page-31-1) are almost identical. Therefore, we present the proof of Theorem [S5](#page-29-0) and omit the proof of Theorem [S7.](#page-31-1) The argument is inspired by the theory of [see, e.g., [Wooldridge, 1994,](#page-64-3) [Hall, 2007\]](#page-61-7) with adaptations to our case where the moment equation might not be centered at each time period.

*Proof of Theorem [S5.](#page-29-0)* Under Conditions [1,](#page-5-0) [3](#page-7-0) and [S14,](#page-28-0) we have that  $\phi_{\infty} = \phi^*$  by the definition of  $\mathcal{G}_{T,t}$  in [\(S12\)](#page-28-2) and Theorem [S4.](#page-26-4) We first prove consistency. By Con-ditions [S3](#page-20-1) and [S7,](#page-21-1) we have that, as  $T \to \infty$ , letting  $V_T(\theta) = \sum_{t=1}^T \frac{G_{T,t}(\theta)}{T}$  and  $\bar{V}_{T}(\theta) = \sum_{t=1}^{T'} E[G_{T',t}(\theta)]/T',$ 

$$
\sup_{\underline{\theta}\in\Theta} \left| V_T(\underline{\theta})^\top \mathfrak{Q}_T V_T(\underline{\theta}) - \lim_{T'\to\infty} \bar{V}_{T'}(\underline{\theta})^\top \mathfrak{Q} \lim_{T'\to\infty} \bar{V}_{T'} \right| \text{ converge to zero in probability } (S22)
$$

Let  $\epsilon > 0$  be an arbitrary positive constant. By [\(S22\)](#page-56-0) and the definition of  $\hat{\ell}_T$ , we have that, with probability tending to one,

$$
\left| V_T(\underline{\theta}_{\infty})^{\top} \mathfrak{Q}_T V_T(\underline{\theta}_{\infty}) - \lim_{T' \to \infty} \bar{V}_{T'}(\underline{\theta}_{\infty})^{\top} \mathfrak{Q} \lim_{T' \to \infty} \bar{V}_{T'}(\underline{\theta}_{\infty}) \right| < \epsilon/2,
$$

$$
\left| V_T(\underline{\theta}_T)^\top \Omega_T V_T(\underline{\theta}_T) - \lim_{T' \to \infty} \bar{V}_{T'}(\hat{\underline{\theta}}_T)^\top \Omega \lim_{T' \to \infty} \bar{V}_{T'}(\hat{\underline{\theta}}_T) \right| < \epsilon/2,
$$
  

$$
V_T(\underline{\theta}_T)^\top \Omega_T V_T(\underline{\theta}_T) \leq V_T(\underline{\theta}_\infty)^\top \Omega_T V_T(\underline{\theta}_\infty).
$$

Combining these three inequalities, we have that, with probability tending to one,

$$
\lim_{T' \to \infty} \bar{V}_{T'}(\hat{\theta}_T)^{\top} \Omega \lim_{T' \to \infty} \bar{V}_{T'}(\hat{\theta}_T) < \lim_{T' \to \infty} \bar{V}_{T'}(\theta_\infty)^{\top} \Omega \lim_{T' \to \infty} \bar{V}_{T'}(\theta_\infty) + \epsilon = \epsilon,\tag{S23}
$$

where the last equality follows from Condition [S14.](#page-28-0)

Let  $N \subseteq \mathcal{Q}$  be an arbitrary open set containing  $\ell_{\infty}$ . By Condition [S4,](#page-21-3)  $\mathcal{Q} \setminus A$  is e compact. By Conditions [S14,](#page-28-0) [S3](#page-20-1) and [S5,](#page-21-0)

<span id="page-57-2"></span><span id="page-57-0"></span>
$$
\inf_{\underline{\theta}\in\Theta\backslash N}\lim_{T'\to\infty}\bar{V}_{T'}(\underline{\theta})^{\top}\Omega\lim_{T'\to\infty}\bar{V}_{T'}
$$

exists and is strictly positive. Taking  $\epsilon$  in [\(S23\)](#page-57-0) to be the above this infimum, we have that, with probability tending to one,

$$
\lim_{T' \to \infty} \bar{V}_{T'}(\hat{\theta}_T)^\top \Omega \lim_{T' \to \infty} \bar{V}_{T'}(\hat{\theta}_T) < \inf_{\underline{\theta} \in \underline{\Theta} \backslash N} \lim_{T' \to \infty} \bar{V}_{T'}(\underline{\theta})^\top \Omega \lim_{T' \to \infty} \bar{V}_{T'}.
$$

This event implies that  $\hat{\theta}_{T} \in N$ . Since N is arbitrary, we have shown that  $\hat{\theta}_{T}$  converges to  $\ell_{\infty}$  in probability as  $T \to \infty$ . We next prove the asymptotic normality of  $\hat{\ell}_{T}$ . Under Condition [S8,](#page-22-0) by a first-order Taylor expansion and the above consistency result, we have that

<span id="page-57-1"></span>
$$
\frac{1}{T} \sum_{t=1}^{T} G_{T,t}(\hat{\theta}_T) = \frac{1}{T} \sum_{t=1}^{T} G_{T,t}(\theta_\infty) + \frac{1}{T} \sum_{t=1}^{T} \nabla_{\theta} G_{T,t}(\theta) |_{\theta = \theta_\infty} (\hat{\theta}_T - \theta_\infty) + o_p(||\hat{\theta}_T - \theta_\infty||). \tag{S24}
$$

The fact that  $\hat{\ell}_{T}$  is a minimizer together with Condition [S8](#page-22-0) implies that

$$
0 = \nabla_{\underline{\theta}} \left\{ V_T(\underline{\theta})^\top \mathfrak{Q}_T V_T(\underline{\theta}) \right\} \Big|_{\underline{\theta} = \hat{\underline{\theta}}_T} = 2 \left\{ \frac{1}{T} \sum_{t=1}^T \nabla_{\underline{\theta}} \mathcal{Q}_{T,t}(\underline{\theta}) \Big|_{\underline{\theta} = \hat{\underline{\theta}}_T} \right\}^\top \mathfrak{Q}_T V_T(\underline{\theta}_T). \tag{S25}
$$

By Conditions [S7–](#page-21-1)[S9,](#page-22-1) using [\(S24\)](#page-57-1) and [\(S25\)](#page-57-2), we have that

$$
0 = T^{1/2} \left\{ \frac{1}{T} \sum_{t=1}^{T} \nabla_{\underline{\theta}} \underline{G}_{T,t}(\underline{\theta}) |_{\underline{\theta} = \hat{\underline{\theta}}_T} \right\}^{\top} \underline{\Omega}_T V_T(\underline{\theta}_T)
$$
  
=  $\underline{R}^{\top} \underline{\Omega} \frac{1}{T^{1/2}} \sum_{t=1}^{T} \underline{G}_{T,t}^q(\underline{\theta}_{\infty}) + \underline{R}^{\top} \underline{\Omega} \underline{R} T^{1/2}(\hat{\underline{\theta}}_T - \underline{\theta}_{\infty}) + o_p(T^{1/2} || \hat{\theta}_T - \theta_{\infty} || + 1).$ 

By Condition [S9](#page-22-1) and Slutsky's Theorem, we have that

$$
T^{1/2}(\hat{\theta}_{T} - \theta_{\infty}) = -(R^{\top} \Omega R)^{-1} R^{\top} \Omega \frac{1}{T^{1/2}} \sum_{t=1}^{T} G_{T,t}^{q}(\theta_{\infty}) + o_{p}(1)
$$

$$
= -A^{-1} R^{\top} \Omega \frac{1}{T^{1/2}} \sum_{t=1}^{T} G_{T,t}^{q}(\theta_{\infty}) + o_{p}(1),
$$

which converges in distribution to  $N(0, A^{-1}BA^{-1})$ . Here, the existence of  $A^{-1}$  follows from Conditions [S3](#page-20-1) and [S6.](#page-21-2)  $\Box$ 

# References

- <span id="page-58-0"></span>Alberto Abadie and Javier Gardeazabal. The economic costs of conflict: A case study of the Basque country. American Economic Review, 93(1):113–132, 2003. ISSN 00028282. doi: 10.1257/000282803321455188.
- <span id="page-58-3"></span>Alberto Abadie and Jérémy L'Hour. A Penalized Synthetic Control Estimator for Disaggregated Data. Journal of the American Statistical Association, 116(536):1817–1834, 2021. ISSN 1537274X. doi: 10.1080/01621459.2021.1971535.
- <span id="page-58-1"></span>Alberto Abadie, Alexis Diamond, and And Jens Hainmueller. Synthetic control methods for comparative case studies: Estimating the effect of California's Tobacco control program. Journal of the American Statistical Association, 105(490):493–505, 2010. ISSN 01621459. doi: 10.1198/jasa.2009.ap08746.
- <span id="page-58-2"></span>Alberto Abadie, Alexis Diamond, and Jens Hainmueller. Comparative Politics and the Synthetic Control Method. American Journal of Political Science, 59(2):495–510, 2015. ISSN 15405907. doi: 10.1111/ajps.12116.
- <span id="page-58-6"></span>Donald W.k. Andrews. Laws of large numbers for dependent non-identically distributed random variables. Econometric Theory, 4(3):458–467, 1988. ISSN 14694360. doi: 10.1017/S0266466600013396.
- <span id="page-58-5"></span>Donald W.K. Andrews. Examples of L2-complete and boundedly-complete distributions. Journal of Econometrics, 199(2):213–220, 2017. ISSN 18726895. doi: 10.1016/j.jeconom.2017.05.011.
- <span id="page-58-7"></span>Miguel Angel Arcones and Bin Yu. Central Limit Theorems for Empirical and U-Processes of Stationary Mixing Sequences. Journal of Theoretical Probability, 7(1):47–71, 1994.
- <span id="page-58-4"></span>Dmitry Arkhangelsky, Susan Athey, David A. Hirshberg, Guido W. Imbens, and Stefan Wager. Synthetic Difference-in-Differences. American Economic Review, 111(12):4088– 4118, 2021. ISSN 0002-8282. doi: 10.1257/aer.20190159.
- <span id="page-59-4"></span>Susan Athey, Mohsen Bayati, Nikolay Doudchenko, Guido Imbens, and Khashayar Khosravi. Matrix Completion Methods for Causal Panel Data Models. Journal of the American Statistical Association, 116(536):1716–1730, 2021. ISSN 1537274X. doi: 10.1080/01621459.2021.1891924.
- <span id="page-59-5"></span>Jushan Bai and Serena Ng. Matrix Completion, Counterfactuals, and Factor Analysis of Missing Data. Journal of the American Statistical Association, 116(536):1746–1763, 2021. ISSN 1537274X. doi: 10.1080/01621459.2021.1967163.
- <span id="page-59-7"></span>Heejung Bang and James M. Robins. Doubly robust estimation in missing data and causal inference models. Biometrics, 61(4):962–973, 2005. ISSN 15410420. doi: 10. 1111/j.1541-0420.2005.00377.x.
- <span id="page-59-3"></span>Eli Ben-Michael, Avi Feller, and Jesse Rothstein. Synthetic Controls with Staggered Adoption. Technical report, 2021a.
- <span id="page-59-0"></span>Eli Ben-Michael, Avi Feller, and Jesse Rothstein. The Augmented Synthetic Control Method. Journal of the American Statistical Association, 116(536):1789–1803, 2021b. ISSN 1537274X. doi: 10.1080/01621459.2021.1929245. URL [https://www.](https://www.tandfonline.com/action/journalInformation?journalCode=uasa20) [tandfonline.com/action/journalInformation?journalCode=uasa20](https://www.tandfonline.com/action/journalInformation?journalCode=uasa20).
- <span id="page-59-1"></span>Carl Bonander, David Humphreys, and Michelle Degli Esposti. Synthetic Control Methods for the Evaluation of Single-Unit Interventions in Epidemiology: A Tutorial. American Journal of Epidemiology, 190(12):2700–2711, 2021. ISSN 14766256. doi: 10.1093/aje/kwab211.
- <span id="page-59-8"></span>Richard C. Bradley and Cristina Tone. A Central Limit Theorem for Non-stationary Strongly Mixing Random Fields. Journal of Theoretical Probability, 30(2):655–674, 2017. ISSN 15729230. doi: 10.1007/s10959-015-0656-2.
- <span id="page-59-2"></span>Christian A.W. Bruhn, Stephen Hetterich, Cynthia Schuck-Paim, Esra Kürüm, Robert J. Taylor, Roger Lustig, Eugene D. Shapiro, Joshua L. Warren, Lone Simonsen, and Daniel M. Weinberger. Estimating the population-level impact of vaccines using synthetic controls. Proceedings of the National Academy of Sciences of the United States of America, 114(7):1524–1529, 2017. ISSN 10916490. doi: 10.1073/pnas.1612833114.
- <span id="page-59-9"></span>Marine Carrasco, Jean Pierre Florens, and Eric Renault. Linear Inverse Problems in Structural Econometrics Estimation Based on Spectral Decomposition and Regularization. Handbook of Econometrics, 6(SUPPL. PART B):5633–5751, 2007. ISSN 15734412. doi: 10.1016/S1573-4412(07)06077-1.
- <span id="page-59-6"></span>Matias D. Cattaneo, Yingjie Feng, and Rocio Titiunik. Prediction Intervals for Synthetic Control Methods. Journal of the American Statistical Association, 116(536):1865–1880, 2021. ISSN 1537274X. doi: 10.1080/01621459.2021.1979561.
- <span id="page-60-9"></span>Gary Chamberlain. Efficiency Bounds for Semiparametric Regression. *Econometrica*, 60 (3):567, 1992. ISSN 00129682. doi: 10.2307/2951584.
- <span id="page-60-7"></span>Xiaohong Chen, Han Hong, and Alessandro Tarozzi. Semiparametric efficiency in GMM models with auxiliary data. Annals of Statistics, 36(2):808–843, 2008. ISSN 00905364. doi: 10.1214/009053607000000947.
- <span id="page-60-1"></span>Victor Chernozhukov, Kaspar Wüthrich, and Yinchu Zhu. An Exact and Robust Conformal Inference Method for Counterfactual and Synthetic Controls. Journal of the American Statistical Association, 116(536):1849–1864, 2021. ISSN 1537274X. doi: 10.1080/01621459.2021.1920957.
- <span id="page-60-4"></span>Yifan Cui, Xu Shi, and Wang Miao. Semiparametric proximal causal inference. arXiv preprint arXiv:2011.08411v1, 2020.
- <span id="page-60-5"></span>Flavio Cunha, James J. Heckman, and Susanne M. Schennach. Estimating the Technology of Cognitive and Noncognitive Skill Formation. Econometrica, 78(3):883–931, 2010. ISSN 0012-9682. doi: 10.3982/ecta6551.
- <span id="page-60-10"></span>James Davidson. A central limit theorem for globally nonstationary near-epoch dependent functions of mixing processes. Econometric Theory, 8(3):313–329, 1992. ISSN 14694360. doi: 10.1017/S0266466600012950.
- <span id="page-60-2"></span>Ben Deaner. Proxy Controls and Panel Data. arXiv preprint arXiv:1810.00283v7, 2018. doi: 10.48550/arxiv.1810.00283.
- <span id="page-60-3"></span>Ben Deaner. Many Proxy Controls. arXiv preprint arXiv:2110.03973v1, 2021.
- <span id="page-60-11"></span>Herold Dehling and Walter Philipp. Empirical Process Techniques for Dependent Data. In *Empirical Process Techniques for Dependent Data*, pages 3–113. Birkhäuser, Boston, MA, 2002. doi: 10.1007/978-1-4612-0099-4<sub>-1</sub>.
- <span id="page-60-0"></span>Nikolay Doudchenko and Guido W. Imbens. Balancing, Regression, Difference-In-Differences and Synthetic Control Methods: A Synthesis. Technical report, National Bureau of Economic Research, Cambridge, MA, 2016.
- <span id="page-60-8"></span>Bruno Ferman and Cristine Pinto. Synthetic controls with imperfect pretreatment fit. Quantitative Economics, 12(4):1197–1221, 2021. ISSN 1759-7323. doi: 10.3982/qe1596.
- <span id="page-60-6"></span>Joachim Freyberger. Non-parametric panel data models with interactive fixed effects. Review of Economic Studies, 85(3):1824–1851, 2018. ISSN 1467937X. doi: 10.1093/ restud/rdx052.
- <span id="page-61-3"></span>Sander Greenland. Interpretation and choice of effect measures in epidemiologic analyses. American Journal of Epidemiology, 125(5):761–768, 1987. ISSN 00029262. doi: 10. 1093/oxfordjournals.aje.a114593.
- <span id="page-61-1"></span>Jinyong Hahn. On the Role of the Propensity Score in Efficient Semiparametric Estimation of Average Treatment Effects. Econometrica, 66(2):315, 1998. ISSN 00129682. doi: 10.2307/2998560.
- <span id="page-61-5"></span>Jinyong Hahn and Ruoyao Shi. Synthetic control and inference. Econometrics, 5(4):52, 2017. ISSN 22251146. doi: 10.3390/econometrics5040052.
- <span id="page-61-7"></span>Alastair R. Hall. Generalized Method of Moments. OUP Oxford, 2007. ISBN 9780470996249. doi: 10.1002/9780470996249.ch12.
- <span id="page-61-6"></span>Lars Peter Hansen. Large Sample Properties of Generalized Method of Moments Estimators. Econometrica, 50(4):1029, jul 1982. ISSN 00129682. doi: 10.2307/1912775.
- <span id="page-61-8"></span>Lars Peter Hansen. A method for calculating bounds on the asymptotic covariance matrices of generalized method of moments estimators. Journal of Econometrics, 30(1-2): 203–238, 1985. ISSN 03044076. doi: 10.1016/0304-4076(85)90138-1.
- <span id="page-61-11"></span>Lars Peter Hansen, John Heaton, and Amir Yaron. Finite-sample properties of some alternative GMM estimators. Journal of Business and Economic Statistics, 14(3):262– 280, 1996. ISSN 15372707. doi: 10.1080/07350015.1996.10524656.
- <span id="page-61-9"></span>Trevor Hastie, Jerome Friedman, and Robert Tibshirani. The elements of statistical learning: data mining, inference, and prediction. Springer Series in Statistics. Springer New York, New York, NY, 2009. ISBN 978-1-4899-0519-2. doi: 10.1007/978-0-387-21606-5.
- <span id="page-61-10"></span>Norbert Herrndorf. A functional central limit theorem for weakly dependent sequences of random variables. The Annals of Probability, 12(1):141–153, 1984.
- <span id="page-61-0"></span>Douglas Holtz-Eakin, Whitney Newey, and Harvey S. Rosen. Estimating Vector Autoregressions with Panel Data. Econometrica, 56(6):1371, 1988. ISSN 00129682. doi: 10.2307/1913103.
- <span id="page-61-4"></span>Cheng Hsiao, H. Steve Ching, and Shui Ki Wan. A panel data approach for program evaluation: Measuring the benefits of political and economic integration of Hong Kong with Mainland China. *Journal of Applied Econometrics*, 27(5):705–740, 2012. ISSN 08837252. doi: 10.1002/jae.1230.
- <span id="page-61-2"></span>Guido W. Imbens. Nonparametric estimation of average treatment effects under exogeneity: A review. Review of Economics and Statistics, 86(1):4–29, 2004. ISSN 00346535. doi: 10.1162/003465304323023651.
- <span id="page-62-10"></span>Piotr Jaworski, Fabrizio Durante, Wolfgang Karl Härdle, and Tomasz Rychlik. Copula Theory and Its Applications, volume 198 of Lecture Notes in Statistics. Springer Berlin Heidelberg, Berlin, Heidelberg, 2010. ISBN 978-3-642-12464-8. doi: 10.1007/ 978-3-642-12465-5.
- <span id="page-62-11"></span>Rainer Kress. Linear integral equations, volume 82. Springer, 2014. doi: 10.1007/ 978-1-4614-9593-2.
- <span id="page-62-1"></span>Kathleen T. Li. Statistical Inference for Average Treatment Effects Estimated by Synthetic Control Methods. Journal of the American Statistical Association, 115(532): 2068–2083, 2020. ISSN 1537274X. doi: 10.1080/01621459.2019.1686986.
- <span id="page-62-6"></span>Marc Lipsitch, Eric Tchetgen Tchetgen, and Ted Cohen. Negative Controls: A tool for detecting confounding and bias in observational studies. Epidemiology, 21(3):383–388, 2010. ISSN 10443983. doi: 10.1097/EDE.0b013e3181d61eeb.
- <span id="page-62-9"></span>D. L. McLeish. On the Invariance Principle for Nonstationary Mixingales. The Annals of Probability, 5(4):616–621, 1977. ISSN 0091-1798. doi: 10.1214/aop/1176995772.
- <span id="page-62-5"></span>Wang Miao and Eric J. Tchetgen Tchetgen. On varieties of doubly robust estimators under missingness not at random with a shadow variable. *Biometrika*,  $103(2)$ :475–482, 2016. ISSN 14643510. doi: 10.1093/biomet/asw016.
- <span id="page-62-2"></span>Wang Miao, Zhi Geng, and Eric J. Tchetgen Tchetgen. Identifying causal effects with proxy variables of an unmeasured confounder. Biometrika, 105(4):987–993, 2018. ISSN 14643510. doi: 10.1093/biomet/asy038.
- <span id="page-62-8"></span>Walter Philipp. The central limit problem for mixing sequences of random variables. Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete, 12(2):155–171, 1969. ISSN 00443719. doi: 10.1007/BF00531648.
- <span id="page-62-7"></span>Benedikt M. Pötscher and Ingmar R. Prucha. Dynamic Nonlinear Econometric Models. Springer Berlin Heidelberg, 1997. doi: 10.1007/978-3-662-03486-6.
- <span id="page-62-4"></span>Joaquin Quiñonero-Candela, Masashi Sugiyama, Neil D Lawrence, and Anton Schwaighofer. Dataset shift in machine learning. Mit Press, 2009.
- <span id="page-62-0"></span>Dan S. Rickman and Hongbo Wang. Two tales of two U.S. states: Regional fiscal austerity and economic performance. Regional Science and Urban Economics, 68:46–55, 2018. ISSN 18792308. doi: 10.1016/j.regsciurbeco.2017.10.008.
- <span id="page-62-3"></span>James Robins. A new approach to causal inference in mortality studies with a sustained exposure period-application to control of the healthy worker survivor effect. Mathematical modelling, 7(9-12):1393–1512, 1986.
- <span id="page-63-3"></span>James Robins and Sander Greenland. The Probability of Causation under a Stochastic Model for Individual Risk. Biometrics, 45(4):1125, 1989. ISSN 0006341X. doi: 10. 2307/2531765.
- <span id="page-63-4"></span>James M. Robins and Sander Greenland. Discussion of "Causal Inference without Counterfactuals" by A.P. Dawid. Journal of the American Statistical Association, 95(450): 477–482, 2000.
- <span id="page-63-6"></span>James M. Robins and Andrea Rotnitzky. Semiparametric efficiency in multivariate regression models with missing data. Journal of the American Statistical Association, 90 (429):122–129, 1995. ISSN 1537274X. doi: 10.1080/01621459.1995.10476494.
- <span id="page-63-5"></span>James M. Robins, Andrea Rotnitzky, and Lue Ping Zhao. Estimation of regression coefficients when some regressors are not always observed. Journal of the American Statistical Association, 89(427):846–866, 1994. ISSN 1537274X. doi: 10.1080/01621459. 1994.10476818.
- <span id="page-63-7"></span>James M. Robins, Andrea Rotnitzky, and Lue Ping Zhao. Analysis of semiparametric regression models for repeated outcomes in the presence of missing data. Journal of the American Statistical Association, 90(429):106–121, 1995. ISSN 1537274X. doi: 10.1080/01621459.1995.10476493.
- <span id="page-63-2"></span>Paul R. Rosenbaum and Donald B. Rubin. The central role of the propensity score in observational studies for causal effects. Biometrika, 70(1):41–55, 1983. ISSN 00063444. doi: 10.1093/biomet/70.1.41.
- <span id="page-63-9"></span>M. Rosenblatt. a Central Limit Theorem and a Strong Mixing Condition. Proceedings of the National Academy of Sciences, 42(1):43–47, 1956. ISSN 0027-8424. doi: 10.1073/ pnas.42.1.43.
- <span id="page-63-1"></span>Daniel O. Scharfstein, Andrea Rotnitzky, and James M. Robins. Adjusting for Nonignorable Drop-Out Using Semiparametric Nonresponse Models. Journal of the American Statistical Association, 94(448):1096–1120, 1999. ISSN 1537274X. doi: 10.1080/01621459.1999.10473862.
- <span id="page-63-8"></span>Thomas A. Severini and Gautam Tripathi. Some identification issues in nonparametric linear models with endogenous regressors. Econometric Theory, 22(2):258–278, 2006. ISSN 02664666. doi: 10.1017/S0266466606060117.
- <span id="page-63-0"></span>Xu Shi, Kendrick Li, Wang Miao, Mengtong Hu, and Eric Tchetgen Tchetgen. Theory for identification and Inference with Synthetic Controls: A Proximal Causal Inference Framework. arXiv preprint arXiv:2108.13935v4, 2023.
- <span id="page-64-1"></span>Heng Shu and Zhiqiang Tan. Improved Estimation of Average Treatment Effects on the Treated: Local Efficiency, Double Robustness, and Beyond. arXiv preprint arXiv:1808.01408v1, 2018. URL <http://arxiv.org/abs/1808.01408>.
- <span id="page-64-0"></span>Eric J Tchetgen Tchetgen, Andrew Ying, Yifan Cui, Xu Shi, and Wang Miao. An Introduction to Proximal Causal Learning. arXiv preprint arXiv:2009.10982v1, 2020.
- <span id="page-64-5"></span>Ramon van Handel. The universal Glivenko-Cantelli property. Probability Theory and Related Fields, 155(3-4):911–934, 2013. ISSN 01788051. doi: 10.1007/s00440-012-0416-5.
- <span id="page-64-2"></span>Tyler J. VanderWeele and James M. Robins. Stochastic counterfactuals and stochastic sufficient causes. Statistica Sinica, 22(1):379–392, 2012. ISSN 10170405. doi: 10.5705/ ss.2008.186.
- <span id="page-64-3"></span>Jeffrey M. Wooldridge. Estimation and inference for dependent processes. In *Handbook of* Econometrics, volume 4, pages 2639–2738. Elsevier, 1994. doi: 10.1016/S1573-4412(05) 80014-5.
- <span id="page-64-4"></span>Andrew Ying, Wang Miao, Xu Shi, and Eric J. Tchetgen Tchetgen. Proximal Causal Inference for Complex Longitudinal Studies. arXiv preprint arXiv:2109.07030v5, 2021.