

**ON HIGHER-DIMENSIONAL SUPERINTEGRABLE SYSTEMS:
A NEW FAMILY OF CLASSICAL AND QUANTUM
HAMILTONIAN MODELS**

MIGUEL A. RODRÍGUEZ AND PIERGIULIO TEMPESTA

ABSTRACT. We introduce a large family of n -dimensional Hamiltonian systems which, contain, as special reductions, several superintegrable systems as the Tremblay-Turbiner-Winternitz system, a generalized Kepler potential and the anisotropic harmonic oscillator with Rosochatius terms. We conjecture that there exist special values in the space of parameters, apart from those leading to known cases, such that this new Hamiltonian family is superintegrable.

CONTENTS

1. Introduction	1
1.1. Classical and Quantum superintegrability	1
1.2. The new model	2
2. On the construction of the new Hamiltonian family	3
3. The quantum version of the model	5
4. Open problems and conjectures	6
Acknowledgements	7
References	7

1. INTRODUCTION

1.1. Classical and Quantum superintegrability. In the last two decades, the theory of superintegrable systems has been one of the most active research areas in mathematical physics [23, 9]. These systems represent a special class of integrable systems which possess, both in the classical and in the quantum case, surprisingly rich algebraic and geometric properties [4], [8], [27], [11], [10], [21], [22]. Essentially, they possess more independent integrals of motion than degrees of freedom. If a system with n degrees of freedom admits $n + 1$ functionally independent integrals of motion, then it is said to be *minimally superintegrable*; if it admits $2n - 1$ integrals (the maximal number allowed), is *maximally superintegrable*.

In classical mechanics, as a consequence of Nekhoroshev theorem [11] bounded orbits of a maximally superintegrable system are closed and periodic.

In quantum mechanics superintegrability is related with the existence of hidden symmetries, responsible for the accidental degeneracy of the spectrum.

Besides, the notion of maximal superintegrability is intimately related with that of *exact solvability*. An exactly solvable quantum mechanical system is a model

whose spectrum and eigenfunctions can be obtained by purely algebraic means. It is characterized by the fact that there exists in its solution space an infinite flag of functional linear spaces, which is preserved by the Hamiltonian.

A particular, well-known example of these flags is given by finite-dimensional representation spaces of semi-simple Lie algebras of first order differential operators. Thus, the Hamiltonian and its quantum integrals of motion turn out to be an element of the universal enveloping algebra of a Lie algebra.

Instead, for a quasi-exactly-solvable system, only few energy levels and eigenfunctions can be obtained algebraically.

A conjecture proposed in [22] states that maximally superintegrable quantum systems are indeed exactly solvable.

Another important issue is that of *separability*, namely the construction of suitable coordinate systems (the *separation variables*) in which dynamics decouple. The formulation of algebraic and geometric techniques allowing to separate variables both for classical and quantum models has been historically one of the most relevant problems in the theory of integrable systems, and has been recently extended to the case of superintegrable models [6]. Typically, superintegrable systems are *multiseparable*, namely separable in more than one coordinate system [16]. However, this property is not necessary: interestingly enough, superintegrable systems, even maximally ones, may or may not be separable [9]. A paradigmatic example is the classical Post-Winternitz system [13], which is indeed maximally superintegrable. However, its Hamilton-Jacobi equation is not additively separable, at least in any orthogonal coordinate system in the configuration space, since the system does not possess integrals of motion quadratic in the momenta, but only higher-degree ones. The same is true for the Schrödinger equation associated, which does not admit multiplicative separability in any coordinate system (See Theorem 1 of [13]). The purpose of this Letter is to introduce in a simple way a new (to the best of our knowledge) n -dimensional generalization of a class of well-known maximally superintegrable systems.

1.2. The new model. Precisely, we propose the following

Definition 1. *We introduce the Hamiltonian*

$$(1.1) \quad H_n^{(cl)} := (x_1^2 + \dots + x_n^2 + \gamma)^{\frac{k-1}{k}} \left\{ \frac{1}{2} \sum_{i=1}^n p_{x_i}^2 + \frac{1}{2k^2} (\omega_1^2 x_1^2 + \dots + \omega_n^2 x_n^2)^{\frac{s-1-k}{k}} + \frac{\alpha_1}{x_1^2} + \dots + \frac{\alpha_n}{x_n^2} \right\},$$

where $\gamma \geq 0$, $k \neq 0$, $s, \alpha_1, \dots, \alpha_n, \omega_1, \dots, \omega_n$ are real parameters.

Our motivation is the fact that, surprisingly, many well-known superintegrable models can be obtained as particular cases of the new family (1.1).

We start our analysis considering first the Hamiltonian (1.1) for $\gamma = 0$.

As we shall prove, for $n = 2$, $s = 1$ and $\omega_1 = \omega_2$ our Hamiltonian (1.1) is equivalent to the well-known Tremblay-Turbiner-Winternitz (TTW) system [24, 25], which has attracted considerable attention in the last decade. The authors conjectured that the system is maximally superintegrable for all rational values of the parameter k . The conjecture was proved for odd k in [14] and subsequently for all rational values of k in the papers [5, 7].

In order to prove the direct relation of system (1.1) with the TTW system, we shall construct a transformation which leads the TTW system to a new Cartesian form.

This allows us immediately to generalize the TTW system to n dimensions. In turn, this form can be readily identified with a particular case of system (1.1).

For $n = 2$, $s = -1/2$ and $\omega_1 = \omega_2$, the Hamiltonian (1.1) reproduces the generalized Kepler system (gKs) [12], which is related with the TTW system via the coupling constant metamorphosis mechanism and is also maximally superintegrable.

Besides, for $k = 1$, $s = 2$, $\omega_1 = \dots = \omega_n$, one readily obtains from system (1.1) the n -dimensional isotropic harmonic oscillator with Rosochatius terms [20]. It is also a maximally superintegrable system, as has been proved independently in [3] and [17].

Another notable reduction of our Hamiltonian, again for $n = 2$, is obtained for $k = 1$, $s = 2$, $\omega_1 = \omega_2$: indeed, in this case we get the system Smorodinsky-Winternitz I [27], one of the four superintegrable systems in the plane with quadratic integrals of motion. Also, for $k = s = 1$, and $\omega_1 = 4\omega^2$, $\omega_2 = \omega^2$ we obtain the system Smorodinsky-Winternitz II.

The Hamiltonian (1.1) is also related with the interesting family of Hamiltonians studied by Ballesteros, Enciso, Herranz and Ragnisco in [1]. This general family describes the motion of a particle on any n -dimensional spherically symmetric curved space and is quasi-maximally superintegrable, namely it possesses $2n - 2$ integrals for any choice of the two arbitrary functions of r appearing in it. Precisely, the particular case of Hamiltonian (1.1) with $\omega_1 = \dots = \omega_n$ and $s = 1$ falls into this class, when the central potential $\mathcal{U}(r)$ vanishes.

Finally, for $\gamma > 0$, $k = 1/2$, $s = 2$, $\omega_1 = \dots = \omega_n$ we recover the class of maximally superintegrable models on an n -dimensional space of nonconstant curvature, introduced in [2].

A quantum version of Hamiltonian (1.1), namely the system (3.3), will also be presented and its properties briefly discussed. We conjecture that for special values of the space of parameters of the quantum system (3.3), we will obtain exactly solvable models.

Clearly, the issue of integrability or superintegrability of our family (1.1), both for the classical and quantum cases, is a very interesting one. Some related conjectures and open problems will be stated in the final section.

The problem of separability of system (1.1) is also completely open, although there are certainly separable subcases. For instance, the TTW system is separable in polar coordinates for any rational value of k .

The paper is organized as follows. In Section 2, we formulate in new variables the standard TTW system. In this way, the link with the new family of systems (1.1) becomes transparent. Also, we propose our generalization of the original TTW system by means of an anisotropic deformation of the harmonic term and by an extra parameter interpolating between the Kepler and the anisotropic cases. In Section 3, we propose a quantum version of our Hamiltonian. We propose several open problems in the final Section 4.

2. ON THE CONSTRUCTION OF THE NEW HAMILTONIAN FAMILY

The classical version of the TTW Hamiltonian in polar coordinates reads:

$$(2.1) \quad H^{TTW} = \frac{1}{2} \left(p_r^2 + \frac{p_\phi^2}{r^2} \right) + \frac{\alpha k^2}{2r^2 \cos^2 k\varphi} + \frac{\beta k^2}{2r^2 \sin^2 k\varphi}$$

We shall prove that for $n = 2$, $s = 2$ and $\omega_1 = \dots = \omega_n = \omega$ Hamiltonian (1.1) reduces to system (2.1). To this aim, first we shall rewrite Hamiltonian (2.1) in Cartesian coordinates. We get

$$(2.2) \quad H_{cart}^{TTW} := \frac{1}{2}(p_x^2 + p_y^2) + \frac{1}{2}\omega^2(x^2 + y^2) + V_k(x, y; \alpha, \beta)$$

where we have introduced the potential

$$(2.3) \quad V_k(x, y; \alpha, \beta) = 2k^2(x^2 + y^2)^{k-1} \left(\frac{\alpha}{((x + iy)^k + (x - iy)^k)^2} - \frac{\beta}{((x + iy)^k - (x - iy)^k)^2} \right).$$

Let us introduce two complex variables:

$$(2.4) \quad z = x + iy, \quad p_z = \frac{1}{2}(p_x - ip_y)$$

and their complex conjugates

$$(2.5) \quad \bar{z} = x - iy, \quad p_{\bar{z}} = \frac{1}{2}(p_x + ip_y)$$

This change of variables preserves the canonical symplectic form in \mathbb{R}^4 . The TTW Hamiltonian can be written in these new coordinates as:

$$(2.6) \quad \mathfrak{H}^{TTW} := 2p_z p_{\bar{z}} + \frac{1}{2}\omega^2 z \bar{z} + V_k(z, \bar{z}; \alpha, \beta)$$

Using de Moivre's formula:

$$(2.7) \quad z = r^{i\varphi}, \quad z^k = r^k(\cos k\varphi + i \sin k\varphi)$$

that is:

$$(2.8) \quad \cos k\varphi = \frac{1}{2|z|^k}(z^k + \bar{z}^k), \quad \sin k\varphi = \frac{1}{2i|z|^k}(z^k - \bar{z}^k)$$

we can express the potential as:

$$(2.9) \quad V_k(z, \bar{z}; \alpha, \beta) = \frac{2\alpha k^2 z^{k-1} \bar{z}^{k-1}}{(z^k + \bar{z}^k)^2} - \frac{2\beta k^2 z^{k-1} \bar{z}^{k-1}}{(z^k - \bar{z}^k)^2}.$$

If $k = 1$ we recover the harmonic oscillator with centrifugal terms:

$$(2.10) \quad V_1(z, \bar{z}; \alpha, \beta) = \frac{2\alpha}{(z + \bar{z})^2} - \frac{2\beta}{(z - \bar{z})^2}$$

and, in fact, (2.9) is the general form of the potential for any k . We can make another change of coordinates, in this case to real coordinates:

$$(2.11) \quad u = \frac{1}{2}(z^k + \bar{z}^k), \quad v = \frac{1}{2i}(z^k - \bar{z}^k)$$

yielding the corresponding transformation for the momenta

$$(2.12) \quad p_z = \frac{k}{2}z^{k-1}(p_u - ip_v), \quad p_{\bar{z}} = \frac{k}{2}\bar{z}^{k-1}(p_u + ip_v).$$

Consequently, the Hamiltonian (2.6) in these new, *Cartesian* coordinates reads

$$(2.13) \quad \mathfrak{H}_k = k^2(u^2 + v^2)^{\frac{k-1}{k}} \left[\frac{1}{2}(p_u^2 + p_v^2) + \frac{\omega^2}{2k^2}(u^2 + v^2)^{\frac{2}{k}-1} + \frac{\alpha}{2u^2} + \frac{\beta}{2v^2} \right].$$

Remark 1. The global factor in eq. (2.13) can be interpreted as a conformally flat metric; the remaining Hamiltonian factor represents a nonlinear oscillator with centrifugal barriers. These barriers can also arise from the reduction of a nonlinear oscillator in a four dimensional space [17, 3]; alternatively, they can come from the construction given in [15].

Let us now introduce a new parameter s , which allows us to treat the Kepler and the harmonic cases on the same footing. Precisely,

$$(2.14) \quad \mathfrak{H}_k^{(s)}(x, y) = \frac{1}{2}(p_x^2 + p_y^2) + \frac{1}{2}\omega^2(x^2 + y^2)^s + V_k(x, y; \alpha, \beta)$$

Using (2.4) and (2.5), we get

$$(2.15) \quad \mathfrak{H}_k^{(s)} = 2p_z p_{\bar{z}} + \frac{1}{2}\omega^2(z\bar{z})^s + V_k(z, \bar{z}; \alpha, \beta)$$

As before, taking into account eqs. (2.11) and (2.12) we get

$$(2.16) \quad (z\bar{z})^s = (u^2 + v^2)^{s/k}, \quad p_z p_{\bar{z}} = \frac{k^2}{4}(z\bar{z})^{k-1}(p_u^2 + p_v^2)$$

and

$$(2.17) \quad V_k(u, v; \alpha, \beta) = \frac{k^2}{2}(u^2 + v^2)^{(k-1)/k} \left(\frac{\alpha}{u^2} + \frac{\beta}{v^2} \right)$$

Thus, we obtain the new Hamiltonian

$$(2.18) \quad \mathfrak{H}_k^{(s)} = \frac{k^2}{2}(u^2 + v^2)^{(k-1)/k}(p_u^2 + p_v^2) + \frac{1}{2}\omega^2(u^2 + v^2)^{s/k} + \frac{k^2}{2}(u^2 + v^2)^{(k-1)/k} \left(\frac{\alpha}{u^2} + \frac{\beta}{v^2} \right)$$

or

$$(2.19) \quad \mathfrak{H}_k^{(s)} = k^2(u^2 + v^2)^{(k-1)/k} \left\{ \frac{1}{2}(p_u^2 + p_v^2) + \frac{\omega^2}{2k^2}(u^2 + v^2)^{(s-k+1)/k} + \frac{\alpha}{2u^2} + \frac{\beta}{2v^2} \right\}$$

Remark 2. The form of the Hamiltonian (2.19) was the inspiration for our Hamiltonian (1.1). This system is more general than system (2.19) for two reasons. The first one is that our Hamiltonian (1.1) is defined in n dimensions. The second one is that we allow anisotropy in the oscillations along distinct axes. This, in turn, relates directly the Hamiltonian (1.1) with well-known anisotropic oscillators.

Finally, we observe that the new parameter s allows us to continuously interpolate between the harmonic and Kepler cases.

3. THE QUANTUM VERSION OF THE MODEL

We wish to propose now the quantum analogue of our Hamiltonian models.

Definition 2. We introduce the family of quantum Hamiltonians

$$(3.1) \quad H_n^{(q)} := (x_1^2 + \dots + x_n^2 + \gamma)^{\frac{k-1}{k}} \cdot \left\{ -\frac{1}{2}(\partial_{x_1}^2 + \dots + \partial_{x_n}^2) + \frac{1}{2k^2}(\omega_1^2 x_1^2 + \dots + \omega_n^2 x_n^2)^{\frac{s-1-k}{k}} + \frac{\alpha_1}{x_1^2} + \dots + \frac{\alpha_n}{x_n^2} \right\},$$

where $\gamma \geq 0$, $k \neq 0$, $s, \alpha_i, \omega_i, i = 1, \dots, n$ are real parameters.

This quantum Hamiltonian has also several interesting reductions. For sake of simplicity, we shall consider $\gamma = 0$.

- For $k = 1$, $s = 2$ it reproduces a quantum anisotropic oscillator with Rosochatius terms, which is maximally superintegrable when all frequencies coincide.
- For $k = 1$, $s = -1/2$ and equal frequencies we get again a Kepler potential with a Rosochatius deformation.
- For $k = 1$, $s = -1/2$ and equal frequencies and $\alpha_n = 0$ we obtain the system

$$(3.2) \quad H = -\frac{1}{2}(\partial_{x_1}^2 + \cdots + \partial_{x_n}^2) - \frac{\gamma}{r} + \frac{\alpha_1}{x_1^2} + \cdots + \frac{\alpha_{n-1}}{x_{n-1}^2},$$

with $r = \sqrt{x_1^2 + \cdots + x_n^2}$, which was proved in [19] to be exactly solvable.

- The two dimensional case

$$(3.3) \quad H_2^{(q)} := (x^2 + y^2)^{\frac{k-1}{k}} \left\{ -\frac{1}{2}(\partial_x^2 + \partial_y^2) + \frac{1}{2k^2}(\omega_1^2 x^2 + \omega_2^2 y^2)^{\frac{s-1-k}{k}} + \frac{\alpha}{x^2} + \frac{\beta}{y^2} \right\},$$

is quite interesting by itself. Notice that for $k = s = 1$, and $\omega_1 = 4\omega^2$, $\omega_2 = \omega^2$, we recover the maximally superintegrable and exactly solvable [22] Smorodinski-Winternitz system II:

$$(3.4) \quad H^{SWII} = -\frac{1}{2}(\partial_x^2 + \partial_y^2) + \omega^2(4x^2 + y^2) + \frac{\alpha}{y^2}$$

once we annul the constant multiplying the linear term in x .

- Again in the case $n = 2$, and for $\omega_1 = \omega_2$, $s = 2$, Hamiltonian (3.3) reduces to the quantum TTW system [24], whereas, for $s = -1/2$, it reduces to the quantum version of the generalized Kepler model [12]. This can be verified by means of variable transformations completely analogous to those illustrated in Section 2.

4. OPEN PROBLEMS AND CONJECTURES

The purpose of this *divertimento* was, in essence, to introduce the Hamiltonian (1.1). It represents a large family of Hamiltonian systems, including as particular cases or reductions several well-known superintegrable models that attracted much attention in the last decades. Many questions concerning Hamiltonian (1.1) remain open. Thus, we propose the following, general problems.

Problem 1. Find all the values of the parameters appearing in the classical Hamiltonian (1.1) which make it integrable or (maximally) superintegrable, in all dimensions. Determine the orbits of the motion.

Problem 2. Find the values of the parameters appearing in the quantum Hamiltonian (3.3) which make it integrable or superintegrable. Find the spectrum and the eigenfunctions associated.

Problem 3. Find, for special values of the parameters, separation variables for the Hamilton-Jacobi equation associated with the classical Hamiltonian (1.1) and for the Schrödinger equation of the quantum case (3.3).

Problem 4. Determine whether the quantum Hamiltonian (3.3) is exactly solvable or quasi-exactly solvable for special values of the space of the parameters.

We conjecture that there exist values of the parameters ensuring classical (maximal) superintegrability and quantum exact solvability, apart from those leading to known reductions.

ACKNOWLEDGEMENTS

We wish to thank G. Gubbiotti and D. Latini for interesting discussions. This work has been partly supported by the Universidad Complutense de Madrid under grant G/6400100/3000, and by the Severo Ochoa Programme for Centres of Excellence in R&D (CEX2019-000904-S), Ministerio de Ciencia, Innovación y Universidades y Agencia Estatal de Investigación, Spain.

P. T. is member of the Gruppo Nazionale di Fisica Matematica (GNFM).

REFERENCES

- [1] A. Ballesteros, A. Enciso, F. Herranz and O. Ragnisco, *Superintegrability on N-dimensional curved spaces: Central potentials, centrifugal terms and monopoles*, Ann. Phys. **324**, 1219-1233 (2009).
- [2] A. Ballesteros, A. Enciso, F. Herranz and O. Ragnisco, *A maximally superintegrable system on an n-dimensional space of nonconstant curvature*, Phys. D **237**, 505-509 (2008).
- [3] N.W. Evans, P.E. Verrier. *Superintegrability of the caged anisotropic oscillator*. J. Math. Phys. **49** 092902 (2008).
- [4] J. Friš, V. Mandrosov, Ya. A. Smorodinsky, M. Uhlir and P. Winternitz, *On higher symmetries in quantum mechanics*, Phys. Lett. **16**, 354-356 (1965).
- [5] E.G. Kalnins, J.M. Kress and W.Jr. Miller, *Families of classical subgroup separable superintegrable systems*, J. Phys. A: Math. Theor. **43** 092001 (2010).
- [6] E.G. Kalnins, J.M. Kress and W.Jr. Miller, *Separation of Variables and Superintegrability*, IOP Publishing, Bristol, UK, 2018.
- [7] E.G. Kalnins, W.Jr. Miller and G.S. Pogosyan, *Superintegrability and higher-order constants for classical and quantum systems*, Phys. At. Nucl. **74** 914-918 (2011).
- [8] A.A. Makarov, J.A. Smorodinsky, Kh. Valiev and P. Winternitz, *A systematic search for nonrelativistic systems with dynamical symmetries*, Nuovo Cimento A **52**, 1061-1084 (1967).
- [9] W. Miller Jr, S. Post and P. Winternitz, *Classical and Quantum Superintegrability with Applications*, J. Phys. A, **46**, 423001 (2013).
- [10] A.S. Mischenko and A. T. Fomenko, *Generalized Liouville method of integration of Hamiltonian systems*, Funct. Anal. Appl. **12**, 113-121 (1978).
- [11] N.N. Nekhoroshev, *Action-angle variables and their generalizations*, Trans. Moscow Math. Soc. **26**, 180-198 (1972).
- [12] S. Post, P. Winternitz, *An infinite family of superintegrable deformations of the Coulomb potential*, J. Phys. A: Math. Theor. **43**, 222001 (2010).
- [13] S. Post, P. Winternitz, *A nonseparable quantum superintegrable system in 2D real Euclidean space*, J. Phys. A: Math. Theor. **44**, 162001 (2010).
- [14] C. Quesne, *Superintegrability of the Tremblay-Turbiner-Winternitz quantum Hamiltonian on a plane for odd k*, J. Phys. A: Math. Theor. **43**, 082001 (2010).
- [15] M.F. Rañada, M.A. Rodríguez, M. Santander. *A new proof of the higher-order superintegrability of a noncentral oscillator with inversely quadratic nonlinearities*. J. Math. Phys. **51** 042901 (2010).
- [16] D. Reyes Nozaleda, P. Tempesta and G. Tondo. *Classical multiseparable Hamiltonian systems, superintegrability and Haantjes geometry*. Commun. Nonl. Sci. Num. Sim. **104**, 106021 (2022).
- [17] M.A. Rodríguez, P. Tempesta and P. Winternitz, *Reduction of superintegrable systems: the anisotropic harmonic oscillator*, Phys. Rev. E **78**, 046608 (2008).
- [18] M.A. Rodríguez, P. Tempesta and P. Winternitz, *Symmetry reduction and superintegrable Hamiltonian systems*, J. Phys. Conf. Ser. **175** 012013 (2009).

- [19] M.A. Rodríguez and P. Winternitz Quantum superintegrability and exact solvability in n dimensions, *J. Math. Phys.* **43**, 1309-1322 (2002).
- [20] E. Rosochatius, Dissertation, Gottingen, Gebr. Unger, Berlin 1877.
- [21] M. Sheftel, P. Tempesta and P. Winternitz, *Superintegrable Systems in Quantum Mechanics and Classical Lie Theory*, *J. Math. Phys.* **42**, 659-673 (2001).
- [22] P. Tempesta, A. Turbiner and P. Winternitz, *Exact Solvability of Superintegrable Systems*, *J. Math. Phys.* **42**, 4248-4257 (2001).
- [23] P. Tempesta, P. Winternitz, J. Harnad, W. Miller, Jr, G. Pogosyan, M.A. Rodríguez (eds), *Superintegrability in Classical and Quantum Systems*, Montréal, CRM proceedings and Lecture Notes, AMS, vol. **37** (2004).
- [24] F. Tremblay, A.V. Turbiner and P. Winternitz *An infinite family of solvable and integrable quantum systems on a plane*, *J. Phys. A: Math. Theor.* **42** 242001 (2009).
- [25] F. Tremblay, A.V. Turbiner, P. Winternitz. *Periodic orbits for an infinite family of classical superintegrable systems*. *J. Phys. A: Math. Theor.* **43** 015202 (2010).
- [26] P.E. Verrier, N.W. Evans. *A new superintegrable Hamiltonian*. *J. Math. Phys.* **49** 022902 (2008).
- [27] P. Winternitz, Ya. Smorodinsky, M. Uhliř and J. Friš, *Symmetry groups in classical and quantum mechanics*, *Yad. Fiz.* **4**, 625-635 (1966), *Sov. J. Nucl. Phys* **4**, 444-450 (1967).

DEPARTAMENTO DE FÍSICA TEÓRICA, FACULTAD DE CIENCIAS FÍSICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, PLAZA DE LAS CIENCIAS, 1, 28040 – MADRID, SPAIN
Email address: `rodrigue@fis.ucm.es`

DEPARTAMENTO DE FÍSICA TEÓRICA, FACULTAD DE CIENCIAS FÍSICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, PLAZA DE LAS CIENCIAS, 1, 28040 – MADRID, SPAIN, AND INSTITUTO DE CIENCIAS MATEMÁTICAS, C/ NICOLÁS CABRERA, NO 13–15, 28049 MADRID, SPAIN
Email address: `piergiulio.tempesta@icmat.es`, `ptempest@ucm.es`