Spectral Properties of the Symmetry Generators of Conformal Quantum Mechanics: A Path-Integral Approach

H. E. Camblong,¹ A. Chakraborty,² P. Lopez Duque,² and C. R. Ordóñez²

¹Department of Physics and Astronomy, University of San Francisco, San Francisco, California 94117-1080, USA

²Department of Physics, University of Houston, Houston, Texas 77024-5005, USA

(Dated: December 7, 2023)

Abstract

A path-integral approach is used to study the spectral properties of the generators of the SO(2,1) symmetry of conformal quantum mechanics (CQM). In particular, we consider the CQM version that corresponds to the weak-coupling regime of the inverse square potential. We develop a general framework to characterize a generic symmetry generator G (linear combinations of the Hamiltonian H, special conformal operator K, and dilation operator D), from which the path-integral propagators follow, leading to a complete spectral decomposition. This is done for the three classes of operators: elliptic, parabolic, and hyperbolic. We also highlight novel results for the hyperbolic operators, with a continuous spectrum, and their quantum-mechanical interpretation. The spectral technique developed for the eigensystem of continuous-spectrum operators can be generalized to other operator problems.

I. INTRODUCTION AND CONTEXT

The path-integral approach, along with associated functional techniques, is a powerful methodology that provides a complete characterization of quantum-mechanical systems. For ordinary quantum mechanics problems, great advances in finding useful solutions have been made in recent decades—and exhaustive lists of path-integral solutions can be found in [1-3]. In this work, we use the path-integral approach to derive the eigenvectors and eigenvalues of some of the operators relevant to conformal quantum mechanics.

Conformal quantum mechanics (CQM) has attracted considerable attention following its initial formulation in the 1970s, when it was first proposed as an example of a scale invariant theory [4], and analyzed in detail by de Alfaro, Fubini, and Furlan (dAFF) [5] as a (0+1)dimensional form of conformal field theory. The dAFF model corresponds to the D = 1limit of the spacetime *D*-dimensional conformally invariant Lagrangian density [5]

$$\mathcal{L} = \frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi - g \phi^{2D/(D-2)} .$$
⁽¹⁾

In the transition to D = 1, one can interpret the resulting theory as standard quantum mechanics, with a "field" Q(t) described as a configuration or position variable subject to an inverse square potential, with Lagrangian

$$L = \frac{1}{2}\dot{Q}^2 - \frac{g}{2Q^2} , \qquad (2)$$

where the dot represents the usual time derivative. This original form of CQM, centered on its SO(2,1) conformal symmetry structure, was subsequently used in seminal papers by Jackiw [6, 7], also including an analysis of a related CQM system of contact interactions [8]. Systems exhibiting this kind of SO(2,1) conformal symmetry have seen renewed interest in a broad range of physical applications over the years. This is primarily due to its simplicity as a model of conformal field theory and its remarkably wide range of applicability to established physical problems that exhibit approximate conformal symmetry in a window of physical scales. While not included in the original dAFF model formulation, outstanding realizations of this type of inverse-square-potential systems have been found where this symmetry generates a quantum anomaly. A list of noteworthy applications of such systems includes molecular physics [9], black hole thermodynamics [10–12] and acceleration radiation [13–16], the Efimov effect [17–20], and graphene [21, 22], among several others [20]. It is noteworthy that, in addition to the quantum symmetry breaking based on a strong-coupling version of Lagrangian (2), another class of quantum anomalies have been found in systems with SO(2,1) conformal symmetry with contact interactions—they involve 1D three-body interactions in 1D and 2D two-body interactions in Fermi systems of ultracold atoms [23–27].

The previous list emphasizes those cases where the CQM interaction is attractive and sufficiently strong, where it leads to a quantum anomaly [9, 28–30] or some kind of renormalization [31–36], or to the fall-to-the-center phenomenon [37, 38]—this regime is called "strong coupling" for short. On the other hand, the form of CQM often discussed in the context of conformal symmetry analyses is based on the dAFF model [5]. The latter strictly applies to the case when the conformal potential is repulsive (or sufficiently weak, even if attractive) to avert the pathologies inherent for strong coupling g in Eq. (2), and the conformal symmetry is maintained. For the sake of simplicity, in this work we solely focus on this version of CQM (in a slightly generalized format), for which we analyze the spectral properties of some of its symmetry generators. We will address elsewhere the more general case, including the presence of anomalies.

Of particular interest for the present work is the fact that the dAFF model (in the original weak-coupling formulation of the CQM symmetry generators) has been recently found to be a CFT_1 realization of the AdS/CFT correspondence [39, 40], leading to renewed interest in this topic [41–46]. In addition, using the operators we discuss in this paper, the dynamic evolution within the dAFF model has been used to study causal diamonds in Minkowski spacetime [47, 48]. The associated physics of finite-lifetime observers, often called diamond observers, was first addressed in Ref. [49] in a remarkable finding that generalizes a similar thermalization of the vacuum for accelerated observers (the celebrated Unruh effect [50]). Accordingly, these observers have access to only a limited region of spacetime, known as the conformal (causal) diamond; and the vacuum they perceive has a diamond temperature inversely proportional to their lifetime. This insightful result on the thermalization of the vacuum of diamond observers has been further confirmed in Refs. [51-56]. As there are a number of parallels with the thermodynamic behavior of black hole horizons, where CQM has been successfully used, it is not surprising that that the dAFF model may also be relevant for causal diamonds. Specifically, in Refs. [47, 48] the dynamic evolution within the dAFF model was shown to be in correspondence with the time evolution of Minkowski observers with a finite lifetime, which is described by the radial conformal Killing fields (RCKF) previously developed in Ref. [57]. This problem is of interest in further characterizing the causal structure of spacetime, and CQM appears to be a promising tool for this purpose.

A. Scope and Main Results of This Paper

In the physical applications mentioned above, and especially in regards to the recent work on CQM as CFT₁ and the physics of causal diamonds, the CQM generators play a central role. For our purposes, the prototypical CQM generators of generalized time evolution can be taken as the operators H, R, and S, where H is the quantum-mechanical Hamiltonian associated with the Lagrangian (2), and R and S are the linear combinations of H and the special conformal generator K involved in setting up the Cartan-Weyl basis. In particular, the spectral decomposition of the operator S, which plays a significant role in causal diamonds [47, 48, 57], is a novel result of our paper. More generally, the dynamical evolution can be described by any of the generalized generators G defined as linear combinations of H, D, and K, where D is the dilation operator.

While the usual operator properties of the set $\{H, R, S\}$ and the generalized generators G have been considered in the literature [5], a path-integral treatment is lacking. It is the purpose of this paper to develop such functional integral approach, with which we compute the propagators and the spectral properties of the generators $\{H, R, S\}$, as well as those of the linear combinations that define a generalized generator G, falling under the three possible classes: elliptic (*R*-like), parabolic (*H*-like), and hyperbolic (*S*-like).

The main results of the paper are summarized below in a convenient format that can help identify the key ideas within the extensive mathematical properties being discussed.

- We have developed a complete Hamiltonian framework for all the generalized generators G of CQM (which proves essential for the path-integral approach).
- We have derived the spectral properties, including the eigenvalues and eigenstates for the three classes of CQM generators G, viz., elliptic (R), parabolic (H), and hyperbolic (S), using a path-integral approach. This involves using different limits of the propagator for the generalized radial harmonic oscillator $K_{l+\nu}^{(RHO)}$ given in Eq. (45). Depending on the classes of the operators, we have used different methods to extract the spectral properties of the operator G from the propagator. Table I acts as a pointer

to the equations describing the spectral properties found in this paper.

CQM Generator	Propagator	Eigenvalues	Eigenstates
Elliptic (R)	$K_{l+ u}^{(RHO)}$ – Eqs. (45), (48)	Discrete – Eq. (51)	Eq. (53)
Parabolic (H)	$\lim_{\omega \to 0} K_{l+\nu}^{(RHO)} - \text{Eq. (56)}$	Continuous in \mathbb{R}^+	Eq. (60)
Hyperbolic (S)	$\lim_{\omega \to -i\omega} K_{l+\nu}^{(RHO)} - \text{Eq. (66)}$	Continuous in \mathbb{R}	Eqs. (81) & (82)

TABLE I: References to the key results about the spectral properties of the CQM generators.

- We have used a novel method based on Fourier transforms (which we call the Fourier Method) to find the eigenstates of the noncompact CQM operators from the propagator. The key equation for this method is succinctly expressed in Eq. (62). Furthermore, we have explored the connection of this method with the retarded and advanced Green's functions in Appendix C, culminating in Eq. (C8).
- In particular, we have derived a complete spectral characterization of the operator S and all hyperbolic generators—a result that has been surprisingly lacking in the literature.
- As a bonus, our detailed analysis of the spectral properties of the propagators via different approaches has uncovered additional mathematical connections and identities, as mentioned at the end of Subsec. VA and in Appendix D.

B. Organization of This Paper

In Section II, we summarize the symmetry properties of CQM and the specifics of the dAFF model; we define the operators R, H, and S that generate the conformal symmetry group, as well as the generic conformal generator G; and we develop the framework that classifies the possible types of G and their role in the dynamical time evolution via their Hamiltonian representation. In Sec. III, we review the path-integral approach for radial problems in quantum mechanics and specifically use it to study a generalized radial harmonic oscillator relevant to CQM. In Sec. IV, we fully characterize the spectral properties of the operator R and its class (elliptic). In Sec. V, we analyze the spectral properties of



FIG. 1: Flowchart of this paper.

the operator H and its class (parabolic); and we develop a novel procedure for spectral properties of operators with continuous spectra. In Sec. VI, we apply this method to the conformal operator S and its class of non-compact hyperbolic generators to fully characterize their spectral data and Green's functions. We conclude the paper in Sec. VII with a brief summary and directions for future work. The appendices cover an alternative dimensional framework for the Hamiltonian description of generators; detailed properties of the relevant path-integral framework; evaluation of the inversion integrals for the spectral decomposition of parabolic and hyperbolic generators; and further analysis of the conformal generators within a differential-equation approach.

The logic of the paper organization is outlined in Fig. 1. The flowchart stresses the need to use an appropriate, alternative Hamiltonian formulation for the path-integral framework before the spectral analysis of the CQM generators is performed; then, from both the Hamiltonian formulation and the path-integral treatment, all the properties of the CQM generators follow systematically, with additional technical details provided by the appendices.

II. SYMMETRY PROPERTIES AND GENERATORS OF CONFORMAL QUAN-TUM MECHANICS (CQM): HAMILTONIAN FORMULATION

As outlined in Sec. I, the dAFF model is the limiting D = 1 field theory of Eq. (1). While Eq. (2) is the simplest form of CQM, in this paper, we consider the more general multicomponent case, where the *d* components correspond to the *d* spatial dimensions of the position coordinates $\mathbf{Q}(t)$ in quantum mechanics, evolving dynamically with respect to the ordinary time *t*. This physical system, with Lagrangian

$$L = \frac{M}{2} \dot{\mathbf{Q}}^2 - \frac{\lambda}{2Q^2} \,, \tag{3}$$

can be described in polar hyperspherical coordinates, with the radial variable $Q = |\mathbf{Q}| \ge 0$ defined in the nonnegative half-line. In the path-integral framework, a system with Lagrangian (3) can be more naturally described within a class of radial functional problems, as we will see in Sec. III. In Eqs. (2) and (3), we are adopting a definition of the coupling constant with an additional factor of 1/2 for convenience, consistent with the numerical factors of Ref. [5], and rescaled in the transition to the quantum theory, with $\lambda = \hbar^2 g/M$, with g dimensionless. In addition, we are inserting an extra dimensional parameter M that can be interpreted as the mass of a nonrelativistic particle in the conventional form of quantum mechanics. Often, the choices M = 1 and $\hbar = 1$ are made for the sake of simplicity (including in Ref. [5]), but we will keep conventional dimensional parameters to help in some of the derivations with the use of appropriate analytic continuations, limiting procedures, and familiar physical interpretations for both the Hamiltonian operator and the other symmetry generators.

A. Conformal Symmetries

Rather than starting directly from the Lagrangian (3), one can begin the problem looking at the more general conditions that define conformal invariance for a generic Lagrangian

$$L = \frac{M}{2} \dot{\mathbf{Q}}^2 - V(\mathbf{Q}) \quad . \tag{4}$$

From Noether's theorem, for systems with Lagrangians of the form (4) the necessary conditions to yield an invariant action under general time reparametrizations [5–7] (with time transformations such that the Lagrangian itself is not necessarily invariant) are: (i) the potential function $V(\mathbf{Q})$ is a homogeneous function of degree -2; (ii) the time transformations appear restricted to any of three independent building blocks: time translations, time dilatations, and inverse time translations. In what follows, $\mathbf{P} \equiv \mathbf{P}_{\mathbf{Q}}$ is the momentum conjugate to \mathbf{Q} . Then, the transformation algebra consists of the three generators (including operator ordering at the quantum level):

• The Hamiltonian

$$H = \frac{\mathbf{P}^2}{2M} + V(\mathbf{Q}) , \qquad (5)$$

associated with time translations (with respect to t).

• The dilation generator

$$D = tH - \frac{1}{4} \left(\mathbf{Q} \cdot \mathbf{P} + \mathbf{P} \cdot \mathbf{Q} \right) , \qquad (6)$$

which enforces scale transformations.

• The special conformal generator

$$K = Ht^2 - \frac{1}{2} (\mathbf{Q} \cdot \mathbf{P} + \mathbf{P} \cdot \mathbf{Q}) t + \frac{1}{2} MQ^2$$
(7)

that enforces inverse time translations.

This class of conformal Hamiltonians includes both the inverse square potential (ISP) $V(\mathbf{Q}) \propto Q^{-2}$ as well as contact interactions—most notably the two-dimensional delta [8], in addition to anisotropic versions of the ISP and the derivative-delta interaction in one spatial dimension [26]. While the symmetry properties of these realizations are generically the same for the whole conformal class, the analytical details of the solutions are specifically dependent on the chosen model. In this paper, we only consider the dAFF model of *non-contact interactions* that corresponds to a (0+1)-dimensional conformal field theory; in this setting, this amounts to the Lagrangian of Eq. (3).

The composition of the three basic types of transformations (5)-(7) generates the linear fractional transformation

$$\tilde{t} = \frac{\alpha t + \beta}{\gamma t + \delta} , \qquad (8)$$

governed by a matrix in the special linear group $SL(2,\mathbb{R})$ (with $\alpha\delta - \beta\gamma = 1$), and with transformed field variable

$$\tilde{\mathbf{Q}}(\tilde{t}) = \frac{\mathbf{Q}(t)}{\gamma t + \delta} \,. \tag{9}$$

At the quantum level, Eq. (9) is written in the Heisenberg picture, and has the same form as its classical counterpart. The associated symmetry group is $SL(2, \mathbb{R})$, which is homomorphic to SO(2, 1), with the generators $\{H, D, K\}$ satisfying the commutator relations [5]

$$[D,H] = -i\hbar H , \qquad [D,K] = i\hbar K , \qquad [H,K] = 2i\hbar D , \qquad (10)$$

which correspond to the one-dimensional case of the algebra of the generic conformal group. To introduce an appropriate Cartan-Weyl basis [58], one can replace H and K by the operators R and S defined by the linear combinations

$$R = \frac{1}{2} \left(\frac{1}{a} K + a H \right) , \qquad (11)$$

$$S \equiv -S' = \frac{1}{2} \left(\frac{1}{a} K - aH \right) , \qquad (12)$$

where a is an arbitrary parameter with dimensions of time (with R and S, as well as D, defined with dimensions of action). The commutators of the Cartan-basis generators are $[S, D] = -i\hbar R$, $[D, R] = i\hbar S$, and $[R, S] = i\hbar D$. The so(2, 1) algebra has rank one and dimension 3, with the Cartan generator R and the ladder operators L_{\pm} defined below, along with their corresponding commutator relations:

$$L_{\pm} = S \pm iD \tag{13}$$

$$[\hbar^{-1}R, L_{\pm}] = \pm L_{\pm} , \qquad [L_{+}, L_{-}] = -2\hbar R .$$
(14)

In this paper, we will use the operator S' = -S (with the sign reversed, as defined above), so that a more transparent physical interpretation can be given to the corresponding effective potential, as in Sec. VI—this choice can be found in the literature, for example, in Ref. [47].

B. Generalized Conformal Generators: Dynamics and Hamiltonian Formulation

The properties of the conformal generators are well-known from the exhaustive analysis of Ref. [5]. The operator R generates a compact subgroup [two-dimensional rotations as the O(2) subgroup of SO(2,1)], and S and D generate non-compact boosts. These distinct behaviors can be characterized by considering the generalized generator [5]

$$G = uH + vD + wK , (15)$$

with discriminant

$$\Delta = v^2 - 4uw \tag{16}$$

that determines its nature: rotations (elliptic type), for $\Delta < 0$, which include R; boosts (hyperbolic type), for $\Delta > 0$, which include both S and D; and parabolic ("lightlike") operators, for $\Delta = 0$, which include the original H and K operators.

A complete analysis [5] in terms of the operators (15) shows that G can be regarded as the generator of time evolution with respect to a modified, effective time τ . For the *d*-component field $\mathbf{Q}(t)$, from the definitions (5)–(7) and (15), the basic equation satisfied in the Heisenberg picture by the action of the operator G on the field variables $\mathbf{Q}(t)$ is

$$\frac{1}{i\hbar} \left[\mathbf{Q}(t), G \right] = f_G(t) \frac{d\mathbf{Q}(t)}{dt} - \frac{1}{2} \frac{df_G(t)}{dt} \mathbf{Q}(t) , \qquad (17)$$

where

$$f_G(t) = u + vt + wt^2 = \sigma |u + vt + wt^2| , \qquad (18)$$

with the sign $\sigma \equiv \sigma_G = \text{sgn}[f_G(t)]$ allowing for arbitrary linear combinations of the generalized generator (15). The dynamical equation (17), just as in the d = 1 case [5], can be simplified to a more standard form by redefining the dynamical time to τ according to

$$d\tau = \frac{dt}{u + vt + wt^2} , \qquad (19)$$

and the field variables to

$$\mathbf{r}(\tau) = \frac{\mathbf{Q}(t)}{|u + vt + wt^2|^{1/2}} \,. \tag{20}$$

In Eq. (20), the absolute value is needed to reproduce Eq. (17), with the auxiliary Eq. (18). Equation (19) can be easily integrated; particular cases (including R, S, D, and K) are listed in Ref. [5]. Thus, the dynamics in the Heisenberg picture is fully described by the equation

$$\frac{1}{i\hbar} \left[\mathbf{r}(\tau), G \right] = \frac{d\mathbf{r}(\tau)}{d\tau} , \qquad (21)$$

which shows that G acts as an effective Hamiltonian for time evolution. One can alternatively describe the dynamics in a generalized Schrödinger picture with respect to G, in terms of

evolving state vectors $|\Psi(\tau)\rangle_{(S)} \equiv |\Psi(\tau)\rangle = U_G(T) |\Psi(\tau_0)\rangle$, where $|\Psi(\tau_0)\rangle$ is the original state in the Heisenberg picture and

$$U_G(T) \equiv U_G(\tau;\tau_0) = \exp\left[-\frac{i}{\hbar}G\left(\tau - \tau_0\right)\right]$$
(22)

acts as an effective evolution operator; thus, the state vector $|\Psi(\tau)\rangle$ satisfies the generalized Schrödinger equation

$$G |\Psi(\tau)\rangle = i\hbar \frac{d |\Psi(\tau)\rangle}{d\tau}$$
 (23)

Then, Eqs. (21) and (23) describe the evolution of the field variables and states under the action of G in the Heisenberg and Schrödinger pictures respectively, with the new effective time τ ; in other words, G formally acts as $i\hbar\partial_{\tau}$. In these transformed variables, the time evolution only depends on the effective time difference $T = \tau - \tau_0$. It should be noted that this is *not* the original Schrödinger picture associated with the Hamiltonian H, and it is G-specific.

In principle, the analysis leading to Eqs. (19)–(23) allows a reformulation of the theory in terms of a new "Hamiltonian" $H_G(\mathbf{r}, \mathbf{p}) \equiv G$. This assignment $H_G \equiv G$, by abuse of notation, simply states that the value of G is to be rewritten in terms of the transformed canonical variables (\mathbf{r}, \mathbf{p}) . By construction, \mathbf{r} is given by Eq. (20), but \mathbf{p} has to be defined consistently in such a way that $|f_G|^{1/2} d\mathbf{r}/d\tau = f_G d\mathbf{Q}/dt - \dot{f}_G \mathbf{Q}/2$ (with the dot notation for derivatives with respect to the original time t)—an expression that can be read off by comparison of Eqs. (17) and (21) or from Eqs. (19)–(20). An obvious choice for \mathbf{p} is the one that satisfies $\mathbf{p} = M\dot{\mathbf{r}}$, whence the transformed momentum is given by

$$\mathbf{p} = \sigma \left| f_G \right|^{1/2} \left(\mathbf{P} - \frac{\dot{f}_G}{2f_G} M \mathbf{Q} \right) \,. \tag{24}$$

From Eq. (15), this implies that the operator $G = uH + v(tH + D_0) + w(t^2H + 2tD_0 + K_0) = f_GH + \dot{f}_GD_0 + wK_0$ [where D_0 and K_0 are the time independent parts in Eqs. (6) and (7)] takes the "normal" Hamiltonian form

$$G \equiv H_G(\mathbf{r}, \mathbf{p}) = \sigma \,\tilde{H}_G(\mathbf{r}, \mathbf{p}) \,, \tag{25}$$

where

$$\tilde{H}_G(\mathbf{r}, \mathbf{p}) = \frac{1}{2M} p^2 + \frac{1}{2} \frac{\lambda}{r^2} + \frac{M}{2} \left(-\frac{\Delta}{4}\right) r^2 .$$

$$\tag{26}$$

In Eq. (26), the usual convention $A = |\mathbf{A}|$ for spatial vectors is used, and the discriminant (16) is shown to be the same as $\Delta = \dot{f_G}^2 - 4wf_G$. The generalized momentum \mathbf{p} in Eq. (24) and the functional form of $H_G(\mathbf{r}, \mathbf{p})$ in Eqs. (25) and (26) can also be verified operationally at the classical level by a straightforward computation of the Hamiltonian from the transformed Lagrangian L_G , using Eqs. (3) and (19)–(20), as shown in Ref. [5].

C. Equivalence Classes of the Generalized Generators and Hamiltonian H_G

Equations (25)-(26) establish the Hamiltonian formulation of the generalized generators G, which are initially defined on the Lie algebra so(2,1). The Hamiltonian H_G is generated by a surjective mapping that endows them with a functional form with respect to the transformed canonical variables (20) and (24), such that its values only depend on the discriminant Δ . As a result, there is a continuous range of linear combinations G, with values of u, v, and w with the same discriminant, that give rise to the same effective Hamiltonian $H_G(\mathbf{r}, \mathbf{p})$. Moreover, the generators G define equivalence classes via the relation $G_1 \simeq G_2$ iff $H_{G_1} = H_{G_2}$, which identifies them by the same values of Δ (i.e., $\Delta_1 = \Delta_2$). This equivalence relation produces surprising connections between apparently distinct operators, such as, for example, H and K, or D and S'. The corresponding effective Hamiltonians are identical and yield equivalent models within the theory, with identical dynamical evolutions with respect to τ , as well as equivalent spectral properties. Therefore, for our main purposes, we can assume that G and H_G are identified without any further qualifications: the Hamiltonian/Lagrangian frameworks directly lead to the path integral and the spectral properties, and the practical use of an equal sign (instead of \simeq) is fully justified for the remainder of this paper.

In summary, the theory is organized in equivalence classes labeled by the value of the discriminant Δ . Moreover, these classes are further categorized in the three types of operator behavior: $\Delta \leq 0$ and $\Delta = 0$ (elliptic, hyperbolic, and parabolic), whose distinct properties can be understood from the change in sign of the quadratic potential in Eq. (26). These behaviors define a coarser equivalence relation leading to the three distinct broad classes of operators.

Regarding the spectral characterization of operators that we are addressing in this paper, the primary goal is to fully determine the eigenstates and eigenvalues of the generalized conformal generator G. We denote its eigenvalues as $\hbar \mathfrak{g}$, where \mathfrak{g} is dimensionless; then, from Eq. (25), it follows that

$$\hbar \mathfrak{g} = E_G = \sigma \tilde{E}_G \,, \tag{27}$$

in terms of the eigenvalues of the associated $H_G(\mathbf{r}, \mathbf{p})$. (It should be noted that, in Ref. [5], the eigenvalues of G are denoted by G'.) In light of Eq. (27), the eigenstates will be labeled interchangeably with \mathfrak{g} or E_G , and the latter may be simplified as E if there is no obvious conflict.

Equation (26) is an exact analog of a standard quantum-mechanical central problem with Hamiltonian and potential

$$\tilde{H}(\mathbf{r}, \mathbf{p}) = \frac{p^2}{2M} + \tilde{V}(r) , \qquad \tilde{V}(r) = \frac{1}{2}M\omega^2 r^2 + \frac{\hbar^2}{2M}\frac{g}{r^2} , \qquad (28)$$

which can be interpreted as representing a harmonic oscillator of squared "frequency"

$$\omega^2 = -\frac{\Delta}{4} \tag{29}$$

(including possibly imaginary frequencies) superimposed with the original inverse square potential. Even though our conformal problem has a different meaning, this analog parametrization is insightful and the "CQM frequency" from Eq. (29) will play an important role in describing the spectral solutions. In what follows, we will also use the functional notation $\tilde{H}[M, g, \omega] \equiv \tilde{H}_{\mathbf{r}, \mathbf{p}}[M, g, \omega]$ for the corresponding parameter dependence, where $\tilde{H}_{\mathbf{r}, \mathbf{p}} \equiv \tilde{H}(\mathbf{r}, \mathbf{p})$, with the appropriate canonical variables. A summary of the different Hamiltonian notations used in the main text is given in Table II.

TABLE II: Here we tabulate the different notations that we have used for specific Hamiltonians. All the expressions are written with position and momentum arguments denoted by \mathbf{r} and \mathbf{p} , respectively.

Notation	Expression	
$H(\mathbf{r},\mathbf{p})$	$\frac{p^2}{2M} + \frac{g}{r^2}$	
$ ilde{H}_G({f r},{f p})$	$\frac{p^2}{2M} + \frac{\lambda}{2r^2} + \frac{M}{2} \left(-\frac{\Delta}{4}\right) r^2$	
$G \equiv H_G$	$\sigma ilde{H}_G({f r},{f p})$	
$\tilde{H}(\mathbf{r},\mathbf{p})\equiv\tilde{H}_{\mathbf{r},\mathbf{p}}\equiv\tilde{H}[M,g,\omega]$	$\frac{p^2}{2M} + \frac{1}{2}M\omega^2 r^2 + \frac{\hbar^2}{2M}\frac{g}{r^2}$	
Ĥ	Generic quantum mechanical Hamiltonian	

The Hamiltonian function of Eq. (26) exhibits a scaling property that is a consequence of the definition (15), which is linear in the coefficients u, v, w, and with Δ given as a quadratic function (16). If this Hamiltonian function is rewritten in the generic analog form of Eq. (28), with frequency (29), then it satisfies the scaling

$$\tilde{H}[M,g,c\,\omega] = c\tilde{H}[M,g,\omega] , \qquad (30)$$

where c > 0 is an arbitrary real and positive scaling constant. In other words, the Hamiltonian is a homogenous function of first degree with respect to the frequency ω . An outline of the proof is as follows: if G is rescaled with the coefficients u' = cu, v' = cv, w' = cw, then it takes the value $G' = cG = \sigma c \tilde{H} [M, g, \omega]$, without a change in σ , but with the discriminant changing as $\Delta' = c^2 \Delta$. At the same time, Eq. (29) gives $\omega' = c \omega$ and this yields $G' = \sigma \tilde{H} [M, g, c \omega]$, whence Eq. (30) follows by comparing the two expressions for G'. This can be also be shown more generally directly at the level of Eq. (28) due to the general scaling properties of this Hamiltonian (as in ordinary dimensional analysis), in terms of the frequency parameter, with \tilde{H} and ω having the same scaling.

From Eq. (26), the spectral properties of the different types of conformal generators can be immediately understood. In this procedure, we can choose the prototypical generators R, H, and S' as representatives of the three main classes of operator behaviors: their general properties are shared by the generators G with $\Delta < 0$, $\Delta = 0$, and $\Delta > 0$, respectively. Then, $\omega = \sqrt{|\Delta|}/2$ for $\Delta < 0$ and $\omega = -i\sqrt{\Delta}/2$ for $\Delta > 0$ (the choice of sign for the latter is discussed in Sec. VI), with $\omega = 0$ for $\Delta = 0$. Moreover, we can spell out the parameter dependence of all the generators, starting with the prototypical operators R, H, and S', whose equivalence classes have discriminants $\Delta = -1$, $\Delta = 0$, and $\Delta = 1$, respectively. Thus,

$$R = \tilde{H}[M, g, \omega = 1/2], \qquad (31)$$

$$H = \tilde{H} \big[M, g, \omega = 0 \big] , \qquad (32)$$

$$S' = \tilde{H}[M, g, \omega = -i/2] .$$
(33)

In addition, as in Ref. [5], the parameters \hbar and M could be set equal to unity (though we will mostly keep the general parametrization for convenience). Most importantly, applying the scaling (30), it follows that $\tilde{H}_G = \sqrt{|\Delta|} \tilde{H}[M, g, \omega = 1/2]$ for elliptic operators, and

 $\tilde{H}_G = \sqrt{\Delta} \tilde{H}[M, g, \omega = -i/2]$ for hyperbolic operators. These scaling relations lead to the following complete characterization of the three classes of operators.

• Elliptic Generators, defined by $\Delta < 0$. These are "generalized R operators," of the form

$$G = \sigma \sqrt{|\Delta|} R , \qquad (34)$$

where R is the prototypical elliptic operator $(\Delta = -1)$ of Eqs. (11) and (31).

• Parabolic Generators, defined by $\Delta = 0$. These are "generalized *H* operators," of the form

$$G = \sigma H av{35}$$

where H is the prototypical parabolic operator of Eqs. (5) and (32).

 Hyperbolic Generators, defined by Δ > 0. These are "generalized S operators," of the form

$$G = \sigma \sqrt{\Delta} S' , \qquad (36)$$

where S' is the prototypical hyperbolic operator ($\Delta = 1$) of Eqs. (12) and (33).

This analysis verifies that the theory is organized in equivalence classes defined by the value of Δ , and where $\sigma = \operatorname{sgn}(f_G)$ just gives the "orientation" of the operator spectrum. Within each class, regardless of the values of u, v, and w, the effective Hamiltonian, dynamical evolution, and spectral properties are identical (except for a possible rescaling of the eigenvalues) but the form of the effective time τ (in terms of t) is specific to each operator. It is also noteworthy that the results of Eqs. (34)–(36) are completely general, and apply even for the parabolic operator K (which, according to this, gives the same propagator as H) and for the hyperbolic operator D (having the same propagator as S'). Incidentally, an alternative form of this theory is presented in Appendix A, which is restricted to the cases with $u \neq 0$ only, but has some advantages in terms of dimensional analysis.

In this section, we have established the general framework for the generators G. Next, we will use the parametrization of Eqs. (25)–(26) for a complete path-integral analysis of the spectral properties of all the conformal operators, for three associated operator classes (elliptic, parabolic and hyperbolic).

III. PATH-INTEGRAL FRAMEWORK: BASIC SETUP FOR CONFORMAL GENERATORS

All properties of a physical system can be completely characterized by using path-integral methods. This includes the spectral properties of operators relevant to the system. We will show how to derive such properties for the CQM operators R, H, and S, and, by extension, according to Eqs. (34)–(36), for the generalized generators G of Eq. (15). Due to the form of the Lagrangian (3) of the dAFF model and the special conformal operator (7), leading to the effective Hamiltonian (26), we need the general d-dimensional path-integral solution for the radial harmonic oscillator, along with the inverse square potential, i.e., for a central problem with Hamiltonian and potential given by Eq. (28). Of course, this is strictly needed for the multicomponent case of CQM, but it is also a useful approach to properly incorporate the inverse square potential with an appropriate analytic continuation (see below and Appendix B), even in the simplest case d = 1.

Remarkably, we are computing the path-integral propagators associated with the operators R, H, and S, as well as their generalized forms G, as generators of dynamical evolution. The specific functional form of these generators involves the generalized canonical variables defined in Eqs. (20) and (24) (or rescalings thereof). For H itself, the effective propagator is indeed the generator of dynamics in ordinary time t, but all the other generators involve dynamical evolutions with respect to their "natural" effective times τ . (In Appendix A, an alternative, dimensional framework is defined for the cases with $u \neq 0$, in terms of a rescaled time $\tilde{\tau}$, which plays a similar role.) Moreover, in addition to describing the dynamics, the propagators play the role of generating functions for the spectral decomposition and will allow us to find the spectral data, namely, the eigenvalues and eigenstates for the family of conformal generators.

A. Path-Integral Framework

We begin setting up the framework with some notational remarks. In the theory outline of this subsection and in Appendix B we typically denote by t the dynamical time, with \hat{H} describing a generic Hamiltonian, as in the usual quantum-mechanical applications. Within this general framework, we use a radial harmonic oscillator (properly generalized to include inverted oscillators and any number of dimensions) as an analog system for a generic conformal generator G. In this sense, when describing the spectral properties and dynamics generated by G, the general framework involves replacing \hat{H} by G or \tilde{H} , and t by τ .

Thus, our main focus will be on a generic Hamiltonian $\hat{H} = p^2/2M + V(\mathbf{r})$, with a time-independent and central potential $V(\mathbf{r}) = V(r)$ in *d* spatial dimensions. We begin our construction by considering the quantum-mechanical propagator

$$K_{(d)}(\mathbf{r}'',\mathbf{r}';t'',t') = \left\langle \mathbf{r}'' \left| \hat{T} \exp\left[-\frac{i}{\hbar} \int_{t'}^{t''} \hat{H} dt \right] \right| \mathbf{r}' \right\rangle , \qquad (37)$$

for a particle of mass M subject to the Hamiltonian \hat{H} . In Eq. (37), \hat{T} is the time-ordering operator. The corresponding path-integral expression

$$K_{(d)}(\mathbf{r}'',\mathbf{r}';t'',t') = \int_{\mathbf{r}(t')=\mathbf{r}'}^{\mathbf{r}(t'')=\mathbf{r}''} \mathcal{D}\mathbf{r}(t) \exp\left\{\frac{i}{\hbar}S\left[\mathbf{r}(t)\right]\left(\mathbf{r}'',\mathbf{r}';t'',t'\right)\right\}$$
(38)

involves the classical action functional $S[\mathbf{r}(t)](\mathbf{r}'', \mathbf{r}'; t'', t')$ for "paths" $\mathbf{r}(t)$ connecting the end points $\mathbf{r}(t') = \mathbf{r}', \mathbf{r}(t'') = \mathbf{r}''$. For time-independent potentials, the propagator $K_{(d)}(\mathbf{r}'', \mathbf{r}'; t'', t') \equiv K_{(d)}(\mathbf{r}'', \mathbf{r}'; T)$ is a function of T = t'' - t' alone and not of the individual endpoint times; we will use this assumption for the remainder of the paper. Specifically, for the generic conformal generator G, the associated propagator is given by $K_{(d)}^{(G)}(\mathbf{r}'', \mathbf{r}'; T) = \langle \mathbf{r}'' | U_G(T) | \mathbf{r}' \rangle$, with $U_G(T)$ as in Eq. (22), in terms of the effective time τ .

In Appendix B, we review the required construction for central potentials in hyperspherical coordinates, and specifically for the radial harmonic oscillator. In general, the propagator $K_{(d)}(\mathbf{r}'', \mathbf{r}'; T)$ can be formally rewritten in hyperspherical polar coordinates [33, 34, 59], which yield a complete set of angular functions, the *d*-dimensional hyperspherical harmonics $Y_{lm}(\Omega)$, where the quantum labels correspond to the *d*-dimensional angular momentum with numbers *l* and *m*, and such that the set *m* takes $g_l = (2l + d - 2)(l + d - 3)!/l!(d - 2)!$ values as multiplicity for a given *l* (Chap. XI in Ref. [59]). The corresponding partial wave expansions of $K_{(d)}(\mathbf{r}'', \mathbf{r}'; T)$ read

$$K_{(d)}(\mathbf{r}'',\mathbf{r}';T) = (r''r')^{-(d-1)/2} \sum_{l=0}^{\infty} \sum_{m=1}^{g_l} Y_{lm}(\mathbf{\Omega}'') Y_{lm}^*(\mathbf{\Omega}') K_{l+\nu}(r'',r';T)$$
(39)

$$= \frac{\Gamma(\nu)}{2\pi^{d/2}} (r''r')^{-(d-1)/2} \sum_{l=0}^{\infty} (l+\nu) C_l^{(\nu)}(\cos\psi_{\mathbf{\Omega}'',\mathbf{\Omega}'}) K_{l+\nu}(r'',r';T) , \qquad (40)$$

where $\nu = d/2 - 1$, $\cos \psi_{\Omega'',\Omega'} = \hat{\mathbf{r}}'' \cdot \hat{\mathbf{r}}'$ (with $\hat{\mathbf{r}} = \mathbf{r}/r$), and the addition theorem for hyperspherical harmonics provides the expression in terms of Gegenbauer polynomials $C_l^{(\nu)}(x)$. In Eqs. (39) and (40), and subsequent expressions, we will use the radial-coordinate symbol $r = |\mathbf{r}|$. For central potentials, the radial propagator $K_{l+\nu}(r'', r'; T)$ is independent of the angular coordinates and quantum numbers \boldsymbol{m} . The chosen normalization of radial prefactors is such that the radial propagator $K_{l+\nu}(r'', r'; T)$ satisfies the composition property

$$K_{l+\nu}(r'',r';t''-t') = \int_0^\infty dr \, K_{l+\nu}(r'',r;t''-t) K_{l+\nu}(r,r';t-t') \,. \tag{41}$$

For the radial propagator associated with G as generalized Hamiltonian with effective time τ , the notation $K_{l+\nu}^{(G)}(r'', r'; T)$ will be used for the remainder of the paper.

An explicit expression for the radial propagator in Eqs. (39)–(40) can be derived in the time-sliced path integral. In short, for a time lattice $t_j = t' + j\epsilon$, for the time interval T = t'' - t' corresponding to the end points $\mathbf{r}_0 \equiv \mathbf{r}'$ and $\mathbf{r}_N \equiv \mathbf{r}''$; in this lattice, $\epsilon = T/N$, with $j = 0, \dots, N$, with $t_0 \equiv t'$ and $t_N \equiv t''$, such that $\mathbf{r}_j = \mathbf{r}(t_j)$. Then, with the restriction to the half-line $r(t) \ge 0$, the radial propagator admits a formal continuum-limit representation [1–3, 60, 61]

$$K_{l+\nu}(r'', r'; T) = \int \mathcal{D}r(t) \ w_{l+\nu}[r^2] \exp\left\{\frac{i}{\hbar} \int_{t'}^{t''} dt \left[\frac{M}{2} \dot{r}^2 - V(r)\right]\right\} , \qquad (42)$$

in terms of the usual one-dimensional path-integral measure $\mathcal{D}r(t)$ (as in Cartesian coordinates), but with a nontrivial radial functional weight $w_{l+\nu}[r^2] = \lim_{N\to\infty} w_{l+\nu}^{(N)}[r^2]$ [see Eq. (B5)], which involves Bessel functions of order $l + \nu$. This has been called a Besselian path integral [1, 2].

The general expressions for Besselian path integrals are given in Appendix B 1, including statements on the associated property of interdimensional dependence [62, 63]. This framework leads to the theorem for the insertion of inverse square potential terms as generalized angular momenta, as proved in Refs. [61, 63, 64]. More precisely:

Theorem. Given a potential $\tilde{V}(r) = V(r) + (\hbar^2/2M)gr^{-2}$, the propagator $\tilde{K}_{l+\nu} \equiv K_{l+\nu}[\tilde{V}]$ associated with $\tilde{V}(r)$ in Eq. (42) is equivalent to the propagator $K_{\mu}[V]$ with the reduced potential V(r) and an effective, typically non-integer, angular momentum variable

$$\mu = \sqrt{(l+\nu)^2 + g} , \qquad (43)$$

such that

$$\tilde{K}_{l+\nu} \equiv K_{l+\nu}[\tilde{V}] = K_{\mu}[V] .$$
(44)

In short, this procedure expresses a practical rule for the insertion of inverse square potential terms by the straightforward analytic continuation (43), which amounts to the introduction of a continuous, effective angular momentum $l_{\text{eff}} = \mu - \nu$. As a result of this asymptotic recombination (see Appendix B1), the theorem defines the potential \tilde{V} to be used in Eq. (28).

B. Propagator for Generalized Radial Harmonic Oscillator

The derivation of the general path integral for the radial propagator of the *d*-dimensional harmonic oscillator involves explicit use of the radial path integral (42). With the inclusion of an additional inverse square potential, i.e., for a Hamiltonian $\tilde{H}(\mathbf{r}, \mathbf{p})$ with an effective potential of the form (28), the propagator takes the form

$$K_{l+\nu}^{(RHO)}(r'',r';T) = \frac{M\omega}{i\hbar\sin\omega T}\sqrt{r'r''} \exp\left[\frac{iM\omega}{2\hbar}\left(r'^2 + r''^2\right)\cot\omega T\right]I_{\mu}\left(\frac{M\omega r'r''}{i\hbar\sin\omega T}\right) ,\quad (45)$$

with a conformal parameter index given by Eq. (43). The path-integral result (45) for the "radial harmonic oscillator" was first derived by Peak and Inomata [64], and has a broad range of applications; it is established by using Weber's second exponential integral for Bessel functions [65] combined with appropriate recursion relations [1–3, 64]. This is briefly reviewed and discussed in Appendix B. Incidentally, while Eq. (45) refers to the problem with an inverse square potential, i.e., of the form (44), we will keep the notation without the tilde for the sake of simplicity.

We will apply the path integral (45) to the characterization of the spectral properties of the generator G of Eqs. (25)–(26), where \tilde{H}_G , given in Eq. (26), is equivalent to Eqs. (28)– (29). This applies to the operator H as well as to R and S' = -S, defined in Eqs. (11) and (12), and related to the generic Hamiltonian of Eq. (28) with the assignments of Eqs. (31) and (33). Moreover, from the analysis of Eqs. (34)–(36), these assignments allow for a thorough characterization of the three operator classes: parabolic, elliptic, and hyperbolic. In this generic framework, the CQM frequency ω to use in Eq. (45) is given through Eq. (29).

Additionally, for some of the applications below, we will compute the path integrals with appropriate analytic continuations. As will be further analyzed in Sec. VI, in Eqs. (31)–(33), the operator S is obtained from R by an analytic continuation $\omega \to -i\omega$ that completely changes the nature of the spectrum (both $\omega \to \mp i\omega$ are equally valid, but the negative sign has operational advantages discussed in Sec. VI). Another such extension is the commonly used continuation to Euclidean time; in this context, for the propagator $K_{l+\nu}(r'', r'; T)$, the following formal replacement is made: $T \rightarrow -iT$. For the propagator of Eq. (45), the corresponding Euclidean-time path integral reads

$$K_{l+\nu}^{(RHO)}(r'',r';-iT) = \frac{M\omega}{\hbar\sinh\omega T}\sqrt{r'r''} \exp\left[-\frac{M\omega}{2\hbar}\left(r'^2+r''^2\right)\coth\omega T\right]I_{\mu}\left(\frac{M\omega r'r''}{\hbar\sinh\omega T}\right).$$
(46)

It should be stressed that the propagators (45)–(46) correspond specifically to the Hamiltonian \tilde{H} of the form of Eq. (28); for CQM, Eq. (25) provides the connection with the generic Hamiltonian H_G associated with the generator G (with possibly an extra sign σ).

In the next sections, for our general description of the generators G, we will use the notations $K_{l+\nu}^{(\text{ell})}$, $K_{l+\nu}^{(\text{par})}$, and $K_{l+\nu}^{(\text{hyp})}$ for the generic propagators of the three classes of generators.

IV. PATH-INTEGRAL SPECTRAL ANALYSIS OF THE GENERIC ELLIPTIC (R-LIKE) GENERATORS

In this section, we analyze the propagators for generic elliptic generators, including the prototypical operator R. The analysis of the spectral decomposition is straightforward for these operators, as it consists of a purely discrete spectrum.

A. Elliptic Generators: Spectral Properties and Path-Integral Propagator

From Eq. (31), it is clear that R can be viewed as an analog harmonic oscillator with an inverse square potential; thus, the propagator is given in Eq. (45). Moreover, a generic elliptic operator G, with a discriminant $\Delta < 0$, inherits the basic spectral properties of R, according to Eq. (34), i.e., $G = \sigma \sqrt{|\Delta|} R$. Thus, it follows that the eigenstates of an elliptic operator G are the same as those of R, with a spectrum of eigenvalues $\hbar \mathfrak{g}$ such that

$$\mathfrak{g} = \sigma \sqrt{|\Delta|} \mathfrak{r} \tag{47}$$

is rescaled from the spectrum $\hbar \mathfrak{r}$ of R. In addition, the functional form of the ellipticgenerator propagator

$$K_{l+\nu}^{(\text{ell})}(r'', r'; T) = K_{l+\nu}^{(RHO)}(r'', r'; T) , \qquad (48)$$

where $K_{l+\nu}^{(RHO)}$ is given in Eq. (45), applies to this entire family, with a CQM frequency $\omega = \sqrt{|\Delta|}/2$, according to Eq. (29).

B. Elliptic Generators: Spectral Analysis

The spectral decomposition of the relevant operators, in terms of eigenfunctions and eigenvalues, can be obtained by performing expansions of the propagator through established identities, with series or integrals that select appropriate discrete or continuous parts of the spectrum. This can be done by identifying expressions of the form $f(A) = \sum_{n} f(a_n)P_n$, and corresponding generalizations with integrals, for a given operator A with eigenvalues a_n , where f is a function of the operator and $P_n = |n\rangle \langle n|$ is the orthogonal projector. The most basic expansion of this kind in the path-integral approach is provided by the propagator itself, due to its basic definition (37) in terms of the time evolution operator. As shown below, this leads to a discrete spectrum for the elliptic operators.

The spectral decomposition for the operator R and all elliptic generators can be made explicit using the bilinear generating function of associated Laguerre polynomials known as the Hille-Hardy formula (pp. 189-190 in Ref. [59]),

$$\frac{1}{(xyz)^{\mu/2}(1-z)} \exp\left[-z\frac{(x+y)}{(1-z)}\right] I_{\mu}\left(\frac{2\sqrt{xyz}}{1-z}\right) = \sum_{n=0}^{\infty} \frac{n!}{\Gamma(n+\mu+1)} L_{n}^{(\mu)}(x) L_{n}^{(\mu)}(y) z^{n} ,$$
(49)

with |z| < 1. The definition of normalization of the generalized Laguerre polynomials used in Eq. (49) is that of Ref. [59], corresponding to its relation $L_n^{(\mu)}(x) = \binom{n+\mu}{n} {}_1F_1(-n,\mu+1;x)$ to the confluent hypergeometric function.

Then, the spectral-series expansion of the Euclidean-time propagator $K_{l+\nu}^{(\text{ell})}(r'', r'; -iT)$, from Eqs.(46) and (48), can be recast in the form

$$K_{l+\nu}(r'', r'; -iT) = \sum_{n} e^{-E_n T/\hbar} \mathcal{U}_{n,l}(r'') \mathcal{U}_{n,l}^*(r') , \qquad (50)$$

with the substitutions $x = M\omega r'^2/\hbar$, $y = M\omega r''^2/\hbar$, and $z = e^{-2\omega T}$, so that $2z/(1-z) = (1+z)/(1-z)-1 = \coth \omega T - 1$. Equation (50) provides the expansion for the radial propagator in terms of reduced radial wave functions $\mathcal{U}_{n,l}(r)$, and a similar expansion involves the *d*dimensional wave functions $\psi_{n,l,m}(\mathbf{r})$ with the full-fledged propagator of Eq. (39). By direct inspection, the eigenvalues $\tilde{E}_{G,n}$ corresponding to \tilde{H}_G are of the form $\tilde{E}_{G,n} = \hbar\omega (1+\mu+2n)$; thus, from Eqs. (27), (29), and (47), and using the eigenvalues $\hbar \mathbf{r}_n$ of R,

$$E_{G,n} = \sigma \tilde{E}_{G,n} = \hbar \mathfrak{g}_n = \sigma \sqrt{|\Delta|} \,\mathfrak{r}_n = \sigma \hbar \omega \left(1 + \mu + 2n\right) \,, \tag{51}$$

where

$$\mathbf{r}_n = \mathbf{r}_0 + n , \quad \mathbf{r}_0 = \frac{1}{2}(1+\mu) ,$$
 (52)

and the parameter μ is specified by Eq. (43); in Ref. [5] this is explicitly written in the form $\mu = 2\mathfrak{r}_0 - 1$. The spectrum (52) is the discrete series $D_{\mathfrak{r}_0}^+$ bounded below of the unitary, irreducible representations of the group SO(2, 1) [58]. Furthermore, their common eigenstates $\mathcal{U}_{n,l}(r)$ are identified as

$$\mathcal{U}_{n,l}(r) = \sqrt{\frac{2\Gamma(n+1)}{\Gamma(1+\mu+n)}} r^{-1/2} \left(\sqrt{\frac{M\omega}{\hbar}} r\right)^{\mu+1} \exp\left(-\frac{M\omega}{2\hbar} r^2\right) L_n^{(\mu)} \left(\frac{M\omega}{\hbar} r^2\right) .$$
(53)

C. Elliptic Generators: Conclusions

Some remarks on the nature of the mathematical results for elliptic operators are in order. The derivation of the spectral properties for elliptic generators involves an expansion of the form (50), which relies on the well-established identity (49). The Euclidean-time framework permits a rigorous procedure for the identification of the corresponding eigenstates and eigenvalues. This is a well-known theorem for the radial harmonic oscillator [1–3], which corresponds to our problem via the analog representation (48).

These rigorous results are valid for arbitrary dimensionality d; in particular, they agree with the corresponding expressions in Sec. 4 of the original dAFF model of Ref. [5] for d = 1. It should be noted that, for d > 1, the value of \mathbf{r}_0 is also dependent on the angular momentum l. The d-dimensional wave functions $\psi_n(\mathbf{r})$ can be assembled by considering the corresponding spectral expansion of the propagator of Eq. (39), whence

$$\psi_{n,l,\boldsymbol{m}}(\mathbf{r}) = r^{-(d-1)/2} \,\mathcal{U}_{n,l}(r) \,Y_{l\boldsymbol{m}}(\boldsymbol{\Omega}) \,, \qquad (54)$$

which only changes the factor $r^{-1/2}$ in Eq. (53) to $r^{-d/2}$, and inserts the angular dependence. As expected, the derived results can be interpreted as a rescaled, ISP-extended version of those for the standard isotropic oscillator in quantum mechanics. For example, the energy levels of the isotropic oscillator are given by $E_n = \hbar \omega (N + d/2)$, with principal quantum number $N = 2n + l + \nu + 1$ (and *n* being the radial quantum number), which is extended with the rule (43), so that $\tilde{E}_n = \hbar \omega (2n + \mu + 1)$, as in Eq. (51). A similar replacement (43) for the wave functions confirms this interpretation for Eq. (53).

V. PATH-INTEGRAL SPECTRAL ANALYSIS OF THE GENERIC PARABOLIC (H-LIKE) GENERATORS

In this section, we analyze the propagators for generic parabolic generators, including the prototypical operator H. The case of H (and parabolic operators by extension) is particularly interesting because it describes the initial Hamiltonian, and a number of approaches can be successfully applied, including a novel technique ("Fourier method," Appendix C) to derive the eigenfunctions of operators with a continuum spectrum.

A. Hamiltonian H and Parabolic Generators: Spectral Properties and Path-Integral Propagator

A generic parabolic operator G, with a discriminant $\Delta = 0$, inherits the basic spectral properties of the original Hamiltonian H, according to Eq. (35), i.e., $G = \sigma H$. Its eigenstates are the same as those of H, with a rescaled spectrum of eigenvalues $\hbar \mathfrak{g}$ such that $\mathfrak{g} = \sigma E/\hbar$ is determined from the spectrum E of H. In particular, the functional form of the propagator $K_{l+\nu}^{(H)}$ applies without any modifications to this family of parabolic generators, with a CQM frequency $\omega = 0$.

Clearly, the Hamiltonian operator H corresponds to a pure inverse square potential. The corresponding path integral could be computed separately (see Appendix B), but it is straightforward to derive it by taking the limit $\omega \to 0$ of the generalized harmonic oscillator propagator of Eq. (45). Thus, the propagator corresponding to the operator H and other members of its parabolic family is given by

$$K_{l+\nu}^{(\text{par})}(r'',r';T) = K_{l+\nu}^{(H)}(r'',r';T) = \lim_{\omega \to 0} K_{l+\nu}^{(RHO)}(r'',r';T)$$
(55)

$$= \frac{M}{i\hbar T} \sqrt{r'r''} \exp\left[\frac{iM}{2\hbar T} (r'^2 + r''^2)\right] I_{\mu}\left(\frac{Mr'r''}{i\hbar T}\right) .$$
(56)

The spectral decomposition and spectral properties of this Hamiltonian model can be analyzed in several ways. We will consider the following 3 methods: (i) the zero-frequency limit of the spectral decomposition of the radial harmonic oscillator, using the Mehler-Heine formula for generalized Laguerre polynomials; (ii) direct evaluation of the spectral decomposition from the propagator (45) using Weber's second exponential integral; (iii) computation of the eigenfunctions with the Fourier transform of the propagator (45), using an integral representation of the product of Bessel functions.

B. Parabolic Generators: Zero-Frequency Limit of the Propagator—Mehler-Heine Formula

The first method uses the Mehler-Heine limit formula for the Laguerre polynomials (p. 191 in Ref. [59]),

$$\lim_{n \to \infty} n^{-\mu} L_n^{(\mu)} \left(\frac{x}{n}\right) = x^{-\mu/2} J_\mu(2\sqrt{x}) , \qquad (57)$$

with the following sequence of formal steps. Starting from the spectral decomposition expansion (50), with Eqs. (51) and (53), and $\sigma = 1$, we can turn the series into an integral by realizing that the eigenvalue level spacing $E_{n+1} - E_n = 2\hbar\omega$ asymptotically approaches zero. Thus, the energy values E_n approach the zero limit except when $n \to \infty$, so that the actual energy levels E of the continuum satisfy the asymptotic approximate replacement $E \approx 2\hbar\omega n$. Correspondingly, the asymptotic conversion into an integral follows the rule $\sum_n \approx \int dE/(2\hbar\omega)$, which amounts to $\sum_n 1/n \approx \int dE/E$. In addition, defining $x/n = Mr^2\omega/\hbar$ to match the argument of the Laguerre polynomials, this gives $x = k^2r^2/4$, with k such that $E = \hbar^2 k^2/2M$, and the argument of the Bessel function in Eq. (57) becomes $2\sqrt{x} = kr$. Then, approximating the eigenfunction coefficient in Eq. (53), for $n \gg 1$, as $c_{n,l} = \sqrt{2\Gamma(n+1)/\Gamma(1+\mu+n)} \approx \sqrt{2/n^{\mu}}$, and $e^{-M\omega r^2/2\hbar} \approx 1$, the asymptotic discrete eigenfunctions become

$$\mathcal{U}_{n,l}(r) \approx \sqrt{2} n^{-1/2} (x/r)^{1/2} J_{\mu}(kr) ,$$

where $x/r = k^2 r/4$. Thus, Eq. (50), for $n \gg 1$, turns into

$$K_{l+\nu}^{(\text{par})}(r'',r';T) \approx \int_0^\infty \frac{dE}{E} n \, e^{-iET/\hbar} \, \mathcal{U}_{n,l}(r'') \, \mathcal{U}_{n,l}^*(r') \\ \approx \int_0^\infty dE e^{-iET/\hbar} \left[\sqrt{\frac{M}{\hbar^2}} \sqrt{r} J_\mu(kr'') \right] \left[\sqrt{\frac{M}{\hbar^2}} \sqrt{r} J_\mu(kr') \right] \,, \qquad (58)$$

which takes the form of the continuous spectral decomposition

$$K_{l+\nu}^{(\text{par})}(r'',r';T) = \int_0^\infty dE e^{-iET/\hbar} \mathcal{U}_{E,l}(r'') \mathcal{U}_{E,l}^*(r') , \qquad (59)$$

where the continuous energy eigenfunctions are given by

$$\mathcal{U}_{E,l}(r) = \sqrt{\frac{M}{\hbar^2}} \sqrt{r} J_{\mu}(kr) , \qquad (60)$$

with $k = \sqrt{2ME/\hbar^2}$. By the normalized form of the expansion (59), the eigenstates (60) are Dirac-normalized with respect to the variable E. Therefore, through the zero-frequency limit, we have derived the continuous analog of the spectral decomposition series (50), for a spectrum with values restricted to the half-energy line $E \in [0, \infty)$ and eigenstates (60).

C. Parabolic Generators: Spectral Decomposition via Weber's Second Exponential Integral

The second method involves writing the spectral decomposition directly in its continuous spectral form (59), which will be expressed in the Euclidean-time version,

$$K_{l+\nu}^{(\text{par})}(r'',r';-iT) = \int_0^\infty dE \, e^{-ET/\hbar} \, \mathcal{U}_{E,l}(r'') \, \mathcal{U}_{E,l}^*(r') \,, \tag{61}$$

where $K_{l+\nu}^{(\text{par})}$ is given by Eq. (56), and the corresponding real-time propagator involves the replacement $T \rightarrow iT$. No additional assumptions are needed as the propagator is expanded in this form by construction from a well-established integral identity for Bessel functions (Sec. 13.31 of Ref. [65] and 10.22.67 of Ref. [66]): Weber's second exponential integral (B11). With the substitutions $E = x^2$, $T/\hbar = c^2$, $\sqrt{2M} r'/\hbar = a$, $\sqrt{2M} r''/\hbar = b$ in Eq. (B11), and insertion of an additional factor $2M\sqrt{r'r''}/\hbar^2$, the identity turns formally into the desired Eq. (61), with the same continuous energy eigenfunctions as displayed in Eq. (60).

D. Parabolic Generators: Spectral Decomposition via a Fourier Method

The third method is based on a simple restatement of the spectral expansion of an operator with a continuous spectrum. From the continuous spectral decomposition (59), the inverse Fourier transform with respect to the variable E converts the propagator $K_{l+\nu}^{(\text{par})}(r'',r';T)$ into the wave-function product

$$\mathcal{U}_{E,l}(r'') \,\mathcal{U}_{E,l}^*(r') = \frac{1}{2\pi\hbar} \,\int_{-\infty}^{\infty} dT \,\exp\left(\frac{iET}{\hbar}\right) \,K_{l+\nu}^{(\text{par})}(r'',r';T) \,. \tag{62}$$

This yields an explicit Fourier-integral representation of the spectral-decomposition wavefunction product. Evaluation of this spectral integral can be established from the integral representation (D1) of the product of Bessel functions (Sec. 13.7 of Ref. [65]). In practice, we are interested in the limit $c \to 0^+$, for which the integral (D1), with s = c + 2it, becomes

$$J_{\mu}(z')J_{\mu}(z'') = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \exp\left(it\right) \exp\left[i\frac{(z'^2 + z''^2)}{4t}\right] I_{\mu}\left(\frac{z'z''}{2it}\right) \frac{dt}{t} .$$
 (63)

In Eq. (63), the following substitutions are made on the right-hand side: z' = kr', z'' = kr'', and $t = ET/\hbar$, with the time interval T as integration variable. Then, the exponential factors become $\exp\left[i(z'^2 + z''^2)/4t\right] = \exp\left[iM(r'^2 + r''^2)/(2\hbar T)\right]$, with the argument of the Bessel function being $z'z''/2it = Mr'r''/i\hbar T$. This shows that the integrand of Eq. (D1) is proportional to the propagator $K_{l+\nu}^{(\text{par})}$ of Eq. (56) after inserting an additional factor $M\sqrt{r'r''}/\hbar^2$, so that

$$\left[\sqrt{\frac{M}{\hbar^2}}\sqrt{r}J_{\mu}(kr'')\right]\left[\sqrt{\frac{M}{\hbar^2}}\sqrt{r}J_{\mu}(kr')\right] = \frac{1}{2\pi\hbar}\int_{-\infty}^{\infty}dT\,\exp\left(\frac{iET}{\hbar}\right)\,K_{l+\nu}^{(\text{par})}(r',r'';T)\,.$$
 (64)

Comparison of Eqs. (62) and (64) shows that the energy eigenfunctions are given by Eq. (60).

Incidentally, the explicit form of the wave function product in Eq. (62) can also be established via the zero-frequency limit of the radial harmonic oscillator propagator. In this limit, the generalized Laguerre polynomials turn into Bessel functions, and the coefficients of the Fourier series (50) (where E_n is linear in n) turn into the required Fourier integral.

E. Parabolic Generators: Conclusions

In conclusion, we have shown by several techniques that the Dirac-normalized continuous eigenstates are given by Eq. (60). All the methods yield the same result, with a purely continuous spectrum restricted to the non-negative energy half-line $E \in [0, \infty)$. This result is valid for arbitrary dimensionality d; in particular, it agrees with the value in Eq. (A.18) of the original dAFF model of Ref. [5] for d = 1, and can be interpreted with the physical insights known for analog elementary problems (e.g., reduces to the free particle when g = 0). For the d-dimensional model, the radial wave function is $\mathcal{U}_{E,l}(r)/r^{(d-1)/2}$.

A number of final remarks on the nature of the mathematical results for parabolic operators are in order. A rigorous characterization of the spectral decomposition as a theorem is provided by the second method, which relies on the direct use of the well-known Weber's second exponential integral; this is similar to the procedure followed for elliptic operators via the Hille-Hardy formula—however, such constructions require an ad hoc guessing of the correct expression that mimics this resolution. By contrast, the third method inverts the process and introduces a systematic calculational procedure whereby the wave functions are obtained by a direct computation, provided that the integrals can be performed. In Appendix C, we elaborate on this novel Fourier method, stating it in more general terms and addressing its basic properties; in its more general form, it will be used in the next section for hyperbolic operators. It should also be noticed that the first method was presented as a formal procedure that shows consistency with the more general result for elliptic operators via an appropriate limit. While these operators, elliptic and parabolic, are of a different nature, they are still related by a continuous transformation with the parameter ω enforcing the asymptotic transition.

Interestingly, the derivations of this section show that there are simple connections between the known mathematical theorems used. Specifically, Weber's second exponential integral has the following two properties: (i) it is a restatement of the continuous limit of the Hille-Hardy formula; (ii) it can be interpreted as a Fourier transform of the integral representation (D1). To our knowledge, these statements are not spelled out in the mathematics literature.

VI. PATH-INTEGRAL SPECTRAL ANALYSIS OF THE GENERIC HYPER-BOLIC (S-LIKE) GENERATORS

In this section, we analyze the propagators for generic hyperbolic generators, including the prototypical operator S'. The analysis of the spectral decomposition is analytically more involved due to the nature of the continuous spectrum.

A. Hyperbolic Generators: Spectral Properties and Path-Integral Propagator

A generic hyperbolic operator G, with a discriminant $\Delta > 0$, inherits the basic spectral properties of S or S', according to Eq. (36), i.e., $G = \sigma \sqrt{\Delta} S'$. Thus, we can proceed as for the other families of operators: the eigenstates of a hyperbolic operator G are the same as those of S', with a spectrum of eigenvalues $\hbar \mathfrak{g}$ such that

$$\mathfrak{g} = \sigma \sqrt{\Delta} \mathfrak{s}' \tag{65}$$

is rescaled from the spectrum $\hbar \mathfrak{s}'$ of S'.

The behavior of the hyperbolic generators falls within the analog model of the generic extended radial harmonic oscillator Hamiltonian \tilde{H} of Eq. (28). Therefore, the functional form of the propagator $K_{l+\nu}^{(hyp)}$ applies to the whole family of hyperbolic generators, with a CQM frequency $\omega = -i\sqrt{\Delta}/2$, from Eq. (29). The imaginary frequency can be obtained with the prescription to analytically continue the potential according to $\omega \rightarrow -i\omega$, with ω a real "frequency." The result of the analytic continuation $\omega \to -i\omega$ is to generate an inverted harmonic oscillator that overlaps with the inverse square potential. By the form of this potential, we can predict that all values of the "energy" are possible, forming a continuum from minus infinity to plus infinity. This physical statement will be verified with the path-integral calculation that follows. It should be noted that systems with such a continuous spectrum, unbounded from below and above, are not common and represent idealized models (as actual physical systems exhibit energy bounds). But the treatment of such systems with a general path-integral technique is of theoretical and practical interest, and the results of the previous subsection can be applied directly. For example, the operator S' (being an inverted radial oscillator) is related to the one-dimensional inverted harmonic oscillator, which is a system that is related to a variety of physical problems of current relevance [67, 68].

By comparison with the propagator equation (45), which is completely general for complex values of the parameter ω , and with the replacement $\omega \to -i\omega$, the propagator associated with S' and all hyperbolic operators is given by

$$K_{l+\nu}^{(\text{hyp})}(r'',r';T) = \frac{M\omega}{i\hbar\sinh(\omega T)}\sqrt{r'r''} \exp\left[\frac{iM\omega}{2\hbar}\left(r'^2 + r''^2\right)\coth(\omega T)\right] I_{\mu}\left(\frac{M\omega r'r''}{i\hbar\sinh(\omega T)}\right).$$
(66)

Incidentally, either extension $\omega \to \pm i\omega$ is equally acceptable. This can be seen directly from the Hamiltonian, or explicitly from the propagator. However, the choice $\omega \to -i\omega$ has the operational advantage that it preserves the boundary condition at infinity corresponding to retarded Green's functions solutions (either bound states or outgoing waves). This statement can be verified in an explorative manner by a simple WKB evaluation, which is asymptotically exact and yields a solution function $\mathcal{U} \sim r^{-1/2+2\kappa} \exp(-M\omega r^2/2\hbar) \xrightarrow{(\omega \to -i\omega)}{\longrightarrow}$ $\mathcal{U} \sim r^{-1/2+2i\kappa} \exp(iM\omega r^2/\hbar)$, where $\kappa = \tilde{E}_G/(2\hbar\omega)$ (see Appendix E); and it is further confirmed by the exact solution in terms of Whittaker functions (see final results in this subsection). The advantage of this rule is that one can directly extrapolate by analytic continuation the correct solutions, including Green's functions, from one sector of the theory to another (for example, as shown below, from the inverted to the regular harmonic oscillator).

B. Hyperbolic Generators: Spectral Analysis

The spectral decomposition can be obtained from the Fourier integral (C2), which, with the substitution $\zeta = \omega T$ takes the explicit form

$$\mathcal{U}_{E,l}(r'') \mathcal{U}_{E,l}^*(r') = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} dT \exp\left(\frac{iET}{\hbar}\right) K_{l+\nu}^{(\text{hyp})}(r'', r'; T) = \frac{1}{2\pi} \frac{M}{\hbar^2} \sqrt{r'r''} \frac{1}{i} \underbrace{\int_{-\infty}^{\infty} \frac{d\zeta}{\sinh\zeta} \exp\left(2i\kappa\zeta\right) \exp\left(i\beta \coth\zeta\right) I_{\mu}\left(\frac{\alpha}{i\sinh\zeta}\right)}_{\mathcal{I}}, \quad (67)$$

where

$$\kappa = \frac{E}{2\hbar\omega} , \quad \beta = \frac{1}{2} \left(x' + x'' \right) , \quad \alpha = \sqrt{x'x''} ,$$
with $x' \equiv \check{r}'^2 = \frac{M\omega}{\hbar} r'^2$ and $x'' \equiv \check{r}''^2 = \frac{M\omega}{\hbar} r''^2 .$
(68)

In all the ensuing equations below, the dimensionless coordinate $\check{r} = \sqrt{M\omega/\hbar} r$ is used.

The integral \mathcal{I} in Eq. (67) can be evaluated by using the approach of Ref. [69] (Sec. 6.1). This involves developing integral representations for the product of two Whittaker functions $M_{\kappa,\mu/2}(z)$ and $W_{\kappa,\mu/2}(z)$ (in different combinations of M and W as well as function indices); the regularized Whittaker function $\mathcal{M}_{\kappa,\mu/2}(z) = M_{\kappa,\mu/2}(z)/\Gamma(1+\mu)$ of Ref. [69] (that removes singular behavior at negative integer values of μ) is also used in the equations below. In Appendix D, we briefly review the basics of these functions, which are related to the confluent hypergeometric functions M(a, b, z) and U(a, b, z), and consider the operational procedure of Ref. [69] that defines the integral representations, with miscellaneous substitutions. For the calculation of \mathcal{I} in Eq. (67), we are going to use two such specific representations. 1. Spectral Decomposition of Hyperbolic Operators via the Fourier Method—Whole-Line Integral

The first computation of \mathcal{I} is based on the direct evaluation of the integral for the entire real line according to the novel identity (D6), which implies

$$\mathcal{I} = e^{\pi\kappa} \frac{\Gamma\left(\frac{1+\mu}{2} + i\kappa\right) \Gamma\left(\frac{1+\mu}{2} - i\kappa\right)}{\left(x'x''\right)^{1/2}} \mathcal{M}_{-i\kappa,\mu/2}(ix'') \mathcal{M}_{i\kappa,\mu/2}(-ix') , \qquad (69)$$

where the limit $c \to 0$ is taken, as in discussed Appendix D. This approach has the distinct advantage of being direct and straightforward [provided that Eq. (D6) is established]. Moreover, due to the symmetry of the propagator with respect to an exchange of r' and r''[thus, an exchange of x' and x'', according to Eq. (68)], the products can be written in either equivalent form

$$\mathcal{M}_{-i\kappa,\mu/2}(ix'') \,\mathcal{M}_{i\kappa,\mu/2}(-ix') = \mathcal{M}_{i\kappa,\mu/2}(-ix'') \,\mathcal{M}_{-i\kappa,\mu/2}(ix') \,, \tag{70}$$

which can also be explicitly verified via a double application of the Whittaker identity of Eq. (74) below.

2. Spectral Decomposition of Hyperbolic Operators via the Fourier Method—Half-Line Integrals

There is a second computation of \mathcal{I} , which uses an alternative (though related) representation from Ref. [69]: Eqs. (D7), and (D8), where the latter is another novel identity. This second approach, while more involved, provides a separate check and additional information, showing the different roles played by the positive and negative energy values, as well as their relationship to the Green's functions. Specifically, in Eq. (67), the integral \mathcal{I} is split into the two contributions from the half-axes (of positive and negative times): $\mathcal{I} = \mathcal{I}_+ + \mathcal{I}_-$, with \mathcal{I}_+ and \mathcal{I}_- defined by

$$\mathcal{I}_{\pm} = \int_{L_{\pm}} \frac{d\zeta}{\sinh \zeta} \exp\left(2i\kappa\zeta\right) \exp\left(i\beta \coth\zeta\right) I_{\mu}\left(\frac{\alpha}{i\sinh\zeta}\right) , \qquad (71)$$

where $L_+ = [0, \infty)$ and $L_- = (-\infty, 0]$. The integrals (71) are explicitly functions of the given parameters: (i) the indices κ and μ ; and (ii) the variables α and β , or alternatively x' and x''. The reversal of the sign of ζ in the integrand of L_- compared to L_+ implies the existence of straightforward symmetries with respect to the parameters of the integrand,

such that \mathcal{I}_{\pm} are not independent but related by

$$\mathcal{I}_{-}(\kappa,\mu;x',x'') = -\mathcal{I}_{+}(-\kappa,\mu;-x',-x'')$$
(72)

[or, by abuse of notation, also $\mathcal{I}_{-}(\kappa,\mu;\alpha,\beta) = -\mathcal{I}_{+}(-\kappa,\mu;-\alpha,-\beta)$]. In Eq. (72), as well as all the equations considered in this paper, the final expressions should be evaluated with the principal values of the multivalued functions involved. With the results of Eq. (D8), we get

$$\mathcal{I}_{\pm} = i \frac{\Gamma_{\mp}}{\sqrt{x'x''}} W_{\pm i\kappa,\mu/2}(\mp ix_{>}) \mathcal{M}_{\pm i\kappa,\mu/2}(\mp ix_{<}) , \qquad (73)$$

where $x_{>}$ and $x_{<}$ are the greater and lesser of the set $\{x', x''\}$ respectively, and the shorthand $\Gamma_{\pm} = \Gamma\left((1+\mu)/2 \pm i\kappa\right)$ is used. When adding the integrals $\mathcal{I} = \mathcal{I}_{+} + \mathcal{I}_{-}$, the following two relations are applied: (i) the semi-circuital analytic continuation [69]

$$\mathcal{M}_{\lambda,\mu/2}(e^{\pm \pi i}z) = e^{\pm(\mu+1)\pi i/2} \mathcal{M}_{-\lambda,\mu/2}(z) .$$
(74)

and (ii) the connection formula [69]

$$\mathcal{M}_{\lambda,\mu/2}(z) = e^{\pm\lambda\pi i} \left[\frac{1}{\Gamma_+} e^{\mp i(\mu+1)\pi i/2} W_{\lambda,\mu/2}(z) + \frac{1}{\Gamma_-} W_{-\lambda,\mu/2}(e^{\pm\pi i}z) \right] .$$
(75)

The Whittaker functions $\mathcal{M}_{\lambda,\mu/2}(z)$ and $W_{\lambda,\mu/2}(z)$ have to be examined to guarantee results valid within their principal branches, where the branch cut is conventionally taken as the negative real half-axis (thus, with the arguments $-\pi < \arg(z) \leq \pi$). For the present calculation, this involves using the lower and upper signs of the identities (74) and (75) (with $\lambda = i\kappa$) respectively, as applied to $\mathcal{I} = \mathcal{I}_+ + \mathcal{I}_+$ in that order, so that

$$\mathcal{I} = i \frac{\Gamma_{+}\Gamma_{-}}{\sqrt{x'x''}} \underbrace{\left[\frac{1}{\Gamma_{+}} W_{i\kappa,\mu/2}(-ix_{>}) \underbrace{\mathcal{M}_{i\kappa,\mu/2}(-ix_{<})}_{e^{-(\mu+1)\pi i/2} M_{-i\kappa,\mu/2}(ix_{<})} + \frac{1}{\Gamma_{-}} W_{-i\kappa,\mu/2}(ix_{>}) \mathcal{M}_{-i\kappa,\mu/2}(ix_{<})}_{e^{\pi\kappa} \mathcal{M}_{i\kappa,\mu/2}(-ix_{>}) \mathcal{M}_{-i\kappa,\mu/2}(ix_{<})} \right]}, \quad (76)$$

which reduces to the same value as before, Eq. (69), for x' > x''; however, due to the symmetry of Eq. (70), this is also true for x' < x'', showing the equality of both results without restriction.

3. Spectral Decomposition of Hyperbolic Operators—Green's Functions

Finally, the integrals \mathcal{I}_{\pm} in Eq. (73) give the retarded/advanced Green's functions

$$G_{l+\nu}^{(\pm)}(r'',r';E) = \mp \frac{i}{\hbar\omega} \frac{\Gamma((1+\mu)/2\mp i\kappa)}{\Gamma(1+\mu)} \frac{1}{\sqrt{r'r''}} W_{\pm i\kappa,\mu/2}(\mp i\check{r}_{>}^{2}) M_{\pm i\kappa,\mu/2}(\mp i\check{r}_{<}^{2}) , \quad (77)$$

where the jump in the Green's functions according to Eq. (C8) provides another, related proof of Eq. (67).

Incidentally, an analytic continuation back to real frequencies can be enforced via $\omega \to i\omega$, i.e., $i\kappa \to \kappa$. Notice the sign reversal compared to the analytic continuation that gave Eq. (66). Then, the Green's functions $G_{l+\nu}^{(\pm)}(r'', r'; E) =$ $\mp (\hbar\omega)^{-1} \Gamma_{\mp} W_{\pm\kappa,\mu/2}(\pm \check{r}_{>}^2) \mathcal{M}_{\pm\kappa,\mu/2}(\pm \check{r}_{<}^2)/\sqrt{r'r''}$ for the elliptic generators or ordinary radial harmonic oscillator are obtained, with \check{r}^2 given in Eq. (68) and $\Gamma_{\pm} = \Gamma((1 + \mu)/2 \pm \kappa)$. This is a familiar result [3] that also allows a rederivation of their spectrum, Eqs. (51)–(52). The critical difference in the spectral behaviors of hyperbolic generators (continuous operators) versus elliptic generators (discrete operators) arises from the asymptotic behavior of $M_{\pm i\kappa,\mu/2}(\mp i\check{r}^2)$ versus $M_{\pm\kappa,\mu/2}(\pm\check{r}^2)$ (both with $\kappa \in \mathbb{R}$), when $r \to \infty$, which forces the reduction of the latter to generalized Laguerre polynomials, with a discrete quantization of the spectrum.

C. Hyperbolic Operators: Conclusions

As a consequence of Eqs. (67), (69) [or Eq. (76)], and (70), the final result of this calculation is the wave function product

$$\mathcal{U}_{E,l}(r'')\mathcal{U}_{E,l}^*(r') = \frac{1}{2\pi\hbar\omega} \Gamma_{-}\Gamma_{+} e^{\pi\kappa} \frac{\mathcal{M}_{i\kappa,\mu/2}(-i\check{r}''^2)\mathcal{M}_{-i\kappa,\mu/2}(i\check{r}'^2)}{\sqrt{r''r'}}$$
(78)

$$= \frac{1}{2\pi\hbar\omega} \Gamma_{-}\Gamma_{+} e^{\pi\kappa} \frac{\mathcal{M}_{-i\kappa,\mu/2}(i\check{r}^{\prime\prime2}) \mathcal{M}_{i\kappa,\mu/2}(-i\check{r}^{\prime\prime2})}{\sqrt{r^{\prime\prime}r^{\prime}}} .$$
(79)

Again, the equivalence of Eqs. (78) and (79) is due to the symmetry of the propagator with respect to the end points, and enforced via the analytic continuation of Eq. (74).

Two important conclusions follow from this result.

(i) The spectrum is indeed a continuum, from minus to plus infinity, as the solution above applies to all such values of the effective energy $\tilde{E}_G \equiv \sigma E_G$, or [from Eqs. (36) and (68)] the parameter

$$\kappa = \sigma \frac{E_G}{2\hbar\omega} = \sigma \frac{\mathfrak{g}}{\sqrt{\Delta}} = \mathfrak{s}' \in (-\infty, \infty) , \qquad (80)$$

which is a dimensionless value covering the whole real axis and equal to the first Whittaker index κ , and such that $\mathfrak{s}' = \kappa$ for the eigenvalues $\hbar \mathfrak{s}'$ of the operator S' = -S.

(ii) The wave functions can be read off from either one of Eqs. (78) or (79), by comparison with Eq. (C2), and are given by

$$\mathcal{U}_{E,l}(r) \equiv \mathcal{U}_{\kappa,\mu}(r) = \frac{e^{\pi\kappa/2}}{\sqrt{2\pi\hbar\omega}} \frac{\Gamma((1+\mu)/2 + i\kappa)}{\Gamma(1+\mu)} \frac{M_{i\kappa,\mu/2}(-i\check{r}^2)}{\sqrt{r}} e^{i\chi}$$
(81)

$$= \frac{e^{\pi\kappa/2}}{\sqrt{2\pi\hbar\omega}} \frac{\Gamma((1+\mu)/2 - i\kappa)}{\Gamma(1+\mu)} \frac{M_{-i\kappa,\mu/2}(i\check{r}^2)}{\sqrt{r}} e^{i\chi'}, \qquad (82)$$

where we have reverted back to the ordinary Whittaker functions $M_{\pm i\kappa,\mu/2}$, and χ and χ' are arbitrary phase factors. The simplest choices are arguably $\chi = 0$ or $\chi' = 0$, but these values are undetermined by the nature of the products (78)–(79); in particular, the choice of gamma function factors is also arbitrary, as $\Gamma_{\pm} = \Gamma_{\mp}^*$. It should be noted that, in the derivation above, $\mathcal{M}_{-i\kappa,\mu/2}(i\tilde{r}^2) \propto \mathcal{M}_{i\kappa,\mu/2}(-i\tilde{r}^2)$, according to Eq. (74); thus, up to a phase factor, the Whittaker functions in Eq. (81)–(82) can take either equivalent form $M_{\mp i\kappa,\mu/2}(\pm i\tilde{r}^2)$.

Moreover, the basic results of Eqs. (78)–(79) and (81)–(82) are also verified by the analysis of Appendix E, based on the associated differential equation.

The following remarks on the nature of the mathematical results for hyperbolic operators are in order. First, these results have been established by a rigorous approach that relies on established integral representations of the product of Whittaker functions. Second, the proof relies on the novel Fourier method for continuous-spectrum operators presented in Appendix C. Third, they provide a remarkable example of a continuous spectrum unbounded from both below and above; and finally, perhaps due to their unusual nature, they have not been addressed in the existing literature: ideally, our paper should contribute to fill that gap.

This concludes our detailed analysis of the most relevant spectral properties of the symmetry operators of CQM, with the added bonus of having established appropriate analytic continuation techniques to compare the different types.

VII. CONCLUSIONS

In this work, we have derived a comprehensive path-integral treatment of the symmetry generators of conformal quantum mechanics (CQM). This analysis is relevant in the context of CQM as a one-dimensional conformal field theory [39–46]. Moreover, it is noteworthy that the symmetry operators in the weak-coupling regime of CQM are of current interest in understanding the nature of spacetime causal structure in the context of causal diamonds and thermal properties of the vacuum [47–49, 51–57].

The main results of our analysis include a complete characterization via path integrals of all the operators from the three families (elliptic, parabolic, and hyperbolic), with distinctly different spectral and time-evolution properties. In addition to establishing appropriate analytic continuations of the path integral for the analog system of a radial harmonic oscillator, we have derived novel expressions and a simple general technique to deal with the spectral characterization of continuous operators. These results can be used to provide further insight into the physical and mathematical properties of CQM, including, inter alia, the physical meaning of this form of SO(2,1) conformal symmetry for near-horizon physics [10, 70] and applications in quantum cosmology [71–73]. Moreover, the general methodology and spectral properties we established for hyperbolic operators are of potential interest in related applications of the inverted harmonic oscillator [67, 68], which we are currently investigating, with impact on problems from the quantum Hall effect to black holes, including issues of thermality and complexity [74–77].

Another line of work for which the CQM generators, as considered in this paper, are relevant is the recent development of Schwarzian mechanics [78–83], which is related to the dAFF model. Most importantly, Schwarzian mechanics is relevant to the low-energy limit of the Sachdev-Ye-Kitaev model [84], as was shown in Refs. [78, 79].

Moreover, the use of the symmetry generators G as alternative Hamiltonians with transformed times τ is still an open question, though some special cases have been suggested in the literature, for example, for the magnetic monopole [6] and the magnetic vortex [7]. In Ref. [85], the roles played by the generators H, R, and S' is considered within a CFT-inspired sine-square deformation, with an interpretation that can be further reexamined within our generalized framework. In addition, the Niederer-Takagi time transformation [86–89] involved in the elliptic generators has been used in the physics of cold atoms [90, 91]. However, the interpretation of the time variable is a subtle notion that finds a more natural context in general-relativistic applications, as has been suggested for matrix models [92]. This interpretation has been partly implemented for the dynamics of harmonic-oscillator-type Unruh-DeWitt detectors in curved spacetimes [93]. More generally, this is an issue of relevance in the near-horizon version of CQM for black hole thermodynamics [10] and causal diamonds [47–49, 51–57], which deserves further investigation.

Finally, the current presentation has been limited to the weak-coupling regime of CQM, as described in Sec. I. An extension of this framework to the strong coupling regime, with additional implications for renormalization and quantum anomalies, is in progress, and will be reported elsewhere.

Acknowledgments

This material is based upon work supported by the Air Force Office of Scientific Research under Grant No. FA9550-21-1-0017 (C.R.O., A.C., and P.L.D.). C.R.O. was partially supported by the Army Research Office (ARO), grant W911NF-23-1-0202. H.E.C. acknowledges support by the University of San Francisco Faculty Development Fund.

This article may be downloaded for personal use only. Any other use requires prior permission of the author and AIP Publishing. This article appeared in H. E. Camblong, A. Chakraborty, P. Lopez Duque, C. R. Ordóñez; Spectral properties of the symmetry generators of conformal quantum mechanics: A path-integral approach. J. Math. Phys. 1 September 2023; 64 (9): 092302, and may be found at https://doi.org/10.1063/5.0150349.

AUTHOR DECLARATIONS

Conflict of Interest

The authors have no conflicts to disclose.

Author Contributions

All authors contributed equally to this work.

DATA AVAILABILITY

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

Appendix A: A Dimensional Form of the Theory of Symmetry Generators and Their Effective Hamiltonian Representation

The framework of Subsec. II B is based on the generators G leading to the effective Hamiltonian (25). It has the definite advantage of being completely general, but the Hamiltonian involves canonical variables and time with reduced dimensions, which are different from those of the original CQM Hamiltonian (5)[(for example, τ is dimensionless and $[\mathbf{r}] = [\mathbf{Q}]/[f]^{1/2}$, with [f] = time). While this poses no serious technical challenges, it may be desirable to develop an alternative method where the relevant variables have their "usual" dimensions, meaning those of the original Hamiltonian H. Specifically, dimensions can be restored to those of H via a characteristic time scale; here, this is naturally provided by the factor u of the generalized generator, Eq. (15), if $u \neq 0$.

In principle, under the assumption $u \neq 0$, the rescaling $G = u\hat{H}_G(\hat{\mathbf{r}}, \hat{\mathbf{p}})$, rather than $G = \sigma \tilde{H}_G(\mathbf{r}, \mathbf{p})$, allows the theory to be redefined consistently in terms of the effective time $\hat{\tau} = u\tau$ and canonical variables $\hat{\mathbf{r}}, \hat{\mathbf{p}}$, such that $\hat{r}^2 = ur^2$ and $\hat{p}^2 = u^{-1}\mathbf{p}^2$. Then, the reduced effective Hamiltonian becomes

$$\frac{1}{u}G \equiv \hat{H}_G(\hat{\mathbf{r}}, \hat{\mathbf{p}}) = \frac{\hat{p}^2}{2M} + \frac{\hbar^2}{2M}\frac{g}{\hat{r}^2} + \frac{M}{2}\left(-\frac{\Delta}{4u^2}\right)\hat{r}^2 \tag{A1}$$

(with the additional assignments $\lambda = \hbar^2 g/M$ and $\hat{p}^2 = \hat{\mathbf{p}}^2$ to be used below). The evolution operator (22) admits the alternative form $U_{\hat{H}}(T) = e^{-i\hat{H}(\hat{\tau}-\hat{\tau}_0)/\hbar}$, and the CQM frequency becomes

$$\hat{\omega}^2 = -\frac{\Delta}{4u^2} \,. \tag{A2}$$

With these redefinitions, the time $\hat{\tau}$, the Hamiltonian $\hat{H}_G(\hat{\mathbf{r}}, \hat{\mathbf{p}})$, and the frequency in Eq. (A2) have the usual dimensions.

In addition, it may prove useful to define the real and positive time-scale parameter

$$a = 2\frac{|u|}{\sqrt{|\Delta|}} , \qquad (A3)$$

which generalizes the original parameter a of Ref. [5] in Eqs. (11) and (12); then, $\hat{\omega}^2 = -\text{sgn}(\Delta)/a^2$, which leads to $\hat{\omega} = 1/a$ for $\Delta < 0$ and $\hat{\omega} = -i/a$ for $\Delta > 0$ (the choice of sign for the latter is discussed in Sec. VI), with $\hat{\omega} = 0$ and $a = \infty$ for $\Delta = 0$. As in Subsec. II B, denoting the corresponding parameter dependence in Eq. (A1) with $\hat{H}[M, g, \hat{\omega}]$ (where the

subscript G is removed for simplicity), we can make the assignments

$$R = \frac{a}{2} \hat{H} \left[M, g, \hat{\omega} = 1/a \right], \qquad (A4)$$

$$H = \hat{H} \left[M, g, \hat{\omega} = 0 \right], \tag{A5}$$

$$S' \equiv \frac{a}{2} \hat{H} \big[M, g, \hat{\omega} = -i/a \big] , \qquad (A6)$$

as u = a/2 for both R and S'. The corresponding description of the three classes of operators follows from $G = u\hat{H}[M, g, \hat{\omega}]$, which also implies that $G = \operatorname{sgn}(u)\sqrt{|\Delta|}(a/2) \hat{H}[M, g, \hat{\omega}]$ for elliptic and hyperbolic operators; consequently,

$$G = \operatorname{sgn}(u)\sqrt{|\Delta|} R$$
 for elliptic generators, (A7)

$$G = u H$$
 for parabolic generators , (A8)

and
$$G = \operatorname{sgn}(u)\sqrt{\Delta} S'$$
 for hyperbolic generators . (A9)

This restricted framework agrees with the more general characterization of Eqs. (34)–(36), where σ is the more general sign, which can be evaluated from $\operatorname{sgn}(u)$ when $\operatorname{sgn}(u) \neq 0$. Despite being somewhat restricted, this approach has some appealing features, i.e., its dimensional-analysis structure, and includes the all-important operators R, H, and S'.

A final remark is in order regarding the restrictions of this particular approach. We have assumed that $u \neq 0$, which basically modifies the original inverse-square-potential Hamiltonian H with the extension (A1). This excludes the family of Hamiltonians H_G with u = 0, for which the discriminant is $\Delta = v^2$, and which encompasses two subclasses: (i) hyperbolic operators with $v \neq 0$, including D by itself, as well as linear combinations of Dand K; (ii) parabolic operators with v = 0, which reduce to simply K (up to an arbitrary multiplication constant w). However, we have seen in Subsec. II B that these operators can be handled with the more general framework, where K and D are described as equivalent to H and S' respectively.

Appendix B: Path-Integral Framework—Summary of Basic Results

In this appendix, we summarize the main results on path integration needed to understand and compute the spectral properties of the operators discussed in the main text. These include the setup of the time-sliced path integral, the transformation to hyperspherical coordinates, and the evaluation of the propagator for the radial harmonic oscillator.

1. Path-Integral Framework Setup

The basic setup starts, as in Sec. III, with the path-integral expression (38) for the propagator, which can be evaluated as the limit of a properly time-sliced integral in Cartesian coordinates,

$$K_{(d)}(\mathbf{r}'',\mathbf{r}';t^{''},t^{'}) = \lim_{N \to \infty} \left(\frac{M}{2\pi i\epsilon\hbar}\right)^{dN/2} \left[\prod_{k=1}^{N-1} \int_{\mathbb{R}^d} d^d \mathbf{r}_k\right] e^{iS^{(N)}/\hbar} .$$
(B1)

As in Subsec. III A, we use t for the dynamical time and \hat{H} for a generic Hamiltonian. The Cartesian form of Eq. (B1) involves a time lattice $t_j = t' + j\epsilon$, for the time interval T = t'' - t' corresponding to the end points $\mathbf{r}_0 \equiv \mathbf{r}'$ and $\mathbf{r}_N \equiv \mathbf{r}''$; in this lattice, $\epsilon = T/N$, with $j = 0, \dots, N$, with $t_0 \equiv t'$ and $t_N \equiv t''$, such that $\mathbf{r}_j = \mathbf{r}(t_j)$. The discrete action in Eq. (B1) is $S^{(N)} = \sum_{j=0}^{N-1} S_j^{(N)}$, with $S_j^{(N)} = M(\mathbf{r}_{j+1} - \mathbf{r}_j)^2/2\epsilon - \epsilon V(\mathbf{r}_j)$, where $V(\mathbf{r})$ is the potential—if assumed to be time-independent, as is the case for the computation of all the CQM operators, the result of this path integral is only a function of T.

The propagator $K_{(d)}(\mathbf{r}'', \mathbf{r}'; T)$ can be formally rewritten in non-Cartesian coordinate systems. It is well-known that caution must be exercised when evaluating Eq. (38) in non-Cartesian coordinates, as this typically leads to extra terms of order \hbar^2 in the action. These arise when nonlinear transformations are performed, both in quantum mechanics [60, 94–97]. and quantum field theory [98]. In general, coordinate transformations of Eq. (B1) can be applied *before* taking the continuum limit, but prescriptions for the choice of position values of the potential are dictated by operator ordering.

2. Path-Integral Framework—Hyperspherical Coordinates and Besselian Path Integrals

For the important case of hyperspherical polar coordinates [33, 34, 59], the *d*-dimensional hyperspherical harmonics $Y_{lm}(\Omega)$ and Gegenbauer polynomials $C_l^{(\nu)}(x)$ provide the necessary framework for separation of variables; see Sec. III and Eqs. (39)–(40) for the partial wave expansions. Then, an explicit expression for the propagator can be derived in the timesliced path integral by transforming Eq. (38) into hyperspherical coordinates before taking the $N \to \infty$ limit. This is especially useful for central potentials: $V(\mathbf{r}) = V(r)$. The critical step in this derivation is rewriting the elements of the discretized action $S_j^{(N)}$ = $M(\mathbf{r}_{j+1} - \mathbf{r}_j)^2/2\epsilon - \epsilon V(r_j)$ by separating the angles $\cos \psi_{j+1,j} = \hat{\mathbf{r}}_{j+1} \cdot \hat{\mathbf{r}}_j$ (with $\hat{\mathbf{r}} = \mathbf{r}/r$) in a radial-angular resolution $\exp\left(iS_j^{(N)}/\hbar\right) = \exp\left[i\frac{M}{2\hbar\epsilon}\left(r_{j+1}^2 + r_j^2\right) - i\frac{\epsilon V_j}{\hbar}\right]\exp\left(z_j\cos\psi_{j+1,j}\right)$, where $z_j = \frac{Mr_jr_{j+1}}{i\epsilon\hbar}$. The product of the last factor $\exp\left(z_j\cos\psi_{j+1,j}\right)$ for all the intervals $(j = 0, \ldots, N-1)$ can be evaluated in the angular integrals of the time-sliced path integral, Eq. (B1), which involves multiple applications of $d^d\mathbf{r}_k = dr_k r_k^{d-1} d\Omega_k$, with *d*-dimensional solid-angle integrations. This is done using the degenerate form of Gegenbauer's addition theorem on partial waves (generalization of the 3D Rayleigh expansion of a plane wave; Sec. 11.5 of Ref. [65]):

$$e^{iz\cos\psi} = (iz/2)^{-\nu}\Gamma(\nu) \sum_{l=0}^{\infty} (l+\nu)I_{l+\nu}(iz)C_l^{(\nu)}(\cos\psi) , \qquad (B2)$$

where $\nu = d/2 - 1$ and $I_p(x)$ is the modified Bessel function of the first kind and order p. With this expansion and performing the angular integrations, along with Eqs. (39)–(40) and the orthonormality of hyperspherical harmonics, the path integral for the radial propagator in Eq. (40) becomes

$$K_{l+\nu}(r'',r';T) = \lim_{N \to \infty} \left(\frac{M}{2\pi i\epsilon\hbar} \right)^{N/2} \prod_{k=1}^{N-1} \left[\int_0^\infty dr_k \right] w_{l+\nu}^{(N)}[r^2] \\ \times \exp\left\{ \frac{i}{\hbar} R^{(N)} \left[r_1, \dots, r_{N-1} \right] \left(r'',r';T \right) \right\} ,$$
(B3)

where the radial action is

$$R^{(N)}[r_1, \dots r_{N-1}](r'', r'; T) = \sum_{j=0}^{N-1} \left[\frac{M(r_{j+1} - r_j)^2}{2\epsilon} - \epsilon V(r_j) \right] .$$
(B4)

In Eq. (B3) a radial functional weight

$$w_{l+\nu}^{(N)}[r^2] = \prod_{j=0}^{N-1} \left[\sqrt{2\pi z_j} e^{-z_j} I_{l+\nu}(z_j) \right]$$
(B5)

has been properly defined with the radial variables appearing through the characteristic dimensionless ratio $z_j = \frac{Mr_jr_{j+1}}{i\epsilon\hbar}$. Equation (B3) [with the restriction to the half-line $r(t) \ge 0$] admits the formal Besselian path-integral representation (42), i.e.,

$$K_{l+\nu}(r'', r'; T) = \int \mathcal{D}r(t) \ w_{l+\nu}[r^2] \exp\left\{\frac{i}{\hbar} \int_{t'}^{t''} dt \left[\frac{M}{2} \dot{r}^2 - V(r)\right]\right\} , \tag{B6}$$

with the angular part of the problem contributing through the nontrivial radial functional weight $w_{l+\nu}[r^2]$. As a result of this hyperspherical resolution, the property known as interdimensional dependence [62, 63] is exhibited by Eqs. (B3)–(B6): d and l appear in the combination $l + \nu$.

The associated radial action R[r(t)] in Eq. (B3) only includes the radial kinetic energy and the interaction potential, and excludes the centrifugal potential, whose role is played instead by the functional weight. The connection with the usual formulation of the classical radial action can be shown by a formal argument, using the asymptotic form of the Bessel function $\sqrt{2\pi z} e^{-z} I_{\mu}(z) \sim e^{-(\mu^2 - 1/4)/2z + O(1/z^2)}$ (for $|z| \to \infty$) in Eq. (B5), where $z \sim 1/\epsilon$. Thus, this leads to the alternative expression

$$K_{l+\nu}(r'',r';T) = \int \mathcal{D}r(t) \exp\left\{\frac{i}{\hbar} \int_{t'}^{t''} dt \left[\frac{M}{2} \dot{r}^2 - \frac{\hbar^2}{2M} \frac{(l+\nu)^2}{r^2} - V(r)\right]\right\} , \qquad (B7)$$

where the action does include the centrifugal term—this formal procedure has been called asymptotic recombination [61]. This equivalence of Eqs. (B6) and (B7) can be established rigorously by the important theorem relating angular momentum and inverse square potential terms—see Eqs. (43) and (44) and associated statement in Subsec. III A. The latter can be inserted in the path integral [as can be seen in Eq. (B7)] by absorbing the inverse-square coupling g as part of an effective angular momentum with $\mu = \sqrt{(l+\nu)^2 + g}$. This was originally shown as above in Ref. [64] for d = 3, with a rigorous proof in Refs. [61, 63].

3. Inverse Square Potential and Radial Harmonic Oscillator

The generic radial path integral (B3) can be used to evaluate the propagator for the inverse square potential and radial harmonic oscillator, by rewriting it in a more convenient form,

$$K_{l+\nu}(r'',r';T) = \lim_{N \to \infty} \left(\frac{\alpha}{i}\right)^N \prod_{k=1}^{N-1} \left[\int_0^\infty dr_k r_k\right] \prod_{j=0}^{N-1} \left[I_{l+\nu}(z_j)e^{iA_j/\hbar}\right]$$
(B8)

where $\alpha = M/(\epsilon\hbar)$ and $A_j = \frac{M}{2\epsilon} \left(r_{j+1}^2 + r_j^2\right) - \epsilon V(r_j)$. The limit $K_{l+\nu}(r'', r'; T) = \lim_{N\to\infty} K_{l+\nu}^{(N)}(r'', r'; T)$ in Eq. (B8) can be taken by evaluating the N-th order term $K_{l+\nu}^{(N)}(r'', r'; T)$ recursively [1–3, 64]. Transforming the path integral into its Euclidean-time form (with the replacements $\epsilon = T/N \to -i\epsilon$, $\alpha \to i\alpha$, and $z_j \to \zeta_j = iz_j$), and defining the

parameter $\beta = \alpha (1 + \omega^2 \epsilon^2 / 2)$, if follows that

$$K_{l+\nu}^{(N)}(r'',r';T) = \exp\left[-\frac{\alpha}{2}\left(r'^{2}+r''^{2}\right)\right] \underbrace{\alpha^{N} \prod_{k=1}^{N-1} \left[\int_{0}^{\infty} dr_{k}r_{k} e^{-\beta r_{k}^{2}}\right] \prod_{j=0}^{N-1} I_{\mu}(\alpha r_{j+1}r_{j})}_{P_{N}(r'',r';\alpha,\beta)} . \tag{B9}$$

In Eq. (B9), the functional form of the propagator is governed by

$$P_N(r_N, r_0; \alpha, \beta) = \frac{\alpha}{\gamma_N} I_\mu \left(\frac{\alpha}{\gamma_N} r_N r_0\right) \exp\left[\frac{\alpha}{2\lambda_N} \left(r_0^2 + r_N^2\right)\right] , \qquad (B10)$$

where the N-dependent parameters γ_N and λ_N are functions of $\eta = \beta/\alpha = 1 + \omega^2 \epsilon^2/2$; (i.e., they depend on the product $\omega\epsilon$). Now, Eq. (B10) can be established by repeated application of Weber's second exponential integral for Bessel functions (Sec. 13.31 of Ref. [65] and 10.22.67 of Ref. [66]),

$$\int_{0}^{\infty} \exp\left(-c^{2}x^{2}\right) J_{\mu}(ax) J_{\mu}(bx) x dx = \frac{1}{2c^{2}} \exp\left(-\frac{a^{2}+b^{2}}{4c^{2}}\right) I_{\mu}\left(\frac{ab}{2c^{2}}\right)$$
(B11)

 $[\operatorname{Re}(\mu) > -1 \text{ and } | \arg(c)| < \pi/4].$ Basically, the repeated integrals in Eq. (B9) give an integral relation between P_N and P_{N-1} , from which Eq. (B11) provides three consistency conditions: the basic recursion relation, $\lambda_N + 1/\lambda_{N-1} = 2\eta$; the relation between the parameters, $\lambda_N = \gamma_N/\gamma_{N-1}$; and the additional recursion relation $1 + \gamma_N\gamma_{N-2} = \gamma_{N-1}^2$. The latter can be shown to be equivalent to the basic recursion relation, and to the corresponding simpler relation for $\gamma_N: \gamma_N + \gamma_{N-2} = 2\eta\gamma_{N-1}$. The initial values for the two-term and three-term relations satisfied by λ_N and γ_N respectively are $\lambda_1 = \infty$ (or $\lambda_2 = 2\eta$) and $\gamma_1 = 1$ with $\gamma_2 = 2\eta$. The consistency of the recursion relations and generic patterns can be proved by mathematical induction, and a closed solution can be derived in the continuum limit $N \to \infty$. An efficient approach [99] to derive the expressions in this limit is via a function $\Phi(T) = \lim_{N\to\infty} \Phi_N$, where $\Phi_N = \epsilon \gamma_N$, with finite differences $\dot{\Phi}_N = (\Phi_{N+1} - \Phi_N)/\epsilon = (\gamma_{N+1} - \gamma_N)$, and $\ddot{\Phi}_N = (\Phi_{N+1} + \Phi_{N-1} - 2\Phi_N)/\epsilon^2 = (\gamma_{N+1} + \gamma_{N-1} - 2\gamma_N)/\epsilon = \omega^2 \Phi_N$. This shows, from Eqs. (B9) and (B10) that

$$K_{l+\nu}(r'',r';T) = \frac{M}{\hbar \Phi(T)} \sqrt{r'r''} \exp\left[-\frac{M}{2\hbar} \frac{\dot{\Phi}(T)}{\Phi(T)} \left(r'^2 + r''^2\right)\right] I_{\mu}\left(\frac{M}{\hbar \Phi(T)} r'r''\right) , \quad (B12)$$

where, from the finite-difference form of the recursion relations and comparison with the free particle, the function

$$\Phi = \frac{1}{\omega} \sinh(\omega T) \tag{B13}$$

(Euclidean-time version) can be found as the solution to the differential equation $\dot{\Phi}(T) = \omega^2 \Phi(T)$ with the initial conditions $\Phi(0) = 0$ and $\dot{\Phi}(0) = 1$. In conclusion, these equations, after conversion to real time, show that the path integral is given by Eq. (45). The propagator (B12) can also be generalized to account for possible time dependence of the oscillator parameters [99].

For the case of the inverse square potential alone, an exact derivation from a perturbative series is possible [100], including a renormalized version for the strong-coupling regime [101] (similar to the case of the renormalized path integral for the two-dimensional delta interaction [102]).

Appendix C: Fourier Method for Continuous-Spectrum Operators

This appendix introduces a novel technique to derive the eigenfunctions of operators with a continuum spectrum. The general properties of this technique are discussed, and then further expanded with its relationship to Green's functions. This topic is critical for the study of the spectral properties of the hyperbolic generators, including S; and it also provides one of the various approaches to understand parabolic generators.

1. Fourier Method

For any operator \tilde{H} with a purely continuous spectrum, the spectral decomposition takes the form

$$K_{l+\nu}(r'',r';T) = \int_{\mathcal{S}} dE e^{-iET/\hbar} \mathcal{U}_{E,l}(r'') \mathcal{U}_{E,l}^*(r') , \qquad (C1)$$

where the "energy" values $E \equiv \tilde{E}$ of \tilde{H} extend over the set S. We assume that \tilde{H} has a purely continuous spectrum, and can be regarded as a sort of Hamiltonian operator associated with an effective time evolution, i.e., defining $K_{(d)}(\mathbf{r}'', \mathbf{r}'; T) = \langle \mathbf{r}'' | e^{-i\tilde{H}T} | \mathbf{r}' \rangle$ and extracting the radial counterpart according to the rules of Sec. III.

Equation (C1) has the form of a Fourier transform restricted to the set S. Its inverse Fourier transform can be obtained by performing the integral with respect to the variable $T \in (-\infty, \infty)$, with the kernel exp (iET/\hbar) . This general inverse Fourier integral converts the propagator $K_{l+\nu}(r'', r'; T)$ into the wave-function product $\mathcal{U}_{E,l}(r'')\mathcal{U}_{E,l}^*(r')$, i.e.,

$$F(E;r'',r') \equiv \mathcal{U}_{E,l}(r'') \,\mathcal{U}_{E,l}^*(r') = \frac{1}{2\pi\hbar} \,\int_{-\infty}^{\infty} dT \,\exp\left(\frac{iET}{\hbar}\right) \,K_{l+\nu}(r'',r';T) \,, \qquad (C2)$$

where the values of the physical energy on the real axis are restricted to the original set $E \in \mathcal{S}$. Indeed, this general theorem simply follows by performing the inverse Fourier integral on the right-hand side of Eq. (C1) written in terms of a variable E', with the familiar auxiliary identity $\int_{-\infty}^{\infty} dT e^{i(E-E')T/\hbar} = 2\pi\hbar\delta(E-E')$.

In conclusion, Eqs. (C1) and (C2) provide the broader context and justification for the results of Subsec. VD and Sec. VI. This "Fourier method" is remarkably simple and elegant, but it has not been explicitly used in the literature. This is partly due to the fact that operators with a purely continuous spectrum are not common (other than the trivial case of a free particle), and partly because it is customary to use Green's functions techniques, which are related but not identical to the result of Eq. (C2).

2. Green's Functions: Definitions and Relation to the Fourier Method

In general, the retarded/advanced Green's functions or resolvents associated with \hat{H} are defined from

$$G_{l+\nu}^{(\pm)}(r'',r';E) = \pm \frac{1}{i\hbar} \int_{-\infty}^{\infty} dT \,\theta(\pm T) \,\exp\left(\frac{iET}{\hbar}\right) \,K_{l+\nu}(r'',r';T) \,, \tag{C3}$$

where θ stands for the Heaviside function, and with the replacement $E \to E \pm i0^+$ that guarantees convergence. Unlike Eq. (C2), the Green's function technique does not give a direct result for the wave function product, though this product can be extracted via an additional step as the residue of the energy poles.

The definitions of Eq. (C3) correspond to the Fourier transform of the (re-tarded/advanced) Green operators

$$G^{(\pm)}(T) = \theta(\pm T) e^{-i\hat{H}T/\hbar} .$$
(C4)

This should be contrasted with the propagator counterparts that involve the time evolution operator $U = e^{-i\hat{H}T/\hbar}$ of Eq. (37) without the Heaviside cutoff. Equation (C4) yields the Fourier-transformed or energy Green operators

$$G^{(\pm)}(E) = \left(E - \hat{H} \pm i0^+\right)^{-1} , \qquad (C5)$$

where the $i0^+$ prescription provides convergence for each case. The corresponding coordinate-space representations give the associated time and energy Green's functions $G_{(d)}^{(\pm)}(r'',r';T)$ and $G_{(d)}^{(\pm)}(r'',r';E)$; in particular,

$$G_{(d)}^{(\pm)}(\mathbf{r}'',\mathbf{r}';E) = \left\langle \mathbf{r}'' \left| \left(E - \hat{H} \pm i0^+ \right)^{-1} \right| \mathbf{r}' \right\rangle .$$
(C6)

Equations (C4)–(C5) can be applied either to the full-fledged multidimensional quantities, or their reduced radial counterparts, which follow from the usual hyperspherical expansion; specifically, the radial energy Green's functions $G_{l+\nu}(r'', r'; E)$ in Eq. (C3) are defined from

$$G_{(d)}^{(\pm)}(\mathbf{r}'',\mathbf{r}';E) = (r''r')^{-(d-1)/2} \sum_{l=0}^{\infty} \sum_{m=1}^{d_l} Y_{lm}(\mathbf{\Omega}'')Y_{lm}^*(\mathbf{\Omega}')G_{l+\nu}^{(\pm)}(r'',r';E) .$$
(C7)

Moreover, with the usual distributional expansion $(A \pm i0^+)^{-1} = \mathcal{P}(A^{-1}) \mp i\pi \delta(A)$ (where \mathcal{P} is the Cauchy principal value), the ratio $-\left(G^{(+)}(E) - G^{(-)}(E)\right)/(2\pi i) = \delta(E - \hat{H})$ is a state density operator. Thus, for a continuous-energy spectral expansion,

$$F(r'', r'; E) \equiv \mathcal{U}_{E,l}(r'') \, \mathcal{U}_{E,l}^*(r') = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} dT \, \exp\left(\frac{iET}{\hbar}\right) \, K_{l+\nu}(r'', r'; T) \\ = -\frac{1}{2\pi i} \underbrace{\left[G_{l+\nu}^{(+)}(r'', r'; E) - G_{l+\nu}^{(-)}(r'', r'; E)\right]}_{\text{disc}[G_{l+\nu}(r'', r'; E)]}, \tag{C8}$$

where $\operatorname{disc}[G_{l+\nu}(E)] = G_{l+\nu}^{(+)}(E) - G_{l+\nu}^{(-)}(E)$ measures the discontinuity of the Green's functions across the branch cut in the complex energy plane. In short, Eq. (C8) summarizes the connection between the straightforward Fourier transform of the propagator and the Green's functions; for the former, the time domain of the Fourier integrals involves the whole time axis, while for the Green's functions, the time domain is restricted to the positive and negative half-axes.

3. Green's Functions—Example: Parabolic Generators

As an example of the relations above, we can revisit the conformal operator H and its parabolic class. The whole time-axis Fourier integral gives directly the wave function product (64), which we derived from Eq. (D1). If we instead use the related integrals of Eq. (D2), the Green's functions become

$$G_{l+\nu}^{(\pm)}(r'',r';E) = \mp \pi i \left(M/\hbar^2 \right) \sqrt{r'r''} J_{\mu}(kr_{<}) H_{\mu}^{(1,2)}(kr_{>}) , \qquad (C9)$$

where $r_{<}(r_{>})$ is the lesser (greater) of r' and r''; then,

$$G_{l+\nu}^{(+)}(r'',r';E) - G_{l+\nu}^{(-)}(r'',r';E) = -\pi i \frac{M}{\hbar^2} \sqrt{r'r''} J_{\mu}(kr_{<}) \left[H_{\mu}^{(1)}(kr_{>}) + H_{\mu}^{(2)}(kr_{>}) \right] ,$$

which, with $H^{(1)}_{\mu}(z) + H^{(2)}_{\mu}(z) = 2J_{\mu}(z)$, gives $-2\pi i$ times the wave function product displayed in Eq. (64), in agreement with the relation (C8).

In the next section, we will apply the technique of Eq. (C2) to derive the spectral decomposition associated with the more involved conformal operator S, and will display the corresponding network of relations developed here.

Appendix D: Relevant Integral Representations for the Product of Bessel and Whittaker Functions as Fourier Transforms of Propagators

In this appendix, we summarize the key results on integral representations of products of the special functions relevant for the propagators discussed in this paper. We will first identify the identities relevant for the operator H and related parabolic generators, i.e., generally for the pure inverse square potential, using products of Bessel functions; and we will then explore a general method and appropriate identities for the hyperbolic operators, corresponding to an inverted radial harmonic oscillator, using products of Whittaker functions. The identification of the relevant identities is a useful addition to the literature of path integrals; and, in the case of the latter class, we establish, inter alia, a new integral identity.

1. Integral Representations for the Product of Bessel Functions

Let us consider the following integral representation of the product of Bessel functions (Sec. 13.7 of Ref. [65]):

$$J_{\mu}(z')J_{\mu}(z'') = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \exp\left(\frac{s}{2} - \frac{z'^2 + z''^2}{2s}\right) I_{\mu}\left(\frac{z'z''}{s}\right) \frac{ds}{s} , \qquad (D1)$$

where c is a real positive constant and $\operatorname{Re}(\mu) > -1$. As shown in Ref. [65]), this representation (D1) can be established by combining an appropriate form of Bessel function Gegenbauer addition theorems with Schläfli's integral $J_{\nu}(z) = \frac{(\frac{1}{2}z)^{\nu}}{2\pi i} \int_{-\infty}^{(0+)} \exp\left(t - \frac{z^2}{4t}\right) \frac{dt}{t^{\nu+1}}$. As shown in Subsec. VA, we consider the limit $c \to 0^+$, that reduces the integral (D1), with s = c + 2it, to the form (63), where $t \to t - i0^+$. Its relevance for our study stems from the fact that the substitution $t \equiv ET/\hbar$ makes it directly applicable to Eq. (62) for the operator H (inverse square potential) as a Fourier transform, leading to Eq. (64).

In a similar way, the counterparts of Eq. (D1) for the half-axis intervals can be found using Hankel functions (for which the corresponding Schläfli's integrals only involve halfaxes), yielding

$$H_{\mu}^{(1,2)}(z_{>}) J_{\mu}(z_{<}) = \pm \frac{1}{\pi i} \int_{0}^{c \pm i\infty} \exp\left[\frac{s}{2} - \frac{(z'^{2} + z''^{2})}{2s}\right] I_{\mu}\left(\frac{z'z''}{s}\right) \frac{ds}{s}$$
(D2)

where c is again a real positive constant and $\operatorname{Re}(\mu) > -1$; and $z_>$ and $z_<$ are the greater and lesser of the set $\{z', z''\}$ respectively. (A related integral is listed in Ref. [103], 6.653-1.)

2. Integral Representations for the Product of Whittaker Functions

We will begin by summarizing and adapting the technique defined in Ref. [69] (Section 6) to set up integral representations of products of Whittaker functions. The method involves writing a single Whittaker function as a confluent hypergeometric function that is related to a Bessel function in integral form. Writing this integral twice for a product of two Whittaker functions gives

$$\mathcal{M}_{\lambda_{1},\mu_{1}/2}(z_{1}) \,\mathcal{M}_{\lambda_{2},\mu_{2}/2}(z_{2}) = \frac{4 \,(z_{1}z_{2})^{1/2} \,e^{(z_{1}+z_{2})/2}}{\Gamma\left(\frac{1+\mu_{1}}{2}+\lambda_{1}\right) \,\Gamma\left(\frac{1+\mu_{2}}{2}-\lambda_{2}\right)} \times \int_{0}^{\infty} dt \int_{0}^{\infty} du \,e^{-t^{2}-u^{2}} t^{2\lambda_{1}} u^{2\lambda_{2}} J_{\mu}(2t\sqrt{z_{1}}) J_{\mu}(2u\sqrt{z_{2}}) ,$$
(D3)

whence a large class of representations can be developed by appropriate variable substitutions. For the important case of products of functions with the same indices, choosing $\lambda_1 = -\lambda_2 \equiv \lambda$ and $\mu_1 = \mu_2 \equiv \mu$, the following representation is obtained by going to "polar coordinates" via $t = \rho \cos \phi$ and $u = \rho \sin \phi$, and evaluating the integral with respect to ρ via Weber's second exponential integral (B11),

$$\mathcal{M}_{i\kappa,\mu/2}(-ix') \,\mathcal{M}_{-i\kappa,\mu/2}(ix'') = \frac{2 \,(x'x'')^{1/2} e^{i(x'-x'')/2}}{\Gamma\left(\frac{1+\mu}{2} + i\kappa\right) \,\Gamma\left(\frac{1+\mu}{2} - i\kappa\right)} \\ \times \int_0^{\frac{\pi}{2}} d\phi \,(\cot\phi)^{2i\kappa} \,e^{i(x'\cos^2\phi - x''\sin^2\phi)} \,I_\mu\left(\sqrt{x'x''}\sin 2\phi\right) \,, \tag{D4}$$

where we are also making the assignments $\lambda = i\kappa$, $z_1 = -ix'$, and $z_2 = ix''$, tailored specifically to the Whittaker functions with imaginary first index needed for the radial inverted oscillator or the conformal operator S. From Eq. (D4), a whole class of integrals can be obtained by appropriate substitutions. In particular, if ζ is defined such that $\sin 2\phi = 1/(i \sinh \zeta)$, then Eq. (D4) turns into a form that appears to match the propagator of Eq. (66), with $\zeta = \omega t$. However, this is a nontrivial substitution that takes the variable ζ into the complex plane and requires further analysis. We will first implement this substitution in two steps: (i) defining a real variable *s* such that $\sin 2\phi = 1/\cosh s$; (ii) replacing *s* by ζ via $\cosh s = i \sinh \zeta$ in the complex plane. Moreover, there are issues with the interpretation of the integral in the complex plane that require an appropriate deformation of the integration contour. This is shown next.

First, from the range of the integral (D4), the substitution $\sin 2\phi = 1/\cosh s$ establishes a one-to-one correspondence that maps $2\phi \in [0, \pi]$ into $s \in (-\infty, \infty)$, only involving real variables. By straightforward algebra, $d\phi = ds/(2\cosh s)$, $\cos 2\phi = \tanh s$, and $\cot \phi = e^s$, which directly yields an integral of the well-known form of Eq. (3a) in Sec. 6.1 of Ref. [69] in the specific variant

$$\mathcal{M}_{i\kappa,\mu/2}(-ix') \mathcal{M}_{-i\kappa,\mu/2}(ix'') = \frac{(x'x'')^{1/2}}{\Gamma\left(\frac{1+\mu}{2} + i\kappa\right) \Gamma\left(\frac{1+\mu}{2} - i\kappa\right)} \times \int_{-\infty}^{\infty} \frac{ds}{\cosh s} \exp\left(2i\kappa s\right) \exp\left[\frac{i}{2}\left(x' + x''\right) \tanh s\right] I_{\mu}\left(\frac{\sqrt{x'x''}}{\cosh s}\right)$$
(D5)

(which agrees with Ref. [103], 6.669-5).

Second, the transformation $\cosh s = i \sinh \zeta$ is evidently a translation $\zeta = s + \sigma$ along the imaginary axis in the complex plane, with displacement σ . The general solution of this equation is $\zeta = \pm s - i\pi/2 + 2\pi n$, with $n \in \mathbb{Z}$, which follows from the $2\pi i$ periodicity of the cosh function (or simply by finding the solution via inversion of the hyperbolic or exponential functions). Moreover, the negative sign can be ignored if we only consider the case of a path without inversion. This shows that the translation displacement is $\sigma = \zeta_n \equiv \zeta_0 + 2\pi n$, with the "principal value" $\zeta = ic$, with $c = -\pi/2$, which we will use below. With this transformation, the following relations are immediately satisfied: $ds = d\zeta$, $\sinh s = i \cosh \zeta$, $\tanh s = \coth \zeta$, $\sinh s = i \cosh \zeta$, and $e^s = e^{i\pi/2}e^{\zeta}$. As a result, the integration path for the transformed version of the integral (D5) is an infinite straight line L parallel to the real ζ axis, with $\operatorname{Im} \zeta = \operatorname{Im} \sigma = c$. With this procedure, we have established a novel identity

$$\mathcal{M}_{i\kappa,\mu/2}(-ix') \mathcal{M}_{-i\kappa,\mu/2}(ix'') = e^{-\pi\kappa} \frac{(x'x'')^{1/2}}{\Gamma\left(\frac{1+\mu}{2} + i\kappa\right) \Gamma\left(\frac{1+\mu}{2} - i\kappa\right)} \times \int_{ic-\infty}^{ic+\infty} \frac{d\zeta}{i\sinh\zeta} \exp\left(2i\kappa\zeta\right) \exp\left[\frac{i}{2}\left(x'+x''\right)\coth\zeta\right] I_{\mu}\left(\frac{\sqrt{x'x''}}{i\sinh\zeta}\right) .$$
(D6)

The integral of direct applicability for the required form of the propagator (66) involves using a path with $c \to 0$. This can be justified from the following properties: (i) the only singularities of the integrand occur at $\zeta = 2\pi ni$, which we will circumvent with appropriate integration contours; and (ii) the behavior is regular at infinity along curves asymptotically parallel to the real axis, with the integral vanishing exponentially for $|\operatorname{Re} \zeta| \to \infty$. Thus, $\oint_C F(\zeta) d\zeta = 0$, for the integrand $F(\zeta)$ in Eq. (D6) with a closed contour C consisting of the line L and a parallel line L' with $c \in (-\pi/2, 0)$. This proves Eq. (D6) for arbitrary c with $-\pi/2 < c < 0$. Moreover, if the restriction to straight lines is relaxed, the line L' can further deformed into any other path with asymptotic limits $ic' - i\infty$ and $ic'' + i\infty$, where c' and c''are arbitrary. For the propagator (66), it suffices to take the limit $c \to 0$. It should be noted that this is an analog for Whittaker functions of the Bessel-function identity (D1). With this auxiliary integral (D6), making the replacements $\zeta = \omega T$, $x' = M\omega r'^2/\hbar$, $x'' = M\omega r''^2/\hbar$, $\kappa = E/(2\hbar\omega)$, and $c = 0^{\mp}$, the main result of Eqs. (78)–(79) in Sec. VI is straightforwardly derived.

An alternative set of representations can be established by performing integrals of the forms (D5) and (D6), but restricted to the half-axis intervals. For example, Eq. (5b) in Sec. 6.1 of Ref. [69] reads

$$W_{\kappa,\mu/2}(a_1t) \mathcal{M}_{\kappa,\mu/2}(a_2t) = \frac{t\sqrt{a_1a_2}}{\Gamma\left((1+\mu)/2 - \kappa\right)} \cdot \int_0^\infty d\xi \, \exp\left[-\frac{1}{2}\left(a_1 + a_2\right)t\cosh\xi\right] I_\mu\left(t\sqrt{a_1a_2}\sinh\xi\right) \left[\coth\left(\frac{\xi}{2}\right)\right]^{2\kappa} \quad (D7)$$

(which agrees with Ref. [103], 6.669-4), where $a_{1,2}$ are real parameters satisfying the critical inequality $a_1 > a_2$, and $\operatorname{Re}((1+\mu)/2-\kappa) > 0$. The variables t, a_1 , and a_2 in Eq. (D7), provide some flexibility of choices; however, as shown in the steps leading to their derivation, with the notation used in Eq. (D3), one can identify $z_1 = ta_1$ and $z_2 = ta_2$. Then, as before, one can make the additional replacements $\kappa \to i\kappa$ and $z_{1,2} = -ix', -ix''$. (This could be done most easily with t = -i and $a_{1,2}$ chosen from the set x', x''.) Finally, with the substitution $\sinh \xi = 1/\sinh \zeta$ (which implies $\cosh \xi = \coth \zeta$, $\coth(\xi/2) = e^{\zeta}$, and $d\xi = -d\zeta/\sinh \zeta$), Eq. (D7) turns into

$$W_{i\kappa,\mu/2}(ix_{>}) \mathcal{M}_{i\kappa,\mu/2}(ix_{<}) = \frac{\sqrt{x'x''}}{\Gamma\left((1+\mu)/2 - i\kappa\right)} \times \int_{0}^{\infty} \frac{d\zeta}{i\sinh\zeta} \exp\left(2i\kappa\zeta\right) \exp\left[\frac{i}{2}\left(x'+x''\right)\coth\zeta\right] I_{\mu}\left(\frac{\sqrt{x'x''}}{i\sinh\zeta}\right) ,$$
(D8)

where $x_{>}$ and $x_{<}$ are the greater and lesser of the set $\{x', x''\}$ respectively.

Incidentally, the known identity of Eq. (D7) is proved, as stated in Ref. [69], by using appropriate substitutions combined with identities (semi-circuital and connection) among Whittaker functions. In fact, this is a reversal of the procedure we used in the main text; in other words, we could just as well use the symmetry property (72), along with the Whittaker identities (74) and (75), and with appropriate limits, to justify the half-axis integrals(s) (D8) from the full integral (D6); and from Eq. (D8) rederive Eq. (D7) via $\sinh \xi = 1/\sinh \zeta$.

Appendix E: Generalized Symmetry Generators of CQM—Differential Equation and Consistency Checks

In this appendix, we consider another aspect of the CQM generalized generator G = uH + vD + wK: its behavior and spectral properties using a differential equation approach. These properties can be deduced by going to the Schrödinger picture for the general multicomponent case of the dAFF model we introduced in Sec. II (*d* components as *d*-dimensional position coordinates in quantum mechanics). This setup for *d* dimensions and for the generic operator *G* extends the particular results of Ref. [5].

Then, in the Schrödinger picture, with the Hamiltonian $\tilde{H}_G(\mathbf{r}, \mathbf{p}) \equiv G/\sigma$ introduced in Eq. (A1), the states $\Psi(\mathbf{r}, \tau)$ evolve according to

$$i\hbar \frac{\partial \Psi(\mathbf{r},\tau)}{\partial \tau} = \tilde{H}_G(\mathbf{r},\mathbf{p}) \ \Psi(\mathbf{r},\tau) \ . \tag{E1}$$

From Eq. (E1), "stationary states" can be defined with respect to τ , such that the eigenvalue equation $\tilde{H}_G |\psi\rangle = \tilde{E}_G |\psi\rangle$ is satisfied; this leads to the radial-coordinate differential equation

$$\frac{1}{2}\frac{\hbar^2}{M} \left[-\frac{d^2}{dr^2} + \frac{g + (l+\nu)^2 - 1/4}{r^2} + \frac{M^2}{\hbar^2} \left(-\frac{\Delta}{4} \right) r^2 \right] \mathcal{U}_{\mathfrak{g},l}(r) = \tilde{E}_G \,\mathcal{U}_{\mathfrak{g},l}(r) \,, \qquad (E2)$$

with the eigenvalues of the operator G being $\hbar \mathfrak{g} = \sigma \tilde{E}_G$ —see Eq. (27) and the discussion therein. In Eq. (E2), the reduced function $\mathcal{U}_{\mathfrak{g},l}(y)$ corresponds to the *d*-dimensional wave function $\psi(\mathbf{r}) = r^{-(d-1)/2} \mathcal{U}_{\mathfrak{g},l}(r) Y_{lm}(\Omega)$, following the usual separation of angular variables.

With the substitutions $A = -i(M/\hbar)\sqrt{\Delta}/2$, $\lambda = i\sigma \mathfrak{g}/\sqrt{\Delta}$, and $\mu = g + (l+\nu)^2$, Eq. (E2) is of the known form (Ref. [69], p. 34)

$$\left[-\frac{d^2}{dr^2} + \frac{\mu^2 - 1/4}{r^2} + A^2 r^2\right] \Phi(r) = 4\lambda A \Phi(r) , \qquad (E3)$$

which can be solved exactly in terms of Whittaker functions. Specifically, the standard Whittaker differential equation for a function P(z) can be transformed into Eq. (E3) with the substitution $z = Ar^2$, such that the solutions are

$$\Phi(r) = r^{-1/2} P_{\lambda,\mu/2}(Ar^2) , \qquad (E4)$$

where, with the notation of Ref. [69], $P_{\lambda,\mu/2}(z)$ stands for either one of the Whittaker functions $\mathcal{M}_{\lambda,\mu/2}(z)$ and $W_{\lambda,\mu/2}(z)$, or a linear combination thereof. The substitutions used in Eq. (E3), which involve the scale A and the index λ , are completely general, covering all classes of operators (elliptic, parabolic, and hyperbolic), according to the value of Δ .

Furthermore, enforcing the regular behavior at the origin, the function $W_{\lambda,\mu/2}(z)$ should be excluded for the wave function solutions—though it is still relevant for the Green's functions (see below). Then, using the values for A, λ , and μ , the ensuing regular choice of the solution (E4) to Eq. (E2) becomes

$$\mathcal{U}(r) = r^{-1/2} \mathcal{M}_{i\sigma\mathfrak{g}/\sqrt{\Delta},\mu/2}(-i\check{r}^2) , \qquad (E5)$$

where

$$\check{r}^2 = \frac{\sqrt{\Delta}}{2} \left(\frac{M}{\hbar}\right) r^2 \equiv \frac{M|\omega|}{\hbar} r^2 .$$
(E6)

The variable \check{r} in Eq. (E6) is written in terms of the general CQM frequency of Eq. (29), i.e., $\omega = -i|\omega|$, which is in agreement with the definitions in Sec. VI: in Eq. (68) and with the analytic extension $\omega \to -i\omega = -i\sqrt{\Delta}/2$ used therein. Moreover, the solution (E5) gives a continuous spectrum with $\mathfrak{g} \in (-\infty, \infty)$. This set of results reproduces the path-integral treatment leading to Eq. (81) [with Eq. (68)] for the hyperbolic-operator eigenfunctions. Most importantly, Eq. (E5) is completely general; when $\Delta < 0$, it can also be applied to the elliptic case—this amounts to the analytic continuation $\kappa = i\sigma \mathfrak{g}/\sqrt{\Delta} \to \sigma \mathfrak{g}/\sqrt{\Delta}$, which (from the asymptotics) requires eigenvalue "quantization" leading to the generalized Laguerre polynomials, as in Eq. (53). For the parabolic case, the limit $\Delta = 0$ leads to Bessel functions $J_{\mu}(kr)$ as regular solutions, as can also be verified directly from Eq. (E2). In the latter case, there is a known subtlety or ambiguity in the choice of the sign of the regular solution from the set $J_{\pm\mu}(kr)$, with a physical argument [37] selecting the positive sign—this is related to the inequivalence of the self-adjoint extension method with physical regularization techniques [104]. In short, the differential-equation approach fully agrees with the path-integral results of the main text for all the conformal generators.

In addition, the Green's functions can be derived from the corresponding differential equation (E2), using the general solutions we already found, Eq. (E4). The usual notational simplification $E \equiv \tilde{E}_G$ is used below [or else, this symbol could be replaced in favor of \mathfrak{g} , according to Eq. (27)]. The radial energy Green's functions $G_{l+\nu}^{(\pm)}(r'', r'; E)$ are defined in Eq. (40), which we will rescale as $\mathcal{G}_{l+\nu}^{(\pm)}(r'', r'; E) = (\hbar^2/2M) G_{l+\nu}^{(\pm)}(r'', r'; E)$. As is well-known, because of the specific form of Eqs. (C5)–(C6), which effectively invert the Schrödinger eigenvalue equation, these coincide with the Green's functions used for the solution of linear differential equations. Then, the radial differential equation for $\mathcal{G}_{l+\nu}(r'', r'; E)$ reads

$$\left\{\frac{d^2}{dr'^2} + \frac{2M}{\hbar^2} \left[E - V(r')\right] - \frac{(l+\nu)^2 - 1/4}{r'^2}\right\} \mathcal{G}_{l+\nu}(r'', r'; E) = \delta(r'' - r') , \qquad (E7)$$

which is a particular case of a one-dimensional Sturm-Liouville problem (with constant coefficient p(r) = 1 for the second-order derivative) [105]. Equation (E2) is the homogeneous form of Eq. (E7), with the obvious identifications; then, defining the the functions $\mathcal{U}_{(<)}^{(\pm)}(r)$ and $\mathcal{U}_{(>)}^{(\pm)}(r)$ that satisfy boundary conditions at the left boundary (here: r = 0) and right boundary (here: $r = \infty$), then [105]

$$\mathcal{G}_{l+\nu}^{(\pm)}(r'',r';E) = \frac{\mathcal{U}_{(<)}^{(\pm)}(r_{<}) \, \mathcal{U}_{(>)}^{(\pm)}(r_{>})}{p(r') \, \mathfrak{W}\left\{\mathcal{U}_{(<)}^{(\pm)} \,, \, \mathcal{U}_{(>)}^{(\pm)}\right\}(r')} \,, \tag{E8}$$

where $r_{<}(r_{>})$ is the lesser (greater) of r' and r'' and $\mathfrak{W}\left\{\mathcal{U}_{(<)}^{(\pm)}, \mathcal{U}_{(>)}^{(\pm)}\right\}$ is the Wronskian of $\mathcal{U}_{(<)}^{(\pm)}(r)$ and $\mathcal{U}_{(>)}^{(\pm)}(r)$. This Green's function technique has also been used for the study of the related strong-coupling inverse square potential regularization [106].

For Eq. (E2), from the general solution (E4), we identify

$$\mathcal{U}_{(<)}^{(\pm)}(r) = r^{-1/2} \,\mathcal{M}_{\pm i\sigma\mathfrak{g}/\sqrt{\Delta},\mu/2}(\mp i\check{r}^2) \tag{E9}$$

as the solutions that satisfy a regular boundary condition $\mathcal{U}(r)|_{r=0} = 0$ at the origin; and

$$\mathcal{U}_{(>)}^{(\pm)}(r) = r^{-1/2} W_{\pm i\sigma \mathfrak{g}/\sqrt{\Delta}, \mu/2}(\mp i \check{r}^2)$$
(E10)

as the solutions that satisfy the appropriate asymptotic boundary condition at infinity (outgoing/incoming waves for $G^{(\pm)}$). The identity (74) shows that $\mathcal{U}_{(<)}^{(\pm)} \propto \mathcal{M}_{\pm i\sigma \mathfrak{g}/\sqrt{\Delta},\mu/2}$ are actually the same function, up to a proportionality constant; however, $\mathcal{U}_{(>)}^{(\pm)} \propto W_{\pm i\sigma \mathfrak{g}/\sqrt{\Delta},\mu/2}$ are distinctly different. The Wronskian can be computed with the identity $\mathfrak{W} \{\mathcal{M}_{\lambda,\mu/2}, W_{\lambda,\mu/2}\}(z) = -[\Gamma((1+\mu)/2-\lambda)]^{-1}$; using Eqs. (E9) and (E10), along with p(r) = 1, and applying the chain rule for $z = \mp i\tilde{r}^2$, the final result for $\mathcal{G}_{l+\nu}^{(\pm)}(r'', r'; E)$ is identical to Eq. (77). This again verifies the equivalence of the path-integral and differentialequation approaches, and provides additional consistency checks for the network of relations defined in this paper.

- C. Grosche and F. Steiner, Handbook of Feynman Path Integrals (Springer-Verlag, Berlin, 1998); and references therein.
- [2] C. Grosche and F. Steiner, How to solve path integrals in quantum mechanics, J. Math. Phys. 36, 2354 (1995).
- [3] H. Kleinert, Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets, 5th ed. (World Scientific, Singapore, 2009); and references therein.
- [4] R. Jackiw, Introducing scale symmetry, Physics Today 25 (1), 23 (1972).
- [5] V. de Alfaro, S. Fubini and G. Furlan, Conformal Invariance in quantum mechanics, Nuovo Cimento A34, 569 (1976).
- [6] R. Jackiw, Dynamical symmetry of the magnetic monopole, Ann. Phys. (N.Y.) 129, 183 (1980).
- [7] R. Jackiw, Dynamical symmetry of the magnetic vortex, Ann. Phys. (N.Y.) 201, 83 (1990).
- [8] R. Jackiw, Delta function potentials in two-dimensional and three-dimensional quantum mechanics, in M. A. B. Bég Memorial Volume, A. Ali and P. Hoodbhoy, eds. (World Scientific, 1991).
- [9] H. E. Camblong, L. N. Epele, H. Fanchiotti, and C. A. García Canal, Quantum anomaly in molecular physics, Phys. Rev. Lett. 87, 220402 (2001).
- [10] H. E. Camblong, and C. R. Ordóñez, Black hole thermodynamics from near-horizon conformal quantum mechanics, Phys. Rev. D 71, 104029 (2005).
- [11] H. E. Camblong and C. R. Ordóñez, Semiclassical methods in curved spacetime and black

hole thermodynamics, Phys. Rev. D 71, 124040 (2005).

- [12] H. E. Camblong, and C. R. Ordóñez, Conformal tightness of holographic scaling in black hole thermodynamics, Classical and Quantum Gravity 30, 175007 (2013).
- [13] H. E. Camblong, A. Chakraborty, and C. R. Ordóñez, Near-horizon aspects of acceleration radiation by free fall of an atom into a black hole, Phys. Rev. D 102, 085010 (2020).
- [14] A. Azizi, H. E. Camblong, A. Chakraborty, C. R. Ordóñez, and M. O. Scully, Acceleration radiation of an atom freely falling into a Kerr black hole and near-horizon conformal quantum mechanics, Phys. Rev. D 104, 065006 (2021).
- [15] A. Azizi, H. E. Camblong, A. Chakraborty, C. R. Ordóñez, and M. O. Scully, Quantum optics meets black hole thermodynamics via conformal quantum mechanics: I. Master equation for acceleration radiation, Phys. Rev. D 104, 084086 (2021).
- [16] Quantum optics meets black hole thermodynamics via conformal quantum mechanics: II. Thermodynamics of acceleration radiation, A. Azizi, H. E. Camblong, A. Chakraborty, C. R. Ordóñez, and M. O. Scully, Phys. Rev. D 104, 084085 (2021).
- [17] V. Efimov, Energy levels arising from resonant two-body forces in a three-body system. Phys. Lett. B 33, 563 (1970).
- [18] V. Efimov, Weakly-bound states of three resonantly-interacting particles. Sov. J. Nucl. Phys 12, 589 (1971).
- [19] P. Naidon and S. Endo, Efimov Physics: a review, Rept. Prog. Phys. 80, 056001 (2017).
- [20] H. E. Camblong and C. R. Ordóñez, Renormalization in conformal quantum mechanics, Physics Letters A 345, 22 (2005).
- [21] A. Gorsky and F. Popov Atomic collapse in graphene and cyclic renormalization group flow, Phys. Rev. D 89, 061702 (2014).
- [22] O. Ovdat, Jinhai Mao, Yuhang Jiang, E. Y. Andrei, and E. Akkermans, Observing a scale anomaly and a universal quantum phase transition in graphene, Nature Communications 8, 507 (2017).
- [23] W. Daza, J. E. Drut, C. Lin, and C. Ordóñez, Virial expansion for the Tan contact and Beth-Uhlenbeck formula from two-dimensional SO(2,1) anomalies, Phys. Rev. A 97, 033630 (2018).
- [24] J. E. Drut, J. R. McKenney, W. S. Daza, C. L. Lin, and C. R. Ordóñez, Quantum Anomaly and Thermodynamics of One-Dimensional Fermions with Three-Body Interactions, Phys.

Rev. Lett. **120**, 243002 (2018).

- [25] J. Maki and C. R. Ordóñez, Virial expansion for a three-component Fermi gas in one dimension: The quantum anomaly correspondence, Phys. Rev. A 100, 063604 (2019).
- [26] H. E. Camblong, A. Chakraborty, W. S. Daza, J. E. Drut, C. L. Lin, and C. R. Ordóñez, Quantum anomaly and thermodynamics of one-dimensional fermions with antisymmetric two-body interactions, Ann. Phys. (N.Y.) 429, 168466 (2021).
- [27] H. Tajima, S. Tsutsui, T. M. Doi, and K. Iida, Cooper triples in attractive three-component fermions: Implication for hadron-quark crossover Phys. Rev. Research 4, L012021 (2022).
- [28] G. N. J. Añaños, H. E. Camblong, C. Gorrichátegui, E. Hernández, and C. R. Ordóñez, Anomalous commutator algebra for conformal quantum mechanics, Phys. Rev. D 67, 045018 (2003).
- [29] G. N. J. Añaños, H. E. Camblong, and C. R. Ordóñez, SO(2,1) conformal anomaly: Beyond contact interactions, Phys. Rev. D 68, 025006 (2003).
- [30] H. E. Camblong and C. R. Ordóñez, Anomaly in conformal quantum mechanics: From molecular physics to black holes, Phys. Rev. D 68, 125013 (2003).
- [31] K.S. Gupta and S.G. Rajeev, Renormalization in quantum mechanics, Phys. Rev. D 48, 5940 (1993).
- [32] H. E. Camblong, L. N. Epele, H. Fanchiotti, and C. A. García Canal, Renormalization of the inverse square potential, Phys. Rev. Lett. 85, 1590 (2000).
- [33] H. E. Camblong, L. N. Epele, H. Fanchiotti, and C. A. García Canal, Dimensional transmutation and dimensional regularization in quantum mechanics: I. General theory, Ann. Phys. (NY) 287, 14 (2001).
- [34] H. E. Camblong, L. N. Epele, H. Fanchiotti, and C. A. García Canal, Dimensional transmutation and dimensional regularization in quantum mechanics: II. Rotational invariance, Ann. Phys. (NY) 287, 57 (2001).
- [35] S. R. Beane et al., Singular potentials and limit cycles, Phys. Rev. A 64, 042103 (2001).
- [36] D. B. Kaplan, J.-W. Lee, D. T. Son, D. T., and M. A. Stephanov, Conformality lost, Phys. Rev. D 80, 125005 (2009).
- [37] L. D. Landau and E. M. Lifshitz, Quantum Mechanics, 3rd ed. (Pergamon, 1977), p. 114.
- [38] H. E. Camblong, L. N. Epele, H. Fanchiotti, C. A. García Canal, and C. R. Ordónez, Multichannel Framework for Singular Quantum Mechanics, Ann. Phys. (NY) 340, 267 (2014).

- [39] C. Chamon, R. Jackiw, S.-Y. Pi, and L. Santos, Conformal quantum mechanics as the CFT₁ dual to AdS₂, Phys. Lett. B **701**, 503 (2011).
- [40] R. Jackiw and S.Y. Pi, Conformal blocks for the 4-point function in conformal quantum mechanics, Phys. Rev. D 86, 045017 (2012) [Erratum ibid. 86 (2012) 089905]
- [41] T. Okazaki, Whittaker vector, Wheeler-DeWitt equation, and the gravity dual of conformal quantum mechanics, Phys. Rev. D 92, 126010 (2015).
- [42] T. Okazaki, Implications of Conformal Symmetry in Quantum Mechanics, Phys. Rev. D 96 (2017) 066030 [1704.00286]
- [43] A. Pinzul and A. Stern, Non-commutative AdS₂/CFT₁ duality: The case of massless scalar fields, Phys. Rev. D 96, 066019 (2017).
- [44] S. Khodaee and D. Vassilevich, Note on correlation functions in conformal quantum mechanics, Mod. Phys. Lett. A 32, 1750157 (2017).
- [45] F. R. de Almeida, A. Pinzul, and A. Stern, Non-commutative AdS₂/CFT₁ duality: The case of massive and interacting scalar fields, Phys. Rev. D 100, 086005 (2019).
- [46] R. L. Ardón, Conformal quantum mechanics as a Floquet-Dirac system, arXiv:2103.15248 (2021).
- [47] M. Arzano, Conformal quantum mechanics of causal diamonds, Journal of High Energy Physics, 2020, 1-14 (2020).
- [48] M. Arzano, Vacuum thermal effects in flat space-time from conformal quantum mechanics, J. High Energ. Phys. 2021, 3 (2021).
- [49] P. Martinetti and C. Rovelli, Diamond's temperature: Unruh effect for bounded trajectories and thermal time hypothesis, Classical and Quantum Gravity 20(22), 4919 (2003).
- [50] W. G. Unruh, Notes on black-hole evaporation, Phys. Rev. D, 14(4), 870 (1976).
- [51] P. Martinetti, Conformal mapping of Unruh temperature, Modern Physics Letters A 24(19), 1473-1483 (2009).
- [52] D. Su and T. C. Ralph, Spacetime diamonds, Phys. Rev. D 93(4), 044023 (2016).
- [53] J. Foo, S. Onoe, M. Zych, and T. C. Ralph, Generating multi-partite entanglement from the quantum vacuum with a finite-lifetime mirror, New Journal of Physics 22(8), 083075 (2020).
- [54] T. De Lorenzo and A. Perez, Light cone thermodynamics, Phys. Rev. D 97(4), 044052 (2018)
- [55] T. Jacobson and M. Visser, Gravitational thermodynamics of causal diamonds in (A)dS, SciPost Phys. 7(6) (2019)

- [56] A. Chakraborty, H. E. Camblong, and C. R. Ordóñez, Thermal effect in a causal diamond: open quantum systems approach, Phys. Rev. D 106, 045027 (2022).
- [57] A. Herrero and J. A. Morales, Radial conformal motions in Minkowski space-time, J. Math. Phys. 40, 3499 (1999).
- [58] B. Wybourne, Classical Groups for Physicists (Wiley, New York, 1974).
- [59] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, eds., *Higher Transcendental Functions* (McGraw-Hill, New York, 1955), Vol. 2.
- [60] C. Grosche and F. Steiner, Path integrals on curved manifolds, Zeitschr. Phys. C 36, 699 (1987).
- [61] A. Inomata, H. Kuratsuji, and C. C. Gerry, Path integrals and coherent states of SU (2) and SU (1, 1) (World Scientific, 1992).
- [62] J. H. Van Vleck, in Wave Mechanics, the First Fifty Years, ed. W. C. Price et al. (Butterworth, London, 1973), p. 26.
- [63] W. Fischer, H. Leschke, and P. Müller, Changing dimension and time: Two well-founded and practical techniques for path integration in quantum physics, J. Phys. A: Math. Gen. 25, 3835 (1992).
- [64] D. Peak and A. Inomata, Summation over Feynman Histories in Polar Coordinates, J. Math. Phys. 10, 1422 (1969).
- [65] G. N. Watson, A Treatise on the Theory of Bessel Functions, 2nd ed. (Cambridge University Press, Cambridge, England, 1944).
- [66] F.W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, NIST Handbook of Mathematical Functions (Cambridge University Press, Cambridge, England, 2010).
- [67] V. Subramanyan, S. Hegde, S. Suraj, Vishveshwara, and B. Bradlyn, Physics of the Inverted Harmonic Oscillator: From the lowest Landau level to event horizons, Ann. Phys. (N.Y.) 435, 168470 (2021).
- [68] G. Barton, Quantum mechanics of the inverted oscillator potential, Ann. Phys. (N.Y.) 166, 322 (1986).
- [69] H. Buchholz, The confluent hypergeometric function (Springer-Verlag, New York, 1969).
- [70] V. Moretti and N. Pinamonti, Aspects of hidden and manifest $SL(2,\mathbb{R})$ symmetry in 2D near-horizon black-hole backgrounds, Nucl. Phys. B **647**,131 (2002).
- [71] B. Pioline and A. Waldron, Quantum cosmology and conformal invariance, Phys. Rev. Lett.

90, 031302 (2003).

- [72] J. Ben Achour and E.R. Livine, Cosmology as a CFT₁, JHEP **12** (2019) 031.
- [73] J. Ben Achour and E. R. Livine, Protected $SL(2,\mathbb{R})$ Symmetry in Quantum Cosmology, JCAP **1909**, 012 (2019).
- [74] S. Dalui, B. Majhi, and P. Mishra, Horizon induces instability locally and creates quantum thermality, Phys. Rev. D. 102, 044006 (2020).
- S. Dalui and B. Majhi, Near-horizon local instability and quantum thermality, Phys. Rev.
 D. 102, 124047 (2020).
- [76] A. Bhattacharyya, W. Chemissany, and S. S. Haque, J. Murugan, and B. Yan, The multifaceted inverted harmonic oscillator: Chaos and complexity, SciPost Phys. 4, 002 (2021).
- [77] L.-C. Qu, J. Chen, and Y.-X. Liu, Chaos and complexity for inverted harmonic oscillators, Phys. Rev. D 105, 126015 (2022).
- [78] T. G. Mertens, G. J. Turiaci, and H. L. Verlinde, Solving the Schwarzian via the conformal bootstrap, JHEP 1708 (2017) 136.
- [79] H.T. Lam, T.G. Mertens, G.J. Turiaci, H. Verlinde, Shockwave S-matrix from Schwarzian quantum mechanics, JHEP 1811 (2018) 182.
- [80] A. Galajinsky, A variant of Schwarzian mechanics, Nucl. Phys. B 936, 661 (2018).
- [81] A. Galajinsky, Schwarzian mechanics via nonlinear realizations, Phys. Lett. B 795, 277 (2019).
- [82] S. Filyukov and I. Masterov, Generalized Schwarzian mechanics, Nucl. Phys. B 964 115316 (2021).
- [83] I. Masterov and M. Masterova, Coupling-constant metamorphosis in SL(2, ℝ)-invariant systems, J. Geom. Phys. 168, 104320 (2021).
- [84] J. Maldacena and D. Stanford, Remarks on the Sachdev-Ye-Kitaev model, Phys. Rev. D 94 106002 (2016).
- [85] T. Tada, Conformal quantum mechanics and sine-square deformation, Progr. Theor. Exp. Phys. 2018, 061B01 (2018).
- [86] U. Niederer, The maximal kinematical invariance group of the harmonic oscillator, Helv. Phys. Acta 46, 191 (1973).
- [87] S. Takagi, Equivalence of a Harmonic Oscillator to a Free Particle, Prog. Theor. Phys. 84, 1019 (1990).

- [88] S. Takagi, Quantum Dynamics and Non-Inertial Frames of Reference. I: Generality, Prog. Theor. Phys. 85, 463 (1991).
- [89] S. Takagi, Quantum Dynamics and Non-Inertial Frames of Reference. II: Harmonic Oscillators, Prog. Theor. Phys. 85, 723 (1991).
- [90] Y. Castin, Exact scaling transform for a unitary quantum gas in a time-dependent harmonic potential, C. R. Phys. 5, 407 (2004).
- [91] E. Wamba, A. Pelster, and J. R. Anglin, Exact quantum field mappings between different experiments on quantum gases, Phys. Rev. A 94, 043628 (2016).
- [92] A. Strominger, A matrix model for AdS_2 , JHEP **03** (2004) 066.
- [93] M. Hotta, A. Kempf, E. Martín-Martínez, T. Tomitsuka, and K. Yamaguchi, Duality in the dynamics of Unruh-DeWitt detectors in conformally related spacetimes, Phys. Rev. D 101, 085017 (2020).
- [94] B. DeWitt, Dynamical Theory in Curved Spaces. I. A Review of the Classical and Quantum Action Principles, Rev. Mod. Phys. 29, 377 (1957).
- [95] S. F. Edwards and Y. V. Gulyaev, Path integrals in polar co-ordinates, Proc. R. Soc. A279, 229 (1964).
- [96] J.-L. Gervais and A. Jevicki, Point canonical transformations in the path integral, Nucl. Phys. B110, 93 (1976).
- [97] K. M. Apfeldorf and C. Ordóñez, Coordinate redefinition invariance and "extra" terms, Nucl. Phys. B479, 515 (1996).
- [98] K. M. Apfeldorf, H. E. Camblong, and C. R. Ordóñez, Field redefinition invariance in quantum field theory, Mod. Phys. Lett. 16, 103 (2001).
- [99] M. J. Goovaerts, Path-integral evaluation of a nonstationary Calogero model, J. Math. Phys. 16, 720 (1975).
- [100] K. V. Bhagwat and S. V. Lawande, A new derivation of the Feynman propagator for the inverse square potential, Phys. Lett. A141, 321 (1989).
- [101] H. E. Camblong and C. R. Ordóñez, Path Integral Treatment of Singular Problems and Bound States, Int. J. of Mod. Phys. A 19, 1413 (2004).
- [102] H. E. Camblong and C. R. Ordóñez, Renormalized Path Integral for the Two-Dimensional Delta-Function Interaction, Phys. Rev. A 65, 052123 (2002).
- [103] I. S. Gradshteyn and I. M. Ryzhik, Table of Integrals, Series, and Products, 6th ed. (Academic

Press, New York, 2000).

- [104] H. E. Camblong, L. N. Epele, H. Fanchiotti, C. A. García Canal, and C. R. Ordónez, On the inequivalence of renormalization and self-adjoint extensions for quantum singular interactions, Phys. Lett. A 364, 458 (2007).
- [105] I. Stakgold, Green's Functions and Boundary Value Problems, 2nd ed. (Wiley, 1998).
- [106] H. E. Camblong and C. R. Ordóñez, Regularized Green's function for the inverse square potential, Mod. Phys. Lett. A17, 817 (2002).