On the new properties of conditional expectations with applications in finance

Ismihan Bayramoglu

Department of Mathematics, Izmir University of Economics Izmir, Turkey

November 7, 2022

Abstract

The concept of conditional expectation is important in applications of probability and statistics in many areas such as reliability engineering, economy, finance, and actuarial sciences due to its property of being the best predictor of a random variable as a function of another random variable. This concept also is essential in the martingale theory and theory of Markov processes. Even though, there has been studied and published many interesting properties of conditional expectations with respect to a sigmaalgebra generated by a random variable it remains an attractive subject having interesting applications in many fields. In this paper, we present some new properties of the conditional expectation of a random variable given another random variable and describe useful applications in problems of pershare-price of stock markets. The copula and dependence properties of conditional expectations as random variables are also studied. We present also some new equalities having interesting applications and results in martingale theory and Markov processes.

Keywords: Conditional expectation, sigma algebra, per-share price, order statistics, prediction **Conflicts of interest statement:** We declare that have no conflicts of interest.

1 Introduction

Let (Ω, \mathcal{F}, P) be a probability space. Consider the random variables X and Y defined in this probability space and having joint distribution function $F_{X,Y}(x,y) = C(F_X(x), F_Y(y)), (x,y) \in \mathbb{R}^2$, where $C(u,v), (u,v) \in I^2 \equiv [0,1]^2$ is a connected copula. Consider functions $\varphi(y) = E(X \mid Y = y), y \in \mathbb{R}$ and $\psi(x) = E(Y \mid X = x), x \in \mathbb{R}$ and random variables $Z_1 \equiv \varphi(Y) = E(X \mid Y)$ and $Z_2 \equiv \psi(X) = E(Y \mid X)$. It is well known that the best predictor of X by Y in the sense of least square distance is $\varphi(Y)$, and the best predictor of Y by X is $\psi(X)$, i.e.

$$\min_{g} E(X - g(Y))^{2} = E(X - \varphi(Y))^{2}$$
$$\min_{h} E(Y - h(X))^{2} = E(Y - \psi(X))^{2},$$

where the min is taken over all measurable functions g and h. The conditional expectation E(X | Y) is a random variable defined in the same probability space (Ω, Γ, P) . The random variable defined as a conditional expectation E(X | Y) is an important classical concept, it is the best predictor for X as a function of Y, and plays a crucial role in many theoretical and practical aspects of probability theory. For example, in practical applications, if we know the joint distribution of X and Y and the value of Y, we can use $\varphi(Y)$ instead of random variable X, whose values are very difficult, expensive, or impossible to measure. A wide description of the concept of conditional expectation and its properties can be found in many books on probability and statistics including Ross (2002), Carlin and Taylor (1975), and Borovkov (1998) among others. In this paper, we aim to consider some unknown and interesting properties of conditional expectations having applications in many areas such as economics, engineering, actuarial sciences, and financial mathematics.

The paper is organized as follows. In chapter 1 we consider the conditional expectation of the random variable Y given X and compare it with the random variable defined as the arithmetic mean of conditional expectations of Y given $X_1, X_2, ..., X_n$ which are the copies (dependent or independent) of X. The application of the results in finance is shown. In Chapter 2 we are interested in the joint distribution of random variables $\varphi(Y)$ and $\psi(X)$ and study the dependence properties and copulas of these random variables. In Section 3 we consider the sequence of any random variables $X_1, X_2, ..., X_n, ...$ and study the properties of the sequence of random variables defined as $Y_1 = X_1, Y_2 = E(X_2 \mid X_1), ..., Y_n = E(X_n \mid X_{n-1}), ...$ and the sequence of random variables defined as $E(X_{n+1} \mid X_{i_1}, X_{i_2}, ..., X_{i_k}), 1 \leq i_1 < i_2 < ... < i_k \leq n, 1 \leq k \leq n$. We present some theorems describing the interesting properties of these sequences and provide examples comparing them with Markov sequences and martingales.

2 Inequalities with application to per-share-price of stock

Let $X_1, X_2, ..., X_n$ be the copies of the random variable X. Let

$$\hat{Y} = E(Y \mid X)$$
$$\bar{Y} = \frac{1}{n} \sum_{i=1}^{n} E(Y \mid X_i).$$

The following theorem compares the predicted value of Y through \hat{Y} with the predicted value of Y through \bar{Y} .

Theorem 1 It is true that

$$E(Y - \bar{Y})^2 \le E(Y - \hat{Y})^2,$$
$$E\left(Y - \frac{1}{n}\sum_{i=1}^n E(Y \mid X_i)\right)^2 \le E\left(Y - E(Y \mid X)\right)^2.$$

Proof. Using Shwarz inequality we can write

$$E\left(Y - \frac{1}{n}\sum_{i=1}^{n}E(Y \mid X_{i})\right)^{2} = E\left(\frac{1}{n}(nY - \sum_{i=1}^{n}E(Y \mid X_{i}))\right)^{2}$$
$$= \frac{1}{n^{2}}E\left(\sum_{i=1}^{n}(Y - E(Y \mid X_{i}))\right)^{2} = \frac{1}{n^{2}}\left\{E\left(\sum_{i=1}^{n}(Y - E(Y \mid X_{i}))^{2}\right)$$
$$+ 2\sum_{1 \le i < j \le n}E\left[Y - E(Y \mid X_{i})(Y - E(Y \mid X_{j}))\right]\right\}$$
$$\leq \frac{1}{n^{2}}E\left(\sum_{i=1}^{n}(Y - E(Y \mid X_{i}))^{2}\right)$$
$$+ 2\sum_{1 \le i < j \le n}\left[E\left(Y - E(Y \mid X_{i})\right)^{2}\right]^{\frac{1}{2}}\left[E\left(Y - E(Y \mid X_{j})\right)^{2}\right]^{\frac{1}{2}}$$
$$= \frac{1}{n^{2}}\left(nE\left(Y - E(Y \mid X)\right)^{2}\right) + 2\frac{n(n-1)}{2}E\left(Y - E(Y \mid X)\right)^{2}$$
$$= \frac{1}{n^{2}}\left((n+n^{2}-n)E(Y - E(Y \mid X))^{2}\right) = E(Y - E(Y \mid X))^{2}.$$

Thus the theorem proved. \blacksquare

This theorem may also be formulated as an important consequence of the following general theorem.

Theorem 2 Let X any Y be any random variables defined on the same propability space and $X_1, X_2, ..., X_n$ be the copies of X, i.e. random variables (dependent or independent) having the same distribution as X. Then

$$E\left(Y - \frac{1}{n}\sum_{i=1}^{n}X_{i}\right)^{2} \leq E(Y - X)^{2}.$$

Proof.

$$\begin{split} &E\left(Y - \frac{1}{n}\sum_{i=1}^{n}X_{i}\right)^{2} \\ &= \frac{1}{n^{2}}E\left(nY - \sum_{i=1}^{n}X_{i}\right)^{2} \\ &= \frac{1}{n^{2}}E\left(\sum_{i=1}^{n}(Y - X_{i})\right)^{2} \\ &= \frac{1}{n^{2}}\left(\sum_{i=1}^{n}E(Y - X_{i})^{2} + 2\sum_{1\leq i < j \leq n}^{n}E(Y - X_{i})E(Y - X_{j})\right) \\ &\leq \frac{1}{n^{2}}\left(\sum_{i=1}^{n}E(Y - X_{i})^{2} + 2\sum_{1\leq i < j \leq n}^{n}\left(E(Y - X_{i})^{2}\right)^{\frac{1}{2}}\left(E(Y - X_{j})^{2}\right)^{\frac{1}{2}}\right) \\ &= \frac{1}{n^{2}}\left(nE(Y - X)^{2} + \frac{n(n-1)}{2}2E(Y - X)^{2}\right) \\ &= \frac{1}{n^{2}}\left((n+n^{2}-n)E(Y - X)^{2}\right) = E(Y - X)^{2}. \end{split}$$

It is clear that if $E(Y \mid X)$ is used instead of X in Theorem 2, then Theorem 1 is obtained.

The simple Theorem 1 may have interesting and important applications. Below we provide an example demonstrating an application of this theorem in the activity of brokers coalition in stock markets.

Example 1 (Brokers in coalition) Assume that n brokers work in the financial market. Each broker has a goal, to buy a fixed number of shares of some popular stock. Let X_i be the suggested price per - share to the market by the *i*th broker respectively, i = 1, 2, ..., n, *i.e.* ith broker suggests that wants to buy this share at a price X_i . Obviously, the traded price in the market is then expected to be $\max(X_1, X_2, ..., X_n) = X_{n:n}$, where $X_{i:n}, 1 \le i \le n$ is the *i*th order statistic of $X_1, X_2, ..., X_n$. Assume that the *i*th broker wants to have a piece of information about whether he wins or loose the came, for a given (suggested by himself) price per share $X_i = x$. Therefore $P(X_i > \max(X_1, X_2, ..., X_{i-1}, X_{i+1}, ..., X_i)$ is the probability that ith broker will certainly have the stock, because he/she suggested more than others. Let $\psi_i(x) = E(X_{n:n} | X_i = x)$. Now, it is known that the conditional r.v. $\psi_i(X_i) = E(X_{n:n} \mid X_i)$ which gives information about the traded-per-share given a suggested price by broker is the predictor for $X_{n:n}$ by X_i in the sense of mean squared error. Then the value $X_{n:n} \mid X_i = x$ may be used as a measure of a successful trading strategy for ith broker. Similarly, one can predict the predicted price per - share X_i , i = 1, 2, ..., n, given the traded price in the market, which is $E(X_i \mid X_{n:n} = y)$, i = 1, 2, ..., n is the best approximate by the ith broker given the traded price y. Note that the historical values of $X_{n:n}$ are known and can be provided from the marked. It is clear that if

$$E(X_{n:n} - E(X_{n:n} \mid X_1))^2 \le E(X_{n:n} - E(X_{n:n} \mid X_2))^2,$$

then X_1 is a better price than X_2 per-share price of a stock. Now assume that n brokers are working in the market as partners in coalition and $X_1, X_2, ..., X_n$ are the suggested prices of these brokers for a stock share put on the market, respectively. Denote by Y the max price suggested by non-coalition brokers. Obviously, $Z = \max(X_1, X_2, ..., X_n, Y)$ is the traded price in the market. Then $E(Z \mid X_i)$ is the information trade price given X_i and $E(Z \mid Y)$ is the information trade price per-share given by non-coalition brokers. Since,

$$\frac{1}{n}\sum_{i=1}^{n} E(Z \mid X_i)$$

is the information trade price per-share average of all brokers in coalition by Theorem 1, we have

$$E\left(Z - \frac{1}{n}\sum_{i=1}^{n} E(Z \mid X_i)\right)^2 \le E(Z - E(Z \mid X_i))^2,$$

$$i = 1, 2, ..., n.$$

i.e. the predictor for a price of the share given by any of the member in the coalition is worse than the average value of predictors of all brokers in the coalition. This means that if a group of brokers is working in coalition they will have better gain than if they work in solitary.

3 Copula and coovariance

Since $E(X \mid Y)$ is the best predictor for X as a function of Y, and $E(Y \mid X)$ is the best predictor of Y as a function of X, it would be interesting to investigate how the dependence structure will change if we replaced X with $E(X \mid Y)$ and Y with $E(Y \mid X)$. For this purpose consider the joint distribution of the random variables $Z_1 \equiv \varphi(Y) = E(X \mid Y)$ and $Z_2 \equiv \psi(X) = E(Y \mid X)$. We are interested in copula of Z_1 and Z_2 . Let $\varphi^{-1}(y) = \inf\{x : \varphi(x) \leq y\}$ and $\psi^{-1}(x) = \inf\{y : \psi(y) \leq x\}$ are the generalized inverses of φ and ψ . Consider the joint distribution function of Z_1 and Z_2 . Let $F_X(x)$ and $F_Y(x)$ be a distribution function of X and Y, respectively. Assuming X and Y have the same support, denote left and right endpoints of the support of X and Y by $a = \inf\{x : F_X(x) > 0\}$ and $b = \sup\{x : F_X(x) < 1\}$, respectively. We allow also the cases $a = -\infty$ and $b = \infty$. We have

$$F_{Z_1,Z_2}(z_1, z_2) = P\{Z_1 \le z_1, Z_2 \le z_2\}$$

= $P\{\varphi(Y) \le z_1, \psi(X) \le z_2\}$
= $P\{Y \le \varphi^{-1}(z_1), X \le \psi^{-1}(z_2)\} = P\{X \le \psi^{-1}(z_2), Y \le \varphi^{-1}(z_1)\}$
= $F_{X,Y}(\psi^{-1}(z_2), \varphi^{-1}(z_1)), (z_1, z_2) \in [a, b]^2$

Hereafter, we assume that $\psi^{-1}(\infty) = \infty$ and $\varphi^{-1}(\infty) = \infty$ and $\psi^{-1}(-\infty) = -\infty$ and $\varphi^{-1}(-\infty) = -\infty$. The marginal distributions are

$$F_{Z_1}(z_1) \equiv P\{Z_1 \le z_1\} = \lim_{z_2 \to \infty} P\{X \le \psi^{-1}(z_2), Y \le \varphi^{-1}(z_1)\} = F_Y(\varphi^{-1}(z_1))$$
$$F_{Z_2}(z_2) \equiv P\{Z_2 \le z_2\} = \lim_{z_1 \to \infty} P\{X \le \psi^{-1}(z_2), Y \le \varphi^{-1}(z_1)\} = F_X(\psi^{-1}(z_2)).$$

Now we are interested in the copula of random vector (Z_1, Z_2) . Denote

$$F_X^{-1}(x) = \inf\{y : F_X(y) \ge x\} \text{ and } F_Y^{-1}(y) = \inf\{x : F_Y(x) \ge y\}.$$

Now, consider

$$F_{Z_1,Z_2}(z_1,z_2) = C_{Z_1,Z_2}(F_{Z_1}(z_1),F_{Z_2}(z_2)),$$
(1)

where $C_{Z_1,Z_2}(t,s)$ is a connecting copula of Z_1 and Z_2 .

Using probability integral transformation

$$F_{Z_1}(z_1) = t \iff F_Y(\varphi^{-1}(z_1)) = t \iff z_1 = \varphi(F_Y^{-1}(t)), \ F_{Z_1}^{-1}(t) = \varphi(F_Y^{-1}(t))$$
$$F_{Z_2}(z_2) = s \iff F_X(\psi^{-1}(z_2)) = s \iff z_2 = \psi(F_X^{-1}(s)), \ F_{Z_2}^{-1}(s) = \psi(F_X^{-1}(s))$$
(2)

we obtain from (1) and (2)

$$C_{Z_1,Z_2}(t,s) = F_{Z_1,Z_2}(F_{Z_1}^{-1}(t), F_{Z_2}^{-1}(s)) = F_{X,Y}(\psi^{-1}(z_2), \varphi^{-1}(z_1))$$
$$= F_{X,Y}(\psi^{-1}(\psi(F_X^{-1}(s)), \varphi^{-1}(\varphi(F_Y^{-1}(t)))$$
$$= F_{X,Y}(F_X^{-1}(s), F_Y^{-1}(t)) = C(s,t)$$

Therefore, we can formulate the following theorem.

Theorem 3 Let the joint distribution function of random variables X and Y be $F_{X,Y}(x,y) = C(F_X(x), F_Y(y)), (x,y) \in [a,b]^2$, where $C(u,v), (u,v) \in I^2 \equiv [0,1]^2$ is a connected copula. Consider functions $\varphi(y) = E(X \mid Y = y), y \in \mathbb{R}$ and $\psi(x) = E(Y \mid X = x), x \in \mathbb{R}$ and random variables $Z_1 \equiv \varphi(Y) = E(X \mid Y)$ and $Z_2 \equiv \psi(X) = E(Y \mid X)$. Assume that $\lim_{t\to\infty} \psi^{-1}(t) = \infty$ and $\lim_{s\to\infty} \varphi^{-1}(s) = \infty$. Then the copula of Z_1 and Z_2 is $C_{Z_1,Z_2}(t,s) = C(s,t), 0 \leq t, s \leq 1$. Therefore, if X and Y are exchangeable then $C_{Z_1,Z_2}(t,s) = C(t,s)$.

Example 2 ()Let (X; Y) be a bivariate normal random vector with joint pdf

$$f(x,y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \exp\{-\frac{1}{2(1-\rho^2)}((\frac{x-\mu_1}{\sigma_1})^2 - 2\rho(\frac{x-\mu_1}{\sigma_1})(\frac{x-\mu_2}{\sigma_2}) + (\frac{y-\mu_2}{\sigma_2})^2\}.$$

Then

$$\psi(x) = E(Y \mid X = x) = \mu_2 + \rho \frac{\sigma_2}{\sigma_1} (x - \mu_1)$$
(3)

$$\varphi(y) = E(X \mid Y = y) = \mu_1 + \rho \frac{\sigma_1}{\sigma_2} (y - \mu_2)$$
(4)

and

$$\psi^{-1}(t) = \frac{t - \mu_2}{\rho \sigma_2} \sigma_1 + \mu_1 \tag{5}$$

$$\varphi^{-1}(s) = \frac{s - \mu_1}{\rho \sigma_1} \sigma_2 + \mu_2 \tag{6}$$

and

$$\lim_{t \to \infty} \psi^{-1}(t) = \infty$$
$$\lim_{s \to \infty} \varphi^{-1}(s) = \infty.$$

Therefore, $C_1(x, y) = C(y, x)$, where C is a copula of (X, Y) and C_1 is a copula of $(\psi(Y), \varphi(X)) = (E(X | Y), E(Y | X))$.

The following well-known property of the conditional expectation state that given Y the conditional expectation of XY is

$$E(XY) = E(E(XY \mid Y)) = E(YE(X \mid Y))$$
$$E(XY) = E(E(XY \mid X)) = E(XE(Y \mid X))$$

Proposition 1

$$Cov(E(X \mid Y), Y) = Cov(E(Y \mid X), X) = Cov(X, Y).$$

Proof. Since

$$E(XY) = E[E(XY \mid Y)] = E[YE(X \mid Y)]$$

and

$$E(Y)E[E(X \mid Y)] = E(Y)E(X)$$

then,

$$Cov(X,Y) = E(XY) - E(X)E(Y) = E(YE(X \mid Y)) - E(Y)E(E(X \mid Y))$$
$$= Cov(Y, E(X \mid Y)) = Cov(E(X \mid Y), Y).$$

Remark 1 It can be observed that Cov(E(X | Y), E(Y | X)) may not be equal to Cov(X, Y).

Indeed,

$$E\psi(Y)\varphi(X) - E\psi(Y)E\varphi(X)$$

= $E\psi(Y)\varphi(X) - E[E(X \mid Y)]E[E(Y \mid X)]$
= $E\psi(Y)\varphi(X) - E(X)E(Y)$
= $E[E(X \mid Y)(E(X \mid Y)] - E(X)E(Y).$

Let for example $\psi(Y) = aY + b$, $\varphi(X) = cX + d$, where a, b, c, d > 0 (see for example (3) and (4)). Then $E\psi(Y)\varphi(X) = acE(XY) + adEY + bcEX + bd$. Therefore, $E\psi(Y)\varphi(X) = acE(XY) + adEY + bcEX + bd = EXY$ only if a = 1, c = 1, b = 0, d = 0. For (3) and (4) this means that it must be $\mu_1 = \mu_2 = 0, \sigma_1 = \sigma_2 = 1, \rho = 1$.

4 Sequences of predicted random variables

Let $X_1, X_2, ..., X_n, ...$ be a sequence of dependent random variables. Let $Y_1 = X_1, Y_2 = E(X_2 \mid X_1), ..., Y_n = E(X_n \mid X_{n-1}), ...$ It is clear that $EY_i = E(E(X_i \mid X_{i-1})) = EX_i, i = 12, ...$ Since $E(Y_1Y_2) = E(X_1 \mid X_2)$.

 $E(X_1E(X_2 | X_1)) = E(E(X_1X_2 | X_1)) = EX_1X_2$, then $Cov(Y_1, Y_2) = Cov(X_1, X_2)$. Furthermore, denoting by $\psi_i(x) = E(X_i | X_{i-1}), i = 2, 3, ...,$ we have

It is well known that the best predictor for X_{n+1} expressed as a function of $X_1, X_2, ..., X_n$ is $E(X_{n+1} | X_1, X_2, ..., X_n) = \Psi(X_1, X_2, ..., X_n)$, i.e.

$$\min_{g} E(X_{n+1} - g(X_1, X_2, ..., X_n))^2 = E(X_{n+1} - E(X_{n+1} \mid X_1, X_2, ..., X_n))^2$$

Theorem 4 Let X, Y and Z be any random variables defined on probability space $\{\Omega, F, P\}$. Then

$$E[X - E(X \mid Y, Z)]^{2} \le \min\left\{E[X - E(X \mid Y)]^{2}, E[X - E(X \mid Z)]^{2}\right\}$$
(7)

Proof. Consider

$$E [(X - E(X | Y, Z))^2 | Y, Z]$$

= $E[(X - E(X | Y) + E(X | Y) - E(X | Y, Z))^2 | Y, Z]$

$$= E[\{X - E(X | Y)\}^{2} | Y, Z] + 2E[\{X - E(X | Y)\}\{E(X | Y) - E(X | Y, Z)\} | Y, Z] + E[\{E(X | Y) - E(X | Y, Z)\}^{2} | Y, Z] = E[\{X - E(X | Y)\}^{2} | Y, Z] + 2[E(X | Y) - E(X | Y, Z)]E[\{X - E(X | Y)\} | Y, Z] + E[\{E(X | Y) - E(X | Y, Z)\}^{2} | Y, Z].$$
(8)

In (8) we take into account the fact that $h(Y,Z) \equiv E(X \mid Y) - E(X \mid Y,Z)$ is Y,Z measurable and behaviors as a constant in conditional expectation with respect to Y,Z and, therefore

$$E [\{ X - E(X \mid Y) \} \{ E(X \mid Y) - E(X \mid Y, Z) \} \mid Y, Z]$$

= $E [\{ X - E(X \mid Y) \} h(Y, Z) \mid Y, Z]$
= $h(Y, Z) E [\{ X - E(X \mid Y) \} h(Y, Z) \mid Y, Z]$
= $[E(X \mid Y) - E(X \mid Y, Z)] E [X - E(X \mid Y) \} \mid Y, Z].$

Since

$$E(E[\{X - E(X | Y)\} | Y, Z] = E(E(X | Y, Z)) - E(E(X | Y) | Y, Z) = EX - E(E(X | Y)) = EX - EX = 0.$$
(9)

Therefore,

$$E[X - E(X | Y, Z)]^{2}$$

= $E[\{X - E(X | Y)\}^{2} | Y, Z]$
+ $E[\{E(X | Y) - E(X | Y, Z)\}^{2} | Y, Z]$ (10)

Applying the operation E to both sides of (10) we obtain

$$E[X - E(X | Y, Z)]^{2}$$

= $E[\{X - E(X | Y)\}^{2}]$
+ $[E(X | Y) - E(X | Y, Z)]^{2}$

which imply

$$E[X - E(X \mid Y, Z)]^2$$

 $\leq E[\{X - E(X \mid Y)\}^2].$

_	_	_	_	

Corollary 1 For any $n \ge 2$, and the sequence of random variables $X_1, X_2, ..., X_n, ...$ it is true that

$$E[X_{n+1} - E(X_{n+1} \mid X_{i_1}, X_{i_2}, \dots, X_{i_l})]^2 \le E[X_{n+1} - E(X_{n+1} \mid X_{i_1}, X_{i_2}, \dots, X_{i_k})]^2,$$

$$1 \le i_1 < i_2 < \dots < i_k < i_l \le n, 1 \le k < l \le n.$$

For example,

$$E [X_{n+1} - E(X_{n+1} | X_1, X_2, ..., X_n)]^2 \le E [X_{n+1} - E(X_{n+1} | X_1, X_2, ..., X_{n-1})]^2$$

$$E [X_{n+1} - E(X_{n+1} | X_1, X_2, ..., X_n)]^2 \le E [X_{n+1} - E(X_{n+1} | X_1, X_2, ..., X_{n-2})]^2$$

$$E [X_{n+1} - E(X_{n+1} | X_1, X_2, ..., X_n)]^2 \le E [X_{n+1} - E(X_{n+1} | X_2, X_3, ..., X_n)]^2$$

$$E [X_{n+1} - E(X_{n+1} | X_2, X_3, X_4]^2 \le E [X_{n+1} - E(X_{n+1} | X_2, X_3)]^2$$

etc.

Example 3 (Martingale) The sequence $X_1, X_2, ..., X_n, ...$ is called a martingale if

$$E(X_{n+1} \mid X_1, X_2, ..., X_n) = X_n.$$

It follows from the corollary that if $X_1, X_2, ..., X_n, ...$ is a martingale then

$$E[X_{n+1} - X_n]^2$$

= $E[X_{n+1} - E(X_{n+1} | X_1, X_2, ..., X_n)]^2$
 $\leq E[X_{n+1} - E(X_{n+1} | X_{i_1}, X_{i_2}, ..., X_{i_k})]^2$
 $1 \leq i_1 < i_2 < ... < i_k \leq n, 1 \leq k \leq n.$

Example 4 (Markov chain) Let $X_1, X_2, ..., X_n, ...$ be a Markov chain, i.e. for an interval A of the real line it is true that

$$P\{X_n \in A \mid X_{i_1}, X_{i_2}, ..., X_{i_k}\} = P\{X_n \in A \mid X_{i_k}\}$$
$$E[X_n - E(X_n \mid X_{i_l})]^2 = E[X_n - E(X_n \mid X_{i_1}, X_{i_2}, ..., X_{i_l})]^2$$
$$\leq E[X_n - E(X_n \mid X_{i_1}, X_{i_2}, ..., X_{i_k})]^2$$
$$= E[X_n - E(X_{n+1} \mid X_{i_k})]^2,$$
$$1 \leq i_1 < i_2 < ... < i_k < i_l \leq n, 1 \leq k < l \leq n$$

Therefore,,

$$E [X_{n+1} - E(X_{n+1} \mid X_l)]^2 \le E [X_{n+1} - E(X_{n+1} \mid X_k)]^2,$$

$$1 \le k < l \le n.$$

A good illustration of this fact can be given with order statistics. Let $X_{1:n} \leq X_{2:n} \leq \cdots \leq X_{n:n}$ be the order statistics of X_1, X_2, \dots, X_n . Then for $1 \leq k < l \leq n$ one can write

$$E[X_{n:n} - E(X_{n:n} \mid X_{l:n})]^2 \le E[X_{n:n} - E(X_{n:n} \mid X_{k:n})]^2,$$
(11)

i.e. $E(X_{n:n} | X_{l:n})$ predicts $X_{n:n}$ better than $E(X_{n:n} | X_{k:n})$.

Example 5 (Order statistics) Let $X_1, X_2, ..., X_n$ be iid random variables and $X_{1:n} \leq X_{2:n} \leq \cdots \leq X_{n:n}$ be the order statistics. It is well known that the order statistics form a Markov chain. Therefore, $E(X_{n:n} | X_{1:n}, ..., X_{n-1:n}, X_{n:n}) = E(X_{n:n} | X_{n-1:n})$. By Markov property,

$$E [X_{n:n} - E(X_{n:n} | X_{1:n}, X_{2:n}, ..., X_{n-1:n})]^2$$

= $E [X_{n:n} - E(X_{n:n} | X_{n-1:n})]^2.$

from the corollary and (11) we have

$$E[X_{n:n} - E(X_{n:n} \mid X_{n-1:n})]^2 \le E[X_{n:n} - E(X_{n:n} \mid X_{n-2:n})]^2.$$

It means that $E(X_{n:n} | X_{n-1:n})$ is better estimation for $X_{n:n}$ than $E(X_{n:n} | X_{n-2:n})$. Let us compute the functions $g_1(x) = E(X_{n:n} | X_{n-1:n} = x)$ and $g_2(x) = E(X_{n:n} | X_{n-2:n} = x)$. From the joint distribution of $X_{r:n}$ and $X_{s:n}$, r < s, we can easily write the conditional pdf's of $X_{n:n} | X_{n-1:n}$ and $X_{n:n} | X_{n-2:n}$ as

$$f_{n|n-1}(z \mid x) = \frac{f(z)}{1 - F(x)}, x < z$$

$$f_{n|n-2}(z \mid x) = \frac{2(F(z) - F(x))}{(1 - F(x))^2} f(z), x < z,$$

respectively. Then for a uniform(0,1) distribution, we can write

$$g_1(x) = E(X_{n:n} \mid X_{n-1:n} = x) = \int_x^1 f_{n|n-1}(z \mid x)dz$$
$$= \frac{1+x}{2}, 0 \le x \le 1$$

and

$$g_2(x) = E(X_{n:n} \mid X_{n-2:n} = x) = 2 \int_x^1 f_{n|n-2}(z \mid x) dz$$
$$= \frac{x+2}{3}, 0 \le x \le 1$$

and it is clear that

$$g_1(x) < g_2(x), 0 \le x \le 1$$

because

$$\frac{x+2}{3} = \frac{1+x}{2} + \frac{1-x}{6}, 0 \le x \le 1.$$

This means that $g_1(X_{n-1:n}) = E(X_{n:n} | X_{n-1:n}) > g_2(X_{n-2:n}) = E(X_{n:n} | X_{n-2:n})$, hence $E(X_{n:n} | X_{n-1:n})$ is better than $E(X_{n:n} | X_{n-2:n})$ as a predictor of $X_{n:n}$. For an exponential distribution $F(x) = C(X_{n:n} | X_{n-2:n})$

 $1 - \exp(-x), x \ge 0$ it can be easily verify that

$$g_{1}(x) = E(X_{n:n} \mid X_{n-1:n} = x)$$

$$= \frac{1}{1 - F(x)} \int_{x}^{\infty} z f_{n|n-1}(z \mid x) dz$$

$$= \frac{1}{1 - F(x)} \int_{x}^{\infty} z f(z) dz$$

$$= e^{x} \int_{x}^{\infty} z e^{-z} dz = e^{x} e^{-x} (x+1) = x + x$$

i.e. $g_1(x) > g_2(x), x \ge 0$ and again $E(X_{n:n} | X_{n-1:n})$ is better than $E(X_{n:n} | X_{n-2:n})$ as a predictor of $X_{n:n}$.

Example 6 (Record Values) Let $X_1, X_2, ..., X_n, ...$ be a sequence of independent identically distributed (i.i.d.) r.v.'s with continuous d.f. F; $X_{1:n} \leq X_{2:n} \leq ... \leq X_{n:n}$ be the order statistics of $X_1, X_2, ..., X_n$. The random variable X_K is called a (upper) record value of the sequence $\{X_n, n \geq 1\}$ if $X_K > \max\{X_1, X_2, ..., X_{K-1}\}$. By convention X_1 is record value. Denote by $\{U(n), n > 1\}$ the sequence of record times:

1

$$U(n) = \min\{j : j > U(n-1), X_j > X_{U(n-1)}\}, n > 1 \quad with \ U(1) = 1.$$

 $X_{U(n)}$ is called n th upper record value. Developments on records have been reviewed by many authors including Nevzorov (1988), Nagaraja (1988), Arnold and Balakrishnan (1989), Arnold, Balakrishnan, Nagaraja (1992), Ahsanullah (1995). The properties of records values of iid random variables have been extensively studied in the literature. Many properties of records can be expressed in terms of the functions $R(x) = -\log \bar{F}(x)$ where $\bar{F}(x) = 1 - F(x)$ and $0 < \bar{F}(x) < 1$. It is well known that, the sequence of record values $X_{U(1)}, X_{U(2)}, ..., X_{U(n)}, ...$ form a Markov chain. From the corollary and (11) we have

$$E \left[X_{U(n)} - E(X_{U(n)} \mid X_{U(1)}, X_{U(2)}, ..., X_{U(n-1)}) \right]^{2}$$

$$\leq E \left[X_{U(n)} - E(X_{U(n)} \mid X_{U(1)}, X_{U(2)}, ..., X_{U(n-2)}) \right]^{2}$$
(12)

By Markov property

$$E\left(X_{U(n)} \mid X_{U(1)}, X_{U(2)}, ..., X_{U(n-1)}\right) = E\left(X_{U(n)} \mid X_{U(n-1)}\right)$$
$$E\left(X_{U(n)} \mid X_{U(1)}, X_{U(2)}, ..., X_{U(n-2)}\right) = E\left(X_{U(n)} \mid X_{U(n-2)}\right).$$
(13)

Then from (12) and (13) for any n > 2 we have

$$E\left[X_{U(n)} - E(X_{U(n)} \mid X_{U(n-1)})\right]^2 \le E\left[X_{U(n)} - E(X_{U(n)} \mid X_{U(n-2)})\right]^2,$$
(14)

i.e. $E(X_{U(n)} | X_{U(n-1)})$ is better than $E(X_{U(n)} | X_{U(n-2)})$ as a predictor of $X_{U(n)}$. It is clear that, (14) can be extended as

$$E \left[X_{U(n)} - E(X_{U(n)} \mid X_{U(l)}) \right]^2 \le E \left[X_{U(n)} - E(X_{U(n)} \mid X_{U(k)}) \right]^2,$$
(15)
$$2 < k < l < n.$$

It is possible to extend the list of examples to the areas where the prediction of random variables with conditional expectations is the subject.

Conclusion 1 This paper investigates the new properties of conditional expectation with respect to a sigma-algebra generated by other random variables. The conditional expectations of the random variable with respect to a sigma-algebra generated by the random variable is its best predictor in the sense of least square distance. Some important inequalities concerning the predictions of random variables are proved. These inequalities can find important applications in many areas such as financial mathematics, actuarial sciences, and reliability engineering. An application of the main inequality having interesting consequences in per-share stock is presented. Considering conditional expectations as random variables, we study also the dependence properties of and copulas of these random variables. Some examples with ordered random variables and martingales are provided.

References

- [1] Ahsanullah, M. (1995) Record Statistics, Nova Science publishers, Inc., Cormack, New York.
- [2] Arnold, B.C., Balakrishnan, N. and Nagaraja, H.N. (1992). A First Course in Order Statistics. Wiley, New York.
- [3] Arnold, B.C., Balakrishnan, N. and Nagaraja, H.N. (1998). Records. Wiley, New York.
- [4] Ahsanullah M and Nevzorov VB (2001) Ordered Random Variables. Nova Science Publishers, Huntington, NY.
- [5] Ahsanullah M and Nevzorov VB (2005) Order Statistics. Examples and Exercises. Nova Science Publishers, Huntington, NY.

- [6] Arnold BC and Balakrishnan N (1989) Relations, Bounds and Approximations for Order Statistics. Lecture Notes in Statistics, 53, Springer, New York.
- [7] Arnold BC, Becker A, Gather U and Zahedi H (1984) On the Markov property of order statistics. Journal of Statistical Planning and Inference 9: 147-154.
- [8] Borovkov, A.A. (2002) Mathematical Statistics. Gordon and Breach Science Publishers.
- [9] David, H.A. and Nagaraja, H.N. (2003). Order Statistics. Third ed., Wiley, New Jersey.
- [10] Karlin, S. and Taylor, H.M. (1975) A first course in Stochastic Processes. Second Edition. Academic Press.
- [11] Ross, S. (2002) A first course in Probability. Prentice Hall.