

Generated using the official AMS L^AT_EX template v6.1 two-column layout. This Work has been submitted for publication to Monthly Weather Review. This Work has not yet been peer-reviewed and is provided by the contributing Author(s) as a means to ensure timely dissemination of scholarly and technical Work on a noncommercial basis. Copyright and all rights therein are maintained by the Author(s) or by other copyright owners. Copyright in this Work may be transferred without further notice, and this version may no longer be accessible.

Supercell low-level mesocyclones: Origins of inflow and vorticity

BRICE E. COFFER,^a MATTHEW D. PARKER,^a JOHN M. PETERS,^b ANDREW R. WADE^c

^a *Department of Marine, Earth, and Atmospheric Sciences, North Carolina State University, Raleigh, NC*

^b *Department of Meteorology and Atmospheric Science, The Pennsylvania State University, University Park, PA*

^c *Cooperative Institute for Severe and High-Impact Weather Research and Operations, University of Oklahoma, Norman, OK*

ABSTRACT: The intensification of low-level mesocyclones in supercell thunderstorms has historically been associated with the development of storm-generated streamwise vorticity along a baroclinic gradient in the forward flank of supercells. However, the ambient streamwise vorticity of the environment (often quantified via storm-relative helicity), especially near the ground, is particularly skillful at discriminating between nontornadic and tornadic supercells. This study investigates whether the origins of the inflow air into supercell low-level mesocyclones, both horizontally and vertically, can help explain the dynamical role of environmental versus storm-generated vorticity in low-level mesocyclone intensification. Simulations of supercells, initialized with wind profiles common to supercell environments observed in nature, show that the air bound for the low-level mesocyclone primarily originates from the undisturbed, ambient environment, rather than from along the forward flank, and from very close to the ground, often in the lowest 200 - 400 m of the atmosphere. Given that the near-ground environmental air comprises the bulk of the inflow into low-level mesocyclones, this likely explains the forecast skill of environmental streamwise vorticity in the lowest few hundred meters of the atmosphere. Contrary to prior conceptual models of low-level mesocyclones, intensification does not appear to require the development of additional horizontal vorticity in the forward flank. Instead, the dominant contributor to vertical vorticity within the low-level mesocyclone is from the environmental horizontal vorticity. This study therefore supports a revised view of low-level mesocyclones in supercells.

SIGNIFICANCE STATEMENT: Supercell thunderstorms produce the majority of tornadoes, and a defining characteristic of supercells is their rotating updraft, known as the “mesocyclone”. When the mesocyclone is stronger at lower altitudes, the likelihood of tornadoes increases. The purpose of this study was to understand if the rotation of the mesocyclone in supercells is due to horizontal spin present in the ambient environment or whether additional horizontal spin generated by the storm itself primarily drives this rotation. Our results suggest that inflow air into supercells, and low-level mesocyclone rotation, is mainly due to the properties of the environmental inflow air, especially near the ground. This hopefully will clarify how our community views the development of low-level mesocyclones in supercells.

1. Introduction

A defining characteristic of supercell thunderstorms is their mesocyclone, a quasi-steady region of vertical vorticity within the storm’s updraft. This persistent feature contributes to a host of severe weather threats, including damaging nontornadic winds (Smith et al. 2012), flash flooding (Nielsen and Schumacher 2020), large individual hail stones (Blair et al. 2017) and/or large accumulations of small hail (Kumjian et al. 2019), and tornadoes (Markowski and Richardson 2009). Conceptual models of supercells have all prominently featured the mesocyclone (Brandes 1978; Lemon and Doswell 1979; Klemp 1987; Doswell

and Burgess 1993), and temporal increases in mesocyclone rotation near the cloud-base have long been identified as a precursor of tornadoes (e.g., Brandes 1993; Burgess et al. 1993). Despite these connections between mesocyclones and tornadoes, the presence of a mesocyclone alone is a poor predictor of supercellular tornadogenesis (Trapp 1999). Although most tornadoes are associated with supercells, perhaps less than 15% of mesocyclones are tornadic (Trapp et al. 2005; Smith et al. 2012).

The process of supercellular tornadogenesis is often described as having three steps (Davies-Jones 2015). First, an updraft needs to require rotation aloft (i.e., the development of a mesocyclone). It is well established that the updrafts of supercells initially acquire their rotation via the tilting of horizontal vorticity associated with the vertical shear of the environmental wind profile (e.g., Rotunno and Klemp 1982; Davies-Jones 1984; Weisman and Rotunno 2000). Specifically, the tilting of streamwise horizontal vorticity (i.e., the component of vorticity aligned parallel to the motion of air in the storm-relative framework) is a necessary requirement for the updraft to acquire net positive vertical vorticity (Davies-Jones 1984, 2017, 2022; Dahl 2017).

The second step in tornadogenesis involves rotation developing at the ground. This is thought to occur primarily through some combination of baroclinic (e.g., Davies-Jones 1982; Davies-Jones and Brooks 1993; Walko 1993; Adlerman et al. 1999; Dahl et al. 2014; Markowski and Richardson 2014; Dahl 2015; Parker and Dahl 2015) and frictional (e.g., Schenkman et al. 2014; Markowski 2016;

Corresponding author: Brice E. Coffey, becoffer@ncsu.edu

Mashiko 2016; Roberts et al. 2016; Fischer and Dahl 2022) generation of horizontal vorticity within downdrafts that is subsequently tilted into the vertical very close to the surface (Rotunno et al. 2017).

Both nontornadic and tornadic supercells readily acquire sub-tornadic rotation near the surface (Parker and Dahl 2015; Coffey et al. 2017). Hence, steps 1 and 2 are necessary, but not sufficient, for tornadogenesis. The third and final step in tornadogenesis is the contraction and intensification of rotation at the ground into a tornadic strength vortex. For this to occur, several conditions apparently need to be met simultaneously. The sub-tornadic rotation needs to be within outflow air that has sufficiently small negative buoyancy so that it does not resist upward acceleration into the low-level updraft (Markowski et al. 2002). The surface rotation also must experience a persistent area of convergence and stretching. Because this is necessarily below the height of the LFC (where air is either neutrally or negatively buoyant), the bulk of the associated vertical accelerations must be provided by the mesocyclone and its associated dynamic lifting (Rotunno and Klemp 1982; Markowski et al. 2012b; Markowski and Richardson 2014; Coffey and Parker 2015, 2017; Goldacker and Parker 2021)¹. This dynamic lifting owes its existence to pressure falls aloft associated with the mesocyclone's circulation. Many studies use the 0 - 1 km vertical perturbation pressure gradient acceleration as a measure of this dynamic lifting provided by the mesocyclone.

Within this chain of processes, the point at which a supercell ultimately succeeds or fails in producing a tornado is strongly linked to the mesocyclone. Hence, low-level mesocyclone intensity appears to exert a substantial influence on tornadogenesis likelihood (e.g., Markowski and Richardson 2014; Coffey and Parker 2017; Peters et al. 2022a). We will refer to the mesocyclone near cloud-base as the “low-level” mesocyclone [in the lower troposphere at approximately 1 km above ground level (AGL)] and consider this level distinct from both the “mid-level” mesocyclone further aloft (in the mid-troposphere between ~3-6 km AGL) and the “near-ground” rotation that develops much closer to the surface (< 250 m AGL and is sometimes referred to as the “tornado cyclone”) based on the dynamical processes that potentially govern the development of rotation at these separate altitudes.

The current thinking regarding the intensification of the mesocyclone at low-levels largely originates from the seminal modeling work of Rotunno and Klemp (1985). Integrating material circuits backward from an area of low-level rotation showed that “upward tilting of baroclinically-generated horizontal vorticity along the cool air boundary situated upstream of the low-level updraft”. The

orientation of the forward-flank baroclinic zone and the storm-relative flow in this region of the storm is such that baroclinically-generated vorticity is predominantly streamwise (Klemp and Rotunno 1983). In simulations with and without rain, rotation in the mid-levels developed consistently through the tilting of environmental horizontal vorticity. On the other hand, low-level rotation was absent in simulations without rain. Many modeling studies since have reinforced the importance of storm-generated streamwise horizontal vorticity along the forward flank of supercells to the development of the low-level mesocyclone (e.g., Wicker and Wilhelmson 1995; Adlerman et al. 1999; Beck and Weiss 2013; Naylor and Gilmore 2014; Marquis et al. 2016). Observations of low-level mesocyclones also seem to point to storm-generated processes aiding in their development. This includes visual cues in low-level cloud features, such as a localized lowering of the wall cloud beneath the main mesocyclone area (often tilted towards the region of forward flank precipitation; Atkins et al. 2014). Additionally, vortex lines around mesocyclones from Doppler radar studies are configured into ‘vortex line arches’, which are highly suggestive of upward tilting of baroclinically generated horizontal vorticity within the low-level mesocyclone (e.g., Straka et al. 2007; Markowski et al. 2008, 2012a), as long as the temperature deficit within a storm's forward flank outflow is not excessively negative (Shabbott and Markowski 2006). As summarized by Markowski and Richardson (2010), our current understanding of low-level mesocyclones assumes the “tilting of baroclinically enhanced low-level horizontal vorticity produces more significant vertical vorticity at low altitudes than does the tilting of environmental vorticity alone” and “formation of low-level mesocyclones usually awaits the development of an extensive forward-flank precipitation region”.

This concept has recently returned to the forefront via a growing interest in the “streamwise vorticity current” (SVC; Orf et al. 2017), a localized region of horizontal streamwise vorticity located parallel to the forward flank outflow boundary. In a very high-resolution simulation of an EF5 tornadic supercell, Orf et al. (2017) stated that the SVC was ingested into the storm's low-level mesocyclone, intensified the updraft, and led to the amplification and maintenance of a long-lived violent tornado. While the idea of storm-generated, streamwise horizontal vorticity production within the forward flank has previously been discussed in the literature (as reviewed above), the unprecedented level of detail in these simulations has inspired subsequent exploration in a number of modeling studies (Markowski 2020; Schueth et al. 2021) and observed cases (Murdzek et al. 2020a,b; Schueth et al. 2021), and has been an explicit emphasis in recent field projects (Weiss et al. 2020). Even so, it appears that not all supercells have SVCs, and the presence (or lack) of an SVC is not a

¹In low LCL environments common to tornadic supercells, buoyant vertical pressure perturbation gradients can also have a slight upward contribution to the total acceleration field (Brown and Nowotarski 2019, see their Fig. 14).

necessary requirement for tornadogenesis success (or failure; Murdzek et al. 2020a). Even when an SVC is present, it is not guaranteed that the augmented, storm-generated streamwise horizontal vorticity within this feature ends up participating in the low-level mesocyclone (Murdzek et al. 2020b).

Notwithstanding the seemingly well-established connection between *storm-generated* horizontal vorticity and low-level mesocyclones, there is accumulating evidence that tornadic environments are distinguished by large *environmental* streamwise vorticity in the lowest 500 - 1000 m AGL (Markowski et al. 2003; Rasmussen 2003; Miller 2006; Esterheld and Giuliano 2008; Nowotarski and Jensen 2013; Parker 2014; Coffey et al. 2019, 2020). Markowski et al. (2012b) asked whether “large environmental vorticity [is] important, especially at low-levels, because its tilting establishes the base of the mid-level mesocyclone at fairly low elevations?”

The vertically integrated storm-relative flux of streamwise vorticity into an updraft is represented by the storm-relative helicity (SRH; Davies-Jones 1984). Because SRH represents the flux of streamwise vorticity into an updraft, an arbitrary layer of inflow is necessarily defined. The choice in the depth of this layer is non-trivial because the properties of inflow air into supercells strongly influences the potential mesocyclone intensity (Peters et al. 2022a). Some studies have attempted to define the layer of inflow based on thermodynamic properties such that only parcels associated with CAPE and minimal CIN are included (the effective inflow layer or EIL; Thompson et al. 2007; Nowotarski et al. 2020). However, model-based proximity soundings show that SRH very near the ground (e.g., 0 - 500 m AGL; SRH500) is the single most predictive parameter in discriminating significantly tornadic supercells from their nontornadic counterparts, in both United States and European severe weather environments (Coffey et al. 2019, 2020). In a pair of simulated supercells (one tornadic, one nontornadic), Coffey and Parker (2017) showed that the environmental inflow parcels bound for the low-level mesocyclone originated *exclusively below 300 m AGL*. The sources of air bound for the low-level mesocyclone across a range of supercells in various environments has not yet been systematically evaluated.

The uncertainty about the importance of environmental versus storm-generated vorticity in low-level mesocyclone intensity leads us to our main research questions:

1. If low-level mesocyclone-genesis intensification is primarily due to storm-generated, forward flank baroclinic generation of horizontal vorticity, then why is near-ground environmental streamwise vorticity such a highly skillful forecast parameter?
2. What role do these competing sources of streamwise vorticity play in modulating the intensity of low-level

dynamic lifting that ultimately can determine whether a supercell fails or succeeds at producing a tornado?

To address these questions, we explore the origins and properties of air parcels that end up in the low-level mesocyclone using a matrix of simulations initialized with wind profiles common to supercell environments observed in nature and representing a spectrum of near-ground horizontal vorticity magnitudes and orientations.

2. Methods

a. CM1 model

Supercell simulations were performed using Cloud Model 1 (CM1; Bryan and Fritsch 2002) release 20.3. Storms were simulated for 3 h on a $200 \times 200 \times 18 \text{ km}^3$ domain with a horizontally homogeneous environment (described in Section 2b). The inner $100 \times 100 \text{ km}^3$ had a horizontal grid-spacing of 80 m during the period of analysis, while the vertical grid-spacing was stretched from 20 m in the lowest 300 m to 280 m at 12 km. The domain was translated with a constant storm-motion, which was determined iteratively by trial and error to keep the storm approximately centered within the domain. A fifth-order advection scheme, utilizing high-order-weighted essentially nonoscillatory finite differencing, was used with no additional artificial diffusion (Wicker and Skamarock 2002). The subgrid-scale turbulence was parameterized by a 1.5-order turbulence kinetic energy closure scheme similar to Deardorff (1980), with separate horizontal and vertical turbulence coefficients. Open-radiative lateral boundary conditions were employed, and where there was an inward mass flux into the domain, the horizontal winds were nudged back towards the base-state fields along the lateral boundaries (i.e., *nudge_{obc}* = 1 in the namelist). The upper-boundary had a rigid, free-slip boundary condition, with a Rayleigh damping sponge applied above 14 km, while the bottom boundary condition was semi-slip with a constant surface drag coefficient (C_d) of 0.0035 to partially capture frictional effects on within-storm processes. The C_d was calculated using the surface layer scheme by Jiménez et al. (2012) based on the thermodynamic profile and the mean of the kinematic profiles discussed in Section 2b. Random perturbations of 0.25 K were added to the initial conditions within the lowest 1000 m AGL. The simulations use a six-category, fully double-moment bulk microphysics scheme from the National Severe Storms Laboratory (NSSL) that explicitly predicts the variable densities of hail and graupel (Ziegler 1985; Mansell 2010; Mansell et al. 2010). Convection is initialized using the heat flux method of Carpenter et al. (1998) and Lasher-Trapp et al. (2021) and is described in more detail below.

1) HEAT FLUX CONVECTIVE INITIALIZATION

We used a Gaussian heat flux based on Lasher-Trapp et al. (2021) and similar to Morrison et al. (2022), that results in a more natural transition from buoyant plumes to a sustained, mature supercell updraft across all the environments used herein (Peters et al. 2022b,c). The heat flux was strongest near the surface and exponentially decreased in magnitude radially and vertically. The horizontal width and height of the Gaussian function were 10000 and 2500 m, respectively. The heat flux linearly increased in time from 0 Wm^{-2} at the start of the simulation to 2000 Wm^{-2} over 1 min. This amplitude was maintained for 28 min, then linearly decreased back to zero over 1 min, for a total of 30 min of active heating. In addition to the heating, a very weak Gaussian-shaped forced convergence [$O(10^{-4})$] was applied to the horizontal wind field over the same spatial and temporal dimensions of the surface heating (similar to Moser and Lasher-Trapp 2018). The combined effect of surface heating and weak convergence resulted in initially shallow thermal-like updrafts that gradually transitioned into a sustained steady supercell updraft (Peters et al. 2022b,c). While the convection organically developed during the first hour of the simulation, the horizontal grid-spacing of the simulation within the inner $100 \times 100 \text{ km}^2$ was 250 m, which Lasher-Trapp et al. (2021) found suitable for simulating overturning circulations and entrainment. Afterwards, the horizontal grid-spacing within the inner domain was decreased to 80 m, as discussed next.

2) DOWNSCALING TO FINER GRID-SPACING

To adequately resolve features important to the development and maintenance of the low-level mesocyclone and tornadoes, the horizontal grid-spacing within the inner domain of the simulations was downscaled to 80 m an hour into the simulation using the technique of Coffey and Parker (2022). The vertical grid-spacing was unchanged. Downscaling was accomplished by fitting a unique gridded interpolation function (cubic spline) to each two- and three-dimensional array in the model’s restart file, essentially creating a one-way nest within CM1.

For comparison, the original 250 m simulations were also run for the full 3 h. Generally the supercells evolved qualitatively similarly between the original 250 m and downscaled 80 m simulations². No simulation that produced a tornado-like vortex at 250 m became nontornadic in the 80 m simulation (and vice-versa). The similarities at different resolutions may be somewhat surprising given the low degree of intrinsic predictability within supercells (Markowski 2020); however, it appears, based on these results, that most of the spread in predictability

occurs within the convection initiation and development phase (similar to predictability challenges in operational convective forecasting, e.g., Galarneau et al. 2022).

b. Kinematic and thermodynamic base-state environments

To best encompass a range of wind profiles with varying magnitudes of near-ground SRH observed in nature, self-organizing maps (SOMs) were used to identify recurring low-level wind profile patterns from 20,194 model-analyzed supercell proximity soundings previously analyzed by Coffey et al. (2019). SOMs have previously been used to identify recurring features in severe weather datasets (Nowotarski and Jensen 2013; Anderson-Frey et al. 2017; Nowotarski and Jones 2018; Warren et al. 2021; Hua and Anderson-Frey 2022). The SOMs were trained on the 0 - 500 m AGL ground-relative u,v wind components, similar to Goldacker and Parker (2021). The 0 - 500 m AGL layer was selected because it represented the most distinct layer between significantly tornadic and nontornadic supercells in Coffey et al. (2019). All soundings were interpolated to a common vertical grid with 50 m grid-spacing. Training was conducted with three nodes in both the x, y direction, using a learning rate of 0.1 and iterated over 10,000 training steps. Wind profiles with layers of anti-streamwise vorticity within the lowest 500 m AGL were excluded, leaving a total of 15,806 of cases used in the SOM training. The cases removed represent environments that are not conducive to low-level mesocyclone development and were predominately associated with nontornadic supercells. While SOMs have many uses, the intention herein was to distill a large dataset of environmental soundings into a handful of archetypal hodographs to initialize a reasonable number of supercell simulations for analysis.

The resulting nine wind profiles from the SOM vary the magnitude and orientation of the near-ground hodograph (Fig. 1), generally increasing the magnitude of 0 - 500 m AGL vertical wind shear vector from left to right and shifting the direction of the shear vector from southwesterly to southerly from top to bottom (Fig. 1). All profiles have at least 20 m s^{-1} of 0 - 6 km bulk vertical wind shear, supportive of supercells. SRH500 increases from $50 \text{ m}^2 \text{ s}^{-2}$ in node 1_0 to $275 \text{ m}^2 \text{ s}^{-2}$ in node 2_2. Each ground-relative wind profile from the SOM was subsequently shifted to the origin of the hodograph at the lowest model level (i.e., no surface wind). Shifting the wind profile minimizes the influence of the semi-slip bottom boundary condition on the wind profile over the course of the 3 h simulation without introducing unnatural, or invented, forces into the model’s equation set (Davies-Jones 2021). The wind profile was also run through a 1 h CM1 single-column simulation in order to let the profile adjust to the semi-slip bottom boundary condition. Differences were essentially

²While not relevant to the current work, the 80 m simulations tended to experience less rightward deviations in the storm motion (Fig. 1) and were less isolated than their 250 m counterparts, perhaps due to overly smooth and steady pressure perturbation fields at a lower resolution.




FIG. 1. Hodograph diagrams, generated via self-organizing maps, for the nine simulations. The black line is the shifted, steady wind profile used to initialize each simulation from 0 to 12 km, n is the number of samples, ST is the percentage of significantly tornadic cases, and NT is the percentage of nontornadic cases. The cyan line is the shifted original 0 – 500 m wind profile outputted directly from the SOM. The ellipses are centered at the surface (yellow), 500 m (green), 1 km (turquoise), 3 km (blue), and 6 km (purple), and the radii are equivalent to the standard deviation of the profiles in each node at those points. The Bunkers and actual storm motion are labeled “b” and “a”, respectively, and the SRH500 displayed uses the Bunkers storm motions.

nonexistent between the initial, adjusted, and final far-field wind profile after the 3 h simulation.

Each kinematic profile from the SOM was paired with the same thermodynamic profile (Fig. 2) in order to isolate the influence of the wind profile on the development of low-level mesocyclones. A modified version of the Weisman and Klemp (1982) sounding was used to generate potential temperature and relative humidity profiles. The modifications are the same as those described by Markowski (2020) and are designed to minimize the appearance of moist absolutely unstable layers and unwanted/uncontrolled convection initiation. This thermodynamic profile is similar to that used by Lasher-Trapp et al. (2021), whose surface heat flux initialization we also used. Mixed-layer CAPE is 3657 J kg^{-1} , CIN is -14 J kg^{-1} , the LCL is approximately at 1050 m AGL, the LFC is near 1500 m AGL, and the top of the EIL is ~ 2700 m AGL (Fig. 2). All parameters were calculated as-in Coffey et al. (2019). These bulk thermodynamic quantities represent the upper-end of the supercell spectrum, and are more similar to the subset of signifi-

cantly tornadic supercells than to nontornadic supercells in the Coffey et al. (2019) dataset.

c. Analysis techniques

1) MESOCYCLONE TRACKING

To objectively analyze each supercell, the low-level mesocyclone was tracked over time. The algorithm tracked the peak of a Gaussian, spatially smoothed product of vertical velocity and vertical vorticity (of at least 0.01 s^{-1}) at 1 km AGL. A temporally smoothed time series of x, y mesocyclone centroids was then computed using a 3rd order polynomial, Savitzky-Golay filter with an 11 point window (i.e., ± 5 minutes of output). This method reliably tracked the right-moving supercell of interest in each simulation bar one. Simulation 1_0, which had the smallest near-ground SRH and weakest storm-relative flow, experienced multiple splits and the right-moving storm began to dissipate two hours into the simulation. For this reason, a subjectively defined right moving updraft area was




FIG. 2. Skew- T log- p diagram of the modified WK82 profile used in each simulation. The environmental T (K) and dewpoint temperature T_d (K) are shown in thick red and thick green, respectively. The solid black line is the pseudoadiabat followed by the mixed-layer parcel, and the area between the temperature curve and the parcel bath (CAPE) is shaded in red. Various environmental thermodynamic parameters are also shown.

instead defined for the simulation 1_0 during a key period of interest (i.e., before the storm began to dissipate).

2) KEY TIME PERIODS TORNADO-GENESIS/FAILURE

In order to analyze supercells at similar points in their evolution, a key time period of tornadogenesis or tornadogenesis failure was determined for each simulation. Similar to the definitions of Coffey et al. (2017), vortices at 10 m AGL (i.e., the lowest bottom model level) were considered tornadoes if they met the following criteria: 1) vertical vorticity $\geq 0.3 \text{ s}^{-1}$ 2) a pressure drop $\leq -10 \text{ hPa}$ throughout the lowest 1 km, and 3) a ground-relative wind speed $\geq 29 \text{ m s}^{-1}$ (i.e., the EF-0 threshold) within 1 km of the position of maximized Okubo-Weiss (OW) parameter ($OW = \zeta^2 - \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x}\right)^2 - \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y}\right)^2$). All three criteria needed to be satisfied for at least two minutes. If these thresholds were not met at any time during a simulation, tornadogenesis failure was defined as the time of maximum

OW at 10 m AGL within a 10 km diameter of the tracked low-level mesocyclone centroid point.

3) TRACERS AND BACKWARD TRAJECTORIES

To visualize the source regions of the low-level mesocyclone, three layers of passive tracers were initialized in CM1 [0 - 500 m AGL, 500 - 1500 m AGL (the approximate height LFC), and 1500 - 2700 m AGL (the top of the EIL)]. The value of the tracer in each layer was initially set to 1. In addition to tracers, backward trajectories were used to determine source regions of the low-level mesocyclone with more precision than tracers can provide. The method of integrating backward trajectories loosely followed that of Gowan et al. (2021), except that between native output intervals (60 s), velocity fields were linearly interpolated into 3 s intervals to ensure that trajectories do not “skip” over entire grid cells during a single integration step. Backward trajectories were initialized within the low-level mesocyclone at 1 km AGL every 60 s between 5 and

10 minutes prior to the defined key time period of tornado-genesis/failure (t_{-10} to t_{-5}) and were tracked backwards for 30 mins. Given that the timescale for tornado formation is roughly 10 mins (Davies-Jones et al. 2001), this time window focuses on the ramp-up of the low-level mesocyclone that proceeds tornado-genesis/failure, not at t_0 when a tornado has potentially already formed. This eludes the potentially problematic issue of determining the exact moment surface rotation should be considered “tornadic” (e.g., Houser et al. 2022). Backward trajectories began at grid points that met or exceeded the 90th percentile of vertical velocity (Table 1) within a 10 km diameter of the low-level mesocyclone centroid, provided that their initial vertical vorticity was at least 0.01 s^{-1} . In other words, the vertical velocity and vertical vorticity thresholds for backward trajectories isolate parcels that make up the most intense upward-moving, cyclonically rotating air in each low-level mesocyclone.

4) BAROTROPIC STENCILS

In order to address the relative contributions of environmental versus storm-generated horizontal vorticity upon the mesocyclone, we complement the tracer and backward trajectory analysis with forward trajectories within CM1. Using the “barotropic stencil” method from Dahl et al. (2014), the initial environmental (i.e., “barotropic”) vorticity of a parcel can be separated from the contribution of vorticity generated by the storm (i.e., “non-barotropic”). Following Dahl et al. (2014), six additional adjacent “stencil” parcels were initialized surrounding a center parcel at distances of 0.5 m. Over time, the stencils are deformed and stretched, and the embedded barotropic vorticity vector is reoriented accordingly. Thus, for parcels that subsequently enter the low-level mesocyclone, the component of the vertical vorticity that is due to the initial ambient horizontal vorticity can be derived based on the final configuration of the stencil. The non-barotropic component is simply the residual between the known vorticity at some final time and the rearranged barotropic vorticity component.

Since the initial parcel locations may not be completely undisturbed base-state air due to far-reaching storm influences on the environment (e.g., Parker 2014; Wade et al. 2018; Coniglio and Parker 2020), we extended the stencil method from Dahl et al. (2014) to parse out two components of barotropic vorticity. The initial stencil vorticity vectors include both the base-state values from model initialization (ω_{env}) and any perturbations that have developed between the model start time and the stencil initialization time (ω_{pert}). Therefore the final vertical vorticity (ζ) of the low-level mesocyclone (LLM)

$$\zeta_{LLM} = \zeta_{BT_{env}} + \zeta_{BT_{pert}} + \zeta_{NBT} \quad (1)$$

where BT is the component due to barotropic rearrangement of the initial stencil vorticity, divided into the base-state environment (env) and pre-existing perturbations ($pert$), and NBT is the residual non-barotropic (storm-generated) component. Because it is unknown whether or not ω_{pert} represents a prior process that is barotropic, baroclinic, or frictional, the three components of ζ_{LLM} will be reported separately. We hope that separating the base-state and perturbation vorticity vectors in this manner helps to address historical concerns that analyzed trajectories were potentially not fully removed from the storm’s influence at the time of initialization.

Forward trajectories were seeded within model restart files 40 minutes before the key time period of tornado-genesis/failure for each simulation and integrated forward natively within CM1 for 35 minutes. This results in parcels with at least 30 minutes of output history during the same five minute composite period (t_{-10} to t_{-5}) highlighted by the backward trajectories. Horizontally, parcels were launched upstream of the low-level mesocyclone within a unique horizontal bounding box for each simulation (shown later) encompassing an estimated 75% of the low-level mesocyclone inflow area based on the backward trajectories, with a 2 km buffer upstream in the $+x$, $+y$, and $-y$ directions. Vertically, parcels were defined over a 1500 m layer starting at 30 m AGL (i.e., the second lowest model level³), except for the 1_0 simulation where parcels were extended up to 2000 m due to higher parcel origin heights (shown later). The center stencil parcels were defined on an isotropic 100 m grid within the bounding box, with the six adjacent stencil parcels initialized surrounding each center parcel at distances of 0.5 m. Each simulation had between 2,000,000 and 12,000,000 total forward trajectories, depending on the areal extent of the inflow and thus the size of the bounding box. Parcel data were saved every 15 s and low-level mesocyclone trajectories were identified using the same thresholds as the backward trajectories, i.e., vertical velocity greater or equal to the 90th percentile (Table 1) and at least 0.01 s^{-1} of vertical vorticity (within $\pm 10 \text{ m}$ of 1 km AGL).

3. Results

All nine wind profiles from the SOM resulted in supercellular convection for the majority of the 3 hour simulation time. Eight of the nine simulations develop quasi-steady, right-moving supercells with persistent (and trackable) low-level mesocyclones of varying intensity (Fig. 3). The lone exception is simulation 1_0, which due to the lack of curvature and storm-relative flow in the hodograph (Fig. 1b), experiences a succession of splits. After two hours, only disorganized multi-cell convection exists

³The second lowest model level was the lowest chosen because surface drag always opposes the local wind field at the lowest model grid point with a “semi-slip” bottom-boundary condition, yielding questionable horizontal vorticity fields (i.e., Wang et al. 2020).




FIG. 3. Horizontal cross sections of 10 m AGL reflectivity (dBZ; shaded) at the key time period of tornado-genesis/failure (see Table 1) for each simulation and the 90th percentile of updraft at 1 km (black contour). Each panel is labeled nontornadic (NT), tornadic (T), or violently tornadic (VT).

throughout the domain in simulation 1_0 and peak vertical velocities drop off substantially compared to the other eight simulations (Fig. 4a). Despite the unsteady nature, at times, the southern-most, right-moving storm in simulation 1_0 periodically displays supercellular features in the reflectivity field, such as a hook echo⁴ (Fig. 3b). Due to the lack of persistent low-level mesocyclone, the 1_0 storm is probably of less interest to the supercell tornadogenesis problem; however, for completeness, we have not excluded any of the analyses for this simulation. In contrast

to 1_0, the other eight simulations experience relatively steady maximum vertical velocities of over 70 m s^{-1} from 1 hour onwards (when the downscaling occurred; Fig. 4), as the initial convection produced by the surface heat flux initialization coalesces into singular, dominant updraft.

By happenstance, the nine simulations can be evenly separated into three groups of three, nontornadic (1_0, 2_0, 0_1), tornadic (0_0, 1_1, 1_2), and violently tornadic (2_1, 0_2, 2_2). The tornadic simulations are found further to right of the SOM (Fig. 1) following trends in the increasing magnitude of SRH500. The three nontornadic simulations all had SRH500 less than $100 \text{ m}^2 \text{ s}^{-2}$, although this threshold did not preclude the 0_0 simulation from

⁴The most prominent of such instances is chosen as the key time period and the centroid of the low-level updraft (albeit weaker than any other simulation; Table 1) was manually defined over time.




FIG. 4. Time series’ of maximum domain w (blue, m s^{-1} , left y axis) and maximum surface ζ (red, s^{-1} , right y axis), with the pre-downscaling period hatched. The key time period for each simulation is shown as a vertical dashed line (note the key time period for nontornadic supercells is not necessarily at the time of peak vertical vorticity in the domain). Also shown as faded lines are the same time series’ from original 250 m simulations ($\zeta_{sf} c_{250m}$ is scaled to account for differences in magnitude due to lower resolution). Each panel is labeled nontornadic (NT), tornadic (T), or violently tornadic (VT). Panels faded yellow correspond to NT simulations.

becoming tornadic (indicating that maybe some amount of storm-generated augmentation was more prominently present in this simulation). Qualitatively, the nontornadic supercells display muted trends in surface vertical vorticity (Fig. 4b-d) and never meet the threshold of a deep, long-lasting vortex underneath the main low-level mesocyclone, whereas the tornadic simulations experience abrupt jumps in the maximum surface vertical vorticity (Fig. 4a,e-i). The six tornadic simulations can be further delineated by their max OW . Vortex intensity increases from $OW \sim 0.2$ in three of the simulations (0_0, 1_1, 1_2) to $OW > 1$ in the violently tornadic ones (2_1, 0_2, 2_2; Table 1). Some of the tornadic simulations experience multiple periods of tornadic activity. In such cases, our analysis on the origins of inflow into the supercells is performed on the low-level mesocyclone that resulted in the most intense tornado, as defined by OW , throughout the simulation.

a. Origins of low-level mesocyclone inflow air

Supercells are largely considered to be products of the environments in which they form. The vertical distribution of quantities such as temperature, moisture, and winds, exert substantial influence over a supercell’s evolution. While previous studies have advanced our understanding of the

inflow properties that favor supercells that produce tornadoes (compared to seemingly similar nontornadic supercells), questions still remain about where, both horizontally and vertically, supercells of varying intensity source most of their inflow air into the low-level updraft and mesocyclone. This in turn may shed light on the comparative importance of environmental air versus air that is modified within the storm.

1) TRACERS

To first broadly understand the height from which the low-level updraft draws most of its inflow, we use tracers initialized within three layers: 0 – 500 m AGL, 500 m – LFC, and the LFC – EIL. Within a 10 km diameter of the 1 km AGL low-level mesocyclone centroid, the progressively stronger parts of the low-level updraft are increasingly made up of air originating from below 500 m during a five minute composite period (t_{-10} to t_{-5}) relative to the key time period of tornado-genesis/failure. (Fig. 5). For the 50th percentile of vertical velocity and above, the supercell’s low-level updraft is a mixture of air from below the LFC, with the largest proportion of air generally originating between 500 m AGL and the LFC (Fig. 5a); however, as the updraft threshold is increased

TABLE 1. Summary of characteristics from each simulation, including the defined key time period of tornado-genesis/failure (min), the threshold of the 90th percentile of vertical velocity (m s^{-1}) at 1 km AGL within a 10 km diameter of the low-level mesocyclone centroid at the key time period, the maximum 1 km updraft speed (m s^{-1}) at the key time period, the maximum ground-relative wind speed (m s^{-1}) at 10 m AGL within 1 km of the position of maximized Okubo-Weiss (OW) parameter for tornadic/nontornadic period of interest (and the corresponding EF rating), the maximum vertical vorticity (s^{-1}) for tornadic/nontornadic period of interest, the maximum OW for tornadic/nontornadic period of interest (s^{-1}).

	key time period	90th percentile 1 km updraft threshold	max 1 km updraft speed	max tornadic surface wind speed	max tornadic surface ζ	max tornadic surface OW
0_0	t = 137	8.7	20.5	51.6 (EF2)	0.81	0.21
1_0	t = 99	3.3	7.6	nontornadic	–	–
2_0	t = 116	6.0	15.1	nontornadic	–	–
0_1	t = 148	6.9	13.1	nontornadic	–	–
1_1	t = 126	6.9	21.5	52.8 (EF2)	1.13	0.29
2_1	t = 86	10.1	27.9	84.7 (EF4)	1.87	1.01
0_2	t = 118	8.4	29.3	96.6 (EF5)	2.15	1.20
1_2	t = 117	6.6	30.5	67.9 (EF3)	1.19	0.22
2_2	t = 143	14.6	41.6	97.2 (EF5)	2.15	1.32

(75th \rightarrow 90th \rightarrow 99th percentile) to isolate the fastest rising air (and thus the largest $\frac{\partial w}{\partial z}$), the concentration of air from the near-ground layer increases substantially across seven out of the nine simulations. At and above the 99th percentile of updraft values within the low-level mesocyclone, most simulations contain essentially pure, undiluted near-surface air with concentration values close to 1 (Fig. 5d), especially in the tornadic simulations. While tracers cannot indicate whether this near-ground air is coming directly from the environment or has passed through the forward flank, concentration values approaching unity within the core of the low-level updraft are likely indicative of air that is unmodified by the storm since re-ingested forward flank air parcels would tend to imply dilution from a history of descent and mixing. Simulations 1_0 and 2_0 (both nontornadic and having the lowest environmental SRH500 values) are noticeable outliers to this trend, although even in 2_0, at least 60% of the low-level updraft air is being fed by the near-ground layer. Compared to other tornadic simulations, the 0_0 supercell has marginally less ($\sim 15\text{--}20\%$) near-ground air within the low-level mesocyclone (an indication that this low-level mesocyclone, in a lower SRH500 environment, perhaps is supplemented by other sources of air).

The explanation for the importance of the near-ground layer in the low-level mesocyclone relates to the ascent angle (ϕ) of the storm-relative wind into an updraft. As inflow air approaches the storm and enters the footprint of the updraft, vertical tilting of horizontal vorticity occurs. From Peters et al. (2022a), the tilting of horizontal vorticity in a perfectly streamwise environment can be expressed as:

$$\zeta_{LLM} = \omega_s \frac{w}{V_{SR}} = \omega_s \tan \phi, \quad (2)$$

where ω_s is streamwise vorticity, w is the vertical velocity of the updraft, and V_{SR} is the storm-relative wind [Davies-Jones (2022) presents a similar equation]. Although the wind profiles in Figure 1 are not perfectly streamwise, Eq. 2 represents a good first order approximation of the tilting of horizontal vorticity into a mature, right-moving supercell’s updraft.

The simplest interpretation of Eq. 2 is that the tilting of ω_s is related to the slope of trajectories entering an updraft (i.e., “rise over run” or $\frac{w}{V_{SR}}$). Peters et al. (2022a) found that ϕ varies with updraft width and storm-relative flow, but the median ϕ for parcels bound for the low-level mesocyclone across a spectrum of supercell wind profiles was approximately 10° . Thus, the slope of the trajectories into the low-level mesocyclone is fairly gentle. Only parcels originating from the near-ground layer are likely to have fully converted ω_s into ζ by 1 km AGL. Thus, as suggested by Markowski et al. (2012b), large near-ground streamwise vorticity establishes the base of the low-level mesocyclone as close to the surface as possible given typical ascent angles, inducing a “dynamical feedback” process of pressure falls and upward directed perturbation pressure gradient accelerations (Goldacker and Parker 2021) needed for lifting and stretching negatively buoyant, circulation-rich air within the supercell’s outflow.

In the current simulations, cross-sections through the low-level updraft display similar trends in ascent angles and tracer concentrations within the supercell’s updraft at varying altitudes. For the 2_2 supercell (Fig. 6), even prior to tornadogenesis, an intense core of vertical velocities greater than 15 m s^{-1} extends down to 500 m AGL (Fig. 6b), with the maximum 1 km updraft exceeding 40 m s^{-1} (Table 1). Within the updraft from 500 m – 1 km AGL, the concentration of air from the near-ground layer is essentially one (Fig. 6c). In fact, this is generally the




FIG. 5. Average tracer concentration in the 0–500 m AGL, 500 m – LFC, and the LFC–EIL layers within a 10 km diameter of the 1 km AGL low-level mesocyclone centroid within an updraft area at and above various vertical velocity thresholds (50, 75, 90, and 99th percentile) during a five minute composite period (t_{-10} to t_{-5}) relative to the key time period of tornado-genesis/failure.

case within the core of the updraft up to 2 km AGL. At this point, a much larger concentration of air from 500 m to the LFC is present (Fig. 6d). Both Nowotarski et al. (2020, their Fig. 5) and Lasher-Trapp et al. (2021, their Fig. 14) show examples of this gentle ascent layer, where air in the upper part of the inflow layer (i.e., 500 m – 2 km AGL) does not contribute to the core of the updraft until much farther aloft (i.e., 2 – 4 km AGL). Below 2 km AGL, what little air that is present from above 500 m is predominately found along the downshear (i.e., the eastern) edge of updraft (Fig. 6d).

Across eight of the nine simulations herein (with the transient 1_0 supercell again being the outlier), for updrafts defined by the 90th percentile of vertical velocity and above, the highest concentration of air is definitively from the 0 – 500 m layer (Fig. 7a,c-i). Where air above this layer contributes most substantially is along the eastern, downshear flank of the updraft (Fig. 7j,l-q). This is consistent with this air stream not yet being fully tilted into the vertical (Eq. 2) and residing along the edge of the updraft footprint. Air from above the LFC is not found with any consistency within the core of the supercells' updraft at any height within the troposphere (not shown), and accord-




FIG. 6. a) Horizontal cross section of 10 dBZ reflectivity for the 2_2 violently tornadic (VT) supercell at the key time period of tornadogenesis. Path of vertical cross-section used in subsequent panels is shown in red from A to A'. b) Vertical cross-section of vertical velocity (shaded) with 1 km AGL labeled as a dashed line and 15 m s^{-1} vertical velocity contour in black. c) As-in b), except shaded is the 0 – 500 m tracer and the 15 m s^{-1} vertical velocity contour is in white. d) As-in c), except shaded is the 500 m – LFC tracer.

ingly is virtually non-existent in the low-level mesocyclone (Fig. 7s-zz). Although this is likely not surprising given the altitude of the LFC ($\sim 1.7 \text{ km}$), there is a historical precedence in tornado forecasting of integrating SRH over depths much greater than the LFC [e.g., 0 – 3 km AGL SRH in Rasmussen and Blanchard (1998) and SRH in the EIL (ESRH) in Thompson et al. (2007)], which implicitly implicates a participation of air parcels in the low-level mesocyclone within this layer.

2) BACKWARD TRAJECTORIES

Next we turn to backward trajectories initialized within the most intense upward-moving, cyclonically rotating air in the low-level mesocyclone. While the three layers of

tracers show that the low-level mesocyclone is predominately made up of near-ground air, tracers alone cannot show the inflow origins of air bound for the low-level mesocyclone (e.g., the undisturbed, ambient environment versus the forward-flank baroclinic zone). To address this, we initialized backward trajectories within the 90th percentile of vertical velocity at 1 km AGL in each simulation, as this area has the highest potential for stretching of sub-tornadic surface vortices into tornadoes.

During a five minute period prior to tornadogenesis/failure (t_{-10} to t_{-5}), the origin height of inflow air into the mesocyclone is highly consistent across eight of the nine supercells. Similar to the tracer analysis, backward trajectories bound for the low-level mesocyclone originate very close to the ground (Table 2). The distributions of the




FIG. 7. Horizontal cross sections of tracer values for each simulation within the 90th percentile of updraft at 1 km (black contour) at the key time period of tornado-genesis/failure (see Table 1) for (top) 0–500 m AGL, (middle) 500 m – LFC, and (bottom) the LFC–EIL tracer layers. Each column is labeled nontornadic (NT), tornadic (T), or violently tornadic (VT).

TABLE 2. Range of low-level mesocyclone trajectory origin heights for both the median trajectory and the trajectory representing the 90th percentile of data.

	median trajectory origin height (m)	90th percentile trajectory origin height (m)
0_0	190.6	336.5
1_0	562.4	1325.3
2_0	368.7	528.7
0_1	161.9	403.8
1_1	274.0	436.6
2_1	245.1	487.4
0_2	161.4	435.1
1_2	310.0	681.5
2_2	321.0	630.9

origin height of trajectories shown in the insets of Figure 8 are generally below 500 m (excluding 1_0). The median origin height for the eight main supercells is less than 400 m AGL and 90% of the parcels in each simulation come from below 700 m AGL (Table 2). Only a few trajectories across the matrix of supercells represent “recycled air”, or air with a history of descent from farther aloft⁵ (Fig. 9). Many of the simulations have median parcel heights less than 300 m (Table 2). The very low altitude of parcels that contribute to the strongest vertical motion within the mesocyclone likely explains the comparative forecast skill of environmental streamwise horizontal vorticity and thus

⁵The exact path of “recycled” low-level mesocyclone trajectories should be treated with caution since the likelihood of errors in the backward trajectory integration is higher for such a flow regime. Regardless, the overwhelming proportion of trajectories that rise into the low-level mesocyclone from the undisturbed inflow compared to the “recycled air” is still qualitatively informative.

environmental SRH in progressively shallower layers (e.g., as shallow as 0 – 250 m AGL in Coffey et al. 2020).

Compared to the vertical extent of the inflow, the horizontal extent of air bound for the low-level mesocyclones across the simulations generally originates from south and east of the low-level updraft (Figs. 8,9), consistent with the orientation of the near-ground hodographs in Figure 3. The trajectory fields in the simulations with higher SRH have a noticeably more expansive inflow region, especially towards the southeast (Fig. 8e-i). The highest density of parcel origins in most of the simulations (Fig. 8d-i) is from well outside of the precipitation field. The paths of these trajectories, coming from the undisturbed, far-field environment into the low-level mesocyclone, appear to traverse the forward flank only minimally (or not at all in some instances, e.g., Fig. 9d,g). Especially for the tornadic supercells, the parcel origins are mostly from the far-field, toward the southeast (Fig. 8e-f-i), with one exception (0_0). In 0_0, the initial locations are primarily due east of the low-level mesocyclone (Fig. 8a) and flow parallel to the forward flank (Fig. 9a), consistent with the orientation of the storm-relative wind in the 0_0 hodograph (Fig. 1a).

At least 65% of parcels bound for the low-level mesocyclone in the eight main supercells spend less than 5 minutes in areas influenced by the storm’s hydrometeor field (Fig. 10) based on the accumulated time within regions characterized by $q_{hyd} > 0.001 \text{ kg kg}^{-1}$ and $\theta'_p < -1 \text{ K}$. In the instances characterized by the weakest forward flank cold pools (Fig. 9f,g,h), this percentage is greater than 90%. For the small percentage of parcels that do interact with baroclinic gradients associated with the forward flank, the experienced deficits in density potential temperature rarely exceed -2 to -3 K, and are generally closer to -1




Fig. 8. Two-dimensional kernel density estimates (KDE) of the final horizontal locations of the backward trajectories initialized from the low-level mesocyclone. Shaded KDE values represent 10, 25, 50, 75, and 90% of the data. The average 10 dBZ reflectivity contour (black) during the five minute composite period (t_{-10} to t_{-5}) relative to the key time period of tornado-genesis/failure is shown for the respective simulations. The blue rectangles represent the bounding box used to initialize forward trajectories. Inset in each panel is a histogram of the final vertical locations of the backward trajectories binned every 10 m from 0 to 2000 m AGL. All data is centered upon the 1 km AGL low-level centroid. Each panel is labeled nontornadic (NT), tornadic (T), or violently tornadic (VT).

K (Fig. 9). In that respect, the density potential temperature fields in the present simulations resemble those of the observed tornadic supercells in Shabbott and Markowski (2006). As a result of the short residence time within the storms' forward flanks, weak deficits in potential temperature, and fast storm-relative winds accelerating towards the supercell, the mean low-level mesocyclone trajectory in the eight main supercells experiences a rather small change in streamwise horizontal vorticity along its inflow path. This is more precisely quantified in the following section.

In summary, inflow air into the low-level mesocyclone originates very close to the ground and overwhelmingly from the undisturbed, far-field environment (toward the southeast). Most parcels bound for the low-level mesocyclone experience minimal effects from the storm's precipitation field. Both of these results would be expected to have a direct effect on the importance of environmental versus storm-generated vorticity contributions to the low-level mesocyclone. We explore this topic directly next.




Fig. 9. Paths of backward trajectories initialized from low-level mesocyclone backward trajectories (every 10th trajectory plotted) in the $x - y$, $x - z$, and $y - z$ planes. In the $x - y$ plane, the average density potential temperature perturbation field (θ'_ρ) at 10 m AGL and the average 10 dBZ reflectivity contour (red) during the five minute composite period (t_{-10} to t_{-5}) relative to the key time period of tornado-gensis/failure is shown for the respective simulations. All data is centered upon the 1 km AGL low-level centroid. Each panel is labeled nontornadic (NT), tornadic (T), or violently tornadic (VT).

b. Contributions of environmental vs. storm-generated vorticity to the low-level mesocyclone

Through both tracers and backward trajectories, we have shown thus far that the air comprising the most intense upward-moving, cyclonically rotating air in the low-level mesocyclone originates from the near-ground layer (<500 m AGL) and predominately from the undisturbed inflow environment, with the highest density of parcels appearing to have very little residence time within the region of precipitation and negative buoyancy associated with the

forward-flank region. On the face of it, these two factors would seemingly implicate the near-ground environmental horizontal streamwise vorticity as the dominant contributor to the overall rotation of the low-level mesocyclone, not the storm-generated streamwise vorticity classically associated with low-level mesocyclone-gensis. To quantitatively evaluate this interpretation, we track forward trajectories bound for the low-level mesocyclone and assess their associated vorticity via stencils of nearby adjacent parcels following the technique described by Dahl et al. (2014). As described in Section 2, forward trajectories




FIG. 10. Histograms of the accumulated time each backward trajectory initialized from low-level mesocyclone spent in regions characterized by $\theta'_\rho < -1$ K (blue) and $q_{hyd} > 0.001$ kg kg⁻¹ (orange) for each simulation. Data is binned in 1 minute increments. The area underneath each histogram sum to 100%. Each panel is labeled nontornadic (NT), tornadic (T), or violently tornadic (VT).

were seeded within model restart files upstream of the low-level mesocyclone within a unique horizontal bounding box for each simulation (the blue boxes in Fig. 8) encompassing roughly 75% of the low-level mesocyclone inflow area based on the origins of the backward trajectories. The seeding of the forward trajectories is meant to represent a majority of the inflow air; computational imitations prevent an exhaustive sample of all possible inflow air.

The forward trajectory parcels that meet the vertical velocity and vertical vorticity thresholds [vertical velocities greater or equal to the 90th percentile at 1 km AGL (Table 1) and at least 0.01 s⁻¹ of vertical vorticity (within ± 10 m of 1 km AGL)]⁶ have similar paths into the low-level mesocyclone and originate from similar locations as the backward trajectories in the previous section (Fig. 11). This provides some quality assurance, which is welcome

⁶An overwhelmingly majority of the forward trajectories released within the inflow region would have qualified as low-level mesocyclone parcels if the parcel output frequency was decreased and/or the depth of the vertical layer surrounding 1 km AGL was increased. These choices simply acted to filter parcels to a reasonable number for analysis given storage and computational constraints.

in light of documented differences in accuracy between forward and backward trajectory techniques (Dahl et al. 2012). Because the forward trajectories were not seeded at the exact terminal locations of the backward trajectories (rather, they were initialized over an isotropic grid covering most of the inflow region), it is not possible to directly compare the two sets of trajectories (as-in Dahl et al. 2012, Gowan et al. 2021); however, many of the details, including the shape, width, depth, and proportion of undisturbed, far-field environment parcels to forward flank parcels, are extremely similar (Figs. 9, 11).

Each seven-parcel stencil tracks $\zeta_{BT_{env}}$ and $\zeta_{BT_{pert}}$ (as described in Section 2) via the deformation and stretching of the initial vorticity vector over time, while ζ_{NBT} is computed as a residual. In eight of the nine simulations (except 1_0), ζ_{BT} is far and away the dominant contributor to the low-level mesocyclone vertical vorticity (ζ_{LLM} ; Fig. 12a). Only in simulation 1_0, the weakest and most transient supercell, is ζ_{NBT} the dominant component of ζ_{LLM} . This is not entirely unexpected considering the entire inflow region of the 1_0 supercell's low-level meso-




FIG. 11. Paths of forward trajectories that end up in the low-level mesocyclone in the $x-y$, $x-z$, and $y-z$ planes. Also shown in the $x-y$ plane, a scatter plot of the initial location of each forward trajectory colored by the ratio of $\zeta_{BT_{env}} / \zeta_{LLM}$ from 0 to 1 and the average 10 dBZ reflectivity contour (red) during the five minute composite period (t_{-10} to t_{-5}) relative to the key time period of tornado-genesis/failure is shown for the respective simulations. All data is centered upon the 1 km AGL low-level centroid. Each panel is labeled nontornadic (NT), tornadic (T), or violently tornadic (VT).

cyclone originates within precipitation of the forward flank (Figs. 8b, 9b, 11b). For the other eight main simulations, $\zeta_{BT_{env}}$ contributes between 65% and 90% of the total ζ_{LLM} indicating that *the environmentally-derived vorticity comprises a much larger percentage of the mesocyclone's vertical vorticity than the storm-generated vorticity* (Fig. 12a). While $\zeta_{BT_{env}}$ is generally quite high for all these eight simulations, there is a trend for the tornadic supercells towards the right of Figure 12a, starting with 1_1, to have a higher percentage of $\zeta_{BT_{env}}$. These simulations also have the most favorable lower tropospheric base-state hodographs

and the highest values of SRH500. The exception to that trend, simulation 0_0, has $\sim 20\%$ lower contribution from $\zeta_{BT_{env}}$ than the other tornadic supercells (Fig. 12a). Not only does the 0_0 supercell have a base-state SRH500 of $67 \text{ m}^2 \text{ s}^{-2}$ (well below the median tornadic SRH500 value from Coffey et al. 2019), but also has the lowest concentration of near-ground tracer among the tornadic low-level mesocyclones (Fig. 5) and highest proportion of parcels that flow into the mesocyclone directly parallel to the forward flank baroclinic gradient (Fig. 9). In total, this potentially suggests the 0_0 supercell required additional augmenta-

tion from within storm baroclinic generation of streamwise vorticity to establish a low-level mesocyclone capable of producing a tornado. Fully fleshing out this hypothesis would require an ensemble of simulations and more additional analysis, which is beyond the scope of this study and will be expanded upon in future work.

The next largest contributor to ζ_{LLM} (either $\zeta_{BT_{pert}}$ or ζ_{NBT}) varies between the individual simulations but is generally less than 20%. There is no discernible trend for the nontornadic or tornadic supercell simulations to have more or less $\zeta_{BT_{pert}}$ than ζ_{NBT} . As a reminder, because we cannot say whether the $\zeta_{BT_{pert}}$ represents prior reorientation or stretching of base-state vorticity versus prior baroclinic (or frictional) generation, it is treated separately. Even if we generously assume $\zeta_{BT_{pert}}$ is entirely attributed to storm-generated effects, their combined contribution would still be less than 35% of the total ζ_{LLM} for the eight main storms.

While the bulk percentages of ζ_{BT} compared to ζ_{NBT} paint a clear picture that most of the low-level mesocyclone rotation is from the environmental vorticity, examining individual parcels and their ratio of $\zeta_{BT_{env}}$ to ζ_{LLM} highlights source regions from which nonbarotropic generation is more prominent, such as the forward flank. There is a trend in some (but not all) of the supercells to have lower percentage of $\zeta_{BT_{env}}$ within parcels that originate closer to the hydrometeor field and forward flank (Fig. 11c,e,i). Those three simulations (2_0, 1_1, 2_2) also represent simulations where the backward trajectories cross through larger density potential temperature gradients (Fig. 9c, e, i) and have a higher frequency of parcels that spend 5 – 25 minutes of accumulated time within the hydrometeor and negative buoyancy fields (Fig. 10c,e,i). Specifically looking at simulation 2_2 (Fig. 11i) as an example, many of the parcels with the lowest percentage of $\zeta_{BT_{env}}$ (and thus highest ζ_{NBT} ; located at approximately $x=30, y=-5$ in Fig. 11i) traverse west-northwestward into forward flank (and larger negative buoyancy gradients; Fig. 9i), before turning back towards the updraft and eventually rising into the low-level mesocyclone. Due to their path through the storm, and exposure to horizontal baroclinity, these isolated parcels likely correspond to the classical conceptual model of forward flank air being re-ingested into the low-level updraft and wall cloud.

1) LOW-LEVEL MESOCYCLONE VORTICITY SOURCES IN SUPPLEMENTAL SIMULATIONS

On one hand, because of the documented forecast skill of near-ground environmental SRH in separating nontornadic from significantly tornadic supercell environments, it seems entirely logical that the environmental vorticity would exert a substantial influence on the total vertical vorticity of the low-level mesocyclone. On the other hand, because the low-level mesocyclone has long been attributed

to baroclinically-generated streamwise horizontal vorticity within the storm (as discussed in the introduction), the degree to which $\zeta_{BT_{env}}$ dominates, and consistency among the simulations, is somewhat surprising. To supplement the results from the simulations presented herein, we also present previously unpublished barotropic stencil analyses from two existing supercell studies in the literature, Wade and Parker (2021) and Peters et al. (2022a).

These studies complement our simulations by virtue of their different thermodynamic and kinematic environmental profiles. Wade and Parker (2021) focused on three high-shear, low-CAPE (HSLC) environments from a VORTEX-SE case in Alabama on 31 March 2016 as well as a companion high-shear, high-CAPE environment from the 3 April 1974 “Super Outbreak” (see their Fig. 6 for Skew T - $\log P$ and hodograph diagrams). Peters et al. (2022a) presented a large number of supercell simulations with “L” and “C” shaped hodographs (see their Figs. 1-3 for Skew T - $\log P$ and hodograph diagrams), independently varying the streamwise vorticity and storm-relative flow to disentangle their influence on low-level mesocyclone characteristic. Each simulation in Peters et al. (2022a) used a constant thermodynamic environment based on the tornadic VORTEX2 composite environment from Parker (2014). The reader is referred to these papers for more details about their simulations. The barotropic stencil analysis was conducted independently amongst the three studies, with varying thresholds of vertical velocity/vertical vorticity and criteria for when/where forward trajectories were seeded. Of note, these studies calculated ζ_{BT} and ζ_{NBT} only; no attempt was made to distinguish $\zeta_{BT_{env}}$ from $\zeta_{BT_{pert}}$. The chief similarity between all three studies is that parcels were filtered to highlight those entering the low-level mesocyclone at 1 km AGL. We believe the modest differences in analysis techniques increase confidence that the results presented in the previous subsection are not unique to our specific methodological choices.

Similar to the eight main supercell simulations in the present study, the low-level mesocyclones of both Wade and Parker (2021) and Peters et al. (2022a) primarily derive vertical vorticity from barotropic sources rather than nonbarotropic (Fig. 12b). In the Wade and Parker (2021) storms, the low-level mesocyclone is almost entirely barotropic for the parcel groups in the high-CAPE storm and low-CAPE 1 and 3 (> 85%). Even in their low-CAPE 2 supercell, which in general presented more analysis challenges than the other three supercells [see Wade (2020) for more details], the barotropic component of the low-level mesocyclone is greater than 65%. Given the differences in storm structures and cold pools amongst the high-CAPE and low-CAPE storms (Wade and Parker 2021), it is remarkable how consistently little the non-barotropic term contributes to the low-level mesocyclone (Fig. 12b, Supplemental Fig. 1). For the Peters et al. (2022a) supercells, the storm-generated, nonbarotropic component comprised




FIG. 12. Percentage of ζ_{LLM} attributable to ζ_{BT} , $\zeta_{BT_{env}}$, $\zeta_{BT_{pert}}$, and ζ_{NBT} for low-level mesocyclone parcels from both a) the supercell simulations presented herein and b) from the supercell simulations by Wade and Parker (2021) and Peters et al. (2022a). In a), ‘x’ marks the median percentage value from all the low-level mesocyclone trajectories for each simulation and the error bars represent the standard deviation of those trajectories, while in b), ‘x’ marks the median percentage from the average trajectory for each simulation [Wade and Parker (2021): $n=4$, Peters et al. (2022a): $n=24$] and the error bars represent the standard deviation of each paper’s simulations.

less than 35% of the total ζ_{LLM} on trajectories in most simulations (Fig. 12b), only exceeding this percentage in the storms with the weakest mesocyclones. Regardless of the differing combinations of storm-relative flow and streamwise vorticity, the environmental, barotropic contribution was generally greater than 65%⁷. In total, these results further demonstrate that *environmentally-derived vorticity comprises a much larger percentage of the low-level mesocyclone than storm-generated vorticity in persistent, mature supercells.*

⁷None of the low-level mesocyclone/updraft attributes systemically evaluated in Peters et al. (2022a), such as updraft and mesocyclone radius, net updraft circulation and rotational velocity, as well as average updraft vertical vorticity and helicity density, displayed any meaningful correlation with the fraction of ζ_{BT} to ζ_{LLM} (Supplemental Fig. 2).

4. Conclusions and discussion

In this article, we sought to address where inflow air bound for the low-level mesocyclone originates from and whether the origins of such air could address the dynamical role of near-ground streamwise vorticity present in the ambient environment versus what is generated in-situ within the forward flank of a supercell. The roles of these competing sources of streamwise vorticity potentially modulate the intensity the low-level mesocyclone and can ultimately determine whether a supercell fails or succeeds at producing a tornado. Using a matrix of nine supercell simulations, initialized with a spectrum of near-ground wind profiles observed in nature, we found the following:

- The air that comprises the core of the mesocyclone at 1 km, where the greatest potential for vertical stretching exists, originates almost exclusively from very close

to the ground, often in the lowest 200 - 400 m AGL. Air originating above 500 m AGL does not tend to contribute to the main updraft until farther aloft.

- Air bound for the low-level mesocyclone primarily originates from the undisturbed, ambient environment, rather than from along the forward flank. In both the nontornadic and tornadic supercells, 60 to 90% of the inflow air into the low-level mesocyclone has little to no residence time within regions characterized by negative buoyancy or hydrometeors in the forward flank.
- The dominant contributor to vertical vorticity within the low-level mesocyclone is from the environmental horizontal vorticity. As much as 90% of the low-level mesocyclone vertical vorticity can be solely attributed to the base-state environment, with storm-generated perturbations accounting for the residual. For the few parcels that do traverse baroclinic gradients within the storm, the contribution of storm-generated vorticity is higher.

We were motivated to understand why near-ground environmental streamwise vorticity is such a highly skillful tornado forecast parameter given the long history linking the low-level mesocyclone intensity to within-storm baroclinic generation of horizontal vorticity (e.g., Klemp and Rotunno 1983; Rotunno and Klemp 1985). Low-level mesocyclone air in our simulated supercells comes from the ambient environment. Thus, low-level mesocyclone intensification does not require the development of additional horizontal vorticity in the forward flank. The ingestion of forward flank parcels may instead be a *symptom* of an intensifying low-level mesocyclone (driven by environmental horizontal vorticity). After all, in order for rain-cooled air to be re-ingested by the storm, substantial dynamic lifting must be present to force negatively (or at least neutrally) buoyant forward flank outflow parcels upwards.

Perhaps some of the apparent prior misattribution can simply be attributed to differences in nomenclature between studies (or lower vertical resolution in prior studies). Many studies of the origins of low-level rotation referred to low-level mesocyclones in the approximately the lowest 250 m (e.g., Rotunno and Klemp 1985; Wicker and Wilhelmson 1995; Adlerman et al. 1999; Beck and Weiss 2013). In actuality, these studies were likely quantifying the development of *near-ground* rotation, which has rather conclusively been attributed to baroclinic gradients and vorticity generation within *downdrafts* in the outflow (Davies-Jones and Brooks 1993; Markowski and Richardson 2014; Dahl et al. 2014; Schenkman et al. 2014; Markowski 2016; Roberts et al. 2016; Fischer and Dahl 2020, 2022). The low-level mesocyclone, on the other hand, is predominately associated with *upward* tilting of

horizontal vorticity⁸. We encourage future research endeavours to use a unified nomenclature when discussing supercell processes in the lower troposphere, as the governing dynamics between the development of rotation near the ground (< 250 m AGL) and near cloud-base (~ 1 km AGL) are very likely distinct (at least prior to tornadogenesis).

The operational ramifications of this study include the explanation that SRH500 is a more skillful tornado parameter than SRH over deeper layers because air near the ground is overwhelmingly more likely to contribute to the low-level mesocyclone than air farther aloft. In turn, horizontal vorticity of the near-storm inflow exerts substantial control over the width and intensity of the low-level mesocyclone (Peters et al. 2022a). These results further underscore the need for more frequent and more numerous observations of the near-ground vertical wind profile than what are currently available. Improvements to forecasters' situational awareness could be realized through high spatial and temporal sampling of the environment near storms (e.g., as shown by Chilson et al. 2019; Bell et al. 2020).

Our simulations also provide three-dimensional context for the observations of forward-flank outflows presented by Shabbott and Markowski (2006), who found tornadic supercells to have small density gradients within the forward flank, whereas streamwise vorticity generation was largest in nontornadic cases. As speculated by Shabbott and Markowski (2006), "large ambient streamwise vorticity might obviate the need for baroclinic streamwise vorticity production", and in fact, "substantial baroclinic vorticity generation in the [forward-flank] outflow might be unfavorable for low-level mesocyclones and tornadogenesis" because excessive negative buoyancy, whether in the rear-flank or the forward-flank is generally unfavorable for tornadoes (Markowski et al. 2002; Grzych et al. 2007; Hirth et al. 2008; Weiss et al. 2015). This seems to be borne out by our simulations. Because of the out-sized role of the ambient environment in low-level mesocyclone-generation herein (regardless of the shape of the wind profile), environments with larger near-ground streamwise vorticity yield more intense low-level mesocyclones. In such cases, additional horizontal vorticity from the forward flank is superfluous to establishing the footprint of dynamic lifting in the low-levels of a supercell.

These results may run counter to the growing interest in SVCs (i.e., Orf et al. 2017; Schueth et al. 2021). The tornadic supercell simulated by Orf et al. (2017) possessed a base-state environment with an incredible amount of near-ground streamwise vorticity ($SRH500 > 300 \text{ m}^2 \text{ s}^{-2}$), and it is not yet known how much the storm-generated vorticity present in the SVC contributed to the low-level meso-

⁸Even if the forward flank contributes more to the overall rotation of the low-level mesocyclone than shown herein, these highly streamwise parcels in the lower troposphere would be tilted into the mesocyclone by an updraft, not a downdraft.

cyclone or whether the absence of an SVC would have hindered tornadogenesis. In a similarly high-resolution tornadic supercell simulation ($\Delta x = 30$ m) in a different environment, Finley et al. (2018) found that a tornado formed prior to forward flank cold pool development, and the SVC seemed to have little influence on the tornadogenesis process, with a mature SVC not developing until well after the tornadogenesis had occurred. Even if an SVC is present, those augmented parcels need to participate in the low-level mesocyclone. This seems not to be the case in our simulations or the recent studies by Murdzek et al. (2020a,b). The orientation of the wind profile, in relation to the baroclinic vorticity produced in the forward flank, might also determine whether these air parcels contribute more or less to the low-level mesocyclone. Future research could focus on the quantitative contribution of the SVC to the low-level mesocyclone and whether the storm-generated SVC is an instigator of tornadogenesis or whether it is primarily a by-product of an environmentally derived low-level mesocyclone.

As with any modeling study, the scope of our results could depend on numerous, untested factors, including (but not limited to) a narrow range of thermodynamic environments, the magnitude of low-level cooling attributable to the microphysics scheme, and the possibility that a single simulation in each wind profile is not representative of what a larger ensemble would produce. To this end, we have presented results from simulations of both Wade and Parker (2021) and Peters et al. (2022a), which used different environments, microphysics schemes, treatments of the lower boundary condition, and sub-grid turbulence closures. Some of the outstanding questions related to this study that remain to be addressed include: Why do the weaker supercells (such as-in 1_0 herein and in Peters et al. (2022a), see Supplemental Fig. 2e,f) have more forward flank baroclinic contributions? Are these storms weaker because the baroclinic contribution is high or is this a symptom of something else wrong in the storm? Additionally, is our finding that the dominant contributor to vertical vorticity within the low-level mesocyclone is from the environmental horizontal vorticity due to the lack of baroclinicity in the forward flank of these storms (and in observed tornadic storms) or is this more to do with the trajectory shapes themselves? Even if there were larger buoyancy gradients and more baroclinic generation of streamwise vorticity, would the additional negative buoyancy still favor inflow parcels from the ambient environment? In the future, consideration should be given to whether the storm generated vorticity contributes more within storms with larger buoyancy gradients within the forward flank than presented herein.

In conclusion, we posit a potential connection between the environmental streamwise vorticity and the potential role of the storm-generated streamwise vorticity. In order for rain-cooled air (such as the SVC) to be re-ingested by

the storm's low-level updraft in the first place, substantial dynamic lifting must be present to force these outflow parcels with negative buoyancy upwards to their levels of free convection. There appears to be a threshold of streamwise vorticity, above which, the dynamical response to streamwise horizontal vorticity increases dramatically (Coffer and Parker 2018; Goldacker and Parker 2021). We hypothesize that the ambient environmental streamwise vorticity establishes a "floor" of low-level mesocyclone intensity (and tornadogenesis potential), which could then be supplemented by within-storm baroclinic generation of streamwise vorticity. In high streamwise vorticity environments, the probability of tornadogenesis is argued to be high because very little (or no) augmentation from the storm is required in order to instigate the dynamical updraft response that favors tornadogenesis. On the other hand, in low streamwise vorticity environments, the probability of tornadogenesis is argued to be lower because much more augmentation from the storm is required. The necessity of within-storm supplementation adds to the possible failure points, as there must be sufficient dynamic lifting to overcome the negative buoyancy associated with the baroclinic vorticity generation. We plan to further explore supercell simulations in both marginal environments and environments with a wider range of thermodynamic profiles to test this hypothesis. Regardless, we believe the results of this study will fundamentally clarify how our community views the development of low-level mesocyclones in supercells.

Acknowledgments. We thank the anonymous peer reviewers for their helpful feedback on this manuscript. B. Coffer's and M. Parker's efforts were supported by the National Science Foundation (NSF) on grant AGS-2130936. J. Peters's efforts were supported by NSF grants AGS-1928666, AGS-1841674, and the Department of Energy Atmospheric System Research (DOE ASR) grant DE-SC0000246356. A. Wade was supported by National Oceanic and Atmospheric Administration (NOAA)–University of Oklahoma Cooperative Agreement NA16OAR4320115. Dr. George Bryan is gratefully acknowledged for providing and supporting the CM1 model. We appreciate the assistance of Dr. Sonia Lasher-Trapp with code for the heat flux initialization in CM1. Dr. Johannes Dahl is acknowledged for general guidance of implementing stencils. Nicholas Goldacker supplied the hodographs from the SOM, and Adam Werkerma developed the mesocyclone tracking algorithm. The base code for backward trajectories was cribbed from Dr. Tom Gowan's GitHub repository (<https://tomgowan.github.io/>). Many figures and analyses were created using open-sourced python packages such as Numpy (Harris et al. 2020), Matplotlib (Hunter 2017), Pandas (Wes McKinney 2010), Xarray (Hoyer and Hamman 2016), Metpy (May et al. 2017), and Jupyter (Kluyver et al. 2016).

Data availability statement. The CM1 model code is available from <https://www2.mmm.ucar.edu/people/bryan/cm1/>. All other namelists, input files, and post-processing scripts are available from the author by request.

References

- Adlerman, E. J., K. K. Droegemeier, and R. Davies-Jones, 1999: A numerical simulation of cyclic mesocyclogenesis. *J. Atmos. Sci.*, **56** (13), 2045–2069.
- Anderson-Frey, A. K., Y. P. Richardson, A. R. Dean, R. L. Thompson, and B. T. Smith, 2017: Self-organizing maps for the investigation of tornadic near-storm environments. *Wea. Forecasting*, **32** (4), 1467–1475.
- Atkins, N. T., E. M. Glidden, and T. M. Nicholson, 2014: Observations of wall cloud formation in supercell thunderstorms during VORTEX2. *Mon. Wea. Rev.*, **142** (12), 4823–4838.
- Beck, J. R., and C. Weiss, 2013: An assessment of low-level baroclinity and vorticity within a simulated supercell. *Mon. Wea. Rev.*, **141** (2), 649–669.
- Bell, T. M., B. R. Greene, P. M. Klein, M. Carney, and P. B. Chilson, 2020: Confronting the boundary layer data gap: Evaluating new and existing methodologies of probing the lower atmosphere. *Atmospheric Measurement Techniques*, **13** (7), 3855–3872.
- Blair, S. F., and Coauthors, 2017: High-resolution hail observations: Implications for NWS warning operations. *Wea. Forecasting*, **32** (3), 1101–1119.
- Brandes, E. A., 1978: Mesocyclone evolution and tornadogenesis: Some observations. *Mon. Wea. Rev.*, **106** (7), 995–1011.
- Brandes, E. A., 1993: Tornadic thunderstorm characteristics determined with doppler radar. *The Tornado: Its Structure, Dynamics, Prediction, and Hazards*, 143–159.
- Brown, M., and C. J. Nowotarski, 2019: The influence of lifting condensation level on low-level outflow and rotation in simulated supercell thunderstorms. *J. Atmos. Sci.*, **76** (5), 1349–1372.
- Bryan, G. H., and J. M. Fritsch, 2002: A benchmark simulation for moist nonhydrostatic numerical models. *Mon. Wea. Rev.*, **130** (12), 2917–2928.
- Burgess, D. W., R. J. Donaldson Jr, and P. R. Desrochers, 1993: Tornado detection and warning by radar. *Washington DC American Geophysical Union Geophysical Monograph Series*, **79**, 203–221.
- Carpenter, R. L., K. K. Droegemeier, and A. M. Blyth, 1998: Entrainment and detrainment in numerically simulated cumulus congestus clouds. Part i: General results. *J. Atmos. Sci.*, **55** (23), 3417–3432.
- Chilson, P. B., and Coauthors, 2019: Moving towards a network of autonomous UAS atmospheric profiling stations for observations in the earth’s lower atmosphere: The 3D mesonet concept. *Sensors*, **19** (12), 2720.
- Coffer, B. E., and M. D. Parker, 2015: Impacts of increasing low-level shear on supercells during the early evening transition. *Mon. Wea. Rev.*, **143** (5), 1945–1969.
- Coffer, B. E., and M. D. Parker, 2017: Simulated supercells in non-tornadic and tornadic VORTEX2 environments. *Mon. Wea. Rev.*, **145** (1), 149–180.
- Coffer, B. E., and M. D. Parker, 2018: Is there a “tipping point” between simulated nontornadic and tornadic supercells in VORTEX2 environments? *Mon. Wea. Rev.*, **146** (8), 2667–2693.
- Coffer, B. E., and M. D. Parker, 2022: Infrasound signals in simulated nontornadic and pre-tornadic supercells. *The Journal of the Acoustical Society of America*, **151** (2), 939–954.
- Coffer, B. E., M. D. Parker, J. M. Dahl, L. J. Wicker, and A. J. Clark, 2017: Volatility of tornadogenesis: An ensemble of simulated non-tornadic and tornadic supercells in VORTEX2 environments. *Mon. Wea. Rev.*, **145** (11).
- Coffer, B. E., M. D. Parker, R. L. Thompson, B. T. Smith, and R. E. Jewell, 2019: Using near-ground storm relative helicity in supercell tornado forecasting. *Wea. Forecasting*, **34** (5), 1417–1435.
- Coffer, B. E., M. Taszarek, and M. D. Parker, 2020: Near-ground wind profiles of tornadic and nontornadic environments in the United States and Europe from ERA5 reanalyses. *Wea. Forecasting*, **35** (6), 2621–2638.
- Coniglio, M. C., and M. D. Parker, 2020: Insights into supercells and their environments from three decades of targeted radiosonde observations. *Monthly Weather Review*, **148** (12), 4893–4915.
- Dahl, J. M., 2015: Near-ground rotation in simulated supercells: On the robustness of the baroclinic mechanism. *Mon. Wea. Rev.*, **143** (12), 4929–4942.
- Dahl, J. M., 2017: Tilting of horizontal shear vorticity and the development of updraft rotation in supercell thunderstorms. *Journal of the Atmospheric Sciences*, **74** (9), 2997–3020.
- Dahl, J. M., M. D. Parker, and L. J. Wicker, 2012: Uncertainties in trajectory calculations within near-surface mesocyclones of simulated supercells. *Mon. Wea. Rev.*, **140** (9), 2959–2966.
- Dahl, J. M., M. D. Parker, and L. J. Wicker, 2014: Imported and storm-generated near-ground vertical vorticity in a simulated supercell. *J. Atmos. Sci.*, **71** (8), 3027–3051.
- Davies-Jones, R., 1982: Observational and theoretical aspects of tornadogenesis. *Intense Atmospheric Vortices*, Springer, 175–189.
- Davies-Jones, R., 1984: Streamwise vorticity: The origin of updraft rotation in supercell storms. *J. Atmos. Sci.*, **41** (20), 2991–3006.
- Davies-Jones, R., 2015: A review of supercell and tornado dynamics. *Atmos. Res.*, **158–159**, 274–291.
- Davies-Jones, R., 2017: Roles of streamwise and transverse partial-vorticity components in steady inviscid isentropic supercell-like flows. *J. Atmos. Sci.*, **74** (9), 3021–3041.
- Davies-Jones, R., 2021: Invented forces in supercell models. *J. Atmos. Sci.*, **78** (9), 2927–2939.
- Davies-Jones, R., 2022: Theory of parcel vorticity evolution in supercell-like flows. *J. Atmos. Sci.*, **79** (5), 1253–1270.
- Davies-Jones, R., and H. Brooks, 1993: Mesocyclogenesis from a theoretical perspective. *The Tornado: Its Structure, Dynamics, Prediction, and Hazards*, 105–114.
- Davies-Jones, R., R. J. Trapp, and H. B. Bluestein, 2001: Tornadoes and tornadic storms. *Severe Convective Storms*, Springer, 167–221.
- Dearhoff, J. W., 1980: Stratocumulus-capped mixed layers derived from a three-dimensional model. *Bound.-Layer Meteor.*, **18** (4), 495–527.

- Doswell, C. A., and D. W. Burgess, 1993: Tornadoes and tornadic storms: A review of conceptual models. *Geophysical Monograph-American Geophysical Union*, **79**, 161–161.
- Esterheld, J. M., and D. J. Giuliano, 2008: Discriminating between tornadic and non-tornadic supercells: A new hodograph technique. *Electronic J. Severe Storms Meteor.*, **3** (2).
- Finley, C. A., L. Orf, B. D. Lee, and R. B. Wilhelmson, 2018: High-resolution simulation of a violent tornado in the 27 April 2011 outbreak environment. *29th Conference on Severe Local Storms*, AMS, [Available online at <https://ams.confex.com/ams/29SLS/webprogram/Paper348812.html>].
- Fischer, J., and J. M. Dahl, 2020: The relative importance of updraft and cold pool characteristics in supercell tornadogenesis using highly idealized simulations. *J. Atmos. Sci.*, **77** (12), 4089–4107.
- Fischer, J., and J. M. Dahl, 2022: Transition of near-ground vorticity dynamics during tornadogenesis. *J. Atmos. Sci.*, **79** (2), 467–483.
- Galarneau, T. J., L. J. Wicker, K. H. Knopfmeier, W. J. Miller, P. S. Skinner, and K. A. Wilson, 2022: Short-term prediction of a nocturnal significant tornado outbreak using a convection-allowing ensemble. *Wea. Forecasting*, **37** (6), 1027–1047.
- Goldacker, N. A., and M. D. Parker, 2021: Low-level updraft intensification in response to environmental wind profiles. *J. Atmos. Sci.*, Submitted.
- Gowan, T. M., W. J. Steenburgh, and J. R. Minder, 2021: Downstream evolution and coastal-to-inland transition of landfalling lake-effect systems. *Mon. Wea. Rev.*, **149** (4), 1023–1040.
- Grzych, M. L., B. D. Lee, and C. A. Finley, 2007: Thermodynamic analysis of supercell rear-flank downdrafts from Project ANSWERS. *Mon. Wea. Rev.*, **135** (1), 240–246.
- Hirth, B. D., J. L. Schroeder, and C. C. Weiss, 2008: Surface analysis of the rear-flank downdraft outflow in two tornadic supercells. *Mon. Wea. Rev.*, **136** (7), 2344–2363.
- Houser, J. L., H. B. Bluestein, K. Thiem, J. Snyder, D. Reif, and Z. Wienhoff, 2022: Additional evaluation of the spatiotemporal evolution of rotation during tornadogenesis using rapid-scan mobile radar observations. *Mon. Wea. Rev.*, **150** (7), 1639–1666.
- Hua, Z., and A. K. Anderson-Frey, 2022: Self-organizing maps for the classification of spatial and temporal variability of tornado-favorable parameters. *Mon. Wea. Rev.*, **150** (2), 393–407.
- Jiménez, P. A., J. Dudhia, J. F. González-Rouco, J. Navarro, J. P. Montávez, and E. García-Bustamante, 2012: A revised scheme for the WRF surface layer formulation. *Mon. Wea. Rev.*, **140** (3), 898–918.
- Klemp, J., 1987: Dynamics of tornadic thunderstorms. *Annual review of fluid mechanics*, **19** (1), 369–402.
- Klemp, J. B., and R. Rotunno, 1983: A study of the tornadic region within a supercell thunderstorm. *J. Atmos. Sci.*, **40** (2), 359–377.
- Kumjian, M. R., Z. J. Lebo, and A. M. Ward, 2019: Storms producing large accumulations of small hail. *J. Appl. Meteor. Climatol.*, **58** (2), 341–364.
- Lasher-Trapp, S., E. Jo, L. R. Allen, B. N. Engelsens, and R. J. Trapp, 2021: Entrainment in a simulated supercell thunderstorm. Part i: The evolution of different entrainment mechanisms and their dilutive effects. *J. Atmos. Sci.*, **78** (9), 2725–2740.
- Lemon, L. R., and C. A. Doswell, III, 1979: Severe thunderstorm evolution and mesocyclone structure as related to tornadogenesis. *Mon. Wea. Rev.*, **107** (9), 1184–1197.
- Mansell, E. R., 2010: On sedimentation and advection in multimoment bulk microphysics. *J. Atmos. Sci.*, **67** (9), 3084–3094.
- Mansell, E. R., C. L. Ziegler, and E. C. Bruning, 2010: Simulated electrification of a small thunderstorm with two-moment bulk microphysics. *J. Atmos. Sci.*, **67** (1), 171–194.
- Markowski, P. M., 2016: An idealized numerical simulation investigation of the effects of surface drag on the development of near-surface vorticity in supercell thunderstorms. *J. Atmos. Sci.*, **73** (11), 4349–4385.
- Markowski, P. M., 2020: What is the intrinsic predictability of tornadic supercell thunderstorms? *Mon. Wea. Rev.*, **148** (8), 3157–3180.
- Markowski, P. M., C. Hannon, J. Frame, E. Lancaster, A. Pietrycha, R. Edwards, and R. L. Thompson, 2003: Characteristics of vertical wind profiles near supercells obtained from the Rapid Update Cycle. *Wea. Forecasting*, **18** (6), 1262–1272.
- Markowski, P. M., and Y. P. Richardson, 2009: Tornadogenesis: Our current understanding, forecasting considerations, and questions to guide future research. *Atmos. Res.*, **93** (1-3), 3–10.
- Markowski, P. M., and Y. P. Richardson, 2010: *Mesoscale Meteorology in Midlatitudes*. Wiley-Blackwell, 372 pp.
- Markowski, P. M., and Y. P. Richardson, 2014: The influence of environmental low-level shear and cold pools on tornadogenesis: Insights from idealized simulations. *J. Atmos. Sci.*, **71** (1), 243–275.
- Markowski, P. M., Y. P. Richardson, E. Rasmussen, J. Straka, R. Davies-Jones, and R. J. Trapp, 2008: Vortex lines within low-level mesocyclones obtained from pseudo-dual-Doppler radar observations. *Mon. Wea. Rev.*, **136** (9), 3513–3535.
- Markowski, P. M., J. M. Straka, and E. N. Rasmussen, 2002: Direct surface thermodynamic observations within the rear-flank downdrafts of nontornadic and tornadic supercells. *Mon. Wea. Rev.*, **130** (7), 1692–1721.
- Markowski, P. M., and Coauthors, 2012a: The pretornadic phase of the Goshen County, Wyoming, supercell of 5 June 2009 intercepted by VORTEX2. Part I: Evolution of kinematic and surface thermodynamic fields. *Mon. Wea. Rev.*, **140** (9), 2887–2915.
- Markowski, P. M., and Coauthors, 2012b: The pretornadic phase of the Goshen County, Wyoming, supercell of 5 June 2009 intercepted by VORTEX2. Part II: Intensification of low-level rotation. *Mon. Wea. Rev.*, **140** (9), 2916–2938.
- Marquis, J., Y. Richardson, P. Markowski, J. Wurman, K. Kosiba, and P. Robinson, 2016: Analysis of the 5 June 2009 Goshen County, Wyoming, tornadic supercell through EnKF assimilation of mobile mesonet and radar observations. Part II: mesocyclone-scale processes affecting tornado formation, maintenance, and decay. *Mon. Wea. Rev.*, **144** (9), 3441–3463.
- Mashiko, W., 2016: A numerical study of the 6 May 2012 Tsukuba City supercell tornado. Part I: Vorticity sources of low-level and midlevel mesocyclones. *Mon. Wea. Rev.*, **144** (3), 1069–1092.
- May, R., S. Arms, P. Marsh, E. Bruning, and J. Leeman, 2017: Metpy: A python package for meteorological data. *Unidata, Accessed*, **31**.

- Miller, D. J., 2006: Observations of low-level thermodynamic and wind shear profiles on significant tornado days. *23rd Conf. on Severe Local Storms*, Amer. Meteor. Soc., St. Louis, MO, [Available online at <http://ams.confex.com/ams/pdfpapers/115403.pdf>].
- Morrison, H., J. M. Peters, K. K. Chandrakar, and S. C. Sherwood, 2022: Influences of environmental relative humidity and horizontal scale of subcloud ascent on deep convective initiation. *J. Atmos. Sci.*, **79** (2), 337–359.
- Moser, D. H., and S. Lasher-Trapp, 2018: Cloud-spacing effects upon entrainment and rainfall along a convective line. *J. Appl. Meteor. Climatol.*, **57** (8), 1865–1882.
- Murdzek, S. S., P. M. Markowski, and Y. P. Richardson, 2020a: Simultaneous dual-Doppler and mobile mesonet observations of streamwise vorticity currents in three supercells. *Mon. Wea. Rev.*, **148** (12), 4859–4874.
- Murdzek, S. S., P. M. Markowski, Y. P. Richardson, and R. L. Tanamachi, 2020b: Processes preventing the development of a significant tornado in a Colorado supercell on 26 May 2010. *Mon. Wea. Rev.*, **148** (5), 1753–1778.
- Naylor, J., and M. S. Gilmore, 2014: Vorticity evolution leading to tornadogenesis and tornadogenesis failure in simulated supercells. *J. Atmos. Sci.*, **71** (3), 1201–1217.
- Nielsen, E. R., and R. S. Schumacher, 2020: Observations of extreme short-term precipitation associated with supercells and mesovortices. *Mon. Wea. Rev.*, **148** (1), 159–182.
- Nowotarski, C. J., and A. A. Jensen, 2013: Classifying proximity soundings with self-organizing maps toward improving supercell and tornado forecasting. *Wea. Forecasting*, **28** (3), 783–801.
- Nowotarski, C. J., and E. A. Jones, 2018: Multivariate self-organizing map approach to classifying supercell tornado environments using near-storm, low-level wind and thermodynamic profiles. *Wea. Forecasting*, **33** (3), 661–670.
- Nowotarski, C. J., J. M. Peters, and J. P. Mulholland, 2020: Evaluating the effective inflow layer of simulated supercell updrafts. *Mon. Wea. Rev.*, **148** (8), 3507–3532.
- Orf, L., R. Wilhelmson, B. Lee, C. Finley, and A. Houston, 2017: Evolution of a long-track violent tornado within a simulated supercell. *Bull. Amer. Meteor. Soc.*, **98** (1), 45–68.
- Parker, M. D., 2014: Composite VORTEX2 supercell environments from near-storm soundings. *Mon. Wea. Rev.*, **142** (2), 508–529.
- Parker, M. D., and J. M. Dahl, 2015: Production of near-surface vertical vorticity by idealized downdrafts. *Mon. Wea. Rev.*, **143** (7), 2795–2816.
- Peters, J. M., B. E. Coffey, M. D. Parker, C. J. Nowotarski, J. P. Mulholland, C. J. Nixon, and J. T. Allen, 2022a: Disentangling the influences of storm-relative flow and horizontal streamwise vorticity on low-level mesocyclones in supercells. *J. Atmos. Sci.*
- Peters, J. M., H. Morrison, T. C. Nelson, J. N. Marquis, J. P. Mulholland, and C. J. Nowotarski, 2022b: The influence of shear on deep convection initiation. Part I: Theory. *J. Atmos. Sci.*, **79** (6), 1669–1690.
- Peters, J. M., H. Morrison, T. C. Nelson, J. N. Marquis, J. P. Mulholland, and C. J. Nowotarski, 2022c: The influence of shear on deep convection initiation. Part II: Simulations. *J. Atmos. Sci.*, **79** (6), 1691–1711.
- Rasmussen, E. N., 2003: Refined supercell and tornado forecast parameters. *Wea. Forecasting*, **18** (3), 530–535.
- Rasmussen, E. N., and D. O. Blanchard, 1998: A baseline climatology of sounding-derived supercell and tornado forecast parameters. *Wea. Forecasting*, **13** (4), 1148–1164.
- Roberts, B., M. Xue, A. D. Schenkman, and D. T. Dawson, II, 2016: The role of surface drag in tornadogenesis within an idealized supercell simulation. *J. Atmos. Sci.*, **73** (9), 3371–3395.
- Rotunno, R., and J. Klemp, 1985: On the rotation and propagation of simulated supercell thunderstorms. *J. Atmos. Sci.*, **42** (3), 271–292.
- Rotunno, R., and J. B. Klemp, 1982: The influence of the shear-induced pressure gradient on thunderstorm motion. *Mon. Wea. Rev.*, **110** (2), 136–151.
- Rotunno, R., P. M. Markowski, and G. H. Bryan, 2017: “near ground” vertical vorticity in supercell thunderstorm models. *J. Atmos. Sci.*, **74** (6), 1757–1766.
- Schenkman, A. D., M. Xue, and M. Hu, 2014: Tornadogenesis in a high-resolution simulation of the 8 May 2003 Oklahoma City supercell. *J. Atmos. Sci.*, **71** (1), 130–154.
- Schueth, A., C. Weiss, and J. M. Dahl, 2021: Comparing observations and simulations of the streamwise vorticity current and the forward flank convergence boundary in a supercell storm. *Mon. Wea. Rev.*
- Shabbott, C. J., and P. M. Markowski, 2006: Surface in situ observations within the outflow of forward-flank downdrafts of supercell thunderstorms. *Mon. Wea. Rev.*, **134** (5), 1422–1441.
- Smith, B. T., R. L. Thompson, J. S. Grams, C. Broyles, and H. E. Brooks, 2012: Convective modes for significant severe thunderstorms in the contiguous United States. Part I: Storm classification and climatology. *Wea. Forecasting*, **27** (5), 1114–1135.
- Straka, J. M., E. N. Rasmussen, R. P. Davies-Jones, and P. M. Markowski, 2007: An observational and idealized numerical examination of low-level counter-rotating vortices in the rear flank of supercells. *Electronic J. Severe Storms Meteor.*, **2** (8).
- Thompson, R. L., C. M. Mead, and R. Edwards, 2007: Effective storm-relative helicity and bulk shear in supercell thunderstorm environments. *Wea. Forecasting*, **22** (1), 102–115.
- Trapp, R. J., 1999: Observations of nontornadic low-level mesocyclones and attendant tornadogenesis failure during VORTEX. *Mon. Wea. Rev.*, **127** (7), 1693–1705.
- Trapp, R. J., G. J. Stumpf, and K. L. Manross, 2005: A reassessment of the percentage of tornadic mesocyclones. *Wea. Forecasting*, **20** (4), 680–687.
- Wade, A. R., 2020: *Dynamics and vorticity evolution in simulated low-CAPE supercells*. North Carolina State University.
- Wade, A. R., M. C. Coniglio, and C. L. Ziegler, 2018: Comparison of near-and far-field supercell inflow environments using radiosonde observations. *Mon. Wea. Rev.*, **146** (8), 2403–2415.
- Wade, A. R., and M. D. Parker, 2021: Dynamics of simulated high-shear, low-CAPE supercells. *Journal of the Atmospheric Sciences*, **78** (5), 1389–1410.
- Walko, R. L., 1993: Tornado spin-up beneath a convective cell: Required basic structure of the near-field boundary layer winds. *Washington*

- DC American Geophysical Union Geophysical Monograph Series*, **79**, 89–95.
- Wang, A., Y. Pan, and P. M. Markowski, 2020: The influence of turbulence memory on idealized tornado simulations. *Mon. Wea. Rev.*, **148 (12)**, 4875–4892.
- Warren, R. A., H. Richter, and R. L. Thompson, 2021: Spectrum of near-storm environments for significant severe right-moving supercells in the contiguous United States. *Mon. Wea. Rev.*, **149 (10)**, 3299–3323.
- Weisman, M. L., and J. B. Klemp, 1982: The dependence of numerically simulated convective storms on vertical wind shear and buoyancy. *Mon. Wea. Rev.*, **110 (6)**, 504–520.
- Weisman, M. L., and R. Rotunno, 2000: The use of vertical wind shear versus helicity in interpreting supercell dynamics. *J. Atmos. Sci.*, **57 (9)**, 1452–1472.
- Weiss, C. C., D. C. Dowell, J. L. Schroeder, P. S. Skinner, A. E. Reinhart, P. M. Markowski, and Y. P. Richardson, 2015: A comparison of near-surface buoyancy and baroclinity across three VORTEX2 supercell intercepts. *Mon. Wea. Rev.*, **143 (7)**, 2736–2753.
- Weiss, C. C., A. Schueth, and A. L. Houston, 2020: TORUS 2019 highlights from the TTUKa mobile Doppler radars. *100th American Meteorological Society Annual Meeting*, AMS.
- Wicker, L. J., and W. C. Skamarock, 2002: Time-splitting methods for elastic models using forward time schemes. *Mon. Wea. Rev.*, **130 (8)**, 2088–2097.
- Wicker, L. J., and R. B. Wilhelmson, 1995: Simulation and analysis of tornado development and decay within a three-dimensional supercell thunderstorm. *J. Atmos. Sci.*, **52 (15)**, 2675–2703.
- Ziegler, C. L., 1985: Retrieval of thermal and microphysical variables in observed convective storms. Part I: Model development and preliminary testing. *J. Atmos. Sci.*, **42 (14)**, 1487–1509.