

Renormalization formalism for superconducting phase transition with inner-Cooper-pair dynamics

Yuehua Su,* Hongyun Wu, Kun Cao, and Chao Zhang

Department of Physics, Yantai University, Yantai 264005, People's Republic of China

As charge carrier of the macroscopic superconductivity, the Cooper pair is a composite particle of two paired electrons, which has both center-of-mass and inner-pair degrees of freedom. In most cases, these two different degrees of freedom can be well described by the macroscopic Ginzburg-Landau theory and the microscopic Bardeen-Cooper-Schrieffer (BCS) theory, respectively. Near the superconducting phase transition where the Cooper pair is fragile and unstable because of the small binding energy, there are non-trivial couplings between these two different degrees of freedom due to such as finite energy and/or momentum transfer. The non-trivial couplings make the original derivation of the Ginzburg-Landau theory from the BCS theory fail in principle as where these two different degrees of freedom should not be decoupled. In this manuscript, we will present a renormalization formalism for an extended Ginzburg-Landau action for the superconducting phase transition where there is finite energy transfer between the center-of-mass and the inner-pair degrees of freedom of Cooper pairs. This formalism will provide a theoretical tool to study the unusual effects of the inner-pair time-retarded physics on the superconducting phase transition.

I. INTRODUCTION

In 1950, Ginzburg and Landau developed a macroscopic phenomenological theory for superconductivity by introducing a complex field as an order parameter to describe the superconducting state as a $U(1)$ symmetry-broken state¹. Later during 1956 to 1957, Bardeen, Cooper and Schrieffer established the well-known microscopic BCS theory for superconductivity with a basic idea of the Cooper-pair formation²⁻⁴. Microscopically, it is the coherent condensation of the Cooper pairs that leads to the macroscopic superconductivity. As the Cooper pair is a composite particle consisting of two electrons, it has both center-of-mass and inner-pair degrees of freedom. The macroscopic Ginzburg-Landau theory only involves the center-of-mass degrees of freedom of Cooper pairs and describes phenomenologically well the macroscopic superconductivity in most cases. The microscopic BCS theory mainly focuses on the inner-pair degrees of freedom for the formation of the Cooper pairs. In the derivation of Gor'kov to link the Ginzburg-Landau theory and the BCS theory^{5,6}, the inner-pair degrees of freedom are integrated out with assumption that the center-of-mass and the inner-pair degrees of freedom of Cooper pairs can be decoupled. This assumption was also taken into account in the derivation of the Ginzburg-Landau theory for an anisotropic superconductor⁷.

Generally, the superconducting pairing gap field Δ should be dependent on both the center-of-mass and the inner-pair degrees of freedom, i.e., $\Delta = \Delta(\mathbf{q}, \nu; \mathbf{k}, \omega)$ with (\mathbf{q}, ν) describing the center-of-mass degrees of freedom and (\mathbf{k}, ω) for the inner-pair ones. The \mathbf{k} variation describes the isotropic or anisotropic inner-pair structure of the pairing gap field. In an electron-phonon interaction driven superconductor, the pairing gap field is mostly \mathbf{k} independent. However, in the case with strong electron-phonon coupling, the pairing gap field is strongly ω dependent.

In the vicinity of the superconducting phase transition, the Cooper pair is very fragile and unstable because the binding energy is very small. In the physics of the superconducting pairing gap field $\Delta(\mathbf{q}, \nu; \mathbf{k}, \omega)$, this implies the non-trivial couplings of the center-of-mass and the inner-pair degrees of freedom due to finite energy transfer. Thus, the finite energy transfer can make the phase transition of the superconducting Cooper pairs different to the superfluidity phase transition of the well-defined bosonic particles, the latter of which have no relevant inner-particle structure. The superconducting phase transition may have unusual physics beyond the Ginzburg-Landau theory in especially the superconductor with strong inner-pair time-retarded physics.

As the center-of-mass and the inner-pair degrees of freedom of Cooper pairs also involve momenta, there is possible finite momentum transfer between these two degrees of freedom. The momentum transfer is one driven mechanism for the occurrence of the Fulde-Ferrell-Larkin-Ovchinnikov (FFLO) phase^{8,9} and the pair-density waves¹⁰. The finite energy and/or momentum transfer between the center-of-mass and the inner-pair degrees of freedom of Cooper pairs comes from the non conservation of the energy and the momentum within the pure center-of-mass channel.

In this manuscript, we will consider a simple case where the superconducting pairing gap field is \mathbf{k} independent and there is finite energy transfer between the center-of-mass and the inner-pair degrees of freedom of Cooper pairs. In this case, the pairing gap field $\Delta = \Delta(\mathbf{q}, \nu; \omega)$. As the macroscopic superconductivity is mostly relevant to the center-of-mass degrees of freedom, we should integrate out the inner-pair degrees of freedom of the pairing gap field. We introduce a renormalization formalism following the idea of the Poor man's renormalization scaling¹¹ to do the functional integral of the inner-pair frequency- ω relevant pairing gap field. This is schematically shown in Fig. 1.

Mathematically, our renormalization formalism can be

expressed as following:

$$\begin{aligned} Z &= \int \mathcal{D}[\Delta, \bar{\Delta}] e^{-S[\Delta, \bar{\Delta}]} \\ &= \int \mathcal{D}[\Delta_c, \bar{\Delta}_c] e^{-S_c[\Delta_c, \bar{\Delta}_c]}, \end{aligned} \quad (1)$$

where the original action $S[\Delta, \bar{\Delta}] = S[\Delta(\mathbf{q}, \nu; \omega), \bar{\Delta}(\mathbf{q}, \nu; \omega)]$ and the renormalized action $S_c[\Delta_c, \bar{\Delta}_c] = S_c[\Delta_c(\mathbf{q}, \nu), \bar{\Delta}_c(\mathbf{q}, \nu)]$ with $\Delta_c(\mathbf{q}, \nu) = \Delta(\mathbf{q}, \nu; \omega)|_{\omega \rightarrow 0}$ and $\bar{\Delta}_c(\mathbf{q}, \nu) = \bar{\Delta}(\mathbf{q}, \nu; \omega)|_{\omega \rightarrow 0}$. During the renormalization process, we integrate out the fields $\Delta(\mathbf{q}, \nu; \omega)$ and $\bar{\Delta}(\mathbf{q}, \nu; \omega)$ within $\delta\omega$ in high- ω range step by step until we arrive at $\Delta_c(\mathbf{q}, \nu)$ and $\bar{\Delta}_c(\mathbf{q}, \nu)$. We thus obtain an effective extended Ginzburg-Landau action S_c with the inner-pair physics mostly integrated out except the ones near the Fermi energy. The non-trivial couplings due to the energy transfer between the center-of-mass and the inner-pair degrees of freedom have been more exactly taken into account. Therefore, the extended Ginzburg-Landau action obtained from the renormalization formalism would include some unusual physics that have not been seriously considered previously in the superconducting phase transition.




FIG. 1: (Color online) Schematic illustration of the renormalization integral of the inner-pair frequency- ω relevant pairing gap field $\Delta(\mathbf{q}, \nu; \omega)$. The red-rectangle relevant fields with $\omega \in [-\frac{\Lambda_\omega}{2}, -\frac{\Lambda_\omega}{2} + \delta\omega]$ and $\omega \in [+ \frac{\Lambda_\omega}{2} - \delta\omega, + \frac{\Lambda_\omega}{2}]$ are first integrated out. The following renormalization process is then done with further $\delta\omega$ rectangles integrated out until we arrive at $\Delta_c(\mathbf{q}, \nu) = \Delta(\mathbf{q}, \nu; \omega)|_{\omega \rightarrow 0}$. Here we shift the inner-pair energy by the Fermi energy.

It should be noted that we have chosen $\Delta_c(\mathbf{q}, \nu)$ and $\bar{\Delta}_c(\mathbf{q}, \nu)$ as the basic variables to describe the superconducting phase transition. This is a theoretical assumption when we focus mainly on the critical superconducting phase transition, where we have assumed that all of the high- ω relevant pairing gap fields are irrelevant to the critical superconducting phase transition. This is different to the definitions $\Delta'_c(\mathbf{q}, \nu) = \sum_{\mathbf{k}\omega} \Delta(\mathbf{q}, \nu; \mathbf{k}, \omega)$ and $\bar{\Delta}'_c(\mathbf{q}, \nu) = \sum_{\mathbf{k}\omega} \bar{\Delta}(\mathbf{q}, \nu; \mathbf{k}, \omega)$, which were introduced by Abrahams and Tsuneto¹² and Gor'kov and Eliashberg¹³ for the non-stationary problems in superconductor. Since $\Delta'_c(\mathbf{q}, \nu) = \Delta(\mathbf{q}, \nu; \mathbf{r}, t)|_{\mathbf{r}=0, t=0}$ and

$\bar{\Delta}'_c(\mathbf{q}, \nu) = \bar{\Delta}(\mathbf{q}, \nu; \mathbf{r}, t)|_{\mathbf{r}=0, t=0}$, they are only well defined for the superconductivity in the ideal case where the attractive pairing interaction is momentum independent and instantaneous without time retarded effects. $\Delta'_c(\mathbf{q}, \nu)$ and $\bar{\Delta}'_c(\mathbf{q}, \nu)$ can be introduced as approximate variables for the superconducting condensate far away from the superconducting phase transition regime, where most of the fermionic excitations are gapped and the center-of-mass and the inner-pair degrees of freedom of the pairing gap fields can be approximately decoupled. This is not the case we will consider in this manuscript.

The manuscript is arranged as below. In Sec. II, we present the action near T_c to fourth order of the pairing gap fields. In Sec. III, we provide a theoretical renormalization formalism to derive an extended Ginzburg-Landau action for the superconductor with non-trivial couplings of the center-of-mass and the inner-pair degrees of freedom of Cooper pairs due to the finite energy transfer between them. Discussion and summary are presented in Sec. IV.

II. ACTION NEAR T_c

In this section we will present a general form of the action for the superconducting phase transition near T_c by using the standard path-integral field theory¹⁴. We assume an attractive pairing interaction which is both space and time dependent, $\hat{V} = -V(x_1 - x_2)$, where $x_l = (\mathbf{r}_l, \tau_l)$, $l = 1, 2$. Here the position vector \mathbf{r} can be two-dimensional (2D) or three-dimensional (3D) and τ is an imaginary time. A minus sign “-” has been explicitly shown out for the attractive character of the pairing interaction.

We consider the following partition function

$$Z = \int \mathcal{D}[c, \bar{c}] e^{-S[c, \bar{c}]}, \quad (2)$$

where the action S is defined as

$$\begin{aligned} S[c, \bar{c}] &= \int dx \sum_{\sigma} \bar{c}_{\sigma}(x) (\partial_{\tau} + H_0) c_{\sigma}(x) \\ &\quad - \int dx_1 dx_2 \bar{c}_{\uparrow}(x_1) \bar{c}_{\downarrow}(x_2) V(x_1 - x_2) c_{\downarrow}(x_2) c_{\uparrow}(x_1). \end{aligned} \quad (3)$$

Here c and \bar{c} are the fermionic Grassmann fields for the electrons, σ is the electron spin defined as $\sigma = \uparrow, \downarrow$. $\int dx = \int_0^{\beta} d\tau d\mathbf{r}$ with $\beta = \frac{1}{k_B T}$ (T is temperature).

Introduce two Hubbard-Stratonovich (HS) fields $\Delta(x_1, x_2)$ and $\bar{\Delta}(x_1, x_2)$, the action can be expressed into the following form:

$$\begin{aligned} S &= \int dx_1 dx_2 \left\{ \bar{\Psi}(x_1) [-g^{-1}(x_1, x_2)] \Psi(x_2) \right. \\ &\quad \left. + \bar{\Delta}(x_1, x_2) V(x_1 - x_2)^{-1} \Delta(x_1, x_2) \right\}. \end{aligned} \quad (4)$$

Here the Nambu spinor field is defined as

$$\Psi(x) = \begin{bmatrix} c_{\uparrow}(x) \\ \bar{c}_{\downarrow}(x) \end{bmatrix}, \quad (5)$$

and the Green's function matrix is given by

$$-g^{-1}(x_1, x_2) = -g_0^{-1}(x_1, x_2) + \widehat{\Delta}(1, 2), \quad (6)$$

where

$$g_0^{-1}(x_1, x_2) = \begin{bmatrix} g_{0\uparrow}^{-1}(x_1, x_2) & 0 \\ 0 & g_{0\downarrow}^{-1}(x_1, x_2) \end{bmatrix}, \quad (7)$$

$$\widehat{\Delta}(x_1, x_2) = \begin{bmatrix} 0 & \Delta(x_1, x_2) \\ \bar{\Delta}(x_2, x_1) & 0 \end{bmatrix}. \quad (8)$$

Here $g_{0\uparrow}^{-1}(x_1, x_2) = -[\partial_{\tau_1} + H_0(x_1)]\delta(x_1 - x_2)$ and $g_{0\downarrow}^{-1}(x_1, x_2) = -[\partial_{\tau_1} - H_0(x_1)]\delta(x_1 - x_2)$. After integrate out the fermionic Grassmann fields c and \bar{c} , we have an action for the HS pairing gap fields Δ and $\bar{\Delta}$ as following:

$$S[\Delta, \bar{\Delta}] = \sum_{n=1} \frac{1}{n} \text{Tr}[(g_0 \widehat{\Delta})^n] + \int dx_1 dx_2 \bar{\Delta}(x_1, x_2) V(x_1 - x_2)^{-1} \Delta(x_1, x_2). \quad (9)$$

Here Tr operation acts on both the spatial-temporal and the spin spaces.

To fourth order of the HS pairing gap fields Δ and $\bar{\Delta}$, the action can be shown to follow

$$S = S^{(2)} + S^{(4)}, \quad (10)$$

where

$$S^{(2)} = \sum_{1,2,p} \bar{\Delta}(1+p, -2-p) [G_0^{(-1)} \delta_{p,0} + V^{-1}(p)] \Delta(1, -2),$$

$$S^{(4)} = \sum_{1,2,3,4} \frac{b_0}{2} \bar{\Delta}(1, -2) \Delta(1, -4) \bar{\Delta}(3, -4) \Delta(3, -2). \quad (11)$$

Here the simplified notations are defined as

$$l \equiv (\mathbf{k}_l, i\omega_l), \quad l = 1, 2, 3, 4, \quad (12)$$

$p = (\mathbf{p}, ip_n)$ with $ip_n = i\frac{2n\pi}{\beta}$ being a bosonic imaginary frequency. $G_0^{-1} = G_0^{-1}(1, 2)$ and $b_0 = b_0(1, 2, 3, 4)$, which are defined by

$$G_0^{-1}(1, 2) = \frac{1}{(i\omega_1 - \varepsilon_{\mathbf{k}_1})(i\omega_2 + \varepsilon_{-\mathbf{k}_2})}, \quad (13)$$

$$b_0(1, 2, 3, 4) = G_0^{-1}(1, 2) \cdot G_0^{-1}(3, 4). \quad (14)$$

$V^{-1}(p)$ is the Fourier transformation of $V(x_1 - x_2)^{-1}$. In (11), the notations $(1+p)$ and $(-2-p)$ are defined as

$$(1+p) \equiv (\mathbf{k}_1 + \mathbf{p}, i\omega_2 + ip_n),$$

$$(-2-p) \equiv (-\mathbf{k}_2 - \mathbf{p}, -i\omega_2 - ip_n). \quad (15)$$

It is noted that we have introduced the following Fourier transformations in the derivation of (11): $\Delta(x_1, x_2) = \frac{1}{\beta V_0} \sum_{1,2} \Delta(1, -2) e^{i(1 \cdot x_1) - i(2 \cdot x_2)}$ where $i(1 \cdot x_1) - i(2 \cdot x_2) \equiv i\mathbf{k}_1 \cdot \mathbf{r}_1 - i\omega_1 \tau_1 - i\mathbf{k}_2 \cdot \mathbf{r}_2 + i\omega_2 \tau_2$ and V_0 is the volume of the system, and $V(x_1, x_2)^{-1} = \sum_p V^{-1}(p) e^{ip \cdot (x_1 - x_2)}$ where $ip \cdot (x_1 - x_2) \equiv i\mathbf{p} \cdot (\mathbf{r}_1 - \mathbf{r}_2) - ip_n(\tau_1 - \tau_2)$.

Introduce the following simplified notations

$$l_c = (\mathbf{q}_{c_l}, i\nu_{c_l}), \quad l_r = (\mathbf{k}_{r_l}, i\omega_{r_l}), \quad l = 1, 2, 3, 4, \quad (16)$$

where the subscript c denotes the center-of-mass degrees of freedom of the pairing gap fields, and r denotes the inner-pair ones. These notations are defined by

$$1_c = [1 - 2], 1_r = \frac{1}{2}[1 + 2],$$

$$3_c = [3 - 4], 3_r = \frac{1}{2}[3 + 4],$$

$$2_c = \frac{1}{2}[1_c + 3_c] + [1_r - 3_r],$$

$$2_r = \frac{1}{4}[1_c - 3_c] + \frac{1}{2}[1_r + 3_r], \quad (17)$$

$$4_c = \frac{1}{2}[1_c + 3_c] - [1_r - 3_r],$$

$$4_r = -\frac{1}{4}[1_c - 3_c] + \frac{1}{2}[1_r + 3_r].$$

Here the algebra equations of the notations l of (12) and l_c of (16) are defined by the following rules:

$$[1 + 2] \equiv (\mathbf{k}_1 + \mathbf{k}_2, i\omega_1 + i\omega_2),$$

$$[1_c + 3_c] \equiv (\mathbf{q}_{c_1} + \mathbf{q}_{c_3}, i\nu_{c_1} + i\nu_{c_3}), \quad (18)$$

$$\frac{1}{2}[1_r + 3_r] \equiv (\frac{1}{2}(\mathbf{k}_{r_1} + \mathbf{k}_{r_3}), \frac{1}{2}(i\omega_{r_1} + i\omega_{r_3})).$$

Following (11) with the new notations, we can reexpress the action $S = S^{(2)} + S^{(4)}$ into the following form:

$$S^{(2)} = \sum_{1_c 1_r p} \bar{\Delta}(1_c, 1_r + p) [G_0^{-1} \delta_{p,0} + V^{-1}(p)] \Delta(1_c, 1_r),$$

$$S^{(4)} = \sum_{l_c l_r} \frac{b_0}{2} \bar{\Delta}(1_c, 1_r) \Delta(2_c, 2_r) \bar{\Delta}(3_c, 3_r) \Delta(4_c, 4_r). \quad (19)$$

Here G_0^{-1} and b_0 are given in (13) and (14). It can be easily seen that the center-of-mass and the inner-pair degrees of freedom of the pairing gap fields are coupled in $S^{(4)}$ but decoupled in $S^{(2)}$, the latter feature has been found previously^{15,16}.

From this action, the saddle-point equation of the pairing gap fields can be obtained from $\frac{\delta S}{\delta \bar{\Delta}(1_c, 1_r)} = 0$, which leads us

$$\sum_p [G_0^{-1} \delta_{p,0} + V^{-1}(p)] \Delta(1_c, 1_r - p) + \sum_{l_c l_r} b_0 \Delta(2_c, 2_r) \bar{\Delta}(3_c, 3_r) \Delta(4_c, 4_r) = 0. \quad (20)$$

It should be noted that when we introduce the center-of-mass and the inner-pair imaginary frequencies $i\nu_{c_l}$ and $i\omega_{r_l}$, the requirement that $i\omega_{r_l}$ are the fermionic imaginary frequencies leads us a constraint for $i\nu_{c_l}$:

$$i\nu_{c_l} = \frac{i2n_{c_l}\pi}{\beta}, \quad n_{c_l} = \text{even integer.} \quad (21)$$

III. RENORMALIZATION FORMALISM FOR AN EXTENDED GINZBURG-LANDAU ACTION

The well-known Ginzburg-Landau action can be obtained from (19) for the simplified case where the attractive pairing interaction is space-time independent, *e.g.*, $V^{-1}(p) = V_g^{-1}\delta_{p,0}$. The detailed investigations are present in Appendix A for the 2D and 3D superconductors.

Let us now consider the case where the role of the attractive pairing interaction in the pairing gap fields is mainly time dynamics but momentum irrelevant. Examples of such attractive pairing interaction can be the electron-phonon induced one or the critical-mode induced one, the former of which is ubiquitous in the traditional metal superconductors and the latter is studied recently by Chubukov *et al.* in their γ model for the novel superconductivity in quantum critical metals¹⁷⁻²². One Einstein-phonon induced pairing interaction can be defined as

$$V^{-1}(ip_n) = -\frac{\beta}{N_\beta g_{ep}} \left[\frac{(ip_n)^2 - \omega_0^2}{\omega_0} \right], \quad (22)$$

where ω_0 is the Einstein-phonon frequency and $g_{ep} \simeq 1(eV)^2$ with reference to the data of boron-doped diamond²³⁻²⁵. Here N_β is the imaginary-frequency number. In the γ model, the pairing interaction has a form as¹⁷⁻²²

$$V^{-1}(ip_n) = g_p^{-1}|p_n|^\gamma, \quad (23)$$

where g_p is an interaction constant with γ dependent dimension.

A. Action for renormalization

With the momentum irrelevant attractive pairing interaction, the superconducting pairing gap fields will also be inner-pair momentum \mathbf{k}_r irrelevant, *i.e.*,

$$\Delta(1_c, 1_r) = \Delta(1_c, i\omega_{r_1}), \quad \bar{\Delta}(1_c, 1_r) = \bar{\Delta}(1_c, i\omega_{r_1}). \quad (24)$$

After do the summation of the inner-pair momenta, we show that the action of (19) is modified into the form:

$$S^{(2)} = \sum_{1_c 1_r p_n} \bar{\Delta}(1_c, 1_r + ip_n) [\mathcal{G}_0^{-1}\delta_{p,0} + V^{-1}(ip_n)] \Delta(1_c, 1_r),$$

$$S^{(4)} = \sum_{l_c l_r} \frac{\mathcal{B}}{2} \bar{\Delta}(1_c, 1_r) \Delta(2_c, 2_r) \bar{\Delta}(3_c, 3_r) \Delta(4_c, 4_r). \quad (25)$$

Here \mathcal{G}_0^{-1} and \mathcal{B} are defined as

$$\mathcal{G}_0^{-1}(1_c, 1_r) = -\frac{i}{2} N \rho_{2D} F(\nu_{c_1}, \omega_{r_1}) \left\langle \frac{1}{i\omega_{r_1} - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}}} \right\rangle,$$

$$\mathcal{B}(l_c, l_r) = \mathcal{G}_0^{-1}(1_c, 1_r) \cdot \mathcal{G}_0^{-1}(3_c, 3_r), \quad (26)$$

where ρ_{2D} is the 2D density of states at the Fermi energy as defined in (A6), and $F(\nu_{c_1}, \omega_{r_1})$ is defined by

$$F(\nu_{c_1}, \omega_{r_1}) = \theta(|\omega_{r_1}| - \frac{1}{2}|\nu_{c_1}|) \cdot \text{sgn}(\omega_{r_1}). \quad (27)$$

Here $\theta(x)$ is the step function and $\text{sgn}(x)$ is the sign function. $A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}} = \frac{\hbar^2}{2m} \mathbf{k}_F \cdot \mathbf{q}_{c_1}$, and the average is defined as

$$\left\langle \frac{1}{i\omega_{r_1} - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}}} \right\rangle = \int_0^{2\pi} d\theta \left[\frac{1}{i\omega_{r_1} - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}}} \right], \quad (28)$$

with θ being the angle between \mathbf{k}_F and \mathbf{q}_{c_1} . Here we have assumed that the energy dispersion near the Fermi energy follows $\varepsilon_{\mathbf{k}} = \frac{\hbar^2 \mathbf{k}^2}{2m} - \mu_F$.

The action in (25) is defined in the imaginary-frequency space. It can be transformed into the real-frequency space, which becomes $S = S^{(2)} + S^{(4)}$ with

$$S^{(2)} = \sum_{1_c 1_r \nu_p} \bar{\Delta}(1_c, 1_r + \nu_p) [\mathcal{G}_0^{-1}\delta_{p,0} + V^{-1}(\nu_p)] \Delta(1_c, 1_r),$$

$$S^{(4)} = \sum_{l_c l_r} \frac{\mathcal{B}}{2} \bar{\Delta}(1_c, 1_r) \Delta(2_c, 2_r) \bar{\Delta}(3_c, 3_r) \Delta(4_c, 4_r). \quad (29)$$

Here the simplified notations in the real-frequency action (29) are similar to that in (16) with all the imaginary frequencies changed into the corresponding real frequencies by the following rules:

$$l_c = (\mathbf{q}_{c_l}, i\nu_{c_l} \rightarrow \nu_{c_l}), l_r = (i\omega_{r_l} \rightarrow \omega_{r_l}), l = 1, 2, 3, 4. \quad (30)$$

\mathcal{G}_0^{-1} and \mathcal{B} are defined as

$$\mathcal{G}_0^{-1}(1_c, 1_r) = \alpha_G \left[n_F(\omega_{r_1} - \frac{1}{2}\nu_{c_1}) \left\langle \frac{1}{\omega_{c_1} - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}} + i\delta^+} \right\rangle + n_F(\omega_{r_1} + \frac{1}{2}\nu_{c_1}) \left\langle \frac{1}{\omega_{c_1} - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}} - i\delta^+} \right\rangle \right],$$

$$\mathcal{B}(l_c, l_r) = \mathcal{G}_0^{-1}(1_c, 1_r) \cdot \mathcal{G}_0^{-1}(3_c, 3_r), \quad (31)$$

where

$$\alpha_G = \frac{\beta N \rho_{2D} \delta \omega_r}{4\pi}. \quad (32)$$

Here $\delta \omega_r$ defines the step value for the real-frequency summation, i.e., $\sum_{\omega_r} \delta \omega_r = \int_{-\Lambda_\omega/2}^{+\Lambda_\omega/2} d\omega_r$.

In the real-frequency action (29), $V^{-1}(ip_n \rightarrow \nu_p)$ with ν_p a continuous real frequency. A detailed derivation of the transformation of the imaginary-frequency action into the real-frequency one is given in Appendix B.

B. Renormalization formalism

In the above section, we have obtained the action of the pairing gap fields which involve both the center-of-mass and the inner-pair degrees of freedom of Cooper pairs. As the macroscopic superconductivity is mostly relevant to the center-of-mass degrees of freedom, we will integrate out the inner-pair ones to derive an extended Ginzburg-Landau action for the macroscopic superconductivity. Because of the finite energy transfer between the center-of-mass and the inner-pair degrees of freedom of Cooper pairs, there are non-trivial couplings between these two different degrees of freedom. We will introduce a renormalization formalism to do the functional integral of the inner-pair frequency relevant pairing gap fields following the idea of the Poor man's renormalization scaling¹¹. The schematic illustration of the renormalization formalism is presented in Fig. 1. By this renormalization formalism, most of the inner-pair degrees of freedom of Cooper pairs except the ones near the Fermi energy can be integrated out and the unusual effects of the inner-pair dynamical physics on the superconducting phase transition can be more exactly taken into account.

Introduce the following simplified vector, matrix and tensor notations:

$$\begin{aligned} m_l &\equiv (l_c, l_r) = (\mathbf{q}_{c_l}, \nu_{c_l}; \omega_{r_l}), \\ \mathcal{G}_{0:m_1 m_1}^{-1} &\equiv \mathcal{G}_0^{-1}(1_c, 1_r), \\ [V^{-1}]_{m_1 m_2} &\equiv V^{-1}(\omega_{r_1} - \omega_{r_2}), \\ \mathcal{B}_{m_1 m_2 m_3 m_4} &\equiv \mathcal{B}(1_c, 1_r; 2_c, 2_r; 3_c, 3_r; 4_c, 4_r). \end{aligned} \quad (33)$$

Separate the pairing gap fields into two types,

$$\begin{aligned} (\Delta, \bar{\Delta}) &\text{ for } S_{<}, \\ (\Delta', \bar{\Delta}') &\text{ for } S_{>}, \end{aligned} \quad (34)$$

where $S_{<}$ and $S_{>}$ are the renormalized and the integrated parts of the action in the renormalization process, respectively. Similarly, we introduce the following notations for $(\Delta, \bar{\Delta})$ and $(\Delta', \bar{\Delta}')$:

$$\begin{aligned} \mathcal{G}^{-1} &= \mathcal{G}_0^{-1} + V^{-1} \quad \text{for } (\Delta, \bar{\Delta}), \\ \mathcal{G}'^{-1} &= \mathcal{G}'_0^{-1} + V'^{-1} \quad \text{for } (\Delta', \bar{\Delta}'). \end{aligned} \quad (35)$$

With these notations, the partition function can be calculated as following:

$$Z = \int \mathcal{D}[\Delta, \bar{\Delta}; \Delta', \bar{\Delta}'] e^{-[S_{<}^{(2)} + S_{<}^{(4)} + S_{>}^{(2)} + S_{>}^{(2)} + S_{>}^{(4)}]}, \quad (36)$$

where

$$\begin{aligned} S_{<}^{(2)} &= \bar{\Delta} \mathcal{G}^{-1} \Delta, \\ S_{<}^{(4)} &= \frac{1}{2} \mathcal{B} \bar{\Delta} \Delta \bar{\Delta} \Delta, \\ S_{>}^{(2)} + S_{>}^{(2)} &= \bar{\Delta}' \mathcal{G}'^{-1} \Delta' + \bar{\Delta}' V^{-1} \Delta + \bar{\Delta} V^{-1} \Delta', \\ S_{>}^{(4)} &= \bar{\Delta}' \hat{\Phi}(\Delta, \bar{\Delta}) \Delta'. \end{aligned} \quad (37)$$

Here we have ignored the contribution from $S_{>}^{(4)}$. All the expressions in (37) are in vector-matrix-tensor forms with the algebra rules defined such as $\bar{\Delta} \mathcal{G}^{-1} \Delta = \sum_{m_1 m_2} \bar{\Delta}_{m_1} \mathcal{G}_{m_1 m_2}^{-1} \Delta_{m_2}$ and $\mathcal{B} \bar{\Delta} \Delta \bar{\Delta} \Delta = \sum_{m_i} \mathcal{B}_{m_1 m_2 m_3 m_4} \bar{\Delta}_{m_1} \Delta_{m_2} \bar{\Delta}_{m_3} \Delta_{m_4}$. $\hat{\Phi}(\Delta, \bar{\Delta})$ is defined as

$$[\hat{\Phi}(\Delta, \bar{\Delta})]_{m'_1 m'_2} = \sum_{m_1 m_2} \mathcal{B}_{m_1 m_2 m'_1 m'_2} \bar{\Delta}_{m_1} \Delta_{m_2}. \quad (38)$$

When integrate out the pairing gap fields Δ' and $\bar{\Delta}'$, we can obtain an effective renormalized partition function as

$$Z^{(RG)} = \int \mathcal{D}[\Delta, \bar{\Delta}] e^{-S_{<}^{(RG)}[\Delta, \bar{\Delta}]}, \quad (39)$$

where the renormalized action follows

$$\begin{aligned} S_{<}^{(RG)} &= S_{<}^{(2)} + S_{<}^{(4)} + \text{Tr}[\mathcal{G}' \hat{\Phi}] - \frac{1}{2} \text{Tr}[(\mathcal{G}' \hat{\Phi})^2] \\ &\quad - \bar{\Delta} V^{-1} \mathcal{G}' (1 - \hat{\Phi} \mathcal{G}') V^{-1} \Delta. \end{aligned} \quad (40)$$

Here the expansion to higher than fourth order of Δ and $\bar{\Delta}$ is neglected. A detailed expansion of the renormalized action leads us its following form:

$$S_{<}^{(RG)} = \bar{\Delta} \mathcal{G}_R^{-1} \Delta + \frac{1}{2} \mathcal{B}_R \bar{\Delta} \Delta \bar{\Delta} \Delta, \quad (41)$$

where the renormalized \mathcal{G}_R^{-1} and \mathcal{B}_R follow

$$\begin{aligned}\mathcal{G}_R^{-1} &= \mathcal{G}^{-1} + \mathcal{A}^{(1)} + \mathcal{A}^{(2)}, \\ \mathcal{B}_R &= \mathcal{B} + \mathcal{B}^{(1)} + \mathcal{B}^{(2)}.\end{aligned}\quad (42)$$

$\mathcal{A}^{(n)}$ and $\mathcal{B}^{(n)}$ with $n = 1, 2$ are defined by

$$\begin{aligned}\mathcal{A}_{m_1 m_2}^{(1)} &= \sum_{m'_1 m'_2} \mathcal{G}'_{m'_1 m'_2} \mathcal{B}_{m_1 m_2 m'_2 m'_1}, \\ \mathcal{A}_{m_1 m_2}^{(2)} &= - \sum_{m'_1 m'_2} [V^{-1}]_{m_1 m'_1} \mathcal{G}'_{m'_1 m'_2} [V^{-1}]_{m'_2 m_2},\end{aligned}\quad (43)$$

and

$$\begin{aligned}\mathcal{B}_{m_1 m_2 m_3 m_4}^{(1)} &= - \sum_{m'_i} \mathcal{G}'_{m'_i m'_i} \mathcal{B}_{m_1 m_2 m'_2 m'_1} \\ &\quad \times \mathcal{G}'_{m'_3 m'_4} \mathcal{B}_{m_3 m_4 m'_4 m'_3}, \\ \mathcal{B}_{m_1 m_2 m_3 m_4}^{(2)} &= \sum_{m'_i} 2[V^{-1}]_{m_1 m'_1} \mathcal{G}'_{m'_1 m'_2} \mathcal{B}_{m_3 m_2 m'_2 m'_3} \\ &\quad \times \mathcal{G}'_{m'_3 m'_4} [V^{-1}]_{m'_4 m_4}.\end{aligned}\quad (44)$$

The schematic illustration of the above renormalization process has been shown in Fig. 1. Step by step we can integrate out the inner-pair high-frequency pairing gap fields and finally, we arrive at $\Delta_c(\mathbf{q}_c, \nu_c) = \Delta(\mathbf{q}_c, \nu_c; \omega_r)|_{\omega_r \rightarrow 0}$ and $\bar{\Delta}_c(\mathbf{q}_c, \nu_c) = \bar{\Delta}(\mathbf{q}_c, \nu_c; \omega_r)|_{\omega_r \rightarrow 0}$. In numerical study, we can set $\Delta_c(\mathbf{q}, \nu) = \sum_{\omega} \Delta(\mathbf{q}, \nu; \omega)|_{\omega \in \delta_{\Lambda}}$ and $\bar{\Delta}_c(\mathbf{q}, \nu) = \sum_{\omega} \bar{\Delta}(\mathbf{q}, \nu; \omega)|_{\omega \in \delta_{\Lambda}}$. Here δ_{Λ} is an infrared energy cut-off for the classical or quantum superconducting phase transition. It can be set as the larger of $k_B T_c$ and δ_E , the latter of which is a characteristic infrared energy scale for the quantum superconducting phase transition. The partition function after the final renormalization step follows

$$Z_c = \int \mathcal{D}[\Delta_c, \bar{\Delta}_c] e^{-S_c[\Delta_c, \bar{\Delta}_c]}, \quad (45)$$

where the renormalized action S_c has a similar form as $S_{<}^{(RG)}$ of (41):

$$S_c = \bar{\Delta}_c \mathcal{G}_R^{-1} \Delta_c + \frac{1}{2} \mathcal{B}_R \bar{\Delta}_c \Delta_c \bar{\Delta}_c \Delta_c. \quad (46)$$

Here \mathcal{G}_R^{-1} and \mathcal{B}_R are set by the final renormalization step.

Compared with the Ginzburg-Landau action (A3) without inner-pair structure, the renormalized action S_c involves the renormalization effects from the couplings of the center-of-mass and the inner-pair degrees of freedom due to finite energy transfer. Therefore, both the superconducting phase transition temperature T_c and the long-wavelength low-energy fluctuations should be renormalized by the inner-pair dynamics. For example, the diagonal part of \mathcal{G}_R^{-1} would renormalize T_c into T_c^* , the latter of which is defined by

$$\mathcal{G}_R^{-1}(\mathbf{q}_c, \nu_c; T_c^*)|_{\mathbf{q}_c=0, \nu_c=0} = 0. \quad (47)$$

The low-energy dynamical fluctuations and the long-wavelength spatial fluctuations of the center-of-mass degrees of freedom, shown in the respective a_1 and a_2 terms of (A3), would also have finite renormalization effects.

It should be noted that all of these renormalization effects from the inner-pair degrees of freedom of Cooper pairs are only part of the whole renormalization effects, as there are other renormalization effects which should be included further for the superconducting phase transition. The latter comes from the center-of-mass relevant fluctuations, the spatial fluctuations and the dynamical and quantum fluctuations, as have been described by the Wilson-Hertz-Millis theories for the critical phase transitions²⁶⁻²⁸. *Therefore, a complete and exact renormalization theory for the superconducting phase transition with non-trivial couplings of the center-of-mass and the inner-pair degrees of freedom of Cooper pairs would involve two renormalization processes, the first one is that we present in this manuscript and the second one is the Wilson-Hertz-Millis renormalization.* The renormalized action (46) obtained in the first renormalization process is a starting point for the second renormalization process. As the Wilson-Hertz-Millis renormalization theories have been well established²⁶⁻³⁰, we will not discuss the Wilson-Hertz-Millis renormalization further in this manuscript, with the relevant renormalization process to be done in future work. Due to the limited memory space of our computers, we can not make a reliable numerical study on the extended Ginzburg-Landau action at present. The numerical investigation on the complete and exact two-process renormalization for the superconducting phase transition will be one task in our future plan.

IV. DISCUSSION AND SUMMARY

The critical phase transition is one long-standing subject in the modern condensed matter field^{17-22,31-38}. In general, it involves two types of problems, one is the microscopic driving mechanism of the critical phase transition and the other is the relevant critical phenomena. In the case of the superconducting phase transition, the microscopic driving mechanism is how the Cooper pairs form microscopically and how the Cooper pairs condense coherently into a macroscopic superconducting state. The issues relevant to the superconducting critical phenomena, similar to other phase transition critical phenomena, mainly focus on the physical effects of the critical fluctuations on thermodynamics, charge transport, magnetic response, etc.

There are two relevant particles in the superconducting phase transition, the Cooper-pair composite particles and the component electrons. However, there are three different degrees of freedom relevant to these two different particles, the center-of-mass and the inner-pair degrees of freedom of the composite Cooper pairs, and the ones of the component electrons. The macro-

scopic Ginzburg-Landau theory¹ and the microscopic BCS theory²⁻⁴ are two main starting points to study the superconducting phase transition. Only the center-of-mass degrees of freedom of the Cooper pairs are relevant in the macroscopic Ginzburg-Landau theory. The BCS theory focuses mainly on the microscopic formation of Cooper pairs, where the inner-pair degrees of freedom of Cooper pairs and the component electrons are involved. In most studies of the superconducting critical phenomena, the superconducting critical fluctuations are assumed to come from bosonic excitations which have no inner structure^{17-22,37,38}. Because the composite Cooper pairs are not rigid point-like particles but have non-trivial inner-pair structure, these theoretical treatments are incomplete and inaccurate in principle for the superconducting phase transition.

In this manuscript, we have presented a theoretical renormalization formalism to derive an extended Ginzburg-Landau action for the superconductor with non-trivial couplings of the center-of-mass and the inner-pair degrees of freedom of Cooper pairs. A following task is to develop a two-process renormalization formalism for the superconducting phase transition. One renormalization process is for the non-trivial couplings of the center-of-mass and the inner-pair degrees of freedom as we have presented in this manuscript, and the other one is for the critical fluctuations in thermal, spatial, dynamical and quantum channels of the center-of-mass degrees of freedom of Cooper pairs, which have been well described by the Wilson-Hertz-Millis theories²⁶⁻²⁸. It should be noted that a complete and well-defined theory for the critical superconducting phase transition should also include the degrees of freedom of the component electrons, which are important and indispensable for the superconducting phase transition and the critical phenomena. How to take into account all the physics of the composite Cooper pairs and the component electrons in the superconducting phase transition is one big issue in the modern condensed matter field.

The renormalization formalism we have presented in the manuscript for the superconductor with non-trivial inner-pair structure may provide a good theoretical tool to study the other phase transitions with composite particles involved. One example is the itinerant magnetic phase transition³⁹, where the itinerant magnetic moment can be regarded from the composite particle in the particle-hole spin channel. Another example is the d-wave bond electronic nematicity^{40,41}, where the d-wave bond nematic order can be regarded as of one composite particle in the particle-hole charge channel. We argue that this renormalization formalism can also be used to study the non-trivial couplings between the center-of-mass of hadron and its sub-particles, quarks and gluons.

In summary, we have presented a theoretical renormalization formalism to derive an extended Ginzburg-Landau action for the superconductor with non-trivial couplings of the center-of-mass and the inner-pair degrees of freedom of Cooper pairs due to finite energy transfer.

The inner-pair dynamical physics of Cooper pairs can be more exactly taken into account in the superconducting phase transition. This extended Ginzburg-Landau action is a starting point for the further renormalization process for the critical superconducting phase transition, which involves further the thermal, the spatial and the dynamical and quantum fluctuations relevant to the center-of-mass degrees of freedom of Cooper pairs. The renormalization formalism we have presented is also a good theoretical tool to study the strong couplings of the other composite particles and their component sub-particles.

ACKNOWLEDGMENTS

We thank Prof. Jun Chang, Prof. Miao Gao, Prof. Yunan Yan and Prof. Tao Li for invaluable discussions. This work was supported by the National Natural Science Foundation of China (Grants No. 11774299 and No. 11874318) and the Natural Science Foundation of Shandong Province (Grants No. ZR2017MA033 and No. ZR2018MA043).

Appendix A: Ginzburg-Landau action

In this Appendix section, we will present the Ginzburg-Landau action for the superconductor with a space-time independent attractive pairing interaction

$$V^{-1}(p) = V_g^{-1} \delta_{p,0}. \quad (\text{A1})$$

In this case, the pairing gap fields will have no inner-pair structure, i.e.,

$$\Delta(1_c, 1_r) = \Delta(1_c), \quad \bar{\Delta}(1_c, 1_r) = \bar{\Delta}(1_c). \quad (\text{A2})$$

Let us first consider the 2D superconductor. By a lengthy and standard derivation, we can show that the Ginzburg-Landau action in the long-wavelength approximation with $|\mathbf{q}_{c1}| \ll 1$ follows

$$S/\beta = \sum_{1_c} [a_0 \cdot \delta\bar{T} + a_1(i\nu_{c1}) + a_2 \cdot \mathbf{q}_{c1}^2] |\Delta(1_c)|^2 + \sum_{1_c} \frac{b}{2} |\Delta(1_c)|^4, \quad (\text{A3})$$

where $\delta\bar{T} = \frac{T-T_c}{T_c}$. The parameters in (A3) are given by

$$\begin{aligned} a_0 &= N\rho_{2D}, \\ a_2 &= \frac{N\rho_{2D}\mu_F\gamma}{16(k_B T_c)^2 m}, \\ b &= \frac{N\rho_{2D}\gamma}{8(k_B T_c)^2}, \end{aligned} \quad (\text{A4})$$

and the function $a_1(i\nu_{c1})$ is defined by

$$a_1(i\nu_{c1}) = N\rho_{2D} I(n_{c1}) \Big|_{n_{c1} = \frac{\nu_{c1}}{4\pi k_B T_c}}. \quad (\text{A5})$$

Here N is the number of the momenta in the first Brillouin zone, $\mu_F = \frac{\hbar^2 \mathbf{k}_F^2}{2m}$ with m being the electron mass and \mathbf{k}_F the Fermi momentum, and ρ_{2D} is the density of states at the Fermi energy defined as

$$\rho_{2D} = \frac{ma^2}{2\pi\hbar^2}, \quad (\text{A6})$$

where a is the lattice constant.

The parameter γ in (A4) comes from the following integrals:

$$\begin{aligned} \gamma_1 &= \int_0^{+\infty} dx \frac{1}{x^2} \left[\frac{\tanh(x)}{x} - \frac{1}{\cosh^2(x)} \right], \\ \gamma_2 &= \int_0^{+\infty} dx \left[\frac{\tanh(x)}{x \cosh^2(x)} \right], \\ \gamma_3 &= \int_0^{+\infty} dx \left[-\frac{\tanh(x)}{x^3} + \frac{1}{x^2 \cosh^2(x)} + \frac{2 \tanh(x)}{x \cosh^2(x)} \right], \end{aligned} \quad (\text{A7})$$

and

$$\gamma = \gamma_1 = \gamma_2 = \gamma_3 = 0.8526. \quad (\text{A8})$$

In (A5), $I(n_c)$ is defined as

$$I(n_c) = \sum_{n=0}^{|n_c|-1} \frac{1}{n + \frac{1}{2}} = \Psi_d\left(\frac{1}{2} + |n_c|\right) - \Psi_d\left(\frac{1}{2}\right), \quad (\text{A9})$$

where $\Psi_d(z)$ is the digamma function. $I(0) = 0$, and for large n_c it can be shown from the properties of the digamma function that $I(n_c) = \ln(|n_c| + \frac{1}{2}) + c_0$ with $c_0 = 1.96351$.

T_c is the mean-field superconducting phase transition temperature defined by

$$\sum_{\mathbf{k}} \left[-\frac{1}{2\varepsilon_{\mathbf{k}}} \tanh\left(\frac{\varepsilon_{\mathbf{k}}}{2k_B T_c}\right) + g_0^{-1} \right] = 0. \quad (\text{A10})$$

Here g_0^{-1} is defined as

$$g_0^{-1} \equiv \frac{\omega_D}{\pi V_g}, \quad (\text{A11})$$

where ω_D is an energy cut-off for the momentum summation, such as the Debye frequency in the electron-phonon interaction driven superconductor. From (A10), T_c is shown to follow

$$k_B T_c = c_0 \omega_D e^{-\frac{1}{\rho^{(0)} g_0}}, \quad (\text{A12})$$

where $c_0 = 1.134$.

It should be noted that in the Ginzburg-Landau action (A3), we have only considered the fluctuations in thermal channel near T_c , the spatial fluctuations near $\mathbf{q}_c = 0$ and the finite dynamical fluctuations. In the $|\Delta|^4$ term we have ignored the \mathbf{q}_c and $i\nu_c$ fluctuations of b .

A detailed derivation of $a_1(i\nu_{c_1})$ is given as following. Consider $\Delta(\mathbf{q}_c, i\nu_c)$ with $\mathbf{q}_c = 0$ for the 2D superconductor. The dynamical fluctuations are described by $S^{(2)}/\beta = \sum_{i\nu_c} a_1(i\nu_c) |\Delta(0, i\nu_c)|^2$, where

$$a_1(i\nu_c) = \frac{1}{\beta} \sum_{\mathbf{k}, i\omega_r} [G_0^{-1} + V_g^{-1}]. \quad (\text{A13})$$

Here G_0^{-1} is defined by (13) with $i\omega_1 = i\omega_r + \frac{1}{2}i\nu_c$, $i\omega_2 = i\omega_r - \frac{1}{2}i\nu_c$ and $\mathbf{k}_1 = \mathbf{k}_2 = \mathbf{k}$. The summation over $i\omega_r$ can be shown to be

$$a_1(i\nu_c) = \sum_{\mathbf{k}} \left[\frac{\tanh\left(\frac{\beta\varepsilon_{\mathbf{k}}}{2}\right)}{2\varepsilon_{\mathbf{k}}} - \frac{\tanh\left(\frac{\beta\varepsilon_{\mathbf{k}}}{2}\right)}{2\varepsilon_{\mathbf{k}} - i\nu_c} \right]. \quad (\text{A14})$$

Here we have used Eq. (A10) to substitute g_0^{-1} for temperature near T_c .




FIG. 2: (Color online) The closed contour for the integral (A15)

Transform the summation of \mathbf{k} into a continuous energy integral and consider the limit $\frac{\omega_D}{2k_B T_c} \rightarrow +\infty$, $a_1(i\nu_c)$ can be reexpressed into a closed-contour integral form in the complex z plane:

$$a_1(i\nu_c) = \frac{N\rho_{2D}}{2} \oint_{ABCD} dz \left[\frac{\tanh(z)}{z} \right], \quad (\text{A15})$$

where the closed contour \overline{ABCD} are defined in Fig. 2 with $A = (-\infty, 0)$, $B = (+\infty, 0)$, $C = (+\infty, -in_c\pi)$, $D = (-\infty, -in_c\pi)$ and $n_c = \frac{\nu_c}{4\pi k_B T_c}$. From the Cauchy's residue theorem, we can show that $a_1(i\nu_{c_1})$ in the action (A3) follows (A5).

It is noted that our result is similar to that obtained by Larkin and Varlamov⁴², whose result Eq. (10.171) in Reference [42] is similar to our (A5) with a finite energy cut-off in their digamma function. Moreover, their final result Eq. (10.177) is based on the power expansion of a small $\frac{\nu_{c_1}}{T}$ of the digamma function⁴².

Following a similar derivation, we can obtain the Ginzburg-Landau action for the 3D superconductor. They follow the same formula to that given in Eq. (A3), with the parameters defined as following:

$$\begin{aligned} a_0 &= N\rho_{3D}, & a_1(i\nu_{c_1}) &= N\rho_{3D} I(n_{c_1}) \Big|_{n_{c_1} = \frac{\nu_{c_1}}{4\pi k_B T_c}}, \\ a_2 &= \frac{N\rho_{3D}\mu_F\gamma}{24(k_B T_c)^2 m}, & b &= \frac{N\rho_{3D}\gamma}{8(k_B T_c)^2}. \end{aligned} \quad (\text{A16})$$

Here ρ_{3D} is the density of states at the Fermi energy defined as

$$\rho_{3D} = \frac{\sqrt{2m\mu_F}ma^3}{2\pi^2\hbar^3}, \quad (\text{A17})$$

and T_c is same defined to (A12).

Appendix B: Transformation from imaginary- to real-frequency actions

Let us give a derivation of the transformation from the imaginary-frequency action (25) to the real-frequency one (29).

Due to the sign function in $F(\nu_{c_1}, \omega_{r_1})$, we separate the action $S^{(2)}$ into two parts, $S^{(2)} = S_+^{(2)} + S_-^{(2)}$, where the former involves the summation of the $i\omega_{r_1}^+$ with $\omega_{r_1} > 0$ and the latter contains the summation of the $i\omega_{r_1}^-$ with $\omega_{r_1} < 0$.




FIG. 3: (Color online) The closed contour for the integral Eq. (B1)

We first ignore the V^{-1} term in action $S^{(2)}$. Introduce two summations I_+ and I_- as below.

$$\begin{aligned} I_+ &= \sum_{i\omega_{r_1}^+} \bar{\Delta}(1_c, 1_r) \mathcal{G}_0^{-1} \Delta(1_c, 1_r) \\ &= \frac{\beta N \rho_{2D}}{4\pi} \oint_C dz \left[\frac{n_F(z)}{z - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}}} \right] \bar{\Delta}(1_c, z) \Delta(1_c, z). \end{aligned} \quad (\text{B1})$$

Here we have ignored the average operation of $\langle \frac{1}{z - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}}} \rangle$ for simplicity. The closed anti-clockwise contour C is schematically shown in Fig. 3 with the radius of the upper half-circle becoming $+\infty$. The contour integral I_+ can thus be reexpressed into

$$\begin{aligned} I_+ &= \frac{\beta N \rho_{2D}}{4\pi} \int_{-\infty}^{+\infty} d\omega_r \left[\frac{n_F(\omega_r)}{\omega_r + \frac{1}{2}i|\nu_{c_1}| - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}}} \right] \\ &\quad \times \bar{\Delta}(1_c, \omega_r + \frac{1}{2}i|\nu_{c_1}|) \Delta(1_c, \omega_r + \frac{1}{2}i|\nu_{c_1}|). \end{aligned} \quad (\text{B2})$$




FIG. 4: (Color online) The closed contour for the integral Eq. (B3)

Similarly, we introduce and calculate I_- as below.

$$\begin{aligned} I_- &= \sum_{i\omega_{r_1}^-} \bar{\Delta}(1_c, 1_r) \mathcal{G}_0^{-1} \Delta(1_c, 1_r) \\ &= \frac{\beta N \rho_{2D}}{4\pi} \oint_{C'} dz \left[\frac{n_F(z)}{z - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}}} \right] \bar{\Delta}(1_c, z) \Delta(1_c, z) \\ &= \frac{\beta N \rho_{2D}}{4\pi} \int_{-\infty}^{+\infty} d\omega_r \left[\frac{n_F(\omega_r)}{\omega_r - \frac{1}{2}i|\nu_{c_1}| - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}}} \right] \\ &\quad \times \bar{\Delta}(1_c, \omega_r - \frac{1}{2}i|\nu_{c_1}|) \Delta(1_c, \omega_r - \frac{1}{2}i|\nu_{c_1}|). \end{aligned} \quad (\text{B3})$$

Here the closed clockwise contour C' is schematically shown in Fig. 4 with the radius of the lower half-circle becoming $+\infty$.

Thus we obtain $I = I_+ + I_-$, which follows

$$\begin{aligned} I(\nu_{c_1}) &= I(i\nu_{c_1} \rightarrow \nu_{c_1} + i\delta^+) \\ &= \frac{\beta N \rho_{2D}}{4\pi} \int_{-\infty}^{+\infty} d\omega_r \left[\frac{n_F(\omega_r - \frac{1}{2}\nu_{c_1})}{\omega_r - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}} + i\delta^+} \right. \\ &\quad \left. + \frac{n_F(\omega_r + \frac{1}{2}\nu_{c_1})}{\omega_r - A_{\mathbf{k}_F \cdot \mathbf{q}_{c_1}} - i\delta^+} \right] \bar{\Delta}(1_c, \omega_r) \Delta(1_c, \omega_r). \end{aligned} \quad (\text{B4})$$

From (B4), we can obtain the transformation from the imaginary-frequency action (25) to the real-frequency one (29).

* Electronic address: suyh@ytu.edu.cn

- 1 V. L. Ginzburg and L. D. Landau, *English translation in Collected papers of L. D. Landau, p. 546-568: 73 On the theory of superconductivity* (Pergamon Press, Oxford, 1965), URL <https://doi.org/10.1016/B978-0-08-010586-4.50078-X>.
- 2 L. N. Cooper, Phys. Rev. **104**, 1189 (1956), URL <https://link.aps.org/doi/10.1103/PhysRev.104.1189>.
- 3 J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. **106**, 162 (1957), URL <https://link.aps.org/doi/10.1103/PhysRev.106.162>.
- 4 J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. **108**, 1175 (1957), URL <https://link.aps.org/doi/10.1103/PhysRev.108.1175>.
- 5 L. P. Gor'kov, Sov. Phys. JETP **36**, 1364 (1959).
- 6 L. P. Gor'kov, Sov. Phys. JETP **37**, 998 (1960).
- 7 L. P. Gor'kov and T. K. Melik-Barkhudarov, Sov. Phys. JETP **18**, 1031 (1964).
- 8 P. Fulde and R. A. Ferrell, Phys. Rev. **135**, A550 (1964), URL <https://link.aps.org/doi/10.1103/PhysRev.135.A550>.
- 9 A. I. Larkin and Y. N. Ovchinnikov, Sov. Phys. JETP **20**, 762 (1965).
- 10 D. F. Agterberg, J. S. Davis, S. D. Edkins, E. Fradkin, D. J. Van Harlingen, S. A. Kivelson, P. A. Lee, L. Radzihovsky, J. M. Tranquada, and Y. Wang, Annual Review of Condensed Matter Physics **11**, 231 (2020), URL <https://doi.org/10.1146/annurev-conmatphys-031119-050711>.
- 11 P. W. Anderson, Journal of Physics C: Solid State Physics **3**, 2436 (1970), URL <https://doi.org/10.1088/0022-3719/3/12/008>.
- 12 E. Abrahams and T. Tsuneto, Phys. Rev. **152**, 416 (1966), URL <https://link.aps.org/doi/10.1103/PhysRev.152.416>.
- 13 L. P. Gor'kov and E. M. Eliashberg, Sov. Phys. JETP **27**, 328 (1968).
- 14 A. Altland and B. Simons, *Condensed matter field theory* (Cambridge University Press, 2006), ISBN 978-0-521-84508-3.
- 15 G. Eilenberger and V. Ambegaokar, Phys. Rev. **158**, 332 (1967), URL <https://link.aps.org/doi/10.1103/PhysRev.158.332>.
- 16 P. Fulde and K. Maki, Phys. kondens. Materie **8**, 371 (1969), URL <https://doi.org/10.1007/BF02422865>.
- 17 A. Abanov and A. V. Chubukov, Phys. Rev. B **102**, 024524 (2020), URL <https://link.aps.org/doi/10.1103/PhysRevB.102.024524>.
- 18 Y.-M. Wu, A. Abanov, Y. Wang, and A. V. Chubukov, Phys. Rev. B **102**, 024525 (2020), URL <https://link.aps.org/doi/10.1103/PhysRevB.102.024525>.
- 19 Y.-M. Wu, A. Abanov, and A. V. Chubukov, Phys. Rev. B **102**, 094516 (2020), URL <https://link.aps.org/doi/10.1103/PhysRevB.102.094516>.
- 20 Y.-M. Wu, S.-S. Zhang, A. Abanov, and A. V. Chubukov, Phys. Rev. B **103**, 024522 (2021), URL <https://link.aps.org/doi/10.1103/PhysRevB.103.024522>.
- 21 Y.-M. Wu, S.-S. Zhang, A. Abanov, and A. V. Chubukov, Phys. Rev. B **103**, 184508 (2021), URL <https://link.aps.org/doi/10.1103/PhysRevB.103.184508>.
- 22 S.-S. Zhang, Y.-M. Wu, A. Abanov, and A. V. Chubukov, Phys. Rev. B **104**, 144509 (2021), URL <https://link.aps.org/doi/10.1103/PhysRevB.104.144509>.
- 23 K.-W. Lee and W. E. Pickett, Phys. Rev. Lett. **93**, 237003 (2004), URL <https://link.aps.org/doi/10.1103/PhysRevLett.93.237003>.
- 24 X. Blase, C. Adessi, and D. Connétable, Phys. Rev. Lett. **93**, 237004 (2004), URL <https://link.aps.org/doi/10.1103/PhysRevLett.93.237004>.
- 25 F. Giustino, M. L. Cohen, and S. G. Louie, Phys. Rev. B **76**, 165108 (2007), URL <https://link.aps.org/doi/10.1103/PhysRevB.76.165108>.
- 26 K. G. Wilson and J. Kogut, Physics Report (Section C of Physics Letters) **12**, 75 (1974).
- 27 J. A. Hertz, Phys. Rev. B **14**, 1165 (1976), URL <https://link.aps.org/doi/10.1103/PhysRevB.14.1165>.
- 28 A. J. Millis, Phys. Rev. B **48**, 7183 (1993), URL <https://link.aps.org/doi/10.1103/PhysRevB.48.7183>.
- 29 M. E. Fisher, Rev. Mod. Phys. **70**, 653 (1998), URL <https://link.aps.org/doi/10.1103/RevModPhys.70.653>.
- 30 H. v. Löhneysen, A. Rosch, M. Vojta, and P. Wölfle, Rev. Mod. Phys. **79**, 1015 (2007), URL <http://link.aps.org/doi/10.1103/RevModPhys.79.1015>.
- 31 Q. Si, S. Rabello, K. Ingersent, and J. L. Smith, Phys. Rev. B **68**, 115103 (2003), URL <http://link.aps.org/doi/10.1103/PhysRevB.68.115103>.
- 32 T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, Science **303**, 1490 (2004), URL <https://science.sciencemag.org/content/303/5663/1490>.
- 33 C. Xu and L. Balents, Phys. Rev. B **84**, 014402 (2011), URL <https://link.aps.org/doi/10.1103/PhysRevB.84.014402>.
- 34 S. Sachdev and B. Keimer, Physics Today **64**, 29 (2011), URL <https://doi.org/10.1063/1.3554314>.
- 35 J. Zaanan, SciPost Phys. **6**, 61 (2019), URL <https://scipost.org/10.21468/SciPostPhys.6.5.061>.
- 36 C. M. Varma, Rev. Mod. Phys. **92**, 031001 (2020), URL <https://link.aps.org/doi/10.1103/RevModPhys.92.031001>.
- 37 S.-S. Lee, Annual Review of Condensed Matter Physics **9**, 227 (2018), URL <https://doi.org/10.1146/annurev-conmatphys-031016-025531>.
- 38 A. V. Chubukov, Journal Club for Condensed Matter Physics **2** (2018), URL https://doi.org/10.36471/JCCM_November_2018_02.
- 39 T. Moriya, *Spin fluctuations in itinerant electron magnetism* (Springer-Verlag Berlin Heidelberg, 1985), ISBN 3-540-15422-1.
- 40 Y. Su, H. Liao, and T. Li, J. Phys.: Condens. Matter **27**, 105702 (2015), URL <https://iopscience.iop.org/article/10.1088/0953-8984/27/10/105702>.
- 41 T. Li and Y. Su, J. Phys.: Condens. Matter **29**, 425603 (2017), URL <https://iopscience.iop.org/article/10.1088/1361-648X/aa85f4>.
- 42 A. I. Larkin and A. A. Varlamov, *Superconductivity: Conventional and unconventional superconductors (Volume 1)* (Springer-Verlag Berlin Heidelberg, 2008), 1st ed., ISBN 978-3-540-73252-5.