# Shadow Celestial Amplitude

Chi-Ming Chang<sup>1,2\*</sup>, Wei Cui<sup>2,1†</sup>, Wen-Jie Ma,<sup>2,1‡</sup> Hongfei Shu<sup>2,1§</sup>, Hao Zou<sup>2,1¶</sup>

<sup>1</sup>Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, China

<sup>2</sup> Yanqi Lake Beijing Institute of Mathematical Sciences and Applications (BIMSA), Huairou District, Beijing 101408, P. R. China

#### Abstract

We study scattering amplitudes in the shadow conformal primary basis, which satisfies the same defining properties as the original conformal primary basis and has many advantages over it. The shadow celestial amplitudes exhibit locality manifestly on the celestial sphere, and behave like correlation functions in conformal field theory under the operator product expansion (OPE) limit. We study the OPE limits for three-point shadow celestial amplitude, and general  $2 \rightarrow n-2$  shadow celestial amplitudes from a large class of Feynman diagrams. In particular, we compute the conformal block expansion of the *s*-channel four-point shadow celestial amplitude of massless scalars at tree-level, and show that the expansion coefficients factorize as products of OPE coefficients.

October 18, 2022

<sup>\*</sup>cmchang@tsinghua.edu.cn

<sup>&</sup>lt;sup>†</sup>cwei@bimsa.cn

<sup>&</sup>lt;sup>‡</sup>wenjia.ma@bimsa.cn

<sup>§</sup>shuphy124@gmail.com

<sup>¶</sup>hzou@bimsa.edu

# Contents

1	Intr	oduction	1
2	Shadow conformal primary basis		4
	2.1	Review on the celestial amplitudes	4
	2.2	Disadvantages of the celestial amplitudes	5
	2.3	A different conformal primary basis	7
	2.4	Translation symmetry	8
3	OPE behaviour of the shadow celestial amplitudes		9
	3.1	OPE analysis for three-point shadow celestial amplitudes	10
	3.2	OPE analysis from the generalized optical theorem	13
	3.3	OPE analysis for a special class of Feynman diagrams	14
	3.4	OPE analysis for four-point shadow celestial amplitudes	15
4	Examples of shadow celestial amplitudes		17
	4.1	Three-point shadow celestial amplitudes	17
	4.2	Four-point shadow celestial amplitudes	18
5	Outlook 2		22
A	Preliminaries		
	A.1	The generalized optical theorem	24
	A.2	Shadow transformation and conformal partial waves	25
В	The	conformal integral	28

# 1 Introduction

Celestial holography is believed to be a concrete realization of holographic principles for quantum gravity in asymptotically flat spacetime (AFS) [1–5]. It relates scattering amplitudes of a quantum field theory or quantum gravity in a four-dimensional AFS to correlation functions of a celestial conformal field theory (CCFT) on the two-dimensional celestial sphere. The Lorentz symmetry of the four-dimensional AFS is realized as the  $SL(2, \mathbb{C})$  conformal symmetry of the celestial sphere. The goal of celestial holography is to study the scattering amplitudes in AFS by using the techniques developed in conformal field theory (CFT). One of the most important achievements in celestial holography is recasting the soft theorems in flat space into Ward identities in two-dimensional CFTs. The currents associated to these Ward identities generate asymptotic symmetries in the four-dimensional spacetime [6–16] and can be re-organized into the  $w_{1+\infty}$  algebra [17–19].

To manifest the  $SL(2, \mathbb{C})$  Lorentz symmetry in the scattering amplitudes, one needs to change the basis of asymptotic states from the standard plane-wave basis to the *conformal primary basis* [20–23]. By definition, the conformal primary basis must satisfy the equations of motion and transform covariantly under  $SL(2, \mathbb{C})$ . The S-matrix elements in the conformal primary basis are referred to as *celestial amplitudes*. The conformal primary basis that is widely used in the literature for massless particles is built from the usual plane-wave basis followed by a Mellin transformation. However, in this basis, the coordinates on the celestial sphere relate directly to the solid angles of the flat space momentum. Thus the corresponding celestial amplitudes are highly constrained by four-dimensional kinematics and do not take the standard form of CFT correlation functions. For example, the four-point celestial amplitudes of massless scalars contain an unfamiliar delta-function  $\delta(\chi - \bar{\chi})$  originated from the momentum conservation. This distributional factor forces the celestial amplitude to live on the equator of the celestial sphere. In addition, depending on the assignments of the incoming and outgoing particles, the celestial amplitudes are only supported in disjoint intervals on the equator.

Finally, in terms of the massless conformal primary basis, the celestial amplitudes do not have proper conformal block expansion. This can be seen by looking at the *s*-channel tree-level celestial amplitude of two incoming, two outgoing massless particles and one massive exchange particle. The imaginary part of the corresponding scattering amplitude in the plane-wave basis is factorized into two three-point scattering amplitudes due to the optical theorem. This leads to a factorization in the conformal partial wave expansion of the celestial amplitudes [24, 25].<sup>1</sup> However, since the integration kernel in the conformal partial wave expansion does not have poles located at the right half-plane, one does not get a conformal block expansion by closing the contour. Again, in the literature, the studies of conformal block expansion of celestial amplitudes are limited to the Klein space [28–31] or three dimensional space [32, 33].

To fix these issues, we consider a different set of conformal primary wavefunctions for massless particles, which are in a different branch of solutions to the two defining properties of conformal primary wavefunction [1], i.e. they satisfy the equations of motion and trans-

<sup>&</sup>lt;sup>1</sup>Partial wave expansion of the celestial amplitudes is also studied in [26, 27].

form covariantly under  $SL(2,\mathbb{C})$ . It turns out that, up to a constant factor, these conformal primary wavefunctions are equivalent to the shadow transformations [34, 35] of the original conformal primary wave functions.<sup>2</sup> We will refer to this basis as the *shadow conformal* primary basis. Expanding the scattering amplitude in the shadow conformal primary basis, we define the *shadow celestial amplitudes*, which can be obtained by performing the shadow transformation of all external operators of the celestial amplitudes.<sup>3</sup> The shadow celestial amplitudes resolve all the abovementioned issues and lead to the standard correlation functions of CFTs. Specifically, the shadow celestial amplitudes of four massless particles no longer have  $\delta(\chi - \bar{\chi})$  and are defined on the entire celestial sphere. In addition, the shadow celestial amplitudes have well-behaved OPE limits. For scattering amplitudes with n external massless real scalars, we consider the OPE limit by making the celestial coordinates of the first two incoming particles close to each other. For amplitudes from a large class of Feynman diagrams, we find that the shadow celestial amplitudes factorize as expected as n-point correlation functions in a CFT. Using the generalized optical theorem, we also obtain the correct factorization of the imaginary part of shadow celestial amplitudes for any Feynman diagrams. What's more, we compute the 4-point shadow celestial amplitudes involving four external massless scalars and one exchange massive scalar and derive its conformal block expansion in the  $12 \leftrightarrow 34$  channel. We find that the coefficients appearing in the conformal block expansion give the correct OPE coefficients obtained from the corresponding 3-point shadow celestial amplitudes matching the results from the OPE analysis.

This paper is organized as follows. In Section 2, we review the celestial amplitudes, discuss their disadvantages, and introduce the shadow celestial amplitudes. In Section 3, we analyze the OPE limits of *n*-point shadow celestial amplitudes involving *n* external real massless scalars and show that the OPE limits are well-defined for the shadow celestial amplitudes. In Section 4, we work out the conformal block expansion on the 4-point shadow celestial amplitudes involving four external massless scalars and one exchange massive scalar. We find the complete agreement between the block expansion coefficients and OPE coefficients. In Section 5, we conclude our work and point out a few future directions. In Appendix A, we review the generalized optical theorem, the shadow transformation and conformal partial waves, that will be used in later parts of this paper.

<sup>&</sup>lt;sup>2</sup>The shadows of the conformal wave functions were previously studied in [20].

<sup>&</sup>lt;sup>3</sup>Implementing the shadow transformation on one of the external operators of the celestial amplitude was discussed in [36–38].

### 2 Shadow conformal primary basis

#### 2.1 Review on the celestial amplitudes

Celestial amplitudes are obtained by expanding the position space amplitudes with respect to the conformal primary wavefunctions [20] instead of the plane-waves, *i.e.*,<sup>4</sup>

$$\mathcal{A}^{\Delta_i}(z_i) = \left(\prod_{j=1}^{k+n} \int d^4 x_j\right) \left(\prod_{j=1}^k \phi^+_{\Delta_j}(z_j; x_j)\right) \left(\prod_{j=k+1}^{k+n} \phi^-_{\Delta_j}(z_j; x_j)\right) \mathcal{M}(x_i) , \qquad (2.1)$$

where  $\mathcal{M}(x_j)$  is the scattering amplitude in position space. The conformal primary wave functions  $\phi_{\Delta}^{\pm}(z;x)$  for massless and massive scalars with mass *m* are given by

$$\phi_{\Delta}^{\pm}(z;x) = \int_{0}^{+\infty} d\omega \,\omega^{\Delta-1} e^{\pm i\omega\hat{q}\cdot x - \epsilon\omega} , \qquad (2.2)$$

and

$$\phi_{\Delta}^{\pm}(z;x) = \int \frac{d^3 \hat{p}'}{\hat{p}'^0} G_{\Delta}(z,\bar{z};\hat{p}') e^{\pm im\hat{p}' \cdot x} , \qquad (2.3)$$

respectively. Here z and  $\bar{z}$  are coordinates on celestial sphere and  $G_{\Delta}(z, \bar{z}; \hat{p})$  is the bulkto-boundary propagator. The coordinates  $(z, \bar{z})$  on the celestial sphere are related to the massless on-shell momenta  $q^{\mu}$  through

$$q^{\mu} = \omega \hat{q}^{\mu} = \omega (1 + z\bar{z}, z + \bar{z}, -i(z - \bar{z}), 1 - z\bar{z})$$
(2.4)

and to the massive on-shell momenta  $p^{\mu}$  through

$$p^{\mu} = m\hat{p}^{\mu} = \frac{m}{2y}(1 + y^2 + z\bar{z}, z + \bar{z}, -i(z - \bar{z}), 1 - y^2 - z\bar{z}).$$
(2.5)

In terms of  $\hat{p}$  in (2.5), the bulk-to-boundary propagator takes the form <sup>5</sup>

$$G_{\Delta}(\hat{q}; \hat{p}') = \frac{1}{(-\hat{q} \cdot \hat{p}')^{\Delta}}$$
 (2.6)

<sup>&</sup>lt;sup>4</sup>Throughout this paper,  $\mathcal{O}_i(z_i)$  should be understood as  $\mathcal{O}_i(z_i, \bar{z}_i)$ . We use this abbreviation to simplify the notation.

<sup>&</sup>lt;sup>5</sup>In this paper, we use the most positive metric in four-dimensional flat space, *i.e.*,  $g_{AB} = \text{diag}(-1, +1, +1, +1)$ .

In terms of the conformal primary basis (2.2), the celestial amplitudes in (2.1) with  $n_1$  massless scalars and  $n_2$  massive scalars can be re-expressed as

$$\mathcal{A}^{\Delta_i}(z_i) = \left(\prod_{j=1}^{n_1} \int d\omega_j \omega_j^{\Delta_j - 1}\right) \left(\prod_{j=1}^{n_2} \int \frac{d^3 \hat{p}'_j}{\hat{p}'^0_j} G_{\Delta_j}(\hat{q}_j; \hat{p}'_j)\right) \mathcal{M}_{k \to n}(q_i, p'_i) , \qquad (2.7)$$

where  $\mathcal{M}(q_i, p'_i)$  is the scattering amplitude in the plane-wave basis.

### 2.2 Disadvantages of the celestial amplitudes

Unfortunately, the celestial amplitudes defined in (2.7) have many disadvantages. For example, the four-point celestial amplitudes  $\mathcal{A}_{12\rightarrow34}^{\Delta_i}$  that describe the two-to-two scattering of four massless particles are exotic. The form of  $\mathcal{A}_{12\rightarrow34}^{\Delta_i}$  is completely fixed by conformal symmetry up to a function of conformal cross-ratios. In other words, we have<sup>6</sup>

$$\mathcal{A}_{12\to34}^{\Delta_i}(z_i) = I_{12-34}(z_i)f(\chi) , \qquad (2.8)$$

where  $f(\chi)$  is a function of cross-ratios

$$\chi = \frac{z_{12}z_{34}}{z_{13}z_{24}} , \quad \bar{\chi} = \frac{\bar{z}_{12}\bar{z}_{34}}{\bar{z}_{13}\bar{z}_{24}} , \qquad (2.9)$$

and  $I_{12-34}$  is given by

$$I_{12-34} \equiv \frac{\left(\frac{z_{24}}{z_{14}}\right)^{h_{12}} \left(\frac{z_{14}}{z_{13}}\right)^{h_{34}}}{z_{12}^{h_1+h_2} z_{34}^{h_3+h_4}} \frac{\left(\frac{\bar{z}_{24}}{\bar{z}_{14}}\right)^{\bar{h}_{12}} \left(\frac{\bar{z}_{14}}{\bar{z}_{13}}\right)^{\bar{h}_{34}}}{\bar{z}_{12}^{\bar{h}_1+\bar{h}_2} \bar{z}_{34}^{\bar{h}_3+\bar{h}_4}} , \quad h_{ij} \equiv h_i - h_j , \quad \bar{h}_{ij} \equiv \bar{h}_i - \bar{h}_j , \quad (2.10)$$

and  $\Delta_i = 2h_i$ . The corresponding scattering amplitude in the plane-wave basis at tree level is given by the diagrams in *s*-, *t*-, and *u*-channels as

$$\mathcal{M}_{2\to 2}(q_i) = (2\pi)^4 \delta^{(4)}(q_1' + q_2' - q_3' - q_4')(-ig)^2 \left(\frac{1}{s - m^2} + \frac{1}{t - m^2} + \frac{1}{u - m^2}\right) , \quad (2.11)$$

<sup>6</sup>Again, we omit the anti-holomorphic dependence to simplify the notation.

where m is the mass of the exchange massive scalar. This leads to the following expression for the function  $f(\chi)$  with  $\chi \ge 1$ 

$$f(\chi) = N_{gm}(\beta)\chi^2(\chi - 1)^{h_{13} - h_{24}}\delta(\chi - \bar{\chi}) \left[ 1 + e^{-i\pi\frac{\beta}{2}} \left(\frac{\chi}{\chi - 1}\right)^{\frac{\beta}{2}} + e^{-i\pi\frac{\beta}{2}}\chi^{\frac{\beta}{2}} \right], \qquad (2.12)$$

where  $h_{ij} \equiv h_i - h_j$  and  $\beta = \sum_{i=1}^4 \Delta_i - 4$  and  $N_{gm}(\beta)$  is given by

$$N_{gm}(\beta) = \frac{g^2 \pi^5 m^{\beta-2}}{2^{\beta-1}} \frac{e^{i\pi\beta/2}}{\sin\pi\beta/2} .$$
 (2.13)

From (2.12), we note that  $\mathcal{A}_{12\to34}^{\Delta_i}(z_i)$  contain a delta-function  $\delta(\chi - \bar{\chi})$  which forces the four-point celestial amplitudes to live on the equator of the celestial sphere and makes the structure of correlation functions in CCFTs very different from those in the standard CFTs. Moreover, (2.12) is for the 12  $\rightarrow$  34 kinematic and only valid when  $\chi \geq 1$ . The other two kinematics 14  $\rightarrow$  23 and 13  $\rightarrow$  24 are defined in distinct intervals  $\chi \leq 0$  and  $0 \leq \chi \leq 1$  on the equator. This also makes the four-point celestial amplitudes exotic. Finally,  $\mathcal{A}_{12\to34}^{\Delta_i}(z_i)$  does not have a proper conformal block expansion. We take the *s*-channel scattering amplitude (the first term in (2.12)) as an example. As we will see in Section 4, the partial wave expansion of the *s*-channel celestial amplitudes  $\mathcal{A}_s^{\Delta_i}$  takes the form<sup>7</sup>

$$\mathcal{A}_{s}^{\Delta_{i}}(z_{i}) = \int_{-\infty}^{+\infty} \frac{d\lambda}{2\pi} \frac{\rho_{h,\bar{h}}^{\Delta_{i}}}{n_{h,\bar{h}}} K_{1-h,1-\bar{h}}^{\Delta_{3},\Delta_{4}} G_{h,\bar{h}}^{\Delta_{i}}(z_{i}) , \qquad (2.14)$$

where  $h = \bar{h} = (1 + i\lambda)/2$  and the spectral density is

$$K_{1-h,1-\bar{h}}^{\Delta_{3},\Delta_{4}} \frac{\rho_{h,\bar{h}}^{\Delta_{i}}}{n_{h,\bar{h}}} = \frac{-g^{2}\pi^{9}m^{\beta-4}}{2^{\beta-6}} \frac{e^{i\pi\beta/2}}{\sin\pi\beta/2} \times B(h+h_{12},h-h_{12})B(h-h_{34},h+h_{34})D^{\Delta\Delta}(m) .$$
(2.15)

When  $\chi \leq 1$ , closing the contour to the right-half  $\lambda$ -plane leads to a vanishing conformal block expansion because the integrand in (2.14) does not have poles located in the right-half  $\lambda$ -plane. On the other hand, when  $\chi > 1$  the integrand in (2.14) does not decay to zero when  $\operatorname{Re}(\lambda)$  approaches infinity and we can not close the contour. This reflects the fact that  $\mathcal{A}_s^{\Delta_i}(z_i)$  only supports on  $\chi > 1$ . Thus, we conclude that  $\mathcal{A}_s^{\Delta_i}(z_i)$  does not have a proper conformal block expansion.

<sup>&</sup>lt;sup>7</sup>See Appendix A.2 for a review on the conformal block and conformal partial wave.

### 2.3 A different conformal primary basis

It is easy to see that the above disadvantages of the celestial amplitudes are due to the fact that the conformal primary wave functions (2.2) for massless scalars are constructed from the plane-waves by performing integral only over the energy  $\omega$ . Thus, we consider a different set of conformal primary wave functions for massless scalars:

$$\widetilde{\phi}_{\Delta}^{\pm}(z;x) = \int \frac{d^3q'}{q'^0} G_{\Delta}(\hat{q};q') e^{\pm i\omega\hat{q}'\cdot x - \epsilon\omega} .$$
(2.16)

Here,  $q^{\mu}$  is defined in (2.4) and  $G_{\Delta}(\hat{q}; q')$  is given by

$$G_{\Delta}(\hat{q};q') = \frac{1}{(-\hat{q}\cdot q')^{\Delta}}, \qquad (2.17)$$

which takes the same form as bulk-to-boundary propagator (2.6) with  $\hat{p}'$  replaced by q'. Using the identity  $q'(z') \cdot \hat{q}(z) = -2\omega |z'-z|^2$  and noting that

$$\int \frac{d^3q}{q^0} = 4 \int d^2z \int_0^{+\infty} d\omega \,\omega \,, \qquad (2.18)$$

the conformal primary waves functions (2.16) can be written as

$$\widetilde{\phi}_{\Delta}^{\pm}(z;x) = \int d^2 z' \frac{2^{1-\Delta}}{|z-z'|^{2\Delta}} \int_0^\infty d\omega \,\omega^{1-\Delta} e^{\pm i\omega \widehat{q}' \cdot x - \epsilon \omega} = \int d^2 z' \frac{2^{1-\Delta}}{|z-z'|^{2\Delta}} \phi_{\Delta}^{\pm}(z';x) , \quad (2.19)$$

which are proportional to the shadows of the original conformal primary wave functions  $\phi_{\Delta}^{\pm}(z;x)$ , that were previously studied in [20]. We dub it *the shadow conformal primary basis*. The shadow conformal primary wave functions (2.16) obey the massless Klein-Gordon equation and transform covariantly under the conformal transformation since the shadow transformation is conformally covariant. In other words, they satisfy the two defining properties for conformal primary wave functions given in [1, 20]. On the other hand, one does not obtain a new wave function by performing the shadow transformation on the massive conformal primary basis (2.3) because the result from the shadow integral takes the same form as (2.3) up to a change of conformal dimension from  $\Delta$  to  $2 - \Delta$  [20].

Using the shadow conformal primary wave functions (2.16), we define the shadow celestial

*amplitudes* as

$$\widetilde{\mathcal{A}}^{\Delta_i}(z_i) = \left(\prod_{j=1}^n \int \frac{d^3 k'_j}{k'^0_j} G_{\Delta_j}(\hat{q}_j; k'_j)\right) \mathcal{M}(k'_i) , \qquad (2.20)$$

where  $k'_j$  can be either  $q'_j$  or  $\hat{p}'_j$  depending on whether the particle is massless or massive.

### 2.4 Translation symmetry

Unlike the scattering amplitudes in the plane-wave basis, translation symmetry in shadow celestial amplitudes  $\tilde{\mathcal{A}}^{\Delta_i}(z_i)$  becomes obscure, while Lorentz symmetry is manifest. In this subsection, we will discuss how the translation generators  $\hat{P}^{\mu}$  act on the shadow celestial amplitude. The action of the translation generators on massive scalars was studied in [39]. Specifically,  $\hat{P}^{\mu}$  act on a massive scalar in CCFT as a differential operator  $\mathcal{P}^{\mu}$ 

$$[\hat{P}^{\mu}, \Phi_a^{\Delta_a}(z_a)] = \mathcal{P}_a^{\mu} \Phi_a^{\Delta_a}(z_a) , \qquad (2.21)$$

where the different operator  $\mathcal{P}^{\mu}_{a}$  acting on massive scalars takes the form

$$\mathcal{P}_{a}^{\mu} = \epsilon_{a} \frac{m_{a}}{2} \left[ \left( \left( \partial_{z_{a}} \partial_{\bar{z}_{a}} \hat{q}_{a}^{\mu} \right) + \frac{\left( \partial_{z_{a}} \hat{q}_{a}^{\mu} \right) \partial_{\bar{z}_{a}} + \left( \partial_{\bar{z}_{a}} \hat{q}_{a}^{\mu} \right) \partial_{z_{a}}}{\Delta_{a} - 1} + \frac{\hat{q}_{a}^{\mu} \partial_{z_{a}} \partial_{\bar{z}_{a}}}{\left( \Delta_{a} - 1 \right)^{2}} \right) e^{-\partial_{\Delta_{a}}} + \frac{\Delta_{a} \hat{q}_{a}^{\mu}}{\Delta_{a} - 1} e^{\partial_{\Delta_{a}}} \right]$$

$$\tag{2.22}$$

with  $\epsilon_k = \pm 1$  labeling incoming and outgoing states, respectively.

To find the action of  $\hat{P}^{\mu}$  on massless shadow celestial amplitudes (2.20), we start with the action in the plane-wave basis

$$\hat{P}^{\mu}_{a}\mathcal{M}(q_{a}) = \epsilon_{a}q^{\mu}_{a}\mathcal{M}(q_{a}) , \qquad (2.23)$$

Substituting (2.23) into (2.20) leads to

$$\hat{P}^{\mu}_{a}\widetilde{\mathcal{A}}^{\Delta_{i}}(z_{i}) = \left(\prod_{j\neq a}^{n} \int \frac{d^{3}k'_{j}}{k'^{0}_{j}} G_{\Delta_{j}}(\hat{q}_{j};k'_{j})\right) \int \frac{d^{3}q'_{a}}{q^{\prime 0}_{a}} \frac{1}{(-\hat{q}_{a}\cdot q'_{a})^{\Delta_{a}}} \epsilon_{a} q^{\prime \mu}_{a} \mathcal{M}(q'_{a}) .$$

$$(2.24)$$

Thus, finding the action of  $\hat{P}^{\mu}$  on massless shadow celestial amplitudes is translated into

finding a differential operator  $\mathcal{P}^{\mu}_{a}$  such that

$$\mathcal{P}_{a}^{\mu} \frac{1}{(-\hat{q}_{a} \cdot q_{a}')^{\Delta_{a}}} = \epsilon_{a} q_{a}'^{\mu} \frac{1}{(-\hat{q}_{a} \cdot q_{a}')^{\Delta_{a}}} .$$
(2.25)

This can be realized by the following differential operator

$$\mathcal{P}_{a}^{\mu} = \frac{1}{2} \epsilon_{a} \left( \left( \partial_{z_{a}} \partial_{\bar{z}_{a}} \hat{q}_{a}^{\mu} \right) + \frac{\left( \partial_{z_{a}} \hat{q}_{a}^{\mu} \right) \partial_{\bar{z}_{a}} + \left( \partial_{\bar{z}_{a}} \hat{q}_{a}^{\mu} \right) \partial_{z_{a}}}{\Delta_{a} - 1} + \frac{\hat{q}_{a}^{\mu} \partial_{z_{a}} \partial_{\bar{z}_{a}}}{(\Delta_{a} - 1)^{2}} \right) e^{-\partial_{\Delta_{a}}} , \qquad (2.26)$$

which coincides with the first term of the massive translation operator (2.22) up to the mass  $m_a$ . In other words, the action of the translation generators  $\hat{P}^{\mu}$  on a massless scalar operator  $\mathcal{O}_a^{\Delta_a}(z_a)$  in CCFT is

$$[\hat{P}^{\mu}, \mathcal{O}_a^{\Delta_a}(z_a)] = \mathcal{P}_a^{\mu} \mathcal{O}_a^{\Delta_a}(z_a) .$$
(2.27)

Thus, the translation symmetries amount to the following differential equations

$$\sum_{a=1}^{n} \mathcal{P}_{a}^{\mu} \widetilde{\mathcal{A}}_{n}^{\Delta_{i}}(z_{i}) = 0 , \qquad (2.28)$$

where  $\mathcal{P}_{a}^{\mu}$  is given in (2.22) when it acts on massive scalar operators and in (2.26) when it acts on massless scalar operators.

### **3** OPE behaviour of the shadow celestial amplitudes

In this section, we will study the OPE behaviour of the shadow celestial amplitudes of 2 incoming massless real scalars, labeled by 1 and 2, and n-2 outgoing massless real scalars. Similar analysis works for the same shadow celestial amplitudes with the incoming and outgoing particles swapped. The shadow celestial amplitudes of interest can be written as<sup>8</sup>

$$\begin{aligned} \widetilde{\mathcal{A}}_{2 \to n-2}^{\Delta_{i}}(z_{i}) = & 2^{1-\Delta_{2}} \int \frac{d^{3}q_{1}'}{q_{1}'^{0}} \frac{1}{(-\hat{q}_{1} \cdot q_{1}')^{\Delta_{1}}} \int d^{2}z_{2}' \frac{1}{|z_{2}' - z_{2}|^{2\Delta_{2}}} \int_{0}^{+\infty} d\omega_{2} \, \omega_{2}^{1-\Delta_{2}} \\ & \times \left(\prod_{k=3}^{n} \int \frac{d^{3}q_{k}'}{q_{k}'^{0}} G_{\Delta_{k}}(\hat{q}_{k}; q_{k}')\right) \mathcal{M}_{2 \to n-2}(q_{i}') , \end{aligned}$$
(3.1)

<sup>8</sup>See Appendix A for the convention of the scattering amplitude in the plane-wave basis.

where we have used (2.18). After changing of variables, we have

$$\widetilde{\mathcal{A}}_{2\to n-2}^{\Delta_{i}}(z_{i}) = 2^{\Delta_{1}+\Delta_{2}} \int d^{4}p \int D^{2}\hat{q}_{2}' \frac{|p|^{2-2\Delta_{1}-2\Delta_{2}}}{(-\hat{q}_{2}\cdot\hat{q}_{2}')^{\Delta_{2}}(-\hat{q}_{2}'\cdot p)^{2-\Delta_{1}-\Delta_{2}}(-\hat{q}_{2}'\cdot Y)^{\Delta_{1}}} \\
\times \left(\prod_{k=3}^{n} \int \frac{d^{3}q_{k}'}{q_{k}'^{0}} G_{\Delta_{k}}(\hat{q}_{k};q_{k}')\right) \mathcal{M}_{2\to n-2}(q_{i}') ,$$
(3.2)

where  $q'_1 + q'_2 \equiv p \equiv M\hat{p}$  with  $\hat{p} \cdot \hat{p} = -1$ ,  $D^2\hat{q}' \equiv d^2z'$  and  $Y^{\mu} \equiv 2(-\hat{q}_1 \cdot \hat{p})\hat{p}^{\mu} - \hat{q}_1^{\mu}$ , and note that  $\omega_2 = \frac{p^2}{2\hat{q}'_2 \cdot p}$ . We have also used the relation

$$\int \frac{d^3 q_1'}{q_1'^0} \int d^2 z_2' \frac{1}{|z_2' - z_2|^{2\Delta_2}} \int_0^{+\infty} d\omega_2 = 2^{\Delta_2} \int d^4 p \int \frac{D^2 \hat{q}_2'}{-\hat{q}_2' \cdot p} \frac{1}{(-\hat{q}_2 \cdot \hat{q}_2')^{\Delta_2}} .$$
(3.3)

In the following subsections, we will start with (3.2) and study the OPE behaviours for different scattering amplitudes  $\mathcal{M}_{2\to n-2}(q'_1, q'_2, \cdots, q'_n)$ .

### 3.1 OPE analysis for three-point shadow celestial amplitudes

We first focus on the three-point shadow celestial amplitudes. Since scattering amplitudes involving three massless scalars cannot satisfy momentum conservation, we will consider the shadow celestial amplitudes for two incoming massless scalars and one outgoing massive scalar. In this case, the scattering amplitudes are

$$i\mathcal{M}_{2\to 1}(q'_1, q'_2, p'_3) = i(2\pi)^4 \delta^{(4)}(p - p'_3)\mathcal{T}_{2\to 1}(p^2) ,$$
 (3.4)

where we defined  $p = q'_1 + q'_2$ . Plugging the above express into (3.2) and changing integral variables from  $p^{\mu}$  to  $M\hat{p}^{\mu}$  then lead to

$$\widetilde{\mathcal{A}}_{2\to1}^{\Delta_i}(z_i) = 2^{\Delta_1 + \Delta_2} \int_0^{+\infty} dM \int \frac{d^3 \hat{p}}{\hat{p}^0} \int D^2 \hat{q}_2' \frac{M^{3-\Delta_1 - \Delta_2}}{(-\hat{q}_2 \cdot \hat{q}_2')^{\Delta_2} (-\hat{q}_2' \cdot \hat{p})^{2-\Delta_1 - \Delta_2} (-\hat{q}_2' \cdot Y)^{\Delta_1}} \times \int \frac{d^3 \hat{p}_3'}{\hat{p}_3'^0} G_{\Delta_3}(\hat{q}_3; \hat{p}_3') (2\pi)^4 \delta^{(4)}(p - p_3') \mathcal{T}_{2\to1}(M^2) .$$
(3.5)

In Appendix B, we compute the integral over  $\hat{q}'_2$  and expand the results around small  $\hat{q}_{12} \equiv -\frac{1}{2}\hat{q}_1 \cdot \hat{q}_2$ . Substituting the expansion (B.6), we get

$$\widetilde{\mathcal{A}}_{2\to1}^{\Delta_{i}} = \frac{(2\pi)^{5}\Gamma[1-\Delta_{1}]\Gamma[1-\Delta_{2}]}{\Gamma[2-\Delta_{1}-\Delta_{2}]} \sum_{n=0}^{\infty} \frac{(\Delta_{1})_{n}(\Delta_{2})_{n}}{(n!)^{2}} \int_{0}^{+\infty} dM M^{3-\Delta_{1}-\Delta_{2}} \mathcal{T}_{2\to1}(M^{2}) \\
\times (\hat{q}_{12})^{n} \int \frac{d^{3}\hat{p}}{\hat{p}^{0}} \int \frac{d^{3}\hat{p}_{3}'}{\hat{p}_{3}'^{0}} G_{\Delta_{1}+n}(\hat{q}_{1};\hat{p}) G_{\Delta_{2}+n}(\hat{q}_{2};\hat{p}) G_{\Delta_{3}}(\hat{q}_{3};\hat{p}_{3}') \delta^{(4)}(p-p_{3}').$$
(3.6)

Rewriting the delta-function as

$$\delta^{(4)}(p - p'_3) = M^{-3}\hat{p}^0\delta(M - m)\delta^{(3)}(\hat{p} - \hat{p}'_3) = M^{-1}p^0\delta(M - m)\delta^{(3)}(p - p'_3)$$
(3.7)

and performing the integral over M and  $\hat{p}_3'$  lead to

$$\widetilde{\mathcal{A}}_{2\to1}^{\Delta_{i}}(z_{i}) = \frac{(2\pi)^{5}\Gamma[1-\Delta_{1}]\Gamma[1-\Delta_{2}]m^{-\Delta_{1}-\Delta_{2}}\mathcal{T}_{2\to1}(m^{2})}{\Gamma[2-\Delta_{1}-\Delta_{2}]}\sum_{n=0}^{\infty}\frac{(\Delta_{1})_{n}(\Delta_{2})_{n}}{(n!)^{2}}(\hat{q}_{12})^{n} \\
\times \int \frac{d^{3}\hat{p}}{\hat{p}^{0}}G_{\Delta_{1}+n}(\hat{q}_{1};\hat{p})G_{\Delta_{2}+n}(\hat{q}_{2};\hat{p})G_{\Delta_{3}}(\hat{q}_{3};\hat{p}) .$$
(3.8)

One can immediately recognize that the second line is exactly the three-point AdS Witten diagram  $W_{\Delta_1+n,\Delta_2+n,\Delta_3}(\hat{q}_i)$ . In the short distance limit  $\hat{q}_1 \rightarrow \hat{q}_2$ ,  $W_{\Delta_1+n,\Delta_2+n,\Delta_3}(\hat{q}_i)$  behaves as

$$W_{\Delta_1+n,\Delta_2+n,\Delta_3}(\hat{q}_i) = \frac{\Gamma[\frac{\Delta_1+\Delta_{23}}{2}+n]\Gamma[\frac{\Delta_{12}+\Delta_3}{2}]\Gamma[\frac{-\Delta_{12}+\Delta_3}{2}]\Gamma[\frac{\sum_{i=1}^3\Delta_i-2}{2}+n]}{\Gamma[\Delta_1+n]\Gamma[\Delta_2+n]\Gamma[\Delta_3]}$$
(3.9)  
 
$$\times (\hat{q}_{12})^{\frac{-\Delta_1-\Delta_2+\Delta_3}{2}-n}(\hat{q}_{23})^{-\Delta_3}[1+O(\hat{q}_{12})] .$$

We arrive

$$\widetilde{\mathcal{A}}_{2\to1}^{\Delta_{i}}(z_{i}) = \frac{(2\pi)^{5}\Gamma[1-\Delta_{1}]\Gamma[1-\Delta_{2}]m^{-\Delta_{1}-\Delta_{2}}\mathcal{T}_{2\to1}(m^{2})}{\Gamma[2-\Delta_{1}-\Delta_{2}]}(\hat{q}_{12})^{\frac{-\Delta_{1}-\Delta_{2}+\Delta_{3}}{2}}(\hat{q}_{23})^{-\Delta_{3}} \times \sum_{n=0}^{\infty} \frac{\Gamma[\frac{\Delta_{1}+\Delta_{23}}{2}+n]\Gamma[\frac{\Delta_{12}+\Delta_{3}}{2}]\Gamma[\frac{-\Delta_{12}+\Delta_{3}}{2}]\Gamma[\frac{-\Delta_{12}+\Delta_{3}}{2}]\Gamma[\frac{\Sigma_{i=1}^{3}\Delta_{i}-2}{2}+n]}{(n!)^{2}\Gamma[\Delta_{1}]\Gamma[\Delta_{2}]\Gamma[\Delta_{3}]}[1+O(\hat{q}_{12})].$$
(3.10)

Finally, computing the summation over n gives

$$\widetilde{\mathcal{A}}_{2\to1}^{\Delta_i} = C_{\Delta_1,\Delta_2,\Delta_3}(m) \hat{q}_{12}^{\frac{-\Delta_1-\Delta_2+\Delta_3}{2}} \hat{q}_{23}^{-\Delta_3} [1+O(\hat{q}_{12})] , \qquad (3.11)$$

where the coefficient  $C_{\Delta_1,\Delta_2,\Delta_3}(m)$  is

$$C_{\Delta_{1},\Delta_{2},\Delta_{3}}(m) = \frac{\mathcal{T}_{2\to1}(m^{2})\Gamma[1-\Delta_{1}]\Gamma[1-\Delta_{2}]\Gamma[\frac{\Delta_{1}+\Delta_{2}}{2}]\Gamma[\frac{\Delta_{1}+\Delta_{3}}{2}]\Gamma[\frac{\Delta_{3}-\Delta_{12}}{2}]\Gamma[\frac{\sum_{i=1}^{3}\Delta_{i}-2}{2}]}{(2\pi)^{-5}m^{\Delta_{1}+\Delta_{2}}\Gamma[\Delta_{1}]\Gamma[\Delta_{2}]\Gamma[\Delta_{3}]\Gamma[\frac{2-\Delta_{1}-\Delta_{23}}{2}]\Gamma[\frac{4-\sum_{i=1}^{3}\Delta_{i}}{2}]}$$
(3.12)

with  $\Delta_{ij} \equiv \Delta_i - \Delta_j$ . From (3.11), we find that in the OPE limit  $\hat{q}_1 \to \hat{q}_2$ ,  $\widetilde{\mathcal{A}}_{2\to 1}^{\Delta_i}$  behave as

$$\lim_{\hat{q}_1 \to \hat{q}_2} \left( \hat{q}_{12}^{\frac{\Delta_1 + \Delta_2 - \Delta_3}{2}} \widetilde{\mathcal{A}}_{2 \to 1}^{\Delta_i}(z_i) \right) = C_{\Delta_1, \Delta_2, \Delta_3}(m) \hat{q}_{23}^{-\Delta_3}.$$
 (3.13)

However, since the OPE coefficient (3.12) diverges when  $\Delta_1 + \Delta_2 - \Delta_3 = -2n$  with  $n \in \mathbb{Z}_{\geq 0}$ , the above limit is only well-defined when  $\Delta_1 + \Delta_2 - \Delta_3 \neq -2n$ .

When  $\Delta_1 + \Delta_2 - \Delta_3 = -2n + 2i\nu$  with  $\nu \in \mathbb{R}$ , we use the following formula

$$\lim_{\hat{q}_1 \to \hat{q}_2} \left( \Gamma[-n+i\nu] \hat{q}_{12}^{-i\nu} \right) = \frac{\Gamma[-n+i\nu]}{\Gamma[i\nu]} \lim_{\hat{q}_1 \to \hat{q}_2} \int_0^\infty dx \; x^{i\nu-1} e^{-x\hat{q}_{12}} = \frac{-2\pi(-1)^n}{n!} \delta(\nu) \;. \tag{3.14}$$

This leads to

$$\lim_{\hat{q}_1 \to \hat{q}_2} \left( \hat{q}_{12}^{-n} \widetilde{\mathcal{A}}_{2 \to 1}^{\Delta_1, \Delta_2, \Delta_1 + \Delta_2 + 2n + i\nu}(z_i) \right) = -2\pi \operatorname{Res}_{\Delta_3 = \Delta_1 + \Delta_2 + 2n} \left( C_{\Delta_1, \Delta_2, \Delta_3}(m) \right) \delta(\nu) \hat{q}_{23}^{-\Delta_3} . \quad (3.15)$$

For later convenience, we define the OPE coefficient  $\mathcal{C}_{\Delta_1,\Delta_2,\Delta_1+\Delta_2+2n}(m)$  as

$$\mathcal{C}_{\Delta_1,\Delta_2,\Delta_3}(m) = \begin{cases}
-2\pi \operatorname{Res}_{\Delta_3 = \Delta_1 + \Delta_2 + 2n} \left( C_{\Delta_1,\Delta_2,\Delta_3}(m) \right) & \text{for } \Delta_3 = \Delta_1 + \Delta_2 + 2n, \\
C_{\Delta_1,\Delta_2,\Delta_3}(m) & \text{for } \Delta_3 \neq \Delta_1 + \Delta_2 + 2n,
\end{cases}$$
(3.16)

with  $n \in \mathbb{Z}_{\geq 0}$ .

From (3.15) and (3.16) we see that apart from the delta-function  $\delta(\nu)$ , the OPE limit of  $\widetilde{\mathcal{A}}_{2\to 1}^{\Delta_1,\Delta_2,\Delta_3}(z_i)$  is finite when  $\Delta_3 = \Delta_1 + \Delta_2$ . In the following subsections, we will focus on the finite parts in the OPE limit and study the OPE behaviour of  $n \geq 3$ -point shadow celestial amplitudes.

### **3.2** OPE analysis from the generalized optical theorem

The leading order OPE behaviour can be obtained through replacing all  $\hat{q}_1$  in (3.2) by  $\hat{q}_2$ , leading to

$$\lim_{\hat{q}_{1} \to \hat{q}_{2}} \widetilde{\mathcal{A}}_{2 \to n-2}^{\Delta_{i}}(z_{i}) = 2^{\Delta_{1}+\Delta_{2}} \int d^{4}p \int D^{2} \hat{q}_{2}' \frac{|p|^{2-2\Delta_{1}-2\Delta_{2}}}{(-\hat{q}_{2} \cdot \hat{q}_{2}')^{\Delta_{2}}(-\hat{q}_{2}' \cdot p)^{2-\Delta_{1}-\Delta_{2}}(-\hat{q}_{2}' \cdot Y')^{\Delta_{1}}} \times \left(\prod_{k=3}^{n} \int \frac{d^{3}q_{k}'}{q_{k}'^{0}} G_{\Delta_{k}}(\hat{q}_{k};q_{k}')\right) (2\pi)^{4} \delta^{(4)}(p-\sum_{k=3}^{n} q_{k}') \mathcal{T}_{2 \to n-2}(q_{i}') ,$$

$$(3.17)$$

where  $Y'^{\mu} \equiv 2(-\hat{q}_2 \cdot \hat{p})\hat{p}^{\mu} - \hat{q}_2^{\mu}$ . For  $n \geq 3$ -point shadow celestial amplitudes, the OPE limit can be studied by the generalized optical theorems. We assume  $\mathcal{M}_{2\to n-2}(q'_1, q'_2, \cdots, q'_n)$  is at tree-level. Using the generalized optical theorem (A.9) and noting that there are only single-particle intermediate states that can appear in tree diagrams, we get <sup>9</sup>

$$\lim_{\hat{q}_{1}\to\hat{q}_{2}} \operatorname{Im} \widetilde{\mathcal{A}}_{2\to n-2}^{\Delta_{i}} = \int d^{4}p \int \frac{D^{2}\hat{q}_{2}' \, 2^{\Delta_{1}+\Delta_{2}-1} |p|^{2-2\Delta_{1}-2\Delta_{2}}}{(-\hat{q}_{2}'\cdot p)^{2-\Delta_{1}-\Delta_{2}} (-\hat{q}_{2}'\cdot Y')^{\Delta_{1}}} \left(\prod_{k=3}^{n} \int \frac{d^{3}q_{k}'}{q_{k}'^{0}} G_{\Delta_{k}}(\hat{q}_{k};q_{k}')\right) \times \sum_{X} \int \frac{d^{3}p_{X}}{(2\pi)^{3}} \frac{1}{2E_{X}} (2\pi)^{4} \delta^{(4)}(p-p_{X}) \mathcal{T}_{2\to X}(p^{2}) \mathcal{M}_{X\to n-2}^{*}(p_{X},q_{i}') .$$
(3.18)

Here X labels all possible physical single-particle states that satisfy the on-shell condition  $p_X^2 = -m_X^2$ . With the help of the delta function, we evaluate the integral over p, leading to

$$\lim_{\hat{q}_{1}\to\hat{q}_{2}} \operatorname{Im} \widetilde{\mathcal{A}}_{2\to n-2}^{\Delta_{1}} = 2^{\Delta_{1}+\Delta_{2}} \pi \sum_{X} \int \frac{d^{3}\hat{p}_{X}}{2\hat{p}_{X}^{0}} \int D^{2}\hat{q}_{2}' \frac{m_{X}^{2-\Delta_{1}-\Delta_{2}}\mathcal{T}_{2\to X}(m_{X}^{2})}{(-\hat{q}_{2}\cdot\hat{q}_{2}')^{\Delta_{2}}(-\hat{q}_{2}'\cdot\hat{p}_{X})^{2-\Delta_{1}-\Delta_{2}}(-\hat{q}_{2}'\cdot Y')^{\Delta_{1}}} \\ \times \left(\prod_{k=3}^{n} \int \frac{d^{3}q_{k}'}{q_{k}'^{0}} G_{\Delta_{k}}(\hat{q}_{k};q_{k}')\right) \mathcal{M}_{X\to n-2}^{*}(m_{X}\hat{p}_{X},q_{i}') .$$

$$(3.19)$$

After evaluating the integral over  $\hat{q}'_2$  by using (B.7), we get

$$\lim_{\hat{q}_1 \to \hat{q}_2} \operatorname{Im} \widetilde{\mathcal{A}}_{2 \to n-2}^{\Delta_i} = \sum_X \frac{\pi^2 \Gamma[1 - \Delta_1] \Gamma[1 - \Delta_2]}{\Gamma[2 - \Delta_1 - \Delta_2]} m_X^{2 - \Delta_1 - \Delta_2} \mathcal{T}_{2 \to X}(m_X^2) \widetilde{\mathcal{A}}_{X \to n-2}^{*\Delta_1 + \Delta_2, \Delta_i}$$

$$= \frac{1}{2} \sum_X \mathcal{C}_{\Delta_1, \Delta_2, \Delta_1 + \Delta_2}(m_X) D^{\Delta_1 + \Delta_2, \Delta_1 + \Delta_2}(m_X) \widetilde{\mathcal{A}}_{X \to n-2}^{*\Delta_1 + \Delta_2, \Delta_i} , \qquad (3.20)$$

<sup>&</sup>lt;sup>9</sup>Here and throughout this paper, we assume that conformal dimensions  $\Delta_i$  are real when taking the imaginary part of  $\widetilde{\mathcal{A}}_{2\to n-2}^{\Delta_i}(z_i)$ .



Figure 1: A particular class of Feynman diagrams which contain a massive propagator  $\frac{1}{-p^2-m^2+i\epsilon}$  with  $p = q'_1 + q'_2$ .

where  $D^{\Delta\Delta}(m) = \frac{(\Delta-1)m^2}{(2\pi)^5}$  is the inverse of coefficient of two-point massive celestial amplitude [28] and  $C_{\Delta_1,\Delta_2,\Delta_1+\Delta_2}(m_X)$  is given in (3.16)

Some comments are made here. The leading OPE behaviour (3.20) works only when the (n-1)-point shadow celestial amplitudes  $\widetilde{\mathcal{A}}_{X\to n-2}^{\Delta_1+\Delta_2,\Delta_i}(m_X;z_i)$  are well-defined. This demands that the integrals appearing in the computation of the (n-1)-point shadow celestial amplitudes  $\widetilde{\mathcal{A}}_{X\to n-2}^{\Delta_1+\Delta_2,\Delta_i}(m_X;z_i)$  must converge. For example, in the case of n = 4, the leading OPE behaviour (3.20) works only when  $\operatorname{Re}(\Delta_3 + \Delta_4 - \Delta_1 - \Delta_2) > 0$ , beyond which the integrals appearing in  $\widetilde{\mathcal{A}}_{1\to 2}^{\Delta_1+\Delta_2,\Delta_3,\Delta_4}(m_X;z_i)$  are divergent. Although one can still define  $\widetilde{\mathcal{A}}_{1\to 2}^{\Delta_1+\Delta_2,\Delta_3,\Delta_4}(m_X;z_i)$  by analytical continuation when  $\operatorname{Re}(\Delta_3 + \Delta_4 - \Delta_1 - \Delta_2) \leq 0$ , (3.20) is no longer dominant in this region.

#### 3.3 OPE analysis for a special class of Feynman diagrams

In the previous subsection, we used the generalized optical theorem to decompose a *n*-point scattering amplitude into a 3-point and a (n-1)-point scattering amplitudes. The shortcoming is that we only obtained the OPE limit of the imaginary part of the shadow celestial amplitude  $\widetilde{\mathcal{A}}_{2\to n-2}^{\Delta_i}$ . In this subsection, we will focus on a special class of Feynman diagrams shown in Figure 1. It allows us to derive a formula for the OPE limit of the shadow celestial amplitude without taking the imaginary part. We note that the scattering amplitudes of the class of Feynman diagrams in Figure 1 take the form

$$\mathcal{M}_{2\to n-2}(q_i') = (2\pi)^4 \delta^{(4)}(p - \sum_{k=3}^n q_k') i\mathcal{T}_{2\to 1}(p^2) \frac{1}{-p^2 - m^2} i\mathcal{T}_{1\to n-2}(p, q_i') , \qquad (3.21)$$

where we defined  $p = q'_1 + q'_2$ . Plugging the above express into (3.2) and changing integral variables from  $p^{\mu}$  to  $M\hat{p}^{\mu}$  with  $\hat{p} \cdot \hat{p} = -1$  then lead to

$$\lim_{\hat{q}_1 \to \hat{q}_2} \widetilde{\mathcal{A}}_{2 \to n-2}^{\Delta_i} = -2^{\Delta_1 + \Delta_2} \int_0^{+\infty} dM \int \frac{d^3 \hat{p}}{\hat{p}^0} \int \frac{D^2 \hat{q}_2' M^{3 - \Delta_1 - \Delta_2}}{(-\hat{q}_2 \cdot \hat{q}_2')^{\Delta_2} (-\hat{q}_2' \cdot \hat{p})^{2 - \Delta_1 - \Delta_2} (-\hat{q}_2' \cdot Y')^{\Delta_1}} \times \left(\prod_{k=3}^n \int \frac{d^3 q_k'}{q_k'^0} G_{\Delta_k}(\hat{q}_k; q_k')\right) \mathcal{T}_{2 \to 1}(M^2) \frac{1}{M^2 - m^2} \mathcal{M}_{1 \to n-2}(p, q_i') .$$
(3.22)

Performing the integral over  $\hat{q}'_2$ , we obtain

$$\lim_{\hat{q}_{1}\to\hat{q}_{2}} \widetilde{\mathcal{A}}_{2\to n-2}^{\Delta_{i}} = \frac{-2\pi\Gamma[1-\Delta_{1}]\Gamma[1-\Delta_{2}]}{\Gamma[2-\Delta_{1}-\Delta_{2}]} \int_{0}^{+\infty} dM \frac{M^{3-\Delta_{1}-\Delta_{2}}\mathcal{T}_{2\to1}(M^{2})}{M^{2}-m^{2}} \widetilde{\mathcal{A}}_{M}^{\Delta_{1}+\Delta_{2},\Delta_{i}} = -\frac{1}{\pi} \int_{0}^{+\infty} dM \frac{M\mathcal{C}_{\Delta_{1},\Delta_{2},\Delta_{1}+\Delta_{2}}(M)D^{\Delta_{1}+\Delta_{2},\Delta_{1}+\Delta_{2}}(M)}{M^{2}-m^{2}} \widetilde{\mathcal{A}}_{M}^{\Delta_{1}+\Delta_{2},\Delta_{i}} .$$
(3.23)

Thus, for the particular class of diagrams in Figure 1, the leading OPE limit of a *n*-point massless shadow celestial amplitudes are given by a superposition of (n - 1)-point shadow celestial amplitudes over a range of mass  $M \in [0, \infty)$ . (3.23) leads to the leading order operator product expansion

$$\mathcal{O}_1(z_1)\mathcal{O}_2(z_2) \sim -\frac{1}{\pi} \int_0^{+\infty} dM \frac{M\mathcal{C}_{\Delta_1,\Delta_2,\Delta_1+\Delta_2}(M)D^{\Delta_1+\Delta_2,\Delta_1+\Delta_2}(M)}{M^2 - m^2} \Phi_M^{\Delta_1+\Delta_2}(z_2) \,. \tag{3.24}$$

We stress here again that (3.23) works only when the (n-1)-point shadow celestial amplitudes  $\widetilde{\mathcal{A}}_{M}^{\Delta_{1}+\Delta_{2},\Delta_{i}}$  are well-defined.

### **3.4 OPE** analysis for four-point shadow celestial amplitudes

As an example, we consider the OPE behaviour of four-point shadow celestial amplitudes  $\widetilde{\mathcal{A}}_{2\to 2}^{\Delta_i}$  involving two incoming and two outgoing massless scalars with a massive scalar exchange. Since only *s*-channel amplitude has an imaginary part and is in our particular class of diagrams shown in Figure 1, we only focus on the *s*-channel amplitude and take the OPE limit  $\hat{q}_1 \to \hat{q}_2$ . The OPE limit  $\hat{q}_3 \to \hat{q}_4$  can be derived in a similar way. In this specific case, (3.20) takes the form as

$$\lim_{\hat{q}_1 \to \hat{q}_2} \operatorname{Im} \widetilde{\mathcal{A}}_s^{\Delta_i}(z_i) = \frac{1}{2} \mathcal{C}_{\Delta_1, \Delta_2, \Delta_1 + \Delta_2}(m) D^{\Delta_1 + \Delta_2, \Delta_1 + \Delta_2}(m) \widetilde{\mathcal{A}}_m^{\Delta_3, \Delta_4, \Delta_1 + \Delta_2} .$$
(3.25)

Here m denotes the mass of the exchange operator. On the other hand, in this case, (3.23) becomes

$$\lim_{\hat{q}_1 \to \hat{q}_2} \widetilde{\mathcal{A}}_{2 \to 2}^{\Delta_i} = -\frac{1}{\pi} \int_0^{+\infty} dM \frac{M \mathcal{C}_{\Delta_1, \Delta_2, \Delta_1 + \Delta_2}(M) D^{\Delta_1 + \Delta_2, \Delta_1 + \Delta_2}(M)}{M^2 - m^2} \widetilde{\mathcal{A}}_M^{\Delta_1 + \Delta_2, \Delta_3, \Delta_4} .$$
(3.26)

We note that in the present case  $\widetilde{\mathcal{A}}_M^{\Delta_1+\Delta_2,\Delta_i}(z_i)$  takes the form as

$$\widetilde{\mathcal{A}}_{M}^{\Delta_{1}+\Delta_{2},\Delta_{3},\Delta_{4}} = -\frac{g}{2^{\Delta_{3}+\Delta_{4}-4}} \int \frac{d^{2}z_{3}'}{|z_{3}-z_{3}'|^{2\Delta_{3}}} \frac{d^{2}z_{4}'}{|z_{4}-z_{4}'|^{2\Delta_{4}}} \int \frac{d^{3}\hat{p}}{\hat{p}^{0}} \frac{1}{(-\hat{q}_{2}\cdot\hat{p})^{\Delta_{1}+\Delta_{2}}} \\ \times \int_{0}^{\infty} d\omega_{3}\,\omega_{3}^{1-\Delta_{3}} \int_{0}^{\infty} d\omega_{4}\,\omega_{4}^{1-\Delta_{4}}(2\pi)^{4}\delta^{(4)}(M\hat{p}-\omega_{3}\hat{q}_{3}'-\omega_{4}\hat{q}_{4}') \,.$$
(3.27)

Rescaling  $\omega_3 \to M\omega_3$  and  $\omega_4 \to M\omega_4$  leads to

$$\widetilde{\mathcal{A}}_{M}^{\Delta_{1}+\Delta_{2},\Delta_{3},\Delta_{4}}(z_{i}) = \frac{-gM^{-\Delta_{3}-\Delta_{4}-2}}{2^{\Delta_{3}+\Delta_{4}-4}} \int \frac{d^{2}z_{3}'}{|z_{3}-z_{3}'|^{2\Delta_{3}}} \frac{d^{2}z_{4}'}{|z_{4}-z_{4}'|^{2\Delta_{4}}} \int \frac{d^{3}\hat{p}}{\hat{p}^{0}} \frac{1}{(-\hat{q}_{2}\cdot\hat{p})^{\Delta_{1}+\Delta_{2}}} \\
\times \int_{0}^{\infty} d\omega_{3}\omega_{3}^{1-\Delta_{3}} \int_{0}^{\infty} d\omega_{4}\omega_{4}^{1-\Delta_{4}}(2\pi)^{4}\delta^{(4)}(\hat{p}-\omega_{3}\hat{q}_{3}'-\omega_{4}\hat{q}_{4}') \qquad (3.28) \\
= M^{-\Delta_{3}-\Delta_{4}}\widetilde{\mathcal{A}}_{M=1}^{\Delta_{1}+\Delta_{2},\Delta_{3},\Delta_{4}}(z_{i}) \,.$$

Plugging the above expression into (3.23) then leads to

$$\lim_{\hat{q}_1 \to \hat{q}_2} \widetilde{\mathcal{A}}_s^{\Delta_i} = -\frac{1}{\pi} \int_0^{+\infty} dM \frac{M^{-\beta-1} \mathcal{C}_{\Delta_1, \Delta_2, \Delta_1 + \Delta_2}(1) D^{\Delta_1 + \Delta_2, \Delta_1 + \Delta_2}(1)}{M^2 - m^2} \widetilde{\mathcal{A}}_{M=1}^{\Delta_1 + \Delta_2, \Delta_3, \Delta_4}(z_i) = \frac{e^{-i\pi\beta/2}}{2\sin(\frac{-\pi\beta}{2})} \mathcal{C}_{\Delta_1, \Delta_2, \Delta_1 + \Delta_2}(m) D^{\Delta_1 + \Delta_2, \Delta_1 + \Delta_2}(m) \widetilde{\mathcal{A}}_m^{\Delta_1 + \Delta_2, \Delta_3, \Delta_4},$$
(3.29)

where we defined  $\beta = 4 - \Delta_1 - \Delta_2 - \Delta_3 - \Delta_4$ . We stress here that (3.25) and (3.29) hold only when  $\widetilde{\mathcal{A}}_m^{\Delta_1 + \Delta_2, \Delta_3, \Delta_4}$  is finite, which constrains the conformal dimensions by  $\operatorname{Re}(\Delta_1 + \Delta_2 - \Delta_3 - \Delta_4) < 0$ .

By further expanding (3.29) in power series of  $\hat{q}_{34}$  and using (3.13), we find

$$\lim_{\hat{q}_{1}\to\hat{q}_{2}} \widetilde{\mathcal{A}}_{s}^{\Delta_{i}} = \frac{e^{-i\pi\beta/2}}{2\sin(\frac{-\pi\beta}{2})} \mathcal{C}_{\Delta_{1},\Delta_{2},\Delta_{1}+\Delta_{2}}(m) D^{\Delta_{1}+\Delta_{2},\Delta_{1}+\Delta_{2}}(m) \mathcal{C}_{\Delta_{3},\Delta_{4},\Delta_{1}+\Delta_{2}}(m) \times \hat{q}_{24}^{-\Delta_{1}-\Delta_{2}} \hat{q}_{34}^{\frac{\Delta_{1}+\Delta_{2}-\Delta_{3}-\Delta_{4}}{2}} (1+\mathcal{O}(\hat{q}_{34}))$$
(3.30)

In the next section, we will compute the s-channel four-point shadow celestial amplitude and derive its conformal block expansion. We will find agreement between the conformal block expansion and the OPE expansion (3.30) when  $\operatorname{Re}(\Delta_1 + \Delta_2 - \Delta_3 - \Delta_4) < 0$ .

### 4 Examples of shadow celestial amplitudes

#### 4.1 Three-point shadow celestial amplitudes

We first consider the shadow celestial amplitudes for two incoming massless scalars and one outgoing massive scalar. The celestial amplitudes  $\mathcal{A}_{2\to 1}^{\Delta_i}$  at tree level is [24]

$$\mathcal{A}_{2\to1}^{\Delta_i}(z_i) = -g \frac{\pi^4 m^{\Delta_1 + \Delta_2 - 4}}{2^{\Delta_1 + \Delta_2 - 5}} \frac{B(\frac{\Delta_1 + \Delta_{12}}{2}, \frac{\Delta_1 - \Delta_{12}}{2})}{|z_{12}|^{\Delta_1 + \Delta_2 - \Delta_3} |z_{13}|^{\Delta_1 - \Delta_2 + \Delta_3} |z_{23}|^{\Delta_2 + \Delta_3 - \Delta_1}}, \qquad (4.1)$$

where  $B(a, b) \equiv \Gamma[a]\Gamma[b]/\Gamma[a+b]$  is the beta-function. With the help of (A.25), we compute the three-point shadow celestial amplitudes  $\widetilde{\mathcal{A}}_{2\to 1}^{\Delta_i}$ . We get

$$\widetilde{\mathcal{A}}_{2\to1}^{\Delta_i}(z_i) = \frac{C_{\Delta_1,\Delta_2,\Delta_3}}{|z_{12}|^{\Delta_1 + \Delta_2 - \Delta_3} |z_{13}|^{\Delta_1 - \Delta_2 + \Delta_3} |z_{23}|^{\Delta_2 + \Delta_3 - \Delta_1}}, \qquad (4.2)$$

where the coefficient  $C_{\Delta_1,\Delta_2,\Delta_3}$  is given in (3.12) with  $\mathcal{T}_{2\to 1}(m^2) = -g$  and  $z_{ij} \equiv z_i - z_j$  and  $\bar{z}_{ij} \equiv \bar{z}_i - \bar{z}_j$ . We mention here that the integral in the shadow transformation from (4.1) to (4.2) converges only when  $\operatorname{Re}(\Delta_1 + \Delta_2 - \Delta_3) > 0$ . However, with (4.2) in hand, we can analytically continue the conformal dimensions to the whole complex plane excluding the poles of  $C_{\Delta_1,\Delta_2,\Delta_3}$ .

The constrains imposed by translation symmetries (2.28) on the shadow celestial amplitude  $\widetilde{\mathcal{A}}_{2\to 1}^{\Delta_i}$  are

$$\begin{split} \left[ (\Delta_{12} + \Delta_3 - 1) (\sum_{i=1}^3 \Delta_i - 3)^2 e^{-\partial_{\Delta_1}} - 8(\Delta_1 - 1)^2 \Delta_3 m e^{\partial_{\Delta_3}} \right] C_{\Delta_1, \Delta_2, \Delta_3}(m) &= 0 , \\ \left[ (-\Delta_{12} + \Delta_3 - 1) (\sum_{i=1}^3 \Delta_i - 3)^2 e^{-\partial_{\Delta_2}} - 8(\Delta_2 - 1)^2 \Delta_3 m e^{\partial_{\Delta_3}} \right] C_{\Delta_1, \Delta_2, \Delta_3}(m) &= 0 , \\ \left[ (\Delta_{12} - \Delta_3 + 1) (\Delta_{12} + \Delta_3 - 1) (\sum_{i=1}^3 \Delta_i - 3)^2 e^{-\partial_{\Delta_3}} + 4\Delta_3 (\Delta_1 + \Delta_2 - \Delta_3 - 1)^2 (\Delta_3 - 1) e^{\partial_{\Delta_3}} \right] C_{\Delta_1, \Delta_2, \Delta_3}(m) = 0 . \end{split}$$
(4.3)

It is then straightforward to confirm that the coefficient (3.12) satisfies the above equality. One can also check that the OPE behaviour of (4.2) agrees with our analysis in (3.13) when  $\Delta_3 \neq \Delta_1 + \Delta_2 + 2n$  and (3.15) when  $\Delta_3 = \Delta_1 + \Delta_2 + 2n$  after setting  $\mathcal{T}_{2\to 1}(m^2) = -g$ .

### 4.2 Four-point shadow celestial amplitudes

In this subsection, we will compute the four-point shadow celestial amplitudes  $\tilde{\mathcal{A}}_{12\to34}^{\Delta_i}$  of four external massless scalars and one exchange massive scalar, and derive its conformal block expansion in the 12 – 34 channel. To achieve this, we start with the corresponding celestial amplitudes  $\mathcal{A}_{12\to34}^{\Delta_i}$  with the explicit formulae given in (2.8), (A.18) and (2.12), and perform the shadow transformation for each external operator.

#### 4.2.1 Shadow transformation for $\mathcal{O}_1$

To get the shadow celestial amplitudes  $\widetilde{\mathcal{A}}_s^{\Delta_i}$ , we first perform the shadow transformation on the operator  $\mathcal{O}_1$ , and denote the result by  $\widehat{\mathcal{A}}_s^{\Delta_i}$ .<sup>10</sup> Using the conformal symmetry, we can fix three of the four coordinates, leading to

$$z_1 = \bar{z}_1 = \chi \in (1, +\infty)$$
,  $z_2 = \bar{z}_2 = 0$ ,  $z_3 = \bar{z}_3 = +\infty$ ,  $z_4 = \bar{z}_4 = 1$ . (4.4)

In this coordinate configuration,  $I_{12-34}$  in (A.18) takes the form as

$$I_{12-34} = |\chi|^{-2h_1 - 2h_2} |\chi - 1|^{2h_{34} - 2h_{12}} \frac{|z_{24}|^{2h_{12}} |z_3|^{-2h_{34}}}{|z_{34}|^{2h_3 + 2h_4}}, \qquad (4.5)$$

where  $\Delta_i = 2h_i$ . Together with (2.8) and (2.12), we simplify the shadow integral as

$$\begin{aligned} \widehat{\mathcal{A}}_{s}^{\widetilde{\Delta}_{1},\Delta_{2},\Delta_{3},\Delta_{4}} &= 2^{\Delta_{1}} \int_{1}^{+\infty} \frac{d^{2}z_{1}}{(z_{1}-z_{1}')^{2\widetilde{h}_{1}}(\bar{z}_{1}-\bar{z}_{1}')^{2\widetilde{h}_{1}}} I_{12-34} f_{s}(\chi) \\ &= 2^{\Delta_{1}} N_{gm}(\beta) \frac{|z_{24}|^{2h_{12}} |z_{3}|^{-2h_{34}}}{|z_{34}|^{2h_{3}+2h_{4}}} \int_{1}^{\infty} \frac{d^{2}\chi \delta(\chi-\bar{\chi})\chi^{2\widetilde{h}_{1}-2h_{2}}(\chi-1)^{h_{34}-h_{12}}}{(\chi-\chi')^{2\widetilde{h}_{1}}(\bar{\chi}-\bar{\chi}')^{2\widetilde{h}_{1}}} \\ &= 2^{\Delta_{1}} N_{gm}(\beta) \frac{|z_{24}|^{2h_{12}} |z_{3}|^{-2h_{34}}}{|z_{34}|^{2h_{3}+2h_{4}}} \int_{0}^{1} \frac{d\chi \chi^{h_{1'}+h_{2}-h_{34}-1}(1-\chi)^{h_{34}-h_{12}}}{(1-\chi\chi')^{2h_{1'}}(1-\chi\bar{\chi}')^{2\overline{h}_{1'}}} , \end{aligned}$$
(4.6)

where  $\widetilde{\Delta}_1 = 2 - \Delta_1$  and  $\widetilde{h}_1 = 1 - h_1$ , and we defined the new conformal cross-ratios  $\chi'$  and  $\overline{\chi}'$  as

$$\chi' = \frac{z_{1'2}z_{34}}{z_{1'3}z_{24}}, \qquad \bar{\chi}' = \frac{\bar{z}_{1'2}\bar{z}_{34}}{\bar{z}_{1'3}\bar{z}_{24}}, \qquad z_{i'j} \equiv z'_i - z_j .$$
(4.7)

<sup>10</sup> See [36, 37] for performing the shadow transformation on one of the operators in four-point gluon amplitudes.

The above integral can be evaluated by using the integral representation of the Appell function  $F_1$ :

$$F_1 \begin{pmatrix} a; b_1, b_2 \\ c \end{pmatrix} = \frac{\Gamma[c]}{\Gamma[a]\Gamma[c-a]} \int_0^1 dt t^{a-1} (1-t)^{c-a-1} (1-tx)^{-b_1} (1-ty)^{-b_2} , \quad (4.8)$$

leading to

$$\begin{aligned} \widehat{\mathcal{A}}_{s}^{\widetilde{\Delta}_{1},\Delta_{2},\Delta_{3},\Delta_{4}} =& 2^{\Delta_{1}} N_{gm}(\beta) \frac{|z_{24}|^{2h_{12}} |z_{3}|^{-2h_{34}}}{|z_{34}|^{2h_{3}+2h_{4}}} B\left(\widetilde{h}_{1}+h_{2}-h_{34},\widetilde{h}_{1}+h_{2}+h_{34}\right) \\ &\times F_{1}\left(\begin{array}{c} \widetilde{h}_{1}+h_{2}-h_{34}; 2\widetilde{h}_{1}, 2\widetilde{h}_{1} \\ 2\widetilde{h}_{1}+2h_{2} \end{array} \middle| \chi', \bar{\chi}' \right) . \end{aligned}$$

$$(4.9)$$

Using the conformal symmetry, one can unfix the coordinates (4.4), and reach the following expression

$$\widehat{\mathcal{A}}_{s}^{\Delta_{i}}(z_{i}) = I_{12-34} \mathcal{G}_{s}(\chi) , \qquad (4.10)$$

where we rename all of the primed (tilde) variables by the corresponding unprimed (untilde) variables to simplify the notation and  $\mathcal{G}_s(\chi)$  is

$$\mathcal{G}_{s}(\chi) = 2^{2-\Delta_{1}} N_{gm}(\beta) B \left(h_{1} + h_{2} - h_{34}, h_{1} + h_{2} + h_{34}\right) \\ \times (\chi \bar{\chi})^{h_{1}+h_{2}} |1-\chi|^{2(h_{13}-h_{24})} F_{1} \left(\begin{array}{c} h_{1} + h_{2} - h_{34}; 2h_{1}, 2h_{1} \\ 2h_{1} + 2h_{2} \end{array} \middle| \chi, \bar{\chi} \right) ,$$

$$(4.11)$$

where we have renamed  $\widetilde{\Delta}_1$ ,  $\widetilde{h}_1$  by  $\Delta_1$ ,  $h_1$ , and in this convention  $\beta = -\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4 - 2$ . To find the conformal block expansion in the 12–34 channel, we use the Burchnall-Chaundy expansion for the Appell function

$$F_{1}\left(\begin{array}{c}a;b_{1},b_{2}\\c\end{array}\middle|x,y\right) = \sum_{n=0}^{+\infty} \frac{(a)_{n}(b_{1})_{n}(b_{2})_{n}(c-a)_{n}}{n!(c+n-1)_{n}(c)_{2n}}x^{n}y^{n} \\ \times {}_{2}F_{1}\left(\begin{array}{c}a+n,b_{1}+n\\c+2n\end{array}\middle|x\right) {}_{2}F_{1}\left(\begin{array}{c}a+n,b_{2}+n\\c+2n\end{array}\middle|y\right),$$
(4.12)

leading to

$$\mathcal{G}_{s}(\chi) = 2^{2-\Delta_{1}} N_{gm}(\beta) \sum_{n=0}^{+\infty} \frac{B(h-h_{34},h+h_{34})(h+h_{12}-n)_{n}(h+h_{12}-n)_{n}}{n!(2h-n-1)_{n}} \times |1-\chi|^{2(h_{13}-h_{24})} (\chi\bar{\chi})^{h}{}_{2}F_{1}\left(\begin{array}{c}h-h_{34},h+h_{12}\\2h\end{array}\right) \chi 2F_{1}\left(\begin{array}{c}\bar{h}-\bar{h}_{34},\bar{h}+\bar{h}_{12}\\2\bar{h}\end{array}\right) \chi,$$

$$(4.13)$$

where we defined

$$h = \bar{h} = h_1 + h_2 + n$$
,  $n \in \mathbb{Z}_{\geq 0}$ . (4.14)

Finally, after using the following identity

$${}_{2}F_{1}\left(\begin{array}{c}a,b\\c\end{array}\middle|x\right) = (1-x)^{c-a-b}{}_{2}F_{1}\left(\begin{array}{c}c-a,c-b\\c\end{array}\middle|x\right),\qquad(4.15)$$

we get the conformal block expansion of  $\mathcal{G}_s(\chi)$  in 12-34 channel,

$$\mathcal{G}_{s}(\chi) = N_{gm}(\beta) \sum_{n=0}^{+\infty} \frac{B(h_{1+2} - h_{34} + n, h_{1+2} + h_{34} + n)\Gamma[2h_{1} + n]^{2}}{2^{\Delta_{1} - 2}n!\Gamma[2h_{1}]\Gamma[1 - 2h_{1}](2h_{1+2} + n - 1)_{n}} G^{h_{i},\bar{h}_{i}}_{h_{1+2} + n,h_{1+2} + n}(\chi) .$$

$$(4.16)$$

Here, we introduced the notation  $h_{i+j} \equiv h_i + h_j$  and the conformal block in s-channel is given by

$$G_{h,\bar{h}}^{h_i,\bar{h}_i}(\chi) = \chi^h_{\ 2} F_1 \left( \begin{array}{c} h - h_{12}, h + h_{34} \\ 2h \end{array} \middle| \chi \right) \bar{\chi}^{\bar{h}}_{\ 2} F_1 \left( \begin{array}{c} \bar{h} - \bar{h}_{12}, \bar{h} + \bar{h}_{34} \\ 2\bar{h} \end{array} \middle| \bar{\chi} \right) .$$
(4.17)

### 4.2.2 Shadow transformation for $\mathcal{O}_2, \mathcal{O}_3$ and $\mathcal{O}_4$

To get the shadow celestial amplitude  $\widetilde{\mathcal{A}}_s^{\Delta_i}$ , we still need perform the shadow transformations on the remaining three operators in  $\widehat{\mathcal{A}}_s^{\Delta_i}$ . However, computing the shadow integral as we did for  $\mathcal{O}_1$  now becomes extremely hard. Instead of computing the shadow integral directly, in this subsection, we will use the conformal partial wave reviewed in the Appendix A.2 to get the conformal block expansion of  $\widetilde{\mathcal{A}}_s^{\Delta_i}$ . Specifically, we first derive the conformal partial expansion of  $\widehat{\mathcal{A}}_s^{\Delta_i}$  from the block expansion (4.16). Then we compute the conformal partial wave expansion of  $\widetilde{\mathcal{A}}_s^{\Delta_i}$  by virtue of (A.28). Finally, we derive the desired conformal block expansion by closing the contour to the right-half plane.

From the block expansion (4.16) and the symmetry property (A.19), we can derive the spectral density  $\hat{\rho}_{h,\bar{h}}^{\Delta_i}$  for  $\hat{\mathcal{A}}_s^{\Delta_i}$ ,

$$K_{1-h,1-\bar{h}}^{\Delta_{3},\Delta_{4}} \frac{\hat{\rho}_{h,\bar{h}}^{\tilde{\Delta}_{1},\Delta_{2},\Delta_{3},\Delta_{4}}}{n_{h,\bar{h}}} = -\frac{g^{2}\pi^{9}m^{\beta-4}}{2^{\beta-6}} \frac{e^{i\pi\beta/2}}{\sin\pi\beta/2} \frac{a_{\Delta}^{\Delta_{1},\Delta_{2}}}{2^{-\Delta_{1}}} \times B(h+h_{12},h-h_{12})B(h-h_{34},h+h_{34})D^{\Delta\Delta}(m) , \qquad (4.18)$$

where  $\Delta = 2h = 2\bar{h}$  and  $a_{\Delta_3}^{\Delta_1,\Delta_2}$  is given in (A.26). Using (A.28), we can also obtain the spectral density  $\tilde{\rho}_{h,\bar{h}}^{\Delta_i}$  of the shadow celestial amplitudes  $\tilde{\mathcal{A}}^{\Delta_i}$  by performing shadow transformation on  $\mathcal{O}_2$ ,  $\mathcal{O}_3$  and  $\mathcal{O}_4$ , giving that

$$\frac{\tilde{\rho}_{h,\bar{h}}^{\Delta_i}}{n_{h,\bar{h}}} K_{1-h,1-\bar{h}}^{2-\Delta_3,2-\Delta_4} = \frac{\tilde{\rho}_{h,\bar{h}}^{\Delta_i}}{n_{h,\bar{h}}} K_{1-h,1-\bar{h}}^{\Delta_3,\Delta_4} = -\pi \frac{e^{-i\pi\beta/2}}{\sin(\frac{-\pi\beta}{2})} C_{\Delta_1,\Delta_2,\Delta}(m) C_{\Delta_3,\Delta_4,\Delta}(m) D^{\Delta\Delta}(m) , \quad (4.19)$$

where  $C_{\Delta_i,\Delta_j,\Delta}(m)$  is given in (3.12) and we used the fact that  $\beta = \sum_{i=1}^{4} \Delta_i - 4 \rightarrow -\beta = 4 - \sum_{i=1}^{4} \Delta_i$  after performing shadow transformations.<sup>11</sup>

The poles of (4.19) located at right-half  $\Delta$ -plane are at  $\Delta = \Delta_1 + \Delta_2 + 2n$  and  $\Delta = \Delta_3 + \Delta_4 + 2n$  with  $n = 0, 1, 2, \cdots$ . This leads to the following conformal block expansion of the shadow celestial amplitude  $\widetilde{\mathcal{A}}_s^{\Delta_i}(z_i)$ 

$$\widetilde{\mathcal{A}}_{s}^{\Delta_{i}}(z_{i}) = I_{12-34} \frac{e^{-i\pi\beta/2}}{2\sin(\frac{-\pi\beta}{2})} \left[ \sum_{n=0}^{\infty} \mathscr{C}_{\Delta_{1}+\Delta_{2}+2n}^{\Delta_{i}}(m) G_{\Delta_{1}+\Delta_{2}+2n}^{\Delta_{i}}(\chi) + \mathscr{C}_{\Delta_{3}+\Delta_{4}+2n}^{\Delta_{i}}(m) G_{\Delta_{3}+\Delta_{4}+2n}^{\Delta_{i}}(\chi) \right]$$

$$(4.20)$$

Here the expanding coefficients  $\mathscr{C}_{\Delta}^{\Delta_i}$  are

$$\mathscr{C}_{\Delta}^{\Delta_i}(m) = \mathcal{C}_{\Delta_1, \Delta_2, \Delta}(m) \mathcal{C}_{\Delta_3, \Delta_4, \Delta}(m) D^{\Delta\Delta}(m) .$$
(4.21)

As we can see from (4.20), conformal block expansion of  $\widetilde{\mathcal{A}}_s^{\Delta_i}$  contains two series of scalar exchange operators with conformal dimension  $\Delta = \Delta_1 + \Delta_2 + 2n$  and  $\Delta = \Delta_3 + \Delta_4 + 2n$ , respectively. It is then natural to expect that these exchange operators are double-trace operators, which take the schematic form as

$$\mathcal{O}_i(\partial\partial)^n \mathcal{O}_j \tag{4.22}$$

<sup>&</sup>lt;sup>11</sup>Using (A.28), one could also obtain the partial wave expansion (2.14) of the celestial amplitudes  $\mathcal{A}_s^{\Delta_i}$  by simply removing the factor  $2^{\Delta_1} a_{\Delta}^{\Delta_1,\Delta_2}$  in (4.18). It is also easy to check that the partial wave expansion of Im  $\mathcal{A}_s^{\Delta_i}$  obtained in this way agrees with the result predicted by the optical theorem [24, 25].

with conformal dimension  $\Delta_i + \Delta_j + 2n$ . After interpreting the exchange operators in (4.20) as double-trace operators, the conformal block expansion (4.20) is then reminiscent of the conformal block expansion of four-point contact Witten diagrams in AdS. Moreover, from the OPE analysis in Section 3, we already saw the double-trace operators  $\mathcal{O}_1\mathcal{O}_2$  and  $\mathcal{O}_3\mathcal{O}_4$ , coming from the leading OPE limit  $\hat{q}_1 \rightarrow \hat{q}_2$  and  $\hat{q}_3 \rightarrow \hat{q}_4$ , respectively. More precisely, we can recognize that the first term in (4.20) recovers the result (3.30) from the  $\hat{q}_1 \rightarrow \hat{q}_2$  OPE limit which is valid when  $\operatorname{Re}(\Delta_1 + \Delta_2 - \Delta_3 - \Delta_4) < 0$ .

Although we can not see the exchange operators with conformal dimension  $\Delta = \Delta_3 + \Delta_4 + 2n$  by looking at the OPE limit  $\hat{q}_1 \rightarrow \hat{q}_2$  at leading order, these exchange operators can be recovered by studying the OPE limit  $\hat{q}_3 \rightarrow \hat{q}_4$ . This makes the four-point shadow celestial amplitude  $\widetilde{\mathcal{A}}_{2\rightarrow 2}^{\Delta_i}$  a bit special since we can implement our OPE analysis for both incoming and outgoing operators and recover all the exchange operators appearing in the conformal block expansion (4.20).

### 5 Outlook

In this paper, we proposed the shadow conformal primary basis (2.16) for massless scalar particles and defined the shadow celestial amplitude in (2.20). The shadow conformal primary basis can be obtained from the conformal primary basis by performing shadow transformations as in (2.19). In terms of the shadow conformal primary basis, the shadow celestial amplitude defined in (2.20) behaves more like a standard CFT correlator. We studied the constraints from the translation symmetry on the shadow celestial amplitudes. Moreover, based on the factorization of the scattering amplitudes in the plane-wave basis, we further showed that the shadow celestial amplitudes enjoy a nice factorization in the leading OPE limit. Since the celestial coordinates in the shadow celestial amplitudes are not directly related to the momenta of the particles, the kinematic constraints like the momentum conservation do not lead to any subtlety when we study the OPE behaviour of the shadow celestial amplitude. We checked the OPE factorization by focusing on the tree-level fourpoint shadow celestial amplitude of four massless external scalars and one massive exchange scalar. Several interesting open questions ensue from our work.

• In this paper, we only studied the OPE limit at leading order in the sense that we set  $\hat{q}_1 = \hat{q}_2$  directly. It would be of great interest to study the OPE limit at sub-leading orders and explore possible exchange operators. As we see, all of the exchange operators in the four-point conformal block expansion (4.20) can be uncovered by looking at the

leading order OPE  $\hat{q}_1 \rightarrow \hat{q}_2$  and  $\hat{q}_3 \rightarrow \hat{q}_4$ . However, we expect that the leading order OPE is not enough to generate all exchange operators in higher-point amplitudes. To get the exchange operators in the higher-point conformal block expansion, one must go beyond the leading OPE.<sup>12</sup>

- The OPE analysis in Section 3.2 works only for scattering amplitude at tree-level. When the scattering amplitude M<sub>2→n</sub> at loop-level is considered, the intermediate states appearing in the generalized optical theorem also include multi-particle states. It would be interesting to see how to translate the factorization of the scattering amplitude in the plane-wave basis into the factorization of the corresponding shadow celestial amplitude when there exist multi-particle states. Moreover, our OPE analysis in Section 3.3 is restricted to a particular class of Feynman diagrams depicted in Figure 1. Extending that analysis to more generic diagrams is necessary to fully understand the OPE behaviour in CCFTs.
- Another avenue would be to explore the flat space correspondence of the double-trace operators appearing in the conformal block expansion (4.20). It is well-known that double-trace operators in AdS correspond to two-particle states. Although some recent progress has been made to relate celestial amplitudes with AdS Witten diagrams [62, 63], it is unclear if this statement still holds in CCFTs. Thus it would be of great interest to study these double-trace operators and explore their holographic dual in flat space.
- Finally, it would be interesting to study the Mellin amplitudes associated with the shadow celestial amplitudes. Celestial Mellin amplitudes in three dimensions have been studied in [33]. For the usual four-dimensional celestial amplitudes, defining the corresponding celestial Mellin amplitudes is subtle due to the existence of delta-function  $\delta(\chi \bar{\chi})$ . In contrast with the celestial amplitudes, the shadow celestial amplitude defined in this paper no longer has this distributional factor and takes the standard form as a CFT correlation function. Thus, following the definition of AdS Mellin amplitudes, one can define Mellin amplitudes associated with the shadow celestial amplitudes, and the techniques developed in AdS Mellin amplitudes can be used to study the shadow celestial Mellin amplitudes.

 $<sup>^{12}</sup>$ Recent progress on the computation of higher-point conformal blocks can be found in [40–61].

### Acknowledgments

We owe our gratitude to Jean-François Fortin, Juntao Wang, Fengjun Xu, and Xinan Zhou for discussions. CC is partly supported by National Key R&D Program of China (NO. 2020YFA0713000). The work of HS is supported in part by the Beijing Postdoctoral Research Foundation.

### **A** Preliminaries

### A.1 The generalized optical theorem

In quantum field theories, one of the most important observables is the S-matrix, which is unitary and defined as

$$\mathcal{S}_{i \to f} \equiv \langle f | S | i \rangle \tag{A.1}$$

where we used i and f to label possible in- and out-states, respectively. To extract the information from the interaction, one can define the T-matrix by

$$S = 1 + iT . (A.2)$$

This leads to the definition of the scattering amplitude  $\mathcal{M}_{i\to f}$  and  $\mathcal{T}_{i\to f}$ 

$$\langle f|iT|i\rangle \equiv i\mathcal{M}_{i\to f} \equiv (2\pi)^4 i\delta^{(4)}(p_f - p_i)\mathcal{T}_{i\to f}.$$
 (A.3)

In perturbation theory, it is the  $i\mathcal{T}_{i\to f}$  that can be computed by Feynman diagrams. The unitarity of the S-matrix implies that

$$i(T^{\dagger} - T) = T^{\dagger}T . \tag{A.4}$$

The above relation can be recast into the matrix form, giving

$$i\langle f|T^{\dagger} - T|i\rangle = \langle f|T^{\dagger}T|i\rangle , \qquad (A.5)$$

After inserting the completeness relation

$$\sum_{X} \prod_{j \in X} \int \frac{d^3 p_j}{(2\pi)^3} \frac{1}{2E_j} = 1 , \qquad (A.6)$$

we get

$$i\langle f|T^{\dagger} - T|i\rangle = \sum_{X} \left(\prod_{j \in X} \int \frac{d^3 q_j}{(2\pi)^3} \frac{1}{2E_j}\right) \langle f|T^{\dagger}|\{q_j\}\rangle\langle\{q_j\}|T|i\rangle .$$
(A.7)

Expressing the T-matrix elements as the scattering amplitude  $\mathcal{M}$  leads to

$$-i(\mathcal{M}_{i\to f} - \mathcal{M}_{f\to i}^*) = \sum_X \left(\prod_{j\in X} \int \frac{d^3 p_j}{(2\pi)^3} \frac{1}{2E_j}\right) \mathcal{M}_{i\to X} \mathcal{M}_{f\to X}^* , \qquad (A.8)$$

which is known as the generalized optical theorem in quantum field theory. In a unitary theory containing only real scalars, the CPT symmetry guarantees that  $\mathcal{M}_{i\to f} = \mathcal{M}_{f\to i}$ . Thus, the generalized optical theorem can be written as

$$\operatorname{Im} \mathcal{M}_{i \to f} = \frac{1}{2} \sum_{X} \left( \prod_{j \in X} \int \frac{d^3 p_j}{(2\pi)^3} \frac{1}{2E_j} \right) \mathcal{M}_{i \to X} \mathcal{M}_{X \to f}^* .$$
(A.9)

### A.2 Shadow transformation and conformal partial waves

In a *d*-dimensional conformal field theory, the shadow operator  $\widetilde{\mathcal{O}}_{\widetilde{\Delta}}^{\mu_1\cdots\mu_J}$  with conformal dimension  $\widetilde{\Delta} \equiv d - \Delta$  and spin-*J* is defined by performing the shadow transformation on the conformal primary operator  $\mathcal{O}_{\Delta}^{\mu_1\cdots\mu_J}$ , *i.e.*,

$$\widetilde{\mathcal{O}}_{d-\Delta}^{\mu_1\cdots\mu_J}(x) = k_{\Delta,J} \int d^d y \frac{\mathcal{I}^{\mu_1\cdots\mu_J,\nu_1\cdots\nu_J}(x-y)}{(x-y)^{2(d-\Delta)}} \mathcal{O}_{\Delta,\nu_1\cdots\nu_J}(y) , \qquad (A.10)$$

where  $\mathcal{I}^{\mu_1\cdots\mu_J,\nu_1\cdots\nu_J}(x)$  is the symmetric traceless inversion tensor that can be constructed from the products of  $I^{\mu\nu}(x) = g^{\mu\nu} - 2x^{\mu}x^{\nu}/x^2$ , and  $k_{\Delta,J}$  is

$$k_{\Delta,J} = \frac{\Gamma[d - \Delta + J]}{\pi^{\frac{d}{2}}(\Delta - 1)_J \Gamma[\Delta - \frac{d}{2}]} .$$
(A.11)

Armed with shadow operators, the following resolution of the identity in Euclidean CFTs can be obtained by the harmonic analysis of the conformal group SO(d + 1, 1)

$$|0\rangle\langle 0| + \sum_{J} \int_{\frac{d}{2}}^{\frac{d}{2}+i\infty} \frac{d\Delta}{2\pi i} \int d^{d}x \mathcal{O}_{\Delta}^{\mu_{1}\cdots\mu_{J}}(x) |0\rangle\langle 0|\widetilde{\mathcal{O}}_{d-\Delta,\mu_{1}\cdots\mu_{J}}(x) + \cdots = 1 , \qquad (A.12)$$

where  $\cdots$  represents complementary and discrete series. Inserting the resolution of the identity into a scalar four-point function in two dimensional Euclidean CFTs then leads to the conformal partial wave expansions,<sup>13</sup>

$$\langle \mathcal{O}_1(z_1)\cdots\mathcal{O}_4(z_4)\rangle = \sum_{J=-\infty}^{+\infty} \int_0^{+\infty} \frac{d\lambda}{2\pi} \frac{\rho_{h,\bar{h}}^{\Delta_i}}{n_{h,\bar{h}}} \Psi_{h,\bar{h}}^{\Delta_i}(z_i) + \cdots, \quad n_{h,\bar{h}} = \frac{-2\pi^3}{(2h-1)(2\bar{h}-1)} ,$$
(A.13)

where  $h \equiv (\Delta + J)/2 = (1 + i\lambda + J)/2$ ,  $\bar{h} \equiv (\Delta - J)/2 = (1 + i\lambda - J)/2$  and the conformal partial wave  $\Psi_{h,\bar{h}}^{\Delta_i}(z_i)$  is defined as

$$\Psi_{h,\bar{h}}^{\Delta_i}(z_i) = \int d^2 z_0 \langle \mathcal{O}_1(z_1) \mathcal{O}_2(z_2) \mathcal{O}_{h,\bar{h}}(z_0) \rangle \langle \tilde{\mathcal{O}}_{1-h,1-\bar{h}}(z_0) \mathcal{O}_3(z_3) \mathcal{O}_4(z_4) \rangle .$$
(A.14)

Evaluating the integral in (A.14) gives an explicit expression for conformal partial waves,

$$\Psi_{h,\bar{h}}^{\Delta_i}(z_i) = I_{12-34}(z_i) \left( K_{1-h,1-\bar{h}}^{\Delta_3,\Delta_4} G_{h,\bar{h}}^{\Delta_i}(\chi) + K_{h,\bar{h}}^{\Delta_1,\Delta_2} G_{1-h,1-\bar{h}}^{\Delta_i}(\chi) \right),$$
(A.15)

where  $G^{\Delta_i}_{h,\bar{h}}(\chi)$  are the conformal blocks, which are functions of conformal cross-ratios

$$\chi = \frac{z_{12}z_{34}}{z_{13}z_{24}} , \quad \bar{\chi} = \frac{\bar{z}_{12}\bar{z}_{34}}{\bar{z}_{13}\bar{z}_{24}} , \qquad (A.16)$$

and the coefficients K are given by

$$K_{h,\bar{h}}^{\Delta_1,\Delta_2} = (-1)^J \frac{\pi \Gamma[2h-1]\Gamma[1-\bar{h}+h_{12}]\Gamma[1-\bar{h}-h_{12}]}{\Gamma[2-2\bar{h}]\Gamma[h+h_{12}]\Gamma[h-h_{12}]} .$$
(A.17)

 $^{13}\mathcal{O}_i(z_i)$  should be understood as  $\mathcal{O}_i(z_i, \bar{z}_i)$ . We use this abbreviation to simplify the notation.

 $I_{12-34}(z_i)$  in (A.15) is

$$I_{12-34} \equiv \frac{\left(\frac{z_{24}}{z_{14}}\right)^{h_{12}} \left(\frac{z_{14}}{z_{13}}\right)^{h_{34}}}{z_{12}^{h_1+h_2} z_{34}^{h_3+h_4}} \frac{\left(\frac{\bar{z}_{24}}{\bar{z}_{14}}\right)^{\bar{h}_{12}} \left(\frac{\bar{z}_{14}}{\bar{z}_{13}}\right)^{\bar{h}_{34}}}{\bar{z}_{12}^{\bar{h}_1+\bar{h}_2} \bar{z}_{34}^{\bar{h}_3+\bar{h}_4}} , \quad h_{ij} \equiv h_i - h_j , \quad \bar{h}_{ij} \equiv \bar{h}_i - \bar{h}_j .$$
(A.18)

Plugging (A.15) into (A.13) and using the following property of the spectral density  $\rho_{h,\bar{h}}^{\Delta_i}$ 

$$\rho_{h,\bar{h}}^{\Delta_i} K_{1-h,1-\bar{h}}^{\tilde{\Delta}_3,\tilde{\Delta}_4} = \rho_{1-h,1-\bar{h}}^{\Delta_i} K_{1-h,1-\bar{h}}^{\tilde{\Delta}_1,\tilde{\Delta}_2} , \qquad (A.19)$$

we get

$$\langle \mathcal{O}_1(z_1)\cdots\mathcal{O}_4(z_4)\rangle = \sum_{J=-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{d\lambda}{2\pi} \frac{\rho_{h,\bar{h}}^{\Delta_i}}{n_{h,\bar{h}}} K^{\Delta_3,\Delta_4}_{1-h,1-\bar{h}} G^{\Delta_i}_{h,\bar{h}}(z_i) + \cdots$$
 (A.20)

The conformal block expansion then can be obtained by enclosing the contour to the righthalf  $\lambda$ -plane in (A.20). After defining

$$\langle \mathcal{O}_1(z_1)\cdots\mathcal{O}_4(z_4)\rangle = I_{12-34}f(\chi) , \qquad \Psi_{h,\bar{h}}^{\Delta_i}(z_i) = I_{12-34}\Psi_{h,\bar{h}}^{\Delta_i}(\chi) , \qquad (A.21)$$

the conformal partial wave expansion then can be written as

$$f(\chi) = \sum_{J=-\infty}^{+\infty} \int_0^{+\infty} \frac{d\lambda}{2\pi} \frac{\rho_{h,\bar{h}}^{\Delta_i}}{n_{h,\bar{h}}} \Psi_{h,\bar{h}}^{\Delta_i}(\chi) + \cdots , \qquad (A.22)$$

The conformal partial wave expansion for four-point functions involving a shadow operator can be easily derived from (A.13) and (A.14) when there are only scalar (J = 0) partial waves. Specifically, starting with (A.13) and performing shadow transformation for the first operator  $\mathcal{O}_1(z_1)$  leads to

$$\langle \widetilde{\mathcal{O}}_{1}(z_{1})\cdots \mathcal{O}_{4}(z_{4})\rangle = \int d^{2}z_{1'}\frac{1}{(z_{11'}^{2})^{\widetilde{\Delta}_{1}}} \int_{0}^{+\infty} \frac{d\lambda}{2\pi} \frac{\rho_{h,\bar{h}}^{\Delta_{i}}}{n_{h,\bar{h}}} \Psi_{h,\bar{h}}^{\Delta_{i}}(z_{1'}, z_{2}, z_{3}, z_{4}) + \cdots$$
 (A.23)

Using the integral representation (A.14) of the conformal partial waves, we find that

$$\int d^{2}z_{1'} \frac{1}{(z_{11'}^{2})^{\tilde{\Delta}_{1}}} \Psi_{h,\bar{h}}^{\Delta_{i}}(z_{1'}, z_{2}, z_{3}, z_{4})$$

$$= \int d^{2}z_{1'} \frac{1}{(z_{11'}^{2})^{\tilde{\Delta}_{1}}} \int d^{2}z_{0} \frac{1}{(z_{1'2}^{2})^{\frac{\Delta_{1}+\Delta_{2}-\Delta}{2}}(z_{01'}^{2})^{\frac{\Delta_{1}+\Delta_{2}-\Delta}{2}}(z_{02}^{2})^{\frac{\Delta_{1}+\Delta_{2}-\Delta_{1}}{2}}} \times \frac{1}{(z_{03}^{2})^{\frac{\tilde{\Delta}+\Delta_{3}-\Delta_{4}}{2}}(z_{04}^{2})^{\frac{\tilde{\Delta}+\Delta_{4}-\Delta_{3}}{2}}(z_{34}^{2})^{\frac{\Delta_{3}+\Delta_{4}-\tilde{\Delta}}{2}}} .$$
(A.24)

The integral over  $z_{1'}$  can be computed by noting that

$$\int d^{2} z_{1'} \frac{1}{(z_{11'}^{2})^{\tilde{\Delta}_{1}}(z_{1'2}^{2})^{\frac{\Delta_{1}+\Delta_{2}-\Delta}{2}}(z_{01'}^{2})^{\frac{\Delta+\Delta_{1}-\Delta_{2}}{2}}(z_{02}^{2})^{\frac{\Delta+\Delta_{2}-\Delta_{1}}{2}}} = a_{\Delta}^{\Delta_{1},\Delta_{2}} \frac{1}{(z_{12}^{2})^{\frac{\tilde{\Delta}_{1}+\Delta_{2}-\Delta}{2}}(z_{01}^{2})^{\frac{\Delta+\tilde{\Delta}_{1}-\Delta_{2}}{2}}(z_{02}^{2})^{\frac{\Delta+\Delta_{2}-\tilde{\Delta}_{1}}{2}}},$$
(A.25)

where  $a_{\Delta}^{\Delta_1,\Delta_2}$  is

$$a_{\Delta}^{\Delta_1,\Delta_2} = \frac{\pi\Gamma[\Delta_1 - 1]\Gamma[\frac{\Delta + \tilde{\Delta}_1 - \Delta_2}{2}]\Gamma[\frac{\tilde{\Delta}_1 + \Delta_2 - \Delta}{2}]}{\Gamma[2 - \Delta_1]\Gamma[\frac{\Delta_1 + \Delta_2 - \Delta}{2}]\Gamma[\frac{\Delta + \Delta_1 - \Delta_2}{2}]} .$$
(A.26)

This leads to the following conformal partial wave expansion for  $\langle \widetilde{\mathcal{O}}_1(z_1) \cdots \mathcal{O}_4(z_4) \rangle$ 

$$\langle \widetilde{\mathcal{O}}_1(z_1)\cdots \mathcal{O}_4(z_4)\rangle = \int_0^{+\infty} \frac{d\lambda}{2\pi} \frac{\rho_{h,\bar{h}}^{\widetilde{\Delta}_1,\Delta_2,\Delta_3,\Delta_4}}{n_{h,\bar{h}}} \Psi_{h,\bar{h}}^{\widetilde{\Delta}_1,\Delta_2,\Delta_3,\Delta_4}(z_i) + \cdots , \qquad (A.27)$$

where the new spectral density  $\rho_{h,\bar{h}}^{\tilde{\Delta}_1,\Delta_2,\Delta_3,\Delta_4}$  is related to the old spectral density  $\rho_{h,\bar{h}}^{\Delta_1,\Delta_2,\Delta_3,\Delta_4}$  through

$$\rho_{h,\bar{h}}^{\tilde{\Delta}_1,\Delta_2,\Delta_3,\Delta_4} = a_{\Delta}^{\Delta_1,\Delta_2} \rho_{h,\bar{h}}^{\Delta_1,\Delta_2,\Delta_3,\Delta_4} .$$
(A.28)

# **B** The conformal integral

In this appendix, we will compute the following conformal integral

$$I(\hat{q}_1, \hat{q}_2, \hat{p}) \equiv \int D^2 \hat{q}'_2 \frac{1}{(-\hat{q}_2 \cdot \hat{q}'_2)^{\Delta_2} (-\hat{q}'_2 \cdot \hat{p})^{2-\Delta_1 - \Delta_2} (-\hat{q}'_2 \cdot Y)^{\Delta_1}}, \qquad (B.1)$$

where  $D^2 \hat{q}' = d^2 z'$ ,  $Y^{\mu} = -2(\hat{q}_1 \cdot \hat{p})\hat{p}^{\mu} - \hat{q}_1^{\mu}$ . Using the Feynman/Schwinger parameterization,  $I(\hat{q}_1, \hat{q}_2, \hat{p})$  can be written as

$$I(\hat{q}_1, \hat{q}_2, \hat{p}) = \frac{1}{\Gamma[\Delta_1]\Gamma[\Delta_2]\Gamma[2 - \Delta_1 - \Delta_2]} \int_0^\infty dx_1 dx_2 \int D^2 \hat{q}_2' \frac{x_1^{\Delta_1 - 1} x_2^{\Delta_2 - 1}}{[-\hat{q}_2' \cdot (\hat{p} + x_1 Y + x_2 \hat{q}_2)]^2} .$$
(B.2)

We evaluate the integral over  $\hat{q}'_2$  by noting that [64]

$$\int D^2 \hat{q}'_2 \frac{1}{(-\hat{q}'_2 \cdot Q)^2} = \frac{2\pi}{-Q^2} , \qquad (B.3)$$

which holds for any timelike vector  $Q^{\mu}$ . This can be seen by going to the rest frame of  $Q^{\mu}$ , *i.e.* we choose  $Q^{\mu} = (1, 0, 0, 0)$ . In this frame, the above integral becomes

$$\int D^2 \hat{q}'_2 \frac{1}{(-\hat{q}'_2 \cdot Q)^2} = \int d^2 z \frac{1}{(1+z\bar{z})^2} = 2\pi .$$
 (B.4)

Recovering the dependence on  $Q^2$  by dimensional analysis then gives (B.3). Plugging (B.3) back into (B.2) leads to

$$I(\hat{q}_{1}, \hat{q}_{2}, \hat{p}) = \frac{2\pi}{\Gamma[\Delta_{1}]\Gamma[\Delta_{2}]\Gamma[2 - \Delta_{1} - \Delta_{2}]} \times \int_{0}^{\infty} \frac{dx_{1}dx_{2} x_{1}^{\Delta_{1} - 1} x_{2}^{\Delta_{2} - 1}}{(1 - 2x_{1}\hat{q}_{1} \cdot \hat{p} - 2x_{2}\hat{q}_{2} \cdot \hat{p} + 4x_{1}x_{2}\hat{q}_{1} \cdot \hat{p}\hat{q}_{2} \cdot \hat{p} - 4x_{1}x_{2}\hat{q}_{12})}$$
(B.5)

Evaluating the integral over  $x_1$ ,  $x_2$  and expanding the results around small  $\hat{q}_{12}$  finally lead to

$$I(\hat{q}_1, \hat{q}_2, \hat{p}) = \sum_{n=0}^{\infty} \frac{2\pi\Gamma[1 - \Delta_1]\Gamma[1 - \Delta_2](\Delta_1)_n (\Delta_2)_n}{2^{\Delta_1 + \Delta_2}\Gamma[2 - \Delta_1 - \Delta_2]\Gamma[n+1]\Gamma[n+1]} \frac{(-\hat{q}_{12})^n}{(\hat{q}_1 \cdot \hat{p})^{\Delta_1 + n}(-\hat{q}_2 \cdot \hat{p})^{\Delta_2 + n}} \cdot$$
(B.6)

Setting  $\hat{q}_1 = \hat{q}_2$  in (B.6) leads to

$$I(\hat{q}_2, \hat{q}_2, \hat{p}) = \frac{2\pi\Gamma[1 - \Delta_1]\Gamma[1 - \Delta_2]}{2^{\Delta_1 + \Delta_2}\Gamma[2 - \Delta_1 - \Delta_2]} .$$
(B.7)

## References

 S. Pasterski, S.-H. Shao, and A. Strominger, "Flat Space Amplitudes and Conformal Symmetry of the Celestial Sphere," *Phys. Rev. D* 96 (2017), no. 6 065026, 1701.00049.

- [2] A. Strominger, "Lectures on the Infrared Structure of Gravity and Gauge Theory," 1703.05448.
- [3] A.-M. Raclariu, "Lectures on Celestial Holography," 2107.02075.
- [4] S. Pasterski, "Lectures on celestial amplitudes," *Eur. Phys. J. C* 81 (2021), no. 12 1062, 2108.04801.
- [5] S. Pasterski, M. Pate, and A.-M. Raclariu, "Celestial Holography," in 2022 Snowmass Summer Study, 11, 2021. 2111.11392.
- [6] S. Weinberg, "Infrared photons and gravitons," *Phys. Rev.* **140** (1965) B516–B524.
- [7] T. He, V. Lysov, P. Mitra, and A. Strominger, "BMS supertranslations and Weinberg's soft graviton theorem," *JHEP* 05 (2015) 151, 1401.7026.
- [8] D. Kapec, P. Mitra, A.-M. Raclariu, and A. Strominger, "2D Stress Tensor for 4D Gravity," *Phys. Rev. Lett.* **119** (2017), no. 12 121601, 1609.00282.
- [9] L. Donnay, A. Puhm, and A. Strominger, "Conformally Soft Photons and Gravitons," JHEP 01 (2019) 184, 1810.05219.
- [10] W. Fan, A. Fotopoulos, and T. R. Taylor, "Soft Limits of Yang-Mills Amplitudes and Conformal Correlators," *JHEP* 05 (2019) 121, 1903.01676.
- [11] M. Pate, A.-M. Raclariu, and A. Strominger, "Conformally Soft Theorem in Gauge Theory," *Phys. Rev. D* 100 (2019), no. 8 085017, 1904.10831.
- [12] T. Adamo, L. Mason, and A. Sharma, "Celestial amplitudes and conformal soft theorems," *Class. Quant. Grav.* **36** (2019), no. 20 205018, 1905.09224.
- [13] A. Puhm, "Conformally Soft Theorem in Gravity," JHEP 09 (2020) 130, 1905.09799.
- [14] A. Guevara, "Notes on Conformal Soft Theorems and Recursion Relations in Gravity," 1906.07810.
- [15] S. Stieberger and T. R. Taylor, "Symmetries of Celestial Amplitudes," *Phys. Lett. B* 793 (2019) 141–143, 1812.01080.
- [16] A. Fotopoulos and T. R. Taylor, "Primary Fields in Celestial CFT," JHEP 10 (2019) 167, 1906.10149.

- [17] A. Strominger, " $w_{1+\infty}$  and the Celestial Sphere," 2105.14346.
- [18] A. Ball, S. A. Narayanan, J. Salzer, and A. Strominger, "Perturbatively exact  $w_{1+\infty}$  asymptotic symmetry of quantum self-dual gravity," *JHEP* **01** (2022) 114, 2111.10392.
- [19] A. Strominger, " $w_{1+\infty}$  Algebra and the Celestial Sphere: Infinite Towers of Soft Graviton, Photon, and Gluon Symmetries," *Phys. Rev. Lett.* **127** (2021), no. 22 221601.
- [20] S. Pasterski and S.-H. Shao, "Conformal basis for flat space amplitudes," *Phys. Rev.* D 96 (2017), no. 6 065022, 1705.01027.
- [21] Y. T. A. Law and M. Zlotnikov, "Massive Spinning Bosons on the Celestial Sphere," *JHEP* 06 (2020) 079, 2004.04309.
- [22] S. A. Narayanan, "Massive Celestial Fermions," *JHEP* **12** (2020) 074, 2009.03883.
- [23] L. Iacobacci and W. Mück, "Conformal Primary Basis for Dirac Spinors," *Phys. Rev.* D 102 (2020), no. 10 106025, 2009.02938.
- [24] H. T. Lam and S.-H. Shao, "Conformal Basis, Optical Theorem, and the Bulk Point Singularity," *Phys. Rev. D* 98 (2018), no. 2 025020, 1711.06138.
- [25] C.-M. Chang, Y.-t. Huang, Z.-X. Huang, and W. Li, "Bulk locality from the celestial amplitude," *SciPost Phys.* 12 (2022), no. 5 176, 2106.11948.
- [26] D. Nandan, A. Schreiber, A. Volovich, and M. Zlotnikov, "Celestial Amplitudes: Conformal Partial Waves and Soft Limits," *JHEP* 10 (2019) 018, 1904.10940.
- [27] Y. T. A. Law and M. Zlotnikov, "Relativistic partial waves for celestial amplitudes," *JHEP* 11 (2020) 149, 2008.02331.
- [28] A. Atanasov, W. Melton, A.-M. Raclariu, and A. Strominger, "Conformal block expansion in celestial CFT," *Phys. Rev. D* 104 (2021), no. 12 126033, 2104.13432.
- [29] Y. Hu, L. Lippstreu, M. Spradlin, A. Y. Srikant, and A. Volovich, "Four-point correlators of light-ray operators in CCFT," *JHEP* 07 (2022) 104, 2203.04255.
- [30] S. Banerjee, R. Basu, and S. A. Bhatkar, "Light transformed gluon correlators in CCFT," 2203.06657.

- [31] S. De, Y. Hu, A. Yelleshpur Srikant, and A. Volovich, "Correlators of Four Light-Ray Operators in CCFT," 2206.08875.
- [32] D. García-Sepúlveda, A. Guevara, J. Kulp, and J. Wu, "Notes on Resonances and Unitarity from Celestial Amplitudes," 2205.14633.
- [33] H. Jiang, "Celestial Mellin Amplitude," 2208.01576.
- [34] S. Ferrara, A. F. Grillo, G. Parisi, and R. Gatto, "The shadow operator formalism for conformal algebra. Vacuum expectation values and operator products," *Lett. Nuovo Cim.* 4S2 (1972) 115–120.
- [35] H. Osborn, "Conformal Blocks for Arbitrary Spins in Two Dimensions," *Phys. Lett. B* 718 (2012) 169–172, 1205.1941.
- [36] W. Fan, A. Fotopoulos, S. Stieberger, T. R. Taylor, and B. Zhu, "Conformal blocks from celestial gluon amplitudes," *JHEP* 05 (2021) 170, 2103.04420.
- [37] W. Fan, A. Fotopoulos, S. Stieberger, T. R. Taylor, and B. Zhu, "Conformal blocks from celestial gluon amplitudes. Part II. Single-valued correlators," *JHEP* 11 (2021) 179, 2108.10337.
- [38] W. Fan, A. Fotopoulos, S. Stieberger, T. R. Taylor, and B. Zhu, "Celestial Yang-Mills amplitudes and D = 4 conformal blocks," *JHEP* 09 (2022) 182, 2206.08979.
- [39] Y. T. A. Law and M. Zlotnikov, "Poincaré constraints on celestial amplitudes," JHEP 03 (2020) 085, 1910.04356.
- [40] K. B. Alkalaev and V. A. Belavin, "From global to heavy-light: 5-point conformal blocks," JHEP 03 (2016) 184, 1512.07627.
- [41] V. Rosenhaus, "Multipoint Conformal Blocks in the Comb Channel," JHEP 02 (2019) 142, 1810.03244.
- [42] S. Parikh, "Holographic dual of the five-point conformal block," JHEP 05 (2019) 051, 1901.01267.
- [43] V. Gonçalves, R. Pereira, and X. Zhou, "20' Five-Point Function from  $AdS_5 \times S^5$ Supergravity," *JHEP* **10** (2019) 247, 1906.05305.

- [44] C. B. Jepsen and S. Parikh, "Propagator identities, holographic conformal blocks, and higher-point AdS diagrams," *JHEP* 10 (2019) 268, 1906.08405.
- [45] S. Parikh, "A multipoint conformal block chain in d dimensions," JHEP 05 (2020) 120, 1911.09190.
- [46] J.-F. Fortin, W. Ma, and W. Skiba, "Higher-Point Conformal Blocks in the Comb Channel," JHEP 07 (2020) 213, 1911.11046.
- [47] J.-F. Fortin, W.-J. Ma, and W. Skiba, "Six-point conformal blocks in the snowflake channel," JHEP 11 (2020) 147, 2004.02824.
- [48] T. Anous and F. M. Haehl, "On the Virasoro six-point identity block and chaos," *JHEP* 08 (2020), no. 08 002, 2005.06440.
- [49] F. M. Haehl, A. Streicher, and Y. Zhao, "Six-point functions and collisions in the black hole interior," *JHEP* 08 (2021) 134, 2105.12755.
- [50] J.-F. Fortin, W.-J. Ma, and W. Skiba, "Seven-point conformal blocks in the extended snowflake channel and beyond," *Phys. Rev. D* 102 (2020), no. 12 125007, 2006.13964.
- [51] S. Hoback and S. Parikh, "Towards Feynman rules for conformal blocks," *JHEP* 01 (2021) 005, 2006.14736.
- [52] J.-F. Fortin, W.-J. Ma, and W. Skiba, "All global one-and two-dimensional higher-point conformal blocks," (2020) arXiv:2009.07674.
- [53] I. Buric, S. Lacroix, J. A. Mann, L. Quintavalle, and V. Schomerus, "From Gaudin Integrable Models to d-dimensional Multipoint Conformal Blocks," *Phys. Rev. Lett.* 126 (2021), no. 2 021602, 2009.11882.
- [54] S. Hoback and S. Parikh, "Dimensional reduction of higher-point conformal blocks," *JHEP* 03 (2021) 187, 2009.12904.
- [55] D. Poland and V. Prilepina, "Recursion relations for 5-point conformal blocks," JHEP 10 (2021) 160, 2103.12092.
- [56] I. Buric, S. Lacroix, J. A. Mann, L. Quintavalle, and V. Schomerus, "Gaudin models and multipoint conformal blocks: general theory," *JHEP* 10 (2021) 139, 2105.00021.

- [57] I. Buric, S. Lacroix, J. A. Mann, L. Quintavalle, and V. Schomerus, "Gaudin models and multipoint conformal blocks. Part II. Comb channel vertices in 3D and 4D," *JHEP* 11 (2021) 182, 2108.00023.
- [58] I. Buric, S. Lacroix, J. A. Mann, L. Quintavalle, and V. Schomerus, "Gaudin models and multipoint conformal blocks III: comb channel coordinates and OPE factorisation," *JHEP* 06 (2022) 144, 2112.10827.
- [59] J.-F. Fortin, S. Hoback, W.-J. Ma, S. Parikh, and W. Skiba, "Feynman Rules for Scalar Conformal Blocks," (2022) 2204.08909.
- [60] C. Bercini, V. Gonçalves, and P. Vieira, "Light-Cone Bootstrap of Higher Point Functions and Wilson Loop Duality," *Phys. Rev. Lett.* **126** (2021), no. 12 121603, 2008.10407.
- [61] A. Antunes, M. S. Costa, V. Goncalves, and J. V. Boas, "Lightcone bootstrap at higher points," *JHEP* 03 (2022) 139, 2111.05453.
- [62] E. Casali, W. Melton, and A. Strominger, "Celestial Amplitudes as AdS-Witten Diagrams," 2204.10249.
- [63] L. Iacobacci, C. Sleight, and M. Taronna, "From Celestial Correlators to AdS, and back," 2208.01629.
- [64] D. Simmons-Duffin, "Projectors, Shadows, and Conformal Blocks," JHEP 04 (2014) 146, 1204.3894.