

Uncertainty Quantification in Synthetic Controls with Staggered Treatment Adoption*

Matias D. Cattaneo[†] Yingjie Feng[‡] Filippo Palomba[§] Rocio Titiunik[¶]

September 7, 2023

Abstract

We propose principled prediction intervals to quantify the uncertainty of a large class of synthetic control predictions (or estimators) in settings with staggered treatment adoption, offering precise non-asymptotic coverage probability guarantees. From a methodological perspective, we provide a detailed discussion of different causal quantities to be predicted, which we call *causal predictands*, allowing for multiple treated units with treatment adoption at possibly different points in time. From a theoretical perspective, our uncertainty quantification methods improve on prior literature by (i) covering a large class of causal predictands in staggered adoption settings, (ii) allowing for synthetic control methods with possibly nonlinear constraints, (iii) proposing scalable robust conic optimization methods and principled data-driven tuning parameter selection, and (iv) offering valid uniform inference across post-treatment periods. We illustrate our methodology with an empirical application studying the effects of economic liberalization in the 1990s on GDP for emerging European countries. Companion general-purpose software packages are provided in `Python`, `R` and `Stata`.

Keywords: causal inference, synthetic controls, staggered treatment adoption, prediction intervals, non-asymptotic inference.

*We thank Alberto Abadie, Simon Freyaldenhoven, and Bartolomeo Stellato for many insightful discussions. Cattaneo and Titiunik gratefully acknowledge financial support from the National Science Foundation (SES-2019432 and SES-2241575), Cattaneo gratefully acknowledges financial support from the National Institute of Health (R01 GM072611-16), and Feng gratefully acknowledges the financial support from the National Natural Science Foundation of China (NSFC) under grants 72203122 and 72133002.

[†]Department of Operations Research and Financial Engineering, Princeton University.

[‡]School of Economics and Management, Tsinghua University.

[§]Department of Economics, Princeton University.

[¶]Department of Politics, Princeton University.

Contents

1	Introduction	1
1.1	Related Literature	3
2	The Effect of Liberalization on GDP for Emerging European Countries	4
3	Causal Inference Framework and Quantities of Interest	6
3.1	Synthetic Control Method	9
4	Prediction Intervals	17
4.1	Discussion of Conditions (i)-(iv)	24
4.2	Simultaneous Prediction Intervals	26
5	Optimization and Tuning Parameter Selection	27
5.1	Conic Programming Approach	28
5.2	Defining Constraints in Simulation	31
5.3	Adjustment for Nonlinear Constraints	32
5.4	Summary of Algorithmic Implementation	33
6	Empirical Application	34
6.1	Empirical Strategy	35
6.2	Results	37
7	Conclusion	42

1 Introduction

The synthetic control method was introduced by [Abadie and Gardeazabal \(2003\)](#), and since then many extensions and generalizations have been proposed in the literature (see [Abadie, 2021](#), and references therein). This methodology is now part of the standard toolkit for program evaluation and treatment effect analysis, offering a complement to traditional difference-in-differences, event studies, and other panel data approaches for causal inference employing longitudinal aggregated data with few treated units. Most of the synthetic control literature concentrates on identification, as well as on prediction or point estimation of treatment effects, under different causal inference frameworks and algorithmic implementations. In contrast, despite its importance for empirical work, principled uncertainty quantification of synthetic control predictions or estimators remains mostly unexplored beyond some specific methods for the canonical single-treatment-unit case.

We develop prediction intervals to quantify the uncertainty of a large class of synthetic control predictions (or estimators) in settings with staggered treatment adoption, offering precise non-asymptotic coverage probability guarantees, scalable robust optimization implementations, and principled tuning parameter selection. Employing a causal inference framework where potential outcomes are assumed to be random, we propose inferential procedures with non-asymptotic probability guarantees; such guarantees are valuable because synthetic control applications often have small sample sizes, limiting the applicability of asymptotic approximations. Conceptually, our proposed prediction intervals capture two sources of uncertainty: one coming from the construction of the synthetic control weights with pre-treatment data, and the other generated by the irreducible sampling variability introduced by the post-treatment outcomes. The proposed prediction intervals also take into account potential misspecification error explicitly, and enjoy other robustness properties due to their non-asymptotic construction.

Our first contribution is methodological in nature due to the complexity added by the staggered treatment adoption setup, which allows for (but does not require) the existence of multiple treatment units changing from control to treatment status at possibly different points in time. In [Section 3](#), we introduce a general causal inference framework that is specifically tailored to synthetic control methods and incorporates staggered treatment adoption. Using this framework, we define different causal quantities to be predicted in the context of synthetic controls, which we refer to as *causal*

predictands, and explain how our inferential methods can be used in each case. Furthermore, our proposed causal framework explicitly allows for misspecification error, multiple covariate features, and cross-equation re-weighting when constructing the synthetic control weights with pre-treatment data.

Building on our general causal inference framework, we present two main theoretical contributions in Section 4. First, we give high-level sufficient conditions leading to valid prediction intervals with precise non-asymptotic guarantees, allowing for stationary and non-stationary data and for synthetic control predictions constructed using multiple re-weighted features and nonlinear constraints. We also provide easy-to-verify primitive conditions, which cover all common synthetic control methods in the literature (e.g., Ridge and elastic net regression). Second, we extend our methods to provide not only pointwise but also joint inference validity across post-treatment time periods. This result allows for the construction of joint “prediction bands” to complement the prediction intervals for different predictands at each point in the post-treatment period.

To complement our methodological and theoretical work, Section 5 discusses scalable robust optimization implementations and principled tuning parameter selection based on our theoretical results. First, we show how to recast our proposed general synthetic control methods for prediction and uncertainty quantification as conic optimization programs (Boyd and Vandenberghe, 2004). This approach gives massive speed improvements for implementation. Second, our proposed methods explicitly employ the non-asymptotic characterizations of the coverage errors associated with the proposed prediction intervals to calibrate the underlying tuning parameters for practice. In particular, we discuss in detail relaxation methods for possibly nonlinear constraints based on our theoretical development. Section 5.4 gives a summary of our algorithmic implementation.

We illustrate our methods with an empirical application investigating the effect of economic liberalization in the 1990s on GDP for emerging European countries. This empirical work is motivated by Billmeier and Nannicini (2013) who, also employing synthetic control methods, studied the same substantive question for countries in other regions of the world. Our main findings suggest that economic liberalization in the 1990s did not have a positive economic impact for emerging European countries. This finding is in line with prior empirical results. The Supplemental Appendix provides additional empirical evidence supporting our main findings, including a re-analysis using alternative synthetic control predictions for different countries and a discussion of specific cases

where the synthetic control method does not appear to be well-suited for the analysis.

We provide general-purpose software implementing all our results in `Python`, `R`, and `Stata`, including detailed documentation and additional replication materials. This software is presented in detail in our companion article [Cattaneo, Feng, Palomba and Titiunik \(2023\)](#), where we discuss several implementation issues related to numerical optimization and tuning parameter selection. To complement the illustration, in Section S.4 of the Supplemental Appendix, we provide more details on how to prepare the data to analyze staggered treatment adoption using synthetic control methods using our companion software.

1.1 Related Literature

Our paper contributes to two strands of the synthetic control literature. First, it contributes to the development of prediction/estimation and inference methods for staggered treatment adoption settings. Putting aside generic linear factor model or matrix completion methods, [Ben-Michael, Feller and Rothstein \(2022\)](#) and [Shaikh and Toulis \(2021\)](#) appear to be the only prior papers that have studied staggered treatment adoption for synthetic controls. The first paper focuses on prediction/estimation in settings where the pre-treatment fit is poor, and develops penalization methods for improving the performance of the canonical synthetic control method. [Ben-Michael, Feller and Rothstein \(2022\)](#) also suggest employing a bootstrap method for assessing uncertainty, but no formalization is provided guaranteeing its (asymptotic) validity. [Shaikh and Toulis \(2021\)](#) focus on uncertainty quantification employing a parametric duration model and propose a permutation-based inferential method under a symmetry assumption. Our paper complements these prior works by offering a nonparametric inference method with demonstrable non-asymptotic coverage guarantees and allowing for misspecification in the construction of the synthetic control weights, which can be applied directly to a large class of synthetic control (possibly penalized) predictions and causal quantities of interest.

Second, from a general methodological perspective, our proposed inference methods are motivated by [Vovk \(2012\)](#), and are most closely related to prior work by [Chernozhukov, Wüthrich and Zhu \(2021a\)](#); [Chernozhukov, Wüthrich and Zhu \(2021b\)](#) on conformal prediction intervals and by [Cattaneo, Feng and Titiunik \(2021\)](#) on non-asymptotic prediction intervals (see [Wainwright \(2019\)](#) for a modern introduction to non-asymptotic statistical learning). We contribute to this second

strand of the literature by developing new prediction interval methods. First, we allow for a large class of causal predictands in staggered adoption settings (prior work covered only the canonical single treated unit case). Second, we cover a large class of synthetic control predictions with possibly nonlinear constraints (prior work allowed for predictions with linear constraints). Third, we develop scalable robust optimization implementations and propose principled data-driven tuning parameter selection (prior work did not provide guidance on these issues). Fourth, we develop valid uniform inference across post-treatment periods (absent in prior work).

There are a few other recent proposals in the literature to quantify uncertainty and conduct inference for synthetic controls. For example, [Li \(2020\)](#) study correctly specified linear factor models, [Masini and Medeiros \(2021\)](#) study high-dimensional penalization methods, [Agarwal, Shah, Shen and Song \(2021\)](#) investigate matrix completion methods, [Shen, Ding, Sekhon and Yu \(2022\)](#) explore panel data methods, and [Shi, Miao, Hu and Tchetgen \(2023\)](#) develop inference methods using a proximal causal inference framework. All these methods rely on asymptotic approximations, in most cases employing standard Gaussian critical values that assume away misspecification errors and other small sample issues. Our work complements these contributions by providing prediction intervals with non-asymptotic coverage guarantees. Finally, all the inferential methods mentioned so far contrast with the original method proposed by [Abadie, Diamond and Hainmueller \(2010\)](#), which relies on design-based permutation of treatment assignment assuming that the potential outcomes are non-random.

2 The Effect of Liberalization on GDP for Emerging European Countries

During the second half of the twentieth century, a considerable number of countries all over the world launched programs of (external) economic liberalization, booming from 22% in 1960 to 73% in the early 2000s ([Wacziarg and Welch, 2008](#)). In the last thirty years, political scientists and economists have investigated the social and economic consequences of such liberalization programs, often reaching conflicting conclusions (see, e.g., [Levine and Renelt, 1992](#); [Sachs, Warner, Åslund and Fischer, 1995](#); [DeJong and Ripoll, 2006](#)).

The impact of liberalization policies on economic welfare has been traditionally investigated

with cross-country analyses (e.g. [Sachs, Warner, Åslund and Fischer, 1995](#)) and individual case studies (e.g. [Bhagwati and Srinivasan, 2001](#)). More recently, scholars have turned to synthetic control methods in the hope of employing a causal inference methodology that allows for the presence of time-varying unobservable confounders. Employing the synthetic control framework originally developed in [Abadie and Gardeazabal \(2003\)](#), [Billmeier and Nannicini \(2013\)](#) analyzed the effects of liberalization in four continents: Africa, Asia, North America, and South America. They used a pre-existing dataset of economic variables (previously used in [Giavazzi and Tabellini, 2005](#)) which includes 180 countries, covers the period 1963–2000, and contains an indicator for economic liberalization originally defined in [Sachs, Warner, Åslund and Fischer \(1995\)](#) and updated in [Wacziarg and Welch \(2008\)](#) (hereafter, the Sachs-Warner indicator). More details on the data and the definition of economic liberalization can be found in the Supplemental Appendix Section S.4.

The main finding of [Billmeier and Nannicini \(2013\)](#) is that economic liberalization—as measured by the Sachs-Warner indicator—has a non-negative effect on GDP per capita. Moreover, the authors document the presence of substantial heterogeneity in the effects depending on the period of time in which the liberalization took place. On the one hand, the predicted effect on real income per capita is positive in those countries that embarked on programs of economic liberalization before the 1980s. On the other hand, in the MENA (Middle East and North Africa) region and in sub-Saharan Africa, where many liberalization episodes occurred after 1985, the magnitude of the predicted effect becomes either statistically insignificant or economically irrelevant.

Our work builds on [Billmeier and Nannicini \(2013\)](#) in three ways. First, we form prediction intervals around causal predictions using our new methodology. Second, rather than constructing a synthetic control independently for each treated unit, we leverage both the presence of multiple treated units and the staggered nature of treatment adoption to jointly form the synthetic controls. Third, by borrowing donors from other continents, we also analyze the impact of economic liberalization events in Europe (see [Table 1](#) below).

As shown in [Table 1](#), we study seven liberalization episodes that occurred in Europe out of the nine available ones. In particular, North Macedonia is not included in the final analysis due to the lack of other variables in the dataset besides GDP per capita before 1994, whereas Slovenia is

Table 1: *Economic liberalization episodes in the Billmeier and Nannicini (2013) dataset.*

Country Name	Event Date	Years Closed	Years Liberalized	Analyzed in this work
Albania	1992	29	9	✓
Bulgaria	1991	28	10	✓
Czech Republic	1991	28	10	✓
Hungary	1990	27	11	✓
North Macedonia	1994	31	7	✗
Poland	1990	27	11	✓
Romania	1992	29	9	✓
Slovak Republic	1991	28	10	✓
Slovenia	1991	28	10	✗

Notes: a comprehensive description of the events that led the Sachs-Warner indicator to switch on for these countries is contained in the Supplemental Appendix Section S.4 together with the full set of donors.

excluded in virtue of the poor pre-treatment fit yielded by the synthetic controls for such country.¹ We discover that the real income per capita trajectory during the 10 years following the liberalization is lower than it would have been in the absence of the liberalization in all countries we analyzed. Similarly, the average post-treatment effect is negative for all the treated units and so is the average post-treatment effect on the treated (see Section 3 for a precise definitions). However, when individual and simultaneous prediction intervals are taken into account, the trajectory of the synthetic control becomes indistinguishable from the realized time series for real income with high probability. Only the average treatment effect on the treated differs from zero with high probability. Results in other continents—fully presented in the Supplemental Appendix Section S.6—confirm in magnitude the ones found in Billmeier and Nannicini (2013), whenever it has been possible to compare them. However, the data do not allow us to draw the conclusion that liberalization events changed the trajectory of GDP per capita in either a favorable or negative way.

3 Causal Inference Framework and Quantities of Interest

We consider the synthetic control (SC) framework with a fixed number of units that may adopt treatment at different times. Specifically, a researcher observes N units for T time periods. Units are indexed by $i = 1, \dots, N$, and time periods are indexed by $t = 1, \dots, T$. Let T_i represent the

¹As detailed in Supplemental Appendix Section S.4 and Supplemental Appendix Section S.8, we still use Slovenia as a donor whenever possible. Supplemental Appendix Section S.8 also presents the results for North Macedonia and Slovenia.

time when unit i receives the treatment, with $T_i = \infty$ denoting that unit i is never treated (at least during the observed T time periods). Each unit i remains untreated in $t = 1, \dots, T_i - 1$ and remains treated since T_i . We assume that there is a (non-empty) set of units that are never treated, i.e., $N_0 := \sum_{i=1}^N \mathbb{1}(T_i = \infty) > 0$, and let $N_1 := N - N_0$ be the number of units that are eventually treated by time period T . Without loss of generality, assume units are ordered in the adoption time: $T_1 \leq T_2 \leq \dots \leq T_{N_1}$. In our empirical application, the treatment of interest is economic liberalization, the adoption time of which is heterogeneous across different countries.

The staggered adoption problem can be analyzed in a multi-valued treatment effects framework. Let $Y_{it}(s)$ denote the potential outcome of unit i in period t that would be observed if unit i had adopted the treatment in period s , for $s = 1, \dots, T, \infty$, and we set $Y_{it}(s) = Y_{it}(\infty)$ for $t < s$. Implicitly, these simplifications impose two standard assumptions: no spillovers (the potential outcomes of unit i depend only on i 's adoption time) and no anticipation (a unit's potential outcomes prior to the treatment are equal to the outcomes it would have had if it had never been treated). Then, the observed outcome can be written as

$$Y_{it} = Y_{it}(\infty)\mathbb{1}(t < T_i) + Y_{it}(T_i)\mathbb{1}(t \geq T_i).$$

A large set of causal predictands can be defined in this context. In particular, for $k \geq 0$, let τ_{ik} be the (individual) treatment effect of unit i in $T_i + k$ (k periods after treatment adoption):

$$\tau_{ik} := Y_{i(T_i+k)}(T_i) - Y_{i(T_i+k)}(\infty).$$

This is the treatment effect of interest in the classical synthetic control analysis with only one treated unit. When multiple treated units or multiple post-treatment periods are available, a researcher might be interested in a variety of other causal predictands. The following are some typical examples:

- (i) Average post-treatment effect on unit i :

$$\tau_i := \frac{1}{T - T_i + 1} \sum_{k=0}^{T-T_i} \tau_{ik}.$$

(ii) Average treatment effect on units treated at time s_0 , k periods after treatment adoption:

$$\tau_{\cdot k, s_0} := \frac{1}{|\{i : T_i = s_0\}|} \sum_{i: T_i = s_0} \tau_{ik} ,$$

where $|\{i : T_i = s_0\}|$ denotes the number of units that get treated at s_0 .

(iii) Average treatment effect on the treated, k periods after treatment adoption:

$$\tau_{\cdot k} := \frac{1}{N_1} \sum_{i=1}^{N_1} \tau_{ik} .$$

Since the observation ends at time T , the number of treated units included in the definition of the average treatment effect $\tau_{\cdot k}$ could vary across k . To avoid this complication, we assume that all treated units are observed at least K periods after the treatment for some $K \geq 1$, i.e., $T_{N_1} \leq T - K$, and attention is restricted to $\tau_{\cdot k}$ for $k \leq K$ only.

The potential outcomes, treatment adoption times, and individual treatment effects are viewed as *random* quantities in general. We assume that there is only a fixed (possibly small) number of treated units and time periods, which is often the case in synthetic control applications. Thus, the various average treatment effects defined above are also random quantities in general, but we continue to refer to them as “treatment effects” to be consistent with analogous quantities defined in the literature (e.g., assuming a fixed, non-random potential outcomes framework). In classical large-sample-based causal analysis, target parameters are often probability or ergodic (non-random) limits of the average effects above as $N_1 \rightarrow \infty$, $T - T_i \rightarrow \infty$, and/or $K \rightarrow \infty$; our results are also valid in such settings. Nevertheless, in this paper we develop statistical inference methods based on prediction intervals that describe a region where a new realization of a (random) causal predictand of interest is likely to be observed, rather than usual confidence intervals giving a region in the parameter space for a non-random parameter of interest.

The canonical synthetic control analysis with one single treated unit can be viewed as a special case of the more general setup described above. Specifically, suppose that unit 1 is the only treated unit who receives the treatment at T_1 , and all other units are *never treated*, i.e., $T_i = \infty$ for all

$i \geq 2$. Then, the observed outcome is

$$Y_{it} = \begin{cases} Y_{it}(\infty) & i = 2, \dots, N \\ Y_{1t}(\infty) & i = 1 \text{ and } t = 1, \dots, T_1 - 1 \\ Y_{1t}(T_1) & i = 1 \text{ and } t = T_1, \dots, T \end{cases}$$

The target causal predictand in this canonical case is usually the individual treatment effect on the treated, i.e., τ_{1k} defined previously.

3.1 Synthetic Control Method

Consider the case where multiple treated units are available, and one would like to find a vector of SC weights possibly different for each treated unit. From now on, we use a superscript $i = 1, \dots, N_1$ in brackets to index the treated units that enter the construction of the desired causal predictand, and a subscript $l = 1, \dots, M$ to denote different features of the treated on which one would like to match.

Let $\mathbf{A}_l^{[i]} = (a_{1,l}^{[i]}, \dots, a_{T_0,l}^{[i]})' \in \mathbb{R}^{T_0}$ be the l th feature of the treated unit i measured in T_0 (user-specified) pre-treatment periods. For each feature l and each treated unit i , there exist $J + K$ variables that are used to predict or match the T_0 -dimensional vector $\mathbf{A}_l^{[i]}$. These $J + K$ variables are separated into two groups denoted by $\mathbf{B}_l^{[i]} = (\mathbf{B}_{1,l}^{[i]}, \mathbf{B}_{2,l}^{[i]}, \dots, \mathbf{B}_{J,l}^{[i]}) \in \mathbb{R}^{T_0 \times J}$ and $\mathbf{C}_l^{[i]} = (\mathbf{C}_{1,l}^{[i]}, \dots, \mathbf{C}_{K,l}^{[i]}) \in \mathbb{R}^{T_0 \times K}$, respectively. More precisely, for each $j = 1, \dots, J$, $\mathbf{B}_{j,l}^{[i]} = (b_{j1,l}^{[i]}, \dots, b_{jT_0,l}^{[i]})'$ corresponds to the l th feature of the j th unit in the donor pool measured in T_0 pre-treatment periods, and for each $k = 1, \dots, K$, $\mathbf{C}_{k,l}^{[i]} = (c_{k1,l}^{[i]}, \dots, c_{kT_0,l}^{[i]})'$ is another vector of control variables used to predict $\mathbf{A}_l^{[i]}$ over the same pre-intervention time span. Stacking the M equations (corresponding to M features) for each treated unit, we define

$$\underbrace{\mathbf{A}^{[i]}}_{T_0 \cdot M \times 1} = \begin{bmatrix} \mathbf{A}_1^{[i]} \\ \vdots \\ \mathbf{A}_M^{[i]} \end{bmatrix}, \quad \underbrace{\mathbf{B}^{[i]}}_{T_0 \cdot M \times J} = \begin{bmatrix} \mathbf{B}_1^{[i]} \\ \vdots \\ \mathbf{B}_M^{[i]} \end{bmatrix}, \quad \underbrace{\mathbf{C}^{[i]}}_{T_0 \cdot M \times K \cdot M} = \begin{bmatrix} \mathbf{C}_1^{[i]} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{C}_2^{[i]} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{C}_M^{[i]} \end{bmatrix}.$$

In our empirical application, $\mathbf{A}^{[i]}$ contains the (log) GDP per capita and the percentage of complete secondary schooling in population ($M = 2$) of a treated economy i (that has ever experienced

liberalization) during the pre-liberalization period, and $\mathbf{B}^{[i]}$ contains the same two features of the donor economies used to match $\mathbf{A}^{[i]}$. For each feature $l = 1, 2$, $\mathbf{C}_l^{[i]}$ contains an intercept and a linear time trend ($K = 2$).

The goal of the synthetic control method is to search for a vector of weights $\mathbf{w} = (\mathbf{w}^{[1]'}, \dots, \mathbf{w}^{[N_1]'})' \in \mathcal{W} \subseteq \mathbb{R}^{JN_1}$ which is common across the M features and a vector of coefficients $\mathbf{r} = (\mathbf{r}^{[1]'}, \dots, \mathbf{r}^{[N_1]'})' \in \mathcal{R} \subseteq \mathbb{R}^{KMN_1}$, such that the linear combination of $\mathbf{B}^{[i]}$ and $\mathbf{C}^{[i]}$ matches $\mathbf{A}^{[i]}$ as closely as possible, for all $1 \leq i \leq N_1$. The feasibility sets \mathcal{W} and \mathcal{R} capture the restrictions imposed. Typical examples include simplex-type, lasso-type and ridge-type constraints (for further details, see [Cattaneo, Feng, Palomba and Titiunik, 2023](#)).

Such SC weights are typically obtained via the following optimization problem: for some $(T_0 \cdot M \cdot N_1) \times (T_0 \cdot M \cdot N_1)$ symmetric weighting matrix \mathbf{V} ,

$$\hat{\boldsymbol{\beta}} := (\hat{\mathbf{w}}', \hat{\mathbf{r}}')' \in \arg \min_{\mathbf{w} \in \mathcal{W}, \mathbf{r} \in \mathcal{R}} (\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r})' \mathbf{V} (\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r}) \quad (3.1)$$

where

$$\underbrace{\mathbf{A}}_{T_0 \cdot M \cdot N_1 \times 1} = \begin{bmatrix} \mathbf{A}^{[1]} \\ \vdots \\ \mathbf{A}^{[N_1]} \end{bmatrix}, \quad \underbrace{\mathbf{B}}_{T_0 \cdot M \cdot N_1 \times J \cdot N_1} = \begin{bmatrix} \mathbf{B}^{[1]} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{B}^{[2]} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{B}^{[N_1]} \end{bmatrix}, \quad \underbrace{\mathbf{C}}_{T_0 \cdot M \cdot N_1 \times K \cdot M \cdot N_1} = \begin{bmatrix} \mathbf{C}^{[1]} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{C}^{[2]} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{C}^{[N_1]} \end{bmatrix}.$$

Accordingly, we write $\hat{\mathbf{w}} = (\hat{\mathbf{w}}^{[1]'}, \dots, \hat{\mathbf{w}}^{[N_1]'})'$ where each $\hat{\mathbf{w}}^{[i]} = (\hat{w}_1^{[i]}, \dots, \hat{w}_J^{[i]})'$ is the SC weights on J control units that are used to predict the counterfactual of the treated unit i . Similarly, write $\hat{\mathbf{r}} = (\hat{\mathbf{r}}^{[1]'}, \dots, \hat{\mathbf{r}}^{[N_1]'})'$ and $\hat{\boldsymbol{\beta}} = (\hat{\boldsymbol{\beta}}^{[1]'}, \dots, \hat{\boldsymbol{\beta}}^{[N_1]'})'$.

Then, the predicted counterfactual outcome of each treated unit is given by

$$\hat{Y}_{it}(\infty) := \mathbf{x}_t^{[i]'} \hat{\mathbf{w}}^{[i]} + \mathbf{g}_t^{[i]'} \hat{\mathbf{r}}^{[i]} = \mathbf{p}_t^{[i]'} \hat{\boldsymbol{\beta}}^{[i]}, \quad \mathbf{p}_t^{[i]} = (\mathbf{x}_t^{[i]'}, \mathbf{g}_t^{[i]'})', \quad i = 1, \dots, N_1, \quad t > T_i,$$

where $\mathbf{x}_t^{[i]}$ is a vector of predictors of the control units measured in time t used to predict the counterfactual of the treated unit i , and $\mathbf{g}_t^{[i]}$ is a vector of predictors that correspond to the additional control variables specified in $\mathbf{C}^{[i]}$. In our empirical application, $\mathbf{x}_t^{[i]}$ is the post-treatment (log) GDP per capita of donor economies, and $\mathbf{g}_t^{[i]}$ includes a constant, a linear time trend and two zeros so that $\mathbf{g}_t^{[i]}$ is conformable with $\hat{\mathbf{r}}^{[i]}$. In general, however, variables included in $\mathbf{x}_t^{[i]}$ and

$\mathbf{g}_t^{[i]}$ need not be the same as those in $\mathbf{B}^{[i]}$ and $\mathbf{C}^{[i]}$. For convenience of later exposition, we write $\mathbf{p}_t = (\mathbf{p}_t^{[1]'}, \dots, \mathbf{p}_t^{[N_1]'})'$.

Any causal predictand τ discussed before can be written as the difference between an observed outcome (possibly a linear combination of outcomes of a few treated units or in different periods) and the corresponding counterfactual outcome. To construct a prediction $\hat{\tau}$ of τ , one only needs to substitute a ‘‘SC prediction’’ for the unobserved counterfactual:

$$\text{SC prediction} := \mathbf{p}'_{\tau} \hat{\boldsymbol{\beta}},$$

where the predictor vector \mathbf{p}_{τ} needs to be defined in context. More details are provided in the following examples.

Example 3.1 (Individual Treatment Effect, τ_{ik}). *Suppose that the individual treatment effect τ_{ik} of the treated unit i with $T_i < \infty$ is of interest for some $0 \leq k \leq T - T_i$. Let the set of pre-treatment periods be $\{t : t \leq T_i - 1\}$. The donor pool consists of units that receive treatment later than $T_i + k$, i.e., $\{j : T_j > T_i + k\}$. The SC weights are constructed using the data of the treated unit i and the control units in the pre-treatment period. Given the prediction of the counterfactual outcome $\hat{Y}_{i(T_i+k)}$, the predicted treatment effect is*

$$\hat{\tau}_{ik} := Y_{i(T_i+k)} - \hat{Y}_{i(T_i+k)}(\infty) = Y_{i(T_i+k)} - \mathbf{p}'_{\tau_{ik}} \hat{\boldsymbol{\beta}}.$$


The predictor vector $\mathbf{p}_{\tau_{ik}}$ in this case is given by

$$\mathbf{p}_{\tau_{ik}} := \underbrace{(\mathbf{0}'_{J+KM}, \dots, \mathbf{0}'_{J+KM})}_{(i-1) \text{ vectors}}, \mathbf{p}_{T_i+k}^{[i]'}, \underbrace{(\mathbf{0}'_{J+KM}, \dots, \mathbf{0}'_{J+KM})}_{(N_1-i) \text{ vectors}}'.$$

where $\mathbf{0}_{J+KM}$ is a $(J + KM)$ -vector of zeros. See Figure 1 for a graphical representation of $\hat{\tau}_{ik}$.

Example 3.2 (Average Post-Treatment Effect, τ_i). *Suppose that the quantity of interest is the treatment effect on the treated unit i averaged across all the post-treatment periods, i.e., τ_i , defined previously. Let the set of pre-treatment periods be $\{t : t \leq T_i - 1\}$, and take the set of all units that are never treated as the donor pool, i.e., $\{j : T_j = \infty\}$. The SC weights in this scenario can be constructed in the same way as in the case of individual treatment effects. The prediction of the*

Figure 1: Graphical illustration of the individual treatment effect, $\hat{\tau}_{ik}$.



Notes: The black line displays the times series of the treated unit's outcome, whereas the blurred gray lines portray the same variable for the donor units. The blue line is the synthetic control constructed out of the donor units using a simplex-type constraint. The pink vertical line represents the quantity $\hat{\tau}_{ik}$ described in Example 3.1, where k is set to 5.

average post-treatment effect is given by

$$\hat{\tau}_i := \frac{1}{T - T_i + 1} \sum_{t=T_i}^T \left(Y_{it} - \hat{Y}_{it}(\infty) \right) = \frac{1}{T - T_i + 1} \sum_{t=T_i}^T Y_{it} - \mathbf{p}'_{\tau_i} \hat{\boldsymbol{\beta}},$$


where the predictor vector in this case is given by

$$\mathbf{p}_{\tau_i} := \left(\underbrace{\mathbf{0}'_{J+KM}, \dots, \mathbf{0}'_{J+KM}}_{(i-1) \text{ vectors}}, \frac{1}{T - T_i + 1} \sum_{t \geq T_i} \mathbf{p}_t^{[i]'}, \underbrace{\mathbf{0}'_{J+KM}, \dots, \mathbf{0}'_{J+KM}}_{(N_1-i) \text{ vectors}} \right)'.$$

See Figure 2 for a graphical representation of $\hat{\tau}_i$.

Example 3.3 (Average Treatment Effect on the Treated at s_0 after k Periods, τ_{k,s_0}). Suppose that the causal predictand of interest is the average treatment effect for units that adopt the treatment at time s_0 , k periods after treatment adoption, i.e., τ_{k,s_0} defined above. Let the set of pre-treatment periods be $\{t : t \leq s_0 - 1\}$, and take the set of units that adopt the treatment later than $s_0 + k$, i.e.,

Figure 2: Graphical illustration of the average post-treatment effect, $\hat{\tau}_i$.



Notes: The black line displays the times series of the treated unit's outcome, whereas the blurred gray lines portray the same variable for the donor units. The blue line is the synthetic control constructed out of the donor units using a simplex-type constraint. The dashed lines represent the post-treatment average of the treated time series (black) and the synthetic control time series (blue). The pink vertical line represents the quantity $\hat{\tau}_i$ described in Example 3.2.

$\{j : T_j > s_0 + k\}$, as the donor pool. We have two different strategies to conduct the SC analysis:

- Implement the procedure described above, which allows for different SC weights on different treated units. Suppose $\{i : T_i = s_0\} = \{i_1, \dots, i_{n_{s_0}}\}$. Then, the predicted effect is given by

$$\hat{\tau}_{\tau,k,s_0} := \frac{1}{n_{s_0}} \sum_{i:T_i=s_0} \left(Y_{i(s_0+k)} - \hat{Y}_{i(s_0+k)}(\infty) \right) = \frac{1}{n_{s_0}} \sum_{i:T_i=s_0} Y_{i(s_0+k)} - \mathbf{p}'_{\tau,k,s_0} \hat{\boldsymbol{\beta}},$$

where the predictor vector in this case is given by

$$\mathbf{p}_{\tau,k,s_0} := \left(\underbrace{\mathbf{0}'_{J+KM}, \dots, \mathbf{0}'_{J+KM}}_{(i_1-1) \text{ vectors}}, \frac{1}{n_{s_0}} \mathbf{p}_{s_0+k}^{[i_1]'}, \dots, \frac{1}{n_{s_0}} \mathbf{p}_{s_0+k}^{[i_{n_{s_0}}]'}, \underbrace{\mathbf{0}'_{J+KM}, \dots, \mathbf{0}'_{J+KM}}_{(N_1-i_{n_{s_0}}) \text{ vectors}} \right)'.$$


- Aggregate different treated units into one single unit, denoted by “ave”, whose potential out-

comes are given by the average of all units treated at time s_0 :

$$Y_t^{ave}(s; s_0) := \frac{1}{n_{s_0}} \sum_{i: T_i = s_0} Y_{it}(s), \quad t = 1, \dots, T, \quad s = 1, \dots, T, \infty.$$

More precisely, $Y_t^{ave}(s; s_0)$ is the potential outcome of the aggregate unit *ave* in period t that would be observed if it had adopted the treatment in period s , while *ave* actually adopted the treatment in period s_0 . Other features of this aggregate unit can be constructed similarly as the average of multiple units treated at s_0 . The SC weights can be obtained using the data of the aggregate unit *ave* and control units in the donor pool from pre-treatment period. Then, the SC analysis proceeds exactly the same way as in Example 3.1. See Figure 3 for a graphical representation of $\hat{\tau}_{k, s_0}$.

Figure 3: Graphical illustration of the average treatment effect on the treated at s_0 after k periods, $\hat{\tau}_{k, s_0}$.



Notes: The black lines display the times series of the treated units' outcome, whereas the blurred gray lines portray the same variable for the donor units. The blue lines are the synthetic controls constructed out of the donor units using a simplex-type constraint. The largest black triangle represents the average of the treated units' outcomes at $s_0 + k$, with s_0 set to 2015 and k set to 5. Similarly, the largest blue circle is the average of the synthetic controls at $s_0 + k$. The pink vertical line represents the quantity $\hat{\tau}_{k, s_0}$ described in Example 3.3. n_{s_0} denotes the number of treated units that received the treatment at time s_0 .

Example 3.4 (Average Treatment Effect on all Treated after k Periods, τ_k). Suppose that the quantity of interest is the average treatment effect for all treated units k periods after treatment adoption, i.e., τ_k defined above. Let the set of pre-treatment periods be $\{t : t \leq T_1 - 1\}$ and the donor pool be $\{j : T_j = \infty\}$. Then, the predicted effect is given by


$$\hat{\tau}_k := \frac{1}{N_1} \sum_{i=1}^{N_1} \left(Y_{i(T_i+k)} - \hat{Y}_{i(T_i+k)} \right) = \frac{1}{N_1} \sum_{i=1}^{N_1} Y_{it} - \mathbf{p}'_{\tau_k} \hat{\boldsymbol{\beta}},$$

where the predictor vector in this case is given by

$$\mathbf{p}_{\tau_k} := \left(\frac{1}{N_1} \mathbf{p}_{T_1+k}^{[1]'}, \dots, \frac{1}{N_1} \mathbf{p}_{T_{N_1}+k}^{[N_1]'} \right)'.$$

See Figure 4 for a graphical representation of $\hat{\tau}_k$.

Figure 4: Graphical illustration of the average treatment effect on all treated after k periods $\hat{\tau}_k$.



Notes: The black lines display the times series of the treated units' outcome, whereas the blurred gray lines portray the same variable for the donor units. The blue lines are the synthetic controls constructed out of the donor units using a simplex-type constraint. The largest black triangle represents the average of the treated units' outcomes k periods after treatment, with k set to 5. Similarly, the largest blue circle is the average of the synthetic controls k periods after treatment. The pink vertical line represents the quantity $\hat{\tau}_{k,s_0}$ described in Example 3.4.

As pointed out by Ben-Michael, Feller and Rothstein (2022), with multiple treated units, the SC

weights could be constructed in two ways: (i) optimizing the separate fit for each treated unit; (ii) optimizing the pooled fit for the average of the treated units. These ideas can be accommodated by choosing a proper weighting matrix \mathbf{V} . For example, taking $\mathbf{V} = \mathbf{I}_{T_0 M N_1}$ yields

$$\hat{\boldsymbol{\beta}} = \arg \min_{\mathbf{w} \in \mathcal{W}, \mathbf{r} \in \mathcal{R}} \sum_{i=1}^{N_1} \sum_{l=1}^M \sum_{t=1}^{T_0} \left(a_{t,l}^{[i]} - \mathbf{b}_{t,l}^{[i]'} \mathbf{w}_l^{[i]} - \mathbf{c}_{t,l}^{[i]'} \mathbf{r}_l^{[i]} \right)^2,$$

where $\mathbf{B}_l^{[i]} := (\mathbf{b}_{1,l}^{[i]}, \dots, \mathbf{b}_{T_0,l}^{[i]})'$ is the l th feature of the J control units in the donor pool, and $\mathbf{C}_l^{[i]} := (\mathbf{c}_{1,l}^{[i]}, \dots, \mathbf{c}_{T_0,l}^{[i]})'$ is the additional K variables used to predict $\mathbf{A}_l^{[i]}$. The objective above is equivalent to minimizing the sum of squared errors of the pre-treatment fit for *each* treated unit and thus is termed “separate fit”.

By contrast, consider the following weighting matrix:

$$\mathbf{V} = \frac{1}{N_1^2} \mathbf{1}_{N_1} \mathbf{1}_{N_1}' \otimes \mathbf{I}_{T_0 M}$$

where \otimes denotes the Kronecker product operator. Then,

$$\hat{\boldsymbol{\beta}} = \arg \min_{\mathbf{w} \in \mathcal{W}, \mathbf{r} \in \mathcal{R}} \sum_{l=1}^M \sum_{t=1}^{T_0} \left[\frac{1}{N_1} \sum_{i=1}^{N_1} \left(a_{t,l}^{[i]} - \mathbf{b}_{t,l}^{[i]'} \mathbf{w}_l^{[i]} - \mathbf{c}_{t,l}^{[i]'} \mathbf{r}_l^{[i]} \right) \right]^2.$$

In this case, the goal is to minimize the sum of squared *averaged* errors across all treated units, which is usually termed “pooled fit”.

To simplify the exposition, we assume that the number of pre-treatment periods T_0 and the number of control units J do not vary when we match on features of different treated units. This accommodates the common practice of using data in all periods before the first unit is treated (i.e., $T_0 = T_1 - 1$) and taking all units that are never treated as donors (i.e., $J = N_0$). However, our setup is general enough to allow for more flexible choices of pre-treatment periods and donor pools that are heterogeneous across different treated units, at the cost of additional cumbersome notation. Selecting the appropriate donor pool for each treatment unit in practice can be done taking into account the causal predictand of interest.

4 Prediction Intervals

Our goal is to use our generic synthetic control framework to construct prediction intervals for the various treatment effects defined in Section 3. Suppose that \mathbf{A} , \mathbf{B} , and \mathbf{C} introduced previously are random quantities defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, and $\mathcal{H} \subseteq \mathcal{F}$ is a sub- σ -field. For some $\alpha, \pi \in (0, 1)$, we say a random interval \mathcal{I} is an (α, π) -valid \mathcal{H} -conditional prediction interval for a causal predictand τ if

$$\mathbb{P}\left\{\mathbb{P}[\tau \in \mathcal{I} \mid \mathcal{H}] \geq 1 - \alpha\right\} \geq 1 - \pi, \quad (4.1)$$

where τ can be thought of as any of the treatment effects defined in Section 3.

If \mathcal{H} is the trivial σ -field over Ω , then \mathcal{I} reduces to an unconditional prediction interval for τ . In the general case, the prediction interval \mathcal{I} is \mathcal{H} -conditionally (α, π) -valid: the conditional coverage probability of \mathcal{I} for τ is at least $(1 - \alpha)$, which holds with probability over \mathcal{H} at least $(1 - \pi)$. In practice, $(1 - \alpha)$ is a desired coverage level chosen by users, say 95%, and π is a “small” number that depends on the sample size and typically goes to zero in some asymptotic sense. In this paper, all results are valid for all T_0 large enough, with the associated probability loss π characterized precisely. Thus, we say that the conditional coverage of the prediction interval \mathcal{I} is at least $(1 - \alpha)$ with high probability, or that the conditional prediction interval offers finite-sample probability guarantees. Our results imply that $\pi \rightarrow 0$ as $T_0 \rightarrow \infty$, but no limits or asymptotic arguments are used in this paper.

Generally, the choice of the conditioning set \mathcal{H} determines the uncertainty that would not be taken into account by the prediction intervals. We consider prediction intervals conditional on all control units as well as on the other predictors used to construct the SC prediction. That is, $\mathcal{H} = \{\mathbf{B}, \mathbf{C}, \mathbf{p}_\tau\}$. Therefore, the uncertainty to be characterized arises from the treated units only. Define the target quantity of the SC weights (conditional on \mathcal{H}) that is analogous to (3.1):

$$\boldsymbol{\beta}_0 := (\mathbf{w}'_0, \mathbf{r}'_0)' = \arg \min_{\mathbf{w} \in \mathcal{W}, \mathbf{r} \in \mathcal{R}} \mathbb{E}\left[(\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r})' \mathbf{V}(\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r}) \mid \mathcal{H}\right]. \quad (4.2)$$

Thus, we can write

$$\mathbf{A} = \mathbf{B}\mathbf{w}_0 + \mathbf{C}\mathbf{r}_0 + \mathbf{U}, \quad \mathbf{w}_0 \in \mathcal{W}, \quad \mathbf{r}_0 \in \mathcal{R}, \quad (4.3)$$

where $\mathbf{U} = (\mathbf{u}^{[1]'}, \dots, \mathbf{u}^{[N_1]'})' \in \mathbb{R}^{T_0 M N_1}$ is the corresponding pseudo-true residual relative to the σ -field \mathcal{H} . Each $\mathbf{u}^{[i]} = (u_{1,1}^{[i]}, \dots, u_{T_0,1}^{[i]}, \dots, u_{1,M}^{[i]}, \dots, u_{T_0,M}^{[i]})' \in \mathbb{R}^{T_0 M}$ is the pseudo-true residual from the M equations (corresponding to the M features) of the treated unit i . Given the pseudo-true value β_0 and a desired causal predictand τ , we have the following decomposition of the predicted effect $\hat{\tau}$:

$$\hat{\tau} - \tau \equiv -\mathbf{p}'_{\tau}(\hat{\beta} - \beta_0) + e_{\tau},$$

where $\mathbf{p}'_{\tau}(\hat{\beta} - \beta_0)$ captures the *in-sample uncertainty* from the SC weights construction using pre-treatment information, and e_{τ} captures the *out-of-sample uncertainty* from the stochastic error in one or a few post-treatment periods. Notice that in-sample uncertainty quantification is necessary in this scenario since the conditioning set $\mathcal{H} \supseteq \{\mathbf{B}, \mathbf{C}\}$, but $\mathcal{H} \not\supseteq \mathbf{A}$.

To construct prediction intervals for a causal predictand τ , we propose to find constants $M_{1,L}$, $M_{1,U}$, $M_{2,L}$ and $M_{2,U}$, possibly depending on $\alpha_1, \alpha_2, \pi_1, \pi_2 \in (0, 1)$ such that

$$\begin{aligned} \mathbb{P}\left\{\mathbb{P}\left[M_{1,L} \leq \mathbf{p}'_{\tau}(\hat{\beta} - \beta_0) \leq M_{1,U} \mid \mathcal{H}\right] \geq 1 - \alpha_1\right\} &\geq 1 - \pi_1, & \text{and} \\ \mathbb{P}\left\{\mathbb{P}\left[M_{2,L} \leq e_{\tau} \leq M_{2,U} \mid \mathcal{H}\right] \geq 1 - \alpha_2\right\} &\geq 1 - \pi_2, \end{aligned}$$

which suffices to guarantee

$$\mathbb{P}\left\{\mathbb{P}\left[\hat{\tau} + M_{1,L} - M_{2,U} \leq \tau \leq \hat{\tau} + M_{1,U} - M_{2,L} \mid \mathcal{H}\right] \geq 1 - \alpha_1 - \alpha_2\right\} \geq 1 - \pi_1 - \pi_2,$$

that is, the prediction interval $\mathcal{I} = [\hat{\tau} + M_{1,L} - M_{2,U}, \hat{\tau} + M_{1,U} - M_{2,L}]$ achieves $(1 - \alpha_1 - \alpha_2)$ \mathcal{H} -conditional coverage probability, which holds with probability at least $1 - \pi_1 - \pi_2$ over \mathcal{H} , as defined in (4.1).

In-sample uncertainty. We propose a simulation-based strategy to bound the in-sample error $\mathbf{p}'_{\tau}(\beta_0 - \hat{\beta})$. Let $\mathbf{Z} = (\mathbf{B}, \mathbf{C})$ and $d = (J + KM)N_1$. Using (3.1) and (4.2), we obtain the following optimization problem characterizing the centered synthetic control weights predictor:

$$\hat{\beta} - \beta_0 = \arg \min_{\beta - \beta_0 \in \Delta} \left\{ (\beta - \beta_0)' \hat{\mathbf{Q}} (\beta - \beta_0) - 2\hat{\gamma}' (\beta - \beta_0) \right\},$$

where $\hat{\mathbf{Q}} = \mathbf{Z}'\mathbf{V}\mathbf{Z}$, $\hat{\gamma}' = \mathbf{U}'\mathbf{V}\mathbf{Z}$, and $\Delta = \{\beta - \beta_0 \in \mathbb{R}^d : \beta \in \mathcal{W} \times \mathcal{R}\}$. We assume that the

constraint sets \mathcal{W} and \mathcal{R} are convex throughout.

Let $\gamma := \mathbb{E}[\hat{\gamma}|\mathcal{H}]$, which is not necessarily equal to $\mathbf{0}$. By optimality of $\hat{\beta}$ and the convexity of \mathcal{W} and \mathcal{R} , it can be shown that $\hat{\beta}$ has to satisfy $\hat{\beta} - \beta_0 \in \Delta$ and $(\hat{\beta} - \beta_0)' \hat{\mathbf{Q}}(\hat{\beta} - \beta_0) - 2(\hat{\gamma} - \gamma)'(\hat{\beta} - \beta_0) \leq 0$. Thus, the minimum and the maximum of $\mathbf{p}'_{\tau}(\beta - \beta_0)$ over the set of β satisfying these restrictions are lower and upper bounds on the in-sample error $\mathbf{p}'_{\tau}(\hat{\beta} - \beta_0)$. Conditional on \mathcal{H} , the uncertainty of these (stochastic) bounds comes from $\hat{\gamma}$ only. We can employ a normal distributional approximation of $\hat{\gamma}$ and set $M_{1,L} = \mathbf{c}_L(\alpha_1/2)$ and $M_{1,U} = \mathbf{c}_U(1 - \alpha_1/2)$ where

$$\begin{aligned} \mathbf{c}_L(\alpha_1/2) &:= (\alpha_1/2)\text{-quantile of } \inf\{\mathbf{p}'_{\tau}\delta : \delta \in \mathcal{M}_{\mathbf{G}}\} \quad \text{and} \\ \mathbf{c}_U(1 - \alpha_1/2) &:= (1 - \alpha_1/2)\text{-quantile of } \sup\{\mathbf{p}'_{\tau}\delta : \delta \in \mathcal{M}_{\mathbf{G}}\} \end{aligned}$$

conditional on \mathcal{H} , with $\mathcal{M}_{\mathbf{G}} = \{\delta \in \Delta : \delta' \hat{\mathbf{Q}}\delta - 2\mathbf{G}'\delta \leq 0\}$, $\mathbf{G}|\mathcal{H} \sim \mathbf{N}(\mathbf{0}, \Sigma)$ and $\Sigma = \mathbb{V}[\hat{\gamma}|\mathcal{H}]$.

However, $\mathbf{c}_L(\alpha_1/2)$ and $\mathbf{c}_U(1 - \alpha_1/2)$ cannot be directly used because they still rely on the infeasible normalized constraint set Δ and the unknown covariance matrix Σ . We thus propose a feasible simulation-based strategy allowing for *possibly nonlinear* constraints, where we replace unknown quantities by plug-in approximations thereof.

First, we need a feasible constraint set Δ^* used in simulation. Specifically, define the distance between a point $\mathbf{a} \in \mathbb{R}^d$ and a set $\Lambda \subseteq \mathbb{R}^d$ by $\text{dist}(\mathbf{a}, \Lambda) = \inf_{\lambda \in \Lambda} \|\mathbf{a} - \lambda\|$, where $\|\cdot\|$ is a generic ℓ_p vector norm on \mathbb{R}^d with $p \geq 1$ (e.g., Euclidean norm or ℓ_1 norm). Intuitively, we require that every point in the original infeasible constraint set Δ be *close* to the feasible constraint set Δ^* in simulation. Consequently, searching for an upper (or lower) bound within the infeasible set Δ can be replaced with doing so within the feasible set Δ^* . This requirement will be formalized as condition (iii) in Theorem 1 below. A proposed strategy for constructing Δ^* is described in (4.6), and more implementation details are discussed in Section 5.2.

Second, we need an estimator $\hat{\Sigma}$ of the covariance matrix Σ . A variety of well-established heteroskedasticity/serial-correlation-robust estimators can be used. We require $\hat{\Sigma}$ to be a “good” approximation of Σ in the sense of condition (iv) in Theorem 1 below. This allows us to approximate the infeasible normal distribution $\mathbf{N}(\mathbf{0}, \Sigma)$ by $\mathbf{N}(\mathbf{0}, \hat{\Sigma})$, which can be simulated using the data.

Once Δ^* and $\hat{\Sigma}$ are available, we can simply draw random vectors from $\mathbf{N}(\mathbf{0}, \hat{\Sigma})$ conditional on

the data, and then set

$$M_{1,L} = \mathbf{c}_L^*(\alpha_1/2) \quad \text{and} \quad M_{1,U} = \mathbf{c}_U^*(1 - \alpha_1/2) \quad (4.4)$$

where

$$\begin{aligned} \mathbf{c}_L^*(\alpha_1/2) &:= (\alpha_1/2)\text{-quantile of } \inf\{\mathbf{p}'_\tau \boldsymbol{\delta} : \boldsymbol{\delta} \in \mathcal{M}_{\mathbf{G}}^*\} \quad \text{and} \\ \mathbf{c}_U^*(1 - \alpha_1/2) &:= (1 - \alpha_1/2)\text{-quantile of } \sup\{\mathbf{p}'_\tau \boldsymbol{\delta} : \boldsymbol{\delta} \in \mathcal{M}_{\mathbf{G}}^*\} \end{aligned}$$

conditional on the data, $\mathcal{M}_{\mathbf{G}}^* = \{\boldsymbol{\delta} \in \Delta^* : \boldsymbol{\delta}' \widehat{\mathbf{Q}} \boldsymbol{\delta} - 2(\mathbf{G}^*)' \boldsymbol{\delta} \leq 0\}$, and $\mathbf{G}^* | \text{Data} \sim \mathbf{N}(\mathbf{0}, \widehat{\boldsymbol{\Sigma}})$.

For example, our main empirical results in Section 6 employ an L1-L2 constrained synthetic control prediction (non-linear constraints) and carefully chosen plug-in approximations $(\Delta^*, \widehat{\boldsymbol{\Sigma}})$ (described precisely below), which exhibits good performance in finite samples. Validity of the resulting non-asymptotic prediction intervals is new to the literature because prior results covered only linear constraints (see Theorem 1) and, in particular, did not provide a principled Δ^* (see Lemma 1).

Out-of-sample uncertainty. To bound the out-of-sample error e_τ , we propose an easy-to-implement approach based on non-asymptotic concentration inequalities. Specifically, assume that $e_\tau - \mathbb{E}[e_\tau | \mathcal{H}]$ is conditional-on- \mathcal{H} sub-Gaussian with parameter $\sigma_{\mathcal{H}}$. Then for any $\varepsilon > 0$,

$$\mathbb{P}\left(|e_\tau - \mathbb{E}[e_\tau | \mathcal{H}]| \geq \varepsilon \mid \mathcal{H}\right) \leq 2 \exp\left(-\frac{\varepsilon^2}{2\sigma_{\mathcal{H}}^2}\right).$$

Consequently, we set

$$M_{2,L} = \mathbb{E}[e_\tau | \mathcal{H}] - \sqrt{2\sigma_{\mathcal{H}}^2 \log(2/\alpha_2)} \quad \text{and} \quad M_{2,U} = \mathbb{E}[e_\tau | \mathcal{H}] + \sqrt{2\sigma_{\mathcal{H}}^2 \log(2/\alpha_2)}, \quad (4.5)$$

which yields a prediction interval $[M_{2,L}, M_{2,U}]$ that covers $e_{T_i+k}^{[i]}$ with at least $(1 - \alpha_2)$ conditional coverage probability. We emphasize that the sub-Gaussianity assumption is one of many possibilities. The above strategy could be applied using other concentration inequalities requiring weaker moment conditions, though the resulting prediction intervals may be wider.

As discussed in the examples below, the generic out-of-sample error e_τ associated with some treatment effect τ is typically a linear combination of individual error terms $e_t^{[i]} = Y_{it}(\infty) - \mathbf{p}_t^{[i]'} \boldsymbol{\beta}_0$

in the post-treatment period, and the sub-Gaussianity of e_τ is implied by assuming $e_t^{[i]}$ is sub-Gaussian. In practice, one could first construct pre-treatment residuals $\widehat{e}_t^{[i]} = Y_{it}(\infty) - \mathbf{p}_t^{[i]'} \widehat{\boldsymbol{\beta}}^{[i]}$, $t = 1, \dots, T_0$, and estimate the conditional moments of $e_t^{[i]}$ employing various parametric or non-parametric regression of $\widehat{e}_t^{[i]}$. Such estimates can then be translated into the necessary estimates of $\mathbb{E}[e_\tau | \mathcal{H}]$ and $\sigma_{\mathcal{H}}^2$ for constructing $M_{2,L}$ and $M_{2,U}$. The unknown conditional moments could also be set using external information, or tabulated across different values to assess the sensitivity of the resulting prediction intervals. See [Cattaneo, Feng, Palomba and Titiunik \(2023\)](#) for more implementation details.

Example 4.1 (Individual Treatment Effect, τ_{ik} , continued). *For the causal predictand τ_{ik} , the out-of-sample error is given by*

$$e_{\tau_{ik}} = Y_{i(T_i+k)}(\infty) - \mathbf{p}_{T_i+k}^{[i]'} \boldsymbol{\beta}_0 = e_{T_i+k}^{[i]},$$

If we assume $e_t^{[i]} - \mathbb{E}[e_t^{[i]} | \mathcal{H}]$ is sub-Gaussian conditional on \mathcal{H} , then the strategy outlined above can be applied.

Example 4.2 (Average Post-Treatment Effect, τ_i , continued). *For the causal predictand τ_i , the out-of-sample error is given by*

$$e_{\tau_i} := \frac{1}{T - T_i + 1} \sum_{t=T_i}^T \left(Y_{it}(\infty) - \mathbf{p}_t^{[i]'} \boldsymbol{\beta}_0^{[i]} \right) = \frac{1}{T - T_i + 1} \sum_{t=T_i}^T e_t^{[i]}.$$

The approach outlined above can still be applied to e_{τ_i} . For example, if $e_t^{[i]} - \mathbb{E}[e_t^{[i]} | \mathcal{H}]$ is conditional-on- \mathcal{H} sub-Gaussian with parameter $\sigma_{\mathcal{H},t} > 0$ for $t = T_i, \dots, T$, it can be shown that e_{τ_i} , as the average of $e_t^{[i]}$ across time, satisfies that for any $\varepsilon > 0$,

$$\mathbb{P}(|e_{\tau_i} - \mathbb{E}[e_{\tau_i} | \mathcal{H}]| \geq \varepsilon | \mathcal{H}) \leq 2 \exp\left(-\varepsilon^2 / (2\bar{\sigma}_{\mathcal{H}}^2)\right), \quad \bar{\sigma}_{\mathcal{H}} = \frac{1}{T - T_i + 1} \sum_{t=T_i}^T \sigma_{\mathcal{H},t}.$$

This inequality holds regardless of the dependence structure of $e_t^{[i]}$. If $e_t^{[i]}$ is independent over t , the

above result can be improved:

$$\mathbb{P}(|e_{\tau_i} - \mathbb{E}[e_{\tau_i} | \mathcal{H}]| \geq \varepsilon | \mathcal{H}) \leq 2 \exp\left(-\varepsilon^2 / (2\tilde{\sigma}_{\mathcal{H}}^2)\right), \quad \tilde{\sigma}_{\mathcal{H}} = \frac{1}{T - T_i + 1} \left(\sum_{t=T_i}^T \sigma_{\mathcal{H},t}^2\right)^{1/2}.$$

In this case, one can characterize each $\sigma_{\mathcal{H},t}$, $t = T_i, \dots, T$, and use the idea outlined above to construct bounds on e_{τ_i} .

Alternatively, one could construct a pre-treatment sequence of errors that is analogous to e_{τ_i} :

$$\tilde{e}_t^{[i]} := \frac{1}{T - T_i + 1} \sum_{t=\ell}^{\ell+T-T_i} e_t^{[i]}, \quad 1 \leq \ell \leq 2T_i - T - 1,$$

and then apply the strategy outlined before to this new sequence of errors, which requires characterizing the (conditional) moments of $\tilde{e}_t^{[i]}$. One caveat is that by construction, $\tilde{e}_t^{[i]}$ could have a different dependence structure from the original sequence $e_t^{[i]}$. For example, even if $e_t^{[i]}$ is independent over t conditional on \mathcal{H} , $\tilde{e}_t^{[i]}$ would be $(T - T_i + 1)$ -dependent conditional on \mathcal{H} in general, that is, $\tilde{e}_t^{[i]}$ is independent of $\tilde{e}_{t+\ell}^{[i]}$ conditional on \mathcal{H} if $\ell \geq T - T_i + 1$.

Example 4.3 (Average Treatment Effect on the Treated at s_0 after k Periods, τ_{k,s_0} , continued).

In this scenario, the out-of-sample error is given by

$$e_{\tau_{k,s_0}} := \frac{1}{n_{s_0}} \sum_{i:T_i=s_0} \left(Y_{i(s_0+k)}(\infty) - \mathbf{p}_{s_0+k}^{[i]'} \boldsymbol{\beta}_0^{[i]} \right) = \frac{1}{n_{s_0}} \sum_{i:T_i=s_0} e_{s_0+k}^{[i]}.$$

The out-of-sample error above is similar to that defined in Example 4.2 except that $e_{\tau_{k,s_0}}$ is a cross-sectional average of $e_t^{[i]}$ rather than a time-series average. The uncertainty quantification strategy outlined in Example 4.2 can still be applied, with the caveat that it is uncommon in SC analysis to assume $e_t^{[i]}$ is stationary and/or independent over i . By contrast, it is reasonable to assume $e_t^{[i]}$ is stationary and/or independent (at least weakly dependent) over time.

Example 4.4 (Average Treatment Effect on all Treated after k Periods, τ_k , continued). In this scenario, the out-of-sample error is given by

$$e_{\tau_k} := \frac{1}{N_1} \sum_{i=1}^{N_1} \left(Y_{i(T_i+k)}(\infty) - \mathbf{p}_{T_i+k}^{[i]'} \boldsymbol{\beta}_0^{[i]} \right) = \frac{1}{N_1} \sum_{i=1}^{N_1} e_{T_i+k}^{[i]}.$$

Since the adoption time T_i may be heterogeneous across i , $e_{\tau,k}$ is an average of the individual errors $e_t^{[i]}$ of different units in different periods. Again, one can characterize the (conditional) moments of each $e_t^{[i]}$ and use the concentration-inequality-based approach to bound $e_{\tau,k}$. By contrast, it would be difficult to implement the other strategy outlined in Example 4.2 that relies on constructed pre-treatment averaged errors analogous to $e_{\tau,k}$, since such averages may include errors that are far away in time and their dependence on \mathcal{H} may be very complex.

In addition to the concentration-based approach described above, other strategies, including location-scale models and quantile regression, were proposed in Cattaneo, Feng and Titiunik (2021) for out-of-sample uncertainty quantification. We briefly review them in Supplemental Appendix S.1.1.

Main theorem. We present our main theorem, which shows that the prediction interval constructed above achieves approximately $(1 - \alpha_1 - \alpha_2)$ conditional coverage probability, which holds with high probability on \mathcal{H} . We use $\|\cdot\|_*$ to denote the dual norm of $\|\cdot\|$ on \mathbb{R}^d , use $\|\cdot\|_F$ to denote the Frobenius matrix norm, and use $\mathcal{B}(\mathbf{0}, \varepsilon)$ to denote an ε -neighborhood around zero for some $\varepsilon > 0$.

Theorem 1. Assume \mathcal{W} and \mathcal{R} are convex, $\hat{\beta}$ in (3.1) and β_0 in (4.2) exist, $\mathcal{H} = \sigma(\mathbf{B}, \mathbf{C}, \mathbf{p}_\tau)$, and $M_{1,L}$, $M_{1,U}$, $M_{2,L}$ and $M_{2,U}$ are specified as in (4.4) and (4.5). In addition, for some finite non-negative constants ϵ_γ , π_γ , ϖ_δ^* , ϵ_δ^* , π_δ^* , ϖ_Δ^* , ϵ_Δ^* , π_Δ^* , $\epsilon_{\gamma,1}^*$, $\epsilon_{\gamma,2}^*$ and π_γ^* , the following conditions hold:

- (i) $\mathbb{P}[\mathbb{P}(\mathbf{p}'_\tau(\hat{\beta} - \beta_0) \in [\mathbf{c}_L(\alpha_0), \mathbf{c}_U(1 - \alpha_0)] | \mathcal{H}) \geq 1 - \alpha_0 - \epsilon_\gamma] \geq 1 - \pi_\gamma$ for any $\alpha_0 \in (0, 1)$;
- (ii) $\mathbb{P}[\mathbb{P}(\sup\{\|\delta\| : \delta \in \mathcal{M}_G\} \leq \varpi_\delta^* | \mathcal{H}) \geq 1 - \epsilon_\delta^*] \geq 1 - \pi_\delta^*$;
- (iii) $\mathbb{P}[\mathbb{P}(\sup_{\mathbf{a} \in \Delta \cap \mathcal{B}(0, \varpi_\delta^*)} \text{dist}(\mathbf{a}, \Delta^* \cap \mathcal{B}(0, \varepsilon)) \leq \varpi_\Delta^* | \mathcal{H}) \geq 1 - \epsilon_\Delta^*] \geq 1 - \pi_\Delta^*$;
- (iv) $\mathbb{P}[\mathbb{P}(\|\Sigma^{-1/2} \hat{\Sigma} \Sigma^{-1/2} - \mathbf{I}_d\|_F \leq 2\epsilon_{\gamma,1}^* | \mathcal{H}) \geq 1 - \epsilon_{\gamma,2}^*] \geq 1 - \pi_\gamma^*$;
- (v) $e_\tau - \mathbb{E}[e_\tau | \mathcal{H}]$ is sub-Gaussian conditional on \mathcal{H} with parameter $\sigma_{\mathcal{H}}$.

Then, for $\epsilon_{\gamma,1}^* \in [0, 1/4]$,

$$\mathbb{P}\left\{\mathbb{P}(\tau \in [\hat{\tau} + M_{1,L} - M_{2,U} - \varepsilon_\Delta, \hat{\tau} + M_{1,U} - M_{2,L} + \varepsilon_\Delta] | \mathcal{H}) \geq 1 - \alpha_1 - \alpha_2 - \epsilon\right\} \geq 1 - \pi,$$

where $\epsilon = \epsilon_\gamma + 2\epsilon_{\gamma,1}^* + \epsilon_{\gamma,2}^* + 2\epsilon_\delta^* + \epsilon_\Delta^*$, $\pi = \pi_\gamma + \pi_\gamma^* + \pi_\delta^* + \pi_\Delta^*$ and $\varepsilon_\Delta = \|\mathbf{p}_\tau\|_* \varpi_\Delta^*$.

Assumptions (i)-(iv) are high-level conditions used for in-sample uncertainty quantification, which can be verified in many practically relevant scenarios. See more detailed discussion in Section 4.1. Condition (v), as we emphasized before, is a moment condition on e_τ that is used to showcase our out-of-sample uncertainty quantification strategy and can be relaxed by utilizing other appropriate concentration inequalities. Moreover, the constant ε_Δ in the theorem is used to adjust the prediction interval for nonlinear constraints. In many SC applications with linear constraints only (e.g., simplex or lasso constraint), such adjustment is *unnecessary* and we can set $\varepsilon_\Delta = 0$. See Section 5.3 for more discussion.

4.1 Discussion of Conditions (i)-(iv)

Theorem 1 relies on the high-level conditions (i)-(iv). We discuss each of them in more detail.

- **Condition (i).** This condition formalizes the idea of distributional approximation of $\widehat{\gamma} - \gamma$ by a Gaussian vector \mathbf{G} . Lemma S.1 in the Supplemental Appendix verifies (i) by assuming the error term $(u_{t,1}^{[1]}, \dots, u_{t,M}^{[1]}, \dots, u_{t,1}^{[N_1]}, \dots, u_{t,M}^{[N_1]})'$ is independent over $1 \leq t \leq T_0$ conditional on \mathcal{H} . In fact, (i) also holds when the errors are only weakly dependent (e.g., β -mixing) conditional on \mathcal{H} . See more discussion in Section S.2.2 of the Supplemental Appendix. Importantly, the features included in \mathbf{A} and \mathbf{B} could be non-stationary, thus covering the cointegration case which is common in SC applications.
- **Condition (ii).** This is a mild condition on the concentration of $\boldsymbol{\delta} \in \mathcal{M}_{\mathbf{G}}$. The requirement $\boldsymbol{\delta}' \widehat{\mathbf{Q}} \boldsymbol{\delta} - 2\mathbf{G}' \boldsymbol{\delta} \leq 0$ is usually known as the *basic inequality* in regression analysis (see Chapter 7 of [Wainwright \(2019\)](#) for the example of lasso). The vector \mathbf{G} is (conditional) Gaussian by construction, making condition (ii) easy to verify based on well-known bounds for Gaussian distributions. This condition holds in a variety of empirically relevant settings, including outcomes-only regression with i.i.d. data, multi-equation regression with weakly dependent data, and cointegrated outcomes and features settings.
- **Condition (iii).** This is a high-level requirement on the “closeness” between Δ and Δ^* . We propose a strategy for constructing Δ^* that can be shown to satisfy (iii) if the constraints specified

in \mathcal{W} and \mathcal{R} are formed by smooth functions. Suppose that

$$\mathcal{W} \times \mathcal{R} = \left\{ \boldsymbol{\beta} \in \mathbb{R}^d : \mathbf{m}_{\text{eq}}(\boldsymbol{\beta}) = \mathbf{0}, \mathbf{m}_{\text{in}}(\boldsymbol{\beta}) \leq \mathbf{0} \right\},$$

where $\mathbf{m}_{\text{eq}}(\cdot) \in \mathbb{R}^{d_{\text{eq}}}$ and $\mathbf{m}_{\text{in}}(\cdot) \in \mathbb{R}^{d_{\text{in}}}$ and d_{eq} and d_{in} denote the number of equality and inequality constraints in $\mathcal{W} \times \mathcal{R}$, respectively. Let the j th constraint in $\mathbf{m}_{\text{in}}(\cdot)$ be $m_{\text{in},j}(\cdot)$. Given tuning parameters $\varrho_j > 0$, $j = 1, \dots, d_{\text{in}}$, let $\mathcal{A} = \{j_1, \dots, j_k\}$ denote the set of indices for the inequality constraints such that $m_{\text{in},j}(\widehat{\boldsymbol{\beta}}) > -\varrho_j$. Then define

$$\Delta^* = \left\{ \boldsymbol{\beta} - \widehat{\boldsymbol{\beta}} : \mathbf{m}_{\text{eq}}(\boldsymbol{\beta}) = \mathbf{0}, m_{\text{in},j}(\boldsymbol{\beta}) \leq m_{\text{in},j}(\widehat{\boldsymbol{\beta}}) \text{ for } j \in \mathcal{A}, m_{\text{in},l}(\boldsymbol{\beta}) \leq \mathbf{0} \text{ for } l \notin \mathcal{A} \right\}. \quad (4.6)$$

The following lemma verifies condition (iii) for this Δ^* . We use $s_{\min}(\mathbf{M})$ to denote the minimum singular value of a matrix \mathbf{M} .

Lemma 1. *Let $\|\cdot\|$ be the Euclidean norm for vectors and the spectral norm for matrices. Assume that with probability over \mathcal{H} at least $1 - \pi_{\Delta}^*$, the following conditions hold: (i) $\mathbb{P}(\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0\| \leq \varpi_{\delta}^* | \mathcal{H}) \geq 1 - \epsilon_{\Delta}^*$; (ii) $\mathbf{m}(\cdot) = (\mathbf{m}_{\text{eq}}(\cdot)', \mathbf{m}_{\text{in}}(\cdot)')$ is twice continuously differentiable on $\mathcal{B}(\boldsymbol{\beta}_0, \varpi_{\delta}^*)$ with $\inf_{\boldsymbol{\beta} \in \mathcal{B}(\boldsymbol{\beta}_0, \varpi_{\delta}^*)} s_{\min}(\frac{\partial}{\partial \boldsymbol{\beta}} \mathbf{m}(\boldsymbol{\beta})) \geq c_{\min}$ for some constant $c_{\min} > 0$; and (iii) for all $1 \leq j \leq d_{\text{in}}$, $\varrho_j \in (\mathfrak{c}\varpi_{\delta}^*, |m_{\text{in},j}(\boldsymbol{\beta}_0)| - \mathfrak{c}\varpi_{\delta}^*)$ for some $\mathfrak{c} > 0$ specified in the proof. Then, for Δ^* defined in (4.6), condition (iii) in Theorem 1 holds with $\varpi_{\Delta}^* = \mathfrak{C}(\varpi_{\delta}^*)^2$ for some constant $\mathfrak{C} > 0$.*

In this lemma, the tuning parameters ϱ_j are introduced to guarantee that with high probability, we can correctly differentiate the binding inequality constraints from the other non-binding ones. In Section 5 below, we provide more practical details about choosing ϱ_j . Also, the concentration requirement for $\widehat{\boldsymbol{\beta}}$ specified in this lemma is usually mild. Since $\widehat{\boldsymbol{\beta}}$ satisfies the basic inequality $(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0)' \widehat{\mathbf{Q}} (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) - 2(\widehat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})' (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq 0$, the concentration of $\widehat{\boldsymbol{\beta}}$ can be shown by combining a distributional approximation of $\widehat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}$ by a Gaussian vector \mathbf{G} and the idea outlined in the previous discussion about condition (ii).

- **Condition (iv).** This is a requirement that $\widehat{\boldsymbol{\Sigma}}$ be a “good” approximation of the unknown covariance matrix $\boldsymbol{\Sigma}$. Many standard covariance estimation strategies such as the family of well-known heteroskedasticity-consistent estimators can be utilized.

4.2 Simultaneous Prediction Intervals

So far we have focused on constructing prediction intervals that have high coverage of the desired treatment effects, in particular, the individual treatment effect in *each* post-treatment period. In some applications, it might be appealing to construct prediction intervals that have high *simultaneous* coverage in multiple post-treatment periods, usually termed *simultaneous prediction intervals* in the literature. They can be employed to test, for example, whether the largest (or smallest) treatment effect across different periods is significantly different from zero.

Specifically, for a particular treated unit $1 \leq i \leq N_1$, we aim to construct a sequence of intervals \mathcal{I}_k for $0 \leq k \leq L$ for some $L \leq T - T_i$ such that

$$\mathbb{P}\left\{\mathbb{P}[\tau_{ik} \in \mathcal{I}_k, \text{ for all } 0 \leq k \leq L \mid \mathcal{H}] \geq 1 - \alpha\right\} \geq 1 - \pi.$$

As described before, the uncertainty of the predicted individual treatment effect $\widehat{\tau}_{ik}$ comes from the in-sample error $\mathbf{p}'_{\tau_{ik}}(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0)$ and the out-of-sample error $e_{T_i+k}^{[i]}$.

Regarding the in-sample error, the following is an immediate generalization of the prediction interval described in (4.4), which enjoys simultaneous coverage in multiple periods. If the constraints imposed in Δ are linear (such as simplex and lasso constraints), set

$$M_{1,L} := (\alpha_1/2)\text{-quantile of } \inf \left\{ \mathbf{p}'_{\tau_{ik}} \boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta^*, \ell^*(\boldsymbol{\delta}) \leq 0, 0 \leq k \leq L \right\} \text{ and}$$

$$M_{1,U} := (1 - \alpha_1/2)\text{-quantile of } \sup \left\{ \mathbf{p}'_{\tau_{ik}} \boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta^*, \ell^*(\boldsymbol{\delta}) \leq 0, 0 \leq k \leq L \right\},$$

which guarantees that with high probability,

$$\mathbb{P}[M_{1,L} \leq \mathbf{p}'_{\tau_{ik}}(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq M_{1,U} \text{ for all } 0 \leq k \leq L \mid \mathcal{H}] \geq 1 - \alpha_1.$$

If the constraints imposed in Δ are nonlinear (such as the ridge-type constraint), further decrease the lower bound $M_{1,L}$ and increase the upper bound $M_{1,U}$ defined above by some $\bar{\varepsilon}_\Delta$. We regard $\bar{\varepsilon}_\Delta$ as a small tuning parameter used to adjust for nonlinear constraints. See more discussion about selecting this parameter in Section 5.3.

Regarding the out-of-sample error, the goal is to find $M_{2,L}$ and $M_{2,U}$ such that with high proba-

bility,

$$\mathbb{P}[M_{2,L} \leq e_{T_i+k}^{[i]} \leq M_{2,U} \text{ for all } 0 \leq k \leq L \mid \mathcal{H}] \geq 1 - \alpha_2.$$

An easy-to-implement strategy analogous to that described in (4.5) is to adjust the intervals based on maximal inequalities. For example, suppose that each $e_{T_i+k}^{[i]} - \mathbb{E}[e_{T_i+k}^{[i]} \mid \mathcal{H}]$, $0 \leq k \leq L$, is conditional sub-Gaussian with parameter $\sigma_{\mathcal{H},k}$ (but is not necessarily independent over k). Then,

$$\mathbb{P}\left(\max_{0 \leq k \leq L} |e_{T_i+k}^{[i]} - \mathbb{E}[e_{T_i+k}^{[i]} \mid \mathcal{H}]| \geq \varepsilon \mid \mathcal{H}\right) \leq 2 \sum_{k=0}^L \exp\left(-\frac{\varepsilon^2}{2\sigma_{\mathcal{H},k}^2}\right).$$

If $\sigma_{\mathcal{H},k} \leq \sigma_{\mathcal{H}}$ for all $0 \leq k \leq T - T_i$, then one can take $M_{2,L} = \mathbb{E}[e_{T_i+k}^{[i]} \mid \mathcal{H}] - \varepsilon$ and $M_{2,U} = \mathbb{E}[e_{T_i+k}^{[i]} \mid \mathcal{H}] + \varepsilon$ with $\varepsilon = \sqrt{2\sigma_{\mathcal{H}}^2 \log(2(L+1)/\alpha_2)}$. Compared with prediction intervals with validity for each period constructed the same way, these simultaneous prediction intervals are slightly wider due to the additional factor $\sqrt{\log(L+1)}$. In practice, one only needs to estimate the conditional mean and variance of $e_t^{[i]}$ using the pre-treatment residuals; flexible parametric or non-parametric estimation methods can be used.

Again, the sub-Gaussianity assumption can be relaxed by using other concentration inequalities requiring weaker moment conditions, though the resulting simultaneous prediction intervals may be wider. Also, there are other strategies to construct prediction intervals that simultaneously cover multiple out-of-sample errors, though they are computationally more cumbersome and usually require more stringent conditions. See Appendix S.1.2 for a brief discussion.

The idea outlined above to achieve simultaneous coverage is general and can also be used to, for example, construct prediction intervals that simultaneously cover treatment effects for multiple treated units rather than for multiple post-treatment periods. In our empirical application, we construct simultaneous prediction intervals for average post-treatment effects across different economies; see details in Section 6.

5 Optimization and Tuning Parameter Selection

This section discusses the scalable robust optimization implementations and principled tuning parameter selection based on our theoretical results. We first show in Section 5.1 that our proposed SC methods for prediction and uncertainty quantification can be recast as conic optimization pro-

grams, which gives massive speed improvement in practice. Then, Sections 5.2 and 5.3 propose easy-to-implement methods for selecting the two kinds of tuning parameters that are used to construct the proposed prediction intervals: $\{\varrho_j : 1 \leq j \leq d_{\text{in}}\}$ and ε_Δ . Recall that ϱ_j s are used to define the constraint set Δ^* in simulation in order to approximately preserve the local geometry of the original constraint set Δ , while ε_Δ is an adjustment of the bounds on the in-sample error that takes into account the “distance” between Δ and Δ^* . In most SC applications, each constraint is imposed on parameters associated with one treated unit rather than multiple treated units, which will be the focus of our discussion below.

5.1 Conic Programming Approach

We first introduce three common types of convex optimization problems: the quadratically constrained linear problem (QCLP), the quadratically constrained quadratic program (QCQP), and the second-order cone program (SOCP). Second, we illustrate the link between these families of convex problems. Finally, the optimization problems underlying the prediction/estimation and uncertainty quantification problems for SC presented in Section 3 are QCQPs and QCLPs, respectively, and thus we show how to represent them as SOCPs to obtain scalable robust implementations. For background knowledge and technical details, see [Boyd and Vandenberghe \(2004\)](#).

The QCQPs and QCLPs are constrained optimization problems of the following form:

$$\begin{aligned} \min_{\mathbf{x}} \quad & \mathbf{x}'\mathbf{P}_0\mathbf{x} + \mathbf{q}'_0\mathbf{x} + w & (5.1) \\ \text{subject to} \quad & \mathbf{x}'\mathbf{P}_j\mathbf{x} + \mathbf{q}'_j\mathbf{x} + r_j \leq 0, \quad j = 1, \dots, m & \text{(Quadratic inequality constraint)} \\ & \mathbf{F}\mathbf{x} = \mathbf{g}, & \text{(Linear equality constraint)} \end{aligned}$$

where $\mathbf{P}_0, \mathbf{P}_1, \dots, \mathbf{P}_m \in \mathcal{M}_{n \times n}(\mathbb{R})$, $\mathbf{q}_0, \mathbf{q}_1, \dots, \mathbf{q}_m \in \mathbb{R}^n$, $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{F} \in \mathbb{R}^{m \times n}$, $\mathbf{g} \in \mathbb{R}^m$, and $r_0, r_1, \dots, r_m, w \in \mathbb{R}$. If all the matrices $\mathbf{P}_0, \mathbf{P}_1, \dots, \mathbf{P}_m$ are positive semi-definite, the QCQP is convex. Moreover, if $\mathbf{P}_0 = \mathbf{0}$ the QCQP becomes a QCLP. For this reason, in what follows we will restrict our attention to QCQPs as they naturally embed QCLPs.

The program (5.1) above can be recast as a SOCP. We use $\|\cdot\|_p$ to denote the ℓ_p vector norm for $p \geq 1$. Let \mathcal{K} be a cone such that $\mathcal{K} = \mathbb{R}_+^m \times \mathcal{K}_1 \times \mathcal{K}_2 \times \dots \times \mathcal{K}_L$ where $\mathcal{K}_l := \{(k_0, \mathbf{k}_1) \in \mathbb{R} \times \mathbb{R}^l : \|\mathbf{k}_1\|_2 \leq k_0\}$, $l = 1, \dots, L$. Let $\preceq_{\mathcal{K}}$ be the *generalized inequality* associated with the cone

\mathcal{K} (see Supplemental Appendix Section S.3.1 for more technical details). An optimization problem is called a *second-order cone program* if it has the following form

$$\begin{aligned} \min_{\mathbf{x}} \quad & \mathbf{c}'\mathbf{x}, & (5.2) \\ \text{subject to} \quad & \mathbf{G}\mathbf{x} \preceq_{\mathcal{K}} \mathbf{h}, & \text{(Second-order cone constraint)} \\ & \mathbf{A}\mathbf{x} = \mathbf{b}. & \text{(Linear equality constraint)} \end{aligned}$$

The generic SC weight construction (3.1) is a QCQP, which can be recast in general as a SOCP. We illustrate the approach for the L1-L2 constraint.² Consider first the prediction/estimation SC optimization problem, which relies on the following program:

$$\begin{aligned} \min_{\mathbf{w}, \mathbf{r}} \quad & (\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r})'\mathbf{V}(\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r}) & (5.3) \\ \text{subject to} \quad & \|\mathbf{w}^{[i]}\|_1 = Q_1^{[i]}, \quad i = 1, \dots, N_1 & \text{(L1 equality constraints)} \\ & \|\mathbf{w}^{[i]}\|_2 \leq Q_2^{[i]}, \quad i = 1, \dots, N_1 & \text{(L2 inequality constraints)} \\ & \mathbf{w} \geq \mathbf{0}, & \text{(non-negativity constraint)} \end{aligned}$$

where, as always, \geq is understood as a component-wise inequality for vectors ($\mathbf{w} \in \mathbb{R}^{J \cdot N_1}$). First, notice that the non-convex constraints $\|\mathbf{w}^{[i]}\|_1 = Q_1^{[i]}, i = 1, \dots, N_1$ can be replaced with the convex constraints $\mathbf{1}'\mathbf{w}^{[i]} = Q_1^{[i]}, i = 1, \dots, N_1$ because of the non-negativity constraint on the elements of \mathbf{w} . Then, we can cast (5.3) as a SOCP as follows

$$\begin{aligned} \min_{\mathbf{w}, \mathbf{r}, v, \{s_i\}_{i=1}^{N_1}} \quad & v \\ \text{subject to} \quad & \mathbf{1}'\mathbf{w}^{[i]} = Q_1^{[i]}, & \text{(L1 equality constraints)} \\ & -\mathbf{w} \preceq_{\mathcal{C}_1} \mathbf{0}, & \text{(cone in } \mathbb{R}^{J \cdot N_1}) \\ & \left[\begin{array}{c} 1 - v \\ 2\mathbf{V}^{1/2}(\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r}) \end{array} \right] \preceq_{\mathcal{C}_2} 1 + v, & \text{(cone in } \mathbb{R}^{2+T_0 \cdot M \cdot N_1}) \\ & s_i \preceq_{\mathcal{C}_3} Q_1^{[i]}, \quad i = 1, \dots, N_1 & (N_1 \text{ cones in } \mathbb{R}) \end{aligned}$$

²Note that simplex, ridge, or least squares are particular cases of L1-L2. In the Supplemental Appendix 2.3, we also illustrate the case when \mathcal{W} is a lasso-type constraint, and provide more general details.

$$\begin{bmatrix} 1 - s_i \\ 2\mathbf{w}^{[i]} \end{bmatrix} \preceq_{\mathcal{C}_4} 1 + s_i, \quad i = 1, \dots, N_1 \quad (N_1 \text{ cones in } \mathbb{R}^{2+J})$$

where $\mathcal{K} = \mathcal{C}_1 \times \mathcal{C}_2 \times \mathcal{C}_3^{N_1} \times \mathcal{C}_4^{N_1} = \mathbb{R}_+^{J \cdot N_1} \times \mathcal{K}_{T_0 \cdot M \cdot N_1 + 1} \times \mathbb{R}_+^{N_1} \times \mathcal{K}_{J+1}^{N_1}$ is the conic constraint for this program.

For uncertainty quantification, we need to solve the optimization problem underlying (4.4). We discuss the lower bound only. Recalling that $\boldsymbol{\beta} = (\mathbf{w}', \mathbf{r}')'$, we have

$$\begin{aligned} \inf_{\boldsymbol{\beta}=(\mathbf{w}',\mathbf{r}')'} \quad & \mathbf{p}'_{\tau}(\boldsymbol{\beta} - \widehat{\boldsymbol{\beta}}) & (5.4) \\ \text{subject to} \quad & \|\mathbf{w}^{[i]}\|_1 = Q_1^{[i]}, \quad i = 1, \dots, N_1 & (\text{L1 equality constraints}) \\ & \|\mathbf{w}^{[i]}\|_2 \leq Q_2^{[i]} + \varrho_1^{[i]}, \quad i = 1, \dots, N_1 & (\text{L2 inequality constraints}) \\ & \mathbf{w} \geq -\boldsymbol{\varrho}_2, & (\text{non-negativity constraint}) \\ & (\boldsymbol{\beta} - \widehat{\boldsymbol{\beta}})' \widehat{\mathbf{Q}} (\boldsymbol{\beta} - \widehat{\boldsymbol{\beta}}) - 2(\mathbf{G}^*)'(\boldsymbol{\beta} - \widehat{\boldsymbol{\beta}}) \leq 0, & (\text{constrained least squares}) \end{aligned}$$

where the scalars $\varrho_1^{[i]}, i = 1, \dots, N_1$ and the vector $\boldsymbol{\varrho}_2$ are regularization parameters used to relax Δ (to Δ^* as discussed in Section 4).

We can cast the SC optimization problem in (5.4) in conic form as follows:

$$\begin{aligned} \min_{\mathbf{w}, \mathbf{r}, \{s_i\}_{i=1}^{N_1}, t} \quad & \mathbf{p}'_{\tau} \boldsymbol{\beta} \\ \text{subject to} \quad & \mathbf{1}' \mathbf{w}^{[i]} = Q_1^{[i]}, \quad i = 1, \dots, N_1 & (\text{L1 equality constraints}) \\ & t + \mathbf{a}' \boldsymbol{\beta} + f \preceq_{\mathcal{C}_1} 0, & (\text{cone in } \mathbb{R}) \\ & -\mathbf{w} \preceq_{\mathcal{C}_2} \boldsymbol{\varrho}_2, & (\text{cone in } \mathbb{R}^{J \cdot N_1}) \\ & s_i \preceq_{\mathcal{C}_3} Q_1^{[i]} + \varrho_1^{[i]}, \quad i = 1, \dots, N_1 & (N_1 \text{ cones in } \mathbb{R}) \\ & \begin{bmatrix} 1 - s_i \\ 2\mathbf{w}^{[i]} \end{bmatrix} \preceq_{\mathcal{C}_4} 1 + s_i, \quad i = 1, \dots, N_1 & (N_1 \text{ cones in } \mathbb{R}^{2+J}) \\ & \begin{bmatrix} 1 - t \\ 2\mathbf{Q}^{1/2} \boldsymbol{\beta} \end{bmatrix} \preceq_{\mathcal{C}_5} 1 + t, & (\text{cone in } \mathbb{R}^{2+(J+KM) \cdot N_1}) \end{aligned}$$

where $\mathcal{K} = \mathcal{C}_1 \times \mathcal{C}_2 \times \mathcal{C}_3 \times \mathcal{C}_4 \times \mathcal{C}_5 = \mathbb{R}_+ \times \mathbb{R}_+^{J \cdot N_1} \times \mathbb{R}_+^{N_1} \times \mathcal{K}_{1+J}^{N_1} \times \mathcal{K}_{1+(J+KM) \cdot N_1}$ is the conic constraint for this program, $\mathbf{a} = -2(' \widehat{\mathbf{Q}} \widehat{\boldsymbol{\beta}} + \mathbf{G}^*)'$, and $f = \widehat{\boldsymbol{\beta}}' \widehat{\mathbf{Q}} \widehat{\boldsymbol{\beta}} + 2\mathbf{G}^* \widehat{\boldsymbol{\beta}}$.

The approaches above are implemented in our companion general-purpose software ([Cattaneo](#),

Feng, Palomba and Titiunik, 2023), where we show that they lead to remarkable speed and scalability improvements. The supplemental appendix gives further discussion and other methodological details.

5.2 Defining Constraints in Simulation

In the proposed procedure, a sequence of tuning parameters ϱ_j , $j = 1, \dots, d_{\text{in}}$, is introduced to determine which inequality constraints are binding. We propose a feasible strategy to select ϱ_j . Consider a (generic) treated unit i , and suppose that it is associated with inequality constraints with indices in $\mathcal{S}^{[i]}$. We start with the idea of condition (ii) in Theorem 1 and use a parameter $\varrho^{[i]}$ to bound the deviation of $\widehat{\boldsymbol{\beta}}^{[i]}$ from $\boldsymbol{\beta}_0^{[i]}$. We use the formula

$$\varrho^{[i]} = \mathcal{C} \frac{\log(T_0)^c}{T_0^{1/2}},$$

where $c = 1/2$ if the data are i.i.d. or weakly dependent, and $c = 1$ if $\mathbf{A}^{[i]}$ and $\mathbf{B}^{[i]}$ form a cointegrated system, and \mathcal{C} is one of the following

$$\mathcal{C}_1 = \frac{\widehat{\sigma}_u}{\min_{1 \leq j \leq J} \widehat{\sigma}_{b_j}}, \quad \mathcal{C}_2 = \frac{\max_{1 \leq j \leq J} \widehat{\sigma}_{b_j} \widehat{\sigma}_u}{\min_{1 \leq j \leq J} \widehat{\sigma}_{b_j}^2}, \quad \mathcal{C}_3 = \frac{\max_{1 \leq j \leq J} \widehat{\sigma}_{b_j u}}{\min_{1 \leq j \leq J} \widehat{\sigma}_{b_j}^2}, \quad (5.5)$$

where $\widehat{\sigma}_{b_j, u}$ is the estimated (unconditional) covariance between the pseudo-true residual $\mathbf{u}^{[i]}$ and the j th column of $\mathbf{B}^{[i]}$ (the features of the j th control unit), and $\widehat{\sigma}_u$ and $\widehat{\sigma}_{b_j}$ are the estimated (unconditional) standard deviation of $\mathbf{u}^{[i]}$ and the j th column of $\mathbf{B}^{[i]}$, respectively. If the synthetic control weights were constructed based on both stationary and non-stationary features, the non-stationary components would govern the precision of the estimation. In such cases, one could ignore the stationary components and set $c = 1$.

Next, we define possibly heterogeneous parameters ϱ_j , $j \in \mathcal{S}^{[i]}$, for different inequality constraints associated with unit i . By the first-order Taylor expansion, if the j th inequality constraint is binding, i.e., $m_{\text{in},j}(\boldsymbol{\beta}_0^{[i]}) = 0$, then

$$m_{\text{in},j}(\widehat{\boldsymbol{\beta}}^{[i]}) \approx \frac{\partial}{\partial \boldsymbol{\beta}'} m_{\text{in},j}(\boldsymbol{\beta}_0^{[i]}) (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0).$$

Then, an intuitive choice of ϱ_j would be

$$\varrho_j = \left\| \frac{\partial}{\partial \boldsymbol{\beta}} m_{\text{in},j}(\widehat{\boldsymbol{\beta}}^{[i]}) \right\|_1 \times \varrho^{[i]}, \quad j \in \mathcal{S}^{[i]}. \quad (5.6)$$

If $m_{\text{in},j}(\widehat{\boldsymbol{\beta}}^{[i]}) > -\varrho_j$, we let the j th constraint be binding in the simulation.

5.3 Adjustment for Nonlinear Constraints

When some constraints in Δ are nonlinear (e.g., ridge-type constraints), we introduce a constant ε_Δ to adjust the bounds on the in-sample error; this constant depends on the distance ϖ_Δ^* between the localized constraint sets Δ and Δ^* specified in condition (iii) of Theorem 1. This adjustment is only necessary for nonlinear constraints; it is not needed when the constraints are linear in parameters (e.g., simplex or lasso constraints).

The distance between Δ and Δ^* typically depends on the the first and second derivatives of the constraint functions $\mathbf{m}_{\text{in}}(\cdot)$. Again, we first focus on the inequality constraints related to one particular treated unit i . Denote by $\mathbf{m}_{\text{in},\mathcal{S}^{[i]}}(\cdot)$ the vector of constraint functions $m_{\text{in},j}(\cdot)$ with $j \in \mathcal{S}^{[i]}$. We propose to set

$$\varepsilon_\Delta^{[i]} = \|\mathbf{p}_\tau^{[i]}\|_1 \times \frac{\sqrt{|\mathcal{S}^{[i]}|}}{2} s_{\min}^{-1} \left(\frac{\partial}{\partial \boldsymbol{\beta}} \mathbf{m}_{\text{in},\mathcal{S}^{[i]}}(\widehat{\boldsymbol{\beta}}^{[i]}) \right) \times \max_{j \in \mathcal{S}^{[i]}} s_{\max} \left(\frac{\partial}{\partial \boldsymbol{\beta} \partial \boldsymbol{\beta}'} m_{\text{in},j}(\widehat{\boldsymbol{\beta}}^{[i]}) \right) \times (\varrho^{[i]})^2,$$

where $\mathbf{p}_\tau^{[i]}$ denotes the subvector of \mathbf{p}_τ that corresponds to $\boldsymbol{\beta}_0^{[i]}$, and $|\mathcal{S}^{[i]}|$ denotes the cardinality of $\mathcal{S}^{[i]}$. Denote by \mathcal{N}_τ the set of treated units to which the causal predictand τ is related. Then set

$$\varepsilon_\Delta = \sum_{i \in \mathcal{N}_\tau} \varepsilon_\Delta^{[i]}.$$

When constraints are linear, the second derivative of $m_{\text{in},j}(\cdot)$ is zero, and the above choice of ε_Δ is exactly 0.

One example of nonlinear constraints that is commonly used in practice is the ridge-type restriction. Specifically, suppose that there is one treated unit i and one constraint

$$\|\boldsymbol{\beta}^{[i]}\|_2^2 = 1.$$

The above choice of ε_Δ simplifies to

$$\varepsilon_\Delta^{[i]} = \|\mathbf{p}_\tau^{[i]}\|_1 \times (2\|\widehat{\boldsymbol{\beta}}^{[i]}\|_2)^{-1} \times (\varrho^{[i]})^2.$$

Remark 1 (Simultaneous Prediction Intervals). In Section 4.2, we discussed how to construct simultaneous prediction intervals. In the case of nonlinear constraints, a tuning parameter $\bar{\varepsilon}_\Delta$ was introduced to adjust the bounds on the in-sample errors. In this context, we can apply the procedure described above to each period $T_i + k$, $0 \leq k \leq L$, which gives us a sequence of constants, denoted by $\varepsilon_{\Delta,k}$, $0 \leq k \leq L$. Then, we let $\bar{\varepsilon}_\Delta = \max_{0 \leq k \leq L} \varepsilon_{\Delta,k}$. \square

5.4 Summary of Algorithmic Implementation

This section summarizes the main procedures underlying the computation of the synthetic control weights $\widehat{\mathbf{w}}$ (Algorithm 1) and the quantification of in-sample and out-of-sample uncertainty (Algorithm 2). We use $\mathfrak{q}_Z(\alpha)$ to denote the α -th quantile of the random variable Z , and assume that the researcher has collected some data denoted by $\{(Y_{it}, \mathbf{X}_{it}, T_i)_{i=1}^N : t = 1, \dots, T\}$, where Y_{it} indicates the outcome of interest, \mathbf{X}_{it} is a vector of other features, and T_i is a variable indicating the treatment timing.

Algorithm 1 Computation of $\widehat{\mathbf{w}}$

Require: data $\{(Y_{it}, \mathbf{X}_{it}, T_i)_{i=1}^N : t = 1, \dots, T\}$

define the conditioning set $\mathcal{H} = \{\mathbf{B}, \mathbf{C}, \mathbf{p}_\tau\}$

a: choose target predictand τ (see Section 3)

b: choose features in $(Y_{it}, \mathbf{X}_{it})$ to include in \mathbf{A} and \mathbf{B}

c: choose adjustment covariates to include in \mathbf{C} (e.g., linear trend, constant)

choose shape of feasibility sets \mathcal{W} and \mathcal{R} (see Section 3.1)

choose a symmetric weighting matrix \mathbf{V}

solve optimization problem in (3.1) and get $\widehat{\boldsymbol{\beta}} = (\widehat{\mathbf{w}}', \widehat{\mathbf{r}})'$

Algorithm 2 Uncertainty Quantification

Require: choose levels $\alpha_1, \alpha_2 \in (0, 1)$ and the number of simulations S

In-sample Uncertainty Quantification

form feasible constraint set Δ^*

a: calculate ϱ using (5.5) and using $\varrho_j, j = 1, \dots, d_{\text{in}}$ (5.6)

b: obtain $\mathcal{A} = \{j : \mathbf{m}_{\text{in},j}(\widehat{\boldsymbol{\beta}}) > -\varrho_j\}$

c: get Δ^* using (4.6)

estimate $\boldsymbol{\Sigma}$

a: estimate conditional moments of the pseudo-residual \mathbf{u} , $\mathbb{E}[\mathbf{u}|\mathcal{H}]$ and $\mathbb{V}[\mathbf{u}|\mathcal{H}]$

b: compute $\widehat{\boldsymbol{\Sigma}} = (\mathbf{Z}'\mathbf{V})\widehat{\mathbb{V}}[\mathbf{u}|\mathcal{H}](\mathbf{V}\mathbf{Z})$ where $\mathbf{Z} = (\mathbf{B}, \mathbf{C})$

for $s \leftarrow 1$ to S **do:**

a: draw $\mathbf{G}^* \sim \mathbf{N}(\mathbf{0}, \widehat{\boldsymbol{\Sigma}})$

b: define $\ell^*(\boldsymbol{\beta}) = \boldsymbol{\beta}'\widehat{\mathbf{Q}}\boldsymbol{\beta} - 2(\mathbf{G}^*)'\boldsymbol{\beta}$

c: solve

$$l_{(s)} := \inf_{\substack{\boldsymbol{\beta} \in \Delta^* \\ \ell^*(\boldsymbol{\beta}) \leq 0}} \mathbf{p}'_{\tau}\boldsymbol{\beta} \quad \text{and} \quad u_{(s)} := \sup_{\substack{\boldsymbol{\beta} \in \Delta^* \\ \ell^*(\boldsymbol{\beta}) \leq 0}} \mathbf{p}'_{\tau}\boldsymbol{\beta}$$

get $M_{1,L}$ as the $(\alpha_1/2)$ -quantile of $\{l_{(s)}\}_{s=1}^S$ and $M_{1,U}$ as the $(1 - \alpha_1/2)$ -quantile of $\{u_{(s)}\}_{s=1}^S$

Out-of-sample Uncertainty Quantification

procedure SUB-GAUSSIAN:

a: use pre-treatment residuals to estimate $\mathbb{E}[e_{\tau}|\mathcal{H}]$ and $\sigma_{\mathcal{H}}^2$

b: get

$$M_{2,L} := \widehat{\mathbb{E}}[e_{\tau}|\mathcal{H}] - \sqrt{2\hat{\sigma}_{\mathcal{H}}^2 \log(2/\alpha_2)} \quad \text{and} \quad M_{2,U} := \widehat{\mathbb{E}}[e_{\tau}|\mathcal{H}] + \sqrt{2\hat{\sigma}_{\mathcal{H}}^2 \log(2/\alpha_2)}$$

end procedure

A more detailed discussion of the procedures, including recommended rules of thumbs for implementation and other practical regularization choices, can be found in [Cattaneo, Feng, Palomba and Titiunik \(2023\)](#), Section 3.1 and Section 4.3).

6 Empirical Application

We illustrate our methodology with a re-analysis of the question studied in [Billmeier and Nannicini \(2013\)](#). For brevity, we only report results for European countries; the interested reader can find complete results for Africa, Asia, North America, and South America in Section S.6 of the Supplemental Appendix. We study the four causal predictands introduced in Section 3, where the i subscript indexes the countries in our sample: (i) individual country treatment effects 1 to

10 periods after the liberalization occurred (Example 3.1, τ_{ik} for $k = 1, \dots, 10$); (ii) average post-treatment effect for each country (Example 3.2, τ_i); (iii) average treatment effect on countries liberalized in 1991, 1 to 10 periods after liberalization (Example 3.3, $\tau_{k,1991}$ for $k = 1, \dots, 10$); (iv) average treatment effect on all liberalized countries, 1 to 10 periods after liberalization (Example 3.4, τ_k for $k = 1, \dots, 10$).

6.1 Empirical Strategy

In our main specification, we match on two features ($M = 2$), logarithm of GDP per capita and percentage of complete secondary schooling attained in population. We obtain the SC weights under the L1-L2 constraint, i.e.,

$$\mathcal{W} = \times_{i=1}^{N_1} \left\{ \mathbf{w}^{[i]} \in \mathbb{R}^J : \|\mathbf{w}^{[i]}\|_1 = 1, \|\mathbf{w}^{[i]}\|_2 \leq Q_2^{[i]} \right\},$$

and conduct covariate adjustment by including a constant term that is common across features, and a constant term and a linear time trend for each feature. We impose no constraint on these additional parameters, i.e., $\mathcal{R} = \mathbb{R}^{KMN_1}$.³ We set the weighting matrix $\mathbf{V} = \mathbf{I}$. The vector of predictors \mathbf{p}_t contains the (log) GDP per capita of countries in the donor pool in each post-treatment period, a constant term, and a linear trend. As described in Section 3, given a particular treatment effect τ , the predictor \mathbf{p}_τ is defined accordingly. Table 2 describes the matrices in detail.

In addition, since political and economic reforms do not happen overnight, we take into account the possibility of anticipation effects up to 1 year before the treatment. For example, Albania underwent a process of economic liberalization in 1992, according to the Sachs-Warner indicator. To address the presence of plausible anticipation effects, we define the pre-treatment period for Albania as 1963-1990 rather than 1963-1991.

In order to quantify the in-sample uncertainty from estimating the SC weights, we need to construct the bounds $M_{1,L}$ and $M_{1,U}$ on $\mathbf{p}'_\tau(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0)$. The following strategy is adopted. First, we treat the synthetic control weights as possibly misspecified, thus estimating both the first and second conditional moments of the pseudo-true residuals \mathbf{u} . The conditional first moment $\mathbb{E}[\mathbf{u} | \mathcal{H}]$ is estimated feature-by-feature using a linear-in-parameters regression of the residual

³In Section S.5 of the Supplemental Appendix, we replicate the whole analysis using the canonical ‘‘simplex’’ constraint, i.e., $\mathcal{W} = \times_{i=1}^{N_1} \{ \mathbf{w}^{[i]} \in \mathbb{R}_+^J : \|\mathbf{w}^{[i]}\|_1 = 1 \}$.

Table 2: *Description of main quantities used for prediction/inference.*

Matrix	Description
$\mathbf{A}^{[i]} = \begin{bmatrix} \mathbf{A}_1^{[i]} \\ \mathbf{A}_2^{[i]} \end{bmatrix}$	pre-treatment (log) GDP per capita and percentage of complete secondary schooling of the i th treated unit
$\mathbf{B}^{[i]} = \begin{bmatrix} \mathbf{B}_1^{[i]} \\ \mathbf{B}_2^{[i]} \end{bmatrix}$	pre-treatment GDP per capita and percentage of complete secondary schooling of the donors for the i th treated unit
$\mathbf{C}^{[i]} = \begin{bmatrix} \mathbf{1} & \mathbf{C}_1^{[i]} & \mathbf{0} \\ \mathbf{1} & \mathbf{0} & \mathbf{C}_2^{[i]} \end{bmatrix}$	additional covariates: a constant and a linear trend in each equation for the i th treated unit and a common constant across equations
$\mathbf{p}_t^{[i]} = \begin{bmatrix} \mathbf{x}_t^{[i]} \\ \mathbf{g}_t^{[i]} \end{bmatrix}$	post-treatment predictors for the i th treated unit: (log) GDP per capita of donors, $\mathbf{x}_t^{[i]}$, and a constant and a linear trend, so $\mathbf{g}_t^{[i]} = (1, t, 0, 0)$

Notes: all quantities in the table are defined for a generic treated unit $i = 1, \dots, N_1$. Feature 1 is GDP per capita and feature 2 is the percentage of complete secondary schooling attained in population. The vectors $\mathbf{0}$ and $\mathbf{1}$ are conformable vectors of zeros and ones, respectively.

$\hat{\mathbf{u}} = \mathbf{A} - \mathbf{B}\hat{\mathbf{w}} - \mathbf{C}\hat{\mathbf{r}}$ on \mathbf{B} and the first lag of \mathbf{B} , whereas the conditional second moment $\mathbb{V}[\mathbf{u} | \mathcal{H}]$ is estimated with an HC1-type estimator. We then draw $S = 200$ i.i.d. random vectors from the Gaussian distribution $\mathbf{N}(0, \hat{\mathbf{\Sigma}})$, conditional on the data, to simulate the criterion function $\ell_{(s)}^*(\boldsymbol{\beta} - \boldsymbol{\beta}_0) := (\boldsymbol{\beta} - \boldsymbol{\beta}_0)' \hat{\mathbf{Q}}(\boldsymbol{\beta} - \boldsymbol{\beta}_0) - 2\mathbf{G}'_{(s)}(\boldsymbol{\beta} - \boldsymbol{\beta}_0)$, $s = 1, \dots, 200$, and solve the following optimization problems

$$l_{(s)} := \inf_{\substack{\boldsymbol{\beta} - \boldsymbol{\beta}_0 \in \Delta^*, \\ \ell_{(s)}^*(\boldsymbol{\beta} - \boldsymbol{\beta}_0) \leq 0}} \mathbf{p}'_{\tau}(\boldsymbol{\beta} - \boldsymbol{\beta}_0) \quad \text{and} \quad u_{(s)} := \sup_{\substack{\boldsymbol{\beta} - \boldsymbol{\beta}_0 \in \Delta^*, \\ \ell_{(s)}^*(\boldsymbol{\beta} - \boldsymbol{\beta}_0) \leq 0}} \mathbf{p}'_{\tau}(\boldsymbol{\beta} - \boldsymbol{\beta}_0),$$

where Δ^* is constructed as explained in Section 4.1. Finally, $M_{1,L}$ is the $(\alpha_1/2)$ -quantile of $\{l_{(s)}\}_{s=1}^S$ and $M_{1,U}$ is the $(1 - \alpha_1/2)$ -quantile of $\{u_{(s)}\}_{s=1}^S$, where α_1 is set to 0.05.

In order to quantify the out-of-sample uncertainty from the stochastic error in the post-treatment period, we need to construct the bounds $M_{2,L}$ and $M_{2,U}$ on e_{τ} . We employ the non-asymptotic bounds described in (4.5), assuming that $e_{\tau} - \mathbb{E}[e_{\tau} | \mathcal{H}]$ is sub-Gaussian conditional on \mathcal{H} . We set $\alpha_2 = 0.05$, and the conditional mean $\mathbb{E}[e_{\tau} | \mathcal{H}]$ and the sub-Gaussian parameter $\sigma_{\mathcal{H}}$ are parametrized and estimated by a linear-in-parameters regression of the pre-treatment residuals on \mathbf{B} .

Finally, the prediction intervals for the counterfactual outcome and the treatment effect of interest are given by

$$\left[\mathbf{p}'_{\tau} \widehat{\boldsymbol{\beta}} - M_{1,U} + M_{2,L}; \mathbf{p}'_{\tau} \widehat{\boldsymbol{\beta}} - M_{1,L} + M_{2,U} \right] \quad \text{and} \quad [\widehat{\tau} + M_{1,L} - M_{2,U}; \widehat{\tau} + M_{1,U} - M_{2,L}],$$

respectively.



6.2 Results

Overall, our point predictions suggest that (external) liberalization episodes in Europe had a negative impact on GDP per capita. These findings might be explained by the nation's net trade position, among other things. For example, a negative effect on GDP can be justified if, following the liberalization event, the difference between imports and exports rises more than it would have in the absence of the treatment. This difference also has to offset the likely rise in consumption and investment. However, once we take uncertainty into account, the prediction intervals for the synthetic control always contain the realized GDP per capita series, implying that there is no strong evidence that the real income trajectory has been altered. In what follows, we report the predictands described in Section 3, Examples 3.1-3.4.

Individual country treatment effect, 1 to 10 periods after liberalization (τ_{ik}). In Figure 5 we show the predicted synthetic control outcomes (panel (a)) and treatment effects (panel (b)) with the corresponding 90% prediction intervals, and the computed weights $\widehat{\mathbf{w}}$ (panel (c)). We clearly see that in all six countries the realized trajectory of GDP per capita (black lines) lies below the synthetic one (blue lines), suggesting that in the absence of the liberalization event, real income per capita would have been higher. However, looking at the 90% prediction intervals (blue vertical bars), we can see that in most cases the distance between the actual GDP series and the counterfactual one is not different from zero with high probability for almost all units and periods. If we consider 90% simultaneous prediction intervals for each unit (blue shaded areas), it is clear that the realized and synthetic trajectories do not simultaneously differ with high probability.

Average post-treatment effect for each liberalized country (τ_i). The second causal predictand of our interest is the average post-treatment effect for each of the six European countries we study. Specifically, we target the average effect over the period following the liberalization up to the year



Figure 5: *Individual Treatment Effects $\hat{\tau}_{ik}$.*



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

2000. Figure 6 shows that in all countries the liberalization episode depressed the real income per capita. However, if we consider individual and simultaneous prediction intervals that possess high simultaneous coverage across treated units, we can see that no treated unit shows a negative average post-treatment effect with high probability.

Figure 6: Average Post-Treatment Effects $\hat{\tau}_i$.




(c) \hat{w}_i

Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals (that simultaneously cover six treated units). In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds. The small number at the bottom-right corner of panel (a) represents the number of periods over which the post-treatment average is computed.

Average treatment effect on countries liberalized in 1991 ($\tau_{k,1991}$). In this third exercise, we analyze the average treatment effect on the treated at 1991, that is, on countries that liberalized in 1991: Bulgaria, Czech Republic, Slovak Republic, and Slovenia. To study this causal predictand, we are again match on real GDP per capita and percentage of complete secondary schooling. In the

Supplemental Appendix Section S.7 we also report the results using a simplex-type constraint and the results using only GDP ($M = 1$). Even in this case, the trajectory for the synthetic real income lies above the realized one, suggesting a negative impact of the liberalization event. However, once again, quantifying uncertainty shows that, with high probability, we can conclude that the two series are not different from each other.

Figure 7: *Average Treatment Effects on the Treated in 1991 $\tau_{k,1991}$.*




Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.

Average treatment effect on all liberalized countries (τ_k). In this last and fourth exercise, we focus on a popular causal predictand: the average treatment effect on all the treated countries, that is, on all countries that liberalized, regardless of when they did so. We focus on the effects up to 10 periods after the adoption of liberalization, which occurs at different times for different countries. Figure 8 reports the results for liberalized countries in Europe. To compute this predictand, we first construct a synthetic control for each treated unit (see panel (c)) and then we pool such


synthetic controls together in the post-treatment period to get a single prediction and a single prediction intervals. Panel (a) and panel (b) show that pooling across the 7 European countries that embarked on liberalization program helps reducing the uncertainty surrounding the synthetic trajectory. Indeed, this predictand is the only one that suggests that the trajectories for synthetic real income and actual real income differ with high probability for the average treated country.

Figure 8: Average Treatment Effects on the Treated τ_k .



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$

(b) $\hat{\tau}_k$



(c) \hat{w}_i

Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.

7 Conclusion

We developed prediction intervals to quantify the uncertainty of a large class of synthetic control predictions or estimators in settings with staggered treatment adoption. Because many synthetic control applications have a limited number of observations, our inference procedures are based on non-asymptotic concentration arguments. The construction of our prediction intervals is designed to capture two sources of uncertainty: the first is the construction or estimation of the synthetic control weights with pre-treatment data, and the second is the variability of the post-treatment outcomes. By combining both sources in a prediction interval, our procedure offers precise non-asymptotic coverage probability guarantees, and allows researchers to implement sensitivity analyses to assess how robust the conclusions of the analysis are to various levels of uncertainty. Our framework is general, allowing for one or multiple treated units, simultaneous or staggered treatment adoption, linear or non-linear constraints, and stationary or non-stationary data. To enhance implementation, we also showed how to recast the methods as conic optimization programs and how to choose the necessary tuning parameters in a principled data-driven way. We illustrated our methods with an empirical application studying the effect of economic liberalization on GDP for European countries in the 1990s, motivated by the work of [Billmeier and Nannicini \(2013\)](#).

All our methods are implemented in `Python`, `R`, and `Stata` software, which is publicly available and discussed in detail in our companion article [Cattaneo, Feng, Palomba and Titiunik \(2023\)](#) and in Section S.4 of the Supplemental Appendix.

References

- Abadie, A. (2021), “Using Synthetic Controls: Feasibility, Data Requirements, and Methodological Aspects,” *Journal of Economic Literature*, 59, 391–425.
- Abadie, A., Diamond, A., and Hainmueller, J. (2010), “Synthetic Control Methods for Comparative Case Studies: Estimating the Effect of California’s Tobacco Control Program,” *Journal of the American Statistical Association*, 105, 493–505.
- Abadie, A., and Gardeazabal, J. (2003), “The Economic Costs of Conflict: A Case Study of the Basque Country,” *American Economic Review*, 93, 113–132.

- Agarwal, A., Shah, D., Shen, D., and Song, D. (2021), “On Robustness of Principal Component Regression,” *Journal of the American Statistical Association*, 116, 1731–1745.
- Ben-Michael, E., Feller, A., and Rothstein, J. (2022), “Synthetic Controls with Staggered Adoption,” *Journal of the Royal Statistical Society, Series B*, 84, 351–381.
- Bhagwati, J., and Srinivasan, T. N. (2001), “Outward-Orientation and Development: Are Revisionists Right?” in *Trade, Development and Political Economy*, Springer, pp. 3–26.
- Billmeier, A., and Nannicini, T. (2013), “Assessing Economic Liberalization Episodes: A Synthetic Control Approach,” *Review of Economics and Statistics*, 95, 983–1001.
- Boyd, S., and Vandenberghe, L. (2004), *Convex Optimization*, Cambridge University Press.
- Cattaneo, M. D., Feng, Y., Palomba, F., and Titiunik, R. (2023), “`scpi`: Uncertainty Quantification for Synthetic Control Methods,” arXiv:2202.05984.
- Cattaneo, M. D., Feng, Y., and Titiunik, R. (2021), “Prediction Intervals for Synthetic Control Methods,” *Journal of the American Statistical Association*, 116, 1865–1880.
- Chernozhukov, V., Wüthrich, K., and Zhu, Y. (2021a), “Distributional Conformal Prediction,” *Proceedings of the National Academy of Sciences*, 118, e2107794118.
- Chernozhukov, V., Wüthrich, K., and Zhu, Y. (2021b), “An Exact and Robust Conformal Inference Method for Counterfactual and Synthetic Controls,” *Journal of the American Statistical Association*, 116, 1849–1864.
- DeJong, D. N., and Ripoll, M. (2006), “Tariffs and Growth: An Empirical Exploration of Contingent Relationships,” *The Review of Economics and Statistics*, 88, 625–640.
- Giavazzi, F., and Tabellini, G. (2005), “Economic and Political Liberalizations,” *Journal of Monetary Economics*, 52, 1297–1330.
- Levine, R., and Renelt, D. (1992), “A Sensitivity Analysis of Cross-Country Growth Regressions,” *American Economic Review*, 82, 942–963.
- Li, K. T. (2020), “Statistical Inference for Average Treatment Effects Estimated by Synthetic Control Methods,” *Journal of the American Statistical Association*, 115, 2068–2083.

- Masini, R., and Medeiros, M. C. (2021), “Counterfactual Analysis with Artificial Controls: Inference, High Dimensions and Nonstationarity,” *Journal of the American Statistical Association*, 116, 1773–1788.
- Sachs, J. D., Warner, A., Åslund, A., and Fischer, S. (1995), “Economic Reform and the Process of Global Integration,” *Brookings Papers on Economic Activity*, 1995, 1–118.
- Shaikh, A. M., and Toulis, P. (2021), “Randomization Tests in Observational Studies with Staggered Adoption of Treatment,” *Journal of the American Statistical Association*, 116, 1835–1848.
- Shen, D., Ding, P., Sekhon, J., and Yu, B. (2022), “A Tale of Two Panel Data Regressions,” arXiv:2207.14481.
- Shi, X., Miao, W., Hu, M., and Tchetgen, E. T. (2023), “Theory for Identification and Inference with Synthetic Controls: A Proximal Causal Inference Framework,” arXiv:2108.13935.
- Vovk, V. (2012), “Conditional Validity of Inductive Conformal Predictors,” in *JMLR: Workshop and Conference Proceedings 25, Asian Conference on Machine Learning*, pp. 475–490.
- Wacziarg, R., and Welch, K. H. (2008), “Trade Liberalization and Growth: New Evidence,” *The World Bank Economic Review*, 22, 187–231.
- Wainwright, M. J. (2019), *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*, Cambridge University Press.

Supplement to “Uncertainty Quantification in Synthetic Controls with Staggered Treatment Adoption”

Matias D. Cattaneo* Yingjie Feng[†] Filippo Palomba[‡] Rocio Titiunik[§]

September 7, 2023

This supplement contains all proofs, additional results, other technical details about data preparation and estimation, and the code used to reproduce the results.

arXiv:2210.05026v3 [econ.EM] 5 Sep 2023

*Department of Operations Research and Financial Engineering, Princeton University.

[†]School of Economics and Management, Tsinghua University.

[‡]Department of Economics, Princeton University.

[§]Department of Politics, Princeton University.

Complete Contents

S.1 Other Strategies for Uncertainty Quantification	3
S.1.1 Out-of-Sample Error	3
S.1.2 Simultaneous Prediction Intervals	3
S.2 Proofs	4
S.2.1 Proof of Theorem 1	4
S.2.2 Verification of Condition (i) in Theorem 1	5
S.2.3 Proof of Lemma 1	6
S.3 Second Order Cone Programming	7
S.3.1 Families of convex optimization problems	8
S.3.2 Link between QCQP and SOCPs.	9
S.4 Data Preparation and Software Implementation	11
S.4.1 Data Description	11
S.4.2 Code	13
S.5 Results with Simplex Constraint	16
S.5.1 Africa	17
S.5.2 Asia	20
S.5.3 Europe	23
S.5.4 North America	27
S.5.5 South America	30
S.6 Results with L1-L2, Other Continents	32
S.6.1 Africa	33
S.6.2 Asia	36
S.6.3 North America	39
S.6.4 South America	42
S.7 Average treatment effect on countries liberalized in 1991	45
S.8 Excluded Countries	48

S.1 Other Strategies for Uncertainty Quantification

S.1.1 Out-of-Sample Error

In Section 4 we discuss the approach for quantifying the uncertainty of $e_t^{[i]}$ based on the non-asymptotic bounds. We briefly describe another two strategies below.

- *Location-scale model.* Suppose that $e_t^{[i]} = \mathbb{E}[e_t^{[i]}|\mathcal{H}] + (\mathbb{V}[e_t^{[i]}|\mathcal{H}])^{1/2}\varepsilon_t^{[i]}$ with $\varepsilon_t^{[i]}$ statistically independent of \mathcal{H} . The bounds on $e_t^{[i]}$ can now be set as $M_{2,L} = \mathbb{E}[e_t^{[i]}|\mathcal{H}] + (\mathbb{V}[e_t^{[i]}|\mathcal{H}])^{1/2}\mathfrak{c}_\varepsilon(\alpha_2/2)$ and $M_{2,U} = \mathbb{E}[e_t^{[i]}|\mathcal{H}] + (\mathbb{V}[e_t^{[i]}|\mathcal{H}])^{1/2}\mathfrak{c}_\varepsilon(1 - \alpha_2/2)$ where $\mathfrak{c}_\varepsilon(\alpha_2/2)$ and $\mathfrak{c}_\varepsilon(1 - \alpha_2/2)$ are $\alpha_2/2$ and $(1 - \alpha_2/2)$ quantiles of $\varepsilon_t^{[i]}$, respectively, and α_2 is the desired pre-specified level.
- *Quantile regression.* We can determine the $\alpha_2/2$ and $(1 - \alpha_2/2)$ conditional quantiles of $e_t^{[i]}|\mathcal{H}$. Consequently, another possibility is to employ quantile regression methods to estimate those quantities using pre-treatment data.

S.1.2 Simultaneous Prediction Intervals

Section 4.2 constructs prediction intervals with simultaneous coverage. We briefly describe two other common approaches below.

- *Bonferroni-type correction.* There has been a large literature on Bonferroni corrections that can be used to construct multiple prediction intervals with simultaneous coverage. For example, consider a simple correction strategy: for each $k = 0, \dots, L$, use any strategy described in Section 4 to construct a prediction interval for $e_{T_i+k}^{[i]}$ that has coverage probability at least $1 - (\alpha_2/(L+1))$. Then, the simultaneous coverage probability of the $L+1$ prediction intervals $\{\tilde{\mathcal{I}}_k : 0 \leq k \leq L\}$ is at least $1 - \alpha_2$. Some other more sophisticated corrections are also available in the literature (see, e.g., [Ravishanker et al., 1987](#)). For instance, the second-order Bonferroni-type bound implies that

$$\mathbb{P}[e_{T_i+k}^{[i]} \in \tilde{\mathcal{I}}_{T_i+k} \text{ for all } 0 \leq k \leq L \mid \mathcal{H}] \geq 1 - \sum_{k=0}^L p_k + \sum_{k=0}^{L-1} p_{k,k+1}, \quad \text{where}$$

$$p_k = \mathbb{P}(e_{T_i+k}^{[i]} \in \tilde{\mathcal{I}}_{T_i+k} \mid \mathcal{H}), \quad p_{k,k+1} = \mathbb{P}(e_{T_i+k}^{[i]} \in \tilde{\mathcal{I}}_{T_i+k}, e_{T_i+k+1}^{[i]} \in \tilde{\mathcal{I}}_{T_i+k+1} \mid \mathcal{H}).$$

Then, one can construct the prediction intervals $\tilde{\mathcal{I}}_k$ with corresponding coverage probabilities p_k and $p_{k,k+1}$ such that $1 - \sum_{k=0}^L p_k + \sum_{k=0}^{L-1} p_{k,k+1} \geq 1 - \alpha_2$. Such bounds are usually sharper, but their implementation requires the modelling of the dependence of $(e_t^{[i]}, e_{t+1}^{[i]})$ conditional on \mathcal{H} and is computationally more burdensome.

- *Scheffé-type intervals.* An alternative approach is to construct Scheffé-type simultaneous prediction intervals, though stronger distributional assumptions need to be made. For instance, assume that $(e_{T_i}^{[i]}, \dots, e_{T_i+L}^{[i]})'$ jointly follows a conditional Gaussian distribution with mean zero and variance $\Sigma_{\mathcal{H}}$. Then,

$$(e_{T_i}^{[i]}, \dots, e_{T_i+L}^{[i]}) \Sigma_{\mathcal{H}}^{-1} (e_{T_i}^{[i]}, \dots, e_{T_i+L}^{[i]})' \sim \chi_{L+1}^2,$$

where χ_{L+1}^2 is χ^2 distribution with $L+1$ degrees of freedom. The sequence of prediction intervals $\tilde{\mathcal{I}}_k = \left[-\sigma_{\mathcal{H},kk} \sqrt{\chi_{L+1}^2(1 - \alpha_2)}, \sigma_{\mathcal{H},kk} \sqrt{\chi_{L+1}^2(1 - \alpha_2)} \right]$ have the simultaneous coverage proba-

bility at least $1 - \alpha_2$, where $\sigma_{\mathcal{H},kk}^2$ is the k th diagonal element of $\Sigma_{\mathcal{H}}$ and $\chi_{L+1}^2(1 - \alpha_2)$ is the $(1 - \alpha_2)$ -quantile of χ^2 distribution with $L + 1$ degrees of freedom.

S.2 Proofs

S.2.1 Proof of Theorem 1

Proof. Let

$$\begin{aligned}\ell(\boldsymbol{\delta}) &= \boldsymbol{\delta}'\widehat{\mathbf{Q}}\boldsymbol{\delta} - 2\mathbf{G}'\boldsymbol{\delta}, \quad \text{and } \mathbf{G}|\mathcal{H} \sim \mathbf{N}(\mathbf{0}, \boldsymbol{\Sigma}) \\ \ell^*(\boldsymbol{\delta}) &= \boldsymbol{\delta}'\widehat{\mathbf{Q}}\boldsymbol{\delta} - 2(\mathbf{G}^*)'\boldsymbol{\delta}, \quad \text{and } \mathbf{G}^*|\text{Data} \sim \mathbf{N}(\mathbf{0}, \widehat{\boldsymbol{\Sigma}})\end{aligned}$$

Accordingly, define

$$\begin{aligned}\zeta_{\mathbb{U}}^* &= \sup \left\{ \mathbf{p}'_{\tau}\boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta^*, \|\boldsymbol{\delta}\| \leq \varpi_{\delta}^*, \ell^*(\boldsymbol{\delta}) \leq 0 \right\}, \\ \zeta_{\mathbb{U}}^{\text{int}} &= \sup \left\{ \mathbf{p}'_{\tau}\boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta, \|\boldsymbol{\delta}\| \leq \varpi_{\delta}^*, \ell^*(\boldsymbol{\delta}) \leq 0 \right\}, \\ \tilde{\zeta}_{\mathbb{U}} &= \sup \left\{ \mathbf{p}'_{\tau}\boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta, \|\boldsymbol{\delta}\| \leq \varpi_{\delta}^*, \ell(\boldsymbol{\delta}) \leq 0 \right\}.\end{aligned}$$

For any $\alpha_0 \in [0, 1]$, let $\tilde{\zeta}_{\mathbb{U}}^*(\alpha_0)$ be the α_0 -quantile of $\zeta_{\mathbb{U}}^*$ conditional on the data and $\tilde{\zeta}_{\mathbb{U}}(\alpha_0)$ be the α_0 -quantile of $\tilde{\zeta}_{\mathbb{U}}$ conditional on \mathcal{H} . Similarly, define

$$\begin{aligned}\zeta_{\mathbb{L}}^* &= \inf \left\{ \mathbf{p}'_{\tau}\boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta^*, \|\boldsymbol{\delta}\| \leq \varpi_{\delta}^*, \ell^*(\boldsymbol{\delta}) \leq 0 \right\}, \\ \zeta_{\mathbb{L}}^{\text{int}} &= \inf \left\{ \mathbf{p}'_{\tau}\boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta, \|\boldsymbol{\delta}\| \leq \varpi_{\delta}^*, \ell^*(\boldsymbol{\delta}) \leq 0 \right\}, \\ \tilde{\zeta}_{\mathbb{L}} &= \inf \left\{ \mathbf{p}'_{\tau}\boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta, \|\boldsymbol{\delta}\| \leq \varpi_{\delta}^*, \ell(\boldsymbol{\delta}) \leq 0 \right\}.\end{aligned}$$

Let $\tilde{\zeta}_{\mathbb{L}}^*(\alpha_0)$ be the α_0 -quantile of $\zeta_{\mathbb{L}}^*$ conditional on the data and $\tilde{\zeta}_{\mathbb{L}}(\alpha_0)$ be the α_0 -quantile of $\tilde{\zeta}_{\mathbb{L}}$ conditional on \mathcal{H} .

Let $\mathbb{P}_1 = \mathbf{N}(\mathbf{0}, \boldsymbol{\Sigma})$ and $\mathbb{P}_2 = \mathbf{N}(\mathbf{0}, \widehat{\boldsymbol{\Sigma}})$. By condition (iv), on an event with \mathbb{P} -probability at least $1 - \pi_{\gamma}^*$, with $\mathbb{P}(\cdot|\mathcal{H})$ -probability at least $1 - \epsilon_{\gamma,2}^*$, the Kullback-Leibler divergence $\mathbb{KL}(\mathbb{P}_1, \mathbb{P}_2) \leq 2(\epsilon_{\gamma,1}^*)^2$, and by Pinsker's inequality, this implies that for any $\kappa' \leq \kappa$,

$$|\mathbb{P}^*(\tilde{\zeta}_{\mathbb{U}}^{\text{int}} \leq \kappa) - \mathbb{P}^*(\tilde{\zeta}_{\mathbb{U}} \leq \kappa)| \leq \epsilon_{\gamma,1}^* \quad \text{and} \quad |\mathbb{P}^*(\tilde{\zeta}_{\mathbb{L}}^{\text{int}} \geq \kappa') - \mathbb{P}^*(\tilde{\zeta}_{\mathbb{L}} \geq \kappa')| \leq \epsilon_{\gamma,1}^*.$$

On the other hand, note that by condition (iii), on an event with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_{\Delta}^*$, with $\mathbb{P}(\cdot|\mathcal{H})$ -probability at least $1 - \epsilon_{\Delta}^*$, the event $\{\tilde{\zeta}_{\mathbb{U}}^* \leq \kappa, \tilde{\zeta}_{\mathbb{L}}^* \geq \kappa'\}$ implies that

$$\begin{aligned}\sup \left\{ \mathbf{p}'_{\tau}\boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta, \|\boldsymbol{\delta}\| \leq \varpi_{\delta}^*, \ell^*(\boldsymbol{\delta}) \leq 0 \right\} &\leq \kappa + \|\mathbf{p}'_{\tau}\|_* \sup_{\mathbf{a} \in \Delta \cap \mathcal{B}(0, \varpi_{\delta}^*)} \text{dist}(\mathbf{a}, \Delta^*) \\ &\leq \kappa + \|\mathbf{p}'_{\tau}\|_* \varpi_{\Delta}^* =: \kappa + \varepsilon_{\Delta}\end{aligned}$$

and

$$\begin{aligned}\inf \left\{ \mathbf{p}'_{\tau}\boldsymbol{\delta} : \boldsymbol{\delta} \in \Delta, \|\boldsymbol{\delta}\| \leq \varpi_{\delta}^*, \ell^*(\boldsymbol{\delta}) \leq 0 \right\} &\geq \kappa' - \|\mathbf{p}'_{\tau}\|_* \sup_{\mathbf{a} \in \Delta \cap \mathcal{B}(0, \varpi_{\delta}^*)} \text{dist}(\mathbf{a}, \Delta^*) \\ &\geq \kappa' - \varepsilon_{\Delta}.\end{aligned}$$

Therefore,

$$\mathbb{P}^*(\tilde{\zeta}_{\mathbb{U}}^* \leq \kappa) \leq \mathbb{P}^*(\tilde{\zeta}_{\mathbb{U}}^{\text{int}} \leq \kappa + \varepsilon_{\Delta}) \quad \text{and} \quad \mathbb{P}^*(\tilde{\zeta}_{\mathbb{L}}^* \geq \kappa') \geq \mathbb{P}^*(\tilde{\zeta}_{\mathbb{L}}^{\text{int}} \geq \kappa' - \varepsilon_{\Delta}).$$

Then, on an event with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_{\gamma}^* - \pi_{\Delta}^*$, with $\mathbb{P}(\cdot|\mathcal{H})$ -probability at least $1 - \epsilon_{\gamma,2}^* - \epsilon_{\Delta}^*$, we have

$$\begin{aligned} 1 - \alpha_1/2 &\leq \mathbb{P}^*\left(\tilde{\zeta}_{\mathbb{U}}^* \leq \mathbf{c}^*(1 - \alpha_1/2)\right) \leq \mathbb{P}^*\left(\tilde{\zeta}_{\mathbb{U}} \leq \mathbf{c}^*(1 - \alpha_1/2) + \varepsilon_{\Delta}\right) + \epsilon_{\gamma,1}^* \quad \text{and} \\ 1 - \alpha_1/2 &\leq \mathbb{P}^*\left(\tilde{\zeta}_{\mathbb{L}}^* \geq \mathbf{c}^*(\alpha_1/2)\right) \leq \mathbb{P}^*\left(\tilde{\zeta}_{\mathbb{L}} \geq \mathbf{c}^*(\alpha_1/2) - \varepsilon_{\Delta}\right) + \epsilon_{\gamma,1}^*. \end{aligned}$$

Also, by condition (ii), we have with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_{\delta}^*$,

$$\tilde{\mathbf{c}}_{\mathbb{U}}(1 - \alpha_1/2 - \epsilon_{\gamma,1}^*) \geq \mathbf{c}_{\mathbb{U}}(1 - \alpha_1/2 - \epsilon_{\gamma,1}^* - \epsilon_{\delta}^*) \quad \text{and} \quad \tilde{\mathbf{c}}_{\mathbb{L}}(\alpha_1/2 + \epsilon_{\gamma,1}^*) \leq \mathbf{c}_{\mathbb{L}}(\alpha_1/2 + \epsilon_{\gamma,1}^* + \epsilon_{\delta}^*).$$

Using conditions (i) and (ii), we have with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_{\gamma} - \pi_{\gamma}^* - \pi_{\Delta}^* - \pi_{\delta}^*$,

$$\begin{aligned} &\mathbb{P}\left(\mathbf{c}_{\mathbb{L}}^*(\alpha_1/2) - \varepsilon_{\Delta} \leq \mathbf{p}'_{\tau}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq \mathbf{c}_{\mathbb{U}}^*(1 - \alpha_1/2) + \varepsilon_{\Delta} \mid \mathcal{H}\right) \\ &\geq \mathbb{P}\left(\tilde{\mathbf{c}}_{\mathbb{L}}(\alpha_1/2 + \epsilon_{\gamma,1}^*) \leq \mathbf{p}'_{\tau}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq \tilde{\mathbf{c}}_{\mathbb{U}}(1 - \alpha_1/2 - \epsilon_{\gamma,1}^*) \mid \mathcal{H}\right) - \epsilon_{\gamma,2}^* - \epsilon_{\Delta}^* \\ &\geq \mathbb{P}\left(\mathbf{c}_{\mathbb{L}}(\alpha_1/2 + \epsilon_{\gamma,1}^* + \epsilon_{\delta}^*) \leq \mathbf{p}'_{\tau}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq \mathbf{c}_{\mathbb{U}}(1 - \alpha_1/2 - \epsilon_{\gamma,1}^* - \epsilon_{\delta}^*) \mid \mathcal{H}\right) - \epsilon_{\gamma,2}^* - \epsilon_{\Delta}^* \\ &\geq 1 - \alpha_1 - 2\epsilon_{\gamma,1}^* - 2\epsilon_{\delta}^* - \epsilon_{\gamma} - \epsilon_{\gamma,2}^* - \epsilon_{\Delta}^*. \end{aligned}$$

Finally, by condition (v), we immediately have $\mathbb{P}(M_{2,\mathbb{L}} \leq e_{\tau} \leq M_{2,\mathbb{U}}) \geq 1 - \alpha_2$. Then the proof is complete. \square

S.2.2 Verification of Condition (i) in Theorem 1

As explained in the main paper, by convexity of the constraint set $\mathcal{W} \times \mathcal{R}$ and the optimality of $\hat{\boldsymbol{\beta}}$,

$$\inf_{\boldsymbol{\delta} \in \mathcal{M}_{\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}}} \mathbf{p}'_{\tau} \boldsymbol{\delta} \leq \mathbf{p}'_{\tau}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq \sup_{\boldsymbol{\delta} \in \mathcal{M}_{\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}}} \mathbf{p}'_{\tau} \boldsymbol{\delta},$$

where $\mathcal{M}_{\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}} = \{\boldsymbol{\delta} \in \Delta : \boldsymbol{\delta}' \hat{\mathbf{Q}} \boldsymbol{\delta} - 2(\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})' \boldsymbol{\delta}\}$. Thus, condition (i) in Theorem 1 indeed requires that $\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}$ can be approximated by a Gaussian vector \mathbf{G} . In general, this can be shown if we assume $(u_{t,1}^{[1]}, \dots, u_{t,M}^{[1]}, \dots, u_{t,1}^{[N_1]}, u_{t,M}^{[N_1]})'$ is independent or weakly dependent conditional on \mathcal{H} . In the following we verify condition (i) by imposing a conditional independence assumption. The extension that allows for weakly dependent errors can be established using the idea of Theorem A in [Cattaneo et al. \(2021\)](#).

Lemma S.1. *Assume \mathcal{W} and \mathcal{R} are convex, $\hat{\boldsymbol{\beta}}$ in Equation (3.1) and $\boldsymbol{\beta}_0$ in Equation (4.2) exist, and $\mathcal{H} = \sigma(\mathbf{B}, \mathbf{C}, \mathbf{p}_{\tau})$. In addition, for some finite nonnegative constants, the following conditions hold:*

(i) $\mathbf{u}_t = (u_{t,1}^{[1]}, \dots, u_{t,M}^{[1]}, \dots, u_{t,1}^{[N_1]}, \dots, u_{t,M}^{[N_1]})$ is independent over t conditional on \mathcal{H} ;

(ii) $\mathbb{P}(\sum_{t=1}^{T_0} \mathbb{E}[\|\sum_{j=1}^{N_1} \sum_{l=1}^M \tilde{\mathbf{z}}_{t,l}^{[j]}(u_{t,l}^{[j]} - \mathbb{E}[u_{t,l}^{[j]}|\mathcal{H}])\|^3 | \mathcal{H}] \geq \epsilon_{\gamma} (84(d^{1/4} + 16))^{-1}) \geq 1 - \pi_{\gamma}$ where $\tilde{\mathbf{z}}_{t,l}^{[j]}$ is the $((j-1)T_0M + (l-1)T_0 + t)$ th column of $\boldsymbol{\Sigma}^{-1/2} \mathbf{Z}'$.

Then, with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_\gamma$,

$$\mathbb{P}\left(\mathbf{c}_L(\alpha_0) \leq \mathbf{p}'_\tau(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq \mathbf{c}_U(1 - \alpha_0) \mid \mathcal{H}\right) \geq 1 - \alpha_0 - \epsilon_\gamma.$$

Proof. Define $\mathcal{M}_\xi = \{\boldsymbol{\delta} \in \Delta : \boldsymbol{\delta}'\widehat{\mathbf{Q}}\boldsymbol{\delta} - 2\xi'\boldsymbol{\delta}\}$. Fix $\widehat{\mathbf{Q}}$ and \mathbf{p}_τ . By Lemma 2 of Cattaneo et al. (2021), for any κ , $\mathcal{A}_\kappa := \{\boldsymbol{\xi} \in \mathbb{R}^d : \sup_{\boldsymbol{\delta} \in \mathcal{M}_\xi} \mathbf{p}'_\tau \boldsymbol{\delta} \leq \kappa\}$ and $\mathcal{A}'_\kappa = \{\boldsymbol{\xi} \in \mathbb{R}^d : \inf_{\boldsymbol{\delta} \in \mathcal{M}_\xi} \mathbf{p}'_\tau \boldsymbol{\delta} \geq \kappa\}$ are convex. By Berry-Esseen Theorem for convex sets Raić (2019),

$$|\mathbb{P}(\widehat{\boldsymbol{\gamma}} - \boldsymbol{\gamma} \in \mathcal{A}_\kappa \mid \mathcal{H}) - \mathbb{P}(\mathbf{G} \in \mathcal{A}_\kappa \mid \mathcal{H})| \leq 42(d^{1/4} + 16) \sum_{t=1}^{T_0} \mathbb{E} \left[\left\| \sum_{j=1}^{N_1} \sum_{l=1}^M \tilde{\mathbf{z}}_{t,l}^{[j]} \tilde{u}_{t,l}^{[j]} \right\|^3 \mid \mathcal{H} \right],$$

where $\tilde{u}_{t,l}^{[j]} = u_{t,l}^{[j]} - \mathbb{E}[u_{t,l}^{[j]} \mid \mathcal{H}]$. By condition (ii), with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_\gamma$,

$$|\mathbb{P}(\widehat{\boldsymbol{\gamma}} - \boldsymbol{\gamma} \in \mathcal{A}_\kappa \mid \mathcal{H}) - \mathbb{P}(\mathbf{G} \in \mathcal{A}_\kappa \mid \mathcal{H})| \leq \epsilon_\gamma/2.$$

Then, for any κ , with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_\gamma$,

$$\mathbb{P}(\mathbf{p}'_\tau(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq \kappa \mid \mathcal{H}) \geq \mathbb{P}(\widehat{\boldsymbol{\gamma}} - \boldsymbol{\gamma} \in \mathcal{A}_\kappa \mid \mathcal{H}) \geq \mathbb{P}(\mathbf{G} \in \mathcal{A}_\kappa \mid \mathcal{H}) - \epsilon_\gamma/2.$$

Similarly, we can show for any κ ,

$$\mathbb{P}(\mathbf{p}'_\tau(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \geq \kappa \mid \mathcal{H}) \geq \mathbb{P}(\widehat{\boldsymbol{\gamma}} - \boldsymbol{\gamma} \in \mathcal{A}'_\kappa \mid \mathcal{H}) \geq \mathbb{P}(\mathbf{G} \in \mathcal{A}'_\kappa \mid \mathcal{H}) - \epsilon_\gamma/2.$$

Therefore, with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_\gamma$,

$$\mathbb{P}\left(\mathbf{c}_L(\alpha_0) \leq \mathbf{p}'_\tau(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) \leq \mathbf{c}_U(1 - \alpha_0) \mid \mathcal{H}\right) \geq 1 - \alpha_0 - \epsilon_\gamma.$$

Then the proof is complete. \square

S.2.3 Proof of Lemma 1

Proof. In this proof, the constant $\mathfrak{C} > 0$ is a generic constant that is independent of T_0 and may be different in different uses. Note that $\mathbf{m}_{\text{eq}}(\boldsymbol{\beta}_0) = \mathbf{0}$ and $\mathbf{m}_{\text{in}}(\boldsymbol{\beta}_0) \leq \mathbf{0}$. If the j th inequality constraint is binding, i.e., $m_{\text{in},j}(\boldsymbol{\beta}_0) = 0$, then $m_{\text{in},j}(\widehat{\boldsymbol{\beta}}) = \frac{\partial}{\partial \boldsymbol{\beta}'} m_{\text{in},j}(\tilde{\boldsymbol{\beta}})(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0)$ for some $\tilde{\boldsymbol{\beta}}$ between $\boldsymbol{\beta}_0$ and $\widehat{\boldsymbol{\beta}}$. Then, for $\mathfrak{c} := \max_{1 \leq j \leq d_{\text{in}}} \sup_{\boldsymbol{\beta} \in \mathcal{B}(\boldsymbol{\beta}_0, \varpi_\delta^*)} \left\| \frac{\partial}{\partial \boldsymbol{\beta}'} m_{\text{in},j}(\boldsymbol{\beta}) \right\|_*$, we have $\max_{1 \leq j \leq d_{\text{in}}} |m_{\text{in},j}(\widehat{\boldsymbol{\beta}})| \leq \mathfrak{c} \varpi_\delta^*$ with $\mathbb{P}(\cdot \mid \mathcal{H})$ -probability at least $1 - \epsilon_\Delta^*$, on an event with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_\Delta^*$. By the condition imposed on the tuning parameter ϱ_{js} , on an event with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_\Delta^*$, with $\mathbb{P}(\cdot \mid \mathcal{H})$ -probability at least $1 - \epsilon_\Delta^*$, \mathcal{A} coincides with the set of indices for the binding inequality constraints. Then, for $\varepsilon \leq \varpi_\delta^*$,

$$\Delta^* \cap \mathcal{B}(0, \varepsilon) = \left\{ \boldsymbol{\beta} - \widehat{\boldsymbol{\beta}} \in \mathcal{B}(0, \varepsilon) : \mathbf{m}_{\text{eq}}(\boldsymbol{\beta}) = \mathbf{0}, \mathbf{m}_{\text{in},j}(\boldsymbol{\beta}) \leq \mathbf{m}_{\text{in},j}(\widehat{\boldsymbol{\beta}}) \text{ for } j \in \mathcal{A} \right\}.$$

Without loss of generality, we assume $\mathbf{m}_{\text{in}}(\boldsymbol{\beta}_0) = \mathbf{0}$ from now on. Otherwise, the non-binding constraints can be dropped and the proof can proceed the same way described below.

Define $\Gamma_{\text{eq}}(\boldsymbol{\beta}) = \frac{\partial}{\partial \boldsymbol{\beta}'} \mathbf{m}_{\text{eq}}(\boldsymbol{\beta})$ and $\Gamma_{\text{in}}(\boldsymbol{\beta}) = \frac{\partial}{\partial \boldsymbol{\beta}'} \mathbf{m}_{\text{in}}(\boldsymbol{\beta})$. Let

$$\Gamma(\boldsymbol{\beta}) = \left(\Gamma'_{\text{eq}}(\boldsymbol{\beta}), \Gamma'_{\text{in}}(\boldsymbol{\beta}), \Gamma'_c(\boldsymbol{\beta}_0) \right)', \quad \Gamma^0 = \Gamma(\boldsymbol{\beta}_0), \quad \Gamma^* = \Gamma(\widehat{\boldsymbol{\beta}}),$$

where $\Gamma_c(\beta_0)$ is chosen such that $\Gamma(\beta_0)$ is non-degenerate. By conditions (i) and (ii) imposed in the lemma, $\|\Gamma^0 - \Gamma^*\| \leq \mathfrak{C}\|\widehat{\beta} - \beta_0\|$ with $\mathbb{P}(\cdot|\mathcal{H})$ -probability at least $1 - \epsilon_\Delta^*$, on an event with \mathbb{P} -probability at least $1 - \pi_\Delta^*$.

Let

$$\mathbf{m}^+(\cdot) = \left(\mathbf{m}'_{\text{eq}}(\cdot), \mathbf{m}'_{\text{in}}(\cdot), (\cdot - \beta_0)' \times \Gamma_c(\beta_0)' \right)'.$$

Then, $\mathbf{m}^+(\beta_0) = \mathbf{0}$. For each β in the neighborhood around β_0 such that $\beta - \beta_0 \in \Delta \cap \mathcal{B}(0, \varepsilon)$, define

$$\lambda^0 = (\Gamma^0)^{-1} \left(\mathbf{m}^+(\beta) - \mathbf{m}^+(\beta_0) \right).$$

Thus, $\Gamma_{\text{eq}}^0 \lambda^0 = \mathbf{0}$, $\Gamma_{\text{in}}^0 \lambda^0 \leq \mathbf{0}$. Note that by Taylor's expansion,

$$\|\lambda^0 - (\Gamma^0)^{-1} \Gamma^0 (\beta - \beta_0)\| \leq \mathfrak{C} \|\beta - \beta_0\|^2,$$

implying that $\|\lambda^0 - (\beta - \beta_0)\| \leq \mathfrak{C}(\varpi_\delta^*)^2$ with $\mathbb{P}(\cdot|\mathcal{H})$ -probability at least $1 - \epsilon_\Delta^*$, on an event with \mathbb{P} -probability at least $1 - \pi_\Delta^*$.

Next, define $\tilde{\mathbf{m}}(\cdot) = \mathbf{m}^+(\widehat{\beta} + \cdot) - \mathbf{m}^+(\widehat{\beta})$ and $\tilde{\beta} = \phi^* + \widehat{\beta}$ for ϕ^* defined below:

$$\phi^* := \tilde{\mathbf{m}}^{-1} \left(\Gamma^* (\lambda^0 - (\Gamma^*)^{-1} (\Gamma^* - \Gamma^0) \lambda^0) \right).$$

By Taylor's expansion,

$$\begin{aligned} \phi^* &= \tilde{\mathbf{m}}^{-1}(\mathbf{0}) + \left[\frac{\partial}{\partial \phi'} \tilde{\mathbf{m}}(\mathbf{0}) \right]^{-1} \Gamma^* \left(\lambda^0 - (\Gamma^*)^{-1} (\Gamma^* - \Gamma^0) \lambda^0 \right) + \mathfrak{R}\mathfrak{e} \\ &= \lambda^0 + (\Gamma^*)^{-1} (\Gamma^* - \Gamma^0) \lambda^0 + \mathfrak{R}\mathfrak{e}, \end{aligned}$$

where $\|(\Gamma^*)^{-1} (\Gamma^* - \Gamma^0) \lambda^0 + \mathfrak{R}\mathfrak{e}\| \leq \mathfrak{C} \|\lambda^0\|^2$ with $\mathbb{P}(\cdot|\mathcal{H})$ -probability at least $1 - \epsilon_\Delta^*$, on an event with \mathbb{P} -probability over \mathcal{H} at least $1 - \pi_\Delta^*$. That is, we actually find $\tilde{\beta}$ such that $\|(\tilde{\beta} - \widehat{\beta}) - \lambda^0\| \leq \mathfrak{C}(\varpi_\delta^*)^2$. Note that

$$\mathbf{m}^+(\widehat{\beta} + \phi^*) - \mathbf{m}^+(\widehat{\beta}) = \tilde{\mathbf{m}}(\phi^*) = \tilde{\mathbf{m}} \left(\tilde{\mathbf{m}}^{-1} \left(\Gamma^* (\lambda^0 - (\Gamma^*)^{-1} (\Gamma^* - \Gamma^0) \lambda^0) \right) \right).$$

Thus, $\mathbf{m}_{\text{eq}}(\tilde{\beta}) = \mathbf{0}$ and $\mathbf{m}_{\text{in}}(\tilde{\beta}) \leq \mathbf{m}_{\text{in}}(\widehat{\beta})$.

In the above, we make use of the fact that Γ^* is non-degenerate, i.e., its smallest eigenvalue is bounded away from zero. Note that by assumptions on the constraints, it is feasible to construct Γ_c such that Γ^0 is non-degenerate (with high probability over \mathcal{H}). Then, by Weyl's inequality,

$$s_{\min}(\Gamma^* (\Gamma^*)') \geq s_{\min}(\Gamma^0 (\Gamma^0)') - \mathfrak{C} \|\widehat{\beta} - \beta_0\|,$$

implying $s_{\min}(\Gamma^*) \geq s_{\min}(\Gamma^0) - \mathfrak{C} \|\widehat{\beta} - \beta_0\| > 0$ with $\mathbb{P}(\cdot|\mathcal{H})$ -probability at least $1 - \epsilon_\Delta^*$, on an event with \mathbb{P} -probability at least $1 - \pi_\Delta^*$, where $s_{\min}(\cdot)$ here denotes the smallest eigenvalue of the symmetric matrix inside. Then, the proof is complete. \square

S.3 Second Order Cone Programming

For the sake of completeness, at the beginning of this section we define again the three types of convex optimization problems presented in the main text in Section 5.1. Second, we illustrate the link between these three families of convex problems. Third, we note that the optimization problems

underlying the prediction/estimation and uncertainty quantification problems for SC presented in Section 3 are QCQPs and QCLPs, respectively, and show how to represent them as SOCPs. Finally, we show how to write the Lasso-type constraint in conic form.

S.3.1 Families of convex optimization problems

QCQPs and QCLPs. A quadratically constrained quadratic program is an optimization problem of with the following form

$$\begin{aligned} \min_{\mathbf{x}} \quad & \mathbf{x}'\mathbf{P}_0\mathbf{x} + \mathbf{q}'_0\mathbf{x} + w && \text{(S.3.1)} \\ \text{subject to} \quad & \mathbf{x}'\mathbf{P}_j\mathbf{x} + \mathbf{q}'_j\mathbf{x} + r_j \leq 0, \quad j = 1, \dots, m && \text{(Quadratic inequality constraint)} \\ & \mathbf{F}\mathbf{x} = \mathbf{g}, && \text{(Linear equality constraint)} \end{aligned}$$

where $\mathbf{P}_0, \mathbf{P}_1, \dots, \mathbf{P}_m \in \mathcal{M}_{n \times n}(\mathbb{R})$, $\mathbf{q}_0, \mathbf{q}_1, \dots, \mathbf{q}_m \in \mathbb{R}^n$, $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{F} \in \mathcal{M}_{m \times n}(\mathbb{R})$, $\mathbf{g} \in \mathbb{R}^m$, and $r_0, r_1, \dots, r_m, w \in \mathbb{R}$. If all the matrices $\mathbf{P}_0, \mathbf{P}_1, \dots, \mathbf{P}_m$ are positive semi-definite the QCQP is convex. Moreover, if $\mathbf{P}_0 = \mathbf{0}$ the QCQP becomes a QCLP. For this reason, in what follows we will restrict our attention to QCQPs as they naturally embed QCLPs.

SOCPs. To define a SOCP, it is necessary to first give the definition of a second-order cone and then introduce the notion of associated *generalized inequality*.

Second-order cone definition. A set \mathcal{C} is called a *cone* if for every $\mathbf{x} \in \mathcal{C}$ and $\alpha \geq 0$ we have $\alpha\mathbf{x} \in \mathcal{C}$. A set \mathcal{C} is a *convex cone* if it is convex and a cone, i.e. if $\forall \mathbf{x}_1, \mathbf{x}_2 \in \mathcal{C}$, and $\forall \alpha_1, \alpha_2 \geq 0$, we have

$$\alpha_1\mathbf{x}_1 + \alpha_2\mathbf{x}_2 \in \mathcal{C}.$$

Now, consider any norm $\|\cdot\|$ defined on the Euclidean space \mathbb{R}^n . The *norm cone* associated with the norm $\|\cdot\|$ is defined to be the set

$$\mathcal{C} = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} : \|\mathbf{x}\| \leq t\}$$

and it is a convex cone by the standard properties of the norms. A *second-order cone* is the associated norm cone for the Euclidean norm and it is typically defined as

$$\mathcal{C} = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} : \|\mathbf{x}\|_2 \leq t\} = \left\{ \begin{bmatrix} \mathbf{x} \\ t \end{bmatrix} : \begin{bmatrix} \mathbf{x} \\ t \end{bmatrix}' \begin{bmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{0} & -1 \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ t \end{bmatrix} \leq 0, t \geq 0 \right\}$$

Generalized inequality. A cone \mathcal{C} is *solid* if it has non-empty interior and it is **pointed** if $\mathbf{x} \in \mathcal{C}$, $-\mathbf{x} \in \mathcal{C}$ implies that $\mathbf{x} = \mathbf{0}$. We say that a cone \mathcal{C} is *proper* if it is convex, closed, solid, and pointed. Proper cones in the Euclidean space \mathbb{R}^n are useful because they induce a partial ordering which enjoys almost all the properties of the basic one in \mathbb{R} . Therefore, given a cone $\mathcal{C} \subseteq \mathbb{R}^n$, we can define the generalized inequality $\preceq_{\mathcal{C}}$ for any two vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$

$$\mathbf{x} \preceq_{\mathcal{C}} \mathbf{y} \iff \mathbf{y} - \mathbf{x} \in \mathcal{C}.$$

From this definition we can see that quadratic constraints such as $\|\mathbf{x}\|_2 \leq h$ can be re-written as second-order cone constraints of the form $\mathbf{x} \preceq_{\mathcal{C}} h$ for some second-order cone \mathcal{C} . Note that if $\mathcal{C} = \mathbb{R}_+^m$ then if $m = 1$, $\preceq_{\mathcal{C}}$ is the standard inequality \leq in \mathbb{R} , whereas if $m > 1$, $\preceq_{\mathcal{C}}$ is the component-wise inequality in \mathbb{R}^m .

Second-order cone program. Let \mathcal{K} be a cone such that $\mathcal{K} = \mathbb{R}_+^m \times \mathcal{K}_1 \times \mathcal{K}_2 \times \dots \times \mathcal{K}_L$ where

$\mathcal{K}_l := \{(k_0, \mathbf{k}_1) \in \mathbb{R} \times \mathbb{R}^l : \|\mathbf{k}_1\|_2 \leq k_0\}, l = 1, \dots, L$. Let $\preceq_{\mathcal{K}}$ be the *generalized inequality* associated with the cone \mathcal{K} . An optimization problem is called *second-order cone program* if it has the following form

$$\begin{aligned} \min_{\mathbf{x}} \quad & \mathbf{c}'\mathbf{x}, & (\text{S.3.2}) \\ \text{subject to} \quad & \mathbf{G}\mathbf{x} \preceq_{\mathcal{K}} \mathbf{h}, & (\text{Second-order cone constraint}) \\ & \mathbf{A}\mathbf{x} = \mathbf{b}. & (\text{Linear equality constraint}) \end{aligned}$$

S.3.2 Link between QCQP and SOCPs.

Any QCQP can be converted to a SOCP (Boyd and Vandenberghe, 2004). In other words, we can always rewrite an optimization problem such as (S.3.1) in the form of (S.3.2). First, we present the general result and then we explain all the necessary steps to reformulate QCQPs as SOCPs. Without loss of generality, assume that $w = 0$ in (S.3.1) and, to ease notation, let $m = 1$ so that there is only a single quadratic inequality constraint. Moreover, given any positive semi-definite matrix \mathbf{P} , let $\mathbf{P}^{1/2}$ be the square root of \mathbf{P} , that is the unique symmetric positive semi-definite matrix \mathbf{R} such that $\mathbf{R}\mathbf{R} = \mathbf{R}'\mathbf{R} = \mathbf{P}$. Then for any QCQP the following two formulations are equivalent

<u>QCQP</u>	<u>SOCP</u>
$\min_{\mathbf{x}} \quad \mathbf{x}'\mathbf{P}_0\mathbf{x} + \mathbf{q}'_0\mathbf{x}$	$\min_{\mathbf{x}, v, t, s} \quad v + \mathbf{q}'_0\mathbf{x}$
subject to $\mathbf{F}\mathbf{x} = \mathbf{g},$	subject to $\mathbf{F}\mathbf{x} = \mathbf{g},$
$\mathbf{x}'\mathbf{P}_1\mathbf{x} + \mathbf{q}'_1\mathbf{x} + r_1 \leq 0.$	$t + \mathbf{q}'_1\mathbf{x} + r_1 \preceq_{\mathbb{R}_+} 0,$
	$\mathbf{P}_0^{1/2}\mathbf{x} \preceq_{K_{1+n}} v,$
	$\mathbf{P}_1^{1/2}\mathbf{x} \preceq_{K_{1+n}} s.$

We can see that the logic beneath the conversion of a QCQP into a SOCP is to “linearize” all the non-linear terms appearing either in the objective function or in the inequality constraints. The “linearization” step does come at a cost, as it requires the introduction of a slack variable every time we rely on it. Indeed, above we linearized the objective function and the quadratic inequality constraint by introducing two auxiliary slack variables.

More formally, let $\mathbf{x}'\mathbf{P}\mathbf{x}$ be a symmetric positive semi-definite quadratic form and consider the constraint $\mathbf{x}'\mathbf{P}\mathbf{x} \leq y$. Then

- (i) Since \mathbf{P} is symmetric positive semi-definite the epigraph $\mathbf{x}'\mathbf{P}\mathbf{x} \leq y$ is a convex set and $\mathbf{P}^{1/2}$ is well-defined.
- (ii) Write the inequality constraint as a constraint involving the Euclidean norm $\|\cdot\|_2$

$$y \geq \mathbf{x}'\mathbf{P}\mathbf{x} = \mathbf{x}'\mathbf{P}^{1/2}\mathbf{P}^{1/2}\mathbf{x} = \|\mathbf{P}^{1/2}\mathbf{x}\|_2^2.$$

- (iii) Note that

$$\|\mathbf{P}^{1/2}\mathbf{x}\|_2^2 \leq y \iff \left\| \begin{bmatrix} 1 - y \\ 2\mathbf{P}^{1/2}\mathbf{x} \end{bmatrix} \right\|_2 \leq 1 + y, \quad (\text{S.3.3})$$

which can be verified by squaring the two sides of the last inequality and expand the norm.

- (iv) More is true, as the right-most inequality in (S.3.3) defines the following second-order cone for given $\mathbf{P}^{1/2}$

$$\mathcal{C} = \left\{ \left(1 - y, 2\mathbf{P}^{1/2}\mathbf{x}, 1 + y \right) : \left\| \begin{bmatrix} 1 - y \\ 2\mathbf{P}^{1/2}\mathbf{x} \end{bmatrix} \right\|_2 \leq 1 + y \right\},$$

which in turn induces the generalized inequality $\mathbf{P}^{1/2}\mathbf{x} \preceq_{\mathcal{C}} y$.

SC problem with lasso-type \mathcal{W} as a SOCP. We show how to write the QCQP as a SOCP when \mathcal{W} has a lasso-type constraint. In this case the SC weight construction (3.1) has the form:

$$\begin{aligned} \min_{\mathbf{w}, \mathbf{r}} \quad & (\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r})' \mathbf{V} (\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r}) & (\text{S.3.4}) \\ \text{subject to} \quad & \|\mathbf{w}^{[i]}\|_1 \leq Q_1^{[i]}, \quad i = 1, \dots, N_1 & (\text{L1 inequality constraints}) \end{aligned}$$

We can write the optimization problem in (S.3.4) as a SOCP of the following form

$$\begin{aligned} \min_{\mathbf{w}, \mathbf{r}, \{\mathbf{z}_i\}_{i=1}^{N_1}, v} \quad & v \\ \text{subject to} \quad & \begin{bmatrix} 1 - v \\ 2\mathbf{V}^{1/2}(\mathbf{A} - \mathbf{B}\mathbf{w} - \mathbf{C}\mathbf{r}) \end{bmatrix} \preceq_{\mathcal{C}_1} 1 + v, & (\text{cone in } \mathbb{R}^{2+T_0 \cdot M \cdot N_1}) \\ & \mathbf{1}' \mathbf{z}_i \preceq_{\mathcal{C}_2} Q_1^{[i]}, \quad i = 1, \dots, N_1 & (N_1 \text{ cones in } \mathbb{R}) \\ & -\mathbf{w} \preceq_{\mathcal{C}_3} \mathbf{z}, & (\text{cone in } \mathbb{R}^{J \cdot N_1}) \\ & \mathbf{w} \preceq_{\mathcal{C}_4} \mathbf{z}, & (\text{cone in } \mathbb{R}^{J \cdot N_1}) \end{aligned}$$

where $\mathcal{K} = \mathcal{C}_1 \times \mathcal{C}_2^{N_1} \times \mathcal{C}_3 \times \mathcal{C}_4 = \mathcal{K}_{1+T_0 \cdot M \cdot N_1} \times \mathbb{R}_+^{N_1} \times \mathbb{R}_+^{J \cdot N_1} \times \mathbb{R}_+^{J \cdot N_1}$ is the conic constraint for this program and $\mathbf{z} := (\mathbf{z}'_1, \dots, \mathbf{z}'_{N_1})'$.

For uncertainty quantification, we need to solve the optimization problem underlying (4.4). Here we discuss the lower bound only for brevity. Recalling that $\boldsymbol{\beta} = (\mathbf{w}', \mathbf{r}')'$, we have

$$\begin{aligned} \inf_{\boldsymbol{\beta}=(\mathbf{w}', \mathbf{r}')'} \quad & \mathbf{p}'_{\tau} (\boldsymbol{\beta} - \widehat{\boldsymbol{\beta}}) & (\text{S.3.5}) \\ \text{subject to} \quad & \|\mathbf{w}^{[i]}\|_1 \leq Q_1^{[i]} + \varrho^{[i]}, \quad i = 1, \dots, N_1 & (\text{L1 inequality constraints}) \\ & (\boldsymbol{\beta} - \widehat{\boldsymbol{\beta}})' \widehat{\mathbf{Q}} (\boldsymbol{\beta} - \widehat{\boldsymbol{\beta}}) - 2(\mathbf{G}^*)' (\boldsymbol{\beta} - \widehat{\boldsymbol{\beta}}) \leq 0, & (\text{constrained least squares}) \end{aligned}$$

where the scalars $\varrho_1^{[i]}, i = 1, \dots, N_1$ are regularization parameters used to relax Δ to Δ^* .

We can cast the SC optimization problem in (S.3.5) in conic form as follows:

$$\begin{aligned} \min_{\mathbf{w}, \mathbf{r}, \{\mathbf{z}_i\}_{i=1}^{N_1}, t} \quad & \mathbf{p}'_{\tau} \boldsymbol{\beta} \\ \text{subject to} \quad & t + \mathbf{a}' \boldsymbol{\beta} + f \preceq_{\mathcal{C}_1} 0, & (\text{cone in } \mathbb{R}) \\ & \mathbf{1}' \mathbf{z}^{[i]} \preceq_{\mathcal{C}_2} Q_1^{[i]} + \varrho^{[i]}, \quad i = 1, \dots, N_1 & (N_1 \text{ cones in } \mathbb{R}) \\ & -\mathbf{w} \preceq_{\mathcal{C}_3} \mathbf{z}, & (\text{cone in } \mathbb{R}^{J \cdot N_1}) \\ & \mathbf{w} \preceq_{\mathcal{C}_4} \mathbf{z}. & (\text{cone in } \mathbb{R}^{J \cdot N_1}) \\ & \begin{bmatrix} 1 - t \\ 2\mathbf{Q}^{1/2} \boldsymbol{\beta} \end{bmatrix} \preceq_{\mathcal{C}_5} 1 + t, & (\text{cone in } \mathbb{R}^{2+(J+KM) \cdot N_1}) \end{aligned}$$

where $\mathcal{K} = \mathcal{C}_1 \times \mathcal{C}_2^{N_1} \times \mathcal{C}_3 \times \mathcal{C}_4 \times \mathcal{C}_5 = \mathbb{R}_+ \times \mathbb{R}_+^{N_1} \times \mathbb{R}_+^{J \cdot N_1} \times \mathbb{R}_+^{J \cdot N_1} \times \mathcal{K}_{1+(J+KM) \cdot N_1}$ is the conic constraint for this program, $\mathbf{a} = -2(\mathbf{Q}\hat{\boldsymbol{\beta}} + \mathbf{G}^*)'$, and $f = \hat{\boldsymbol{\beta}}' \mathbf{Q} \hat{\boldsymbol{\beta}} + 2\mathbf{G}^* \hat{\boldsymbol{\beta}}$.

S.4 Data Preparation and Software Implementation

In this section, we first describe the variables in the [Billmeier and Nannicini \(2013\)](#) (BN, henceforth) dataset and explain all the changes and additions we made. Then, we illustrate how to use the companion R package `scpi` to prepare the data for the analysis and we briefly explain how to replicate the results.

S.4.1 Data Description

The original BN dataset contains data on some economic and political variables for 180 countries, over a period of time spanning from 1960 to 2005.¹ In detail, the variables available in the dataset are:

- real GDP per capita in 2002 US dollars
- enrollment rate in secondary schooling
- population growth
- yearly inflation rate
- the investment ratio (the investment of a country as a percentage of GDP)
- an indicator that captures whether the country is a democracy (1) or not (0)
- an indicator that captures whether the economy of the country is considered closed (0) or not (1) as developed in [Sachs, Warner, Åslund and Fischer \(1995\)](#) (henceforth, Sachs-Warner indicator). In particular, the indicator takes value 0 if any of the following condition is verified:
 - i) the average tariff is above 40%
 - ii) non-tariff barriers are imposed on a volume of imports larger than 40%
 - iii) the country has a socialist economic system
 - iv) the exchange rate black market premium is above 20%
 - v) state monopolies control most of the country exports

Despite having six candidate variables to match on, we end up matching on two variables because of missing data. Specifically, in the case of Europe - analyzed in the main article - at least one of these variables is completely missing before 1990 for Albania, Bulgaria, Romania, Slovak Republic, and Slovenia. Moreover, the original schooling variable has missing data for our countries of interest. Thus we replace it with the percentage of complete secondary schooling attained in population from the Barro-Lee dataset on educational attainment.²

A second difference of our final dataset from the original one used in BN is the final pool of countries - treated and donors - on which we conduct the analysis. In particular, we adopt the following data cleaning criteria to select the countries to be included in our final dataset:

¹We downloaded the dataset from the Harvard Dataverse at <https://dataverse.harvard.edu/dataset.xhtml?persistentId=doi:10.7910/DVN/28699>.

²source: <http://www.barrolee.com/>

1. At least 35 data entries for real GDP per capita from 1963 to 2000. This excludes the following economies (number of periods when real GDP per capita is observed in parenthesis): Armenia (8), Azerbaijan (8), Belarus (8), Dominican Republic (21), Ecuador (31), El Salvador (30), Estonia (8), Georgia (8), Kazakhstan (8), Kyrgyzstan (8), Latvia (8), Lithuania (8), Moldova (8), Russia (8), Tajikistan (8), Turkmenistan (8), Ukraine (8), and Uzbekistan (8).
2. At least 15 pre-treatment periods, that is, the economy is observed for at least 15 periods before it embarks on the liberalization program in the Sachs-Warner. This excludes the following economies (year in which the economy was first declared liberalized in parenthesis): Australia (1964), Austria (≤ 1963), Barbados (1966), Belgium (≤ 1963), Bolivia (1964), Canada (≤ 1963), Chile (1976), Cyprus (1964), Denmark (1966), Finland (≤ 1963), France (≤ 1963), Germany (≤ 1963), Greece (≤ 1963), Hong Kong (≤ 1963), Indonesia (1970), Ireland (1966), Italy (≤ 1963), Jamaica (≤ 1963), Japan (1964), Jordan (1965), Kenya (1965), Luxembourg (≤ 1963), Malaysia (≤ 1963), Mauritius (1968), Morocco (≤ 1963), Netherlands (≤ 1963), Norway (≤ 1963), Peru (≤ 1963), Portugal (≤ 1963), Singapore (1965), South Korea (1968), Spain (≤ 1963), Sri Lanka (1977), Sweden (≤ 1963), Switzerland (≤ 1963), Syria (≤ 1963), Taiwan (≤ 1963), Thailand (≤ 1963), United Kingdom (≤ 1963), United States (≤ 1963), Yemen (1965).
3. At least one entry for the percentage of complete secondary schooling attained in population. This excludes the following economies: Angola, Burkina Faso, Chad, Ethiopia, Guinea, Guinea-Bissau, Madagascar, Nigeria, and North Macedonia.
4. The liberalization is an absorbing state, that is, whenever the country embarks on a liberalization program, it remains so for the whole remaining time this country is observed. This excludes Venezuela.

Table [S.1](#) shows the final set of countries we select, together with their treatment date.

Table S.1: *List of all countries included in our analysis*

Africa				Asia			
Algeria	∞	Madagascar	1996	Bangladesh	1996	Nepal	1991
Angola	∞	Malawi	∞	China	∞	Pakistan	∞
Benin	1990	Mali	1988	India	∞	Philippines	1988
Botswana	1979	Mauritania	1995	Iran	∞	Turkey	1989
Burkina Faso	1998	Mozambique	1995	Israel	1985		
Burundi	1999	Niger	1994				
Cameroon	1993	Nigeria	∞				
Chad	∞	Rwanda	∞				
Congo	∞	Senegal	∞				
Egypt	1995	Sierra Leone	∞	Europe			
Ethiopia	1996	South Africa	1991	Albania	1992	Malta	∞
Gabon	∞	Tanzania	1995	Bulgaria	1991	North Macedonia	1994
Gambia	1985	Togo	∞	Croatia	∞	Poland	1990
Ghana	1985	Tunisia	1989	Czech Republic	1991	Romania	1992
Guinea	1986	Uganda	1988	Hungary	1990	Slovak Republic	1991
Guinea-Bissau	1987	Zambia	1993	Iceland	∞	Slovenia	1991
Ivory Coast	1994	Zimbabwe	∞				
Lesotho	∞						
North America				South America			
Costa Rica	1986	Mexico	1986	Argentina	1991	Guyana	1988
Guatemala	1988	Nicaragua	1991	Brazil	1991	Paraguay	1989
Haiti	∞	Panama	1996	Colombia	1986	Uruguay	1990
Honduras	1991						

Notes: ∞ denotes that a country has never experienced liberalization during the time span of observation.

To be precise, not all treated units in Table S.1 are analyzed in our work; we only study our target causal predictands for those units having at least 10 post-treatment periods. In other words, units treated after 1992 are not analyzed. Despite this fact, we still include units treated after 1992 because the staggered nature of this design, together with the fact that we focus on effects up to 10 years after liberalization events, make it possible to use such economies as donor units. To better illustrate this concept, take the case of Botswana that gets the treatment in 1979 and for which we study the effects of such treatment up to 1988 (see Figure S.1c). This allows us to use all countries treated from 1989 (included) onwards as donor units. In the same spirit, all countries treated after 1994 can be used as donor units for Israel (see Figure S.4c).

S.4.2 Code

In what follows we briefly explain how to replicate the results presented in the main paper and in this supplemental appendix using our R software package `scpi`. In particular, we focus on the two functions `scdataMulti()` and `scplotMulti()`, which have to be considered as the multiple treated units counterpart of `scdata()` and `scplot()` that we introduce and thoroughly describe in Cattaneo, Feng, Palomba and Titiunik (2023).

The next snippet shows how to prepare the data to predict the individual treatment effect τ_{it} and the average post-treatment effect τ_i . To accomplish this task, we leverage the function `scdataMulti()`, which receives as an input an object of class `DataFrame` and outputs the matrices **A**, **B**, **C**, and **P** together with other useful objects that are described in detail in the help files.

```
# Load packages and set seed
```

```

library(scp)
library(haven)
set.seed(8894)

# Load dataset
data <- haven::read_dta(paste0(path.data, "final_data.dta"))
data <- subset(data, restricted2 == 1)
data$lgdp <- log(data$rgdppp)

# Features to be matched
features <- list(c("lgdp", "lsc"))

# Covariates for adjustment
covs.adj <- list("lgdp" = c("constant", "trend"),
               "lsc" = c("constant", "trend"))

# Select treated units to be analyzed
units <- unique(subset(data, continent == "Europe" & treated == 1 & trDate <= 1992)$
               countryname)
units <- units[!(units %in% c("Czech Republic", "Poland"))]

# Set number of post-estimation periods
post.est <- 10

# Prepare data for Individual treatment effect
df <- scdataMulti(data, id.var = "countryname", outcome.var = "lgdp",
                 treatment.var = "liberalization", time.var = "year",
                 constant = TRUE, cointegrated.data = TRUE, post.est = post.est,
                 units.est = units, features = features, cov.adj = covs.adj, anticipation = 1)

# Prepare data for Average post-treatment effect
df.unit <- scdataMulti(data, id.var = "countryname", outcome.var = "lgdp", effect = "unit",
                     treatment.var = "liberalization", time.var = "year", constant = TRUE,
                     cointegrated.data = TRUE, post.est = post.est, units.est = units,
                     features = features, cov.adj = covs.adj, anticipation = 1)

# Prepare data for Average treatment effect on the treated
df.time <- scdataMulti(data, id.var = "countryname", outcome.var = "lgdp", effect = "time",
                     treatment.var = "liberalization", time.var = "year",
                     constant = TRUE, cointegrated.data = TRUE, post.est = post.est,
                     units.est = units[[cont]], features = features, cov.adj = covs.adj,
                     anticipation = 1)

```

First of all, it is important to notice the presence of the option `effect = "unit"` when it comes to predict the average post-treatment effect and of the option `effect = "time"` when the predictand of interest is the average treatment effect on the treated. Second, the most important input variable for `scdataMulti()` is the treatment variable `treatment.var`. In our case, `treatment.var` is the Sachs-Warner indicator which *de facto* has 1 as the absorbing state (once a country is described as open in our data, it remains so for all the period of analysis). Third, `scdataMulti()` has two options that allow users to choose which treated units to study and how many post-treatment periods should be considered in the analysis. These options are `units.est` and `post.est`, respectively. Finally, all the other options—`cointegrated.data`, `constant`, `anticipation`, `features`, `cov.adj`, `id.var` and `time.var`—are identical to those of `scdata()`.

Once the data matrices **A**, **B**, **C**, and **P** have been prepared, uncertainty quantification proceeds in the standard way. Specifically, the user has to call the function `scpi()` with the appropriate input object.

```

res <- scpi(df, sims = sims, cores = cores, w.constr = list("name" = "L1-L2"),
           u.order = 1, u.lags = 1)

res.unit <- scpi(df.unit, sims = sims, cores = cores, w.constr = list("name" = "L1-L2"),
                u.order = 1, u.lags = 1)

```

```
res.time <- scpi(df.time, sims = sims, cores = cores, w.constr = list("name" = "L1-L2"),
  u.order = 1, u.lags = 1)
```

Finally, it is possible to visualize the results through the function `scplotMulti()`. This function allows the user to either plot the synthetic control and the actual time series (`type = "series"`) or directly the treatment effect time series (`type = "treatment"`). This latter is simply the difference between the actual time series and the synthetic control one. In addition, simultaneous prediction intervals are readily available through the option `joint = TRUE`.

```
scplotMulti(res[, type = "series", joint = TRUE, ncols = 2)
scplotMulti(res.unit[, type = "series", joint = TRUE, ncols = 2)
scplotMulti(res.time[, type = "series", joint = TRUE, ncols = 2)

scplotMulti(res[, type = "treatment", joint = TRUE, ncols = 2)
scplotMulti(res.unit[, type = "treatment", joint = TRUE, ncols = 2)
scplotMulti(res.time[, type = "treatment", joint = TRUE, ncols = 2)
```

The next and final snippet showcases how to prepare the data for the third causal quantity analyzed in this paper: the average treatment effect on the treated at s_0 .

```
# Load packages and set seed
library(scp)
library(haven)
set.seed(8894)

data <- haven::read_dta(paste0(path.data, "final_data.dta"))
data <- subset(data, restricted2 == 1)
data$lgdp <- log(data$rgdppp)
post.est <- 10

# One Feature
eu.tr.91 <- unique(subset(data, trDate %in% c(1990,1991,1992) & continent == "Europe")$
  countryname)

data.co <- subset(data, !(countryname %in% eu.tr.91))
data.tr <- subset(data, countryname %in% eu.tr.91)

data.tr.agg <- aggregate(data.tr[c("lgdp", "liberalization")],
  by=list(data.tr$year), FUN=mean, na.rm = TRUE)
names(data.tr.agg) <- c("year", "lgdp", "liberalization")
data.tr.agg$countryname <- "Average Unit"
data.tr.agg$liberalization <- 1*(data.tr.agg$liberalization > 0)
data.agg <- rbind(data.tr.agg, data.co[names(data.tr.agg)])

features <- list(c("lgdp"))
covs.adj <- list("lgdp" = c("constant", "trend"))

df.agg <- scdataMulti(data.agg, id.var = "countryname", outcome.var = "lgdp",
  treatment.var = "liberalization", time.var = "year",
  constant = FALSE, cointegrated.data = T, post.est = 10,
  units.est = "Average Unit", features=features, cov.adj=covs.adj)

res.avg.l112 <- scpi(df.agg, sims = sims, cores = 1, w.constr = list("name" = "L1-L2"),
  u.order = 1, u.lags = 1)
```

Data preparation follows exactly the lines of Example 3.3, and thus the selected treated units are aggregated into a single one—termed “ave”—whose potential outcomes are simply

$$Y_t^{\text{ave}}(s) := \frac{1}{n_{s_0}} \sum_{i:T_i=s_0} Y_{it}(s), \quad t = 1, \dots, T.$$

The predictand of interest can be computed using “ave” as the only treated unit and control units from pre-treatment periods. Then, everything else proceeds as in the individual treatment effect case.


S.5 Results with Simplex Constraint

In this Section we report the results for Africa, Asia, Europe, North America, and South America using the simplex constraint in place of the L1-L2 constraint used in the main analysis, that is,

$$\mathcal{W} = \times_{i=1}^{N_1} \left\{ \mathbf{w}^{[i]} \in \mathbb{R}_+^J : \|\mathbf{w}^{[i]}\|_1 = 1 \right\}.$$


S.5.1 Africa

Figure S.1: Individual Treatment Effects $\hat{\tau}_{it}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$



(b) $\hat{\tau}_{it}$



(c) \hat{w}_i



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.2: Average Post-Treatment Effects $\hat{\tau}_i$.



Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.


Figure S.3: Average Treatment Effects on the Treated τ_k .



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.


S.5.2 Asia

Figure S.4: Individual Treatment Effects $\hat{\tau}_{it}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$



(b) $\hat{\tau}_{it}$



(c) \hat{w}_i



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.5: Average Post-Treatment Effects $\hat{\tau}_i$.



Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.


Figure S.6: Average Treatment Effects on the Treated τ_k .



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.


S.5.3 Europe

Figure S.7: Individual Treatment Effects $\hat{\tau}_{it}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$



(b) $\hat{\tau}_{it}$



(c) \hat{w}_i


Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.8: Average Post-Treatment Effects $\hat{\tau}_i$.




Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.9: Average Treatment Effect on the Treated in 1990 and 1991.




Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 500 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.

Figure S.10: Average Treatment Effects on the Treated τ_k .



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$

(b) $\hat{\tau}_k$




(c) \hat{w}_i

Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.


S.5.4 North America

Figure S.11: Individual Treatment Effects $\hat{\tau}_{it}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$



(b) $\hat{\tau}_{it}$



(c) \hat{w}_i



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.12: Average Post-Treatment Effects $\hat{\tau}_i$.



Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.


Figure S.13: Average Treatment Effects on the Treated τ_k .



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.


S.5.5 South America

Figure S.14: Individual Treatment Effects $\hat{\tau}_{it}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$



(b) $\hat{\tau}_{it}$



(c) \hat{w}_i


Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.15: Average Post-Treatment Effects $\hat{\tau}_i$.




Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.16: Average Treatment Effects on the Treated $\tau_{.k}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$

(b) $\hat{\tau}_{.k}$



(c) \hat{w}_i

Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.


S.6 Results with L1-L2, Other Continents

In this Section we report the results for Africa, Asia, North America, and South America using the L1-L2 constraint as in the main analysis for Europe, that is,

$$\mathcal{W} = \prod_{i=1}^{N_1} \left\{ \mathbf{w}^{[i]} \in \mathbb{R}^J : \|\mathbf{w}^{[i]}\|_1 = 1, \|\mathbf{w}^{[i]}\|_2 \leq Q_2^{[i]} \right\}.$$


S.6.1 Africa

Figure S.17: Individual Treatment Effects $\hat{\tau}_{it}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$



(b) $\hat{\tau}_{it}$



(c) \hat{w}_i



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.18: Average Post-Treatment Effects $\hat{\tau}_i$.



Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.19: Average Treatment Effects on the Treated τ_k .





(c) \hat{w}_i

Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.



S.6.2 Asia

Figure S.20: Individual Treatment Effects $\hat{\tau}_{it}$.





Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.21: Average Post-Treatment Effects $\hat{\tau}_i$.



Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.


Figure S.22: Average Treatment Effects on the Treated τ_k .



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.


S.6.3 North America

Figure S.23: Individual Treatment Effects $\hat{\tau}_{it}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$



(b) $\hat{\tau}_{it}$



(c) \hat{w}_i



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.24: Average Post-Treatment Effects $\hat{\tau}_i$.



Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.


Figure S.25: Average Treatment Effects on the Treated τ_k .



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.


S.6.4 South America

Figure S.26: Individual Treatment Effects $\hat{\tau}_{it}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$



(b) $\hat{\tau}_{it}$



(c) \hat{w}_i


Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.27: Average Post-Treatment Effects $\hat{\tau}_i$.




Notes: Blue bars report 90% prediction intervals, whereas red bars report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty is quantified using sub-Gaussian bounds.

Figure S.28: Average Treatment Effects on the Treated $\tau_{.k}$.



(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$

(b) $\hat{\tau}_{.k}$




(c) \hat{w}_i

Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.

S.7 Average treatment effect on countries liberalized in 1991


In this section we present alternative predictions of the average treatment effect on those European countries that underwent a liberalization program in 1991. First, we replicate the main analysis presented in Section 6 using a simplex-type constraint, and then we conduct the same exercise using two features: the logarithm of GDP per capita and the percentage of complete secondary schooling attained in population.

Figure S.29: *Average Treatment Effect on the Treated in 1991: Simplex constraint, $M = 1$.*




Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.

Figure S.30: Average Treatment Effect on the Treated in 1991: L1-L2 constraint, $M = 1$.



Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.

Figure S.31: Average Treatment Effect on the Treated in 1991: Simplex constraint, $M = 2$.




Notes: Blue bars report 90% prediction intervals, whereas blue shaded areas report 90% simultaneous prediction intervals. In-sample uncertainty is quantified using 200 simulations, whereas out-of-sample uncertainty using sub-Gaussian bounds.

S.8 Excluded Countries

In this section we present point predictions of individual treatment effects for all those treated units that have been excluded from the main analysis. Three different reasons lead us to remove a country from our study: (i) missing additional features; (ii) poor pre-treatment fit; (iii) treatment date between 1993 and 2000.


Case I: Missing Additional Features. In the case of North Macedonia, we only have data on real GDP per capita, as we do not observe the percentage of complete secondary schooling attained in population. Figure S.32 shows the estimated synthetic control for North Macedonia when matching exclusively on real income.

Figure S.32: Individual Treatment Effect $\hat{\tau}_{it}$: North Macedonia.




Case II: Poor pre-treatment fit. We excluded Slovenia from the main analysis that uses the L1-L2 constraint. This decision is motivated by an evident discrepancy between the realized series for real income and the synthetic control (Figure S.33a). When we use a simplex-type constraint (Figure S.33b), the number of excluded countries rises to 4—Czech Republic, Poland, Slovak Republic, and Slovenia. This happens because the synthetic control obtained under the simplex constraint is a convex combinations of unexposed units, i.e., each weight is forced to be within the closed interval $[0, 1]$ (Figure S.33d). The L1-L2 constraint relies on the same constraint but on top of that adds a ridge-type constraint. In this case, it is evident that this leads to a better pre-treatment matching for at least three countries (Czech Republic, Poland, and Slovak Republic).


Figure S.33: *Individual Treatment Effect $\hat{\tau}_{it}$: Excluded European countries.*




(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$, *L1-L2*



(b) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$, *simplex*




(c) \hat{w}_i , *L1-L2*




(d) \hat{w}_i , *simplex*

Case III: treatment date after 1992. Since our main analysis requires at least 10 post-treatment periods, our dataset ranges until the year 2000, and we consider one year of possible anticipation effects, units treated in 1993 or afterwards are not included. Indeed, such units are observed for less than 10 years in the treated status. Figure S.34 reports the point predictions for all these countries.


Figure S.34: Individual Treatment Effect $\hat{\tau}_{it}$: Treated after 1992.




(a) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$, L1-L2



(b) $Y_{it}(T_i)$ and $\hat{Y}_{it}(\infty)$, simplex



(c) \hat{w}_i , L1-L2



(d) \hat{w}_i , simplex

Supplement References

- Billmeier, A., and Nannicini, T. (2013), “Assessing Economic Liberalization Episodes: A Synthetic Control Approach,” *Review of Economics and Statistics*, 95, 983–1001.
- Boyd, S., and Vandenberghe, L. (2004), *Convex Optimization*, Cambridge University Press.
- Cattaneo, M. D., Feng, Y., Palomba, F., and Titiunik, R. (2023), “`scpi`: Uncertainty Quantification for Synthetic Control Methods,” arXiv:2202.05984.
- Cattaneo, M. D., Feng, Y., and Titiunik, R. (2021), “Prediction Intervals for Synthetic Control Methods,” *Journal of the American Statistical Association*, 116, 1865–1880.
- Raič, M. (2019), “A Multivariate Berry–Esseen Theorem with Explicit Constants,” *Bernoulli*, 25, 2824–2853.
- Ravishanker, N., Hochberg, Y., and Melnick, E. L. (1987), “Approximate Simultaneous Prediction Intervals for Multiple Forecasts,” *Technometrics*, 29, 371–376.
- Sachs, J. D., Warner, A., Åslund, A., and Fischer, S. (1995), “Economic Reform and the Process of Global Integration,” *Brookings Papers on Economic Activity*, 1995, 1–118.