arXiv:2210.05541v1 [hep-th] 11 Oct 2022

Radiation Reaction and Gravitational Waves at Fourth Post-Minkowskian Order

Christoph Dlapa,¹ Gregor Kälin,¹ Zhengwen Liu,^{2, 1} Jakob Neef,^{3, 4} and Rafael A. Porto¹

¹Deutsches Elektronen-Synchrotron DESY, Notkestr. 85, 22607 Hamburg, Germany

²Niels Bohr International Academy, Niels Bohr Institute,

University of Copenhagen, Blegdamsvej 17, DK-2100 Copenhagen, Denmark

³School of Mathematics and Statistics, University College Dublin, Belfield, Dublin 4, Ireland, D04 V1W8

⁴Humboldt-Universität zu Berlin, Zum Grossen Windkanal 2, D-12489 Berlin, Germany

We obtain the total impulse in the scattering of non-spinning binaries in general relativity at fourth Post-Minkowskian order, i.e. $\mathcal{O}(G^4)$, including linear, nonlinear, and hereditary radiation-reaction effects. We derive the total radiated spacetime momentum as well as the associated energy flux. The latter can be used to compute gravitational-wave observables for generic (un)bound orbits. We employ the ('in-in') Schwinger-Keldysh worldline effective field theory framework in combination with modern 'multi-loop' integration techniques from collider physics. The complete results are in agreement with various partial calculations in the Post-Newtonian/Minkowskian expansion.

Introduction. Waveform models are an essential ingredient in data analysis and characterization of gravitational wave (GW) signals from compact binaries [1]. The level of accuracy plays a critical role, in particular for future detectors such as LISA [2] and ET [3]. In order to benefit the most from the anticipated observational reach [2–9], the modelling of GW sources must therefore continue to develop—both through analytic methodologies [10–18] and numerical simulations [19–21]—in parallel with the expected increase in sensitivity with nextgeneration GW interferometers.

Motivated by the *Effective-One-Body* (EOB) formalism [22–27], the *Boundary-to-Bound* (B2B) dictionary between unbound and bound observables [28–30], and benefiting from powerful 'multi-loop' integrations tools [31–61], significant progress has been achieved in recent years in our analytic understanding of (classical) gravitational scattering in the Post-Minkowskian (PM) expansion in powers of *G* (Newton's constant); both via effective field theory (EFT) [62–78] and amplitudebased [79–98] methodologies. The PM regime incorporates an infinite tower of Post-Newtonian (PN) corrections at a given order in *G* that may increase the accuracy of phenomenological waveform models [99, 100].

However, despite some notable exceptions [26, 73– 77, 79, 84, 97, 98, 101], the majority of the PM computations have so far impacted our knowledge of the *conser*vative sector, with potential interactions [66, 92] as well as 'tail effects' [67, 93] known to 4PM order. Yet, until now, complete results had not been obtained at the same level of accuracy. The purpose of this letter is therefore to report the total change of (mechanical) momentum, a.k.a. the impulse, for the gravitational scattering of non-spinning bodies—including all the hitherto unknown linear, nonlinear and hereditary radiation-reaction dissipative effects—at $\mathcal{O}(G^4)$, from which we derive the total radiated spacetime momentum and GW energy flux.

Building on pioneering developments in the PN regime [102–109], the derivation proceeds via the EFT approach

in a PM scheme [62], extended in [76] to simultaneously incorporate conservative and dissipative effects via the 'in-in' Schwinger-Keldysh formalism [110-114]. As discussed in [76], the in-in impulse resembles the 'in-out' counterpart used in the conservative sector [66, 67], except for its causal structure which entails the use of retarded Green's functions [76]. After adapting integration tools to our problem, the calculation of the impulse is mapped to a series of 'three-loop' mass-independent integrals. As in previous derivations [62, 63, 66, 67], the latter are solved via the methodology of differential equations [32–38]. The relativistic two-body problem is then reduced to obtaining the necessary boundary conditions in the near-static limit. The boundary integrals are computed using the method of regions [45], involving potential (off-shell) and radiation (on-shell) modes [102]. The full solution is thus *bootstrapped* to all orders in the velocities from the same type of calculations needed in the EFT approach with PN sources [102, 109, 115-118]. As a nontrivial check, by rewriting retarded Green's functions as Feynman propagators plus a *reactive* term [76], we recover the value in [66, 67] for the Feynman-only (conservative) part. Agreement is also found in the overlap with various PN derivations [26, 27, 119–124] and partial PM results [98] obtained using the relations in [22].

The B2B dictionary [28–30] allows us to connect scattering data to observables for bound states via analytic continuation. However, similarly to the lack of periastron advance at 3PM [28, 29, 63], the symmetries of the problem yield a vanishing coefficient for the radiated energy integrated over a period of elliptic-like motion at 4PM, trivially recovered by the B2B map. Nevertheless, since nonlinear radiation-reaction effects do not contribute to the integrated radiated energy at $\mathcal{O}(G^4)$, we can then derive the GW flux in an adiabatic approximation [30]. This allows for the computation of radiative observables for generic (un)bound orbits through balance equations, as in the EOB approach [22], thus including an infinite series of velocity corrections.



Figure 1. In-in Feynman topologies to 4PM order. The arrows indicate the flow of (retarded) time. The crosses represent the location of the derivative in the impulse in (3). The last two diagrams are the only 'self-energies' needed at 4PM [76].

The EFT *in-integrand*. Following the Schwinger-Keldysh formalism [110–114] adapted to the EFT approach in [76, 104], the effective action is obtained via a *closed-time-path* integral involving a doubling of the metric perturbation $(h_{\mu\nu}^{\pm})$ as well as the worldline $(x_{a,\pm}^{\alpha})$ degrees of freedom, schematically,

$$e^{i\mathcal{S}_{\rm eff}[x_{a,\pm}]} = \int \mathcal{D}h^+ \mathcal{D}h^- e^{i\left\{S_{\rm EH}[h^{\pm}] + S_{\rm pp}[h^{\pm}, x_{a,\pm}]\right\}}, \quad (1)$$

with $S_{\rm EH}$ and $S_{\rm pp}$ the closed-path version of the Einstein-Hilbert and point-particle worldline actions, respectively. We ignore here spin degrees of freedom and finite-size effects (see [62, 65]). We also restrict ourselves to the classical regime and therefore the path integral in (1) is computed in the saddle-point approximation—keeping only connected 'tree-level' Feynman diagrams of the gravitational field(s)—with the compact objects treated as external nonpropagating sources.

In this scenario, the matrix of (causal) propagators is given by the Keldysh representation:

$$K^{AB}(x-y) = i \begin{pmatrix} 0 & -\Delta_{\mathrm{adv}}(x-y) \\ -\Delta_{\mathrm{ret}}(x-y) & 0 \end{pmatrix}, \quad (2)$$

with $A, B \in \{+, -\}$ and $\Delta_{\text{ret/adv}}$ the standard retarded/advanced Green's functions. The impulse, e.g. for particle 1, then follows from [76]

$$\Delta p_{1}^{\mu} = -\eta^{\mu\nu} \int_{-\infty}^{\infty} \mathrm{d}\tau_{1} \frac{\delta \mathcal{S}_{\mathrm{eff}}[x_{a,\pm}]}{\delta x_{1,-}^{\nu}(\tau_{1})} \bigg|_{\mathrm{PL}} = \sum_{n} G^{n} \Delta^{(n)} p_{1}^{\mu} \,, \quad (3)$$

to all PM orders, where the subscript 'PL' stands for the *Physical Limit*: $\{x_{a,-} \rightarrow 0, x_{a,+} \rightarrow x_a\}$ [103]. As for the conservative sector [66, 67], we must also include *iterations* from lower order solutions to the equations of motion. The latter are obtained from the effective action in the same physical limit. The diagrams needed to $\mathcal{O}(G^4)$ are depicted in Fig. 1. Mirror images (not shown) must also be computed. See [76] for details.

The impulse is further decomposed into scalar integrals in the perpendicular and longitudinal directions, i.e. for particle 1 (and likewise for particle 2)

$$\Delta^{(n)} p_1^{\mu} = c_{1b}^{(n)} \frac{\hat{b}^{\mu}}{b^n} + \frac{1}{b^n} \sum_a c_{1\tilde{u}_a}^{(n)} \check{u}_a^{\mu} , \qquad (4)$$

with $b^{\mu} \equiv b_1^{\mu} - b_2^{\mu}$ the impact parameter, $b \equiv \sqrt{-b^{\mu}b_{\mu}}$, and $\hat{b}^{\mu} \equiv b^{\mu}/b$. We use the notation [97]

$$\check{u}_{1}^{\mu} \equiv \frac{\gamma u_{2}^{\mu} - u_{1}^{\mu}}{\gamma^{2} - 1} , \quad \check{u}_{2}^{\mu} \equiv \frac{\gamma u_{1}^{\mu} - u_{2}^{\mu}}{\gamma^{2} - 1} , \quad \gamma \equiv u_{1} \cdot u_{2} , \quad (5)$$

with u_a 's the incoming velocities, $b \cdot u_a = 0$, $u_a^2 = 1$, and $\check{u}_a \cdot u_b = \delta_{ab}$. Ignoring absorption, the preservation of the on-shell condition, $p_a^2 = m_a^2$, implies $2p_a \cdot \Delta p_a = -(\Delta p_a)^2$, which serves as a nontrivial consistency check.

Integration. Similarly to the derivations in [66, 67], but incorporating the key distinction between Feynman and retarded propagators, the components of the impulse can be reduced to different families of integrals,

$$\int \prod_{i=1}^{3} \frac{\mathrm{d}^{d}\ell_{i}}{\pi^{d/2}} \frac{\delta(\ell_{i} \cdot u_{a_{i}})}{(\pm \ell_{i} \cdot u_{\phi_{i}} - i0)^{n_{i}}} \prod_{k=1}^{9} \frac{1}{D_{k}^{\nu_{k}}}, \qquad (6)$$

restricted by Dirac- δ functions. Following [66, 67], the $\ell_{i=1,2,3}$'s are the loop momenta, n_i, ν_k are integers, and $a_i \in \{1,2\}$, with $u_{\underline{1}} = u_2, u_{\underline{2}} = u_1$. In contrast to the conservative part, the D_k 's are now various sets of retarded/advanced propagators, e.g. $\{(\ell^0 \pm i0)^2 - \ell^2, \ldots\}$, consistent with causality. The same constraints as before apply on the external data [63], such that the relevant integrals can only depend on γ . As in our previous calculations [66, 67], we conveniently introduce the parameter x, defined through the relation $\gamma \equiv (x^2+1)/2x$ [95], and compute these integrals by using dimensional regularization (in $d \equiv 4 - 2\epsilon$ dimensions) and the method of differential equations [32–38].

The integration problem then resembles the steps already performed for the computation in [66, 67], except for a few notable differences. First of all, as before we use integration-by-parts (IBP) relations [39–44] and reduce (6) to a basis of master integrals. Because of the fewer number of symmetries of the in-in integrand, the algebraic manipulations become a bit more involved than with Feynman-only propagators. But more importantly, the boundary conditions in the near-static limit $\gamma \simeq 1$ must be computed in terms of retarded/advanced Green's functions. For this purpose, we resort to the method of regions and the expansion into potential and radiation modes. The potential-only part was obtained in [66], and recovered here from the full solution. For radiation modes, the same type of integrals appearing in PN derivations [109], combined with leftover integrals over potential-only modes at one and two loops, are sufficient to bootstrap the entire answer. See [61] for more details.

Total impulse. Inputing the values of the boundary master integrals and translating from x to γ space, we find

$$\begin{split} \frac{c_{14}^{(1)tot}}{\pi} &= -\frac{3h_1m_1m_2(m_1^3 + m_2^3)}{64(\gamma^2 - 1)^{5/2}} + m_1^2m_2^2(m_1 + m_2) \Big[\frac{21h_2E^2\left(\frac{\gamma+1}{\gamma+1}\right)}{32(\gamma - 1)\sqrt{\gamma^2 - 1}} + \frac{3h_3K^2\left(\frac{\gamma+1}{\gamma+1}\right)}{16(\gamma^2 - 1)^{3/2}} - \frac{3h_4E\left(\frac{\gamma+1}{\gamma+1}\right)}{16(\gamma^2 - 1)^{3/2}} + \frac{\pi^2h_5}{8\sqrt{\gamma^2 - 1}} + \frac{h_6\log\left(\frac{\gamma+1}{2}\right)}{16(\gamma^2 - 1)^{5/2}} \\ &+ \frac{3h_7Li_2\left(\frac{\sqrt{\gamma+1}}{\gamma+1}\right)}{4(\gamma - 1)(\gamma + 1)^2} - \frac{3h_3Li_2\left(\frac{\gamma+1}{\gamma+1}\right)}{4(\gamma - 1)(\gamma + 1)^2} \Big] + m_1^2m_2^2 \Big[\frac{h_8}{48(\gamma^2 - 1)^3} + \frac{\sqrt{\gamma^2 - 1}h_9}{768(\gamma - 1)^{3/9}(\gamma + 1)^4} + \frac{h_{10}\log\left(\frac{\gamma+1}{2}\right)}{8(\gamma^2 - 1)^2} - \frac{h_{11}\log\left(\frac{\gamma+1}{2}\right)}{32(\gamma^2 - 1)^{5/2}} + \frac{h_{12}\log(\gamma)}{16(\gamma^2 - 1)^{5/2}} \\ &- \frac{h_{13}\operatorname{arccosh}(\gamma)}{8(\gamma - 1)(\gamma + 1)^4} + \frac{h_{14}\operatorname{arccosh}(\gamma)}{16(\gamma^2 - 1)^{7/2}} - \frac{3h_{15}\log\left(\frac{\gamma+1}{2}\right)}{8\sqrt{\gamma^2 - 1}} + \frac{3h_{16}\log\left(\frac{\gamma+1}{2}\right)}{8\sqrt{\gamma^2 - 1}} + \frac{3h_{16}\log\left(\frac{\gamma+1}{2}\right)}{16(\gamma^2 - 1)^2} - \frac{3h_{17}Li_2\left(\frac{\gamma+1}{\gamma+1}\right)}{64(\gamma^2 - 1)^2} - \frac{3}{2}\sqrt{\gamma^2 - 1}h_{18}Li_2\left(\frac{1 - \gamma}{\gamma + 1}\right)} \Big] \\ &+ m_1^2m_2^2 \Big[\frac{3h_{15}\log\left(\frac{\gamma+1}{2}\right)}{8\sqrt{\gamma^2 - 1}} + \frac{3h_{16}\log\left(\frac{\gamma+1}{2}\right)}{16(\gamma^2 - 1)^2} + \frac{h_{19}}{4k_{19}(\gamma^2 - 1)^3} + \frac{h_{20}\log\left(\frac{\gamma+1}{2}\right)}{192\gamma^7(\gamma^2 - 1)^{5/2}} + \frac{h_{21}\log\left(\frac{\gamma+1}{2}\right)}{8(\gamma^2 - 1)^2} + \frac{h_{22}\log\left(\frac{\gamma+1}{2}\right)}{2(\gamma^2 - 1)^{3/2}} + \frac{h_{23}\log(\gamma)}{16(\gamma^2 - 1)^{3/2}} \\ &- \frac{h_{24}\operatorname{arccosh}(\gamma)}{16(\gamma^2 - 1)^{7/2}} - \frac{3h_{26}\operatorname{arccosh}(\gamma)}{3k_{20}(\gamma^2 - 1)^{7/2}} + \frac{3h_{20}\log\left(\frac{\gamma+1}{2}\right)}{2(\gamma^2 - 1)^{5/2}} + \frac{h_{21}\log\left(\frac{\gamma+1}{2}\right)}{16(\gamma^2 - 1)^2} + \frac{h_{22}\log\left(\frac{\gamma+1}{2}\right)}{16(\gamma^2 - 1)^{3/2}} + \frac{h_{23}\log(\gamma)}{16(\gamma^2 - 1)^{3/2}} \\ &- \frac{h_{24}\operatorname{arccosh}(\gamma)}{16(\gamma^2 - 1)^{7/2}} - \frac{3h_{26}\operatorname{arccosh}(\gamma)}{3k_{20}(\gamma^2 - 1)^{7/2}} + \frac{3h_{26}\log\left(\frac{\gamma+1}{2}\right)}{2(\gamma^2 - 1)^{3/2}} + \frac{h_{22}\log\left(\frac{\gamma+1}{2}\right)}{16(\gamma^2 - 1)^{3/2}} + \frac{h_{23}\log(\gamma)}{16(\gamma^2 - 1)^{3/2}} \\ &- \frac{h_{24}\operatorname{arccosh}(\gamma)}{(\gamma^2 - 1)^{3/2}} + \frac{2h_{23}\operatorname{arccosh}(\gamma)}{(\gamma^2 - 1)^{3/2}} + \frac{3h_{24}\operatorname{arccosh}(\gamma)}{16(\gamma^2 - 1)^{2}} + \frac{h_{24}\log\left(\frac{\gamma+1}{2}\right)}{16(\gamma^2 - 1)^{3/2}} \\ &- \frac{h_{24}\operatorname{arccosh}(\gamma)}{(\gamma^2 - 1)^{3/2}} + \frac{h_{24}\operatorname{arccosh}(\gamma)}{(\gamma^2 - 1)^{3/2}} + \frac{h_{24}\operatorname{arccosh}(\gamma)}{16(\gamma^2 - 1)^{3/2}} + \frac{h_{24}\operatorname{arccosh}($$

See Table I for the list of $h_i(\gamma)$ polynomials.

The impulse for the second particle follows by exchanging $1 \leftrightarrow 2$ in the masses, incoming velocities, and impact parameter. As expected from the calculations in [66, 67], the complete results feature dilogarithms (Li₂(z)), and complete elliptic integrals of the first (K(z)) and second (E(z)) kind.

Conservative. As shown explicitly in [76], a (timesymmetric) conservative contribution can be identified by rewriting retarded Green's functions in terms of Feynman propagators plus a reactive term, and keeping the real part of the Feynman-only piece, i.e. $\Delta p_{\rm cons}^{\mu} \equiv \mathbb{R} \Delta p_{\rm F}^{\mu}$, with imaginary terms canceling out against counter-parts from the reactive terms. Performing these steps in the full in-in integrand, and associated boundary conditions entering in the total impulse, we readily recover the conservative results in [66, 67] including potential and radiation-reaction tail effects.

Dissipative. As discussed in [76], the terms stemming off of the mismatch between Feynman and retarded propagators incorporate dissipative effects. Needless to say, these terms can also be read off directly from the total result by subtracting the conservative part. Following the analysis in [66, 67], we disentangle the various pieces according to factors of $v_{\infty}^{-2\epsilon}$, with $v_{\infty} \equiv \sqrt{\gamma^2 - 1}$, which signal the presence of an on-shell mode.

Starting with a single radiation mode we encounter instantaneous dissipative effects at linear order in the radiation-reaction. The latter are odd under time reversal and contribute to the \hat{b} and \check{u}_a directions. We find agreement in the overlap with known partial results in linear-response theory in the PN literature [26, 27, 119– 123], as well as with the (odd) contribution to the $c_{1b}^{(4)}$ coefficient recently derived in [98]. All of the remaining radiative terms involve two radiation modes. After removing the Feynman-only (radiative) conservative pieces [66, 67], the leftovers contain hereditary as well as nonlinear radiation-reaction dissipative effects. The former enters both in the longitudinal and perpendicular directions, whereas the latter contributes only to the total radiated perpendicular momentum at this order.¹ We also find perfect consistency with known nonlinear and hereditary results in the PN expansion [120–124].

See the supplemental material and ancillary file for explicit expressions.

¹ At this point, however, we cannot distinguish whether nonlinear radiation-reaction terms are due to either effects at second order in the linear radiation-reaction force or truly nonlinear gravitational corrections.



Table I. Polynomials entering the total impulse as well as conservative/dissipative parts (see the supplemental material).

Radiated energy/momentum. From the impulse we derive the change in the mechanical momentum of the system (in the incoming center-of-mass), which gives us the total radiated momentum, $P_{\rm rad}^{\mu} = -(\Delta p_1^{\mu} + \Delta p_2^{\mu})$. The radiated energy for hyperbolic motion at 4PM, given by $\Delta E_{\rm hyp} \equiv P_{\rm rad} \cdot \frac{m_1 u_1 + m_2 u_2}{M\Gamma}$ with $\Gamma \equiv \frac{E}{M} (M, E$ the total mass and energy and $\nu \equiv \frac{m_1 m_2}{M^2}$), then becomes:

$$\begin{split} \Delta E_{\text{hyp}}^{4\text{PM}} &= -\frac{G^4 M^5 \nu^2}{b^4 \Gamma} \Biggl\{ \frac{15\pi^2 \left(\gamma^2 - 1\right) \left(27 \left(\gamma^2 - 1\right) h_{31} + 2h_{50}\right) + 64 \left(45h_{32} - h_{48}\right)}{1440 \left(\gamma^2 - 1\right)^3} + \frac{h_{49}}{1440 \gamma^7 \left(\gamma^2 - 1\right)^{5/2}} - \operatorname{arccosh}^2(\gamma) \left(\frac{16h_{53}}{\left(\gamma^2 - 1\right)^2} + \frac{32h_{54}}{\left(\gamma^2 - 1\right)^{7/2}} \right) \\ &- \frac{h_{55} \log(2) \operatorname{arccosh}(\gamma)}{4 \left(\gamma^2 - 1\right)^2} + \frac{h_{57} \log \left(\frac{2}{\gamma + 1}\right)}{4 \left(\gamma^2 - 1\right)^2} - \frac{h_{58} \log(\gamma) \operatorname{arccosh}(\gamma)}{4 \left(\gamma^2 - 1\right)^2} + \operatorname{arccosh}(\gamma) \left(\frac{h_{51}}{480 \gamma^8 \left(\gamma^2 - 1\right)^3} - \frac{16h_{52}}{5 \left(\gamma^2 - 1\right)^{3/2}} \right) \\ &- \frac{h_{56} \text{Li}_2 \left(\sqrt{\frac{\gamma^2 - 1}{\gamma + 1}} \right)}{8 \left(\gamma^2 - 1\right)^2} + \frac{h_{57} \text{Li}_2 \left(\sqrt{\gamma^2 - 1} - \gamma \right)}{2 \left(\gamma^2 - 1\right)^2} + \frac{h_{58} \text{Li}_2 \left(- \left(\gamma - \sqrt{\gamma^2 - 1} \right)^2 \right)}{8 \left(\gamma^2 - 1\right)^2} \\ &+ \nu \left[\frac{4 \left(-45h_{32} + 30h_{33} - 30h_{37} + h_{48} \right)}{45 \left(\gamma^2 - 1\right)^3} + \frac{\pi^2 \left(54 \left(\gamma^2 - 1 \right) h_{31} + h_{39} - 4h_{50} \right)}{96 \left(\gamma^2 - 1\right)^2} - \operatorname{arccosh}^2(\gamma) \left(\frac{16 \left(h_{42} - 2h_{53}\right)}{\left(\gamma^2 - 1\right)^2} - \frac{64 \left(h_{43} + h_{54}\right)}{\left(\gamma^2 - 1\right)^{7/2}} \right) \\ &+ \frac{h_{38} - 490\gamma \left(3840\gamma^7 h_{34} + h_{49} \right)}{45 \left(\gamma^2 - 1\right)^3} + \frac{\left(3h_{46} + 2h_{57} \right) \log \left(\frac{\gamma^2 + 1}{2} \right) \operatorname{arccosh}(\gamma)}{4 \left(\gamma^2 - 1\right)^2} + \frac{\left(h_{44} + 2h_{55}\right) \log(2) \operatorname{arccosh}(\gamma)}{4 \left(\gamma^2 - 1\right)^2} + \frac{\left(h_{47} + 2h_{58}\right) \log(\gamma) \operatorname{arccosh}(\gamma)}{4 \left(\gamma^2 - 1\right)^2} \\ &+ \operatorname{arccosh}(\gamma) \left(\frac{53760\gamma^9 h_{35} - 14\gamma h_{51} + h_{40}}{3360\gamma^9 \left(\gamma^2 - 1\right)^3} - \frac{32 \left(15h_{36} - 10h_{41} - 3h_{52}\right)}{15 \left(\gamma^2 - 1\right)^{3/2}} \right) + \frac{\left(h_{56} - 6h_{45}\right) \operatorname{Li}_2 \left(\sqrt{\frac{\gamma^2 - 1}{\gamma + 1}} \right)}{4 \left(\gamma^2 - 1\right)^2} \\ &- \frac{\left(3h_{46} + 2h_{57}\right) \operatorname{Li}_2 \left(\sqrt{\gamma^2 - 1} - \gamma \right)}{2 \left(\gamma^2 - 1\right)^2} - \frac{\left(h_{47} + 2h_{58}\right) \operatorname{Li}_2 \left(- \left(\gamma - \sqrt{\gamma^2 - 1} \right)^2 \right)}{16 \left(\gamma^2 - 1\right)^2} \\ &- \frac{\left(3h_{46} + 2h_{57}\right) \operatorname{Li}_2 \left(\sqrt{\gamma^2 - 1} - \gamma \right)}{2 \left(\gamma^2 - 1\right)^2} - \frac{\left(h_{47} + 2h_{58}\right) \operatorname{Li}_2 \left(- \left(\gamma - \sqrt{\gamma^2 - 1} \right)^2 \right)}{16 \left(\gamma^2 - 1\right)^2} \\ \\ &- \frac{\left(3h_{46} + 2h_{57}\right) \operatorname{Li}_2 \left(\sqrt{\gamma^2 - 1} - \gamma \right)}{2 \left(\gamma^2 - 1\right)^2} - \frac{\left(h_{47} + 2h_{58}\right) \operatorname{Li}_2 \left(- \left(\gamma - \sqrt{\gamma^2 - 1} \right)^2 \right)}{16 \left(\gamma^2 - 1\right)^2} \\ \\ &- \frac{\left(h_{47} + 2h_{58}\right) \operatorname{Li}_2 \left(- \left(\gamma - \sqrt{\gamma^2 - 1} \right)^2 \right)}{16 \left(\gamma^2 -$$

Expanding in small velocities $(v_{\infty}^2 = \gamma^2 - 1 \ll 1)$ we find,

$$\frac{b^4 \Delta E_{\text{hyp}}^{4\text{PM}}}{G^4 M^5 \nu^2} = \frac{1568}{45 v_{\infty}} + \left(\frac{18608}{525} - \frac{1136\nu}{45}\right) v_{\infty} + \frac{3136 v_{\infty}^2}{45} \\
+ \left(\frac{764\nu^2}{45} - \frac{356\nu}{63} + \frac{220348}{11025}\right) v_{\infty}^3 + \left(\frac{1216}{105} - \frac{2272\nu}{45}\right) v_{\infty}^4 \\
+ \left(-\frac{622\nu^3}{45} + \frac{3028\nu^2}{1575} - \frac{199538\nu}{33075} - \frac{151854}{13475}\right) v_{\infty}^5 \\
+ \left(\frac{1528\nu^2}{45} - \frac{8056\nu}{1575} + \frac{117248}{1575}\right) v_{\infty}^6 + O\left(v_{\infty}^7\right) .$$
(9)

A notable feature of the 4PM momentum is the emergence of a total recoil along the perpendicular direction. The full result is rather lengthy, however, performing a velocity expansion we find (with $\Delta_m \equiv \frac{m_1 - m_2}{M}$)

$$\frac{b^4 P_{b,\text{rad}}^{4\text{PM}}}{\pi \Delta_m G^4 M^5 \nu^2} = \frac{37}{30} + \frac{1661 v_\infty^2}{50} + \frac{1491 v_\infty^3}{400} + \frac{23563 v_\infty^4}{10080} - \frac{26757 v_\infty^5}{5600} + \frac{700793 v_\infty^6}{506880} + O(v_\infty^7) \,. \tag{10}$$

Both expressions in (9) and (10) are consistent with the state-of-the-art in the PN expansion [119–124]. On the other hand, in the opposite limit, as γ gets large,

$$\frac{b^4 \Gamma \Delta E_{\rm hyp}^{\rm 4PM}}{G^4 M^5 \nu^2} \to \frac{13696}{105} \gamma^3 \nu \log(2\gamma) \,, \tag{11}$$

which signals the presence of (logarithmic) mass singularities. We return to these limits in the conclusions.

GW energy flux. The B2B map allows us to connect the radiated energy for the scattering process to its counterpart over a period of an elliptic-like orbit via $\Delta E_{\rm ell}(j) = \Delta E_{\rm hyp}(j) - \Delta E_{\rm hyp}(-j)$ [30], where $j \equiv \frac{p_{\infty}b}{GM^2\nu}$, with $p_{\infty} \equiv \frac{M\nu}{\Gamma} v_{\infty}$, is the (reduced) angular momentum. However, similarly to the periastron advance at $\mathcal{O}(G^3)$ [28, 29], this expression vanishes at 4PM. To obtain radiative observables for generic orbits we derived instead the energy flux. Since nonlinear radiation-reaction terms do not contribute to the energy loss at this order we can resort to an adiabatic expansion. By writing the PM-expanded energy flux in an isotropic gauge as [30]

$$\frac{dE}{dt} = \frac{M}{r} \sum_{n} \mathcal{F}_{E}^{(n)}(\gamma) \left(\frac{GM}{r}\right)^{(n+3)}, \qquad (12)$$

we find at 4PM order (see [30] for the 3PM term)

$$M\pi\xi \mathcal{F}_{E}^{(1)} = \frac{3\pi\Gamma^{2}\nu}{4(\gamma^{2}-1)^{3/2}} \Delta E_{j\,\rm hyp}^{(1)} - \frac{2\nu^{3}\Delta E_{j\,\rm hyp}^{(0)}}{(\gamma^{2}-1)^{2}\Gamma^{6}\xi^{2}} \times \left[(\gamma-1)^{3}\left(10\gamma^{3}-10\gamma^{2}-9\gamma+5\right)\nu^{2} + 4\left(5\gamma^{5}-8\gamma^{4}+\gamma^{3}+4\gamma^{2}-3\gamma+1\right)\nu + 8\gamma^{4}-4\gamma^{2}-1\right],$$
(13)

with $\Delta E(j) = \sum_{n=0}^{\infty} \frac{\Delta E_j^{(n)}}{j^{n+3}}, \ \xi \equiv \frac{E_1 E_2}{E^2}, \ E_a \equiv \sqrt{p_{\infty}^2 + m_a^2}.$

Conclusions. We completed the knowledge of the total relativistic impulse in the scattering of non-spinning bodies at 4PM order, including linear, nonlinear, and hereditary radiation-reaction effects. We also derived the total radiated spacetime momentum at $\mathcal{O}(G^4)$, from which we extracted the energy flux. In combination with conservative results in [66, 67], the GW flux can be used to compute observables for generic (un)bound orbits, incorporating an infinite series of velocity corrections. The most intricate part of the calculation involves a series of master integrals with retarded propagators, which we are able to compute to all orders in the velocity through the methodology of differential equations, without resorting to PN resummations. The boundary conditions in the near-static limit are obtained via the method of regions, thus making direct contact with derivations in the PN regime with potential and radiation modes. We find perfect agreement with various partial calculations in the literature. Explicit values can be found in the supplemental material and ancillary file.

There are still some key aspects of the structure of the impulse at 4PM order which deserve further study. Firstly, concerning the high-energy limit, although many nontrivial cancelations occur we find that the impulse does not transition smoothly, but rather it diverges when $m_a \rightarrow 0$ while $\gamma m_1 m_2$ is held fixed. Yet, we find that all integrals of the form in (6) vanish in dimensional regularization for the case of incoming velocities obeying the null condition $(u_a^2 = 0)$. This implies that the divergent terms in the $m_a \rightarrow 0$ limit are associated with the enforcement from the onset of timelike worldlines $(u_a^2 = 1)$ in the massive theory. Moreover, similarly to what happens at 3PM, the radiated energy diverges, this time with a factor of $\log(2\gamma)$ in the $\gamma \gg 1$ limit at 4PM order. We expect this behavior to be tamed in the nonperturbative solution (see e.g. [125, 126] and references therein).

Secondly, there is the issue of the mass scaling of the impulse [27], and violations thereof, e.g. [117, 118]. As we mentioned, some of the radiative contributions affect only the total radiated momentum in the \hat{b} -direction, while conserving energy. Moreover, they are even under time reversal, see (10). To gain intuition about these terms, from the impulse and total recoil we derived the relative deflection angle (see supplemental material). We find perfect consistency with the (odd-in-velocity) PN values in [27]. Yet, starting at 5PN order, we encounter conservative-like (even-in-velocity) $\mathcal{O}(\nu^2)$ contributions beyond the Feynman-only part.² In principle, depending on their origin, these terms could be incorporated into a *relative* Hamiltonian. We will discuss these issues in more detail elsewhere.

² The deflection angle is however in tension with the two distinct values reported in [117, 118].

In summary, in addition to the natural connections to GW science, e.g. [99, 100], the solution of the (classical) relativistic scattering problem at $\mathcal{O}(G^4)$ presented here demonstrates how the worldline EFT approach [15, 18], combined with the methodology of differential equations and integration by regions—already successfully implemented both in the conservative [62–67] and dissipative [72–76] sectors—are very powerful tools to tackle the entire two-body dynamics in general relativity within the PM expansion. Complete results at higher PM orders, including spin and tidal effects, are underway.

Acknowledgements. We thank Donato Bini, Johannes Blümlein, Ruth Britto, Gihyuk Cho, Walter Goldberger, Carlo Heissenberg, Francois Larrouturou, Massimiliano Riva, Ira Rothstein, Rodolfo Russo, and Chia-Hsien Shen for useful discussions. We are particularly grateful to Donato Bini, Thibault Damour and Andrea Geralico for sharing their results on higher order radiative effects in the PN expansion prior to submission. We would like to thank also the participants of the program "High-Precision Gravitational Waves" at the KITP (supported in part by the National Science Foundation under Grant No. NSF PHY-1748958) for discussions. ZL is grateful to the Mainz Institute for Theoretical Physics of the DFG Cluster of Excellence PRISMA⁺ (Project ID 39083149), for its hospitality and support. This work received support from the ERC-CoG Precision Gravity: From the LHC to LISA provided by the European Research Council (ERC) under the European Union's H2020 research and innovation programme (grant No. 817791). ZL is supported partially by DFF grant 1026-00077B, the Carlsberg foundation, and the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 847523 'INTERACTIONS'.

- R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA), GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run, (2021), arXiv:2111.03606 [gr-qc].
- [2] P. Amaro-Seoane *et al.* (LISA), Laser Interferometer Space Antenna, (2017), arXiv:1702.00786 [astroph.IM].
- [3] M. Punturo *et al.*, The einstein telescope: A third-generation gravitational wave observatory, Class. Quant. Grav. 27, 194002 (2010).
- [4] A. Buonanno and B. Sathyaprakash, Sources of Gravitational Waves: Theory and Observations, (2014), arXiv:1410.7832.
- [5] R. A. Porto, The Tune of Love and the Nature(ness) of Spacetime, Fortsch. Phys. 64, 723 (2016), arXiv:1606.08895.
- [6] R. A. Porto, The Music of the Spheres: The Dawn of Gravitational Wave Science, (2017), arXiv:1703.06440 [physics.pop-ph].

- [7] M. Maggiore *et al.*, Science Case for the Einstein Telescope, JCAP **03**, 050, arXiv:1912.02622 [astro-ph.CO].
- [8] E. Barausse *et al.*, Prospects for Fundamental Physics with LISA, Gen. Rel. Grav. **52**, 81, arXiv:2001.09793 [gr-qc].
- [9] S. Bernitt *et al.*, Fundamental Physics in the Gravitational-Wave Era, Nucl. Phys. News **32**, 16 (2022).
- [10] T. Damour, Introductory lectures on the Effective One Body formalism, Int. J. Mod. Phys. A 23, 1130 (2008), arXiv:0802.4047 [gr-qc].
- [11] L. Blanchet, Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries, Living Rev.Rel. 17, 2 (2014), arXiv:1310.1528 [gr-qc].
- [12] G. Schäfer and P. Jaranowski, Hamiltonian formulation of general relativity and post-Newtonian dynamics of compact binaries, Living Rev. Rel. 21, 7 (2018), arXiv:1805.07240.
- [13] W. D. Goldberger, Les Houches lectures on effective field theories and gravitational radiation, in Les Houches Summer School - Session 86 (2007) arXiv:hepph/0701129.
- [14] I. Rothstein, Progress in Effective Field Theory Approach to the Binary Inspiral Problem, Gen. Rel. Grav. 46, 1726 (2014).
- [15] R. A. Porto, The effective field theorist's approach to gravitational dynamics, Phys. Rept. 633, 1 (2016), arXiv:1601.04914.
- [16] L. Barack and A. Pound, Self-force and radiation reaction in general relativity, Rept. Prog. Phys. 82, 016904 (2019), arXiv:1805.10385 [gr-qc].
- [17] A. Buonanno, M. Khalil, D. O'Connell, R. Roiban, M. P. Solon, and M. Zeng, Snowmass White Paper: Gravitational Waves and Scattering Amplitudes, in 2022 Snowmass Summer Study (2022) arXiv:2204.05194 [hep-th].
- [18] W. D. Goldberger, Effective field theories of gravity and compact binary dynamics: A Snowmass 2021 whitepaper, (2022), arXiv:2206.14249 [hep-th].
- [19] P. Ajith *et al.*, The NINJA-2 catalog of hybrid post-Newtonian/numerical-relativity waveforms for nonprecessing black-hole binaries, Class. Quant. Grav. 29, 124001 (2012), [Erratum: Class.Quant.Grav. 30, 199401 (2013)], arXiv:1201.5319 [gr-qc].
- [20] B. Szilágyi, J. Blackman, A. Buonanno, A. Taracchini, H. P. Pfeiffer, M. A. Scheel, T. Chu, L. E. Kidder, and Y. Pan, Approaching the Post-Newtonian Regime with Numerical Relativity: A Compact-Object Binary Simulation Spanning 350 Gravitational-Wave Cycles, Phys. Rev. Lett. **115**, 031102 (2015), arXiv:1502.04953 [gr-qc].
- [21] T. Dietrich *et al.*, CoRe database of binary neutron star merger waveforms, Class. Quant. Grav. **35**, 24LT01 (2018), arXiv:1806.01625 [gr-qc].
- [22] D. Bini and T. Damour, Gravitational radiation reaction along general orbits in the effective onebody formalism, Phys. Rev. D 86, 124012 (2012), arXiv:1210.2834 [gr-qc].
- [23] T. Damour, Gravitational scattering, post-Minkowskian approximation and Effective One-Body theory, Phys. Rev. D94, 104015 (2016), arXiv:1609.00354.
- [24] T. Damour, High-energy gravitational scattering and the general relativistic two-body problem, Phys. Rev. D97, 044038 (2018), arXiv:1710.10599.
- [25] T. Damour, Classical and quantum scattering in post-

Minkowskian gravity, Phys. Rev. D **102**, 024060 (2020), arXiv:1912.02139 [gr-qc].

- [26] T. Damour, Radiative contribution to classical gravitational scattering at the third order in G, Phys. Rev. D 102, 124008 (2020), arXiv:2010.01641 [gr-qc].
- [27] D. Bini, T. Damour, and A. Geralico, Radiative contributions to gravitational scattering, Phys. Rev. D 104, 084031 (2021), arXiv:2107.08896 [gr-qc].
- [28] G. Kälin and R. A. Porto, From Boundary Data to Bound States, JHEP 01, 072, arXiv:1910.03008.
- [29] G. Kälin and R. A. Porto, From boundary data to bound states. Part II: Scattering angle to dynamical invariants (with twist), JHEP 02, 120, arXiv:1911.09130.
- [30] G. Cho, G. Kälin, and R. A. Porto, From Boundary Data to Bound States III: Radiative Effects, (2021), arXiv:2112.03976 [hep-th].
- [31] V. A. Smirnov, Analytic tools for Feynman integrals (Springer, 2012).
- [32] A. Kotikov, Differential equation method: The Calculation of N point Feynman diagrams, Phys. Lett. B 267, 123 (1991), [Erratum: Phys.Lett.B 295, 409–409 (1992)].
- [33] E. Remiddi, Differential equations for Feynman graph amplitudes, Nuovo Cim. A 110, 1435 (1997), arXiv:hepth/9711188.
- [34] J. M. Henn, Multiloop integrals in dimensional regularization made simple, Phys. Rev. Lett. 110, 251601 (2013), arXiv:1304.1806.
- [35] M. Prausa, epsilon: A tool to find a canonical basis of master integrals, Comput. Phys. Commun. 219, 361 (2017), arXiv:1701.00725 [hep-ph].
- [36] R. N. Lee, Libra: A package for transformation of differential systems for multiloop integrals, Comput. Phys. Commun. 267, 108058 (2021), arXiv:2012.00279 [hepph].
- [37] R. N. Lee, Reducing differential equations for multiloop master integrals, JHEP 04, 108, arXiv:1411.0911 [hepph].
- [38] L. Adams and S. Weinzierl, The ε -form of the differential equations for Feynman integrals in the elliptic case, Phys. Lett. B **781**, 270 (2018), arXiv:1802.05020 [hepph].
- [39] K. Chetyrkin and F. Tkachov, Integration by Parts: The Algorithm to Calculate beta Functions in 4 Loops, Nucl. Phys. B 192, 159 (1981).
- [40] F. Tkachov, A Theorem on Analytical Calculability of Four Loop Renormalization Group Functions, Phys. Lett. B 100, 65 (1981).
- [41] A. V. Smirnov and F. S. Chuharev, FIRE6: Feynman Integral REduction with Modular Arithmetic, Comput. Phys. Commun. 247, 106877 (2020), arXiv:1901.07808 [hep-ph].
- [42] A. Smirnov and V. Smirnov, How to choose master integrals, Nucl. Phys. B 960, 115213 (2020), arXiv:2002.08042 [hep-ph].
- [43] R. Lee, Presenting LiteRed: a tool for the Loop InTEgrals REDuction, (2012), arXiv:1212.2685.
- [44] R. N. Lee, LiteRed 1.4: a powerful tool for reduction of multiloop integrals, J. Phys. Conf. Ser. 523, 012059 (2014), arXiv:1310.1145 [hep-ph].
- [45] M. Beneke and V. A. Smirnov, Asymptotic expansion of Feynman integrals near threshold, Nucl. Phys. B522, 321 (1998), arXiv:hep-ph/9711391.
- [46] B. Jantzen, A. V. Smirnov, and V. A. Smirnov, Ex-

pansion by regions: revealing potential and Glauber regions automatically, Eur. Phys. J. C **72**, 2139 (2012), arXiv:1206.0546 [hep-ph].

- [47] A. V. Smirnov, FIESTA4: Optimized Feynman integral calculations with GPU support, Comput. Phys. Commun. 204, 189 (2016), arXiv:1511.03614 [hep-ph].
- [48] C. Meyer, Evaluating multi-loop Feynman integrals using differential equations: automatizing the transformation to a canonical basis, PoS LL2016, 028 (2016).
- [49] C. Meyer, Transforming differential equations of multiloop Feynman integrals into canonical form, JHEP 04, 006, arXiv:1611.01087 [hep-ph].
- [50] J. Broedel, C. Duhr, F. Dulat, R. Marzucca, B. Penante, and L. Tancredi, An analytic solution for the equal-mass banana graph, JHEP 09, 112, arXiv:1907.03787 [hepth].
- [51] A. Primo and L. Tancredi, Maximal cuts and differential equations for Feynman integrals. An application to the three-loop massive banana graph, Nucl. Phys. B 921, 316 (2017), arXiv:1704.05465 [hep-ph].
- [52] M. Hidding, DiffExp, a Mathematica package for computing Feynman integrals in terms of one-dimensional series expansions, Comput. Phys. Commun. 269, 108125 (2021), arXiv:2006.05510 [hep-ph].
- [53] A. Goncharov, Multiple polylogarithms and mixed Tate motives, (2001), arXiv:math/0103059.
- [54] K.-T. Chen, Iterated path integrals, Bull. Am. Math. Soc. 83, 831 (1977).
- [55] C. Duhr, Mathematical aspects of scattering amplitudes, (2014), arXiv:1411.7538 [hep-ph].
- [56] C. Duhr and F. Dulat, PolyLogTools polylogs for the masses, JHEP 08, 135, arXiv:1904.07279 [hep-th].
- [57] C. Dlapa, J. Henn, and K. Yan, Deriving canonical differential equations for Feynman integrals from a single uniform weight integral, JHEP 05, 025, arXiv:2002.02340 [hep-ph].
- [58] A. V. Smirnov, N. D. Shapurov, and L. I. Vysotsky, FI-ESTA5: numerical high-performance Feynman integral evaluation, (2021), arXiv:2110.11660 [hep-ph].
- [59] R. N. Lee, A. V. Smirnov, V. A. Smirnov, and M. Steinhauser, Four-loop quark form factor with quartic fundamental colour factor, JHEP 02, 172, arXiv:1901.02898 [hep-ph].
- [60] J. Blümlein, Analytic Integration Methods in Quantum Field Theory: An Introduction, in Antidifferentiation and the Calculation of Feynman Amplitudes (2021) arXiv:2103.10652 [hep-th].
- [61] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, Bootstrapping the relativistic two-body problem, *To appear*.
- [62] G. Kälin and R. A. Porto, Post-Minkowskian Effective Field Theory for Conservative Binary Dynamics, JHEP 11, 106, arXiv:2006.01184 [hep-th].
- [63] G. Kälin, Z. Liu, and R. A. Porto, Conservative Dynamics of Binary Systems to Third Post-Minkowskian Order from the Effective Field Theory Approach, Phys. Rev. Lett. **125**, 261103 (2020), arXiv:2007.04977 [hep-th].
- [64] G. Kälin, Z. Liu, and R. A. Porto, Conservative Tidal Effects in Compact Binary Systems to Next-to-Leading Post-Minkowskian Order, Phys. Rev. D 102, 124025 (2020), arXiv:2008.06047 [hep-th].
- [65] Z. Liu, R. A. Porto, and Z. Yang, Spin Effects in the Effective Field Theory Approach to Post-Minkowskian Conservative Dynamics, JHEP 06, 012, arXiv:2102.10059 [hep-th].

- [66] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, Dynamics of Binary Systems to Fourth Post-Minkowskian Order from the Effective Field Theory Approach, Phys. Lett. B 831, 137203 (2022), 2106.08276 [hep-th].
- [67] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, Conservative Dynamics of Binary Systems at Fourth Post-Minkowskian Order in the Large-Eccentricity Expansion, Phys. Rev. Lett. **128**, 161104 (2022), arXiv:2112.11296 [hep-th].
- [68] G. Mogull, J. Plefka, and J. Steinhoff, Classical black hole scattering from a worldline quantum field theory, JHEP 02, 048, arXiv:2010.02865 [hep-th].
- [69] G. U. Jakobsen, G. Mogull, J. Plefka, and J. Steinhoff, Classical Gravitational Bremsstrahlung from a Worldline Quantum Field Theory, Phys. Rev. Lett. **126**, 201103 (2021), arXiv:2101.12688 [gr-qc].
- [70] G. U. Jakobsen and G. Mogull, Conservative and Radiative Dynamics of Spinning Bodies at Third Post-Minkowskian Order Using Worldline Quantum Field Theory, Phys. Rev. Lett. **128**, 141102 (2022), arXiv:2201.07778 [hep-th].
- [71] G. U. Jakobsen, G. Mogull, J. Plefka, and J. Steinhoff, SUSY in the sky with gravitons, JHEP 01, 027, arXiv:2109.04465 [hep-th].
- [72] S. Mougiakakos, M. M. Riva, and F. Vernizzi, Gravitational Bremsstrahlung in the post-Minkowskian effective field theory, Phys. Rev. D 104, 024041 (2021), arXiv:2102.08339 [gr-qc].
- [73] M. M. Riva and F. Vernizzi, Radiated momentum in the post-Minkowskian worldline approach via reverse unitarity, JHEP 11, 228, arXiv:2110.10140 [hep-th].
- [74] S. Mougiakakos, M. M. Riva, and F. Vernizzi, Gravitational Bremsstrahlung with tidal effects in the post-Minkowskian expansion, (2022), arXiv:2204.06556 [hep-th].
- [75] M. M. Riva, F. Vernizzi, and L. K. Wong, Gravitational Bremsstrahlung from Spinning Binaries in the Post-Minkowskian Expansion, (2022), arXiv:2205.15295 [hep-th].
- [76] G. Kälin, J. Neef, and R. A. Porto, Radiation-Reaction in the Effective Field Theory Approach to Post-Minkowskian Dynamics, (2022), arXiv:2207.00580 [hep-th].
- [77] G. U. Jakobsen, G. Mogull, J. Plefka, and B. Sauer, All Things Retarded: Radiation-Reaction in Worldline Quantum Field Theory, (2022), arXiv:2207.00569 [hepth].
- [78] R. Jinno, G. Kälin, Z. Liu, and H. Rubira, Machine Learning Post-Minkowskian Integrals, (2022), arXiv:2209.01091 [hep-th].
- [79] D. Amati, M. Ciafaloni, and G. Veneziano, Higher Order Gravitational Deflection and Soft Bremsstrahlung in Planckian Energy Superstring Collisions, Nucl. Phys. B 347, 550 (1990).
- [80] D. Neill and I. Z. Rothstein, Classical Space-Times from the S Matrix, Nucl. Phys. B877, 177 (2013), arXiv:1304.7263.
- [81] V. Vaidya, Gravitational spin Hamiltonians from the S matrix, Phys. Rev. D91, 024017 (2015), arXiv:1410.5348.
- [82] W. D. Goldberger and A. K. Ridgway, Radiation and the classical double copy for color charges, Phys. Rev. D 95, 125010 (2017), arXiv:1611.03493.
- [83] C. Cheung, I. Z. Rothstein, and M. P. Solon, From Scat-

tering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion, Phys. Rev. Lett. **121**, 251101 (2018), arXiv:1808.02489.

- [84] M. Ciafaloni, D. Colferai, and G. Veneziano, Infrared features of gravitational scattering and radiation in the eikonal approach, Phys. Rev. D 99, 066008 (2019), arXiv:1812.08137 [hep-th].
- [85] A. Guevara, A. Ochirov, and J. Vines, Scattering of Spinning Black Holes from Exponentiated Soft Factors, JHEP 09, 056, arXiv:1812.06895.
- [86] D. A. Kosower, B. Maybee, and D. O'Connell, Amplitudes, Observables, and Classical Scattering, JHEP 02, 137, arXiv:1811.10950.
- [87] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order, Phys. Rev. Lett. **122**, 201603 (2019), arXiv:1901.04424.
- [88] K. Haddad and A. Helset, Tidal effects in quantum field theory, JHEP 12, 024, arXiv:2008.04920 [hep-th].
- [89] A. Brandhuber, G. Chen, G. Travaglini, and C. Wen, Classical gravitational scattering from a gauge-invariant double copy, JHEP 10, 118, arXiv:2108.04216 [hep-th].
- [90] R. Aoude, K. Haddad, and A. Helset, Classical gravitational spinning-spinless scattering at $\mathcal{O}(G^2 S^{\infty})$, (2022), arXiv:2205.02809 [hep-th].
- [91] N. E. J. Bjerrum-Bohr, P. H. Damgaard, L. Planté, and P. Vanhove, The amplitude for classical gravitational scattering at third Post-Minkowskian order, JHEP 08, 172, arXiv:2105.05218 [hep-th].
- [92] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, Scattering Amplitudes and Conservative Binary Dynamics at $\mathcal{O}(G^4)$, Phys. Rev. Lett. **126**, 171601 (2021), arXiv:2101.07254 [hep-th].
- [93] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, Scattering Amplitudes, the Tail Effect, and Conservative Binary Dynamics at O(G4), Phys. Rev. Lett. **128**, 161103 (2022), arXiv:2112.10750 [hep-th].
- [94] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, The eikonal approach to gravitational scattering and radiation at $\mathcal{O}(G^3)$, JHEP **07**, 169, arXiv:2104.03256 [hep-th].
- [95] J. Parra-Martinez, M. S. Ruf, and M. Zeng, Extremal black hole scattering at $\mathcal{O}(G^3)$: graviton dominance, eikonal exponentiation, and differential equations, JHEP **11**, 023, arXiv:2005.04236 [hep-th].
- [96] F. Febres Cordero, M. Kraus, G. Lin, M. S. Ruf, and M. Zeng, Conservative Binary Dynamics with a Spinning Black Hole at \$\mathcal{O}(G^3)\$ from Scattering Amplitudes, (2022), arXiv:2205.07357 [hep-th].
- [97] E. Herrmann, J. Parra-Martinez, M. S. Ruf, and M. Zeng, Radiative Classical Gravitational Observables at $\mathcal{O}(G^3)$ from Scattering Amplitudes, (2021), arXiv:2104.03957 [hep-th].
- [98] A. V. Manohar, A. K. Ridgway, and C.-H. Shen, Radiated Angular Momentum and Dissipative Effects in Classical Scattering, (2022), arXiv:2203.04283 [hep-th].
- [99] M. Khalil, A. Buonanno, J. Steinhoff, and J. Vines, Energetics and scattering of gravitational two-body systems at fourth post-Minkowskian order, (2022), arXiv:2204.05047 [gr-qc].
- [100] S. Hopper, A. Nagar, and P. Rettegno, Strong-field scat-

tering of two spinning black holes: Numerics versus Analytics, (2022), arXiv:2204.10299 [gr-qc].

- [101] S. J. Kovacs and K. S. Thorne, The Generation of Gravitational Waves. 4. Bremsstrahlung, Astrophys. J. 224, 62 (1978).
- [102] W. D. Goldberger and I. Z. Rothstein, An Effective field theory of gravity for extended objects, Phys. Rev. D73, 104029 (2006), arXiv:hep-th/0409156.
- [103] C. R. Galley, Classical Mechanics of Nonconservative Systems, Phys. Rev. Lett. **110**, 174301 (2013), arXiv:1210.2745 [gr-qc].
- [104] C. R. Galley and M. Tiglio, Radiation reaction and gravitational waves in the effective field theory approach, Phys. Rev. D 79, 124027 (2009).
- [105] C. R. Galley, A. K. Leibovich, and I. Z. Rothstein, Finite size corrections to the radiation reaction force in classical electrodynamics, Phys. Rev. Lett. **105**, 094802 (2010), arXiv:1005.2617 [gr-qc].
- [106] C. R. Galley and A. K. Leibovich, Radiation reaction at 3.5 post-Newtonian order in effective field theory, Phys. Rev. D 86, 044029 (2012), arXiv:1205.3842 [gr-qc].
- [107] N. T. Maia, C. R. Galley, A. K. Leibovich, and R. A. Porto, Radiation reaction for spinning bodies in effective field theory I: Spin-orbit effects, Phys. Rev. D96, 084064 (2017), arXiv:1705.07934.
- [108] N. T. Maia, C. R. Galley, A. K. Leibovich, and R. A. Porto, Radiation reaction for spinning bodies in effective field theory II: Spin-spin effects, Phys. Rev. D96, 084065 (2017), arXiv:1705.07938.
- [109] C. Galley, A. Leibovich, R. A. Porto, and A. Ross, Tail Effect in Gravitational Radiation Reaction: Time Nonlocality and Renormalization Group Evolution, Phys. Rev. D 93, 124010 (2016), arXiv:1511.07379.
- [110] J. S. Schwinger, Brownian motion of a quantum oscillator, J. Math. Phys. 2, 407 (1961).
- [111] L. V. Keldysh, Diagram technique for nonequilibrium processes, Zh. Eksp. Teor. Fiz. 47, 1515 (1964).
- [112] E. Calzetta and B. L. Hu, Closed Time Path Functional Formalism in Curved Space-Time: Application to Cosmological Back Reaction Problems, Phys. Rev. D 35, 495 (1987).
- [113] E. Calzetta and B. L. Hu, Nonequilibrium Quantum Fields: Closed Time Path Effective Action, Wigner Function and Boltzmann Equation, Phys. Rev. D 37, 2878 (1988).
- [114] R. D. Jordan, Effective Field Equations for Expectation Values, Phys. Rev. D 33, 444 (1986).
- [115] S. Foffa and R. Sturani, Dynamics of the gravitational two-body problem at fourth post-Newtonian order and at quadratic order in the Newton constant, Phys. Rev. D 87, 064011 (2013), arXiv:1206.7087 [gr-qc].
- [116] S. Foffa, R. A. Porto, I. Rothstein, and R. Sturani, Conservative dynamics of binary systems to fourth Post-Newtonian order in the EFT approach II: Renormalized Lagrangian, Phys. Rev. D100, 024048 (2019), arXiv:1903.05118.
- [117] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach, (2021), arXiv:2110.13822 [gr-qc].
- [118] G. L. Almeida, S. Foffa, and R. Sturani, Gravitational radiation contributions to the two-body scattering angle, (2022), arXiv:2209.11594 [gr-qc].
- [119] G. Cho, S. Dandapat, and A. Gopakumar, Third or-

der post-Newtonian gravitational radiation from twobody scattering: Instantaneous energy and angular momentum radiation, Phys. Rev. D **105**, 084018 (2022), arXiv:2111.00818 [gr-qc].

- [120] G. Cho, Third post-Newtonian gravitational radiation from two-body scattering. II. Hereditary energy radiation, Phys. Rev. D 105, 104035 (2022), arXiv:2203.10872 [gr-qc].
- [121] D. Bini and A. Geralico, Higher-order tail contributions to the energy and angular momentum fluxes in a two-body scattering process, Phys. Rev. D 104, 104020 (2021), arXiv:2108.05445 [gr-qc].
- [122] D. Bini and A. Geralico, Momentum recoil in the relativistic two-body problem: Higher-order tails, Phys. Rev. D 105, 084028 (2022), arXiv:2202.03037 [gr-qc].
- [123] D. Bini and A. Geralico, Multipolar invariants and the eccentricity enhancement function parametrization of gravitational radiation, Phys. Rev. D 105, 124001 (2022), arXiv:2204.08077 [gr-qc].
- [124] D. Bini, T. Damour, and A. Geralico, Private Communication.
- [125] A. Gruzinov and G. Veneziano, Gravitational Radiation from Massless Particle Collisions, Class. Quant. Grav. 33, 125012 (2016), arXiv:1409.4555 [gr-qc].
- [126] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, The eikonal operator at arbitrary velocities I: the soft-radiation limit, JHEP 07, 039, arXiv:2204.02378 [hep-th].

SUPPLEMENTAL MATERIAL

We give explicit values for the conservative and dissipative contributions to the impulse as well as the (relative) scattering angle. Ready-to-use results can also be found in the ancillary file.

Impulse. For the conservative part we find

$$\begin{aligned} \frac{c_{1b}^{(4)\text{cons}}}{\pi} &= -\frac{3h_1m_1m_2\left(m_1^3 + m_2^3\right)}{64\left(\gamma^2 - 1\right)^{5/2}} + m_1^2m_2^2\left(m_1 + m_2\right) \left[\frac{2!\sqrt{\gamma^2 - 1}h_2\text{E}^2\left(\frac{\gamma - 1}{\gamma + 1}\right)}{32(\gamma - 1)^2(\gamma + 1)} + \frac{3h_3\text{K}^2\left(\frac{\gamma - 1}{\gamma + 1}\right)}{16\left(\gamma^2 - 1\right)^{3/2}} - \frac{3h_4\text{E}\left(\frac{\gamma - 1}{\gamma + 1}\right)}{16\left(\gamma^2 - 1\right)^{3/2}} + \frac{\pi^2h_5}{8\sqrt{\gamma^2 - 1}} \\ &+ \frac{h_6\log\left(\frac{\gamma - 1}{2}\right)}{16\left(\gamma^2 - 1\right)^{3/2}} + \frac{3h_7\text{Li}_2\left(\sqrt{\frac{\gamma - 1}{\gamma + 1}}\right)}{(\gamma - 1)(\gamma + 1)^2} - \frac{3h_7\text{Li}_2\left(\frac{\gamma - 1}{\gamma + 1}\right)}{4(\gamma - 1)(\gamma + 1)^2} - \frac{3h_{15}\log\left(\frac{\gamma - 1}{2}\right)\log\left(\frac{\gamma + 1}{2}\right)}{8\sqrt{\gamma^2 - 1}} + \frac{3h_{16}\log\left(\frac{\gamma - 1}{2}\right)\arctan\left(\gamma\right)}{16\left(\gamma^2 - 1\right)^2} + \frac{h_{20}}{192\gamma^7\left(\gamma^2 - 1\right)^{5/2}} \\ &+ \frac{h_{22}\log\left(\frac{\gamma + 1}{2}\right)}{16\left(\gamma^2 - 1\right)^{3/2}} + \frac{h_{23}\log(\gamma)}{2\left(\gamma^2 - 1\right)^{3/2}} - \frac{h_{24}\arccos(\gamma)}{16\left(\gamma^2 - 1\right)^3} - \frac{3h_{26}\arccos(\gamma)}{32\left(\gamma^2 - 1\right)^{7/2}} + \frac{3h_{27}\log^2\left(\frac{\gamma + 1}{2}\right)}{2\sqrt{\gamma^2 - 1}} + \frac{3h_{28}\log\left(\frac{\gamma + 1}{2}\right) \arccos(\gamma)}{16\left(\gamma^2 - 1\right)^2} + \frac{h_{29}\text{Li}_2\left(\frac{1 - \gamma}{\gamma + 1}\right)}{4\sqrt{\gamma^2 - 1}} \\ &+ \frac{3\sqrt{\gamma^2 - 1}h_{30}\text{Li}_2\left(\frac{\gamma - 1}{\gamma + 1}\right)}{16\left(\gamma^2 - 1\right)^{3/2}} - \frac{h_{24}\arccos(\gamma)}{16\left(\gamma^2 - 1\right)^3} - \frac{3h_{26}\arccos(\gamma)}{32\left(\gamma^2 - 1\right)^{7/2}} + \frac{3h_{27}\log\left(\frac{\gamma + 1}{2}\right)}{2\sqrt{\gamma^2 - 1}} + \frac{3h_{28}\log\left(\frac{\gamma + 1}{2}\right)\operatorname{arccosh}(\gamma)}{16\left(\gamma^2 - 1\right)^2} + \frac{h_{29}\text{Li}_2\left(\frac{1 - \gamma}{\gamma + 1}\right)}{4\sqrt{\gamma^2 - 1}} \\ &+ \frac{3\sqrt{\gamma^2 - 1}h_{30}\text{Li}_2\left(\frac{\gamma - 1}{\gamma + 1}\right)}{16\left(\gamma^2 - 1\right)^3} - \frac{h_{24}\arccos(\gamma)}{16\left(\gamma^2 - 1\right)^3} - \frac{3h_{26}\operatorname{arccosh}(\gamma)}{32\left(\gamma^2 - 1\right)} + \frac{3h_{28}\log\left(\frac{\gamma + 1}{2}\right)}{16\left(\gamma^2 - 1\right)^2} + \frac{h_{29}\text{Li}_2\left(\frac{1 - \gamma}{\gamma + 1}\right)}{4\sqrt{\gamma^2 - 1}} \\ &+ \frac{3\sqrt{\gamma^2 - 1}h_{30}\text{Li}_2\left(\frac{\gamma - 1}{\gamma + 1}\right)}{16\left(\gamma^2 - 1\right)^2} - \frac{2h_{32}m_1m_2^2\left(m_1^2 + m_2^2\right)}{(\gamma^2 - 1)^3} + m_1^2m_2^2\left[\frac{4h_{33}}{3\left(\gamma^2 - 1\right)^3} - \frac{16h_{36}\operatorname{arccosh}(\gamma)}{(\gamma^2 - 1)^{3/2}}\right] \\ &+ \frac{3\sqrt{\gamma^2 - 1}h_{32}\left(m_1 + m_2\right)^2}{32\left(\gamma^2 - 1\right)} - \frac{2h_{32}m_1m_2^2\left(m_1^2 + m_2^2\right)}{(\gamma^2 - 1)^3} + m_1^3m_2^2\left[-\frac{4h_{33}}{3\left(\gamma^2 - 1\right)^3} + \frac{16h_{36}\operatorname{arccosh}(\gamma)}{(\gamma^2 - 1)^{3/2}}\right] \\ &+ \frac{3h_{32}\left(m_1^2 - 1\right)^2}{32\left(\gamma^2 - 1\right)} - \frac{2h_{32}m_1m_2^2\left(m_1^2 + m_2^2\right)}{(\gamma^2 - 1)^3} + m_1^3m_2^2\left[-\frac{4h_{33}}{3\left(\gamma^2 - 1\right)^3}\right] + \frac{3h_{32}\left(m_1^2 - 1\right)^2}{(\gamma^2 - 1)^{3/$$

which reproduces the result in [66, 67]. For the dissipative part we split the result into two terms. For the (instantaneous) contribution involving a single radiation mode we have

$$\begin{aligned} \frac{c_{15,1rad}^{c}}{\pi} &= m_1^3 m_2^2 \bigg[\frac{h_8}{48 (\gamma^2 - 1)^3} + \frac{h_{10} \log \left(\frac{\gamma + 1}{2}\right)}{8 (\gamma^2 - 1)^2} + \frac{h_{14} \operatorname{arccosh}(\gamma)}{16 (\gamma^2 - 1)^{7/2}} \bigg] + m_1^2 m_2^3 \bigg[\frac{h_{19}}{48 (\gamma^2 - 1)^3} + \frac{h_{21} \log \left(\frac{\gamma + 1}{2}\right)}{8 (\gamma^2 - 1)^2} + \frac{h_{25} \operatorname{arccosh}(\gamma)}{16 (\gamma^2 - 1)^{7/2}} \bigg], \\ c_{141,1rad}^{(4)diss} &= m_1^3 m_2^2 \bigg[-\frac{8h_{34}}{3 (\gamma^2 - 1)^{5/2}} + \frac{8h_{35} \operatorname{arccosh}(\gamma)}{(\gamma^2 - 1)^3} \bigg], \\ c_{142,1rad}^{(4)diss} &= m_1^3 m_2^2 \bigg[\frac{h_{38}}{705600\gamma^8 (\gamma^2 - 1)^{5/2}} + \frac{\pi^2 \left(108h_{31}(\gamma^2 - 1) + h_{39}\right)}{192 (\gamma^2 - 1)^2} + \frac{h_{40} \operatorname{arccosh}(\gamma)}{6720\gamma^9 (\gamma^2 - 1)^3} + \frac{32h_{43} \operatorname{arccosh}^2(\gamma)}{(\gamma^2 - 1)^{7/2}} + \frac{h_{44} \log(2) \operatorname{arccosh}(\gamma)}{8 (\gamma^2 - 1)^2} \\ &\quad + \frac{3h_{45} \left(\operatorname{Li}_2 \left(\frac{\gamma - 1}{(\gamma + 1)} - 4\operatorname{Li}_2 \left(\sqrt{\frac{\gamma - 1}{(\gamma + 1)}}\right)\right)}{16 (\gamma^2 - 1)^2} + \frac{3h_{46} \left(\log \left(\frac{\gamma + 1}{2}\right) \operatorname{arccosh}(\gamma) - 2\operatorname{Li}_2 \left(\sqrt{\gamma^2 - 1} - \gamma\right)\right)}{8 (\gamma^2 - 1)^2} \\ &\quad - \frac{h_{47} \left(\operatorname{Li}_2 \left(-\left(\gamma - \sqrt{\gamma^2 - 1}\right)^2\right) - 2\log(\gamma) \operatorname{arccosh}(\gamma)}{16 (\gamma^2 - 1)^2} + \frac{8h_{60} \operatorname{arccosh}^2(\gamma)}{(\gamma^2 - 1)^2}\right)}{16 (\gamma^2 - 1)^2} + \frac{8h_{60} \operatorname{arccosh}^2(\gamma)}{(\gamma^2 - 1)^2} + \frac{8h_{60} \operatorname{arccosh}(\gamma)}{(\gamma^2 - 1)^2} \\ &\quad + \frac{\pi^2 \left(27h_{31}(\gamma^2 - 1) + 2h_{50}\right)}{96 (\gamma^2 - 1)^2} + \frac{h_{51} \operatorname{arccosh}(\gamma)}{480\gamma^8 (\gamma^2 - 1)^3} - \frac{32h_{54} \operatorname{arccosh}^2(\gamma)}{(\gamma^2 - 1)^{7/2}} - \frac{h_{55} \log(2) \operatorname{arccosh}(\gamma)}{4 (\gamma^2 - 1)^2} + \frac{h_{56} \operatorname{Li}_2 \left(\frac{\gamma - 1}{\gamma + 1}\right)}{32 (\gamma^2 - 1)^2} \\ &\quad - \frac{h_{57} \left(\log \left(\frac{\gamma + 1}{2}\right) \operatorname{arccosh}(\gamma) - 2\operatorname{Li}_2 \left(\sqrt{\gamma^2 - 1} - \gamma\right)\right)}{4 (\gamma^2 - 1)^2} + \frac{h_{58} \left(\operatorname{Li}_2 \left(-\left(\gamma - \sqrt{\gamma^2 - 1}\right)^2\right) - 2\log(\gamma) \operatorname{arccosh}(\gamma)}{8 (\gamma^2 - 1)^2} + \frac{h_{56} \operatorname{arccosh}^2(\gamma)}{8 (\gamma^2 - 1)^2} - \frac{h_{56} \operatorname{arccosh}^2(\gamma)}{(\gamma^2 - 1)^2} - \frac{h_{56} \operatorname{secsh}(\gamma)}{8 (\gamma^2 - 1)^2} + \frac{h_{56} \operatorname{secsh}^2(\gamma)}{(\gamma^2 - 1)^2} - \frac{h_{56} \operatorname{secsh}(\gamma)}{(\gamma^2 - 1)^2} + \frac{h_{56} \operatorname{secsh}^2(\gamma)}{(\gamma^2 - 1)^2} - \frac{h_{56} \operatorname{secsh}(\gamma)}{(\gamma^2 - 1)^2} + \frac{h_{56} \operatorname{secsh}^2(\gamma)}{(\gamma^2 - 1)^2} - \frac{h_{56} \operatorname{secsh}(\gamma)}{(\gamma^2 - 1)^2} + \frac{h_{56} \operatorname{secsh}^2(\gamma)}{(\gamma^2 - 1)^2} - \frac{h_{56} \operatorname{secsh}(\gamma)}{(\gamma^2 - 1)^2} + \frac{h_{56} \operatorname{secsh}^2(\gamma)}{(\gamma^2 - 1)^2} + \frac{h_{56} \operatorname{se$$

whereas the contribution involving two radiation modes going on-shell is given by

$$\frac{c_{1b,2rad}^{(4)diss}}{\pi} = m_1^3 m_2^2 \left[\frac{\sqrt{\gamma^2 - 1} \left(h_9 - 4\gamma^2(\gamma + 1)h_{20}\right)}{768(\gamma - 1)^3\gamma^9(\gamma + 1)^4} + \frac{\log(\gamma) \left(h_{12} - 8\left(\gamma^2 - 1\right)h_{23}\right)}{16\left(\gamma^2 - 1\right)^{5/2}} + \frac{\arccos(\gamma) \left((\gamma + 1)h_{24} - 2(\gamma - 1)^2h_{13}\right)}{16(\gamma - 1)^3(\gamma + 1)^4} + \frac{3h_{26} \operatorname{arccosh}^2(\gamma)}{32\left(\gamma^2 - 1\right)^{3/2}} + \frac{3\left(h_{15} - 4h_{27}\right)\log^2\left(\frac{\gamma + 1}{2}\right)}{8\sqrt{\gamma^2 - 1}} + \log\left(\frac{\gamma + 1}{2}\right)\left(\frac{-2\left(\gamma^2 - 1\right)h_{22} - h_{11}}{32\left(\gamma^2 - 1\right)^{5/2}} - \frac{3\left(h_{16} + h_{28}\right)\operatorname{arccosh}(\gamma)}{16\left(\gamma^2 - 1\right)^2}\right) + \frac{\left(-3\left(\gamma^2 - 1\right)h_{18} - 8h_{29}\right)\operatorname{Li}_2\left(\frac{1 - \gamma}{\gamma + 1}\right)}{32\sqrt{\gamma^2 - 1}} - \frac{3\left(h_{17} + 8h_{30}\right)\operatorname{Li}_2\left(\frac{\gamma - 1}{\gamma + 1}\right)}{64\sqrt{\gamma^2 - 1}} \right], \quad (16)$$

$$\begin{aligned} r_{1\dot{u}_{1},2\text{rad}}^{(4)\text{diss}} &= m_{1}^{3}m_{2}^{2} \bigg[\frac{4\left(h_{33}-h_{37}\right)}{3\left(\gamma^{2}-1\right)^{3}} - \frac{16\left(3h_{36}-2h_{41}\right)\operatorname{arccosh}(\gamma)}{3\left(\gamma^{2}-1\right)^{3/2}} - \frac{8\left(h_{42}+h_{60}\right)\operatorname{arccosh}^{2}(\gamma)}{\left(\gamma^{2}-1\right)^{2}} \bigg] \\ &+ m_{1}^{2}m_{2}^{3} \bigg[\frac{2\left(45h_{32}-h_{48}\right)}{45\left(\gamma^{2}-1\right)^{3}} - \frac{16h_{52}\operatorname{arccosh}(\gamma)}{5\left(\gamma^{2}-1\right)^{3/2}} - \frac{16\left(h_{53}+h_{59}\right)\operatorname{arccosh}^{2}(\gamma)}{\left(\gamma^{2}-1\right)^{2}} \bigg] \end{aligned}$$

We find perfect agreement with various partial results reported in [26, 27, 119–124].

Scattering angle. By performing a Lorentz boost using the value of the total recoil we can derive the relative spacelike impulse, Δp , via (see [26])

$$\Delta \boldsymbol{p} = \Delta \boldsymbol{p}_1 + \frac{E_1}{E} \boldsymbol{P}_{\text{rad}} + \mathcal{O}(\boldsymbol{P}_{\text{rad}}^2).$$
(17)

From here we can obtain a relative scattering angle, given by (recall $j \equiv p_{\infty} b/(GM^2 \nu)$)

$$\frac{\chi_{\text{rel}}}{2} \equiv \frac{1}{2} \arccos\left(\frac{\boldsymbol{p}_{+} \cdot \boldsymbol{p}_{-}}{|\boldsymbol{p}_{-}||\boldsymbol{p}_{+}|}\right) = \sum_{n=1}^{\infty} \chi_{b,\text{rel}}^{(n)} \left(\frac{GM}{b}\right)^{n} = \sum_{n=1}^{\infty} \frac{\chi_{j,\text{rel}}^{(n)}}{j^{n}},\tag{18}$$

with p_{-} and $p_{+} \equiv p_{-} + \Delta p$, the relative incoming/outgoing three-momentum at infinity.

We find, in impact parameter space,

$$\chi_{b,\text{rel}}^{(4)}(\gamma) = \chi_{b,\text{rel}}^{(4)\text{cons}}(\gamma) + \delta\chi_{b,\text{rel}}^{(4)\text{rr}}(\gamma)$$
(19)

$$\delta\chi_{b,\text{rel}}^{(4)\text{rr}}(\gamma) = \chi_{b,\text{rel}}^{(4)\text{1rad}}(\gamma) + \chi_{b,\text{rel}}^{(4)\text{2rad}}(\gamma), \qquad (20)$$

where

$$\frac{\chi_{b,\text{rel}}^{(4)\text{cons}}(\gamma)}{\pi\Gamma} = \frac{3h_{61}}{128\left(\gamma^2 - 1\right)^3} + \nu \left[-\frac{3h_3 \text{K}^2\left(\frac{\gamma-1}{\gamma+1}\right)}{32\left(\gamma^2 - 1\right)^2} + \frac{3h_4 \text{E}\left(\frac{\gamma-1}{\gamma+1}\right) \text{K}\left(\frac{\gamma-1}{\gamma+1}\right)}{32\left(\gamma^2 - 1\right)^2} + \frac{\pi^2 h_5}{16\left(1 - \gamma^2\right)} + \frac{3h_{27} \log^2\left(\frac{\gamma+1}{2}\right)}{4\left(1 - \gamma^2\right)} - \frac{h_6 \log\left(\frac{\gamma-1}{2}\right)}{32\left(\gamma^2 - 1\right)^2} + \frac{3h_{15} \log\left(\frac{\gamma-1}{2}\right) \log\left(\frac{\gamma+1}{2}\right)}{16\left(\gamma^2 - 1\right)} - \frac{h_{22} \log\left(\frac{\gamma+1}{2}\right)}{32\left(\gamma^2 - 1\right)^2} - \frac{h_{23} \log(\gamma)}{4\left(\gamma^2 - 1\right)^2} + \frac{3h_{26} \arccos^2(\gamma)}{64\left(\gamma^2 - 1\right)^4} + \frac{h_{24} \operatorname{accosh}(\gamma)}{32\left(\gamma^2 - 1\right)^{7/2}} - \frac{3h_{16} \log\left(\frac{\gamma-1}{2}\right) \operatorname{accosh}(\gamma)}{32\left(\gamma^2 - 1\right)^{5/2}} - \frac{3h_{28} \log\left(\frac{\gamma+1}{2}\right) \operatorname{accosh}(\gamma)}{32\left(\gamma^2 - 1\right)^{5/2}} - \frac{h_{62}}{344\gamma^7 \left(\gamma^2 - 1\right)^3} - \frac{21h_2 \text{E}^2\left(\frac{\gamma-1}{\gamma+1}\right)}{64\left(\gamma - 1\right)^2\left(\gamma + 1\right)} - \frac{3\sqrt{\gamma^2 - 1}h_7 \text{Li}_2\left(\sqrt{\frac{\gamma-1}{\gamma+1}}\right)}{2\left(\gamma - 1\right)^2\left(\gamma + 1\right)^3} + \frac{h_{29} \text{Li}_2\left(\frac{1-\gamma}{\gamma+1}\right)}{8\left(1 - \gamma^2\right)} + \left(\frac{3\sqrt{\gamma^2 - 1}h_7}{8\left(\gamma - 1\right)^2\left(\gamma + 1\right)^3} + \frac{3h_{30}}{16 - 16\gamma^2}\right) \text{Li}_2\left(\frac{\gamma - 1}{\gamma+1}\right)\right],$$

$$(21)$$

which agrees with the Feynman-only part computed in [66, 67] (including potential and tail terms), and

$$\frac{\Gamma\chi_{b,\text{rel}}^{(4)\text{Irad}}(\gamma)}{\pi\nu} = \frac{h_{64}}{96\left(\gamma^2 - 1\right)^{7/2}} + \frac{h_{65}\log\left(\frac{\gamma+1}{2}\right)}{16\left(\gamma^2 - 1\right)^{5/2}} + \frac{h_{63}\operatorname{arcsinh}\left(\frac{\sqrt{\gamma-1}}{\sqrt{2}}\right)}{8\left(\gamma^2 - 1\right)^4} - \frac{h_{25}\operatorname{arccosh}(\gamma)}{32\left(\gamma^2 - 1\right)^4} + \frac{h_{66}\operatorname{arcsinh}\left(\frac{\sqrt{\gamma-1}}{\sqrt{2}}\right)}{16\left(\gamma^2 - 1\right)^{5/2}} + \frac{h_{63}\log\left(\frac{\gamma+1}{2}\right)}{16\left(\gamma^2 - 1\right)^{5/2}} - \frac{\operatorname{arccosh}(\gamma)\left((\gamma+1)h_{14}+(\gamma-3)h_{25}\right)}{32\left(\gamma^2 - 1\right)^4} + \frac{h_{66}\operatorname{arcsinh}\left(\frac{\sqrt{\gamma-1}}{\sqrt{2}}\right)}{8\left(\gamma-1\right)^2\left(\gamma+1\right)^4}\right], \\ \frac{\Gamma\chi_{b,\text{rel}}^{(4)2\text{rad}}(\gamma)}{\pi\nu^2} = \frac{\log\left(\frac{\gamma+1}{2}\right)\left(2\left(\gamma^2 - 1\right)h_{22} + h_{11}\right)}{64(\gamma-1)^3(\gamma+1)^2} - \frac{\log(\gamma)\left(h_{12} - 8\left(\gamma^2 - 1\right)h_{23}\right)}{32(\gamma-1)^3(\gamma+1)^2} + \frac{\operatorname{arccosh}(\gamma)\left(2(\gamma-1)^2h_{13}-(\gamma+1)h_{24}\right)}{32\left(\gamma^2 - 1\right)^{7/2}} + \frac{3\sqrt{\gamma^2 - 1}\left(h_{16} + h_{28}\right)\log\left(\frac{\gamma+1}{2}\right)\operatorname{arccosh}(\gamma)}{32(\gamma-1)^3(\gamma+1)^2} - \frac{h_9 - 4\gamma^2(\gamma+1)h_{20}}{1536\gamma^9\left(\gamma^2 - 1\right)^3} - \frac{3\left(h_{15} - 4h_{27}\right)\log^2\left(\frac{\gamma+1}{2}\right)}{16(\gamma-1)} + \frac{3h_{26}\operatorname{arccosh}^2(\gamma)}{64(\gamma-1)^4(\gamma+1)^3} + \left(\frac{3}{64}(\gamma+1)h_{18} + \frac{h_{29}}{8(\gamma-1)}\right)\operatorname{Li}_2\left(\frac{1-\gamma}{\gamma+1}\right) + \frac{3\left(h_{17} + 8h_{30}\right)\operatorname{Li}_2\left(\frac{\gamma-1}{\gamma+1}\right)}{128(\gamma-1)}, \end{aligned}$$

where (for the sake of notation) we have introduced a few additional polynomials:

$$\begin{split} h_{61} &= 35(\gamma - 1)(\gamma + 1) \left(33\gamma^4 - 18\gamma^2 + 1 \right) \\ h_{62} &= 3600\gamma^{16} + 4320\gamma^{15} - 35360\gamma^{14} + 33249\gamma^{13} + 27952\gamma^{12} - 25145\gamma^{11} - 15056\gamma^{10} - 32177\gamma^9 + 64424\gamma^8 - 38135\gamma^7 + 13349\gamma^6 - 1471\gamma^4 \\ &\quad + 207\gamma^2 - 45 \\ h_{63} &= \gamma^2 \left(2\gamma^2 - 3 \right) \left(2\gamma^2 - 1 \right) \left(35\gamma^4 - 30\gamma^2 + 11 \right) \\ h_{64} &= -4140\gamma^8 + 702\gamma^7 + 15018\gamma^6 - 8491\gamma^5 - 9366\gamma^4 + 10052\gamma^3 - 6210\gamma^2 + 2681\gamma - 102 \\ h_{65} &= 210\gamma^7 - 240\gamma^6 - 755\gamma^5 + 216\gamma^4 + 1200\gamma^3 - 508\gamma^2 - 295\gamma + 124 \\ h_{66} &= \gamma \left(2\gamma^2 - 3 \right) \left(2\gamma^2 - 1 \right) \left(35\gamma^4 - 30\gamma^2 + 11 \right) \\ h_{67} &= - \left((\gamma - 1) \left(420\gamma^9 + 7596\gamma^8 - 2040\gamma^7 - 30840\gamma^6 + 21667\gamma^5 + 18929\gamma^4 - 26710\gamma^3 + 14910\gamma^2 - 3177\gamma - 947 \right) \right) \\ h_{68} &= (\gamma - 1) \left(490\gamma^7 - 290\gamma^6 - 1725\gamma^5 + 189\gamma^4 + 2632\gamma^3 - 952\gamma^2 - 661\gamma + 253 \right) \end{split}$$

Due to the different properties under time reversal, the distinction between the two radiation-reaction terms is straightforward. This is also manifest after performing a PN expansion in small $v_{\infty}^2 \ll 1$. In order to compare with

the literature it is convenient to transform to j space. We find at $\mathcal{O}(G^4)$:

$$\frac{\delta\chi_{j,\mathrm{rel}}^{(4)\mathrm{rr}}(\gamma)}{\pi} = \frac{121\nu v_{\infty}}{30} + \left(\frac{23111\nu}{1680} - \frac{437\nu^2}{60}\right)v_{\infty}^3 + \left(\frac{511\nu^3}{48} - \frac{75253\nu^2}{3360} + \frac{44759\nu}{2240}\right)v_{\infty}^5 + \frac{1491}{400}\nu^2 v_{\infty}^6 + \left(-\frac{455\nu^4}{32} + \frac{26367\nu^3}{896} - \frac{288007\nu^2}{10080} + \frac{1350131\nu}{1182720}\right)v_{\infty}^7 + \mathcal{O}\left(v_{\infty}^8\right),$$

$$(24)$$

in complete agreement with the result in Table XI of [27], except for the $\mathcal{O}(\nu^2 v_{\infty}^6)$ term. Likewise, we can PN expand the full result using the same convention as in Eq. (9.2) of [27]. Keeping only even-in-velocity terms, we find

$$\frac{\Gamma^{3}\chi_{j,\mathrm{rel}}^{(4)}(\gamma) - \chi_{j,\mathrm{Sch}}^{(4)}(\gamma)}{\pi} \bigg|_{\mathrm{even}} = -\frac{15\nu}{4} + \left(\frac{123\pi^{2}\nu}{256} - \frac{557\nu}{16}\right)v_{\infty}^{2} + \left(\frac{33601\pi^{2}\nu}{16384} - \frac{6113\nu}{96} - \frac{37}{5}\nu\log\left(v_{\infty}/2\right)\right)v_{\infty}^{4} \\
+ \left(\frac{1491\nu^{2}}{400} + \frac{93031\pi^{2}\nu}{32768} - \frac{615581\nu}{19200} - \frac{1357}{280}\nu\log\left(v_{\infty}/2\right)\right)v_{\infty}^{6} + \mathcal{O}\left(v_{\infty}^{8}\right),$$
(25)

in agreement with Eq. (9.2) except for the same $\mathcal{O}(\nu^2 v_{\infty}^6)$ contribution, which first enters at 5PN order. (The expression in (25) also disagrees with the two distinct results given in [117, 118].)

We will discuss the implications in more detail elsewhere. Let us, however, add a few remarks. First of all, although one could in principle incorporate these extra terms into a relative Hamiltonian, it is not entirely clear from our calculation alone which portion corresponds to nonlinear gravitational radiation-reaction forces (and/or hereditary terms), which are evaluated on the unperturbed (conservative) solution; or instead produced by effects at second order in the linear radiation-reaction force, and thus evaluated on the leading non-conservative trajectory. While we could in principle include the former the latter clearly does not belong there. Secondly, these extra terms radiate linear momentum, see (10). Therefore, they are only 'conservative' from the point of view of the relative motion. Finally, following [27] we have used the relative impulse to obtain the deflection angle. Although the full analysis in [27] is valid at linear order in the radiation-reaction force, we have only used the Lorentz boost to the outgoing center-of-mass frame, see (17). In principle, this holds up to $\mathcal{O}(P_{\rm rad}^2)$, namely 6PM order. However, it is still possible that (17) may have to be modified to fully incorporate non-linear radiation-reaction effects in the relative motion.