# Large sample behavior of the least trimmed squares estimator

Yijun Zuo

Department of Statistics and Probability

Michigan State University, East Lansing, MI 48824

zuo@msu.edu

October 29, 2024

### Abstract

The least trimmed squares (LTS) estimator is popular in location, regression, machine learning, and AI literature. Despite the empirical version of least trimmed squares (LTS) being repeatedly studied in the literature, the population version of the LTS has never been introduced and studied. The lack of the population version hinders the study of the large sample properties of the LTS utilizing the empirical process theory.

Novel properties of the objective function in both empirical and population settings of the LTS and other properties, are established for the first time in this article. The primary properties of the objective function facilitate the establishment of other original results, including the influence function and Fisher consistency. The strong consistency is established with the help of a generalized Glivenko-Cantelli Theorem over a class of functions for the first time. Differentiability and stochastic equicontinuity promote the establishment of asymptotic normality with a concise and novel approach.

AMS 2000 Classification: Primary 62J05, 62G36; Secondary 62J99, 62G99

Key words and phrase: trimmed squares of residuals, continuity and differentiability of objective function, influence function, Fisher consistency, asymptotics.

Running title: Asymptotics of the least trimmed squares.

### 1 Introduction

In classical multiple linear regression analysis, it is assumed that there is a relationship for a given data set  $\{(\boldsymbol{x}_i^{\top}, y_i)^{\top}, i \in \{1, \cdots, n\}\}$ :

<span id="page-2-0"></span>
$$
y_i = (1, \mathbf{x}_i^{\top})\beta_0 + e_i, \quad i \in \{1, \cdots, n\},\tag{1}
$$

where  $y_i$  and  $e_i$  (an error term, a random variable, and is assumed to have a zero mean and unknown variance  $\sigma^2$  in the classic regression theory) are in  $\mathbb{R}^1$ ,  $\top$  stands for the transpose,  $\beta_0 = (\beta_{01}, \cdots, \beta_{0p})^\top$ , the true unknown parameter, and  $\mathbf{x_i} = (x_{i1}, \cdots, x_{i(p-1)})^\top$  is in  $\mathbb{R}^{p-1}$  $(p \geq 2)$  and could be random. It is seen that  $\beta_{01}$  is the intercept term. Writing  $w_i = (1, x_i^{\top})^{\top}$ , one has  $y_i = \boldsymbol{w}_i^{\top} \boldsymbol{\beta}_0 + e_i$ . The classic assumptions such as linearity and homoscedasticity are implicitly assumed here. Others will be assumed later when they are needed.

One wants to estimate the  $\beta_0$  based on the given sample  $\mathbf{z}^{(n)} := \{(\mathbf{x}_i^{\top}, y_i)^{\top}, i \in \{1, \cdots, n\}\}.$ (hereafter it is implicitly assumed that they are i.i.d. from parent  $(x, y)$ ) For a candidate coefficient vector  $\beta$ , call the difference between  $y_i$  (observed) and  $w_i^{\dagger} \beta$  (predicted), the ith residual,  $r_i(\boldsymbol{\beta})$ , ( $\boldsymbol{\beta}$  is often suppressed). That is,

$$
r_i := r_i(\boldsymbol{\beta}) = y_i - \boldsymbol{w}_i^{\top} \boldsymbol{\beta}.
$$
\n(2)

To estimate  $\beta_0$ , the classic *least squares* (LS) minimizes the sum of squares of residuals,

$$
\widehat{\beta}_{ls} := \arg\min_{\beta \in \mathbb{R}^p} \sum_{i=1}^n r_i^2.
$$

Alternatively, one can replace the square above by the absolute value to obtain the least absolute deviations estimator (aka,  $L_1$  estimator, in contrast to the  $L_2$  (LS) estimator).

Due to its great computability and optimal properties when the error  $e_i$  follows a Gaussian distribution, the LS estimator is popular in practice across multiple disciplines. It, however, can behave badly when the error distribution slightly departs from Gaussian assumption, particularly when the errors are heavy-tailed or contain outliers. In fact, both  $L_1$  and  $L_2$ estimators have the worst 0% asymptotic breakdown point, in sharp contrast to the 50% of the least trimmed squares estimator (Rousseeuw and Leroy (1987) (RL87)). The latter is one of the most robust alternatives to the LS estimator. Robust alternatives to the LS estimator are abundant in the literature. The most popular are, M-estimators (Huber (1964)), least median squares (LMS) and least trimmed squares (LTS) estimators (Rousseeuw (1984) (R84)), S-estimators (Rousseeuw and Yohai (1984)), MM-estimators (Yohai (1987)),  $\tau$ -estimators (Yohai and Zamar (1988)), and maximum depth estimators (Rousseeuw and Hubert (1999)), and Zuo  $(2021 \text{ a}, \text{ b})$ , Zuo and Zuo  $(2023)$   $(ZZ23)$ , among others.

Due to its cube-root consistency of LMS in R84 and its other drawbacks, LTS is preferred over LMS (see Rousseeuw and Van Driessen (2006) (RVD06)). LTS is popular in the literature in view of its fast computibility and high robustness and often severing as the initial estimator for many high breakdown point iterative procedures (e.g., S- and MM- estimators). The LTS is defined as the minimizer of the sum of  $h$  trimmed squares of residuals. Namely,

<span id="page-2-1"></span>
$$
\widehat{\beta}_{lts} := \arg \min_{\beta \in \mathbb{R}^p} \sum_{i=1}^h r_{i:n}^2,\tag{3}
$$

where  $r_{1:n}^2 \leq r_{2:n}^2 \leq \cdots \leq r_{n:n}^2$  are the ordered squared residuals and  $\lceil n/2 \rceil \leq h < n$  and  $\lceil x \rceil$ is the ceiling function.

There are copious studies on the LTS in the literature. Most focused on its computation, e.g., RL87, Stromberg (1993), Hawkins (1994), Hössjer (1995), RVD99, RVD06, Hawkins and Oliver (1999), Agulló (2001), Hofmann et al. (2010), and Klouda (2015).

Others sporadically addressed the asymptotics, e.g., RL87, and Mašíček (2004) addressed the asymptotic normality of the LTS, but limited to the location case, that is, when  $p = 1$ . Víšek (2006a b c) also addressed the asymptotics of the LTS without employing advanced technical tools in a series (three) of lengthy articles for the consistency, root-n consistency, and asymptotic normality, respectively. The analysis is technically demanding and with difficult verified assumptions  $A, B, C$ . Furthermore, the analysis is limited to the non-random vectors  $x_i$  case. In this article, without those assumptions and limitation, those results are established in a concise manner with the help of advanced empirical process theory.

Replacing  $(1, x_i^{\top})\beta_0$  by a unspecified nonlinear function  $h(x_i, \beta_0)$ , Chen, at al. (1997)  $(CSZ97)$  and Cížek  $(2004, 2005)$   $(C04, C05)$  discussed the asymptotics of the LTS in a nonlinear regression setting. Now that more general non-linear case has been addressed, one might wonder is there any merits to discuss the special linear case in this article?

There are at least these merits: (i) the nonlinear function  $h(x_i, \beta_0)$  cannot always cover the linear case of  $(1, x_i^{\top})\beta_0$  for the usual LTS (e.g., in the exponential and power regression cases); (ii) many assumptions for the nonlinear case (see A1, A2, A3, A4 in CSZ97; H1, H2, H3, H4, H5, H6; D1, D2; I1, I2 in C04 and C05 ) (which are usual difficult to verify) can be dropped for the linear case as demonstrated in this article. (iii) A key assumption that  $\{h(x, \beta), \beta \in \Theta\}$  form a VC class of functions over a compact parameter space  $\Theta$  (see CSZ97, C04, C05) can be verified directly in this article.

To avoid all the drawbacks and limitations above and take advantage of the standard results of the empirical process theory, this article defines the population version of the LTS (Sec[.2.1\)](#page-4-0), introduces the novel partition of the parameter space (Sec[.2.2\)](#page-5-0), and investigates the primary properties of the objective function for the LTS both in the empirical and population settings (Sec[.2\)](#page-4-1) for the first time. The obtained novel results facilitate the versification of some fundamental assumptions conveniently made in the literature. The major contributions of this article thus include

- (a) introducing a novel partition of the parameter space and defining an original population version of the LTS for the first time;
- (b) investigating primary properties of the sample and population versions of the objective function for the LTS, obtaining original results;
- (c) first time obtaining the influence function  $(p \geq 2)$  and Fisher consistency for the LTS;
- (d) first time establishing the strong consistency of the sample LTS via a generalized Glivenko-Cantelli Theorem without artificial assumptions; and
- (e) first time employing a novel and concise approach based on the empirical process theory to establish asymptotic normality of the sample LTS.

The rest of the article is organized as follows. Section [2](#page-4-1) introduces first time the population version of LTS and addresses the properties of the LTS estimator in both empirical and population settings, including the global continuity and local differentiability and convexity of its objective function; its influence function (in  $p > 2$  cases) and Fisher consistency are established for the first time. Section [3](#page-9-0) establishes the strong consistency via a generalized Glivenko-Cantelli Theorem and the asymptotic normality of the estimator is re-established in a very different and concise approach (via stochastic equicontinuity) rather than the previous ones in the literature. Section [4](#page-13-0) addresses the asymptotic inference procedures based on the asymptotic normality and bootstrapping. Concluding remarks in Section [5](#page-15-0) end the article. Major proofs are deferred to an Appendix.

# <span id="page-4-1"></span>2 Definition and Properties of the LTS

### <span id="page-4-0"></span>2.1 Definition

Denote by  $F_{(\mathbf{x}^{\top},y)}$  the joint distribution of  $\mathbf{x}^{\top}$  and y in model [\(1\)](#page-2-0). Throughout  $F_{\mathbf{Z}}$  stands for the distribution function of the random vector  $\mathbf{Z}^{\top}$ . For a given  $\boldsymbol{\beta} \in \mathbb{R}^p$  and an  $\alpha \in [1/2, c]$ ,  $1/2 < c < 1$ , let  $q(\mathcal{A}, \alpha) = F_W^{-1}(\alpha)$  be the  $\alpha$ th quantile of  $F_W$  with  $W := W(\mathcal{A}) = (y - \mathbf{w}^\top \mathcal{A})^2$ , where  $w^{\top} = (1, x^{\top})$ . The constant  $c = 1$  case is excluded to avoid unbounded  $q(\beta, \alpha)$  and the LS cases. Define an objective function

<span id="page-4-3"></span>
$$
O(F_{(\mathbf{x}^{\top},y)}, \boldsymbol{\beta}, \alpha) = \int (y - \boldsymbol{w}^{\top} \boldsymbol{\beta})^2 \mathbb{1} \left( (y - \boldsymbol{w}^{\top} \boldsymbol{\beta})^2 \le q(\boldsymbol{\beta}, \alpha) \right) dF_{(\mathbf{x}^{\top},y)}, \tag{4}
$$

and a regression functional

<span id="page-4-4"></span>
$$
\beta_{lts}(F_{(\mathbf{x}^{\top},y)},\alpha) = \arg\min_{\boldsymbol{\beta}\in\mathbb{R}^p} O(F_{(\mathbf{x}^{\top},y)},\boldsymbol{\beta},\alpha),\tag{5}
$$

where  $\mathbb{1}(A)$  is the indicator of A (i.e., it is one if A holds and zero otherwise). Let  $F_{(\mathbf{x}^{\top},y)}^n$ be the sample version of the  $F_{(\mathbf{x}^\top, y)}$  based on a sample  $\mathbf{z}^{(n)} := \{(\mathbf{x}_i^\top, y_i)^\top, i \in \{1, 2, \cdots, n\}\}.$ The  $F_{(\mathbf{x}^{\top},y)}^n$  and  $\mathbf{z}^{(n)}$  will be used interchangeably. Using the  $F_{(\mathbf{x}^{\top},y)}^n$ , one obtains the sample versions

<span id="page-4-5"></span>
$$
O(F_{(\mathbf{x}^{\top},y)}^n, \boldsymbol{\beta}, \alpha) = \frac{1}{n} \sum_{i=1}^{\lfloor \alpha n \rfloor + 1} r_{i:n}^2,
$$
\n(6)

where  $|x|$  is the floor function. Further

<span id="page-4-2"></span>
$$
\widehat{\boldsymbol{\beta}}_{lts}^n := \boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top,y)}^n, \alpha) = \arg\min_{\boldsymbol{\beta} \in \mathbb{R}^p} O(F_{(\mathbf{x}^\top,y)}^n, \boldsymbol{\beta}, \alpha). \tag{7}
$$

It is readily seen that the  $\hat{\beta}_{lts}^n$  above is identical to the  $\hat{\beta}_{lts}$  in [\(3\)](#page-2-1) with  $h = \lfloor \alpha n \rfloor + 1$ . Henceforth we prefer to treat the  $\hat{\beta}_{lts}^n$  rather than the  $\hat{\beta}_{lts}$  in [\(3\)](#page-2-1).

The first natural question is the existence of the minimizer in the right-hand side (RHS) of [\(7\)](#page-4-2), or the existence of the  $\hat{\beta}_{lts}^n$ . Does it always exist? If it exists, will it be unique? Unique existence is a key precondition for the study of asymptotics of an estimator.

One might take the existence for granted since the objective is non-negative and has a finite infimum which can be approximated by objective values of a sequence of  $\beta$ s. There is a sub-sequence of  $\beta$ s with their objective values converging to the infimum which is minimum due the continuity of the objective function. The sub-sequence of  $\beta$ s converges to a point  $\beta^*$  which is the minimizer of the RHS. There are multiple issues with the arguments above. The existence and the convergence of the  $\beta$  sub-sequence (to a minimum) and continuity of objective function need to be proved. In the sequel, we take a different approach.

### <span id="page-5-0"></span>2.2 Properties in the empirical case

Write  $O^n(\beta)$  and  $\mathbb{1}_i$  for the  $O(F_{(\mathbf{x}',y)}^n, \beta, \alpha)$  and the  $\mathbb{1}(r_i^2 \leq r_{h:n}^2)$ , respectively. It is seen that

$$
O^{n}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^{n} r_i^2 \mathbb{1} \left( r_i^2 \le r_{h:n}^2 \right) = \frac{1}{n} \sum_{i=1}^{n} r_i^2 \mathbb{1}_i,
$$
\n(8)

where  $h = |\alpha n| + 1$ . The fraction  $1/n$  will often be ignored in the following discussion.

#### Existence and uniqueness

**Partition parameter space** For a given sample  $z^{(n)}$ , an  $\alpha$  (or h), and any  $\beta^1 \in \mathbb{R}^p$ , let  $r_{i:n}^2(\boldsymbol{\beta}^1) = r_{k_i}^2(\boldsymbol{\beta}^1)$  for an integer  $k_i$ . Note that  $r_j$  and  $k_i$  depend on  $\boldsymbol{\beta}^1$ , i.e.,  $r_j := r_j(\boldsymbol{\beta}^1)$ ,  $k_i := k_i(\boldsymbol{\beta}^1)$ . Obviously  $r_{k_1}^2 \leq r_{k_2}^2 \leq \cdots \leq r_{k_n}^2$ . Call  $\{k_i, 1 \leq i \leq h\}$   $\boldsymbol{\beta}^1$ -h-integer set. If  $r_i^2 \neq r_j^2$  for any distinct i and j, then the h-integer set is unique. Hereafter we assume (A0): W has a density for any given  $\beta$ . Then almost surely (a.s.), the h-integer set is unique.

Consider the unique cases. There are other  $\beta s$  in  $\mathbb{R}^p$  that share the same h-integer set as that of the  $\beta^1$ . Denote the set of such points that have the same h integers as  $\beta^1$  by

<span id="page-5-1"></span>
$$
S_{\beta^1} := \left\{ \beta \in \mathbb{R}^p : k_i(\beta) = k_i(\beta^1) = k_i, \ i \in \{1, 2, \cdots, h\} \ \{k_i, 1 \le i \le h\} \text{ is unique.} \right\}
$$
 (9)

If (A0) holds, then  $S_{\beta^1} \neq \emptyset$  (a.s.). If it is  $\mathbb{R}^p$ , then we have a trivial case (see Remarks 2.1 below). Otherwise, there are only finitely many such sets (for a fixed  $n$ ) that partition  $\mathbb{R}^p$ . Let  $\cup_{l=1}^L \overline{S}_{\beta^l} = \mathbb{R}^p$ , where  $S_{\beta^l}$ s are defined similarly to [\(9\)](#page-5-1) and are disjoint for different  $l, 1 \leq l \leq L := \binom{n}{h}$  $\mathbf{F}_h^n$ , and  $\overline{A}$  is the closure of the set A. Write  $\mathbf{X}_n = (\boldsymbol{w}_1, \cdots, \boldsymbol{w}_n)'$ , an  $n \times p$ matrix. Assume (A1):  $X_n$  and any its h rows have a full rank p. As in the R package ltsReg (by Valentin Todorov), hereafter we assume that  $p < n/2$ .

**Lemma 2.1** Assume that  $(A0)$  and  $(A1)$  hold, then

- (i) (a) For any  $l$   $(1 \leq l \leq L)$ ,  $r_k^2$  $k_1(\beta^l) < r_{k_2(\beta^l)}^2 < \cdots < r_{k_h(\beta^l)}^2$  over  $S_{\beta^l}$ .
	- (b) For any  $\eta \in S_{\beta^l}$ , there exists an open ball  $B(\eta, \delta)$  centered at  $\eta$  with a radius  $\delta > 0$ such that for any  $\beta \in B(\eta, \delta)$

<span id="page-5-2"></span>
$$
O^{n}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^{h} r_{k_{i}(\boldsymbol{\beta})}^{2}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^{h} r_{k_{i}(\boldsymbol{\eta})}^{2}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^{h} r_{k_{i}(\boldsymbol{\beta}^{l})}^{2}(\boldsymbol{\beta}), \ (a.s.)
$$
 (10)

(ii) The graph of  $O^n(\mathcal{B})$  over  $\mathcal{B} \in \mathbb{R}^p$  is composed of the L closures of graphs of the quadratic function of  $\beta$ :  $\frac{1}{n}$  $\frac{1}{n}\sum_{i=1}^h r_k^2$  $\sum_{k_i(\boldsymbol{\beta}^l)}^2 (\boldsymbol{\beta})$  for  $O^n(\boldsymbol{\beta}^l)$  and any  $l$   $(1 \leq l \leq L)$ , joined together.

(iii)  $O^n(\beta)$  is continuous in  $\beta \in \mathbb{R}^p$ .

(iv)  $O<sup>n</sup>(\beta)$  is differentiable and strictly convex over each  $S_{\beta}$  for any  $1 \leq l \leq L$ .

**Proof:** see the Appendix.

### Remarks 2.1

(a) If  $S_{\beta^0} = \mathbb{R}^p$ , then  $O^n(\beta)$  is a twice differentiable and strictly convex quadratic function of  $\beta$ , the existence and the uniqueness of  $\widehat{\beta}_{lts}^n$  are trivial as long as  $\mathbf{X}_n$  has a full rank.

(b) Replacing  $(1, x_i)$ ' $\beta$  by a nonlinear  $h(x_i, \beta)$ , C04 and C05 also addressed the continuity and differentiability of the objective function of the LTS. However, they assumed that (i)  $F_W$ is twice differentiable around the points corresponding to the square roots of the  $\alpha$ -quantiles of W, (ii)  $h(x_i, \beta)$  is continuous over parameter space B, (iii)  $h(x_i, \beta)$  is twice differentiable in  $\beta$  for  $\beta \in B(\beta_0, \delta)$  a.s., and (iv)  $\partial h(x_i, \beta)/\partial \beta$  is continuous in  $\beta$ , All assumptions were never verified in C04 and C05 though, however they are not required in Lemma 2.1.

(c) Continuity and differentiability inferred just based on  $O<sup>n</sup>(\beta)$  being the sum of h continuous and differentiable functions (squares of residual) without (i) or [\(10\)](#page-5-2) might be inadequate (flawed). In general,  $O<sup>n</sup>(\beta)$  is not differentiable nor convex in  $\beta$  globally.  $\Box$ 

 $\text{Let } \mathbf{y}_n := (y_1, \cdots, y_n)^\top,\, \bm{M}_n := \bm{M}(\mathbf{y}_n, \mathbf{X}_n, \bm{\beta}, \alpha) = \sum_{i=1}^n \bm{w}_i \bm{w}_i^\top \mathbb{1}_i = \sum_{i=1}^h \bm{w}_{k_i(\bm{\beta})} \bm{w}_{k_i(\bm{\beta})}^\top.$ Note that  $\mathbb{1}_i$  depends on  $\beta$ .

Theorem 2.1 Assume that (A0) and (A1) hold. Then

- (i)  $\hat{\beta}_{lts}^n$  exists and is the local minimum of  $O^n(\beta)$  over  $S_{\beta^{l_0}}$  for some  $l_0$   $(1 \leq l_0 \leq L)$ .
- (ii) Over  $S_{\beta^{l_0}}$ ,  $\widehat{\beta}^n_{lts}$  is the solution of the system of equations

<span id="page-6-0"></span>
$$
\sum_{i=1}^{n} (y_i - \boldsymbol{w}_i^{\top} \boldsymbol{\beta}) \boldsymbol{w}_i \mathbb{1}_i = \mathbf{0},
$$
\n(11)

(iii) Over  $S_{\beta^{l_0}}$ , the unique solution is

$$
\widehat{\boldsymbol{\beta}}_{lts}^n = \boldsymbol{M}_n(\mathbf{y}_n, \mathbf{X}_n, \widehat{\boldsymbol{\beta}}_{lts}^n, \alpha)^{-1} \sum_{i=1}^h y_{k_i(\boldsymbol{\beta}^{l_0})} \boldsymbol{w}_{k_i(\boldsymbol{\beta}^{l_0})}. \tag{12}
$$

**Proof:** The given conditions and Lemma 2.1 allow one to focus on a piece  $S_{\beta^l}$ ,  $1 \leq l \leq L$ , all results follows in a straightforward fashion, for more details, see the Appendix.  $\Box$ 

### Remarks 2.2

(a) Unique existence, which is often implicitly assumed or ignored in the literature, is central for the discussion of asymptotics of  $\hat{\beta}_{lts}^n$ . Existence of  $\hat{\beta}_{lts}^n$  could also be established under the assumption that there are no  $\lfloor (n+1)/2 \rfloor$  sample points of  $z^{(n)}$  contained in any  $(p-1)$ -dimensional hyperplane, similarly to that of Theorem 2.2 for LST in Zuo and Zuo [\[40\]](#page-18-0). It is established here without such assumption nevertheless.

(b) A sufficient condition for the invertibility of  $M_n$  is that any h rows of  $X_n$  form a full rank sub-matrix. The latter is true if  $(A1)$  holds.

(c) Víšek (2006a) also addressed the existence of  $\hat{\beta}_{lts}^n$  (Assertion 1) for non-random covariates (carriers) satisfying many demanding assumptions  $(A, B)$ . The uniqueness was left unaddressed though. unaddressed though.

### 2.3 Properties in the population case

The best breakdown point of the LTS (see p. 132 of RL87) reflects its global robustness. We now examine its local robustness via the influence function to depict its complete robust picture.

### Definition of influence function

For a distribution F on  $\mathbb{R}^p$  and an  $\varepsilon \in (0, 1/2)$ , the version of F contaminated by an  $\varepsilon$ amount of an arbitrary distribution G on  $\mathbb{R}^p$  is denoted by  $F(\varepsilon, G) = (1 - \varepsilon)F + \varepsilon G$  (an  $\varepsilon$ ) amount deviation from the assumed F).  $F(\varepsilon, G)$  is actually a convex contamination of F. There are other types of contamination such as contamination by total variation or Hellinger distances. We cite the definition of influence function from Hampel et al. [\[7\]](#page-16-0).

**Definition 2.2** [\[7\]](#page-16-0) The *influence function* (IF) of a functional **t** at a given point  $x \in \mathbb{R}^p$  for a given  $F$  is defined as

IF
$$
(\mathbf{x}; \mathbf{t}, F) = \lim_{\varepsilon \to 0^+} \frac{\mathbf{t}(F(\varepsilon, \delta_{\mathbf{x}})) - \mathbf{t}(F)}{\varepsilon},
$$
 (13)

where  $\delta_{\mathbf{x}}$  is the point-mass probability measure at  $\mathbf{x} \in \mathbb{R}^p$ .

The function IF( $x; t, F$ ) describes the relative effect (influence) on t of an infinitesimal point-mass contamination at  $x$  and measures the local robustness of  $t$ .

To establish the IF for the functional  $\beta_{lts}(F_{(\mathbf{z}^{\top},y)},\alpha)$ , we need to first show its existence and uniqueness with or without point-mass contamination. To that end, write

$$
F_{\varepsilon}(\mathbf{z}) := F(\varepsilon, \delta_{\mathbf{z}}) = (1 - \varepsilon) F_{(\mathbf{z}^{\top}, y)} + \varepsilon \delta_{\mathbf{z}},
$$

with  $\mathbf{u} = (\mathbf{s}^\top, t)^\top \in \mathbb{R}^p$ ,  $\mathbf{s} \in \mathbb{R}^{p-1}$ ,  $t \in \mathbb{R}^1$  being the corresponding random vector (i.e.  $F_{\mathbf{u}} = F_{\varepsilon}(\mathbf{z}) = F(\varepsilon, \delta_{\mathbf{z}})$ . The versions of [\(4\)](#page-4-3) and [\(5\)](#page-4-4) at the contaminated  $F(\varepsilon, \delta_{\mathbf{z}})$  are respectively

$$
O(F_{\varepsilon}(\mathbf{z}), \boldsymbol{\beta}, \alpha) = \int (t - \mathbf{v}^{\top} \boldsymbol{\beta})^2 \mathbb{1} \left( (t - \mathbf{v}^{\top} \boldsymbol{\beta})^2 \le q_{\varepsilon}(\mathbf{z}, \boldsymbol{\beta}, \alpha) \right) dF_{\mathbf{u}}(\mathbf{s}, t), \tag{14}
$$

with  $q_{\varepsilon}(\mathbf{z}, \boldsymbol{\beta}, \alpha)$  being the  $\alpha$ th quantile of the distribution function of  $(t - \boldsymbol{v}^\top \boldsymbol{\beta})^2$ ,  $\boldsymbol{v} = (1, \boldsymbol{s}^\top)^\top$ , and

$$
\beta_{lts}(F_{\varepsilon}(\mathbf{z}), \alpha) = \arg\min_{\boldsymbol{\beta} \in \mathbb{R}^p} O(F_{\varepsilon}(\mathbf{z}), \boldsymbol{\beta}, \alpha).
$$
\n(15)

For  $\beta_{lts}(F_{(\mathbf{z}^\top,y)},\alpha)$  defined in [\(5\)](#page-4-4) and  $\beta_{lts}(F_\varepsilon(\mathbf{z}),\alpha)$  above, we have an analogous result to Theorem 2.1. (Assume that the counterpart of model [\(1\)](#page-2-0) is  $y = (1, \mathbf{x}^{\top})\boldsymbol{\beta}_0 + e = \mathbf{w}^{\top}\boldsymbol{\beta}_0 + e$ . Before we derive the influence function, we need to establish existence and uniqueness.

### Existence and Uniqueness

Write  $O(\beta)$  for  $O(F_{(\boldsymbol{x}^\top,y)}, \beta, \alpha)$  in [\(4\)](#page-4-3). To have a counterpart of Lemma 2.1, we need

(A2) W has a positive density around a small neighborhood of  $q(\beta,\alpha)$  for the given  $\alpha, \beta$ .

**Lemma 2.2** Assume (A2) holds and  $E(\boldsymbol{w}\boldsymbol{w}^{\top})$  exists. Then

- (i)  $O(\beta)$  is continuous in  $\beta \in \mathbb{R}^p$ ;
- (ii)  $O(\beta)$  is twice differentiable in  $\beta \in \mathbb{R}^p$ ,

$$
\frac{\partial^2 O(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}^2} = 2E(\boldsymbol{w}\boldsymbol{w}^\top \mathbb{1}((y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 \leq q(\boldsymbol{\beta}, \alpha));
$$

(iii)  $O(\beta)$  is strictly convex in  $\beta \in \mathbb{R}^p$ .

**Proof:** The boundedness of the integrand in  $(4)$ , given conditions and Lebesgue dominated convergence theorem leads to the desired results, for details, see the Appendix.

Note that (ii) and (iii) above are global in  $\beta$ , stronger than the empirical counterparts, all are attributed to boundary of  $S_{\beta^l}$  issue. We now treat the existence and uniqueness of  $\beta_{lts}$ , which is central for the asymptotics study.

**Theorem 2.3** Assume (A2) holds and  $E(\boldsymbol{w}\boldsymbol{w}^{\top})$  exists,  $q_{\varepsilon}(\mathbf{z}, \boldsymbol{\beta}, \alpha)$  is continuous in  $\boldsymbol{\beta}$ , and  $P((t - \boldsymbol{v}^\top \boldsymbol{\beta})^2 = q_{\varepsilon}(\mathbf{z}, \boldsymbol{\beta}, \alpha)) = 0$  for any  $\boldsymbol{\beta} \in \mathbb{R}^p$  and the given  $\alpha$ . Then

(i)  $\beta_{lts}(F_{(\mathbf{x}^{\top},y)},\alpha)$  and  $\beta_{lts}(F_{\varepsilon}(\mathbf{z}),\alpha)$  exist.

(ii) Furthermore, they are the solution of system of equations, respectively

$$
\int (y - \boldsymbol{w}^\top \boldsymbol{\beta}) \boldsymbol{w} \mathbb{1} \left( (y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 \le q(\boldsymbol{\beta}, \alpha) \right) dF_{(\boldsymbol{x}^\top, y)}(\boldsymbol{x}, y) = \mathbf{0},\tag{16}
$$

$$
\int (t - \mathbf{v}^\top \boldsymbol{\beta}) \mathbf{v} \mathbb{1} \left( (t - \mathbf{v}^\top \boldsymbol{\beta})^2 \le q_\varepsilon(\mathbf{z}, \boldsymbol{\beta}, \alpha) \right) dF_\mathbf{u}(\mathbf{s}, t) = \mathbf{0}.
$$
 (17)

(iii)  $\beta_{lts}(F_{(\mathbf{z}^\top,y)},\alpha)$  and  $\beta_{lts}(F_{\varepsilon}(\mathbf{z}),\alpha)$  are unique provided that

$$
\int \boldsymbol{w}\boldsymbol{w}^{\top} \mathbb{1}\left( (y - \boldsymbol{w}^{\top}\boldsymbol{\beta})^2 \le q(\boldsymbol{\beta}, \alpha) \right) dF_{(\boldsymbol{x}^{\top}, y)}(\boldsymbol{x}, y), \tag{18}
$$

<span id="page-8-2"></span><span id="page-8-1"></span><span id="page-8-0"></span>
$$
\int \boldsymbol{v} \boldsymbol{v}^{\top} \mathbb{1} \left( (t - \boldsymbol{v}^{\top} \boldsymbol{\beta})^2 \le q_{\varepsilon}(\mathbf{z}, \boldsymbol{\beta}, \alpha) \right) dF_{\mathbf{u}}(\mathbf{s}, t)
$$
(19)

are invertible for  $\beta$  in a small neighborhood of  $\beta_{lts}(F_{(\mathbf{z}^{\top},y)},\alpha)$  and  $\beta_{lts}(F_{\varepsilon}(\mathbf{z}),\alpha)$  respectively. **Proof:** In light of Lemma 2.2, the proof is straightforward, see the Appendix.  $\Box$ 

Continuity of  $q_{\varepsilon}(\mathbf{z}, \boldsymbol{\beta}, \alpha)$  in  $\boldsymbol{\beta}$  is necessary for the differentiability of  $O(F_{\varepsilon}(\mathbf{z}), \boldsymbol{\beta}, \alpha)$ . In the non-contaminated case, continuity of  $q(\beta, \alpha)$  is guaranteed by  $(A2)$ .

Does the population version of the LTS,  $\beta_{lts}$ , defined in [\(5\)](#page-4-4), have something to do with  $\beta_0$ ? It turns out under some conditions, they are identical, which is called Fisher consistency.

### Fisher consistency

**Theorem 2.4** Assume (A2) holds and  $E(\bm{w}\bm{w}^{\top})$  exists, then  $\beta_{lts}(F_{(\bm{x}^{\top},y)},\alpha) = \beta_0$  provided that (i)  $E_{(\boldsymbol{x}^\top,y)}\left(\boldsymbol{w}\boldsymbol{w}^\top\mathbb{1}(r(\boldsymbol{\beta})^2\leq F_{r(\boldsymbol{\beta})}^{-1}\right)$  $\binom{n-1}{r(\beta)^2}(\alpha)$  is invertible, and (ii)  $E_{(\mathbf{x}^\top,y)}(e\mathbf{w}1(e^2 \leq F_{e^2}^{-1}))$  $e^{-1}_{e^2}(\alpha)$  = 0, where  $r(\beta) = y - \boldsymbol{w}^\top \beta$ .

**Proof:** Theorem 2.3 leads directly to the desired result, see the Appendix. □

### Influence function

**Theorem 2.5** Assume that the assumptions in Theorem 2.3 hold. Set  $\beta_{lts} := \beta_{lts}(F_{(\mathbf{x}^{\top},y)},\alpha)$ . Then for any  $\mathbf{z}_0 := (\mathbf{s}_0^{\top}, t_0)^{\top} \in \mathbb{R}^p$ , we have that

IF
$$
(\mathbf{z}_0; \boldsymbol{\beta}_{lts}, F_{(\boldsymbol{x}^\top, y)}) = \begin{cases} \mathbf{0}, & \text{if } (t_0 - \boldsymbol{v}_0^\top \boldsymbol{\beta}_{lts})^2 > q(\boldsymbol{\beta}_{lts}, \alpha), \\ M^{-1}(t_0 - \boldsymbol{v}_0^\top \boldsymbol{\beta}_{lts})\mathbf{v}_0, & \text{otherwise}, \end{cases}
$$

provided that  $\boldsymbol{M} = E_{(\boldsymbol{x}^\top,y)} \Big( \boldsymbol{w} \boldsymbol{w}^\top \boldsymbol{1} \left( r(\boldsymbol{\beta}_{lts})^2 \leq q(\boldsymbol{\beta}_{lts}, \alpha) \right) \Big)$  is invertible, where  $\boldsymbol{v}_0 = (1, \boldsymbol{s}_0^\top)^\top$ .

**Proof:** The connection to the derivative of a functional is the key, see the Appendix.  $\Box$ 

#### Remarks 2.3

(a) When  $p = 1$ , the problem in our model [\(1\)](#page-2-0) becomes a location problem (see p. 158 of RL87) and the IF of the LTS estimation functional has been given on p. 191 of RL87. In the location setting, Tableman [\[27\]](#page-17-0) also studied the IF of the LTS. When  $p = 2$ , namely in the simple regression case, Ollerer et al.  $(2015)$  studied IF of the sparse-LTS functional under the assumption that  $x$  and  $e$  are independent and normally distributed. Under stringent assumptions on the error terms  $e_i$  and on x, Mašíček (2004) also addressed the IF of LTS for any p, but the point mass at  $(x^{\top}, z)$  with z being the error term, an unusual contaminating point. The above result is much more general and valid for any  $p \geq 1$ ,  $x'$ , and e.

(b) The influence function of  $\beta_{lts}$  remains bounded if the contaminating point  $(s'_0, t_0)$ does not follow the model (i.e., its residual is extremely large), in particular for bad leverage points and vertical outliers. This shows the good robust properties of the LTS.

(c) The influence function of  $\beta_{lts}$ , unfortunately, might be unbounded (in  $p > 1$  case), sharing the drawback of the sparse-LTS (in the  $p = 2$  case). The latter was shown in Ollerer et al. (2015). Trimming based on the residuals (or squared residuals) will have this type of drawback since the term  $w^\top \beta$  can be bounded, but  $||x||$  might not.

# <span id="page-9-0"></span>3 Asymptotic properties

Víšek  $(2006a, b, c)$  (V06a, b, c) rigorously addressed the consistency, root-n consistency, and normality of the LTS under a restrictive setting  $(x_i)$  are non-random covariates) plus many assumptions on  $x_i$ s and on the distribution of  $e_i$  in three lengthy series papers.

 $Cizek$  (2004, 2005) (C04, C05) also addressed asymptotic properties of an extended LTS under  $\beta$ -mixing conditions for  $x_i$  with nonlinear regression function  $h(x_i, \beta)$ , a more general setting, but with the price of numerous artificial assumptions (H1, H2, H3, H4, H5, H6; D1,D2; I1,I2) that are never verified in any concrete example nor for the special case of linear LTS. That is, C04 and C05 cannot trivially be applied for the linear LTS in [\(3\)](#page-2-1).

Strong consistency has been addressed in CSZ97 in nonlinear setting with key assumptions that are never verified even for the special linear LTS. We now rigorously establish strong consistency without any artificial assumptions.

### 3.1 Strong consistency

Following the notations of Pollard (1984) (P84), write

$$
O(\beta, P) := O(F_{(\boldsymbol{x}^\top, y)}, \beta, \alpha) = P[(y - \boldsymbol{w}^\top \beta)^2 \mathbb{1}(r(\beta)^2 \le F_{r(\beta)^2}^{-1}(\alpha))] = Pf,
$$
  

$$
O(\beta, P_n) := O(F_{(\boldsymbol{x}^\top, y)}^n, \beta, \alpha) = \frac{1}{n} \sum_{i=1}^n r_i^2 \mathbb{1}(r_i^2 \le (r)_{n:n}^2) = P_n f,
$$

where  $f := f(x, y, \beta, \alpha) = (y - \boldsymbol{w}^\top \beta)^2 \mathbb{1}(r(\beta)^2 \leq F_{r(\beta)}^{-1})$  $r_{r(\boldsymbol{\beta})^2}^{r-1}(\alpha)$ ,  $\alpha$  and  $h = \lfloor \alpha n \rfloor + 1$  are fixed.

Under corresponding assumptions in Theorems 2.1 and 2.3,  $\hat{\beta}_{lts}^{n}$  and  $\beta_{lts}$  are unique minimizers of  $O(\beta, P_n)$  and  $O(\beta, P)$ , respectively.

To show that  $\hat{\beta}_{lts}^n$  converges to  $\beta_{lts}$  a.s., one can take the approach given in Section 4.2 of Zuo [\[40\]](#page-18-0). But here we take a different and more direct approach.

To show that  $\widehat{\beta}_{lts}^n$  converges to  $\beta_{lts}$  a.s., it will suffice to prove that  $O(\widehat{\beta}_{lts}^n, P) \to O(\beta_{lts}, P)$ a.s., because  $O(\beta, P)$  is bounded away from  $O(\beta_{lts}, P)$  outside each neighborhood of  $\beta_{lts}$  in light of continuity and compactness (also see Lemma 4.3 of Zuo [\[40\]](#page-18-0)). Let  $\Theta$  be a closed ball centered at  $\beta_{lts}$  with a radius  $r > 0$ . Define a class of functions

$$
\mathscr{F}(\boldsymbol{\beta}, \alpha) = \left\{ f(\boldsymbol{x}, y, \boldsymbol{\beta}, \alpha) = (y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 \mathbb{1} (r(\boldsymbol{\beta})^2 \le F_{r(\boldsymbol{\beta})^2}^{-1}(\alpha)) : \boldsymbol{\beta} \in \Theta, \alpha \in [1/2, c] \right\}
$$

If we prove uniform almost sure convergence of  $P_n$  to P over  $\mathscr F$  (see Lemma 3.1 below), then we can deduce that  $O(\widehat{\beta}_{lts}^n, P) \to O(\beta_{lts}, P)$  a.s. from

$$
O(\widehat{\beta}_{lts}^n, P_n) - O(\widehat{\beta}_{lts}^n, P) \to 0 \text{ (in light of Lemma 3.1), and}
$$
  

$$
O(\widehat{\beta}_{lts}^n, P_n) \le O(\beta_{lts}, P_n) \to O(\beta_{lts}, P) \le O(\widehat{\beta}_{lts}^n, P).
$$

The above discussions and arguments have led to

Theorem 3.1. Under corresponding assumptions in Theorems 2.1 and 2.3 for uniqueness of  $\widehat{\beta}_{lts}^n$  and  $\beta_{lts}$  respectively, we have  $\widehat{\beta}_{lts}^n$  converges a.s. to  $\beta_{lts}$  (i.e.  $\widehat{\beta}_{lts}^n - \beta_{lts} = o(1)$ , a.s.).

The above is based on the following generalized Glivenko-Cantelli Theorem.

**Lemma 3.1.** sup $_{f \in \mathcal{F}} |P_n f - Pf| \to 0$  a.s., provided that (A2) holds.

**Proof:** Verifying two requirements in the Theorem 24 in II.5 of P84 leads to the result. Showing the covering number for functions in  $\mathscr F$  is bounded is challenging. Essentially one needs to show that the graphs of functions in  $\mathscr F$  form a VC class of sets (this was avoided in the literature, e.g., in C04, C05, CSZ97, and V06a, b, c). For details, see the Appendix.  $\Box$ 

### 3.2 Root-n consistency and asymptotic normality

Instead of treating the root-n consistency separately as in V06b c, we will establish asymptotic normality of  $\hat{\beta}_{lts}^{n}$  directly via stochastic equicontinuity (see p. 139 of P84 or the supplementary of Zuo (2020)).

Stochastic equicontinuity refers to a sequence of stochastic processes  $\{Z_n(t): t \in T\}$ whose shared index set T comes equipped with a semi-metric  $d(\cdot, \cdot)$ .

**Definition 3.1** [IIV. 1, Def. 2 of P84]. Call  $Z_n$  stochastically equicontinuous at  $t_0$  if for each  $\eta > 0$  and  $\epsilon > 0$  there exists a neighborhood U of  $t_0$  for which

$$
\limsup P\left(\sup_U |Z_n(t) - Z_n(t_0)| > \eta\right) < \epsilon.
$$
\n(20)

It is readily seen (see p.140 of P84) that if  $\tau_n$  is a sequence of random elements of T that converges in probability to  $t_0$ , then

$$
Z_n(\tau_n) - Z_n(t_0) \to 0 \text{ in probability},\tag{21}
$$

because, with probability tending to one,  $\tau_n$  will belong to each U. The form above will be easier to apply, especially when behavior of a particular  $\tau_n$  sequence is under investigation.

Suppose  $\mathscr{F} = \{f(\cdot, t) : t \in T\}$ , with T a subset of  $\mathbb{R}^k$ , is a collection of real, P-integrable functions on the set S where P (probability measure) lives. Denote by  $P_n$  the empirical measure formed from n independent observations on  $P$ , and define the empirical process  $E_n$ as the signed measure  $n^{1/2}(P_n - P)$ . Define

$$
F(t) = Pf(\cdot, t),
$$
  

$$
F_n(t) = P_n f(\cdot, t).
$$

Suppose  $f(\cdot, t)$  has a linear approximation near the  $t_0$  at which  $F(\cdot)$  takes on its minimum value:

<span id="page-11-0"></span>
$$
f(\cdot, t) = f(\cdot, t_0) + (t - t_0)^\top \nabla f(\cdot, t) + |t - t_0| r(\cdot, t). \tag{22}
$$

For completeness set  $r(\cdot, t_0) = 0$ , where  $\nabla$  (differential operator) is a vector of k real functions on S. We cite theorem 5 in VII.1 of P84 (p. 141) for the asymptotic normality of  $\tau_n$ .

**Lemma 3.2** [P84]. Suppose  $\{\tau_n\}$  is a sequence of random vectors converging in probability to the value  $t_0$  at which  $F(\cdot)$  has its minimum. Define  $r(\cdot, t)$  and the vector of functions  $\nabla(\cdot, t)$  by [\(22\)](#page-11-0). If

(i)  $t_0$  is an interior point of the parameter set T;

(ii)  $F(\cdot)$  has a non-singular second derivative matrix V at  $t_0$ ;

(iii)  $F_n(\tau_n) = o_p(n^{-1}) + \inf_t F_n(t);$ 

(iv) the components of  $\nabla f(\cdot, t)$  all belong to  $\mathscr{L}^2(P)$ ;

(v) the sequence  $\{E_n r(\cdot, t)\}\$ is stochastically equicontinuous at  $t_0$ ;

then

$$
n^{1/2}(\tau_n - t_0) \stackrel{d}{\longrightarrow} N(\mathbf{0}, V^{-1}[P(\nabla \nabla^{\top}) - (P\nabla)(P\nabla)^{\top}]V^{-1}).
$$

By Theorems 2.1 and 3.1, assume, without loss of generality (w.l.o.g.), that  $\hat{\beta}_{lts}^{n}$  and  $\beta_{lts}$  belong to a ball centered at  $\beta_{lts}$  with a large enough radius  $r_0$ ,  $B(\beta_{lts}, r_0)$ , and that  $\Theta = B(\beta_{lts}, r_0)$  is our parameter space of  $\beta$  hereafter. In order to apply the Lemma, we first realize that in our case,  $\hat{\beta}_{lts}^n$  and  $\beta_{lts}$  correspond to  $\tau_n$  and  $t_0$  (assume, w.l.o.g. that  $\beta_{lts} = 0$  in light of regression equivariance, see Section 4);  $\beta$  and  $\Theta$  correspond to t and T;  $f(\cdot,t) := f(\boldsymbol{x}^\top,y,\boldsymbol{\beta},\alpha) = (y - \boldsymbol{w}^\top\boldsymbol{\beta})^2 \mathbb{1}(r(\boldsymbol{\beta})^2 \leq F_{r(\boldsymbol{\beta})}^{-1}$  $r_{(\beta)^2}^{-1}(\alpha)$ ). In our case (see the proof (ii) of Lemma 2.2 in the Appendix),

$$
\nabla f(\cdot,t) := \nabla f(\boldsymbol{x},y,\boldsymbol{\beta},\alpha) = \frac{\partial}{\partial \boldsymbol{\beta}} f(\boldsymbol{x},y,\boldsymbol{\beta},\alpha) = -2(y - \boldsymbol{w}^\top \boldsymbol{\beta}) \boldsymbol{w} \mathbb{1} \left( r(\boldsymbol{\beta})^2 \leq F_{r(\boldsymbol{\beta})^2}^{-1}(\alpha) \right).
$$

We will have to assume that  $P(\nabla_i^2) = P(4(y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 w_i^2 1\mathbb{1}(r(\boldsymbol{\beta})^2 \leq F_{r(\boldsymbol{\beta})}^{-1})$  $\binom{n-1}{r(\boldsymbol{\beta})^2}(\alpha)$ ) exists to meet (iv) of the lemma, where  $i \in \{1, \dots, p\}$  and  $\mathbf{w}^{\top} = (1, \mathbf{x}^{\top}) = (1, x_1, \dots, x_{p-1})$ . It is readily seen that a sufficient condition for this assumption to hold is the existence of  $P(x_i^2)$ . In our case,  $V = 2P(\boldsymbol{w}\boldsymbol{w}^{\top}\boldsymbol{\mathbb{1}}(r(\boldsymbol{\beta})^2 \leq F_{r(\boldsymbol{\beta})}^{-1})$  $r_{(0)}^{-1}(\alpha)$ ), and we will have to assume that it is invertible when  $\beta$  is replaced by  $\beta_{lts}$  (this is covered by [\(18\)](#page-8-0)) to meet (ii) of the Lemma. In our case,

$$
r(\cdot, t) = \left(\frac{\boldsymbol{\beta}^{\top}}{\|\boldsymbol{\beta}\|} V/2 \frac{\boldsymbol{\beta}}{\|\boldsymbol{\beta}\|}\right) \|\boldsymbol{\beta}\|.
$$

We will assume that  $\lambda_{min}$  and  $\lambda_{max}$  are the minimum and maximum eigenvalues of positive semidefinite matrix V over all  $\beta \in \Theta$  and  $\alpha \in [1/2, c]$ .

#### Theorem 3.2 Assume that

(i) the uniqueness assumptions for  $\hat{\beta}_{lts}^{n}$  and  $\beta_{lts}$  in Theorems 2.1 and 2.3 hold respectively;

(ii)  $P(x_i^2)$  exists with  $\boldsymbol{x} = (x_1, \dots, x_{p-1});$ 

then

$$
n^{1/2}(\widehat{\beta}_{lts}^n - \beta_{lts}) \xrightarrow{d} N(\mathbf{0}, V^{-1}[P(\nabla \nabla^{\top}) - (P\nabla)(P\nabla)^{\top}]V^{-1}),
$$

where  $\beta$  in V and  $\nabla$  is replaced by  $\beta_{lts}$  (which could be assumed to be zero, or is  $\beta_0$  under Theorem 2.4).

**Proof:** The key to apply Lemma is to verify  $(v)$ , for details see the Appendix.

#### Remarks 3.1

(a) In the case of  $p = 1$ , that is in the location case, the asymptotic normality of the LTS has been studied in Butler (1982), RL87, Bednarski and Clarke (1993), and Mašíček (2004).

(b) Hössjer (1994), under the rank-based optimization framework and stringent assumptions on error term  $e_i$  (even density that is strictly decreasing for positive value, bounded absolute first moment) and on  $x_i$  (bounded fourth moment), covers the asymptotic normality of the LTS. V06c also treated the general case  $p \geq 1$  and obtained the asymptotic normality of the LTS under many stringent conditions on the non-random covariates  $x_i$ s and the distributions of  $e_i$  in a twenty-seven pages long article. The assumption  $\mathcal C$  is quite artificial and never verified there. On the other hand, C04 and C05 also addressed the asymptotic normality of LTS in the nonlinear regression under a dependence setting. For these extensions, many artificial assumptions  $(D1, D2, H1, H2, H3, H4, H5, H6, I1, I2)$  are imposed, but they are never verified for the linear LTS case. So those results do not cover the LTS in [\(3\)](#page-2-1).

(c) Furthermore, since there was no population version like [\(4\)](#page-4-3) and [\(5\)](#page-4-4) before, empirical process theory could not be employed to verify the VC class of functions in V06a, b, c, and C04, C05. Our approach here is quite different from former classical analyses and much more neat and concise (the standard empirical process theory was asserted to be non-applicable in C04 and C05 due to the lack of a population version of the LTS).  $\Box$ 

### <span id="page-13-0"></span>4 Inference procedures

In order to utilize the asymptotic normality result in Theorem 3.2, we need to figure out the asymptotic covariance. For simplicity, assume that  $\boldsymbol{z} = (\boldsymbol{x}^\top, y)^\top$  follows elliptical distributions  $E(g; \mu, \Sigma)$  with density

$$
f_{\boldsymbol{z}}(\boldsymbol{x}^{\top},y) = \frac{g(((\boldsymbol{x}^{\top},y)^{\top} - \boldsymbol{\mu})^{\top} \boldsymbol{\Sigma}^{-1}((\boldsymbol{x}^{\top},y)^{\top} - \boldsymbol{\mu}))}{\sqrt{\det(\boldsymbol{\Sigma})}},
$$

where  $\mu \in \mathbb{R}^p$  and  $\Sigma$  is a positive definite matrix of size p which is proportional to the covariance matrix if the latter exists. We assume the  $f_{\mathbf{z}}$  is unimodal.

#### Equivariance

A regression estimation functional  $t(\cdot)$  is said to be *regression*, scale, and affine equivariant (see Zuo (2021a)) if respectively

$$
t(F_{(\mathbf{w}, y+\mathbf{w}^{\top}b)}) = t(F_{(\mathbf{w}, y)}) + \mathbf{b}, \forall \mathbf{b} \in \mathbb{R}^{p};
$$
  
\n
$$
t(F_{(\mathbf{w}, sy)}) = st(F_{(\mathbf{w}, y)}), \forall \mathbf{s} \in \mathbb{R};
$$
  
\n
$$
t(F_{(\mathbf{A}^{\top}\mathbf{w}, y)}) = \mathbf{A}^{-1}t(F_{(\mathbf{w}, y)}), \forall \text{ nonsingular } \mathbf{A} \in \mathbb{R}^{p \times p};
$$

**Theorem 4.1**  $\beta_{lts}(F_{(\mathbf{x}',y)}, \alpha)$  is regression, scale, and affine equivariant.

**Proof:** See the empirical version treatment given in RL87 (p. 132).  $\Box$ 

**Transformation** Assume the Cholesky decomposition of  $\Sigma$  yields a non-singular lower triangular matrix  $\boldsymbol{L}$  of the form

$$
\left(\begin{array}{cc}A&0\\v^\top&c\end{array}\right)
$$

with  $\Sigma = LL^{\top}$ . Hence  $\det(A) \neq 0 \neq c$ . Now transfer  $(x^{\top}, y)$  to  $(s^{\top}, t)$  with  $(s^{\top}, t)^{\top} =$  $\boldsymbol{L}^{-1}((\boldsymbol{x}^\top,y)^\top-\boldsymbol{\mu})$ . It is readily seen that the distribution of  $(\boldsymbol{s}^\top,t)^\top$  follows  $E(g; \boldsymbol{0}, \boldsymbol{I}_{\boldsymbol{p}\times\boldsymbol{p}})$ .

Note that  $(\boldsymbol{x}^\top, y)^\top = \boldsymbol{L}(\boldsymbol{s}^\top, t)^\top + (\boldsymbol{\mu}_1^\top, \boldsymbol{\mu}_2)^\top$  with  $\boldsymbol{\mu} = (\boldsymbol{\mu}_1^\top, \boldsymbol{\mu}_2)^\top$ . That is,

$$
x = As + \mu_1,\tag{23}
$$

$$
y = \mathbf{v}^\top \mathbf{s} + ct + \mu_2. \tag{24}
$$

Equivalently,

$$
(1, \mathbf{s}^{\top})^{\top} = \mathbf{B}^{-1}(1, \mathbf{x}^{\top})^{\top},\tag{25}
$$

<span id="page-14-1"></span><span id="page-14-0"></span>
$$
t = \frac{y - (1, \mathbf{s}^{\top})(\mu_2, \mathbf{v}^{\top})^{\top}}{c},\tag{26}
$$

where

$$
B = \begin{pmatrix} 1 & \mathbf{0}^\top \\ \mu_1 & A \end{pmatrix}, \quad B^{-1} = \begin{pmatrix} 1 & \mathbf{0}^\top \\ -A^{-1} \mu_1 & A^{-1} \end{pmatrix},
$$

It is readily seen that  $(25)$  is an affine transformation on w and  $(26)$  is first an affine transformation on  $w$  then a regression transformation on  $y$  followed by a scale transformation on y. In light of Theorem 4.1, we can assume hereafter, w.l.o.g. that  $(x^{\top}, y)$  follows an  $E(g; 0, I_{p \times p})$  (spherical) distribution and  $I_{p \times p}$  is the covariance matrix of  $(\boldsymbol{x}^{\top}, y)$ .

**Theorem 4.2** Assume that  $e \sim \mathcal{N}(0, \sigma^2)$ , e and x are independent. Then

- (1)  $P\nabla = \mathbf{0}$  and  $P(\nabla \nabla') = 8\sigma^2 C \mathbf{I}_{p \times p}$ , with  $C = \Phi(c) - 1/2 - ce^{-c^2/2}/\sqrt{2\pi}$ , where  $c = \sqrt{F_{\chi^2(1)}^{-1}(\alpha)}$  and  $\Phi(x)$  is the CDF of  $\mathcal{N}(0,1)$ ,  $\chi^2(1)$  is a chi-square random variable with one degree of freedom.
- (2)  $V = 2C_1I_{p\times p}$  with  $C_1 = 2\Phi(c) 1$ .
- (3)  $n^{1/2}(\widehat{\beta}_{lts}^n \beta_{lts}) \stackrel{d}{\longrightarrow} \mathcal{N}(\mathbf{0}, \frac{2C\sigma^2}{C_1^2})$  $\frac{C\sigma^2}{C_1^2}I_{p\times p}$ , where C and C1 are defined in (1) and (2) above.

**Proof:** See the Appendix.

Approximate  $100(1 - \gamma)\%$  confidence region

(i) Based on the asymptotic normality Under the setting of Theorem 4.2, an approximate  $100(1-\gamma)$ % confidence region for the unknown regression parameter  $\beta_0$  is:

$$
\Big\{\boldsymbol{\beta}\in\mathbb{R}^p:\ \ \|\boldsymbol{\beta}-\widehat{\boldsymbol{\beta}}^n_{lts}\|\leq \sqrt{\frac{2C\sigma^2}{C_1^2n}}F^{-1}_{\chi^2(p)}(\gamma)\Big\},
$$

where  $\|\cdot\|$  stands for the Euclidean distance. Without asymptotic normality, one can appeal to the next procedure.

(ii) Based on bootstrapping scheme and depth-median and depth-quantile Here no assumptions on the underlying distribution are needed. This approximate procedure first

 $r_{\text{e}}$  re-samples *n* points with replacement from the given original sample points and calculates an  $\widehat{\beta}_{lts}^n$ . Repeat this m (a large number, say  $10^4$ ) times and obtain m such  $\widehat{\beta}_{lts}^n$ .

The next step is to calculate the depth, with respect to a location depth function (e.g., halfspace depth (Zuo (2018)) or projection depth (Zuo (2003) and Shao et al. (2022)), of these m points in the parameter space of  $\beta$ . Trimming  $|\gamma m|$  of the least deepest points among the m points, the left points form a convex hull, that is an approximate  $100(1 - \gamma)\%$ confidence region for the unknown regression parameter  $\beta_0$  (see Zuo (2009, 2010)) in the location and low dimensions cases).

# <span id="page-15-0"></span>5 Concluding remarks

Without the population version for the LTS (see [\(5\)](#page-4-4)), it will be difficult to apply the empirical process theory to study the asymptotics of the LTS, e.g, to verify the key result, the VC-class property of regression function class (indexed by  $\beta$ ) will be challenging. Without an explicit regression function (unlike the linear case), one can avoid this challenge, and simply assume this VC-class property. This is what exactly done in nonlinear setting LTS in C04, C05. The latters even believed that the standard empirical process theory is not applicable for the asymptotics of LTS while V06a, b, c, addressed the asymptotics, without any advanced tools, employed elementary tools with numerous artificial assumptions and lengthy articles.

By partitioning the parameter space and introducing the population version for the LTS, this article establishes some fundamental and primary properties for the objective function of the LTS in both empirical and population settings. These newly obtained original results verify some key facts and facilitate the application of standard empirical process theory to the establishment of asymptotic normality for the sample LTS in a concise and neat fashion. Some of newly obtained results, such as Fisher and strong consistency, and influence function, are original and obtained as by-products.

The asymptotic normality is applied in Theorem 4.2 for the practical inference procedure of confidence regions of the regression parameter  $\beta_0$ . There are open problems left here; one is the estimation of the variance of e, which is now unrealistically assumed to be known, and the other is the testing of hypothesis on  $\beta_0$ .

# Acknowledgments

Insightful comments and useful suggestions from Profs. Wei Shao and Derek Young have significantly improved the manuscript and are highly appreciated. Special thanks go to Prof. Derek Young for making the technical report of Chen, Stromberg, and Zhou available.

# Declarations

### Funding

This author declares that there is no funding received for this study.

### Conflicts of interests/Competing interests

This author declares that there is no conflict of interests/Competing interests.

# References

- $[1]$  Agullö, J. (2001), "New algorithms for computing the least trimmed squares regression estimator", Computational Statistics & Data Analysis 36 (2001) 425– 439
- [2] Bednarski, T. and Clarke, B.R. (1993), "Trimmed likelihood estimation of location and scale of the normal distribution". Austral. J. Statist. 35, 141–153.
- [3] Butler, R.W. (1982), "Nonparametric interval point prediction using data trimmed by a Grubbs type outlier rule". Ann. Statist. 10, 197–204.
- [4] Chen Y., Stromberg A., and Zhou M. (1997), "The least trimmed squares estimate in nonlinear regression". Technical report, 1997/365, Department of statistics, University of Kentucky.
- [5] Cížek, P. (2004), "Asymptotics of least trimmed squares regression", CentER Discussion Paper 2004-72, Tilburg University, The Netherlands
- [6] Čížek, P. (2005), "Least Trimmed Squares in nonlinear regression under dependence", Journal of Statistical Planning and Inference, 136, 3967-3988.
- <span id="page-16-0"></span>[7] Hampel, F. R., Ronchetti, E. M., Rousseeuw, P. J., and Stahel, W. A. (1986), Robust Statistics: The Approach Based on Influence Functions, John Wiley & Sons, New York.
- [8] Hawkins, D. M. (1994), "The feasible solution algorithm for least trimmed squares regression", Computational Statistics & Data Analysis, 17, 185-196.
- [9] Hawkins, D. M. and Olive, D. J. (1999), "Improved feasible solution algorithms for high breakdown estimation " Computational Statistics & Data Analysis,  $30(1)$ , 1-11.
- [10] Hofmann, M., Gatu, C., and Kontoghiorghes, E.J. (2010), "An Exact Least Trimmed Squares Algorithm for a Range of Coverage Values," Journal of Computational and Graphical Statistics, 19, 191-204
- [11] Hofmann, M., and Kontoghiorghes, E.J. (2010), "Matrix strategies for computing the least trimmed squares estimation of the general linear and SUR models", Computational Statistics & Data Analysis, 54, 3392-3403
- [12] Hössjer, O. (1994), "Rank-Based Estimates in the Linear Model with High Breakdown Point", J. Amer. Statist. Assoc., 89, 149-158.
- [13] Hössjer, O. (1995), "Exact computation of the least trimmed squares estimate in simple linear regression", Comput. Statist. Data Anal., 19, pp. 265-282
- [14] Huber, P. J. (1964), "Robust estimation of a location parameter", Ann. Math. Statist., 35 73-101.
- [15] Klouda, K. (2015), "An Exact Polynomial Time Algorithm for Computing the Least Trimmed Squares Estimate," Computational Statistics & Data Analysis, 84, 27-40.
- [16] Mašíček, L. (2004), "Optimality of the Least Weighted Squares Estimator," Kybernetika, 40, 715-734.
- $[17]$  Ollerer, V., Croux, C., and Alfons, A.  $(2015)$  "The influence function of penalized regression estimators", Statistics, 49:4, 741-765
- [18] Pollard, D. (1984), Convergence of Stochastic Processes, Springer, Berlin.
- [19] Rousseeuw, P. J. (1984), "Least median of squares regression", J. Amer. Statist. Assoc. 79, 871-880.
- [20] Rousseeuw, P. J., and Hubert, M. (1999), "Regression depth (with discussion)", J. Amer. Statist. Assoc., 94, 388–433.
- [21] Rousseeuw, P.J., and Leroy, A. Robust regression and outlier detection. Wiley New York. (1987).
- [22] Rousseeuw, P. J. and Van Driessen, K. (1999), "A fast algorithm for the minimum covariance determinant estimator", Technometrics, 41(3), 212-223.
- [23] Rousseeuw, P. J. and Van Driessen, K. (2006), "Computing LTS Regression for Large Data Sets", Data Mining and Knowledge Discovery 12, 29-45.
- [24] Rousseeuw, P. J. and Yohai, V. J. (1984), "Robust regression by means of S-estimators". In Robust and Nonlinear Time Series Analysis. Lecture Notes in Statist. Springer, New York. 26 256-272
- [25] Shao, W., Zuo, Y. and Luo, J. (2022), "Employing the MCMC Technique to Compute the Projection Depth in High Dimensions", Journal of Computational and Applied Mathematics, 411, 114278
- [26] Stromberg, A. J. (1993), "Computation of High Breakdown Nonlinear Regression Parameters", Journal of the American Statistical Association, 88:421, 237–244.
- <span id="page-17-0"></span>[27] Tableman, M. (1994), "The influence functions for the least trimmed squares and the least trimmed absolute deviations estimators", Statistics & Probability Letters 19 (1994) 329-337.
- [28] Víšek, J. Á. (2006a), The least trimmed squares. Part I: Consistency.  $Kybernetika$ , 42, 1-36.
- [29] Víšek, J. Á. (2006b) The least trimmed squares. Part II:  $\sqrt{n}$ -consistency. Kybernetika, 42, 181-202.
- [30] Víšek, J. A. (2006c), The least trimmed squares. Part III: Asymptotic normality. Kybernetika, 42, 203-224.
- [31] Yohai, V.J. (1987), "High breakdown-point and high efficiency estimates for regression", Ann. Statist., 15, 642–656.
- [32] Yohai, V.J. and Zamar, R.H. (1988), "High breakdown estimates of regression by means of the minimization of an efficient scale", J. Amer. Statist. Assoc., 83, 406-413.
- [33] Zuo, Y. (2003), "Projection-based depth functions and associated medians", Ann. Statist., 31, 1460-1490.
- [34] Zuo, Y. (2009), "Data depth trimming procedure outperforms the classical  $t$  (or  $T^2$ ) one", Journal of Probability and Statistics, Volume 2009, Article ID 373572, 9 pages.
- [35] Zuo, Y. (2010), "Is t procedure  $\bar{x} \pm t_{\alpha}(n-1)s/\sqrt{n}$  optimal?" The American Statistician, 64(2), 170-173.
- [36] Zuo, Y. (2018), "A new approach for the computation of halfspace depth in high dimensions". Communications in Statistics - Simulation and Computation, 48(3): 900-921.
- [37] Zuo, Y. (2020), "Large sample properties of the regression depth induced median", Statistics and Probability Letters, November 2020 166, arXiv1809.09896.
- [38] Zuo, Y. (2021a), "On general notions of depth for regression" Statistical Science 2021, Vol. 36, No. 1, 142–157, [arXiv:1805.02046.](http://arxiv.org/abs/1805.02046)
- <span id="page-18-0"></span>[39] Zuo, Y. (2021b), "Robustness of the deepest projection regression depth functional", Statistical Papers, vol. 62(3), pages 1167-1193.
- [40] Zuo, Y. and Zuo, H. (2023), "Least sum of squares of trimmed residuals regression", Electronic Journal of Statistics. 17 (2), 2447-2484, (2023) DOI: 10.1214/23-EJS2154, [arXiv:2202.10329](http://arxiv.org/abs/2202.10329)

# Appendix: Proofs

### Proof of Lemma 2.1

**Proof:** (i) Based on the definition in [\(9\)](#page-5-1), over  $S_{\beta}$ , there is no tie among the smallest h squared residuals. The assertion (a) follows straightforwardly.

The first and the last equality in (b) is trivial, it suffices to focus on the middle one. Let  $k_i := k_i(\boldsymbol{\eta}) \ (= k_i(\boldsymbol{\beta}^l)$  in light of [\(9\)](#page-5-1)). By [\(6\)](#page-4-5) we have that

$$
O^{n}(\eta) = \frac{1}{n} \sum_{i=1}^{h} r_{k_i}^{2}(\eta).
$$
 (27)

Let  $r_i := r_i(\eta) = y_i - \mathbf{w}_i^{\top} \eta$  and  $\gamma = \min\{\min_{1 \le i \ne j \le n} \{|r_i^2 - r_j^2|\}, 1\}$ . Then  $1 \ge \gamma > 0$  (a.s.).

Based on the continuity of  $r_{k_i}^2(\boldsymbol{\beta})$  in  $\boldsymbol{\beta}$ , for any  $1 \leq i \leq h$  and for any given  $\varepsilon \in (0,1)$ , we can fix a small  $\delta > 0$  so that  $|r_{k_i}^2(\boldsymbol{\beta}) - r_{k_i}^2(\boldsymbol{\eta})| < \gamma \varepsilon/4h$  for any  $\boldsymbol{\beta} \in B(\boldsymbol{\eta}, \delta)$ . Now we have for any  $\beta \in B(\eta, \delta)$ , (assume below  $2 \leq i \leq h$ )

$$
r_{k_i}^2(\boldsymbol{\beta}) - r_{k_{(i-1)}}^2(\boldsymbol{\beta}) > r_{k_i}^2(\boldsymbol{\eta}) - \frac{\gamma \varepsilon}{4h} - [r_{k_{(i-1)}}^2(\boldsymbol{\eta}) + \frac{\gamma \varepsilon}{4h}]
$$

$$
= r_{k_i}^2(\boldsymbol{\eta}) - r_{k_{(i-1)}}^2(\boldsymbol{\eta}) - \frac{\gamma \varepsilon}{2h}
$$

$$
\geq \gamma - \frac{\gamma \varepsilon}{2h} > 0 \ (a.s.),
$$

Thus,  $k_i = k_i(\eta)$ ,  $1 \leq i \leq h$  forms the h-integer set for any  $\beta \in B(\eta, \delta)$ . Part (b) follows.

(ii) The domain of function  $O^n(\beta)$  is the union of the pieces of  $\overline{S}_{\beta^l}$  and the function of  $O^n(\beta)$  over  $S_{\beta^l}$  is a quadratic function of  $\beta$ :  $O^n(\beta) = \sum_{j=1}^h r_k^2$  $\mathcal{L}_{k_j(\boldsymbol{\beta}^l)}^2(\boldsymbol{\beta}),$  the statement follows.

(iii) By (ii), it is clear that  $O^n(\beta)$  is continuous in  $\beta$  over each piece of  $S_{\beta}$ . We only need to show that this holds true for any  $\beta$  that is on the boundary of  $S_{\beta}$ .

Let  $\eta$  lie on the common boundary of  $S_{\beta^s}$  and  $S_{\beta^t}$ , then  $O^n(\beta) = \frac{1}{n} \sum_{i=1}^h r_{k_i(\beta^s)}^2$  for any  $\beta \in S_{\beta^s}$  [this is obviously true if  $\beta \in S_{\beta^s}$ , it is also true if  $\beta$  on the boundary of  $S_{\beta^s}$ since in this case the  $\beta$ -h-integer set is not unique, there are at least two, the one of them is  $k_1(\boldsymbol{\beta}^s), \cdots, k_h(\boldsymbol{\beta}^s)$  and  $O^n(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^h r_k^2$  $\mathcal{L}_{k_i(\boldsymbol{\beta}^t)}^2$  for any  $\boldsymbol{\beta} \in S_{\boldsymbol{\beta}^t}$ . Let  $\{\boldsymbol{\beta}_j\}$  be a sequence approaching  $\eta$ , where  $\beta_j$  could be on  $\overline{S}_{\beta^s}$  or on  $\overline{S}_{\beta^t}$ . We show that  $O^n(\beta_j)$  approaches to  $O^{n}(\eta)$ . Note that  $O^{n}(\eta) = \frac{1}{n} \sum_{i=1}^{h} r_{k_i(\beta^s)}^{2}(\eta) = \frac{1}{n} \sum_{i=1}^{h} r_{k_i}^{2}$  $\mathcal{L}^2_{k_i(\boldsymbol{\beta}^t)}(\boldsymbol{\eta}).$  Partition  $\{\boldsymbol{\beta}_j\}$  into  $\{\boldsymbol{\beta}_{j_s}\}$ and  $\{\beta_{j_t}\}\$  so that all members of the former belong to  $S_{\beta^s}$  where the latter are all within  $S_{\beta^t}$ . By continuity of the sum of h squared residuals in  $\beta$ , both  $O^n(\beta_{j_s})$  and  $O^n(\beta_{j_t})$  approach to  $O^n(\eta)$  since both  $\{\beta_{j_s}\}\$  and  $\{\beta_{j_t}\}\$ approach  $\eta$  as  $\min\{s,t\}\to\infty$ .

(iv) Note that for any  $l, 1 \leq l \leq L$ , over  $S_{\beta}l$ , one has a least squares problem with n reduced to h,  $O<sup>n</sup>(\beta)$  is a quadratic function and hence is twice differentiable and strictly convex in light of the following

$$
n\frac{\partial}{\partial \beta}O^n(\beta) = -2\sum_{i=1}^n r_i \mathbb{1}_i \mathbf{w}_i = -2\mathbf{X}_n^\top \mathbf{D} \mathbf{R},
$$
  

$$
n\frac{\partial^2}{\partial \beta^2}O^n(\beta) = 2\mathbf{X}_n^\top \mathbf{D} \mathbf{X}_n = 2\mathbf{X}^*_{n\top} \mathbf{X}^*_{n\top = 2\sum_{i=1}^h \mathbf{w}_{k_i(\beta^l)} \mathbf{w}_{k_i(\beta^l)}^\top,
$$

where  $\mathbf{R} = (r_1, r_2, \dots, r_n)^\top$ ,  $\mathbf{D} = \text{diag}(\mathbb{1}_i)$ , and  $\mathbf{X}_{\top}^* n = \mathbf{D}\mathbf{X}_n$ . Strict convexity follows from the positive definite of the Hessian matrix:  $\frac{2}{n}X^*_{n}^{\top}X^*_{n}$  (an invertible matrix due to (A1), see (iii) in the proof of Theorem 2.1).

### Proof of Theorem 2.1

**Proof**: (i) Over each  $S_{\beta}$ , an open set,  $O^{n}(\beta)$  is twice differentiable and strictly convex in light of given condition, hence it has a unique minimizer (otherwise, one can show that by openness and strictly convexity there is a third point in  $S_{\beta^l}$  that attains a strictly smaller objective value than the two minimizers). Since there are only finitely many  $S_{\beta^l}$ , the assertion follows if we can prove that the minimum does not reach at a boundary point of some  $S_{\beta}l$ .

Assume it is otherwise. That is,  $O<sup>n</sup>(\beta)$  reaches its global minimum at point  $\beta_1$  which is a boundary point of  $S_{\beta^l}$  for some l. Assume that over  $S_{\beta^l}$ ,  $O^n(\beta)$  attains its local minimum value at the unique point  $\beta_2$ . Then,  $O^n(\beta_1) \leq O^n(\beta_2)$ . If equality holds then we have the desired result (since there are points besides  $\beta_2$  in  $S_{\beta^l}$  which also attain the minimum value as  $\beta_2$ , a contradiction). Otherwise, there is a point  $\beta_3$  in the small neighborhood of  $\beta_1$  so that  $O^n(\beta_3) \leq O^n(\beta_1) + (O^n(\beta_2) - O^n(\beta_1))/2 < O^n(\beta_2)$ . A contradiction appears.

(ii) It is seen from (i) that  $O<sup>n</sup>(\beta)$  is twice continuously differentiable, hence its first derivative evaluated at the global minimum must be zero. By  $(i)$ , we have the equation [\(11\)](#page-6-0).

(iii) This part directly follows from (ii) and the invertibility of  $M_n$ . The latter follows from  $(A1)$  that implies that the p columns of matrix  $X_n$  are linearly independent and also implies that any h sub-rows of  $\mathbf{X}_n$  has a full rank.

#### Proof of Lemma 2.2

**Proof:** Denote the integrand in [\(4\)](#page-4-3) as  $G(\boldsymbol{\beta}) := (y - \boldsymbol{w}^{\top}\boldsymbol{\beta})^2 \mathbb{1} ((y - \boldsymbol{w}^{\top}\boldsymbol{\beta})^2 \leq q(\boldsymbol{\beta}, \alpha))$  for a given point  $(\boldsymbol{x}^\top, y) \in \mathbb{R}^p$ . Write  $G(\boldsymbol{\beta}) := (y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 \big(1 - \mathbb{1} \left( (y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 > q(\boldsymbol{\beta}, \alpha) \right) \big).$ 

(i) By the strictly monotonicity of  $F_W$  around  $q(\beta, \alpha)$ , we have the continuity of the  $q(\boldsymbol{\beta}, \alpha)$ . Consequently,  $G(\boldsymbol{\beta})$  is obvious continuous and so is  $O(\boldsymbol{\beta})$  in  $\boldsymbol{\beta} \in \mathbb{R}^p$ .

(ii) For arbitrary points  $(x^{\top}, y)$  and  $\beta$  in  $\mathbb{R}^p$ , there are three cases for the relationship between the squared residual and its quantile: (a)  $(y - w^{\top} \beta)^2 > q(\beta, \alpha)$  (b)  $(y - w^{\top} \beta)^2 <$  $q(\boldsymbol{\beta}, \alpha)$  and (c)  $(y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 = q(\boldsymbol{\beta}, \alpha)$ . Case (c) happens with probability zero, we thus skip this case and treat (a) and (b) only. By the continuity in  $\beta$ , there is a small neighborhood of  $\beta$ :  $B(\beta, \delta)$ , centered at  $\beta$  with radius  $\delta$  such that (a) (or (b)) holds for all  $\beta \in B(\beta, \delta)$ . This implies that

$$
\frac{\partial}{\partial \boldsymbol{\beta}} \mathbb{1} \left( (y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 > q(\boldsymbol{\beta}, \alpha) \right) = \mathbf{0}, \ (a.s.)
$$

and

$$
\frac{\partial}{\partial \boldsymbol{\beta}} G(\boldsymbol{\beta}) = -2(y - \boldsymbol{w}^\top \boldsymbol{\beta}) \boldsymbol{w} \mathbb{1} \left( (y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 \le q(\boldsymbol{\beta}, \alpha) \right), (a.s).
$$

Hence, we have that

$$
\frac{\partial^2}{\partial \boldsymbol{\beta}^2} G(\boldsymbol{\beta}) = 2 \mathbf{w} \mathbf{w}^\top \mathbb{1} \left( (y - \mathbf{w}^\top \boldsymbol{\beta})^2 \le q(\boldsymbol{\beta}, \alpha) \right), \ (a.s).
$$

Note that  $E(\boldsymbol{w}\boldsymbol{w}^{\top})$  exists. Then, by the Lebesgue dominated convergence theorem, the desired result follows.

(iii) The strict convexity follows from the twice differentiability and the positive definite of the second order derivative of  $O(\beta)$ .

### Proof of Theorem 2.3

**Proof:** We will treat  $\beta_{lts}(F_{(\mathbf{z}^{\top}, y)}, \alpha)$ , the counterpart for  $\beta_{lts}(F_{\varepsilon}(z), \alpha)$  can be treated analogously.

(i) Existence follows from the positive semi-definiteness of the Hessian matrix (see proof of (ii) of Lemma 2.2) and the convexity of  $O(\beta)$ .

(ii) The equation follows from the differentiability and the first order derivative of  $O(\beta)$ given in the proof (ii) of Lemma 2.2.

(iii) The uniqueness follows from the positive definite of the Hessian matrix based on the given condition (invertibility).  $\Box$ 

### Proof of Theorem 2.4

**Proof:** By theorem 2.3, (i) and given conditions guarantee the existence and the uniqueness of  $\beta_{lts}(F_{(\mathbf{x}^{\top},y)},\alpha)$  which is the unique solution of the system of the equations

$$
\int (y - \boldsymbol{w}^\top \boldsymbol{\beta}) \boldsymbol{w} \mathbb{1} \left( (y - \boldsymbol{w}^\top \boldsymbol{\beta})^2 \le q(\boldsymbol{\beta}, \alpha) \right) dF_{(\boldsymbol{x}^\top, y)}(\boldsymbol{x}, y) = \mathbf{0}.
$$

Notice that  $y - \boldsymbol{w}^\top \boldsymbol{\beta} = -\boldsymbol{w}^\top (\boldsymbol{\beta} - \boldsymbol{\beta}_0) + e$ . Inserting this into the above equation we have

$$
\int (-\boldsymbol{w}^\top(\boldsymbol{\beta}-\boldsymbol{\beta}_0)+e)\boldsymbol{w}1\left((-\boldsymbol{w}^\top(\boldsymbol{\beta}-\boldsymbol{\beta}_0)+e)^2\leq F_{(-\boldsymbol{w}^\top(\boldsymbol{\beta}-\boldsymbol{\beta}_0)+e)^2}^{-1}(\alpha)\right)dF_{(\boldsymbol{x}^\top,y)}(\boldsymbol{x},y)=\mathbf{0}.
$$

By (ii) it is readily seen that  $\beta = \beta_0$  is a solution of the above system of equations. Uniqueness leads to the desired result.

### Proof of Theorem 2.5

**Proof:** Write  $\beta_{lts}^{\varepsilon}(\mathbf{z}_0)$  for  $\beta_{lts}(F_{\varepsilon}(\mathbf{z}_0), \alpha)$  and insert it for  $\beta$  into [\(17\)](#page-8-1) and take derivative with respect to  $\varepsilon$  in both sides of [\(17\)](#page-8-1) and let  $\varepsilon \to 0$ , we obtain (in light of dominated theorem)

$$
\left(\int \frac{\partial}{\partial \beta_{lts}^{\varepsilon}(\mathbf{z}_0)} r(\beta_{lts}^{\varepsilon}(\mathbf{z}_0)) \mathbf{v} \mathbb{1} \left(r(\beta_{lts}^{\varepsilon}(\mathbf{z}_0))^2 \leq q_{\varepsilon}(\mathbf{z}, \beta_{lts}^{\varepsilon}(\mathbf{z}_0), \alpha)\right) \Big|_{\varepsilon \to 0} dF_{(\mathbf{x}^\top y)}\right) \text{IF}(\mathbf{z}_0, \beta_{lts}, F_{(\mathbf{x}^\top, y)})
$$
  
+ 
$$
\int r(\beta_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha))) \mathbf{w} \mathbb{1} \left(r(\beta_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha)))\right)^2 \leq q(\beta_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha), \alpha)\right) d(\delta_{\mathbf{z}_0} - F_{(\mathbf{x}^\top, y)})
$$
  
= **0**, (28)

where  $r(\beta) = t - v^\top \beta$  in the first term on the LHS and  $r(\beta) = y - w^\top \beta$  in the second term on the LHS. Call the two terms on the LHS as  $T_1$  and  $T_2$  respectively and call the integrand in  $T_1$  as  $T_0$ , then it is seen that (see the proof (i) of Theorem 2.1)

<span id="page-21-0"></span>
$$
T_0 = \frac{\partial}{\partial \beta_{lts}^{\varepsilon}(\mathbf{z}_0)} (t - \mathbf{v}^\top \beta_{lts}^{\varepsilon}(\mathbf{z}_0)) \mathbf{v} \mathbb{1} \left( (t - \mathbf{v}^\top \beta_{lts}^{\varepsilon}(\mathbf{z}_0))^2 \le q_{\varepsilon}(\mathbf{z}, \beta_{lts}^{\varepsilon}(\mathbf{z}_0), \alpha) \right) \Big|_{\varepsilon \to 0}
$$
  
= - $\mathbf{w} \mathbf{w}^\top \mathbb{1} \left( (y - \mathbf{w}^\top \beta_{lts})^2 \le q(\beta_{lts}, \alpha) \right).$ 

Focus on the  $T_2$ , it is readily seen that

$$
T_2 = \int (y - \mathbf{w}^\top \boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha)) \mathbf{w} \mathbb{1} \left( (y - \mathbf{w}^\top \boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha))^2 \leq q(\boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha), \alpha) \right) d\delta_{\mathbf{z}_0}
$$

$$
- \int (y - \mathbf{w}^\top \boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha)) \mathbf{w} \mathbb{1} \left( (y - \mathbf{w}^\top \boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha))^2 \leq q(\boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha), \alpha) \right) dF_{(\mathbf{x}^\top, y)},
$$

In light of [\(16\)](#page-8-2) we have

$$
T_2 = \int (y - \mathbf{w}^\top \boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha)) \mathbf{w} \mathbb{1} ((y - \mathbf{w}^\top \boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha))^2 \le q(\boldsymbol{\beta}_{lts}(F_{(\mathbf{x}^\top, y)}, \alpha), \alpha)) d\delta_{\mathbf{z}_0}
$$
  
= 
$$
\begin{cases} \mathbf{0}, & \text{if } (t_0 - \mathbf{v}_0^\top \boldsymbol{\beta}_{lts})^2 > q(\boldsymbol{\beta}_{lts}, \alpha), \\ (t_0 - \mathbf{v}_0^\top \boldsymbol{\beta}_{lts}) \mathbf{v}_0, & \text{otherwise.} \end{cases}
$$

This,  $T_0$ , and display [\(28\)](#page-21-0) lead to the desired result.  $\Box$ 

### Proof of Lemma 3.1

Proof: We invoke Theorem 24 in II.5 of Pollard (1984) (P84). The first requirement of the theorem is the existence of an envelope of  $\mathscr{F}$ . The latter is  $\sup_{\beta \in \Theta} F^{-1}_{r(\beta)}$  $r^{-1}(\beta)^2(c)$ , which is bounded since  $\Theta$  is compact and  $F_W^{-1}$  is continuous in  $\beta$ , and  $F_W^{-1}(\alpha)$  is non-decreasing in

 $\alpha \in [1/2, c]$ . To complete the proof, we only need to verify the second requirement of the theorem.

For the second requirement, that is, to bound the covering numbers, it suffices to show that the graphs of functions in  $\mathcal{F}(\beta,\alpha)$  have only polynomial discrimination (see Theorem 25 and Example 26 in II.5 of P84).

The graph of a real-valued function f on a set S is defined as the subset (see p. 27 of P84)

$$
G_f = \{(s, t) : 0 \le t \le f(s) \text{ or } f(s) \le t \le 0, s \in S\}.
$$

The graph of a function in  $\mathscr{F}(\beta,\alpha)$  contains a point  $(\mathbf{x}^{\top}(\omega),y(\omega),t)$  if and only if  $0 \le t \le$  $f(x, y, \beta, \alpha)$  or  $f(x, y, \beta, \alpha) \le t \le 0$ . The latter case could be excluded since the function is always nonnegative (and equals 0 case covered by the former case). The former case happens if and only if  $0 \le \sqrt{t} \le y - w^\top \beta$  or  $0 \le \sqrt{t} \le -y + w^\top \beta$ .

Given a collection of n points  $(x_i^{\top}, y_i, t_i)$   $(t_i \geq 0)$ , the graph of a function in  $\mathscr{F}(\beta, \alpha)$ picks out only points that belong to  $\{\sqrt{t_i} \geq 0\} \cap \{y_i - \beta^\top w_i - \sqrt{t_i} \geq 0\} \cup \{\sqrt{t_i} \geq 0\} \cap \{-y_i + \beta^\top w_i\}$  $\mathcal{B}^\top w_i - \sqrt{t_i} \geq 0$ . Introduce n new points  $(w_i^\top, y_i, z_i) := ((1, x_i^\top), y_i, \sqrt{t_i})$  in  $\mathbb{R}^{p+2}$ . On  $\mathbb{R}^{p+2}$ define a vector space  $\mathscr G$  of functions

$$
g_{a,b,c}(\boldsymbol{w},y,z) = \mathbf{a}^\top \boldsymbol{w} + b y + c z,
$$

where  $a \in \mathbb{R}^p$ ,  $b \in \mathbb{R}^1$ , and  $c \in \mathbb{R}^1$  and  $\mathscr{G} := \{g_{a,b,c}(\boldsymbol{w},y,z) = \mathbf{a}^\top \boldsymbol{w} + by + cz, a \in \mathbb{R}^p, b \in \mathbb{R}^1$ , and  $c \in \mathbb{R}^1$  which is a  $\mathbb{R}^{p+2}$ -dimensional vector space.

It is clear now that the graph of a function in  $\mathcal{F}(\beta,\alpha)$  picks out only points that belong to the sets of  $\{g \geq 0\}$  for  $g \in \mathscr{G}$  (ignoring the union and intersection operations at this moment). By Lemma 18 in II.4 of P84 (p. 20), the graphs of functions in  $\mathscr{F}(\beta,\alpha)$  pick only polynomial numbers of subsets of  $\{\bm p_i:=(\bm w_i^\top,y_i,z_i), i\in\{1,\cdots,n\}\}$ ; those sets corresponding to  $g \in \mathscr{G}$  with  $a \in \{0, -\beta, \beta\}, b \in \{0, 1, -1\}, \text{ and } c \in \{1, -1\}$  pick up even few subsets from  $\{p_i, i \in \{1, \dots, n\}\}\.$  This in conjunction with Lemma 15 in II.4 of P84 (p. 18), yields that the graphs of functions in  $\mathcal{F}(\beta,\alpha)$  have only polynomial discrimination.

By Theorem 24 in II.5 of P84 we have completed the proof.

### Proof of Theorem 3.2

**Proof**: To apply Lemma 3.2, we need to verify the five conditions, among them only (iii) and (v) need to be addressed, all others are satisfied trivially. For (iii), it holds automatically since our  $\tau_n = \widehat{\beta}_{lts}^n$  is defined to be the minimizer of  $F_n(t)$  over  $t \in T(=\Theta)$ .

So the only condition that needs to be verified is the  $(v)$ , the stochastic equicontinuity of  $\{E_n r(\cdot, t)\}\$ at  $t_0$ . For that, we will appeal to the Equicontinuity Lemma (VII.4 of P84, p. 150). To apply the Lemma, we will verify the condition for the random covering numbers satisfy the uniformity condition. To that end, we look at the class of functions

$$
\mathscr{R}(\boldsymbol{\beta},\alpha) = \left\{ r(\cdot,\cdot,\alpha,\boldsymbol{\beta}) = \left( \frac{\boldsymbol{\beta}^{\top}}{\|\boldsymbol{\beta}\|} V/2 \frac{\boldsymbol{\beta}}{\|\boldsymbol{\beta}\|} \right) \|\boldsymbol{\beta}\| : \ \boldsymbol{\beta} \in \Theta, \alpha \in [1/2,c] \right\}.
$$

Obviously,  $\lambda_{max} r_0/2$  is an envelope for the class  $\mathscr{R}$  in  $\mathscr{L}^2(P)$ , where  $r_0$  is the radius of the ball  $\Theta = B(\beta_{lt}, r_0)$ . We now show that the covering numbers of  $\mathscr R$  is uniformly bounded, which amply suffices for the Equicontinuity Lemma. For this, we will invoke Lemmas II.25 and II.36 of P84. To apply Lemma II.25, we need to show that the graphs of functions in  $\mathscr R$ have only polynomial discrimination. The graph of  $r(x, y, \alpha, \beta)$  contains a point  $(x^{\top}, y, t)$ ,  $t \geq 0$  if and only if  $\left(\frac{\boldsymbol{\beta}^{\top}}{\|\boldsymbol{\beta}\|}\right)$  $\frac{\boldsymbol{\beta}^\top}{\|\boldsymbol{\beta}\|}V/2\frac{\boldsymbol{\beta}}{\|\boldsymbol{\beta}\|}$  $\|\beta\|$  $\Big|\|\beta\|\geq t$  for all  $\beta\in\Theta$  and  $\alpha\in[1/2,c].$ 

Equivalently, the graph of  $r(x, y, \alpha, \beta)$  contains a point  $(x^{\top}, y, t)$ ,  $t \geq 0$  if and only if  $\lambda_{min}/2\|\boldsymbol{\beta}\| \geq t$ . For a collection of n points  $(\boldsymbol{x}_i^{\top}, y_i, t_i)$  with  $t_i \geq 0$ , the graph picks out those points satisfying  $\lambda_{min}/2\|\boldsymbol{\beta}\| - t_i \geq 0$ . Construct from  $(\boldsymbol{x}_i^{\top}, y_i, t_i)$  a point  $z_i = t_i$  in  $\mathbb{R}$ . On  $\mathbb{R}$ define a vector space  $\mathscr G$  of functions

$$
g_{a,b}(x) = ax + b, \quad a, \ b \in \mathbb{R}.
$$

By Lemma 18 of P84, the sets  $\{g \geq 0\}$ , for  $g \in \mathscr{G}$ , pick out only a polynomial number of subsets from  $\{z_i\}$ ; those sets corresponding to functions in  $\mathscr G$  with  $a = -1$  and  $b = \lambda_{min}/2||\boldsymbol{\beta}||$ pick out even fewer subsets from  $\{z_i\}$ . Thus the graphs of functions in  $\mathcal R$  have only polynomial discrimination discrimination.

### Proof of Theorem 4.2:

**Proof:** In order to invoke Theorem 3.2, we only need to check the uniqueness of  $\hat{\beta}_{lts}^{n}$  and  $\beta_{lts}$ . The former is guaranteed by the (iii) of Theorem 2.1 since (A1) holds true a.s.. This is because that any p columns of the  $X_n$  or any its h rows could be regarded as a sample from an absolutely continuous random vector with dimension n or  $h$ . The probability that these p points lie in a  $(p-1)$  dimensional non-degenerated hyperplane (the normal vector is non-zero) is zero.

The latter is guaranteed by the (iii) of Theorem 2.3 since  $W = (y - \boldsymbol{w}^\top \boldsymbol{\beta})^2$  is the square of a normal distribution with mean  $-\beta_1$  and hence has a positive density and furthermore (18) becomes  $2(\Phi(c/\sigma) - 1/2)I_{p\times p}$  hence is invertible, where c is defined in Theorem 4.2. By Theorems 4.1 and 2.4, we can assume, w.l.o.g., that  $\beta_{lts} = \beta_0 = 0$ . Utilizing the independence between e and  $x$  and Theorem 3.2, a straightforward calculation leads to the results.  $\Box$