A new matrix equation expression for the solution of non-autonomous linear systems of ODEs

Stefano Pozza[†] Niel Van Buggenhout[†]

Abstract

The solution of systems of non-autonomous linear ordinary differential equations is crucial in a variety of applications, such us nuclear magnetic resonance spectroscopy. A new method with spectral accuracy has been recently introduced in the scalar case. The method is based on a product that generalizes the convolution. In this work, we show that it is possible to extend the method to solve systems of non-autonomous linear ordinary differential equations (ODEs). In this new approach, the ODE solution can be expressed through a linear system that can be equivalently rewritten as a matrix equation. Numerical examples illustrate the method's efficacy and the low-rank property of the matrix equation solution.

1 Introduction

Systems of non-autonomous linear ordinary differential equations (ODEs) appear in a variety of applications, and its numerical computation is often challenging, particularly for large-to-huge size systems. For instance, in nuclear magnetic resonance spectroscopy (NMR) [1], the system solution describes the dynamics of the nuclear spins of a sample in a time-varying magnetic field. The size of such systems is $2^k \times 2^k$ for a sample with k spins and is usually sparse. In [2], we proposed a new method with spectral accuracy for solving *scalar* non-autonomous ordinary differential equations. In the present work, we extend this method to the case of systems of non-autonomous ODEs.

Consider a matrix $\tilde{A}(t) \in \mathbb{C}^{N \times N}$ composed of elements from $C^{\infty}(\mathcal{I})$, i.e., the set of functions infinitely differentiable (smooth) over \mathcal{I} , with \mathcal{I} a closed and bounded interval in \mathbb{R} . The system

$$\frac{d}{dt}U_s(t) = \tilde{A}(t)U_s(t), \quad U_s(s) = I_N, \quad \text{for } t \ge s, \quad t, s \in \mathcal{I},$$
(1)

has a unique solution $U_s(t) \in \mathbb{C}^{N \times N}$; I_N stands for the $N \times N$ identity matrix. Note that the condition $U_s(s) = I_N$ is not restrictive, since, given a matrix $B \in \mathbb{C}^{N \times N}$, the matrix-valued function $V_s(t) := U_s(t)B$ solves the ODE

$$\frac{d}{dt}V_s(t) = \tilde{A}(t)V_s(t), \quad V_s(s) = B \quad \text{for } t \ge s, \quad t, s \in \mathcal{I}.$$

At the heart of the new method for solving (1) is a non-commutative convolution-like product, denoted by \star , defined between certain distributions [3]. Thanks to this product, the solution of (1) can be expressed through the \star -product inverse and its formulation as a sequence of integrals and differential equations; see [4–8]. In [2], we illustrated that, by discretizing the \star -product with orthogonal functions, the solution of a scalar ODE is accessible by solving a linear system. In this work, we extend the results in [2], showing that, following the same principles, we can solve (1) through a linear system. Moreover, we show that the linear system solution can be expressed as the solution of a matrix equation with a rank one right-hand side. Numerical experiments illustrate that the solution of the matrix equation can also be low-rank.

In Section 2, we recall the \star -product definition and the related expression for the solution of an ODE. The expression is then discretized and approximated by the solution of a linear system. Section 3 shows how to transform the linear system into a matrix equation, and Section 4 concludes the paper.

[†]Charles University, Sokolovská 83 186, 75 Praha 8, Czech Republic. (pozza@karlin.mff.cuni.cz, buggenhout@karlin.mff.cuni.cz)

This work was supported by Charles University Research programs No. PRIMUS/21/SCI/009 and UNCE/SCI/023, and by the Magica project ANR-20-CE29-0007 funded by the French National Research Agency.

2 Solution of an ODE by the *-product

We use the Heaviside theta function

$$\Theta(t-s) = \begin{cases} 1, & t \ge s \\ 0, & t < s \end{cases}$$

to rewrite (1) in the following equivalent form

$$\frac{d}{dt}U(t,s) = \tilde{A}(t)\Theta(t-s)U(t,s), \quad U(s,s) = I_N, \quad \text{for } t, s \in \mathcal{I}.$$
(2)

Note that $\Theta(t-s)$ endows the condition $t \ge s$ in equation (2) and that U(t,s) is the bivariate function expressing the solutions of (1) for every initial time $s \in \mathcal{I}$, with U(t,s) = 0 for t < s. From now on, we will denote with a tilde all the bivariate functions that are infinitely differentiable in both t and s over \mathcal{I} , i.e., $\tilde{f} \in C^{\infty}(\mathcal{I} \times \mathcal{I})$. Moreover, we define the following class of functions

$$C^{\infty}_{\Theta}(\mathcal{I}) := \left\{ f : f(t,s) = \tilde{f}(t,s)\Theta(t-s), \quad \tilde{f} \in C^{\infty}(\mathcal{I} \times \mathcal{I}) \right\}.$$

Consider now the $N \times N$ matrices $A_1(t, s), A_2(t, s) \in (C_{\Theta}^{\infty}(\mathcal{I}))^{N \times N}$, i.e., matrices composed of elements from $C_{\Theta}^{\infty}(\mathcal{I})$. Then, the \star -product is defined as

$$(A_2 \star A_1)(t,s) := \int_{\mathcal{I}} A_2(t,\tau) A_1(\tau,s) \,\mathrm{d}\tau.$$
(3)

The *-product can be extended to a larger class of matrices composed of elements from the class $D(\mathcal{I}) \supset C_{\Theta}^{\infty}(\mathcal{I})$, that is, the class of the superpositions of $\Theta(t-s)$, Dirac delta distribution $\delta(t-s)$, and Dirac delta derivatives described in [6]. In such a class, $\delta(t-s)I_N$ is the *-product identity, i.e., $A(t,s) \star \delta(t-s)I_N = \delta(t-s)I_N \star A(t,s) = A(t,s)$. Moreover, in the larger class $\mathcal{D}(\mathcal{I})$, the *-product admits inverses under certain conditions [6], i.e., for certain $f(t,s) \in C_{\Theta}^{\infty}$, there exists $f(t,s)^{-*}$ such that $f(t,s) \star f(t,s)^{-*} = f(t,s)^{-*} \star f(t,s) = \delta(t-s)$.

Following [4], the solution of (2) can be expressed as

$$U(t,s) = \Theta(t-s) \star R_{\star}(A)(t,s), \tag{4}$$

where $A(t,s) = \tilde{A}(t)\Theta(t-s)$ and $R_{\star}(A)$ is the \star -resolvent of A, i.e.,

$$R_{\star}(A)(t,s) = \delta(t-s)I_N + \sum_{k=1}^{\infty} A(t,s)^{k\star},$$

with $A(t,s)^{k*} = A \star \cdots \star A$, the kth power of the *-product. Note that the series $\sum_{k=1}^{\infty} A(t,s)^{*k}$ converges for every $A \in (C_{\Theta}^{\infty}(\mathcal{I}))^{N \times N}$. Expression (4) hides an infinite series of nested integrals. However, as shown in [2], it is possible to approximate the *-product by the usual matrix-matrix product in the scalar case. This approximation allows us to compute (4) more simply and cheaply. We recall its basics below.

Without loss of generality, we set $\mathcal{I} = [0, 1]$. Moreover, we consider the family of orthonormal shifted Legendre polynomials $\{p_k\}_k$. Then, any $f(t, s) \in C^{\infty}_{\Theta}(\mathcal{I})$ can be expanded into the following series (e.g., [9])

$$f(t,s) = \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} f_{k,\ell} p_k(t) p_\ell(s), \ t \neq s, \ t, s \in \mathcal{I}, \quad f_{k,\ell} = \int_{\mathcal{I}} \int_{\mathcal{I}} f(\tau,\rho) p_k(\tau) p_\ell(\rho) \ d\rho \ d\tau.$$
(5)

By defining the *coefficient matrix* F_M and the vector $\phi_M(t)$ as

$$F_M := \begin{bmatrix} f_{0,0} & f_{0,1} & \dots & f_{0,M-1} \\ f_{1,0} & f_{1,1} & \dots & f_{1,M-1} \\ \vdots & \vdots & & \vdots \\ f_{M-1,0} & f_{M-1,1} & \dots & f_{M-1,M-1} \end{bmatrix}, \quad \phi_M(t) := \begin{bmatrix} p_0(s) \\ p_1(s) \\ \vdots \\ p_{M-1}(s) \end{bmatrix},$$
(6)

the truncated expansion series can be written in the matrix form:

$$f_M(t,s) := \sum_{k=0}^{M-1} \sum_{\ell=0}^{M-1} f_{k,\ell} \, p_k(t) p_\ell(s) = \phi_M(t)^T F_M \, \phi_M(s).$$

Let us consider the functions $f, g, h \in C^{\infty}_{\Theta}(\mathcal{I})$ so that $h = f \star g$, and the related coefficient matrices (6), respectively, F_M, G_M, H_M . Following [6], H_M can be approximated by the expression

$$H_M \approx \dot{H}_m := F_M G_M. \tag{7}$$

Therefore, there is a connection between the \star -algebra over $\mathcal{D}(\mathcal{I})$ and the usual matrix algebra. The elements and operations which form the \star -algebra and the related elements and operations forming the usual matrix algebra are given in Table 1 (in the first two columns for the scalar case); for more details, we refer to [6].

The approximation in the scalar case can be easily extended to the matrix one. Indeed, if $A(t,s) = [a_{ij}(t,s)]_{i,j=1}^N$ is an $N \times N$ matrix with elements $a_{ij}(t) \in C_{\Theta}^{\infty}(\mathcal{I})$, then for each a_{ij} , we can compute the related coefficient matrices $F_M^{(i,j)}$ (6) obtaining the block matrix

$$\mathcal{A}_{M} = \begin{bmatrix} F_{M}^{(1,1)} & \cdots & F_{M}^{(1,N)} \\ \vdots & \ddots & \vdots \\ F_{M}^{(N,1)} & \cdots & F_{M}^{(N,N)} \end{bmatrix} \in \mathbb{C}^{MN \times MN}.$$

$$(8)$$

Let us define the $N \times N$ matrices $A(t,s), B(t,s), C(t,s) \in (C_{\Theta}^{\infty}(\mathcal{I}))^{N \times N}$ so that $C(t,s) = A(t,s) \star B(t,s)$ and let their coefficient matrices (8) be, respectively, $\mathcal{A}_M, \mathcal{B}_M, \mathcal{C}_M$. Then, analogously to the scalar case, \mathcal{C}_M is approximated by

$$\mathcal{C}_M \approx \hat{\mathcal{C}}_M := \mathcal{A}_M \mathcal{B}_M.$$

As a consequence, also in the matrix case, the \star -algebra can be approximated by the usual matrix algebra, as summarized in the last two columns of Table 1.

$f(t,s) \in C^{\infty}_{\Theta}(\mathcal{I})$	$F_M \in \mathbb{C}^{M \times M}$	$A(t,s) \in (C^{\infty}_{\Theta}(\mathcal{I}))^{N \times N}$	$\mathcal{A}_M \in \mathbb{C}^{MN imes MN}$
*-operation/elements	matrix operation/elements	*-operation/elements	matrix operation/elements
$q = f \star g$	$Q_M = F_M G_M$	$C = A \star B$	$\mathcal{C}_M = \mathcal{A}_M \mathcal{B}_M$
f + g	$F_M + G_M$	A + B	$\mathcal{A}_M + \mathcal{B}_M$
$1_\star := \delta(t-s)$	I_M , identity matrix	$1_{\star} := \delta(t-s)I_N$	I_{MN} , identity matrix
$f^{\star-1}$	F_M^{-1}	$A^{\star -1}$	\mathcal{A}_M^{-1}
$R_{\star}(f) := (1_{\star} - f)^{\star - 1}$	$R(F_M) := (I_M - F_M)^{-1}$	$R_{\star}(f) := (1_{\star} - A)^{\star - 1}$	$R(\mathcal{A}_M) := (I_{MN} - \mathcal{A}_M)^{-1}$

Table 1: The *-algebra operations and the corresponding matrix algebra operation after discretization, scalar case (first two columns), matrix case (last two columns).

The matrix-valued function U(t, s) in (2) is composed of elements from $C_{\Theta}^{\infty}(\mathcal{I})$. Therefore, we can define the related coefficient matrix \mathcal{U}_M as in (8). Then, expression (4) can be approximated by

$$\mathcal{U}_M \approx (I_N \otimes T_M)(I_{MN} - \mathcal{A}_M)^{-1},$$

where \otimes is the Kronecker product, T_M is the coefficient matrix of $\Theta(t-s)$, and \mathcal{A}_M is the coefficient matrix of $\tilde{A}(t)\Theta(t-s)$, with $\tilde{A}(t)$ from (2). Moreover, we can approximate the solution of (2) for s = 0 by the formula:

$$U(t,0) \approx \phi_M(t)^T \mathcal{U}_M \phi_M(0) = (I_N \otimes \phi_M(t)^T) (I_N \otimes T_M) (I_{MN} - \mathcal{A}_M)^{-1} (I_N \otimes \phi_M(0))$$
$$= (I_N \otimes \phi_M(t)^T T_M) (I_{MN} - \mathcal{A}_M)^{-1} (I_N \otimes \phi_M(0)).$$

Note that, as explained in [2], the approximation converges quickly enough to the solution only when s is the left endpoint of the interval \mathcal{I} , i.e., s = 0.

In practical situations, the initial time s of the evolution is fixed (s = 0), and the initial condition is given as a vector $v \in \mathbb{C}^N$. Then, we get the simpler problem,

$$\frac{d}{dt}u(t) = \tilde{A}(t)\Theta(t-s)u(t), \quad u(0) = v, \quad \text{for } t, s \in \mathcal{I},$$
(9)

where the solution u(t) is an N-size vector. Thus, u(t) is approximated by:

$$u(t) \approx (I_N \otimes \phi_M(t)^T T_M) (I_{MN} - \mathcal{A}_M)^{-1} (I_N \otimes \phi_M(0)) v$$

$$\approx (I_N \otimes \phi_M(t)^T T_M) (I_{MN} - \mathcal{A}_M)^{-1} (v \otimes \phi_M(0)).$$

Then, solving the linear system

$$(I_{MN} - \mathcal{A}_M)x = v \otimes \phi_M(0), \tag{10}$$

one can approximate the solution of (9) in terms of its expansion coefficients $u_M := (I_N \otimes T_M)x$, that is,

$$u(t) \approx \hat{u}(t) := (I_N \otimes \phi_M(t)^T) u_M.$$
(11)

2.1 Numerical examples

Given a random vector v with elements in [0, 1], we aim to compute the bilinear form $v^T u(t)$ obtained by solving the following ODE system

$$\frac{d}{dt}u(t) = -2\sqrt{-1}\pi\tilde{H}(t)u(t), \quad u(0) = v, \quad \text{for } t \in [0,T].$$
(12)

This system of ODEs comes from Experiment 2 (Strong coupling) in [10], and $v^T u(t)$ represents an NMR experiment with a magic angle spinning (MAS) for k spins; see, e.g., [1]. The so-called Hamiltonian $\tilde{H}(t)$ is a $2^k \times 2^k$ matrix-valued function and has the form

$$H(t) = D + B(\cos(2\pi\nu t) + \cos(4\pi\nu t)),$$
(13)

with D, B sparse matrices described in [10]. In our experiments, we set $T = 10^{-3}$, $\nu = 10^4$, and k = 4, 7, 10, so obtaining three systems with exponentially increasing sizes.

The approximated solution $\hat{u}(t)$ (11) is computed by solving the linear system (10)* with M = 1000. The numerical experiments were performed using MatLab R2022a, and the linear systems were solved by the MatLab GMRES method implementation, gmres, with tolerance set to 1e - 15. In Figure 1, we compare the approximated bilinear form $v^T \hat{u}(t)$ with the solution obtained by the MatLab function ode45 with relative and absolute tolerance set to 3e - 14. Figure 2 reports the corresponding relative and absolute errors over the interval [0, T] (the reference for the error is again the ode45 solution). In all the experiments, GMRES stopped after a maximum of 27 iterations (for the cases k = 7, 10 due to residual stagnation). The numerical results show that the method is able to compute the solution with accuracy comparable with a well-established method.



Figure 1: Real and imaginary parts of $v^T u(t)$ approximations, with u(t) the solution of (12). The red circles represent approximation $v^T \hat{u}(t)$ from (11), while the blue line represents the ode45 approximation. From left to right, k = 4, 7, 10.

3 Matrix equation formulation

The matrix-valued function $\tilde{A}(t)$ in (1) can always be written in the form

$$\tilde{A}(t) = \sum_{k=1}^{d} A_k \tilde{f}_k(t), \tag{14}$$

^{*}The matrices $F_M^{(i,j)}$ in the block coefficient matrix (8) are numerically banded with bandwidth $b_{i,j}$. In order to avoid error accumulation, the last $b_{i,j}$ rows of each $F_M^{(i,j)}$ have been set equal to zero; see [2].



Figure 2: Absolute (blue circles) and relative (red crosses) errors of approximation $v^T \hat{u}(t)$, with $\hat{u}(t)$ from (11). From left to right, k = 4, 7, 10.

with $\tilde{f}_1, \ldots, \tilde{f}_d$ distinct scalar functions and A_1, \ldots, A_d constant matrices. In many applications, d is small. For instance, in the examples from Section 2.1, we have d = 2. Then, exploiting expression (14), the (block) coefficient matrix (8) of $A(t, s) = \tilde{A}(t)\Theta(t - s)$ becomes

$$\mathcal{A}_M = \sum_{k=1}^d A_k \otimes F_M^{(k)},$$

with $F_M^{(k)}$ the coefficient matrix (6) of $\tilde{f}_k(t)$. The solution x of the linear system (10) can, hence, be rewritten in terms of the solution X of the following matrix equation

$$X - \sum_{k=1}^{d} F_{M}^{(k)} X A_{k}^{T} = \phi_{M}(0) b^{T}, \quad x = vec(X),$$
(15)

where vec(X) denotes the vectorization of X, i.e., the vector obtained by stacking the columns of X into a single vector. The matrix equation (15) has a rank 1 right-hand side $\phi_M(0)b^T$. This suggests that the solution X may have a low numerical rank. Figure 3 reports the computed singular values of X, where x = vec(X) is the linear system solution of each of the experiments performed in Section 2.1. For k = 4, the solution X is full rank, while for k = 7, 10, the numerical rank of X is, respectively, 12, 72 (we consider as numerical rank the index of the last singular value before the stagnation visible in the plots). Clearly, this preliminary study shows that the numerical rank of X increases slowly with the size of X.



Figure 3: Singular values of the matrix X, with x = vec(X) the solution of (10) for the examples in Section 2.1. From left to right, k = 4, 7, 10.

4 Discussions and conclusion

In this work, we present a new method for solving systems of non-autonomous linear ODEs. The method is based on the solution of a linear system that can be rewritten as a matrix equation. Several examples illustrate that the method is able to compute the solution with accuracy comparable to the well-established Runge-Kutta method implemented by the MatLab function ode45. Moreover, the experiments show that the solution of the matrix equation is a numerical low-rank matrix when the ODE system is large enough. This may be exploited

using projection methods with low-rank techniques (see, e.g., [11, 12]). In [10], we also show that matrix A_M in (8) can be compressed by the Tensor Train decomposition (note that [10] uses a different family of orthogonal functions instead of the Legendre polynomials). A Tensor Train approach may further reduce the memory and computational cost of the method. Another possible approach could be extrapolation methods able to exploit the dependence of equation (10) on *s*; see, e.g., [13, 14].

Overall, the results suggest that the presented method may be an effective solver for large-to-huge systems of ODEs once we are able to exploit the solution's low-rank structure and the other mentioned properties. We are currently investigating these possible approaches.

References

- [1] S. Hafner and H. W. Spiess, Concepts Magn. Reson. 10, 99–128 (1998).
- [2] S. Pozza and N. Van Buggenhout, The *-product approach for linear odes: a numerical study of the scalar case, 2022.
- [3] L. Schwartz, Théorie Des Distributions (Hermann, Paris, 1978).
- [4] P.L. Giscard, K. Lui, S.J. Thwaite, and D. Jaksch, J. Math. Phys. 56(5), 053503 (2015).
- [5] P. L. Giscard and C. Bonhomme, Phys. Rev. Research 2(Apr), 023081 (2020).
- [6] P.L. Giscard and S. Pozza, Appl. Math. 65(6), 807–827 (2020).
- [7] P.L. Giscard and S. Pozza, Linear Algebra Appl. 624, 153–173 (2021).
- [8] P.L. Giscard and S. Pozza, Boll Unione Mat Ital (2022).
- [9] R. A. Silverman et al., Special functions and their applications (Courier Corporation, 1972).
- [10] S. Cipolla, S. Pozza, M. Redivo-Zaglia, and N. Van Buggenhout, Numer. Algorithms (2022).
- [11] V. Simoncini, SIAM J. Sci. Comput. 29(3), 1268–1288 (2007).
- [12] D. Palitta and P. Kürschner, Numer. Algorithms 88(3), 1383–1417 (2021).
- [13] C. Brezinski and M. Redivo Zaglia, Extrapolation methods, Studies in Computational Mathematics, Vol. 2 (North-Holland Publishing Co., Amsterdam, 1991), Theory and practice, With 1 IBM-PC floppy disk (5.25 inch).
- [14] D. Buoso, A. Karapiperi, and S. Pozza, Appl. Numer. Math. 90, 38–54 (2015).