Schrödinger cat states prepared by logical gate with non-Gaussian resource state: effect of finite squeezing and efficiency versus monotones

A. V. Baeva and I. V. Sokolov^{*}

Saint Petersburg State University, Universitetskaya nab. 7/9, 199034 Saint Petersburg, Russia

Quantum measurement-induced gate based on entanglement with ideal cubic phase state used as a non-Gaussian resource is able to produce Shrödinger cat state in the form of two high fidelity "copies" of the target state on phase plane [N.I. Masalaeva, I.V. Sokolov, Phys. Lett. A **424**, 127846 (2022)]. In this work we examine the effect of finite initial squeezing of the resource state on the gate performance. We present exact solution for the gate output state and demonstrate that there exists a degree of squeezing, available in experiment, such that the output cat state quality almost does not impove with the further increase of squeezing. On the other hand, the probability of the expected ancilla measurement outcome decreases with squeezing. Since an overall efficiency of the conditional scheme should account for the probability of success, we argue that such measures of non-Gaussianity of the resource state, as Wigner logarithmic negativiy and non-Gaussianity, may not be directly applicable to assess the efficiency of non-Gaussian gates based on quantum entanglement and subsequent projective measurement.

I. INTRODUCTION

The continuous-variable (CV) quantum information schemes based on Gaussian resource states were extensively explored both theoretically and experimentally [1-4], including their essentially multimode implementation [5-9]. While the continuous-variable Gaussian cluster schemes are able to perform Gaussian transformations of the input states, in order to achieve universal quantum computing there is a need to introduce the non-Gaussian logical gates [1, 2].

A minimal nonlinearity sufficient to prepare non-Gaussian resource states is cubic. The cubic phase state based on cubic nonlinearity was first considered in [10, 11], and some approaches to the implementation of such states, as well as of the cubic (or higher) phase gates, were explored both theoretically and experimentally [12-17].

Recently, we demonstrated [18, 19] that continuous-variable measurement-induced two-node logical gate is able to prepare Schrödinger cat-like quantum superpositions, if Gaussian (e. g. squeezed) resource state of an ancillary oscillator is substituted with non-Gaussian one, and standard homodyne measurement is applied. This was shown for the cubic phase state used as a resource.

A variety of methods to prepare CV Schrödinger cat states were discussed and implemented experimentally so far. This can be directly achieved by unitary evolution assisted by a strong enough non-linear interaction [20]. The schemes based on a hybrid measurement-induced evolution also can create cat-like states. The optical Schrödinger cat states were generated using photon subtraction in a low-photon regime [21] and from wide-band CV squeezed light [22], the homodyne detection with photon number state as a resource [23], and the iterative schemes which allow an incremental enlargement of cat states [24–26]. The Schrödinger cat states can be prepared by using a photonic even-parity detector and CV entanglement [27], by photon number measurement on two-mode [28] or multimode [29] Gaussian state.

A key feature of the gate [18, 19] is that the Schrödinger cat state emerges when the ancillary oscillator measurement is compatible not with one, but with two different values of its physical variables. As far as these values are imprinted by the entanglement into the target oscillator observables, the two-component cat-like quantum superposition is created at the output of the gate. This feature does not appear if Gaussian (squeezed) resource state is used, and needs the measurement which is consistent with this criterion.

Since the experimentally attainable degree of squeezing is limited, in this work we focus on the effect of finite initial squeezing of the resource state on the gate performance. The exact solution for the gate output state in terms of the Airy function is presented for vacuum input state. We demonstrate that there exists a degree of squeezing, available in experiment, such that the output superposition quality almost does not impove with the further increase of squeezing. This quality is estimated by means of the

^{*}Electronic address: i.sokolov@spbu.ru, sokolov.i.v@gmail.com

fidelity between the output state and the superposition of two coherent states well spaced on the phase plane, which is of interest for some error correction protocols.

On the other hand, we find that the probability of the expected ancilla measurement outcome decreases with squeezing. It is natural to admit that an overall efficiency of the conditional scheme should account for the probability of success. We argue that such measures of non-Gaussianity of the resource state, as Wigner logarithmic negativity and non-Gaussianity, may not be directly applicable to assess the efficiency of non-Gaussian gates, which are based on quantum entanglement and subsequent projective measurement.

II. CAT STATES FROM CV GATE WITH CUBIC RESOURCE STATE BASED ON FINITE INITIAL SQUEEZING

The non-Gaussian gate based on perfect cubic phase state was introduced in [18, 19]. The measurementinduced two-node gate uses the cubic phase state of the ancillary oscillator as an elementary non-Gaussian resource, the C_Z operation which entangles an input signal with the ancilla, and the projecting homodyne measurement. Under optimal conditions, the gate output state is close to "perfect" Schrödinger cat state, that is, to the superposition of two symmetrically displaced undistorted copies of the input state.

A key feature of the gate is that the ancilla measurement outcome provides multivalued information about the output state canonical variables, which results in the preparation of a cat-like state. This feature is easily interpreted [18, 19] in terms of a clear pictorial representation which is extendable to some other measurement-induced schemes based on the non-Gaussian resources.

Let us outline briefly the gate operation in a more realistic configuration, when the ancillary resource oscillator is subject to a finite initial squeezing before the cubic nonlinear perturbation is applied to the ancilla. In general, the target oscillator may be initially prepared in an arbitrary state

$$|\psi_1\rangle = \int dx_1 \psi(x_1) |x_1\rangle,$$

which is assumed to occupy a limited range $\{\Delta x_1, \Delta y_1\}$ of the coordinate and momentum.

As the initial state of the ancillary oscillator, prepared before the cubic evolution is applied, we consider the squeezed state whose uncertainty region on the phase plane is squeezed along the momentum axis by the factor $s \leq 1$,

$$\psi^{(sq)}(x_2) = \frac{\sqrt{s}}{\pi^{1/4}} e^{-(sx_2)^2/2}.$$
(1)

In order to prepare the non-Gaussian cubic resource state of ancilla, one applies the unitary evolution operator $\exp(i\gamma q_2^3)$ to the state (1). In the following, we use the notation $\{q, p\}$ for the coordinate and momentum operators.

Next, the C_Z entangling unitary evolution operator $\exp(iq_1q_2)$ is applied, which prepares the state

$$|\psi_{12}\rangle = \frac{\sqrt{s}}{\pi^{1/4}} \int dx_1 dx_2 \psi(x_1) e^{ix_2(x_1 + \gamma x_2^2)} e^{-(sx_2)^2/2} |x_1\rangle |x_2\rangle, \tag{2}$$

and the ancillary oscillator momentum is measured with the outcome y_m .

Projecting the state (2) on the homodyne detector eigenstate $|y_m\rangle_{p_2}$, one arrives at the target oscillator output state wave function (unnormalized),

$$\tilde{\psi}^{(out)}(x) = \psi_1(x)\tilde{\varphi}(x - y_m),\tag{3}$$

where the target oscillator input state wave function is multiplied by the factor which is expressed [30] in terms of the Airy function,

$$\tilde{\varphi}(x-y_m) = \frac{\sqrt{s}}{\pi^{3/4}\sqrt{2}} \int dx' \ e^{ix'(x-y_m+\gamma x'^2)} e^{-(sx')^2/2} = \frac{\sqrt{2s}\pi^{1/4}}{(3\gamma)^{1/3}} \exp\left[\frac{s^2}{6\gamma}\left(x-y_m+\frac{s^4}{18\gamma}\right)\right] \operatorname{Ai}\left[\frac{1}{(3\gamma)^{1/3}}\left(x-y_m+\frac{s^4}{12\gamma}\right)\right].$$
(4)

The probability density of the ancilla momentum measurement outcome y_m is

$$P(y_m) = \langle \tilde{\psi}^{(out)} | \tilde{\psi}^{(out)} \rangle = \int dx \left| \tilde{\psi}^{(out)}(x) \right|^2.$$
(5)

The state conditionally prepared by the gate finally is given by

$$\psi^{(out)}(x) = \frac{1}{\sqrt{P(y_m)}} \tilde{\psi}^{(out)}(x).$$
(6)

Previously [19], we have explored in detail the performance of the gate based on the ideal initial resource state (that is, in the limit where instead of the squeezed state (1) one uses the momentum eigenstate $|0\rangle_{p_2}$) for a representative set of Fock states of the target oscillator. Here we focus on the effect of finite initial squeezing. To be specific, we assume the target oscillator to be in the vacuum state which is given by (1), where $x_2 \rightarrow x_1$, s = 1.

As seen from the exact result (4), in the limit of perfect squeezing, $s \to 0$, the added factor vanishes, and $P(y_m) \to 0$. In this limit, the probability density to observe any given value y_m of the ancilla momentum also vanishes. Therefore, in our previous work we were able to explore in detail quantum statistics of the emerging Schrödinger cat-like states in dependence on the cubic interaction parameter γ and on the measurement outcome y_m , but could not discuss the probability density and propose an optimal choice of the initial ancilla squeezing.

Let us first consider how a finite initial squeezing affects the quality of the emerging superpositions.

In [19] we compared the exact solution obtained under the assumption of perfect squeezing, and the state closest to it, which corresponds to the Schrödinger cat state in the form of a superposition of two undistorted copies of the vacuum state of the target oscillator (i. e., the Glauber states), symmetrically spaced on the phase plane along the momentum quadrature,

$$|\psi_{cat}^{(out)}\rangle = \frac{\left(e^{i\theta}|\alpha\rangle + e^{-i\theta}| - \alpha\rangle\right)}{\sqrt{2\left(1 + \cos(2\theta)e^{-2|\alpha|^2}\right)}},\tag{7}$$

where

$$\alpha = ip^{(+)}, \quad p^{(+)} = \sqrt{y_m/3\gamma}, \quad \theta = \frac{\pi}{4} - \frac{2}{3\sqrt{3\gamma}}y_m^{3/2}.$$
 (8)

We have shown that in the limit of perfect squeezing and within a relevant range of the parameters $\{\gamma, y_m\}$, the proposed gate prepares with high fidelity the superpositions (7) of Glauber states. Since the experimentally available squeezing is limited, it remained unclear to which degree a finite squeezing can affect the possibility to obtain the Schrödinger cat states close to (7).



FIG. 1: Infidelity $1 - F_{cat}$ for the ancilla momentum measurement outcome $y_m = 3, 3.6, 4.5, 6, 9, 15$ respectively in dependence on the initial ancilla state squeezing, where $1/s \ge 1$ is the stretching factor of the ancilla coordinate quadrature. In these plots, the cubic deformation coefficient is chosen as $\gamma = y_m/30$, which provides the cat states with the same spacing $2p^{(+)} = 3.16$ (see (8)) between the copies along the momentum axis.

In Fig. 1 we represent the effect of initial squeezing on the infidelity $1 - F_{cat}$ between the gate output state (6) and the cat state (7), where

$$F_{cat} = \left| \int dx \psi^{(out)*}(x) \psi^{(out)}_{cat}(x) \right|^2.$$
(9)

Fig. 1 shows that starting from some threshold squeezing, which is achievable in experiment, a further increase in squeezing does not lead to a significant increase in fidelity. The fidelity difference from unity

in this limit is due to local deformations of the input state copies on the phase plane, as seen from the Figs. 2 and 3. This deformation is the smaller, the larger the parameter γ of cubic nonlinearity.



FIG. 2: Output state Wigner function for the cubic deformation coefficient $\gamma = 0.1$ and the ancilla momentum measurement outcome $y_m = 3$ (low fidelity configuration). The ancilla initial squeezing is 5 dB (a), 9 dB (b), and 14 dB (c), which corresponds to 1/s = 1.78, 2.82, and 5.01, correspondingly.



FIG. 3: Same as in Fig. 2 for the cubic deformation coefficient $\gamma = 0.5$ and the ancilla momentum measurement outcome $y_m = 15$ (high fidelity configuration).



FIG. 4: Probability density $P(y_m)$ of the measurement outcome $y_m = 3, 6, 9, 12, 15$ respectively in dependence on the initial ancilla state squeezing, where $1/s \ge 1$ is the stretching factor of the ancilla coordinate quadrature. For any measurement outcome in these plots, the cubic deformation coefficient is assumed to be $\gamma = y_m/30$, which provides the spacing $2p^{(+)} = 3.16$ between the centers of copies along the momentum axis.

However, as seen from Fig. 4, the degree of initial squeezing significantly affects the probability to obtain the expected measurement result y_m . This is due to the fact that the probability density per unit momentum interval of the ancillary oscillator, initially prepared in cubic state, essentially depends on squeezing, as illustrated in Fig. 5.



FIG. 5: A visual representation of the ancillary oscillator support region on the phase plane before and after the cubic deformation with $\gamma = 0.1$ for the initial squeezing of 5 and 14 dB respectively. Two distinct areas on the phase plane of the ancilla observables compatible with the same momentum measurement outcome $y_m = 3$ are marked by the asterisks. Due to entanglement, both values of the ancilla coordinate are transferred to the target oscillator observables, thus generating the two-component cat state.

For a qualitative interpretation of this feature of the gate under consideration, we introduce in Fig. 5 the support regions of cubic phase state for 5 and 14 dB squeezing, built in the semiclassical approximation, i. e. when the evolution of points on the phase plane caused by the cubic Hamiltonian $-\gamma q_2^3$, is described via a semiclassical mapping of the form $x_2 \to x_2$, $y_2 \to y_2 + 3\gamma q_2^2$.

At low squeezing, the resource quasi-cubic state has a low probability density near $p_2 = y_m$, as shown in Fig. 5 for 5 dB squeezing. For an arbitrarily large squeezing, the support region of the resource state is stretched infinitely into an arbitrarily "thin" parabola, and the probability of the expected momentum measurement result vanishes.

As is well known, a correct quantum description of the statistical distribution, which corresponds to a given quantum state, is represented by its Wigner function. In Fig. 6 we show the Wigner function of the cubic phase state built for the same values of the parameters of cubic interaction and squeezing, as in the figures 2, 3, and 5.



FIG. 6: Wigner function of the non-ideal cubic phase state for the cubic deformation coefficient $\gamma = 0.1$, generated from the initial ancilla state with squeezing of 5 dB (a), 9 dB (b) and 14 dB (c).

One can note a good agreement between the results that follow from the exact form (6) of the gate output state, and the qualitative picture introduced above. This means that despite a more complex behavior of the cubic phase state Wigner function in comparison with its semiclassical interpretation, some features present in the behavior of the Wigner function (e. g., oscillations) may average out and not manifest themselves in the output state.

Hence, in turn, we can conclude that our approach [18, 19] to qualitative analysis of the gates that use non-Gaussian resource states and measurement can be useful for the search and evaluation of other logical devices.

III. EFFICIENCY OF THE GATE VERSUS MONOTONES

Recently, the problem of universal measures of non-Gaussianity of the states that can serve as a resource for universal quantum computing has been widely discussed [31]. One could expect that a non-Gaussian resource state, for which a well-chosen measure is larger, would be able to take more efficiently the result of quantum evolution out of the class of Gaussian schemes when used optimally. Among the measures discussed so far in this context are the non-Gaussianity and Wigner logarithmic negativity. For finite squeezing, the non-Gaussianity and Wigner logarithmic negativity of the cubic phase state, which our gate uses as a non-Gaussian resource, are [32] monotonous functions of the parameter γ/s^3 (in the notation of our work, $1/s \ge 1$ is the stretching factor of the coordinate quadrature). This parameter increases with the degree of squeezing.

Our results demonstrate that for large enough parameter γ of the cubic non-linearity, our gate can conditionally produce the output state which is very close to the cat-like superposition (7) of the Glauber states almost independently on the degree of squeezing (after some threshold value, see Fig. 1). If the measured ancilla momentum y_m fits to the range specified before the measurement, the "amount of non-Gaussianity" in the output cat-like state becomes, for large enough squeezing, almost independent on the squeezing present in the parameter γ/s^3 , no matter which measure is used.

However, from the point of view of a real use of the measurement-assisted gates, the concept of efficiency should include not only the quality of the prepared states, but also the probability of obtaining the necessary measurement result. In our scheme, the "amount of non-Gaussianity" present in the output state, weighted by its probability shown in Fig. 4, vanishes in the limit of perfect squeezing.

That is, the use of cubic phase state with the larger value of the parameter γ/s^3 and, hence, with the larger non-Gaussianity and Wigner logarithmic negativity, may render the gate less efficient in the sense mentioned above.

One can conclude from this example that the non-Gaussianity measures based on global properties of the resource states may be generally not suitable for the schemes based on the entanglement and projective measurements. Physically speaking, the efficiency of such schemes may depend more on the resource state behavior in the region of phase space, where the resource state overlaps with the state detected by the measuring device, than on its global properties.

Conclusion

We have investigated quantum statistical properties of two-component Schrödinger cats created at the output of the conditional non-Gaussian logical gate that we have proposed previously. The gate is based on quantum entanglement of the target oscillator with the ancillary one, and the subsequent projective measurement of ancilla. A key feature of the regime where cat-like states arise is that the measurement provides multivalued information about the target system physical variables.

As a non-Gaussian resource, we considered here cubic phase state generated from the initial squeezed state of ancilla with an arbitrary finite initial squeezing.

We showed that in the relevant range of parameters, an increase in the degree of initial squeezing above some threshold range achievable in experiment has negligible effect on the fidelity between the gate output state and the Schrödinger cat state in the form of superposition of two coherent Glauber states. On the other hand, we demonstrated that the degree of initial squeezing essentially affects the probability of the projective measurement outcome that leads to the desired output state. This probability vanishes for both too small and too large degree of squeezing, and reaches its maximum value at a certain degree of squeezing, which depends on the degree of cubic non-linearity used in the preparation of the resource state and on the output state parameters. We provided a simple physical interpretation of these results using both the exact and semiclassical representation of the resource state properties and of the measurement procedure on the phase plane.

Based on our results, we also draw attention to the fact that such measures of non-Gaussianity of the resource state, as Wigner logarithmic negativity and non-Gaussianity, may be not applicable to assess the overall efficiency of non-Gaussian gates which use quantum entanglement and subsequent projective measurement.

S. Lloyd, S. L. Braunstein, Quantum computation with continuous variables, Phys. Rev. Lett. 82 (1999) 1784.

 ^[2] S.L. Braunstein, P. van Loock, Quantum information with continuous variables, Rev. Mod. Phys. 77 (2005) 513.

^[3] A. Furusawa, J.L. Sørensen, S.L. Braunstein, C. A. Fuchs, H. J. Kimble, E. S. Polzik, Unconditional quantum teleportation, Science 282 (1998) 706.

^[4] X. Li, Q. Pan, J. Jing, J. Zhang, C. Xie, K. Peng, Quantum dense coding exploiting a bright Einstein-Podolsky-Rosen beam, Phys. Rev. Lett. 88 (2002) 047904.

- [5] M. Gu, C. Weedbrook, N.C. Menicucci, T. C. Ralph, P. van Loock, Quantum computing with continuousvariable clusters, Phys. Rev. A 79 (2009) 062318.
- [6] C. Weedbrook, S. Pirandola, R. Garcia-Patron, N. J. Cerf, T. C. Ralph, J. H. Shapiro, S. Lloyd, Gaussian quantum information, Rev. Mod. Phys. 84 (2012) 621.
- [7] R. Ukai, Multi-Step Multi-Input One-Way Quantum Information Processing with Spatial and Temporal Modes of Light, Springer, Tokyo, 2015.
- [8] S. Yokoyama, R. Ukai, S.C. Armstrong, C. Sornphiphatphong, T. Kaji, S. Suzuki, J. Yoshikawa, H. Yonezawa, N. C. Menicucci, A. Furusawa, Ultra-large-scale continuous-variable cluster states multiplexed in the time domain, Nat. Photonics 7 (2013) 982.
- [9] J. Roslund, R. Medeiros de Araújo, S. Jiang, C. Fabre, N. Treps, Wavelength-multiplexed quantum networks with ultrafast frequency combs, Nat. Photonics 8 (2014) 109.
- [10] D. Gottesman, A. Kitaev, J. Preskill, Encoding a qubit in oscillator, Phys. Rev. A 64 (2001) 012310.
- [11] S.D. Bartlett, B.C. Sanders, Universal continuous-variable quantum computation: Requirement of optical nonlinearity for photon counting, Phys. Rev. A 65 (2002) 042304.
- [12] S. Ghose, B. C. Sanders, Non-Gaussian ancilla states for continuous variable quantum computation via Gaussian maps, J. Mod. Opt. 54 (2007) 855.
- [13] P. Marek, R. Filip, A. Furusawa, Deterministic implementation of weak quantum cubic nonlinearity, Phys. Rev. A 84 (2011) 053802.
- [14] M. Yukawa, K. Miyata, H. Yonezawa, P. Marek, R. Filip, A. Furusawa, Emulating quantum cubic nonlinearity, Phys. Rev. A 88 (2013) 053816.
- [15] K. Marshall, R. Pooser, G. Siopsis, C. Weedbrook, Repeat-until-success cubic phase gate for universal continuous-variable quantum computation, Phys. Rev. A 91 (2015) 032321.
- [16] K. Miyata, H. Ogawa, P. Marek, R. Filip, H. Yonezawa, J.-I. Yoshikawa, A. Furusawa, Implementation of a quantum cubic gate by adaptive non-Gaussian measurement, Phys. Rev. A 93 (2016) 022301.
- [17] P. Marek, R. Filip, H. Ogawa, A. Sakaguchi, S. Takeda, J.-I. Yoshikawa, A. Furusawa, General implementation of arbitrary nonlinear quadrature phase gates, Phys. Rev. A 97 (2018) 022329.
- [18] I. V. Sokolov, Schrödinger cat states in continuous variable non-Gaussian networks, Phys. Lett. A 384 (2020) 126762.
- [19] N. I. Masalaeva, I. V. Sokolov, Quantum statistics of Schrodinger cat states prepared by logical gate with non-Gaussian resource state, Phys. Lett. A 424 (2022) 127846.
- [20] B. Yurke, D. Stoler, Generating quantum mechanical superpositions of macroscopically distinguishable states via amplitude dispersion, Phys. Rev. Lett. 57 (1986) 13.
- [21] A. Ourjoumtsev, R. Tualle-Brouri, J. Laurat, P. Grangier, Generating optical Schrödinger kittens for quantum information processing, Science 312, (2006) 83.
- [22] K. Takase, A. Kawasaki, B. K. Jeong, M. Endo, T. Kashiwazaki, T. Kazama, K. Embutsu, K. Watanabe, T. Umeki, S. Miki, H. Terai, M. Yabuno, F. China, W. Asavanant, J. Yoshikawa, A. Furusawa, Generation of Schrödinger cat states with Wigner negativity using a continuous-wave low-loss waveguide optical parametric amplifier, Opt. Expr. **30**(9) (2022) 14161.
- [23] A. Ourjoumtsev, H. Jeong, R. Tualle-Brouri, P. Grangier, Generation of optical "Schrödinger cats" from photon number states, Nature 448 (2007) 784.
- [24] J. Etesse, R. Blandino, B. Kanseri, R. Tualle-Brouri, Proposal for a loophole-free violation of Bell's inequalities with a set of single photons and homodyne measurements, New J. Phys. 16 (2014) 053001.
- [25] J. Etesse, M. Bouillard, B. Kanseri, R. Tualle-Brouri, Experimental generation of squeezed cat states with an operation allowing iterative growth, Phys. Rev. Lett. 114 (2015) 193602.
- [26] D.V. Sychev, A.E. Ulanov, A.A. Pushkina, M. W. Richards, I. A. Fedorov, A. I. Lvovsky, Enlargement of optical Schrödinger cat states, Nat. Photonics 11 (2017) 379.
- [27] G.S. Thekaddath, B.A. Bell, I.A.Walmsley, A. I. Lvovsky, Engineering Schrodinger cat states with a photonic even-parity detector, Quantum 4 (2020) 239.
- [28] K. Takase, J. Yoshikawa, W. Asanavant, M. Endo, A. Furusawa, Generation of optical Schrödinger cat states by generalized photon subtraction, Phys. Rev. A 103 (2021) 013710.
- [29] D. Su, C.R. Myers, K. K. Sabapathy, Conversion of Gaussian states to non-Gaussian states using photonnumber-resolving detectors, Phys. Rev. A 100 (2019) 052301.
- [30] O. Vallée, M. Soares, Airy functions and applications to physics, World Scientific, 2010.
- [31] M. Walschaers, Non-Gaussian quantum states and where to find them, PRX Quantum 2 (2021) 030204.
- [32] F. Albarelli, M. G. Genoni, M. G. A. Paris, A. Ferraro, Resource theory of quantum non-Gaussianity and Wigner negativity, Phys. Rev. A 98 (2018) 052350.