LAGUERRE-FREUD EQUATIONS FOR THREE FAMILIES OF HYPERGEOMETRICAL DISCRETE ORTHOGONAL POLYNOMIALS

ITSASO FERNÁNDEZ-IRISARRI¹ AND MANUEL MAÑAS²

ABSTRACT. The Cholesky factorization of the moment matrix is considered for discrete orthogonal polynomials of hypergeometrical type. We derive the Laguerre–Freud equations when the first moments of the weights are given by the $_1F_2$, $_2F_2$ and $_3F_2$ generalized hypergeometrical functions.

1. INTRODUCTION

Discrete orthogonal polynomials is an important and active area of the theory of orthogonal polynomials see [44, 17] as well as [32, 33, 18, 47]. When a discrete Pearson equation is fulfilled by the weight we are dealing with semiclassical discrete orthogonal polynomials, see [24, 25, 22, 23]. For some specific type of weights of generalized Charlier and Meixner types, the corresponding Freud–Laguerre type equations for the coefficients of the three term recurrence has been studied, see for example [20, 27, 28, 29, 46].

This paper is the final one in our series of works on the of the Gauss-Borel factorization of the corresponding moment matrix, see [39], in standard discrete orthogonality. In [42] the Cholesky factorization of the moment matrix was used to study discrete orthogonal polynomials on the homogeneous lattice. For weights subject to a discrete Pearson equation, so that the moments where logarithmic derivatives of generalized hypergeometric functions, a banded semi-infinite matrix Ψ , the so called Laguerre–Freud structure matrix, that models the shifts by ± 1 in the independent variable of the sequence of orthogonal polynomials was given. Contiguous relations for the generalized hypergeometric functions translate into symmetries for the corresponding moment matrix, and the 3D Nijhoff–Capel discrete Toda lattice [43, 31] describes the corresponding contiguous shifts for the squared norms of the orthogonal polynomials. In [40] we gave an interpretation for the contiguous transformations for the generalized hypergeometric functions in terms of simple Christoffel and Geronimus transformations. Using the Geronimus–Uvarov perturbations we got determinantal expressions for the shifted orthogonal polynomials. Then, in [26] we discussed the generalized Charlier, Meixner and type I Hahn discrete orthogonal polynomials (according to the nomenclature in [24]), analyzed the Laguerre–Freud structure matrix Ψ , and derive from its banded structure and its compatibility with the Toda equation and the Jacobi matrix, a number of nonlinear equations for the coefficients $\{\beta_n, \gamma_n\}$ of the three term recursion relations $zP_n(z) = P_{n+1}(z) + \beta_n P_n(z) + \gamma_n P_{n-1}(z)$ satisfied by the orthogonal polynomial sequence. These Laguerre–Freud equations are of the form $\gamma_{n+1} = \mathscr{G}(n, \gamma_n, \gamma_{n-1}, \dots, \beta_n, \beta_{n-1} \dots)$ and $\beta_{n+1} = \mathscr{B}(n, \gamma_{n+1}, \gamma_n, \dots, \beta_n, \beta_{n-1}, \dots)$, for some functions \mathscr{B}, \mathscr{G} . Magnus [35, 36, 37, 38] named

¹Departamento de Física Teórica, Universidad Complutense de Madrid, Plaza Ciencias 1, 28040-Madrid, Spain

²Departamento de Física Teórica, Universidad Complutense de Madrid, Plaza Ciencias 1, 28040-Madrid, Spain & Instituto de Ciencias Matematicas (ICMAT), Campus de Cantoblanco UAM, 28049-Madrid, Spain

E-mail addresses: ¹itsasofe@ucm.es, ²manuel.manas@ucm.es.

²⁰²⁰ Mathematics Subject Classification. 42C05,33C45,33C47.

Key words and phrases. Hypergeometrical discrete orthogonal polynomials, Laguerre–Freud equations, generalized hypergeometrical functions, Pearson equations, semiclassical discrete orthogonal polynomials, Jacobi matrix, recursion relations, banded matrices.

The authors thanks financial support from the Spanish "Agencia Estatal de Investigación" research project [PGC2018-096504-B-C33], Ortogonalidad y Aproximación: Teoría y Aplicaciones en Física Matemática and research project [PID2021- 122154NB-I00], Ortogonalidad y aproximación con aplicaciones en machine learning y teoría de la probabilidad.

these type of relations, attending to [34, 30], as Laguerre–Freud relations. There are several papers discussing Laguerre–Freud relations for the generalized Charlier, generalized Meixner and type I generalized Hahn cases, see [46, 20, 27, 28, 29, 22].

In this paper we conclude our studies on these themes, and extend these methods to three families of discrete orthogonal polynomials with first moments given by the generalized hypergeometric functions ${}_{1}F_{2}$, ${}_{2}F_{2}$ and ${}_{3}F_{2}$. Notice that generalized Charlier, Meixner and type Hahn correspond to ${}_{0}F_{1}$, ${}_{1}F_{1}$ and ${}_{2}F_{1}$, respectively. For the three families we explicitly compute the Laguerre-Freud structure matrix which happens to be a pentadiagonal, hexadiagonal or heptadiagonal banded matrices, respectively. We also compute the Laguerre-Freud equations and give explicit expressions for the leading nontrivial coefficients of the orthogonal polynomials in terms of the recursion coefficients.

The layout of the paper is as follows. We first complete this introduction by discussing, without proof but referring to the appropriate sources, some preliminary material. Then, we have three sections devoted each of them to the corresponding hypergeometrical function, $_1F_2$, $_2F_2$ and $_3F_2$, finding in each case the Laguerre–Freud structure matrix and corresponding Laguerre–Freud equations.

1.1. Pearson equations and discrete orthogonal polynomials. Let us recall some relevant facts of the theory of orthogonal polynomials. Given a linear functional $\rho_z \in \mathbb{C}^*[z]$, the corresponding moment matrix is

$$G = (G_{n,m}),$$
 $G_{n,m} = \rho_{n+m},$ $\rho_n = \langle \rho_z, z^n \rangle,$ $n, m \in \mathbb{N}_0 := \{0, 1, 2, \dots\},$

with ρ_n being the *n*-th moment of the linear functional ρ_z . Let us assume that the moment matrix is such that all its truncations, which are Hankel matrices, $G_{i+1,j} = G_{i,j+1}$,



are nonsingular; i.e. the Hankel determinants $\Delta_k := \det G^{[k]}$ do not cancel, $\Delta_k \neq 0$, $k \in \mathbb{N}_0$. Then, there are monic polynomials

(1)
$$P_n(z) = z^n + p_n^1 z^{n-1} + \dots + p_n^n, \qquad n \in \mathbb{N}_0,$$

satisfying the following orthogonality relations $\langle \rho, P_n(z)z^k \rangle = 0$, for $k \in \{0, \ldots, n-1\}$ and $\langle \rho, P_n(z)z^n \rangle = H_n \neq 0$, and $\{P_n(z)\}_{n \in \mathbb{N}_0}$ is a sequence of orthogonal polynomials, i.e., $\langle \rho, P_n(z)P_m(z) \rangle = \delta_{n,m}H_n$ for $n, m \in \mathbb{N}_0$. The symmetric bilinear form $\langle F, G \rangle_{\rho} := \langle \rho, FG \rangle$, is such that the moment matrix is the Gram matrix of this bilinear form and $\langle P_n, P_m \rangle_{\rho} := \delta_{n,m}H_n$.

Introducing he monomial sequence $\chi(z) := \begin{pmatrix} 1 & z & z^2 & \cdots \end{pmatrix}^\top$ the moment matrix is $G = \langle \rho, \chi \chi^\top \rangle$, and χ is an eigenvector of the *shift matrix*, $\Lambda \chi = x\chi$, where

$$\Lambda := \begin{bmatrix} 0 & 1 & 0 & \cdots & \cdots \\ & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \end{bmatrix}.$$

Hence $\Lambda G = G\Lambda^{\top}$, this equation is equivalent to the fact that the moment matrix is a Hankel. As the moment matrix is symmetric its Borel–Gauss factorization is a Cholesky factorization, i.e.

(2)
$$G = S^{-1}HS^{-+},$$

where S is a lower unitriangular matrix that can be written as



and $H = \text{diag}(H_0, H_1, \dots)$ is a diagonal matrix, with $H_k \neq 0$, for $k \in \mathbb{N}_0$. The Cholesky factorization does hold whenever the principal minors of the moment matrix; i.e., the Hankel determinants Δ_k , do not cancel.

The entries $P_n(z)$ of the polynomial sequence

$$P(z) := S\chi(z),$$

are the monic orthogonal polynomials of the functional ρ .

We have the determinantal expressions $H_k = \frac{\Delta_{k+1}}{\Delta_k}$ and $p_k^1 = -\frac{\Delta_k}{\Delta_k}$, with



The lower Hessenberg semi-infinite matrix

(4)

has the polynomial sequence P(z) as eigenvector with eigenvalue z. The Hankel condition $\Lambda G = G \Lambda^{\top}$ and the Cholesky factorization gives

 $J = S\Lambda S^{-1}$

As the Hessenberg matrix JH is symmetric the Jacobi matrix J is tridiagonal, i.e.

$$J = \begin{bmatrix} \beta_0 & 1 & 0 \\ & \ddots & \\ \gamma_1 & \beta_1 & 1 & \ddots \\ 0 & \gamma_2 & \beta_2 & 1 & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \end{bmatrix}$$

and we find the three term recursion relation $zP_n(z) = P_{n+1}(z) + \beta_n P_n(z) + \gamma_n P_{n-1}(z)$, that with the initial conditions $P_{-1} = 0$ and $P_0 = 1$ completely determines the sequence of orthogonal polynomials $\{P_n(z)\}_{n \in \mathbb{N}_0}$ in terms of the recursion coefficients β_n, γ_n . The recursion coefficients are given by

(6)
$$\beta_n = p_n^1 - p_{n+1}^1 = -\frac{\Delta_n}{\Delta_n} + \frac{\Delta_{n+1}}{\Delta_{n+1}}, \qquad \gamma_{n+1} = \frac{H_{n+1}}{H_n} = \frac{\Delta_{n+1}\Delta_{n-1}}{\Delta_n^2}, \qquad n \in \mathbb{N}_0.$$

We introduce the following diagonal matrices $\gamma := \operatorname{diag}(\gamma_1, \gamma_2, \dots)$ and $\beta := \operatorname{diag}(\beta_0, \beta_1, \dots)$. In general, given any semi-infinite matrix A, we will write $A = A_{-} + A_{+}$, where A_{-} is a strictly lower triangular matrix and A_+ an upper triangular matrix. Moreover, A_0 will denote the diagonal part of A.

The lower Pascal matrix is

(7)
$$B = (B_{n,m}),$$
 $B_{n,m} := \begin{cases} \binom{n}{m}, & n \ge m, \\ 0, & n < m. \end{cases}$

so that

(8)
$$\chi(z+1) = B\chi(z).$$

The inverse of this Pascal matrix is

$$B^{-1} = [\tilde{B}_{n,m}], \qquad \qquad \tilde{B}_{n,m} := \begin{cases} (-1)^{n+m} \binom{n}{m}, & n \ge m, \\ 0, & n < m. \end{cases}$$

and $\chi(z-1) = B^{-1}\chi(z)$. The *dressed Pascal matrices* are $\Pi := SBS^{-1}$ and $\Pi^{-1} := SB^{-1}S^{-1}$ and give he connection between shifted and non shifted polynomials, i.e.

(9)
$$P(z+1) = \Pi P(z),$$
 $P(z-1) = \Pi^{-1} P(z).$

The lower Pascal matrix can be expressed as follows

$$B^{\pm 1} = I \pm \Lambda^{\top} D + (\Lambda^{\top})^2 D^{[2]} \pm (\Lambda^{\top})^3 D^{[3]} + \cdots,$$

where D = diag(1, 2, 3, ...) and $D^{[k]} := \frac{1}{k} \text{diag}(k^{(k)}, (k+1)^{(k)}, (k+2)^{(k)} \cdots)$, in terms of the falling factorials $x^{(k)} := x(x-1)(x-2) \cdots (x-k+1)$. That is,

$$D_n^{[k]} = \frac{(n+k)\cdots(n+1)}{k}, \qquad k \in \mathbb{N}, \qquad n \in \mathbb{N}_0.$$

The lower unitriangular factor can be written as $S = I + \Lambda^{\top} S^{[1]} + (\Lambda^{\top})^2 S^{[2]} + \cdots$ with $S^{[k]} = \text{diag} (S_0^{[k]}, S_1^{[k]}, \dots)$. We will use the *shift operators* T_{\pm} acting over the diagonal matrices as follows

$$T_{-} \operatorname{diag}(a_0, a_1, \dots) := \operatorname{diag}(a_1, a_2, \dots), \qquad T_{+} \operatorname{diag}(a_0, a_1, \dots) := \operatorname{diag}(0, a_0, a_1, \dots)$$

For any diagonal matrix $A = diag(A_0, A_1, ...)$ we have

(10)
$$\Lambda A = (T_{-}A)\Lambda, \qquad A\Lambda = \Lambda(T_{+}A), \qquad A\Lambda^{\top} = \Lambda^{\top}(T_{-}A), \qquad \Lambda^{\top}A = (T_{+}A)\Lambda^{\top}$$

In terms of these shift operators we find

$$2D^{[2]} = (T_{-}D)D, \qquad 3D^{[3]} = (T_{-}^{2}D)(T_{-}D)D = 2(T_{-}D^{[2]})D = 2D^{[2]}(T_{-}^{2}D).$$

For the inverse matrix

$$S^{-1} = I + \Lambda^{\top} S^{[-1]} + (\Lambda^{\top})^2 S^{[-2]} + \cdots,$$

we have

$$\begin{split} S^{[-1]} &= -S^{[1]}, \\ S^{[-2]} &= -S^{[2]} + (T_{-}S^{[1]})S^{[1]}, \\ S^{[-3]} &= -S^{[3]} + (T_{-}S^{[2]})S^{[1]} + (T_{-}^{2}S^{[1]})S^{[2]} - (T_{-}^{2}S^{[1]})(T_{-}S^{[1]})S^{[1]}. \end{split}$$

Corresponding expansions for the dressed Pascal matrices are

$$\Pi^{\pm 1} = I + \Lambda^{\top} \pi^{[\pm 1]} + (\Lambda^{\top})^2 \pi^{[\pm 2]} + \cdots$$

with $\pi^{[\pm n]} = \text{diag}(\pi_0^{[\pm n]}, \pi_1^{[\pm n]}, \dots)$. We have

(11)
$$\pi_n^{[\pm 1]} = \pm (n+1), \quad \pi_n^{[\pm 2]} = \frac{(n+2)(n+1)}{2} \pm p_{n+2}^1(n+1) \mp (n+2)p_{n+1}^1 = \frac{(n+2)(n+1)}{2} \mp (n+1)\beta_{n+1} \mp p_{n+1}^1.$$

(12)
$$\pi_n^{[\pm3]} = \pm \frac{(n+3)(n+2)(n+1)}{3} + \frac{(n+2)(n+1)}{2} p_{n+3}^1 - \frac{(n+3)(n+2)}{2} p_{n+1}^1 \\ \pm (n+1)p_{n+3}^2 \mp (n+3)p_{n+2}^2 \pm (n+3)p_{n+2}^1 p_{n+1}^1 \mp (n+2)p_{n+3}^1 p_{n+1}^1$$

Moreover, the following relations are fulfilled

(13)
$$\pi^{[1]} + \pi^{[-1]} = 0, \quad \pi^{[2]} + \pi^{[-2]} = 2D^{[2]}, \quad \pi^{[3]} + \pi^{[-3]} = 2((T_-^2 S^{[1]})D^{[2]} - (T_- D^{[2]})S^{[1]}).$$

We now recall some facts regarding discrete orthogonality. Hence, we focus on measures with support on the homogeneous lattice \mathbb{N}_0 , i.e. $\rho = \sum_{k=0}^{\infty} \delta(z-k)w(k)$, with moments given by

(14)
$$\rho_n = \sum_{k=0}^{\infty} k^n w(k),$$

and, in particular, with 0-th moment given by

(15)
$$\rho_0 = \sum_{k=0}^{\infty} w(k).$$

The weights we consider in this paper satisfy the following discrete Pearson equation

(16)
$$\nabla(\sigma w) = \tau w,$$

that is $\sigma(k)w(k) - \sigma(k-1)w(k-1) = \tau(k)w(k)$, for $k \in \mathbb{N}$, with $\sigma(z), \tau(z) \in \mathbb{R}[z]$. If we write $\theta := \tau - \sigma$, the previous Pearson equation reads

(17)
$$\theta(k+1)w(k+1) = \sigma(k)w(k), \qquad k \in \mathbb{N}_0.$$

If $N + 1 := \deg \theta(z)$ and $M := \deg \sigma(z)$, and the zeros of these polynomials are $\{-b_i + 1\}_{i=1}^N$ and $\{-a_i\}_{i=1}^M$ we write $\theta(z) = z(z+b_1-1)\cdots(z+b_N-1)$ and $\sigma(z) = \eta(z+a_1)\cdots(z+a_M)$. According to (15) the 0-th moment

$$\rho_0 = \sum_{k=0}^{\infty} w(k) = \sum_{k=0}^{\infty} \frac{(a_1)_k \cdots (a_M)_k}{(b_1+1)_k \cdots (b_N+1)_k} \frac{\eta^k}{k!} = {}_M F_N(a_1, \dots, a_M; b_1, \dots, b_N; \eta) = {}_M F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix},$$

is the generalized hypergeometric function, where we are using the two standard notations, see [16, 45]. Then, according to (14), for $n \in \mathbb{N}$, the corresponding higher moments $\rho_n = \sum_{k=0}^{\infty} k^n w(k)$, are

$$\rho_n = \vartheta_\eta^n \rho_0 = \vartheta_\eta^n \Big({}_M F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix} \Big), \qquad \qquad \vartheta_\eta \coloneqq = \eta \frac{\partial}{\partial \eta}.$$

Given a function $f(\eta)$, we consider the Wronskian

$$\mathscr{W}_k(f) := egin{pmatrix} f & artheta_\eta f & artheta_\eta^2 f & \cdots & artheta_\eta^k f \ artheta_\eta f & artheta_\eta^2 f & \cdots & artheta_\eta^k f \ artheta_\eta^2 f & \ddots & artheta_\eta^k f \ artheta_z^2 f & artheta_\eta^{k+1} f & \cdots & artheta_\eta^{2k} f \ artheta_z^k f & artheta_\eta^{k+1} f & \cdots & artheta_\eta^{2k} f \ artheta_z^{k+1} f & artheta_\eta^{k+1} f & \cdots & artheta_\eta^{2k} f \ artheta_z^{k+1} f & artheta_z^{k+1} f$$

Let us introduce the following tau-functions

$$\tau_k := \mathscr{W}_k \Big({}_M F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix} \Big),$$

i.e. Wronskians of generalized hypergeometric functions. Then, we have that

$$\Delta_k = \tau_k, \qquad \qquad \tilde{\Delta}_k = \vartheta_\eta \tau_k, \qquad \qquad H_k = \frac{\tau_{k+1}}{\tau_k}, \qquad \qquad p_k^1 = -\vartheta_\eta \log \tau_k.$$

In [42] it was shown that:

Theorem 1 (Hypergeometric symmetries). Let the weight w be subject to a discrete Pearson equation of the type (17), where the functions θ , σ are polynomials, with $\theta(0) = 0$. Then,

i) The moment matrix fulfills

(18)
$$\theta(\Lambda)G = B\sigma(\Lambda)GB^{\top}.$$

ii) The Jacobi matrix satisfies

(19)

$$\Pi^{-1}H\theta(J^{\top}) = \sigma(J)H\Pi^{\top}$$

and the matrices $H\theta(J^{\top})$ and $\sigma(J)H$ are symmetric.

Theorem 2 (Laguerre–Freud structure matrix). Let us assume that the weight w solves the discrete Pearson equation (17) with θ, σ polynomials such that $\theta(0) = 0$, deg $\theta(z) = N + 1$, deg $\sigma(z) = M$. Then,:

i) The Laguerre–Freud structure matrix

(20)
$$\Psi := \Pi^{-1} H \theta(J^{\top}) = \sigma(J) H \Pi^{\top} = \Pi^{-1} \theta(J) H = H \sigma(J^{\top}) \Pi^{\top}$$

(21)
$$= \theta(J+I)\Pi^{-1}H = H\Pi^{\top}\sigma(J^{\top}-I),$$

has only N + M + 2 possibly nonzero diagonals (N + 1 superdiagonals and M subdiagonals)

$$\Psi = (\Lambda^{\top})^{M} \psi^{(-M)} + \dots + \Lambda^{\top} \psi^{(-1)} + \psi^{(0)} + \psi^{(1)} \Lambda + \dots + \psi^{(N+1)} \Lambda^{N+1},$$

for some diagonal matrices $\psi^{(k)}$. In particular, the lowest subdiagonal and highest superdiagonal are given by

(22)
$$\begin{cases} (\Lambda^{\top})^{M}\psi^{(-M)} = \eta(J_{-})^{M}H, \quad \psi^{(-M)} = \eta H \prod_{k=0}^{M-1} T_{-}^{k}\gamma = \eta \operatorname{diag}\left(H_{0} \prod_{k=1}^{M} \gamma_{k}, H_{1} \prod_{k=2}^{M+1} \gamma_{k}, \ldots\right), \\ \psi^{(N+1)}\Lambda^{N+1} = H(J_{-}^{\top})^{N+1}, \quad \psi^{(N+1)} = H \prod_{k=0}^{N} T_{-}^{k}\gamma = \operatorname{diag}\left(H_{0} \prod_{k=1}^{N+1} \gamma_{k}, H_{1} \prod_{k=2}^{N+2} \gamma_{k}, \ldots\right). \end{cases}$$

ii) The vector P(z) of orthogonal polynomials fulfill the following structure equations

(23)
$$\theta(z)P(z-1) = \Psi H^{-1}P(z), \qquad \sigma(z)P(z+1) = \Psi^{\top}H^{-1}P(z).$$

iii) The following compatibility conditions for the Laguerre-Freud and Jacobi matrices hold

(24a)
$$[\Psi H^{-1}, J] = \Psi H^{-1}$$

(24b)
$$[J, \Psi^{\top} H^{-1}] = \Psi^{\top} H^{-1}.$$

We now recall some facts regarding the Toda flows appearing in this discrete orthogonal polynomial setting. Let us define the strictly lower triangular matrix $\Phi := (\vartheta_{\eta}S)S^{-1}$. Then, see [42], we have that the orthogonal polynomial sequence P fulfills $\vartheta_{\eta}P = \Phi P$, and the Sato–Wilson equations $-\Phi H + \vartheta_{\eta}H - H\Phi^{\top} = JH$ are satisfied. Consequently, $\Phi = -J_{-}$ and, for $n \in \mathbb{N}_{0}$, we have $\vartheta_{\eta} \log H_{n} = J_{n,n}$. Moreover, $\Phi = (\vartheta_{\eta}S)S^{-1} = -\Lambda^{\top}\gamma$ and $(\vartheta_{\eta}H)H^{-1} = \beta$ are satisfied. The functions $q_{n} := \log H_{n}, n \in \mathbb{N}$, satisfy the Toda equations

(25)
$$\vartheta_n^2 q_n = e^{q_{n+1}-q_n} - e^{q_n-q_{n-1}}$$

For $n \in \mathbb{N}$, we also have $\vartheta_{\eta}P_n(z) = -\gamma_n P_{n-1}(z)$. Additionally, the Lax equation $\vartheta_{\eta}J = [J_+, J]$ is fulfilled, and recursion coefficients satisfy the following Toda system, $\vartheta_{\eta}\beta_n = \gamma_{n+1} - \gamma_n$, and $\vartheta_{\eta}\log\gamma_n = \beta_n - \beta_{n-1}$, for $n \in \mathbb{N}_0$ and $\beta_{-1} = 0$. Thus, we get $\vartheta_{\eta}^2\log\gamma_n + 2\gamma_n = \gamma_{n+1} + \gamma_{n-1}$.

From the compatibility of $P(z+1) = \Pi P(z)$ and $\vartheta_{\eta}(P(z)) = \Phi P(z)$ we get $\vartheta_{\eta}\Pi = [\Phi, \Pi]$. In the general case, the dressed Pascal matrix Π is a lower unitriangular semi-infinite matrix, that possibly has an infinite number of subdiagonals. However, when the weight $w(z) = v(z)\eta^z$ satisfies the Pearson equation (17), with v independent of η , that is $\theta(k+1)v(k+1)\eta = \sigma(k)v(k)$, the situation improves as we have the banded semi-infinite matrix Ψ that models the shift in the z variable as in (23). From the previous discrete Pearson equation we see that $\sigma(z) = \eta \kappa(z)$ with κ, θ η -independent polynomials in z

(26)
$$\theta(k+1)v(k+1) = \eta \kappa(k)v(k).$$

Proposition 1 ([42]). Let us assume a weight w satisfying the Pearson equation (17). Then, the Laguerre–Freud structure matrix Ψ given in (20) satisfies

(27a)
$$\vartheta_{\eta}(\eta^{-1}\Psi^{\top}H^{-1}) = [\Phi, \eta^{-1}\Psi^{\top}H^{-1}],$$

(27b)
$$\vartheta_{\eta}(\Psi H^{-1}) = [\Phi, \Psi H^{-1}].$$

Relations (27a) and (27b) are gauge equivalent.

2. The $_1F_2$ hypergeometrical family

We choose $\sigma(z) = \eta(z + a_1)$ and $\theta(z) = z(z + b_1)(z + b_2)$ with corresponding weight given by

$$w(z) = \frac{(a_1)_z}{(b_1+1)_z(b_2+1)_z} \frac{\eta^z}{z!}$$

with first moment given by the generalized hypergeometric function: $\rho_0 = {}_1F_2\begin{bmatrix}a_1\\b_1+1,b_2+1\\;\eta\end{bmatrix}$, and subsequent moments $\rho_n = \vartheta_{\eta}^n F_2\begin{bmatrix}a_1\\b_1+1,b_2+1\\;\eta\end{bmatrix}$. The corresponding moments exist for any $\eta \in \mathbb{C}$.

Theorem 3 (The generalized Laguerre–Freud structure matrix). For $\sigma(z) = \eta(z + a_1)$ and $\theta(z) = z(z + b_1)(z + b_2)$ we find for the subleading coefficients the following expression

$$p_n^1 = \frac{n(n+1)}{2} - n\beta_n - \frac{1}{\eta + \gamma_{n+1}}(\gamma_{n+1}(\gamma_n + \gamma_{n+1} + \gamma_{n+2} + (\beta_{n+1} + b_1)(\beta_{n+1} + b_2) + \beta_n(\beta_{n+1} + \beta_n + b_1 + b_2) - n(2\beta_n + \beta_{n+1} + b_1 + b_2 - n) - \eta) - \eta(n+1)(\beta_n + a_1))$$

and we also find

$$\pi_{n-1}^{[2]} = \frac{1}{\eta + \gamma_{n+1}} \Big(\gamma_{n+1} \big(\gamma_n + \gamma_{n+1} + \gamma_{n+2} + (\beta_{n+1} + b_1)(\beta_{n+1} + b_2) + \beta_n (\beta_{n+1} + \beta_n + b_1 + b_2) - n(2\beta_n + \beta_{n+1} + b_1 + b_2 - 2n) - \eta \Big) - \eta (n+1)(a_1 + a_2) \Big).$$

The Laguerre–Freud structure matrix is the following pentadiagonal matrix

$$\Psi = \begin{bmatrix} \psi_0^{(0)} \ \psi_0^{(1)} \ \psi_0^{(2)} \ H_3 & 0 \cdots \\ \eta H_1 \ \psi_1^{(0)} \ \psi_1^{(1)} \ \psi_1^{(2)} \ H_4 & \ddots \\ 0 & \eta H_2 \ \psi_2^{(0)} \ \psi_2^{(1)} \ \psi_2^{(2)} \ H_5 & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \end{bmatrix},$$

with

$$\begin{split} \psi_n^{(0)} &= \eta H_n(n+\beta_n+a_1), \\ \psi_n^{(1)} &= \eta \Big(H_{n+1} + H_n \big(\pi_{n-1}^{[2]} + (n+1)(\beta_n+a_1) \big) \Big), \\ \psi_n^{(2)} &= H_{n+2}(\beta_n+\beta_{n+1}+\beta_{n+2}+b_1+b_2-n). \end{split}$$

Proof. As deg $\sigma = 1$ and deg $\theta = 3$ for the Lagure–Freud matrix we have

$$\Psi = \Lambda^{\top} \psi^{(-1)} + \psi^{(0)} + \psi^{(1)} \Lambda + \psi^{(2)} \Lambda^2 + \psi^{(3)} \Lambda^3.$$

Form $\Psi = \sigma(J)H\Pi^{\top}$ we get the first subdiagonal, the main diagonal and an expression for the first superdiagonal depending upon $\pi^{[2]}$:

$$\psi^{(-1)} = \eta T_- H,$$

$$\psi^{(0)} = \eta H(T_+D + \beta + a_1),$$

$$\psi^{(1)} = \eta \Big(T_-H + H \big(T_+ \pi^{[2]} + D(\beta + a_1) \big) \Big).$$

From $\Psi = \Pi^{-1} H \theta(J^{\top})$ we get an expression for the first superdiagonal depending on de $T_{+}^{2} \pi^{[-2]}$ and the other two nonzero superdiagonals

$$\begin{split} \psi^{(1)} &= T_{-}H[T_{+}\gamma + \gamma + T_{-}\gamma + (T_{-}\beta + b_{1})(T_{-}\beta + b_{2}) + \beta(T_{-}\beta + \beta + b_{1} + b_{2}) - T_{+}D(T_{+}\beta + \beta + \beta + b_{1} + b_{2}) + T_{+}\beta + \beta + b_{1} + b_{2} + T_{-}\beta) + T_{+}^{2}\pi^{[-2]}], \\ \psi^{(2)} &= T_{-}^{2}H[\beta + T_{-}\beta + T_{-}^{2}\beta + b_{1} + b_{2} - T_{+}D], \\ \psi^{(3)} &= T_{-}^{3}H. \end{split}$$

To obtain the diagonal matrix $\pi^{[\pm 2]}$ we combine $T_+^2 \pi^{[-2]} - T_+ \pi^{[-2]} = T_+ D(T_+\beta - \beta - 1)$ with $\pi^{[-2]} = 2D^{[2]} - \pi^{[2]}$ to get

$$T_{+}^{2}\pi^{[-2]} = T_{+}D(T_{+}\beta - \beta - 1) + 2T_{+}D^{[2]} - T_{+}\pi^{[2]}.$$

Now, equating the two formulas for $\psi^{(1)}$ and using the previous results we find

$$T_{+}\pi^{[2]}(\eta H + T_{-}H) = T_{-}H[T_{+}\gamma + \gamma + T_{-}\gamma + (T_{-}\beta + b_{1})(T_{-}\beta + b_{2}) + \beta(T_{-}\beta + \beta + b_{1} + b_{2}) - T_{+}D(\beta + T_{-}\beta + b_{1} + b_{2}) - T_{+}D(\beta + 1) + 2T_{+}D^{[2]} - \eta I] - \eta HD(\beta + a_{1}).$$

Consequently, we derive the given expression for p_n^1 .

We now explore the consequences of the compatibility relation $[\Psi H^{-1}, J] = \Psi H^{-1}$. We get trivial relations and $T_+\pi^{[2]} - T_+^2\pi^{[2]} = T_+D(T_+\beta - \beta + I)$. However, a new formula appears from the first superdiagonal:

$$\begin{split} \eta(\beta + T_{-}\beta - 1) &= T_{-}\gamma(T_{+}D - 2T_{-}\beta - T_{-}^{2}\beta + b_{1} + b_{2} + 1) + \gamma(\beta + 1 - T_{-}\beta) \\ &+ T_{+}\gamma(T_{+}\beta + 2\beta + b_{1} + b_{2} - T_{+}^{2}D + 1) + \beta^{3} - (T_{-}\beta)^{3} + (T_{-}\beta)^{2} + \beta^{2} + \beta T_{-}\beta + \\ &(b_{1} + b_{2})(\beta^{2} - (T_{-}\beta)^{2} + T_{-}\beta + \beta) - T_{+}D(2\beta^{2} - \beta T_{-}\beta - (T_{-}\beta)^{2} + 3\beta) \\ &+ (\beta + 1 - T_{-}\beta)[2T_{+}D^{[2]} - T_{+}D(b_{1} + b_{2}) + b_{1}b_{2} - T_{+}\pi^{[2]}]. \end{split}$$

No new relations are obtained form the second compatibility (27a). Using this compatibility, we seek for Laguerre–Freud equations

(28)
$$\beta_{n+1} = \mathscr{B}_n(\gamma_{n+1}, \gamma_n, \dots, \beta_n, \beta_{n-1}, \dots), \qquad \gamma_{n+2} = \mathscr{G}_n(\gamma_{n+1}, \gamma_n, \dots, \beta_{n+1}, \beta_n, \dots).$$

Theorem 4. We find the following Laguerre–Freud equations

$$\begin{split} \gamma_{n+2} &= \frac{\eta + \gamma_{n+1}}{\gamma_{n+1}} n(\beta_{n-1} - \beta_n + 1) - \left(\gamma_n + \gamma_{n+1} + (\beta_{n+1} + b_1)(\beta_{n+1} + b_2) + \beta_n(\beta_{n+1} + \beta_n + b_1 + b_2) - n(2\beta_n + \beta_{n+1} + b_1 + b_2 - n) - \eta\right) + \frac{\eta}{\gamma_{n+1}} (n+1)(\beta_n + a_1) \\ &+ \frac{\eta + \gamma_{n+1}}{(\eta + \gamma_n)\gamma_{n+1}} \left(\gamma_n(\gamma_{n-1} + \gamma_n + \gamma_{n+1} + (\beta_n + b_1)(\beta_n + b_2) + \beta_{n-1}(\beta_n + \beta_{n-1} + b_1 + b_2) - (n-1)(2\beta_{n-1} + \beta_n + b_1 + b_2 - n + 1) - \eta\right) - \eta n(\beta_{n-1} + a_1) \Big), \end{split}$$

$$\square$$

$$\beta_{n+2} = n - 2\beta_{n+1} + b_1 + b_2 + 1 + \frac{1}{\gamma_{n+2}} \Big(\eta (1 - \beta_n - \beta_{n+1}) + \gamma_{n+1} (\beta_n + 1 - \beta_{n+1}) \\ + \gamma_n (\beta_{n-1} + 2\beta_n + b_1 + b_2 - n + 2) + \beta_n^3 - \beta_{n+1}^3 + \beta_{n+1}^2 + \beta_n^2 + \beta_n \beta_{n+1} \\ + (b_1 + b_2) (\beta_n^2 - \beta_{n+1}^2 + \beta_{n+1} + \beta_n) - n(2\beta_n^2 - \beta_n \beta_{n+1} - \beta_{n+1}^2 + 3\beta_n) \\ + (\beta_n - \beta_{n+1} + 1) \big(n(n+1) - n(b_1 + b_2) + b_1 b_2 \\ - \frac{1}{\eta + \gamma_{n+1}} (\gamma_{n+1} (\gamma_n + \gamma_{n+1} + \gamma_{n+2} + (\beta_{n+1} + b_1) (\beta_{n+1} + b_2) \\ + \beta_n (\beta_{n+1} + \beta_n + b_1 + b_2) - n(2\beta_n + \beta_{n+1} + b_1 + b_2 - n) - \eta) - \eta (n+1) (\beta_n + a_1)) \Big) \Big).$$

Proof. To get $\gamma_{n+2} = F_1(n, \gamma_{n+1}, \gamma_n, \dots, \beta_{n+1}, \beta_n, \dots)$ we consider the compatibility $T_+\pi^{[2]} - T_+^2\pi^{[2]} = T_+D(T_+\beta + 1 - \beta)$ and solve γ_{n+2} . Also, we obtain $\beta_{n+2} = F_2(n, \gamma_{n+2}, \gamma_{n+1}, \dots, \beta_{n+1}, \beta_n, \dots)$ by solving for β_{n+2} in the first subdiagonal of the first type of compatibility.

3. The $_2F_2$ hypergeometrical family

Let us take $\sigma(z) = \eta(z+a_1)(z+a_2)$ and $\theta(z) = z(z+b_1)(z+b_2)$ and consider the corresponding Pearson equation

$$(k+1)(k+1+b_1)(k+1+b_2)w(k+1) = \eta(k+a_1)(k+a_2)w(k)$$

whose solutions are proportional to $w(z) = \frac{(a_1)_z(a_2)_z}{(b_1+1)_z(b_2+1)_z} \frac{\eta^z}{z!}$, so that he discrete orthogonality measure is

$$\rho_z = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k}{(b_1+1)_k (b_2+1)_k} \frac{\eta^k}{k!}$$

with first moment given by the generalized hypergeometric function: $\rho_0 = {}_2F_2\begin{bmatrix} a_1, a_2\\ b_1+1, b_2+1 \end{bmatrix}$, and subsequent moments $\rho_n = \vartheta_{\eta 2}^n F_2\begin{bmatrix} a_1, a_2\\ b_1+1, b_2+1 \end{bmatrix}$. The corresponding moments exist for any $\eta \in \mathbb{C}$.

Definition 1. Let us introduce

$$A_{n} := -\eta^{2} \Big(\frac{n(n+1)}{2} + \beta_{n-1} + \beta_{n} + \gamma_{n+2} + \gamma_{n+1} + (n+1)(\beta_{n+1} + a_{1} + a_{2}) + (\beta_{n+1} + a_{1})(\beta_{n+1} + a_{2}) \Big) \\ + \eta \Big(4\gamma_{n} + \gamma_{n+1}(a_{1} + a_{2} - b_{1} - b_{2} + 2(n+1) + \beta_{n-1} - \beta_{n+1}) + \gamma_{n+2}(n-1 - 2\beta_{n+1} - \beta_{n+2} - b_{1} - b_{2}) \\ + \frac{n(n+1)}{2}(a_{1} + a_{2} - b_{1} - b_{2} - \beta_{n+1}) + (\beta_{n+1} + b_{1} + b_{2} + a_{1} + a_{2} - n^{2})(\beta_{n} + \beta_{n-1}) + 2(\beta_{n}^{2} + \beta_{n-1}^{2}) \\ - (\beta_{n+1} + b_{1})(\beta_{n+1} + b_{2})(\beta_{n+1} - n - 1) \Big) \\ = 2(-\alpha \Big(\frac{n(n-1)}{2} + \beta_{n+2} + \beta_{n+1} + \beta_{n+1} + \beta_{n+1} + b_{2}) + (\beta_{n+1} - n - 1) \Big)$$

$$-\gamma_{n+1}\Big(\frac{n(n-1)}{2}+\gamma_{n+2}+\gamma_{n+1}+\gamma_n+\beta_{n-1}+(\beta_{n+1}+b_1)(\beta_{n+1}+b_2)+(\beta_n-n)(\beta_{n+1}+\beta_n+b_1+b_2)\Big),$$

and

(29)
$$B_n := \eta^2 + \eta (2\beta_{n-1} + 2\beta_n + \beta_{n+1} + a_1 + a_2 + b_1 + b_2 + 2n) + \gamma_{n+1}.$$

Theorem 5 (The generalized Laguerre–Freud structure matrix). For $\sigma(z) = \eta(z + a_1)(z + a_2)$ and $\theta(z) = z(z + c_1)(z + c_2)$ we find for the subleading coefficients the following expression

(30)
$$p_{n-1}^{1} = \frac{A_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \beta_{n+2}, \gamma_{n}, \gamma_{n+1}, \gamma_{n+2})}{B_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \gamma_{n+1})}.$$

The Laguerre–Freud structure matrix is the following hexadiagonal matrix

$$\Psi = \begin{bmatrix} \psi_0^{(0)} & \psi_0^{(1)} & \psi_0^{(2)} & H_3 & 0 \cdots \\ \psi_0^{(-1)} & \psi_1^{(0)} & \psi_1^{(1)} & \psi_1^{(2)} & H_4 & \ddots \\ \eta H_2 & \psi_1^{(-1)} & \psi_2^{(0)} & \psi_2^{(1)} & \psi_2^{(2)} & H_5 & \ddots \\ 0 & \eta H_3 & \psi_2^{(-1)} & \psi_3^{(0)} & \psi_3^{(1)} & \psi_3^{(2)} & H_6 & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \vdots & \ddots \\ \end{bmatrix}$$

with

$$\begin{split} \psi_n^{(2)} &= (\beta_n + \beta_{n+1} + \beta_{n+2} + b_1 + b_2 - n)H_{n+2}, \\ \psi_n^{(0)} &= \eta \Big(\frac{n(n-1)}{2} + \beta_{n-1} + n(\beta_n + a_1 + a_2) + \gamma_n + \gamma_{n+1} + (\beta_n + a_1)(\beta_n + a_2) - p_{n-1}^1\Big)H_n, \\ \psi_n^{(1)} &= \Big(\frac{n(n-1)}{2} + \gamma_n + \gamma_{n+1} + \gamma_{n+2} + (\beta_{n+1} + b_1)(\beta_{n+1} + b_2) \\ &+ (\beta_n - n)(\beta_n + \beta_{n+1} + b_1 + b_2) - \beta_{n-1} + p_{n-1}^1\Big)H_{n+1}, \end{split}$$

 $\psi_n^{(-1)} = \eta(\beta_n + \beta_{n+1} + a_1 + a_2 + n)H_{n+1}.$

Proof. We write (20) and (21) as

(31)
$$\Psi = \sigma(J)H\Pi^{\top},$$

(32)
$$\Psi = \Pi^{-1} H \theta(J^{\top}).$$

The Laguerre–Freud matrix Ψ is a hexadiagonal matrix

$$\Psi = (\Lambda^{\top})^2 \psi^{(-2)} + \Lambda^{\top} \psi^{(-1)} + \psi^{(0)} + \psi^{(1)} \Lambda + \psi^{(2)} \Lambda^2 + \psi^{(3)} \Lambda^3,$$

with $\psi^{(n)}$ diagonal matrices. From Equation (31) we get

$$\psi^{(-2)} = \eta T_{-}^{2} H,$$
 $\psi^{(-1)} = \eta T_{-} H (T_{+} D + \beta + T_{-} \beta + a_{1} + a_{2}).$

From Equation (32) we obtain

$$\psi^{(3)} = T_{-}^{3}H,$$
 $\psi^{(2)} = T_{-}^{2}H(T_{-}^{2}\beta + \beta + T_{-}\beta - T_{+}D + b_{1} + b_{2}).$

We also have expressions for the diagonal matrices $\psi^{(0)}, \psi^{(1)}$ in terms of $\pi^{[\pm 2]}, \pi^{[\pm 3]}$. From Equation (31) we get

$$\begin{aligned} &(33)\\ \psi^{(0)} &= \eta H[T_+^2 \pi^{[2]} + T_+ D(T_+ \beta + \beta + a_1 + a_2) + \gamma + T_+ \gamma + (\beta + a_1)(\beta + a_2)], \\ &(34)\\ \psi^{(1)} &= \eta H[\gamma(\beta + T_- \beta + a_1 + a_2) + D(\gamma + T_+ \gamma + (\beta + a_1)(\beta + a_2)) + T_+^2 \pi^{[3]} + T_+ \pi^{[2]}(T_+ \beta + \beta + a_1 + a_2)], \\ &\text{and Equation (32) leads to} \end{aligned}$$

(35)

$$\psi^{(0)} = H[(\beta + b_1)(\beta + b_2)(\beta - T_+D) + \gamma(2\beta + T_-\beta + b_1 + b_2 - T_+D) + T_+\gamma(2\beta + T_+\beta + b_1 + b_2 - T_+D) - T_+D[T_+^2\gamma + T_+\beta(\beta + T_+\beta + b_1 + b_2)] + T_+^2\pi^{[-2]}(\beta + T_+\beta + T_+\beta + b_1 + b_2) - T_+^3\pi^{[-3]}],$$

(36)

$$\psi^{(1)} = T_{-}H[T_{+}^{2}\pi^{[-2]} - T_{+}D(T_{-}\beta + \beta + T_{+}\beta + b_{1} + b_{2}) + T_{-}\gamma + \gamma + T_{+}\gamma + (T_{-}\beta + b_{1})(T_{-}\beta + b_{2}) + \beta(T_{-}\beta + \beta + b_{1} + b_{2})].$$

We get a system of two equations by equating the RHS of (33) and (35) and the RHS of (34) and (36). Using the shift $n \to n+1$ in the first relation obtained from the two possible expressions for $\psi^{(0)}$, we now show that the π matrices involved in the system can be expressed uniquely in terms of p_n^2, p_{n-1}^1 . From the relations

(37)
$$p_n^1 = p_{n-1}^1 - \beta_{n-1},$$

(38)
$$p_{n+1}^1 = p_{n-1}^1 - \beta_{n-1} - \beta_n,$$

(39)
$$p_{n+1}^2 = p_n^2 - \gamma_n - \beta_n p_{n-1}^1 + \beta_n \beta_{n-1},$$

we get

Then, we solve the equation for p_n^2 in the equation obtained for the two expressions (33) and (35) for $\psi^{(0)}$ and substitute in the Equation gotten from the two expressions (34) and (36) for $\psi^{(1)}$ to deduce

$$(40) \quad p_{n-1}^{1} \left(\eta^{2} + \eta (2\beta_{n-1} + 2\beta_{n} + \beta_{n+1} + a_{1} + a_{2} + b_{1} + b_{2} + 2n) + \gamma_{n+1} \right)$$

$$= -\eta^{2} \left(\frac{n(n+1)}{2} + \beta_{n-1} + \beta_{n} + \gamma_{n+2} + \gamma_{n+1} + (n+1)(\beta_{n+1} + a_{1} + a_{2}) + (\beta_{n+1} + a_{1})(\beta_{n+1} + a_{2}) \right)$$

$$+ \eta \left(\gamma_{n+1}(a_{1} + a_{2} - b_{1} - b_{2} + 2(n+1) + \beta_{n-1} - \beta_{n+1}) + \gamma_{n+2}(n-1 - 2\beta_{n+1} - \beta_{n+2} - b_{1} - b_{2}) + n(\beta_{n-1} + \beta_{n}) + (a_{1} + a_{2})\beta_{n-1} + 4\gamma_{n} + 2(\beta_{n}^{2} + \beta_{n-1}^{2}) + \beta_{n}(\beta_{n+1} + a_{1} + a_{2} + b_{1} + b_{2}) + (n+1)a_{1}a_{2} + \beta_{n-1}(\beta_{n+1} + b_{1} + b_{2}) - (\beta_{n+1} + b_{1})(\beta_{n+1} + b_{2})(\beta_{n+1} - n - 1)$$

$$+ \frac{n(n+1)}{2}(a_{1} + a_{2} - b_{1} - b_{2} - \beta_{n+1}) - n^{2}(\beta_{n} + \beta_{n-1}) \right)$$

$$- \gamma_{n+1} \left(\frac{n(n-1)}{2} + \gamma_{n+2} + \gamma_{n+1} + \gamma_{n} + \beta_{n-1} - n(\beta_{n+1} + \beta_{n} + b_{1} + b_{2}) + (\beta_{n+1} + b_{1})(\beta_{n+1} + b_{2}) + \beta_{n}(\beta_{n+1} + \beta_{n} + b_{1} + b_{2}) \right).$$

That is, we have proven Equation (30) for p_n^1 . It is not necessary at this point to find p_n^2 in order to get the Freud–Laguerre matrix, as we can get $\psi^{(0)}$ from (33) and $\psi^{(1)}$ from (36). The entries in the diagonal matrices $\psi^{(0)}, \psi^{(1)}$ can be written in terms of p_{n-1}^1 as:

$$\begin{split} \psi_n^{(0)} &= \eta H_n \Big(\frac{n(n-1)}{2} + \beta_{n-1} + n(\beta_n + a_1 + a_2) + \gamma_n + \gamma_{n+1} + (\beta_n + a_1)(\beta_n + a_2) - p_{n-1}^1 \Big), \\ \psi_n^{(1)} &= \gamma_{n+1} H_n \Big(\frac{n(n-1)}{2} + \gamma_{n+1} + \gamma_{n+2} + \gamma_n + (\beta_{n+1} + b_1)(\beta_{n+1} + b_2) \\ &+ (\beta_n - n)(\beta_n + \beta_{n+1} + b_1 + b_2) - \beta_{n-1} + p_{n-1}^1 \Big). \end{split}$$

Definition 2. Let us consider

(41)

$$C_{n} := \eta \left((n + \beta_{n} + \beta_{n+1} + a_{1} + a_{2})(\beta_{n} - \beta_{n+1} - 1) - \beta_{n} - \beta_{n+1} + \gamma_{n} \right) \\ - n(\beta_{n+1} - \beta_{n} - 1)(\beta_{n+1} + \beta_{n} + \beta_{n-1}) - \beta_{n}(\beta_{n+1} + \beta_{n} + b_{1} + b_{2}) \\ - \gamma_{n}(\beta_{n-1} + 2\beta_{n} + b_{1} + b_{2} - n + 2) + \gamma_{n+2}(2\beta_{n+1} + b_{1} + b_{2} - n - 1),$$

,

(42)
$$D_n := \frac{n(n-1)}{2} + \beta_{n+1}^2 + (\beta_{n+1} - n)(b_1 + b_2) + b_1b_2 + (n-1)\beta_{n-1} + \gamma_{n+1}$$
and

$$(43) \quad E_n := -\eta \Big(\frac{n(n-1)}{2} + \beta_{n-1} + n(\beta_n + a_1 + a_2) + (\beta_n + a_1)(\beta_n + a_2) \\ - \gamma_n(n-2 + \beta_{n-1} + \beta_n + a_1 + a_2) + \gamma_{n+1}(n+1 + \beta_n + \beta_{n+1} + a_1 + a_2) \Big) \\ + \gamma_{n+1} \Big(\frac{n(n-1)}{2} + \gamma_{n+1} - \beta_{n-1} + (\beta_n - n)(\beta_{n+1} + \beta_n + b_1 + b_2) + (\beta_{n+1} + b_1)(\beta_{n+1} + b_2) \Big) \\ - \gamma_n \Big(\frac{(n-1)(n-2)}{2} + \gamma_n + \gamma_{n-1} + (\beta_n + b_1)(\beta_n + b_2) + (\beta_{n-1} - n + 1)(\beta_n + \beta_{n-1} + b_1 + b_2) \Big) \Big)$$

Theorem 6 (Consequences of compatibility conditions). The compatibility conditions lead to two alternative relations expressions for the subleading coefficient p_{n-1}^1 of the orthogonal polynomials:

(44)
$$p_{n-1}^{1} = \frac{C_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \gamma_{n}) + \gamma_{n+2}(\beta_{n+2} - \eta)}{1 + \beta_{n} - \beta_{n+1}} - D_{n}(\beta_{n-1}, \beta_{n+1}, \gamma_{n+1}),$$

(45)
$$p_{n-1}^{1} = \frac{E_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \gamma_{n-1}, \gamma_{n}, \gamma_{n+1}) + \gamma_{n+1}\gamma_{n+2}}{\gamma_{n} - \gamma_{n+1} - \eta}.$$

where p_n^1 is given in Equation (30).

Proof. We compute explicitly the compatibility equation $[\Psi H^{-1}, J] = \Psi H^{-1}$. In the one hand ΨH^{-1} can be expressed as

$$\Psi H^{-1} = (\Lambda^{\top})^2 M^{(-2)} + \Lambda^{\top} M^{(-1)} + M^{(0)} + M^{(1)} \Lambda + M^{(2)} \Lambda^2 + M^{(3)} \Lambda^3$$

with $M^{(n)}$ being the following diagonal matrices

$$M^{(0)} = \eta \Big(T_{+}^{2} \pi^{[2]} + T_{+} D(T_{+}\beta + \beta + a_{1} + a_{2}) + \gamma + T_{+}\gamma + (\beta + a_{1})(\beta + a_{2}) \Big),$$

or alternatively

$$\begin{split} M^{(0)} &= \beta^3 + (b_1 + b_2)\beta^2 + b_1b_2\beta - T_+D[\beta^2 + (b_1 + b_2)\beta + b_1b_2 + T_+\beta(\beta + T_+\beta + b_1 + b_2)] \\ &+ \gamma(2\beta + T_-\beta + b_1 + b_2 - T_+D) + T_+\gamma(2\beta + T_+\beta + b_1 + b_2 - T_+D) \\ &- T_+^2\gamma T_+D + T_+^2\pi^{[-2]}(\beta + T_+\beta + T_+^2\beta + b_1 + b_2) - T_+^3\pi^{[-3]}, \end{split}$$

 $M^{(-1)} = \eta \gamma (T_{+}D + \beta + T_{-}\beta + a_{1} + a_{2}),$ $M^{(-2)} = \eta \gamma (T_{-}\gamma).$ In the other hand, the commutator can be written as follows

$$[\Psi H^{-1}, J] = (\Lambda^{\top})^2 \tilde{M}^{(-2)} + \Lambda^{\top} \tilde{M}^{(-1)} + \tilde{M}^{(0)} + \tilde{M}^{(1)} \Lambda + \tilde{M}^{(2)} \Lambda^2 + \tilde{M}^{(3)} \Lambda^3,$$

where the diagonal matrices $M^{(n)}$ are

$$\begin{split} \tilde{M}^{(3)} &= I, \\ \tilde{M}^{(2)} &= T_{+}^{2} \pi^{[-2]} - T_{+} \pi^{[-2]} + T_{+} D(\beta - T_{+}\beta) + T_{-}^{2}\beta + T_{-}\beta + \beta + b_{1} + b_{2}, \\ \tilde{M}^{(1)} &= \eta \left(T_{+}^{2} \pi^{[2]} - T_{+} \pi^{[2]} - (\beta + a_{1} + a_{2}) + T_{+} (D\beta) - DT_{-}\beta \right) \\ &+ (T_{-}\beta - \beta) \left(T_{+}^{2} \pi^{[-2]} - T_{+} D(T_{+}\beta + b_{1} + b_{2}) + \gamma + (T_{-}\beta + b_{1})(T_{-}\beta + b_{2}) - \eta(a_{1} + a_{2}) \right) \\ &+ (\beta^{2} - (T_{-}\beta)^{2})(T_{+}D + \eta) + T_{-}\gamma (T_{-}^{2}\beta + 2T_{-}\beta - T_{+}D + b_{1} + b_{2} - \eta) \\ &- T_{+}\gamma (2\beta + T_{+}\beta - T_{+}^{2}D + b_{1} + b_{2} - \eta), \end{split}$$

$$\begin{split} \tilde{M}^{(0)} &= \eta T_{+} \gamma \left(T_{+}^{2} D + T_{+} \beta + \beta + a_{1} + a_{2} \right) \\ &+ \gamma \left(T_{+}^{2} \pi^{[-2]} - T_{+} D (T_{-} \beta + \beta + T_{+} \beta + b_{1} + b_{2}) + T_{-} \gamma + \gamma + (T_{-} \beta + b_{1}) (T_{-} \beta + b_{2}) \right) \\ &+ \beta (T_{-} \beta + \beta + b_{1} + b_{2}) \right) - T_{+} \gamma \left(T_{+}^{3} \pi^{[-2]} - T_{+}^{2} D (\beta + T_{+} \beta + T_{+}^{2} \beta + b_{1} + b_{2}) \right) \\ &+ T_{+} \gamma + T_{+}^{2} \gamma + (\beta + b_{1}) (\beta + b_{2}) \mathbb{T}_{+} \beta (\beta + T_{+} \beta + b_{1} + b_{2}) - \eta \gamma (T_{+} D + \beta + T_{-} \beta + a_{1} + a_{2}), \\ \tilde{M}^{(-1)} &= \eta \gamma (T_{+} \pi^{[2]} - T_{+}^{2} \pi^{[2]} + T_{+} D (-T_{-} \beta - T_{+} \beta) + D (\beta + T_{-} \beta) + (a_{1} + a_{2})), \\ \tilde{M}^{(-2)} &= \eta \gamma (T_{-} \gamma). \end{split}$$

From the equations derived from second superdiagonal and first subdiagonal one obtains, respectively, the following equations

$$T_{+}^{2}\pi^{[-2]} - T_{+}\pi^{[-2]} = T_{+}D(T_{+}\beta - \beta - 1), \qquad T_{+}\pi^{[2]} - T_{+}^{2}\pi^{[2]} = T_{+}D(1 - \beta + T_{+}\beta).$$

Hence , they can be gotten directly from the expressions for the diagonal matrices $\pi^{[\pm 2]}$.

From the first superdiagonal we get (44) and using the alternative expression of the main diagonal it can be seen that it corresponds to the identity:

$$\pi_{n-2}^{[-3]} - \pi_{n-3}^{[-3]} = \pi_{n-2}^{[-2]} (1 + \beta_n - \beta_{n-2}) - (n-1)\gamma_n + n\gamma_{n-1}.$$

Compatibility in the main diagonal is (45).

We discuss now about Laguerre-Freud equations (28).

Definition 3. Let us introduce (46)

$$\begin{split} \hat{A}_n &:= -\eta^2 \Big(\frac{n(n+1)}{2} + \beta_{n-1} + \beta_n + \gamma_{n+1} + (n+1)(\beta_{n+1} + a_1 + a_2) + (\beta_{n+1} + a_1)(\beta_{n+1} + a_2) \Big) \\ &+ \eta \Big(4\gamma_n + \gamma_{n+1}(a_1 + a_2 - b_1 - b_2 + 2(n+1) + \beta_{n-1} - \beta_{n+1}) \\ &+ \frac{n(n+1)}{2}(a_1 + a_2 - b_1 - b_2 - \beta_{n+1}) + (\beta_{n+1} + b_1 + b_2 + a_1 + a_2 - n^2)(\beta_n + \beta_{n-1}) + 2(\beta_n^2 + \beta_{n-1}^2) \\ &- (\beta_{n+1} + b_1)(\beta_{n+1} + b_2)(\beta_{n+1} - n - 1) \Big) \end{split}$$

$$-\gamma_{n+1}\Big(\frac{n(n-1)}{2}+\gamma_{n+1}+\gamma_n+\beta_{n-1}+(\beta_{n+1}+b_1)(\beta_{n+1}+b_2)+(\beta_n-n)(\beta_{n+1}+\beta_n+b_1+b_2)\Big),$$

(47)

$$F_n := -\frac{E_n}{\gamma_n - \gamma_{n+1} - \eta} + \frac{C_n}{1 + \beta_n - \beta_{n+1}} - D_n + \frac{1}{\eta(1 + \beta_n - \beta_{n+1})} \Big(\hat{A}_n - \frac{E_n}{\gamma_n - \gamma_{n+1} - \eta} B_n \Big),$$

(48)

$$G_n := \frac{\gamma_{n+1}}{\gamma_n - \gamma_{n+1} - \eta} + \frac{\eta}{1 + \beta_n - \beta_{n+1}} + \frac{\eta^2 - \eta(n - 1 - 2\beta_{n+1} - b_1 - b_2) + \gamma_{n+1} + \frac{\gamma_{n+1}B_n}{\gamma_n - \gamma_{n+1} - \eta}}{\eta(1 + \beta_n - \beta_{n+1})}.$$

Then, we have the following result

Theorem 7 (Laguerre–Freud equations). Given the functions $\hat{A}_n, B_n, C_n, D_n, E_n, F_n$ and G_n , as in (46), (29),(41),(42),(43),(47) and (48), respectively, the following nonlinear equations of Laguerre–Freud type are satisfied (49)

$$\beta_{n+2} = \frac{\hat{A}_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_n, \gamma_{n+1}) - (\eta^2 - \eta(n-1-2\beta_{n+1}-b_1-b_2) + \gamma_{n+1})\gamma_{n+2}}{\eta\gamma_{n+2}} - \frac{\frac{\eta\gamma_{n+2}}{2}}{\frac{1}{\gamma_n - \gamma_{n+1} - \eta}} - \frac{E_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_{n-1}, \gamma_n, \gamma_{n+1}) + \gamma_{n+1}\gamma_{n+2}}{\gamma_n - \gamma_{n+1} - \eta} - \frac{B_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_{n+1})}{\eta\gamma_{n+2}},$$

(50)

$$\gamma_{n+2} = \frac{F_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_{n-1}, \gamma_n, \gamma_{n+1})}{G_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_n, \gamma_{n+1})}$$

Proof. From (30) we get

(51)
$$\beta_{n+2} = \frac{\hat{A}_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_n, \gamma_{n+1}, \gamma_{n+2}) - p_{n-1}^1 B_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_{n+1})}{\eta \gamma_{n+2}},$$

with

$$\tilde{A}_n := \hat{A}_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_n, \gamma_{n+1}) - (\eta^2 - \eta(n-1-2\beta_{n+1}-b_1-b_2) + \gamma_{n+1})\gamma_{n+2}.$$

Therefore, recalling (44), we get the Laguerre–Freud equation (49).

To prove the second Laguerre-Freud equation (50) we first notice that (49) can be written as

$$2\eta\beta_{n+2}\gamma_{n+2} = A_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_n, \gamma_{n+1}, \gamma_{n+2}) - \frac{E_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_{n-1}, \gamma_n, \gamma_{n+1}) + \gamma_{n+1}\gamma_{n+2}}{\gamma_n - \gamma_{n+1} - \eta} B_n(\eta, \beta_{n-1}, \beta_n, \beta_{n+1}, \gamma_{n+1}).$$

Consequently, recalling (44) we find

$$p_{n-1}^{1} = \frac{C_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \gamma_{n}) - \eta\gamma_{n+2}}{1 + \beta_{n} - \beta_{n+1}} - D_{n}(\beta_{n-1}, \beta_{n+1}, \gamma_{n+1}) + \frac{\gamma_{n+2}\beta_{n+2}}{1 + \beta_{n} - \beta_{n+1}}$$

$$= \frac{C_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \gamma_{n}) - \eta\gamma_{n+2}}{1 + \beta_{n} - \beta_{n+1}} - D_{n}(\beta_{n-1}, \beta_{n+1}, \gamma_{n+1})$$

$$+ \frac{1}{\eta(1 + \beta_{n} - \beta_{n+1})} \Big(\tilde{A}_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \gamma_{n}, \gamma_{n+1}, \gamma_{n+2}) - \frac{E_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \gamma_{n-1}, \gamma_{n}, \gamma_{n+1}) + \gamma_{n+1}\gamma_{n+2}}{\gamma_{n} - \gamma_{n+1} - \eta} B_{n}(\eta, \beta_{n-1}, \beta_{n}, \beta_{n+1}, \gamma_{n+1}) \Big).$$

Thus, we see that the following relation is fulfilled

$$p_{n-1}^{1} = \frac{C_n - \eta \gamma_{n+2}}{1 + \beta_n - \beta_{n+1}} - D_n + \frac{1}{\eta(1 + \beta_n - \beta_{n+1})} \Big(\hat{A}_n - (\eta^2 - \eta(n - 1 - 2\beta_{n+1} - b_1 - b_2) + \gamma_{n+1}) \gamma_{n+2} - \frac{E_n + \gamma_{n+1} \gamma_{n+2}}{\gamma_n - \gamma_{n+1} - \eta} B_n \Big).$$

14

Now, from Equation (44) we get

$$\begin{aligned} \frac{E_n + \gamma_{n+1}\gamma_{n+2}}{\gamma_n - \gamma_{n+1} - \eta} &= \frac{C_n - \eta\gamma_{n+2}}{1 + \beta_n - \beta_{n+1}} - D_n \\ &+ \frac{1}{\eta(1 + \beta_n - \beta_{n+1})} \Big(\hat{A}_n - (\eta^2 - \eta(n - 1 - 2\beta_{n+1} - b_1 - b_2) + \gamma_{n+1})\gamma_{n+2} \\ &- \frac{E_n + \gamma_{n+1}\gamma_{n+2}}{\gamma_n - \gamma_{n+1} - \eta} B_n \Big). \end{aligned}$$
Hence, we find the second Laguerre–Freud equation (50).

Hence, we find the second Laguerre–Freud equation (50).

4. The $_{3}F_{2}$ hypergeometrical case

Let us take $\sigma(z) = \eta(z+a_1)(z+a_2)(z+a_3)$ and $\theta(z) = z(z+b_1)(z+b_2)$ and consider the corresponding Pearson equation

$$(k+1)(k+1+b_1)(k+1+b_2)w(k+1) = \eta(k+a_1)(k+a_2)(k+a_3)w(k)$$

whose solutions are proportional to $w(z) = \frac{(a_1)_z(a_2)_z(a_3)_z}{(b_1+1)_z(b_2+1)_z} \frac{\eta^z}{z!}$, so that he discrete orthogonality measure is

$$\rho_z = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k (a_3)_k}{(b_1 + 1)_k (b_2 + 1)_k} \frac{\eta^k}{k!}$$

with first moment given by the generalized hypergeometric function: $\rho_0 = {}_{3}F_2 \begin{bmatrix} a_1, a_2, a_3 \\ b_1 + 1, b_2 + 1 \end{bmatrix}$, and subsequent moments $\rho_n = \vartheta_{\eta 3}^n F_2 \begin{bmatrix} a_1, a_2, a_3 \\ b_1 + 1, b_2 + 1 \end{bmatrix}$. The corresponding moments exist for any $\eta \in \mathbb{C}$, whenever one of the a's is a non-positive integer, when $|\eta| < 1$ or when $|\eta| = 1$ and

$$\operatorname{Re}(b_1 + b_2 - a_1 - a_2 - a_3) > 0$$

Definition 4. We consider

$$\begin{split} A_n &:= n(n+1) + n(\beta_{n-1} + \beta_n + \beta_{n+1}) + (a_1 + a_2 + a_3)(n + \beta_n + \beta_{n+1}) + a_1a_2 + a_2a_3 + a_3a_1 \\ &+ \gamma_n + \gamma_{n+1} + \gamma_{n+2} + \beta_n^2 + \beta_{n+1}^2 + \beta_n\beta_{n+1}, \end{split}$$

$$B_n &= -n(\beta_n^2 - \beta_{n+1}^2 + \beta_{n+1} + \beta_n - \gamma_{n+2}) + (\beta_{n+1} - \beta_n - 1)(n(b_1 + b_2 - \beta_{n-1}) - b_1b_2) \\ &- (b_1 + b_2)(\beta_{n+1}^2 - \beta_n^2 - \beta_n - \beta_{n+1}) - \gamma_{n+2}(\beta_{n+2} + 2\beta_{n+1} + b_1 + b_2 - 1) - \gamma_{n+1}(\beta_{n+1} - \beta_n - 1) \\ &+ \gamma_n(\beta_{n-1} + 2\beta_n + b_1 + b_2 - (n-2)) - \beta_{n+1}^3 + \beta_n^3 + \beta_n^2 + \beta_{n+1}^2 + \beta_n\beta_{n+1}, \end{split}$$

$$\begin{split} C_n &= \eta + \beta_{n+1} - \beta_{n-2}, \\ D_n &= \beta_n^3 + (b_1 + b_2) \left(\beta_n^2 - \beta_{n-1} - \beta_{n-2} + \frac{n(n-1)}{2} \right) + b_1 b_2 \beta_n \\ &\quad - n[\beta_n^2 - \beta_{n-1}\beta_{n-2} + (b_1 + b_2)\beta_n + \gamma_{n-1}] - \beta_n(\beta_{n-1} + \beta_{n-2}) - 2\beta_{n-1}\beta_{n-2} - \beta_{n-1}^2 - \beta_{n-2}^2 \\ &\quad + \gamma_n [b_1 b_2 + (b_1 + b_2)(\beta_n + \beta_{n-1} - n + 1) - n + \beta_{n-1} + 2\beta_n + b_1 + b_2] \\ &\quad - \gamma_{n+1} [(b_1 + b_2)(\beta_n + \beta_{n+1} - n) + b_1 b_2 + n - (\beta_{n+1} + 2\beta_n + b_1 + b_2)], \\ E_n &= n\gamma_{n+1} (\beta_n + \beta_{n+1}) + \gamma_{n+1} (\beta_{n-1} + \beta_{n-2}) - \gamma_n \beta_{n-2} - (n-1)\gamma_n (\beta_{n-1} + \beta_n), \\ F_n &= \gamma_{n+1} \left(\frac{n(n-1)}{2} + \gamma_{n+1} + \gamma_{n+2} + \beta_n^2 + \beta_{n+1}^2 + \beta_n \beta_{n+1} \right) \\ &\quad - \gamma_n \left(\frac{(n-1)(n-2)}{2} + \gamma_{n-1} + \gamma_n + \beta_n^2 + \beta_{n-1}^2 + \beta_n \beta_{n-1} \right), \end{split}$$

 $G_n = (\gamma_{n+1} - \gamma_n)(a_1a_2 + a_2a_3 + a_3a_1 + (a_1 + a_2 + a_3)(\beta_n + n + \gamma_{n+1}\beta_{n+1} - \gamma_n(\beta_{n-1} - 1)).$

Theorem 8. (The Laguerre-Freud structure matrix) The Laguerre-Freud structure matrix is the following heptadiagonal matrix

where,

$$\psi_{n}^{(-2)} = \eta H_{n+2}(\beta_{+} + \beta_{n+1} + \beta_{n+2} + a_{1} + a_{2} + a_{3} + n),$$

$$\psi_{n}^{(-1)} = \eta H_{n+1}(\pi_{n-2}^{[2]} + n(\beta_{n-1} + \beta_{n} + \beta_{n+1} + a_{1} + a_{2} + a_{3}) + \gamma_{n} + \gamma_{n+1} + \gamma_{n+2} + \beta_{n+1}^{2} + \beta_{n}^{2} + \beta_{n+1}\beta_{n} + (\beta_{n} + \beta_{n+1})(a_{1} + a_{2} + a_{3}) + a_{1}a_{2} + a_{2}a_{3} + a_{3}a_{1})$$

$$\psi^{(0)} = H_n(\beta_n(\beta_n+b_1)(\beta_n+b_2) + \gamma_n(\beta_{n-1}+2\beta_n+b_1+b_2) + \gamma_{n+1}(\beta_{n+1}+2\beta_n+b_1+b_2) - n(\beta_n^2+\beta_{n-1}^2+\beta_n\beta_{n-1}+(\beta_n+\beta_{n-1})(b_1+b_2)b_1b_2 + \gamma_{n-1}+\gamma_n+\gamma_{n+1}) + \pi_{n-2}^{[-2]}(\beta_n+\beta_{n-1}+\beta_{n-2}+b_1+b_2) - \pi_{n-3}^{[-3]}),$$

or, alternatively,

$$\begin{split} \psi_n^{(0)} &= \eta H_n(\pi_{n-3}^{[3]} + \pi_{n-2}^{[2]}(\beta_{n-2} + \beta_{n-1} + \beta_n + a_1 + a_2 + a_3) + n(\beta_{n-1}^2 + \beta_n^2 + \beta_n \beta_{n-1} + a_1 a_2 + a_2 a_3 + a_3 a_1 \\ &+ (\beta_n + \beta_{n-1})(a_1 + a_2 + a_3) + \gamma_{n-1} + \gamma_n + \gamma_{n+1}) + \gamma_n(\beta_{n-1} + 2\beta_n + a_1 + a_2 + a_3) \\ &+ \gamma_{n+1}(\beta_{n+1} + 2\beta_n + a_1 + a_2 + a_3) + (\beta_n + a_1)(\beta_n + a_2)(\beta_n + a_3)), \end{split}$$
$$\psi_n^{(1)} &= H_{n+1}(\beta_{n+1}^2 + \beta_n^2 + \beta_n \beta_{n+1} + (\beta_n + \beta_{n+1})(b_1 + b_2) + b_1 b_2 + \gamma_{n+2} + \gamma_{n+1} + \gamma_n \end{split}$$

$$\varphi_{n}^{(2)} = H_{n+1}(\beta_{n+1} + \beta_n + \beta_n \beta_{n+1} + (\beta_n + \beta_{n+1})(b_1 + b_2) + b_1 b_2 + \gamma_{n+2} + \gamma_{n+1} + \gamma_n - n(\beta_{n+1} + \beta_N + \beta_{n-1} + b_1 + b_2) + \pi_{n-2}^{[-2]}),$$

$$\psi_n^{(2)} = H_{n+2}(\beta_n + \beta_{n+1} + \beta_{n+2} + b_1 + b_2 - n).$$

For the diagonal matrix $\pi^{[-2]}$ the nontrivial entries are

(52)
$$\pi_{n-2}^{[-2]} = \frac{\eta A(\beta_{n-1}, \beta_n, \beta_{n+1}, \beta_{n+2}, \gamma_n, \gamma_{n+1}, \gamma_{n+2}) + B(\beta_{n-1}, \beta_n, \beta_{n+1}, \beta_{n+2}, \gamma_n, \gamma_{n+1}, \gamma_{n+2})}{C(\beta_n, \beta_{n+1}, \eta)}$$

and, the sub-leading coefficient of the corresponding hypergeometrical discrete orthogonal polynomials is

$$p_{n-2}^{1} = \frac{\eta A(\beta_{n-1}, \beta_{n}, \beta_{n+1}, \beta_{n+2}, \gamma_{n}, \gamma_{n+1}, \gamma_{n+2}) + B(\beta_{n-1}, \beta_{n}, \beta_{n+1}, \beta_{n+2}, \gamma_{n}, \gamma_{n+1}, \gamma_{n+2})}{C(\beta_{n}, \beta_{n+1}, \eta)} + \beta_{n-2} - (n-1)\beta_{n-1} - \frac{(n-3)(n-4)}{2},$$

and the
$$\pi^{[-3]}$$
 matrix entries are
(53)

$$\pi_{n-3}^{[-3]} = p_{n-2}^{1}[(\eta+1)(\gamma_{n}-\gamma_{n+1})+\beta_{n}+\beta_{n-1}+\beta_{n-2}+b_{1}+b_{2}] + D_{n}(\beta_{n-2},\beta_{n-1},\beta_{n},\beta_{n+1},\gamma_{n-1},\gamma_{n},\gamma_{n+1}) + (\eta+1)E_{n}(\beta_{n-2},\beta_{n-1},\beta_{n},\beta_{n+1},\gamma_{n},\gamma_{n+1}) + (\eta-1)F_{n}(\beta_{n-1},\beta_{n},\beta_{n+1},\gamma_{n-1},\gamma_{n},\gamma_{n+1},\gamma_{n+2}) + \eta G_{n}(\beta_{n-1},\beta_{n},\beta_{n+1},\gamma_{n},\gamma_{n+1})$$

Proof. Matrix elements of $\psi^{(-3)}, \psi^{(-2)}, \psi^{(-1)}, \psi^{(0)}$ are obtained using $\Psi = \sigma(J)H\Pi^{\top}$. On the other hand, the entries of $\psi^{(3)}, \psi^{(2)}, \psi^{(1)}$ and the alternative expression of $\psi^{(0)}$ are gotten from $\Psi = \Pi^{-1}H\theta(J^{\top})$. The matrix $\pi^{[-2]}$ entries follow from the equation from the first subdiagonal obtained from $[\Psi H^{-1}, \Lambda^{\top} \gamma] = \vartheta_{\eta}(\Psi H^{-1})$. Applying ϑ_{η} to $\psi^{(0)}H^{-1}$, where $\psi^{(0)}$ is the first expression for the main diagonal, and equating to $\psi^{(-1)}H^{-1}$, after cleaning it is obtained $\pi^{[-2]}$. From $\pi^{[-2]}$ using $p_{n-2}^1 = \pi_{n-2}^{[-2]} - \frac{n(n-1)}{2} - (n-1)\beta_{n-1} + \beta_{n-2}$ it is obtained p_{n-2}^1 . Finally, the matrix $\pi^{[-3]}$ is obtained from the main diagonal of $[\Psi H^{-1}, J] = \Psi H^{-1}$.

Theorem 9. (Compatibility conditions) The following relations are satisfied

(54)
$$\pi_{n-1}^{[2]} - \pi_{n-2}^{[2]} = n(1 - \beta_n + \beta_{n-1})$$

(55)
$$\pi_{n-2}^{[3]} - \pi_{n-3}^{[3]} = \pi_{n-2}^{[2]}(\beta_{n-2} - \beta_n + 1) + n\gamma_{n-1} - (n-1)\gamma_n$$

(56)
$$\vartheta_{\eta} \pi_{n-2}^{[-2]} = (n-1)\gamma_n - n\gamma_{n-1}$$

(57)
$$\vartheta_{\eta} \pi_{n-3}^{[-3]} = (\gamma_n - \gamma_{n-2}) \pi_{n-2}^{[-2]} + (n-1)(\beta_{n-2} - \beta_{n-1} - 1)\gamma_n$$

Proof. Equation (54) is obtained from the second subdiagonal and the second superdiagonal of the resulting matrix of $[\Psi H^{-1}, J] = \Psi H^{-1}$, it is also obtained from the second subdiagonal of $[\Psi H^{-1}, \Lambda^{\top} \gamma] = \vartheta_{\eta}(\Psi H^{-1})$. Equation (55) is obtained from the first superdiagonal and from the first subdiagonal of $[\Psi H^{-1}, J] = \Psi H^{-1}$, it is also obtained from the first subdiagonal of $[\Psi H^{-1}, \Lambda^{\top} \gamma] = \vartheta_{\eta}(\Psi H^{-1})$ using the alternative expression of $\psi^{(0)}$. Then, (56) is a derived from the first superdiagonal of $[\Psi H^{-1}, \Lambda^{\top} \gamma] = \vartheta_{\eta}(\Psi H^{-1})$ and Equation (57) is gotten from the main diagonal of $[\Psi H^{-1}, \Lambda^{\top} \gamma] = \vartheta_{\eta}(\Psi H^{-1})$. \Box

Remark 1. i) The first three equations are equivalent to

$$\pi_{n-2}^{[-2]} - \pi_{n-1}^{[-2]} = n(\beta_{n-1} - \beta_n - 1),$$

$$\pi_{n-2}^{[-3]} - \pi_{n-3}^{[-3]} = \pi_{n-2}^{[-2]}(1 + \beta_n - \beta_{n-2}) - (n-1)\gamma_n + n\gamma_{n-1},$$

$$\vartheta_n p_{n-1}^1 = -\gamma_{n-1}.$$

- ii) Notice that compatibility conditions do not depend of hypergeometric parameters a's or b's.
- iii) Equations (52) and (58) combine and give a constraint among $(\beta_{n-1}, \beta_n, \beta_{n+1}, \beta_{n+2}, \gamma_n, \gamma_{n+1}, \gamma_{n+2})$, while Equations (53) and (59) leads to a ligature between $(\beta_{n-2}, \beta_{n-1}, \beta_n, \beta_{n+1}, \beta_{n+2}, \gamma_{n-1}, \gamma_n, \gamma_{n+1}, \gamma_{n+2})$. With these two equations at hand two Laguerre-Freud equations of the type (28) can be derived.

CONCLUSIONS AND OUTLOOK

Adler and van Moerbeke have throughly used the Gauss–Borel factorization of the moment matrix, see [1, 2, 3], In their studies of integrable systems and orthogonal polynomials. We have applied this ideas in different contexts, CMV orthogonal polynomials, matrix orthogonal polynomials, multiple orthogonal polynomials and multivariate orthogonal, see [5, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14]. For a general overview see [39].

Recently [42] we we applied that approach to study the consequences of the Pearson equation on the moment matrix and Jacobi matrices. For that description a new banded matrix is required, the Laguerre–Freud structure matrix that encodes the Laguerre–Freud relations for the recurrence coefficients. We have also found that the contiguous relations fulfilled generalized hypergeometric functions determining the moments of the weight described for the squared norms of the orthogonal polynomials a discrete Toda hierarchy known as Nijhoff–Capel equation, see [43]. In [40] we studied the role of Christoffel and Geronimus-Christoffel transformations to characterize the shifts in the spectral independent variable of the orthogonal polynomials. In [26] we deepened in that program and searched further for the discrete semi-classical

cases, finding Laguerre–Freud relations for the recursion coefficients for three types of discrete orthogonal polynomials of generalized Charlier, generalized Meixner and generalized Hahn of type I cases. In this paper we concluded this program by getting the Laguerre–Freud structure matrices and equations for three families of hypergeometrical discrete orthogonal polynomials.

A question of general character that remains open is the possibility of explicit determination (using the methods of [26] and this paper) of Laguerre–Freud equations for arbitrary hypergeometrical families ${}_{p}F_{q}$ of discrete orthogonal polynomials. In [26] and this paper we have shown that is possible for the cases (p,q) = (0,1), (1,1), (2,1), (1,2), (2,2), (3,2). For the future, we will also extend these techniques to multiple discrete orthogonal polynomials [15] and study its relations with the transformations presented in [19] and quadrilateral lattices [21, 41].

References

- [1] Mark Adler and Pierre van Moerbeke, Vertex operator solutions to the discrete KP hierarchy, Communications in Mathematical Physics 203 (1999) 185-210
- [2] ——, Generalized orthogonal polynomials, discrete KP and Riemann–Hilbert problems, Communications in Mathematical Physics 207 (1999) 589-620.
- [3] _____, Darboux transforms on band matrices, weights and associated polynomials, International Mathematics Research Notices 18 (2001) 935-984.
- [4] Carlos Álvarez-Fernández, Ulises Fidalgo Prieto, and Manuel Mañas, Multiple orthogonal polynomials of mixed type: Gauss-Borel factorization and the multi-component 2D Toda hierarchy, Advances in Mathematics 227 (2011) 1451–1525.
- [5] Carlos Álvarez-Fernández and Manuel Mañas, Orthogonal Laurent polynomials on the unit circle, extended CMV ordering and 2D Toda type integrable hierarchies, Advances in Mathematics 240 (2013) 132-193
- [6] Carlos Álvarez-Fernández, Gerardo Ariznabarreta, Juan C. García-Ardila, Manuel Mañas, and Francisco Marcellán, Christoffel transformations for matrix orthogonal polynomials in the real line and the non-Abelian 2D Toda lattice hierarchy, International Mathematics Research Notices 2017 nº5 (2017) 1285-1341, DOI:10.1093/imrn/rnw027.
- [7] Gerardo Ariznabarreta, Juan C. García-Ardila, Manuel Mañas, and Francisco Marcellán, Matrix biorthogonal polynomials on the real line: Geronimus transformations, Bulletin of Mathematical Sciences 9 (2019) 195007 (68 pages) DOI:10.1142/S1664360719500073 (World Scientfic) or DOI:10.1007/s13373-018-0128-y3 (Springer-Verlag).
- [8] Gerardo Ariznabarreta, Juan C. García-Ardila, Manuel Mañas, and Francisco Marcellán, Non-Abelian integrable hierarchies: matrix biorthogonal polynomials and perturbations Journal of Physics A: Mathematical & Theoretical **51** (2018) 205204.
- [9] Gerardo Ariznabarreta and Manuel Mañas, Matrix orthogonal Laurent polynomials on the unit circle and Toda type integrable systems Advances in Mathematics **264** (2014) 396-463.
- [10] —, Multivariate orthogonal polynomials and integrable systems, Advances in Mathematics **302** (2016) 628–739.
- [11] _____, Christoffel transformations for multivariate orthogonal polynomials, Journal of Approximation Theory 225 (2018) 242–283.
 [12] _____, Multivariate orthogonal Laurent polynomials and integrable systems, Publications of RIMS Kyoto University 56 (2022)
- 79–195,DOI:10.4171/PRIMS/58-1-4.
- [13] —, Multivariate Toda hierarchies and biorthogonal polynomials, Physica D434 (2022) 33202, DOI:10.1016/j.physd.2022.133202.
- [14] Gerardo Ariznabarreta, Manuel Mañas, and Alfredo Toledano, CMV Biorthogonal Laurent Polynomials: Perturbations and Christoffel Formulas, Studies in Applied Mathematics 140 (2018) 333–400.
- [15] Jorge Arvesú, Jonathan Coussement, and Walter Van Assche, Some discrete multiple orthogonal polynomials, Journal of Computational and Applied Mathematics 153 (2003).
- [16] R. A. Askey and Adri B. Olde Daalhuis, *Generalized hypergeometric function* (2010), in Olver, Frank W. J.; Lozier, Daniel M.; Boisvert, Ronald F.; Clark, Charles W. (eds.), NIST Handbook of Mathematical Functions, Cambridge University Press.
- [17] Jinho Baik, Thomas Kriecherbauer, Kenneth T.-R. McLaughlin, and Peter D. Miller, *Discrete Orthogonal Polynomials*, Annals of Mathematics Studies 164, Princeton University Press, 2007.
- [18] Richard Beals and Roderick Wong, Special functions and orthogonal polynomials, Cambridge Studies in Advanced Mathematics 153, Cambridge University Press, 2016.
- [19] Amílcar Branquinho, Ana Foulquié-Moreno, and Manuel Mañas, Multiple orthogonal polynomials: Pearson equations and Christoffel formulas, Analysis and mathematical Physics 12 (2022) 129 DOI:/10.1007/s13324-022-00734-1.
- [20] Peter A. Clarkson, Recurrence coefficients for discrete orthonormal polynomials and the Painlevé equations, Journal of Physics A: Mathematical & Theoretical 46 (2013) 185205.
- [21] Adam Doliwa, Paolo Maria Santini, and Manuel Mañas, *Transformations of quadrilateral lattices*, Journal of Mathematical Physics 41 (2000) 944–990.

- [22] Diego Dominici, Laguerre-Freud equations for generalized Hahn polynomials of type I, Journal of Difference Equations and Applications 24 (2018) 916-940.
- [23] _____, Matrix factorizations and orthogonal polynomials, Random Matrices Theory Applications 9 (2020) 2040003, 33 pp.
- [24] Diego Dominici and Francisco Marcellán, Discrete semiclassical orthogonal polynomials of class one, Pacific Journal of Mathematics 268 nº2 (2012) 389-411.
- [25] ——, Discrete semiclassical orthogonal polynomials of class 2 in Orthogonal Polynomials: Current Trends and Applications, edited by E. Huertas and F. Marcellán, SEMA SIMAI Springer Series, 22 (2021) 103-169, Springer.
- [26] Itsaso Fernández-Irisarri and Manuel Mañas, Pearson Equations for Discrete Orthogonal Polynomials: II. Generalized Charlier, Meixner and Hahn of type I cases, arXiv:2107.02177 [math.CA].
- [27] Galina Filipuk and Walter Van Assche, *Recurrence coefficients of generalized Charlier polynomials and the fifth Painlevé equation*, Proceedings of American Mathematical Society **141** (2013) 551–62.
- [28] _____, Recurrence Coefficients of a New Generalization of the Meixner Polynomials, Symmetry, Integrability and Geometry: Methods and Applications (SIGMA) 7 (2011), 068, 11 pages.
- [29] _____, Discrete Orthogonal Polynomials with Hypergeometric Weights and Painlevé VI, Symmetry, Integrability and Geometry: Methods and Applications (SIGMA) 14 (2018), 088, 19 pages.
- [30] Géza Freud. On the coefficients in the recursion formulae of orthogonal polynomials, Proceedings of the Royal Irish Academy Section A 76 nº1 (1976) 1-6.
- [31] Jarmo Hietarinta, Nalini Joshi and Frank W. Nijhoff, *Discrete Systems and Integrability*, Cambridge Texts in Applied Mathematics, Cambridge University Press, 2016.
- [32] Mourad E. H.Ismail, Classical and Quantum Orthogonal Polynomails in One Variable, Encyclopedia of Mathematics and its Applications 98, Cambridge University Press, 2009.
- [33] Mourad E. H. Ismail and Walter Van Assche, Encyclopedia of Special Functions: The Askey-Bateman Project. Volume I: Univariate Orthogonal Polynomials, Edited by Mourad Ismail, Cambridge University Press, 2020.
- [34] Edmond Laguerre, Sur la réduction en fractions continues d'une fraction qui satisfait à une équation différentialle linéaire du premier ordre dont les coefficients sont rationnels. Journal de Mathématiques Pures et Appliquées 4^e série, tome 1 (1885) 135-165.
- [35] Alphonse P. Magnus, A proof of Freud's conjecture about the orthogonal polynomials related to $|x|\rho \exp(-x^{2m})$, for integer m, in "Orthogonal polynomials and applications (Bar-le-Duc, 1984)", Lecture Notes in Mathematics **1171** 362–372, Springer, 1985.
- [36] ——, On Freud's equations for exponential weights, Journal of Approximation Theory 46(1) (1986) 65–99.
- [37] —, Painlevé-type differential equations for the recurrence coefficients of semi-classical orthogonal polynomials, Journal of Computational and Applied Mathematics 57 (1995) 215–237.
- [38] ——, Freud's equations for orthogonal polynomials as discrete Painlevé equations, in "Symmetries and integrability of difference equations (Canterbury, 1996)", London Mathematical Society Lecture Note Series 255 228–243, Cambridge University Press, 1999.
- [39] Manuel Mañas, Revisiting Biorthogonal Polynomials. An LU factorization discussion in Orthogonal Polynomials: Current Trends and Applications, edited by E. Huertas and F. Marcellán, SEMA SIMAI Springer Series, 22 (2021) 273-308, Springer.
- [40] —, Pearson Equations for Discrete Orthogonal Polynomials: III. Christoffel and Geronimus transformations, Revista Real Academia Ciencias Exactas, Físicas y Naturales. Serie A-Matemáticas 116 (2022) 116, DOI:10.1007/s13398-022-01296-4.
- [41] Manuel Mañas, Adam Doliwa, and Paolo Maria Santini, Darboux transformations for multidimensional quadrilateral lattices. I, Physics Letters A 232 (1997) 99–105.
- [42] Manuel Mañas, Itsaso Fernández-Irrisarri, and Omar González-Fernández, Pearson Equations for Discrete Orthogonal Polynomials: I. Generalized Hypergeometric Functions and Toda Equations, Studies inn Applied Mathematics 148 (2022) 1141-1179, DOI:10.1111/sapm.124712.
- [43] Frank W. Nijhoff and Hans W. Capel, *The direct linearisation approach to hierarchies of integrable PDEs in 2 + 1 dimensions: I. Lattice equations and the differential-difference hierarchies.* Inverse Problems 6 (1990) 567-590.
- [44] Arthur F. Nikiforov, Sergei K. Suslov, and Vasilii B. Uvarov, *Classical Orhogonal Polynomials of a Discrete Variable*, Springer Series in Computational Physics, Springer, 1991.
- [45] Lucy Joan Slater, Generalized Hypergeometric Functions, Cambridge University Press, 1966, Cambridge.
- [46] Christophe Smet and Walter Van Assche, Orthogonal polynomials on a bi-lattice, Constructive Approximation 36 (2012) 215-242.
- [47] Walter Van Assche, Orthogonal Polynomials and Painlevé Equations, Australian Mathematical Society Lecture Series 27, Cambridge University Press, 2018.