K-THEORY OF NON-COMMUTATIVE BERNOULLI **SHIFTS**

SAYAN CHAKRABORTY, SIEGFRIED ECHTERHOFF, JULIAN KRANZ, AND SHINTARO NISHIKAWA

ABSTRACT. For a large class of C^* -algebras A, we calculate the K theory of reduced crossed products $A^{\otimes G} \rtimes_r G$ of Bernoulli shifts by groups satisfying the Baum–Connes conjecture. In particular, we give explicit formulas for finite-dimensional C^* -algebras, UHF-algebras, rotation algebras, and several other examples. As an application, we obtain a formula for the K -theory of reduced C^* -algebras of wreath products $H \wr G$ for large classes of groups H and G. Our methods use a generalization of techniques developed by the second named author together with Joachim Cuntz and Xin Li, and a trivialization theorem for finite group actions on UHF algebras developed in a companion paper by the third and fourth named authors.

CONTENTS

1. INTRODUCTION

Let G be a countable discrete group and let A be a separable unital C^* algebra. Equip the infinite tensor product $A^{\otimes G}$ with the natural Bernoulli shift action (see Section [2](#page-4-0) for the definition). The objective of this paper is

Date: October 18, 2022.

²⁰¹⁰ *Mathematics Subject Classification.* Primary 46L80, 19K35; Secondary 46L55.

Key words and phrases. Bernoulli shifts, K-theory, KK-theory, the Going–Down principle, the Baum–Connes conjecture.

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project-ID 427320536 SFB 1442 and under Germany's Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics, Geometry, Structure. The first author was also supported by DST, Government of India under the *DST-INSPIRE Faculty Scheme* with Faculty Reg. No. IFA19-MA139.

to compute the K-theory group of the reduced crossed product $A^{\otimes G} \rtimes_r G$ in as many cases as possible.

Building up on the work in [\[CEL13\]](#page-29-1), Xin Li [\[Li19\]](#page-29-2) computed this when A is a finite-dimensional C^{*}-algebra of the form $\mathbb{C} \oplus \overline{\bigoplus}_{1 \leq k \leq N} M_{p_k}$, assuming the Baum–Connes conjecture with coefficients (referred to as BCC below) for G [\[BCH94\]](#page-29-3). His motivation was to compute the K -theory of the reduced group C^* -algebra of the wreath product $H \wr G$ for an arbitrary finite group H. For $A = \mathbb{C} \oplus \bigoplus_{1 \leq k \leq N} M_{p_k}$, the K-theory groups $K_*\left(A^{\otimes G} \rtimes_r G\right)$ are computed in [\[Li19\]](#page-29-2) as

(1.1)
$$
K_*\left(A^{\otimes G} \rtimes_r G\right) \cong \bigoplus_{[F] \in G \backslash \text{FIN}} \bigoplus_{[S] \in G_F \backslash \{1, \ldots, N\}^F} K_*\left(C_r^*(G_S)\right)
$$

where FIN is the set of all the finite subsets of G equipped with the lefttranslation G-action and $G_F := \text{Stab}_G(F)$, resp. $G_S := \text{Stab}_G(S)$, are the stabilizer groups at F , resp. at S , for the action of G on FIN. Note that when G is torsion-free, [\(1.1\)](#page-1-0) is the direct sum of $K_*(C_r^*(G))$ and infinitely many copies of $K_*(\mathbb{C})$. For wreath products H \ G with respect to a finite group H, we have $C_r^*(H \wr G) \cong C_r^*(H)^{\otimes G} \rtimes_r G$. In this case, the number N in [\(1.1\)](#page-1-0) corresponds to the number of non-trivial conjugacy classes of H. Our results were also motivated in part by the paper [\[Ohh15\]](#page-30-0) by Issei Ohhashi, where he gives K-theory computations for crossed products $A^{\otimes Z} \rtimes \mathbb{Z}$ for Bernoulli shifts by the integer group Z.

Our first result computes K_* $(A^{\otimes Z} \rtimes_r G)$ for an arbitrary finite-dimensional C^* -algebra A and for an arbitrary infinite countable G-set Z with the action of G on $A^{\otimes Z}$ induced by the G-action on Z. In what follows we let FIN(Z) denote the collection of finite subsets of Z and we put $FIN^{\times}(Z) :=$ $FIN(Z) \setminus \{\emptyset\}.$ In the case $Z = G$ we shall simply write FIN and FIN[×] as above.

Theorem A (Theorem [3.6\)](#page-14-1). Let G be a discrete group satisfying BCC and let Z be a countably infinite G-set. Let $A = \bigoplus_{0 \leq j \leq N} M_{k_j}$ where k_0, \ldots, k_N $(N \geq 1)$ are positive integers with $gcd(k_0, \ldots, k_N) = n$. We have an explicit isomorphism

$$
K_{*}(A^{\otimes Z} \rtimes_{r} G) \cong \bigoplus_{[F] \in G \backslash \mathrm{FIN}(Z)} \bigoplus_{[S] \in G_F \backslash (\{1, \ldots, N\}^F)} K_{*}(C_{r}^{*}(G_S))[1/n].
$$

For $N = 0$, we have

$$
K_*\left(M_n^{\otimes Z} \rtimes_r G\right) \cong K_*(C^*_r(G))[1/n].
$$

Our strategy of the proof is as follows: First, we consider the case $n = 1$ and construct a unit-preserving KK-equivalence $\mathbb{C}^{N+1} \sim_{KK} A$ using an arithmetic argument about matrices in $SL(n, \mathbb{Z})$ with non-negative entries. By an argument borrowed from [\[Izu19,](#page-29-4) [Sza18\]](#page-30-1), this allows us to replace A by \mathbb{C}^{N+1} and apply Xin Li's formula [\(1.1\)](#page-1-0). Second, we reduce the general

case to the case $n = 1$ with the help of a trivialization theorem for finite group actions on UHF algebras developed in the companion paper [\[KN22\]](#page-29-5).

One of our main techniques is the following theorem which is inspired by the regular basis technique from [\[CEL13\]](#page-29-1).

Theorem B (Theorem [2.8,](#page-8-0) Corollary [2.10\)](#page-10-0). Let G be a discrete group satisfying BCC. Let A be a separable, unital C^* -algebra that satisfies the Universal Coefficient Theorem (UCT) and such that the unital inclusion $\iota: \mathbb{C} \to A$ induces a split-injection $K_0(\mathbb{C}) \to K_0(A)$. Then $K_*\left(A^{\otimes G} \rtimes_r G\right)$ only depends on G and the cokernel $\tilde{K}_*(A)$ of $\iota_* \colon K_*(\mathbb{C}) \to K_*(A)$. For any countable G-set Z we have

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right) \cong \bigoplus_{[F] \in G \backslash \mathrm{FIN}(Z)} K_*\left(B^{\otimes F} \rtimes_r G_F\right),
$$

where B is any C^* -algebra satisfying UCT with K-theory isomorphic to $\tilde{K}_{*}(A)^{1}$ $\tilde{K}_{*}(A)^{1}$ $\tilde{K}_{*}(A)^{1}$ and where $G_F = \text{Stab}_G(F)$. In particular, if G is torsion-free and the action of G on Z is free, we have

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \text{FIN}^{\times}(Z)} K_*\left(B^{\otimes F}\right).
$$

Note that the second formula can be computed more explicitly using the Künneth theorem. In the important case that $Z = G$ with the left translation action (or, more generally, if G acts properly on Z) the stabilizers G_F for $F \in FIN^{\times}(Z)$ are all finite and the formula becomes more explicit once we can compute K_* $(B^{\otimes H} \rtimes_r H)$ for finite groups H and for relevant building blocks for B like C, $C_0(\mathbb{R})$, the Cuntz-algebras \mathcal{O}_n , and $C_0(\mathbb{R}) \otimes \mathcal{O}_n$. For $H = \mathbb{Z}/2$, these computations have been done by Izumi [\[Izu19\]](#page-29-4). For general H , computing these K -groups is a non-trivial, indeed challenging task. For $B = C_0(\mathbb{R})$ however, they are nothing but the equivariant topological K-theory K_H^* (\mathbb{R}^H) for the H-Euclidean space \mathbb{R}^H , i.e., $\mathbb{R}^{|H|}$ with H-action induced by translation of coordinates. The groups K_H^* (\mathbb{R}^H) are quite well-studied([\[Kar02\]](#page-29-6) [\[EP09\]](#page-29-7)). Using these results, we give a more explicit formula for $A = C(S^1)$ (Example [4.2\)](#page-15-0) and for the rotation algebras (or noncommutative tori) $A = A_{\theta}$ (Example [4.4\)](#page-16-0). More explicitly, we have

Theorem C (Example [4.2\)](#page-15-0). Let G be a discrete group satisfying BCC. We have

$$
K_*\left(C(S^1)^{\otimes G} \rtimes_r G\right)
$$

\n
$$
\cong K_*(C_r^*(G)) \oplus \left(\bigoplus_{\substack{[F] \in G \backslash \mathrm{FIN}^{\times}, \\[G_F \backslash F] \text{ even}}} K_*(C_r^*(G_F))\right) \oplus \left(\bigoplus_{\substack{[F] \in G \backslash \mathrm{FIN}^{\times}, \\[G_F \backslash F] \text{ odd}}} K_{G_F}^*\left(\mathbb{R}^{G_F}\right)\right)
$$

.

¹Note that this determines B uniquely up to KK -equivalence.

As another application of Theorem [B,](#page-2-1) we obtain a formula for the Ktheory of reduced C^* -algebras of many wreath products $H \wr G$:

Theorem D. (Theorem [4.14\)](#page-20-0) Let G be a discrete group satisfying BCC and let H be a discrete group such that $C_r^*(H)$ satisfies the UCT and such that the inclusion $\mathbb{C} \to C_r^*(H)$ induces a split injection on K_0 . Then we have

$$
K_*(C_r^*(H \wr G)) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \mathrm{FIN}^{\times}} K_*\left(B^{\otimes F} \rtimes_r G_F\right),
$$

where B is any C^* -algebra satisfying the UCT with K-theory isomorphic to $\tilde{K}_*(C^*_r(H))$. In particular, if G is torsion-free, we have

$$
K_*(C_r^*(H \wr G)) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \setminus \mathrm{FIN}^{\times}} K_*\left(B^{\otimes F}\right).
$$

We note that both the UCT assumption on $C^*_{r}(H)$ and the split-injectivity of the map $K_0(\mathbb{C}) \to K_0(C^*_r(H))$ hold for every a-T-menable (in particular every amenable) group H (see Remark [4.15](#page-20-1) below).

In Section [4,](#page-14-0) we also obtain formulas for several C^* -algebras A that are not covered in Theorem [B,](#page-2-1) in particular for Cuntz algebras.

Theorem E (Corollary [4.10,](#page-18-0) Proposition [4.11\)](#page-18-1). Let G be a discrete group satisfying BCC and let $n \geq 2$. Then we have

$$
K_*\left(\mathcal{O}_{n+1}^{\otimes G}\rtimes_r G\right)[1/n]=0,
$$

where \mathcal{O}_{n+1} is the Cuntz algebra on $n+1$ generators. If G is finite and $n = p$ prime, then $K_*\left(\mathcal{O}_{p+1}^{\otimes G} \rtimes_r G\right)$ is a finitely generated p-group, that is a group of the form $\bigoplus_{1 \leq j \leq N} \mathbb{Z}/p^{k_j}\mathbb{Z}$.

In Section [5,](#page-21-0) we obtain a K-theory formula for Bernoulli shifts on unital AF-algebras in terms of colimits over the orbit category $Or_{\mathcal{FIN}}(G)$. The result also applies to more general examples, in particular to many unital ASH-algebras (see Remark [5.5\)](#page-23-0).

Theorem F (Theorem [5.4\)](#page-23-1). Let A be a unital AF-algebra, let G be an infinite discrete group satisfying BCC, let Z be a countable proper G-set, and let $S \subseteq \mathbb{Z}$ be the set of all positive integers n such that $[1_A] \in K_0(A)$ is divisible by n. Then, the natural inclusions

$$
C_r^*(H) \to C_r^*(G), \ C_r^*(H) \to A^{\otimes Z} \rtimes_r H,
$$

$$
C_r^*(G) \to A^{\otimes Z} \rtimes_r G, \ A^{\otimes Z} \rtimes_r H \to A^{\otimes Z} \rtimes_r G,
$$

induce the following pushout diagram

$$
\begin{aligned}\n\text{colim} & K_*(C_r^*(H))[S^{-1}] \longrightarrow \text{colim} & K_*\left(A^{\otimes Z} \rtimes_r H\right) \\
& \downarrow & \downarrow & \\
K_*(C_r^*(G))[S^{-1}] \longrightarrow K_*\left(A^{\otimes Z} \rtimes_r G\right).\n\end{aligned}
$$

In particular, if G is torsion-free, this pushout diagram reads

$$
\tilde{K}_*(C_r^*(G))[S^{-1}] \oplus K_*\left(A^{\otimes Z}\right)_G \cong K_*(A^{\otimes Z} \rtimes_r G),
$$

where $\tilde{K}_*(C^*_r(G))$ denotes the cokernel of the map $K_*(\mathbb{C}) \to K_*(C^*_r(G))$ induced from the unital inclusion $\mathbb{C} \hookrightarrow C_r^*(G)$.

The paper is structured as follows: In Section [2,](#page-4-0) we develop our main machinery, including Theorem [B.](#page-2-1) We apply this machinery in Section [3](#page-11-0) to prove Theorem [A.](#page-1-1) In Section [4,](#page-14-0) we compute the K -theory of many more examples, including Theorems [C,](#page-2-2) [D,](#page-3-0) and [E.](#page-3-1) Bernoulli shifts on unital AFalgebras are investigated in Section [5](#page-21-0) where we prove Theorem [F.](#page-3-2) Based on similar ideas, we obtain in Section [6](#page-26-0) some very general K -theory formulas up to inverting an integer k or up to tensoring with the rationals $\mathbb Q$ (see Theorem [6.3](#page-27-0) and Theorem [6.4\)](#page-27-1). The rational K-theory computations apply to all unital stably finite C^* -algebras satisfying the UCT.

Acknowledgements. The authors would like to thank Arthur Bartels and Eusebio Gardella for helpful discussions and N. Christopher Phillips for pointing out Corollary [6.6](#page-28-0) to them.

2. General strategy

To avoid technical complications with KK -theory, we assume throughout that all C^* -algebras are separable except for the algebra $\mathcal{B}(H)$ of bounded operators on a Hilbert space H . For a C^* -algebra A and a finite set F , we write $A^{\otimes F}$ to denote the minimal tensor product $\otimes_{x\in F}A$. If A is moreover unital and Z is a (not necessarily finite) countable set, we denote by $A^{\otimes Z}$ the filtered colimit^{[2](#page-4-1)} $\underline{\text{colim}} A^{\otimes F}$ taken over all finite subsets $F \subseteq Z$ with respect to the connecting maps $A^{\otimes F} \ni x \mapsto x \otimes 1 \in A^{\otimes F} \otimes A^{\otimes F'-F} \cong A^{\otimes F'}$, for finite sets F, F' with $F \subseteq F'$. Hence $A^{\otimes Z}$ is the closed linear span of the elementary tensors $\bigotimes_{z\in Z}a_z$, where $a_z\in A$ for all $z\in Z$, and $a_z=1$ for all but finitely many z. If G is a discrete group acting on Z , we call the G-action on $A^{\otimes Z}$ given by permutation of the tensor factors the *Bernoulli shift.* More explicitly, $g\left(\bigotimes_{z\in Z} a_s\right)$ \setminus $=\bigotimes_{z\in Z}a_{g^{-1}z}$ for an elementary tensor $\bigotimes_{z\in Z}a_z\in A^{\otimes Z}.$

²We use the term 'filtered colimit' (which is the standard categorical notion) instead of the terms 'direct limit' or 'inductive limit' which seem to be more commonly used in C^* -algebra theory. We do this to be consistent with the the use of more general colimits of functors in Section [5.](#page-21-0)

For a countable discrete group G and a $G-C^*$ -algebra A , the $Baum$ Connes conjecture with coefficients (BCC) predicts a formula for the Ktheory $K_*(A \rtimes_r G)$ of the reduced crossed product $A \rtimes_r G$, see [\[BCH94\]](#page-29-3). The precise formulation of the conjecture is not too important for us. We mostly need one of its consequences recalled in Theorem [2.3](#page-5-0) below. Note that the Baum–Connes conjecture with coefficients has been verified for many groups, including all a-T-menable groups [\[HK01\]](#page-29-8) and all hyperbolic groups [\[Laf12\]](#page-29-9).

We refer the reader to [\[Kas88\]](#page-29-10) for the definition and basic properties of the equivariant KK-groups $KK^G(A, B)$. Recall from [\[MN06\]](#page-30-2) that KK^G can be organized into a triangulated category^{[3](#page-5-1)} with $G-C^*$ -algebras as objects, the groups $KK^G(A, B)$ as morphism sets, and composition given by the Kasparov product. The construction of KK^G -elements from G - $*$ homomorphisms may be interpreted as a functor from the category of G- C^* -algebras to the category KK^G . Furthermore, $A \mapsto K_*(A \rtimes_r G)$ factorizes through this functor.

Definition 2.1. A morphism ϕ in $KK^G(A, B)$ is a weak K-equivalence in KK^G if its restrictions to $KK^H(A, B)$ induce isomorphisms

$$
K_*(A \rtimes_r H) \cong K_*(B \rtimes_r H)
$$

for all finite subgroups H of G .

Remark 2.2. Weak K-equivalences are in general weaker than weak equivalences in KK^G in the sense of [\[MN06\]](#page-30-2). The latter ones are required to induce a KK^H -equivalence for all finite subgroups H of G .

The following theorem has been shown in [\[CEOO04\]](#page-29-11) in the setting of locally compact groups (see also [\[MN06\]](#page-30-2)). A detailed proof in the (easier) discrete case is given in [\[CELY17,](#page-29-12) Section 3.5].

Theorem 2.3. Suppose that the Baum–Connes conjecture holds for G with coefficients in A and B. Then, any weak K-equivalence ϕ in $KK^G(A, B)$ induces an isomorphism

$$
K_*(A \rtimes_r G) \cong K_*(B \rtimes_r G).
$$

Let A be a unital C^* -algebra and let G be a countable group satisfying BCC. Our general strategy to compute $K_*(A^{\otimes Z} \rtimes_r G)$ is to replace $A^{\otimes Z}$ by a weakly K -equivalent $G-C^*$ -algebra with computable K -theory and then apply Theorem [2.3](#page-5-0) above.

The following lemma and corollary are straightforward generalizations of [\[Izu19,](#page-29-4) Theorem 2.1] and [\[Sza18,](#page-30-1) Corollary 6.9].

Lemma 2.4 (see [\[Izu19,](#page-29-4) Theorem 2.1]). Let A and B be not necessarily unital C^* -algebras, let G be a countable discrete group, and let F be a finite G-set. Then there is a map from $KK(A, B)$ to $KK^G(A^{\otimes F}, B^{\otimes F})$ which

³Recently, KK^G was even refined to a stable ∞ -category [\[BEL21\]](#page-29-13).

sends the class of a $*$ -homomorphism $\phi: A \rightarrow B$ to the class of $\phi^{\otimes F}$. Furthermore, this map is compatible with compositions (i.e. Kasparov products) and in particular sends KK -equivalences to a KK^G -equivalences. In particular, the Bernoulli shifts on $\hat{A}^{\otimes F}$ and $B^{\otimes F}$ are KK^G -equivalent if \hat{A} and \hat{B} are KK -equivalent. The analogous statement holds if we replace the minimal tensor product \otimes by the maximal one.

Proof. We recall the description of KK^G in terms of asymptotic morphisms which admit an equivariant c.c.p. lift [\[Tho99\]](#page-30-3) (see also Appendix in [\[KS03\]](#page-29-14)). For any $G-C^*$ -algebras A_0 and B_0 , let $[[A_0, B_0]]_{cp}^G$ be the set of homotopy equivalence classes of completely positive equivariant asymptotic homomorphisms from A_0 to B_0 , see [\[Tho99,](#page-30-3) Section 2]. The usual composition law for asymptotic homomorphisms restricts to $[[A_0, B_0]]_{\text{cp}}^G$, [\[Tho99,](#page-30-3) Theorem 2]. Let S be the suspension functor $SA_0 = C_0(\mathbb{R}) \otimes A_0$. Denote by \tilde{K}_G the C^{*}-algebra of compact operators on $\ell^2(G\times\mathbb{N})$. It is a $G-C^*$ algebra with respect to the regular representation on $\ell^2(G)$. The assignment $(A_0, B_0) \mapsto \widetilde{KK}^G(A_0, B_0) = [[SA_0 \otimes \widetilde{\mathcal{K}}_G, SB_0 \otimes \widetilde{\mathcal{K}}_G]]^G_{\mathrm{cp}}, \ (\phi: A_0 \to$ $B_0 \rightarrow S\phi \otimes id_{\tilde{\mathcal{K}}_G}$ defines a bifunctor from $G-C^*$ -algebras to abelian groups. Thomsen showed that there is a natural isomorphism $\widetilde{KK}^G(A_0, B_0) \cong$ $KK^G(A_0, B_0)$ of bifunctors that sends the composition product of asymptotic morphisms to the Kasparov product, see [\[Tho99,](#page-30-3) Theorem 4.8]. Now, given $\phi \in KK(A, B)$, we represent ϕ as a completely positive asymptotic homomorphism from $SA \otimes \mathcal{K}(\ell^2(\mathbb{N}))$ to $SB \otimes \mathcal{K}(\ell^2(\mathbb{N}))$. Taking the (pointwise) minimal tensor product of ϕ with itself over F, we obtain a completely positive equivariant asymptotic homomorphism $\phi^{\otimes F}$ from $(SA \otimes$ $\mathcal{K}(\ell^2(\mathbb{N})))^{\otimes F}$ to $(SB\otimes\mathcal{K}(\ell^2(\mathbb{N})))^{\otimes F}$. This construction clearly respects homotopy equivalences. Therefore, $[\phi] \mapsto [\phi^{\otimes F}]$ defines a map $KK(A, B) \rightarrow$ $KK^G((SA\otimes \mathcal{K}(\ell^2(\mathbb{N})))^{\otimes F},(SB\otimes \mathcal{K}(\ell^2(\mathbb{N})))^{\otimes F})$. By construction, this map is compatible with compositions and sends a $*$ -homomorphism $\phi: A \rightarrow B$ to $(S\phi \otimes id_{\mathcal{K}(\ell^2(\mathbb{N}))})^{\otimes F}$. By the stabilization theorem and by Kasparov's Bott-periodicity theorem^{[4](#page-6-0)} (see [\[Bla98,](#page-29-15) Theorem 20.3.2]), the exterior tensor product by the identity on $C_0(\mathbb{R})^{\otimes F} \otimes \mathcal{K}(\ell^2(\mathbb{N}))^{\otimes F}$ induces a natural isomorphism

$$
KK^G(A_0, B_0)
$$

\n
$$
\cong KK^G(A_0 \otimes C_0(\mathbb{R})^{\otimes F} \otimes \mathcal{K}(\ell^2(\mathbb{N}))^{\otimes F}, B_0 \otimes C_0(\mathbb{R})^{\otimes F} \otimes \mathcal{K}(\ell^2(\mathbb{N}))^{\otimes F})
$$

for all $G-C^*$ -algebras A_0 and B_0 . Therefore, the map $[\phi] \mapsto [\phi^{\otimes F}]$ can be naturally regarded as a map from $KK(A, B)$ to $KK^G(A^{\otimes F}, B^{\otimes F})$. By construction, this map is compatible with compositions and sends the class of a *-homomorphism $\phi: A \to B$ to the class of $\phi^{\otimes F}$. The case of the maximal tensor product is proven in the exact same way. \Box

⁴Since the G-action on F factors through the finite group $Sym(F)$, the theorem is applicable even if G itself is not finite.

We emphasize that the map $\phi \mapsto \phi^{\otimes F}$ is not a group homomorphism. For example, it maps n in $KK(\mathbb{C}, \mathbb{C}) \cong \mathbb{Z}$ to the class $[\pi_n]$ in $KK^G(\mathbb{C}, \mathbb{C})$ of the finite dimensional unitary representation π_n of G on $\ell^2(\{1,\ldots,n\}^F)$ defined by permutation on the set $\{1, \ldots, n\}^F$.

Corollary 2.5 (see [\[Sza18,](#page-30-1) Corollary 6.9]). Let G be a discrete group and let $\phi: A \rightarrow B$ be a unital *-homomorphism which is a KK-equivalence. Then for any countable G-set Z, the map

$$
\phi^{\otimes Z}\colon A^{\otimes Z}\to B^{\otimes Z}
$$

is a weak K-equivalence in KK^G . In particular, $\phi^{\otimes Z}$ induces an isomorphism

$$
K_*(A^{\otimes Z} \rtimes_r G) \to K_*(B^{\otimes Z} \rtimes_r G)
$$

whenever G satisfies BCC.

Proof. For the first statement we may assume that G is finite. Since $K(-\rtimes_r$ G) commutes with filtered colimits, we may as well assume that Z is finite and apply Lemma [2.4.](#page-5-2) The second statement follows from Theorem [2.3.](#page-5-0) \Box

Definition 2.6. Let G be a discrete group, let Z be a countable G-set and let A_0 and B be C^* -algebras with A_0 unital. We define a $G-C^*$ -algebra $\mathcal{J}_{A_0,B}^{Z}$ as

(2.1)
$$
\mathcal{J}_{A_0,B}^Z \coloneqq \bigoplus_{F \in \text{FIN}(Z)} A_0^{\otimes Z - F} \otimes B^{\otimes F}.
$$

The G-action is defined so that a group element $g \in G$ sends $A_0^{\otimes Z-F} \otimes B^{\otimes F}$ to $A_0^{\otimes Z-gF} \otimes B^{\otimes gF}$ by the obvious *-homomorphism.

In the above definition, A_0 should be thought of as either the complex numbers $A_0 = \mathbb{C}$ or a UHF-algebra (see Section [6\)](#page-26-0). For $A_0 = \mathbb{C}$, we just write $\mathcal{J}_B^Z \coloneqq \mathcal{J}_{\mathbb{C},B}^Z$. Note that $\mathcal{J}_{A_0,B}^Z$ is a $G\text{-}C_0(\text{FIN}(Z))$ -algebra in a natural way. The following lemma is crucial for our computations:

Lemma 2.7. Let G, Z, A_0 and B be as above. Then $\mathcal{J}_{A_0,B}^Z$ can naturally be identified with the filtered colimit $\underbrace{\text{colim}}_{S} A_0^{\otimes Z-S} \otimes (A_0 \oplus B)^{\otimes S}$ taken over all finite subsets $S \subseteq Z$, with respect to the obvious connecting maps.^{[5](#page-7-0)}

Proof. Just observe that for all finite subsets $S \subseteq Z$ we have canonical isomorphisms

$$
A_0^{\otimes Z-S}\otimes (A_0\oplus B)^{\otimes S}\cong A_0^{\otimes Z-S}\otimes \left(\bigoplus_{F\subseteq S}A_0^{\otimes S-F}\otimes B^{\otimes F}\right)\cong \bigoplus_{F\subseteq S}A_0^{\otimes Z-F}\otimes B^{\otimes F}.
$$

⁵If $S \subseteq S'$ the connecting map is the tensor product of the identity on $(A_0 \oplus B)^{\otimes S}$ with the canonical inclusion $A_0^{\otimes Z-S} = A_0^{\otimes Z-S'} \otimes A_0^{\otimes S'-S} \hookrightarrow A_0^{\otimes Z-S'} \otimes (A_0 \oplus B)^{\otimes S'-S}.$

Theorem 2.8 (Theorem [B\)](#page-2-1). Let A, A_0 and B be C^* -algebras with A and A_0 unital and let $\iota: A_0 \to A$ be a unital \ast -homomorphism. Let $\phi \in KK(B, A)$ be an element such that $\iota \oplus \phi \in KK(A_0 \oplus B, A)$ is a KK-equivalence. Then for each countable G-set Z, there is a weak K-equivalence in $KK^G(\mathcal{J}_{A_0,B}^Z,A^{\otimes Z})$. If G moreover satisfies BCC, there is an isomorphism

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right) \cong K_*\left(\mathcal{J}_{A_0,B}^Z \rtimes_r G\right)
$$

\n
$$
\cong \bigoplus_{[F] \in G \backslash \text{FIN}(Z)} K_*\left(\left(A_0^{\otimes Z-F} \otimes B^{\otimes F}\right) \rtimes_r G_F\right),
$$

where $\text{FIN}(Z)$ denotes the set of finite subsets of Z and where G_F denotes the stabilizer of F for the action of G on $\text{FIN}(Z)$.

For the proof, we need the following well-known lemma:

Lemma 2.9. Let Z be a countable G-set and let A_Z be a $G-C_0(Z)$ -algebra. For any G-C^{*}-algebra D and any choice of representatives z for $[z] \in G \setminus Z$, there is a natural isomorphism

$$
\Psi = (\Psi_z)_{[z]} : KK^G(A_Z, D) \xrightarrow{\cong} \prod_{[z] \in G \setminus Z} KK^{G_z}(A_z, D)
$$

where G_z is the stabilizer of z and A_z is the fiber of A_z at $z \in Z$. The map Ψ is independent of the choice of representatives in the sense that for any $z \in Z$ and $g \in G$, the diagram

$$
KK^G(A_Z, D) \xrightarrow{\Psi_z} KK^{G_z}(A_z, D)
$$

\n
$$
\cong \downarrow g
$$

\n
$$
KK^{G_{gz}}(A_{gz}, D)
$$

\n
$$
KK^{G_{gz}}(A_{gz}, D)
$$

commutes where $g: KK^{G_z}(A_z, D) \to KK^{G_{gz}}(A_{qz}, D)$ is given by conjugation with g.

Proof. Recall that for a subgroup $H \subseteq G$, and an $H-C^*$ -algebra B, the induced $G-C^*$ -algebra $\text{Ind}_{H}^{G}B$ is defined as

$$
\operatorname{Ind}_{H}^{G} B := \left\{ f \in C_{b}(G, B) \middle| \begin{array}{c} h(f(s)) = f(sh^{-1}), s \in G, h \in H \\ \text{and } sH \mapsto ||f(s)|| \in C_{0}(G/H) \end{array} \right\}
$$

equipped with the left-translation G-action (see [\[CE01,](#page-29-16) Section 2] for example). Since Z is discrete, we have a natural isomorphism

(2.2)
$$
A_Z = \bigoplus_{z \in Z} A_z \cong \bigoplus_{[z] \in G \setminus Z} \text{Ind}_{G_z}^G A_z,
$$

where the last isomorphism follows from [\[CELY17,](#page-29-12) Theorem 3.4.13]. Now let

$$
\Psi = (\Psi_z)_{[z]} \colon KK^G(A_Z, D) \to \prod_{[z] \in G \setminus Z} KK^{G_z}(A_z, D)
$$

be the map with components

$$
\Psi_z: KK^G(A_Z, D) \xrightarrow{\text{Res}_G^{G_z}} KK^{G_z}(A_Z, D) \xrightarrow{\iota_z^*} KK^{G_z}(A_z, D),
$$

where $\iota_z: A_z \hookrightarrow A_z$ is the inclusion. The second part of the lemma follows from the construction of Ψ and the fact that conjugation by any $g \in G$ acts trivially on $KK^G(A_Z, D)$. We show that Ψ is an isomorphism. By [\(2.2\)](#page-8-1) and [\[Kas88,](#page-29-10) Theorem 2.9], we can identify Ψ with the product, taken over all $[z] \in G \backslash Z$, of the compression maps

$$
\text{comp}_{G_z}^G: KK^G(\text{Ind}_{G_z}^G A_z, D) \to KK^{G_z}(A_z, D).
$$

These are isomorphisms by [\[CE01,](#page-29-16) Proposition 5.14] (see also [\[MN06,](#page-30-2) (20)]). \Box

Proof of Theorem [2.8.](#page-8-0) For each orbit [F] in $G\$ FIN(Z) with stabilizer G_F , we define an element

$$
\Phi_F \in KK^{G_F}(A_0^{\otimes Z-F}\otimes B^{\otimes F},A^{\otimes Z})
$$

as the composition

$$
A_0^{\otimes Z-F}\otimes B^{\otimes F} \to A_0^{\otimes Z-F}\otimes (A_0\oplus B)^{\otimes F} \to A_0^{\otimes Z-F}\otimes A^{\otimes F} \to A^{\otimes Z}.
$$

Here the first and the third map are the obvious maps and the second map is the tensor product of the identity on $A_0^{\otimes Z-F}$ and the KK^{G_F} -equivalence $(\iota \oplus \phi)^{\otimes F}$ obtained in Lemma [2.4.](#page-5-2) When F is the empty set, we define Φ_F as the unital map $\iota^{\otimes Z} \colon A_0^{\otimes Z} \to A^{\otimes Z}$. By Lemma [2.9](#page-8-2) the family $\{\Phi_F : [F] \in$ $G\backslash{\rm FIN}(Z)\}$ defines an element $\Phi\in KK^G(J_{A_0,B}^Z,A^{\otimes Z})$ that does not depend on the choice of representatives F for each $[\check{F}]$ (since we have $\Phi_{gF} = g(\Phi_F)$ for every $g \in G$).

We show that Φ is a weak K-equivalence. By construction, for any finite subgroup H of G and for any finite H-subset S of Z, the element Φ may be restricted to

$$
\Phi_S\colon \bigoplus_{F\in{\rm FIN}(Z), F\subset S} A_0^{\otimes Z-F}\otimes B^{\otimes F} \to A_0^{\otimes Z-S}\otimes A^{\otimes S}
$$

in KK^H . Using Lemma [2.7,](#page-7-1) Φ_S can be identified with the tensor product of the identity on $A_0^{\otimes Z-S}$ and the KK^H -equivalence $(\iota \oplus \phi)^{\otimes S} \in KK^H((A_0 \oplus$ $(B)^{\otimes S}, A^{\otimes S}$ via the isomorphism

$$
(A_0 \oplus B)^{\otimes S} \cong \bigoplus_{F \in \text{FIN}(Z), F \subset S} A_0^{\otimes S - F} \otimes B^{\otimes F}.
$$

Thus each Φ_S induces an isomorphism

$$
K_*\left(\left(A_0^{\otimes Z-S}\otimes (A_0\oplus B)^{\otimes S}\right)\rtimes H\right)\cong K_*\left(\left(A_0^{\otimes Z-S}\otimes A^{\otimes S}\right)\rtimes H\right).
$$

By taking the filtered colimit over finite H-subsets $S \subseteq Z$, we see that Φ is a weak K-equivalence.

The K-theory computation follows from the fact that $\mathcal{J}_{A_0,B}^Z$ is a G- $C_0(FIN(Z))$ -algebra for the discrete G-set FIN(Z) together with Green's imprimitivity Theorem (e.g., see [\[CEL13,](#page-29-1) Remark 3.13] for details). \square

The following special case of Theorem [2.8](#page-8-0) will be used to compute examples in Section [4.](#page-14-0) Its main advantage is that we do not have to construct the element ϕ in order to apply it.

Corollary 2.10. Let G be a discrete group satisfying BCC. Let Z be a countable G-set and let A be a unital C^* -algebra satisfying the UCT such that the unital inclusion $\mathbb{C} \to A$ induces a split injection $K_*(\mathbb{C}) \to K_*(A)$. Denote by $\tilde{K}_*(A)$ its cokernel and let B be any C^* -algebra satisfying the UCT with $K_*(B) \cong \tilde{K}_*(A)$. Then we have

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right) \cong K_*({\mathcal J}_B^Z \rtimes_r G) \cong \bigoplus_{[F] \in G \backslash \mathrm{FIN}(Z)} K_*\left(B^{\otimes F} \rtimes_r G_F\right).
$$

In particular, if $Z = G$ equipped with the left translation, then $K_*\left(A^{\otimes G} \rtimes_r G\right)$ only depends on G and $\tilde{K}_{*}(A)$.

Note that B exists and is uniquely determined up to KK -equivalence (see [\[Bla98,](#page-29-15) Corollary 23.10.2]).

Proof. By the UCT, the inclusion map $K_*(B) \cong \tilde{K}_*(A) \to K_*(A)$ is induced by an element $\phi \in KK(B,A)$. By construction, $\iota \oplus \phi \in KK(\mathbb{C} \oplus B, A)$ is a KK -equivalence, so that we can apply Theorem [2.8.](#page-8-0)

As another special case of Theorem [2.8,](#page-8-0) we recover

Corollary 2.11 (CEL13, Example 3.17), [\[Li19,](#page-29-2) Proposition 2.4]). Let G be a discrete group satisfying BCC and let Z be a countable G-set. Then for $n \geq 1$, we have

$$
K_*(C(\{0,1,\ldots,n\}^Z)\rtimes_r G)\cong \bigoplus_{[F]\in G\backslash \mathrm{FIN}(Z)}\bigoplus_{[S]\in G_F\backslash (\{1,\ldots,n\}^F)}K_*(C_r^*(G_S)).
$$

Moreover, if $Z = G$ with the left translation action, we get

$$
K_*(C(\{0,1,\ldots,n\}^G) \rtimes_r G)
$$

\n
$$
\cong K_*(C_r^*(G)) \oplus \bigoplus_{[C] \in C} \bigoplus_{[X] \in N_C \backslash F(C)} \bigoplus_{[S] \in C \backslash \{1,\ldots,n\}^{C \cdot X}} K_*(C_r^*(C_S)).
$$

Here $\mathcal C$ denotes the set of all conjugacy classes of finite subgroups of G , $F(C)$ the nonempty finite subsets of $C\backslash G$, $N_C = \{g \in G : gCg^{-1} = C\}$ the normalizer of C in G, and $C_S = G_S \cap C$ the stabilizer of S in C.

Proof. Let $A = C({0, \ldots, n})$ and $B = C({1, \ldots, n})$ and let $\phi: B \to A$ be the canonical inclusion. The first isomorphism follows from Theorem [2.8.](#page-8-0) The second isomorphism is obtained by analyzing the orbit structure of $\bigsqcup_{F \in \text{FIN}(G)} \{1, \ldots, n\}^F$ (see [\[Li19,](#page-29-2) Proposition 2.4] for details).

12 S. CHAKRABORTY, S. ECHTERHOFF, J. KRANZ, AND S. NISHIKAWA

3. Finite-dimensional algebras

In this section we compute the K -theory of crossed products of the form $A^{\otimes Z} \rtimes_r G$ where $A = \bigoplus_{0 \leq j \leq N} M_{k_j}$ is a finite-dimensional C^* -algebra, Z is a countable G-set, and where G is a group satisfying BCC. This generalizes the case $k_0 = 1$ from [\[Li19\]](#page-29-2). We denote by $gcd(k_0, \ldots, k_N)$ the greatest common divisor of k_0, \ldots, k_N . We believe that the following theorem is known to experts. In lack of a reference, we give a detailed proof here.

Theorem 3.1. Let k_0, \ldots, k_N be positive integers with $gcd(k_0, \ldots, k_N) = 1$. Then there is a unital ∗-homomorphism

$$
\phi \colon \mathbb{C}^{N+1} \to \bigoplus_{0 \le j \le N} M_{k_j}
$$

that induces a KK-equivalence. Moreover when $N = 1$, there are exactly two such ∗-homomorphisms up to unitary equivalence.

The main ingredient for the proof is the following arithmetic fact:

Proposition 3.2. For any pair of positive integers $k_1, k_2 \in \mathbb{N}$, there is a unique matrix X in $SL(2, \mathbb{Z})$ with non-negative entries such that

$$
X\begin{bmatrix}n\\n\end{bmatrix}=\begin{bmatrix}k_1\\k_2\end{bmatrix}
$$

where $n = \gcd(k_1, k_2)$. If we allow X to be in $GL(2, \mathbb{Z})$, then there are exactly two such X: the one X_0 in $SL(2, \mathbb{Z})$ and X_0 $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$.

Proof. Existence: Let $f: \mathbb{N}_+^2 \to \mathbb{N}_+^2$ be given by

$$
\begin{bmatrix} k_1 \\ k_2 \end{bmatrix} \mapsto \begin{cases} \begin{bmatrix} k_1 - k_2 \\ k_2 \end{bmatrix}, & \text{if } k_1 > k_2 \\ \begin{bmatrix} k_1 \\ k_2 - k_1 \end{bmatrix}, & \text{if } k_1 < k_2 \\ \begin{bmatrix} k_1 \\ k_2 \end{bmatrix}, & \text{if } k_1 = k_2. \end{cases}
$$

The Euclidian algorithm precisely says that there exists a $k \in \mathbb{N}$ such that f^k $\begin{bmatrix} k_1 \\ k_2 \end{bmatrix}$ k_2 1 = $\lceil n \rceil$ n 1 . Let $A =$ $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $B =$ $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$. Then the map f defined above is given by

$$
\begin{bmatrix} k_1 \\ k_2 \end{bmatrix} \mapsto \begin{cases} A^{-1} \begin{bmatrix} k_1 \\ k_2 \end{bmatrix}, & \text{if } k_1 > k_2 \\ B^{-1} \begin{bmatrix} k_1 \\ k_2 \end{bmatrix}, & \text{if } k_1 < k_2 \\ \begin{bmatrix} k_1 \\ k_2 \end{bmatrix}, & \text{if } k_1 = k_2. \end{cases}
$$

Now the above formulation of the Euclidean algorithm gives us integers $a_1, a_2, \dots, a_l, b_1, b_2, \dots, b_l \ge 1$ such that

$$
A^{a_1}B^{b_1}A^{a_2}B^{b_2}\cdots A^{a_l}B^{b_l}\begin{bmatrix}n\\n\end{bmatrix}=\begin{bmatrix}k_1\\k_2\end{bmatrix}.
$$

Then $X := A^{a_1} B^{b_1} A^{a_2} B^{b_2} \cdots A^{a_l} B^{b_l} \in SL(2, \mathbb{Z})$ is the required matrix.

Uniqueness: Suppose there are two such matrices X_1 and X_2 . Then, $Y = X_2^{-1}X_1 \in SL(2, \mathbb{Z})$ satisfies

$$
Y\begin{bmatrix}1\\1\end{bmatrix}=\begin{bmatrix}1\\1\end{bmatrix}.
$$

From this, we have

$$
Y = \begin{bmatrix} a & 1 - a \\ a - 1 & 2 - a \end{bmatrix}
$$

for $a \in \mathbb{Z}$. Now, we get

$$
X_1=X_2\begin{bmatrix}a&1-a\\a-1&2-a\end{bmatrix},
$$

but since both X_1 and X_2 have non-negative entries and since they are nonsingular, it is not hard to see that a must be 1. Thus, $X_1 = X_2$. The last assertion is immediate. $\hfill \square$

Remark 3.3. It follows from the proof that the subsemigroup in $SL(2, \mathbb{Z})$ consisting of matrices with non-negative entries is the free monoid of two generators A and B.

Corollary 3.4. For any positive integers k_0, \ldots, k_N with $gcd(k_0, \ldots, k_N) =$ n, there is a matrix X in $SL(N + 1, \mathbb{Z})$ with non-negative entries such that

$$
X\begin{bmatrix} n \\ \vdots \\ n \end{bmatrix} = \begin{bmatrix} k_0 \\ \vdots \\ k_N \end{bmatrix}.
$$

Proof. We give the proof by induction on N. The case $N = 0$ is clear. Let $N \geq 1$ and let $k_0, \ldots, k_N \geq 1$ be positive integers with $gcd(k_0, \ldots, k_N) = n$. Set $l := \gcd(k_0, \ldots, k_{N-1})$. By induction, we may assume that there exists a

matrix $\tilde{X} \in SL(N, \mathbb{Z})$ with non-negative entries such that \tilde{X} $\sqrt{ }$ $\overline{}$ l . . . l 1 $\Big\} =$ $\sqrt{ }$ $\overline{}$ k_{0} . . . k_{N-1} 1 $\vert \cdot$ Since $gcd(l, k_N) = gcd(l, n) = n$, it follows from Proposition [3.2](#page-11-1) that there are matrices $X_0, X_N \in SL(2, \mathbb{Z})$ with non-negative entries such that X_0 $\lceil n \rceil$ n 1 = $\lceil l \rceil$ n 1 and X_N $\lceil n \rceil$ n 1 = \int l k_N T . Now for $i \in \{1, \ldots, N-1\}$, let Y_i denote the matrix with X_0 as the $(i, i + 1)$ -th diagonal block, with ones in all other diagonal entries and with zeros elsewhere, and let $Y_N =$ $\begin{bmatrix} I_{N-1} & 0 \end{bmatrix}$ $0\quad X_N$ 1 . Write $Y \coloneqq$ $\begin{bmatrix} \tilde{X} & 0 \\ 0 & 1 \end{bmatrix}$. Then $X := Y \cdot Y_N \cdots Y_1 \in SL(N + 1, \mathbb{Z})$ has non-negative entries and

$$
X\begin{bmatrix} n \\ \vdots \\ n \end{bmatrix} = Y\left(Y_N \cdots Y_1 \begin{bmatrix} n \\ \vdots \\ n \end{bmatrix}\right) = Y\begin{bmatrix} l \\ \vdots \\ l \\ k_N \end{bmatrix} = \begin{bmatrix} k_0 \\ \vdots \\ k_N \end{bmatrix}
$$

as desired. \Box

Proof of Theorem [3.1.](#page-11-2) Unital *-homomorphisms from \mathbb{C}^{N+1} to $\bigoplus_{0 \leq j \leq N} M_{k_j}$ are classified up to unitary equivalence by their induced maps on ordered K_0 groups with units. If we identify the K_0 -groups of \mathbb{C}^{N+1} and $\bigoplus_{0 \leq j \leq N} M_{k_j}$ with \mathbb{Z}^{N+1} , we may represent the induced maps on K_0 -groups by $(N + 1)$ square matrices with non-negative integer entries that send Ī $\overline{}$ 1 . . . 1 T to $\sqrt{ }$ $\overline{}$ k_0 . . . k_N Ť $\vert \cdot$

Such a matrix is an isomorphism on K-theory if and only if it is in $GL(N + \mathbb{R})$ $1, \mathbb{Z}$). All the assertions now follow from Proposition [3.2](#page-11-1) and Corollary $3.4.$

We are now ready to prove Theorem [A.](#page-1-1) For the proof, we need a result from [\[KN22\]](#page-29-5). We formulate it in full generality here since this will be needed in later sections of the paper. Recall that a supernatural number is a formal product $\mathfrak{n} = \prod_p p^{n_p}$ of prime powers with $n_p \in \{0, 1, \ldots, \infty\}$. We denote by $M_{\mathfrak{n}} \coloneqq \otimes_p M_p^{\otimes n_p}$ the associated UHF-algebra. $M_{\mathfrak{n}}$ and \mathfrak{n} are called of infinite type if $n_p \in \{0,\infty\}$ for all p and $n_p \neq 0$ for at least one p. For an abelian group L , we denote by

$$
L[1/\mathfrak{n}] := \mathrm{colim}(L \xrightarrow{\cdot p_1} L \xrightarrow{\cdot p_2} \cdots)
$$

the *localization at* **n** where (p_1, p_2, \dots) is a sequence of primes containing every p with $n_p \geq 1$ infinitely many times. In other words, $L[1/n]$ is the localization at the set of all primes dividing n. If n is finite, this definition recovers the usual localization at a natural number.

Theorem 3.5 ([\[KN22,](#page-29-5) Corollary 2.11]). Let G be a discrete group satisfying BCC, let Z a countable G-set, let A a G-C^{*}-algebra and let M_n a UHFalgebra. Assume that Z is infinite or that $\mathfrak n$ is of infinite type. Then the inclusion $A \to A \otimes M_n^{\otimes Z}$ induces an isomorphism

$$
K_*(A\rtimes_r G)[1/\mathfrak{n}]\cong K_*\left(\left(A\otimes M_\mathfrak{n}^{\otimes Z}\right)\rtimes_r G\right).
$$

In particular, the right-hand side is a $\mathbb{Z}[1/\mathfrak{n}]$ -module.

Theorem 3.6 (Theorem [A\)](#page-1-1). Let G be a discrete group satisfying BCC. Let Z be a countably infinite G-set and let $A = \bigoplus_{0 \leq j \leq N} M_{k_j}$ where k_0, \ldots, k_N $(N \geq 1)$ are positive integers with $gcd(k_0, \ldots, k_N) = n$. Then

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right) \cong K_*\left(C\left(\{0,\ldots,N\}^Z\right) \rtimes_r G\right)[1/n] \cong \bigoplus_{[F] \in G \backslash \text{FIN}(Z) \ S] \in G_F \backslash (\{1,\ldots,N\}^F)} K_*(C_r^*(G_S))[1/n].
$$

Proof. Write $B = \bigoplus_{0 \leq j \leq N} M_{k_j/n}$ so that B satisfies the assumptions of Theorem [3.1](#page-11-2) and so that $A \cong B \otimes M_n$. By Theorem [3.5,](#page-14-2) the inclusion $B^{\otimes Z} \hookrightarrow A^{\otimes Z}$ induces an isomorphism

$$
K_*\left(B^{\otimes Z} \rtimes_r G\right)[1/n] \cong K_*\left(A^{\otimes Z} \rtimes_r G\right).
$$

By Theorem [3.1](#page-11-2) and Corollary [2.5,](#page-7-2) we furthermore have

$$
K_*\left(B^{\otimes Z} \rtimes_r G\right) \cong K_*\left(C\left(\{0,\ldots,N\}^Z\right) \rtimes_r G\right).
$$

This proves the isomorphism in the first line of the theorem. The isomor-phism in the second line follows from Corollary [2.11.](#page-10-1)

4. More examples

In this section, we compute the K -theory of Bernoulli shifts in more examples. We mostly restrict ourselves to the case $Z = G$ with the left translation action, but some of the results have straightforward generalizations to arbitrary countable G-sets. Recall that for $Z = G$ we write $FIN = FIN(G)$ for the collection of finite subsets of G and we put $\mathcal{J}_B := \mathcal{J}^G_{\mathbb{C},B}$ as in Definition [2.6](#page-7-3) for any C^* -algebra B . We start with some easy applications of Corollary [2.5.](#page-7-2)

Example 4.1. Let A be the Fibonacci algebra [\[Dav96,](#page-29-17) Example III 2.6], which is the filtered colimit of $(M_{m_k} \oplus M_{n_k})_{k \in \mathbb{N}}$ where $m_1 = n_1 = 1$ and where the connecting maps are given by repeated use of the partial embedding matrix $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$. Since this matrix belongs to GL(2, Z), the unital embedding $\mathbb{C} \oplus \bar{\mathbb{C}} \hookrightarrow \bar{\mathcal{A}}$ is a KK-equivalence.

Let K^+ be the unitization of the algebra of compact operators on $\ell^2(\mathbb{N})$ and let $p \in \mathcal{K}$ be a rank-1 projection. Then the unital embedding $\mathbb{C} \oplus \mathbb{C} \hookrightarrow \mathcal{K}^+$ given by $(\lambda, \mu) \mapsto \lambda(1 - p) + \mu p$ is a KK-equivalence.

Now let A be either A or K^+ and let G be a discrete group satisfying BCC. Then by Corollaries [2.5](#page-7-2) and [2.11,](#page-10-1) we have

$$
K_*\left(A^{\otimes G} \rtimes_r G\right) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \mathrm{FIN}^{\times}} K_*(C_r^*(G_F)).
$$

In particular, if G is torsion free, we have

$$
K_*\left(A^{\otimes G} \rtimes_r G\right) \cong K_*(C^*_r(G)) \oplus \bigoplus_{[F] \in G \backslash \mathrm{FIN}^{\times}} K_*(\mathbb{C}).
$$

Example 4.2 (Theorem [C\)](#page-2-2). Consider $A = C(S^1)$. Note that the canonical inclusion $\phi: C_0(\mathbb{R}) \to C(S^1)$ together with the unital inclusion $\iota: \mathbb{C} \to C(S^1)$ induces a KK-equivalence $\iota \oplus \phi \in KK(\mathbb{C} \oplus C_0(\mathbb{R}), A)$. By Theorem [2.8,](#page-8-0) we obtain a weak K-equivalence $\Phi \colon \mathcal{J}_{C_0(\mathbb{R})} \to A^{\otimes G}$. We can compute the K-theory of $\mathcal{J}_{C_0(\mathbb{R})} \rtimes_r G$ as

$$
K_*\left(\mathcal{J}_{C_0(\mathbb{R})} \rtimes_r G\right) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \mathrm{FIN}^{\times}} K_*\left(C_0(\mathbb{R})^{\otimes F} \rtimes_r G_F\right).
$$

In this expression, each nonempty finite subset $F \subseteq G$ can be written as $F = G_F \cdot L_F$ where L_F is a complete set of representatives for $G_F \backslash F$. If the cardinality of L_F is even, the G_F -action on $C_0(\mathbb{R})^{\otimes F} = C_0(\mathbb{R}^F)$ is KK^{G_F} equivalent to the trivial action on C by Kasparov's Bott-periodicity theorem (see [\[Bla98,](#page-29-15) Theorem 20.3.2]). When the cardinality of L_F is odd, then

$$
C_0(\mathbb{R})^{\otimes F} \cong \left(C_0(\mathbb{R})^{\otimes G_F}\right)^{\otimes |L_F|-1} \otimes C_0(\mathbb{R})^{\otimes G_F}
$$

is KK^{G_F} -equivalent to $C_0(\mathbb{R})^{\otimes G_F} = C_0(\mathbb{R}^{G_F})$. Therefore, we have

$$
K_*\left(C_0(\mathbb{R})^{\otimes F} \rtimes_r G_F\right) \cong \begin{cases} K_*(C^*_r(G_F)), & \text{if } |G_F \backslash F| \text{ is even} \\ K_*\left(C_0\left(\mathbb{R}^{G_F}\right) \rtimes_r G_F\right), & \text{if } |G_F \backslash F| \text{ is odd} \end{cases}
$$

.

In general, for any finite group H and for any orthogonal representation $H \to O(V)$ on a finite-dimensional Euclidean space V, the K-theory of $C_0(V) \rtimes_r H$ is the well-studied equivariant topological K-theory $K_H^*(V)$ of V (see [\[Kar02\]](#page-29-6), [\[EP09\]](#page-29-7)). It is known to be a finitely generated free abelian group with $\mathrm{rank}_\mathbb{Z} K_H^*(V)$ equal to the number of conjugacy classes $\langle g \rangle$ of H which are oriented and even/odd respectively (see [\[Kar02,](#page-29-6) Theorem 1.8]). Here, a conjugacy class $\langle g \rangle$ of H is oriented if the centralizer C_g of g acts on the g-fixed points V^g of V by oriented automorphisms. The class $\langle g \rangle$ is even/odd if the dimension of V^g is even/odd respectively. For example for any cyclic group $\mathbb{Z}/m\mathbb{Z}$, using [\[EP09,](#page-29-7) Example 4.2], we have

$$
K_{\mathbb{Z}/m\mathbb{Z}}^*(\mathbb{R}^m) = \begin{cases} \mathbb{Z}^m, & * = 1 \\ 0, & * = 0 \end{cases}
$$

for odd $m \geq 1$, and

$$
K_{\mathbb{Z}/m\mathbb{Z}}^*(\mathbb{R}^m) = \begin{cases} \mathbb{Z}^{m/2}, & * = 1\\ 0, & * = 0 \end{cases}
$$

for even $m \geq 2$. We summarize our discussion as follows:

Theorem 4.3. Let G be a discrete group satisfying BCC. We have

$$
K_*\left(C(S^1)^{\otimes G} \rtimes_r G\right)
$$

$$
\cong K_*(C_r^*(G)) \oplus \left(\bigoplus_{\substack{[F] \in G \backslash \mathrm{FIN}^{\times}, \\[G_F \backslash F] \text{ even}}} K_*(C_r^*(G_F))\right) \oplus \left(\bigoplus_{\substack{[F] \in G \backslash \mathrm{FIN}^{\times}, \\[G_F \backslash F] \text{ odd}}} K_{G_F}^*\left(\mathbb{R}^{G_F}\right)\right).
$$

Example 4.4. Let A_{θ} be the rotation algebra for $\theta \in \mathbb{R}$, the universal C^* algebra generated by two unitaries u and v satisfying $uv = vue^{2\pi i\theta}$. It is well-known that $K_0(A_\theta) \cong \mathbb{Z}^2$, that $K_1(A_\theta) \cong \mathbb{Z}^2$, and that the unital inclusion $\iota: \mathbb{C} \to A_{\theta}$ induces a split injection on K_0 . Since A_{θ} satisfies the UCT, we can apply Corollary [2.10](#page-10-0) to $B = \mathbb{C} \oplus C_0(\mathbb{R}) \oplus C_0(\mathbb{R})$ and obtain

$$
K_*\left(A_{\theta}^{\otimes G} \rtimes_r G\right)
$$

\n
$$
\cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \text{FIN}^{\times}} K_*\left((\mathbb{C} \oplus C_0(\mathbb{R}) \oplus C_0(\mathbb{R}))^{\otimes F} \rtimes_r G_F\right).
$$

Using Theorem [2.8](#page-8-0) for $B = C_0(\mathbb{R}) \oplus C_0(\mathbb{R})$, each summand for $F \in FIN^{\times}$ may be computed as

$$
K_*\left((\mathbb{C}\oplus C_0(\mathbb{R})\oplus C_0(\mathbb{R}))^{\otimes F}\rtimes_r G_F\right)
$$

\n
$$
\cong \bigoplus_{[X_1,X_2]} K_*\left(\left(C_0\left(\mathbb{R}^{X_1}\right)\otimes C_0\left(\mathbb{R}^{X_2}\right)\right)\rtimes_r G_{X_{1,2}}\right),
$$

where the sum is taken over all ordered equivalence classes $[X_1, X_2]$ of pairs of ordered disjoint subsets X_1, X_2 of F modulo the action of the stabilizer G_F of F, and where we set $G_{X_{1,2}} := G_F \cap G_{X_1} \cap G_{X_2}$. As in Example [4.2,](#page-15-0) the computation of $K_*\left((C_0 \left(\mathbb{R}^{X_1} \right) \otimes C_0 \left(\mathbb{R}^{X_2} \right) \right) \rtimes_r G_{X_{1,2}} \right)$ either reduces to $K_*(C_r^*(G_{X_{1,2}}))$ or to that of $K_*\left(C_0\left(\mathbb{R}^{G_{X_{1,2}}}\right) \times_r G_{X_{1,2}}\right) = K_{G_{X_{1,2}}}^*$ $\left(\mathbb{R}^{G_{X_{1,2}}}\right)$ in general. We summarize the discussion as follows.

Theorem 4.5. Let G be a discrete group satisfying BCC and let $\theta \in \mathbb{R}$. We have

$$
K_*\left(A_{\theta}^{\otimes G} \rtimes_r G\right) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \text{FIN}^{\times}} K_F
$$

where for all $F \in \text{FIN}^{\times}$ we set

$$
K_F \coloneqq \left(\bigoplus_{\substack{[X_1,X_2],X_1\sqcup X_2\subset F,\\ |G_{X_{1,2}}\backslash (X_1\sqcup X_2)|\text{ even}}} K_*(C^*_r(G_{X_{1,2}})) \right) \oplus \left(\bigoplus_{\substack{[X_1,X_2],X_1\sqcup X_2\subset F,\\ |G_{X_{1,2}}\backslash (X_1\sqcup X_2)|\text{ odd}}} K^*_{G_{X_{1,2}}} \left(\mathbb{R}^{G_{X_{1,2}}}\right) \right)
$$

.

Cuntz algebras. For $n \in \{2, 3, ..., \infty\}$ we denote by \mathcal{O}_n the Cuntz algebra on n generators.

Example 4.6. [\[Sza18,](#page-30-1) Corollary 6.9] Let G be a discrete group satisfying BCC. Then the unital inclusion $\mathbb{C} \to \mathcal{O}_{\infty}$ induces an isomorphism

$$
K_* (C_r^* (G)) \cong K_* \left(\mathcal{O}_\infty^{\otimes G} \rtimes_r G \right).
$$

Proof. Combine Corollary [2.5](#page-7-2) and the fact that the unital inclusion $\mathbb{C} \to \mathcal{O}_{\infty}$ is a KK-equivalence.

Example 4.7. Let G be a discrete group satisfying BCC. Then we have

$$
K_*\left(\mathcal{O}_2^{\otimes G} \rtimes_r G\right) = 0.
$$

Proof. Combine Lemma [2.4,](#page-5-2) Theorem [2.3](#page-5-0) and the fact that \mathcal{O}_2 is KKequivalent to 0. \Box

Example 4.8. Let G be a discrete group satisfying BCC. Let $A = \mathbb{C} \oplus \mathcal{O}_n$ for $n \geq 3$. By Theorem [2.8,](#page-8-0) we have

$$
K_*\left(A^{\otimes G} \rtimes_r G\right) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \mathrm{FIN}^\times} K_*\left(\mathcal{O}_n^{\otimes F} \rtimes_r G_F\right).
$$

Each summand for $F = G_F \cdot L_F$ becomes $K_* \left((\mathcal{O}_n^{\otimes L_F})^{\otimes G_F} \rtimes_r G_F \right)$. It follows from the UCT and an inductive application of the Künneth theorem that $\mathcal{O}_n^{\otimes L_F}$ is KK-equivalent to $\mathcal{O}_n \otimes (C_0(\mathbb{R}) \oplus \mathbb{C})^{|L_F|-1}$. Thus, by Lemma [2.4,](#page-5-2) we may express $K_*(\mathcal{O}_n^{\otimes F} \rtimes_r G_F)$ explicitly in terms of

$$
K_*(\mathcal{O}_n^{\otimes H} \rtimes_r H)
$$
 and $K_*((C_0(\mathbb{R}) \otimes \mathcal{O}_n)^{\otimes H} \rtimes_r H)$

for finite subgroups H of G_F . These groups for $H = \mathbb{Z}/2$ are nicely computed in [\[Izu19\]](#page-29-4). According to [\[Izu19\]](#page-29-4), we have

$$
K_*\left(\mathcal{O}_{n+1}^{\otimes \mathbb{Z}/2} \rtimes_r \mathbb{Z}/2\right) = \begin{cases} \mathbb{Z}/n \oplus \mathbb{Z}/n, & * = 0\\ 0, & * = 1 \end{cases}
$$

for odd n ,

$$
K_*\left(\mathcal{O}_{n+1}^{\otimes \mathbb{Z}/2} \rtimes_r \mathbb{Z}/2\right) = \begin{cases} \mathbb{Z}/\frac{n}{2} \oplus \mathbb{Z}/2n, & * = 0, \\ 0 & * = 1 \end{cases}
$$

for even n , and

$$
K_*\left((C_0(\mathbb{R}) \otimes \mathcal{O}_{n+1})^{\otimes \mathbb{Z}/2} \rtimes_r \mathbb{Z}/2\right) = \begin{cases} 0, & * = 0 \\ \mathbb{Z}/n \oplus \mathbb{Z}/n, & * = 1 \end{cases}
$$

for odd n,

$$
K_*\left((C_0(\mathbb{R}) \otimes \mathcal{O}_{n+1})^{\otimes \mathbb{Z}/2} \rtimes_r \mathbb{Z}/2\right) = \begin{cases} 0, & * = 0\\ \mathbb{Z}/\frac{n}{2} \oplus \mathbb{Z}/2n, & * = 1 \end{cases}
$$

for even *n*. Using similar methods as in $[Iau19]$, we can compute the case $H = \mathbb{Z}/3$. We omit the very technical computations. For general H, the computations become increasingly complicated as the order of H increases.

Question 4.9. Is $K_*\left(\mathcal{O}_n^{\otimes H} \rtimes_r H\right)$ computable for all finite groups H or at least for all cyclic groups?

Although we do not know how to compute $K_*\left(\mathcal{O}_n^{\otimes H} \rtimes_r H\right)$ in general, we can say something about its structure. The following Corollary is a combination of Theorem [3.5](#page-14-2) and Corollary [2.5.](#page-7-2)

Corollary 4.10. Let G be a discrete group satisfying BCC, Z a countable G set, and A a C^* -algebra. Let M_n be a UHF-algebra of infinite type such that $A \otimes M_{\mathfrak{n}}$ is KK-equivalent to zero (for instance $A = \mathcal{O}_{n+1}$ and $M_{\mathfrak{n}} = M_{n} \otimes$ for some $n \geq 2$). Then

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right)[1/\mathfrak{n}] \cong 0.
$$

For $A = \mathcal{O}_{n+1}$ and a finite group H, we can say a bit more:

Proposition 4.11. Let H be a finite group, let Z be a finite H -set and let $n \geq 2$. Then $K_*\left(\mathcal{O}_{n+1}^{\otimes Z} \rtimes_r H\right)$ is a finitely generated abelian group L such that $L[1/n] = 0$. That is, any element in L is annihilated by n^k for some k. In particular, if $n = p$ is prime, then L is isomorphic to the direct sum of finitely many p-groups $\mathbb{Z}/p^k\mathbb{Z}$.

Proof. The method used in [\[Izu19\]](#page-29-4), at an abstract level, tells us that $K_*\left(\mathcal{O}_{n+1}^{\otimes Z} \rtimes_r H\right)$ is finitely generated. To see this, let \mathcal{T}_{n+1} be the universal C^* -algebra generated by isometries $s_1 \cdots, s_{n+1}$ with mutually orthogonal range projections $q_i = s_i s_i^*$ and we let $p = 1 - \sum_{1 \leq j \leq n+1} q_i$. The ideal generated by p is isomorphic to $\mathcal K$ and we have the following exact sequences

(4.1)
$$
0 \longrightarrow I_1 \rtimes H \longrightarrow \mathcal{T}_{n+1}^{\otimes Z} \rtimes H \longrightarrow \mathcal{O}_{n+1}^{\otimes Z} \rtimes H \longrightarrow 0,
$$

$$
(4.2) \qquad 0 \longrightarrow I_{m+1} \rtimes H \longrightarrow I_m \rtimes H \longrightarrow (I_m/I_{m+1}) \rtimes H \longrightarrow 0,
$$

where for $1 \leq m \leq |Z|$, an H-ideal I_m of $\mathcal{T}_{n+1}^{\otimes Z}$ is defined as the ideal generated by $\mathcal{K}^{\otimes F} \otimes \mathcal{T}_{n+1}^{\otimes Z-F}$ for subsets F of H with $|F| = m$. In particular $I_{|Z|} =$ $\mathcal{K}^{\otimes Z}$. By Lemma [2.4,](#page-5-2) or by the stabilization theorem, $K_*\left(\mathcal{K}^{\otimes Z} \rtimes H\right) \cong$ $K_*(C^*(H))$. Moreover, for each $1 \leq m < |Z|$, the quotient I_m/I_{m+1} is the direct sum of $\mathcal{K}^{\otimes F} \otimes \mathcal{O}_{n+1}^{\otimes Z-F}$ over the subsets F of Z with $|F| = m$. Thus, $I_m/I_{m+1}\rtimes H$ is Morita-equivalent to the direct sum of $\mathcal{K}^{\otimes F}\otimes \mathcal{O}_{n+1}^{\otimes Z-F}\rtimes H_F$ over $[F] \in H \backslash \{F \subset Z \mid |F| = m\}$ where H_F is the stabilizer of F in H. By induction on the size |Z| of Z (for all finite groups H at the same time), $K_* \left(\mathcal{K}^{\otimes F} \otimes \mathcal{O}_{n+1}^{\otimes Z-F} \rtimes H_F \right) \cong K_* \left(\mathcal{O}_{n+1}^{\otimes Z-F} \rtimes H_F \right)$ is finitely generated. Using the six-term exact sequences on K -theory associated to [\(4.2\)](#page-18-2), we see that $K_*(I_m \rtimes H)$ are all finitely-generated. By Lemma [2.4,](#page-5-2)

 $K_*\left(\mathcal{T}_{n+1}^{\otimes Z} \rtimes H\right) \cong K_*(C^*(H))$ since \mathcal{T}_{n+1} is KK -equivalent to \mathbb{C} by [\[Pim97\]](#page-30-4). Using the six-term exact sequence on K -theory associated to (4.1) , we now see that $K_*\left(\mathcal{O}_{n+1}^{\otimes Z} \rtimes H\right)$ is finitely-generated.

On the other hand, we have

$$
K_*\left(\left(\mathcal{O}_{n+1}^{\otimes Z}\otimes M_{n^{\infty}}^{\otimes Z}\right)\rtimes_r H\right)\cong K_*\left(\mathcal{O}_{n+1}^{\otimes Z}\rtimes_r H\right)[1/n]
$$

by Theorem [3.5.](#page-14-2) The assertion follows from Lemma [2.4](#page-5-2) since $\mathcal{O}_{n+1}\otimes M_n\infty$ is KK -equivalent to 0 by the UCT.

For an infinite G-set Z, Corollary [4.10](#page-18-0) for $A = \mathcal{O}_{n+1}$ may also be deduced from the combination of Theorem [3.5](#page-14-2) and the following result (for $A =$ $\mathcal{O}_{n+1}\otimes M_n$).

Theorem 4.12. Let G be a discrete group satisfying BCC and let Z be a countably infinite G-set. Let A be any unital C^* -algebra such that $[1_{A\otimes r}] =$ $0 \in K_0(A^{\otimes r})$ for some $r \geq 1$. Then we have

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right) \cong 0.
$$

Proof. By Theorem [2.3](#page-5-0) it is enough to show $K_*(A^{\otimes Z} \rtimes_r H) \cong 0$ for all finite subgroups H of G . Since Z is infinite, it contains infinitely many orbits of type H/H_0 for some fixed subgroup $H_0 \subset H$. Denote Z_0 the union of orbits of type H/H_0 . Let L be a (necessarily infinite) complete set of representatives for $H\backslash Z_0$. We have

$$
A^{\otimes Z_0} \cong (A^{\otimes L})^{\otimes H/H_0}.
$$

By assumption, the unital inclusion $\mathbb{C} \to A^{\otimes r}$ induces the zero element in $K_0(A^{\otimes r}) = KK(\mathbb{C}, A^{\otimes r})$. In particular, the maps

$$
A^{\otimes N} \to A^{\otimes N+r},
$$

induce the zero map in KK -theory since, on the level of KK -theory, they are given by the exterior Kasparov product with $[0] = [1] \in KK(\mathbb{C}, A^{\otimes r})$. It follows from Lemma [2.4,](#page-5-2) that the unital inclusions

$$
\left(A^{\otimes N}\right)^{\otimes H/H_0} \to \left(A^{\otimes N+r}\right)^{\otimes H/H_0}
$$

are zero in KK^H . Writing $A^{\otimes Z} = A^{\otimes Z_0} \otimes A^{\otimes Z - Z_0}$, we have

$$
K_*\left(A^{\otimes Z} \rtimes_r H\right) \cong \underset{N}{\text{colim}} K_*\left(\left(\left(A^{\otimes N}\right)^{\otimes H/H_0} \otimes A^{Z-Z_0}\right) \rtimes_r H\right).
$$

The right-hand side is zero by the preceding argument. \Box

Remark 4.13. Theorem [4.12](#page-19-0) can be applied whenever A is Morita equivalent to \mathcal{O}_{n^r+1} such that $[1_A] = n \in K_0(A) \cong \mathbb{Z}/n^r$, or whenever $K_0(A) \cong \mathbb{Q}/\mathbb{Z}$. In the second case, this is due to the fact that $[1]^{\otimes 2} \in K_0(A^{\otimes 2})$ is in the image of $K_0(A) \otimes_{\mathbb{Z}} K_0(A) \cong \mathbb{Q}/\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Q}/\mathbb{Z} = 0.$

Wreath products. For discrete groups G and H, the wreath product $H \wr G$ is defined as the semi-direct product $(\bigoplus_{g\in G} H)\rtimes G$, where G acts by left translation. We have a canonical isomorphism

$$
C_r^*(H \wr G) \cong C_r^*(H)^{\otimes G} \rtimes_r G.
$$

An application of Corollary [2.10](#page-10-0) gives the following result which generalizes [\[Li19\]](#page-29-2).

Theorem 4.14 (Theorem [D\)](#page-3-0). Let G be a group satisfying BCC and let H be a group for which $C_r^*(H)$ satisfies the UCT and for which the unital inclusion $\mathbb{C} \to C_r^*(H)$ induces a split injection $K_*(\mathbb{C}) \to K_*(C_r^*(H))$. Denote its cokernel by $\tilde{K}_*(C^*_r(H))$. Then we have

$$
K_*(C_r^*(H \wr G)) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \text{FIN}^{\times}} K_*\left(B^{\otimes F} \rtimes_r G_F\right)
$$

where B is any C^{*}-algebra satisfying UCT with $K_*(B) \cong \tilde{K}_*(C_r^*(H))$. In particular, if G is torsion-free, we have

$$
K_*(C_r^*(H \wr G)) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \mathrm{FIN}^{\times}} K_*\left(B^{\otimes F}\right).
$$

Remark 4.15. The assumptions on H in the above theorem are not very restrictive: If H is a discrete group for which the Baum–Connes assembly map is split-injective (for instance if H satisfies BCC, or if H is exact by [\[Hig00,](#page-29-18) Theorem 1.1]), then the unital inclusion $\mathbb{C} \to C_r^*(H)$ induces a split injection $K_*(\mathbb{C}) \to K_*(C_r^*(H))$ since the corresponding map $K_*^{\text{top}}(\lbrace e \rbrace) \to$ $K_*^{\text{top}}(H)$ always splits.

On the other hand, it follows from Tu's [\[Tu99,](#page-30-5) Proposition 10.7] that $C_r^*(H)$ satisfies the UCT for every a-T-menable (in particular every amenable) group H , or, more generally, if H satisfies the strong Baum–Connes conjecture in the sense of [\[MN06\]](#page-30-2).

Example 4.16. Let G be a group which satisfies BCC and consider the wreath product $\mathbb{F}_n \wr G$ with \mathbb{F}_n the free group in n generators. Since \mathbb{F}_n is known to be a-T-menable it follows that Theorem [4.14](#page-20-0) applies. Since $K_0(C_r^*(\mathbb{F}_n)) = \mathbb{Z}$ and $K_1(C_r^*(\mathbb{F}_n)) = \mathbb{Z}^n$ we may choose $B = \bigoplus_{i=1}^n C_0(\mathbb{R})$ so that Theorem [4.14](#page-20-0) implies

$$
K_*(C_r^*(\mathbb{F}_n \wr G)) \cong K_*(C_r^*(G)) \oplus \bigoplus_{F \in G \backslash \mathrm{FIN}^{\times}} K_*(B^{\otimes F} \rtimes_r G_F),
$$

where each summand $K_*(B^{\otimes F} \rtimes_r G_F)$ decomposes into a direct sum of equivariant K-theory groups of the form $K_H^*(V)$ for certain subgroups H of G_F and certain euclidean H-spaces V. A more precise analysis can be done, at least for $n = 2$, along the lines of Example [4.4.](#page-16-0)

In particular, if G is torsion free, we get

$$
K_*(C_r^*(\mathbb{F}_n \wr G)) \cong K_*(C_r^*(G)) \oplus \bigoplus_{[F] \in G \backslash \text{FIN}^{\times}} K_*(B^{\otimes F})
$$

with $B^{\otimes F} \cong \bigoplus_{i=1}^{n^{|F|}} C_0(\mathbb{R}^{|F|})$. Therefore each G-orbit of a nonempty finite set $F \subseteq G$ provides $n^{|F|}$ copies of $\mathbb Z$ as direct summands of K_0 if $|F|$ is even and of K_1 if $|F|$ is odd.

We close this section with

Corollary 4.17. Suppose that G and H are as as in Theorem $\frac{1}{4}$, such that the K-theory of both $C_r^*(G)$ and $C_r^*(H)$ is free abelian. Then the Ktheory of $C_r^*(H \wr G)$ is free abelian as well.

Proof. Note that $\tilde{K}_*(C^*_r(H))$ is free abelian as it is the direct summand of the free abelian group $K_*(C^*_r(H))$. Therefore, in Theorem [4.14,](#page-20-0) B can be taken as the direct sum of, possibly infinitely many, $\mathbb C$ and $C_0(\mathbb R)$. The assertion follows from the fact that equivariant K-theory $K^*_{G_0}(\mathbb{R}^{G_0})$ (more generally $K^*_{G_0}(V)$ for any G_0 -Euclidean space V) is a (finitely-generated) free abelian group for any finite group G_0 by [\[Kar02\]](#page-29-6) (or [\[EP09\]](#page-29-7)).

5. AF-algebras

Let $A = \operatorname{colim}_{A} A_n$ be a unital AF-algebra (with unital connecting maps) and let G be a discrete group satisfying BCC. If Z is a countable G -set, then in principle, we can try to compute

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right) \cong \underline{\operatorname{colim}}_n K_*\left(A_n^{\otimes Z} \rtimes_r G\right)
$$

using the decomposition from Theorem [3.6.](#page-14-1) In general, such a computation can be challenging, even in relatively simple cases like $A = M_2 \oplus M_3 \infty$. Instead of trying to compute the connecting maps, we now provide an abstract approach to calculating $K_*(A^{\otimes Z} \rtimes_r G)$ for an arbitrary unital AF-algebra A, a discrete group G satisfying BCC and a countable proper G-set Z (for example $Z = G$ with the left translation action).

We first need some preparation. Let $\mathcal F$ be a family of subgroups, i.e. a non-empty set of subgroups of G closed under taking conjugates and subgroups. The orbit category $\text{Or}_{\mathcal{F}}(G)$ has as objects homogeneous Gspaces G/H for each $H \in \mathcal{F}$ and as morphisms G-maps (see [\[DL98,](#page-29-19) Definition 1.1.). We will mainly use the family \mathcal{FIN} of finite subgroups. For any $G-C^*$ -algebra A, we have a functor from $Or_{\mathcal{F}}(G)$ to the category of graded abelian groups that sends G/H to $K_*(A \rtimes_r H)$ and a morphism $G/H_0 \to G/H_1$ given by $H_0 \to gH_1$ (so that $H_0 \subseteq gH_1g^{-1}$) to the map $K_*(A \rtimes_r H_0) \to K_*(A \rtimes_r H_1)$ induced by the composition

$$
A \rtimes_r H_0 \hookrightarrow A \rtimes_r gH_1g^{-1} \cong A \rtimes_r H_1,
$$

where the second map is given by conjugation with g^{-1} inside $A \rtimes_r G$.^{[6](#page-22-0)} We denote by

$$
\operatorname*{colim}_{G/H \in \operatorname{Or}_{\mathcal{F}}(G)} K_*(A \rtimes_r H)
$$

the colimit of the functor $G/H \mapsto K_*(A\rtimes_r H)$ from $\text{Or}_{\mathcal{F}}(G)$ to the category of graded abelian groups. The inclusions $A\rtimes_r H \to A\rtimes_r G$ induce a natural homomorphism

(5.1)
$$
\underset{G/H \in \text{Or}_{\mathcal{F}}(G)}{\text{colim}} K_*(A \rtimes_r H) \to K_*(A \rtimes_r G).
$$

Remark 5.1. The map (5.1) should not be confused with the assembly map

$$
a\text{smb}_{\mathcal{F}}\colon H_*^G(E_{\mathcal{F}}G,\mathbb{K}_A^{\text{top}})\to K_*(A\rtimes_r G),
$$

which corresponds to taking the homotopy colimit at the level of K -theory spectra instead of taking the ordinary colimit at the level of K-theory groups, see [\[DL98,](#page-29-19) Section 5.1]. By the involved universal properties, there is a natural commuting diagram

but the vertical map is neither injective nor surjective in general. One can think of the map [\(5.1\)](#page-22-1) as the best approximation of $K_*(A \rtimes_r G)$ using 0-dimensional $G-F-CW$ -complexes. Likewise, the best approximation by r -dimensional $G-F-CW$ -complexes can be defined by taking the colimit of $H_*^G(X, \mathbb{K}_A^{\text{top}})$ $_A^{\text{top}}$) over the category of *r*-dimensional *G*-*F*-CW-complexes.

Let us give two examples of how to compute $\underset{G/H \in \text{Or}_{\mathcal{TM}}(G)}{\text{colim}} K_*(A \rtimes_r H)$ when the structure or $Or_{\mathcal{FIN}}(G)$ is understood. Recall that the *coinvariants* of a G -module L are given by

$$
L_G \coloneqq \mathrm{colim}_G L \cong L/\langle x - gx \mid g \in G, x \in L \rangle.
$$

Here the group G is considered as a category with one object with morphisms given by the elements of G , and L is considered as a functor from G to the category of (graded) abelian groups.

Example 5.2. If G is torsion-free, then $\text{colim}_{G/H \in \text{Or}_{\mathcal{FIN}}(G)} K_*(A \rtimes_r H)$ can be identified with the coinvariants $K_*(A)_G$.

 6 This is a well-defined functor since different choices for g give the same map on Ktheory. On the other hand, $G/H \mapsto A \rtimes_r H$ does not define a functor. This is why it requires more care to upgrade this to a functor taking values in the category of spectra, see [\[DL98,](#page-29-19) Section 2].

Example 5.3. If G has only one non-trivial conjugacy class [H] of finite subgroups and if $N_G(H)$ denotes the normalizer of H in G, then the colimit colim $G/H \in \text{Corr}_\mathcal{I}(\mathcal{G})$ and $K_*(A \rtimes_r H)$ is given by the pushout

$$
K_*(A) \longrightarrow K_*(A \rtimes_r H)_{N_G(H)}
$$

\n
$$
\downarrow \qquad \qquad \downarrow
$$

\n
$$
K_*(A)_G \longrightarrow \underset{G/H \in \text{Or}_{\mathcal{FIN}}(G)}{\text{colim}} K_*(A \rtimes_r H).
$$

For a G -module M and a set S of positive integers, we denote by

$$
M[S^{-1}] := \underline{\text{colim}}(M \xrightarrow{\cdot s_1} M \xrightarrow{\cdot s_2} \cdots) \cong M \otimes_{\mathbb{Z}} \mathbb{Z}[S^{-1}]
$$

its localization at S, where (s_1, s_2, \dots) is a sequence containing every element of S infinitely many times. Note that localization at S commutes with taking coinvariants since both constructions are colimits.

Theorem 5.4 (Theorem [F\)](#page-3-2). Let A be a unital C^* -algebra of the form $A =$ $\underset{\epsilon}{\text{colim}} M_{k_n}(A_n)$ (with unital connecting maps) where $(k_n)_{n\in\mathbb{N}}$ is a sequence of positive integers and where each A_n is a unital C^* -algebra satisfying the assumptions of Theorem [2.8](#page-8-0) for the unital inclusion $\mathbb{C} \hookrightarrow A_n$. Let G be an infinite discrete group satisfying BCC , let Z be a countable proper G -set, and let $S \subseteq \mathbb{Z}$ be the set of all positive integers n such that $[1_A] \in K_0(A)$ is divisible by n. Then, the natural inclusions

(5.2)
$$
C_r^*(H) \to C_r^*(G), \ C_r^*(H) \to A^{\otimes Z} \rtimes_r H,
$$

$$
C_r^*(G) \to A^{\otimes Z} \rtimes_r G, \ A^{\otimes Z} \rtimes_r H \to A^{\otimes Z} \rtimes_r G,
$$

induce the following pushout diagram

$$
\begin{aligned}\n\text{colim}_{G/H \in \text{Or}_{\mathcal{FI}\mathcal{N}}(G)} & K_*(C_r^*(H))[S^{-1}] \longrightarrow \text{colim}_{G/H \in \text{Or}_{\mathcal{FI}\mathcal{N}}(G)} K_*(A^{\otimes Z} \rtimes_r H) \\
& \downarrow \\
& K_*(C_r^*(G))[S^{-1}] \longrightarrow K_*(A^{\otimes Z} \rtimes_r G).\n\end{aligned}
$$

In particular, if G is torsion-free, this pushout diagram reads

$$
\tilde{K}_*(C_r^*(G))[S^{-1}] \oplus K_*\left(A^{\otimes Z}\right)_G \cong K_*(A^{\otimes Z} \rtimes_r G),
$$

where $\tilde{K}_*(C^*_r(G))$ denotes the cokernel of $K_*(\mathbb{C}) \to K_*(C^*_r(G))$ induced from the unital inclusion $\mathbb{C} \hookrightarrow C_r^*(G)$.

Remark 5.5. By Theorem [3.1,](#page-11-2) Theorem [5.4](#page-23-1) applies to all unital AF-algebras. More generally, Theorem [5.4](#page-23-1) applies whenever A is of the form $\mathop{\rm colim}\limits_{n} M_{k_n}(A_n)$ (with unital connecting maps) where each A_n is one of the following examples:

(1) The unitization B^+ of a C^* -algebra B , e.g. $\mathbb{C} \oplus B$ for unital B ;

- (2) A C^* -algebra of the form $\bigoplus_{1 \leq j \leq N} C(X_j) \otimes M_{k_j}$ for nonempty compact metric spaces X_j and $\gcd(\overline{k}_1, \ldots, k_N) = 1$ (use Theorem [3.1\)](#page-11-2);
- (3) A reduced group C^* -algebra $C_r^*(\Gamma)$ of a countable group Γ that satisfies the UCT and such that the map $K_*(\mathbb{C}) \to K_*(C_r^*(\Gamma))$ is a split-injection (see Remark [4.15](#page-20-1) and Theorem [4.14\)](#page-20-0).

It would be interesting to know if Theorem [5.4](#page-23-1) also holds for unital ASHalgebras, i.e. when A is a filtered colimit of algebras of the form $p(C(X) \otimes$ M_n)p for a projection $p \in C(X) \otimes M_n$.

For the proof of Theorem [5.4,](#page-23-1) we need the following Lemma.

Lemma 5.6. Let G be any discrete group, let F be a family of subgroups of G, let $H_i \in \mathcal{F}$, and let A_i be H_i -C^{*}-algebras for $i \in I$. Denote by $A = \bigoplus_{i \in I} \text{Ind}_{H_i}^G(A_i)$ the direct sum of the induced $G-C^*$ -algebras. Then, the natural map

$$
\underset{G/H \in \text{Or}_{\mathcal{F}}(G)}{\text{colim}} K_*(A \rtimes_r H) \to K_*(A \rtimes_r G)
$$

is an isomorphism.

Proof. By additivity we may assume $A = \text{Ind}_{H_0}^G A_0$ for $H_0 \in \mathcal{F}$ and an H_0 -C^{*}-algebra A_0 . Write

$$
A=\bigoplus_{[z]\in G/H_0}A_z
$$

where A_z is the fiber of A at $[z] \in G/H_0$. Note that A_z is a zH_0z^{-1} -C^{*}algebra, in particular, A_e is the H_0 -C^{*}-algebra A_0 . By [\[CELY17,](#page-29-12) Proposition 2.6.8], the inclusion $A_e \rightarrow A$ induces an isomorphism

$$
K_*(A_e \rtimes_r H_0) \cong K_*(A \rtimes_r G).
$$

This isomorphism factors through colim_{G/H∈Or}_{π (G)} K_{*}(A $\rtimes_r H$) as

$$
K_*(A_e \rtimes_r H_0) \to K_*(A \rtimes_r H_0) \to \underset{G/H \in \text{Or}_{\mathcal{F}}(G)}{\text{colim}} K_*(A \rtimes_r H) \to K_*(A \rtimes_r G).
$$

Our claim follows once we show that the map

(5.3)
$$
K_*(A_e \rtimes_r H_0) \to \underset{G/H \in \text{Or}_{\mathcal{F}}(G)}{\text{colim}} K_*(A \rtimes_r H)
$$

is surjective. Fix a subgroup $H \in \mathcal{F}$. Decomposing G/H_0 into H-orbits and using the decomposition of [\(2.2\)](#page-8-1), we obtain a decomposition

(5.4)
$$
K_*(A \rtimes_r H) \cong \bigoplus_{[z] \in H \backslash G/H_0} K_*(A_z \rtimes_r H_z),
$$

where $H_z := H \cap zH_0z^{-1} \in \mathcal{F}$. In [\(5.4\)](#page-24-0), the inclusions $K_*(A_z \rtimes_r H_z) \subseteq$ $K_*(A \rtimes_r H)$ are induced by the natural inclusions $A_z \rtimes_r H_z \subseteq A \rtimes_r H$. In the colimit colim_{G/H∈Or_F(G)} $K_*(A \rtimes_r H)$, the summand

$$
K_*(A_z \rtimes_r H_z) \subseteq K_*(A \rtimes_r H)
$$

gets via conjugation with z^{-1} identified with the summand

$$
K_*(A_e \rtimes_r H'_z) \subset K_*(A \rtimes_r H'_z)
$$

corresponding to $[e] \in G/H_0$ and $H'_z = z^{-1}Hz \cap H_0$. But the elements in colim_{G/H∈Or_F(G)} $K_*(A \rtimes_r H)$ coming from $K_*(A_e \rtimes_r H'_z)$ are certainly in the image of the map in [\(5.3\)](#page-24-1) since

$$
K_*(A_e \rtimes_r H'_z) \to \mathrm{colim}_{G/H \in \mathrm{Or}_{\mathcal{F}}(G)} K_*(A \rtimes_r H)
$$

factors through the K-theory map of the inclusion $A_e \rtimes_r H'_z \subseteq A_e \times_r H_0$. \Box

As a direct consequence of Lemma [5.6,](#page-24-2) we get

Corollary 5.7. Let G be any discrete group, let Z be a countable G-set, and let A_0 and B be C^* -algebras with A_0 unital. Let $\dot{\mathcal{J}}_{A_0,B}^Z$ be the G - C^* -algebra

$$
\dot{\mathcal{J}}^Z_{A_0,B} = \bigoplus_{F \in \text{FIN}^\times(Z)} A_0^{\otimes Z-F} \otimes B^{\otimes F},
$$

so that $\mathcal{J}_{A_0,B}^Z = A_0^{\otimes Z} \oplus \dot{\mathcal{J}}_{A_0,B}^Z$ as in Definition [2.6.](#page-7-3) Then the natural map

$$
\operatornamewithlimits{colim}_{G/H \in \operatorname{Or}_{\mathcal{F}}(G)} K_* \left(\dot{\mathcal{J}}^Z_{A_0,B} \rtimes_r H \right) \to K_* \left(\dot{\mathcal{J}}^Z_{A_0,B} \rtimes_r G \right)
$$

is an isomorphism for any family $\mathcal F$ of subgroups of G containing all the stabilizers of the G-action on $\text{FIN}^{\times}(Z)$. This applies in particular when Z is a proper G-set and $\mathcal{F} = \mathcal{F} \mathcal{I} \mathcal{N}$ is the family of finite subgroups.

Proof of Theorem [5.4.](#page-23-1) We prove Theorem [5.4](#page-23-1) by considering three successively more general cases:

Case 1. A satisfies the assumptions of Theorem [2.8](#page-8-0) for the unital inclusion $\iota\colon \mathbb{C} \to A.$

By assumption, there is a C^* -algebra B and an element $\phi \in KK(B,A)$ such that $\iota \oplus \phi \in KK(\mathbb{C} \oplus B, A)$ is a KK-equivalence. Using the weak K-equivalence of \mathcal{J}_B^Z and $A^{\otimes Z}$ constructed in Theorem [2.8,](#page-8-0) we may identify the maps in [\(5.2\)](#page-23-2) with the natural maps

(5.5)
$$
C_r^*(G) \to \mathcal{J}_B^Z \rtimes_r G, \quad \mathcal{J}_B^Z \rtimes_r H \to \mathcal{J}_B^Z \rtimes_r G,
$$

where the first map is induced from the (non-unital) inclusion $\mathbb{C} \hookrightarrow \mathcal{J}_{B}^{Z}$ corresponding to $F = \emptyset$. We may therefore replace $A^{\otimes Z}$ by \mathcal{J}_B^Z throughout the proof. The corresponding statement for \mathcal{J}_{B}^{Z} then follows from Corollary [5.7](#page-25-0) since by the decomposition $\mathcal{J}_B^Z = \mathbb{C} \oplus \dot{\mathcal{J}}_B^Z$, we have

$$
K_*(\mathcal{J}_B^Z \rtimes_r G) \cong K_*(C_r^*(G)) \oplus \underset{G/H \in \text{Or}_{\mathcal{F} \mathcal{I} \mathcal{N}}(G)}{\text{colim}} K_*(\dot{\mathcal{J}}_B^Z \rtimes_r H)
$$

and consequently a pushout diagram

$$
\begin{array}{ccc}\n\operatorname{colim}_{G/H \in \operatorname{Or}_{\mathcal{FI}\mathcal{N}}(G)} & K_*(C_r^*(H)) & \longrightarrow & \operatorname{colim}_{G/H \in \operatorname{Or}_{\mathcal{FI}\mathcal{N}}(G)} & K_*(\mathcal{J}_B^Z \rtimes_r H) \\
\downarrow & & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \\
K_*(C_r^*(G)) & \longrightarrow & K_*(\mathcal{J}_B^Z \rtimes_r G).\n\end{array}
$$

In the torsion free case this becomes

$$
K_*(\mathcal{J}_B^Z \rtimes_r G) \cong K_*(C_r^*(G)) \oplus_{K_*(\mathbb{C})} K_*(\mathcal{J}_B^Z)_G \cong \tilde{K}_*(C_r^*(G)) \oplus K_*(\mathcal{J}_B^Z)_G.
$$

Case 2. $A = M_n \otimes D$ where D is as in Case [1.](#page-25-1)

By applying Case [1](#page-25-1) and localizing at n, we see that the maps in (5.2) (for D instead of A) induce a pushout diagram

$$
\begin{aligned}\n\text{colim} & K_*(C_r^*(H))[1/n] \longrightarrow \text{colim} & K_*\left(D^{\otimes Z} \rtimes_r H\right)[1/n] \\
& \downarrow \\
& K_*(C_r^*(G))[1/n] \longrightarrow K_*\left(D^{\otimes Z} \rtimes_r G\right)[1/n],\n\end{aligned}
$$

and in the torsion-free case an isomorphism

$$
\tilde{K}_*(C^*_r(G))[1/n] \oplus K_*\left(D^{\otimes Z}\right)[1/n]_G \cong K_*\left(D^{\otimes Z} \rtimes_r G\right)[1/n].
$$

Here we have used that localization at n commutes with taking colimits. By Theorem [3.5,](#page-14-2) the unital inclusion $D \to M_n \otimes D = A$ induces an isomorphism $K_*\left(D^{\otimes Z} \rtimes_r H\right)[1/n] \cong K_*\left(A^{\otimes Z} \rtimes_r H\right)$ for every subgroup H of G. This finishes the proof of Case [2.](#page-26-1)

Case 3. $A = \underline{\text{colim}}_k A_k$ where each A_k is as in Case [2.](#page-26-1)

For each k, denote by $S_k \subseteq S$ the set of positive integers n such that the unit $[1] \in K_0(A_k)$ is divisible by n. Note that we have $S = \bigcup_k S_k$. By Case [2,](#page-26-1) the conclusion of the theorem holds if we replace A by A_k and S by S_k . Now the general case follows by taking the filtered colimit along k .

6. RATIONAL AND k -ADIC COMPUTATIONS

In this section we give systematic tools to compute the K -theory of noncommutative Bernoulli shifts up to localizing at a supernatural number n (i.e. at the set of prime factors of $\mathfrak n$). The results apply to unital C^* algebras A for which the inclusion $\iota: \mathbb{C} \to A$ does not induce a split injection $K_0(\mathbb{C}) \to K_0(A)$ integrally, but a split injection

(6.1)
$$
K_0(\iota)[1/\mathfrak{n}]: K_0(\mathbb{C})[1/\mathfrak{n}] \hookrightarrow K_0(A)[1/\mathfrak{n}]
$$

after localizing at a supernatural number n. Important special cases are $\mathfrak{n} = k^{\infty}$ for $k \in \mathbb{N}$ (Example [6.1\)](#page-26-2), or when $\mathfrak{n} = \prod_{p} p^{\infty}$ (Example [6.2\)](#page-27-2). The latter case amounts to rational K -theory computations since in this case we have $L[1/\mathfrak{n}] \cong L \otimes_{\mathbb{Z}} \mathbb{Q}$ for any abelian group L. We give two examples of when one of these situations naturally occurs:

Example 6.1 ($M_n = M_k \infty$). Let A be a unital C^{*}-algebra that admits a finite-dimensional representation $A \to M_k(\mathbb{C})$ for some k. Then the unital inclusion $\iota: \mathbb{C} \to A$ induces a split-injection $K_*(\mathbb{C})[1/k] \to K_*(A)[1/k]$. Concrete examples are unital continuous trace C^* -algebras or subhomogeneous C ∗ -algebras.

Example 6.2 ($M_n = Q$). Let A be a unital C^{*}-algebra for which the unit $[1] \in K_0(A)$ is not torsion, for instance let A be unital and stably finite. Then the inclusion $\iota: \mathbb{C} \to A$ induces a split injection $K_*(\mathbb{C}) \otimes_{\mathbb{Z}} \mathbb{Q} \to K_*(A) \otimes_{\mathbb{Z}} \mathbb{Q}$.

Theorem 6.3 (c.f. Theorem [2.8\)](#page-8-0). Let G be a discrete group satisfying BCC , let Z be a countable G-set and let $\mathfrak n$ be a supernatural number. Let A be a unital C^* -algebra satisfying the UCT such that the unital inclusion $\mathbb{C} \to A$ induces a split injection $K_*(\mathbb{C})[1/\mathfrak{n}] \to K_*(A)[1/\mathfrak{n}]$. Denote by $\tilde{K}_{*}(A)$ the cokernel of the injection $K_{*}(\mathbb{C}) \to K_{*}(A)$ and let B be any C^{*} algebra satisfying the UCT with $K_*(B) \cong \tilde{K}_*(A)$. Then we have

$$
K_* \left(A^{\otimes Z} \rtimes_r G \right) [1/n] \cong K_* \left(\mathcal{J}_B^Z \rtimes_r G \right) [1/n] \cong \bigoplus_{[F] \in G \backslash \text{FIN}(Z)} K_* \left(B^{\otimes F} \rtimes_r G_F \right) [1/n].
$$

Proof. Replacing **n** by \mathfrak{n}^{∞} , we may assume that **n** is of infinite type. By the UCT, there is a UCT C^* -algebra B whose K -theory is isomorphic to the cokernel $\tilde{K}_*(A)$ of $K_*(\iota)$, and an element $\phi \in KK(B\otimes M_{\mathfrak{n}}, A\otimes M_{\mathfrak{n}})$ which together with the inclusion $\iota_{\mathfrak{n}}: M_{\mathfrak{n}} \to A \otimes M_{\mathfrak{n}}$ induces a KK-equivalence

$$
\iota_{\mathfrak{n}} \oplus \phi \in KK((\mathbb{C} \oplus B) \otimes M_{\mathfrak{n}}, A \otimes M_{\mathfrak{n}}).
$$

We can thus apply Theorem [2.8](#page-8-0) for $A \otimes M_n$ in place of A, for $A_0 = M_n$ and for $B \otimes M_{\mathfrak{n}}$ in place of B. We get

$$
K_*\left(A^{\otimes Z} \rtimes_r G\right)[1/\mathfrak{n}] \cong K_*\left((A \otimes M_{\mathfrak{n}})^{\otimes Z} \rtimes_r G\right)
$$

\n
$$
\cong K_*\left(\mathcal{J}_{M_{\mathfrak{n}},B\otimes M_{\mathfrak{n}}}^Z \rtimes_r G\right)
$$

\n
$$
\cong \bigoplus_{[F]\in G\backslash \text{FIN}(Z)} K_*\left((M_{\mathfrak{n}}^{\otimes Z-F} \otimes (B \otimes M_{\mathfrak{n}})^{\otimes F}) \rtimes_r G_F\right)
$$

\n
$$
\cong \bigoplus_{[F]\in G\backslash \text{FIN}(Z)} K_*\left(B^{\otimes F} \rtimes_r G_F\right)[1/\mathfrak{n}]
$$

where the first and last isomorphisms are obtained from Theorem [3.5.](#page-14-2) \Box

Theorem [5.4](#page-23-1) has a counter-part as well:

Theorem 6.4. Let G be a discrete group satisfying BCC, let Z be a countable, proper G -set, and let $\mathfrak n$ a supernatural number. Let A be a unital C^* -algebra which is a unital filtered colimit of C^* -algebras A_k satisfying the assumptions of Theorem [6.3.](#page-27-0) Then, the natural inclusions

$$
C_r^*(H) \to C_r^*(G), \quad C_r^*(H) \to A^{\otimes Z} \rtimes_r H,
$$

$$
C_r^*(G) \to A^{\otimes Z} \rtimes_r G, \quad A^{\otimes Z} \rtimes_r H \to A^{\otimes Z} \rtimes_r G,
$$

induce a pushout diagram

$$
\begin{array}{ccc}\n\operatorname{colim}_{G/H \in \operatorname{Or}_{\mathcal{FI}\mathcal{N}}(G)} & K_*(C_r^*(H))[1/\mathfrak{n}] & \longrightarrow & \operatorname{colim}_{G/H \in \operatorname{Or}_{\mathcal{FI}\mathcal{N}}(G)} K_* \left(A^{\otimes Z} \rtimes_r H \right)[1/\mathfrak{n}] \\
& & \downarrow & & \downarrow & \\
& K_*(C_r^*(G))[1/\mathfrak{n}] & \longrightarrow & K_* \left(A^{\otimes Z} \rtimes_r G \right)[1/\mathfrak{n}].\n\end{array}
$$

In particular, if G is torsion-free, this pushout diagram reads

$$
\tilde{K}_*(C^*_r(G))[1/\mathfrak{n}]\oplus K_*\left(A^{\otimes Z}\right)_G[1/\mathfrak{n}]\cong K_*\left(A^{\otimes Z}\rtimes_r G\right)[1/\mathfrak{n}].
$$

Proof. Without loss of generality we may assume that $\mathfrak n$ is of infinite type and A satisfies the assumption of Theorem [6.3.](#page-27-0) Let B, ϕ and ι_n be as in the proof of Theorem [6.3.](#page-27-0) In particular, Theorem [2.8](#page-8-0) applies so that $(A \otimes M_{\mathfrak{n}})^{\otimes Z}$ is weakly K-equivalent to $\mathcal{J}_{M_n,B\otimes M_n}^Z$. As in the proof of Theorem [5.4,](#page-23-1) the Theorem now follows from the combination of Corollary [5.7](#page-25-0) and Theorem $3.5.$

Example 6.5. Let $p \in C(X) \otimes M_n$ be a rank-k projection over $C(X)$ corresponding to some vector bundle on X. Let $A = p(C(X) \otimes M_n)p$ be the associated homogeneous C^* -algebra. Of course, A is Morita-equivalent to $C(X)$ but the unit-inclusion $\iota: \mathbb{C} \to A$ is not split-injective in general; it corresponds to the inclusion $\mathbb{Z}[p] \to K^0(X)$. On the other hand, A has a k-dimensional irreducible representation and therefore satisfies the assumptions of Theorem [6.3.](#page-27-0) Therefore, we get

$$
K_*\left(A^{\otimes G} \rtimes_r G\right)[1/k] \cong \bigoplus_{[F] \in G \backslash \mathrm{FIN}} K_*\left(B^{\otimes F} \rtimes_r G_F\right)[1/k]
$$

where B is any UCT C^* -algebra whose K-theory is isomorphic to the cokernel of $K_*(\iota)$.

As an application to Theorem [6.3,](#page-27-0) we obtain a proof of the fact that Bernoulli shifts by finite groups rarely have the Rokhlin property (see [\[Izu04,](#page-29-20) Definition 3.1]). This result is possibly known to experts and we would like to thank N. C. Phillips for mentioning it to us.

Corollary 6.6. Let A be a unital C^* -algebra satisfying the UCT such that $[1] \in K_0(A)$ is not torsion (for instance, suppose that A is stably finite). Let $G \neq \{e\}$ be a finite group and let Z a G-set. Then the Bernoulli shift of G on $A^{\otimes Z}$ does not have the Rokhlin property.

Proof. It follows from Theorem [6.3](#page-27-0) that the inclusion $C_r^*(G) \hookrightarrow A^{\otimes Z} \rtimes_r G$ induces a split injection

(6.2)
$$
K_*(C_r^*(G)) \otimes_{\mathbb{Z}} \mathbb{Q} \hookrightarrow K_* (A^{\otimes Z} \rtimes_r G) \otimes_{\mathbb{Z}} \mathbb{Q}.
$$

Now assume that the action of G on $A^{\otimes Z}$ has the Rokhlin property. Using [\[Phi11,](#page-30-6) Theorem 2.6], we can find a unital equivariant ∗-homomorphism $C(G) \to A^{\otimes Z}$. But then we can factor the map in [\(6.2\)](#page-28-1) as the composition

$$
K_*(C_r^*(G))\otimes_{\mathbb{Z}}\mathbb{Q}\to K_*(C(G)\rtimes_r G)\otimes_{\mathbb{Z}}\mathbb{Q}\to K_*\left(A^{\otimes Z}\rtimes_r G\right)\otimes_{\mathbb{Z}}\mathbb{Q},
$$

which is never injective unless $G = \{e\}.$

REFERENCES

- [BCH94] Paul Baum, Alain Connes, and Nigel Higson. Classifying space for proper actions and K-theory of group C ∗ -algebras. In C ∗ *-algebras: 1943–1993 (San Antonio, TX, 1993)*, volume 167 of *Contemp. Math.*, pages 240–291. Amer. Math. Soc., Providence, RI, 1994.
- [BEL21] Ulrich Bunke, Alexander Engel, and Markus Land. A stable ∞-category for equivariant KK-theory. *arXiv:2102.13372*, 2021.
- [Bla98] Bruce Blackadar. K*-theory for operator algebras*, volume 5 of *Mathematical Sciences Research Institute Publications*. Cambridge University Press, Cambridge, second edition, 1998.
- [CE01] Jérôme Chabert and Siegfried Echterhoff. Permanence properties of the Baum-Connes conjecture. *Doc. Math.*, 6:127–183, 2001.
- [CEL13] Joachim Cuntz, Siegfried Echterhoff, and Xin Li. On the K-theory of crossed products by automorphic semigroup actions. *Q. J. Math.*, 64(3):747–784, 2013.
- [CELY17] Joachim Cuntz, Siegfried Echterhoff, Xin Li, and Guoliang Yu. K*-theory for group* C ∗ *-algebras and semigroup* C ∗ *-algebras*, volume 47 of *Oberwolfach Seminars*. Birkhäuser/Springer, Cham, 2017.
- [CEOO04] J. Chabert, S. Echterhoff, and H. Oyono-Oyono. Going-down functors, the Künneth formula, and the Baum-Connes conjecture. *Geom. Funct. Anal.*, 14(3):491–528, 2004.
- [Dav96] Kenneth R. Davidson. C ∗ *-algebras by example*, volume 6 of *Fields Institute Monographs*. American Mathematical Society, Providence, RI, 1996.
- [DL98] James F. Davis and Wolfgang Lück. Spaces over a category and assembly maps in isomorphism conjectures in K- and L-theory. K*-Theory*, 15(3):201– 252, 1998.
- [EP09] Siegfried Echterhoff and Oliver Pfante. Equivariant K-theory of finite dimensional real vector spaces. *Münster J. Math.*, 2:65–94, 2009.
- [Hig00] N. Higson. Bivariant K-theory and the Novikov conjecture. *Geom. Funct. Anal.*, 10(3):563–581, 2000.
- [HK01] Nigel Higson and Gennadi Kasparov. E-theory and KK-theory for groups which act properly and isometrically on Hilbert space. *Invent. Math.*, 144(1):23–74, 2001.
- [Izu04] Masaki Izumi. Finite group actions on C^* -algebras with the Rohlin property. I. *Duke Math. J.*, 122(2):233–280, 2004.
- [Izu19] Masaki Izumi. The K-theory of the flip automorphisms. In *Operator algebras and mathematical physics*, volume 80 of *Adv. Stud. Pure Math.*, pages 123–137. Math. Soc. Japan, Tokyo, 2019.
- [Kar02] Max Karoubi. Equivariant K-theory of real vector spaces and real projective spaces. *Topology Appl.*, 122(3):531–546, 2002.
- [Kas88] G. G. Kasparov. Equivariant KK-theory and the Novikov conjecture. *Invent. Math.*, 91(1):147–201, 1988.
- [KN22] Julian Kranz and Shintaro Nishikawa. K-theory of Bernoulli shifts of finite groups on UHF-algebras. *arXiv:2210.00061*, 2022.
- [KS03] Gennadi Kasparov and Georges Skandalis. Groups acting properly on "bolic" spaces and the Novikov conjecture. *Ann. of Math. (2)*, 158(1):165–206, 2003.
- [Laf12] Vincent Lafforgue. La conjecture de Baum-Connes à coefficients pour les groupes hyperboliques. *J. Noncommut. Geom.*, 6(1):1–197, 2012.
- [Li19] Xin Li. K-theory for generalized Lamplighter groups. *Proc. Amer. Math. Soc.*, 147(10):4371–4378, 2019.

- [MN06] Ralf Meyer and Ryszard Nest. The Baum-Connes conjecture via localisation of categories. *Topology*, 45(2):209–259, 2006.
- [Ohh15] Issei Ohhashi. K-theory for the crossed products of infinite tensor product of C ∗ -algebras by the shift. *arXiv:1502.04290*, 2015.
- [Phi11] N. Christopher Phillips. Equivariant semiprojectivity. *arXiv:1112.4584*, 2011.
- [Pim97] Michael V. Pimsner. A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by Z. In *Free probability theory (Waterloo, ON, 1995)*, volume 12 of *Fields Inst. Commun.*, pages 189–212. Amer. Math. Soc., Providence, RI, 1997.
- [Sza18] Gábor Szabó. Equivariant Kirchberg-Phillips-type absorption for amenable group actions. *Comm. Math. Phys.*, 361(3):1115–1154, 2018.
- [Tho99] Klaus Thomsen. Asymptotic homomorphisms and equivariant KK-theory. *J. Funct. Anal.*, 163(2):324–343, 1999.
- [Tu99] Jean-Louis Tu. La conjecture de Baum-Connes pour les feuilletages moyennables. K*-Theory*, 17(3):215–264, 1999.

S.C.: STAT-MATH UNIT, INDIAN STATISTICAL INSTITUTE, 203 BARRACKPORE TRUNK Road, Kolkata 700 108, India.

sayan2008@gmail.com

S.E & J.K.& S.N.: Mathematisches Institut, Fachbereich Mathematik und INFORMATIK DER UNIVERSITÄT MÜNSTER, EINSTEINSTRASSE 62, 48149 MÜNSTER, GER-MANY.

echters@uni-muenster.de, j kran05@uni-muenster.de, snishika@uni-muenster.de.