Martin Beneke,* Patrick Hager[†] and Robert Szafron

Abstract This chapter reviews the construction of "soft-collinear gravity", the effective field theory which describes the interaction of collinear and soft gravitons with matter (and themselves), to all orders in the soft-collinear power expansion, focusing on the essential concepts. Among them are an emergent soft background gauge symmetry, which lives on the light-like trajectories of energetic particles and allows for a manifestly gauge-invariant representation of the interactions in terms of a soft covariant derivative and the soft Riemann tensor, and a systematic treatment of collinear interactions, which are absent at leading power in gravity. The gravitational soft theorems are derived from soft-collinear gravity at the Lagrangian level. The symmetries of the effective theory provide a transparent explanation of why soft graviton emission is universal to sub-sub-leading power, but gauge boson emission is not and suggest a physical interpretation of the form of the universal soft factors in terms of the charges corresponding to the soft symmetries. The power counting of soft-collinear gravity further provides an understanding of the structure of loop corrections to the soft theorems.

Keywords

Gravitation, soft-collinear effective field theory, effective Lagrangian, soft and collinear divergences, soft theorem, power corrections, Einstein-Hilbert theory

R. Szafron Department of Physics, Brookhaven National Laboratory, Upton, N.Y., 11973, U.S.A.

TUM-HEP-1425/22, 14 October 2022

M. Beneke, P. Hager

Physik Department T31, James-Franck-Straße 1, Technische Universität München, D-85748 Garching, Germany

^{*} Corresponding author

[†] Address after October 1st, 2022: PRISMA⁺ Cluster of Excellence & Mainz Institute for Theoretical Physics, Johannes Gutenberg University, D–55099 Mainz, Germany

"My reasons for now attacking this question are: (1) Because I can. [...] (2) Because something might go wrong and this would be interesting. Unfortunately, nothing does go wrong."

S. Weinberg, Ref. [38]

1 Introduction

The gravitational force is widely perceived to be fundamentally different from the gauge forces that govern the other microscopic interactions of the elementary particles. The gravitational interactions are inevitably non-renormalisable, calling for a modification at very short distances (or a non-trivial ultraviolet fixed point). Their underlying gauge symmetry is related to space-time transformations, in contrast to the internal $SU(3) \times SU(2) \times U(1)$ gauge symmetries operating on fields in rigid Minkowski space-time.

Yet, from the low-energy perspective and applying the basic principles of quantum field theory, the Lagrangian of weak-field gravity on Minkowski space follows from the desire to construct a consistent theory for a massless spin-2 particle in very much the same way as gauge theories do for the case of a massless spin-1 particle. The universality and space-time symmetry of gravitation then arises from the requirement that interacting massless fields with spin larger than $\frac{1}{2}$ must couple to a conserved current, which is the energy-momentum tensor for spin-2. This motivates a closer inspection of the relation between gauge theory and gravitational scattering in Minkowski space.

The study of gravitational scattering amplitudes in quantised weak-field gravity has attracted much attention after the discovery of remarkable relations between graviton and gluon scattering amplitudes [12, 13], which state that tree-level amplitudes of the former can be obtained by "squaring" Yang-Mills tree-level amplitudes and replacing colour factors by kinematic ones. The simplest example is the three-point amplitude in spinor-helicity notation (reviewed in [27]):

Numerous extensions of such "double copy" or "colour-kinematics duality" relations have been found to different gauge/gravity theories, to non-trivial classical backgrounds, and to the one-loop level.



Fig. 1 Emission of a boson with momentum k from an external energetic particle with momentum p.

Much can be learnt by looking at the behaviour of quantum amplitudes in the infrared (IR). When an energetic, massless particle with momentum p^{μ} emits another massless particle with momentum k^{μ} , see Figure 1, the internal propagator $1/(p+k)^2$ becomes singular in the *soft* limit $k^{\mu} \rightarrow 0$, and in the *collinear* limit $p^{\mu} || k^{\mu}$. When such configurations are integrated over the phase space of the emitted particle, or appear inside loops, the result is a logarithmic divergence. It has been recognised from the early days of quantum field theory that the soft and collinear limits exhibit universal, process-independent features [16, 28, 30] and that a precise definition of quantum-mechanical observables is required to obtain sensible, IR-finite results. The study of these limits in quantum electrodynamics and non-abelian gauge theories accordingly has a long history.

In QCD, which is strongly interacting in the infrared, understanding the soft and collinear limit is of paramount importance to make predictions for high-energy scattering, and culminates in powerful factorisation theorems [23]. In this way, one isolates the infrared physics in some well-defined functions, while leaving the more process-dependent features for perturbative calculations. Given the presence of several momentum scales, it is logical to apply effective Lagrangians to capture the IR physics, especially as Lagrangians are better suited than amplitudes to uncover the gauge-invariance and recursive structure of multi-scale problems. Soft-collinear effective theory (SCET) [2, 3, 5, 6] has therefore emerged as an important conceptual and calculational tool for factorisation in gauge theories. It is designed to precisely reproduce Feynman amplitudes in their soft and collinear limits. Moreover, at least in principle, it can do so beyond the leading power in the expansion in small scale ratios. Although the case of high-energy gravitational scattering appears to be of less practical relevance, in view of the above mentioned relations between graviton and Yang-Mills scattering amplitudes, which is presently not understood at Lagrangian level, it is suggestive to apply effective Lagrangian techniques to at least the soft and collinear limits of graviton amplitudes-which is the subject of this chapter.

These limits already exhibit interesting similarities and differences between massless spin-1 and spin-2 particles (gauge bosons and gravitons, respectively) coupled to matter. It has been noted long ago [38] that the "eikonal" or leading soft limit of gravity is very similar to gauge theory. The long-wavelength radiation "sees" only the direction of motion (classical trajectory) and charge of energetic particles. Thus, in the eikonal approximation, the amplitude for radiating a single soft graviton from energetic particles with momenta p_i^{μ} emerging from a hard scattering process,

Martin Beneke, Patrick Hager and Robert Szafron

$$\mathscr{A}_{\rm rad}(p_i;k) = \frac{\kappa}{2} \sum_i \frac{p_i^{\mu} p_i^{\nu} \varepsilon_{\mu\nu}(k)}{p_i \cdot k} \mathscr{A}(p_i), \qquad (2)$$

is obtained from its gauge-theory correspondent,

$$\mathscr{A}_{\rm rad}(p_i;k) = -g_s \sum_i t_i^a \frac{p_i \cdot \boldsymbol{\varepsilon}^a(k)}{p_i \cdot k} \mathscr{A}(p_i), \qquad (3)$$

by simply replacing the gauge charge (generator) by the gravitational charge, momentum, $t_i^a \rightarrow p_i^v$ and adjusting the coupling g_s and polarisation vector $\varepsilon_{\mu}^a(k)$ to the gravitational coupling, $\kappa = \sqrt{32\pi G_N}$, and polarisation tensor, $\varepsilon_{\mu\nu}(k)$. Since eikonalised propagators $1/(p_i \cdot k)$ are closely related to semi-infinite Wilson line operators, we expect soft graviton Wilson lines to play a similar role for soft graviton physics [32, 40] as they do in gauge theories. Eqs. (2), (3) represent the leading terms in the so-called *soft theorems*, to which we shall return in a later section of this chapter.

The collinear limit of graviton amplitudes is, however, very different from the one of gauge amplitudes. In fact, in gravity, collinear enhancements and singularities are absent altogether. As a consequence, even if the gravitational coupling was not minuscule, i.e. near Planckian scattering energies, energetic particles do not produce gravitational jets, which in QCD constitute the most visible footprints of the non-abelian charges of the quarks and gluons. The absence of collinear singularities for graviton emission was first shown in [38] in the simultaneous eikonal limit. Weinberg also noted that it would be rather troublesome, if this was not the case, since it would prevent the existence of massless particles with gravitational charges, that is, any non-vanishing four momentum. However, while massless particles with gauge charges do not exist in Nature, and hence there is no conflict with the existence of collinear singularities in gauge theories, there *are* massless particles which gravitate, such as the photons and the gravitons themselves.

There is a simple qualitative explanation for the absence of collinear graviton singularities based on the classical radiation pattern [11]. When an energetic particle with virtuality much less than its three-momentum squared \mathbf{p}^2 emits a graviton with momentum \mathbf{k} with small angle θ between \mathbf{p} and \mathbf{k} , the near mass-shell singularity of the emitting particle propagator $1/(|\mathbf{p}||\mathbf{k}|(1 - \cos \theta))$ yields a factor θ^{-2} for the splitting amplitude. Quantising the radiation field in the spherical basis with single-particle states $|\mathbf{k}jm;\lambda\rangle$, where λ denotes helicity (± 2 for gravitons and ± 1 for gauge bosons) and *jm* the angular momentum quantum numbers with respect to the quantisation axis \mathbf{p} , this implies that the emitted graviton must be in a state $|\mathbf{k}j0;\lambda\rangle$, where m = 0 due to angular momentum and helicity conservation. The angular dependence of this state is given by the spin-weighted spherical harmonic or Wigner function $D_{\pm\lambda,0}^j(\mathbf{k}) \propto \sin^{|\lambda|} \frac{\theta}{2}$, which tends to zero as $\theta^{|\lambda|}$ in the $\theta \rightarrow 0$ limit. Thus, the splitting amplitude has no singularity in the collinear limit for graviton emission ($\lambda = \pm 2$) in contrast to the case of gauge bosons.

The above argument refers to the physical polarisation states of the graviton and thus does not cover the properties of individual Feynman amplitudes in general, in particular in covariant gauges, which do have collinear divergences. The formal demonstration of the absence of collinear divergences without the restriction to the eikonal limit adopted in [38] has been presented only relatively recently [1] with diagrammatic factorisation methods. This fact is made evident in the construction of the soft-collinear effective Lagrangian for gravity ("soft-collinear gravity") [11]: the leading effective Lagrangian describing collinear graviton self-interactions and their interactions with matter is a free theory. This motivates the investigation of collinear gravitational physics at *sub-leading order* in the collinear expansion, where it is non-trivial, and naturally leads to the systematic construction of soft-collinear effective gravity beyond the leading power in both, the collinear and soft limits [10].

The present chapter starts with a review of basic ideas and methods for softcollinear Lagrangians, assuming no prior familiarity with the subject. We then provide a technically light-weight discussion of soft-collinear gravity, focusing on the exposition of the principles of the construction, the structure and emergent symmetries of the result at the expense of many technical details, for which we refer to [10]. By definition, soft-collinear gravity builds an extension of the so-called soft theorems to all orders in the loop and soft expansion. It is nevertheless of interest to rederive them from the effective Lagrangian [9]. In the last section of this chapter, we briefly cover how this provides an understanding of why in gravity the soft theorem extends to the next-to-next-to-soft order (but does not in gauge theory) and how the form of the universal terms is related to the (emergent) soft gauge symmetries of the effective Lagrangian. The chapter concludes with a discussion of loop corrections to the soft theorem.

2 Basic ideas and concepts

The following section sets up the notation and introduces a number of concepts that arise in effective field theory (EFT) and SCET in particular.

2.1 Perturbative gravity

The full theory, from which SCET gravity is constructed, is the Einstein-Hilbert theory with action

$$S_{\rm EH} = -\frac{2}{\kappa^2} \int d^4 x \sqrt{-g} R \,, \tag{4}$$

coupled to matter, here a minimally-coupled scalar field φ in the curved space-time with metric tensor $g_{\mu\nu}$. The matter part is described by the action

$$S_{\varphi} = \int d^4 x \sqrt{-g} \, \frac{1}{2} g^{\mu\nu} \partial_{\mu} \varphi \partial_{\nu} \varphi \,, \tag{5}$$

where g denotes the metric determinant.³ At this point, there is an important notion to clarify: the Einstein-Hilbert action is not renormalisable in the strict sense. Instead, it should be treated as the first term in a low-energy EFT of gravity. This idea was pioneered in [25], and is also well explained in [26]. If gravitational loops are included, it is necessary to introduce additional terms to render the theory finite. These higher-order terms correspond to a derivative expansion, and they can be expressed as products of Riemann tensors. Schematically, the full action then takes the form

$$S_{\rm grav,EFT} = -\int d^4x \sqrt{-g} \left(\Lambda + \frac{2}{\kappa^2} R - c_1 R^2 - c_2 R_{\mu\nu} R^{\mu\nu} + \dots \right),$$
(7)

where Λ is the cosmological constant and $R_{\mu\nu} = R^{\alpha}_{\mu\alpha\nu}$ is the Ricci tensor. This subtlety is important if one wants to consider higher-loop orders in gravity, though this is not very relevant for the following discussion. While the detailed form of the soft-collinear effective theory is determined by the full theory from which it is derived, the *construction* of the effective theory does not depend on the precise loop order that is considered. All these higher-order terms respect the same gauge symmetry that is already present in the leading term, the diffeomorphism invariance, and this symmetry forms the guiding principle of the SCET construction.

The diffeomorphisms can be arranged in the form of local translations

$$x^{\mu} \to x^{\prime \mu}(x) = x^{\mu} + \mathcal{E}^{\mu}(x), \qquad (8)$$

where $\mathcal{E}^{\mu}(x)$ is some (not necessarily small) vector field. Under such a transformation, scalar fields behave as

$$\boldsymbol{\varphi}(x) \to \boldsymbol{\varphi}'(x') \stackrel{!}{=} \boldsymbol{\varphi}(x) \,. \tag{9}$$

The transformed scalar field $\varphi'(x')$ can be expressed as

$$\varphi'(x + \varepsilon(x)) \equiv T_{\varepsilon} \varphi'(x), \tag{10}$$

where the translation operator T_{ε} is defined as

$$T_{\varepsilon}f(x) = f(x) + \varepsilon^{\alpha}(x)\partial_{\alpha}f(x) + \frac{1}{2}\varepsilon^{\alpha}(x)\varepsilon^{\beta}(x)\partial_{\alpha}\partial_{\beta}f(x) + \mathscr{O}(\varepsilon^{3}).$$
(11)

In the following, this active point of view is adopted. That is, the local translations correspond to *purely internal* transformations, acting on the field space, instead of actually transforming the coordinates. In practice, this means that the dynamical fields $\varphi(x) \rightarrow \varphi'(x)$ transform, but never the coordinates *x* themselves. A scalar

$$R^{\mu}_{\ \nu\alpha\beta} = \partial_{\alpha}\Gamma^{\mu}_{\ \beta\nu} - \partial_{\beta}\Gamma^{\mu}_{\ \alpha\nu} + \Gamma^{\mu}_{\ \alpha\lambda}\Gamma^{\lambda}_{\ \beta\nu} - \Gamma^{\mu}_{\ \beta\lambda}\Gamma^{\lambda}_{\ \alpha\nu} \tag{6}$$

³ In the following, the convention

for the Riemann tensor and the metric signature (+, -, -, -) are employed.

field then transforms as

$$\varphi(x) \to \varphi'(x) = [U(x)\varphi(x)],$$
 (12)

where due to (10) U(x) is the inverse translation, $U(x) = T_{\varepsilon}^{-1.4}$. The metric field transforms as

$$g_{\mu\nu}(x) \to \left[U(x) \left(U_{\mu}^{\ \alpha}(x) U_{\nu}^{\ \beta}(x) g_{\alpha\beta}(x) \right) \right], \tag{13}$$

with Jacobi matrices

$$U^{\mu}_{\ \alpha}(x) = \frac{\partial x'^{\mu}}{\partial x^{\alpha}}(x), \quad U_{\mu}^{\ \alpha}(x) = \frac{\partial x^{\alpha}}{\partial x'^{\mu}}(x).$$
(14)

It is convenient to adopt this active point of view for two reasons: first, it emphasises the formal similarity to gauge theories. Note that the transformation (12) looks formally the same as the transformation of a matter field $\phi^a(x)$ with respect to a non-abelian gauge symmetry,

$$\phi^a(x) \to U^{ab}(x)\phi^b(x), \tag{15}$$

which then exposes the similarities and differences between these transformations in gravity and gauge theory. Second, in this active point of view, one never has to worry about transformations of the coordinates, the integral measure d^4x as well as derivatives $\frac{\partial}{\partial x^{\mu}}$, and only needs to keep track of the transformation of the dynamic fields. Note that this point of view does not change the *form* of invariant objects. In the passive point of view, $\int d^4x \sqrt{-g} \mathcal{L}$ is diffeomorphism-invariant, since the invariant measure $d^4x \sqrt{-g}$ appears in combination with a scalar quantity \mathcal{L} . In the active point of view, the relevant object is the scalar density $\sqrt{-g}\mathcal{L}$, which is gaugeinvariant up to total derivatives, rendering the integral manifestly invariant. This provides a guiding principle for the required manipulations, such as the construction of gauge-invariant or covariant fields.

To construct perturbative gravity, one assumes small fluctuations of the metric field, and performs a weak-field expansion of

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + \kappa h_{\mu\nu}(x), \qquad (16)$$

in $h_{\mu\nu}(x)$ around Minkowski-space with metric $\eta_{\mu\nu}$. The action then turns into an infinite series in $h_{\mu\nu}$, resp. κ :

$$S = \sum_{k=0}^{\infty} \kappa^k S^{(k)} \,, \tag{17}$$

where the precise form of $S^{(k)}$ at higher orders depends on which terms one considers to be part of the "full theory", i.e. if one only considers Einstein-Hilbert (4), or also takes into account higher-order Riemann terms as in (7).

⁴ The square-bracket notation emphasises that U(x) is a derivative operator that acts only on $\varphi(x)$.

In this weak-field expansion, the residual gauge transformations correspond to the translations⁵

$$x^{\mu} \to x^{\mu} + \kappa \varepsilon^{\mu}(x) \,. \tag{18}$$

In this expansion, the action of U(x) is given by

$$[U(x)\varphi(x)] = \varphi(x) - \kappa \varepsilon^{\alpha}(x)\partial_{\alpha}\varphi(x) + \frac{\kappa^{2}}{2}\varepsilon^{\alpha}(x)\varepsilon^{\beta}(x)\partial_{\alpha}\partial_{\beta}\varphi(x) + \kappa^{2}\varepsilon^{\alpha}(x)\partial_{\alpha}\varepsilon^{\beta}(x)\partial_{\beta}\varphi(x) + \mathscr{O}(\varepsilon^{3}).$$
(19)

To first order, the transformations (12), (13) reproduce the well-known results

$$\begin{aligned} h_{\mu\nu} &\to h_{\mu\nu} - \partial_{\mu} \varepsilon_{\nu} - \partial_{\nu} \varepsilon_{\mu} + \mathscr{O}(\varepsilon^{2}, \varepsilon h) \,, \\ \varphi &\to \varphi - \kappa \varepsilon^{\alpha} \partial_{\alpha} \varphi + \mathscr{O}(\varepsilon^{2}) \,. \end{aligned}$$
 (20)

An immediate consequence of these truncated diffeomorphisms is that objects which are homogeneous in h, that is, are monomials in h, cannot be gauge-invariant at the same time. Gauge-invariant objects, such as the Riemann tensor, are represented as a series order-by-order in h or κ . When working with a theory expanded in h, sub-leading terms, that is, higher-order terms in h, appear in precise combinations to yield a gauge-invariant theory. This is a generic feature of non-linearly realised symmetries.

2.2 Basic concepts of SCET

SCET is the theory describing the (self-)interactions of soft and collinear particles [2, 3, 5, 6]. It is one of the more complicated effective theories in modern particle physics, and is used to great success in high-energy collider physics. Despite its technical nature, one can intuitively understand the construction with only a few key concepts. It is this intuition that is paramount in constructing the gravitational analogue. This section serves as an introduction into the underlying concepts of SCET. While the section aims to be self-contained, it will focus on exposition rather than derivations. Detailed discussions and computations can be found in [5, 7, 9, 10].

2.2.1 Kinematics and power-counting

The general kinematics underlying SCET consist of an energetic scattering, characterised by its large energy scale Q of some hard process, which creates a number of energetic particles, as well as some low-energy, soft radiation. These energetic particles are called "collinear" and develop into jets by collinear radiation. The jets

8

⁵ We extract a factor of κ from ε^{μ} , so that $h_{\mu\nu}$ has the linear gauge transformation (20) that does not contain κ explicitly.



Fig. 2 The generic kinematic situation of a scattering process described by SCET. Blue lines denote collinear particles forming jets, emanating from a point-like hard interaction. Red lines represent soft modes connecting jets.

are assumed to be well-separated in angle from each other. The situation is depicted in Figure 2.

To be precise, the energetic particles of jet *i* are taken to be ultrarelativistic (light-like), and characterised by the light-like direction n_{i-}^{μ} of the jet with respect to which they have small transverse momentum. Corresponding to each n_{i-}^{μ} , there is a n_{i+}^{μ} such that $n_{i+} \cdot n_{i-} = 2$, $n_{i\pm}^2 = 0$. These two reference vectors, as well as the two remaining transverse directions, form a basis, with metric tensor

$$\eta^{\mu\nu} = \frac{1}{2} n^{\mu}_{i+} n^{\nu}_{i-} + \frac{1}{2} n^{\nu}_{i+} n^{\mu}_{i-} + \eta^{\mu\nu}_{\perp i}.$$
⁽²¹⁾

One can decompose a collinear momentum p^{μ} as

$$p^{\mu} = n_{i+} p \frac{n_{i-}^{\mu}}{2} + p_{\perp i}^{\mu} + n_{i-} p \frac{n_{i+}^{\mu}}{2}, \qquad (22)$$

where the subscript $\perp i$ denotes the transverse components with respect to $n_{i+}^{\mu}, n_{i-}^{\mu}$. Introducing the power-counting parameter $\lambda \sim p_{\perp i}/(n_{i+}p) \ll 1$, the components of the collinear momentum *p* scale as

$$(n_{i+}p, p_{\perp i}, n_{i-}p) \sim (1, \lambda, \lambda^2)Q, \qquad (23)$$

and have virtuality $p^2 = \lambda^2 Q^2$. It is a convention to set Q = 1, which is adopted in the following. The condition $\lambda \ll 1$ means that the momentum flows in the direction n_{i-}^{μ} , and only small fractions are deposited in the transverse directions. For multiple particles in the same collinear sector, it means that these are separated by a small angle, thus they constitute a jet. Besides the collinear modes, also soft modes can

be present, which can interact with collinear modes without changing their collinear nature. This implies isotropic momentum scaling $k^{\mu} \sim \lambda^2$ and virtuality $k^2 \sim \lambda^4$, parametrically smaller than collinear virtuality. Soft modes can be exchanged between the energetic particles in different directions, as shown in Figure 2. This kinematic situation, consisting only of collinear and soft modes of different virtuality, is usually denoted as SCET_I. In this chapter, we refer exclusively to this situation.

2.2.2 Field content

The goal is to construct an EFT that provides a systematic expansion for the fulltheory scattering amplitudes in the soft and collinear limits. The construction differs considerably from traditional EFTs, such as the Fermi theory of weak decays. In these theories, one is interested in "light physics" and integrates out the "heavy fields", with masses above some scale Λ . This gives rise to an effective Lagrangian wherein only the light fields are dynamical, which now contains (local) higherdimensional operators, that describe the short-distance physics. The dimension of these operators serves as power-counting parameter for the expansion in $1/\Lambda$.

In SCET, however, the soft and collinear regions of *all* relevant full-theory particles can contribute to scattering amplitudes. Instead of integrating out heavy fields, one integrates out certain fluctuations of the fields, or, equivalently, certain *regions* of momenta. Specifically, one integrates out the *hard regions* of momenta, while keeping *collinear* and *soft* ones as the degrees of freedom in the EFT. In order to achieve this systematically at the Lagrangian level, one needs to split the full-theory fields ϕ_J into hard, soft and collinear modes $\phi_{J,h}$, $\phi_{J,s}$, ϕ_{J,c_i} .

To construct a systematic expansion, these modes must be *homogeneous* in λ , that is, they must scale with a unique power of λ , in order to label them as hard, collinear, and soft. For this reason, the EFT uses the different fields ϕ_{J,c_i} , $\phi_{J,s}$, which specifically describe the fluctuations of the original field in the respective kinematic regions. For SCET, the relevant modes are soft and collinear ones, as depicted in Figure 2. In addition, each term in the Lagrangian must be fully expanded to also be homogeneous, in the sense that it does not contain any further sub-leading terms.

Since the components n_{i+p} of collinear fluctuations are of the order of the hard scale, one cannot expand in n_{i+p}/Q , and the effective Lagrangian cannot be local. To see this, note that any operator can feature an arbitrary number of large derivatives $n_{i+}\partial$, which all scale as $n_{i+}\partial \sim 1$ and are thus in principle present at any order in λ . However, this tower of derivatives can be traded for the non-locality in the n_{i+}^{μ} direction. This is possible simply by rewriting the derivatives as a translation, using

$$\phi_{J,c_i}(x+tn_{i+}) = \sum_{k=0}^{\infty} \frac{t^k}{k!} (n_{i+}\partial)^k \phi_{J,c_i}(x) \,. \tag{24}$$

Thus, instead of keeping track of large derivatives $n_{i+}\partial$, one allows for non-localities in the n_{i+}^{μ} direction for collinear objects. When several energetic particles scatter at large angles, the theory is therefore non-local along the directions n_{i+}^{μ} , i.e.

orthogonal to the classical trajectories of the collinear particles. It is, however, local along these trajectories, resp. the direction n_{i-}^{μ} of their momentum.

In conventional effective theories, the importance of a local operator is tied to its mass dimension. In SCET, however, the power-counting is not related to the mass dimension. In particular, different components of a field can acquire different scaling in λ . To determine the power-counting of these fields, one considers the two-point function

$$\langle 0|T\varphi(x)\varphi(0)|0\rangle = \int \underbrace{\frac{d^4p}{(2\pi)^4}}_{\lambda^4(\lambda^8)} e^{-ipx} \underbrace{\frac{i}{p^2 + i0}}_{\lambda^{-2}(\lambda^{-4})} \sim \lambda^2(\lambda^4), \qquad (25)$$

where the power-counting of collinear (soft) momenta is given explicitly. One obtains

$$\varphi_{c_i}(x) \sim \lambda, \quad \varphi_s(x) \sim \lambda^2,$$
(26)

for the collinear and soft scalar field, respectively. For the graviton field $h_{\mu\nu}(x)$, one first performs the weak-field expansion and fixes a general de-Donder gauge with parameter *b*. Then, the two-point function reads

$$\langle 0|Th_{\mu\nu}(x)h_{\alpha\beta}(0)|0\rangle = i\kappa^2 \int \frac{d^4p}{(2\pi)^4} \frac{e^{-ipx}}{p^2 + i0} \left(P_{\mu\nu,\alpha\beta} + \frac{1-b}{b}S_{\mu\nu,\alpha\beta}\right), \quad (27)$$

where

4

$$P_{\mu\nu,\alpha\beta} = \frac{1}{2} \left(\eta_{\mu\alpha} \eta_{\nu\beta} + \eta_{\mu\beta} \eta_{\nu\alpha} - \eta_{\mu\nu} \eta_{\alpha\beta} \right),$$

$$S_{\mu\nu,\alpha\beta} = \frac{1}{2p^2} \left(\eta_{\mu\alpha} p_{\nu} p_{\beta} + \eta_{\mu\beta} p_{\nu} p_{\alpha} + p_{\mu} p_{\alpha} \eta_{\nu\beta} + p_{\mu} p_{\beta} \eta_{\nu\alpha} \right).$$
(28)

Inserting the collinear momentum scaling, one obtains $P_{\mu\nu,\alpha\beta} \sim 1$ if it is non-vanishing, since it does not depend on momenta. The other combination, $S_{\mu\nu,\alpha\beta}$, has non-trivial λ -scaling. For example, for the \perp + and \perp \perp modes,⁶ one obtains

$$S_{\perp+,\perp+} \sim \frac{1}{\lambda^2}, \quad P_{\perp+,\perp+} = 0,$$

$$S_{\perp\perp,\perp\perp} \sim 1, \quad P_{\perp\perp,\perp\perp} \sim 1.$$
(29)

Then, the scaling of the components of the collinear graviton field is easily determined from (27) to be [11]

$$\begin{array}{ll} h_{++} \sim \lambda^{-1}, & h_{+\perp} \sim 1, & h_{+-} \sim \lambda, \\ h_{--} \sim \lambda^3, & h_{-\perp} \sim \lambda^2, & h_{\perp\perp} \sim \lambda, \end{array}$$
(30)

⁶ For a tensor index μ , +, - means $T_{\pm} \equiv n_{\pm}^{\mu}T_{\mu}$, while \perp stands for $T_{\mu_{\perp}}$. In the remainder of Section 2, we drop the collinear direction label *i* whenever referring to a single collinear sector.

which implies $h^{\mu}_{\mu} \equiv h \sim \lambda$ for the trace. Note that the theory contains the $\mathscr{O}(1)$ field component $h_{\mu\perp+} \sim 1$, and even a power-enhanced component $h_{++} \sim \lambda^{-1}$. This is problematic for the λ expansion in the effective theory and must be addressed in the construction of the theory. For now, observe that coupling $h_{\mu\nu}$ to a vector V^{μ} of the same collinearity yields

$$h_{\mu\nu}V^{\nu} = \frac{1}{2}(h_{\mu+}V_{-} + h_{\mu-}V_{+}) + h_{\mu\nu_{\perp}}V^{\nu_{\perp}} \sim \lambda V_{\mu}, \qquad (31)$$

so index contractions within the same collinear sector are suppressed by a power of λ . This argument does not hold for couplings between different sectors, where h_{++} could give rise to power-enhancement. For the soft graviton, it is straightforward to derive the isotropic scaling $s_{\mu\nu} \sim \lambda^2$ from its propagator.

2.2.3 Gauge symmetry

Next, it is useful to discuss the gauge symmetry of SCET. In the full theory, there is only one gauge symmetry, diffeomorphism invariance. Each full-theory field furnishes some representation of the diffeomorphism group, and comes with its own gauge transformation. For the scalar field and the graviton, the linear transformation is given in (20). Next, one performs the mode split, that is, for each full-theory field $h_{\text{full},\mu\nu}(x)$ and $\varphi(x)$, one obtains collinear modes $h_{\mu\nu}(x)$, $\varphi_c(x)$ and soft modes $s_{\mu\nu}(x)$, $\varphi_s(x)$. This has implications for the gauge symmetry in the EFT. Once the split is implemented, e.g. naively as $h_{\text{full},\mu\nu} = h_{\mu\nu} + s_{\mu\nu}$, the right-hand side has to transform like the full-theory graviton on the left. This gives constraints on the allowed gauge symmetry, since the two fields on the right-hand side are modes with homogeneous power-counting by construction. The soft field can never transform with a gauge parameter that contains collinear fluctuations, since this would turn the soft field into a collinear field. But such gauge parameters are allowed in the transformation of h_{uv} . The solution is to *extend* the gauge symmetry to two separate, collinear and soft, gauge symmetries such that the collinear fields take the role of *fluctuations* on top of the *soft background* $g_{s\mu\nu} = \eta_{\mu\nu} + \kappa s_{\mu\nu}$. The technical details will be presented in Section 4. The collinear gauge transformation then reads

$$\kappa h_{\mu\nu} \rightarrow \left[U_c \left(U_{c\mu}^{\ \alpha} U_{c\nu}^{\ \beta} (g_{s\alpha\beta} + \kappa h_{\alpha\beta}) \right) \right] - g_{s\mu\nu} ,$$

$$g_{s\mu\nu} \rightarrow g_{s\mu\nu} .$$
(32)

The intuition behind these transformations is clear: the fluctuation $h_{\mu\nu}$ comes with its own gauge symmetry, but the background $g_{s\mu\nu}$ is unaffected by this collinear transformation, as it must be. The fluctuation $h_{\mu\nu}$ transforms in the same way as the $h_{\mu\nu}$ of the weak field expansion in the full theory, except that the rigid Minkowski background $\eta_{\mu\nu}$ is replaced by $g_{s\mu\nu}$ in (32). The soft background field is itself dynamic, with $s_{\mu\nu}$ transforming under the soft gauge symmetry $U_s(x)$ (to be discussed in Section 4). Let us stress the main message: to consistently implement the split into soft and collinear modes, one treats the collinear fields as fluctuations on top of a soft background.⁷

2.2.4 Light-front multipole expansion

Due to the power-counting of their momenta (and thus coordinate arguments), products containing both soft and collinear fields are not homogeneous in λ . In Fourier space, one finds, for example,

$$\varphi_{c}(x)\varphi_{s}(x) = \int \frac{d^{4}p}{(2\pi)^{4}} \frac{d^{4}k_{s}}{(2\pi)^{4}} e^{-i(p+k_{s})\cdot x} \tilde{\varphi}_{c}(p)\tilde{\varphi}_{s}(k_{s}).$$
(33)

Here, the product in the exponent reads

$$(p+k_s)\cdot x = \frac{1}{2}\underbrace{(n_+p+n_+k_s)}_{1+\lambda^2}n_-x + \underbrace{(p_\perp+k_{s\perp})}_{\lambda+\lambda^2}\cdot x_\perp + \frac{1}{2}\underbrace{(n_-p+n_-k_s)}_{\lambda^2+\lambda^2}n_+x.$$
 (34)

Only the combination

$$n_{-}p + n_{-}k_{s} \sim \lambda^{2} \tag{35}$$

scales homogeneously as $\mathcal{O}(\lambda^2)$, while the $k_{s\perp}$ and n_+k_s components are suppressed with respect to p_{\perp} and n_+p . The physical reason for this is that soft fields can only resolve large distances

$$x_{-}^{\mu} \equiv n_{+} x \frac{n_{-}^{\mu}}{2}, \qquad (36)$$

whereas collinear fields fluctuate also over smaller distances. Therefore, one must expand the exponential in (33) as

$$e^{-i(p+k_s)\cdot x} = e^{-i(p+n_-k_s\frac{n_+}{2})\cdot x} \left(1 - ik_{s\perp} \cdot x_{\perp} - \frac{i}{2}n_+k_sn_-x + \dots\right), \quad (37)$$

keeping only the homogeneous soft momentum n_k in the exponent. This expansion is equivalent to the *light-front multipole expansion* [5, 6]

$$\varphi_s(x) = \varphi_s(x_-) + (x - x_-)^{\mu} \left[\partial_{\mu} \varphi_s \right](x_-) + \mathscr{O}(\lambda^2 \varphi_s).$$
(38)

of the soft field on the left-hand side of (33). This expansion of soft fields about the large coordinate n_+x of collinear fields must be applied whenever soft and collinear fields appear in a product. It generates an infinite tower of sub-leading in λ soft-collinear interactions, which precisely reproduces the expansion in the small soft momenta $n_{i+}k_s, k_{s\perp}$ of the momentum-space amplitude. One may note the similarities to the standard multipole expansion for spatially localised systems, which also appears in non-relativistic effective theories.

⁷ Since the background field method may be more familiar in gauge theories, we invite the reader to compare the gauge transformations in SCET for QCD [6] with those for gravity discussed here.

This has an important implication for the gauge symmetry, since the gauge transformation (32) of $h_{\mu\nu}$ contains products of collinear and soft fields. Furthermore, the soft gauge transformation $U_s(x)$ of collinear fields is a product of soft and collinear fields, which still mixes different powers in λ , and therefore must be multipoleexpanded. For example, the soft gauge transformation of the collinear matter field follows from (20) and reads

$$\varphi_c(x) \to U_s(x)\varphi_c(x) = U_s(x_-)\varphi_c(x) + x_{\perp}^{\alpha} \left[\partial_{\alpha} U_s\right](x_-)\varphi_c(x) + \dots$$
(39)

The additional terms beyond the leading one imply that the transformation mixes different orders in λ in a way that is incompatible with the multipole expansion of the Lagrangian. To alleviate this, one needs to find a way to obtain collinear fields that have a *homogeneous* gauge transformation, respecting the multipole expansion. The resolution of this subtlety is technically quite involved, but intuitively very simple: recall that the multipole expansion is necessary, because soft fields cannot resolve the small-scale fluctuations of collinear modes. However, in the naive split $g_{\mu\nu}(x) = g_{s\mu\nu}(x) + \kappa h_{\mu\nu}(x)$, one implicitly assumes that this is possible, since the full $s_{\mu\nu}(x)$ appears inside $g_{s\mu\nu}(x)$. Therefore, the soft gauge symmetry, which is the symmetry of the background metric $g_{suv}(x)$, is constructed with respect to the wrong background field. Instead, one needs to identify the appropriate background field $\hat{g}_{s\mu\nu}(x_{-})$, whose constituent soft fields can only depend on x_{-}^{μ} . In other words, these soft background fields live only on the classical trajectory of the collinear particles. This proper background field comes with its residual transformation, and it is this residual transformation that is "homogeneous" in λ , respecting the multipole expansion.

A trace of this homogeneous background field can already be seen in the leadingpower Lagrangian. By performing the multipole expansion and keeping only the leading terms in (5), one obtains the soft-collinear interaction

$$\mathscr{L}^{(0)} = -\frac{\kappa}{8} n_{-}^{\mu} n_{-}^{\nu} s_{\mu\nu}(x_{-}) \partial_{+} \varphi_{c} \partial_{+} \varphi_{c} \,. \tag{40}$$

When inserting this expression into a soft-emission diagram, the combination $\varphi_c \partial_+ \varphi_c$ generates the eikonal propagator $\frac{in_+p}{2p\cdot k} = \frac{i}{n_-k}$. Therefore, the leading-power interaction from the homogeneous background field $s_{--}(x_-)$ yields a term proportional to $\varepsilon_{--}p_+$. Finding the full expression of the soft background field is more involved, and is explained in Section 4. Once this proper background field is identified, it will be straightforward to construct the EFT systematically to all orders.

2.2.5 Basic features of the effective Lagrangian

The effective theory then takes a simple structure. The Lagrangian \mathscr{L}_{SCET} splits into a soft-collinear and a purely-soft Lagrangian

$$\mathscr{L}_{\text{SCET}} = \sum_{i} \mathscr{L}_{c_i} \left[h_{i\mu\nu}(x), s_{\mu\nu}(x_{i-}) \right] + \mathscr{L}_s \left[s_{\mu\nu}(x) \right].$$
(41)

In the collinear part, there is a sum over all collinear sectors defined by the directions n_{i-}^{μ} of the jets. This sum arises because one needs a *hard* scattering to generate particles of different collinear sectors. Since the Lagrangian does not describe hard scattering, there is no direct interaction vertex between different collinear sectors. Instead, these processes are allocated to the so-called *N*-jet operators, which generate energetic and soft particles from hard scattering.

The soft-collinear Lagrangian contains also purely-collinear terms. This purelycollinear Lagrangian, as well as the purely-soft Lagrangian, is then completely equivalent to the original full theory (in weak-field expansion). This is due to the fact that if only one scale is present, e.g. by only considering purely-collinear or purely-soft modes without any external sources, then there is no Lorentz-invariant notion of soft or collinear. One could simply perform a Lorentz boost and collinear modes would become soft, and vice-versa. It is the presence of a source that provides meaning to the notion of soft and collinear in the first place.

Therefore, all the non-trivial physics is contained inside the soft-collinear interaction vertices as well as the "sources", which are described by *N*-jet operators. These vertices stem from the terms in the collinear Lagrangian that are covariant with respect to the non-trivial soft background.

From this perspective, the following structure of the theory arises, shown in Fig. 3. The collinear sector *i* is constructed to be covariant with respect to a soft background metric denoted by $\hat{g}_{s_i\mu\nu}(x_{i-})$, which is constructed from the dynamical soft field $\eta_{\mu\nu} + \kappa s_{\mu\nu}(x)$, restricted to the classical light-like trajectories x_{i-}^{μ} of the energetic particles. Since due to the multipole expansion, collinear fields interact with soft fields at x_{i-}^{μ} only, these are effectively $i = 1, \ldots, N$ separate soft gauge symmetries $U_s(x_{i-})$. Note that the interactions with the soft field are blind to the non-locality of the collinear sector in the n_{i+}^{μ} direction since $(x^{\mu} + tn_{i+}^{\mu})_{-} = x_{-}^{\mu}$. On the other hand, the soft fields have self-interactions, which pervade all of space-time, and are controlled by the soft Lagrangian \mathscr{L}_s .

2.3 Gravity vs QCD: a comparison

At this point, it is instructive to compare the gravitational situation to the gaugetheory one, where the SCET construction is well understood. The basic aspects of the construction, as discussed in Section 2.2, are the same for both theories, replacing the background field $\hat{g}_{s\mu\nu}(x_-) \rightarrow n_-A_s(x_-)$.

The first main difference lies in the nature of the "full theory" itself. In QCD, the starting point for the EFT construction is Yang-Mills theory, a renormalisable field theory. Contrast this with gravity: here, the natural starting point, Einstein-Hilbert theory, is not renormalisable. Instead, one takes the effective action (7), up to a desired order in the loop-expansion (or curvature, respectively). In addition, one then performs the weak-field expansion, since the relevant degree of freedom is the fluctuation $h_{\mu\nu}$, the graviton field. Therefore, the "full theory" underlying SCET gravity is the weak-field expansion of an effective extension of the Einstein-Hilbert



Fig. 3 An intuitive picture of the form of SCET. The soft modes (red) are described by the purelysoft Lagrangian \mathcal{L}_s , which takes the same form as the full theory. Each collinear sector (blue) is described by its own Lagrangian \mathcal{L}_{c_i} , which contains purely-collinear and soft-collinear interactions (pink). These soft-collinear terms are covariant with respect to a homogeneous background field $g_{i}^{\mu\nu}(x_{i-})$, living only on the classical trajectory x_{i-}^{μ} of the collinear particles, and describe soft-collinear physics to all orders.

action, and thus already defined as an infinite series in κ that must be truncated at some order. This truncated theory is then expanded in λ . The λ -expansion shares many features with the κ -expansion but is not identical to it.

Another major difference is the nature of the underlying gauge symmetry. In QCD, the gauge symmetry is purely internal, and it has a colour charge with a corresponding generator $t^a \sim \lambda^0$ that is unrelated to the kinematics that define the SCET expansion. In gravity, however, the charge is related to momenta P^{μ} and therefore the gauge symmetry is connected to kinematics. In particular the collinear momentum components have different λ -scaling (23). This implies that the gauge transformation parameters in gravity also have non-trivial power-counting, and are not homogeneous in λ . For example, a collinear matter field transforms under an infinitesimal collinear transformation as

$$\varphi_c \to \varphi_c - \kappa \varepsilon_c^{\alpha} \partial_{\alpha} \varphi_c \,, \tag{42}$$

where the second term is suppressed by $\mathcal{O}(\lambda)$ relative to the first due to the scaling of ∂_{μ} and the gauge parameter (see (45) below). Therefore, gauge transformations mix different powers of λ , and no homogeneous scaling can be achieved when manifest gauge-invariance is imposed. An object can either be homogeneous in λ , or gauge-invariant, but never both at the same time.

These formal differences aside, one can now take a look at the physical content of both theories. In QCD, the collinear gluon field A_c contains two physical and two unphysical components. The large component $n_+A_c \sim \lambda^0$ as well as the small $n_{-}A_{c} \sim \lambda^{2}$ are unphysical, while the transverse components $A_{c\perp} \sim \lambda$ are physical. The large component n_+A_c is problematic: if this field was an allowed building block, one could add arbitrarily many such fields to any operator while keeping its λ -counting fixed. Thus, there would not exist a finite operator basis, and the powercounting would be meaningless, since one would have to perform an infinite number of matching computations already at leading power. This is alleviated by noting that in light-cone gauge, $n_+A_c = 0$, thus it is a gauge-artifact. Therefore, one introduces a Wilson line W_c (definition in (47) below), which controls these n_+A_c to all orders. These Wilson lines are homogeneous in λ , but an infinite series in the gauge coupling g_s . In gravity, a similar situation arises, but worse with respect to the powercounting: the collinear graviton $h_{\mu\nu}$ contains modes $h_{++} \sim \lambda^{-1}$, $h_{+\perp} \sim \lambda^0$, where h_{++} is even power-counting enhanced. Clearly these must be controlled. Therefore, one employs a similar concept as in QCD, by introducing the analogue of the collinear Wilson line. In this way, the unphysical $h_{+\mu}$ components can be controlled to all orders. Just as in QCD, this "Wilson line" is an infinite series in the coupling κ . Due to the aforementioned inhomogeneity of the gauge symmetry, however, this implies that these gravitational "Wilson lines" are also an infinite series in λ , and no longer homogeneous. These "Wilson lines" can be used to implement a covariant version of light-cone gauge, in the sense that one defines gauge-invariant composite objects that satisfy the light-cone gauge properties.

The suppressed unphysical degrees of freedom, $n_{-}A_c$ in QCD and $h_{-\perp}, h_{--}$ in gravity, can be eliminated using the equations of motion. This leaves only the two transverse polarisations $A_{c\perp} \sim \lambda$ and $h_{\perp\perp} \sim \lambda$ as physical degrees of freedom in the effective theory. Since they count with a positive power of λ , a sensible operator basis exists.

The soft sector is slightly different from the collinear one. Here, one first needs to identify the appropriate homogeneous background field, and then one organises the sub-leading terms in gauge-covariant objects. The background field comes with a covariant derivative, which can be eliminated in the sources and the sub-leading Lagrangian terms using the equations of motion. For the operator basis, only the subleading gauge-covariant objects are relevant. In gauge theory, this gauge-covariant object is the field-strength tensor $F_{s\mu\nu} \sim \lambda^4$, which is the first derivative of the gluon field. In gravity, the first sensible gauge-covariant object is the Riemann tensor, $R^{\mu}_{\ \nu\alpha\beta} \sim \lambda^{6}$, which is the second derivative of the metric fluctuation. These gauge-covariant objects can appear in the sources and mediate process-dependent soft emissions. Therefore, already at this stage, one can anticipate that in a softemission process, there are universal terms and the non-universality in gravity is more strongly suppressed compared to gauge theory. The previously discussed features are summarised in Table 1. While there are essential differences between both theories, the construction of effective theories proceeds similarly. This will guide the following sections.

	QCD	Gravity
gauge symmetry	<i>SU</i> (3)	$\operatorname{Diff}(M)$
gauge charge	$t^a \sim \lambda^0$	$P^{\mu} \sim (\lambda^0, \lambda, \lambda^2)$
dimensionful coupling	no	yes
fundamental degree of freedom	$A_{\mu} \sim p_{\mu}$	$h_{\mu u}\simrac{p_{\mu}p_{ u}}{\lambda}$
unsuppressed components	$n_+A_c \sim 1$	$h_{++}\sim\lambda^{-1},h_{+\perp}\sim1$
physical degrees of freedom	$A_{c\perp}\sim\lambda$	$h_{\perp\perp}\sim\lambda$
redundant degrees of freedom	$nA_c \sim \lambda^2$	$h_{\perp -} \sim \lambda^2, h_{} \sim \lambda^3$
field-strength / curvature	$F_{\mu\nu}\sim\partial A$	$R^{\mu}_{\ \nulphaeta}\sim\partial^{2}h$

 Table 1 A comparison of the main features of gauge-theory (QCD) and gravity from the SCET perspective.

3 Collinear gravity

It is convenient to first consider only collinear modes in order to familiarise oneself with the construction. The purely-collinear EFT for gravity is obtained after splitting a generic field $\phi_J = \phi_{J,h} + \sum_i \phi_{J,c_i}$. After integrating out the hard modes, there are no left-over interactions of the collinear fields ϕ_{J,c_i} of different sectors, as the sum of the collinear momenta belonging to different directions has hard scaling, and corresponds to off-shell degrees of freedom which have already been eliminated. The Lagrangian is simply the sum

$$\mathscr{L}_{\text{SCET}} = \sum_{i} \mathscr{L}_{c_i} \,. \tag{43}$$

For this reason it is possible to focus only on a single collinear direction. To simplify the notation, in the following the subscripts c, i are omitted. In addition to the Lagrangian, the theory contains so-called *currents* (aka "sources", or "*N*-jet operators"), which contain products of fields in more than one collinear direction. Their matching coefficients absorb the hard modes that have been integrated out. These are discussed in more detail in Section 5. Here, we focus on the effective Lagrangian.

At this stage, the gauge symmetry consists of *N*-copies of collinear gauge symmetry, such that each collinear sector transforms under its own gauge symmetry and is invariant with respect to all the remaining collinear symmetries. Collinear gauge transformations must not distort the scaling of the collinear gauge fields. In contrast with SCET QCD, in SCET gravity, the gauge transformation parameter ε^{μ} acquires λ -scaling. Enforcing homogeneity of the infinitesimal gauge transformation

$$h_{\mu\nu} \to h_{\mu\nu} - \partial_{\mu} \varepsilon_{\nu} - \partial_{\nu} \varepsilon_{\mu} \sim \mathscr{O}(h_{\mu\nu}) , \qquad (44)$$

 ε^{μ} scales as

$$n_{+}\varepsilon \sim \frac{1}{\lambda}, \quad n_{-}\varepsilon \sim \lambda, \quad \varepsilon^{\mu_{\perp}} \sim 1.$$
 (45)

Note that ε^{μ} scales as a small translation λx^{μ} . Beyond the linear order the transformation of $h_{\mu\nu}$ takes the form

$$h'_{\mu\nu} = h_{\mu\nu} - \partial_{\mu}\varepsilon_{\nu} - \partial_{\nu}\varepsilon_{\mu} - \kappa \left[\partial_{\mu}\varepsilon^{\alpha}h_{\alpha\nu} - \partial_{\nu}\varepsilon^{\alpha}h_{\alpha\mu} - \varepsilon^{\alpha}\partial_{\alpha}h_{\mu\nu} \right] + \partial_{\mu}\varepsilon^{\alpha}\partial_{\alpha}\varepsilon_{\nu} + \partial_{\nu}\varepsilon^{\alpha}\partial_{\alpha}\varepsilon_{\mu} + \partial_{\mu}\varepsilon_{\alpha}\partial_{\nu}\varepsilon^{\alpha} + \varepsilon^{\alpha}\partial_{\alpha}(\partial_{\mu}\varepsilon_{\nu} + \partial_{\nu}\varepsilon_{\mu}) + \mathcal{O}(\lambda^{3}).$$
(46)

Collinear gauge-invariance is not only required from a formal point of view, but also ensures that the EFT power-counting is meaningful, i.e. one cannot generate an infinite number of operators with the help of the unsuppressed collinear fields. For this reason, it is beneficial to introduce the concept of *gauge-invariant building blocks*. An arbitrary current contains fields in multiple collinear directions. Since each sector has its own gauge symmetry, the invariance of the complete operator under each collinear symmetry implies that each sector is separately gauge-invariant. This is automatically achieved if the operator is built from the gauge-invariant building blocks defined as the collinear field dressed with "Wilson lines", chosen such that the building block is a gauge-singlet that is always suppressed by at least one power of λ . In gauge theory, this idea leads to the concept of the collinear Wilson line W_c

$$W_c(x) = P \exp\left(ig_s \int_{-\infty}^0 ds \, n_+ A_c(x+sn_+)\right),\tag{47}$$

which fulfills the following identity

$$W_c^{\dagger} n_+ D_c W_c = n_+ \partial , \qquad (48)$$

such that the collinear gauge-invariant gluon building block is $\mathscr{A}_{\mu} = \frac{1}{g_s} W_c^{\dagger} [iD_{c\mu} W_c]$. This leads to the notion of "covariant light-cone gauge", since the composite operator \mathscr{A}_{μ} built from $A_{c\mu}$ satisfies $n_+ \mathscr{A} = 0$ in *any* gauge.

In analogy to QCD, where the collinear-gauge invariant gluon field satisfies $\mathscr{A}_{+} = 0$, in SCET gravity one constructs the manifestly gauge-invariant graviton building block $\mathfrak{h}_{\mu\nu}$ [10, 24, 34], which depends on the collinear field $h_{\mu\nu}$, and satisfies $\mathfrak{h}_{\mu+} = 0$, that is, it coincides with the elementary graviton field in light-cone gauge. Just like in QCD, where the collinear Wilson line connects the collinear gluon field to the gauge invariant object, in gravity, the collinear graviton field is related to the gauge invariant graviton building block via

$$\eta_{\mu\nu} + \kappa \mathfrak{h}_{\mu\nu}(x) = W^{\alpha}_{\ \mu} W^{\beta}_{\ \nu} \left[W^{-1}_c g_{\alpha\beta}(x) \right], \tag{49}$$

where $g_{\alpha\beta}(x) = \eta_{\alpha\beta} + \kappa h_{\alpha\beta}(x)$ and the gravitational collinear "Wilson line" is

$$W_{c}^{-1} = T_{\theta_{c}}[h] = 1 + \kappa \theta_{c}^{\alpha} \partial_{\alpha} + \frac{\kappa^{2}}{2} \theta_{c}^{\alpha} \theta_{c}^{\beta} \partial_{\alpha} \partial_{\beta} + \mathscr{O}(\theta_{c}^{3}), \qquad (50)$$

19

with parameter $\theta_c[h] \equiv \theta_c[h_{\mu\nu}(x)]$ chosen such that, given $h_{\mu\nu}$, the invariant field $\mathfrak{h}_{\mu\nu}$ satisfies $\mathfrak{h}_{\mu+} = 0$, and $W^{\alpha}_{\ \mu}$ is the Jacobian for the transformation (50), defined as in (14). The above translation corresponds to the transformation to the coordinate system, in which the metric fluctuation satisfies light-cone gauge. Note, however, that we do not actually fix the gauge. Rather, W_c is employed to define the composite operator $\mathfrak{h}_{\mu\nu}(x)$ built from the elementary field $h_{\mu\nu}$ for which no special coordinate system or gauge-fixing is assumed. The "Wilson line" transforms as [24]

$$W_c^{-1} \to W_c^{-1} U^{-1}(x),$$
 (51)

and this transformation precisely cancels the collinear gauge transformation of $h_{\mu\nu}$ in (49). It is also used to define the collinear gauge-invariant matter field χ_c ,

$$\boldsymbol{\chi}_c = \begin{bmatrix} W_c^{-1} \boldsymbol{\varphi}_c \end{bmatrix} = \boldsymbol{\varphi}_c + \mathscr{O}(\lambda \boldsymbol{\varphi}_c) \,. \tag{52}$$

Note that $W_c^{-1} = 1 + \mathcal{O}(\lambda)$ in stark contrast to (47), which is closely related to the absence of purely-collinear gravitational interactions at leading power in λ as will be seen below.

The explicit construction of the collinear "Wilson line" requires determining the parameter $\theta_c^{\mu}[h]$. The closed-form version does not exist in gravity, but instead it can be defined perturbatively in the weak-field expansion

$$\boldsymbol{\theta}_c^{\mu} = \boldsymbol{\theta}_c^{\mu(0)} + \boldsymbol{\theta}_c^{\mu(1)} + \dots$$
 (53)

Then, one needs to perform the expansion of (49),

$$\mathfrak{h}_{\mu\nu} = h_{\mu\nu} + \partial_{\mu}\theta_{c\nu} + \partial_{\nu}\theta_{c\mu} + \mathscr{O}(\theta_c^2, \theta_c h), \qquad (54)$$

and insert (53). Requiring $\mathfrak{h}_{\mu+} = 0$ determines

$$\theta_{c\mu}^{(0)} = -\frac{1}{n_{+}\partial} \left(h_{\mu+} - \frac{1}{2} \frac{\partial_{\mu}}{n_{+}\partial} h_{++} \right).$$
(55)

The sub-leading (non-linear) terms can be obtained iteratively [10]. The inverse derivative is defined as

$$\frac{1}{in_+\partial + i\varepsilon}f(x^{\mu}) = -i\int_{-\infty}^0 ds f(x^{\mu} + sn_+^{\mu}).$$
(56)

Inserting back the leading term in the expansion of θ_c (55) into $\mathfrak{h}_{\mu\nu}$ (54) leads to the explicit formula for the gauge-invariant graviton building block expressed in terms of the collinear graviton

$$\mathfrak{h}_{\mu\nu} = h_{\mu\nu} - \frac{\partial_{\mu}}{n_{+}\partial} \left(h_{\nu+} - \frac{1}{2} \frac{\partial_{\nu}}{n_{+}\partial} h_{++} \right) - \frac{\partial_{\nu}}{n_{+}\partial} \left(h_{\mu+} - \frac{1}{2} \frac{\partial_{\mu}}{n_{+}\partial} h_{++} \right) + \mathscr{O}(\lambda h_{\mu\nu}),$$
(57)

which resembles the analogous expression for the gauge-invariant collinear gluon field, 8

$$\mathscr{A}_{c\mu_{\perp}} = \frac{1}{g_s} W_c^{\dagger} \left[i D_{c\mu_{\perp}} W_c \right] = A_{c\mu_{\perp}} - \frac{\partial_{\mu_{\perp}}}{n_{+} \partial} A_{c+} + \mathscr{O}(g_s A_{c\mu_{\perp}}) \,. \tag{58}$$

Having understood the basic building blocks of the theory, it is straightforward to derive the collinear Lagrangian. The first step involves performing the weak-field expansion of (5). Then one re-expresses the collinear fields in terms of the gaugeinvariant building blocks by inserting the gravitational collinear "Wilson line" (50). Since the "Wilson line" corresponds to a collinear gauge transformation, one can simply put $h_{\mu\nu} \rightarrow h_{\mu\nu}$ and $\varphi_c \rightarrow \chi_c$. Then, one can set $h_{\mu+} = 0$. Finally, grouping the terms by their order in the λ -expansion yields

$$\mathscr{L}^{(0)} = \frac{1}{2} \partial_{\mu} \chi_c \partial^{\mu} \chi_c \,, \tag{59}$$

$$\mathscr{L}^{(1)} = -\frac{\kappa}{2} (\mathfrak{h}_{\mu\nu} \partial^{\mu} \chi_{c} \partial^{\nu} \chi_{c} - \frac{1}{2} \mathfrak{h} \partial_{\alpha} \chi_{c} \partial^{\alpha} \chi_{c}), \qquad (60)$$

$$\mathscr{E}^{(2)} = \frac{\kappa^2}{2} \left(\mathfrak{h}^{\mu\alpha} \mathfrak{h}^{\nu}_{\alpha} - \frac{1}{2} \mathfrak{h} \mathfrak{h}^{\mu\nu} + \frac{1}{8} (\mathfrak{h}^2 - 2\mathfrak{h}^{\alpha\beta} \mathfrak{h}_{\alpha\beta}) \eta^{\mu\nu} \right) \partial_{\mu} \chi_c \partial_{\nu} \chi_c \,. \tag{61}$$

The leading-power Lagrangian $\mathscr{L}^{(0)}$ describes a free theory of gauge-invariant building blocks. It follows from this simple observation that there cannot be collinear singularities in gravity. Interactions with collinear gravitons start at sub-leading power and all terms are manifestly gauge-invariant.

The Lagrangian above is expressed entirely in terms of the gauge-invariant building blocks χ_c , $\mathfrak{h}_{\mu\nu}$. This form makes it clear that there is a well-defined expansion in λ , since the metric components unsuppressed in λ do not appear due to $\mathfrak{h}_{\mu+} = 0$. Since the Lagrangian is collinear gauge-invariant, it takes the same form when expressed in terms of the original fields φ_c , $h_{\mu\nu}$. However, since h_{++} and $h_{+\perp}$ do not vanish, this form hides the gauge cancellations that occur between the unsuppressed metric components and lacks manifest power counting.

The sub-leading components $\mathfrak{h}_{\mu-}$ can be eliminated from the Lagrangian by using equations of motion. To their respective leading order in λ , one finds [10]

$$\mathfrak{h}_{\mu_{\perp}-} = -2\frac{\partial^{\alpha_{\perp}}}{\partial_{+}}\mathfrak{h}_{\mu_{\perp}\alpha_{\perp}} + \mathscr{O}(\lambda^{3}), \qquad (62)$$

$$\mathfrak{h}_{--} = 4 \frac{\partial^{\alpha_{\perp}} \partial^{\beta_{\perp}}}{\partial_{+}^{2}} \mathfrak{h}_{\alpha_{\perp}\beta_{\perp}} + \mathscr{O}(\lambda^{4}), \qquad (63)$$

where the first equation arises from the field equation for $\mathfrak{h}_{\mu_{\perp}-}$, and the second from tracing the one for $\mathfrak{h}_{\mu_{\perp}\nu_{\perp}}$. The Lagrangian is linear in \mathfrak{h}_{--} , and the field equation for \mathfrak{h}_{--} results in the trace constraint

⁸ Despite the fact that $W_c^{-1} = 1 + \mathcal{O}(\lambda)$, $\mathfrak{h}_{\mu\nu}$ differs from $h_{\mu\nu}$ at $\mathcal{O}(\lambda^0 h_{\mu\nu})$ due to the Jacobian terms in (49) multiplying the $\eta_{\alpha\beta}$ term in $g_{\alpha\beta}(x)$.

Martin Beneke, Patrick Hager and Robert Szafron

$$\mathfrak{h} = \frac{\kappa}{2} \left(\mathfrak{h}_{\alpha_{\perp}\beta_{\perp}} \mathfrak{h}^{\alpha_{\perp}\beta_{\perp}} - \frac{1}{\partial_{+}^{2}} \left[\partial_{+} \mathfrak{h}_{\alpha_{\perp}\beta_{\perp}} \partial_{+} \mathfrak{h}^{\alpha_{\perp}\beta_{\perp}} - \partial_{+} \chi_{c} \partial_{+} \chi_{c} \right] \right) + \mathscr{O}(\lambda^{3}), \quad (64)$$

which shows that \mathfrak{h} counts as $\mathscr{O}(\lambda^2)$. The trace terms in $\mathscr{L}^{(2)}$ can therefore be dropped at this order, while the second interaction term in $\mathscr{L}^{(1)}$ is $\mathscr{O}(\lambda^2)$. These simplifications can be made manifest by expressing the Lagrangian in terms of $\mathfrak{h}_{\mu_{\perp}\nu_{\perp}}$ only.

The collinear Einstein-Hilbert Lagrangian for the graviton self-interactions, as well as for the scalar field with φ^4 self-interactions are discussed in detail in [10]. Interestingly, the trilinear graviton self-interaction when expressed in terms of $\mathfrak{h}_{\mu_{\perp}\nu_{\perp}}$ (equivalently, light-cone gauge) has a manifest double-copy structure at the Lagrangian level [8].

4 Soft-collinear gravity

The full soft-collinear EFT can be constructed separately for each collinear sector as seen from (41), therefore the index i will be dropped. The first step is to introduce the soft and collinear modes. One way to implement the mode split consistently is to treat the collinear modes as small-distance fluctuations on top of the slowly-varying soft background. For the metric field, the decomposition takes the form

$$g_{\mu\nu} = \eta_{\mu\nu} + \kappa h_{\mu\nu} + \kappa s_{\mu\nu} \equiv g_{s\mu\nu} + \kappa h_{\mu\nu} \,, \tag{65}$$

where by definition all components of the soft field $s_{\mu\nu}$ have isotropic scaling

$$s_{\mu\nu} \sim \lambda^2$$
. (66)

It is convenient to work with the soft background in terms of a soft metric tensor $g_{s\mu\nu}$ to leverage the geometric intuition. Therefore, this first step is fully equivalent to a standard weak-field expansion in $h_{\mu\nu}$ about a non-trivial dynamic background $g_{s\mu\nu}$.

The gauge symmetry is modified by this decomposition, and changes into the gauge symmetry of the background (the soft gauge symmetry), and the gauge symmetry of the collinear fluctuations, which are covariant with respect to this background. The soft gauge symmetry is the standard diffeomorphism applied to the slowly varying background field. The transformations of the soft and collinear fluctuations read

$$h_{\mu\nu} \rightarrow \left[U_s \left(U_{s\mu}^{\ \alpha} U_{s\nu}^{\ \beta} h_{\alpha\beta} \right) \right],$$

$$\kappa s_{\mu\nu} \rightarrow \left[U_s (U_{s\mu}^{\ \alpha} U_{s\nu}^{\ \beta} (\eta_{\alpha\beta} + \kappa s_{\alpha\beta}) \right] - \eta_{\mu\nu}.$$
(67)

It is understood that the *x*-dependence of $U_s(x)$ is the one of a soft field with variations over distances $\mathcal{O}(1/\lambda^2)$. The soft fluctuation $s_{\mu\nu}$ has the standard transformation of a metric perturbation, whereas the transformation of the collinear graviton

22

 $h_{\mu\nu}$ is the one of a standard rank-2 tensor, i.e. covariant like an ordinary matter field. Note that the soft metric tensor $g_{s\mu\nu}$ also transforms like a rank-2 tensor (13)

$$g_{s\mu\nu} \to \left[U_s \left(U_{s\mu}^{\ \alpha} U_{s\nu}^{\ \beta} g_{s\alpha\beta} \right) \right]. \tag{68}$$

The interpretation of these transformations is clear: the background geometry is described by $g_{s\mu\nu}$, which transforms like an ordinary metric tensor, and $s_{\mu\nu}$ inherits a non-linear transformation from it. From the soft point of view, the collinear gravitons transform just like *ordinary matter fields*, i.e. like a standard rank-2 tensor, and *not* non-linearly like a metric perturbation. Therefore, from the soft perspective, any collinear field, regardless if gauge or matter, transforms like a matter field, i.e. according to its tensor representation.

The collinear gauge transformations are the transformations of the fields defined on the background metric and were already introduced in (32). They can be cast in the form

$$\kappa h_{\mu\nu} \rightarrow \left[U_c \left(U_{c\mu}^{\ \alpha} U_{c\nu}^{\ \beta} \kappa h_{\alpha\beta} \right) \right] + \left[U_c \left(U_{c\mu}^{\ \alpha} U_{c\nu}^{\ \beta} g_{s\alpha\beta} \right) \right] - g_{s\mu\nu} , \qquad (69)$$

$$g_{s\mu\nu} \rightarrow g_{s\mu\nu} ,$$

which renders the transformation of $h_{\mu\nu}$ reminiscent of an ordinary gauge-field transformation with a covariant and an inhomogeneous term. The fluctuation $h_{\mu\nu}$ has a non-linear transformation, similar to the standard weak-field expansion around flat space (20), but with respect to the soft background $g_{s\mu\nu}$, which appears in the inhomogeneous term instead of $\eta_{\mu\nu}$. Observe here that the soft background is necessarily invariant under the collinear transformations, since $U_c(x)$ is a collinear field.

Both gauge symmetries, soft and collinear, have the property that the full metric tensor $g_{\mu\nu} = g_{s\mu\nu} + \kappa h_{\mu\nu}$ transforms like the standard metric tensor. The transformations of matter fields follow the same logic. They can be found in [10].

However, as already discussed in Section 2.2.4, the transformations do not satisfy completely homogeneous λ -scaling, since they contain multiplications of collinear and soft fields at point *x*. The following construction identifies the background metric and the collinear field redefinition that renders the gauge symmetries compatible with a manifestly gauge-covariant form of the light-front multipole expansion.

4.1 Riemann normal coordinates

To gain some intuition for this procedure in the gravitational context, a simpler example is considered first: the static multipole expansion around the space-time point x = 0. The metric tensor in this theory should be multipole-expanded as

$$g_{\mu\nu}(x) = g_{\mu\nu}(0) + x^{\alpha} \left[\partial_{\alpha}g_{\mu\nu}\right](0) + \frac{1}{2}x^{\alpha}x^{\beta} \left[\partial_{\alpha}\partial_{\beta}g_{\mu\nu}\right](0) + \mathscr{O}(x^{3}).$$
(70)

In gauge theories, the corresponding expansion of $A_{\mu}(x)$ is conveniently performed using Fock-Schwinger gauge $x^{\mu}A_{\mu}(x) = 0$. This gauge condition sets $A_{\mu}(0) = 0$, and the higher-order terms in the Taylor-expansion are expressed in terms of the field-strength tensor and its derivatives. Since the resulting expansion is manifestly gauge-covariant, the gauge-fixing can easily be removed at the end. This technique has found numerous applications [33] to non-perturbative techniques in QCD, in which case short-distance fluctuations are treated in the background of the soft QCD vacuum.

In gravity, the analogous gauge corresponds to the familiar Riemann normal coordinates. The "gauge condition" reads

$$x^{\mu}x^{\nu}\Gamma^{\alpha}_{\ \mu\nu}(x) = 0. \tag{71}$$

Compared to Fock-Schwinger gauge $x^{\mu}A_{\mu}(x) = 0$, one immediately notices that in gravity, the gauge condition starts at second order in *x*, and restricts the derivative of the metric tensor at x = 0, but not $g_{\mu\nu}(0)$, since $\Gamma \sim \partial g$. Indeed, the effect of Riemann normal coordinates is to set the first derivative $[\partial_{\alpha}g_{\mu\nu}](0) = 0$, and to express the higher derivatives via the Riemann tensor and its derivatives at x = 0. The metric tensor near x = 0 is then expressed as

$$\tilde{g}_{\mu\nu}(x) = g_{\mu\nu}(0) - \frac{1}{3} x^{\alpha} x^{\beta} R_{\alpha\mu\beta\nu}(0) + \mathcal{O}(x^3).$$
(72)

The gauge condition (71) does not fix the diffeomorphism symmetry completely, but only up to *global* linear transformations $A_{\mu}^{\alpha} \in GL(1,3)$,

$$x^{\mu} \to A_{\alpha}{}^{\mu} x^{\alpha} \,. \tag{73}$$

Using this residual symmetry, one can transform the leading term $g_{\mu\nu}(0)$ to the Minkowski metric $\eta_{\mu\nu}$, and the metric tensor in these coordinates reads

$$\tilde{g}_{\mu\nu}(x) = \eta_{\mu\nu} - \frac{1}{3} x^{\alpha} x^{\beta} R_{\alpha\mu\beta\nu}(0) + \mathcal{O}(x^3), \qquad (74)$$

which is the standard form of the metric in Riemann normal coordinates. The remaining residual symmetries of the full diffeomorphisms are now the ones of the Minkowski metric $\eta_{\mu\nu}$, i.e. global Lorentz transformations.

The lesson of this section is the following: using Riemann normal coordinates, one can express the multipole expansion of the metric tensor in terms of the metric at the origin and Riemann tensor terms. Then, one can exploit the residual symmetry to diagonalise this metric and change it to the flat-space one at x = 0. The result is a covariant multipole expansion that features global Poincaré transformations as residual symmetries.

4.2 Fixed-line normal coordinates

The above discussion motivates a generalisation of the Riemann normal coordinates adapted to the situation where the physical system is not localised around a spacetime point, but around the light-like trajectory $x_{-}^{\mu} = n_{+}x\frac{n_{-}^{\mu}}{2}$, which allows for the light-front multipole expansion of soft fluctuations. The metric tensor is now expanded as

$$g_{s\mu\nu}(x) = g_{s\mu\nu}(x_{-}) + x_{\perp}^{\alpha} \left[\partial_{\alpha} g_{s\mu\nu} \right](x_{-}) + \frac{1}{2} n_{-}x \left[n_{+} \partial g_{s\mu\nu} \right](x_{-}) + \frac{1}{2} x_{\perp}^{\alpha} x_{\perp}^{\beta} \left[\partial_{\alpha} \partial_{\beta} g_{s\mu\nu} \right](x_{-}) + \mathcal{O}(\lambda^{3} g_{s\mu\nu}),$$
(75)

which differs from the previous one (70) in two aspects: first, the expansion is only in the directions orthogonal to the light-like line x_{-}^{μ} , and second, the coefficients in the Taylor-expansion are dynamical fields and depend on x_{-}^{μ} . In gauge theory, the generalisation of Fock-Schwinger gauge is "fixed-line" gauge [5] $(x-x_{-})^{\mu}A_{\mu}(x_{-}) = 0$. In gravity, this suggests the *fixed-line normal coordinate* condition [10]

$$(x - x_{-})^{\alpha} (x - x_{-})^{\beta} \Gamma^{\mu}_{\ \alpha\beta} (x) = 0, \qquad (76)$$

which replaces (71). Suppose that $\Gamma^{\mu}_{\alpha\beta}(x)$ does not satisfy this condition, then one can find new coordinates \tilde{x}^{μ} , related to x^{μ} by

$$\tilde{x}^{\mu} = x^{\mu} + \frac{1}{2} (x - x_{-})^{\alpha} (x - x_{-})^{\beta} \Gamma^{\mu}_{\ \alpha\beta}$$

$$+ \frac{1}{6} (x - x_{-})^{\alpha} (x - x_{-})^{\beta} (x - x_{-})^{\nu} \left(\Gamma^{\mu}_{\ \alpha\tau} \Gamma^{\tau}_{\ \beta\nu} + \left[\partial_{\nu} \Gamma^{\mu}_{\ \alpha\beta} \right] \right) + \mathcal{O}((x - x_{-})^{4}),$$
(77)

such that $\tilde{\Gamma}^{\mu}_{\ \alpha\beta}(x)$ fulfills (76). In the above and for the remainder of Section 4, the following convention is adopted: whenever a soft field appears without explicit position argument, the field is evaluated at x^{μ}_{-} .

Once again, the fixed-line normal coordinate gauge condition (76) does not fix the gauge completely. For example, (76) leaves the components $\tilde{\Gamma}_{--}^{\mu}(\tilde{x})$ unconstrained and (77) does not affect $g_{s\mu\nu}(x_{-})$, since it is second order in $(x - x_{-})^{\mu}$. The remaining diffeomorphisms form a residual symmetry, which is the analogue of the global symmetry transformations from before (73), but since the parameters depend on x_{-}^{μ} , it is a *gauge* symmetry. As for the Riemann normal coordinates, the residual transformations can be used to further simplify the residual metric tensor $g_{s\mu\nu}(x_{-})$ by performing a linear coordinate transformation local in x_{-}^{μ} in the components orthogonal to x_{-}^{μ} . After this step, one can identify the homogeneous soft background field. To this end, introduce the vierbein

$$g_{s\mu\nu}(x_{-}) \equiv e_{\mu}^{\ \alpha}(x_{-})e_{\nu}^{\ \beta}(x_{-})\eta_{\alpha\beta}, \qquad (78)$$

and its inverse

Martin Beneke, Patrick Hager and Robert Szafron

$$E^{\mu}_{\ \alpha}(x_{-})e^{\alpha}_{\ \nu}(x_{-}) = \delta^{\mu}_{\nu}.$$
(79)

These objects can be thought of as matrices that locally diagonalise the metric tensor to express it in terms of the flat-space metric $\eta_{\mu\nu}$. Performing the linear transformation $\tilde{x}^{\mu} \to \check{x}^{\mu}(\tilde{x})$ defined by

$$n_{+}\check{x} = n_{+}\tilde{x}, \quad \check{x}_{\perp}^{\mu} = e_{\alpha}^{\ \mu_{\perp}}\tilde{x}^{\alpha}, \quad n_{-}\check{x} = n_{-\rho}e_{\alpha}^{\ \rho}\tilde{x}^{\alpha}, \tag{80}$$

which leaves x_{-}^{μ} invariant, yields a new coordinate system \check{x} , the fixed-line normal coordinates (FLNC). The original coordinate can be expressed in terms of the FLNC \check{x}^{μ} as

$$x^{\mu} = \check{x}^{\mu} + \theta^{\mu}_{\text{FLNC}}(\check{x}), \qquad (81)$$

with parameter

$$\theta_{\text{FLNC}}^{\mu}(x) = (E^{\mu}{}_{\rho} - \delta^{\mu}_{\rho})(x - x_{-})^{\rho} - \frac{1}{2}(x - x_{-})^{\rho}(x - x_{-})^{\sigma}E^{\alpha}{}_{\rho}E^{\beta}{}_{\sigma}\Gamma^{\mu}{}_{\alpha\beta}$$
(82)

$$+\frac{1}{6}(x-x_{-})^{\rho}(x-x_{-})^{\sigma}(x-x_{-})^{\lambda}E^{\alpha}_{\ \rho}E^{\beta}_{\ \sigma}E^{\nu}_{\ \lambda}\left(2\Gamma^{\mu}_{\ \alpha\tau}\ \Gamma^{\tau}_{\ \beta\nu}\ -\left[\partial_{\nu}\Gamma^{\mu}_{\ \alpha\beta}\ \right]\right)+\mathscr{O}(x^{4}).$$

Note that every $(x - x_{-})^{\mu}$ generates a contraction with the inverse vierbein as a result of this linear transformation. Returning now to the active point of view, the translation operator $T_{\theta_{\text{FLNC}}}(x)$ corresponding to the coordinate change to \check{x}^{μ} defines a new "Wilson line" $R_{\text{FLNC}}(x)$ via

$$R_{\rm FLNC}^{-1}(x) = T_{\theta_{\rm FLNC}}(x).$$
(83)

As an intermediate result, one can compute the metric tensor in fixed-line normal coordinates, denoted by $\check{g}_{s\mu\nu}$, using $R_{\rm FLNC}$ as

$$\check{g}_{s\mu\nu}(x) \equiv R^{\alpha}_{\ \mu}(x)R^{\beta}_{\ \nu}(x)\left[R^{-1}_{\text{FLNC}}(x)g_{s\alpha\beta}(x)\right],\tag{84}$$

with

$$R^{\alpha}{}_{\mu}(x) = \frac{\partial x^{\alpha}}{\partial \check{x}^{\mu}}(x) \tag{85}$$

the Jacobian of the transformation. As was the case for Riemann normal coordinates, the metric in FLNC can now be split into a background metric (simply $\eta_{\mu\nu}$ for the former), which forms the leading term, and the multipole series of manifestly covariant terms expressed via the Riemann tensor. Therefore, it is convenient to split the metric tensor as

$$\check{g}_{s\mu\nu}(x) \equiv \hat{g}_{s\mu\nu}(x) + \mathfrak{g}_{s\mu\nu}(x), \qquad (86)$$

where $g_{s\mu\nu}$ are the manifestly gauge-covariant terms, i.e. the Riemann tensor terms, and the background field is $\hat{g}_{s\mu\nu}(x)$. It is given by

$$\hat{g}_{s+-}(x) = e_{-+} - (x - x_{-})^{\alpha} \left[\Omega_{-}\right]_{\alpha+}, \qquad (87)$$

$$\hat{g}_{s\mu_{\perp}-}(x) = e_{-\mu_{\perp}} - (x - x_{-})^{\alpha} \left[\Omega_{-}\right]_{\alpha\mu_{\perp}},$$
(88)

$$\hat{g}_{s--}(x) = \left(e_{-}^{\alpha} - (x - x_{-})^{\rho} \left[\Omega_{-}\right]_{\rho}^{\alpha}\right) \left(e_{-}^{\beta} - (x - x_{-})^{\sigma} \left[\Omega_{-}\right]_{\sigma}^{\beta}\right) \eta_{\alpha\beta}, \quad (89)$$

$$\hat{g}_{s\mu_{\perp}\nu_{\perp}}(x) = \eta_{\mu_{\perp}\nu_{\perp}}, \qquad (90)$$

$$\hat{g}_{s+\perp}(x) = \hat{g}_{s+\perp}(x) = 0.$$
 (91)

Here, the fields on the right-hand side implicitly depend on x_- , i.e. $e_{-+} \equiv e_{-+}(x_-)$ and $[\Omega_-]_{\alpha\beta}$ is the spin-connection, defined as⁹

$$[\Omega_{-}]^{\alpha\beta} = e_{\nu}^{\ \alpha} \left[\partial_{-} E^{\nu\beta} \right] + e_{\nu}^{\ \alpha} \Gamma^{\nu}_{\ \sigma-} E^{\sigma\beta} .$$
⁽⁹³⁾

27

Note that the components orthogonal to x_{-}^{μ} coincide with the flat-space metric, as in these directions the light-front multipole expansion is the same as the static one.

The remainder term $\mathfrak{g}_{s\mu\nu}$ in (86) is manifestly gauge-covariant and depends only on the Riemann tensor. Up to $\mathcal{O}(\lambda^4)$, it is given by

$$\mathfrak{g}_{s\mu\nu}(x) = -\frac{n_{+\mu}n_{+\nu}}{4}x_{\perp}^{\alpha}x_{\perp}^{\beta}R_{\alpha-\beta-} - \frac{n_{+\mu}}{2}\frac{2}{3}x_{\perp}^{\alpha}x_{\perp}^{\beta}R_{\alpha\nu_{\perp}\beta-} - \frac{n_{+\nu}}{2}\frac{2}{3}x_{\perp}^{\alpha}x_{\perp}^{\beta}R_{\alpha\mu_{\perp}\beta-} -\left(\frac{n_{+\mu}n_{-\nu}}{4} + \frac{n_{+\nu}n_{-\mu}}{4}\right)\frac{2}{3}x_{\perp}^{\alpha}x_{\perp}^{\beta}R_{\alpha+\beta-} - \frac{1}{3}x_{\perp}^{\alpha}x_{\perp}^{\beta}R_{\alpha\mu_{\perp}\beta\nu_{\perp}}$$
(94)
$$-\frac{n_{-\mu}}{2}\frac{1}{3}x_{\perp}^{\alpha}x_{\perp}^{\beta}R_{\alpha\nu_{\perp}\beta+} - \frac{n_{-\nu}}{2}\frac{1}{3}x_{\perp}^{\alpha}x_{\perp}^{\beta}R_{\alpha\mu_{\perp}\beta+} - \frac{n_{-\mu}n_{-\nu}}{4}\frac{1}{3}x_{\perp}^{\alpha}x_{\perp}^{\beta}R_{\alpha+\beta+}.$$

The explicit expressions of $\hat{g}_{s\mu\nu}$ and $g_{s\mu\nu}$ obtained from this construction are important results, while the intermediate $\check{g}_{s\mu\nu}$ is no longer employed in the following.

The result for $\hat{g}_{s\mu\nu}$ contains two gauge fields appearing only in the n_{-}^{μ} components: the vierbein and the spin connection. They can be expanded in terms of the soft graviton field as

$$e_{-}^{\alpha} = \delta_{-}^{\alpha} + \frac{\kappa}{2}s_{-}^{\alpha} - \frac{\kappa^{2}}{8}s_{-\beta}s^{\beta\alpha} + \mathcal{O}(s^{3}), \qquad (95)$$

$$[\Omega_{-}]_{\alpha\beta} = -\frac{\kappa}{2} \left(\left[\partial_{\alpha} s_{\beta-} \right] - \left[\partial_{\beta} s_{\alpha-} \right] \right) + \mathscr{O}(s^{2}) \,. \tag{96}$$

Note that the spin connection is defined as the derivative of the metric, resp. vierbein. In the EFT, the soft metric and vierbein are always evaluated at x_{-}^{μ} . Thus, in the EFT, $\partial_{\perp}s_{\mu\nu}(x_{-}) = 0$ if the derivative were taken after setting $x = x_{-}$. This means that the vierbein and the spin-connection above should be interpreted as *two truly*

$$\left[\Omega_{\mu}\right]^{\alpha\beta}(x) = e_{\nu}^{\alpha} \left[\partial_{\mu} E^{\nu\beta}\right](x) + e_{\nu}^{\alpha} \Gamma^{\nu}_{\ \sigma\mu} \ E^{\sigma\beta}(x) \tag{92}$$

⁹ The definitions (78), (79) can be extended from the light-cone x_{-}^{μ} to x^{μ} , in which case

coincides with the usual spin-connection. The background-field construction only needs $[\Omega_{-}]^{\alpha\beta}(x_{-})$, which is well-defined on the light-cone, since only the derivative $n_{-}\partial$ along the light-cone appears in (93).

independent gauge fields from the EFT perspective, since the derivative contains additional information that cannot be obtained from the vierbein in the EFT.

Corresponding to these two independent gauge fields, the soft gauge transformations induce two *emergent* soft gauge symmetries that take the form of local translations and local Lorentz transformations on the light-cone x_{-}^{μ} of the collinear modes, and global transformations on the hyper-planes of fixed x_{-}^{μ} . More precisely, performing a gauge transformation $U_s(x) = T_{\varepsilon}^{-1}$ with local translation parameter $\varepsilon_{\mu}(x)$, $\hat{g}_{s\mu\nu}$ does not transform with T_{ε}^{-1} , but instead with $T_{\varepsilon+\omega}^{-1}$, defined by the parameters $\varepsilon_{\mu}(x_{-})$ and $\omega_{\mu\nu}(x_{-}) = -\frac{1}{2} \left(\left[\partial_{\mu} \varepsilon_{\nu} \right] - \left[\partial_{\nu} \varepsilon_{\mu} \right] \right)(x_{-})$, which again should be regarded as two independent transformations. The corresponding infinitesimal coordinate transformation is given by

$$x^{\mu} \to x^{\mu} + \kappa \varepsilon^{\mu}(x_{-}) + \kappa \omega^{\mu}{}_{\nu}(x_{-})(x - x_{-})^{\nu}.$$
⁽⁹⁷⁾

The first term stems from the transformation of the vierbein, while the second one, proportional to $(x - x_-)^{\mu}$, arises from the spin-connection transformation. The two gauge symmetries of the collinear fields in the effective theory are *emergent*, since they have a different physical meaning than the original one. The emergent soft gauge symmetries, and consequently the soft fields, live only along the classical trajectory of the collinear particles x_-^{μ} and are "visible" only to the collinear fields in direction *i*. Other collinear directions see their own soft gauge symmetry. The symmetries of the different collinear sectors are induced by the gauge symmetry of the full theory, and connected through the purely-soft part of the Lagrangian, which continues to transform under the full soft symmetry parameter $\varepsilon_{\mu}(x)$.

The effective soft gauge fields $e_{-\alpha}(x_{-})$, $[\Omega_{-}]_{\alpha\beta}(x_{-})$ are non-linear functions of $s_{\mu\nu}$. Since the residual background metric has the property that only $\hat{g}_{s\mu-}$ is non-vanishing, it is possible to define a soft-covariant derivative

$$n_{-}D_{s} \equiv \hat{g}_{s}^{\mu-}\partial_{\mu}$$

$$= \partial_{-} - \frac{\kappa}{2}s_{-\mu}\partial^{\mu} + \frac{\kappa^{2}}{8}s_{+-}s_{--}\partial_{+} + \frac{\kappa^{2}}{16}s_{-\alpha_{\perp}}s_{-}^{\alpha_{\perp}}\partial_{+} + \frac{\kappa}{2}\left[\Omega_{-}\right]_{\mu\nu}J^{\mu\nu} + \mathscr{O}(\lambda^{3}),$$

$$(98)$$

where

$$J^{\mu\nu} = (x - x_{-})^{\mu} \partial^{\nu} - (x - x_{-})^{\nu} \partial^{\mu}$$
(99)

is the angular momentum (Lorentz generator) operator. The form of (98) is significant and revealing. Usually, a scalar field coupled to gravity does not have a covariant derivative. However, in SCET gravity, the residual symmetry corresponds to local Lorentz transformations that live only on the classical trajectory of the energetic particles. A scalar field has a non-trivial transformation under this symmetry, and therefore requires the introduction of a covariant derivative. Namely, if a collinear scalar field transforms under (97) infinitesimally as

$$T_{\varepsilon+\omega}^{-1}\varphi_c(x) = 1 - \kappa\varepsilon^{\alpha}\partial_{\alpha}\varphi_c(x) - \kappa\omega_{\alpha\beta}\left(x - x_{-}\right)^{\beta}\partial^{\alpha}\varphi_c(x) + \mathscr{O}(\varepsilon^2), \qquad (100)$$

one can check that the derivative (98) has the covariant transformation

$$n_{-}D_{s}\varphi_{c}(x) \to T_{\varepsilon+\omega}^{-1}[n_{-}D_{s}\varphi_{c}(x)] - \kappa\omega_{-\alpha}D_{s}^{\alpha}\varphi_{c}(x) + \mathscr{O}(\varepsilon^{2}).$$
(101)

The transverse partial derivatives ∂_{\perp} and $n_+\partial$ already transform as in (101), and no modification is necessary.

The soft-covariant derivative (98) contains the two independent gauge fields. At the linear order in $s_{\mu\nu}$, corresponding to the minimal-coupling terms in the Lagrangian, the first gauge field $e_{-\mu}$ couples to the momentum $P^{\mu} \equiv -i\partial^{\mu}$, while the second gauge field, the spin-connection, couples to the angular momentum of the scalar field. For representations with non-vanishing spin, the total angular momentum would appear in (98). In the case of several directions, the object $n_{i-}D_s$ appears inside the *i*-collinear Lagrangian \mathcal{L}_{c_i} . In this sense, one has $i = 1, \ldots, N$ gauge symmetries that are restricted to their respective collinear trajectories, all ultimately tied to the same background soft field through the purely soft Lagrangian $\mathcal{L}_s[s_{\mu\nu}(x)]$ in (41).

It is important to stress that, as seen in (98) and (94), the fields $\hat{g}_{s\mu\nu}$ and $g_{s\mu\nu}$ are expressed in terms of the *original* soft graviton field $s_{\mu\nu}$, where no gauge is fixed. This is where the residual gauge symmetry stems from. The FLNC coordinates are merely a tool to construct the covariant light-front multipole expansion of the EFT, and one does not fix this gauge in the final Lagrangian, which is a function of $s_{\mu\nu}$. Similarly to the case of light-cone gauge in the collinear sector, using the analogue of Wilson lines, the FLNC condition can be implemented "covariantly" by redefining the collinear fields, without restricting the gauge freedom of the soft field.

4.3 Soft-collinear Lagrangian

Equipped with these concepts, it is not difficult to derive the effective Lagrangian by purely algebraic manipulations. As a first step, one redefines the collinear fields using the collinear "Wilson line" W_c (50) and the previously defined *R*-"Wilson line" (83)¹⁰ to obtain collinear fields that have gauge transformations *compatible* with the multipole expansion. These new fields are denoted by $\hat{\varphi}_c$ and $\hat{h}_{\mu\nu}$, and are given by¹¹

$$\varphi_{c} = \left[R_{\text{FLNC}} W_{c}^{-1} \hat{\varphi}_{c} \right] = \left[R_{\text{FLNC}} \hat{\chi}_{c} \right], \qquad (102)$$

$$\kappa h_{\mu\nu} = \left[R_{\text{FLNC}} R_{\mu}^{\ \alpha} R_{\nu}^{\ \beta} \left(W_{\ \alpha}^{\rho} W_{\ \beta}^{\sigma} \left[W_{c}^{-1} (\kappa \hat{h}_{\rho\sigma} + \hat{g}_{s\rho\sigma}(x)) \right] - \hat{g}_{s\alpha\beta}(x) \right) \right]$$

$$= \left[R_{\text{FLNC}} R_{\mu}^{\ \alpha} R_{\nu}^{\ \beta} \hat{\mathfrak{h}}_{\alpha\beta} \right]. \qquad (103)$$

¹⁰ In the following, R_{FLNC} , its Jacobians R_{μ}^{α} and its determinant det(<u>R</u>) are understood to be evaluated at *x* if no argument is given.

¹¹ There is one subtlety: the collinear "Wilson line" in the soft-collinear theory is different from the purely-collinear one at the non-linear level, as the soft background field has a non-vanishing \hat{g}_{s+-} component, and therefore appears in the transformation of \hat{h}_{+-} . This is explained in [10].

Here the collinear fields on the left-hand side are assumed to be in light-cone gauge, but the gauge is not fixed on the right-hand side. The right-hand sides also identify the collinear gauge-invariant building blocks $\hat{\chi}_c$ and $\hat{\mathfrak{h}}_{\mu\nu}$ after the field redefinition, i.e. they are invariant with respect to the compatible gauge symmetries.

To illustrate the calculation of the effective Lagrangian, consider the collinear matter field in the soft background $g_{s\mu\nu}(x)$ as defined in (65),¹²

$$\mathscr{L}_{\varphi} = \frac{1}{2} \sqrt{-g_s} g_s^{\mu\nu} \partial_{\mu} \varphi_c \partial_{\nu} \varphi_c \,. \tag{104}$$

Inserting (102) results in

$$\mathscr{L}_{\varphi} = \frac{1}{2} \sqrt{-g_s} g_s^{\mu\nu} \left[\partial_{\mu} (R_{\text{FLNC}} \hat{\chi}_c) \right] \left[\partial_{\nu} (R_{\text{FLNC}} \hat{\chi}_c) \right]$$
$$= \frac{1}{2} \det(\underline{R}) \left[R_{\text{FLNC}}^{-1} \sqrt{-g_s} \right] R_{\mu}^{\ \alpha} R_{\nu}^{\ \beta} \left[R_{\text{FLNC}}^{-1} g_s^{\mu\nu} \right] \partial_{\alpha} \hat{\chi}_c \partial_{\beta} \hat{\chi}_c.$$
(105)

The metric $\check{g}^{\alpha\beta}(x)$ from (84) can now be identified, together with the corresponding metric determinant. Inserting (86) and dropping the Riemann tensor terms, which are sub-leading in the λ -expansion gives

$$\mathcal{L}_{\varphi} = \frac{1}{2} \sqrt{-\hat{g}_s} \hat{g}_s^{\mu\nu} \partial_{\mu} \hat{\chi}_c \partial_{\nu} \hat{\chi}_c$$
$$= \sqrt{-\hat{g}_s} \left(\frac{1}{2} n_+ \partial \hat{\chi}_c n_- D_s \hat{\chi}_c + \frac{1}{2} \partial_{\alpha_\perp} \hat{\chi}_c \partial^{\alpha_\perp} \hat{\chi}_c \right).$$
(106)

The last line follows from (98) and shows that after the multipole expansion, the leading soft interactions are collected into the covariant derivative.

Applying this procedure to the background-field Lagrangian, now including nonlinear terms in the collinear graviton field and keeping the Riemann tensor terms in (86), one finds the soft-collinear Lagrangian in the form

$$\mathscr{L} = \sqrt{-\hat{g}_s} \left(\mathscr{L}^{(0)} + \mathscr{L}^{(1)} + \mathscr{L}^{(2)} + \dots \right), \tag{107}$$

where

$$\mathscr{L}^{(0)} = \frac{1}{2} n_{+} \partial \hat{\chi}_{c} n_{-} D_{s} \hat{\chi}_{c} + \frac{1}{2} \partial_{\alpha_{\perp}} \hat{\chi}_{c} \partial^{\alpha_{\perp}} \hat{\chi}_{c} , \qquad (108)$$

$$\mathscr{L}^{(1)} = -\frac{\kappa}{2} \hat{\mathfrak{h}}^{\mu\nu} \partial_{\mu} \hat{\chi}_{c} \partial_{\nu} \hat{\chi}_{c} + \frac{\kappa}{4} \hat{\mathfrak{h}}^{\beta_{\perp}}_{\beta_{\perp}} \left(n_{+} \partial \hat{\chi}_{c} n_{-} D_{s} \hat{\chi}_{c} + \partial_{\alpha_{\perp}} \hat{\chi}_{c} \partial^{\alpha_{\perp}} \hat{\chi}_{c} \right) , \qquad (109)$$

$$\mathscr{L}^{(2)} = -\frac{1}{8} x_{\perp}^{\alpha} x_{\perp}^{\beta} R_{\alpha-\beta-} (n_{+}\partial\hat{\chi}_{c})^{2} + \frac{\kappa}{2} \hat{\mathfrak{h}}^{\mu\alpha} \hat{\mathfrak{h}}_{\alpha}{}^{\nu} \partial_{\mu}\hat{\chi}_{c} \partial_{\nu}\hat{\chi}_{c} - \frac{\kappa^{2}}{4} \hat{\mathfrak{h}}^{\alpha\perp}{}_{\alpha\perp} \hat{\mathfrak{h}}^{\mu\nu} \partial_{\mu}\hat{\chi}_{c} \partial_{\nu}\hat{\chi}_{c} + \frac{\kappa^{2}}{16} \left((\hat{\mathfrak{h}}^{\alpha\perp}{}_{\alpha\perp})^{2} - 2\hat{\mathfrak{h}}^{\alpha\beta} \hat{\mathfrak{h}}_{\alpha\beta} \right) \left(n_{+}\partial\hat{\chi}_{c} n_{-} D_{s}\hat{\chi}_{c} + \partial_{\mu\perp}\hat{\chi}_{c} \partial^{\mu\perp}\hat{\chi}_{c} \right) .$$
(110)

¹² In (104) – (107), $g_{s\mu\nu}$ and its determinant are evaluated at x, not x₋.

The superscript on \mathscr{L} denotes the lowest order in the λ -expansion at which the respective terms contribute. By expanding out the implicit collinear Wilson lines and n_D_s , each term generates an infinite tower of higher-order terms in λ . The derivation of the above Lagrangian is presented in detail in [10], where one can also find the explicit λ -expansion of the Lagrangian up to $\mathscr{O}(\lambda^2)$ in terms of the elementary fields $\hat{\varphi}_c$, $\hat{h}_{\mu\nu}$, $s_{\mu\nu}$. In general, there is also the soft matter field φ_s , however, in the absence of scalar self-interactions, it does not contribute to the soft-collinear Lagrangian at $\mathscr{O}(\lambda^2)$.

The soft graviton field $s_{\mu\nu}$ appears above only inside the soft-covariant derivative n_D_s , inside the Riemann tensor and the gauge-invariant building blocks. This highlights a formal similarity of the gravitational soft-collinear interactions to the respective gauge-theory result [6], featuring a leading interaction via a covariant derivative, and sub-leading interactions starting from the quadrupole term (in contrast to the dipole term in gauge theory). The main difference in gravity is that the covariant derivative contains not one but *two* gauge fields.

Moreover, the Lagrangian above is expressed in terms of the gauge-invariant building blocks $\hat{\chi}_c$, $\hat{\mathfrak{h}}_{\mu\nu}$. Similar to the purely-collinear Lagrangian (59) in Section 3, the theory is invariant under the redefined hatted-collinear transformations, and one can decide to employ either the gauge-invariant or non-invariant fields $\hat{\varphi}_c$, $\hat{h}_{\mu\nu}$. With the latter choice one finds that the collinear Wilson line W_c^{-1} drops out of all terms that do not contain the soft Riemann tensor. Above, this applies to all but the first term in (110), since this term is not minimally-coupled to the homogeneous background field [10] and would consequently contain explicit factors of W_c^{-1} .

In the purely-collinear sector (setting all soft fields to zero), gravity is distinct from gauge theory. There is no collinear-covariant derivative D_c^{μ} in gravity, and correspondingly, leading-power collinear interactions are absent. In combination with the operator basis discussed below, where the first graviton building block starts at $\mathcal{O}(\lambda)$, this implies that one cannot generate collinear emissions at leading power. In consequence, there are no jets in gravity and each additional collinear emission costs a power of λ . This immediately shows that gravity does not exhibit any collinear divergences, and indeed *nothing does go wrong* in gravity [38].

Moreover, in the soft-collinear sector the covariant derivative can be eliminated iteratively from the sub-leading power Lagrangians by making use of the field equation, such that it appears only in $\mathscr{L}^{(0)}$. Therefore, gravity features an *extended eikonal* interaction compared to gauge theory, related to the two independent gauge fields in $n_D s$. This fundamental result follows quite naturally from the SCET gravity construction.

Figure 4 displays the space-time structure of SCET gravity interactions, showing for simplicity only a single collinear direction and suppressing the transverse directions. The classical trajectory of a massless, energetic (collinear) particle passing through $\vec{x} = 0$ at t = 0 is the line $x_{-}^{\mu} = (n_{+}x)\frac{n_{-}^{\mu}}{2}$ (dashed red). Collinear field operators describing these particles are non-local in the opposite light-like direction n_{+}^{μ} (wavy blue) as discussed around (24) and Section 5, but interact with soft fields only at x_{-}^{μ} via the extended eikonal interaction and the manifestly covariant Riemann tensor terms as a consequence of the light-front multipole expansion.

Martin Beneke, Patrick Hager and Robert Szafron



Fig. 4 See text for explanation.

4.4 Gauge-invariant building blocks

In the construction of the Lagrangian, we identified a set of gauge-invariant operators that can be used as ingredients in the hard matching. The only collinear gauge-invariant building blocks in the operator basis are (cf. (102), (103))

$$\hat{\boldsymbol{\chi}}_{c} = \begin{bmatrix} W_{c}^{-1} \hat{\boldsymbol{\varphi}}_{c} \end{bmatrix}, \\ \kappa \hat{\boldsymbol{\mathfrak{h}}}_{\mu\nu} = W^{\alpha}_{\ \mu} W^{\beta}_{\ \nu} \begin{bmatrix} W_{c}^{-1} \kappa \hat{\boldsymbol{h}}_{\alpha\beta} \end{bmatrix} + \left(W^{\alpha}_{\ \mu} W^{\beta}_{\ \nu} \begin{bmatrix} W_{c}^{-1} \hat{\boldsymbol{g}}_{s\alpha\beta}(\boldsymbol{x}) \end{bmatrix} - \hat{\boldsymbol{g}}_{s\mu\nu}(\boldsymbol{x}) \right),$$
(111)

where $\mu \nu \in \{\perp \perp, \perp -, --\}$ thanks to the condition $\hat{\mathfrak{h}}_{\mu+} = 0$ defining the collinear Wilson line. The sub-leading minus-components $\hat{\mathfrak{h}}_{\perp-}$ and $\hat{\mathfrak{h}}_{--}$ can be eliminated using the collinear equation of motion, which, at leading-order, express these components in terms of derivatives of $\hat{\mathfrak{h}}_{\perp\perp}$, similar to (62) in the purely collinear theory. The building blocks $\hat{\chi}_c$, $\hat{\mathfrak{h}}_{\perp\perp}$ both scale as $\mathscr{O}(\lambda)$. Additional suppression in λ is obtained by acting with ∂^{μ}_{\perp} on collinear building blocks. Each additional derivative increases the λ -power by one unit. The ordinary derivative is the appropriate one, because one operates on gauge-invariant field products.

While the building blocks are inherently inhomogeneous in λ , one can use the power-counting of the leading term to characterise these operators. The effect of the infinite tower of terms sub-leading in λ is simply to render the leading term gauge-invariant. If one fixes collinear light-cone gauge, the entire sub-leading tower disappears and one obtains homogeneously-scaling operators.

For the soft sector, one employs gauge-*covariant* building blocks, like the soft scalar φ_s , the derivative n_-D_s or the Riemann tensor $R^{\mu}_{\ \nu\alpha\beta}$. However, the soft-covariant derivative can be eliminated from the building blocks by the collinear field equations [10]. For the scalar field, at leading power

$$n_{-}D_{s}\hat{\chi}_{c} = -\frac{\partial_{\perp}^{2}}{n_{+}\partial}\hat{\chi}_{c}, \qquad (112)$$

and a similar equation can be derived from the Einstein-Hilbert Lagrangian. This shows that $n_-D_s\mathfrak{h}_{\perp\perp}$ is redundant [10] and can be traded for (non-local) combinations of the other collinear and soft building blocks, which will contribute to the equation of motion at higher powers. This implies that the soft-covariant derivative is not a relevant building block for the *N*-jet operators (sources). When n_-D_s is eliminated systematically, the remaining soft-graviton building block is the Riemann tensor and its derivatives. Therefore, soft gauge-covariance implies that softgraviton building blocks are suppressed by at least λ^6 .¹³

5 Sources and hard matching

The aim of the previous sections was to construct the Lagrangian of an effective theory that reproduces the gravitational scattering amplitudes in the soft and collinear limits, that is when some of the external momenta k_i form small invariants¹⁴ $k_i \cdot k_j \ll Q^2$, where Q is the scale of the hard scattering. The soft-collinear effective Lagrangian captures only part of the full scattering amplitude: it describes purely-collinear interactions within a single collinear direction and the soft-collinear interactions. To recover the full scattering amplitude, one also needs to include the effect of the highly virtual propagators and loops, the "hard" region. Hard lines and loops always connect different directions and are the source of large-angle scattering of energetic particles in the first place. In the pictorial representation of Figure 2, the entire hard process is shrunk to the central point.

In technical terms, in the scattering of multiple collinear sectors, the intermediate highly-virtual states are hard modes due to the presence of the different large momentum components $n_{i+}p_i, n_{j+}p_j, \ldots$, from which one can form hard invariants

$$(n_{i+}p_i)(n_{j+}p_j)(n_{i-}n_{j-}) \sim \lambda^0.$$
(113)

The hard region is integrated out in the EFT by construction and is then encoded in the hard matching coefficients of the operators containing fields of multiple collinear directions. This gives rise to the so-called *N*-jet operators or "currents" or "sources".¹⁵ The full scattering amplitude is then the sum of matrix elements of *N*jet operators evaluated with the soft-collinear effective Lagrangian. The construction

33

¹³ There is an implicit dependence on the soft metric field in the definition of the collinear building blocks (111) through W_c^{-1} and $\hat{g}_{s\mu\nu}$. This dependence is already constrained by collinear gauge-invariance and does not give rise to soft building blocks. The soft-graviton vertices generated by this dependence always contain collinear gravitons as well.

¹⁴ Lorentz-invariant products of different momenta.

 $^{^{15}}$ The reason why the *N*-jet operators do not appear in the Lagrangian as interactions connecting different collinear sectors is that they can appear in a hard process only once, i.e. the hard subgraph must be connected, provided that there are no collinear particles in the same direction in the initial

of the effective Lagrangian is a prerequisite to the construction of N-jet operators, because it defines what is the field basis from which the hard sources are constructed and the symmetries they must respect. After having established this in the previous sections, the admissible sources can now be easily specified. Their matching coefficients depend on the particular scattering process, and must of course be calculated in a given context.

The *N*-jet operators are constructed from the collinear and soft building blocks. It corresponds to a purely-hard scattering that produces the constituent buildingblock particles of the operator. For example, the tree-level scattering of *N* widelyseparated energetic scalar particles is encoded in a current operator consisting of *N* copies of the collinear gauge-invariant scalar $\hat{\chi}_{c_i}$, one for each direction, combined with a hard matching coefficient. One can then think of the hard matching coefficient as the *non-radiative amplitude of the hard process*, while the building blocks correspond to the external legs of the amplitude.

This notion is made more precise in the following. More details for the gauge theory situation can be found in [7] and for the gravitational case in [10]. The generic N-jet operator in gauge-theory is a light-ray operator [7]

$$\mathscr{J}(0) = \int [dt]_N \widetilde{C}(t_{i_1}, t_{i_2}, \dots) J_s(0) \prod_{i=1}^N J_i(t_{i_1}, t_{i_2}, \dots), \qquad (114)$$

 $[dt]_N = \prod_{i_k} dt_{i_k}$ and "0" refers to the origin x = 0, where the hard-scattering is supposed to take place. Here, $\widetilde{C}(t_i)$ is the hard matching coefficient, J_i are composite operators of *i*-collinear fields, and J_s of soft fields. The variables t_{i_k} correspond to the non-locality of collinear operators along the light-cone, for example, $J_i(t_{i_1}, t_{i_2}, ...)$ might refer to the product $\hat{\chi}_{c_i}(t_{i_1}n_{i_+})\hat{\chi}_{c_i}(t_{i_2}n_{i_+})\dots$ of fields in the same collinear direction. In momentum space, $t_{i_k} \to n_{i_k} \cdot p_{i_k}$, which can be expressed in terms of the fraction of total momentum in direction *i* carried by the factor *k* in J_i , which provides the relation to the intuitive notion of the non-radiative amplitude.

Like the full-theory scattering amplitude, these *N*-jet operators must be gauge invariant. In the effective theory, this means invariance under both the soft and the collinear gauge symmetries. To ensure collinear gauge-invariance, one uses the gauge-invariant building blocks in the collinear current J_i , explicitly given in (111) in the previous section. Soft fields are automatically collinear gauge-invariant. In addition, the collinear building blocks are soft gauge-covariant, as they are constructed from the hatted fields. For J_s , one should employ soft-covariant building blocks like φ_s or the Riemann tensor $R^{\mu}_{\ \ \nu\alpha\beta}$, as discussed in Section 4.4. The *N*-jet operator as defined in (114) is now collinear gauge-invariant and soft gauge-covariant, transforming under global soft Poincaré transformations $U_s(0)$. The *N*-jet operator as defined in (114) is located at x = 0, however, and hence not translation-invariant. To render it manifestly gauge-invariant under both soft and collinear gauge transformations, one defines the translation- and therefore gauge-invariant current

and final state of the scattering. In this case, collinear modes in different directions can never "meet again" in a hard process as a consequence of the Coleman-Norton theorem [22].

$$\mathscr{J} = \int d^4 x T_x \mathscr{J}(0) T_x^{-1}, \qquad (115)$$

where $T_x = e^{ix\hat{p}}$ is the translation operator. In practical computations, the integral and translation operators reduces to the momentum conserving δ -function once the amplitude is evaluated. Therefore, one can choose the simpler *N*-jet operator (114) located at x = 0 and impose momentum and angular-momentum conservation by hand, similar to imposing colour-neutrality in QCD.

As said above, the hard coefficient is obtained from a process-specific matching computation. To perform this matching, one follows the standard method of evaluating a suitable on-shell Green function in the EFT and in the full theory and demanding that they are equal up to a desired power in the λ -expansion. One typically chooses the simplest possible external states that give non-vanishing matrix elements in both theories. Once the matching coefficients are determined, one can use the EFT operators to evaluate arbitrarily complex low-energy matrix elements. Note that these sources are the only place in the effective theory where renormalisation takes place. The Lagrangian is not renormalised [5], in the sense that no further renormalisation is required, when the SCET Lagrangian is expressed in terms of the renormalised full-theory couplings.¹⁶ The matching coefficients, however, receive loop corrections.

6 Soft theorem

If the radiative amplitude for the emission of (usually) a single soft particle from a hard scattering process can be expressed in terms of the non-radiative one without detailed knowledge of the latter, one refers to the corresponding result as a "soft theorem". The best known and earliest example of a soft theorem is the abelian version of (3), which expresses the soft-photon emission amplitude in QED in terms of a universal "eikonal factor". In QED, the absence of photon self-interactions leads to the exponentiation of multiple soft photon emissions. Of particular relevance to this chapter is the further fact, known as the Low-Burnett-Kroll theorem [31, 19], that the soft theorem extends to next-to-leading order in the soft expansion.

Almost 50 years after Weinberg's first discussion of the gravitational soft theorem (2), Cachazo and Strominger [20] made the remarkable observation that its universality extends by *two* orders in the soft expansion. If $\mathscr{A}(\{p_i\})$ denotes the amplitude for a *N*-particle hard-scattering process with momenta p_1, \ldots, p_N , the single graviton emission amplitude at tree-level reads

$$\mathscr{A}_{\rm rad}(\{p_i\};k) = \frac{\kappa}{2} \sum_{i=1}^{N} \bar{u}(p_i) \left(\frac{\boldsymbol{\varepsilon}_{\mu\nu}(k) p_i^{\mu} p_i^{\nu}}{p_i \cdot k} + \frac{\boldsymbol{\varepsilon}_{\mu\nu}(k) p_i^{\mu} k_{\rho} J_i^{\nu\rho}}{p_i \cdot k} \right)$$

35

¹⁶ In gravity, one has to keep in mind that the renormalised full theory is constructed up to a certain loop order by introducing additional higher-derivative operators, as explained in Section 2.1.

Martin Beneke, Patrick Hager and Robert Szafron

$$+\frac{1}{2}\frac{\varepsilon_{\mu\nu}(k)k_{\rho}k_{\sigma}J_{i}^{\rho\mu}J_{i}^{\sigma\nu}}{p_{i}\cdot k}+\mathscr{O}(k^{2})\right)\mathscr{A}(\{p_{i}\}).$$
 (116)

Here

$$J_i^{\mu\nu} = L_i^{\mu\nu} + \Sigma_i^{\mu\nu} = p_i^{\mu} \frac{\partial}{\partial p_{i\nu}} - p_i^{\nu} \frac{\partial}{\partial p_{i\mu}} + \Sigma_i^{\mu\nu}$$
(117)

refers to generators of the Lorentz group, in this context usually referred to as the "angular momentum operator", with $L_i^{\mu\nu}$ the orbital angular momentum operator of particle *i* from which the graviton is emitted, and $\Sigma_i^{\mu\nu}$ the spin operator in the representation of emitter ("matter") particle *i*.¹⁷ $\bar{u}(p_i)$ denotes the polarisation functions of the matter particles, and $\varepsilon_{\mu\nu}(k)$ the graviton polarisation tensor. In the following subsections, we shall focus on scalar matter, in which case $J_i^{\mu\nu} = L_i^{\mu\nu}$ and the $\bar{u}(p_i)$ are trivial. It is worth noting that one recovers the Low-Burnett-Kroll theorem for the emission of a photon, and its non-abelian generalisation [21], from (116) by substituting $p_i^{\mu} \rightarrow t_i^a$, $\varepsilon_{\mu\nu}(k) \rightarrow \varepsilon_{\nu}(k)$ and $\kappa/2 \rightarrow -g_s$, and dropping the next-to-next-to-soft term in the second line of (116), which is unique to gravitons. The above substitutions are suggestive of colour-kinematics duality, but we shall see below that regarding the next-to-soft terms in gauge theory and gravity as analogues of each other does not correspond to the proper interpretation of their physics origin.

The discovery of the sub-sub-leading gravitational soft theorem (116) was inspired [36, 37] by a relation between the leading soft theorem and the Bondi, van der Burg, Metzner and Sachs symmetries [17, 35], which consist of a subgroup of diffeomorphisms operating on the asymptotically flat regions of space-times, the significance of which has not yet become completely clear. The actual derivation of (116) in [20] used the spinor-helicity formalism, which is particularly elegant for matter particles with spin, as the angular momentum operator in spinor-helicity variables combines the orbital and spin parts in a simple way. Subsequently, the next-to-next-to-soft theorem was derived in various other ways [18, 15, 41].

In the spirit of the article, we ask what the effective field theory can say about the soft theorem. Obviously, the EFT must reproduce it as the special case of single-soft emission, but can it provide additional insight? In previous sections, the important role of the emergent gauge symmetries in the construction of soft-collinear gravity has become apparent. The multipole expansion of the Lagrangian naturally leads to structures that already resemble the angular momentum operator, which shows up explicitly in the coupling to one of the gauge fields. In the following we therefore focus on the fundamental questions: Why are there three and exactly three universal terms in the soft theorem? What is the origin and interpretation of the angular momentum factors? Is the soft theorem corrected by loop effects?

Before approaching these questions from soft-collinear gravity, the essence of the derivation of (116) from the explicit expansion of the scattering amplitude will be briefly reviewed, following closely [15].

¹⁷ All up to a factor of *i*, omitted for the convenience of writing (116) without factors of *i*.



Fig. 5 Emission from the external leg gives the first term in the decomposition of the amplitude in (118).

6.1 Soft theorem from graviton amplitudes

Since amplitudes factorise over their poles, the N + 1-point amplitude with emission of a single soft graviton at tree-level from a hard *N*-particle scattering process with amplitude $\mathscr{A}(\{p_i\})$ can be written as

$$\mathscr{A}_{\rm rad}^{\mu\nu}(\{p_i\};k) = \sum_{i=1}^{N} \frac{p_i^{\mu} p_i^{\nu}}{p_i \cdot k} \mathscr{A}(p_1, \dots, p_i + k, \dots, p_N) + \mathscr{B}^{\mu\nu}(\{p_i\};k).$$
(118)

The polarisation tensor $\varepsilon_{\mu\nu}(k)$ of the graviton has been stripped off, as well as the overall gravitational coupling factor $\kappa/2$. The first term on the right-hand side contains the non-radiative amplitude with one momentum shifted by the graviton momentum k and fully accounts for the singular $\mathcal{O}(1/k)$ term of the amplitude, which arises from the emission off an external leg, as shown in Figure 5. The second term starts at the next-to-soft order $\mathcal{O}(k^0)$. For simplicity of presentation, it is assumed that the N hard particles have spin 0.

One now makes use of the invariance of on-shell amplitudes under the gauge transformations of weak-field gravity, which implies that $\mathcal{E}_{\mu\nu}(k)\mathscr{A}_{rad}^{\mu\nu}(\{p_i\};k)$ does not change if the graviton polarisation tensor is replaced by

$$\varepsilon_{\mu\nu}(k) \to \varepsilon_{\mu\nu}(k) + k_{\mu}\alpha_{\nu}(\{p_i\};k), \qquad (119)$$

where $\alpha_v(\{p_i\};k)$ is a function of momenta satisfying $k^v \alpha_v(\{p_i\};k) = 0$. Since $\alpha_v(\{p_i\};k)$ is arbitrary, it follows that

$$0 = k_{\mu} \mathscr{A}_{rad}^{\mu\nu}(\{p_i\};k)$$

= $\sum_{i=1}^{N} p_i^{\nu} \mathscr{A}(p_1, \dots, p_i + k, \dots, p_N) + k_{\mu} \mathscr{B}^{\mu\nu}(\{p_i\};k).$ (120)

The amplitudes in this equation are analytic in k for tree-level emission. Expanding in k provides relations between (derivatives of) $\mathscr{B}^{\mu\nu}$ and $\mathscr{A}(\{p_i\})$. The vanishing

of (120) at $\mathcal{O}(k^0)$ is guaranteed by momentum conservation, $\sum_{i=1}^{N} p_i^{\nu} = 0$. The next two orders result in

$$\sum_{i=1}^{N} p_i^{\nu} \frac{\partial}{\partial p_{i\mu}} \mathscr{A}(\{p_i\}) + \mathscr{B}^{\mu\nu}(\{p_i\}; 0) = 0, \qquad (121)$$

$$\sum_{i=1}^{N} p_{i}^{\nu} \frac{\partial^{2}}{\partial p_{i\mu} \partial p_{i\rho}} \mathscr{A}(\{p_{i}\}) + \left[\frac{\partial \mathscr{B}^{\mu\nu}}{\partial k_{\rho}} + \frac{\partial \mathscr{B}^{\rho\nu}}{\partial k_{\mu}}\right] (\{p_{i}\}; 0) = 0.$$
(122)

The expansion of (120) does not give the above two equations directly, but multiplied by k_{μ} and $k_{\mu}k_{\rho}$, respectively. Removing these vectors is justified, because any local gauge-invariant term in $\mathscr{B}^{\mu\nu}$ that would be missed in this operation can appear only at order k^2 [15].

Gauge invariance thus allows to express the $k \to 0$ limit of $\mathscr{B}^{\mu\nu}$ in terms of the derivative of the non-radiative amplitude, as well as a symmetric first-order derivative of $\mathscr{B}^{\mu\nu}$ in terms of the second derivative. Inserting (121), (122) into the expansion of (118) in k gives

$$\mathscr{A}_{\rm rad}^{\mu\nu}(\{p_i\};k) = \sum_{i=1}^{N} \frac{p_i^{\nu}}{p_i \cdot k} \left[p_i^{\mu} + k_{\rho} J_i^{\mu\rho} \right] \mathscr{A}(\{p_i\}) + \frac{1}{2} k_{\rho} k_{\sigma} \sum_{i=1}^{N} \frac{p_i^{\nu}}{p_i \cdot k} J_i^{\mu\rho} \frac{\partial}{\partial p_{i\sigma}} \mathscr{A}(\{p_i\}) + \frac{1}{2} k_{\rho} \left[\frac{\partial \mathscr{B}^{\mu\nu}}{\partial k_{\rho}} - \frac{\partial \mathscr{B}^{\rho\nu}}{\partial k_{\mu}} \right] (\{p_i\};0) + \mathscr{O}(k^2).$$
(123)

The first line of this equation already shows that the next-to-soft term is universal and proves the corresponding term in (116), while the next-to-next-to-soft $\mathcal{O}(k)$ terms in the second and third line still contain the anti-symmetric first-order derivatives of the non-singular part $\mathscr{B}^{\mu\nu}$ of the radiative amplitude.

It is worth noting that the previous equation can be adapted to gauge-boson rather than graviton emission by simply replacing $p_i^{\nu} \rightarrow t_i^a$ and removing the index ν on $\mathscr{A}_{rad}^{\mu\nu}$ and $\mathscr{B}^{\mu\nu}$, since the derivation goes through for non-abelian gauge invariance under shifts

$$\varepsilon_{\mu}(k) \to \varepsilon_{\mu}(k) + k_{\mu}\alpha(\{p_i\};k)$$
 (124)

of the emitted soft gauge boson polarisation vector. The key point about gravitation is that the radiative amplitude *also* satisfies $k_V \mathscr{A}_{rad}^{\mu\nu}(\{p_i\};k) = 0$, which can be verified for (123), but has not yet been used explicitly. This fixes the so far undetermined first derivatives in terms of the non-radiative amplitude:

$$\left[\frac{\partial \mathscr{B}^{\mu\nu}}{\partial k_{\rho}} - \frac{\partial \mathscr{B}^{\rho\nu}}{\partial k_{\mu}}\right] (\{p_i\}; 0) = \sum_{i=1}^{N} J_i^{\rho\mu} \frac{\partial}{\partial p_{i\nu}} \mathscr{A}(\{p_i\}).$$
(125)

Substituting into (123) yields the soft theorem (116) including three universal terms, that is, terms that can be expressed in terms of (derivatives of) the non-radiative

38

amplitude. The authors of [15] further checked that there are not enough constraints from gauge invariance to determine all second derivatives of $\mathscr{B}^{\mu\nu}$, hence the next $\mathscr{O}(k^2)$ term in the soft expansion is no longer universal.

The diagrammatic derivation of the soft theorem is remarkably simple on the one hand, and highlights the crucial role of gauge invariance. On the other hand, the number of universal terms appears from a rather technical argument and the derivation provides no physical understanding of why the derivative expansion organises itself such that the orbital angular momentum operator arises.

6.2 Soft theorem from SCET gravity

The soft-collinear effective Lagrangian by construction allows one to generate from its Feynman rules the soft and collinear limits of an amplitude to a desired accuracy in the soft and collinear expansion, and the loop expansion. However, it organises the derivation of the soft theorem in a different form from the above, since the soft factors arise entirely from the emission from external legs. For the case of gauge theory this was noted first in [7].¹⁸

To begin, it is instructive to formulate Weinberg's leading eikonal graviton emission amplitude (2) in the EFT language by recalling from (108), (98) that an energetic scalar with its large momentum p_i^{μ} directed along the light-like vector n_{i-}^{μ} interacts with a soft graviton through the effective Lagrangian¹⁹

$$\mathscr{L}_{i}^{(0)} = [n_{i+}\partial\chi_{c_{i}}]^{\dagger} \left[-\frac{\kappa}{4} n_{i-}^{\mu} n_{i-}^{\nu} s_{\mu\nu}(x_{i-}) i n_{i+}\partial \right] \chi_{c_{i}}.$$
 (126)

The structure of (2) is already manifest in this Lagrangian, which couples the energetic particle to the soft gauge field and graviton only proportionally to the large momentum $p_i^{\mu} \propto n_{i-}^{\mu}$. The content of the leading term in the soft theorem can now be stated in operatorial form as

$$\sum_{i} i \int d^4 x \, T\left\{\chi_{c_i}^{\dagger}(0), \mathscr{L}_i^{(0)}(x)\right\}\Big|_{\text{tree}},\tag{127}$$

where the sum over *i* runs over the energetic particles created in the hard process. The entire hard, non-radiative *N*-particle scattering process is sourced by a gaugeinvariant product of $N \chi_{c_i}^{\dagger}$ fields²⁰ as described in Section 5. Contracting $\chi_{c_i}^{\dagger}(0)$ with $\chi_{c_i}(x)$ in $\mathcal{L}_i^{(0)}$ to form the collinear matter propagator i/p^2 , and taking the

¹⁸ This statement holds in the position-space SCET framework. For a discussion of the LBK theorem in the so-called label formalism, see [29].

¹⁹ In the following, we drop the hats of the collinear fields. Furthermore, we consider a *complex* scalar field. This has the advantage that in the convention where all external particles are outgoing, the *N*-jet operator contains one field χ_c^{\dagger} for each particle. Thus, one can always identify χ_c^{\dagger} in the Lagrangian as the outgoing field, simplifying the following derivation.

²⁰ Employing the convention that all particles are out-going.



Fig. 6 Possible contributions to the radiative amplitude in SCET. The first diagram represents the time-ordered product of the current and the Lagrangian. Both current and Lagrangian can be suppressed by a single power of λ , or there can be a contribution from the λ^2 -suppressed Lagrangian together with the leading-power current. There are no soft building blocks at order λ^2 and λ^4 ; hence, the second diagram does not contribute to the amplitude in SCET.

matrix element with N matter particles and a soft graviton, immediately yields the amplitude (2). At this point, it is essential that soft gauge bosons and gravitons cannot be emitted directly from the hard vertex at this order in the soft expansion, since there are no source operators containing soft fields that would be invariant under the soft gauge symmetry. The entire radiative amplitude originates from the time-ordered product with the universal Lagrangian interaction. This guarantees the universality of the soft theorem, that is, its form is independent of the non-radiative, hard process. Briefly,

Lagrangian insertions \Rightarrow universal terms,

while the source operators in the effective theory have process-dependent matching coefficients to the full theory. Thus, when there is a source operator with N collinear fields and a soft graviton, it will contribute a non-universal term to the soft expansion of the radiative amplitude.

Unlike for amplitudes, in soft-collinear effective theory the separation of the amplitude into emission from the external leg and the hard vertex (see Figure 6) is gauge-invariant and corresponds to the separation into gauge-invariant Lagrangian insertions and gauge-invariant source operators with soft graviton fields. The number of universal terms in the soft expansion follows from this observation without the need for an explicit computation, since, as noted in Section 4.4, the leading gaugeinvariant soft building block for gravitons is the Riemann tensor, which scales as $\mathcal{O}(\lambda^6)$, corresponding to a third-order correction in the soft expansion. It is this simple consequence of the soft gauge symmetry of the effective Lagrangian, which implies that there is some form of universal soft theorem including a universal nextto-next-to-soft term for gravity, as any soft emission up these orders must arise from universal Lagrangian terms, independent of the source for the energetic particles. The additional universal term compared to gauge boson emission follows from the observation that the leading soft building block for soft gauge fields is the field strength tensor $F_{\mu\nu}^s$, which scales as $\mathcal{O}(\lambda^4)$, hence there are non-universal contributions already at the next-to-next-to-soft order.

This answers the first question. To gain a better understanding of the detailed form of the soft theorem, in particular the appearance of the angular momentum operator in the sub-leading terms, one must investigate the structure of the effective Lagrangian. As before, the specific case of scalar matter will be considered. Since the soft theorem is a statement about tree-level, single graviton emission, two simplifications can be made in the general soft-collinear gravity Lagrangian:

- The collinear graviton field can be set to zero, since there are no internal or external collinear graviton lines.
- Only linear interactions in the soft graviton field need to be retained.

In the remainder of this subsection, the basic ideas will be explained for the nextto-soft term, which counts as $\mathcal{O}(\lambda^2)$. For the derivation of the sub-sub-leading soft term and more technical detail, we refer to [9]. This reference also discusses the case of non-abelian gauge theory, including the case of matter with spin $\frac{1}{2}$ and 1, which demonstrates how the spin term $\Sigma^{\mu\nu}$ in $J^{\mu\nu}$ is encoded in the effective Lagrangian.

The soft-collinear effective Lagrangian for the complex gauge-invariant scalar field χ_c up to $\mathcal{O}(\lambda^2)$, with simplifications as stated above already applied, can be expressed in terms of the energy-momentum tensor (dropping the index *i* when referring to the Lagrangian, since its form is the same for all *i*)

$$T^{\mu\nu} = \left[\partial^{\mu}\chi_{c}\right]^{\dagger}\partial^{\nu}\chi_{c} + \left[\partial^{\nu}\chi_{c}\right]^{\dagger}\partial^{\mu}\chi_{c} - \eta^{\mu\nu}\left[\partial_{\alpha}\chi_{c}\right]^{\dagger}\partial^{\alpha}\chi_{c}$$
(128)

as

$$\mathscr{L} = \frac{1}{2} \left[n_+ \partial \chi_c \right]^{\dagger} n_- \partial \chi_c + \frac{1}{2} \left[n_- \partial \chi_c \right]^{\dagger} n_+ \partial \chi_c + \left[\partial_{\mu_\perp} \chi_c^{\dagger} \right] \partial^{\mu_\perp} \chi_c$$
(129)
$$- \frac{\kappa}{4} s_{-\mu} T^{\mu_+} - \frac{\kappa}{4} \left[\partial_{[\mu} s_{\nu]-} \right] (x - x_-)^{\mu} T^{\nu_+} - \frac{1}{8} x_{\perp}^{\alpha} x_{\perp}^{\beta} R_{\alpha-\beta-}^s T^{++} + \mathscr{O}(x^3).$$

When no space-time argument is specified, all soft fields are evaluated at x_{-}^{μ} after derivatives are taken. In this form, the coupling of $s_{\mu-}$ to the energy-momentum tensor $T^{\mu\nu}$ becomes transparent, as well as the coupling of its derivative $\partial_{[\alpha}s_{\beta]-}$, which is an independent gauge field in the effective theory, to the angular momentum density

$$\mathscr{J}^{\alpha\beta\mu} = (x - x_{-})^{\alpha} T^{\beta\mu} - (x - x_{-})^{\beta} T^{\alpha\mu} .$$
(130)

The Lagrangian (129) is not yet homogeneous in λ . The λ expansion is made manifest by decomposing the contraction of the indices μ, ν in light-cone components. Then

$$\mathscr{L} = \mathscr{L}^{(0)} + \mathscr{L}^{(1)} + \mathscr{L}^{(2)} + \mathscr{O}(\lambda^3), \qquad (131)$$

where

$$\mathscr{L}^{(0)} = \frac{1}{2} \left[n_+ \partial \chi_c \right]^{\dagger} n_- \partial \chi_c + \frac{1}{2} \left[n_- \partial \chi_c \right]^{\dagger} n_+ \partial \chi_c + \left[\partial_{\mu_\perp} \chi_c \right]^{\dagger} \partial^{\mu_\perp} \chi_c - \frac{\kappa}{8} s_{--} T^{++},$$
(132)

Martin Beneke, Patrick Hager and Robert Szafron

$$\mathscr{L}^{(1)} = -\frac{\kappa}{4} s_{-\mu_{\perp}} T^{\mu_{\perp}+} - \frac{\kappa}{8} \left[\partial_{[\mu} s_{-]-} \right] x_{\perp}^{\mu} T^{++}, \qquad (133)$$

$$\mathscr{L}^{(2)} = -\frac{\kappa}{8} s_{+-} T^{+-} - \frac{\kappa}{4} \left[\partial_{[\mu_{\perp}} s_{\nu_{\perp}]-} \right] x_{\perp}^{\mu} T^{\nu_{\perp}+} - \frac{\kappa}{16} \left[\partial_{[+} s_{-]-} \right] n_{-} x T^{++} - \frac{1}{8} x_{\perp}^{\alpha} x_{\perp}^{\beta} R^{s}_{\alpha-\beta-} T^{++} .$$
(134)

The source operator $\hat{\mathscr{A}}$ that represents the hard scattering process also needs to be expanded in λ :

$$\hat{\mathscr{A}} = \hat{\mathscr{A}}^{(0)} + \hat{\mathscr{A}}^{(1)} + \hat{\mathscr{A}}^{(2)} + \mathscr{O}(\lambda^3 \hat{\mathscr{A}}^{(0)}).$$
(135)

The $\mathscr{A}^{(n)}$ up to this order are operators containing $i\partial_{\perp i}^{\mu_1} \dots i\partial_{\perp i}^{\mu_n} \chi_{c_i}^{\dagger}$, which corresponds to the expansion of the non-radiative momentum-space hard-scattering amplitude in the transverse momentum $p_{i\perp}^{\mu}$ of particle *i*. The operatorial version of the soft theorem then amounts to the expansion in λ of the right-hand side of

$$\hat{\mathcal{A}}_{\text{rad}} = i \int d^4 x \, T\left\{\hat{\mathcal{A}}, \mathcal{L}\right\}$$
(136)

in the tree-level approximation and under the assumption that the operator is evaluated in the state $\langle \{p_i\}; k | \hat{\mathcal{A}}_{rad} | 0 \rangle$.

It is always possible to define the *N* light-cone vectors n_{i-}^{μ} such that the *external* energetic matter particle momenta of the radiative amplitude satisfy $p_{i\perp}^{\mu} = 0$ for all *i* with respect to their light-cone vectors. This choice simplifies the calculations considerably. Many terms vanish, which would otherwise contribute to restoring Lorentz-invariant scalar products of the external momenta. These scalar products do not depend on the specific choice of the reference vectors that break manifest Lorentz invariance. In particular, one finds that

$$T\left\{\hat{\mathscr{A}}^{(n)},\mathscr{L}^{(0)}\right\}, T\left\{\hat{\mathscr{A}}^{(0)},\mathscr{L}^{(1)}\right\}$$
(137)

evaluate to zero on the right-hand side of (136), leaving²¹

$$\widehat{\mathscr{A}}_{\mathrm{rad}}^{(2)} \stackrel{\circ}{=} i \int d^4 x \, T\left\{\widehat{\mathscr{A}}^{(1)}, \mathscr{L}^{(1)}\right\} + i \int d^4 x \, T\left\{\widehat{\mathscr{A}}^{(0)}, \mathscr{L}^{(2)}\right\}$$
(138)

for the next-to-soft term.

Examining (133), (134) one finds that the Riemann-tensor term in the last line of (134) does not contribute to the time-ordered product with $\hat{\mathscr{A}}^{(0)}$ in the frame $p_{i\perp}^{\mu} = 0$, while the remaining terms and $\mathscr{L}^{(1)}$ can be rewritten after integrating by parts into

42

²¹ The symbol $\stackrel{\circ}{=}$ means "equal up to terms that do not contribute to the specific tree-level matrix elements considered here, i.e. with single soft emission, no collinear emissions, and \perp -component of the external collinear particle momenta set to zero."

$$\mathscr{L}^{(1)} \stackrel{\scriptscriptstyle \triangle}{=} \frac{\kappa}{2} \left[\partial_{\mu} s_{\nu} \right] \chi_{c}^{\dagger} \stackrel{\leftarrow}{L}_{+\perp}^{\mu\nu} n_{+} \partial \chi_{c} , \qquad (139)$$

$$\mathscr{L}^{(2)} \stackrel{\scriptscriptstyle \triangle}{=} \frac{\kappa}{2} \left[\partial_{\mu} s_{\nu} \right] \chi_{c}^{\dagger} \stackrel{\leftarrow}{L}_{+-}^{\mu\nu} n_{+} \partial \chi_{c} , \qquad (140)$$

where

$$L^{\mu\nu} = x^{[\mu}\partial^{\nu]} = \frac{1}{2}x^{[\mu}n_{-}^{\nu]}n_{+}\cdot\partial + \dots = \underbrace{\frac{1}{4}n_{-}\cdot xn_{+}\cdot\partial n_{+}^{[\mu}n_{-}^{\nu]}}_{L_{+-}^{\mu\nu}} + \underbrace{\frac{1}{2}x_{\perp}^{[\mu}n_{-}^{\nu]}n_{+}\cdot\partial}_{L_{++}^{\mu\nu}} \quad (141)$$

is the orbital angular momentum operator in position space and the dots after the first equality represent terms that vanish in the $p_{\perp}^{\mu} = 0$ frame. Both time-ordered products in (138) can now be combined into

$$\hat{\mathscr{A}}_{\rm rad}^{(2)} = \int d^4 x \, T \left\{ \hat{\mathscr{A}}, \frac{\kappa}{2} \chi_c^{\dagger} \overleftarrow{L}^{\mu\nu} \left[\partial_{\mu} s_{\nu-} \right] i n_+ \partial \chi_c \right\}, \tag{142}$$

which is precisely the next-to-soft term in (116). This answers the second question: the angular momentum operator is seen to appear since the soft-collinear expansion of the Lagrangian together with the light-cone multipole expansion of soft interactions with energetic particles naturally provide the right structures. At the next-to-soft order, the appearance of the angular momentum is already evident in (129) and related to the soft gauge symmetry of the effective Lagrangian.

This observation provides a deeper understanding of the gravitational soft theorem. Unlike the gauge theory case, where the next-to-soft term in the LBK theorem can be shown to be related to the soft field-strength tensor $F_{\mu\nu}^s$ [9], and is therefore gauge-invariant without requiring a conserved quantity in the scattering process, the next-to-soft term in gravity takes the form of an *eikonal term*, just as the leading term. The leading term

$$\varepsilon_{\mu-} p^{\mu} \frac{n_+ p}{p \cdot k} \tag{143}$$

is related to the coupling of the first soft gauge field $s_{-\mu}$ to the energy momentum tensor $T^{\mu+}$ in (129). It is gauge-invariant due to the conservation of momentum. The first sub-leading order in the soft theorem,

$$k_{\rho}\varepsilon_{\mu-}J^{\rho\mu}\frac{n_{+}p}{p\cdot k},\qquad(144)$$

is generated by the next term in (129). It involves the second soft gauge field, $\partial_{[\mu}s_{\nu]-} \equiv \partial_{\mu}s_{\nu-} - \partial_{\nu}s_{\mu-}$ (or $[\Omega_{-}]_{\mu\nu}$, see (96)), which couples to the energymomentum density. The next-to-soft term for gravity is only gauge-invariant once angular momentum conservation of the scattering process is imposed. Hence it is the two-fold emergent soft gauge symmetry that implies two eikonal terms in the soft theorem, consistent with the covariant derivative

$$n_{-}D_{s} = \partial_{-} - \frac{\kappa}{2}s_{-\mu}\partial^{\mu} - \frac{\kappa}{4}\partial_{[\mu}s_{\nu]-}J^{\mu\nu} + \dots$$
(145)

43

Finally, the Riemann-tensor terms, which are present in the sub-leading softcollinear interaction Lagrangians $\mathscr{L}^{(2)}, \mathscr{L}^{(3)}$ and $\mathscr{L}^{(4)}$, generate the next-to-nextto-soft term [9]

$$\frac{1}{2}\varepsilon_{\mu\nu}k_{\rho}k_{\sigma}J^{\rho\mu}\frac{J^{\sigma\nu}}{p\cdot k}$$
(146)

in (116). Here, one factor of $J^{\mu\nu}$ is related to the charge of one of the soft gauge symmetries, while the other arises from the kinematic multipole expansion. Originating from the Riemann tensor, this term is manifestly gauge-invariant without requiring further (non-existent) conserved charges. Thus, despite appearance, the *next-to-next-to soft term* in gravity has the same physical origin as the *next-to-soft term* in the LBK theorem, while the soft *and* next-to-soft term should be viewed as the gravitational analogues of the familiar eikonal factor in gauge theories.

6.3 Loop corrections to the soft theorem

While the soft theorem is agnostic about the details of the hard non-radiative amplitude that produces the N energetic particles (in widely separated directions), it holds only at tree-level in the interactions of the energetic particles with soft gravitons. Soft-collinear gravity generates the soft and collinear loops in gravitational scattering to any order, so it is natural to ask whether it provides insight on the loop corrections to the soft theorem. In this subsection, we will show that:

The leading soft factor is not modified by loop effects. The sub-leading factor is only corrected by one-loop, and the sub-sub-leading factor is only modified by one- and two-loop contributions. Higher-order loop corrections do not affect the gravitational soft theorem.

This should be compared to non-abelian gauge theory, where already the leading term receives loop corrections of any order. The above statement was already made in [14], but the reasoning from EFT provided in [9] and below is somewhat different, and relies only on 1) power counting, 2) the eikonal identity, and 3) the necessity to form a soft invariant from a given soft loop integral, as otherwise the integral is scaleless and vanishes. It should be noted that the radiative amplitude is infrared-divergent at loop level, and a regularisation is needed. The above statement holds when singularities are regulated dimensionally in $d = 4 - 2\varepsilon$ dimensions.

In SCET, loop contributions arise from three different loop momentum regions, the hard, the collinear and the soft region, the latter two corresponding to collinear and soft modes in the effective theory. The hard modes are integrated out, thus the contributions of the hard loops are inside the matching coefficients $\tilde{C}^{X}(t_i)$ and consequently part of the non-radiative amplitudes $\langle p_1, \ldots, p_N | \mathcal{J}(0) | 0 \rangle$ defined in Section 5. Hence, hard loops never affect the soft theorem directly—they modify the underlying non-radiative process.



Fig. 7 Collinear one-loop corrections. Left: Collinear graviton from the source in one of the N given directions. Middle and right: Collinear loop corrections to an external leg. Two further diagrams similar to the first two but with the soft graviton attached to the collinear graviton line are not shown.

Soft-collinear gravity differs from gauge theory in the purely soft and collinear sectors, ultimately due to the dimensionful gravitational coupling:

- *i*) There are no collinear singularities. In the purely-collinear sector, that is, in the Lagrangian terms containing only collinear but no soft fields, there are no leading-power interactions. The λ -expansion corresponds to the weak-field expansion, and the first collinear interaction appears in $\mathcal{O}(\lambda)$. Purely-collinear gravity is an expansion in collinear momenta $p_{\perp} \sim \lambda$.
- *ii*) In the purely-soft sector, that is, in the Lagrangian terms containing only soft but no collinear fields, there are also no leading-power interactions. Here, the weak-field expansion agrees with the λ^2 -expansion, corresponding to an expansion in soft momenta $k \sim \lambda^2$. Purely-soft interaction vertices thus start at $\mathcal{O}(\lambda^2)$.

Hence, whenever a purely-collinear or a purely-soft interaction takes place, the contribution is already suppressed by at least one order of λ or λ^2 , respectively. In gravity, only soft-collinear interactions of the eikonal type exist at leading power.

Consider first adding only collinear loops to the single-soft emission amplitude. At the one-loop level, there exist two possibilities, shown in Figure 7. First, a collinear graviton field $\mathfrak{h}_{\perp\perp}$ in a given collinear direction n_{i-}^{μ} can be added to the source, which comes with an extra power of λ for the hard amplitude. The collinear graviton must be attached to the external leg *i* through a collinear interaction vertex, which costs at least another power of λ , resulting in at least $\mathcal{O}(\lambda^2)$ suppression of the collinear loop. Second, the collinear loop correction can be on the external leg *i* only with no connection to the hard vertex, in which case one either needs two purely-collinear vertices of at least $\mathcal{O}(\lambda)$ or a four-point vertex, which is already $\mathcal{O}(\lambda^2)$. By the same argument, adding any further collinear loop incurs at least another factor $\mathscr{O}(\lambda^2)$ per loop. It follows that the leading term in the soft theorem can never be corrected, while a collinear *n*-loop correction can contribute only at the *n*th sub-leading order to the soft theorem, consistent with the claim. Note that the argument holds for every collinear direction *i* separately, since the collinear modes from different directions cannot interact other than through soft modes, as evident from (41), which implies soft loops.

The case of soft loops is less straightforward. Since there are no hard vertices with soft fields up to sub-sub-leading order, all soft loops to this order must be built



Fig. 8 Diagram classes modifying the soft emission process at the one-loop order. The softcollinear interactions in the first two diagrams are present at leading power in λ . However, due to multipole expansion, soft-collinear vertices are only sensitive to the n_{i-k} component, and the soft loop vanishes unless a soft scale, provided by the injection of the full soft momentum k, is present. This can only happen by a purely-soft interaction vertex. Hence, only the right-most diagram is non-vanishing, but is at least $\mathcal{O}(\lambda^2)$ suppressed, since it contains a purely-soft interaction.

from Lagrangian interactions. We first consider the case that all loops are soft and none is collinear.

Beginning with one-loop soft corrections, the soft loop can connect at most two collinear directions with topologies as shown in Figure 8, and a similar set of diagrams, where the soft loop is attached to a single leg only. These contributions vanish unless the external soft graviton is connected to the loop through a purely-soft interaction, as shown in the right-most diagram in the figure.²² The key point is that soft fields are always multipole-expanded in interactions with collinear fields, which implies that for soft momentum p_s , only $(n_i - p_s) n_{i+}^{\mu}/2$ enters the momentum-conservation delta-function at a soft-collinear vertex and hence the collinear propagators in direction *i*. The loop depicted in Figure 8, with the soft graviton emission removed, is given by the (dimensionally-regulated) integral

$$I \propto \int \frac{d^d l}{(2\pi)^d} \frac{1}{p_i^2 + n_{i+} p_i n_{i-} l + i0} \frac{1}{p_j^2 - n_{j+} p_j n_{j-} l + i0} \frac{1}{l^2 + i0}, \qquad (147)$$

which for on-shell external particles, $p_i^2 = p_j^2 = 0$, simplifies to

$$I \propto \int \frac{d^d l}{(2\pi)^d} \frac{1}{l^2 + i0} \frac{1}{n_{i-}l + i0} \frac{1}{n_{j-}l + i0} \,. \tag{148}$$

This integral is evidently scaleless, and vanishes. Note that the diagrams of Figure 8 provide numerators for this integral, which are polynomial in the external and the loop momentum. They have not been written explicitly, since whether an integral is scaleless or not is independent of such numerators.

If one now attaches the external soft graviton to one of the collinear lines (see first three diagrams in Figure 8), one can always route the external soft momentum *k* such that it appears in the eikonal propagators of only one of the legs, say *i*. In this way, the loop integral (148) may be modified to include eikonal propagators of the form $(n_{i-}(l+k)+i0)^{-1}$. Since only the $n_{i-}kn_{i+}^{\mu}/2$ component of the

²² See Appendix B of [4] for a similar discussion for soft interactions in gauge theory at next-toleading power.



Fig. 9 Examples of soft two-loop diagrams. Left: Diagram with soft emission from a collinear line. Middle: Two diagrams with soft-collinear interactions and a purely-soft vertex. Right: A non-vanishing soft two-loop diagram. In order for the soft scale *k* to be present in both loops, the loops must be connected via purely-soft interactions to the emitted graviton, which results in a contribution at sub-sub-leading order due to λ^4 suppression.

soft momentum can ever appear in the denominator, one cannot form an invariant scalar product containing k with the required soft scaling $\mathcal{O}(\lambda^4)$ (as $n_{i+}^2 = 0$) and the soft loop integral will remain scaleless and vanishing. In order for soft loops to yield a non-zero contribution, one needs to bring the full external soft momentum k^{μ} into the loop integral. This requires the external soft graviton to couple to the loop through a purely-soft interaction (as in the right-most diagram in the figure). Such interaction vertices involve the full momentum conservation delta function and lead to propagators $1/(l+k)^2$, which allows the soft integral to depend on the soft invariant $(n_{i-k})(n_{j-k})(n_{i+n_{j+1}})$ and be non-zero. However, by point *ii*) above, such a purely-soft vertex comes at the cost of power-suppression by at least λ^2 . Hence, soft one-loop corrections also cannot affect the leading term in the soft theorem, however, the next-to-soft term can be modified by diagrams with the external soft graviton attached to a purely-soft vertex.

The above argument generalises to the following all-order statement: In soft loopcorrections to the soft theorem, contrary to the tree-level case, the emitted soft graviton must always attach to a purely-soft vertex, and never directly to any of the energetic particle lines. The reason is that soft-collinear interactions involve the soft field at the multipole-expanded point x_{-}^{μ} to any order in the λ -expansion. Hence, if the emitted graviton couples directly to an energetic line, one can always route its momentum such that the entire loop integral will depend only on $n_{i-k}n_{i+}^{\mu}/2$ of a single collinear direction, *i*, and no soft invariant can be formed to provide a scale to the loop diagram.

Continuing with two soft loops, whenever the diagram contains a second purelysoft vertex (as in the right-most diagram of Figure 9), there is another $\mathcal{O}(\lambda^2)$ suppression factor from this vertex and the diagram can at best contribute to the subsub-leading soft factor, in agreement with the assertion. This conclusion can potentially be evaded by coupling the soft lines to the energetic lines with $\mathcal{O}(\lambda^0)$ or $\mathcal{O}(\lambda^1)$ vertices, which adds only linear eikonal-type propagators, possibly raised to higher powers due to the multipole expansion. The argument in the previous paragraph excludes coupling the emitted soft graviton to an energetic line (as in the left-most diagram of Figure 9), but the middle two graphs in the figure are still an option. We next show that these graphs obtain another $\mathcal{O}(\lambda^2)$ suppression from the soft-collinear vertices on the energetic lines and they can only contribute to the sub-sub-leading term in the soft theorem consistent with the claim.²³

Consider first the case that all soft-graviton couplings to the energetic particle lines are through the leading-power soft-collinear interaction $\frac{\kappa}{4}s_{--}\chi_c^{\dagger}(n_+\partial^2\chi_c)$ from (132). This gives rise to the eikonal interaction, which has the well-known property [39] that multiple emissions take the form of *independent* emissions after summing over all possible attachments to the energetic lines. For i = 1, ..., N such lines with external momenta $\{p_i\}$, the emission of an arbitrary number of soft gravitons with momenta $\{q_k\}$ is given by the non-radiative amplitude multiplied by the eikonal factor:

$$\mathscr{A}(\{p_i\})\sum_{m=0}^{\infty}\frac{1}{m!}\prod_{k=1}^{m}\left(\sum_{i=1}^{N}\frac{\kappa}{2}\frac{p_i^{\mu_k}p_i^{\nu_k}\varepsilon_{\mu_k\nu_k}(k)}{p_i\cdot q_k+i\varepsilon}\right).$$
(149)

To construct a loop diagram with a single emitted soft graviton, one must tie together the q_k amongst each other with a soft graviton propagator or with the purelysoft vertex from which the external graviton is emitted. The key observation is that due to the factorised form of the denominator of the eikonal amplitude (149), the external momentum k does not enter the loop integral of those pairs of q_k tied together directly, which therefore can only depend on the energetic momenta p_i . Now, a dimensionally-regulated soft loop integral must be proportional to $(S/\mu^2)^{\varepsilon}$, where S is a soft invariant, that is, a scalar product of momenta, which scales as λ^4 , and has mass dimension two. Since no such invariant can be built from the available momenta as $p_i \cdot p_j \sim \lambda^0$ and $p_i^2 = 0$, these integrals must be scaleless and therefore the entire graph vanishes. This applies to the two middle diagrams in Figure 9 when all soft-collinear vertices are from $\mathscr{L}^{(0)}$, which would otherwise contribute at $\mathcal{O}(\lambda^2)$. Note that this vanishing also holds if some of the q_k are tied together through purely-soft vertices, as long as the emitted graviton is not attached to them, since in this case k^{μ} never enters the corresponding subgraph. This results in the following all-order statement: If a loop integral involves only the leading-power eikonal interactions from $\mathscr{L}^{(0)}$, it vanishes unless all soft propagators are connected to the external graviton line through purely-soft vertices. As a consequence, any non-vanishing soft *n*-loop graph has a suppression of at least $\mathcal{O}(\lambda^{2n})$ from the combination of the required purely-soft vertices.

There is still the possibility that the emitted graviton couples to an energetic line through some of the sub-leading in λ soft-collinear interactions from $\mathscr{L}^{(1)}$ (109) and $\mathscr{L}^{(2)}$ (110). The vertices from $\mathscr{L}^{(1)}$ contain a single transverse index. In the frame $p_{i\perp}^{\mu} = 0$, there is no transverse external vector available, hence diagrams with

²³ This and the following arguments fill a loop-hole in the discussion of soft loops in [14], where it is assumed that the dimensionful coupling κ that comes with a loop must be compensated by a soft momentum $k \sim \mathcal{O}(\lambda)$, so a k-soft-loop diagram must be $\mathcal{O}(\lambda^{2k})$. However, in soft-collinear interactions, the factor of κ can be compensated by powers of the $\mathcal{O}(1)$ collinear momenta as there exists a $\mathcal{O}(\lambda^0)$ vertex. In other words, for soft-collinear interactions the loop-expansion in κ is not equivalent to a λ^2 -expansion.



Fig. 10 $\mathscr{O}(\lambda^4)$ diagram with one purely-soft vertex and sub-leading soft-collinear interactions on line *j*. Except for the emitted graviton with momentum *k*, all other wavy lines have to be tied together amongst each other.

only a single $\mathscr{L}^{(1)}$ insertion on an energetic line vanish. It already follows that one cannot evade the λ^2 suppression per loop by the use of sub-leading soft-collinear interactions and the two-loop diagrams are all at least of order $\mathscr{O}(\lambda^4)$. However, these diagrams actually vanish. At the two-loop order, the only non-vanishing topologies contain two purely-soft vertices connected by a soft propagator as in the right-most diagram of Figure 9, or a purely-soft four-graviton vertex, which also scales at least as $\mathscr{O}(\lambda^4)$.

To see this, assume that the single purely-soft three-graviton vertex connects the emitted graviton to the energetic legs *i* and *j*. Since the sub-leading eikonal vertices that provide further $\mathcal{O}(\lambda^2)$ suppression reside on a single energetic leg, one can further assume without loss of generality that line *i* has only leading-power eikonal vertices attached to it, and then route the external momentum k^{μ} through this energetic leg. Leg *i* is therefore an eikonal leg, that is, after summing over all permutations of attached momenta, the amplitude for this leg takes the form of independent emissions. All other external energetic lines also have this property, except for the one with sub-leading soft-collinear vertices, which may or may not be leg *j*. The situation and momentum assignments are illustrated in Figure 10.

At the two-loop order there is only one pair of open lines to be tied up with a soft graviton propagator, but for later purposes it is convenient to proceed with the general situation shown in the figure. Because leg i is an eikonal leg, the momenta k and q appear in the entire diagram only in denominators

$$\frac{1}{(k+q)^2 q^2} \frac{1}{n_{i-} \cdot (k+q)} \times \prod_k \frac{1}{n_{j-} \cdot L_k(q, l_j)},$$
(150)

where $L_k(q, l_j)$ is the linear sum of a subset (or all) of the momenta attached to leg *j*. Note that this expression is general, since it was not assumed that leg *j* is eikonal. Let *l* be one of the momenta l_j . The loop integral over *l* is then of the form Martin Beneke, Patrick Hager and Robert Szafron

$$\int \frac{d^d l}{(2\pi)^d} \frac{1}{l^2} \frac{1}{n_{a-} \cdot l} \prod_k \frac{1}{n_{j-} \cdot L_k(q, l_j)},$$
(151)

where $a \neq j$ refers to the eikonal leg to which the graviton line *l* is attached and we used that the emissions are independent from this leg. After integrating over *l*, the result can only depend on $(n_{j-q})n_{j+}^{\mu}/2$ and $(n_{j-l_j})n_{j+}^{\mu}/2$, from which one cannot form a non-vanishing invariant, hence the integral is scaleless, and consequently the entire diagram is zero. If a = j, the expression (151) without the factor $1/(n_{a-} \cdot l)$ holds, and the same conclusion is reached.

To complete the proof of the main assertion for soft-loop corrections, it remains to show that three- and higher-loop diagrams are inevitably more suppressed than $\mathscr{O}(\lambda^4)$, so that they cannot affect the soft theorem to the sub-sub-leading order. The preceding two all-order statements imply that a diagram that is not already excluded by them must 1) have exactly one purely-soft vertex of $\mathscr{O}(\lambda^2)$, to which the external soft graviton is attached, and 2) $\mathscr{O}(\lambda^2)$ suppression from sub-leading soft-collinear interactions on a single energetic leg, say *j*, plus 3) an arbitrary number of additional leading-power eikonal interactions. The property of independent soft emission therefore applies to all energetic lines except *j*. However, this case is covered by the arguments of the previous paragraph, which holds to any loop order.

Having established suppression of either only collinear or only soft loop corrections in accordance with the claim, we finally turn to arbitrary loop diagrams that can have both, which start from two loops. It is easy to see that two-loop diagrams must be at least $\mathcal{O}(\lambda^4)$, since the external soft graviton must still attach to a purelysoft collinear vertex by the same argument as before, while the collinear loop still costs a factor of λ^2 . The case of higher-loop diagrams is covered by the argument of the previous paragraph, which continues to hold when the leg with the sub-leading soft-collinear Lagrangian insertions is replaced by the leg, to which a collinear loop is attached. Any mixed collinear-soft loop diagram with more than two collinear loops is already beyond the next-to-next-to-soft order and does not have to be considered. This concludes the characterisation of the structure of loop corrections to the soft theorem, as formulated at the beginning of this subsection.

The one- and two-loop corrections to the soft theorem are largely universal, since they arise almost exclusively from effective Lagrangian terms. The non-universal part can be parameterised in terms of the matching coefficients $\tilde{C}^{X}(t_{i})$ of SCET source operators with up to two additional collinear transverse graviton fields. An explicit calculation of the loop-corrected soft factors has, however, not yet been performed.

Acknowledgement

This work was supported by the Excellence Cluster ORIGINS funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Grant No. EXC - 2094 - 390783311. RS is supported by the United States Department of Energy under Grant Contract DE-SC0012704. MB thanks the Albert

Einstein Center at the University of Bern for hospitality while this work was finalised.

References

- R. Akhoury, R. Saotome and G. Sterman, Phys. Rev. D 84, 104040 (2011) [arXiv:1109.0270 [hep-th]].
- [2] C. W. Bauer, S. Fleming, D. Pirjol and I. W. Stewart, Phys. Rev. D 63, 114020 (2001) [arXiv:hep-ph/0011336].
- [3] C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D 65, 054022 (2002) [arXiv:hep-ph/0109045].
- [4] M. Beneke, A. Broggio, S. Jaskiewicz and L. Vernazza, JHEP 07, 078 (2020) [arXiv:1912.01585 [hep-ph]].
- [5] M. Beneke, A. P. Chapovsky, M. Diehl and T. Feldmann, Nucl. Phys. B 643, 431-476 (2002) [arXiv:hep-ph/0206152].
- [6] M. Beneke and T. Feldmann, Phys. Lett. B 553, 267-276 (2003) [arXiv:hepph/0211358].
- [7] M. Beneke, M. Garny, R. Szafron and J. Wang, PoS RADCOR2017, 048 (2017) [arXiv:1712.07462 [hep-ph]].
- [8] M. Beneke, P. Hager and A. F. Sanfilippo, JHEP 02, 083 (2022) [arXiv:2106.09054 [hep-th]].
- [9] M. Beneke, P. Hager and R. Szafron, JHEP 03, 199 (2022) [arXiv:2110.02969 [hep-th]].
- [10] M. Beneke, P. Hager and R. Szafron, JHEP 03, 080 (2022) [arXiv:2112.04983 [hep-ph]].
- [11] M. Beneke and G. Kirilin, JHEP 09, 066 (2012) [arXiv:1207.4926 [hep-ph]].
- [12] Z. Bern, J. J. M. Carrasco and H. Johansson, Phys. Rev. D 78, 085011 (2008) [arXiv:0805.3993 [hep-ph]].
- [13] Z. Bern, J. J. Carrasco, M. Chiodaroli, H. Johansson and R. Roiban, [arXiv:1909.01358 [hep-th]].
- [14] Z. Bern, S. Davies and J. Nohle, Phys. Rev. D 90, no.8, 085015 (2014) [arXiv:1405.1015 [hep-th]].
- [15] Z. Bern, S. Davies, P. Di Vecchia and J. Nohle, Phys. Rev. D 90, no.8, 084035 (2014) [arXiv:1406.6987 [hep-th]].
- [16] F. Bloch and A. Nordsieck, Phys. Rev. 52, 54-59 (1937)
- [17] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, Proc. Roy. Soc. Lond. A 269, 21-52 (1962)
- [18] J. Broedel, M. de Leeuw, J. Plefka and M. Rosso, Phys. Rev. D 90, no.6, 065024 (2014) [arXiv:1406.6574 [hep-th]].
- [19] T. H. Burnett and N. M. Kroll, Phys. Rev. Lett. 20, 86 (1968)
- [20] F. Cachazo and A. Strominger, [arXiv:1404.4091 [hep-th]].
- [21] E. Casali, JHEP 08, 077 (2014) [arXiv:1404.5551 [hep-th]].
- [22] S. Coleman and R. E. Norton, Nuovo Cim. 38, 438-442 (1965)

- [23] J. C. Collins, D. E. Soper and G. F. Sterman, Adv. Ser. Direct. High Energy Phys. 5, 1-91 (1989) [arXiv:hep-ph/0409313 [hep-ph]].
- [24] W. Donnelly and S. B. Giddings, Phys. Rev. D 93, no.2, 024030 (2016) [erratum: Phys. Rev. D 94, no.2, 029903 (2016)] [arXiv:1507.07921 [hep-th]].
- [25] J. F. Donoghue, Phys. Rev. D 50, 3874-3888 (1994) [arXiv:gr-qc/9405057].
- [26] J. F. Donoghue, M. M. Ivanov and A. Shkerin, [arXiv:1702.00319 [hep-th]].
- [27] H. Elvang and Y. t. Huang, [arXiv:1308.1697 [hep-th]].
- [28] T. Kinoshita, J. Math. Phys. 3, 650-677 (1962)
- [29] A. J. Larkoski, D. Neill and I. W. Stewart, JHEP 06, 077 (2015) [arXiv:1412.3108 [hep-th]].
- [30] T. D. Lee and M. Nauenberg, Phys. Rev. 133, B1549-B1562 (1964)
- [31] F. E. Low, Phys. Rev. 110, 974-977 (1958)
- [32] S. G. Naculich and H. J. Schnitzer, JHEP **05**, 087 (2011) [arXiv:1101.1524 [hep-th]].
- [33] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Fortsch. Phys. 32, 585 (1984)
- [34] T. Okui and A. Yunesi, Phys. Rev. D **97**, 066011 (2018) [arXiv:1710.07685 [hep-th]].
- [35] R. K. Sachs, Proc. Roy. Soc. Lond. A 270, 103-126 (1962)
- [36] A. Strominger, JHEP 07, 152 (2014) [arXiv:1312.2229 [hep-th]].
- [37] A. Strominger, "Lectures on the Infrared Structure of Gravity and Gauge Theory," [arXiv:1703.05448 [hep-th]].
- [38] S. Weinberg, Phys. Rev. 140, B516-B524 (1965)
- [39] S. Weinberg, "The Quantum theory of fields. Vol. 1: Foundations," Cambridge University Press, 2005.
- [40] C. D. White, JHEP 05, 060 (2011) [arXiv:1103.2981 [hep-th]].
- [41] M. Zlotnikov, JHEP 10, 148 (2014) [arXiv:1407.5936 [hep-th]].