

Efficient surrogate-assisted inference for patient-reported outcome measures with complex missing mechanism

Jaeyoung Park^{*} Muxuan Liang[†] Ying-Qi Zhao[‡] Xiang Zhong[§]

Abstract

Patient-reported outcome (PRO) measures are increasingly collected as a means of measuring healthcare quality and value. The capability to predict such measures enables patient-provider shared decision making and the delivery of patient-centered care. However, due to their voluntary nature, PRO measures often suffer from a high missing rate, and the missingness may depend on many patient factors. Under such a complex missing mechanism, statistical inference of the parameters in prediction models for PRO measures is challenging, especially when flexible imputation models such as machine learning or nonparametric methods are used. Specifically, the slow convergence rate of the flexible imputation model may lead to non-negligible bias, and the traditional missing propensity, capable of removing such a bias, is hard to estimate due to the complex missing mechanism. To efficiently infer the parameters of interest, we propose to use an informative surrogate that can lead to a flexible imputation model lying in a low-dimensional subspace. To remove the bias due to the flexible imputation model, we identify a class of weighting functions as alternatives to the traditional propensity score and estimate the low-dimensional one within the identified function class. Based on the estimated low-dimensional weighting function, we construct a one-step debiased estimator without using any information of the true missing propensity. We establish the asymptotic normality of the one-step debiased estimator. Simulation and an application to real-world data demonstrate the superiority of the proposed method.

Keywords: Patient-reported Outcome; Dimension Reduction; Semiparametric Inference; Semi-supervised Learning; Double Machine Learning.

^{*}Booth School of Business, University of Chicago

[†]Department of Biostatistics, University of Florida

[‡]Public Health Sciences Divisions, Fred Hutchinson Cancer Center

[§]Department of Industrial and Systems Engineering, University of Florida

1 Introduction

Patient-reported outcome (PRO) measures are increasingly collected before and after an intervention or a treatment as a means of measuring healthcare quality and value, which is an important step toward patient-centered care. Knowing the measure goes up or down alone might not be sufficient to determine the effectiveness of the intervention. More importantly, whether the measure has changed with a sufficiently large margin, known as the minimally clinically important difference (MCID), needs to be evaluated. If the intervention is an elective surgery, identifying patients at risk of not achieving an MCID, particularly before the surgery, is important for pre-surgical decisions. There is a growing interest in applying machine learning techniques to identify factors associated with post-surgical PRO measures or predict whether a patient is likely to achieve an MCID before their surgery. In this work, our objective is to infer risk factors associated with not achieving an MCID under predictive modeling defined through minimizing the discrepancy between the prediction and the observed outcome measures. Specifically, we aim at obtaining the inference of the coefficients minimizing the deviance of a generalized linear model (GLM) with possible model misspecification.

The increasing adoption of electronic health record (EHR) systems has provided unprecedented opportunities to learn a model for predicting PRO measures using massive observational data. Although the volume of observational data is large, the quality of such observational data may be uncertain. One of the major difficulties is missing data, especially missing the outcome data. In our motivating example, the MCIDs can only be observed from the participants who take both pre- and post-surgical surveys. The participants who completed both surveys may only account for a small portion (e.g., 1/3) of the participants whose EHR data is available, according to the response rate reported in literature (Ho et al., 2019; Pronk et al., 2019) and from our own data. Unfortunately, low survey response rates are not uncommon in healthcare and other service industries.

Many approaches have been developed to deal with missing outcomes under the assumption of missing at random (MAR) (Kang and Schafer, 2007). One seminal work is the propensity inverse weighting approach (Rosenbaum and Rubin, 1983; Horvitz and Thompson, 1952). For this approach, one first estimates the probability of missing w.r.t the covariate (also called the propensity) and then uses the inverse of the estimated propensity to adjust for the selection bias. When the propensity is poorly estimated, the propensity inverse weighting methods may not perform well. Another major type of approach is known as imputation. This approach first learns an imputation model using the fully observed part of the data; then, imputes the missing outcomes with the predicted values; and finally, refits the predictive model based on the imputed outcomes (Rubin, 2004). When the estimated

imputation model is biased, the refitted predictive model may also be biased. To maintain robustness against the possible misspecification in the propensity and the imputation models, one possible solution is to use the doubly robust methods (Robins et al., 1994). The doubly robust methods that incorporate both the propensity score and the imputation models can lead to a consistent estimate for the outcome as long as either model is correctly specified (Tan, 2006, 2010; Qin et al., 2008; Qin and Zhang, 2007; Rubin and van der Laan, 2008; Cao et al., 2009; Han, 2012; Rotnitzky et al., 2012; Han et al., 2016).

Statistical inference for the parameters in predictive modeling with outcome missingness is also challenging. In particular, when the missing mechanism is dependent on multiple covariates through a nonlinear relationship, an unbiased estimator for the missing propensity with a fast convergence rate may be infeasible. For the inverse weighting approaches and the doubly robust methods, a parametric model for the propensity may not capture the potential non-linearity. To ensure an unbiased propensity estimate, nonparametric regressions and machine learning methods have been adopted. These methods may lead to a slower convergence rate and hinder the inference of the parameters in the predictive model, especially when the number of the covariates is large. When the number of the covariates is small, to address the slow convergence rate, the double machine learning approach was proposed in Chernozhukov et al. (2018). They adopted a cross-fitting algorithm using a doubly robust formulation and proposed to estimate both the propensity and the imputation model using nonparametric or machine learning methods. They proved that, as long as the product of the convergence rates of the propensity and imputation estimates is smaller than $n^{-1/2}$, a valid inference procedure for the parameters in the predictive model is possible, where n represents the sample size. However, the large number of the covariates and not meeting the smoothness condition on the true propensity may negate the required rate condition.

To help address the above statistical inference challenge, one possible strategy is to leverage a surrogate outcome. The surrogate outcomes herein are defined as alternative clinical outcomes that are likely to predict the clinical benefit of primary interest. In our motivating example, the MCID of the global physical health T-score in the Patient-Reported Outcomes Measurement Information System (PROMIS) survey is a well acknowledged measurement for evaluating surgery benefit. There are other PRO measures collected that represent different but related mental or physical health performances that can be considered as surrogate outcomes. In many applications, a surrogate outcome can help improve the efficiency or overcome the difficulties due to complex missing mechanisms. In the application of causal inference (Prentice, 1989; Frangakis and Rubin, 2002; Fleming et al., 1994; Cheng et al., 2018; Anderer et al., 2021), a surrogate can be used to improve the efficiency of estimating the average treatment effect (ATE). In the application of semi-supervised inference, under the

assumption of missing completely at random (MCAR), Hou et al. (2021) showed that a surrogate can help infer the predicted risk derived from a high-dimensional working model even when the true risk prediction model depends on multiple covariates. However, their approach cannot be applied under the assumption of missing at random (MAR), which is the setting we need to deal with.

In this work, we focus on how to use surrogate outcomes to facilitate the inference of model parameters with outcome missingness. The parameter of interest herein is defined as the minimizer of the deviance under a GLM with possible model misspecification. We propose a concept of an informative surrogate, defined as a surrogate outcome that enables a low-dimensional imputation model conditional on the surrogate and the covariates (i.e., the imputation model lies in a low-dimensional subspace generated by the surrogate and covariates). Under the MAR assumption, we exploit the role of this informative surrogate to 1) allow for a low-dimensional imputation model under a large number of covariates; 2) avoid estimating the complex missing propensity; 3) improve the efficiency in parameter estimation. To harvest the potential benefit brought by informative surrogate outcomes, we propose the following procedure. First, we estimate a flexible imputation model (e.g., using kernel regression) in a reduced subspace that is constructed by leveraging the information from informative surrogate outcomes. Subsequently, we can impute the missing outcomes and obtain an initial estimator for the parameters of interest. Then, we bypass the estimation of the complex missing propensity entirely, and instead estimate a low-dimensional weighting function based on the reduced subspace to adjust for the possible bias due to the estimated imputation model. Finally, a one-step debiased estimator for the parameters in the predictive model can be constructed. Both the point and interval estimates of the parameters can be obtained from the proposed procedure. We show that the proposed method can provide a valid inference procedure for the parameters of interest without requiring a consistent propensity estimation. In addition, when the true propensity lies in the same subspace as the imputation model, the proposed method leads to a semiparametric efficient estimator for the parameters in the predictive model. Extensive simulation and an analysis of real-world data are provided to demonstrate the superior performance of the proposed method.

The remainder of the paper is organized as follows. In Section 2, we define the parameter of interest and introduce our proposed inference procedure. In Section 3, we demonstrate the theoretical validity of the proposed inference procedure. In Section 4, we provide numerical studies to bolster the superiority of the proposed methods over other existing methods and methods without information of the surrogate. In Section 5, we apply the proposed method to derive a predictive rule to infer post-surgery improvement for joint replacement surgery patients. In Section 6, we discuss possible future works.

2 Method

Let \mathbf{X} be a p -dimensional covariate and Y be a binary, categorical, or continuous outcome of interest. Without loss of generality, we choose a GLM as a working model for $E[Y | \mathbf{X}]$, i.e., $E[Y | \mathbf{X}] = b(\mathbf{X}^\top \boldsymbol{\beta})$, where $b(\cdot)$ is a known link function. The parameter of interest, $\boldsymbol{\beta}$ is defined as the minimizer of the deviance under the working model, i.e., $\boldsymbol{\beta}^* = \arg \min E[\ell(\boldsymbol{\beta})]$, where $\ell(\boldsymbol{\beta}) = b(\mathbf{X}^\top \boldsymbol{\beta}) - Y \mathbf{X}^\top \boldsymbol{\beta}$. In the dataset, we collect the covariate \mathbf{X} , the outcome Y , the informative surrogate outcome Z , and the missing indicator R from all samples. The missing indicator R indicates whether Y is observed ($R = 1$) or not ($R = 0$). We also assume that the surrogate Z can be fully observed. Collectively, the observed data can be denoted as (\mathbf{X}, Z, R, RY) . If the working model is misspecified, i.e., $E[Y | \mathbf{X}] \neq b(\mathbf{X}^\top \boldsymbol{\beta}^*)$, $\boldsymbol{\beta}^*$ that minimizes the deviance, a goodness-of-fit statistic, is still meaningful.

2.1 A doubly robust formulation for estimation

In this section, we derive the semiparametric lower bound of estimating $\boldsymbol{\beta}^*$ with and without informative surrogate outcomes and then discuss how the informative surrogate can be used to construct a doubly robust estimation for $\boldsymbol{\beta}^*$. To ensure the identifiability of $\boldsymbol{\beta}^*$ given the observed data defined above, we request that $Y \perp R | \mathbf{X}, Z$. Without the surrogate outcome, the strong ignorability condition assumes that $Y \perp R | \mathbf{X}$, which implies that \mathbf{X} captures all confounders between the missing indicator R and the outcome Y . Direct comparison of the efficiency in estimating $\boldsymbol{\beta}^*$ under the strong ignorability and the requested condition may not be meaningful if the conditions are mutually exclusive. To enable a meaningful comparison, we further assume that $R \perp Z | \mathbf{X}$ in addition to $Y \perp R | \mathbf{X}, Z$. In this case, the strong ignorability condition is equivalent to the requested condition $Y \perp R | \mathbf{X}, Z$.

Under these assumptions, we obtain Theorem 1 that, an estimator achieving the semiparametric lower bound has guaranteed efficiency gain from using the surrogate outcome when $E[Y | \mathbf{X}] \neq E[Y | Z, \mathbf{X}]$. The proof of Theorem 1 can be found in the Online Supporting Information.

Theorem 1 *Assume that $R \perp Z | \mathbf{X}$ and $Y \perp R | \mathbf{X}, Z$. No matter whether the distribution of (\mathbf{X}, Z) is known or unknown, the semiparametric lower bound for β_1^* using the surrogate outcome Z is lower than using only X if $E[Y | \mathbf{X}] \neq E[Y | Z, \mathbf{X}]$, where β_1^* is the first coordinate of $\boldsymbol{\beta}^*$.*

Remark 1 *When the marginal missing rate goes to 1, compared with the conditional distribution $Y | \mathbf{X}, Z$ being only accessible on the fully observed part, the distribution of (\mathbf{X}, Z) is accessible on the entire dataset. In the semiparametric lower bound, the variance related to the conditional distribution $Y | \mathbf{X}, Z$ will dominate the variance related to the distribution of (\mathbf{X}, Z) . This means that when the*

marginal missing rate goes to 1, the variance of a semiparametric efficient estimator will share the same format of the semiparametric lower bound where the distribution of (\mathbf{X}, Z) can be considered as known. Thus, Theorem 1 shows the possible efficiency gain under both cases where the marginal missing rate goes to 1 and is bounded away from 1.

To derive the estimator of β^* that achieves the semiparametric lower bound, motivated by the form of the augmented inverse propensity weighted estimator (AIPW) (Robins et al., 1994; Robins and Rotnitzky, 1995; Rotnitzky et al., 2012), we consider the estimating equation

$$\widehat{E}_n [S(\beta; \pi, Q)] = 0,$$

where $\widehat{E}_n [\cdot]$ is the empirical mean,

$$\begin{aligned} & S(\beta; \pi, Q) \\ &= [\pi^{-1}(\mathbf{X}, Z)R \{b'(\mathbf{X}^\top \beta) - Y\} - \pi^{-1}(\mathbf{X}, Z)\{R - \pi(\mathbf{X}, Z)\} \{b'(\mathbf{X}^\top \beta) - Q(Z, \mathbf{X})\}] \mathbf{X}, \end{aligned}$$

$\pi(Z, \mathbf{X})$ is a working model for $P(R = 1 | Z, \mathbf{X})$, and $Q(Z, \mathbf{X})$ is a working model for $E[Y | Z, \mathbf{X}]$. This estimating equation enjoys doubly robustness, i.e., the solution of the estimating equations is a consistent estimator of β^* if either $\pi(Z, \mathbf{X}) = P(R = 1 | Z, \mathbf{X})$ or $Q(Z, \mathbf{X}) = E[Y | Z, \mathbf{X}]$. Utilizing this double robust formulation, we propose a two-step algorithm with a cross-fitting procedure.

2.2 First step: dimension reduction through informative surrogate

To start with, we formally define the concept of informative surrogate outcomes. An surrogate outcome Z is informative if there exists a $(p+1) \times d$ matrix, $\mathbf{\Gamma}$, with orthogonal columns satisfying $Y \perp \widetilde{\mathbf{X}} | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}}$ and $d < p$, where $\widetilde{\mathbf{X}}^\top = (Z, \mathbf{X}^\top)$. This definition implies that, conditioning on the surrogate outcome, the dimension of the space constructed by the covariates and the surrogate can be reduced to d , which is expected to be much smaller than p . The columns of $\mathbf{\Gamma}$ represent the reduced subspace. Thus, if the surrogate is informative, $Q(Z, \mathbf{X}) = E[Y | Z, \mathbf{X}]$ is a function lying in a low-dimensional subspace, i.e., there exists an unknown link function g such that $Q(Z, \mathbf{X}) = g(\mathbf{\Gamma}^\top \widetilde{\mathbf{X}})$. Consequently, an efficient estimator to this low-dimensional imputation model may have a faster convergence rate than directing using the kernel regression to estimate $E[Y | \mathbf{X}]$ which is p -dimensional.

Given an informative surrogate outcome, various existing methods can be employed to estimate the reduced subspace. The assumption $Y \perp \widetilde{\mathbf{X}} | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}}$ is closely related to the dimension reduction literature. When there are multiple surrogate outcomes, the methods in the dimension reduction literature selecting the true reduced dimension can be used to select candidate informative surrogate outcomes. For example, we can select the surrogate outcome that leads to the lowest reduced dimension. In the literature (Cook, 2007; Xia et al., 2002; Xia, 2007; Ma and Zhu, 2012, 2013), the smallest

space generated by the columns of $\mathbf{\Gamma}$ is referred to as the central subspace. When the data is fully observed, the dimension reduction methods, such as the minimum average variance estimation (MAVE) (Xia et al., 2002), sliced-inverse regression (SIR) (Cook, 2007), or the semiparametric approach in Ma and Zhu (2012, 2013), can be directly applied to estimate the central subspace $\mathbf{\Gamma}$, and the estimated subspace is asymptotically normal. For our setting with missing outcome data, if $Y \perp \tilde{\mathbf{X}} \mid \mathbf{\Gamma}^\top \tilde{\mathbf{X}}$, then $Q(Z, \mathbf{X}) = E[Y \mid Z, \mathbf{X}] = E[Y \mid \mathbf{\Gamma}^\top \tilde{\mathbf{X}}] = E[Y \mid \mathbf{\Gamma}^\top \tilde{\mathbf{X}}, R = 1]$. This implies that the conditional mean of Y restricted to $R = 1$ shares the same subspace with the unrestricted conditional mean. Thus, to estimate $\mathbf{\Gamma}$, we only need to apply these dimension reduction methods to the fully observed part of the data.

After we obtain $\hat{\mathbf{\Gamma}}$, we can use nonparametric regressions or machine learning methods to fit Y w.r.t $\hat{\mathbf{\Gamma}}^\top \tilde{\mathbf{X}}$ to derive the unknown link function g and estimate $\hat{Q}(Z, \mathbf{X}) = \hat{g}(\hat{\mathbf{\Gamma}}^\top \tilde{\mathbf{X}})$. Then, we obtain an initial estimator for $\boldsymbol{\beta}^*$ by minimizing $\hat{E}_n \left[b(\mathbf{X}^\top \boldsymbol{\beta}) - \hat{g}(\hat{\mathbf{\Gamma}}^\top \tilde{\mathbf{X}}) X^\top \boldsymbol{\beta} \right]$. Denote the minimizer as $\hat{\boldsymbol{\beta}}$. Due to the slow convergence rate of nonparametric regressions or machine learning methods, the convergence rate of \hat{Q} is dominated by that of \hat{g} . Subsequently, $\hat{\boldsymbol{\beta}}$ may suffer from the slow convergence rate of \hat{g} . Thus, to obtain an estimator with a faster convergence rate, we need to remove the bias due to the estimation error of \hat{g} , which will be discussed in the following section.

2.3 Second step: debias using a low-dimensional weighting function

We propose to remove the bias due to the estimation error of \hat{g} using a low-dimensional weighting function. For ease of exposition, we consider the inference of β_1^* , which is the first coefficient in $\boldsymbol{\beta}^*$; the proposed method can be extended to infer $\mathbf{u}^\top \boldsymbol{\beta}^*$ for any \mathbf{u} .

We first construct an estimating equation for β_1^* . Let \mathbf{w}^* be the minimizer of

$$E \left[b''(\mathbf{X}^\top \boldsymbol{\beta}^*) (X_1 - X_{-1}^\top \mathbf{w})^2 \right],$$

where X_1 is the first covariate in \mathbf{X} and \mathbf{X}_{-1} is the covariate vector of \mathbf{X} excluding X_1 . Consider the following estimating equation for β_1^* ,

$$E \left[\{S(\boldsymbol{\beta}; \pi, Q)\}^\top \mathbf{v} \right], \tag{1}$$

where $\mathbf{v}^\top = (1, -\mathbf{w}^{*\top})$ and $\boldsymbol{\beta}_{-1}$ is fixed at $\boldsymbol{\beta}_{-1}^*$.

For any choice of π , the first-order bias of the proposed estimating Equation (1) with Q replaced by \hat{Q} is $E \left[\left\{ R/\pi(\tilde{\mathbf{X}}) - 1 \right\} \left\{ \hat{g}(\hat{\mathbf{\Gamma}}^\top \tilde{\mathbf{X}}) - g(\mathbf{\Gamma}^\top \tilde{\mathbf{X}}) \right\} \mathbf{X}^\top \mathbf{v} \right]$. In order to remove this bias, one possible strategy is to choose $\pi(\tilde{\mathbf{X}})$ such that

$$E \left[\left\{ R/\pi(\tilde{\mathbf{X}}) - 1 \right\} f(\mathbf{\Gamma}^\top \tilde{\mathbf{X}}) \mathbf{X}^\top \mathbf{v} \right] = 0, \text{ for any } f \in L_2(\mathbf{\Gamma}^\top \tilde{\mathbf{X}}). \tag{2}$$

Let $P(R = 1) = \rho > 0$ and $\eta(\cdot | R)$ be the conditional density function of $\mathbf{\Gamma}^\top \widetilde{\mathbf{X}}$ given R . Define $J_r(\mathbf{\Gamma}^\top \tilde{\mathbf{x}}) = E \left[\mathbf{X}^\top \mathbf{v} | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}} = \mathbf{\Gamma}^\top \tilde{\mathbf{x}}, R = r \right] \eta(\mathbf{\Gamma}^\top \tilde{\mathbf{x}} | R = r)$ for $r = 0$ and 1 . Theorem 2 characterizes the solution to Equation (2). The proof of Theorem 2 can be found in the Online Supporting Information.

Theorem 2 *We assume the following regularity condition: for almost all $\tilde{\mathbf{x}}$, if $J_1(\mathbf{\Gamma}^\top \tilde{\mathbf{x}}) = 0$, then $J_0(\mathbf{\Gamma}^\top \tilde{\mathbf{x}}) = 0$. Then $\pi_*^{-1}(\tilde{\mathbf{x}}) = 1 + \{\rho J_1(\mathbf{\Gamma}^\top \tilde{\mathbf{x}})\}^{-1} J_0(\mathbf{\Gamma}^\top \tilde{\mathbf{x}})(1 - \rho)$ is well-defined. Further, the solution set of Equation (2) can be characterized as all the functions of the form $\pi_*^{-1}(\tilde{\mathbf{x}}) + \mathcal{T}h(\tilde{\mathbf{x}})$ on event $\{\tilde{\mathbf{x}} : J_1(\mathbf{\Gamma}^\top \tilde{\mathbf{x}}) \neq 0\}$, where \mathcal{T} is a linear operator defined as*

$$\mathcal{T}h = h - E \left[h \mathbf{X}^\top \mathbf{v} | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}}, R = 1 \right] / E \left[\mathbf{X}^\top \mathbf{v} | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}}, R = 1 \right],$$

and h is an arbitrary function in $L_2(\widetilde{\mathbf{X}})$.

Remark 2 *Equation (2) is solvable without the regularity condition. This can be verified by taking π as the true propensity $P(R = 1 | Z, \mathbf{X})$ in Equation (2). The regularity condition is required to ensure that π_*^{-1} as well as the characterization in Theorem 2 are both well-defined. This condition can be easily satisfied. For example, assume $\eta(\mathbf{\Gamma}^\top \widetilde{\mathbf{X}} | R = 1) > 0$. If $E \left[\mathbf{X}^\top \mathbf{v} | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}}, R = 1 \right]$ has a continuous distribution, then we have $P \left[J_1(\mathbf{\Gamma}^\top \widetilde{\mathbf{X}}) = 0 \right] = 0$, and the regularity condition is naturally satisfied. In addition, when $P(R = 1 | Z, \mathbf{X}) = P(R = 1 | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}})$, the regularity condition is also satisfied because $E \left[\mathbf{X}^\top \mathbf{v} | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}}, R = 1 \right] = E \left[\mathbf{X}^\top \mathbf{v} | \mathbf{\Gamma}^\top \widetilde{\mathbf{X}}, R = 0 \right]$.*

We then use $\pi_*^{-1}(\widetilde{\mathbf{X}})$ to replace the true propensity in estimating equation (1). The weighting function $\pi_*^{-1}(\widetilde{\mathbf{X}})$ only depends on $\mathbf{\Gamma}^\top \widetilde{\mathbf{X}}$ and thus is a low-dimensional function. To estimate the weighting function $\pi_*^{-1}(\widetilde{\mathbf{X}})$, we consider the trimmed kernel estimates. First, we estimate \mathbf{w} by minimizing $\widehat{E}_n \left[b''(\mathbf{X}^\top \widehat{\boldsymbol{\beta}})(X_1 - X_{-1}^\top \mathbf{w})^2 \right]$, and construct $\widehat{\mathbf{v}}^\top = (1, -\widehat{\mathbf{w}}^\top)$. Then, using kernel regressions, we consider the following estimator for π_*^{-1} :

$$\widehat{\pi}^{-1}(\tilde{\mathbf{x}}; \widehat{\boldsymbol{\Gamma}}, \widehat{\mathbf{v}}) = \begin{cases} 1 + \left\{ \widehat{J}_1(\widehat{\boldsymbol{\Gamma}}^\top \tilde{\mathbf{x}}) \widehat{\rho} \right\}^{-1} \widehat{J}_0(\widehat{\boldsymbol{\Gamma}}^\top \tilde{\mathbf{x}})(1 - \widehat{\rho}) & \left| \widehat{J}_1(\widehat{\boldsymbol{\Gamma}}^\top \tilde{\mathbf{x}}) \right| > c_n, \\ \widehat{\rho}^{-1}, & \left| \widehat{J}_1(\widehat{\boldsymbol{\Gamma}}^\top \tilde{\mathbf{x}}) \right| \leq c_n, \end{cases}$$

where $\widehat{J}_r(\widehat{\boldsymbol{\Gamma}}^\top \tilde{\mathbf{x}}) = \widehat{E}_n \left[\mathbf{X}^\top \widehat{\mathbf{v}} K_h(\widehat{\boldsymbol{\Gamma}}^\top \widetilde{\mathbf{X}} - \widehat{\boldsymbol{\Gamma}}^\top \tilde{\mathbf{x}}) | R = r \right]$, $\widehat{\rho} = \widehat{E}_n[R]$, $K_h(\cdot) = K(\cdot)/h^d$, and $\widehat{E}_n[\cdot | R = r]$ is the empirical mean over the samples with $R = r$. The function $K(\cdot)$ is a kernel function with the order of ν , and the bandwidth parameter h is selected according to Theorem 3. The proposed estimator equals to the kernel regression when $\left| \widehat{J}_1(\widehat{\boldsymbol{\Gamma}}^\top \tilde{\mathbf{x}}) \right|$ is far from 0, and equals to $\widehat{\rho}^{-1}$, when $\left| \widehat{J}_1(\widehat{\boldsymbol{\Gamma}}^\top \tilde{\mathbf{x}}) \right|$ is close to 0. The term c_n is used to trim possible extremities of the kernel regression estimates.

After obtaining the estimator $\hat{\pi}^{-1}$ for $\pi_*^{-1}(\tilde{\mathbf{X}})$, we construct the estimating equation by incorporating $\hat{\pi}^{-1}$ to remove the bias, i.e., $\hat{E}_n \left[\left\{ S(\beta; \hat{Q}, \hat{\pi}) \right\}^\top \hat{\mathbf{v}} \right]$ with a constraint $\beta_{-1} = \hat{\beta}_{-1}$. However, directly solving this equation may lead to multiple solutions or numerically unstable solutions (Chapter 5 in Vaart (1998)). To avoid possible computational issues, we use its first-order expansion and construct a one-step debiased estimator $\tilde{\beta}_1 = \hat{\beta}_1 - \bar{I}^{-1}S$, where $S = \hat{E}_n \left[\left\{ S(\hat{\beta}; \hat{Q}, \hat{\pi}) \right\}^\top \hat{\mathbf{v}} \right]$, and $\bar{I} = \hat{E}_n \left[b''(\mathbf{X}^\top \hat{\beta}) X_1 \mathbf{X}^\top \hat{\mathbf{v}} \right]$. Another challenge in constructing the debiased estimator is the correlation between the estimation errors of \hat{Q} , $\hat{\pi}$ and the samples used to construct the estimator. We adopt the cross-fitting procedure proposed in Chernozhukov et al. (2018) in the implementation.

2.4 Implementation

The entire procedure can be separated into two steps. In the first step, we obtain $\hat{\Gamma}$ using all fully observed data. Then, using all fully observed data, we regress Y on $\hat{\Gamma}^\top \tilde{\mathbf{X}}$ using kernel regressions and denote the estimated link function as \hat{g} . Using the estimated imputation model, $\hat{Q}(Z, \mathbf{X}) = \hat{g}(\hat{\Gamma}^\top \tilde{\mathbf{X}})$, we obtain an initial estimate $\hat{\beta}$. Using the initial estimate, we solve $\min_{\mathbf{w}} \hat{E}_n \left[b''(\mathbf{X}^\top \hat{\beta})(X_1 - X_{-1}^\top \mathbf{w})^2 \right]$, and denote its minimizer as $\hat{\mathbf{w}}$. Then we can construct $\hat{\mathbf{v}}^\top = (1, -\hat{\mathbf{w}}^\top)$. In the second step, we estimate the identified weighting function and use it to form a one-step debiased estimator for β_1^* . First, we split the entire data into K subsets (I_1, \dots, I_K) with equal sample sizes. For a specific set k , the estimated link function denoted as $\hat{g}_{(-k)}(\cdot)$ is obtained through kernel regression of Y w.r.t $\hat{\Gamma} \tilde{\mathbf{X}}$ using the data excluding I_k . The estimated weighting function denoted as $\hat{\pi}_{(-k)}^{-1}(\tilde{\mathbf{X}}; \hat{\Gamma}, \hat{\mathbf{v}})$ is obtained through the truncated kernel regression using the data excluding I_k . Specifically,

$$\hat{\pi}_{(-k)}^{-1}(\tilde{\mathbf{x}}; \hat{\Gamma}, \hat{\mathbf{v}}) = \begin{cases} 1 + \left\{ \hat{J}_1^{(-k)}(\hat{\Gamma}^\top \tilde{\mathbf{x}}) \hat{\rho} \right\}^{-1} \hat{J}_0^{(-k)}(\hat{\Gamma}^\top \tilde{\mathbf{x}})(1 - \hat{\rho}) & \left| \hat{J}_1^{(-k)}(\hat{\Gamma}^\top \tilde{\mathbf{x}}) \right| > c_n, \\ \hat{\rho}^{-1}, & \left| \hat{J}_1^{(-k)}(\hat{\Gamma}^\top \tilde{\mathbf{x}}) \right| \leq c_n, \end{cases}$$

where $\hat{J}_r^{(-k)}(\hat{\Gamma}^\top \tilde{\mathbf{x}}) = \hat{E}_n^{(-k)} \left[\mathbf{X}^\top \hat{\mathbf{v}} K_h(\hat{\Gamma}^\top \tilde{\mathbf{X}} - \hat{\Gamma}^\top \tilde{\mathbf{x}}) \mid R = r \right]$ and $\hat{E}_n^{(-k)}[\cdot \mid R = r]$ is the empirical average over the samples with $R = r$ and excluding those in I_k . Then, the one-step debiased estimator is $\tilde{\beta}_1 = \hat{\beta}_1 - \bar{I}^{-1} \bar{S}$, where

$$\bar{S} = \sum_{k=1}^K S^{(k)} / K, \quad S^{(k)} = \hat{E}_n^{(k)} \left[\left\{ S(\hat{\beta}; \hat{Q}_{(-k)}, \hat{\pi}_{(-k)}) \right\}^\top \hat{\mathbf{v}} \right].$$

A summary of the entire algorithm can be found in the Online Supporting Information. To estimate the asymptotic variance of $\tilde{\beta}_1$, we bootstrap based on the entire sample for B times; for the b th bootstrapped dataset, we implement the algorithm and obtain $\tilde{\beta}_1^{(b)}$, where $b = 1, \dots, B$. We use the variance of $\left\{ \tilde{\beta}_1^{(b)} \right\}_{b=1}^B$ as the estimate for the asymptotic variance to construct interval estimations.

3 Theoretical properties

In this section, we provide the asymptotic property of the proposed estimator. To accommodate the situation where the marginal missing rate may be close to 1, we assume that the distribution of (\mathbf{X}, Z) and the conditional distribution $Y | Z, \mathbf{X}$ do not depend on n ; the missing propensity $P(R = 1 | Z, \mathbf{X})$ may depend on n . Specifically, we consider two scenarios: 1) the missing propensity $P(R = 1 | Z, \mathbf{X})$ does not change with n ; 2) $P(R = 1 | Z, \mathbf{X}) = \rho_n w(\tilde{\mathbf{X}})$ with $\rho_n \rightarrow 0$, where $w(\tilde{\mathbf{X}})$ does not depend on n , $w(\tilde{\mathbf{X}})$ is always non-negative and bounded away from ∞ , and $E[w(\tilde{\mathbf{X}})] = 1$. Notice that for both scenarios, we will not assume the positivity assumption, i.e., $P(R = 1 | Z, \mathbf{X})$ is bounded away from 0. This is a benefit of not using the inverse of the true propensity in the estimation and inference procedure. For simplicity, we focus on the required conditions and theoretical results for Scenario 1) in the main text, and leave those for Scenario 2) in the Online Supporting Information. In addition, the proofs of the theorems can be found in the Online Supporting Information. For Scenario 1), the following conditions are required.

Assumption 1 *The covariate \mathbf{X} 's and the surrogate outcome Z are bounded, and the function $b''(\cdot)$ is continuously differentiable; $\max\{\|\beta^*\|_2, \|\mathbf{v}\|_2\}$ is bounded. In addition, the smallest eigenvalue of $E[\mathbf{X}\mathbf{X}^\top]$ is bounded away from 0.*

Assumption 2 *There is a positive constant $\gamma_d > 1/4$ such that $\|\hat{g}(\hat{\Gamma}^\top \tilde{\mathbf{X}}) - Q(Z, \mathbf{X})\|_\infty = O_p(n^{-\gamma_d})$, and $\{\text{vec}(\hat{\Gamma}) - \text{vec}(\Gamma)\} = n^{-1} \sum_{i=1}^n 1\{R_i = 1\} \psi(\mathbf{X}_i, Z_i, Y_i) + o_p(n^{-1/2})$, where $\psi(\mathbf{X}_i, Z_i, Y_i)$ is bounded and $\text{vec}(\cdot)$ represents the vectorization of the matrix. In addition, we assume that*

$$\sup_{\tilde{\mathbf{x}}} \left| (\hat{g} - g)(\hat{\Gamma}^\top \tilde{\mathbf{x}}) - (\hat{g} - g)(\Gamma^\top \tilde{\mathbf{x}}) \right| = o_p(n^{-1/2}).$$

Assumption 3 *Function $J_r(\Gamma^\top \tilde{\mathbf{x}})$'s are ν th order differentiable w.r.t $\Gamma^\top \tilde{\mathbf{x}}$ with bounded derivatives. Define $G(\Gamma^\top \tilde{\mathbf{x}}) = J_1^{-1}(\Gamma^\top \tilde{\mathbf{x}}) J_0(\Gamma^\top \tilde{\mathbf{x}})$. We assume that $G(\Gamma^\top \tilde{\mathbf{x}})$ is bounded away from 0 and $+\infty$ and continuously differentiable on the open set $\{\tilde{\mathbf{x}} : J_1(\Gamma^\top \tilde{\mathbf{x}}) \neq 0\}$. We also assume that the density function of $\mathbf{A}^\top \tilde{\mathbf{X}}$, $\eta(\mathbf{A}^\top \tilde{\mathbf{X}})$, is bounded away from 0 and $+\infty$, and ν th order differentiable with bounded derivatives. Define*

$$H_1(\text{vec}(\Gamma); \mathbf{v}) = E \left[\left\{ R/\pi_*(\tilde{\mathbf{X}}) - 1 \right\} \left\{ \nabla g(\Gamma^\top \tilde{\mathbf{X}}) \otimes \mathbf{X} \right\} \mathbf{X}^\top \mathbf{v} \right],$$

where \otimes is the Kronecker product. The $H_1(\text{vec}(\Gamma); \mathbf{v})$ is bounded for any $\|\mathbf{v}\|_2 = 1$.

Assumption 4 *When $t > 0$ is small enough, there exist positive constants A_0 and γ_m such that $P\{0 \neq |J_1(\Gamma^\top \tilde{\mathbf{X}})| \leq t\} \leq A_0 t^{\gamma_m}$.*

Assumption 5 Take $c_n = \tilde{\delta}_n^{2/(2+\gamma_m)}$, where $\tilde{\delta}_n = (n\hbar^d/\log n)^{-1/2} + \hbar^\nu + n^{-\gamma_d}$. We assume that $n^{-\gamma_d}\tilde{\delta}_n^{\gamma_m/(2+\gamma_m)} = o(n^{-1/2})$.

In Condition 1, for ease of exposition, we assume a bounded design for each \mathbf{X} and Z . Condition 2 includes the requirement for the chosen dimension reduction method to estimate $\mathbf{\Gamma}$ and the chosen method to estimate g . Specifically, we assume that the uniform convergence rate of \hat{Q} is $O_p(n^{-\gamma_d})$, and the estimated subspace $\hat{\mathbf{\Gamma}}$ is asymptotically linear. In addition, we assume that $\sup_{\tilde{\mathbf{x}}} |(\hat{g} - g)(\hat{\mathbf{\Gamma}}^\top \tilde{\mathbf{x}}) - (\hat{g} - g)(\mathbf{\Gamma}^\top \tilde{\mathbf{x}})| = o_p(n^{-1/2})$. Many nonparametric methods such as kernel regressions satisfy Condition 2 (Silverman, 1978). Condition 3 assumes the regularity conditions including the smoothness of $J_r(\mathbf{\Gamma}^\top \tilde{\mathbf{x}})$'s to ensure that π_*^{-1} can be well-estimated and the asymptotic variances are well-defined. Condition 4 restricts the concentration near $J_1(\mathbf{\Gamma}^\top \tilde{\mathbf{x}}) = 0$ by the parameter γ_m . When $J_1(\mathbf{\Gamma}^\top \tilde{\mathbf{x}})$ is bounded away from 0, we have $\gamma_m = +\infty$; when $J_1(\mathbf{\Gamma}^\top \tilde{\mathbf{X}})$ has a continuous distribution, we have $\gamma_m \geq 1$. Condition 5 specifies the condition on γ_d and γ_m , which requires that $\gamma_d > (2 + \gamma_m)/4(1 + \gamma_m)$.

Under these conditions, Theorem 3 shows that the one-step debiased estimator is asymptotically normal.

Theorem 3 Under Conditions 1- 5, we have $\sqrt{n}(\tilde{\beta}_1 - \beta_1^*) \rightarrow N(0, \sigma^2)$, where the formula of the asymptotic variance σ^2 can be found in the Online Supporting Information.

Theorem 3 implies that we can construct a valid confidence interval if we can consistently estimate σ^2 . To estimate the asymptotic variance σ^2 , we can adopt a plug-in approach. When the asymptotic variance of $\hat{\mathbf{\Gamma}}$ is explicitly known, we can construct estimators for the unknown parameters in the asymptotic variance formula and construct a plug-in estimator for σ^2 . In this work, we choose the bootstrap procedure, which has been shown to have a better numerical performance for regression with semi-nonparametric nuisance models (Liu et al., 2020).

In addition, Corollary 1 shows a sufficient condition that the semiparametric lower bound can be achieved.

Corollary 1 When the true propensity $P(R = 1 | Z, \mathbf{X}) = P(R = 1 | \mathbf{\Gamma}^\top \tilde{\mathbf{X}})$, the one-step debiased estimator $\tilde{\beta}_1$ obtains the semiparametric lower bound.

4 Simulations

In this section, we conduct simulations and compare the proposed method with other methods to demonstrate 1) the advantage of avoiding complex propensity estimation; and 2) the efficiency gain

from incorporating the surrogate outcome. To show the advantage of avoiding modeling the complex propensity, we compare our proposed method with two baseline approaches. Baseline 1 follows the double machine learning procedure proposed in Chernozhukov et al. (2018), which estimates both the propensity and the imputation model using kernel regressions. When using the kernel regressions to estimate the propensity and the imputation model, we first implement dimension reduction and then conduct the kernel regression. Another baseline approach (Baseline 2) adopts the same procedure as Baseline 1 but uses a logistic regression to estimate the missing propensity. For both Baselines 1 and 2, we implement a threshold of 0.001 for the estimated propensities to avoid extreme value. To show the efficiency gain from incorporating the surrogate outcome, besides the proposed procedure using the surrogate outcome (denoted as “Proposed with Z ”), we implement another approach (denoted as “Proposed w/o Z ”) following the same procedure but only using \mathbf{X} (no surrogate outcome Z) in the dimension reduction, imputation model estimation, and weighting function estimation. For both proposed procedures, we specify c_n following the results in Theorem 3. For the dimension reduction adopted in all these approaches, we use the kernel sliced regression method and choose the reduced dimension using cross-validation.

To compare with the proposed method, we consider 8 simulation scenarios in total with varying missing rates, varying sample sizes, and different types of outcomes, i.e., continuous outcomes and binary outcomes. For each type of outcome, we consider a moderate marginal missing rate of 50% and a high marginal missing rate of 90%. For both scenarios, we change the sample size from 500 to 1000. To generate the data under each scenario, we first generate the missing indicator R following a Bernoulli distribution with the success probabilities of 0.5 (moderate marginal missing rate) or 0.1 (high marginal missing rate). Then, we generate the covariate based on $R = 1$ or $R = 0$. When the outcome is missing ($R = 0$), the covariates, i.e., $\mathbf{X} \mid R = 0$, follow a standard multivariate Gaussian distribution with zero means and the identity covariance matrix; when the outcome is observed ($R = 1$), the covariates follow a mixture of two multivariate Gaussian distributions. With a probability of 0.7, the covariates are generated following a standard multivariate Gaussian distribution; otherwise, the covariates are generated following a multivariate Gaussian distribution $N(1, 1.5\mathbf{I}_p)$, i.e., $\mathbf{X} \mid R = 1 \sim \xi N(0, \mathbf{I}_p) + (1 - \xi)N(1, 1.5\mathbf{I}_p)$, where ξ follows a Bernoulli distribution with a success probability of 0.7. To generate the outcome Y given the covariates, for the scenario with continuous outcomes, we consider $Y = \mathbf{X}^\top \boldsymbol{\beta}_0 + \sum_{j=5}^8 |X_j|/4 + \epsilon$, where $\boldsymbol{\beta}_0 = (1, 1, -1, -1, 0, 0, 0, 0)^\top$ and $\epsilon \sim N(0, 2)$. For the scenario with binary outcomes, we consider $Y = 1\{0.5\mathbf{X}^\top \boldsymbol{\beta}_0 + \sum_{j=5}^8 |X_j|/4 + \epsilon > 0\}$. The surrogate outcome is generated from $Z = \delta \mathbf{X}^\top \boldsymbol{\beta}_0 + \sum_{j=5}^8 |X_j|/4 + \epsilon_z$ where $\epsilon_z \sim N(0, 1)$. For the scenario with continuous outcomes, δ is fixed at 0.5; for the scenario with binary outcomes, δ is fixed

at 0.25.

To evaluate the proposed methods, we compare the coverage and deviance in all the scenarios. For each scenario, we run 500 replicates. For each replicate, we estimate the coefficients and use bootstrapping (500 bootstraps) to construct a 95%-confidence interval using the training samples. We also calculate the deviance using the coefficients estimated by each approach on an independently generated testing dataset with a sample size of 10^4 .

Table 1 exhibits the coverage metrics of the first four coefficients. From Table 1, the proposed methods (Proposed w/o and with Z) achieve the nominal coverage in all the scenarios, whereas Baseline 1 and Baseline 2 do not, especially in high missing rate settings. This suggests both Baseline 1 and Baseline 2 are incapable of handling the complex missing mechanism. Figure 1 summarizes the deviance metrics for all different scenarios. At the moderate missing rate (the left two columns), the method Proposed with Z performs comparably to Baseline 2 but outperforms the other methods. In the high missing rate scenarios, Proposed with Z achieves the minimum deviance. When comparing the two methods, Proposed with Z surpasses Proposed w/o Z in all the scenarios due to the efficiency gain from Z . In summary, considering both coverage and deviance metrics, the proposed method with the surrogate outcome is dominant.




Figure 1: Deviance under different missing rates, sample sizes, and outcome types.

Table 1: Coverage of the 95% confidence interval for the coefficients.

	$n = 500$				$n = 1000$			
	Continuous, Missing rate of 50%							
	β_1	β_2	β_3	β_4	β_1	β_2	β_3	β_4
Baseline 1	0.872	0.840	0.886	0.864	0.847	0.769	0.861	0.817
Baseline 2	0.896	0.912	0.906	0.928	0.885	0.873	0.893	0.875
Proposed w/o Z	0.920	0.938	0.910	0.922	0.901	0.901	0.891	0.893
Proposed with Z	0.952	0.964	0.958	0.968	0.954	0.954	0.956	0.954
	Continuous, Missing rate of 90%							
	β_1	β_2	β_3	β_4	β_1	β_2	β_3	β_4
Baseline 1	0.866	0.840	0.844	0.854	0.740	0.798	0.752	0.762
Baseline 2	0.904	0.876	0.922	0.920	0.910	0.898	0.894	0.902
Proposed w/o Z	0.954	0.966	0.968	0.946	0.942	0.944	0.938	0.952
Proposed with Z	0.946	0.968	0.966	0.962	0.946	0.964	0.954	0.956
	Binary, Missing rate of 50%							
	β_1	β_2	β_3	β_4	β_1	β_2	β_3	β_4
Baseline 1	0.908	0.918	0.916	0.932	0.892	0.882	0.918	0.900
Baseline 2	0.928	0.932	0.916	0.930	0.920	0.898	0.894	0.912
Proposed w/o Z	0.968	0.970	0.956	0.950	0.962	0.964	0.964	0.964
Proposed with Z	0.970	0.972	0.948	0.962	0.958	0.968	0.956	0.946
	Binary, Missing rate of 90%							
	β_1	β_2	β_3	β_4	β_1	β_2	β_3	β_4
Baseline 1	0.950	0.940	0.948	0.962	0.892	0.942	0.964	0.940
Baseline 2	0.932	0.930	0.920	0.956	0.924	0.912	0.896	0.926
Proposed w/o Z	0.956	0.950	0.948	0.964	0.954	0.962	0.944	0.950
Proposed with Z	0.966	0.966	0.976	0.948	0.942	0.962	0.966	0.974

5 Application to PROMIS global physical health T-score

In this section, we applied our proposed method to predict whether the improvement of the PROMIS global physical health T-score will exceed the MCID after receiving total joint replacement using the information obtained before scheduling the surgery. In addition to making prediction, we also aimed to identify the driving factors of not achieving the MCID in the presence of the high missing-rate outcome. The dataset includes 1044 University of Florida Health patients, who participated in the pre-surgical survey and underwent the total joint replacement surgery. In the analysis, we incorporated many baseline covariates including demographics, socioeconomic characteristics, medical history, and care characteristics before surgery (e.g., 30 days before admission for surgery). According to the convention of constructing MCID, the outcome of interest is obtained based on the one-half standard deviation of the difference between pre- and post-surgical PROMIS global health T-scores (Fontana et al., 2019; Katakam et al., 2022).

For our data, the difference between pre- and post-surgical scores has an average of 9.5 and a standard deviation of 8.1, and consequently, the MCID is 4.1. Thus, the outcome $Y = 1$ if the difference is less than 4.1, and $Y = 0$, otherwise. In terms of the missing proportion, all the 1044 identified patients took the pre-surgical survey, but only 261 patients (25%) responded to the post-surgical survey (the missing-rate is 75%). Of the patients who took both surveys, 67(25.7%) patients did not meet the MCID. Since the PROMIS global physical T-score is derived from the ten survey questionnaire items, individual items can be considered as candidates for the surrogate outcome. To construct an informative surrogate, we relied on actual data and regressed the target outcome w.r.t the pre-surgical survey responses, and used the predicted values as a single informative surrogate outcome.

We conducted two analyses to investigate the performance of the proposed method. In the first analysis, we compared the proposed methods with the baseline methods in terms of the deviance $E \left[\ell(\hat{\beta}) \right]$, where $\ell(\hat{\beta}) = b(\mathbf{X}^\top \hat{\beta}) - \hat{g}(\hat{\Gamma}^\top \tilde{\mathbf{X}}) \mathbf{X}^\top \hat{\beta}$. Specifically, we randomly split the entire dataset into a training dataset and a testing dataset with equal sample sizes. We estimated coefficients using the training dataset, and then we calculated the deviance on the testing dataset. This procedure was repeated 1000 times. In the second analysis, we fitted the model on the entire dataset and compare the variables selected by different methods.

Table 2 shows that the proposed method with Z achieves the lowest deviance. In terms of the selected variables, compared with other methods, the proposed method with Z uniquely revealed that geriatric patients were less likely to achieve the MCID (estimated coefficient is 0.367; 95%-CI

Table 2: Comparison of averaged deviances (standard deviations) in real data example

Baseline1	Baseline2	Proposed w/o Z	Proposed w/ Z
0.714 (0.105)	0.696 (0.094)	0.685 (0.091)	0.670 (0.094)

is $[0.020, 0.714]$). This is in accordance with the existing research finding that elderly patients were more likely to have post-operative adverse clinical outcomes than younger patients in total joint replacement (Higuera et al., 2011; Malkani et al., 2017). The coefficients with confidence intervals for other covariates are presented in the Online Supporting Information.

6 Discussion

In this work, we propose a debias approach to infer the parameters of interest under a possibly misspecified GLM. This approach uses an informative surrogate outcome which leads to a low-dimensional flexible imputation model, and estimates a low-dimensional weighting function instead of the complex propensity score. When the true propensity happens to enjoy the same low-dimensional structure, the proposed method achieves the semi-parametric efficiency lower bound. Compared with the double machine learning method, the proposed approach relaxes the requirement on the propensity estimation and maintains almost the same flexibility/requirement on the imputation model estimation.

There are multiple future directions to extend the proposed approach. First, we can consider extending the proposed approach to high-dimensional settings where $p/n \rightarrow +\infty$. In a high-dimensional setting, the $\hat{\Gamma}$ may not be asymptotic normal due to the possible penalization. In this case, in order to achieve an asymptotic normal estimator, an additional debias procedure is needed to adjust for the bias due to the estimation error of $\hat{\Gamma}$. Second, we can investigate more choices of the function h . In this work, to pursue a low-dimensional weighting function, we choose h such that $\mathcal{T}h = 0$ when constructing the weighting function. However, we can choose other potential alternatives to mitigate certain deficiencies such as a choice to minimize the asymptotic variance of the debiased estimator; a choice to avoid possible negative weights. Third, we can combine the proposed approach with the augmented minimax linear estimation (Hirshberg and Wager, 2021) to avoid the computation of the Riesz representer. Especially in high-dimensional setting, an explicit form of the weights to debias $\hat{\Gamma}$ might be intractable.

Supplemental Materials

Proofs of all theorems and additional simulation results are contained in the online supplemental materials.

References

- Anderer, A., Bastani, H., and Silberholz, J. (2021). Adaptive clinical trial designs with surrogates: When should we bother? *Management Science*.
- Cao, W., Tsiatis, A. A., and Davidian, M. (2009). Improving efficiency and robustness of the doubly robust estimator for a population mean with incomplete data. *Biometrika*, 96(3):723–734.
- Cheng, D., Ananthakrishnan, A., and Cai, T. (2018). Efficient and robust semi-supervised estimation of average treatment effects in electronic medical records data. *arXiv e-prints*, pages arXiv–1804.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Dufo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1):C1–C68.
- Cook, R. D. (2007). Fisher lecture: dimension reduction in regression. *Statist. Sci.*, 22(1):1–26.
- Fleming, T. R., Prentice, R. L., Pepe, M. S., and Glidden, D. (1994). Surrogate and auxiliary endpoints in clinical trials, with potential applications in cancer and aids research. *Statistics in medicine*, 13(9):955–968.
- Fontana, M. A., Lyman, S., Sarker, G. K., Padgett, D. E., and MacLean, C. H. (2019). Can machine learning algorithms predict which patients will achieve minimally clinically important differences from total joint arthroplasty? *Clinical orthopaedics and related research*, 477(6):1267.
- Frangakis, C. E. and Rubin, D. B. (2002). Principal stratification in causal inference. *Biometrics*, 58(1):21–29.
- Han, P. (2012). A note on improving the efficiency of inverse probability weighted estimator using the augmentation term. *Statistics & Probability Letters*, 82(12):2221–2228.
- Han, P., Wang, L., and Song, P. X.-K. (2016). Doubly robust and locally efficient estimation with missing outcomes. *Statistica Sinica*, pages 691–719.
- Higuera, C. A., Elsharkawy, K., Klika, A. K., Brocone, M., and Barsoum, W. K. (2011). 2010 mid-america orthopaedic association physician in training award: predictors of early adverse outcomes after knee and hip arthroplasty in geriatric patients. *Clinical Orthopaedics and Related Research*®[®], 469(5):1391–1400.

- Hirshberg, D. A. and Wager, S. (2021). Augmented minimax linear estimation. *The Annals of Statistics*, 49(6):3206–3227.
- Ho, A., Purdie, C., Tirosh, O., and Tran, P. (2019). Improving the response rate of patient-reported outcome measures in an australian tertiary metropolitan hospital. *Patient Related Outcome Measures*, 10:217.
- Horvitz, D. G. and Thompson, D. J. (1952). A generalization of sampling without replacement from a finite universe. *Journal of the American statistical Association*, 47(260):663–685.
- Hou, J., Guo, Z., and Cai, T. (2021). Surrogate assisted semi-supervised inference for high dimensional risk prediction. *arXiv preprint arXiv:2105.01264*.
- Kang, J. D. and Schafer, J. L. (2007). Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data. *Statistical science*, 22(4):523–539.
- Katakam, A., Karhade, A. V., Collins, A., Shin, D., Bragdon, C., Chen, A. F., Melnic, C. M., Schwab, J. H., and Bedair, H. S. (2022). Development of machine learning algorithms to predict achievement of minimal clinically important difference for the koos-ps following total knee arthroplasty. *Journal of Orthopaedic Research®*, 40(4):808–815.
- Liu, M., Zhang, Y., Liao, K. P., and Cai, T. (2020). Augmented transfer regression learning with semi-non-parametric nuisance models.
- Ma, Y. and Zhu, L. (2012). A semiparametric approach to dimension reduction. *Journal of the American Statistical Association*, 107(497):168–179.
- Ma, Y. and Zhu, L. (2013). Efficient estimation in sufficient dimension reduction. *Annals of Statistics*, 41(1):250–268.
- Malkani, A. L., Dilworth, B., Ong, K., Baykal, D., Lau, E., Mackin, T. N., and Lee, G.-C. (2017). High risk of readmission in octogenarians undergoing primary hip arthroplasty. *Clinical Orthopaedics and Related Research®*, 475(12):2878–2888.
- Prentice, R. L. (1989). Surrogate endpoints in clinical trials: definition and operational criteria. *Statistics in medicine*, 8(4):431–440.
- Pronk, Y., Pilot, P., Brinkman, J. M., van Heerwaarden, R. J., and van der Weegen, W. (2019). Response rate and costs for automated patient-reported outcomes collection alone compared to combined automated and manual collection. *Journal of patient-reported outcomes*, 3(1):1–8.

- Qin, J., Shao, J., and Zhang, B. (2008). Efficient and doubly robust imputation for covariate-dependent missing responses. *Journal of the American Statistical Association*, 103(482):797–810.
- Qin, J. and Zhang, B. (2007). Empirical-likelihood-based inference in missing response problems and its application in observational studies. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(1):101–122.
- Robins, J. M. and Rotnitzky, A. (1995). Semiparametric efficiency in multivariate regression models with missing data. *Journal of the American Statistical Association*, 90(429):122–129.
- Robins, J. M., Rotnitzky, A., and Zhao, L. P. (1994). Estimation of regression coefficients when some regressors are not always observed. *Journal of the American statistical Association*, 89(427):846–866.
- Rosenbaum, P. R. and Rubin, D. B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55.
- Rotnitzky, A., Lei, Q., Sued, M., and Robins, J. M. (2012). Improved double-robust estimation in missing data and causal inference models. *Biometrika*, 99(2):439–456.
- Rubin, D. B. (2004). *Multiple imputation for nonresponse in surveys*, volume 81. John Wiley & Sons.
- Rubin, D. B. and van der Laan, M. J. (2008). Empirical efficiency maximization: improved locally efficient covariate adjustment in randomized experiments and survival analysis. *The International Journal of Biostatistics*, 4(1).
- Silverman, B. W. (1978). Weak and strong uniform consistency of the kernel estimate of a density and its derivatives. *The Annals of Statistics*, pages 177–184.
- Tan, Z. (2006). A distributional approach for causal inference using propensity scores. *Journal of the American Statistical Association*, 101(476):1619–1637.
- Tan, Z. (2010). Bounded, efficient and doubly robust estimation with inverse weighting. *Biometrika*, 97(3):661–682.
- Vaart, A. W. v. d. (1998). *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- Xia, Y. (2007). A constructive approach to the estimation of dimension reduction directions. *The Annals of Statistics*, 35(6):2654–2690.

Xia, Y., Tong, H., Li, W. K., and Zhu, L.-X. (2002). An adaptive estimation of dimension reduction space. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 64(3):363–410.