

On the distribution of heat in fibered magnetic fields

Theodore D. Drivas, Daniel Ginsberg, and Hezekiah Grayer II

ABSTRACT. We study the equilibrium temperature distribution in a model for strongly magnetized plasmas in dimension two and higher. Provided the magnetic field is sufficiently structured (integrable in the sense that it is fibered by co-dimension one invariant tori, on most of which the field lines ergodically wander) and the effective thermal diffusivity transverse to the tori is small, it is proved that the temperature distribution is well approximated by a function that only varies across the invariant surfaces. The same result holds for “nearly integrable” magnetic fields up to a “critical” size. In this case, a volume of non-integrability is defined in terms of the temperature defect distribution and related the non-integrable structure of the magnetic field, confirming a physical conjecture of Paul–Hudson–Helander [6]. Our proof crucially uses a certain quantitative ergodicity condition for the magnetic field lines on full measure set of invariant tori, which is automatic in two dimensions for magnetic fields without null points and, in higher dimensions, is guaranteed by a Diophantine condition on the rotational transform of the magnetic field.

1. Introduction

The heat conduction in strongly magnetized plasmas is influenced locally by the direction of the magnetic field $B : \mathbb{R}^d \rightarrow \mathbb{R}^d$. Braginskii [3] (see also [4, 12]) derived an effective anisotropic diffusion equation for the temperature T in such an environment which, in steady state and free of heat sources, reads

$$\operatorname{div}(b\nabla_b T + \varepsilon\nabla_b^\perp T) = 0 \quad (1.1)$$

where, assuming the magnetic field has no null points $|B| \neq 0$, we introduced

$$b = \frac{B}{|B|}, \quad \nabla_b = b \cdot \nabla, \quad \nabla_b^\perp = \nabla - b\nabla_b. \quad (1.2)$$

This equilibrium equation captures macroscopically the phenomenon that charged particle dynamics strongly influenced by B favors collisions aligned with b . In (1.1), the parameter $\varepsilon > 0$ represents the ratio $\kappa_\perp/\kappa_\parallel$ of the transverse diffusion coefficient to the longitudinal. In general it is a scalar function of local density and field magnitude $|B|$, however its magnitude is small in many applications of interest where $|B|$ is large. In our work, we treat ε as a constant and study the limit $\varepsilon \rightarrow 0$.

For arbitrary B , it is not immediate what emerges in the limit $\varepsilon \rightarrow 0$ of (1.1), given some fixed boundary conditions. We focus on toroidal “Arnold fibered” fields B . These are solenoidal vector fields B having the property that there is a smooth function $\psi : D \rightarrow \mathbb{R}$ defined in a bounded region $D \subset \mathbb{R}^d$ with $|\nabla\psi| \neq 0$ in D , whose level sets S_ψ are $(d-1)$ -dimensional tori such that ψ is a first integral

$$B \cdot \nabla\psi = 0. \quad (1.3)$$

We shall term these fields (*toroidally fibered*). In two dimensions, if B is divergence-free and sufficiently smooth, then $B = \nabla^\perp A$ for a “streamfunction” $A : D \rightarrow \mathbb{R}$ where $\nabla^\perp = (-\partial_y, \partial_x)$. If B has no nulls, then $|\nabla A| > 0$, so any non-vanishing divergence-free field in two dimensions is fibered by its streamfunction, e.g. $\psi = A$. See Figure 1 (a). In three dimensions, its straightforward to

write down explicit fibered fields, see (1.13) and Figure 1 (b). Moreover, as we will later discuss, non-degenerate magnetohydrostatic equilibria have this property.




FIGURE 1. Examples of fibered magnetic fields. Left: a 2d magnetic field without null points on a (topologically) annular domain – the periodic channel. Integral curves are levels of the streamfunction. Right: a 3d toroidal magnetic field; depicted in grayscale are distinct level surfaces of the first integral, ψ , the flux function.

The temperature equation (1.1) is to be solved in a toroidal shell D with boundaries S_{\pm} that are level sets of the first integral ψ . Call $\psi_- := \inf_D \psi$ and $\psi_+ := \sup_D \psi$. Since ψ is non-degenerate by assumption, S_{\pm} are the levels corresponding to the values ψ_{\pm} . To complete the problem, we impose Dirichlet boundary conditions for the temperature field $T : D \rightarrow \mathbb{R}$ on these surfaces. Overall, we consider the system

$$\begin{aligned} \operatorname{div}(b\nabla_b T + \varepsilon\nabla_b^{\perp} T) &= 0 && \text{in } D, \\ T &= T_{\pm}, && \text{on } S_{\pm}, \end{aligned} \quad (1.4)$$

for constants T_-, T_+ . The 1.4 system is used in practice as an efficient method to visualize the flux surfaces of the magnetic field [4, 5, 6].

To state our main result concerning the convergence of T_{ε} , we use some notions from mixing to characterize the behavior of B via its trajectories on the flux surfaces S_{ψ} . Denoting $I = [\psi_-, \psi_+]$, we say that S_{ψ} is an “ergodic” surface if ψ is in the set

$$E(\gamma, M) := \left\{ \psi \in I : \|u\|_{\dot{H}^{-\gamma}(S_{\psi})} \leq M \|\nabla_B u\|_{L^2(S_{\psi})}, \text{ for all } u \in H^1(S_{\psi}) \right\} \quad (1.5)$$

for some nonnegative γ and M , where $\dot{H}^{\gamma}(S_{\psi})$ denotes the homogeneous Sobolev space of index γ on S_{ψ} . The sets $E(\gamma, M)$ of ergodic values may be empty or may have full measure, depending on B . The definition of these sets is motivated by a Diophantine condition, see (4.1). We then define the collection

$$N(\gamma, M) = I \setminus E(\gamma, M), \quad (1.6)$$

of “non-ergodic” values of ψ . Note that if $M > M'$ then $N(\gamma, M) \subseteq N(\gamma, M')$.

DEFINITION 1. We say that B satisfies the “ergodicity condition” if, with $N(\gamma, M)$ defined as in (1.6), for some $c, \gamma > 0$, we have

$$\lim_{M \rightarrow \infty} M^c \mu(N(\gamma, M)) = 0, \quad (1.7)$$

where μ denotes the one-dimensional Lebesgue measure.

Our main result below roughly states that, provided B is ergodic on almost all of the surfaces S_{ψ} such that the ergodicity condition holds, the temperatures profiles T_{ε} indeed converge (in $H^1(D)$) to the effective temperature T_0 . A consequence of our theorem is that the limiting temperature profile T_0 itself fibers B . This fact partially motivated the work of Paul–Hudson–Helander [6].

THEOREM 1.1. *Let $d \geq 2$ and let B be toroidally fibered by ψ , and let D be the region bounded by two level sets S_{\pm} . For $\varepsilon > 0$, let $T_{\varepsilon} : D \rightarrow \mathbb{R}$ be the solution of system (1.4) for constants T_{-} and T_{+} . If the ergodicity condition from Definition 1 holds, then*

$$T_{\varepsilon} \rightarrow T_0 := \Theta(\psi) \quad \text{in } H^1(D) \quad (1.8)$$

where $\Theta(\psi)$ is the solution of the one-dimensional boundary-value problem on $\psi \in [\psi_{-}, \psi_{+}]$:

$$\frac{d}{d\psi} \left(\frac{d\Theta}{d\psi} \int_{S_{\psi}} |\nabla\psi| d\mathcal{H}^{(d-1)} \right) = 0, \quad \Theta(\psi_{\pm}) = T_{\pm}. \quad (1.9)$$

In fact, there is $C := C(D, B) > 0$ such that

$$\|T_{\varepsilon} - T_0\|_{H^1(D)} \leq C\varepsilon^{\frac{c}{2+c}}, \quad (1.10)$$

where c is the largest so that there is a $\gamma > 0$ making condition (1.7) of Definition 1 hold.

The proof is found in §2. Briefly, if each integral curve of $B|_{S_{\psi}}$ covers S_{ψ} densely for some ψ , (that is, if S_{ψ} is an ‘‘irrational torus’’), then B ‘‘nearly’’ spans the tangent space at each point. On such a torus, one encounters a small divisors problem; the operator ∇_B is bounded below on S_{ψ} but the lower bound may be arbitrarily small. However, for $\psi \in E(\gamma, M)$, this lower bound cannot be less than $1/M$. On the complement $N(\gamma, M)$, the operator ∇_B is not bounded below, but the ergodicity condition (1.7) ensures that the measures of the sets $N(\gamma, M)$ go to zero as M increases. The net result is one of homogenization to a one-dimensional limit profile adapted to the geometry of the invariant tori that satisfies an effective diffusion equation. See §2 for further discussion.

In the upcoming Corollaries 1.1, 1.2, we show that this condition holds for a large family of physically-relevant vector fields B . Whenever $d = 2$, the sets $N(\gamma, M)$ are empty for large enough M ; that is, every surface S_{ψ} is ergodic in this setting (in three and higher-dimensions, the ergodicity condition need not be true in general). Thus c in bound (1.10) may be taken to ∞ for any $\gamma \geq 0$. It follows from our main theorem that, in this case, we have convergence of T_{ε} to the effective temperature T_0 . More quantitatively:

COROLLARY 1.1. *Let $d = 2$ and let B be a non-vanishing divergence-free vector field. Then*

$$\|T_{\varepsilon} - T_0\|_{H^1(D)} \leq C\varepsilon, \quad (1.11)$$

where $T_0 = \Theta(\psi)$ where Θ is given by (2.6).

In three dimensions, an important example of fibered fields are the smooth solutions of the magnetohydrostatic equations

$$(\text{curl } B) \times B = \nabla p, \quad \text{div } B = 0, \quad \text{in } D \subset \mathbb{R}^3, \quad (1.12)$$

having the property that the pressure satisfies $\nabla p \neq 0$. As noted by Arnold [1, 2] since $|\nabla p|$ is nonvanishing by assumption, each surface S_p is a smooth two-dimensional surface which admits two everywhere transverse non-vanishing tangent vector fields ($\text{curl } B$ and B) and are thus two-dimensional tori or cylinders. In this setting B is fibered by its pressure, $\psi = p$. It is straightforward to construct fields B of this type which are axisymmetric, see e.g. [11] and [14]. It is an open problem (see [14], [15]) to construct such smooth magnetohydrostatic equilibria with $|\nabla p| > 0$ outside of Euclidean symmetry.

More generally, in three dimensions, given a non-degenerate function $\psi : D \rightarrow \mathbb{R}$ whose level sets are tori along with functions $\theta, \phi : D \rightarrow \mathbb{R}$, any vector field of the form

$$B = \nabla\psi \times \nabla\theta + \nabla\phi \times \nabla\chi \quad (1.13)$$

is divergence-free. If χ is chosen so that $\chi = \chi(\psi, \phi)$, B is fibered by ψ , and known as “integrable” because the integral curves of B obey a Hamiltonian system with Hamiltonian χ , and this Hamiltonian is integrable in the usual sense¹ when $\partial_\theta \chi = 0$ (see (1.15)). See Figure 1, right panel. Fields of this form play an important role in the problem of confining a plasma with a magnetic field [13]. Such fields may sometimes be regarded as MHS solutions held steady by external forcing (e.g. by current carrying coils in some particular geometry) [9, 10].

Suppose θ, ϕ form a coordinate system on S_ψ and so we write $u = u(\psi, \theta, \phi)$. Then if B is as in (1.13) with $\partial_\theta \chi = \partial_\phi \chi = 0$, it follows after writing $\iota(\psi) = \chi'(\psi)$,

$$(B \cdot \nabla)u = [\partial_\phi u + \iota(\psi)\partial_\theta u] J, \quad (1.16)$$

where $J = \nabla\psi \times \nabla\theta \cdot \nabla\phi$. Generally, by a theorem of Sternberg [21], if B is *any* nonvanishing divergence-free vector field fibered by a function ψ , (in particular, this includes the case $\chi = \chi(\psi, \phi)$ of (1.13)) then on each S_ψ there are coordinates θ, ϕ and a number $\iota = \iota(\psi)$ so that, expressed in these coordinates, B takes the form (1.16) for a function $J = J(\psi, \theta, \phi) > 0$. We call the function ι from (1.16) the rotational transform. Our main result in three dimensions, proven in §4, is that provided ι is invertible with Lipschitz inverse, we have convergence $T_\varepsilon \rightarrow T_0$ in $H^1(D)$.

COROLLARY 1.2. *Suppose that B is a nonvanishing divergence-free vector field fibered by a function ψ . Suppose that the rotational transform ι from (1.16) is invertible and for some $L > 0$*

$$|\psi_1 - \psi_2| \leq L|\iota(\psi_1) - \iota(\psi_2)| \quad (1.17)$$

holds for all $\psi_1, \psi_2 \in I$. Then condition (1.7) holds for any $\gamma > 2$ with $c = 1$. Consequently,

$$\|T_\varepsilon - T_0\|_{H^1(D)} \leq C\varepsilon^{\frac{1}{3}}, \quad (1.18)$$

where $T_0 = \Theta(\psi)$ where Θ is given by (2.6).

In other words, we show that the ergodic condition holds for integrable Arnol’d fibered fields B with monotone rotational transform. Such fields are of specific interest in the plasma physics community, see the discussion in [9]. However such plasma equilibria, if they exist, may be unstable. Thus, it is important to also understand the behavior of non-integrable fields $\partial_\theta \chi \neq 0$. There is an obstruction: the behavior of particle transport (and thus of heat) in non-integrable fields can be quite complicated because non-integrable Hamiltonian systems may exhibit chaos.

In [6], the authors consider non-integrable magnetic fields taking the form (1.13) where

$$\chi_\varepsilon(\psi, \theta, \phi) = \chi_0(\psi) + \varepsilon^a \chi_1(\psi, \theta, \phi), \quad (1.19)$$

and $a \geq 1/2$. A modification of the proof of Theorem 1.1 (see Section 5) gives the following generalization of Corollary 1.2 to fields of this type which are “weakly nonintegrable.” We require

¹We suppose that with B as in (1.13), the functions θ, ϕ, ψ together form a coordinate system in D . Then for any smooth $u : D \rightarrow \mathbb{R}$, we have $\nabla u = \partial_\psi u \nabla\psi + \partial_\phi u \nabla\phi + \partial_\theta u \nabla\theta$ and so, writing $J = \nabla\psi \times \nabla\theta \cdot \nabla\phi$, which is nonvanishing by our assumption, we have the formula

$$(B \cdot \nabla)u = [\partial_\phi u + \iota(\psi, \theta, \phi)\partial_\theta u + \tau(\psi, \theta, \phi)\partial_\psi u] J, \quad (1.14)$$

where $\tau(\psi, \theta, \phi) := -\partial_\theta \chi(\psi, \theta, \phi)$ and where we have introduced the rotational transform $\iota(\psi, \theta, \phi) := \partial_\psi \chi(\psi, \theta, \phi)$. There is a simple interpretation of the function χ . Consider any integral curve of B , parametrized by ϕ . That is, we consider $\Psi(\phi), \vartheta(\phi)$ defined by

$$\frac{d}{d\phi} \Psi = \frac{B \cdot \nabla\psi}{B \cdot \nabla\phi} = -\partial_\theta \chi, \quad \frac{d}{d\phi} \vartheta = \frac{B \cdot \nabla\theta}{B \cdot \nabla\phi} = \partial_\psi \chi, \quad (1.15)$$

with the understanding that the quantities on the right-hand sides are evaluated at $(\psi, \theta, \phi) = (\Psi(\phi), \vartheta(\phi), \phi)$. Thus the integral curves of B satisfy a Hamiltonian system with Hamiltonian χ . Note that if $\partial_\theta \chi = 0$, the above system is integrable (has a conserved quantity) since ψ is constant along the flow. This also be seen from the formula (1.14).

the following anisotropic Sobolev spaces tailored to the invariant tori: $f \in L^2(D)$ which are finite in the norm

$$\|f\|_{H^{(0,\gamma)}}^2 := \int_{\psi_-}^{\psi_+} \|f(\psi, \cdot)\|_{H^\gamma(S_\psi)}^2 d\psi. \quad (1.20)$$

THEOREM 1.2. *Suppose that B is as in (1.13) where χ_ε is given by (1.19) for $a \geq 1/2$ and satisfies $\|\partial_\theta \chi_1\|_{L^\infty(D)} < 1$ and $\partial_\theta \chi_1|_{\partial D} = 0$. Moreover, assume that $\iota = \chi'_0$ satisfies the condition from Corollary 1.2. Then letting B_0 be defined by (1.13) with $\varepsilon = 0$, there is a constant $C = C(L)$ such that*

$$\|\nabla_{b_0}^\perp (T_\varepsilon - T)\|_{L^2} + \|T_\varepsilon - T\|_{L^2}^2 \leq C\varepsilon^{2a-1} \|\partial_\theta \chi_1\|_{L^2}^2 + C\varepsilon^{\frac{1}{3}} (\|\Delta T_0\|_{H^{(0,\gamma)}}^2 + \|\Delta T_0\|_{L^\infty}^2), \quad (1.21)$$

$$\|\nabla_{b_0} (T_\varepsilon - T)\|_{L^2} \leq C\varepsilon^{2a} \|\partial_\theta \chi_1\|_{L^2}^2 + C\varepsilon^{\frac{4}{3}} (\|\Delta T_0\|_{H^{(0,\gamma)}}^2 + \|\Delta T_0\|_{L^\infty}^2). \quad (1.22)$$

When $a > 1/2$, the estimate (1.21) implies that $T_\varepsilon \rightarrow T_0$ almost everywhere. When $a = 1/2$, we show the same result is true, and in this case, even though $\rho^* = \lim_{\varepsilon \rightarrow 0} T_\varepsilon - T_0 \in H^1(D)$ may not vanish almost everywhere, we find $\rho^*|_{S_\psi} = 0$ for all ψ except possibly for a family of ψ lying in a set of measure zero. Here, $\rho^*|_{S_\psi}$ denotes the trace of the function ρ^* on the surface S_ψ , and this quantity is well-defined whenever $\rho^* \in H^1(D)$, by the trace theorem. In fact, the support of $\rho^* = \lim_{\varepsilon \rightarrow 0} \rho_\varepsilon$ is contained in the collection of non-ergodic surfaces $N(\gamma) = I \setminus E(\gamma)$, which has measure zero, where $E(\gamma)$ denote the family of ergodic surfaces

$$E(\gamma) = \bigcup_{M>0} E(\gamma, M). \quad (1.23)$$

COROLLARY 1.3. *Under the hypotheses of Corollary 1.2 with $a = 1/2$, the sequence $\rho_\varepsilon = T_\varepsilon - T_0$ converges weakly in H^1 to a distribution ρ^* in $H^1(D)$ with the property that*

$$\rho^*|_{S_\psi} = 0, \quad \text{whenever } \psi \in E(\gamma). \quad (1.24)$$

That is, the support of ρ^ is contained in $N(\gamma) = I \setminus E(\gamma)$.*

Our final result relates directly to the work of [6]. In that paper, the authors consider the sets

$$\mathcal{N}(\varepsilon) = \{(\psi, \theta, \phi) : |\nabla_b T(\psi, \theta, \phi)|^2 \geq \varepsilon |\nabla_b^\perp T(\psi, \theta, \phi)|^2\}, \quad (1.25)$$

where $b = B/|B|$ with B as in (1.13), and study them as a proxy for the ‘‘non-integrability’’ of the field B . Using the estimates from the above section, we can get an upper bound on the measure set in the limit $\varepsilon \rightarrow 0$.

PROPOSITION 1.1. *Define B as in (1.13) and $\mathcal{N}(\varepsilon)$ as in (1.25). Suppose that the boundary values T_\pm from (1.9) satisfy $T_+ \neq T_-$. Under the hypotheses of Corollary 1.2, there is a constant C depending continuously on $\|T'_0\|_{L^\infty}$, $\|1/T'_0\|_{L^\infty}$, $\|T''_0\|_{L^\infty}$, $(1 - \|\nabla \chi_1\|_{L^\infty})^{-1}$ and $(2 - \gamma)^{-1}$ so that*

$$\mu(\mathcal{N}(\varepsilon)) \leq C \left(\varepsilon^{2a-1} \|\partial_\theta \chi_1\|_{L^\infty}^2 + \varepsilon^{1/3} \|\Delta T_0\|_{H^{(0,\gamma)}}^2 \right). \quad (1.26)$$

Note that the ‘‘integrable’’ case corresponds to taking $\chi_1 = 0$ and it follows that in this case

$$\lim_{\varepsilon \rightarrow 0} \mu(\mathcal{N}(\varepsilon)) = 0. \quad (1.27)$$

Since now $\chi = \chi_0$ is integrable, this agrees with the fact that the effective volume of non-integrability is zero. If χ_1 is nonzero, we get the same result with $a > 1/2$ but if $a = 1/2$ we instead have

$$\lim_{\varepsilon \rightarrow 0} \mu(\mathcal{N}(\varepsilon)) \leq C \|\partial_\theta \chi_1\|_{L^\infty}^2. \quad (1.28)$$

This exhibits a relationship between the volume of the set (1.25) and the non-integrability of the Hamiltonian χ , captured by the θ -dependence of the perturbation χ_1 . Our proof of the Proposition (see Section 5) partially confirms a conjecture announced in [6].

2. Proof of Theorem 1.1

The limiting profile $\Theta(\psi)$ is a function of the flux function ψ only, and is determined from the following heuristic. If B is fibered, expanding $T_\varepsilon = T_0 + \varepsilon T_1$ leads to

$$\operatorname{div}(b\nabla_b T_0) = 0, \quad (2.1)$$

$$\operatorname{div}(\nabla_b^\perp T_0) = -\operatorname{div}(b\nabla_b T_1). \quad (2.2)$$

Equation (2.1) is underdetermined for T_0 ; indeed noting that $b \cdot \nabla\psi = 0$ one sees that any function

$$T_0 = \Theta(\psi) \quad (2.3)$$

will satisfy (2.1). This arbitrariness of Θ may be eliminated by considering the second condition. Indeed, note that since b is tangent to each surface S_ψ , we have

$$\frac{\operatorname{div}(b\nabla_b T_1)}{|\nabla\psi|} \Big|_{S_\psi} = \operatorname{div}_{S_\psi} \left(\left[\frac{b\nabla_b T_1}{|\nabla\psi|} \right] \Big|_{S_\psi} \right) \quad (2.4)$$

where $\operatorname{div}_{S_\psi}$ denotes the divergence operator on S_ψ . See Lemma A.2. In light of this, equation (2.2) comes with the following compatibility condition: on each invariant torus S_ψ ,

$$\int_{S_\psi} \frac{\Delta T_0}{|\nabla\psi|} d\mathcal{H}^{(d-1)} = 0, \quad (2.5)$$

where $d\mathcal{H}^{(d-1)}$ is the $(d-1)$ dimensional Hausdorff measure on S_ψ . Here we used that $\nabla_b^\perp T_0 = \nabla T_0$ since $b \cdot \nabla T_0 = 0$. Because T_0 is constant on S_ψ , it follows from Lemma (A.1) (see (A.5)) that the solvability requirement (2.5) and the boundary conditions of (1.4) are satisfied if Θ is the unique solution to (1.9). From (1.9), we deduce that the effective temperature distribution is give explicitly by

$$\Theta(\psi) = T_- + (T_+ - T_-) \frac{H(\psi; \psi_-)}{H(\psi_+; \psi_-)}, \quad \text{where} \quad H(\psi; \psi_-) := \int_{\psi_-}^{\psi} \frac{ds}{\Gamma(s)}. \quad (2.6)$$

Note that in two-dimensions, $\Gamma(\psi)$ is simply the circulation of the vector field B on the circle S_ψ :

$$\Gamma(\psi) = \int_{S_\psi} |\nabla\psi| d\mathcal{H} = \int_{S_\psi} B \cdot d\ell \quad \text{in two dimensions.} \quad (2.7)$$

Observing (2.5), we start with the following simple lemma. We first introduce the homogenous fractional Sobolev seminorms on S_ψ . For each S_ψ we pick coordinates $\theta_1, \dots, \theta_{d-1}$ on S_ψ such that θ_j maps S_ψ to $[0, 2\pi]$. For $k \in \mathbb{Z}^{d-1}$, we define

$$\widehat{u}(k) = \frac{1}{(2\pi)^{d-1}} \int_{[0, 2\pi]^{d-1}} \prod_{j=1}^{d-1} e^{ik_j \theta_j} u(\theta_1, \dots, \theta_{d-1}) d\theta_1 \cdots d\theta_{d-1}, \quad (2.8)$$

and then for $\gamma \in \mathbb{R}$, we define $\|\cdot\|_{\dot{H}^\gamma}$ by

$$\|u\|_{\dot{H}^\gamma(S_\psi)}^2 = \sum_{k \in \mathbb{Z}^{d-1} \setminus 0} |k|^{2\gamma} |\widehat{u}(k)|^2. \quad (2.9)$$

Recall also that (1.20) defines the anisotropic spaces $H^{(0,\gamma)}$ tailored to the tori. The result is then

LEMMA 2.1. *Suppose that $F \in H^{(0,\gamma)}(D) \cap L^\infty(D)$ for some $\gamma \geq 0$ and that F satisfies*

$$\int_{S_\psi} \frac{F}{|\nabla\psi|} d\mathcal{H}^{(d-1)} = 0, \quad (2.10)$$

for all $\psi \in [\psi_-, \psi_+]$. There is a constant C depending only on D so that for any $M > 0$ we have

$$\left| \int_D F u d\mu \right| \leq CM \|F\|_{H^{(0,\gamma)}} \|\nabla_b u\|_{L^2} + C\mu(N(\gamma, M))^{1/2} \|F\|_{L^\infty} \|u\|_{L^2}. \quad (2.11)$$

PROOF. By the co-area formula (A.1), we have

$$\int_D Fu \, d\mu = \int_{\psi_-}^{\psi_+} \int_{S_\psi} \frac{F}{|\nabla\psi|} u \, d\mathcal{H}^{(d-1)} \, d\psi. \quad (2.12)$$

Now, for each ψ , we write

$$\int_{S_\psi} \frac{F}{|\nabla\psi|} u \, d\mathcal{H}^{(d-1)} = \int_{[0,2\pi]^{d-1}} G(\psi, \theta_1, \dots, \theta_{d-1}) u(\psi, \theta_1, \dots, \theta_{d-1}) \, d\theta_1 \cdots d\theta_{d-1}, \quad (2.13)$$

where $G = \frac{F}{|\nabla\psi|} |h|^{1/2}$, where we are writing the metric on S_ψ as $h = h_{\alpha\beta} d\theta^\alpha d\theta^\beta$ and $|h| = \det h_{\alpha\beta}$. By Parseval's theorem,

$$\int_{[0,2\pi]^{d-1}} G(\psi, \theta_1, \dots, \theta_{d-1}) u(\psi, \theta_1, \dots, \theta_{d-1}) \, d\theta_1 \cdots d\theta_{d-1} = (2\pi)^{d-1} \sum_{k \in \mathbb{Z}^{d-1}} \widehat{G}(\psi, k) \widehat{u}(\psi, -k), \quad (2.14)$$

where \widehat{F}, \widehat{u} are defined as in (2.8). Now we note that

$$\widehat{G}(\psi, 0) = \int_{[0,2\pi]^{d-1}} \frac{F}{|\nabla\psi|} |h|^{1/2} \, d\theta_1 \cdots d\theta_{d-1} = \int_{S_\psi} \frac{F}{|\nabla\psi|} \, d\mathcal{H}^{(d-1)} = 0, \quad (2.15)$$

by assumption. We therefore have

$$\int_{S_\psi} \frac{F}{|\nabla\psi|} u \, d\mathcal{H}^{(d-1)} = (2\pi)^{d-1} \sum_{k \in \mathbb{Z}^{d-1} \setminus \{0\}} \widehat{G}(\psi, k) \widehat{u}(\psi, -k), \quad (2.16)$$

and it follows that

$$\left| \int_D Fu \, d\mu \right| \leq C \int_{\psi_-}^{\psi_+} \sum_{|k| \neq 0} |\widehat{G}(\psi, k) \widehat{u}(\psi, -k)| \, d\psi. \quad (2.17)$$

Now we split $[\psi_-, \psi_+] = E(\gamma, M) \cup N(\gamma, M)$ and bound

$$\begin{aligned} & \int_{\psi \in E(\gamma, M)} \sum_{|k| \neq 0} |\widehat{G}(\psi, k) \widehat{u}(\psi, -k)| \, d\psi \\ & \leq \int_{\psi \in E(\gamma, M)} \|G\|_{H^\gamma(S_\psi)} \|u\|_{\dot{H}^{-\gamma}(S_\psi)} \, d\psi \leq M \int_{\psi \in E(\gamma, M)} \|G\|_{H^\gamma(S_\psi)} \|\nabla_B u\|_{L^2(S_\psi)} \, d\psi \\ & \leq M \|G\|_{H^{(0,\gamma)}(D)} \|\nabla_B u\|_{L^2(D)}, \end{aligned} \quad (2.18)$$

by the definition of $E(\gamma, M)$ and

$$\begin{aligned} & \int_{\psi \in N(\gamma, M)} \sum_{|k| \neq 0} |\widehat{G}(\psi, k) \widehat{u}(\psi, -k)| \, d\psi \leq \|G\|_{L^2(N(\gamma, M))} \|u\|_{L^2(D)} \\ & \leq \mu(N(\gamma, M))^{1/2} \|G\|_{L^\infty(D)} \|u\|_{L^2(D)}. \end{aligned} \quad (2.19)$$

This gives the result. \square

Our main result, Theorem 1.1, will be a direct result of the following estimate.

PROPOSITION 2.1. *Define $T_0 = \Theta(\psi)$ where Θ is given by (2.6) and let $\rho = T_\varepsilon - T_0$. Under the hypotheses of Theorem 1.1, there is a constant C depending only on the domain D so that for each $\varepsilon > 0$ and $M > 0$, we have*

$$\|\nabla_b \rho\|_{L^2}^2 + \varepsilon \|\nabla_b^\perp \rho\|_{L^2}^2 + \varepsilon \|\rho\|_{L^2}^2 \leq C (M^2 \varepsilon^2 + \varepsilon \mu(N(\gamma, M))) (\|\Delta T_0\|_{H^{(0,\gamma)}}^2 + \|\Delta T_0\|_{L^\infty}^2) \quad (2.20)$$

PROOF. The remainder ρ satisfies

$$\operatorname{div}(b\nabla_b\rho) + \varepsilon \operatorname{div}(\nabla_b^\perp\rho) = -\varepsilon\Delta T_0, \quad \text{in } D, \quad (2.21)$$

$$\rho|_{S_\pm} = 0, \quad (2.22)$$

where we wrote $\Delta T_0 = \operatorname{div}(\nabla_b^\perp T_0)$ since $b \cdot \nabla T_0 = 0$. If we multiply (2.21) by ρ and integrate over D , using the co-area formula (A.1) we find

$$\int_D \left(|\nabla_b\rho|^2 + \varepsilon |\nabla_b^\perp\rho|^2 \right) d\mu = \varepsilon \int_D \Delta T_0 \rho d\mu = \varepsilon \int_{\psi_-}^{\psi_+} \int_{S_\psi} \frac{\Delta T_0}{|\nabla\psi|} \rho d\mathcal{H}^{(d-1)} d\psi. \quad (2.23)$$

Recall that we have defined T_0 so that $F = \Delta T_0$ satisfies the condition (2.10). We can therefore apply Lemma 2.1, and by (2.11) we have

$$\begin{aligned} \varepsilon \left| \int_{\psi_0}^{\psi_1} \int_{S_\psi} \frac{\Delta T_0}{|\nabla\psi|} \rho d\mathcal{H}^{(d-1)} d\psi \right| & \leq M\varepsilon \|\Delta T_0\|_{H(0,\gamma)} \|\nabla_b\rho\|_{L^2} + \varepsilon\mu(N(\gamma, M))^{1/2} \|\Delta T_0\|_{L^\infty} \|\rho\|_{L^2} \\ & \leq \frac{1}{2} \left(M^2\varepsilon^2 \|\Delta T_0\|_{H(0,\gamma)}^2 + \frac{1}{\delta} \varepsilon\mu(N(\gamma, M)) \|\Delta T_0\|_{L^\infty}^2 \right) + \frac{1}{2} \|\nabla_b\rho\|_{L^2}^2 + \frac{\delta}{2} \varepsilon \|\rho\|_{L^2}^2 \end{aligned} \quad (2.24)$$

for any $\delta > 0$. Since $\rho|_{\partial D} = 0$, by Poincaré's inequality we have

$$\|\rho\|_{L^2}^2 \leq C_P \|\nabla\rho\|_{L^2}^2, \quad (2.25)$$

where C_P is the Poincaré constant for D . Taking δ so that δC_P is sufficiently small, we see from (2.23) and (2.24) that there is a constant $C > 0$ depending only on C_P so that

$$\|\nabla_b\rho\|_{L^2}^2 + \varepsilon \|\nabla_b^\perp\rho\|_{L^2}^2 + \varepsilon \|\rho\|_{L^2}^2 \leq C (M^2\varepsilon^2 + \varepsilon\mu(N(\gamma, M))) (\|\Delta T_0\|_{H(0,\gamma)}^2 + \|\Delta T_0\|_{L^\infty}^2), \quad (2.26)$$

after using (2.25) again to bound $\varepsilon \|\rho\|_{L^2}^2 \leq C' \|\nabla_b\rho\|_{L^2}^2 + C' \varepsilon \|\nabla_b^\perp\rho\|_{L^2}^2$ for another constant C' . \square

PROOF OF THEOREM 1.1. If we take $M = \varepsilon^{-\frac{1}{2+c}}$, then writing $N_\varepsilon = N(\gamma, \varepsilon^{-\frac{1}{2+c}})$, (2.20) gives

$$\|\nabla_b^\perp\rho\|_{L^2} + \|\rho\|_{L^2}^2 \leq C\varepsilon^{\frac{c}{2+c}} \left(1 + \varepsilon^{-\frac{c}{2+c}} \mu(N_\varepsilon) \right) (\|\Delta T_0\|_{H(0,\gamma)}^2 + \|\Delta T_0\|_{L^\infty}^2), \quad (2.27)$$

$$\|\nabla_b\rho\|_{L^2} \leq C\varepsilon^{1+\frac{c}{2+c}} \left(1 + \varepsilon^{-\frac{c}{2+c}} \mu(N_\varepsilon) \right) (\|\Delta T_0\|_{H(0,\gamma)}^2 + \|\Delta T_0\|_{L^\infty}^2). \quad (2.28)$$

By assumption, $\lim_{\varepsilon \rightarrow 0} \varepsilon^{-\frac{c}{2+c}} \mu(N_\varepsilon) = 0$ and the result follows. \square

3. Proof of Corollary 1.1: The 2d case

If $|B| > 0$ in D , B is fibered by its streamfunction ψ , $B = \nabla^\perp\psi$. We bound

$$\|u\|_{\dot{H}^0(S_\psi)}^2 = \sum_{k \in \mathbb{Z} \setminus 0} |\widehat{u}(k)|^2 \leq \sum_{k \in \mathbb{Z}} |k|^2 |\widehat{u}(k)|^2 \leq \frac{C}{\inf_{S_\psi} |B|^2} \|\nabla_B u\|_{L^2(S_\psi)}^2 \quad (3.1)$$

for a constant $C > 0$. Here we have used that B spans the tangent space to S_ψ at each point. Therefore, $E(0, M) = D$ whenever $M \geq \frac{C}{\inf_D |B|}$, and so $N(0, M)$ is empty in this case. Thus (1.7) holds for any $c \geq 0$ and the result follows.

4. Proof of Corollary 1.2: The 3d integrable case

We first show that if $\gamma > 2$, under the hypotheses of Corollary 1.2, the condition (1.7) holds with $c = 1$. We start by relating this condition to the ‘‘Diophantine’’ condition.

Let $I = [\psi_-, \psi_+]$. Fix $\iota = \iota(\psi)$ with $\iota \in L^\infty(I)$. Let $|(m, n)| = \sqrt{m^2 + n^2}$. We define

$$D(\gamma, M) = \left\{ \psi \in I : |m + \iota(\psi)n| \geq \frac{1}{M|(m, n)|^\gamma} \text{ for all } (m, n) \in \mathbb{Z}^2 \setminus \{0\} \right\}. \quad (4.1)$$

If $\psi \in D(\gamma, M)$ for some γ, M , we will say that ψ is ‘‘Diophantine’’ and that the surface S_ψ is a ‘‘Diophantine surface’’. With $m + \iota(\psi)n$ replaced by $\omega \cdot (m, n)$ for $\omega \in \mathbb{R}^2$, these sets play a fundamental role in the proof of the celebrated KAM theorem [19].

At least when $\iota(\psi) = \psi$, these sets are empty if $\gamma < 2$ (see [18]), but it turns out that if ι is bi-Lipschitz and $\gamma > 2$ they have positive measure; in fact the complement of $\cup_{M>0} D(\gamma, M)$ has zero measure, as the next result shows.

LEMMA 4.1. *Let $\iota : I \rightarrow \mathbb{R}$ be an invertible function and suppose there is $L > 0$ so that*

$$\frac{1}{L} |\psi_1 - \psi_2| \leq |\iota(\psi_1) - \iota(\psi_2)| \leq L |\psi_1 - \psi_2|. \quad (4.2)$$

Define $D(\gamma, M)$ as in (4.1) and let μ denote the one-dimensional Lebesgue measure. If $\gamma > 2$ and $M > 0$, there is a constant K depending only on $\mu(I)$, L , and $1/(\gamma - 2)$ so that

$$\mu(I \setminus D(\gamma, M)) \leq K \left(\frac{1}{M^{\frac{1}{1+\gamma}}} + \frac{1}{M} \right). \quad (4.3)$$

Taking $M \rightarrow \infty$ in (4.3) shows that the set of ψ which fails the condition (4.1) for *all* $M > 0$ has zero measure. Equivalently, the set of ψ the condition in (4.1) for *some* M has full measure, though in general the complement may be nonempty.

PROOF. This result follows from a straightforward modification of the argument from e.g. [18]. We include the details here for the convenience of the reader. For $(m, n) \in \mathbb{Z}^2 \setminus (0, 0)$, we define

$$\Pi_{(m,n)}(\gamma, M) = \left\{ \psi \in I : |m + \iota(\psi)n| < \frac{1}{M|(m, n)|^\gamma} \right\}, \quad (4.4)$$

so that

$$I \setminus D(\gamma, M) \subseteq \bigcup_{(m,n) \neq (0,0)} \Pi_{(m,n)}(\gamma, M). \quad (4.5)$$

If $n \neq 0$, $\Pi_{(m,n)}(\gamma, M)$ is contained in the interval $[\psi_1, \psi_2] \subset I$ where ψ_1 are such that $\iota(\psi_1) = \frac{M}{n} \frac{1}{|(m,n)|^\gamma} + \frac{m}{n}$ and $\iota(\psi_2) = -\iota(\psi_1)$ (such ψ_1, ψ_2 exist and are unique since by (4.2) ι is invertible). These are the maximal and minimal values of ψ such that (4.4) holds for a given (m, n) . Therefore

$$\mu(\Pi_{(m,n)}(\gamma, M)) \leq |\psi_1 - \psi_2| \leq \frac{L}{2M|n|||(m, n)|^\gamma}. \quad (4.6)$$

If $n = 0$, $\Pi_{(m,0)}(\gamma, M) = I$ when $|m| < M^{-1/(1+\gamma)}$ and it is empty otherwise. Therefore

$$\begin{aligned} \sum_{(m,n) \neq (0,0)} \mu(\Pi_{(m,n)}(\gamma, M)) &\leq \sum_{m \neq 0} \mu(\Pi_{(m,0)}(\gamma, M)) + \sum_{(m,n) \neq (0,0), n \neq 0} \mu(\Pi_{(m,n)}(\gamma, M)) \\ &\leq |I| \frac{1}{M^{1/(1+\gamma)}} + \frac{L}{M} + \frac{L}{2M} \sum_{|m|, |n| \geq 1} \frac{1}{|n|||(m, n)|^\gamma}, \end{aligned} \quad (4.7)$$

since there are two terms in the second sum on the first line with $m = 0$. The last term here is bounded by

$$\begin{aligned} 2 \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n} \frac{1}{(m^2 + n^2)^{\gamma/2}} &\leq 2 \sum_{m=1}^{\infty} \int_1^{\infty} \frac{dz}{z(m^2 + z^2)^{\gamma/2}} = 2 \sum_{m=1}^{\infty} \frac{1}{m^{\gamma}} \int_{1/m}^{\infty} \frac{dw}{w(1+w^2)^{\gamma/2}} \\ &\leq 2 \sum_{m=1}^{\infty} \frac{1}{m^{\gamma-1}} \int_0^{\infty} \frac{dw}{(1+w^2)^{\gamma/2}} \end{aligned} \quad (4.8)$$

which is finite if $\gamma > 2$. In this case we therefore have

$$\mu(I \setminus D(\gamma, M)) \leq \mu \left(\bigcup_{(m,n) \neq (0,0)} \Pi_{m,n}(\gamma, M) \right) \leq C_1 \left(\frac{1}{M^{1/(1+\gamma)}} + \frac{1}{M} \right), \quad (4.9)$$

for a constant C_1 depending only on $|I|$, L and $1/(\gamma - 2)$, which proves (1.26). \square

The values of $\psi \in D(\gamma, M)$ are sometimes called ‘‘strongly non-resonant’’, and the surfaces S_{ψ} with $\psi \in D(\gamma, M)$ are called ‘‘non-resonant flux surfaces’’.

PROOF OF COROLLARY 1.2. We first consider a field of the form (1.13) when $\chi = \chi(\psi)$ and where θ, ϕ form a coordinate system on each S_{ψ} . With $\iota(\psi) = \chi'(\psi)$, in this setting we have

$$(B \cdot \nabla)u = J[\partial_{\phi} + \iota(\psi)\partial_{\theta}]u, \quad J = \nabla\psi \times \nabla\theta \cdot \nabla\phi \quad (4.10)$$

where $J \neq 0$ by assumption. We claim that the Diophantine surfaces are ergodic, in the sense that there is a constant C with

$$D(\gamma, CM) \subseteq E(\gamma, M) \quad (4.11)$$

for any $M > 0$. It follows from this claim that $N(\gamma, M) \subseteq I \setminus D(\gamma, CM)$ and so the condition (1.7) holds for any $c < 1$, since (4.3) then implies that

$$\mu(N(\gamma, M)) \leq \frac{C}{M}, \quad M \geq 1. \quad (4.12)$$

We now prove (4.11). Whenever $\psi \in D(\gamma, M)$, for any smooth function $u : S_{\psi} \rightarrow \mathbb{R}$, we have

$$\|u\|_{\dot{H}^{-\gamma}(S_{\psi})}^2 = \sum_{(m,n) \in \mathbb{Z}^2 \setminus 0} \frac{|\widehat{u}(m,n)|^2}{(m^2 + n^2)^{\gamma}} \leq M^2 \sum_{(m,n) \in \mathbb{Z}^2 \setminus 0} |m + \iota(\psi)n|^2 |\widehat{u}(m,n)|^2. \quad (4.13)$$

Now we note that by (4.10),

$$\widehat{(\nabla_{B/J} u)}(\psi, m, n) = -2\pi i(m + \iota(\psi)n)\widehat{u}(\psi, m, n). \quad (4.14)$$

It follows that if $\psi \in D(\gamma, M)$, there are constants C_1, C_2 so that

$$\|u\|_{\dot{H}^{-\gamma}(S_{\psi})}^2 \leq C_1 M^2 \|\nabla_{B/J} u\|_{L^2(S_{\psi})}^2 \leq C_2 M^2 \|\nabla_b u\|_{L^2(S_{\psi})}^2, \quad (4.15)$$

and this gives (4.11).

We now show how to get the same result for any non-vanishing divergence-free fibered field provided the rotational transform satisfies the bound (1.17). We first show that the rotational transform is well-defined in this setting; that is, that we can find coordinates so that B takes the form (1.16).

In light of (A.6), since $\operatorname{div} B = 0$, it follows that $\frac{1}{|\nabla\psi|}$ is an integral invariant of $B|_{S_{\psi}}$ (namely $U = \frac{1}{|\nabla\psi|}$ is a conserved density along the flow of $B|_{S_{\psi}}$ on S_{ψ}) and so by Sternberg’s theorem (Theorem 1 of [21]), B is orbitally conjugate to a linear operator on S_{ψ} . That is, there are coordinates (θ, ϕ) on S_{ψ} so that in these coordinates, there is a nonvanishing function $J = J(\theta, \phi)$ so that on S_{ψ} , B takes the form

$$(B \cdot \nabla^T)u = J(\partial_{\phi} + \iota\partial_{\theta})u, \quad u \in C^{\infty}(S_{\psi}) \quad (4.16)$$

for a real number ι (compare with (1.14)), where ∇^T denotes the tangential gradient on S_ψ , given by $\nabla^T u = (\nabla - \frac{\nabla\psi}{|\nabla\psi|^2} \nabla\psi \cdot \nabla)u$ when u is a function defined in a neighborhood of S_ψ . Applying this theorem on each S_ψ then gives $\iota = \iota(\psi)$.

The above proof goes through with a minor change, which is that we want to replace the fractional Sobolev norm $\|u\|_{\dot{H}^\gamma}$, which was defined relative to a fixed coordinate system (because the Fourier coefficients $\hat{u}(k)$ depend on the choice of coordinates in (2.8)), with a fractional Sobolev norm $\|u\|_{\dot{H}^\gamma}$ defined relative to the coordinates guaranteed by Sternberg's theorem. This means we want to modify the definition of $E(\gamma, M)$ and define

$$\tilde{E}(\gamma, M) = \left\{ \psi \in I : \|u\|_{\dot{H}^{-\gamma}(S_\psi)} \leq M \|\nabla_B u\|_{L^2(S_\psi)}, \text{ for all } u \in H^1(S_\psi) \right\}. \quad (4.17)$$

It is clear that the proof of Theorem 1.1 goes through without change if we replace the sets $E(\gamma, M)$ with $\tilde{E}(\gamma, M)$. It is also clear from the formula (4.16) and the above argument that there is a constant $C > 0$ so that (4.11) holds with E replaced with \tilde{E} , if the rotational transform $\iota = \iota(\psi)$ from (4.16) satisfies (1.17). \square

5. Proof of Theorem 1.2, Corollary 1.3 and Proposition 1.1: the non-integrable case

We now consider the non-integrable case $\chi_1 \neq 0$ of (1.19). With B as in (1.13)-(1.19), we set

$$B_0 = \nabla\psi \times \nabla\theta + \iota(\psi)\nabla\phi \times \nabla\psi, \quad B_1 = \nabla\phi \times \nabla\chi_1 \quad (5.1)$$

where $\iota(\psi) = \chi_0'(\psi)$. We then write $b = B/|B|$ in the form

$$b = b_0 + \varepsilon^a b_1, \quad b_0 = \frac{B_0}{|B|}, \quad b_1 = \frac{B_1}{|B|}. \quad (5.2)$$

5.1. Proof of Theorem 1.2. We start by recording a simple estimate.

LEMMA 5.1. *If $\|\nabla\chi_1\|_{L^\infty} < 1$ and $a \geq 1/2$, there is a constant $C > 0$ so that for any function $u \in H^1(D)$, we have*

$$\frac{1}{C} \left(\|\nabla_b u\|_{L^2}^2 + \varepsilon \|\nabla_b^\perp u\|_{L^2}^2 \right) \leq \|\nabla_{b_0} u\|_{L^2}^2 + \varepsilon \|\nabla_{b_0}^\perp u\|_{L^2}^2 \leq C \left(\|\nabla_b u\|_{L^2}^2 + \varepsilon \|\nabla_b^\perp u\|_{L^2}^2 \right). \quad (5.3)$$

PROOF. This follows after writing $\nabla_{b_0} = \nabla_b - \varepsilon^a \nabla_{b_1}$, noting that $|b_1| \leq C|\nabla\chi_1|$, and

$$|\nabla_{b_0} u| \leq |\nabla_b u| + \varepsilon^a |\nabla\chi_1| |\nabla u|, \quad |\nabla_{b_0}^\perp u| = |(\nabla - b_0 \nabla_{b_0})u| \leq |\nabla_b^\perp u| + \varepsilon^a |\nabla\chi_1| |\nabla u| \quad (5.4)$$

for smooth u . It follows that for $u \in H^1(D)$,

$$\|\nabla_{b_0} u\|_{L^2}^2 + \varepsilon \|\nabla_{b_0}^\perp u\|_{L^2}^2 \leq \|\nabla_b u\|_{L^2}^2 + \varepsilon \|\nabla_b^\perp u\|_{L^2}^2 + \varepsilon^{2a} \|\nabla\chi_1\|_{L^\infty}^2 \|\nabla u\|_{L^2}^2. \quad (5.5)$$

Provided $2a \geq 1$, this gives

$$(1 - \varepsilon \|\nabla\chi_1\|_{L^\infty}) \|\nabla_{b_0} u\|_{L^2}^2 + \varepsilon (1 - \|\nabla\chi_1\|_{L^\infty}^2) \|\nabla_{b_0}^\perp u\|_{L^2}^2 \leq \|\nabla_b u\|_{L^2}^2 + \varepsilon \|\nabla_b^\perp u\|_{L^2}^2, \quad (5.6)$$

which is the second bound in (5.3). The first bound is nearly identical. \square

PROOF OF COROLLARY 1.2. Since $\nabla_{b_0} T_0 = 0$ we have

$$\begin{aligned} \operatorname{div}(b \nabla_b T_0) + \varepsilon \operatorname{div}(\nabla_b^\perp T_0) &= \varepsilon^a \operatorname{div}(b \nabla_{b_1} T_0) + \varepsilon \Delta T_0 - \varepsilon \operatorname{div}(b \nabla_b T_0) \\ &= (\varepsilon^a - \varepsilon^{1+a}) \operatorname{div}(b \nabla_{b_1} T_0) + \varepsilon \operatorname{div}(\nabla T_0). \end{aligned} \quad (5.7)$$

Then $\rho = T_\varepsilon - T_0$ satisfies

$$\operatorname{div}(b \nabla_b \rho) + \varepsilon \operatorname{div}(\nabla_b^\perp \rho) = (\varepsilon^{1+a} - \varepsilon^a) \operatorname{div}(b \nabla_{b_1} T_0) - \varepsilon \operatorname{div}(\nabla T_0) \quad \text{in } D, \quad (5.8)$$

$$\rho|_{S_\pm} = 0. \quad (5.9)$$

Since $\partial_\theta \chi_1|_{\partial D} = 0$ by assumption, b is tangent to ∂D and so if we multiply this by ρ and integrate over D , we find

$$\int_D |\nabla_b \rho|^2 + \varepsilon |\nabla_b^\perp \rho|^2 d\mu = \varepsilon \int_D \Delta T_0 \rho d\mu + (\varepsilon^a - \varepsilon^{1+a}) \int_D \nabla_{b_1} T_0 \nabla_b \rho d\mu, \quad (5.10)$$

after integrating by parts in the second term on the right-hand side. Since $|\nabla_{b_1} T_0| \leq |\partial_\theta \chi_1| |T'_0|$, for any $\delta > 0$ we have

$$\|\nabla_b \rho\|_{L^2}^2 + \varepsilon \|\nabla_b^\perp \rho\|_{L^2}^2 \leq \frac{\varepsilon^{2a}}{2\delta} \|\partial_\theta \chi_1\|_{L^2}^2 \|T'_0\|_{L^\infty}^2 + \frac{\delta}{2} \|\nabla_b \rho\|_{L^2}^2 + \varepsilon \left| \int_{\psi_-}^{\psi_+} \int_{S_\psi} \frac{\operatorname{div} \nabla T_0}{|\nabla \psi|} \rho d\mathcal{H}^{(d-1)} d\psi \right|. \quad (5.11)$$

Arguing as in (2.24)-(2.26) and using (5.3), this implies that there is a constant $C > 0$ so that for any $M > 0$,

$$\begin{aligned} & \|\nabla_{b_0} \rho\|_{L^2}^2 + \varepsilon \|\nabla_{b_0}^\perp \rho\|_{L^2}^2 + \varepsilon \|\rho\|_{L^2}^2 \\ & \leq C \varepsilon^{2a} \|\partial_\theta \chi_1\|_{L^2}^2 \|T'_0\|_{L^2}^2 + C (M^2 \varepsilon^2 + \varepsilon \mu(N(\gamma, M))) (\|\Delta T_0\|_{H(0,\gamma)}^2 + \|\Delta T_0\|_{L^\infty}^2). \end{aligned} \quad (5.12)$$

If we take $M = \varepsilon^{-1/3}$ and use the estimate (4.12) for $\mu(N(\gamma, M))$, we find

$$\|\nabla_{b_0}^\perp \rho\|_{L^2} + \|\rho\|_{L^2}^2 \leq C \varepsilon^{2a-1} \|\partial_\theta \chi_1\|_{L^2}^2 + C \varepsilon^{\frac{1}{3}} (\|\Delta T_0\|_{H(0,\gamma)}^2 + \|\Delta T_0\|_{L^\infty}^2), \quad (5.13)$$

$$\|\nabla_{b_0} \rho\|_{L^2} \leq C \varepsilon^{2a} \|\partial_\theta \chi_1\|_{L^2}^2 + C \varepsilon^{\frac{4}{3}} (\|\Delta T_0\|_{H(0,\gamma)}^2 + \|\Delta T_0\|_{L^\infty}^2). \quad (5.14)$$

□

5.2. Proof of Corollary 1.3. Before proving Corollary 1.3, we collect some preliminary results. First, from the uniform bounds (1.21) and (1.22), it follows that the sequence $T_\varepsilon - T_0$ has weak limit ρ^* in H^1 and that $\nabla_b(T_\varepsilon - T_0)$ converges strongly to 0 in L^2 .

If we knew that $\nabla_B \rho^*$ was smooth, it would follow that $\nabla_B \rho^* = 0$ everywhere. We however only know that $\nabla_B \rho^*$ is in L^2 and in particular the restriction $\nabla_B \rho^*|_{S_\psi}$ need not be defined. The following result shows that $\nabla_B \rho^*|_{S_\psi} = 0$ in a weak sense.

LEMMA 5.2. *Let $\rho^* = \lim_{\varepsilon \rightarrow 0} T_\varepsilon - T_0$. For any ψ and any $v \in C^2(S_\psi)$, with $B' = B|\nabla \psi|^{-1}$,*

$$\int_{S_\psi} \rho^* \operatorname{div}_{S_\psi}(B'v) d\mathcal{H}^{(d-1)} = 0. \quad (5.15)$$

REMARK. The statement of (5.15) holds with B' replaced by B (or indeed any multiple of B) but it is more convenient for our purposes to use B' since $\operatorname{div}_{S_\psi}(B'v) = |\nabla \psi| \operatorname{div}(Bv) = |\nabla \psi| \nabla_B v$, by (A.6) and the fact that $\operatorname{div} B = 0$. See Lemma 5.3.

PROOF. The idea is to use Stokes' theorem to write (5.15) in terms of an integral in the interior of D and to integrate by parts in the interior to exploit the fact that $\nabla_B \rho^* = 0$ almost everywhere.

Fix a surface $S_{\psi'}$. Define a cutoff function $Q \in C^\infty(\mathbb{R})$ with $Q(z) = 1$ when $|z| \leq 1$ and $Q(z) = 0$ when $|z| > 2$. Define $Q_\delta(\psi) = Q((\psi - \psi')/\delta)$, so that $Q_\delta(\psi)$ vanishes when $|\psi - \psi'| < 2\delta$ and $Q_\delta(\psi) \equiv 1$ when $|\psi - \psi'| \leq \delta$. Let $D_-(\psi') = D \cap \{\psi \leq \psi'\}$. Then the boundary of $D_-(\psi')$ is $S_{\psi'} \cup S_{\psi_-}$, and if δ is sufficiently small, Q_δ vanishes identically on S_{ψ_-} .

Let V denote the constant extension of v to $D_-(\psi')$, $V(\psi, \theta, \phi) = v(\psi', \theta, \phi)$ for all $(\psi, \theta, \phi) \in D_-(\psi')$. By Stokes' theorem, since the outer unit normal to $S_{\psi'}$ is $n = \frac{\nabla \psi}{|\nabla \psi|}$, we have

$$\begin{aligned} \int_{S_{\psi'}} \rho^* \operatorname{div}_{S_{\psi'}}(B'u) d\mathcal{H}^{(d-1)} &= \int_{S_{\psi'}} (n \cdot n) \rho^* \operatorname{div}_{S_{\psi'}}(B'V) Q_\delta d\mathcal{H}^{(d-1)} \\ &= \int_{D_-} Q_\delta \operatorname{div} \left(\rho^* \operatorname{div}_{S_{\psi'}}(B'V) \frac{\nabla \psi}{|\nabla \psi|} \right) d\mu + \int_{D_-} \rho^* \operatorname{div}_{S_{\psi'}}(B'V) \frac{\nabla \psi}{|\nabla \psi|} \cdot (\nabla Q_\delta) d\mu. \end{aligned} \quad (5.16)$$

The first term here is bounded by

$$\|\operatorname{div}_{S_{\psi'}}(B'V)\|_{H^1(|\psi-\psi'|\leq\delta)}\|\rho^*\|_{H^1(|\psi-\psi'|\leq\delta)}\leq C\delta^{1/2}\|v\|_{C^2(S_{\psi'})}\|\rho^*\|_{H^1(D)}. \quad (5.17)$$

As for the second term, we use the identity (A.6) to write

$$\operatorname{div}_{S_{\psi}}(B'V)=\frac{1}{|\nabla\psi|}\operatorname{div}(BV), \quad (5.18)$$

and since $\frac{\nabla\psi}{|\nabla\psi|}\cdot(\nabla Q_\delta)=\frac{1}{\delta}|\nabla\psi|Q'_\delta$ and b is tangent to $S_{\psi'}$, the second term in (5.16) is

$$\frac{1}{\delta}\int_{D_-}\rho^*\operatorname{div}_{S_{\psi'}}(B'V)|\nabla\psi|Q'_\delta d\mu=\frac{1}{\delta}\int_{D_-}\rho^*\operatorname{div}(BV)Q'_\delta d\mu=-\frac{1}{\delta}\int_{D_-}\nabla_B(\rho^*Q'_\delta)V d\mu, \quad (5.19)$$

after integrating by parts. We have therefore shown that for any $\delta > 0$,

$$\begin{aligned} \left|\int_{S_{\psi'}}\rho^*\operatorname{div}_{S_{\psi}}(B'V)d\mathcal{H}^{(d-1)}\right| &\leq C\delta^{1/2}\|v\|_{C^2(S_{\psi'})}\|\rho^*\|_{H^1(D)}+\frac{1}{\delta}C(\|\nabla_B\rho^*\|_{L^2}+\|\nabla_BQ'_\delta\|_{L^2})\|V\|_{L^2} \\ &= C\delta^{1/2}\|v\|_{C^2(S_{\psi'})}\|\rho^*\|_{H^1(D)}, \end{aligned} \quad (5.20)$$

where we used that $\nabla_B u = 0$ whenever $u = u(\psi)$ and that $\nabla_B \rho^* = 0$ in L^2 . Taking $\delta \rightarrow 0$ gives the claim. \square

The condition (5.15) nearly says that $\rho^* = 0$ on each S_ψ in the sense of distributions, but in order to conclude this we would need to know that any test function v can be written in the form $v = \operatorname{div}_{S_\psi}(Bw)$ for some test function w . This need not be possible on an arbitrary surface S_ψ , but by the next Lemma it is possible provided S_ψ is a Diophantine surface. We set

$$D(\gamma)=\bigcup_{M>0}D(\gamma,M). \quad (5.21)$$

Note that by (4.11), $D(\gamma) \subseteq E(\gamma)$ where $E(\gamma) = \cup_{M>0}E(\gamma, M)$ denotes the collection of all ergodic values of ψ . Note also that by Lemma 4.1, the complement $I \setminus D(\gamma)$ has zero measure when $\gamma > 2$.

LEMMA 5.3. *Fix $\gamma > 2$ and define $D(\gamma)$ as in (5.21). If $\psi \in D(\gamma)$ and $v \in H^{s+\gamma}(S_\psi)$ for some $s \geq 0$, there is $w \in H^s(S_\psi)$ satisfying*

$$\operatorname{div}_{S_\psi}(B'w)=v, \quad B'=B|\nabla\psi|^{-1}. \quad (5.22)$$

REMARK. The equation (5.22) is sometimes known as the ‘‘magnetic differential equation’’. Lemma 5.3 simply says that you can solve this equation on good (sufficiently ergodic) flux surfaces.

PROOF. We start by using (A.6) to write

$$\operatorname{div}_{S_\psi}(B'w)=\frac{1}{|\nabla\psi|}\operatorname{div}(Bw)=\frac{1}{|\nabla\psi|}\nabla_Bw, \quad (5.23)$$

and so (5.22) takes the form

$$\nabla_{B/J}w=V, \quad V=J|\nabla\psi|v, \quad (5.24)$$

where recall $J=|g|^{-1/2}=|(\nabla\psi\times\nabla\theta)\cdot\nabla\phi|$. We now define

$$\widehat{w}(m,n)=\frac{i}{2\pi}\frac{1}{m+i(\psi)n}\widehat{V}(m,n). \quad (5.25)$$

Because $\psi \in D(\gamma)$, for some $M > 0$, we have the bound

$$|\widehat{w}(m,n)|^2\leq M^2(m^2+n^2)^\gamma|\widehat{V}(m,n)|^2, \quad (5.26)$$

so that in particular $w \in H^s(S_\psi)$ whenever $v = V/(J|\nabla\psi|) \in H^{s+\gamma}(S_\psi)$. By the identity (4.14), it follows that w satisfies (5.22). \square

PROOF OF COROLLARY 1.3. Fix $\gamma > 2$ and define $D(\gamma)$ as in (5.21). It follows from Lemma 5.3 that given $v \in C^\infty(S_\psi)$, there is $w \in C^\infty(S_\psi)$ so that with $\operatorname{div}_{S_\psi}(B|\nabla\psi|w) = v$. It then follows from Lemma 5.15 that

$$\int_{S_\psi} \rho^* v \, d\mathcal{H}^{(d-1)} = \int_{S_\psi} \rho^* \operatorname{div}_{S_\psi}(B|\nabla\psi|w) \, d\mathcal{H}^{(d-1)} = 0, \quad (5.27)$$

as required. \square

5.3. Proof of Proposition 1.1: effective volume of non-integrability. It follows immediately from the equation (1.9) (or, equivalently, (2.6)) for $T_0 = \Theta(\psi)$ that either T_0 is constant or else T'_0 is nonvanishing. Since $T_0|_{S_+} = T_+ \neq T_- = T_0|_{S_-}$ we have that T'_0 is nonvanishing and in particular T'_0 is bounded below. With $\lambda = \min_D \frac{1}{|T'_0|}$, for any set $N \subseteq D$ we therefore have

$$\mu(N) \leq \lambda^2 \int_N |T'_0|^2 \, d\mu \leq C \left(\lambda^2 \int_N |\nabla_b^\perp T|^2 \, d\mu + \lambda^2 \int_N |\nabla_b^\perp \rho|^2 \, d\mu \right), \quad (5.28)$$

where we used that $T'_0 = |\nabla\psi|^{-2} \nabla T_0 \cdot \nabla\psi$ and that $|\nabla T_0| \leq |\nabla_b^\perp T| \leq |\nabla_b^\perp T| + |\nabla_b^\perp \rho|^2$. In particular, with $N = N(\varepsilon)$, since $\|\nabla_b^\perp T\|_{L^2(N(\varepsilon))}^2 \leq \varepsilon^{-1} \|\nabla_b T\|_{L^2(N(\varepsilon))}^2$,

$$\begin{aligned} \mu(N(\varepsilon)) &\leq C\lambda^2 \left(\varepsilon^{-1} \int_{N(\varepsilon)} |\nabla_b T|^2 \, d\mu + \int_{N(\varepsilon)} |\nabla_b^\perp \rho|^2 \, d\mu \right) \\ &\leq C\lambda^2 \left(\varepsilon^{-1} \int_{N(\varepsilon)} |\nabla_b T_0|^2 \, d\mu + \varepsilon^{-1} \int_D |\nabla_b \rho|^2 \, d\mu + \int_D |\nabla_b^\perp \rho|^2 \, d\mu \right) \\ &\leq C\lambda^2 \varepsilon^{-1} \left(\int_{N(\varepsilon)} |\nabla_b T_0|^2 \, d\mu + \int_D |\nabla_b \rho|^2 \, d\mu + \varepsilon \int_D |\nabla_b^\perp \rho|^2 \, d\mu \right). \end{aligned} \quad (5.29)$$

Since $b = b_0 + \varepsilon^a b_1$ and $\nabla_{b_0} T_0 = 0$, we have

$$\|\nabla_b T_0\|_{L^2(D)}^2 \leq \varepsilon^{2a} \|\nabla_{b_1} T_0\|_{L^2(D)}^2 \leq \varepsilon^{2a} \|\partial_\theta \chi_1\|_{L^2(D)}^2 \|T'_0\|_{L^\infty(D)}^2. \quad (5.30)$$

To handle the second and third terms in (5.29), we use (1.21)-(1.22) combined with (5.3). By (5.30) we therefore have

$$\mu(N(\varepsilon)) \leq C\lambda^2 \left(\varepsilon^{2a-1} \|\partial_\theta \chi_1\|_{L^2(D)}^2 \|T'_0\|_{L^\infty(D)}^2 + \varepsilon^{1/3} \|\Delta T_0\|_{H^{(0,\gamma)}(D)}^2 \right). \quad (5.31)$$

Appendix A. Geometric identities from the co-area formula

In this section we collect some geometric formulas that we will use repeatedly. In what follows, we fix $\psi : D \rightarrow \mathbb{R}$ such that $|\nabla\psi| \neq 0$ on D and so that the level surfaces S_ψ are codimension one manifolds which foliate D . Let $\psi_- = \inf_D \psi$ and $\psi_+ = \sup_D \psi$. We will use the co-area formula

$$\int_D u \, d\mu = \int_{\psi_-}^{\psi_+} \int_{S_\psi} \frac{u}{|\nabla\psi|} \, d\mathcal{H}^{(d-1)} \, d\psi, \quad (A.1)$$

see e.g. [22]. We start with a simple result that generalizes Lemma E.2 from [8].

LEMMA A.1. *If $F \in H^2(D)$, we have*

$$\frac{d}{d\psi} \int_{S_\psi} F \, d\mathcal{H}^{(d-1)} = \int_{S_\psi} \operatorname{div} \left(\frac{\nabla\psi}{|\nabla\psi|} F \right) \, d\mathcal{H}^{(d-1)}. \quad (A.2)$$

PROOF. We start with the observation that

$$\int_{S_{\psi_2}} F \, d\mathcal{H}^{(d-1)} - \int_{S_{\psi_1}} F \, d\mathcal{H}^{(d-1)} = \int_{D_{\psi_1, \psi_2}} \operatorname{div} \left(\frac{\nabla\psi}{|\nabla\psi|} F \right) \, d\mu, \quad (A.3)$$

where $D_{\psi_1, \psi_2} = \cup_{\psi_1 \leq \psi' \leq \psi_2} S_{\psi'}$ denotes the region bounded by the surfaces S_{ψ_1}, S_{ψ_2} . Indeed, by the divergence theorem,

$$\int_{D_{\psi_1, \psi_2}} \operatorname{div} \left(\frac{\nabla \psi}{|\nabla \psi|} F \right) d\mu = \int_{S_{\psi_2}} n^{S_{\psi_2}} \cdot \frac{\nabla \psi}{|\nabla \psi|} F d\mathcal{H}^{(d-1)} + \int_{S_{\psi_1}} n^{S_{\psi_1}} \cdot \frac{\nabla \psi}{|\nabla \psi|} F d\mathcal{H}^{(d-1)}, \quad (\text{A.4})$$

where n^{S_ψ} denotes the outward-pointing unit normal to S_ψ . Then $n^{S_{\psi_2}} = \frac{\nabla \psi}{|\nabla \psi|}|_{S_{\psi_2}}$ and $n^{S_{\psi_1}} = -\frac{\nabla \psi}{|\nabla \psi|}|_{S_{\psi_1}}$, so (A.4) gives (A.3). Dividing (A.3) by $\psi_2 - \psi_1$ and taking the limit gives (A.2). \square

In particular, if $F = F(\psi)$ is constant on S_ψ , writing $\Delta F = \operatorname{div}(\nabla F) = \operatorname{div}(\nabla \psi F')$ we have

$$\int_{S_\psi} \Delta F d\mathcal{H}^{(d-1)} = \frac{d}{d\psi} \left(\int_{S_\psi} |\nabla \psi| F' d\mathcal{H}^{(d-1)} \right) = \frac{d}{d\psi} \left(\left[\int_{S_\psi} |\nabla \psi| d\mathcal{H}^{(d-1)} \right] F' \right). \quad (\text{A.5})$$

Another consequence of the formula (A.1) is the following

LEMMA A.2. *Let X be a vector field defined in D with the property that $X|_{S_\psi}$ is tangent to S_ψ . Then the divergence $\operatorname{div} X$ in D is related to the divergence operator $\operatorname{div}_{S_\psi}$ on S_ψ by*

$$\frac{\operatorname{div} X}{|\nabla \psi|} \Big|_{S_\psi} = \operatorname{div}_{S_\psi} \left(\left[\frac{X}{|\nabla \psi|} \right] \Big|_{S_\psi} \right). \quad (\text{A.6})$$

In particular, if X is divergence-free in D , then $\varrho := \frac{1}{|\nabla \psi|} \Big|_{S_\psi}$ is a density conserved by X on S_ψ .

PROOF. This can be seen by working in local coordinates but it is simpler to use (A.1) and note that if $u \in C^\infty(D)$ is any test function then

$$\begin{aligned} - \int_D \operatorname{div} X u d\mu &= \int_D (X \cdot \nabla) u d\mu = \int_{\psi_-}^{\psi_+} \int_{S_\psi} (X \cdot \nabla) u \frac{d\mathcal{H}^{(d-1)}}{|\nabla \psi|} \\ &= \int_{\psi_-}^{\psi_+} \int_{S_\psi} (X \cdot \nabla^T) u \frac{d\mathcal{H}^{(d-1)}}{|\nabla \psi|} = - \int_{\psi_-}^{\psi_+} \int_{S_\psi} \operatorname{div}_{S_\psi} \left(\frac{X}{|\nabla \psi|} \right) u d\mathcal{H}^{(d-1)}, \end{aligned}$$

where we used that $X \cdot \nabla = X \cdot \nabla^T$ on S_ψ , where ∇^T denotes the tangential gradient on S_ψ , $\nabla^T U = (\nabla - \frac{\nabla \psi}{|\nabla \psi|^2} \nabla \psi \cdot \nabla) u$ whenever u is an extension of U from S_ψ to a neighborhood of S_ψ . By (A.1), the left-hand side is $-\int_{\psi_-}^{\psi_+} \int_{S_\psi} \frac{\operatorname{div} X}{|\nabla \psi|} u d\mathcal{H}^{(d-1)}$. Then (A.6) follows since u is arbitrary. \square

Acknowledgements. We thank P. Constantin, P. Helander, S. Hudson, E. Paul, and P. Torres de Lizaar for numerous insightful discussions. The research of TD was partially supported by NSF-DMS grant 2106233. The work of DG was partially supported by the Simons Center for Hidden Symmetries and Fusion Energy award # 601960. The studies of HG are supported by Ford Foundation and the NSF Graduate Research Fellowship under grant DGE-2039656.

References

- [1] Arnold, V. I. (1966). On the topology of three-dimensional steady flows of an ideal fluid. In Vladimir I. Arnold- Collected Works (pp. 25-28). Springer, Berlin, Heidelberg.
- [2] Arnold, V. I., and Khesin, B. A. (2021). Topological methods in hydrodynamics (Vol. 125). Springer Nature.
- [3] Braginskii, S. I. (1965). Transport Processes in Plasma, ed. MA Leontovich (New York, USA: Consultants Bureau), 201.
- [4] Hudson, S. R., and Breslau, J. (2008). Temperature contours and ghost surfaces for chaotic magnetic fields. Physical review letters, 100(9), 095001.
- [5] Helander, P., Hudson, S. R., and Paul, E. J. (2022). On heat conduction in an irregular magnetic field. Part 1. Journal of Plasma Physics, 88(1).
- [6] Paul, E. J., Hudson, S. R., and Helander, P. (2022). Heat conduction in an irregular magnetic field. Part 2. Heat transport as a measure of the effective non-integrable volume. Journal of Plasma Physics, 88(1).
- [7] Braginskii, S.I., 1965. Transport processes in a plasma. Reviews of plasma physics, 1.

- [8] Constantin, P., Drivas, T. D., and Ginsberg, D. (2021). Flexibility and rigidity in steady fluid motion. *Communications in Mathematical Physics*, 385(1), 521-563.
- [9] Constantin, P., Drivas, T. D., and Ginsberg, D. (2021). On quasisymmetric plasma equilibria sustained by small force. *Journal of Plasma Physics*, 87(1).
- [10] Constantin, P., Drivas, T. D., and Ginsberg, D. (2022). Flexibility and rigidity of free boundary MHD equilibria. *Nonlinearity*, 35(5), 2363.
- [11] Freidberg, J. P. 1987. "Ideal magnetohydrodynamics". United States.
- [12] Helander, P. and Sigmar, D., *Collisional Transport in Magnetized Plasmas*. United Kingdom: Cambridge University Press, 2005.
- [13] Helander, P. Theory of plasma confinement in non-axisymmetric magnetic fields, *Reports on Progress in Physics*.
- [14] Grad, H., Toroidal containment of a plasma. *The Physics of Fluids*, 10(1), 137-154, (1967)
- [15] Grad, H., Theory and applications of the nonexistence of simple toroidal plasma equilibrium. *International Journal of Fusion Energy* 3.2, 33-46, (1985)
- [16] L. Ryzhik, A. Novikov, G. Papanicolaou, Boundary layers for cellular flows at high Peclet numbers, *Communications in Pure and Applied Mathematics*, 58, 2005, 867-922.
- [17] Bensoussan, A., Lions, J. L., & Papanicolaou, G. (2011). *Asymptotic analysis for periodic structures* (Vol. 374). American Mathematical Soc.
- [18] Treschev, D., Zubelevich, O. (2009). *Introduction to the Perturbation Theory of Hamiltonian Systems*. Germany: Springer Berlin Heidelberg.
- [19] Arnold, Vladimir I. "Proof of a theorem of AN Kolmogorov on the invariance of quasi-periodic motions under small perturbations of the Hamiltonian." *Russian Mathematical Surveys* 18.5 (1963): 9.
- [20] Fitzpatrick, R. Helical temperature perturbations associated with tearing modes in tokamak plasmas *Cite as: Physics of Plasmas* 2, 825 (1995);
- [21] Sternberg, S., *American Journal of Mathematics* , Apr., 1957, Vol. 79, No. 2 (Apr., 1957), pp. 397-402
- [22] Gariepy, R. F., Evans, L. C. (2015). *Measure Theory and Fine Properties of Functions*, Revised Edition. United Kingdom: CRC Press.

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY, 11794

Email address: `tdrivas@math.stonybrook.edu`

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544

Email address: `dg42@princeton.edu`

PROGRAM IN APPLIED AND COMPUTATIONAL MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544

Email address: `hgrayer@math.princeton.edu`