

ABELIAN SUBGROUPS OF HELLY GROUPS

MOTIEJUS VALIUNAS

ABSTRACT. We observe that abelian subgroups of Helly groups are finitely generated, and consequently, soluble subgroups of Helly groups are virtually abelian.

A simplicial graph Γ is called *Helly* if for any collection $\{B_i \mid i \in I\}$ of balls in Γ , if $B_i \cap B_j \neq \emptyset$ for all $i, j \in I$ then $\bigcap_{i \in I} B_i \neq \emptyset$. We say a group G is *Helly* if it acts geometrically on a Helly graph.

The class of Helly groups has been introduced and studied in [CCG⁺20]. It is a large class of groups, including all hyperbolic, cocompactly cubulated, and $C(4)$ – $T(4)$ small cancellation groups [CCG⁺20, Theorem 1.1]. On the other hand, all Helly groups are biautomatic and have some other non-positive-curvature-like behaviour [CCG⁺20, Theorem 1.5].

Motivated by this, we may ask whether all abelian subgroups of Helly groups are finitely generated [CCG⁺20, Question 9.3], which is a special case of the long-standing open question on whether all abelian subgroups of biautomatic groups are finitely generated [GS91, §7]. In this note, we observe that the results of Descombes & Lang [DL16] can be combined with those of Haettel & Osajda [HO21] to give an affirmative answer to the former question. We expect this result to be well-known to experts, but it does not seem to appear anywhere in the literature.

Theorem. *Abelian subgroups of Helly groups are finitely generated.*

Before we prove the Theorem, let us note one immediate consequence.

Corollary. *Soluble subgroups of Helly groups are virtually abelian.*

Proof. Let G be a Helly group. Since, by the Theorem, abelian subgroups of G are finitely generated, it follows that soluble subgroups of G are polycyclic [Seg83, Theorem 2 on p. 25]. On the other hand, as G is biautomatic [CCG⁺20, Theorem 1.5(1)], polycyclic subgroups of G are virtually abelian [GS91, Theorem 6.15]. \square

Proof of the Theorem. Let G be a Helly group, so that G acts geometrically on a Helly graph Γ . By [CCG⁺20, Theorem 6.3], G also acts geometrically on the *injective hull* $E\Gamma$ of Γ —a locally finite polyhedral complex of dimension $N < \infty$.

Let $H \leq G$ be an abelian subgroup, and let $n = \dim_{\mathbb{Q}}(H \otimes \mathbb{Q}) \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$ be the torsion-free rank of H . Then H (and so G) has a subgroup isomorphic to \mathbb{Z}^k for any non-negative integer $k \leq n$. On the other hand, $E\Gamma$ admits a G -equivariant consistent bicombing [Lan13, Proposition 3.8], implying by the Flat Torus Theorem [DL16, Theorem 1.2] that $E\Gamma$ contains an isometrically embedded k -dimensional normed real vector space, and in particular that $k \leq N$, whenever G contains a free abelian subgroup of rank k . Therefore, we have $n \leq N$, and in particular $n < \infty$.

Now consider the translation length function on H with respect to the action on Γ , defined by $\|h\| := \lim_{m \rightarrow \infty} \frac{d_{\Gamma}(x, h^m \cdot x)}{m}$ for any $h \in H$, where $x \in \Gamma$ is a choice of a basepoint. It is clear that $\|h^m\| = |m| \cdot \|h\|$ and (since H is abelian) $\|gh\| \leq \|g\| + \|h\|$ for any $g, h \in H$ and $m \in \mathbb{Z}$, implying that $\|\cdot\|$ induces a semi-norm on $H \otimes \mathbb{Q} \cong \mathbb{Q}^n$. On the other hand, for any infinite order element $h \in H$, the translation length $\|h\|$ is non-zero and rational with denominator bounded above by $2N$ [HO21, Theorem O], implying that $\|h\| \geq \frac{1}{2N}$. Therefore, the image of the map $p: H \rightarrow H \otimes \mathbb{Q}$ is a discrete subgroup of $H \otimes \mathbb{Q} \cong \mathbb{Q}^n$, and therefore isomorphic to a lattice in \mathbb{R}^n , implying that $p(H) \cong \mathbb{Z}^n$.

Finally, $\ker(p)$ is a torsion subgroup of the Helly group G , implying by [HO21, Corollary I] that $\ker(p)$ is finite. This shows that H is finitely generated, as required. \square

Acknowledgement. *The author would like to thank Damian Osajda for a valuable discussion.*

REFERENCES

- [CCG⁺20] J. Chalopin, V. Chepoi, A. Genevois, H. Hirai, and D. Osajda, *Helly groups*, preprint, available at arXiv:2002.06895 [math.GR], 2020.
- [DL16] D. Descombes and U. Lang, *Flats in spaces with convex geodesic bicomings*, *Anal. Geom. Metr. Spaces* **4** (2016), 68–84.
- [GS91] S. M. Gersten and H. B. Short, *Rational subgroups of biautomatic groups*, *Ann. Math.* **134** (1991), no. 1, 125–158.
- [HO21] T. Haettel and D. Osajda, *Locally elliptic actions, torsion groups, and nonpositively curved spaces*, preprint, available at arXiv:2110.12431 [math.GR], 2021.
- [Lan13] U. Lang, *Injective hulls of certain discrete metric spaces and groups*, *J. Topol. Anal.* **5** (2013), no. 3, 297–331.
- [Seg83] D. Segal, *Polycyclic groups*, *Cambridge Tracts in Mathematics*, vol. 82, Cambridge University Press, 1983.

INSTYTUT MATEMATYCZNY, UNIwersYTET WROCLAWSKI, PLAC GRUNWALDZKI 2/4, 50-384 WROCLAW,
POLAND

Email address: Motiejus.Valiunas@math.uni.wroc.pl