

Extraction of ωn , ωp and ϕN scattering lengths from ω and ϕ differential photoproduction cross sections on the deuterium target

Chengdong Han,^{1,2,*} Wei Kou,^{1,2,†} Rong Wang,^{1,2,‡} and Xurong Chen^{1,2,3,§}

¹*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

²*University of Chinese Academy of Sciences, Beijing 100049, China*

³*Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China*

(Dated: November 1, 2022)

In this study, we try to extract ωn , ωp and ϕN scattering lengths from the differential cross-section data of near-threshold ω and ϕ photoproductions not only to the energy at threshold t_{thr} , but also in t to $t = 0$. The incoherent data of ω and ϕ photoproductions on deuterium target for scattering length extractions are provided by CBELSA/TAPS Collaboration and LEPS Collaboration respectively, where the deuteron is usually approximated as a weakly bound state of a quasi-free neutron plus a quasi-free proton. Under the Vector Meson Dominance model and the assumption of energy independence of the differential cross-section, we obtained the $|\alpha_{\omega n^*}|$ and $|\alpha_{\omega p^*}|$ of loosely bound neutron and proton in the deuteron to be 0.709 ± 0.078 fm and 0.621 ± 0.022 fm respectively at threshold t_{thr} (1.258 ± 0.130 fm and 1.056 ± 0.032 fm respectively in t to $t = 0$) for the first time by fitting the near-threshold data of $\gamma d \rightarrow \omega n(p)$ and $\gamma d \rightarrow \omega p(n)$. For a comparison study, we also extracted the scattering length $|\alpha_{\omega p}|$ of the free proton from the ω photoproduction on the hydrogen target by CBELSA/TAPS Collaboration. With the near-threshold and incoherent ϕ photoproduction data of $\gamma d \rightarrow \phi pn$ by LEPS Collaboration, we obtained scattering length $|\alpha_{\phi N}|$ of the bound nucleon inside the deuteron to be 0.109 ± 0.008 fm at threshold t_{thr} and 0.276 ± 0.010 fm in t to $t = 0$, respectively. The obtained $|\alpha_{\omega p}|$ and $|\alpha_{\phi N}|$ are basically in agreement with the experimental indications and the theoretical predictions within the current uncertainties.

I. INTRODUCTION

Vector meson-nucleon interactions are the extremely important research topics in the non-perturbation domain of quantum chromodynamics (QCD). Since the discovery of the vector mesons, they have become attractive probes for the study of hadronic interaction. Due to the behavior of the near threshold cross section is related to the vector meson-nucleon scattering length [1], experimentally, the vector meson-nucleon interaction can be investigated using vector meson photoproduction within the Vector Meson Dominance (VMD) model. The ωp , ϕp , $J/\psi - p$, $\psi(2S)p$ and $\rho^0 p$ scattering lengths have been fully analysed in Refs. [2–7]. Based on the VMD model, the absolute value of the scattering length may be determined from the total near-threshold vector-meson photoproduction cross-section [1] or the differential cross-section of vector-meson photoproduction at threshold [8].

We try to extract the scattering length of the vector meson-neutron and vector meson-proton with current available experimental data. In Ref. [3], they estimate the $J/\psi - p$ scattering length using the VMD model with the new GlueX data to the threshold. Within the VMD model, the total $\gamma N \rightarrow XN$ cross-section is related to both the total $XN \rightarrow XN$ cross-section at threshold en-

ergy and the scattering length $|\alpha_{XN}|$ by [8]:

$$\sigma^{\gamma N}(s_{thr}) = \frac{\alpha\pi}{\gamma_X^2} \frac{q_{XN}}{k_{\gamma N}} \cdot \sigma^{XN}(s_{thr}) = \frac{\alpha\pi}{\gamma_X^2} \frac{q_{XN}}{k_{\gamma N}} \cdot 4\pi\alpha_{XN}^2, \quad (1)$$

where $\alpha = 1/137$ denotes the fine structure constant, and X is a index which represents the vector mesons (e.g. ω , ϕ , J/ψ , etc.) The $k_{\gamma N}$ and q_{XN} in the above equation are the momenta in the center-of-mass of the initial and final state particles, respectively, and γ_X is the photon-vector meson coupling constant obtained from the $X \rightarrow e^+e^-$ decay width. The photon- ω and photon- ϕ coupling constants used in this work are 8.53 and 6.71 [9], respectively. Eq. (1) is taken at the threshold energy, where $s_{thr} = (M + m)^2$ with M and m being the masses of the vector meson and nucleon, respectively.

To estimate the scattering length with the experimental measurements of the differential photoproduction cross-section, in this study, we adopted the relation between the total cross-section and the differential cross-section at the threshold [5]. The relationship between the scattering length and the differential cross-section is expressed as:

$$\begin{aligned} \frac{d\sigma^{\gamma X}}{dt}(s_{thr}, t = t_{thr}) &= \frac{\alpha\pi}{\gamma_X^2} \frac{\pi}{k_{\gamma N}^2} \cdot \alpha_{XN}^2, \\ \frac{d\sigma^{\gamma X}}{dt}(s_{thr}, t = 0) &= \frac{\alpha\pi}{\gamma_X^2} \frac{\pi}{k_{\gamma N}^2} \cdot \alpha_{XN}^2. \end{aligned} \quad (2)$$

A key problem in determining the scattering length at threshold t_{thr} is to extrapolate the cross section to the point of $t \rightarrow t_{thr}$ or $s \rightarrow s_{thr}$. In addition, the

* chdhan@impcas.ac.cn

† kouwei@impcas.ac.cn

‡ rwang@impcas.ac.cn

§ xchen@impcas.ac.cn (Corresponding author)

$d\sigma^{\gamma X}/dt(s_{thr}, t=0)$ at left-hand side of Eq. (2) is not a directly measurable quantity, as it requires extrapolation of the energy to the threshold and extrapolation of t from the physical region ($t_{min} < t < t_{max}$) to the non-physical point $t=0$. Therefore, when the vector meson nucleon scattering length is extracted from the differential cross-section data, we can not only extrapolate the energy at the threshold t_{thr} , but also extrapolate t to $t=0$.

In this work, the following exponential function was used to fit the differential cross-section data of near-threshold ω and ϕ photoproductions on deuterium target or hydrogen target.

$$\frac{d\sigma}{dt} = Ae^{-bt}, \quad (3)$$

where $A = d\sigma/dt|_{t=0}$ denotes the forward differential cross-section and b describes the slope parameter. Combining Eq. (2) and Eq. (3), we can obtain the forward differential cross-section value $d\sigma/dt|_{t=t_{thr}}$ and $d\sigma/dt|_{t=0}$ and extract the vector meson-nucleon scattering length $|\alpha_{XN}|$.

Great progress has been made in meson-proton scattering length measurements, but little is known about the vector meson-neutron scattering length. As the vector meson-proton scattering length has already been fixed by the data of ω , ϕ , J/ψ , $\psi(2S)$ and ρ^0 vector meson photoproductions near threshold in the previous analyses [3, 5–8, 10], We want to try whether we can extract the vector meson-neutron scattering length using the deuterium target data that the neutron and the proton are loosely bound. The differential cross-section data of near-threshold ω and ϕ photoproductions on deuterium target were used for this analysis, including the incoherent photoproduction of vector meson (ω and ϕ vector mesons) [11, 12] where the neutron or the proton is knocked out. The incoherent data of ω and ϕ photoproductions for scattering length extraction are provided by CBELSA/TAPS Collaboration and LEPS Collaboration, respectively. In this work, we extract ωn , ωp and ϕN scattering length from the differential cross-section data of near-threshold ω and ϕ photoproductions on deuterium target, where the deuteron is usually approximated as a weakly bound state of a quasi-free neutron plus a quasi-free proton. For comparison study, we also extracted the scattering length $|\alpha_{\omega p}|$ of the free proton from the ω photoproduction on the hydrogen target by CBELSA/TAPS Collaboration.

II. ωn , ωp AND ϕN SCATTERING LENGTHS FROM ω AND ϕ DIFFERENTIAL PHOTOPRODUCTION CROSS SECTIONS

A. ωn^* scattering length from ω differential photoproduction cross sections on the deuterium target

Figure 1 shows the differential cross sections of the ω meson photoproductions produced off the bound neutron




FIG. 1. Differential cross sections of the near threshold photoproduction of ω meson produced off the quasi-free neutron versus the momentum transfer $-t$ in the deuterium target [11]. The three incident photon energies E_γ (1.175, 1.215 and 1.240 GeV) near the threshold of ω mesons produced off the quasi-free neutron are marked in the figure. Some cross sections are scaled using the coefficients shown in the figure.

as a function of momentum transfer $-t$, with three incident photon energies E_γ . This reaction with exactly four neutral hits ($\omega \rightarrow \gamma\gamma\gamma$ and neutron) of the experiment at ELSA [11] is detected where the ω meson was produced off the bound neutron in the liquid deuterium target giving the quasi-free reaction $\gamma d \rightarrow \omega n(p)$. The differential cross sections of the $|t|$ -dependence are fitted with an exponential function. We determined the parameters A and the b at three incident photon energies E_γ (1.175, 1.215 and 1.240 GeV) through fitting to the ω mesons produced off the bound neutron with deuterium target data [11] with Eq. (3). Then the values of $d\sigma^{\gamma n^*}/dt(s_{thr}, t_{thr})$ and $d\sigma^{\gamma n^*}/dt(s_{thr}, 0)$ are calculated by fitting parameters A and b obtained by fitting the $|t|$ -dependent differential photoproduction cross-section data. The scattering length $|\alpha_{\omega n^*}|$ at the threshold t_{thr} and in t to $t=0$ are then calculated by Eq. (2). The extracted scattering length $|\alpha_{\omega n^*}|$ are listed in Table I. The averaged scattering length $|\alpha_{\omega n^*}|$ for the three extracted values at different E_γ energies at threshold is calculated to be 0.709 ± 0.078 fm. The averaged scattering length $|\alpha_{\omega n^*}|$ for the three extracted values at different E_γ energies in t to $t=0$ is calculated to be 1.258 ± 0.130 fm. Here we use the following formula to calculate the weighted average: $\bar{x} \pm \delta \bar{x} = \sum_i \omega_i x_i / \sum_i \omega_i \pm (\sum_i \omega_i)^{-1/2}$ with $\omega_i = 1/(\delta x_i)^2$ [13]. It is worth noting that the weighted average values of scattering length at different energies obtained here are consistent with the simultaneous fitting results of all data sets.

B. ωp^* scattering length from ω differential photoproduction cross sections on the deuterium target




FIG. 2. Differential cross sections of the photoproduction of ω meson produced off the quasi-free proton versus the momentum transfer $-t$ in deuterium near the thresholds [11]. The three incident photon energies E_γ (1.175, 1.215 and 1.240 GeV) near the threshold of ω meson produced off the quasi-free proton are marked in the figure. Some cross sections are scaled using the coefficients shown in the figure.

Figure 2 shows the differential cross sections of the ω meson photoproductions produced off the bound proton as a function of momentum transfer $-t$. This reaction with exactly three neutral hits ($\omega \rightarrow \gamma\gamma\gamma$) and one charged hit (proton) of the experiment at ELSA [11] is detected where the ω meson was produced off the bound proton in the liquid deuterium target giving the quasi-free reaction $\gamma d \rightarrow \omega p(n)$. The differential cross sections of the $|t|$ -dependence are fitted with an exponential function. We determined the parameters A and the b at three incident photon energies E_γ (1.175, 1.215 and 1.240 GeV) by fitting the ω mesons produced off the bound proton on the deuterium target data [11] with Eq. (3). The values of $d\sigma^{\gamma p^*}/dt(s_{thr}, t_{thr})$ and $d\sigma^{\gamma p^*}/dt(s_{thr}, 0)$ are calculated by fitting parameters A and b obtained by fitting the $|t|$ -dependent differential photoproduction cross-section data. The scattering length $|\alpha_{\omega p^*}|$ at the threshold t_{thr} and in t to $t=0$ are then calculated by Eq. (2). The extracted scattering length $|\alpha_{\omega p^*}|$ are listed in Table II. The averaged scattering length $|\alpha_{\omega p^*}|$ for the three extracted values at different E_γ energies at threshold is calculated to be 0.621 ± 0.022 fm. The averaged scattering length $|\alpha_{\omega p^*}|$ for the three extracted values at different E_γ energies in t to $t=0$ is calculated to be 1.056 ± 0.032 fm.

C. ωp scattering length from ω differential photoproduction cross sections on the hydrogen target




FIG. 3. Differential cross sections of the photoproductions of ω meson produced off the free proton near the threshold versus the momentum transfer $-t$ in the hydrogen target [11]. The four incident photon energies E_γ (1.138, 1.163, 1.188 and 1.225 GeV) near the threshold of ω meson are marked in the figure. The exclusive experimental data results by CBELSA/TAPS Collaboration are used for the extraction of the ωp scattering length. Some cross sections are scaled using the coefficients shown in the figure.

Figure 3 and Figure 4 show the differential cross sections of the ω meson photoproductions produced off the free proton on the liquid hydrogen target as a function of momentum transfer $-t$ of the exclusive analysis and the inclusive analysis [11], respectively. In the exclusive analysis, the recoil nucleon was identified in coincidence. In the inclusive analysis, there is no condition for detection of the recoil nucleon. We determined the parameters A and the b at four incident photon energies E_γ (1.175, 1.215 and 1.240 GeV) by fitting the ω mesons produced off the free proton in liquid hydrogen target with Eq. (3). The values of $d\sigma^{\gamma p}/dt(s_{thr}, t_{thr})$ and $d\sigma^{\gamma p}/dt(s_{thr}, 0)$ are calculated by fitting parameters A and b obtained by fitting the $|t|$ -dependent differential photoproduction cross-section data. The scattering length $|\alpha_{\omega p}|$ at the threshold t_{thr} and in t to $t=0$ are then calculated by Eq. (2). The extracted scattering length $|\alpha_{\omega p}|$ of the exclusive analysis and the inclusive analysis are listed in Table III and IV, respectively. The averaged scattering lengths $|\alpha_{\omega p}|$ of the exclusive analysis and inclusive analysis at different E_γ energies at threshold are calculated to be 0.811 ± 0.019 fm and 0.752 ± 0.021 fm, respectively. The averaged scattering length $|\alpha_{\omega p}|$ of the exclusive analysis and inclusive analysis at different E_γ energies in t to $t=0$ are calculated to be 1.293 ± 0.032 fm and 1.192 ± 0.030 fm, respectively. We see that the ωp scattering




FIG. 4. Differential cross sections of the photoproduction of ω meson produced off the free proton near the threshold versus the momentum transfer $-t$ in the hydrogen target [11]. The four incident photon energies E_γ (1.138, 1.163, 1.188 and 1.225 GeV) near the threshold of ω meson are marked in the figure. The inclusive experimental data by CBELSA/TAPS Collaboration are used for the extraction of the ωp scattering length. Some cross sections are scaled using the coefficients shown in the figure.

length of the free proton based on the exclusive data and the inclusive data agree with each other. We also find that the free proton ωp scattering length may be larger than the bound proton in the deuteron, but more precise data are needed to further check this. The A2 Collaboration at MAMI conducted an experimental of ω photoproduction on the proton and estimated that the ωp scattering length was equal to 0.82 ± 0.03 fm [2] based on the VMD model, which is consistent with the result of our extraction through incoherent experimental data of ω photoproductions provided by CBELSA/TAPS Collaboration [11]. The obtained scattering length $|\alpha_{\omega p}|$ in this work are consistent with the experimental measurements of photoproduction of the ω meson on the proton near the threshold by Ishikawa $|\alpha_{\omega p}| = 0.97 \pm 0.16$ fm [10]. In short, our results of the ωp scattering length are consistent with previous experimental measurements and extration results based on some theoretical models.

D. ϕN scattering length from ϕ differential photoproduction cross sections on the deuterium target

In this following analysis, the scattering length $|\alpha_{\phi N}|$ refers to the averaged value of the $|\alpha_{\omega n^*}|$ and $|\alpha_{\omega p^*}|$ of quasi-free neutron and proton in the deuteron. Figure 5 shows the differential cross sections of incoherent ϕ meson photoproductions from the deuteron as a function of momentum transfer $-t$. For the incoherent ϕ meson pho-




FIG. 5. Differential cross sections of the incoherent ϕ meson photoproduction ($\gamma d \rightarrow \phi p n$) near the threshold as a function of the momentum transfer $-t$ off the quasi-free nucleon in the deuteron target [12]. The five incident photon energies E_γ (1.62, 1.72, 1.82, 1.92 and 2.02 GeV) near the threshold of incoherent ϕ meson photoproduction are marked in the figure. Some cross sections are scaled using the coefficients shown in the figure.

toproduction from the deuteron in the LEPS spectrometer [12], a cut for the ϕ meson selection is performed on the missing mass spectra, to ensure the ϕ meson was interacting with the individual nucleon rather than the whole deuteron. The differential cross sections of the $|t|$ -dependence are still fitted with an exponential function. We determined the parameters A and the b with five incident photon energies E_γ (1.62, 1.72, 1.82, 1.92 and 2.02 GeV) by fitting the ϕ meson produced off the quasi-free proton or quasi-free neutron in the deuteron target with Eq. (3). The value of $d\sigma^{\gamma N}/dt(s_{thr}, t_{thr})$ and $d\sigma^{\gamma N}/dt(s_{thr}, 0)$ are calculated by fitting parameters A and b obtained by fitting the $|t|$ -dependent differential photoproduction cross-section data. The scattering length $|\alpha_{\phi N}|$ at the threshold t_{thr} and in t to $t = 0$ are then calculated by Eq. (2). The extracted scattering length $|\alpha_{\phi N}|$ are listed in Table V. The averaged scattering length $|\alpha_{\phi N}|$ of incoherent ϕ meson photoproduction in deuteron at different E_γ energies at threshold is calculated to be 0.109 ± 0.008 fm at threshold t_{thr} and 0.276 ± 0.010 fm in t to $t = 0$, respectively, The scattering length $|\alpha_{\phi N}|$ at threshold t_{thr} obtained in this work is a little larger than the estimated value ϕp at threshold t_{thr} extracted from the ϕ -meson photoproduction data from the CLAS experiment in Hall B of Jefferson Laboratory [4]. Furthermore, the value for $|\alpha_{\phi N}|$ at threshold t_{thr} as determined in this paper is smaller than the $|\alpha_{\phi N}|$ result 0.15 fm given from forward coherent ϕ -meson pho-

ACKNOWLEDGMENTS

This work is supported by the Strategic Priority Research Program of Chinese Academy of Sciences under

the Grant No. XDB34030301, the National Natural Science Foundation of China No. 12005266 and Guangdong Major Project of Basic and Applied Basic Research No. 2020B0301030008.

-
- [1] M. Gell-Mann and F. Zachariasen, *Phys. Rev.* **124**, 953 (1961).
- [2] I. I. Strakovsky *et al.*, *Phys. Rev. C* **91**, 045207 (2015), arXiv:1407.3465 [nucl-ex].
- [3] I. Strakovsky, D. Epifanov, and L. Pentchev, *Phys. Rev. C* **101**, 042201 (2020), arXiv:1911.12686 [hep-ph].
- [4] I. I. Strakovsky, L. Pentchev, and A. Titov, *Phys. Rev. C* **101**, 045201 (2020), arXiv:2001.08851 [hep-ph].
- [5] L. Pentchev and I. I. Strakovsky, *Eur. Phys. J. A* **57**, 56 (2021), arXiv:2009.04502 [hep-ph].
- [6] X.-Y. Wang, F. Zeng, and I. I. Strakovsky, *Phys. Rev. C* **106**, 015202 (2022), arXiv:2205.07661 [hep-ph].
- [7] X.-Y. Wang, F. Zeng, Q. Wang, and L. Zhang, (2022), arXiv:2206.09170 [nucl-th].
- [8] A. I. Titov, T. Nakano, S. Date, and Y. Ohashi, *Phys. Rev. C* **76**, 048202 (2007), arXiv:hep-ph/0703227.
- [9] R. L. Workman and Others (Particle Data Group), *PTEP* **2022**, 083C01 (2022).
- [10] T. Ishikawa *et al.*, *Phys. Rev. C* **101**, 052201 (2020), arXiv:1904.02797 [nucl-ex].
- [11] F. Dietz *et al.* (CBELSA/TAPS), *Eur. Phys. J. A* **51**, 6 (2015).
- [12] W. C. Chang *et al.* (LEPS), *Phys. Lett. B* **684**, 6 (2010), arXiv:0907.1705 [nucl-ex].
- [13] P. A. Zyla *et al.* (Particle Data Group), *PTEP* **2020**, 083C01 (2020).
- [14] T. Mibe *et al.* (LEPS), *Phys. Rev. Lett.* **95**, 182001 (2005), arXiv:nucl-ex/0506015.
- [15] W. C. Chang *et al.*, *Phys. Lett. B* **658**, 209 (2008), arXiv:nucl-ex/0703034.
- [16] Y. Koike and A. Hayashigaki, *Prog. Theor. Phys.* **98**, 631 (1997), arXiv:nucl-th/9609001.
- [17] H. Gao, T. S. H. Lee, and V. Marinov, *Phys. Rev. C* **63**, 022201 (2001), arXiv:nucl-th/0010042.
- [18] I. I. Strakovsky, W. J. Briscoe, L. Pentchev, and A. Schmidt, *Phys. Rev. D* **104**, 074028 (2021), arXiv:2108.02871 [hep-ph].
- [19] N. Baillie *et al.* (CLAS), *Phys. Rev. Lett.* **108**, 142001 (2012), [Erratum: *Phys.Rev.Lett.* 108, 199902 (2012)], arXiv:1110.2770 [nucl-ex].
- [20] S. Tkachenko *et al.* (CLAS), *Phys. Rev. C* **89**, 045206 (2014), [Addendum: *Phys.Rev.C* 90, 059901 (2014)], arXiv:1402.2477 [nucl-ex].
- [21] W. R. Armstrong *et al.*, (2017), arXiv:1708.00835 [nucl-ex].
- [22] W. Armstrong *et al.*, (2017), arXiv:1708.00888 [nucl-ex].
- [23] W. Armstrong *et al.*, (2017), arXiv:1708.00891 [nucl-ex].
- [24] A. Accardi *et al.*, *Eur. Phys. J. A* **52**, 268 (2016), arXiv:1212.1701 [nucl-ex].
- [25] X. Chen, *Proceedings, 26th International Workshop on Deep Inelastic Scattering and Related Subjects (DIS 2018): Port Island, Kobe, Japan, April 16-20, 2018*, PoS **DIS2018**, 170 (2018), arXiv:1809.00448 [nucl-ex].
- [26] X. Chen, F.-K. Guo, C. D. Roberts, and R. Wang, *Few Body Syst.* **61**, 43 (2020), arXiv:2008.00102 [hep-ph].
- [27] D. P. Anderle *et al.*, (2021), arXiv:2102.09222 [nucl-ex].

TABLE I. The absolute value of the extracted ωn^* scattering length obtained from the differential cross-section data of ω meson produced off the bound neutron in deuteron near threshold at different photon incident energies using different extrapolating methods.

E_γ (GeV)	1.175	1.215	1.240
$ \alpha_{\omega n^*} $ (fm) ($d\sigma^{\gamma n^*}/dt(s_{thr}, t_{thr})$)	0.683 ± 0.163	0.717 ± 0.125	0.715 ± 0.124
$ \alpha_{\omega n^*} $ (fm) ($d\sigma^{\gamma n^*}/dt(s_{thr}, 0)$)	1.337 ± 0.226	1.263 ± 0.224	1.176 ± 0.222

TABLE II. The absolute value of the extracted ωp^* scattering length obtained from the differential cross-section data of ω meson produced off the bound proton in deuteron near threshold at different photon incident energies using different extrapolating methods.

E_γ (GeV)	1.175	1.215	1.240
$ \alpha_{\omega p^*} $ (fm) ($d\sigma^{\gamma p^*}/dt(s_{thr}, t_{thr})$)	0.608 ± 0.048	0.616 ± 0.033	0.631 ± 0.035
$ \alpha_{\omega p^*} $ (fm) ($d\sigma^{\gamma p^*}/dt(s_{thr}, 0)$)	1.110 ± 0.063	0.973 ± 0.049	1.125 ± 0.056

TABLE III. The absolute value of the extracted ωp scattering length obtained from the differential cross-section data of ω meson produced off the free proton near threshold with exclusive analysis at different photon energies using different extrapolating methods.

E_γ (GeV)	1.138	1.163	1.188	1.225
$ \alpha_{\omega p} $ (fm) ($d\sigma^{\gamma p}/dt(s_{thr}, t_{thr})$)	0.827 ± 0.091	0.816 ± 0.043	0.835 ± 0.029	0.770 ± 0.034
$ \alpha_{\omega p} $ (fm) ($d\sigma^{\gamma p}/dt(s_{thr}, 0)$)	2.077 ± 0.158	1.581 ± 0.081	1.229 ± 0.044	1.155 ± 0.057

TABLE IV. The absolute value of the extracted ωp scattering length obtained from the differential cross-section data of ω meson produced off the free proton near threshold with inclusive analysis at different photon energies using different extrapolating methods.

E_γ (GeV)	1.138	1.163	1.188	1.225
$ \alpha_{\omega p} $ (fm) ($d\sigma^{\gamma p}/dt(s_{thr}, t_{thr})$)	0.740 ± 0.192	0.722 ± 0.076	0.787 ± 0.054	0.748 ± 0.024
$ \alpha_{\omega p} $ (fm) ($d\sigma^{\gamma p}/dt(s_{thr}, 0)$)	1.277 ± 0.237	1.200 ± 0.125	1.150 ± 0.081	1.197 ± 0.033

TABLE V. The absolute value of the extracted ϕN scattering length obtained from from the differential cross sections of incoherent ϕ mesons photonproduction from the deuteron near threshold at different photon energies using different extrapolating methods.

E_γ (GeV)	1.62	1.72	1.82	1.92	2.02
$ \alpha_{\phi N} $ (fm) ($d\sigma^{\gamma N}/dt(s_{thr}, t_{thr})$)	0.105 ± 0.114	0.066 ± 0.041	0.103 ± 0.024	0.114 ± 0.015	0.110 ± 0.011
$ \alpha_{\phi N} $ (fm) ($d\sigma^{\gamma N}/dt(s_{thr}, 0)$)	0.171 ± 0.110	0.397 ± 0.135	0.290 ± 0.034	0.285 ± 0.018	0.271 ± 0.012

TABLE VI. The ratio of the scattering length $|\alpha_{\omega n^*}|$ to the scattering length $|\alpha_{\omega p^*}|$ with the deuterium target data and the ratio of the scattering length $|\alpha_{\omega p}|$ of the hydrogen target data from the exclusive analysis and from the inclusive analysis to the scattering length $|\alpha_{\omega p^*}|$ with the deuterium target data.

Category	$ \alpha_{\omega n^*} / \alpha_{\omega p^*} $	$ \alpha_{\omega p} / \alpha_{\omega p^*} $
Ratio ($d\sigma^{\gamma p}/dt(s_{thr}, t_{thr})$)	1.142 ± 0.116	1.263 ± 0.040
Ratio ($d\sigma^{\gamma p}/dt(s_{thr}, 0)$)	1.191 ± 0.108	1.174 ± 0.035
Average Ratio	1.168 ± 0.079	1.213 ± 0.026