Quantum Reflective Kinks

Jarah Evslin^{1,2} * and Hui Liu^{3,2,4} \dagger

 Institute of Modern Physics, NanChangLu 509, Lanzhou 730000, China
 University of the Chinese Academy of Sciences, YuQuanLu 19A, Beijing 100049, China
 School of Fundamental Physics and Mathematical Sciences, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou 310024, China
 Arnold Sommerfeld Center, Ludwig-Maximilians-Universität, Theresienstraße 37, 80333 München, Germany

Abstract

We scatter a meson off of a scalar kink in quantum field theory, at leading order in perturbation theory. We calculate the full quantum state, at leading order, at all times and also check that the reflection and transmission coefficients agree with those which would be obtained in relativistic quantum mechanics.

1 Introduction

Classical kink-(anti)kink scattering has been a major industry since the discovery of a fractal pattern of resonance windows in the ϕ^4 double well model in Ref. [1]. For example, there has recently been a flurry of activity trying to understand whether the resonances are caused by the shape modes of individual kinks or collective bound modes of the combined kink-antikink system [2, 3, 4].

In contrast, the much simpler process of kink-meson scattering has received relatively little attention [5, 6, 7, 8, 9, 10]. In classical ϕ^4 field theory, Refs. [11, 12] discovered that the mesons apply a negative pressure to the kinks. The reason for this negative pressure is quite simple. Inside of the kink, n mesons of frequency ω can fuse into one meson of frequency $n\omega$, which as a result of the mass shell condition has more momentum than the sum of the initial n mesons. Conservation of momentum demands that this excess momentum comes from the kink, forcing the kink to move in the opposite direction.

This meson fusion appears to be the only phenomenon which occurs in meson-kink scattering in the classical ϕ^4 model. However in the case of reflective kinks, the mesons may

^{*}jarah@impcas.ac.cn

[†]hui.liu@campus.lmu.de

also reflect, leading to a positive pressure [13]. Thus, in general, two processes are allowed in classical kink-meson scattering, fusion and reflection.

In the quantum theory, we expect a much richer phenomenology. For one, an infinite tower of unstable excited states corresponding to multiple shape mode excitations is expected. These should appear as narrow resonances in elastic kink-meson scattering. Inelastic kink-meson scattering is also expected to be rich. For example, Raman spectroscopy will be possible, in which a monochromatic meson scatters off of the kink, exciting an internal shape mode. The kink's excitation spectrum can be read off of the observed energy decreases of the scattered meson. Deexcitation of a shape mode is also possible, in the inverse process. Finally, in the quantum theory, one expects not only meson fusion, but also meson fission.

All of these processes are expected to occur at tree level in the kink Hamiltonian, introduced in Ref. [14]. However, in traditional approaches which allow multiloop calculations, such as the collective coordinate method of Refs. [15, 16], they have been prohibitively difficult to calculate. Recently a new approach, linearized kink perturbation theory, has been introduced at one-loop in Ref. [17] and beyond in Ref. [18]. This new formalism is particularly simple¹ in the one-kink sector, which contains a single kink and any number of mesons and impurities. It is thus well suited to the scattering of a meson with a nonrelativistic kink.

We plan to investigate all of these new tree-level phenomena in the near future. However, so far applications of the formalism have suffered from two limitations. First, all explicit calculations have considered only kinks described by reflectionless potentials² and second, all calculations so far have been time-independent.

In the present note, we treat kink-meson scattering using the free kink Hamiltonian. As will be reviewed below, the free kink Hamiltonian describes a free theory in the sense that it is a sum of quantum harmonic oscillators, but it has a position-dependent mass term which leads to some dynamics. Clearly, this is a calculation which needs to be performed before our general study of tree level interactions. In doing so, we treat the two shortcomings noted above. In particular, we treat not only reflectionless but also reflective kinks. We evolve the system in time, and so we observe that our treatment of reflective kinks, in the free theory, reproduces the same reflection and transmission coefficients that would be calculated in relativistic quantum mechanics.

¹It requires a choice of base point in moduli space. If the kink moves too far from the base point, as will happen in kink-(anti)kink scattering, then one must compose the evolutions calculated at distinct base points.

²An exception to this is Ref. [19], however the treatment of the particular features of reflective normal modes was handled inside of the code that performed the numerical integrals, and only described roughly in an Appendix. The treatment here is related to that which was used inside the numerical code.

2 Review

Consider a (1+1)-dimensional Schrodinger picture quantum field theory of a scalar meson field $\phi(x)$ with conjugate $\pi(x)$ defined by the Hamiltonian H

$$\mathcal{H}(x) = \frac{1}{2} : \pi(x)\pi(x) :_{a} + \frac{1}{2} : \partial_{x}\phi(x)\partial_{x}\phi(x) :_{a} + \frac{1}{g^{2}} : V(g\phi(x)) :_{a}$$
$$H = \int dx \mathcal{H}(x), \qquad m^{2} = V^{(2)}(gf(\pm\infty)).$$
(2.1)

Here $::_a$ is the usual normal-ordering with mass m, $V^{(n)}(gf(x))$ is the *n*th derivative of V with respect to its argument and f(x) is a kink solution

$$\phi(x,t) = f(x), \qquad -gf''(x) + V^{(1)}(gf(x)) = 0.$$
 (2.2)

If the two sign choices in the definition of m^2 lead to different values, then the quantumcorrected vacuum energies will differ on the two sides of the kink and so the kink will accelerate [20]. In this case the kink states are never Hamiltonian eigenstates and so we will not consider this case further.

In terms of the defining Hamiltonian, the kink states are nonperturbative. In classical field theory this corresponds to the fact that the classical field $\phi(x)$ is far from a minimum of the potential, and it would be fixed by writing the Hamiltonian in terms of $\phi(x) - f(x)$. In the quantum theory such a replacement is ill-defined as a result of the regularization, which in this case is achieved via normal ordering. Indeed a naive application of the replacement can change the spectrum of the Hamiltonian [21] and so lead to wrong answers. To ensure that the spectrum of the Hamiltonian is not changed, we transform it to the kink Hamiltonian H' by conjugating with a unitary displacement operator

$$\mathcal{D}_f = \operatorname{Exp}\left[-i\int dx f(x)\pi(x)\right], \qquad H' = \mathcal{D}_f^{\dagger}H\mathcal{D}_f.$$
 (2.3)

This displacement operator plays the same role in the quantum theory as the shift in the classical theory

$$: F[\phi(x), \pi(x)] :_{a} \mathcal{D}_{f} = \mathcal{D}_{f} : F[\phi(x) + f(x), \pi(x)] :_{a} .$$
(2.4)

In summary, we have constructed a kink Hamiltonian H' which has the same spectrum as the defining Hamiltonian H. While the defining Hamiltonian generates time evolution in the defining frame of the Hilbert space, the unitary transformation \mathcal{D}_f^{\dagger} takes this defining frame to the kink frame. Masses may be measured and time may be evolved in the kink frame using H'. The isospectral property means that these masses will be the correct ones

$$H|K\rangle = E|K\rangle \Rightarrow H'\mathcal{D}_f^{\dagger}|K\rangle = E\mathcal{D}_f^{\dagger}|K\rangle$$
 (2.5)

and an application of \mathcal{D}_f after time evolution yields the time evolution that would be calculated using the defining Hamiltonian [22]

$$\mathcal{D}_f e^{-iH't} \mathcal{D}_f^{\dagger} = e^{-iHt}.$$
(2.6)

Thus nonperturbative problems using H can be transformed into problems which, in the one-kink sector, are perturbative using H'.

One can use Eq. (2.3) to calculate the kink Hamiltonian H'. Let us decompose it into terms H'_j which, when normal ordered, have j factors of ϕ and its conjugate π . These terms will be of order g^{j-2} . Then H'_0 is the classical kink mass Q_0 , while the tadpole H'_1 vanishes as a result of the classical equations of motion (2.2). The first nontrivial term is the free kink Hamiltonian

$$H_2' = \frac{1}{2} \int dx \left[:\pi^2(x) :_a + :(\partial_x \phi(x))^2 :_a + V^{(2)}(gf(x)) :\phi^2(x) :_a \right].$$
(2.7)

It is free, in the sense that it is quadratic in the fields, but notice that the mass term depends on x. This means that the constant frequency solutions of its classical equations of motion

$$V^{(2)}(gf(x))\mathfrak{g}(x) = \omega^2\mathfrak{g}(x) + \mathfrak{g}''(x), \qquad \phi(x,t) = e^{-i\omega t}\mathfrak{g}(x)$$
(2.8)

are not plane waves, but rather are normal modes.

In general there are three kinds of normal mode, classified by their frequencies ω . There is always a zero mode $\mathfrak{g}_B(x) = f'(x)/\sqrt{Q_0}$ with zero frequency $\omega_B = 0$. Sometimes there will be real shape modes $\mathfrak{g}_S(x)$ with frequencies $0 < \omega_S < m$. For each real k there will be a continuum mode with frequency ω_k

$$\omega_k = \sqrt{m^2 + k^2}.\tag{2.9}$$

As $\omega_k = \omega_{-k}$ one needs to define the decomposition into these two modes. We will require

$$\mathfrak{g}_{-k}(x) = \mathfrak{g}_k^*(x) \tag{2.10}$$

and we will fix the normalization and decomposition of all modes using the completeness relations³ in k space

$$\int dx |\mathbf{g}_B(x)|^2 = 1, \ \int dx \mathbf{g}_{k_1}(x) \mathbf{g}_{k_2}^*(x) = 2\pi \delta(k_1 - k_2), \ \int dx \mathbf{g}_{S_1}(x) \mathbf{g}_{S_2}^*(x) = \delta_{S_1 S_2}$$
(2.11)

and in x space

$$\mathfrak{g}_B(x)\mathfrak{g}_B(y) + \sum \frac{dk}{2\pi}\mathfrak{g}_k(x)\mathfrak{g}_k^*(y) = \delta(x-y), \qquad \sum \frac{dk}{2\pi} = \int \frac{dk}{2\pi} + \sum_S$$
(2.12)

³Completeness follows from the fact that (2.8) is a Sturm-Liouville equation.

where we have defined Σ to be an integral over k together with a sum over shape modes S.

As the normal modes are a complete basis of bounded functions of x, and we are working in the Schrödinger picture where the fields only depend on x, we may follow Ref. [23] and decompose the fields in terms of normal modes

$$\phi(x) = \phi_0 \mathfrak{g}_B(x) + \sum \frac{dk}{2\pi} \left(B_k^{\ddagger} + \frac{B_{-k}}{2\omega_k} \right) \mathfrak{g}_k(x)$$

$$\pi(x) = \pi_0 \mathfrak{g}_B(x) + i \sum \frac{dk}{2\pi} \left(\omega_k B_k^{\ddagger} - \frac{B_{-k}}{2} \right) \mathfrak{g}_k(x)$$
(2.13)

where $B_k^{\ddagger} = B_k^{\dagger}/(2\omega_k)$ and $B_{-S} = B_S$. Thus any operator in the theory may be expanded in terms of the operators ϕ_0 , π_0 , B_S , B_S^{\ddagger} , B_k and B_k^{\ddagger} which, as a result of the canonical commutation relations satisfied by $\phi(x)$ and $\pi(x)$, satisfy the algebra

$$[\phi_0, \pi_0] = i,$$
 $[B_{S_1}, B_{S_2}^{\ddagger}] = \delta_{S_1 S_2},$ $[B_{k_1}, B_{k_2}^{\ddagger}] = 2\pi\delta(k_1 - k_2).$

Inserting the decomposition (2.13) into the formula (2.7) for the free kink Hamiltonian, we find an enormous simplification

$$H_{2}' = Q_{1} + \frac{\pi_{0}^{2}}{2} + \int \frac{dk}{2\pi} \omega_{k} B_{k}^{\ddagger} B_{k}$$
(2.14)

where

$$Q_1 = -\frac{1}{4} \sum \frac{dk}{2\pi} \int \frac{dp}{2\pi} \frac{(\omega_p - \omega_k)^2}{\omega_p} \tilde{\mathfrak{g}}_k^2(p) - \frac{1}{4} \int \frac{dp}{2\pi} \omega_p \tilde{\mathfrak{g}}_B(p) \tilde{\mathfrak{g}}_B(p), \qquad \tilde{\mathfrak{g}}(p) = \int dx \mathfrak{g}(x) e^{ipx}.$$
(2.15)

We have found that the free kink Hamiltonian H'_2 is the sum of three terms. The first, Q_1 , is the one-loop kink mass in the form found in Ref. [23]. The second describes the quantum mechanics of a free particle, in this case the center of mass of the kink. The third is an infinite sum of quantum harmonic oscillators, one for each continuum mode $\mathfrak{g}_k(x)$ and also one for each shape mode $\mathfrak{g}_S(x)$ if there are any. The ground state $|0\rangle_0$ of the free kink Hamiltonian therefore corresponds to the ground state of each of these quantum mechanical systems

$$\pi_0|0\rangle_0 = B_k|0\rangle_0 = B_S|0\rangle_0 = 0.$$
(2.16)

This state is the first approximation in a semiclassical expansion of the kink ground state, in the kink frame. At this order, continuum normal modes and shape modes can be excited in the kink frame by acting with B_k^{\ddagger} and B_S^{\ddagger} respectively. We remind the reader that the corresponding states in the defining frame of the Hilbert space are then obtained by acting with \mathcal{D}_f , so that $\mathcal{D}_f|0\rangle_0$ is the leading term in the kink ground state.

3 Normal Modes

Consider normal modes with asymptotic behavior

$$\mathfrak{g}_{k}(x) = \begin{cases} B_{k}e^{ikx} + C_{k}e^{-ikx} & \text{if } x \ll -1/m \\ D_{k}e^{ikx} + E_{k}e^{-ikx} & \text{if } x \gg 1/m \end{cases}$$

$$B_{k}^{*} = B_{-k}, \quad C_{k}^{*} = C_{-k}, \quad D_{k}^{*} = D_{-k}, \quad E_{k}^{*} = E_{-k}.$$
(3.1)

The coefficients B, C, D and E are constrained by the completeness relations (2.11) and (2.12). Normal modes at distinct values of |k| will automatically be orthogonal as they satisfy the same Sturm-Liouville equation with a distinct eigenvalue. However some care needs to be taken to ensure that $\mathfrak{g}_k(x)$ and $\mathfrak{g}_{-k}(x)$ are orthogonal and correctly normalized. As the nonvanishing integrals are infinite, we only consider the large |x| region.

Let us impose the k-space completeness relation (2.11) for the modes \mathfrak{g}_k . Since we are only interested in large |x|, it is sufficient to impose

$$\lim_{L \to \infty} \left[\int_{x=-2L}^{x=-L} + \int_{x=L}^{x=2L} \right] dx \mathfrak{g}_{k_1}(x) \mathfrak{g}_{k_2}(x) = \begin{cases} 2L & \text{if } k_1 = -k_2 \\ 0 & \text{if } k_1 \neq -k_2. \end{cases}$$
(3.2)

This is easily evaluated

$$\lim_{L \to \infty} \left[\int_{x=-2L}^{x=-L} + \int_{x=L}^{x=2L} \right] \frac{dx}{L} \mathfrak{g}_{k_1}(x) \mathfrak{g}_{k_2}(x) = \begin{cases} |B_k|^2 + |C_k|^2 + |D_k|^2 + |E_k|^2 & \text{if } k = k_1 = -k_2 \\ 2B_k C_k + 2D_k E_k & \text{if } k = k_1 = k_2 \\ 0 & \text{if } k_1 \neq \pm k_2. \end{cases}$$

$$(3.3)$$

Then we learn that

$$|B_k|^2 + |C_k|^2 + |D_k|^2 + |E_k|^2 = 2, \qquad B_k C_k + D_k E_k = 0.$$
(3.4)

Next we impose the completeness relation in position space (2.12). When |x|, $|y| \gg 1/m$ only the continuum modes contribute, leaving

$$\int \frac{dk}{2\pi} \mathfrak{g}_k(x) \mathfrak{g}_{-k}(y) = \delta(x-y).$$
(3.5)

Evaluating the left hand side we find

$$\int \frac{dk}{2\pi} \mathfrak{g}_k(x) \mathfrak{g}_{-k}(y) = \int \frac{dk}{2\pi} \begin{cases} (|B_k|^2 + |C_k|^2) e^{ik(x-y)} & \text{if } x, \ y \ll -1/m \\ (|D_k|^2 + |E_k|^2) e^{ik(x-y)} & \text{if } x, \ y \gg 1/m \\ (B_k^* E_k + C_k D_k^*) e^{-ik(x+y)} & \text{if } x \ll -1/m, \ 1/m \ll y. \end{cases}$$
(3.6)

We are interested in the case in which |x| and |y| are very large and so B_k , C_k and D_k and E_k vary much more slowly than the plane wave factors with respect to k. To avoid an unwanted contribution at x = -y we impose the constraint

$$B_k^* E_k + C_k D_k^* = 0. (3.7)$$

In addition, to arrive at the correct normalization of the delta function in the limit of large |x| and |y|, one finds the constraints

$$|B_k|^2 + |C_k|^2 = |D_k|^2 + |E_k|^2 = 1.$$
(3.8)

We can summarize all of these constraints with the condition that the matrix

$$U = \begin{pmatrix} B_k & C_k^* \\ E_k & D_k^* \end{pmatrix}, \qquad U^{\dagger}U = 1$$
(3.9)

is unitary.

4 Propagating Wave Packets

Consider a moving wave packet

$$\Phi(x) = \exp\left[-\frac{(x-x_0)^2}{4\sigma^2} + ixk_0\right], \qquad x_0 \ll -1/m, \qquad k_0 \gg \frac{1}{\sigma}, \qquad \sigma \ll |x_0|.$$
(4.1)

This corresponds to a meson beginning far to the left of a kink, at $x = x_0$, and moving to the right with momentum k_0 . We will assume that it moves fast enough that smearing can be neglected.

The completeness of the normal modes implies that, at $|x| \gg 1/m$, any wave packet may be decomposed

$$\Phi(x) = \int \frac{dk}{2\pi} \alpha_k \mathfrak{g}_k(x), \qquad \alpha_k = \int dx \Phi(x) \mathfrak{g}_k^*(x).$$
(4.2)

The wave packet (4.1) is supported at $x \ll -1/m$ and so one may insert the asymptotic formula for $\mathfrak{g}_k^*(x)$ valid at $x \ll -1/m$. Therefore

$$\alpha_k = 2\sigma\sqrt{\pi} \left(B_k^* e^{-i(k-k_0)x_0} e^{-(k-k_0)^2 \sigma^2} + C_k^* e^{i(k+k_0)x_0} e^{-(k+k_0)^2 \sigma^2} \right).$$
(4.3)

In Dirac notation, in the kink frame, the wave packet corresponds to the one-meson state

$$|\Phi\rangle = \int dx \Phi(x) |x\rangle = \int \frac{dk}{2\pi} \alpha_k |k\rangle, \qquad |k\rangle = B_k^{\ddagger} |0\rangle_0, \qquad |x\rangle = \int \frac{dk}{2\pi} \mathfrak{g}_k^*(x) |k\rangle \tag{4.4}$$

which is part of the meson Fock space decomposition of the one-kink sector. Thus we begin with a meson at $x = x_0$ moving towards a kink at $x \sim 0$. We remind the reader that $|0\rangle_0$, defined in Eq. (2.16), is the lowest order approximation to the ground state of a single kink, as expressed in the kink frame of the Hilbert space.

At time t the wave packet evolves to

$$|\Phi(t)\rangle = \int dx \Phi(x,t)|x\rangle, \qquad \Phi(x,t) = \int \frac{dk}{2\pi} e^{-i\omega_k t} \alpha_k \mathfrak{g}_k(x). \tag{4.5}$$

With σ large enough, k is always close to k_0 or $-k_0$. Therefore we will expand in powers of $(k \pm k_0)$, out to first order. Wave packet smearing appears only at second order and so will be missed by this approximation. At first order

$$\omega_k = \omega_{k_0} + (\pm k - k_0) \frac{k_0}{\omega_{k_0}}.$$
(4.6)

Inserting (4.3) and (4.6) into Eq. (4.5) we find the wave packet evolution.

At $x \ll 0$ we obtain the wave packet

$$\Phi(x,t) = 2\sigma\sqrt{\pi}e^{-i\omega_{k_0}t} \int \frac{dk}{2\pi} \left(B_k^* e^{-i(k-k_0)\left(x_0 + \frac{k_0}{\omega_{k_0}}t\right)} e^{-(k-k_0)^2\sigma^2} + C_k^* e^{i(k+k_0)\left(x_0 + \frac{k_0}{\omega_{k_0}}t\right)} e^{-(k+k_0)^2\sigma^2} \right) \left(B_k e^{ikx} + C_k e^{-ikx} \right).$$
(4.7)

For σ large enough, the first Gaussian is supported at $k = k_0$ and the second at $k = -k_0$ and so we approximate the coefficients B_k and C_k by their values at $k = \pm k_0$, yielding

$$\Phi(x,t) = e^{-i\omega_{k_0}t} \left(\left(|B_{k_0}|^2 + |C_{k_0}|^2 \right) \exp\left[-\frac{\left(-x + x_0 + \frac{k_0}{\omega_{k_0}}t \right)^2}{4\sigma^2} + ik_0 x \right] + 2B_{k_0}^* C_{k_0} \exp\left[-\frac{\left(x + x_0 + \frac{k_0}{\omega_{k_0}}t \right)^2}{4\sigma^2} - ik_0 x \right] \right).$$
(4.8)

The two Gaussian factors are supported, respectively, when the position x is equal to plus or minus

$$x_t = x_0 + \frac{k_0}{\omega_{k_0}}t.$$
 (4.9)

Here we recognize k_0/ω_{k_0} as the group velocity of the wave packet and t as the propagation time. We are now considering the case $x \ll 0$, and so the first Gaussian is only supported when $x_t \ll 0$ and the second when $x_t \gg 0$.

Thus at early times $t \ll \omega_{k_0} |x_0| / k_0$, only the first Gaussian factor may contribute and we find

$$\Phi(x,t) = e^{-i\omega_{k_0}t + ik_0x} \operatorname{Exp}\left[-\frac{\left(-x + x_0 + \frac{k_0}{\omega_{k_0}}t\right)^2}{4\sigma^2}\right] = e^{-i\left(m^2/\omega_{k_0}\right)t} \Phi\left(x - \frac{k_0}{\omega_{k_0}}t\right)$$
(4.10)

where we have used (3.8). The interpretation is that, to this order in the $(k - k_0)$ expansion, the wave packet simply moves rigidly to the right before arriving at the kink. At times $t \gg \omega_{k_0}|x_0|/k_0$, after the meson and kink have interacted, only the second Gaussian may contribute. Thus, at $x \ll -1/m$, the wave function becomes

$$\Phi(x,t) = 2e^{-i\omega_{k_0}t - ik_0x} B_{k_0}^* C_{k_0} \operatorname{Exp}\left[-\frac{\left(x + x_0 + \frac{k_0}{\omega_{k_0}}t\right)^2}{4\sigma^2}\right].$$
(4.11)

We see that the kink's momentum has changed sign, now the part of the wave packet at x < 0 is traveling in the opposite direction. Also the position is $x = -x_t$, it has reflected from the kink. However the entire wave packet has not reflected. Here we have only calculated the contribution to the wave packet at x < 0 and the amplitude has been damped by the factor $2B^*C$. We interpret this factor as the reflection coefficient.

What happens on the right side of the kink? One need only insert $\mathfrak{g}_k(x)$ from Eq. (3.1), at $x \gg 0$, into Eq. (4.5) to obtain

$$\Phi(x,t) = e^{-i\omega_{k_0}t} \left(\left(B_{k_0}^* D_{k_0} + C_{k_0} E_{k_0}^* \right) \operatorname{Exp} \left[-\frac{\left(-x + x_0 + \frac{k_0}{\omega_{k_0}}t \right)^2}{4\sigma^2} + ik_0 x \right] + \left(B_{k_0}^* E_{k_0} + C_{k_0} D_{k_0}^* \right) \operatorname{Exp} \left[-\frac{\left(x + x_0 + \frac{k_0}{\omega_{k_0}}t \right)^2}{4\sigma^2} - ik_0 x \right] \right).$$
(4.12)

The second line vanishes as a result of the constraint Eq. (3.7). The Gaussian on the first line has support when $x = x_t$.

Since we are considering x > 0, this only occurs when $x_t > 0$ and so at $t \gg \omega_{k_0} |x_0|/k_0$, after the meson has interacted with the kink. This is reasonable, the meson cannot get to the right of the kink before they have interacted. At these late times, the position is x_t , implying that at this order the meson has continued to move past the kink at its initial velocity. We interpret the $(B^*D + CE^*)$ factor as the transmission coefficient.

5 Comparison with Quantum Mechanics

The above calculation was equivalent to one in relativistic quantum mechanics. This is because the kink frame of quantum field theory reduces the leading order problem of meson-kink scattering in the full quantum field theory to a simple quantum mechanical exercise. Of course the above calculation has the advantage that it can be generalized to higher orders, revealing for example the backreaction of the meson on the kink, which is invisible in quantum mechanics where the kink is replaced by a potential well.

But is the reflection coefficient really $2B^*C$? Usually in quantum mechanics one does not use a normal mode basis for such a problem, as it complicates the derivation and obscures the physics. It was only used here because of the simple connection to the kink-frame meson Fock space in Eq. (4.4).

For simplicity, let us consider quantum mechanics with a symmetric potential. This corresponds to a symmetric mass term in H_2 , which arises from an antisymmetric kink in a symmetric potential in quantum field theory. Beyond such cases, in general the quantum corrections to the vacuum energies on the two sides of the kink will disagree and so the kink will accelerate [24], and will not correspond to a Hamiltonian eigenstate. While such problems are invisible at the low order of perturbation theory considered here, and anyway do not prevent the application of our formalism, nonetheless they justify our interest in symmetric potentials.

In quantum mechanics, the kink at this order is replaced by a potential well and scattering is described by a symmetric and unitary S-matrix

$$S = \left(\begin{array}{cc} t & r \\ r & t \end{array}\right) \tag{5.1}$$

where t and r are the complex transmission and reflection coefficients such that, as a result of the unitarity of S,

$$|t|^2 + |r|^2 = 1, \qquad \operatorname{Arg}(r) = \operatorname{Arg}(t) \pm \frac{\pi}{2}.$$
 (5.2)

Consider k > 0. Then B_k and E_k have the interpretations of particles incident on the well from the left and right respectively, while C_k and D_k correspond to particles moving away from the well on the left and right. The S matrix relates these

$$\left(\begin{array}{c} C_k\\ D_k\end{array}\right) = S \left(\begin{array}{c} E_k\\ B_k\end{array}\right).$$
(5.3)

The vectors (C_k, D_k) and (B_k, E_k) are orthogonal (3.4). This fixes (C_k, D_k) up to a phase ϕ

$$S\begin{pmatrix} E_k\\ B_k \end{pmatrix} = \begin{pmatrix} C_k\\ D_k \end{pmatrix} = e^{i\phi} \begin{pmatrix} -E_k\\ B_k \end{pmatrix}.$$
(5.4)

This linear set of equations is solved by

$$B_k = -\frac{t + e^{i\phi}}{r}E_k, \qquad E_k = \frac{e^{i\phi} - t}{r}B_k.$$
 (5.5)

Writing

$$t = |t|e^{i\theta}, \qquad r = i\epsilon_1|r|e^{i\theta}, \qquad \epsilon_1 = \pm 1$$
(5.6)

and combining the two equations (5.5) one finds

$$E_{k} = \frac{e^{2i\phi} - |t|^{2} e^{2i\theta}}{|r|^{2} e^{2i\theta}} E_{k}$$
(5.7)

implying

$$e^{i\phi} = \epsilon_2 e^{i\theta}, \qquad \epsilon_2 = \pm 1$$
 (5.8)

and so

$$B_k = i\epsilon_1\epsilon_2 \frac{1+\epsilon_2|t|}{|r|} E_k = i\epsilon_1\epsilon_2 \sqrt{\frac{1+\epsilon_2|t|}{1-\epsilon_2|t|}} E_k.$$
(5.9)

The condition $|B_k|^2 + |E_k|^2 = 1$ then leads to

$$|B_k| = \sqrt{\frac{1+\epsilon_2|t|}{2}}, \qquad |E_k| = \sqrt{\frac{1-\epsilon_2|t|}{2}}.$$
 (5.10)

Then, for some phase λ ,

$$B_k = e^{i\lambda} \sqrt{\frac{1+\epsilon_2|t|}{2}}, \qquad E_k = -ie^{i\lambda} \epsilon_1 \epsilon_2 \sqrt{\frac{1-\epsilon_2|t|}{2}}.$$
(5.11)

Now using the original equation (5.4) we find

$$C_{k} = -e^{i\phi}E_{k} = ie^{i(\lambda+\phi)}\epsilon_{1}\epsilon_{2}\sqrt{\frac{1-\epsilon_{2}|t|}{2}}, \qquad D_{k} = e^{i\phi}B_{k} = e^{i(\lambda+\phi)}\sqrt{\frac{1+\epsilon_{2}|t|}{2}}.$$
 (5.12)

Therefore

$$2B_k^* C_k = 2ie^{i\phi} \epsilon_1 \epsilon_2 \sqrt{\frac{1+\epsilon_2|t|}{2}} \sqrt{\frac{1-\epsilon_2|t|}{2}} = ie^{i\phi} \epsilon_1 \epsilon_2 |r| = r.$$
(5.13)

Thus the quantum field theory coefficient for reflection, $2B^*C$, agrees with the usual result from quantum mechanics. Similarly

$$B_k^* D_k + C_k E_k^* = e^{i\phi} \left(|B_k|^2 - |E_k|^2 \right) = e^{i\phi} \epsilon_2 |t| = t$$
(5.14)

and so the transmission coefficient also agrees with the usual result from quantum mechanics.

6 Remarks

This calculation has been trivial. It is a calculation that has been performed countless times in relativistic quantum mechanics over the past century, corresponding to the scattering of a particle through a symmetric barrier or well. The interpretation of the calculation is somewhat novel, as the particle plays the role of the fundamental meson field $\phi(x)$ and the well is a leading approximation to a quantum kink. The marvel of the kink frame is that it transforms a nonperturbative quantum field theory computation into an old exercise in quantum mechanics. However it is likely expected that kink-meson scattering can, at leading order in the kink Hamiltonian, be treated in quantum mechanics.

So what has been gained by repeating this old calculation? We have extended the formalism of Refs. [17, 18] to include both time-evolution and also reflective potentials. Reflective potentials are obviously interesting because they describe most kinks. However, more to the point, we expect reflection at tree level even for kinks which are reflectionless in the free theory, and so this development is necessary in our opinion even for a perturbative treatment of reflectionless kinks such as those of the ϕ^4 theory and the Sine-Gordon model.

More importantly, we have finally introduced time evolution into the formalism of linearized kink perturbation theory. It has always used the Schrodinger picture, where operators are time-independent. In addition, except for Ref. [19], it has restricted attention to Hamiltonian eigenstates, so that even the states do not evolve. Even in Ref. [19], only time t = 0was considered, although instantaneous accelerations were computed. The present note, on the other hand, describes for the first time how to apply this formalism to finite-time time evolution.

The leading order state in quantum field theory has been found at each time t, even during the interaction itself although the asymptotic formula for $\mathfrak{g}_k(x)$ cannot be applied there. The analytic formula for $\mathfrak{g}_k(x)$ are anyway known in the Sine-Gordon and ϕ^4 models, and very good numerical approximations have been found in many other models. Therefore Eqs.(4.3) and (4.5) indeed give the full state, at leading order in the semiclassical expansion, at any time during the evolution. This is progress with respect to Euclidean time methods, which are generally applied to the calculation of the S-matrix and so are essentially restricted to infinite-time evolution. These leading order expressions for the states at each time, as well as the general formalism for kink-frame time evolution introduced here, of course are necessary to compute the perturbative corrections, which include the new physics that we intend to investigate in the near future.

Acknowledgement

JE is supported by NSFC MianShang grants 11875296 and 11675223. HL acknowledges the support from CAS-DAAD Joint Fellowship Programme for Doctoral students of UCAS.

References

- [1] D. K. Campbell, J. F. Schonfeld and C. A. Wingate, "Resonance structure in kinkantikink interactions in ϕ^4 theory," Physica **D9** (1983) 1.
- [2] P. Dorey, K. Mersh, T. Romanczukiewicz and Y. Shnir, "Kink-antikink collisions in the ϕ^6 model," Phys. Rev. Lett. **107** (2011), 091602 doi:10.1103/PhysRevLett.107.091602 [arXiv:1101.5951 [hep-th]].
- [3] P. Dorey, A. Gorina, I. Perapechka, T. Romańczukiewicz and Y. Shnir, "Resonance structures in kink-antikink collisions in a deformed sine-Gordon model," JHEP 09 (2021), 145 doi:10.1007/JHEP09(2021)145 [arXiv:2106.09560 [hep-th]].
- [4] C. Adam, P. Dorey, A. Garcia Martin-Caro, M. Huidobro, K. Oles, T. Romanczukiewicz,
 Y. Shnir and A. Wereszczynski, "Multikink scattering in the φ⁶ model revisited," [arXiv:2209.08849 [hep-th]].
- [5] L. D. Faddeev and V. E. Korepin, "Quantum Theory of Solitons: Preliminary Version," Phys. Rept. 42 (1978), 1-87 doi:10.1016/0370-1573(78)90058-3
- [6] M. Lowe, "BOSON SOLITON SCATTERING IN THE SINE-GORDON MODEL," Nucl. Phys. B 159 (1979), 349-362 doi:10.1016/0550-3213(79)90339-0
- [7] J. A. Parmentola and I. Zahed, "MESON SOLITON SCATTERING WITH SOLITON RECOIL," Print-87-0301 (STONY BROOK).
- [8] M. S. Swanson, "SOLITON-PARTICLE SCATTERING AND BERRY'S PHASE," Phys. Rev. D 38 (1988), 3122-3127 doi:10.1103/PhysRevD.38.3122
- [9] M. Uehara, A. Hayashi and S. Saito, "Meson soliton scattering with full recoil in standard collective coordinate quantization," Nucl. Phys. A 534 (1991), 680-696 doi:10.1016/0375-9474(91)90466-J
- [10] A. M. H. H. Abdelhady and H. Weigel, "Wave-Packet Scattering off the Kink-Solution," Int. J. Mod. Phys. A 26 (2011), 3625-3640 doi:10.1142/S0217751X11054012 [arXiv:1106.3497 [nlin.PS]].

- [11] T. Romanczukiewicz, "Interaction between kink and radiation in phi^{**}4 model," Acta Phys. Polon. B **35** (2004), 523-540 [arXiv:hep-th/0303058 [hep-th]].
- [12] T. Romanczukiewicz, "Interaction between topological defects and radiation," Acta Phys. Polon. B 36 (2005), 3877-3887
- [13] P. Forgacs, A. Lukacs and T. Romanczukiewicz, "Negative radiation pressure exerted on kinks," Phys. Rev. D 77 (2008), 125012 doi:10.1103/PhysRevD.77.125012 [arXiv:0802.0080 [hep-th]].
- R. F. Dashen, B. Hasslacher and A. Neveu, "Nonperturbative Methods and Extended Hadron Models in Field Theory 2. Two-Dimensional Models and Extended Hadrons," Phys. Rev. D 10 (1974) 4130. doi:10.1103/PhysRevD.10.4130
- [15] J. L. Gervais, A. Jevicki and B. Sakita, "Collective Coordinate Method for Quantization of Extended Systems," Phys. Rept. 23 (1976), 281-293 doi:10.1016/0370-1573(76)90049-1
- [16] J. L. Gervais and A. Jevicki, "Point Canonical Transformations in Path Integral," Nucl. Phys. B 110 (1976), 93-112 doi:10.1016/0550-3213(76)90422-3
- [17] J. Evslin, "Manifestly Finite Derivation of the Quantum Kink Mass," JHEP 11 (2019), 161 doi:10.1007/JHEP11(2019)161 [arXiv:1908.06710 [hep-th]].
- [18] J. Evslin and H. Guo, "Two-Loop Scalar Kinks," Phys. Rev. D 103 (2021) no.12, 125011 doi:10.1103/PhysRevD.103.125011 [arXiv:2012.04912 [hep-th]].
- [19] J. Evslin, C. Halcrow, T. Romanczukiewicz and A. Wereszczynski, "Spectral walls at one loop," Phys. Rev. D 105 (2022) no.12, 125002 doi:10.1103/PhysRevD.105.125002 [arXiv:2202.08249 [hep-th]].
- [20] H. Weigel, "Quantum Instabilities of Solitons," AIP Conf. Proc. 2116 (2019) no.1, 170002 doi:10.1063/1.5114153 [arXiv:1907.10942 [hep-th]].
- [21] A. Rebhan and P. van Nieuwenhuizen, "No saturation of the quantum Bogomolnyi bound by two-dimensional supersymmetric solitons," Nucl. Phys. B 508 (1997) 449 doi:10.1016/S0550-3213(97)00625-1, 10.1016/S0550-3213(97)80021-1 [hep-th/9707163].
- [22] Rob Pisarski, Private communication.
- [23] K. E. Cahill, A. Comtet and R. J. Glauber, "Mass Formulas for Static Solitons," Phys. Lett. B 64 (1976), 283-285 doi:10.1016/0370-2693(76)90202-1

[24] T. Romańczukiewicz, "Could the primordial radiation be responsible for vanishing of topological impuritys?," Phys. Lett. B 773 (2017), 295-299 doi:10.1016/j.physletb.2017.08.045 [arXiv:1706.05192 [hep-th]].