Classification of degenerate non-homogeneous Hamiltonian operators

Marta Dell'Atti¹ Pierandrea Vergallo^{2,3}

 ¹ School of Mathematics & Physics University of Portsmouth marta.dell'atti@port.ac.uk
 ² Department of Mathematical, Computer, Physical and Earth Sciences University of Messina,
 V.le F. Stagno D'Alcontres 31, I-98166 Messina, Italy pierandrea.vergallo@unime.it

³ Istituto Nazionale di Fisica Nucleare, Sez. Lecce

Abstract

We investigate non-homogeneous Hamiltonian operators composed of a first order Dubrovin-Novikov operator and an ultralocal one. The study of such operators turns out to be fundamental for the inverted system of equations associated with a class of Hamiltonian scalar equations. Often, the involved operators are degenerate in the first order term. For this reason a complete classification of the operators with degenerate leading coefficient in systems with two and three components is presented.

1 Introduction

The Hamiltonian formalism for Partial Differential Equations (PDEs) is one of the leading tools to study nonlinear systems [24, 6], following the well known developed theory for finite dimensional ones. As shown in [14, 17], Hamiltonian operators link conserved quantities with symmetries of the system, mapping the former onto the latter, and then leading to a deeper investigation of the structure of the solutions. This formalism represents a strong theoretical and practical connection between geometry and mathematical physics [22, 5, 2]. Dubrovin and Novikov introduced differential-geometric Poisson brackets as a natural extension of finite dimensional symplectic structures in traditional Hamiltonian mechanics that turned out to arise in several examples in nonlinear PDEs. The characterisation of these structures is related to (pseudo-) Riemannian geometry and algebraic geometry, especially for systems in 1 + 1 dimensions (or in independent variables x, t).

More in general, geometrical methods are well established tools widely used to find solutions to systems, such as the generalized hodograph method introduced by Tsarev [28], valid for strictly hyperbolic systems. The Hamiltonian formalism is also used to discuss the integrability. In particular, finding two compatible Hamiltonian structures is strictly related to the existence of infinitely many commuting symmetries and conservation laws, as proved by Magri [19]. In the context of hydrodynamic type systems, an approach to describe integrability is based on the analysis of geometrical elements in the so called method of hydrody-

namic reductions, developed for systems in 2 + 1 dimensions [10, 9] and then extended to systems in 1 + 1 dimensions with infinitely many components [8].

First, we recall some basic notions concerning the Hamiltonian formalism [22]. Let us consider a system described by *n* field variables

$$u^{i} = u^{i}(t, x), \qquad i = 1, \dots, n$$
 (1)

depending on the independent variables *t*, *x*, and let u_{σ}^{i} denote the *x*-derivatives of *u* σ times. A Hamiltonian operator is a linear operator $A^{ij} = a^{ij\sigma}D_{\sigma}$ such that the associated bracket for functionals *f*, *g*

$$\{f,g\} = \int \frac{\delta f}{\delta u^i} A^{ij} \frac{\delta g}{\delta u^j} dx,$$
(2)

is a Poisson bracket, i.e. it is bilinear, skew-symmetric and satisfies the Jacobi identity

$$\{f,g\} = -\{g,f\},\$$

$$\{f,\{g,h\}\} + \{g,\{h,f\}\} + \{h,\{f,g\}\} = 0.$$

An evolutionary system

$$u_t^i = F^i(x, u, u_\sigma), \tag{3}$$

with $u = \{u^i\}_{i=1}^n$ is Hamiltonian if it admits the following representation

$$u_t^i = F^i(x, u, u_\sigma) = A^{ij} \frac{\delta H}{\delta u^j},\tag{4}$$

where δ is the variational derivative, *H* is the Hamiltonian functional $H = \int h(u) dx$, written in terms of the Hamiltonian density *h*, and *A* is a Hamiltonian operator.

In [4, 5], Dubrovin and Novikov present a class of Hamiltonian operators which are homogeneous in the order of derivation and are also known as *homogeneous Hamiltonian operators*. They prove [4] that first order homogeneous operators of the form

$$g^{ij}\partial_x + \Gamma_k^{ij} u_x^k, \tag{5}$$

with det $g \neq 0$, are Hamiltonian if and only if

$$g_{ij} = (g^{ij})^{-1}, \qquad \Gamma_k^{ij} = -g^{is} \Gamma_{sk}^j,$$
 (6)

i.e. g^{ij} is a flat metric and the coefficients Γ_{jk}^i are Christoffel symbols for the metric tensor g. Operators of this type naturally arise in homogeneous quasilinear systems of first order PDEs, also known as *hydrodynamic type systems*

$$u_t^i = v_i^i(u) u_x^j, \qquad i = 1, \dots, n,$$
 (7)

where $v(u) = (v_j^i)_{1 \le i,j \le n}$ is the coefficient matrix depending on the field variables. Indeed, if (7) is Hamiltonian with a Dubrovin–Novikov operator (5), it can be expressed as

$$u_t^i = v_j^i(u) u_x^j = A^{ij} \frac{\partial h}{\partial u^j} = \left(g^{ij}\partial_x + \Gamma_k^{ij} u_x^k\right) \frac{\partial h}{\partial u^j} = \left(\nabla^i \nabla_j h\right) u_x^j,$$
(8)

where h = h(u) is the hydrodynamic Hamiltonian density and ∇_i the covariant derivative.

In the case of higher order homogeneous operators of degree *m*, the Dubrovin–Novikov operator generalizes as

$$A^{ij} = g^{ij} D_x^m + b_k^{ij} u_x^k D_x^{m-1} + \left(C_k^{ij} u_{xx}^k + C_{kl}^{ij} u_x^k u_x^l \right) D_x^{m-2} + \dots + \left(d_k^{ij} u_{nx}^k + \dots + d_{k_1\dots k_m}^{ij} u_x^{k_1} \dots u_x^{k_m} \right),$$
(9)

where the coefficients b_k^{ij} , C_k^{ij} , ... depend on the field variables. Dubrovin and Novikov present also an extension of homogeneous structures[5], introducing *non-homogeneous* Hamiltonian operators as sum of two or more homogeneous ones. A leading example in this context is offered by the Korteweg-De Vries equation

$$u_t = 6uu_x + u_{xxx}, \tag{10}$$

which possesses a Hamiltonian structure through the operator

$$A = \partial_x^3 + 2u\partial_x + u_x, \tag{11}$$

given by the sum of the third order operator ∂_x^3 and the first order operator $2u\partial_x + u_x$.

Following the notation used by Dubrovin and Novikov, if an operator is given by the sum of two homogeneous operators of order k and m respectively, we denote the order of the non-homogeneous operator via the sum k + m.

Let us consider the simplest case, with k = 1 and m = 0 for the so-called *non-homogeneous operators of hydrodynamic type* 1 + 0. They naturally arise in non-homogeneous quasilinear systems of first-order PDEs. Let $C^{ij} = A^{ij} + \omega^{ij}$, where A^{ij} is homogeneous of order 1 and ω^{ij} is a symplectic structure of order 0. One can easily generalise (8) to systems of this type

$$u_t^i = \left(g^{ij}\partial_x + \Gamma_k^{ij}u_x^k + \omega^{ij}\right)\frac{\partial h}{\partial u^j}$$

= $\left(\nabla^i \nabla_j h\right)u_x^j + \tilde{\nabla}^i h,$ (12)

where $\tilde{\nabla}^i = \omega^{is} \nabla_s$ and ∇_s is the standard gradient.

A remarkable example of a non-homogeneous quasilinear system possessing such a construction is given by the 3-waves equation [22]

$$\begin{cases}
u_t^1 = -c_1 u_x^1 - 2(c_2 - c_3) u^2 u^3 \\
u_t^2 = -c_2 u_x^2 - 2(c_1 - c_3) u^1 u^3 \\
u_t^3 = -c_3 u_x^3 - 2(c_2 - c_1) u^1 u^2
\end{cases}$$
(13)

that is a Hamiltonian with operator

$$C^{ij} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \partial_x + \begin{pmatrix} 0 & -2u^3 & 2u^2 \\ 2u^3 & 0 & 2u^1 \\ -2u^2 & -2u^1 & 0 \end{pmatrix}.$$
 (14)

Finally, following the approach introduced by S. I. Tsarev [27], we will see how non-homogeneous hydrodynamic operators arise in a class of systems obtained by the inversion of an evolutionary Hamiltonian equation (see Theorem 4.1). Often, the inversion of such equations leads to a degeneration of the leading coefficient g^{ij} in the first order operator. This is a strong motivation for the investigation of degenerate 1+0 structures.

In this paper, we present a complete classification of Hamiltonian operators for systems in two and three components of the form

$$C^{ij} = g^{ij}\partial_x + b^{ij}_k u^k_x + \omega^{ij}, \qquad (15)$$

focusing on the case when the leading coefficient is degenerate (i.e. its rank is lower than the number of components of the system) and some related remarkable examples of systems in 1+1 dimensions exhibiting this feature. The importance of a deeper study of such operators has been remarked by O. I. Mokhov in [22], who finds Hamiltonian structures of this type in the study of the real reduction of 2-waves interaction system, but also by Dubrovin and Novikon themselves [5].

In section 2, we introduce the conditions for non-homogeneous operators of hydrodynamic type to be Hamiltonian, either with non-degenerate or degenerate assumptions. We establish the connection between such operators and non-homogeneous systems of first order PDEs, introducing the corresponding inverted systems and their associated Hamiltonian structures. In section 3, we show a complete classification, up to diffeomorphisms of the manifold defined by the field variables, of degenerate operators of type 1 + 0 for systems with two and three components. In section 4, we provide several examples with Hamiltonian structures fitting the above mentioned classification, with particular emphasis on inverted Hamiltonian systems.

2 Non-homogeneous hydrodynamic operators

In this section we review non-homogeneous operators of hydrodynamic type, as originally introduced in [5] and further investigated in [23, 16].

Non-homogeneous operators of hydrodynamic type are introduced as the natural generalization of homogeneous Hamiltonian operators (9)

$$C^{ij} = g^{ij}\partial_x + b^{ij}_k u^k_x + \omega^{ij}, \qquad (16)$$

where g^{ij} , b_k^{ij} and ω^{ij} depend on the field variables u. Then, the underlying non-homogeneous local Poisson structure of hydrodynamic type is defined as

$$\{u^{i}(x), u^{j}(y)\} = g^{ij}(u(x))\delta_{x}(x-y) + b_{k}^{ij}(u(x))u_{x}^{k}\delta(x-y) + \omega^{ij}(u(x))\delta(x-y).$$
(17)

Notice that operators of type 1 + 0 are composed of two homogeneous operators $A^{ij} = g^{ij}\partial_x + b^{ij}_k u^k_x$ of order 1, and ω^{ij} of order 0. The conditions for C^{ij} to be Hamiltonian are given by the constraints for each of its homogeneous part to be Hamiltonian (Theorem 2.2 and Theorem 2.1 respectively) and an additional compatibility condition between the two (Theorem 2.3).

We recall that operators of order zero, also known as *ultralocal operators*, are Hamiltonian if the following conditions are satisfied.

Theorem 2.1. [22] The operator $\omega^{ij}(u)$ is Hamiltonian if and only if it forms a finite-dimensional Poisson

structure, i.e. it satisfies the conditions

$$\omega^{ij}(u) = -\omega^{ji}(u), \tag{18}$$

$$\omega^{is} \frac{\partial \omega^{jk}}{\partial u^s} + \omega^{js} \frac{\partial \omega^{ki}}{\partial u^s} + \omega^{ks} \frac{\partial \omega^{ij}}{\partial u^s} = 0.$$
(19)

We remark that in the non-degenerate case, i.e. det $\omega^{ij} \neq 0$, conditions (18) and (19) are respectively skew-symmetry and closedness of the 2-form ω .

In the case of operators of first order the following result holds.

Theorem 2.2.[22] The operator A^{*ij*} is Hamiltonian if and only if

$$g^{ij} = g^{ji} \tag{20}$$

$$\frac{\partial g^{ij}}{\partial u^k} = b_k^{ij} + b_k^{ji} \tag{21}$$

$$g^{is}b_s^{jk} - g^{js}b_s^{ik} = 0 \tag{22}$$

$$g^{is}\left(\frac{\partial b_s^{jr}}{\partial u^k} - \frac{\partial b_k^{jr}}{\partial u^s}\right) + b_s^{ij}b_k^{sr} - b_s^{ir}b_k^{sj} = 0$$
⁽²³⁾

$$g^{is}\frac{\partial b_q^{jr}}{\partial u^s} - b_s^{ij}b_q^{sr} - b_s^{ir}b_q^{js} = g^{js}\frac{\partial b_q^{ir}}{\partial u^s} - b_s^{ji}b_q^{sr} - b_q^{is}b_s^{jr},$$
(24)

$$\sum_{(q,k)} \left\{ \frac{\partial}{\partial u^q} \left(g^{is} \left(\frac{\partial b_s^{jr}}{\partial u^k} - \frac{\partial b_k^{jr}}{\partial u^s} \right) + b_s^{ij} b_k^{sr} - b_s^{ir} b_k^{sj} \right) + \sum_{(i,j,k)} \left(b_q^{si} \left(\frac{\partial b_k^{jr}}{\partial u^s} - \frac{\partial b_s^{jr}}{\partial u^k} \right) \right) \right\} = 0$$
(25)

with the sum over (q, k) and (i, j, k) is on cyclic permutations of the indices.

Let us remark that here there is no assumption about the non-degeneracy properties of metric. The conditions for non-homogeneous operators of hydrodynamic type to be Hamiltonian are shown in the following theorem.

Theorem 2.3.[23, 22] The operator (16) is Hamiltonian if and only if $g^{ij} \partial_x + b^{ij}_k u^k_x$ is Hamiltonian, ω^{ij} is Hamiltonian, and the compatibility conditions are satisfied

$$\Phi^{ijk} = \Phi^{kij},\tag{26}$$

$$\frac{\partial \Phi^{ijk}}{\partial u^r} = \sum_{(i,j,k)} b_r^{si} \frac{\partial \omega^{jk}}{\partial u^s} + \left(\frac{\partial b_r^{ij}}{\partial u^s} - \frac{\partial b_s^{ij}}{\partial u^r}\right) \omega^{sk},$$
(27)

where Φ^{ijk} is the (3,0)-tensor

$$\Phi^{ijk} = g^{is} \frac{\partial \omega^{jk}}{\partial u^s} - b_s^{ij} \,\omega^{sk} - b_s^{ik} \,\omega^{js}.$$
⁽²⁸⁾

3 Classification for systems in two and three components

Savoldi[26] presents a complete classification of degenerate first order homogeneous operators for systems with two and three components. Starting from these results, in this section we provide a novel complete classification of degenerate operators of type 1+0. To obtain an explicit form of ω^{ij} by means of Theorem

2.3 it is sufficient to solve conditions (26) and (27) with fixed tensors g^{ij} and b_k^{ij} , giving rise to an overdetermined system of PDEs. In addition, we require the ultralocal operator ω^{ij} to be Hamiltonian imposing (18) and (19), via Theorem 2.1. In appendix A we report the details of the computations.

The following computations are carried out with the support of computer algebra methods, implemented in Maple, Reduce and Mathematica. The use of symbolic computation for integrable systems and Hamiltonian structures is itself an ongoing topic of research[17, 31].

3.1 Systems in n = 2 components

Let us consider systems with two components, with field variables u, v. In general, given n the number of components of the hydrodynamic system, in the degenerate case the operator g^{ij} can be classified by its rank, with rank $(g^{ij}) = m < n$. In the following, we explicit the number of components n for the operator $C_{n,k}^{ij}$ while the index k is used to distinguish between different operators.

For n = 2, rank (g^{ij}) is in $\{0, 1\}$. The only solution for the case rank $(g^{ij}) = 0$ is the trivial one, then the operator reduces to a symplectic form. In the case rank $(g^{ij}) = 1$, we can construct two different operators,

$$C_{2,1}^{ij} = \begin{pmatrix} \partial_x & 0\\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & f(v)\\ -f(v) & 0 \end{pmatrix},$$
(29)

$$C_{2,2}^{ij} = \begin{pmatrix} \partial_x & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & -\frac{v_x}{u} \\ \frac{v_x}{u} & 0 \end{pmatrix} + \begin{pmatrix} 0 & \frac{f(v)}{u} \\ -\frac{f(v)}{u} & 0 \end{pmatrix},$$
(30)

where f(v) is an arbitrary function depending only on the variable v.

Theorem 3.1. Every degenerate operator of type 1 + 0 in two components can be mapped either onto an ultralocal Hamiltonian operator or onto one between $C_{2,1}^{ij}$ and $C_{2,2}^{ij}$.

Proof. Considering Theorem 2.3, we compute the symplectic structure satisfying (26) and (27) for each degenerate operator of the classification introduced by Savoldi in two components.

3.2 Systems in n = 3 components

Let us consider the case of systems with three components u, v, w, for which the degenerate metric has rank (g^{ij}) in $\{0,1,2\}$. We denote with f, g, h, l arbitrary functions, specifying the explicit dependence on the variables, and with c arbitrary constants.

• rank
$$(g^{ij}) = 0$$

$$C_{3,1}^{ij} = \begin{pmatrix} 0 & w_x & 0 \\ -w_x & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & f(u,v,w) & 0 \\ -f(u,v,w) & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$
(31)

• rank $(g^{ij}) = 1$

$$C_{3,2}^{ij} = \begin{pmatrix} \partial_x & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & f(v,w) & g(v,w) \\ -f(v,w) & 0 & h(v,w) \\ -g(v,w) & -h(v,w) & 0 \end{pmatrix},$$
(32)

where the function f(v, w) is expressed in terms of the functions h(v, w) and g(v, w) as

$$f(v,w) = h(v,w) \left(l(w) + \int_{1}^{v} \frac{g(s,w)\partial_{w}h(s,w) - h(s,w)\partial_{w}g(s,w)}{h(s,w)^{2}} \, ds \right).$$
(33)

$$C_{3,3}^{ij} = \begin{pmatrix} \partial_x & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & w_x & 0 \\ -w_x & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & f(v,w) & 0 \\ -f(v,w) & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$
(34)

$$C_{3,4}^{ij} = \begin{pmatrix} \partial_x & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & -\frac{w_x}{u} \\ 0 & 0 & 0 \\ \frac{w_x}{u} & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & \frac{f(v,w)}{u} \\ 0 & 0 & 0 \\ -\frac{f(v,w)}{u} & 0 & 0 \end{pmatrix},$$
(35)

$$C_{3,5}^{ij} = \begin{pmatrix} \partial_x & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & -\frac{v_x}{u} & -\frac{w_x}{u} \\ \frac{v_x}{u} & 0 & 0 \\ \frac{w_x}{u} & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & \frac{f(v,w)}{u} & \frac{g(v,w)}{u} \\ -\frac{f(v,w)}{u} & 0 & \frac{h(v,w)}{u} \\ -\frac{g(v,w)}{u} & -\frac{h(v,w)}{u} & 0 \end{pmatrix},$$
(36)

with f(v, w) given in (33).

• rank $(g^{ij}) = 2$

$$C_{3,6}^{ij} = \begin{pmatrix} \partial_x & 0 & 0 \\ 0 & \partial_x & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & f(w) & g(w) \\ -f(w) & 0 & cg(w) \\ -g(w) & -cg(w) & 0 \end{pmatrix},$$
(37)

$$C_{3,7}^{ij} = \begin{pmatrix} \partial_x & 0 & 0 \\ 0 & \partial_x & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -\frac{w_x}{v} \\ 0 & \frac{w_x}{v} & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & cf(w) \\ 0 & 0 & \frac{(1-cu)f(w)}{v} \\ -cf(w) & -\frac{(1-cu)f(w)}{v} & 0 \end{pmatrix},$$
(38)

$$C_{3,8}^{ij} = \begin{pmatrix} \partial_x & 0 & 0 \\ 0 & \partial_x & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & -\frac{ww_x}{uw-v} \\ 0 & 0 & \frac{w_x}{uw-v} \\ \frac{ww_x}{uw-v} & -\frac{w_x}{uw-v} & 0 \end{pmatrix}$$

$$+ (1+w^2) f(w) \begin{pmatrix} 0 & \frac{1}{(1+w^2)} & \frac{(w-cv\sqrt{1+w^2})}{uw-v} \\ -\frac{1}{(1+w^2)} & 0 & -\frac{(1-cu\sqrt{1+w^2})}{uw-v} \\ -\frac{(w-cv\sqrt{1+w^2})}{uw-v} & \frac{(1-cu\sqrt{1+w^2})}{uw-v} & 0 \end{pmatrix},$$
(40)

$$C_{3,9}^{ij} = \begin{pmatrix} 0 & \partial_x & 0 \\ \partial_x & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & f(w) & cg(w) \\ -f(w) & 0 & g(w) \\ -cg(w) & -g(w) & 0 \end{pmatrix},$$
(41)

$$C_{3,10}^{ij} = \begin{pmatrix} 0 & \partial_x & 0 \\ \partial_x & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & -\frac{w_x}{v} \\ 0 & 0 & 0 \\ \frac{w_x}{v} & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & f(w) & \frac{h(w) - ug(w)}{v} \\ -f(w) & 0 & g(w) \\ -\frac{h(w) - ug(w)}{v} & -g(w) & 0 \end{pmatrix},$$
(42)

with the additional condition

$$h(w)g'(w) - g(w)(f(w) + h'(w)) = 0,$$
(43)

$$C_{3,11}^{ij} = \begin{pmatrix} 0 & \partial_x & 0 \\ \partial_x & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & \frac{w_x}{uw - v} \\ 0 & 0 & -\frac{ww_x}{uw - v} \\ -\frac{w_x}{uw - v} & \frac{ww_x}{uw - v} & 0 \end{pmatrix}$$

$$+ f(w) \begin{pmatrix} 0 & \frac{c}{\sqrt{w}} & \frac{(uw - 2c\sqrt{w})}{uw - v} \\ -\frac{c}{\sqrt{w}} & 0 & -\frac{w(v - 2c\sqrt{w})}{uw - v} \\ -\frac{(uw - 2c\sqrt{w})}{uw - v} & w \frac{w(v - 2c\sqrt{w})}{uw - v} & 0 \end{pmatrix}$$
(44)

Remark 3.1. Condition (43) can be explicitly solved with respect to any function among f, g and h.

Theorem 3.2. Every degenerate operator of type 1 + 0 in three components can be mapped either onto an ultralocal operator satisfying the closure relation, or onto one among $C_{3,k}^{ij}$ with k = 1,...,11.

Proof. Imposing the conditions on the operators to be Hamiltonian, we obtain the extension of the classification for degenerate first order operators presented by Savoldi in three components [26]. See Appendix A for more details.

Remark 3.2. In the proposed classification, we have considered three arbitrary functions for the sake of generality and in view of possible relevance for applications. However, we emphasise that changes of variables can simplify the form of operators. To do so one should look for those changes of variables that leave the order 1 operator invariant, then apply them to the order 0 one.

4 Applications

In this section we present some examples of non-homogeneous quasilinear systems with degenerate Hamiltonian structure of order 1 + 0 in two and three components.

Example 4.1 (2-wave interaction system). Mokhov[22] studies the real reduction of 2-waves interaction system formulated in terms of the system of hydrodynamic equations in two field variables u = u(x, t) and v = v(x, t)

$$\begin{cases} u_t = auv\\ v_t = av_x + u^2 \end{cases}, \tag{45}$$

with a constant. The system admits a Hamiltonian formulation, with the operator

$$C^{ij} = \begin{pmatrix} 0 & 0 \\ 0 & \partial_x \end{pmatrix} + \begin{pmatrix} 0 & -u \\ u & 0 \end{pmatrix},$$
(46)

and the Hamiltonian functional

$$H = \frac{1}{2} \int \left(av^2 - u^2 \right) dx.$$
 (47)

The 1 + 0 operator (46) is degenerate, since the rank of the order 1 term is lower than the number of components of the system. Moreover, by applying the exchange $u \leftrightarrow v$ it is evident that the operator found by Mokhov is of type $C_{2,1}^{ij}$ in (29).

Example 4.2 (Sinh-Gordon equation). Let us consider the Sinh-Gordon equation

$$\varphi_{\tau\xi} = \sinh \varphi. \tag{48}$$

Applying the change of variables $\varphi = 2 \log u$, we have

$$\left(2\frac{u_{\tau}}{u}\right)_{\xi} = \frac{1}{2}\left(u^2 - \frac{1}{u^2}\right).$$
(49)

Introducing $v = 2u_{\tau}/u$ and considering the light-cone coordinates $\tau = t, \xi = t - x$

$$\begin{cases} u_t = \frac{1}{2}uv \\ v_t = v_x + \frac{1}{2}\left(u^2 - \frac{1}{u^2}\right) \end{cases},$$
(50)

we show that the system is Hamiltonian with the non-homogeneous hydrodynamic operator of shape (29) with the exchange of variables $u \leftrightarrow v$ and the function f(u) = u/2

$$C^{ij} = \begin{pmatrix} 0 & 0 \\ 0 & \partial_x \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 & u \\ -u & 0 \end{pmatrix}.$$
 (51)

The corresponding Hamiltonian density is

$$h(u,v) = \frac{1}{2} \left(v^2 - u^2 + \frac{1}{u^2} \right).$$
(52)

4.1 Inverted Hamiltonian systems

In this section we show the connection between degenerate operators of type 1 + 0 and scalar equations possessing a local Hamiltonian structure.

Let us briefly recall that the momentum of a Hamiltonian equation $u_t = A^{ij} \delta H / \delta u^j$ is a functional defining a *x*-translation

$$u_x^i = A^{ij} \frac{\delta P}{\delta u^j}, \quad \text{with } P = \int p(u, u_\sigma) dx,$$

for i = 1,...,n. Tsarev[27] proves that under the inversion of the independent variables x and t, the Hamiltonian property is preserved by the system. It is well known that the momentum is a conserved quantity in a Hamiltonian system, hence there exists $q(u, u_{\sigma})$ such that $p_t = q_x$. Then, one can choose $H' = \int q(u, u_{\sigma}) dt$ as the Hamiltonian functional of the inverted system.

Non-homogeneous operators of hydrodynamic type are related to the study of scalar evolutionary equations possessing a *local* Hamiltonian structure. Indeed, by introducing the new set of variables

$$u^{1} = u, \ u^{2} = u_{x}, \ u^{3} = u_{xx}, \ \dots,$$
 (53)

it is in some cases possible to write an equivalent non-homogeneous quasilinear system that can be seen as evolutionary with respect to the independent variable *x*, obtaining the *inverted system*.

Remark 4.1. Let us observe that every invertible system of order k has the form

$$u_t = F_1(u, u_x, \dots, u_{(k-1)x}) + F_2(u, u_x, \dots, u_{(k-1)x})u_{kx},$$
(54)

where F_1 , F_2 are arbitrary functions. Note that this is the case of KdV and many other examples in nonlinear phenomena. Indeed, considering the lower derivatives as parameters, we need the system to be linear in u_{kx} in order to conserve linearity in u_t once inverted.

The following result offers an explicit connection between non-homogeneous hydrodynamic operators and inverted systems.

Proposition 4.1. Let us consider the evolutionary equation $u_t = F(u, u_\sigma)$ endowed with a local Hamiltonian structure and a momentum density p in (4.1) depending on u only. Then, if the inverted system in the set of variables (53) admits a local Hamiltonian structure, this is given in terms of a non-homogeneous operator of hydrodynamic type.

Proof. We observe the following

$$q_x = p_t = p_u(u)u_t = p_u(u)F(u,u_\sigma), \quad \sigma \le k,$$
(55)

where p_t is of order $\leq k$, at most equal to the order of the equation, and so is q_x . Hence, $q(u, u_\sigma)$ is of order at most k - 1. This implies that the Hamiltonian $H' = \int q(u^1, \dots, u^{k-1}) dt$ is of hydrodynamic type for the inverted system in the new variables. In [27, 15], it has been proved that the Hamiltonian property is preserved after a change of dependent variables and an inversion of t and x. Then, the inverted system is

quasilinear of first order and already Hamiltonian. The operator B^{ij} in

$$u_x^i = B^{ij} \frac{\delta H'}{\delta u^j},\tag{56}$$

being local, must be of type 1 + 0, i.e. a non-homogeneous operator of hydrodynamic type.

The Proposition 4.1 justifies a deeper investigation of such operators, for which KdV offers a leading example, as follows. We emphasise the previous theorem does not guarantee that the operator is in general non-degenerate.

Example 4.3 (KdV equation - I). Let us consider the KdV equation

$$u_t = 6uu_x + u_{xxx}, \tag{57}$$

which is widely known to be Hamiltonian. Inverting the equation, we obtain the evolutionary system with respect to x in three components $u^1(x,t)$, $u^2(x,t)$, $u^3(x,t)$ defined as $u = u^1$, $u_x = u^2$, $u_{xx} = u^3$, yielding the following non-homogeneous system of hydrodynamic type

$$\begin{cases} u_x^1 = u^2 \\ u_x^2 = u^3 \\ u_x^3 = u_t^1 + 6u^1 u^2 \end{cases}$$
(58)

This system is Hamiltonian with the following non-homogeneous hydrodynamic type operator [27]

$$C^{ij} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \partial_t + \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 6u^1 \\ 0 & -6u^1 & 0 \end{pmatrix},$$
(59)

with the leading coefficient g^{ij} being degenerate. It is easy to see that applying the change of variables $u^1 = w$, we obtain again the operator (32), where

$$g(v, w) = 0,$$
 $f(v, w) = h(v, w) l(w),$
 $l(w) = 6w,$ $h(v, w) = -1.$

Example 4.4 (KdV equation - II). Mokhov [21] finds a transformation of variables (also known as *local quadratic unimodular change*)

$$u^{1} = \frac{w^{1} - w^{3}}{\sqrt{2}}, \qquad u^{2} = w^{2}, \qquad u^{3} = \frac{w^{1} + w^{3}}{\sqrt{2}} + \left(w^{1} - w^{3}\right)^{2},$$
 (60)

such that the KdV equation reads as

$$\begin{cases} w_x^1 = -\frac{1}{2} \left(w^1 - w^3 \right)_t + w^2 \left(w^1 - w^3 \right) + \frac{1}{\sqrt{2}} w^2 \\ w_x^2 = \left(w^1 - w^3 \right)^2 + \frac{1}{\sqrt{2}} \left(w^1 + w^3 \right) \\ w_x^3 = -\frac{1}{2} \left(w^1 - w^3 \right)_t + w^2 \left(w^1 - w^3 \right) - \frac{1}{\sqrt{2}} w^2 \end{cases}$$
(61)

After this local change, the KdV is a bi-Hamiltonian system with respect to two non-homogeneous operators 1 + 0 of hydrodynamic type, one of these being the operator

$$C^{ij} = \frac{1}{2} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} \partial_t + \begin{pmatrix} 0 & w^1 - w^3 + \frac{1}{\sqrt{2}} & 0 \\ w^3 - w^1 - \frac{1}{\sqrt{2}} & 0 & w^3 - w^1 + \frac{1}{\sqrt{2}} \\ 0 & w^1 - w^3 - \frac{1}{\sqrt{2}} & 0 \end{pmatrix},$$
 (62)

which is degenerate, since rank $(g^{ij}) = 1$. The Hamiltonian given in terms of the new variables is

$$H = \int \left((w^1)^2 - (w^2)^2 - (w^3)^2 \right) dx.$$
(63)

To show that the obtained operator is indeed one of those classified above, we consider a new change of variables

$$w^{1} = \frac{\bar{u}^{1} - \bar{u}^{3}}{\sqrt{2}}, \qquad w^{2} = \bar{u}^{2}, \qquad w^{3} = \frac{\bar{u}^{3} - \bar{u}^{1}}{\sqrt{2}}.$$
 (64)

The degenerate first order operator is written with the leading coefficient $\bar{g} = d\bar{u}^1 \otimes d\bar{u}^1$ and the skew-symmetric bivector

$$\bar{\omega} = -\sqrt{2}\bar{u}^3 \left(d\bar{u}^1 \wedge d\bar{u}^2 - d\bar{u}^2 \wedge d\bar{u}^3 \right).$$

The operator (62) is of type $C_{2,2}^{ij}$ in three components showed in (32). In particular,

$$g(v,w) = 0, \qquad f(v,w) = l(w)h(v,w),$$
$$l(w) = -1, \qquad h(v,w) = \sqrt{2}w.$$

Example 4.5 (Generalised KdV equation). Let us consider the generalised KdV equation

$$u_t + 3(n+1)u^n u_x + u_{xxx} = 0 ag{65}$$

where *n* is a positive integer. It is known that (65) is Hamiltonian with the operator ∂_x for any *n*. The case n = 2 corresponds to the modified KdV equation (mKdV), it is integrable and it has a second Hamiltonian structure, with operator $\partial_x^3 + 6 \partial_x u \partial_x^{-1} u \partial_x$. The Hamiltonians associated with mKdV are

$$H_1 = \int \left(\frac{3}{4}u^4 + \frac{1}{2}u_x^2\right) dx, \qquad H_2 = \int \frac{1}{2}u^2 dx.$$
(66)

In (65) we introduce the variables $u^1 = u$, $u^2 = u_x$, $u^3 = u_{xx}$, so that the equation reads as a quasilinear system of first order PDEs

$$u_x^1 = u^2, \qquad u_x^2 = u^3, \qquad u_x^3 = -u_t^1 - 3(n+1)(u^1)^n u^2.$$
 (67)

The Hamiltonian structure is still conserved after the scalar equation is transformed into a system, i.e. (67) has Hamiltonian structure with the operator

$$C^{ij} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \partial_t + \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & -3(n+1)(u^1)^{n-1} \\ 0 & 3(n+1)(u^1)^{n-1} & 0 \end{pmatrix}$$
(68)

and the Hamiltonian functional

$$H = \int \left(3(u^1)^{n+1} - u^1 u^3 + \frac{(u^2)^2}{2} \right) dx.$$
 (69)

The operator (68) is $C_{3,2}^{ij}$ in (32) with the exchange $u^1 \leftrightarrow u^3$ and

$$g(u^{1}, u^{2}) = 0, \qquad f(u^{1}, u^{2}) = h(u^{1}, u^{2}) l(u^{1}),$$

$$l(u^{1}) = 3(n+1)(u^{1})^{n-1}, \qquad h(u^{1}, u^{2}) = -1.$$

Remark 4.2. Let us observe that for n > 2 the generalised KdV equation is not integrable, even if it is Hamiltonian as proved in the previous example. We emphasise that this feature is more general than the integrability property.

We finally present two examples violating the hypothesis of locality, either in terms of the momentum or of the operator defined for the inverted Hamiltonian structure.

Example 4.6. We consider the linearised KdV equation

$$u_t = u_{xxx}, \tag{70}$$

for which the inverted system is easily given in the new variables by

$$u_x^1 = u^2 \quad u_x^2 = u^3 \quad u_x^3 = u_t^1.$$
(71)

The associated momentum is given in terms of the density $p(u) = \partial_x^{-2} u$. Here again, it is not possible to write the resulting system with Hamiltonian operator of type 1 + 0 [27].

Example 4.7. We consider the Harry-Dym equation [16, 25]

$$u_t = \left(\frac{1}{\sqrt{u}}\right)_{xxx} = -\frac{15}{8}u^{-7/2} (u_x)^3 + \frac{9}{4}u^{-5/2} u_x u_{xx} - \frac{1}{2}u^{-3/2} u_{xxx},$$
(72)

admitting the Hamiltonian structures

$$u_t = A_1 \frac{\delta H_1}{\delta u} = -\frac{1}{2} \partial_x^3 \frac{\delta H_1}{\delta u}, \quad \text{with } H_1 = -\int 4\sqrt{u} \, dx, \tag{73}$$

$$u_t = A_2 \frac{\delta H_2}{\delta u} = -(2u\partial_x - u_x) \frac{\delta H_2}{\delta u}, \quad \text{with } H_2 = -\int \left(\frac{15}{32}u^{-7/2}u_x - \frac{1}{16}u^{-5/2}u_{xx}\right) dx.$$
(74)

Introducing the variables $u_x = u^2$, $u_{xx} = u^3$, the inverted system is

$$u_x^1 = u^2$$
, $u_x^2 = u^3$, $u_x^3 = -2(u^1)^{3/2}u_t^1 - \frac{15}{4}(u^1)^{-2}(u^2)^3 + \frac{9}{2}u^1u^2u^3$. (75)

The momentum *P* associated with the operator A_2 is

$$u_x = -(2u\partial_x - u_x)\frac{\delta P}{\delta u}, \qquad P = \int p(u)dx = -\int u\,dx,$$

and following the above procedure the Hamiltonian H' as a functional in the new variables is

$$H' = -\int \left(\frac{3}{4}(u^1)^{-5/2}(u^2)^2 - \frac{1}{2}(u^1)^{-3/2}u^3\right) dx.$$
 (76)

With this Hamiltonian it is not possible to build a local operator of the form 1 + 0 for the inverted system, hence this operator will be non-local.

5 Conclusions

The study of non-homogeneous quasilinear systems of first order PDEs is an ongoing research topic in integrable systems and Hamiltonian PDEs. To the authors' knowledge, a general criterion to discuss integrability for this kind of systems is not currently known, unlike the homogeneous systems [28]. This paper represents a first step towards the investigation of integrability of non-homogeneous systems, focusing on the Hamiltonian property. The study of possible bi-Hamiltonian structures associated with these type will be the subject of a future paper. Non-homogeneous operators of order k + m play an important role in non-linear phenomena and their investigation represents another interesting topic [18, 16, 7, 13, 20]. Even in the simplest case where k = 1, and m = 0 we showed how the conditions for the operator to be Hamiltonian lead to a specific form, this being exactly solvable. Higher order operators require a more general study, especially for what concerns the degenerate case.

As future perspectives, the authors emphasise the necessity to further investigate the integrability of non-homogeneous quasilinear systems, the compatibility conditions between systems and operators in the sense of [29], but also their associated geometric structure, following the lead of the homogeneous case, where both operators and systems are linked to projective algebraic geometry [1, 11, 12, 30] and differential Riemannian geometry. Finally, the discrete analogous of non-homogeneous operators were introduced by Dubrovin in [3], letting the classification method described in this paper suitable for discrete operators as well.

Acknowledgements The authors thank C. Benassi, F. Coppini, E. V. Ferapontov, A. Moro, M. Pavlov and R. Vitolo for stimulating discussions and interesting remarks.

PV also acknowledges the financial support of GNFM of the Istituto Nazionale di Alta Matematica, of PRIN 2017 "Multiscale phenomena in Continuum Mechanics: singular limits, off-equilibrium and transitions", project number 2017YBKNCE and the research project Mathematical Methods in Non Linear Physics (MMNLP) by the Commissione Scientifica Nazionale – Gruppo 4 – Fisica Teorica of the Istituto Nazionale di Fisica Nucleare (INFN).

MD would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme "Dispersive hydrodynamics: mathematics, simulation and experiments, with applications in nonlinear waves" (HDY2) where work on this paper was undertaken. This work was supported by EPSRC grant no EP/R014604/1.

References

- [1] Agafonov, S. I. and Ferapontov, E. V. Systems of conservation laws in the context of the projective theory of congruences. *Izvestiya: Mathematics*, 60(6):1097–1122, dec 1996.
- [2] Dorfman, I. Y. and Mokhov, O. I. Local symplectic operators and structures related to them. *Journal of Mathematical Physics*, 32(12):3288–3296, 1991.
- [3] Dubrovin, B. A. Differential-geometric Poisson brackets on a lattice. *Functional Analysis and Its Applications*, 23(2):131–133, 1989.
- [4] Dubrovin, B. A. and Novikov, S. P. Hamiltonian formalism of one-dimensional systems of hydrodynamic type and the Bogolyubov–Whitham averaging method. *Soviet Math. Dokl*, 3(27):665–669, 1983.
- [5] Dubrovin, B. A. and Novikov, S. P. Poisson brackets of hydrodynamic type. *Soviet Math. Dokl*, 30:651–654, 1984.
- [6] Dubrovin, B. A., Krichever, I. M. and Novikov, S. P. Integrable systems. I. *Dynamical Systems IV: Symplectic Geometry and its Applications, Encyclopaedia of Mathematical Sciences*, 4:173–280, 2001.
- [7] Falqui, G. On a Camassa–Holm type equation with two dependent variables. *Journal of Physics A: Mathematical and General*, 39(2):327–342, dec 2005.
- [8] Ferapontov, E. and Marshall, D. Differential-geometric approach to the integrability of hydrodynamic chains: the Haantjes tensor. *Math. Ann.*, 339, 2007.
- [9] Ferapontov, E., Moro, A., Sokolov, V. Hamiltonian systems of hydrodynamic type in 2+1 dimensions. *Commun. Math. Phys.*, 285:31–65, 2008.
- [10] Ferapontov, E. V., Khusnutdinova, K. R. On integrability of (2+1)-dimensional quasilinear systems. *Commun. Math. Phys.*, 248:187–206, 2004.
- [11] Ferapontov, E. V., Pavlov, M. V. and Vitolo, R. F. Projective-geometric aspects of homogeneous thirdorder Hamiltonian operators. *Journal of Geometry and Physics*, 85:16–28, 2014.
- [12] Ferapontov, E. V., Pavlov, M. V. and Vitolo, R. F. Systems of conservation laws with third-order Hamiltonian structures. *Letters in Mathematical Physics*, 108(06):1525–1550, 2018.
- [13] Jannelli, A., Manganaro, N. and Rizzo, A. Riemann problems for the nonhomogeneous Aw-Rascle model. *Communications in Nonlinear Science and Numerical Simulation*, 118:107010, 2023.
- [14] Kersten, P., Krasil'shchik, I. S. and Verbovetsky, A. M. Hamiltonian operators and *l**-coverings. *Journal of Geometry and Physics*, 50:273–302, 2004.
- [15] Kersten, P., Krasil'shchik, I. S., Verbovetsky, A. M. and Vitolo, R. F. Systems of conservation laws with third-order Hamiltonian structures. *Differential Equations - Geometry, Symmetries and Integrability: The Abel Symposium 2008, Kruglikov, B., Lychagin, V. and Straume, E.,* 108:187–198, 2009.
- [16] Konyaev, A. Geometry of inhomogeneous Poisson brackets, multicomponent Harry Dym hierarchies and multicomponent Hunter-Saxton equations. arXiv:2112.05635, 2021.

- [17] Krasil'shchik, I. S., Verbovetsky, A. M. and Vitolo, R. F. The symbolic computation of integrability structures for partial differential equations. *Texts and Monographs in Symbolic Computation. Springer. ISBN 978-3-319-71654-1; see http://gdeq.org/Symbolic_Book for downloading program files that are discussed in the book, 2018.*
- [18] Lorenzoni, P., Savoldi, A. and Vitolo, R. F. Bi-Hamiltonian structures of KdV type. *Journal of Physics* A: Mathematical and Theoretical, 51(4), 2017.
- [19] Magri, F. A simple model of the integrable Hamiltonian equation. *Journal of Mathematical Physics*, 19(5):1156–1162, 1978.
- [20] Manganaro, N., Rizzo, A. Riemann Problems and Exact Solutions for the p-System. *Mathematics*, 10(6), 2022.
- [21] Mokhov, O. I. Symplectic and Poisson Geometry on Loop Spaces of Manifolds and Nonlinear Equations, arXiv:hep-th/9503076. *High Energy Physics - Theory*, 1995.
- [22] Mokhov, O. I. Symplectic and Poisson geometry on loop spaces of smooth manifolds, and integrable equations. *Russian Mathematical Surveys*, 53(3):515–622, 1998.
- [23] Mokhov, O. I. and Ferapontov, E. V. Hamiltonian pairs associated with skew-symmetric Killing tensors on spaces of constant curvature. *Functional Analysis and Its Applications*, 28(2):123–125, 1994.
- [24] Novikov, S., Manakov, S.V., Pitaevskii, L.P. and Zakharov, V.E. Theory of Solitons: The Inverse Scattering Method. Monographs in Contemporary Mathematics. Springer US, 1984.
- [25] Pedroni, M., Sciacca, V. and Zubelli, J. On the Bi-Hamiltonian Theory for the Harry Dym Equation. *Theoretical and Mathematical Physics*, 133, 01 2002.
- [26] Savoldi, A. On deformations of one-dimensional Poisson structures of hydrodynamic type with degenerate metric. *Journal of Geometry and Physics*, 104:246–276, 2016.
- [27] Tsarev, S. P. The hamiltonian property of stationary and inverse equations of condensed matter mechanics and mathematical physics. *Mathematical notes of the Academy of Sciences of the USSR*, 46(1):569–573, 1989.
- [28] Tsarev, S. P. The geometry of Hamiltonian systems of hydrodynamic type. The generalised hodograph method. *Mathematics of the USSR-Izvestiya*, 37(2):397–419, 1991.
- [29] Vergallo, P. and Vitolo, R. F. Homogeneous Hamiltonian operators and the theory of coverings. *Differential Geometry and its Applications*, 75:101713, 2021.
- [30] Vergallo, P. and Vitolo, R. F. Projective geometry of homogeneous second order Hamiltonian operators. arXiv:2203.04237., 2022.
- [31] Vitolo, R. F. Computing with Hamiltonian operators. *Computer Physics Communications*, 244:228–245, 2019.

A Derivation of the operators in Theorem 3.1 and Theorem 3.2

We give the details of the procedure followed to compute the classifications of Section 3. The computations are carried out with the support of computer algebra systems (Maple, Reduce and Mathematica) and finally checked by hand.

For the sake of simplicity, we describe the first nontrivial operator obtained in three components, in the text identified as $C_{3,2}$. We start by considering the degenerate operator of order 1 for a system in three components in the Savoldi's classification [26]

$$g^{ij} = \begin{pmatrix} \partial_x & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad b^{ij}_k = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad \forall k \in \{1, 2, 3\}.$$
(77)

We add to the operator an order 0 operator

$$\begin{pmatrix} \partial_x & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} \omega^{11} & \omega^{12} & \omega^{13} \\ \omega^{21} & \omega^{22} & \omega^{23} \\ \omega^{31} & \omega^{32} & \omega^{33} \end{pmatrix}.$$
 (78)

Being ω an ultralocal tail, its elements ω^{ij} are functions at most depending on the three variables of the system u, v, w.

The operator (78) as a whole is Hamiltonian if its parts are Hamiltonian and the compatibility conditions established in Theorem II.3 are satisfied. In particular, the ultralocal term ω is Hamiltonian if it fulfills the Theorem II.1. We reduce the number of free functions in (78) by using the skew-symmetry property

$$\begin{pmatrix} \partial_x & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & \omega^{12} & \omega^{13} \\ -\omega^{12} & 0 & \omega^{23} \\ -\omega^{13} & -\omega^{23} & 0 \end{pmatrix},$$
(79)

so that the unknown functions are

$$\omega^{12}(u,v,w), \quad \omega^{13}(u,v,w), \quad \omega^{23}(u,v,w).$$
 (80)

To implement the constraints expressed in Theorem II.3, we evaluate the tensor Φ^{ijk} for the case (77)

$$\Phi^{ijk} = g^{is} \frac{\partial \omega^{jk}}{\partial u^s} - b_s^{ij} \omega^{sk} - b_s^{ik} \omega^{js} = g^{is} \frac{\partial \omega^{jk}}{\partial u^s},$$
(81)

where the sum over *s* is intended via repeated indices. For three components s = 1, 2, 3 and we use the notation for the variables $u^1 = u$, $u^2 = v$, $u^3 = w$. The only non-zero element in *g* is $g^{11} = 1$, hence the constraint (26) on the non-zero elements of the tensor Φ^{ijk} takes the form $\Phi^{1jk} = \Phi^{k1j}$, with

$$\Phi^{1jk} = \begin{pmatrix} 0 & \frac{\partial \omega^{12}}{\partial u} & \frac{\partial \omega^{13}}{\partial u} \\ -\frac{\partial \omega^{12}}{\partial u} & 0 & \frac{\partial \omega^{23}}{\partial u} \\ -\frac{\partial \omega^{13}}{\partial u} & -\frac{\partial \omega^{23}}{\partial u} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ \frac{\partial \omega^{12}}{\partial u} & 0 & 0 \\ \frac{\partial \omega^{13}}{\partial u} & 0 & 0 \end{pmatrix} = \Phi^{k1j}.$$
(82)

The constraints on the field variables are

$$\frac{\partial \omega^{ij}(u,v,w)}{\partial u} = 0 \implies \omega^{ij}(u,v,w) = \omega^{ij}(v,w), \tag{83}$$

i.e. they do not depend on the variable *u*. We introduce the notation

$$\begin{cases} \omega^{12}(v,w) = f(v,w) \\ \omega^{13}(v,w) = g(v,w) \\ \omega^{23}(v,w) = h(v,w) \end{cases}$$
(84)

The constraint (27) yields

$$\frac{\partial^2 \omega^{ij}}{\partial u^2} = 0, \quad \frac{\partial^2 \omega^{ij}}{\partial u \partial v} = 0, \quad \frac{\partial^2 \omega^{ij}}{\partial u \partial w} = 0, \tag{85}$$

not producing any further restriction for the form of the functions, given (83). Finally, the closure requirement (19) is

$$\omega^{12} \frac{\partial \omega^{23}}{\partial v} - \omega^{23} \frac{\partial \omega^{12}}{\partial v} + \omega^{13} \frac{\partial \omega^{23}}{\partial w} - \omega^{23} \frac{\partial \omega^{13}}{\partial w} = 0,$$
(86)

with the notation (84) this becomes

$$f(v,w)\frac{\partial h(v,w)}{\partial v} - h(v,w)\frac{\partial f(v,w)}{\partial v} + g(v,w)\frac{\partial h(v,w)}{\partial w} - h(v,w)\frac{\partial g(v,w)}{\partial w} = 0.$$
(87)

We solve the last constraint with respect to the field f(v, w). Observing that

$$\frac{\partial}{\partial v} \left(\frac{f}{h} \right) = \frac{1}{h} \frac{\partial f}{\partial v} - \frac{f}{h^2} \frac{\partial h}{\partial v}, \tag{88}$$

we obtain the expression for f(v, w) given in (33). The operator is then

$$C_{3,2}^{ij} = \begin{pmatrix} \partial_x & f(v,w) & g(v,w) \\ -f(v,w) & 0 & h(v,w) \\ -g(v,w) & -h(v,w) & 0 \end{pmatrix}.$$

We remark that for fixed rank of g^{ij} the resulting conditions strongly depend on the structure of b_k^{ij} . For instance, for the operator $C_{3,3}^{ij}$ we have the same operator g^{ij} as in (77), but different b_k^{ij}

$$b_1^{ij} = b_2^{ij} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad b_3^{ij} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$
(89)

We look for the corresponding operator ω after considering the skew-symmetry property, hence we have three field variables ω^{12} , ω^{13} , ω^{23} .

The first conditions are imposed by comparing the tensors

$$\Phi^{1jk} = \begin{pmatrix} 0 & -\omega^{13} + \frac{\partial \omega^{12}}{\partial u} & \frac{\partial \omega^{13}}{\partial u} \\ \omega^{13} - \frac{\partial \omega^{12}}{\partial u} & 0 & \frac{\partial \omega^{23}}{\partial u} \\ -\frac{\partial \omega^{13}}{\partial u} & -\frac{\partial \omega^{23}}{\partial u} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ -\omega^{13} + \frac{\partial \omega^{12}}{\partial u} & \omega^{23} & 0 \\ \frac{\partial \omega^{13}}{\partial u} & 0 & 0 \end{pmatrix} = \Phi^{k1j}.$$
(90)

At this stage we can already reduce the number of free functions, since $\omega^{23} = 0$. Solving the remaining equations, we find the dependence of the fields ω^{12} , ω^{13} on the variables *u*, *v*, *w*

$$\omega^{12}(u, v, w) = f(v, w) + ug(v, w)$$

$$\omega^{13}(u, v, w) = g(v, w).$$
(91)

The constraints from the relation (19) further reduce the number of free functions, in particular g(v, w) = 0. The corresponding operator is then

$$C_{3,3}^{ij} = \begin{pmatrix} \partial_x & w_x + f(v,w) & 0 \\ -w_x - f(v,w) & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The same procedure has been carried out for all the possible forms of operators g^{ij} and b_k^{ij} , obtaining the above mentioned classification.