Estimation of Heterogeneous Treatment Effects Using a Conditional

Moment Based Approach

Xiaolin Sun¹

¹Department of Economics, Simon Fraser University

October 4, 2024

Abstract

We propose a new estimator for heterogeneous treatment effects in a partially linear model (PLM) with multiple exogenous covariates and a potentially endogenous treatment variable. Our approach integrates a Robinson transformation to handle the nonparametric component, the Smooth Minimum Distance (SMD) method to leverage conditional mean independence restrictions, and a Neyman-Orthogonalized first-order condition (FOC). By employing regularized model selection techniques like the Lasso method, our estimator accommodates numerous covariates while exhibiting reduced bias, consistency, and asymptotic normality. Simulations demonstrate its robust performance with diverse instrument sets compared to traditional GMM-type estimators. Applying this method to estimate Medicaid's heterogeneous treatment effects from the Oregon Health Insurance Experiment reveals more robust and reliable results than conventional GMM approaches.

Keywords: Endogeneity; Conditional mean independence; Neyman-orthogonal moments; Lasso.

JEL Classification: C13; C14.

1 Introduction

Heterogeneous treatment effects have been a central focus in the literature since Imbens and Angrist (1994). These models explore how the effects of policies, programs, or other interventions on a specific outcome may vary across individuals with distinct characteristics. To quantify heterogeneous treatment effects, researchers often augment standard linear models by including interaction terms between the treatment variable and relevant covariates (as discussed in Section 21.4, Chapter 21 of Wooldridge (2010) and Section 7.6, Chapter 7 of Imbens and Rubin (2015), Page 125). This extended model becomes linear in the treatment variable, the interaction term, and the covariates, with the parameters of interest typically encompassing the effects of the treatment and its interaction with some covariates. Furthermore, significant attention has been devoted to addressing unobserved heterogeneity (Wooldridge (1997), Wooldridge (2004)). In this paper, we adopt a standard and widely used semi-parametric model to demonstrate the advantages of integrating a conditional moment-based approach into the estimation of the observed heterogeneous treatment effects, that is, the parameters in front of the treatment and the interaction terms.

Our work introduces a novel estimator, D-RSMD, extending the R-SMD estimator proposed by Antoine and Sun (2021) to accommodate big data settings. Specifically, the D-RSMD estimator combines the debiased method with R-SMD, integrating these approaches for improved estimation procedures. This new estimator offers two key properties essential for applied researchers. While versatile and applicable across various scenarios, we specifically focus on estimating heterogeneous treatment effects to illustrate those properties. We look into the case where the outcome variable adopts a classic partially linear structure encompassing linear and non-parametric components. Within the linear part, the treatment variable is endogenous, and the interaction terms are the treatment multiplied by exogenous covariates.

The primary advantage of the D-RSMD estimator is its reliance on a single valid instrument, bypassing the need to explicitly model the relationship between treatment and the instrument. This eliminates the requirement to generate additional exogenous variables, assume specific functional forms for the first stage (as in 2SLS), or select moments for GMM estimation. The choice of generated variables, model specifications in the first stage, or moments significantly impacts estimation results and the asymptotic properties of treatment effects. This vital advantage is shared by Smooth Minimum Distance (SMD) estimators like the SMD estimator proposed and examined in Lavergne and Patilea (2013) and the R-SMD estimator introduced by Antoine and Sun (2021). Such an approach is crucial for avoiding model specification issues and ensuring robust estimation of treatment effects.

For example, consider a scenario where the treatment variable is endogenous and interacts with a covariate. The traditional GMM method typically requires two moments: a valid instrument and an additional generated variable, for instance, the interaction between the instrument and covariate, to estimate both parameters. If we employ 2SLS, we need a model specification assumption. However, if the covariate in the interaction term exhibits limited variation or lacks information, GMM estimates can become inconsistent with high variance, resembling a weak identification problem. In contrast, our D-RSMD method relies solely on a single valid instrument, eliminating the need for additional generated variables, first-stage model specification, or moment selection.

Another approach to addressing this issue brought about by interacting with the covariate is splitting the data set, which is effective when dealing with binary covariates and a known outcome model with a substantial number of observations. However, the D-RSMD method offers distinct advantages by leveraging all information from a single conditional moment restriction across all observations, producing competitive results with smaller standard errors (SEs), sometimes even smaller SEs than those from traditional GMM with the correct model specification. We illustrate this idea using simulations in the appendix.

The second advantage of the D-RSMD estimator is its applicability in big data settings. The R-SMD estimator introduced by Antoine and Sun (2021) combines Robinson transformation with SMD estimation to estimate parameters in the linear portion of a partially linear model¹. In their work, Antoine and Sun (2021) employ the Nadaraya-Watson estimator to estimate conditional

¹see Robinson (1988) and Li and Racine (2006) for combining Robinson transformation and 2SLS estimators.

means, restricting the number of covariates to fewer than four (refer to Chapter 7 of Li and Racine (2006)). However, when dealing with models involving a large number of covariates, as in their empirical application, they resort to dimension reduction techniques such as Principal Component Analysis (PCA).

In contrast, the D-RSMD method is tailored for big data scenarios, thus relaxing restrictions on the number of covariates. This is achieved by applying a method that works in big data settings, then incorporating the debiased method proposed by Chernozhukov et al. (2018). In their work, Chernozhukov et al. (2018) employ various techniques such as Lasso, random forests, neural networks, boosted regression trees, ridge regression, and so on. Then they employ the debiased method to reduce the effect of bias introduced by those techniques, which involves Neyman-Orthogonalization. This method yields an orthogonalized first-order condition (FOC) akin to the Robinson transformation proposed in Robinson (1988) for partially linear models.

Our work employs the Neyman-Orthogonalized method following Robinson transformation. This adaptation is necessary due to the additional layer introduced by the SMD estimation procedure within the objective function. Orthogonalizing the FOC with this layer helps mitigate bias arising from employing the method in big data settings. Consequently, our D-RSMD estimator extends the debiased method proposed by Chernozhukov et al. (2018) into the settings of SMD and U-statistics. Recent work by ? provides general results for the Neyman orthogonalization of U-statistic type moments, offering a general construction of orthogonal quadratic moment functions for obtaining debiased estimators and valid inference in settings with U-statistics. While our objective function resembles a special case of the function in ?, it incorporates a special weighting component based on the Fourier Transform using two observations from SMD, adding another layer of complexity to the construction of the Neyman-orthogonal FOC. With this extra weighting component, our identifying moment cannot be linearized. Hence, the methods and the inference results of ? assuming the existence of a linearized component in Equation (2.8) of ? are not applicable in our work.

Our D-RSMD focuses on using the Lasso method. The Lasso method is not the only option we

have; as discussed in Chernozhukov et al. (2018), there are lots of available estimation methods. Here, we use Lasso as an illustration. Some researchers use sieves for the non-parametric part of the model, and they need to specify how to select the number of sieves in their estimation procedure in practice. In comparison, our new method does not need to choose the number of polynomials for the sieve method. When we use the Lasso method, the tuning parameter is inside the penalty term for the estimation of the nuisance parameter. In practice, there are several ways to choose Lasso tuning parameters (see van de Geer (2016)), and we use cross-validation (Chernozhukov et al. (2018)). When we know there are fewer than four covariates, the Nadaraya-Watson estimator is an alternative method. This estimation procedure also needs one tuning parameter, namely the bandwidth. The bandwidth can be chosen by rule of thumb or cross-validation in practice. We show that our D-RSMD estimator is consistent and \sqrt{n} asymptotically normal under mild regularity conditions. Monte-Carlo simulations show that the D-RSMD estimator outperforms the R-SMD and GMM-type estimators in terms of much smaller bias and lower standard errors.

To demonstrate the benefits of our method, we examine the impact of Medicaid using data from the Oregon Health Insurance Experiment.² Our estimator, using a single valid instrument in the Oregon Health Insurance Experiment, yields statistically significant results for heterogeneous treatment effects, unlike the GMM estimator. Moreover, our approach produces more reliable results without requiring a generated instrument variable. This section presents an empirical analysis of the new D-RSMD estimator, emphasizing its advantages in estimating treatment effects and associated standard errors. Furthermore, it contributes to the literature on the Oregon Health Insurance Experiment by uncovering heterogeneous treatment effects that traditional methods fail to detect.

The paper is organized as follows: Section 2 introduces our framework, motivation, and the estimator. Section 3 states the large sample properties for our estimator. Section 4 presents the simulation results for finite samples with a large number of covariates. Section 5 uses the D-RSMD

²I would like to acknowledge A. Colin Cameron for drawing my attention to the Oregon Health Insurance Experiment and providing access to the dataset. The dataset is sourced from his and Pravin K. Trivedi's book in 2022 and is also available on the website https://www.stata-press.com/data/mus2.html.

estimator to estimate the effects of enrollment in Medicaid on outcomes of interest, based on the Oregon Medicaid health experiment. Additional results and proofs are in the Appendix.

2 Framework and Motivation

Our framework is based on a partially linear model featuring a binary treatment variable W_i . Our focus lies in understanding the treatment's impact on the outcome variable. However, it is important to note that the treatment is not always exogenous. Take, for instance, the Oregon Medicaid health experiment, where treatment corresponds to enrollment in Medicaid—a decision influenced by the choice of individual lottery winner, rendering enrollment endogenous. Moreover, the treatment effect may not be homogeneous; it can vary depending on certain covariate variables in X_i , a vector of q_X covariates $(X_i = [X'_{i1}, X'_{i2}]')$. Thus, we seek to examine the treatment effect conditional on specific groups determined by the values of the covariates X_{i1} . The remaining part of the model consists of an unknown function of exogenous X_i . While we maintain a fixed dimension for X_{i1} for illustrative purposes, the dimension of X_i is unrestricted and can be either high or low-dimensional. Additionally, the outcome variable y_i is not constrained to be binary. For studies involving binary outcome variables, a large number of covariates, and similar restricted dimension settings for estimating heterogeneous treatment effects, see Nekipelov et al. (2018). Generally, we focus solely on scenarios where we have knowledge of the functional form of the treatment and its heterogeneity, representing an initial step toward utilizing conditional moment restriction to estimate unobserved heterogeneity.

$$y_i = \theta_{w0} W_i + W_i \cdot X'_{i1} \theta_{wx0} + f_{0,1}(X_i) + \epsilon_i$$
(1)

The formal definition for $W_i \cdot X'_{i1} \theta_{wx0}$ is $\sum_{k=1}^{q_{X_1}} W_i X_{i1,k} \times \theta_{wx0,k}$ where $X_{i1,k}$ is a k-th element inside the vector X_{i1} . The treatment part of interest is $W_i + W_i \cdot X_{i1}$. The parameters that measure heterogeneous treatment effects, i.e. θ_{w0} and θ_{wx0} , are our key parameters.

As in the previous discussion, W_i and the interaction term between W_i and X_{i1} are endogenous.

To estimate the key parameters, a vector³ of instruments Z_i is introduced. For instance, in the Oregon Medicaid health experiment, the instrument is the lottery outcome, that is, the random assignment⁴. We maintain the conditional mean independence assumption for the instrument. With valid instrument and exogenous control variables, the conditional moment restriction is

$$E(\epsilon_i | X_i, Z_i) = 0 \ a.s. \tag{2}$$

We consider the single treatment case, where the covariates may be correlated with the instruments. Traditional estimators, such as GMM or 2SLS, typically require $1 + q_{X_1}$ moments or exogenous variables to identify (and estimate) $1 + q_{X_1}$ parameters within θ_{w0} and θ_{wx0} . These moments or exogenous variables include functions of Z_i and X_i . For instance, the interaction terms $Z_i \cdot X_{i1}$ are often employed as generated instruments or in constructing moments.

However, as discussed in Section 1, if there is little variation in X_{i1} , the $Z_i \cdot X_{i1}$ interaction terms become less informative. In such cases, traditional GMM methods will fail to provide reliable estimates. For example, if X_{i1} is binary, limited variation means that the majority of observations in X_{i1} are either 0 or 1. With only one valid instrument, GMM encounters the problem of generating new moments when we need to estimate more than one parameter, leading to under-identification. While splitting the sample in this case can solve the problem, it decreases the number of observations, thus resulting in larger standard errors, as discussed in the appendix.

This dilemma is resolved by directly employing the conditional moment restriction. This restriction encapsulates all the information of each instrument, regardless of the number of instruments used. Consequently, this approach allows us to utilize just one instrument, such as the random assignment Z_i , to identify and estimate parameters associated with both W_i and $W_i \cdot X_{i1}$. Similar approaches have been employed in the construction of other Bierens-type estimators, such as the Integrated Conditional Moment (ICM) estimators. Notable references include Bierens (1982),

 $^{{}^{3}}Z_{i}$ denotes the general vector of instruments. If we only use one instrument, Z_{i} will be the random assignment.

If we use a vector of instruments, Z_i is the vector including the random assignment and other instruments.

⁴In the experiment, the lottery winners are allowed to enroll in the Medicaid program. Not every lottery winner enrolled. The treatment variable is the actual enrollment status.

Antoine and Lavergne (2014), Antoine and Sun (2021), and ?. We will demonstrate that our estimation strategy, relying solely on a single valid instrument (e.g., the random assignment Z_i), yields reliable inference for both parameters.

The function $f_{0,1}(.)$ represents the nuisance parameter, which is not of interest in our analysis. The unknown form of $f_{0,1}(.)$ allows X_i to enter the model flexibly. The initial step in constructing our estimator involves employing a Robinson transformation, as described in Robinson (1988). This transformation involves subtracting the conditional expectation of y_i with respect to the controls X_i . As a result, the term $f_{0,1}(X_i)$ vanishes, eliminating the need to assume any specific form for $f_{0,1}(X_i)$. With the Robinson transformation and the exogeneity assumption on X_i , Equation 1 simplifies to:

$$y_i - E(y_i|X_i) = (P_i - E(P_i|X_i))'\theta_0 + \epsilon_i$$

where $P_i = [W_i, W_i \cdot X'_{i1}]'$ and $\theta_0 = [\theta_{w0}, \theta'_{wx0}]'$. Denote

$$\epsilon_j(\theta, g_0) = \tilde{y}_j - \tilde{P}'_j \theta \tag{3}$$

with $\tilde{y}_j \equiv y_j - E(y_j|X_j)$ and $\tilde{P}_j \equiv P_j - E(P_j|X_j)$.

This equation also illustrates that P_j may encompass any known function form of the treatment, interactions, covariates, and so forth. The symbol g_0 represents all the unknown real-valued functions, i.e., nuisance parameters. At this stage, g_0 comprises a vector of $g_{0,y}(X_j)$ (or $E(y_j|X_j)$) and $g_{0,P}(X_j)$ (or $E(P_j|X_j)$). Through this transformation, we exclude $f_{0,1}(X_i)$ from the model and introduce g_0 . This is also the first step in Antoine and Sun (2021).

Even though we only have a conditional moment restriction, we can exploit all the information in it by using an infinite number of unconditional moments. This equivalence between a conditional moment restriction and an infinite number of unconditional moment restrictions is from Bierens (1982) and used in constructing SMD-type or ICM-type estimators. Based on Equation (3) and the equivalence, we have the following results:

$$E[\epsilon_j(\theta_0, g_0)e^{it'Z_j}] = 0 \ \forall t \in \mathbb{R}^{q_z} \qquad \Longleftrightarrow \qquad E(\epsilon_j(\theta_0, g_0)|Z_j) = 0 \ a.s. \tag{4}$$

t is a vector of any real number.⁵ Next, we define the population objective function in Equation (5). Inside Equation (5), $\mu(t)$ is a strictly positive measure on the vector t.

$$M_{\infty}(\theta, g_0) = \int_{\mathbb{R}^{q_z}} |E[\epsilon_j(\theta, g_0)e^{it'Z_j}]|^2 d\mu(t)$$
(5)

The objective function involves the norm of a complex function. To estimate θ_0 , we need to find the derivative of the objective function, which is difficult to compute. Under the independent assumption for the population, the objective function has an alternative expression. In the Appendix, we show the equivalence between these two objective functions.

$$\forall j \neq l, \ M_{\infty}(\theta, g_0) = E[\epsilon_j(\theta, g_0)\epsilon_l(\theta, g_0)\kappa_{j,l}] \text{ with } \kappa_{j,l} = \int_{\mathbb{R}^{q_z}} e^{it'(Z_j - Z_l)} d\mu(t)$$
(6)

The objective function defined in Equation (6) depends on the parameters θ and the nuisance parameters g_0 , as specified in Equation (3) after the Robinson transformation. Under the regularity assumptions provided subsequently, θ_0 serves as the unique minimizer of the objective function $M_{\infty}(\theta, g_0)$, where g_0 represents a vector of $g_{0,y}(X_j)$ and $g_{0,P}(X_j)$ at this stage. Antoine and Sun (2021) similarly define an objective function incorporating $\kappa_{j,l}$ as the special weighting component from the SMD method, summarizing all the information from the conditional moment. While this may resemble a special case of the general moment considered in ?, we find that with $\kappa_{j,l}$, it is not possible to identify a function and constants satisfying Equation (2.8) of ?. Consequently, the methodology and inference results of ? do not apply. Therefore, we make assumptions and derive theorems tailored to our specific setting.

The FOC of $M_{\infty}(\theta, g_0)$ with respect to θ is

$$E[\tilde{P}_j(\tilde{y}_l - \tilde{P}_l'\theta)\kappa_{j,l}] = 0 \tag{7}$$

When we do not know g_0 , the FOC of $M_{\infty}(\theta, g)$ is as follows:

$$E((P_j - g_P(X_j)) [y_l - g_y(X_l) - (P_l - g_P(X_l))'\theta]\kappa_{j,l}) = 0$$
(8)

⁵Using higher order Fourier terms, for instance, $E[\epsilon_j(\theta_0, g_0)e^{it^2 Z_j}]$ will lead to implicit choice of $\mu(t)$ and assumptions about k(.) introduced in the later discussion and in practice. We choose to work with $e^{it'Z_j}$ because of its explicit and easy-to-use practice results. This is also the choice of Antoine and Sun (2021).

The FOC defined in Equation (7) provides an explicit form for θ_0 . This FOC extends the framework established by Antoine and Sun (2021) to incorporate interaction terms within the parametric component of the model. Thus, the estimator, derived as the sample analog of θ_0 , is a direct extension of Antoine and Sun (2021). It also represents a specialized instance of the methodology proposed by Lavergne and Patilea (2013) when the bandwidth within $\kappa_{j,l}$ is held constant.

Assumption 1. (Regularity assumptions)

- (a) $E(\epsilon_i | X_i, Z_i) = 0.$
- (b) $E(\tilde{P}_i|Z_i) \neq 0$ a.s. (with probability 1) with $\tilde{P}_i = P_i E(P_i|X_i)$.
- (c) $E(\tilde{P}_i\tilde{P}'_i)$ is nonsingular.

(d) Let $f_Z(.)$ denote the density function of Z_j . We assume that $E(\tilde{P}_j|Z_j = .)f_Z(.)$ is L_q for some $1 \le q \le 2$.

(e) (y_l, W_l, X_l, Z_l) is an independent and identical copy of (y_j, W_j, X_j, Z_j) with $l, j \in \{1, ..., n\}$.

(f) Let μ be a given strictly positive measure on \mathbb{R}^{q_z} . Let k(.) be the Fourier inversion integral induced by μ , $k(Z_j - Z_l) = \int_{\mathbb{R}^{q_z}} e^{it'(Z_j - Z_l)} d\mu(t)$. We assume that k(.) is a symmetric bounded density function on \mathbb{R}^{q_z} and that its Fourier transform is strictly positive.

Assumption 1(**a**) is the exogeneity assumption for the controls and instruments. In the potential outcomes setting, Assumption 1(**a**) is also the Random Assignment Assumption. The model form implies the Exclusion Restriction Assumption as in Imbens and Rubin (2015), Page 558, that is, the value of the instrument does not affect the potential outcomes directly. Assumption 1(**b**) is the relevant instrument assumption. Assumption 1(**c**) and the corresponding Assumption in Robinson (1988) are not the same. This assumption is made in a high-dimensional setting. For not invertible $E(\tilde{P}_i\tilde{P}'_i)$, see the solution provided in Chernozhukov et al. (2018).

Assumption 1(**f**) is for the measure $\mu(.)$. These conditions in Assumption 1(**f**) are not very restrictive. μ has a \mathbb{R}^{q_z} support and is symmetric. This leads us to have a symmetric $k(Z_j - Z_l)$. Also, the fourier transform of $k(Z_j - Z_l)$ is μ . Thus, the fourier transform of $k(Z_j - Z_l)$ is strictly positive. There are many different available measures. Measures after Fourier inversion have the respective results. For the simplicity of computation, we focus on those k(.) with explicit formulas. Specifically, we use the CDF of the Gaussian distribution in simulations and applications.

Under Assumption 1, θ_0 is the unique minimizer of $M_{\infty}(\theta, g)$ when $g = g_0$. Specifically, assumptions 1(c) and 1(d) guarantee the identification of the parameters that we are interested in under the FOC defined in Equation (7). This is established in Antoine and Sun (2021) and is easily extended to our setting.

However, given that the nuisance parameters in g_0 are unknown, Antoine and Sun (2021) consider the sample analog of θ_0 defined in Equation (7) as an infeasible estimator. To obtain a feasible estimator, Antoine and Sun (2021) estimate the nuisance parameters as the first step. They achieve this by estimating their nuisance parameters using Nadaraya-Watson estimators. They demonstrate that, under appropriate assumptions on the bandwidth, the bias introduced by the Nadaraya-Watson estimators on the estimation of the key parameters is sufficiently small, ensuring that the feasible and infeasible estimators of the key parameter share the same asymptotic properties. However, the Nadaraya-Watson estimator imposes a constraint on the number of covariates to satisfy these assumptions. This constraint may not be suitable when there are a large number of available covariates. Moreover, even with a small number of covariates, when $g \neq g_0$, the bias resulting from estimating the nuisance parameters affects the estimation of the key parameters, particularly when the sample size is small. This is because the FOC defined in Equation (8) is not orthogonal to the nuisance parameter, leading to a more pronounced effect of the bias on the estimate of the key parameter. This is illustrated in the proof of Appendix.

To address this bias issue, we propose a new FOC that extends the Neyman-orthogonal estimator from Chernozhukov et al. (2018) to the SMD and U-statistic settings. The Neyman-Orthogonal method constructs an FOC that is orthogonal to the nuisance parameters. This new FOC ensures that all partial derivatives with respect to the nuisance parameters are zero, effectively making it orthogonal to the bias introduced by estimators. The concept of partial derivatives with respect to a function is elucidated in Chernozhukov et al. (2018).

Following Neyman-orthogonalization, the new FOC for θ_0 becomes less sensitive to the bias of

the estimator for the nuisance parameters. It is, however, sensitive to the square of the bias. As long as the bias is of the order $o_p(n^{-1/4})$, we still obtain a \sqrt{n} -asymptotically normally distributed estimator for the key parameter. It is important to note that under this order, there are numerous estimation methods to choose from, including Lasso, Sieves, and Random Forest, among others.

Our new FOC for θ_0 corresponding to Equation (7) is

$$E[\Psi(D;\theta,g_0)] \equiv E\left[\left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\right) [\tilde{y}_l - \tilde{P}'_l\theta]\kappa_{j,l}\right] = 0$$
(9)

with $g_{0,\tilde{P}_m}(X_l) \equiv E[(P_m - g_{0,P}(X_m))\kappa_{m,l}|X_l]$ and $g_{0,\kappa_{m,l}}(X_l) \equiv E[\kappa_{m,l}|X_l]$. $g_{0,\tilde{P}_m}(X_l)$ has the same dimension as \tilde{P}_j , and $g_{0,\kappa_{m,l}}(X_l)$ is a function of X_l : $\mathbb{R}^{q_X} \to \mathbb{R}$.

 $g_{0,\tilde{P}_m}(X_l)$ and $g_{0,\kappa_{m,l}}(X_l)$ are two additional parameters inside the nuisance parameter vector. The FOC defined in Equation (9) has a partial derivative with respect to all nuisance parameters equal to zero. This is shown in the Appendix.

Assumption 2. The singular values of the matrix $E\left[\kappa_{j,l}\left(\tilde{P}_j - g_{0,\tilde{P}_m}(X_l)g_{0,\kappa_{m,l}}^{-1}(X_l)\right)\tilde{P}_l'\right]$ are between c_0 and c_1 with $0 < c_0 < c_1$.

Assumption 2 is the identification assumption. This is the similar assumption as Assumption 3.1 (e) in Chernozhukov et al. (2018). As a contrary, ? and ? do not impose the identification assumption, because the structure of the $\Psi(.; \theta, g)$ of their works guarantee the identification. Here, due to the fact that our $\Psi(D; \theta, g)$ is constructed under the SMD method with the special weight $\kappa_{j,l}$, we need to assume additional identification assumption as in Chernozhukov et al. (2018). Additionally, this assumption can be examined. We show this detailed information in the identification section of the Appendix, Section D⁶. In this section, we show that with the information of the true Data Generating Process (DGP), the determinant of the matrix $E\left[\kappa_{j,l}\left(\tilde{P}_j - g_{0,\tilde{P}_m}(X_l)g_{0,\kappa_{m,l}}^{-1}(X_l)\right)\tilde{P}_l^{\prime}\right]$ is deducible. Also, in the applied setting, if the matrix is singular, the resulting standard error of the estimator will be huge and abnormal. This provides an intuitive way to determine whether the matrix is invertible.

⁶In the Appendix, we illustrate the identification issue for the group of SMD estimators, such as SMD, RSMD, and D-RSMD. We consider cases involving various types of instruments and models.

Equation (9) gives us the identification of the true key parameter and the explicit forms of the infeasible and feasible estimators.

Proposition 1. (Identification of θ_0 using the orthogonalized FOC) Under Assumptions 1-2 and FOC defined in Equation (9)

$$\theta_0^* = E\left[\kappa_{j,l}\left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\right)\tilde{P}_l'\right]^{-1}E\left[\kappa_{j,l}\left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\right)\tilde{y}_l\right]$$

If we plug Equation (3) into the formula for θ_0^* with error term ϵ_l , we will have $\theta_0^* = \theta_0$. Thus, the sample analog of θ_0^* delivers an estimator for θ_0 . Because we have two distinct individuals inside the expectation (e.g., j and l), we need to replace the expectation by the average of a double summation to obtain the infeasible estimator under Equation (9). The closed-form expression for the infeasible estimator, $\tilde{\theta}_{n,o}$, is:

$$\tilde{\theta}_{n,o} = \left[\sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right) \tilde{P}_{l}'\right]^{-1} \left[\sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right) \tilde{y}_{l}\right]$$

The infeasible estimator $\tilde{\theta}_{n,o}$ depends on unknown nuisance parameters: $g_{0,\tilde{P}_m}(X_l)$, $g_{0,\kappa_{m,l}}(X_l)$, $E(y_l|X_l)$, and $E(P_l|X_l)$. All of the nuisance parameters are conditional expectation functions on covariates. Hence, in practice, we need to find estimators for these conditional expectation functions to obtain a feasible estimator. Replacing every nuisance parameter g_0 with estimators \hat{g} such that \hat{g} converges to g_0 at a rate of $o_p(n^{-1/4})$ will deliver the feasible estimator on θ_0 with \sqrt{n} asymptotic normality.

The nuisance parameters are functions of X involving q_X variables. If q_X is very large, such as $q_X \ge n$, there will be overfitting. To mitigate this, we need to reduce the number of variables included in the regression. Using Lasso will be one of the solutions. In order to apply Lasso, we assume sparsity for the nuisance parameters; that is, the conditional means can be described with only a few non-zero parameters in front of X. The number of non-zero parameters, s_0 , is allowed to grow at the rate of $o_p(n^{1/2}/log(q_X))$. Under the assumed rate for s_0 , the Lasso estimation has the following property (as Equation (2.1) of Chapter 2 in van de Geer (2016)) on the order of mean square error:

$$||X(\hat{\beta} - \beta_0)||_2^2/n = \mathcal{O}_p\left(\frac{s_0 \log(q_X)}{n}\right)$$

where $\hat{\beta}$ is the Lasso estimator (Chernozhukov et al. (2018) and van de Geer (2016)), and $||.||_2$ is the \mathcal{L}_2 norm. When $3 < q_X < n$, we can still use the Lasso method with a higher assumed rate for s_0 to select variables. If $q_X \leq 3$, we can still use the Nadaraya-Watson estimator to estimate the nuisance parameter with a second-degree kernel. It is worth noting that $q_X \leq 3$ is a constraint that is seldom realistic in practice. Nevertheless, our estimation procedure allows us to accommodate larger values of q_X . Upon replacing every nuisance parameter with its estimate, we obtain the feasible estimator $\hat{\theta}_{n,o}$, referred to as the D-RSMD estimator.

$$\hat{\theta}_{n,o} = \left[\sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}}(X_{l})}{\widehat{g_{0,\kappa_{m,l}}}(X_{l})}\right) \hat{\tilde{P}}_{l}^{\prime}\right]^{-1} \left[\sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}}(X_{l})}{\widehat{g_{0,\kappa_{m,l}}}(X_{l})}\right) \hat{\tilde{y}}_{l}\right]$$
(10)
since $E[\kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}}(X_{l})}{\widehat{g_{0,\kappa_{m,l}}}(X_{l})}\right) \hat{\tilde{P}}_{l}^{\prime}]$ is invertible.

This leads to the algorithm of our D-RSMD estimation procedure.

Algorithm 2.1. (Implementation of the D-RSMD estimator)

- 1. Conduct the Robinson transformation. This step involves estimating the conditional means inside $\hat{\tilde{P}}_j$ and $\hat{\tilde{y}}_l$. Any estimator \hat{g} that converges to g_0 at a rate of $o_p(n^{-1/4})$ can be applied.
- 2. Calculate $\kappa_{j,l}$. Within $\kappa_{j,l}$, $\mu(.)$ represents the CDF of the Gaussian distribution. Given that the Fourier transform of the Gaussian distribution is also Gaussian, computing $\kappa_{j,l}$ is straightforward.
- 3. Estimate the nuisance parameters $\widehat{g_{0,\tilde{P}_m}}(X_l)$ and $\widehat{g_{0,\kappa_{m,l}}}(X_l)$ within the orthogonal FOC.
- 4. Calculate the estimate based on Equation (10) for $\theta_{n,o}$.

All of the nuisance parameters for the D-RSMD estimator in the simulation and empirical application are estimated by the Lasso method with cross validation. The maximum degree of the polynomial for the nuisance parameters is 5, which guarantees that the nuisance parameters can be approximated by 5 degree polynomials in all controls. Lasso with cross validation helps us select the controls and their polynomials.

3 Large Sample Theory

References such as Lavergne and Patilea (2013) provide a general framework for analyzing the asymptotic properties of SMD estimators, even in cases where explicit forms of these estimators are not available. Building upon this work, Antoine and Sun (2021) extend the methodology to accommodate semiparametric models and derive the asymptotic properties specific to the RSMD estimator. In this section, we establish the asymptotic properties of the D-RSMD estimator, demonstrating that the feasible estimator $\hat{\theta}_{n,o}$ and the infeasible estimator $\tilde{\theta}_{n,o}$ are outlined as follows:

Proposition 2. (Consistency and Asymptotic normality of $\tilde{\theta}_{n,o}$)

Under Assumption 1 and iid assumption for the sample, $\tilde{\theta}_{n,o}$ is consistent for θ_0 , that is $\tilde{\theta}_{n,o} \xrightarrow{p} \theta_0$, and asymptotically normally distributed,

$$\sqrt{n}(\tilde{\theta}_{n,o}-\theta_0) \xrightarrow{d} N\left(0, A^{-1}\Sigma\left(A^{-1}\right)'\right)$$

where $A = E\left[\kappa_{j,l}\left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\right)\tilde{P}'_l\right]$, and Σ stands for the variance-covariance matrix involving terms of $(\tilde{P}_i, \epsilon_i, Z_i, X_i)$, which is defined in the proof section of the Appendix.

The asymptotic properties of $\tilde{\theta}_{n,o}$ are based on the corresponding properties of U-statistics. This is thoroughly discussed in Hoeffding (1948), particularly in Theorem 7.1 on Page 320 of the book.

Assumption 3. \hat{g} converges to g_0 at a rate of $o_p(n^{-1/4})$. For the Lasso method, the number of non-zero parameters s_0 grows at the rate of $o_p(n^{1/2}/\log(q_X))$. For the Nadaraya-Watson estimator, $\sqrt{n}\left(\sum_{s=1}^{q_z} h_s^4 + \left[\frac{1}{nh_1...h_{q_z}}\right]\right) = o(1)$ where h is the bandwidth.

Theorem 3.1. (Consistency and Asymptotic normality of the D-RSMD estimator: $\hat{\theta}_{n,o}$) Under

Assumptions 1 - 3, $\hat{\theta}_{n,o}$ is consistent, and has an asymptotically normal distribution, that is,

$$\sqrt{n}(\hat{\theta}_{n,o} - \theta_0) \xrightarrow{d} N\left(0, A^{-1}\Sigma\left(A^{-1}\right)'\right),$$

Notably, the asymptotic variances of the D-RSMD and R-SMD estimators proposed by Antoine and Sun (2021) differ, primarily due to additional terms introduced through the debiasing process in A and Σ . Analytical comparison of these asymptotic variances is challenging, given the complexity introduced by these extra terms. Furthermore, we provide the explicit form of the estimator for variance under heteroskedasticity in the following:

$$[n(n-1)C_n]^{-1} \sum_{j=1}^n ((\sum_{l=1}^n \kappa_{j,l} \left(\widehat{\tilde{P}}_l - \frac{\widehat{g_{0,\tilde{P}_l}}(X_j)}{\widehat{g_{0,\kappa_{m,j}}}(X_j)} \right)) (\sum_{l=1}^n \kappa_{j,l} \left(\widehat{\tilde{P}}_l - \frac{\widehat{g_{0,\tilde{P}_l}}(X_j)}{\widehat{g_{0,\kappa_{m,j}}}(X_j)} \right)') \hat{\epsilon}_j^2) [n(n-1)C_n']^{-1}$$
(11)

with $C_n = \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\widehat{\tilde{P}}_j - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \widehat{\tilde{P}}_l'.$

Based on Theorem 3.1, the feasible and infeasible estimators exhibit the same asymptotic distribution and demonstrate consistency. This is from Assumption 3 on the convergence rate, which ensures that the bias introduced during nuisance parameter estimation has negligible impact on subsequent steps. For example, when the number of covariates (q_X) and the sample size are small, the D-RSMD estimator exhibits less bias compared to the R-SMD estimator, especially when both use the same Nadaraya-Watson estimator for nuisance parameter estimation. In Chernozhukov et al. (2018), simulations illustrate a comparison between Neyman orthogonal estimators and non-orthogonal estimators. Further details on bias analysis are presented in the subsequent section, specifically in the simulation section.

4 Simulation Study

We consider a partially linear model with one endogenous variable and its interaction in the linear part.

$$y_{i} = \theta_{w0}W_{i} + W_{i} \cdot X_{i1}\theta_{wx0} + \sum_{q=1}^{q_{X}}\beta_{1q}X_{qi} + \sum_{q=1}^{q_{X}}\beta_{2q}X_{qi}^{2} + \epsilon_{i}$$
(12)
$$W_{i} = I(\theta_{z0}Z_{i} + \theta_{z3}Z_{i}^{3} + \sum_{q=1}^{q_{X}}\alpha_{1q}X_{qi} + \sum_{q=1}^{q_{X}}\alpha_{3q}X_{qi}^{3} > -v_{i})$$

I(.) is the indicator function. It will take the value one if the statement inside is true and zero otherwise. In the simulations, the key parameters θ_{w0} and θ_{wx0} are 2 and 3 respectively. The other parameters are in the following. $\theta_{z0} = 3$, $\theta_{z3} = 4$, $\alpha_{3q} = 2$, $\beta_{2q} = -3$, $\alpha_{1q} = \beta_{1q} = 1$ for $q \leq S$ with S the number of non-zero parameters, and q_X ($q_X = 30$) the number of covariates. There are nonlinear terms in the model, so S is chosen to be a half of s_0 , the sparsity level. $\alpha_{1q} = \alpha_{3q} = \beta_{1q} = \beta_{2q} = 0$ when q > S. Here, S = 5. If β_{2q} is 0 for all q, y_i is linear in X_i . Otherwise, the model is partially linear.

We use a binary instrument to create a treatment variable and explore simulation outcomes. The results when we use a categorical instrument in the DGP are in the appendix. Z_i is the instrument with $E(Z_i) = 0.318$. The covariate X_i is modeled as $X_i^* + 0.4Z_i$, where X_i^* follows a multivariate standard normal distribution. This implies a correlation between X_i and Z_i . The errors (ϵ, v) are bivariate normally distributed with mean 0, variance 1, and covariance $\frac{4}{9}$. Consequently, the treatment variable W_i is endogenous.

To assess the performance of our D-RSMD estimators, we conduct 5,000 Monte Carlo replications for each scenario, where $q_X \in \{3, 30\}$ and the sample size $n \in \{3000, 5000\}$. Our benchmark scenario involves 3000 observations and 30 control variables, which mirrors our empirical example. For instance, when examining the Oregon Health Insurance Experiment, we segment the dataset into three age-based groups, each containing around 6,000 observations and 21 covariates.

We report and compare simulation results for several estimators in our study. These include the D-RSMD estimator proposed in this paper (referred to as DRSMD-Lasso in Section 5), where we use the Lasso method for estimating nuisance parameters. We also consider the R-SMD estimator by Antoine and Sun (2021) using Lasso, the R-GMM estimator combining Robinson Transformation with GMM (GMM-Lasso in Section 5), a GMM estimator treating $f_{0,1}(X_i)$ as linear in X_i , and a GMM (Oracle) estimator utilizing the true $f_{0,1}(X_i)$. We present results for the D-RSMD estimator with one and two instruments, along with results for the R-SMD estimator under both conditions. GMM-type estimators require at least two instruments to estimate two parameters. We list the available instrument sets used for estimation in the third column of each table.

All nuisance parameters for the D-RSMD estimator in both the simulation and empirical application sections are estimated using the Lasso method with cross-validation. We employ a maximum polynomial degree of 5 for the nuisance parameters. This involves generating 5-degree polynomials for each covariate and using the Lasso method with cross-validation to select relevant controls and their respective polynomials, resulting in predicted conditional expectations. This approach provides a non-parametric estimation similar to the sieve method by selecting polynomial degrees. Cross-validation is utilized to select the penalty level and mitigate the risk of overfitting. For RSMD and RGMM estimators with $q_X = 30$, we also adopt the same Lasso method for polynomial selection to ensure comparability across estimators.

We present the results in Table 1. Firstly, focusing on the first panel with $q_X = 3$, we observe that D-RSMD using Z_1 achieves the lowest MAD and Med.SE compared to all estimators. The RRs are close to 5% for θ_{w0} and slightly oversized for θ_{wx0} . Using the instrument Z_2 (not in the DGP) for D-RSMD results in higher MAD and Med.SE compared to utilizing the correct instrument. The D-RSMD estimator also performs well when employing (Z_1, Z_1X_1) and (Z_2, Z_2X_1) . Comparing D-RSMD estimators, those using Z_1 or the pair of instruments including the correct instrument tend to yield lower MADs and Med.SEs, which is reasonable because working with the correct variable increases estimation precision. The Med.Bias column indicates that for θ_{w0} , D-RSMD estimators using (Z_1, Z_1X_1) and (Z_2, Z_2X_1) exhibit higher bias than other D-RSMD estimators, suggesting that using Z_1 leads to lower bias in this DGP. This property of D-RSMD also holds when there are 30 controls in the model. However, using one instrument may not always bring a lower bias. This is shown in the appendix when Z_2 is the instrument used in the DGP.

The second panel of Table 1 presents results for $q_X = 30$. Similar to the scenario with $q_X = 3$, among D-RSMD estimators, using Z_1 as the instrument yields better results, as evidenced by lower Med.Bias, MAD, and Med.SE values. Comparing the results obtained with Z_1 versus Z_2 , we notice a slight distortion in the RR when Z_1 is employed. However, this size distortion diminishes when using a sample size of 5,000 observations per replication, as shown in the last panel of the table. Comparing D-RSMD with RSMD underscores the advantage of D-RSMD in producing less biased

				θ_w	0			$ heta_{wx0}$		
	Estimator	Instrument	Med.Bias	MAD	Med.SE	RR	Med.Bias	MAD	Med.SE	RR
n = 3,000	D-RSMD	Z_1	-0.006	0.146	0.215	0.048	-0.005	0.039	0.057	0.052
$q_X = 3$	D-RSMD	Z_2	-0.004	0.212	0.314	0.043	-0.004	0.110	0.182	0.050
	D-RSMD	(Z_1, Z_1X_1)	-0.026	0.164	0.247	0.048	-0.004	0.124	0.186	0.047
	D-RSMD	(Z_2, Z_2X_1)	-0.035	0.230	0.340	0.049	-0.001	0.200	0.301	0.040
	GMM (Oracle)	(Z_1, Z_1X_1)	-0.004	0.164	0.250	0.048	-0.001	0.126	0.191	0.047
n = 3,000	D-RSMD	Z_1	0.028	0.288	0.427	0.045	-0.012	0.045	0.059	0.064
$q_X = 30$	D-RSMD	Z_2	0.033	0.524	0.773	0.030	-0.012	0.126	0.206	0.037
	D-RSMD	(Z_1, Z_1X_1)	-0.259	0.291	0.431	0.082	0.005	0.111	0.163	0.049
	D-RSMD	(Z_1, Z_2)	0.041	0.333	0.493	0.041	-0.013	0.181	0.287	0.039
	RSMD	Z_1	0.128	0.408	0.563	0.072	-0.020	0.053	0.065	0.105
	RSMD	Z_2	0.154	0.555	0.790	0.039	-0.024	0.147	0.302	0.018
	RSMD	(Z_1, Z_1X_1)	0.230	0.463	0.572	0.110	-0.337	0.137	0.273	0.144
	RSMD	(Z_1, Z_2)	0.140	0.438	0.605	0.060	-0.020	0.195	0.328	0.033
	GMM	(Z_1, Z_1X_1)	7.471	2.968	4.368	0.356	-6.735	0.953	1.430	0.999
	GMM	(Z_1, Z_2)	6.857	5.691	11.948	0.043	-4.444	21.525	47.007	0.000
	GMM (Oracle)	$(Z_1, Z_1 X_1)$	0.007	0.266	0.392	0.048	-0.004	0.107	0.157	0.049
	GMM (Oracle)	(Z_1, Z_2)	-0.011	0.635	1.424	0.003	0.114	2.502	5.548	0.000
n = 5,000	D-RSMD	Z_1	0.003	0.209	0.314	0.052	-0.005	0.029	0.040	0.062
$q_X = 30$	D-RSMD	(Z_1, Z_1X_1)	-0.158	0.210	0.308	0.083	0.006	0.078	0.120	0.045
	GMM (Oracle)	(Z_1, Z_1X_1)	0.001	0.203	0.304	0.051	0.002	0.080	0.121	0.045

Table 1: Performance of Estimators with Z_1 as the Instrument in the DGP

Note: Simulation Results for θ_{w0} and θ_{wx0} in the benchmark model using D-RSMD estimator 5,000 replications. We report the Monte-Carlo Median Bias (Med.Bias), Median Absolute Deviation (MAD), median of asymptotic standard error under heteroskedasticity (Med.SE), and Rejection Rate (RR) using a 5% t-test. Z_1 : binary instrument. Z_2 : variable representing the sum of Z_1 and another binary variable, resulting in three possible values for Z_2 . (Z_1, Z_1X_1) : pair consisting of Z_1 and the interaction term Z_1X_1 (where X_1 is a covariate). (Z_2, Z_2X_1) : pair comprising Z_2 and the interaction term Z_2X_1 . (Z_1, Z_2) : pair including both Z_1 and Z_2 . The correlation between Z_1 and Z_2 is around 0.7.

estimates, particularly when employing the Lasso method for nuisance parameter estimation.

Furthermore, comparing D-RSMD results with the GMM (Oracle) estimator reveals that D-

RSMD estimates using Z_1 closely approximate the GMM (Oracle) estimates for θ_{w0} in terms of MAD, Med.SE, and RR. For θ_{wx0} , D-RSMD results with Z_1 outperform the GMM (Oracle) estimator using (Z_1, Z_1X_1) , exhibiting lower MAD and Med.SE values. These findings suggest that D-RSMD with Z_1 achieves satisfactory performance due to its utilization of an infinite number of unconditional moments, equivalent to the conditional moment, unlike the GMM (Oracle) estimator, which relies solely on two unconditional moments.

Analyzing outcomes from instrument sets (Z_1, Z_1X_1) and (Z_1, Z_2) reveals notable differences between D-RSMD and GMM (Oracle) estimators. Specifically, for GMM (Oracle) estimators, (Z_1, Z_1X_1) emerges as the preferred choice, exhibiting the lowest Med.Bias for both θ_{w0} and θ_{wx0} . Conversely, GMM estimators using (Z_1, Z_2) demonstrate scattered results and numerous extreme cases, with the RR approaching 0 for θ_{wx0} , highlighting the challenges posed by instrument correlation within (Z_1, Z_2) for GMM type estimators.

In contrast, the choice of instrument sets has a smaller impact on D-RSMD results compared to its effect on GMM (Oracle) estimators. This observation suggests that D-RSMD exhibits more stable performance when utilizing at least one valid instrument or an instrument closely related to the valid instrument in the instrument set, compared with GMM-type estimators.

5 Empirical Application

In this section, we investigate the heterogeneous treatment effects of Medicaid enrollment from a Randomized Controlled Trial (RCT) conducted in Oregon. In early 2008, Oregon expanded Medicaid enrollment through a lottery system for low-income, uninsured adults. This lottery provided random selection, enabling researchers to analyze the effects of health insurance on various outcomes, including medical, financial, and labour market impacts. This experiment has been extensively studied in various articles, such as Baicker et al. (2014) and Finkelstein et al. (2012). Detailed information and datasets related to this RCT are available on the public website.⁷ In

 $^{^7} See \ \texttt{https://www.nber.org/research/data/oregon-health-insurance-experiment-data}$

this paper, we utilize a dataset derived from the original datasets.⁸ The dataset comprises 18,572 observations.

Many studies have focused on the homogeneous treatment effects of Medicaid. Finkelstein et al. (2012) assumed homogeneity in their analysis but also explored the potential for heterogeneous treatment effects through regression analysis. Estimates from 2SLS rule out the possibility of heterogeneous treatment effects. However, this may be due to the misspecification of the first stage or problems with the generated instrument variables.

To address this problem, we employ the D-RSMD estimator to estimate heterogeneous treatment effects. We compare the results between the new and traditional estimation methods, both with and without the generated instrument variable, to obtain reliable inference.

Here, the heterogeneity arises from interaction terms involving indicators (X_1) for household income above 50% of the federal poverty line in 2008 (income), household income (hhincome), TANF (cash welfare assistance to low-income families), or cigarette smoking level (smoke). The lottery serves as the instrumental variable. We adopt similar controls to those used by Baicker et al. (2014) and Finkelstein et al. (2012), including household controls, lottery and survey wave indicators, and individual characteristics. The dependent variables include the current employment indicator, constructed from three indicators: hours of employment (employment), total out-of-pocket spending on medical care (out-of-pocket cost), and whether the individual currently owes money to a healthcare provider (debt for health). All dependent variables are derived from a mail survey conducted between July 2009 and March 2010, approximately one year after treatment.

Given that health conditions, employment status, and other outcomes are closely linked to individuals' ages, we identify age as a key source of heterogeneity. We derive an age group variable from the year of birth data. First, we calculate the age of each individual, resulting in ages ranging from 21 to 64 years. Next, we partition the original dataset into three or five subsets based on age to

⁸Special thanks to A. Colin Cameron for providing the dataset and bringing the Oregon Health Insurance Experiment to my attention through his and Pravin K. Trivedi's book in 2022. The dataset is also accessible at https://www.stata-press.com/data/mus2.html

create the age-group variable. Each subsample contains approximately 5900 to 6900 observations. Individuals within the same age group share more similarities. In Section 5.1, we present results categorized into three age groups (e.g., 21 to 35, 35 to 50, and 50 to 64). Here, the 'agegroup' variable represents a categorical variable with three values, mirroring the age groups.

To provide a clearer comparison, we contrast the D-RSMD estimator with traditional estimators, such as the GMM estimator. The D-RSMD estimator is designed to cope with a nonparametric first stage and a partially linear second stage. The framework for the D-RSMD estimator is outlined as follows:

$$\begin{split} Debt_i &= \theta_{w0} Medicaid_i + \theta_{wx0} Medicaid_i \times income_i + \theta_{x10} income_i \\ &+ \theta_{x20} agegroup_i + \theta_{x30} Medicaid_i \times agegroup_i + f_{0,1}(X_i) + \epsilon_i \\ Medicaid_i &= I(f_{0,2}(X_i, Lottery_i) > v_i) \end{split}$$

The framework for GMM estimators includes a linear first stage within the indicator function as well as a linear second stage for the dependent variable.

$$\begin{split} Debt_i &= \theta_{w0} Medicaid_i + \theta_{wx0} Medicaid_i \times income_i + \theta_{x10} income_i \\ &+ \theta_{x20} agegroup_i + \theta_{x30} Medicaid_i \times agegroup_i + X'_i \beta_x + \epsilon_i \\ Medicaid_i &= I(\alpha_z Lottery_i + X'_i \alpha_x > v_i) \end{split}$$

The model for GMM-Lasso estimators includes a linear first stage for the indicator function and a partially linear second stage for the dependent variable.

$$\begin{split} Debt_{i} &= \theta_{w0} Medicaid_{i} + \theta_{wx0} Medicaid_{i} \times income_{i} + \theta_{x10} income_{i} \\ &+ \theta_{x20} agegroup_{i} + \theta_{x30} Medicaid_{i} \times agegroup_{i} + f_{0,1}(X_{i}) + \epsilon_{i} \\ Medicaid_{i} &= I(\alpha_{z} Lottery_{i} + X_{i}'\alpha_{x} > v_{i}) \end{split}$$

In Table 2, we present preliminary results for three estimators of θ . D-RSMD stands out as the sole estimator utilizing the lottery for estimating multiple parameters. We report D-RSMD results in two columns: using only the lottery (Z_1) and incorporating the lottery and its interaction term

 (Z_1, Z_1X_1) . This setup allows for a direct comparison of results across the two instrument sets and three estimators.

Using the DRSMD-Lasso estimator with the lottery as the sole instrument (Z_1) , we obtain statistically significant estimates for Medicaid and the interaction between Medicaid and income at a 5% significance level. Similarly, we observe statistically significant estimates for income and the interaction term between Medicaid and age at a 15% significance level with the lottery. These findings corroborate the conclusions drawn in Section 5.1, suggesting heterogeneous treatment effects in Medicaid related to age and income when employing the valid instrument alone. Conversely, analyzing outcomes using the GMM estimator or GMM-Lasso with the instrument set (Z_1, Z_1X_1) would erroneously lead to the conclusion of no heterogeneity in treatment effects, underscoring the problems associated with using generated instruments or moments.

Heterogeneous treatment effects										
Debt for Health		GMM		MM-Lasso	DRSMD-Lasso					
Variables	Z_1	$(Z_1, Z_1 X_1)$	Z_1	(Z_1, Z_1X_1)	Z_1	$(Z_1, Z_1 X_1)$				
Nr. 1 1		-0.148***		-0.206*	-0.187***	-0.178***				
Medicaid		(0.044)		(0.115)	(0.037)	(0.046)				
M. 1		0.003		-0.045	0.074***	-0.018				
Medicaid "income		(0.053)		(0.068)	(0.020)	(0.056)				
		0.003		0.014	-0.025.	0.005				
income		(0.020)		(0.062)	(0.016)	(0.025)				
		-0.007		-0.010	-0.006	-0.007				
agegroup		(0.010)		(0.021)	(0.005)	(0.009)				
λτ 1· · 1ψ		-0.019		-0.016	-0.018.	-0.016				
Medicaid*agegroup		(0.032)		(0.054)	(0.011)	(0.028)				

Table 2: Heterogeneous Treatment Effects for Robustness Check (5 parameters)

Note: *** Significant at 1%, ** at 5%, * at 10%, . at 15%.

5.1 Results

In this section, we examine the potential heterogeneous treatment effects of Medicaid on healthrelated debt, with a focus on household income as a source of heterogeneity. We partition the dataset into three subsets based on age to facilitate intuitive interpretation. The complete table presenting all treatment effects is provided at the end of this section. Additional results for robustness checks are in Appendix Section B.1.

We consider two sets of instruments: the lottery (used exclusively for D-RSMD) and the pair of the lottery along with its interaction with X_1 (indicating income greater than 50% of the poverty line). To illustrate, the model for debt is specified as:

$$Debt_i = \theta_{w0} Medicaid_i + \theta_{wx0} Medicaid_i \times X_{i1} + f_{0,1}(X_i) + \epsilon_i$$

Table 3 presents results for three age groups: 21–35, 35–50, and 50–64 years old. For the age group 35–60, the total number of observations is 6693. Heterogeneous treatment effects are observed among individuals. The estimates obtained by D-RSMD (using the lottery as an instrument or the lottery and its interaction with X_1 as instruments) are very similar to each other and to the GMM estimator (using the lottery and its interaction with X_1 as instruments). This indicates a strong and reliable interaction between the lottery and X_1 . Notably, only the D-RSMD estimation procedure reveals statistically significant heterogeneous effects. The age group exhibits significant results for both θ_{w0} and θ_{wx0} when employing our new estimator with the valid lottery instrument, suggesting that the effect of Medicaid on debt for health varies depending on income level, thus indicating heterogeneity.

For the age group 35–60, the interpretation of the D-RSMD estimate (using Z_1 or (Z_1, Z_1X_1)) for θ_{w0} is that for people with an income 50% below the federal poverty line, Medicaid enrollment decreases their probability of owing money to health providers by 23.2 log points on average, ceteris paribus. For people with an income above 50% of the federal poverty line, Medicaid enrollment reduces their probability of owing money, but not by that much. This is reasonable because enrollment in Medicaid is not that critical to reducing debt for individuals with higher incomes compared with those with lower incomes.

Now focus on the θ_{wx0} . Comparing the results of D-RSMD between the two instrument sets $(Z_1 \text{ or } (Z_1, Z_1X_1))$, we find that the estimates are close, but the standard errors using Z_1 are substantially lower, suggesting that using one valid instrument provides a different result for statistical significance.

For the age group 21–35, all estimators generate similar results using (Z_1, Z_1X_1) . The coefficients for θ_{w0} are not statistically significant, and the ones for θ_{wx0} are statistically significant. It suggests that when households' incomes are above the 50% federal poverty line, individuals with Medicaid will be less likely to owe money to their health providers. It also suggests that Medicaid helps people with higher income levels more than it helps people with lower incomes. Using only the lottery as the instrument, our new procedure generates the opposite results. The interpretation is that for people with lower incomes, Medicaid enrollment decreases their probability of owing money to health providers by 19.9 log points on average, holding other variables constant. For people with higher incomes, the effect of Medicaid decreases.

Based on the D-RSMD results using only the lottery, we do not find support in the data to say that there are heterogeneous treatment effects for individuals between 21 and 35 years old. Furthermore, we discover that the results for individuals aged 21–35 differ significantly between the two instrument sets. It suggests that the generated interaction $Lottery \times X_1$ is invalid.

Next, we re-estimate a homogeneous model. In a homogeneous model, the interaction term is not included in the regression model. The traditional estimation method only requires one instrument to estimate one parameter. When we examine the homogeneous treatment effects using both instrument sets in Table 3, all estimators yield similar results to the DRSMD-Lasso method using only the lottery as the instrument in the heterogeneous treatment effects panel. This suggests that using a valid instrument to estimate both parameters produces more reliable outcomes. In practical applications where the homogeneity of the model is uncertain, the DRSMD-Lasso method proves to be highly valuable. Its ability to provide reliable estimates under both homogeneous and heterogeneous conditions, coupled with its smaller standard errors compared to traditional methods, makes it a preferred choice. In particular, the smaller standard errors enhance the significance of results obtained through t-tests.

The estimates and standard errors of average treatment effects in Table 4 are calculated based on Table 3. We conduct the robustness check in the Appendix.

Panel A: Heterogeneous treatment effects									
Debt for Health	Age:	21 - 35	Age:	36 - 50	Age:	51 - 64			
Estimator for θ_{w0}	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)			
CMM		-0.069		-0.213***		-0.193***			
GMM		(0.055)		(0.041)		(0.046)			
CODIT		-0.091		-0.259***		-0.245***			
GMM-Lasso		(0.093)		(0.078)		(0.090)			
	-0.199***	-0.053	-0.232***	-0.232***	-0.182**	-0.190***			
DRSMD-Lasso	(0.071)	Age: $21 - 35$ Z_1 Z_1 $(Z_1, Z_1 X_1)$ Z_1 -0.069 (0.055) -0.091 (0.093) 0.093 -0.232 $.071$ (0.065) (0.062) 2071 (0.065) (0.062) 21 $(Z_1, Z_1 X_1)$ Z_1 -0.204^{**} (0.096) -0.184 (0.131) 0.058 -0.252^{**} 0.091 $.036$ (0.105) (0.03) (0.046) $.036$ (0.105) (0.046) (0.046) $.049$ (0.046) (0.046) (0.046) $.180^*$ -0.113 -0.256 $.104$ (0.092) (0.08) $.063$ (0.066) (0.03) (0.066) (0.03)	(0.062)	(0.050)	(0.072)	(0.056)			
Estimator for θ_{wx0}	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)			
0.04		-0.204**		0.108		0.069			
GMM		(0.096)		(0.085)		(0.094)			
CMALL		-0.184		0.008		-0.081			
GMM-Lasso		(0.131)		(0.128)		(0.153)			
DDCMDI	0.058	-0.252**	0.091***	0.088	0.076*	0.096			
DRSMD-Lasso	(0.036)	** -0.053 -0.232*** -0.232***) (0.065) (0.062) (0.050) $(Z_1, Z_1 X_1)$ Z_1 $(Z_1, Z_1 X_1)$ -0.204** 0.108 (0.096) (0.085) -0.184 0.008 (0.131) (0.128) -0.252** 0.091*** 0.088) (0.105) (0.031) (0.096) reatment effects ** -0.130*** -0.170*** -0.189**) (0.046) (0.040) (0.037)	(0.096)	(0.040)	(0.103)				
Panel B: Homoge	eneous trea	tment effec	ts						
CMM	-0.161***	-0.130***	-0.170***	-0.189***	-0.164***	-0.177***			
GMM	(0.049)	(0.046)	(0.040)	(0.037)	(0.044)	(0.041)			
CMM L	-0.180*	-0.113	-0.256***	-0.260***	-0.279**	-0.244***			
GMM-Lasso	(0.104)	(0.092)	(0.097)	(0.078)	(0.112)	(0.089)			
DDCMDI	-0.172***	-0.187***	-0.198***	-0.198***	-0.150**	-0.145**			
DRSMD-Lasso	(0.063)	tment effects 1 - 35 Age: $36 - 50$ $(Z_1, Z_1 X_1)$ Z_1 $(Z_1, Z_1 Z_1)$ -0.069 -0.213* (0.055) (0.041) -0.091 -0.259* (0.093) (0.078) -0.053 -0.232*** (0.065) (0.062) (0.065) (0.062) (0.065) (0.062) (0.065) (0.062) (0.065) (0.062) (0.065) (0.062) (0.066) (0.062) (0.096) (0.085) -0.104^{**} 0.108 (0.131) (0.128) -0.252^{**} 0.091^{***} (0.105) (0.031) (0.096) ment effects -0.170^{***} -0.189^{**} (0.046) (0.040) (0.037) -0.130^{***} -0.198^{***} -0.260^{**} (0.092) (0.097) (0.078) -0.187^{***} -0.198^{***} -0.198^{**} (0.066) (0.056) (0.057)	(0.057)	(0.063)	(0.064)				
N	59	962	6	393	5	917			

Table 3: Heterogeneous Treatment Effects of Medicaid on Debt for Health

Note: *** Significant at 1%, ** at 5%, * at 10%. Each row shows the estimates and robust standard errors for the same type of estimator. In the columns, we present the instruments these estimators used and the age groups. The interaction term is Medicaid \times Above 50% Federal Poverty Line.

Average Treatment	Effects					
Debt for Health	Age:	21 - 35	Age:	36 - 50	Age:	51 - 64
Estimator for LATE	Z_1	(Z_1, Z_1X_1)	Z_1	$(Z_1, Z_1 X_1)$	Z_1	(Z_1, Z_1X_1)
CDDL		-0.190***		-0.150***		-0.149***
GMM		(0.055)		Age: $36 - 50$ Age: $51 - 64$ Z_1 $(Z_1, Z_1 X_1)$ Z_1 (Z_1, Z_1) -0.150^{***} -0.149^3 (0.048) (0.054) -0.255^{**} -0.296 (0.110) $(0.133)^{**}$ $.179^{***}$ -0.181^{***} -0.135^{**} $0.054)$ (0.067) (0.060) 6693 5917	(0.055)	
CLOUT		-0.201*		-0.255**		-0.296**
GMM-Lasso		(0.112)		(0.110)		(0.133)
	-0.164***	-0.202***	-0.179***	-0.181***	-0.135**	-0.129*
DRSMD-Lasso	(0.061)	(0.069)	(0.054)	Age: $36 - 50$ Age: $51 - 64$ $(Z_1, Z_1 X_1)$ Z_1 $(Z_1, Z_1 Z_1)$ -0.150^{***} -0.149^* (0.048) $(0.055$ -0.255^{**} -0.296° (0.110) $(0.133)^{***}$ $*^{***}$ -0.181^{***} -0.135^{**} 6693 5917	(0.075)	
N	59	962	6	693	5	917

Table 4: Heterogeneous Treatment Effects of Medicaid on Debt for Health

Note: *** Significant at 1%, ** at 5%, * at 10%. Each row shows the estimates and robust standard errors for the same type of estimator. In the columns, we present the instruments these estimators used and the age groups. The interaction term is Medicaid × Above 50% Federal Poverty Line. The expression for LATE is $\theta_{w0} + \theta_{wx0}E(X)$.

References

- Ai, C. and X. Chen (2003). Efficient estimation of models with conditional moment restrictions containing unknown functions. *Econometrica* 71(6), 1795–1843.
- Angrist, J. D. and A. B. Keueger (1991). Does compulsory school attendance affect schooling and earnings? *The Quarterly journal of economics* 106(4), 979–1014.
- Antoine, B. and P. Lavergne (2014). Conditional moment models under semi-strong identification. Journal of Econometrics 182(1), 59 – 69. Causality, Prediction, and Specification Analysis: Recent Advances and Future Directions.
- Antoine, B. and X. Sun (2021). Partially Linear Models with Endogeneity: a conditional moment based approach. *Accepted at The Econometrics Journal*, to appear.
- Ashenfelter, O. and C. Rouse (1998). Income, schooling, and ability: Evidence from a new sample of identical twins. *The Quarterly journal of economics* 113(1), 253–284.
- Baicker, K., A. Finkelstein, J. Song, and S. Taubman (2014). The impact of medicaid on labor market activity and program participation: Evidence from the oregon health insurance experiment. *The American economic review 104*(5), 322–328.
- Bierens, H. (1982). Consistent model specification tests. Journal of Econometrics 20(1), 105–134.
- Bose, A. and S. Chatterjee (2018). U-Statistics, Mm-Estimators and Resampling. Springer Singapore.
- Card, D. (1993, October). Using geographic variation in college proximity to estimate the return to schooling. Working Paper 4483, National Bureau of Economic Research.
- Champeney, D. C. (1987). A Handbook of Fourier Theorems. Cambridge University Press.
- Chen, X. (2007). Large sample sieve estimation of semi-nonparametric models. In J. Heckman andE. Leamer (Eds.), *Handbook of Econometrics* (1 ed.), Volume 6B, Chapter 76. Elsevier.

- Chernozhukov, V., D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins (2018). Double/debiased machine learning for treatment and structural parameters. *The econometrics journal* 21(1), C1–C68.
- Dieterle, S. G. and A. Snell (2016). A simple diagnostic to investigate instrument validity and heterogeneous effects when using a single instrument. *Labour Economics* 42(C), 76–86.
- Dominguez, M. and I. Lobato (2004). Consistent estimation of models defined by conditional moment restrictions. *Econometrica* 72(5), 1601–1615.
- Engle, R. F., C. W. J. Granger, J. Rice, and A. Weiss (1986). Semiparametric estimates of the relation between weather and electricity sales. *Journal of the American Statistical Association* 81(394), 310–320.
- Finkelstein, A., S. Taubman, B. Wright, M. Bernstein, J. Gruber, J. P. Newhouse, H. Allen, and K. Baicker (2012). The oregon health insurance experiment: Evidence from the first year. *The Quarterly journal of economics* 127(3), 1057–1106.
- Florens, J.-P., J. Johannes, and S. V. Bellegem (2012, September). Instrumental regression in partially linear models. *The Econometrics Journal* 15(2), 304–324.
- Garen, J. (1984). The returns to schooling: A selectivity bias approach with a continuous choice variable. *Econometrica* 52(5), 1199–1218.
- Hansen, L. (1982). Large sample properties of generalized method of moments estimators. *Econo*metrica 50(4), 1029–54.
- Hayashi, F. (2000). *Econometrics*. Princeton, NJ [u.a.]: Princeton Univ. Press.
- Heckman, J. and E. Vytlacil (1998). Instrumental variables methods for the correlated random coefficient model: Estimating the average rate of return to schooling when the return is correlated with schooling. *The Journal of Human Resources* 33(4), 974–987.

- Heckman, J. J., H. Ichimura, and P. E. Todd (1997). Matching as an econometric evaluation estimator: Evidence from evaluating a job training programme. *The Review of economic studies* 64(4), 605–654.
- Hoeffding, W. (1948). A class of statistics with asymptotically normal distribution. The Annals of mathematical statistics 19(3), 293–325.
- Imbens, G. W. and J. D. Angrist (1994). Identification and estimation of local average treatment effects. *Econometrica* 62(2), 467–475.
- Imbens, G. W. and D. B. Rubin (2015). Causal Inference for Statistics, Social, and Biomedical Sciences: An Introduction. Cambridge University Press.
- Kitagawa, T. (2015). A test for instrument validity. *Econometrica* 83(5), 2043–2063.
- Lavergne, P. and V. Patilea (2013). Smooth minimum distance estimation and testing with conditional estimating equations: Uniform in bandwidth theory. *Journal of Econometrics* 177, 47–59.
- Li, Q. (1996). On the root-n-consistent semiparametric estimation of partially linear models. *Economics Letters* 51(3), 277–285.
- Li, Q. and J. S. Racine (2006). *Nonparametric Econometrics: Theory and Practice*. Princeton University Press.
- Nekipelov, D., V. Semenova, and V. Syrgkanis (2018). Regularized orthogonal machine learning for nonlinear semiparametric models.
- Otsu, T. (2011). Empirical likelihood estimation of conditional moment restriction models with unknown functions. *Econometric Theory* 27(1), 8–46.
- Pakes, A. and D. Pollard (1989). Simulation and the asymptotics of optimization estimators. *Econometrica* 57(5), 1027–1057.

- Patilea, V. and P. Lavergne (2013). Smooth minimum distance estimation and testing with conditional estimating equations: Uniform in bandwidth theory. *Econometrics* 177(1), 47–59.
- Robinson, P. (1988, July). Root-n-consistent semiparametric regression. *Econometrica* 56(4), 931–954.
- Serfling, R. J. (1980). Approximation theorems of mathematical statistics. Wiley series in probability and mathematical statistics. John Wiley and Sons, Inc.
- Sherman, R. P. (1994, 03). Maximal inequalities for degenerate u-processes with applications to optimization estimators. The Annals of Statistics 22(1), 439–459.
- Syrgkanis, V., V. Lei, M. Oprescu, M. Hei, K. Battocchi, and G. Lewis (2019). Machine learning estimation of heterogeneous treatment effects with instruments.
- van de Geer, S. (2016). Estimation and Testing Under Sparsity École d'Été de Probabilités de Saint-Flour XLV 2015 / by Sara van de Geer. (1st ed. 2016. ed.). École d'Été de Probabilités de Saint-Flour, 2159. Springer International Publishing : Imprint: Springer.
- Wooldridge, J. (1997). On two stage least squares estimation of the average treatment effect in a random coefficient model. *Economics Letters* 56(2), 129–133.
- Wooldridge, J. M. (2004). Estimating average partial effects under conditional moment independence assumptions. cemmap working paper CWP03/04, London.
- Wooldridge, J. M. (2010). Econometric analysis of cross section and panel data / Jeffrey M. Wooldridge. (2nd ed. ed.). MIT Press.
- Yanagi, T. (2019). Inference on local average treatment effects for misclassified treatment. *Econo*metric reviews 38(8), 938–960.

A Appendix: Additional Results of Monte Carlo Simulation

A.1 The Advantage of the D-RSMD Estimator vs. Splitting the Sample

Table 5 presents simulation results using a binary covariate in the interaction term. This simulation follows a similar DGP as the one in the simulation section, except that X_1 is a binary variable. From the table, the D-RSMD estimator provides much better estimation in terms of MAD and Med.SE than the GMM with the correct model specification (GMM (Oracle)). However, without the correct model specification, RGMM (which incorporates Lasso to approximate the unknown nonparametric part and then GMM) does not perform as effectively. The situation worsens with fewer observations and when the mean of X_1 is closer to 0 or 1. Here, $E(X_1) = 0.2$. This is shown when we use the subsets of the dataset, for instance, the rows when $X_1 = 1$.

n = 3,000 and there are 30 covariates												
				θ_w	0		θ_{wx0}					
Dataset	Estimator	Instrument	Med.Bias	MAD	Med.SE	RR	Med.Bias	MAD	Med.SE	RR		
Full dataset	D-RSMD	Z_1	0.056	0.219	0.327	0.049	-0.053	0.120	0.180	0.058		
Rows when $X_1 = 1$	RGMM	Z_1	-0.293	0.413	0.962	0.003						
Rows when $X_1 = 0$	RGMM	Z_1	0.676	0.295	0.885	0.003						
Full dataset	RGMM	(Z_1, Z_1X_1)	0.592	0.326	0.817	0.003	-0.822	0.180	0.412	0.527		
Rows when $X_1 = 1$	GMM (Oracle)	Z_1	-0.006	0.409	0.575	0.044						
Rows when $X_1 = 0$	GMM (Oracle)	Z_1	0.004	0.265	0.396	0.046						
Full dataset	GMM (Oracle)	(Z_1, Z_1X_1)	0.005	0.234	0.347	0.048	-0.003	0.155	0.228	0.053		

Table 5: Performance of Estimators with and without Splitting the Sample

Note: Simulation Results for θ_{w0} and θ_{wx0} in the benchmark model with Z_1 as the Instrument inside DGP. There are 5,000 replications. We report the Monte-Carlo Median Bias (Med.Bias), Median Absolute Deviation (MAD), median of asymptotic standard error under heteroskedasticity (Med.SE), and Rejection Rate (RR) using a 5% t-test. Z_1 is the binary instrument. X_1 is the binary covariate with $P_r(X_1 = 1) = 0.2$. The dataset is splitted for different X_1 values when employing GMM and RGMM to handle the heterogeneity of the treatment.

A.2 The Advantage of the D-RSMD Estimator vs. Using Another Instrument Set

Next, we demonstrate the performance of the GMM (Oracle) estimator using a pair of the instrument and the interaction term between a predicted endogenous variable and a covariate, while avoiding the use of the interaction term between the instrument and the covariate. This approach yields favourable results; however, the D-RSMD estimator still achieves competitive performance. In scenarios where the model specification is unknown, the RGMM estimator using $(Z_1, \hat{E}(W|X)X_1)$ exhibits higher bias, MAD, Med.SE, and a larger rejection rate (RR), as illustrated in Table 6.

				0	$ heta_{wx0}$					
	Estimator	Instrument	Med.Bias	MAD	Med.SE	RR	Med.Bias	MAD	Med.SE	RR
n = 3,000	D-RSMD	Z_1	0.028	0.288	0.427	0.045	-0.012	0.045	0.059	0.064
P = 30	RGMM	$(Z_1, \hat{E}(W X)X_1)$	0.196	0.358	0.812	0.002	-0.166	0.092	0.178	0.130
	GMM (Oracle)	$(Z_1, \hat{E}(W X)X_1)$	0.010	0.252	0.372	0.050	-0.001	0.049	0.073	0.051
	GMM (Oracle)	(Z_1, Z_2)	-0.011	0.635	1.424	0.003	0.114	2.502	5.548	0.000

Table 6: Performance of Estimators with and without using an additional instrument variable

Note: Simulation Results for θ_{w0} and θ_{wx0} in the benchmark model with Z_1 as the Instrument inside DGP. There are 5,000 replications. We report the Monte-Carlo Median Bias (Med.Bias), Median Absolute Deviation (MAD), median of asymptotic standard error under heteroskedasticity (Med.SE), and Rejection Rate (RR) using a 5% t-test. Z_2 : the sum of Z_1 and another binary variable. There are three values in Z_2 . $(Z_1, \hat{E}(W|X)X_1)$: pair consisting of Z_1 and the interaction term $\hat{E}(W|X)X_1$ (where $\hat{E}(W|X)$ is the estimated conditional expectation and X_1 is a covariate). (Z_1, Z_2) : pair including both Z_1 and Z_2 . The correlation between Z_1 and Z_2 is around 0.7.

A.3 Results for the Model with a Categorical Instrument

In this section, we introduce a categorical instrument Z with three values. Specifically, for the benchmark model, we generate the endogenous treatment using Z_2 .

When the number of covariates is small, for example, $q_X = 1$, we apply the Nadaraya-Watson estimator with a rule-of-thumb bandwidth $h = \sigma_x n^{-0.2}$ to estimate the nuisance parameters for the R-SMD and R-GMM estimators. In our setting with n = 3,000 and $q_X = 1$, the Nadaraya-Watson estimators for nuisance parameters use a bandwidth of $0.202\hat{\sigma}_x$. For the D-RSMD, we use the Lasso method to estimate the nuisance parameters. Both D-RSMD and R-SMD estimators utilize the CDF of a standard Gaussian distribution as $\mu(\cdot)$ inside $\kappa(Z_j - Z_l) = \int_{\mathbb{R}^{q_z}} e^{it'(Z_j - Z_l)} d\mu(t)$.

Table 7 presents the simulation results of D-RSMD, R-SMD, and GMM-type estimators for θ_{w0} and θ_{wx0} . The sample size is 3,000 with 5,000 replications and only one covariate ($q_X = 1$). The table displays results for D-RSMD and R-SMD estimators using one instrument, either Z_1 or Z_2 , and two instruments, namely (Z_1, Z_1X_1) or (Z_2, Z_2X_1). GMM-type estimators utilize pairs of instruments: (Z_1, Z_1X_1) or (Z_2, Z_2X_1).

Recall that Z_2 is the instrument used to generate the treatment variable. Intuitively, estimators that employ Z_2 or (Z_2, Z_2X_1) are expected to yield better results compared to those using Z_1 or (Z_1, Z_1X_1) . This is indeed evident from Table 7, particularly for R-GMM, GMM, and GMM (Oracle) estimators. However, for D-RSMD and RSMD, this is not consistently observed. For instance, the MAD and Med.SE for D-RSMD with Z_1 are smaller than those with Z_2 for both parameters, although this property may be case-specific and does not persist in the subsequent case with $q_X = 30$. SMD-type estimators generally use all available information from the instrument. These estimators remain effective when using a closely related instrument.

Comparing D-RSMD and GMM (Oracle), when both employ (Z_2, Z_2X_1) , we find that their results are similar. This suggests that D-RSMD with (Z_2, Z_2X_1) performs comparably to GMM (Oracle). Notably, D-RSMD exhibits a lower median bias than R-SMD with (Z_2, Z_2X_1) , indicating the effectiveness of bias reduction in D-RSMD. The Med.SE for D-RSMD is also smaller than that for R-SMD, and the rejection rate (RR) for D-RSMD is closer to 5% compared to R-SMD.

Next, we consider the scenario with $q_X = 30$. We set the sparsity level to 10, where $\alpha_{3q} = 2$, $\beta_{2q} = -3$, and $\alpha_{1q} = \beta_{1q} = 1$ for $q \leq 5$. We continue to use the benchmark model in our simulation study.

Table 8 presents simulation results for D-RSMD, RSMD, and GMM type estimators with sample sizes $n \in \{2000, 3000, 5000\}$, each based on 5000 replications.

In the second panel of the table, D-RSMD estimators using Z_2 generally exhibit better perfor-

			θ_w	0					
Estimator	Instrument	Med.Bias	MAD	Med.SE	RR	Med.Bias	MAD	Med.SE	RR
D-RSMD	Z_1	-0.007	0.069	0.103	0.047	0.019	0.036	0.052	0.067
D-RSMD	Z_2	-0.017	0.129	0.211	0.035	0.034	0.313	0.523	0.035
D-RSMD	(Z_1, Z_1X_1)	0.002	0.077	0.112	0.050	-0.008	0.089	0.130	0.044
D-RSMD	(Z_2, Z_2X_1)	0.001	0.045	0.065	0.048	-0.006	0.055	0.077	0.054
R-SMD	Z_1	-0.207	0.072	0.236	0.007	0.424	0.065	0.449	0.000
R-SMD	Z_2	0.027	0.172	0.280	0.022	-0.060	0.416	0.693	0.018
R-SMD	(Z_1, Z_1X_1)	0.064	0.077	0.127	0.052	-0.112	0.083	0.220	0.003
R-SMD	(Z_2, Z_2X_1)	0.028	0.039	0.072	0.031	-0.053	0.048	0.076	0.079
R-GMM	(Z_1, Z_1X_1)	-0.044	0.076	0.182	0.003	-0.239	0.089	0.245	0.026
R-GMM	$(\mathbb{Z}_2, \mathbb{Z}_2 \mathbb{X}_1)$	-0.015	0.042	0.105	0.001	-0.172	0.053	0.156	0.043
GMM	(Z_1, Z_1X_1)	2.571	0.318	0.472	1.000	-6.003	0.415	0.618	1.000
GMM	$(\mathbb{Z}_2, \mathbb{Z}_2 \mathbb{X}_1)$	2.176	0.231	0.322	1.000	-5.233	0.319	0.454	1.000
GMM (Oracle)	(Z_1, Z_1X_1)	0.000	0.076	0.113	0.048	-0.003	0.093	0.137	0.045
GMM (Oracle)	(Z_2, Z_2X_1)	0.000	0.042	0.061	0.049	0.000	0.053	0.078	0.057

Table 7: Performance of Estimators with a Categorical Instrument in the DGP when $q_X = 1$

Note: Simulation Results for θ_{w0} and θ_{wx0} in the benchmark model with Z_2 in the DGP. There are 5,000 replications. The sample size is 3,000. We report the Monte-Carlo Median Bias (Med.Bias), Median Absolute Deviation (MAD), the median of asymptotic standard error under heteroskedasticity (Med.SE), and Rejection Rate (RR) using a 5% t-test. Z_1 : binary instrument. Z_2 : variable representing the sum of Z_1 and another binary variable, resulting in three possible values for Z_2 . (Z_1, Z_1X_1) : pair consisting of Z_1 and the interaction term Z_1X_1 (where X_1 is a covariate). (Z_2, Z_2X_1) : pair comprising Z_2 and the interaction term Z_2X_1 . (Z_1, Z_2) : pair including both Z_1 and Z_2 . The correlation between Z_1 and Z_2 is around 0.7.

mance than those using Z_1 , showing smaller Med.Bias, MAD, and Med.SE when estimating θ_{w0} . However, for θ_{wx0} , D-RSMD with Z_1 yields competitive estimates.

Comparing D-RSMD and GMM type estimators in the 3000 observation panel of Table 8, GMM (Oracle) demonstrates relatively better performance with the instrument set (Z_2, Z_2X_1) , characterized by lower Med.Bias and MAD. Nonetheless, D-RSMD using the same instrument set still produces promising results.

It's worth noting that the GMM (Oracle) estimator encounters challenges with the instrument

				θ_w	0			$ heta_{wx0}$			
	Estimator	Instrument	Med.Bias	MAD	Med.SE	RR	Med.Bias	MAD	Med.SE	RR	
n = 2,000	D-RSMD	Z_2	-0.041	0.115	0.146	0.064	-0.019	0.106	0.158	0.064	
	D-RSMD	$(\mathbb{Z}_2,\mathbb{Z}_2\mathbb{X}_1)$	-0.059	0.119	0.144	0.072	-0.033	0.079	0.101	0.064	
	GMM (Oracle)	$(\mathbb{Z}_2, \mathbb{Z}_2 \mathbb{X}_1)$	-0.006	0.110	0.160	0.050	0.001	0.059	0.087	0.053	
n = 3,000	D-RSMD	Z_1	0.139	0.765	1.140	0.012	-0.013	0.091	0.158	0.020	
	D-RSMD	Z_2	-0.018	0.119	0.158	0.044	-0.019	0.145	0.236	0.048	
	D-RSMD	(Z_1, Z_1X_1)	-0.115	0.557	0.843	0.025	-0.015	0.105	0.154	0.037	
	D-RSMD	$(\mathbb{Z}_2, \mathbb{Z}_2 \mathbb{X}_1)$	-0.028	0.100	0.118	0.055	-0.023	0.068	0.083	0.059	
	D-RSMD	(Z_1, Z_2)	-0.019	0.189	0.267	0.036	-0.018	0.274	0.435	0.034	
	RSMD	Z_1	-0.476	0.773	1.173	0.024	-0.049	0.101	0.161	0.052	
	RSMD	Z_2	-0.134	0.178	0.313	0.083	-0.052	0.231	0.421	0.058	
	RSMD	(Z_1, Z_1X_1)	0.082	0.790	1.158	0.015	-0.279	0.135	0.260	0.071	
	RSMD	$(\mathbb{Z}_2,\mathbb{Z}_2\mathbb{X}_1)$	-0.202	0.111	0.190	0.157	-0.074	0.054	0.085	0.129	
	RSMD	(Z_1, Z_2)	-0.128	0.267	0.411	0.073	-0.058	0.374	0.602	0.045	
	GMM	$(\mathbb{Z}_2,\mathbb{Z}_2\mathbb{X}_1)$	3.379	0.897	1.354	0.715	-5.957	0.509	0.726	1.000	
	GMM	(Z_1, Z_2)	12.082	4.798	7.355	0.230	-26.653	9.961	15.090	0.312	
	GMM (Oracle)	$(\mathbb{Z}_2, \mathbb{Z}_2 \mathbb{X}_1)$	0.000	0.089	0.130	0.056	0.000	0.048	0.071	0.052	
	GMM (Oracle)	(Z_1, Z_2)	0.003	0.506	0.802	0.004	0.015	1.065	1.712	0.002	
n = 5,000	D-RSMD	Z_2	-0.005	0.397	0.694	0.021	-0.028	0.714	1.267	0.022	
	D-RSMD	$(\mathbb{Z}_2, \mathbb{Z}_2 \mathbb{X}_1)$	-0.016	0.082	0.096	0.053	-0.013	0.054	0.066	0.056	
	GMM (Oracle)	$(\mathbb{Z}_2, \mathbb{Z}_2 \mathbb{X}_1)$	-0.001	0.068	0.101	0.050	-0.001	0.037	0.055	0.049	

Table 8: Performance of Estimators with a Categorical Instrument in the DGP when $q_X = 30$

Note: Simulation Results for θ_{w0} and θ_{wx0} in the benchmark model with Z_2 in the DGP. There are 5,000 replications with the sample size $n \in \{2000, 3000, 5000\}$. We report the Monte-Carlo Median Bias (Med.Bias), Median Absolute Deviation (MAD), median of asymptotic standard error under heteroskedasticity (Med.SE), and Rejection Rate (RR) using a 5% t-test. Z_1 : binary instrument. Z_2 : variable representing the sum of Z_1 and another binary variable, resulting in three possible values for Z_2 . (Z_1, Z_1X_1) : pair consisting of Z_1 and the interaction term Z_1X_1 (where X_1 is a covariate). (Z_2, Z_2X_1) : pair comprising Z_2 and the interaction term Z_2X_1 . (Z_1, Z_2) : pair including both Z_1 and Z_2 . The correlation between Z_1 and Z_2 is around 0.7.

set (Z_1, Z_2) , which is less problematic for the D-RSMD estimator. This is the same observation as that for the previous case with $q_X = 1$.

With a categorical instrument in the DGP, the D-RSMD estimator generates a smaller bias
compared to RSMD estimators. D-RSMD's flexibility allows for estimation with various instrument sets. Notably, D-RSMD remains competitive compared to GMM (Oracle) estimators.

With a categorical instrument in the DGP, the D-RSMD estimators have the proper size and a smaller bias than the RSMD estimators. As the sample size grows, the size distortions decrease for (Z_2, Z_2X_1) . D-RSMD allows us to use any possible instrument set to estimate heterogeneous treatment effects. Compared to GMM (Oracle), the D-RSMD estimator still generates competitive results.

B Appendix: Additional Results of Empirical Application

B.1 Robustness Check

In Section 5.1, the interaction term includes a covariate indicating whether household income exceeds 50% of the Federal Poverty Line. To offer further insights into these regressions, we conduct various robustness checks. The first subsection examines estimation results when the covariate is 1 for household income exceeding 100% or 150% of the Federal Poverty Line. The second subsection provides a brief overview of other robustness checks.

• Estimation Results with Other Covariates

The results are included in Table 9 when X_1 is an indicator for "Income Above 100% Federal Poverty Line". Comparing Table 9 with Table 3, we find that the results are quite similar. For individuals between 36 and 50 years old, there are heterogeneous treatment effects. The estimates show that they benefit more from Medicaid health coverage when they have lower incomes. We do not find support in the data to say that there are heterogeneous treatment effects for individuals between 21 and 35 years old, and the DRSMD-Lasso esimator using only one valid instrument variable generates more reliable results.

Table 10 reports the outcomes when X_1 is an indicator for "Income Above 150% Federal Poverty Line". Results in Tables 10 and 9 are similar in estimates but different in standard

Debt for Health	Age:	21 - 35	Age:	36 - 50	Age: 51 - 64		
Estimator for θ_{w0}	Z_1	(Z_1, Z_1X_1)	Z_1	$(Z_1, Z_1 X_1)$	Z_1	$(Z_1, Z_1 X_1)$	
CLOU		-0.123**		-0.179***		-0.142***	
GMM		(0.048)		(0.038)		(0.041)	
	-0.178***	-0.123**	-0.202***	-0.200***	-0.164**	-0.131	
DRSMD-Lasso	(0.063)	(0.061)	(0.057)	(0.049)	(0.065)	(0.053)	
Estimator for θ_{wx0}	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	
CMM		-0.200		0.072		-0.156	
GMM		(0.148)		(0.159)		(0.185)	
	0.023	-0.212	0.102***	0.032	0.096	-0.154	
DRSMD-Lasso	(0.033)	(0.154)	(0.038)	(0.192)	(0.044)	(0.217)	
Ν	5	962	6	693	5917		

Heterogeneous Treatment Effects of Medicaid on Debt for Health

Table 9: Income above 100% Federal Poverty Line

Note: *** Significant at 1%, ** at 5%, * at 10%.

errors. In the table, estimation results from using a dummy for income above 150% Federal Poverty Line show that the treatment effects of Medicaid on debt for people between 36 and 50 are not supported by the data to be heterogeneous because the estimate for the parameter in front of the interaction term is not statistically significant at the 5% significance level.

The effects of Medicaid on debt for people between 21 and 35 are heterogeneous because the estimate is statistically significant. The difference in the results using two instrument sets for the DRSMD-Lasso estimators suggests that using only the lottery variable as an instrument generates more reliable results. The difference between Tables 10 and 9 shows that the covariate inside the interaction is important for the estimation results. It can be explained by the fact that there are only 808 individuals with incomes above 150% Federal Poverty Line for people between 36 and 50 and 794 individuals for people between 21 and 35 in this data set.

Debt for Health	Age:	21 - 35	Age:	36 - 50	Age: 51 - 64		
Estimator for θ_{w0}	Z_1	$(Z_1, Z_1 X_1)$	Z_1	$(Z_1, Z_1 X_1)$	Z_1	(Z_1, Z_1X_1)	
CLOU		-0.145***		-0.179***		-0.145***	
GMM		(0.048)		(0.038)		(0.042)	
	-0.181***	-0.149**	-0.207***	-0.205***	-0.154**	-0.121**	
DRSMD-Lasso	(0.062)	(0.062)	(0.056)	(0.053)	(0.063)	(0.057)	
Estimator for θ_{wx0}	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	
CMM		-0.204		0.195		-0.521	
GMM		(0.245)		(0.340)		(0.426)	
	0.095*	-0.254	0.067	0.030	0.078	-0.620	
DRSMD-Lasso	(0.055)	(0.292)	(0.061)	(0.453)	(0.067)	(0.559)	
Ν	5	962	6	693	5917		

Heterogeneous Treatment Effects of Medicaid on Debt for Health

Table 10: Income above 150% Federal Poverty Line

Note: *** Significant at 1%, ** at 5%, * at 10%.

• Other Robustness Checks

We present additional robustness checks. One check focuses on the variable TANF, which exhibits limited variation with only 2% of individuals receiving it. We examine whether TANF contributes to heterogeneity. Table 11 displays the results for the Debt for Health variable across five age groups. The DRSMD-Lasso estimator with Z_1 yields statistically significant estimates of -0.215 for θ_{w0} and 0.307 for θ_{wx0} at a 5% significance level in the first column. This suggests that, for individuals aged 21 to 29, TANF mitigates the negative effect of Medicaid on the likelihood of having debt. The new method, utilizing the valid instrument Z_1 continues to produce reliable results with the lowest standard error.

Tables 12 and 13 display the effects of treatment on employment and out-of-pocket costs, respectively. Both tables indicate heterogeneous treatment effects among young individuals (aged 21 to 29) when X_1 is a binary variable representing "Income Above 50% of the Federal Poverty Line."

Panel A: Heterogeneous treatment effects										
Debt for Health	Age:	21 - 29	Age: 30 - 38		Age: 39 - 47		Age: 48 - 56		Age: 57 - 64	
Estimator for θ_{w0}	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	$(Z_1, Z_1 X_1)$	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)
CLDI		-0.193***		-0.199***		-0.113**		-0.196***		-0.157**
GMM		(0.065)		(0.061)		(0.049)		(0.046)		(0.068)
CADIT		-0.270*		-0.216		-0.211*		-0.351***		-0.295
GMM-Lasso		(0.147)		(0.136)		(0.110)		(0.112)		(0.182)
	-0.215**	-0.217**	-0.137*	-0.010	-0.148**	-0.135**	-0.199***	-0.192***	-0.139	-0.141
DRSMD-Lasso	(0.087)	(0.103)	(0.082)	(0.337)	(0.068)	(0.068)	(0.062)	(0.062)	(0.107)	(0.107)
Estimator for θ_{wx0}	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	$(Z_1, Z_1 X_1)$	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)
CLDI		0.212		1.959		-0.791		2.059		3.074
GMM		(0.900)		(1.353)		(0.652)		(2.040)		(3.827)
0.011		0.983		1.904		-0.785		1.083		0.818
GMM-Lasso		(1.099)		(1.348)		(0.956)		(1.844)		(1.164)
DDCMD I	0.307**	0.777	0.042	-9.902	-0.320***	-1.448	0.470	-1.725	-0.048	-0.281
DRSMD-Lasso	(0.142)	(4.239)	(0.153)	(19.627)	(0.085)	(1.554)	(0.417)	(1.821)	(0.522)	(0.590)
N	3	504		3596	3	941	4	599		2932

Table 11: Heterogeneous Treatment Effects of Medicaid on Debt for Health

Note: *** Significant at 1%, ** at 5%, * at 10%. Each row shows the estimates and robust standard errors for the same type of estimator. In the columns, we present the instruments these estimators used and the age groups. The interaction term is $Medicaid \times TANF$.

Employ	Age:	21 - 29	Age: 30 - 38		Age: 39 - 47		Age: 48 - 56		Age: 57 - 64	
Estimator for θ_{w0}	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	$(Z_1, Z_1 X_1)$	Z_1	(Z_1, Z_1X_1)
CMM		0.134^{*}		-0.042		0.007		-0.063		0.022
GMM		(0.071)		(0.067)		(0.048)		(0.040)		(0.064)
CMML		0.143		-0.125		-0.069		-0.105		-0.063
GMM-Lasso		(0.112)		(0.118)		(0.089)		(0.077)		(0.123)
DDCMD Lassa	0.255***	0.159**	0.014	-0.067	0.010	-0.001	0.033	-0.058	0.086	0.072
DRSMD-Lasso	(0.093)	(0.081)	(0.087)	(0.085)	(0.071)	(0.060)	(0.066)	(0.049)	(0.103)	(0.086)
Estimator for θ_{wx0}	Z_1	$(\mathbb{Z}_1,\mathbb{Z}_1\mathbb{X}_1)$	Z_1	(Z_1, Z_1X_1)	Z_1	$(\mathbb{Z}_1,\mathbb{Z}_1\mathbb{X}_1)$	Z_1	$(\mathbb{Z}_1,\mathbb{Z}_1\mathbb{X}_1)$	Z_1	(Z_1, Z_1X_1)
CMM		-0.070		0.066		-0.025		0.159		-0.046
GIMIN		(0.128)		(0.120)		(0.100)		(0.104)		(0.127)
CMM Losso		-0.014		0.198		0.120		0.304^{*}		0.157
GIMM-Lasso		(0.179)		(0.145)		(0.150)		(0.162)		(0.193)
DRSMD Loggo	-0.233***	-0.003	-0.092**	0.073	0.006	0.040	-0.083**	0.169	-0.018	0.012
DRSMD-Lasso	(0.043)	(0.145)	(0.039)	(0.135)	(0.037)	(0.113)	(0.036)	(0.113)	(0.041)	(0.153)
Ν	3	504	3	3596	3941		4	1599	2932	

Panel A: Heterogeneous treatment effects

Table 12: Heterogeneous Treatment Effects of Medicaid on Employ

Note: *** Significant at 1%, ** at 5%, * at 10%. Each row shows the estimates and robust standard errors for the same type of estimator. In the columns, we present the instruments these estimators used and the age groups. The interaction term is Medicaid \times Above 50% Federal Poverty Line.

i anei 21. Heterog	cheous treat	ment encets								
Out of Pocket Cost	Age: 2	21 - 29	Age:	30 - 38	Age:	39 - 47	Age:	48 - 56	Age:	57 - 64
Estimator for θ_{w0}	Z_1	(Z_1, Z_1X_1)	Z_1	$(Z_1, Z_1 X_1)$	Z_1	(Z_1, Z_1X_1)	Z_1	$(Z_1, Z_1 X_1)$	Z_1	(Z_1, Z_1X_1)
0.01		-238.166**		-294.692***		-32.148		-149.770^{***}		-153.110*
GMM		(94.947)		(92.333)		(69.362)		(53.074)		(87.341)
0.011		-545.537***		-569.684***		-276.342**		-321.387***		-465.763**
GMM-Lasso		(148.534)		(141.359)		(131.711)		(91.886)		(189.849)
DDCMD L	-373.155***	-250.372**	-153.656	-376.780***	-55.406	18.780	-162.227	-185.345***	262.158	-49.106
DRSMD-Lasso	(136.072)	(119.078)	(142.101)	(126.053)	(99.185)	(84.691)	(101.883)	(68.418)	(197.082)	(114.506)
Estimator for θ_{wx0}	Z_1	(Z_1, Z_1X_1)	Z_1	$(\mathbb{Z}_1,\mathbb{Z}_1\mathbb{X}_1)$	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)	Z_1	(Z_1, Z_1X_1)
CUDI		-101.939		384.290**		-269.238*		155.502		173.800
GMM		(193.546)		(181.372)		(156.752)		(165.046)		(211.235)
CADALT		226.006		558.392**		-77.050		307.058		477.385
GMM-Lasso		(245.546)		(243.326)		(209.610)		(276.655)		(370.165)
DDONDI	104.666^{*}	-179.667	46.142	496.881**	-72.394*	-242.417	6.897	72.042	-111.454*	502.058*
DRSMD-Lasso	(62.096)	(214.131)	(53.675)	(214.081)	(43.318)	(167.287)	(45.935)	(181.569)	(57.602)	(295.269)
Ν	35	04	3	3596	3	941	4599		2932	

Panel A: Heterogeneous treatment effects

Table 13: Heterogeneous Treatment Effects of Medicaid on Out of Pocket Cost

Note: *** Significant at 1%, ** at 5%, * at 10%. Each row shows the estimates and robust standard errors for the same type of estimator. In the columns, we present the instruments these estimators used and the age groups. The interaction term is Medicaid \times Above 50% Federal Poverty Line.

C Appendix: Proofs of the Theoretical Results

C.1 Equivalence between the Objective Functions (5) and (6)

Proof. The objective function (5) can be written as

$$M_{\infty}(\theta,g) = \int_{\mathbb{R}^{q_z}} E[\epsilon_j(\theta,g)e^{it'Z_j}]E[\epsilon_l(\theta,g)e^{-it'Z_l}]d\mu(t)$$

From Assumption 1(e), for all $j \neq l$, $Cov(\epsilon_j(\theta, g)e^{it'Z_j}, \epsilon_l(\theta, g)e^{-it'Z_l}) = 0$. Thus, for all $j \neq l$, we have:

$$M_{\infty}(\beta) = \int_{\mathbb{R}^{q_z}} E(\epsilon_j(\theta, g) e^{it'Z_j} \epsilon_l(\theta, g) e^{-it'Z_l}) d\mu(t)$$

$$= \int_{\mathbb{R}^{q_z}} E(\epsilon_j(\theta, g) \epsilon_l(\theta, g) e^{it'(Z_j - Z_l)}) d\mu(t)$$

$$= E(\int_{\mathbb{R}^{q_z}} \epsilon_j(\theta, g) \epsilon_l(\theta, g) e^{it'(Z_j - Z_l)} d\mu(t))$$

Thus, the objective function becomes

$$M_{\infty}(\beta) = E(\epsilon_j(\theta, g)\epsilon_l(\theta, g)\kappa_{j,l})$$

where $\kappa_{j,l} = k(Z_j - Z_l) = \int_{\mathbb{R}^{q_z}} e^{it'(Z_j - Z_l)} d\mu(t)$. And k(u) is the inverse Fourier transform of $d\mu(t)$ with $u = Z_j - Z_l$.

C.2 Proof of Proposition 1

The orthogonal FOC in Equation (9) implies that

$$E\left[\left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right)(\tilde{y}_{l} - \tilde{P}_{l}^{\prime}\theta)\kappa_{j,l}\right] = 0$$

$$E\left[\kappa_{j,l}\left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right)\tilde{y}_{l} - \kappa_{j,l}\left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right)\tilde{P}_{l}^{\prime}\theta\right] = 0$$

$$\theta_{0} = E\left[\kappa_{j,l}\left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right)\tilde{P}_{l}^{\prime}\right]^{-1}E\left[\kappa_{j,l}\left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right)\tilde{y}_{l}\right]$$

The proof for invertibility of $E\left[\kappa_{j,l}\left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\right)\tilde{P}'_l\right]$ follows the same steps of proofs for identification in Antoine and Sun (2021).

C.3 Proof of Orthogonal Properties of Equation (8) and (9)

In this section, we check the orthogonal properties of the two FOCs.

$$M_{\infty}(\theta,g) = -\frac{1}{2}E[\epsilon_j(\theta)\epsilon_l(\theta)\kappa_{j,l}] = -\frac{1}{2}E[(\tilde{y}_j - \tilde{P}'_j\theta)(\tilde{y}_l - \tilde{P}'_l\theta)\kappa_{j,l}]$$

where g stands for all nuisance parameters.

The FOC defined in Equation (8) is not orthogonal

The key parameter is θ , the true value is θ_0 . The nuisance parameters are $g_P(X_i)$ and $g_y(X_i)$. Their true values are $E(P_i|X_i) = g_{0,P}(X_i)$ and $E(y_i|X_i) = g_{0,y}(X_i)$.

The first order condition is written as follows:

$$E[(P_j - g_P(X_j))(y_l - g_y(X_l) - [P_l - g_P(X_l)]'\theta)\kappa_{j,l}] = 0$$

If $P_i = [W_i, f(W_i, X_i)]'$ and there are only one variable in $X_i, \partial_\theta \varphi(D; \theta, g)$ becomes:

$$\partial_{\theta}\varphi(D;\theta,g) = \begin{bmatrix} (P_{1j} - g_{P_1}(X_j))[y_l - g_y(X_l) - (P_{1l} - g_{P_1}(X_l))\theta_1 - (P_{2l} - g_{P_2}(X_l))\theta_2]\kappa_{j,l} \\ (P_{2j} - g_{P_2}(X_j))[y_l - g_y(X_l) - (P_{1l} - g_{P_1}(X_l))\theta_1 - (P_{2l} - g_{P_2}(X_l))\theta_2]\kappa_{j,l} \end{bmatrix}$$

Prove that $\partial_g E[\partial_\theta \varphi(D; \theta_0, g_0)] \neq 0.$

Proof. The first row of $\partial_{\theta}\varphi(D;\theta_0,g_0+r(g-g_0))$ is defined as I where

$$I = [P_{1j} - g_{0,P_1}(X_j) - r(g_{P_1}(X_j) - g_{0,P_1}(X_j))]$$

$$[y_l - g_{0,y}(X_l) - r(g_y(X_l) - g_{0,y}(X_l))$$

$$- (P_{1l} - g_{0,P_1}(X_l) - r(g_{P_1}(X_l) - g_{0,P_1}(X_l)))\theta_1$$

$$- (P_{2l} - g_{0,P_2}(X_l) - r(g_{P_2}(X_l) - g_{0,P_2}(X_l)))\theta_2]\kappa_{j,l}$$

According to the definition for $\partial_g E[\partial_\theta \varphi(D; \theta_0, g_0)] \neq 0$ in Chernozhukov et al. (2018), to show that $\partial_g E[\partial_\theta \varphi(D; \theta_0, g_0)] \neq 0$ we need to show $\partial_r E[I]|_{r=0} \neq 0$. Under a regularity condition, for instance, $\int_\theta \int_r |\partial_r E[I]|_{r=0} |dr d\theta < \infty$, we have the following:

$$\partial_r E[I]|_{r=0} = -I_1 - I_2 + I_3 + I_4$$

$$\begin{split} I_1 &= E[(g_{P_1}(X_j) - g_{0,P_1}(X_j))(y_l - g_{0,y}(X_l) \\ &- (P_{1l} - g_{0,P_1}(X_l))\theta_{0,1} - (P_{2l} - g_{0,P_2}(X_l))\theta_{0,2})\kappa_{j,l}] \\ &= E\left[(g_{P_1}(X_j) - g_{0,P_1}(X_j))\epsilon_l\kappa_{j,l}\right] \\ &= \int_{\mathbb{R}^{q_z}} E[(g_{P_1}(X_j) - g_{0,P_1}(X_j))\epsilon_l e^{it'(Z_j - Z_l)}]d\mu(t) \\ &= \int_{\mathbb{R}^{q_z}} E[(g_{P_1}(X_j) - g_{0,P_1}(X_j))e^{it'Z_j}]E[\epsilon_l e^{-it'Z_l}]d\mu(t) = 0 \end{split}$$

The first equation sign is a result of notation. The second one is derived based on the outcomes following the Robinson Transformation. The third equality arises from the definition of $\kappa_{j,l}$. The fourth one is inferred from Assumption 1(e). The final equal sign is a consequence of Assumption 1(a).

$$I_{2} = E [(P_{1j} - g_{0,P_{1}}(X_{j}))(g_{y}(X_{l}) - g_{0,y}(X_{l}))\kappa_{j,l}] \neq 0$$

$$I_{3} = E [(P_{1j} - g_{0,P_{1}}(X_{j}))(g_{P_{1}}(X_{l}) - g_{0,P_{1}}(X_{l}))\theta_{1}\kappa_{j,l}] \neq 0$$

$$I_{4} = E [(P_{1j} - g_{0,P_{1}}(X_{j}))(g_{P_{2}}(X_{l}) - g_{0,P_{2}}(X_{l}))\theta_{2}\kappa_{j,l}] \neq 0$$

Hence, $\partial_g E[\partial_\theta \varphi(D; \theta_0, g_0)]$ is probably not 0.

-		

The FOC defined in Equation (9) is orthogonal

With
$$E[(P_{1m} - g_{0,P_1}(X_m))\kappa_{m,l}|X_l] = g_{0,\tilde{P}_{1,j}}(X_l), E[\kappa_{m,l}|X_l] = g_{0,\kappa_{m,l}}(X_l),$$

 $\Psi(D;\theta,g)$

$$= \begin{bmatrix} \left(P_{1j} - g_{P_1}(X_j) - \frac{g_{\tilde{P}_{1,j}}(X_l)}{g_{\kappa_{j,l}}(X_l)}\right) [y_l - g_y(X_l) - (P_{1l} - g_{P_1}(X_l))\theta_1 - (P_{2l} - g_{P_2}(X_l))\theta_2]\kappa_{j,l} \\ \left(P_{2j} - g_{P_2}(X_j) - \frac{g_{\tilde{P}_{2,j}}(X_l)}{g_{\kappa_{j,l}}(X_l)}\right) [y_l - g_y(X_l) - (P_{1l} - g_{P_1}(X_l))\theta_1 - (P_{2l} - g_{P_2}(X_l))\theta_2]\kappa_{j,l} \end{bmatrix}$$

Prove that $\partial_g E[\Psi(D; \theta_0, g_0)] = 0.$

Proof. The first row of $\Psi(D; \theta_0, g_0 + r(g - g_0))$ is I'.

$$I' = \begin{pmatrix} P_{1j} - g_{0,P_1}(X_j) - r(g_{P_1}(X_j) - g_{0,P_1}(X_j)) - \frac{g_{0,\tilde{P}_{1,j}}(X_l) + r[g_{\tilde{P}_{1,j}}(X_l) - g_{0,\tilde{P}_{1,j}}(X_l)]}{g_{0,\kappa_{m,l}}(X_l) + r[g_{\kappa_{j,l}}(X_l) - g_{0,\kappa_{m,l}}(X_l)]} \end{pmatrix}$$

$$= \begin{pmatrix} P_{1j} - g_{0,y}(X_l) - r(g_y(X_l) - g_{0,y}(X_l)) \\ - (P_{1l} - g_{0,P_1}(X_l) - r(g_{P_1}(X_l) - g_{0,P_1}(X_l))) \\ - (P_{2l} - g_{0,P_2}(X_l) - r(g_{P_2}(X_l) - g_{0,P_2}(X_l))) \\ \end{pmatrix}$$

$$\partial_r E[I']|_{r=0} = -I_1 - I'_2 + I'_3 + I'_4 - I_5$$

$$I_{1} = E[(g_{P_{1}}(X_{j}) - g_{0,P_{1}}(X_{j}))(y_{l} - g_{0,y}(X_{l}) - (P_{1l} - g_{0,P_{1}}(X_{l}))\theta_{0,1}$$

- $(P_{2l} - g_{0,P_{2}}(X_{l}))\theta_{0,2})\kappa_{j,l}]$
= $E[(g_{P_{1}}(X_{j}) - g_{0,P_{1}}(X_{j}))\epsilon_{l}\kappa_{j,l}]$
= 0

The I_1 term is the same as the I_1 in the previous proof, which demonstrates that the FOC defined in Equation (8) is not orthogonal. Therefore, it remains 0.

$$\begin{split} I'_{2} &= E[(P_{1j} - g_{0,P_{1}}(X_{j}) - \frac{E[(P_{1m} - g_{0,P_{1}}(X_{m}))\kappa_{m,l}|X_{l}]}{E[\kappa_{m,l}|X_{l}]})(g_{y}(X_{l}) - g_{0,y}(X_{l}))\kappa_{j,l}] \\ &= E(E\left[\left(P_{1j} - g_{0,P_{1}}(X_{j}) - \frac{E[(P_{1m} - g_{0,P_{1}}(X_{m}))\kappa_{m,l}|X_{l}]}{E[\kappa_{m,l}|X_{l}]}\right)\kappa_{j,l}|X_{l}\right] \\ &= E(E\left[(P_{1j} - g_{0,P_{1}}(X_{j}))\kappa_{j,l} - \frac{E[(P_{1m} - g_{0,P_{1}}(X_{m}))\kappa_{m,l}|X_{l}]}{E[\kappa_{m,l}|X_{l}]}\kappa_{j,l}|X_{l}\right] \\ &= g_{y}(X_{l}) - g_{0,y}(X_{l})) \\ &= 0 \end{split}$$

because $E\left[\left(P_{1j} - g_{0,P_1}(X_j)\right)\kappa_{j,l} - \frac{E\left[\left(P_{1m} - g_{0,P_1}(X_m)\right)\kappa_{m,l}|X_l\right]}{E\left[\kappa_{m,l}|X_l\right]}\kappa_{j,l}|X_l\right] = 0$. To see this, notice that after conditional on X_l , it becomes $E\left[\tilde{P}_{1j}\kappa_{j,l}|X_l\right] - E\left[\frac{E\left[\tilde{P}_{1m}\kappa_{m,l}|X_l\right]}{E\left[\kappa_{m,l}|X_l\right]}\kappa_{j,l}|X_l\right]$, which is 0 with $\tilde{P}_{1j} = 0$.

$$\begin{split} &(P_{1j} - g_{0,P_1}(X_j)). \text{ Following the same idea, we have the next equations for } I'_3 \text{ and } I'_4: \\ &I'_3 = E\left[(P_{1j} - g_{0,P_1}(X_j) - \frac{E[(P_{1m} - g_{0,P_1}(X_m))\kappa_{m,l}|X_l]}{E[\kappa_{m,l}|X_l]})(g_{P_1}(X_l) - g_{0,P_1}(X_l))\theta_1\kappa_{j,l}\right] \\ &= 0 \\ &I'_4 = E\left[(P_{1j} - g_{0,P_1}(X_j) - \frac{E[(P_{1m} - g_{0,P_1}(X_m))\kappa_{m,l}|X_l]}{E[\kappa_{m,l}|X_l]})(g_{P_2}(X_l) - g_{0,P_2}(X_l))\theta_2\kappa_{j,l}\right] \\ &= 0 \\ &I_5 = E\left[\frac{\left(g_{\tilde{P}_{1,j}}(X_l) - g_{0,\tilde{P}_{1,j}}(X_l)\right)g_{0,\kappa_{m,l}}(X_l) - \left(g_{\kappa_{j,l}}(X_l) - g_{0,\kappa_{m,l}}(X_l)\right)g_{0,\tilde{P}_{1,j}}(X_l)}{(g_{0,\kappa_{m,l}}(X_l))^2}\epsilon_l\kappa_{j,l}\right] \\ &= 0 \\ &= 0 \end{split}$$

For I_5 , the first equal sign is a result of notation. The second one serves a notation purpose, summarizing the fraction into a function of X_l . The third equality arises from the definition of $\kappa_{j,l}$. The fourth one is inferred from Assumption 1(e). The final equal sign is a consequence of Assumption 1(a).

Hence, $\partial_g E[\Psi(D; \theta_0, g_0)] = 0.$

11		

C.4 Proof of Proposition 2

Proof.

Denote $A_n =$

$$\begin{split} \tilde{\theta}_{n,o} &= \left[\frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \tilde{P}_{l}' \right]^{-1} \\ & \left[\frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \left(\tilde{P}_{l}\theta_{0} + \epsilon_{l} \right) \right] \\ &= \theta_{0} + \left[\frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \tilde{P}_{l}' \right]^{-1} \\ & \left[\frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \epsilon_{l} \right] \\ \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \tilde{P}_{l}' \end{split}$$

 $B_n = \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right) \epsilon_l.$ We first show that A_n is a U-statistic and find its probability limit. Then we show that B_n is also a U-statistic and find its probability limit.

To show that A_n is a U-statistic, notice that

$$A_{n} = \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \tilde{P}_{l}'$$
$$= \frac{1}{2} \frac{2}{n(n-1)} \sum_{j$$

According to WLLN for U-statistics under Assumption 1,

$$A_n \xrightarrow{p} A$$
 with $A \equiv E\left[\kappa_{j,l}\left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\right)\tilde{P}'_l\right]$

and under Assumption 2, A is nonsingular.

To show that B_n is also a U-statistic, notice that:

$$B_n = \frac{1}{n(n-1)} \sum_{j$$

Define $h(\tilde{p}_1, e_1, z_1, x_1; \tilde{p}_2, e_2, z_2, x_2) = \kappa_{1,2} \tilde{p}_1 e_2 + \kappa_{2,1} \tilde{p}_2 e_1 - \kappa_{1,2} \frac{g_{0,\tilde{P}_1}(x_2)}{g_{0,\kappa_{1,2}}(x_2)} e_2 - \kappa_{2,1} \frac{g_{0,\tilde{P}_2}(x_1)}{g_{0,\kappa_{2,1}}(x_1)} e_1$. Since

h is a symmetric function of observations 1 and 2, a U-statistic with kernel h is defined as

$$B'_{n} = \frac{2}{n(n-1)} \sum_{j < l}^{n} h(\tilde{P}_{j}, \epsilon_{j}, Z_{j}, X_{j}; \tilde{P}_{l}, \epsilon_{l}, Z_{l}, X_{l}) \quad \text{and} \quad B_{n} = \frac{1}{2} B'_{n}$$

And we have:

$$\begin{split} E(B'_{n}) &= E(\kappa_{j,l}\tilde{P}_{j}\epsilon_{l} + \kappa_{l,j}\tilde{P}_{l}\epsilon_{j}) - \left(E\left(\kappa_{j,l}\frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\epsilon_{l}\right) + E\left(\kappa_{l,j}\frac{g_{0,\tilde{P}_{m}}(X_{j})}{g_{0,\kappa_{m,j}}(X_{j})}\epsilon_{j}\right)\right) \\ &= 2E(\kappa_{j,l}\tilde{P}_{j}\epsilon_{l}) - 2E\left(\kappa_{j,l}\frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\epsilon_{l}\right) \\ &= 2\int_{\mathbb{R}^{q_{2}}} E[\tilde{P}_{j}\epsilon_{l}e^{it'(Z_{j}-Z_{l})}]d\mu(t) - 2\int_{\mathbb{R}^{q_{2}}} E[\frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\epsilon_{l}e^{it'(Z_{j}-Z_{l})}]d\mu(t) \\ &= 2\int_{\mathbb{R}^{q_{2}}} E[\tilde{P}_{j}e^{it'Z_{j}}\epsilon_{l}e^{-it'Z_{l}}]d\mu(t) - 2\int_{\mathbb{R}^{q_{2}}} E[\frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}e^{it'Z_{j}}\epsilon_{l}e^{-it'Z_{l}}]d\mu(t) \\ &= 2\int_{\mathbb{R}^{q_{2}}} E[\tilde{P}_{j}e^{it'Z_{j}}]E[\epsilon_{l}e^{-it'Z_{l}}]d\mu(t) - 2\int_{\mathbb{R}^{q_{2}}} E[e^{it'Z_{j}}]E[\frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\epsilon_{l}e^{-it'Z_{l}}]d\mu(t) \\ &= 0 \qquad \text{since } E[\epsilon'_{l}e^{-it'Z_{l}}] = 0 \text{ and } E[\epsilon_{l}|X_{l},Z_{l}] = 0. \end{split}$$

Hence, $E(B_n) = 0$. According to WLLN for U statistics, we have $B_n \xrightarrow{p} 0$, and we conclude that $\tilde{\beta}_n$ is a consistent estimator of β_0 .

To derive the asymptotic normality, we first need to compute the asymptotic variance for the U-statistic B'_n , which means that we need to find the variance for the following:

$$E(h(\tilde{P}_1, \epsilon_1, Z_1, X_1; \tilde{P}_2, \epsilon_2, Z_2, X_2) | \tilde{P}_1 = \tilde{p}_1, \epsilon_1 = e_1, Z_1 = z_1, X_1 = x_1).$$

Let $h_1(\tilde{p}_1, e_1, z_1, x_1) \equiv E(h(\tilde{P}_1, \epsilon_1, Z_1, X_1; \tilde{P}_2, \epsilon_2, Z_2, X_2) | \tilde{P}_1 = \tilde{p}_1, \epsilon_1 = e_1, Z_1 = z_1, X_1 = x_1)$. We have:

$$h(\tilde{p}_1, e_1, z_1, x_1; \tilde{p}_2, e_2, z_2, x_2) = \kappa_{1,2} \tilde{p}_1 e_2 + \kappa_{2,1} \tilde{p}_2 e_1 - \kappa_{1,2} \frac{g_{0,\tilde{P}_1}(x_2)}{g_{0,\kappa_{1,2}}(x_2)} e_2 - \kappa_{2,1} \frac{g_{0,\tilde{P}_2}(x_1)}{g_{0,\kappa_{2,1}}(x_1)} e_1$$

and

$$\begin{split} h_1(\tilde{p}_1, e_1, z_1, x_1) &= E[\int_{\mathbb{R}^{q_z}} e^{it'(z_1 - Z_2)} d\mu(t) \tilde{p}_1 \epsilon_2 + \int_{R^{q_z}} e^{it'(Z_2 - z_1)} d\mu(t) \tilde{P}_2 e_1 \\ &- \int_{\mathbb{R}^{q_z}} e^{it'(z_1 - Z_2)} d\mu(t) \frac{g_{0, \tilde{P}_1}(X_2)}{g_{0, \kappa_{1,2}}(X_2)} \epsilon_2 - \int_{R^{q_z}} e^{it'(Z_2 - z_1)} d\mu(t) \frac{g_{0, \tilde{P}_2}(x_1)}{g_{0, \kappa_{2,1}}(x_1)} e_1] \\ &= E[\int_{\mathbb{R}^{q_z}} e^{it'(z_1 - Z_2)} d\mu(t) \tilde{p}_1 \epsilon_2] + E[\int_{R^{q_z}} e^{it'(Z_2 - z_1)} d\mu(t) \tilde{P}_2 e_1] \\ &- E[\int_{\mathbb{R}^{q_z}} e^{it'(z_1 - Z_2)} d\mu(t) \frac{g_{0, \tilde{P}_1}(X_2)}{g_{0, \kappa_{1,2}}(X_2)} \epsilon_2] - E[\int_{R^{q_z}} e^{it'(Z_2 - z_1)} d\mu(t) \\ &\quad \frac{g_{0, \tilde{P}_2}(x_1)}{g_{0, \kappa_{2,1}}(x_1)} e_1] \end{split}$$

The first element of the right hand side is

$$\begin{split} E[\int_{\mathbb{R}^{q_z}} e^{it'(z_1 - Z_2)} d\mu(t) \tilde{p}_1 \epsilon_2] &= \int_{\mathbb{R}^{q_z}} E[e^{it'z_1} e^{-it'Z_2} \tilde{p}_1 \epsilon_2] d\mu(t) \\ &= \int_{\mathbb{R}^{q_z}} e^{it'z_1} \tilde{p}_1 E[e^{-it'Z_2} \epsilon_2] d\mu(t) = 0 \end{split}$$

The third term is

$$\begin{split} E[\int_{\mathbb{R}^{q_z}} e^{it'(z_1 - Z_2)} d\mu(t) \frac{g_{0,\tilde{P}_1}(X_2)}{g_{0,\kappa_{1,2}}(X_2)} \epsilon_2] &= \int_{\mathbb{R}^{q_z}} E[e^{it'z_1} e^{-it'Z_2} \frac{g_{0,\tilde{P}_1}(X_2)}{g_{0,\kappa_{1,2}}(X_2)} \epsilon_2] d\mu(t) \\ &= \int_{\mathbb{R}^{q_z}} e^{it'z_1} \tilde{p}_1 E[e^{-it'Z_2} \frac{g_{0,\tilde{P}_1}(X_2)}{g_{0,\kappa_{1,2}}(X_2)} \epsilon_2] d\mu(t) = 0 \end{split}$$

The second term of the right hand side is

$$\begin{split} E[\int_{R^{q_z}} e^{it'(Z_2 - z_1)} d\mu(t) \tilde{P}_2 e_1] &= \int_{\mathbb{R}^{q_z}} E[e^{it'Z_2} e^{-it'z_1} \tilde{P}_2 e_1] d\mu(t) \\ &= \int_{\mathbb{R}^{q_z}} e^{-it'z_1} e_1 E[e^{it'Z_2} \tilde{P}_2] d\mu(t) \end{split}$$

The fourth term is

$$\begin{split} E[\int_{R^{q_z}} e^{it'(Z_2 - z_1)} d\mu(t) \frac{g_{0,\tilde{P}_2}(x_1)}{g_{0,\kappa_{2,1}}(x_1)} e_1] &= \int_{\mathbb{R}^{q_z}} E[e^{it'Z_2} e^{-it'z_1} \frac{g_{0,\tilde{P}_2}(x_1)}{g_{0,\kappa_{2,1}}(x_1)} e_1] d\mu(t) \\ &= \int_{\mathbb{R}^{q_z}} e^{-it'z_1} e_1 \frac{g_{0,\tilde{P}_2}(x_1)}{g_{0,\kappa_{2,1}}(x_1)} E[e^{it'Z_2}] d\mu(t) \end{split}$$

Hence, $h_1(\tilde{p}_1, e_1, z_1, x_1) = \int_{\mathbb{R}^{q_z}} e^{-it'z_1} e_1 E[e^{it'Z_2} \tilde{P}_2] d\mu(t) - \int_{\mathbb{R}^{q_z}} e^{-it'z_1} e_1 \frac{g_{0,\tilde{P}_2}(x_1)}{g_{0,\kappa_{2,1}}(x_1)} E[e^{it'Z_2}] d\mu(t).$ $h_1(\tilde{p}_1, e_1, z_1, x_1) = \int_{\mathbb{R}^{q_z}} e^{-it'z_1} e_1 \left(E[e^{it'Z_2} \tilde{P}_2] - \frac{g_{0,\tilde{P}_2}(x_1)}{g_{0,\kappa_{2,1}}(x_1)} E[e^{it'Z_2}] \right) d\mu(t)$ Since $E(h_1(\tilde{P}_1, \epsilon_1, Z_1, X_1)) = 0$, we have:

$$\begin{aligned} Var[h_{1}(\tilde{P}_{1},\epsilon_{1},Z_{1},X_{1})] &= Var\left[\int_{\mathbb{R}^{q_{z}}} e^{-it'Z_{1}}\epsilon_{1}\left(E[e^{it'Z_{2}}\tilde{P}_{2}] - \frac{g_{0,\tilde{P}_{2}}(X_{1})}{g_{0,\kappa_{2,1}}(X_{1})}E[e^{it'Z_{2}}]\right)d\mu(t)\right] \\ &= Var\left[\int_{\mathbb{R}^{q_{z}}} e^{-it'Z_{j}}\epsilon_{j}\left(E[e^{it'Z_{l}}\tilde{P}_{l}] - \frac{g_{0,\tilde{P}_{m}}(X_{j})}{g_{0,\kappa_{m,j}}(X_{j})}E[e^{it'Z_{l}}]\right)d\mu(t)\right]\end{aligned}$$

Following Hoeffding (1948), the asymptotic distribution for U-statistics yields:

$$\sqrt{n}(B'_n-0) \xrightarrow{d} \mathcal{N}(0, 4Var[h_1(\tilde{P}_1, \epsilon_1, Z_1, X_1)])$$

Thus,

$$\sqrt{n}(\tilde{\theta}_{n,o} - \theta_0) \xrightarrow{d} N\left(0, A^{-1}\Sigma\left(A^{-1}\right)'\right)$$

with $A = E\left[\kappa_{j,l}\left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\right)\tilde{P}_l'\right]$ and $\Sigma = Var[h_1(\tilde{P}_1, \epsilon_1, Z_1, X_1)].$

C.5 Proof of Theorem 3.1

Proof. Recall that

$$\hat{\tilde{y}}_i = y_i - \hat{g}_y(X_i) = y_i - E(y_i|X_i) + E(y_j|X_i) - \hat{g}_y(X_i)$$

For the first two terms of the right hand side, we have

$$y_{i} - E(y_{i}|X_{i}) = (P_{i} - E(P_{i}|X_{i}))'\theta_{0} + \epsilon_{i}$$

= $(P_{i} - \hat{g}_{P}(X_{i}))'\theta_{0} + (\hat{g}_{P}(X_{i}) - E(P_{i}|X_{i}))'\theta_{0} + \epsilon_{i}$

In matrix form, the feasible estimator writes:

$$\hat{\theta}_{n,o} = \left[\frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}}(X_{l})}{\widehat{g_{0,\kappa_{m,l}}}(X_{l})}\right) \hat{\tilde{P}}_{l}'\right]^{-1} \\ \left[\frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}}(X_{l})}{\widehat{g_{0,\kappa_{m,l}}}(X_{l})}\right) \hat{\tilde{y}}_{l}\right]$$

Define $C_n = \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\widehat{\tilde{P}}_j - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \widehat{\tilde{P}}'_l$. We get:

$$\begin{split} \hat{\theta}_{n,o} &= [C_n]^{-1} [\frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\hat{\tilde{P}}_j - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \\ &= [y_l - E(y_l | X_l) + E(y_l | X_l) - \hat{g}_y(X_l)]] \\ &= [C_n]^{-1} [\frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\hat{\tilde{P}}_j - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \\ &= [(P_l - \hat{g}_P(X_l))' \theta_0 + (\hat{g}_P(X_l) - E(P_l | X_l))' \theta_0 + \epsilon_l + E(y_l | X_l) - \hat{g}_y(X_l)]] \\ &= \theta_0 + [C_n]^{-1} [\frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\hat{\tilde{P}}_j - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \\ &= [(\hat{g}_P(X_l) - E(P_l | X_l))' \theta_0 + \epsilon_l + E(y_l | X_l) - \hat{g}_y(X_l)]] \end{split}$$

Consider now,

$$\begin{split} A_n - C_n &= \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l \neq j}^n \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right) \tilde{P}'_l \\ &- \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l \neq j}^n \kappa_{j,l} \left(\tilde{P}_j - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \tilde{P}'_l \\ &= \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l \neq j}^n \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right) \tilde{P}'_l \\ &- \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l \neq j}^n \kappa_{j,l} \\ \left(\tilde{P}_j + \tilde{P}_j - \tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \tilde{P}'_l \\ &= \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l \neq j}^n \kappa_{j,l} \left[\left(\tilde{\tilde{P}}_j - \tilde{P}_j + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \tilde{P}'_l \\ &+ \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right) (-\tilde{\tilde{P}}_l + \tilde{P}_l) \right] \\ & \xrightarrow{P} = 0 \end{split}$$

Hence, we have $\operatorname{Plim}C_n = A$, since we showed in the proof of Proposition 2 that $\operatorname{Plim}A_n = A$. Define now the following quantities:

$$D_{n} = \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}(X_{l})}}{\widehat{g_{0,\kappa_{m,l}}(X_{l})}} \right)$$

$$[(\hat{g}_{P}(X_{l}) - E(P_{l}|X_{l}))'\theta_{0} + \epsilon_{l} + E(y_{l}|X_{l}) - \hat{g}_{y}(X_{l})]$$

$$E_{n} = \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}(X_{l})}}{\widehat{g_{0,\kappa_{m,l}}(X_{l})}} \right) (\hat{g}_{P}(X_{l}) - E(P_{l}|X_{l}))'\theta_{0}$$

$$F_{n} = \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}(X_{l})}}{\widehat{g_{0,\kappa_{m,l}}(X_{l})}} \right) \epsilon_{l}$$

$$G_{n} = \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \frac{\widehat{g_{0,\tilde{P}_{m}}(X_{l})}{\widehat{g_{0,\kappa_{m,l}}(X_{l})}} \right) [E(y_{l}|X_{l}) - \hat{g}_{y}(X_{l})]$$

We have, $D_n = E_n + F_n + G_n$. For consistency, we show that the probability limits for E_n , F_n , and G_n are all zero.

$$\begin{split} E_n &= \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\hat{\tilde{P}}_j - \frac{\widehat{g_{0,\tilde{P}_m}(X_l)}}{\widehat{g_{0,\kappa_{m,l}}(X_l)}} \right) (\hat{g}_P(X_l) - E(P_l|X_l))' \theta_0 \\ &= \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\tilde{P}_j + \hat{\tilde{P}}_j - \tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}(X_l)}}{\widehat{g_{0,\kappa_{m,l}}(X_l)}} \right) \\ &\quad (\hat{g}_P(X_l) - E(P_l|X_l))' \theta_0 \\ &= \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} + \hat{\tilde{P}}_j - \tilde{P}_j + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}(X_l)}}{\widehat{g_{0,\kappa_{m,l}}(X_l)}} \right) \\ &\quad (\hat{g}_P(X_l) - E(P_l|X_l))' \theta_0 \\ &= \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right) (\hat{g}_P(X_l) - E(P_l|X_l))' \theta_0 \\ &+ \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\hat{\tilde{P}}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}(X_l)}} \right) \\ &\quad (\hat{g}_P(X_l) - E(P_l|X_l))' \theta_0 \\ &+ \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\hat{\tilde{P}}_j - \tilde{P}_j + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \\ &\quad (\hat{g}_P(X_l) - E(P_l|X_l))' \theta_0 \end{array}$$

$$F_{n} = \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} + \tilde{\tilde{P}}_{j} - \tilde{P}_{j} + \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} - \frac{\widehat{g_{0,\tilde{P}_{m}}}(X_{l})}{\widehat{g_{0,\kappa_{m,l}}}(X_{l})} \right) \epsilon_{l}$$

$$= \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \epsilon_{l}$$

$$+ \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\hat{\tilde{P}}_{j} - \tilde{P}_{j} + \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} - \frac{\widehat{g_{0,\tilde{P}_{m}}}(X_{l})}{\widehat{g_{0,\kappa_{m,l}}}(X_{l})} \right) \epsilon_{l}$$

$$\stackrel{P}{\to} 0$$

$$\begin{aligned} G_n &= \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l \neq j}^n \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right) [E(y_l | X_l) - \hat{g}_y(X_l)] \\ &+ \frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l \neq j}^n \kappa_{j,l} \left(\hat{\tilde{P}}_j - \tilde{P}_j + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) [E(y_l | X_l) - \hat{g}_y(X_l)] \\ &\stackrel{P}{\to} 0 \end{aligned}$$

All in all, we have $\text{Plim}D_n = \text{Plim}(E_n + F_n + G_n) = 0$, so $\hat{\theta}_{n,o} \xrightarrow{P} \theta_0$. In addition, we have:

$$\sqrt{n}(\hat{\theta}_{n,o} - \theta_0) = [C_n]^{-1}\sqrt{n}[E_n + F_n + G_n]$$

And we study each term separately:

$$\sqrt{n}F_n = \sqrt{n}B_n + \sqrt{n}\frac{1}{n(n-1)}\sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l}$$
$$\left(\hat{\tilde{P}}_j - \tilde{P}_j + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)}\right)\epsilon_l$$

The second term of $\sqrt{n}F_n$ is $o_p(1)$ because $\left(\hat{\tilde{P}}_j - \tilde{P}_j + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)}\right)$ is $o_p(1)$ under Assumption 3, and $\sqrt{n}\frac{1}{n(n-1)}\sum_{j=1}^n\sum_{l\neq j}^n\kappa_{j,l}\epsilon_l$ is $O_p(1)$.

$$\begin{split} \sqrt{n}E_n &= \sqrt{n}\frac{1}{n(n-1)}\sum_{j=1}^n\sum_{l\neq j}^n\kappa_{j,l}\left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\right)(\hat{g}_P(X_l) - E(P_l|X_l))'\theta_0 \\ &+ \sqrt{n}\frac{1}{n(n-1)}\sum_{j=1}^n\sum_{l\neq j}^n\kappa_{j,l}\left(\tilde{\tilde{P}}_j - \tilde{P}_j + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)}\right) \\ &\quad (\hat{g}_P(X_l) - E(P_l|X_l))'\theta_0 \end{split}$$

$$\sqrt{n}G_n = \sqrt{n}\frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right) [E(y_l|X_l) - \hat{g}_y(X_l)] \\
+ \sqrt{n}\frac{1}{n(n-1)} \sum_{j=1}^n \sum_{l\neq j}^n \kappa_{j,l} \left(\hat{\tilde{P}}_j - \tilde{P}_j + \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\tilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)} \right) \\
[E(y_l|X_l) - \hat{g}_y(X_l)]$$

The second terms of $\sqrt{n}E_n$ and $\sqrt{n}G_n$ are $o_p(1)$ under Assumption 3, because

$$\left(\widehat{\tilde{P}}_j - \widetilde{P}_j + \frac{g_{0,\widetilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} - \frac{\widehat{g_{0,\widetilde{P}_m}}(X_l)}{\widehat{g_{0,\kappa_{m,l}}}(X_l)}\right)$$

and

$$(\hat{g}_P(X_l) - E(P_l|X_l))$$
 or $[E(y_l|X_l) - \hat{g}_y(X_l)]$

are both $o_p(n^{-1/4})$. $o_p(n^{-1/4})o_p(n^{-1/4})$ is $o_p(n^{-1/2})$. Thus, the second terms are $o_p(1)$.

Therefore, the task is to demonstrate that both of these first terms are $o_p(1)$, meaning their magnitudes do not grow too fast relative to the sample size n. Once we establish this, we can conclude that the first terms of $\sqrt{n}E_n$ and $\sqrt{n}G_n$ are $o_p(1)$, completing the proof.

In the first term of $\sqrt{nE_n}$, we have $(\hat{g}_P(X_l) - E(P_l|X_l))'\theta_0$, which is $o_p(n^{-1/4})$. Recall that $E[\tilde{P}_j\kappa_{j,l} - \frac{E[\tilde{P}_m\kappa_{m,l}|X_l]}{E[\kappa_{m,l}|X_l]}\kappa_{j,l}|X_l] = 0$. That is, $E[\tilde{P}_j\kappa_{j,l} - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\kappa_{j,l}|X_l] = 0$. This is discussed in the proof that shows the FOC defined in Equation (9) is orthogonal, specifically the proof for I'_2 being 0.

The first term of $\sqrt{n}E_n$ is:

$$\begin{split} \sqrt{n} \frac{1}{n(n-1)} \sum_{j=1}^{n} \sum_{l\neq j}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) (\hat{g}_{P}(X_{l}) - E(P_{l}|X_{l}))' \theta_{0} \\ &= \frac{1}{n} \sum_{l=1}^{n} (\hat{g}_{P}(X_{l}) - E(P_{l}|X_{l}))' \theta_{0} \sqrt{n} \frac{1}{n-1} \sum_{j\neq l}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \\ &= \frac{1}{n} \sum_{l=1}^{n} o_{p}(n^{-1/4}) \frac{\sqrt{n}}{\sqrt{n-1}} \frac{\sqrt{n-1}}{n-1} \sum_{j\neq l}^{n} \kappa_{j,l} \left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})} \right) \end{split}$$

The first equality arises because we sum across j and l except for cases when j = l and rearranging the order of summation does not affect the outcome. The second equality holds under Assumption 3. Additionally, $\frac{\sqrt{n-1}}{n-1} \sum_{j\neq l}^{n} \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right)$ is $O_p(1)$ because $\frac{1}{n-1} \sum_{j\neq l}^{n} \kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right)$ represents the sample average of $\kappa_{j,l} \left(\tilde{P}_j - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)} \right)$ when X_l is fixed. The conditional expectation $E[\tilde{P}_j\kappa_{j,l} - \frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\kappa_{j,l}|X_l]$ is 0, and the corresponding variance is bounded.

Hence, the first term of $\sqrt{n}E_n$ becomes $\frac{1}{n}\sum_{l=1}^n o_p(n^{-1/4})O_P(1)$. That is, $o_p(1)$. Based on a similar analysis, the first term of $\sqrt{n}G_n$ is also $o_p(1)$.

From Assumption 3, $\sqrt{n}[E_n + F_n + G_n] = \sqrt{n}B_n + o_p(1).$

Hence,

$$\sqrt{n}(\hat{\theta}_{n,o} - \theta_0) = [C_n]^{-1}\sqrt{n}B_n + o_p(1)$$

 $\hat{\theta}_{n,o}$ and $\tilde{\theta}_{n,o}$ share the same asymptotic distribution.

C.6 Consistent Estimator of the Asymptotic Variance (11)

$$\begin{aligned} & \operatorname{Var}\left[\int_{\mathbb{R}^{q_{z}}} e^{-it'Z_{j}} \epsilon_{j} \left(E[e^{it'Z_{l}}\tilde{P}_{l}] - \frac{g_{0,\tilde{P}_{m}}(X_{j})}{g_{0,\kappa_{m,j}}(X_{j})}E[e^{it'Z_{l}}]\right) d\mu(t)\right] \\ &= \operatorname{Var}\left[E_{l}\left[\int_{\mathbb{R}^{q_{z}}} e^{it'(Z_{l}-Z_{j})} \epsilon_{j} \left(\tilde{P}_{l} - \frac{g_{0,\tilde{P}_{m}}(X_{j})}{g_{0,\kappa_{m,j}}(X_{j})}\right) d\mu(t)\right]\right] \\ &= \operatorname{Var}\left[E_{l}\left[k(Z_{l} - Z_{j}) \left(\tilde{P}_{l} - \frac{g_{0,\tilde{P}_{m}}(X_{j})}{g_{0,\kappa_{m,j}}(X_{j})}\right) \epsilon_{j}\right]\right] \\ &= E[\left(E_{l}\left[k(Z_{l} - Z_{j}) \left(\tilde{P}_{l} - \frac{g_{0,\tilde{P}_{m}}(X_{j})}{g_{0,\kappa_{m,j}}(X_{j})}\right)\right]\right) \\ &\left(E_{l}\left[k(Z_{l} - Z_{j}) \left(\tilde{P}_{l} - \frac{g_{0,\tilde{P}_{m}}(X_{j})}{g_{0,\kappa_{m,j}}(X_{j})}\right)'\right]\right) [\epsilon_{j}]^{2}]\end{aligned}$$

Under heteroskedasticity, the estimator for variance of the feasible D-RSMD is

$$[n(n-1)C_n]^{-1} \sum_{j=1}^n ((\sum_{l=1}^n \kappa_{j,l} \left(\widehat{\tilde{P}}_l - \frac{\widehat{g_{0,\tilde{P}_l}}(X_j)}{\widehat{g_{0,\kappa_{m,j}}}(X_j)} \right)) (\sum_{l=1}^n \kappa_{j,l} \left(\widehat{\tilde{P}}_l - \frac{\widehat{g_{0,\tilde{P}_l}}(X_j)}{\widehat{g_{0,\kappa_{m,j}}}(X_j)} \right)') \widehat{\epsilon}_j^2) \\ [n(n-1)C'_n]^{-1}$$

D Appendix: Identification

D.1 Summary

In this section, we provide a summary to illustrate the identification issue for D-RSMD. Due to the complexity of the estimator introduced by the debiasing process, we cannot assume higher-order assumptions for the identification, as in Antoine and Sun (2021). Hence, in this section, we will discuss the issue step by step. If there are constants in the model, the conditions for identification will be slightly different, but the logic will remain the same.

We will start with the simplest model first. Then move on to the complicated models. Models are provided with coherent simulation results. Those results are included in the Table 14.

1. Model (1):

$$y_i = \beta_1 W_{i,1} + \beta_2 W_{i,2} + u_i$$

 $W_{i,1} = \pi_1 Z_i + v_{i,1}$
 $W_{i,2} = \pi_2 Z_i + v_{i,2}$

SMD cannot identify β_1 and β_2 with continuous or discrete Z_i . To identify β_1 and β_2 with SMD, the model need to have a nonlinear part of Z_i (continuous) or another control variable X_i inside one of the equations for W_i .

2.

$$y_i = \beta_1 W_{i,1} + \beta_2 W_{i,2} + u_i$$
$$W_{i,1} = \pi_{1,1} Z_{i,1} + \pi_{1,2} Z_{i,2} + v_{i,1}$$
$$W_{i,2} = \pi_{2,1} Z_{i,1} + \pi_{2,2} Z_{i,2} + v_{i,2}$$

When (continuous or discrete) $Z_{i,1}$ and $Z_{i,2}$ are independent, we are unable to identify both parameters using SMD with only one instrument $Z_{i,1}$ under some specific conditions for the mean of $E(Z_{i,2})$, $\pi_{1,1}$, $\pi_{1,2}$, $\pi_{2,1}$, and $\pi_{2,2}$. We are able to identify both parameters β_1 and β_2 when these conditions are not met. 3. Model (3):

$$y_{i} = \beta_{1}W_{i,1} + \beta_{2}W_{i,1} * X_{i} + u_{i}$$
$$W_{i,1} = \pi_{1}Z_{i} + v_{i,1}$$
$$W_{i,1} * X_{i} = \pi_{1}Z_{i} * X_{i} + v_{i,1} * X_{i}$$

If Z_i and X_i are dependent, when Z_i is continuous, both parameters are identified, and when Z_i is binary, using SMD we can identify 0 or 1 parameter based on the value of $E(X_i|Z_i)$. If Z_i and X_i are independent, using SMD we can identify 0 or 1 parameter based on the value of $E(X_i)$. This is the same case for a continuous or discrete Z_i .

4. Model (4):

$$y_{i} = \beta_{1}W_{i,1} + \beta_{2}W_{i,1} * X_{i} + \gamma_{1}X_{i} + u_{i}$$
$$W_{i,1} = \pi_{1}Z_{i} + v_{i,1}$$
$$W_{i,1} * Z_{i} = \pi_{1}Z_{i} * X_{i} + v_{i,1} * X_{i}$$

If Z_i and X_i are dependent, RSMD identifies both parameters using a continuous or discrete Z_i . If Z_i and X_i are independent, we can use RSMD to identify 0 or 1 parameter based on the value of $E(X_i)$ with continuous or discrete Z_i .

5. D-RSMD has extra terms in its formula and it is not symmetric. It will identify both parameters when RSMD cannot.

D.2 Proofs and Simulation Examples

Following that, we move on to sections with more detailed explanations, proofs, and simulation studies to provide more information on the identification issue for methods incorporating SMD type estimators, such as SMD, RSMD, and D-RSMD.

1. Prove the identification of the key parameters using the SMD method for the simplest model in the following (Model 1).

$$y_i = \beta_1 W_{i,1} + \beta_2 W_{i,2} + u_i$$
$$W_{i,1} = \pi_1 Z_i + v_{i,1}$$
$$W_{i,2} = \pi_2 Z_i + v_{i,2}$$

Proof. The SMD estimator is

$$\begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = [E(\kappa_{j,l}W_jW_l')]^{-1}E(\kappa_{j,l}W_jy_l)$$

$$= \left[E\left(\kappa_{j,l}\begin{pmatrix} W_{j,1} \\ W_{j,2} \end{pmatrix}\begin{pmatrix} W_{l,1} \\ W_{l,2} \end{pmatrix}'\right)\right]^{-1}E(\kappa_{j,l}W_jy_l)$$

We need to prove that $E(\kappa_{j,l}W_jW'_l)$ is nonsingular. To show that $E(\kappa_{j,l}W_jW'_l)$ is nonsingular, we consider the associated quadratic form, and show that it is positive definite. For any *a* real vector of size *p*, we have:

$$\begin{split} E(a'W_{j}W_{l}'a\kappa_{j,l}) &= E(\kappa_{j,l}a'E(W_{j}|Z_{j})E(W_{l}'a|Z_{l})) \\ &= E(\int_{R^{q_{w}}} e^{it'(Z_{j}-Z_{l})}d\mu(t)a'E(W_{j}|Z_{j})E(W_{l}'|Z_{l})a) \\ &= \int_{R^{q_{w}}} E[e^{it'(Z_{j}-Z_{l})}a'E(W_{j}|Z_{j})E(W_{l}'|Z_{l})a]d\mu(t) \\ &= \int_{R^{q_{w}}} E[e^{it'Z_{j}}a'E(W_{j}|Z_{j})E(W_{l}'|Z_{l})ae^{-it'Z_{l}}]d\mu(t) \\ &= \int_{R^{q_{w}}} E[a'e^{it'Z_{j}}E(W_{j}|Z_{j})]E[E(W_{l}'|Z_{l})ae^{-it'Z_{l}}]d\mu(t) \\ &= \int_{R^{q_{w}}} E[a'e^{it'Z_{j}}E(W_{j}|Z_{j})]E[E(W_{j}'|Z_{j})ae^{-it'Z_{j}}]d\mu(t) \\ &= \int_{R^{q_{w}}} \left| \left(\int_{R^{q_{w}}} a'e^{it'Z_{j}}E(W_{j}|Z_{j})f_{Z}(Z_{j})dZ_{j} \right) \right|^{2}d\mu(t) \\ &= (2\pi)^{2q_{w}} \int_{R^{q_{w}}} \left| \left(\mathcal{F}[a'E(W_{j}|Z_{j})f_{Z}(Z_{j})](t) \right) \right|^{2}d\mu(t) \\ &\geq 0 \end{split}$$

with μ strictly positive on \mathbb{R}^p and $\mathcal{F}[g]$ the Fourier transform of a well-defined function g(.)on \mathbb{R}^{q_w} is formally defined as,

$$\mathcal{F}[g](t) = \frac{1}{(2\pi)^{q_w}} \int \exp^{it'u} g(u) du \,. \tag{13}$$

We then have:

$$\begin{aligned} a'E(\kappa_{j,l}W_jW_l')a &= 0 &\Leftrightarrow & \exists \ a \neq 0 \ s.t. \ a'E(W_j|Z_j)f(W_j) = 0 \ a.s. \\ &\Leftrightarrow & \exists \ a \neq 0 \ s.t. \ a'E(W_j|Z_j) = 0 \ a.s. \end{aligned}$$

In the specific case,

$$\Leftrightarrow \exists a \neq 0 \ s.t. \ a' E(W_j | Z_j) = a_1 \pi_1 Z_i + a_2 \pi_2 Z_i = (a_1 \pi_1 + a_2 \pi_2) Z_i = 0 \ a.s.$$

 β_1 and β_2 are not identified. To identify β_1 and β_2 , the model must contain a nonlinear part of Z_i or another control variable X_i inside one of the equations for W_i .

2. Prove the identification of the key parameters using the SMD method for the model with an interaction term (Model (3)).

$$y_{i} = \beta_{1}W_{i,1} + \beta_{2}W_{i,1} * X_{i} + u_{i}$$
$$W_{i,1} = \pi_{1}Z_{i} + v_{i,1}$$
$$W_{i,1} * X_{i} = \pi_{1}Z_{i} * X_{i} + v_{i,1} * X_{i}$$

Proof. We need to prove that $E(\kappa_{j,l}W_jW'_l)$ is nonsingular. Follow the same logic as in the previous proof.

$$a' E(\kappa_{j,l} W_j W_l) a = 0 \Leftrightarrow \exists a \neq 0 \ s.t. \ a' E(W_j | Z_j) = 0 \ a.s.$$

 $E(W_{i,1} | Z_i) = \pi_1 Z_i$

$$E(W_{i,1} * X_i | Z_i) = E((\pi_1 Z_i + v_{i,1}) * X_i | Z_i)$$

= $E(\pi_1 Z_i * X_i + v_{i,1} * X_i | Z_i)$
= $\pi_1 Z_i E(X_i | Z_i) + E(v_{i,1} * X_i | Z_i)$

$$a'E(W_i|Z_i) = a_1\pi_1Z_i + a_2\pi_1Z_iE(X_i|Z_i) + a_2E(v_{i,1} * X_i|Z_i)$$

If $v_{i,1} \perp (X_i, Z_i)$,

$$E(v_{i,1} * X_i | Z_i) = 0$$
$$a' E(W_i | Z_i) = a_1 \pi_1 Z_i + a_2 \pi_1 Z_i E(X_i | Z_i)$$

When Z_i and X_i are dependent, and Z_i is continuous, there are no $a_1 \neq 0$ and $a_2 \neq 0$, s.t. $a'E(W_j|Z_j) = 0$ a.s.. Both parameters are identified.

When Z_i is binary, there will be a problem.

For instance,

$$a'E(W_i|Z_i = 0) = a_1\pi_1 * 0 + a_2\pi_1 * 0 * E(X_i|Z_i = 0) = 0$$
$$a'E(W_i|Z_i = 1) = a_1\pi_1 * 1 + a_2\pi_1 * 1 * E(X_i|Z_i = 1) = 0$$

When $E(X_i|Z_i=1) \neq 0$ we can find a $a_1 \neq 0$ and $a_2 \neq 0$, s.t. $a'E(W_j|Z_j) = 0$ a.s.

When W_i is 3-value instrument, both parameters will be identified.

Here are 2 examples for simulation illustration when using the SMD estimator. We consider the case that W_i and Z_i are independent for the second example.

(a)

$$y_i = 2W_{i,1} + 3W_{i,2} + u_i$$
$$W_{i,1} = 4Z_{i,1} + Z_{i,2} + v_{i,1}$$

$$W_{i,2} = Z_{i,1} + 3Z_{i,2} + v_{i,2}$$

When $Z_{i,1}$ and $Z_{i,2}$ are independent, we are not able to identify both parameters inside the model for y_i using SMD with only one instrument $Z_{i,1}$ or $Z_{i,2}$ when $E(Z_{i,2})$ or $E(Z_{i,1})$ are zero. Indeed, identification of SMD with $Z_{i,1}$ yields the following equation.

$$a'E(W_i|Z_{i,1}) = a_1E(4Z_{i,1} + Z_{i,2} + v_{i,1}|Z_{i,1}) + a_2E(Z_{i,1} + 3Z_{i,2} + v_{i,2}|Z_{i,1})$$
$$= a_14Z_{i,1} + a_1E(Z_{i,2}|Z_{i,1}) + a_2Z_{i,1} + a_2E(3Z_{i,2}|Z_{i,1})$$

When $Z_{i,1}$ and $Z_{i,2}$ are independent,

$$a'E(W_i|Z_{i,1}) = a_1 4Z_{i,1} + a_1 E(Z_{i,2}) + a_2 Z_{i,1} + a_2 E(3Z_{i,2})$$
$$a'E(W_i|Z_{i,1}) = (4a_1 + a_2)Z_{i,1} + (a_1 + 3a_2)E(Z_{i,2})$$

i. When $Z_{i,1}$ and $Z_{i,2}$ are continuous, we need $E(Z_{i,2}) \neq 0$ to identify both parameters.

ii. When $E(Z_{i,2}) = 0$, there are infinite possible combinations for a_1 and a_2 to have

$$a'E(W_i|Z_{i,1}) = 0$$

. In this case, neither parameter is identified.

iii. When $Z_{i,1}$ and $Z_{i,2}$ are binary 0, 1, $E(Z_{i,2}) \neq 0$. $a_1 = 0$ and $a_2 = 0$ to have

$$a'E(W_i|Z_{i,1}) = 0$$

. Thus, we will identify both parameters.

(b)

$$y_i = 2W_{i,1} + 3W_{i,1} * X_i + u_i$$

 $W_{i,1} = 2Z_i + v_{i,1}$

$$W_{i,1} * X_i = 2Z_i * X_i + v_{i,1} * X_i$$

When Z_i and X_i are independent and $E(X_i) = 0$, we are able to identify the first parameter. When Z_i and X_i are independent and $E(X_i) \neq 0$, there is no identification for both parameters. Indeed,

$$a'E(W_i|Z_i) = a_1E(2Z_i + v_{i,1}|Z_i) + a_2E(2Z_i * X_i + v_{i,1} * X_i|Z_i)$$
$$= 2a_1Z_i + 2a_2Z_i * E(X_i|Z_i)$$

When Z_i and X_i are independent,

$$a'E(W_i|Z_i) = 2a_1Z_i + 2a_2Z_i * E(X_i)$$

- i. When $E(X_i) = 0$, a_1 must be 0 for a continuous or discrete Z_i . a_2 can be any value. Thus, we can only identify the first parameter.
- ii. When $E(X_i) = 1$, there are infinite combinations for (a_1, a_2) to make

$$a'E(W_i|Z_i) = 0$$

. Both parameters are not identified.

All of these examples are verified used simulation studies. The simulation results are included in the Table 14.

3. Prove the identification of the key parameters using the RSMD method for the following model (Model (4)).

$$y_{i} = \beta_{1}W_{i,1} + \beta_{2}W_{i,1} * X_{i} + X_{i} + u_{i}$$
$$W_{i,1} = \pi_{1}Z_{i} + v_{i,1}$$
$$W_{i,1} * X_{i} = \pi_{1}Z_{i} * X_{i} + v_{i,1} * X_{i}$$

Proof. We need

$$a'E(\kappa_{j,l}\tilde{W}_j\tilde{W}_l)a = 0 \Leftrightarrow \exists a \neq 0 \ s.t. \ a'E(\tilde{W}_j|Z_j) = 0 \ a.s.$$

to show that the parameters are not identified.

$$E(\tilde{W}_{i,1}|Z_i) = E(W_{i,1} - E(W_{i,1}|X_i)|Z_i)$$

= $E(\pi_1 Z_i + v_{i,1} - E(\pi_1 Z_i + v_{i,1}|X_i)|Z_i)$
= $E(\pi_1 Z_i - E(\pi_1 Z_i|X_i)|Z_i)$
= $\pi_1 Z_i - \pi_1 E(E(Z_i|X_i)|Z_i)$

$$E(W_{i,1}X_i - E(W_{i,1}X_i|X_i)|Z_i) = E(W_{i,1}X_i - E(W_{i,1}|X_i)X_i|Z_i)$$

$$= E(\pi_1 Z_i X_i + v_{i,1}X_i - E(W_{i,1}|X_i)X_i|Z_i)$$

$$= E(\pi_1 Z_i X_i - E(\pi_1 Z_i + v_{i,1}|X_i)X_i|Z_i)$$

$$= E(\pi_1 Z_i X_i - E(\pi_1 Z_i|X_i) * X_i|Z_i)$$

$$= \pi_1 Z_i E(X_i|Z_i) - \pi_1 E(E(Z_i|X_i)X_i|Z_i)$$

$$a'E(\tilde{W}_{i}|Z_{i}) = a_{1}(\pi_{1}Z_{i} - \pi_{1}E(E(Z_{i}|X_{i})|Z_{i})) + a_{2}(\pi_{1}Z_{i}E(X_{i}|Z_{i}) - \pi_{1}E(E(Z_{i}|X_{i})X_{i}|Z_{i}))$$

Thus, when $a'E(\tilde{W}_i|Z_i) = 0$,

$$a_1 Z_i - a_1 E(E(Z_i|X_i)|Z_i) + a_2 Z_i E(X_i|Z_i) - a_2 E(E(Z_i|X_i)X_i|Z_i) = 0$$
(14)

(a) When the instrument Z_i is binary, if $Z_i = 0$, Equation 14 becomes:

$$-a_1 E(E(Z_i|X_i)|Z_i = 0) - a_2 E(E(Z_i|X_i)X_i|Z_i = 0) = 0$$

If $Z_i = 1$, Equation 14 becomes:

$$a_1 - a_1 E(E(Z_i|X_i)|Z_i = 1) + a_2 E(X_i|Z_i = 1) - a_2 E(E(Z_i|X_i)X_i|Z_i = 1) = 0$$

We need to solve for a_1 and a_2 . If both of them are 0, both parameters will be identified. If one of them is 0, only one parameter will be identified.

$$\begin{pmatrix} E(E(Z_i|X_i)|Z_i=0) & E(E(Z_i|X_i)X_i|Z_i=0) \\ 1 - E(E(Z_i|X_i)|Z_i=1) & (E(X_i|Z_i=1) - E(E(Z_i|X_i)X_i|Z_i=1)) \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

We will have a problem if $Z_i \perp X_i$. The determinant of the above matrix is 0. In that scenario, there are many possible solutions for a_1 and a_2 . We cannot identify both parameters.

Another straightforward way to illustrate that is in the following.

$$a'E(\tilde{W}_{i}|Z_{i}) = a_{1}E(\tilde{W}_{i,1}|Z_{i}) + a_{2}E(\tilde{W}_{i,1} * X_{i}|Z_{i})$$

$$a'E(\tilde{W}_{i}|Z_{i} = 0) = a_{1}E(\tilde{W}_{i,1}|Z_{i} = 0) + a_{2}E(\tilde{W}_{i,1} * X_{i}|Z_{i} = 0) = 0$$

$$a'E(\tilde{W}_{i}|Z_{i} = 1) = a_{1}E(\tilde{W}_{i,1}|Z_{i} = 1) + a_{2}E(\tilde{W}_{i,1} * X_{i}|Z_{i} = 1) = 0$$

$$\begin{pmatrix} E(\tilde{W}_{i,1}|Z_{i} = 0) & E(\tilde{W}_{i,1} * X_{i}|Z_{i} = 0) \\ E(\tilde{W}_{i,1}|Z_{i} = 1) & E(\tilde{W}_{i,1} * X_{i}|Z_{i} = 1) \end{pmatrix} \begin{pmatrix} a_{1} \\ a_{2} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$A\begin{pmatrix} a_{1} \\ a_{2} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

If matrix A is invertible or det(A) is not 0, then a_1 and a_2 are zeros. Both parameters are identified. If det(A) is 0, we cannot identify both parameters. $det(A) = E(\tilde{W}_{i,1}|Z_i = 0)E(\tilde{W}_{i,1} * X_i|Z_i = 1) - E(\tilde{W}_{i,1} * X_i|Z_i = 0)E(\tilde{W}_{i,1}|Z_i = 1)$. Additionally, with a binary instrument, the RSMD identifies 2 or less than 2 parameters. For instance, if we want to identify 3 parameters with 2 parameters for 2 interaction terms, RSMD can only identify 2 parameters at most.

$$a'E(\tilde{W}_i|Z_i) = a_1E(\tilde{W}_i|Z_i) + a_2E(\tilde{W}_iX_{i,1}|Z_i) + a_3E(\tilde{W}_iX_{i,2}|Z_i)$$
$$a'E(\tilde{W}_i|Z_i=0) = a_1E(\tilde{W}_i|Z_i=0) + a_2E(\tilde{W}_iX_{i,1}|Z_i=0) + a_3E(\tilde{W}_iX_{i,2}|Z_i=0)$$
$$a'E(\tilde{W}_i|Z_i=1) = a_1E(\tilde{W}_i|Z_i=1) + a_2E(\tilde{W}_iX_{i,1}|Z_i=1) + a_3E(\tilde{W}_iX_{i,2}|Z_i=1)$$

4. Prove the identification of the key parameters using the D-RSMD method for Model (4).

The D-RSMD estimator relies on the invertibility of the following matrix. Because it is not symmetric, we cannot use the same method to prove the identification problem. There is an alternative proof.

$$E\left[\kappa_{j,l}\left(\tilde{P}_{j}-\frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right)\tilde{P}_{l}'\right]=E\left[\kappa_{j,l}\tilde{P}_{j}\tilde{P}_{l}'\right]-E\left[\kappa_{j,l}\frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\tilde{P}_{l}'\right]$$

The invertibility of the matrix is illustrated by a simple example. X_i and W_i are random variables. $P_i = [W_i, W_i X_i]'$. W is the treatment. X is the control. Z is the instrument.

$$\begin{split} \tilde{P}_{j} &\equiv \begin{pmatrix} \tilde{W}_{j} \\ \tilde{W}_{j}X_{j} \end{pmatrix} = \begin{pmatrix} W_{j} - E(W_{j}|X_{j}) \\ W_{j}X_{j} - E(W_{j}|X_{j})X_{j} \end{pmatrix} \\ g_{0,\tilde{P}_{1,m}}(X_{l}) &= E[(P_{1m} - g_{0,P_{1}}(X_{m}))\kappa_{m,l}|X_{l}] = E[\tilde{W}_{j}\kappa_{j,l}|X_{l}] = a(X_{l}), \\ g_{0,\tilde{P}_{2,j}}(X_{l}) &= E[(P_{2j} - g_{0,P_{2}}(X_{j}))\kappa_{j,l}|X_{l}] = E[\tilde{W}_{j}X_{j}\kappa_{j,l}|X_{l}] = b(X_{l}), \\ g_{0,\kappa_{m,l}}(X_{l}) &= E[\kappa_{m,l}|X_{l}] = c(X_{l}). \end{split}$$

Based on the definition of \tilde{P}_j , $E(\tilde{P}_j) = 0$.

$$E(\kappa_{j,l}\tilde{P}_{j}\tilde{P}_{l}') = E \begin{bmatrix} \kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l} & \kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}X_{l} \\ \kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}X_{j} & \kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}X_{l}X_{j} \end{bmatrix}$$

$$E\left[\kappa_{j,l}\frac{g_{0,\tilde{P}_m}(X_l)}{g_{0,\kappa_{m,l}}(X_l)}\tilde{P}_l'\right] = E\left[\kappa_{j,l}\frac{a(X_l)}{c(X_l)}\tilde{W}_l \quad \kappa_{j,l}\frac{a(X_l)}{c(X_l)}\tilde{W}_l X_l\right]$$

$$\begin{split} E(\kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}) &= E[\kappa_{j,l}E(\tilde{W}_{j}|Z_{j})E(\tilde{W}_{l}|Z_{l})] \\ &= E[\kappa_{j,l}E(\tilde{W}_{j}|Z_{j})E(\tilde{W}_{l}|Z_{l})|Z_{j} = 1, Z_{l} = 1]P_{r}(Z_{j} = 1, Z_{l} = 1) \\ &+ E[\kappa_{j,l}E(\tilde{W}_{j}|Z_{j})E(\tilde{W}_{l}|Z_{l})|Z_{j} = 1, Z_{l} = 0]P_{r}(Z_{j} = 1, Z_{l} = 0) \\ &+ E[\kappa_{j,l}E(\tilde{W}_{j}|Z_{j})E(\tilde{W}_{l}|Z_{l})|Z_{j} = 0, Z_{l} = 1]P_{r}(Z_{j} = 0, Z_{l} = 1) \\ &+ E[\kappa_{j,l}E(\tilde{W}_{j}|Z_{j})E(\tilde{W}_{l}|Z_{l})|Z_{j} = 0, Z_{l} = 0]P_{r}(Z_{j} = 0, Z_{l} = 0) \end{split}$$

In the simulations, we use Gaussian CDF for $\mu(t)$ inside $\kappa_{j,l}$. $\kappa_{j,l} = 1$ when instrument variables $Z_j = Z_l$ and $\kappa_{j,l} = a \sim 0.61$ when $Z_j \neq Z_l$.

$$\begin{split} E(\kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}) &= E(\tilde{W}_{j}|Z_{j}=1)E(\tilde{W}_{l}|Z_{l}=1)P_{r}(Z_{j}=1)P_{r}(Z_{l}=1)\\ &+ aE(\tilde{W}_{j}|Z_{j}=1)E(\tilde{W}_{l}|Z_{l}=0)P_{r}(Z_{j}=1)P_{r}(Z_{l}=0)\\ &+ aE(\tilde{W}_{j}|Z_{j}=0)E(\tilde{W}_{l}|Z_{l}=1)P_{r}(Z_{j}=0)P_{r}(Z_{l}=1)\\ &+ E(\tilde{W}_{j}|Z_{j}=0)E(\tilde{W}_{l}|Z_{l}=0)P_{r}(Z_{j}=0)P_{r}(Z_{l}=0) \end{split}$$

Because
$$E(\tilde{W}_j | Z_j = 1) E(\tilde{W}_l | Z_l = 1) P_r(Z_j = 1) P_r(Z_l = 1)$$

+ $E(\tilde{W}_j | Z_j = 1) E(\tilde{W}_l | Z_l = 0) P_r(Z_j = 1) P_r(Z_l = 0)$
= $E(\tilde{W}_j | Z_j = 1) P_r(Z_j = 1) [E(\tilde{W}_l | Z_l = 1) P_r(Z_l = 1) + E(\tilde{W}_l | Z_l = 0) P_r(Z_l = 0)]$
= 0,

$$E(\kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}) = (1-a)E(\tilde{W}_{j}|Z_{j}=1)E(\tilde{W}_{l}|Z_{l}=1)P_{r}(Z_{j}=1)P_{r}(Z_{l}=1)$$
$$+(1-a)E(\tilde{W}_{j}|Z_{j}=0)E(\tilde{W}_{l}|Z_{l}=0)P_{r}(Z_{j}=0)P_{r}(Z_{l}=0)$$

Denote $E(\tilde{W}_l|Z_l=1)$ as θ_1 , $E(\tilde{W}_l|Z_l=0)$ as θ_2 , and $P_r(Z_j=1)$ as b. We have

$$E(\kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}) = (1-a)(\theta_{1}^{2}b^{2} + \theta_{2}^{2}(1-b)^{2})$$

Follow the same steps,

$$E[\kappa_{j,l}\tilde{W}_j\tilde{W}_lX_l] = (1-a)E(\tilde{W}_j|Z_j = 1)E(X_l\tilde{W}_l|Z_l = 1)P_r(Z_j = 1)P_r(Z_l = 1)$$
$$+(1-a)E(\tilde{W}_j|Z_j = 0)E(X_l\tilde{W}_l|Z_l = 0)P_r(Z_j = 0)P_r(Z_l = 0)$$

Denote $E(X_l \tilde{W}_l | Z_l = 1)$ as θ_3 , $E(X_l \tilde{W}_l | Z_l = 0)$ as θ_4 .

$$E[\kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}X_{l}] = (1-a)(\theta_{1}\theta_{3}b^{2} + \theta_{2}\theta_{4}(1-b)^{2})$$

$$E[\kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}X_{l}X_{j}] = (1-a)E(X_{j}\tilde{W}_{j}|Z_{j} = 1)E(X_{l}\tilde{W}_{l}|Z_{l} = 1)P_{r}(Z_{j} = 1)P_{r}(Z_{l} = 1)$$

$$+(1-a)E(X_{j}\tilde{W}_{j}|Z_{j} = 0)E(X_{l}\tilde{W}_{l}|Z_{l} = 0)P_{r}(Z_{j} = 0)P_{r}(Z_{l} = 0)$$

$$E[\kappa_{j,l}\tilde{W}_{j}\tilde{W}_{l}X_{l}X_{j}] = (1-a)(\theta_{3}^{2}b^{2} + \theta_{4}^{2}(1-b)^{2})$$

$$E(\kappa_{j,l}\tilde{P}_{j}\tilde{P}'_{l}) = \begin{bmatrix} (1-a)(\theta_{1}^{2}b^{2} + \theta_{2}^{2}(1-b)^{2}) & (1-a)(\theta_{1}\theta_{3}b^{2} + \theta_{2}\theta_{4}(1-b)^{2}) \\ (1-a)(\theta_{1}\theta_{3}b^{2} + \theta_{2}\theta_{4}(1-b)^{2}) & (1-a)(\theta_{3}^{2}b^{2} + \theta_{4}^{2}(1-b)^{2}) \end{bmatrix}$$

$$E\left[\kappa_{j,l}\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}\right] = E\left[\kappa_{j,l}E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}\right)\right]$$

$$= E\left[\kappa_{j,l}E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}\right)|Z_{j} = 1, Z_{l} = 1\right]P_{r}(Z_{j} = 1, Z_{l} = 1)$$

$$+ E\left[\kappa_{j,l}E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}\right)|Z_{j} = 1, Z_{l} = 0\right]P_{r}(Z_{j} = 1, Z_{l} = 0)$$

$$+ E\left[\kappa_{j,l}E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}\right)|Z_{j} = 0, Z_{l} = 1\right]P_{r}(Z_{j} = 0, Z_{l} = 1)$$

$$+ E\left[\kappa_{j,l}E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}\right)|Z_{j} = 0, Z_{l} = 0\right]P_{r}(Z_{j} = 0, Z_{l} = 0)$$

$$E\left[\kappa_{j,l}\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}\right] = E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=1\right)P_{r}(Z_{j}=1)P_{r}(Z_{l}=1)$$

$$+ aE\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=0\right)P_{r}(Z_{j}=1)P_{r}(Z_{l}=0)$$

$$+ aE\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=1\right)P_{r}(Z_{j}=0)P_{r}(Z_{l}=1)$$

$$+ E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=0\right)P_{r}(Z_{j}=0)P_{r}(Z_{l}=0)$$

$$\begin{split} E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}\right) &= E\left(\frac{a(X_{l})}{c(X_{l})}E(\tilde{W}_{l}|X_{l})\right) = 0\\ \text{with } \tilde{P}_{j} &= [\tilde{W}_{j},\tilde{W}_{j}X_{j}]' = [W_{j} - E(W_{j}|X_{j}), W_{j}X_{j} - E(W_{j}|X_{j})X_{j}]\\ E\left[\kappa_{j,l}\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}\right] &= P_{r}(Z_{j}=1)(1-a)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=1\right)P_{r}(Z_{l}=1)\\ &+ aP_{r}(Z_{j}=1)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=1\right)P_{r}(Z_{l}=1)\\ &+ aP_{r}(Z_{j}=1)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=0\right)P_{r}(Z_{l}=0)\\ &+ aE\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=1\right)P_{r}(Z_{l}=0)P_{r}(Z_{l}=1)\\ &+ E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=0\right)P_{r}(Z_{l}=0)\\ &= P_{r}(Z_{j}=1)(1-a)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=1\right)P_{r}(Z_{l}=1)\\ &+ P_{r}(Z_{j}=0)(1-a)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}|Z_{l}=0\right)P_{r}(Z_{l}=0) \end{split}$$

When we have $P_r(Z_j = 1) = 0.5, d_1 = 0.$

Denote $E\left(\frac{a(X_l)}{c(X_l)}\tilde{W}_l|Z_l=1\right)$ as δ_1 and $E\left(\frac{a(X_l)}{c(X_l)}\tilde{W}_l|Z_l=0\right)$ as δ_2 .

$$E\left[\kappa_{j,l}\frac{a(X_l)}{c(X_l)}\tilde{W}_l\right] = (1-a)(b^2\delta_1 + (1-b)^2\delta_2)$$

Follow the same steps,

$$\begin{split} E\left[\kappa_{j,l}\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}\right] &= P_{r}(Z_{j}=1)(1-a)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}|Z_{l}=1\right)P_{r}(Z_{l}=1) \\ &+ aP_{r}(Z_{j}=1)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}|Z_{l}=1\right)P_{r}(Z_{l}=1) \\ &+ aP_{r}(Z_{j}=1)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}|Z_{l}=0\right)P_{r}(Z_{l}=0) \\ &+ aE\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}|Z_{l}=1\right)P_{r}(Z_{j}=0)P_{r}(Z_{l}=1) \\ &+ E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}|Z_{l}=0\right)P_{r}(Z_{l}=0) \\ &= P_{r}(Z_{j}=1)(1-a)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}|Z_{l}=1\right)P_{r}(Z_{l}=1) \\ &+ P_{r}(Z_{j}=0)(1-a)E\left(\frac{a(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}|Z_{l}=0\right)P_{r}(Z_{l}=0) \end{split}$$

Denote
$$E\left(\frac{a(X_l)}{c(X_l)}\tilde{W}_l X_l | Z_l = 1\right)$$
 as δ_3 and $E\left(\frac{a(X_l)}{c(X_l)}\tilde{W}_l X_l | Z_l = 0\right)$ as δ_4 .

$$E\left[\kappa_{j,l}\frac{a(X_l)}{c(X_l)}\tilde{W}_lX_l\right] = (1-a)(b^2\delta_3 + (1-b)^2\delta_4)$$

Denote
$$E\left(\frac{b(X_l)}{c(X_l)}\tilde{W}_l|Z_l=1\right)$$
 as δ_5 and $E\left(\frac{b(X_l)}{c(X_l)}\tilde{W}_l|Z_l=0\right)$ as δ_6 .
 $E\left[\kappa_{j,l}\frac{b(X_l)}{c(X_l)}\tilde{W}_l\right] = (1-a)(b^2\delta_5 + (1-b)^2\delta_6)$

Denote $E\left(\frac{b(X_l)}{c(X_l)}\tilde{W}_l X_l | Z_l = 1\right)$ as δ_7 and $E\left(\frac{b(X_l)}{c(X_l)}\tilde{W}_l X_l | Z_l = 0\right)$ as δ_8 .

$$E\left[\kappa_{j,l}\frac{b(X_{l})}{c(X_{l})}\tilde{W}_{l}X_{l}\right] = (1-a)(b^{2}\delta_{7} + (1-b)^{2}\delta_{8})$$

$$E\left[\kappa_{j,l}\left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right)\tilde{P}_{l}'\right] \text{ becomes:}$$

$$\begin{bmatrix} (1-a)(\theta_{1}^{2}b^{2} + \theta_{2}^{2}(1-b)^{2}) - (1-a)(b^{2}\delta_{1} + (1-b)^{2}\delta_{2}) & e_{1} \\ (1-a)(\theta_{1}\theta_{3}b^{2} + \theta_{2}\theta_{4}(1-b)^{2}) - (1-a)(b^{2}\delta_{5} + (1-b)^{2}\delta_{6}) & e_{2} \end{bmatrix}$$

$$e_{1} = (1-a)(\theta_{1}\theta_{3}b^{2} + \theta_{2}\theta_{4}(1-b)^{2}) - (1-a)(b^{2}\delta_{3} + (1-b)^{2}\delta_{4})$$

$$e_{2} = (1-a)(\theta_{3}^{2}b^{2} + \theta_{4}^{2}(1-b)^{2}) - (1-a)(b^{2}\delta_{7} + (1-b)^{2}\delta_{8})$$
Since a is not 1, the determinant of $E\left[\kappa_{j,l}\left(\tilde{P}_{j} - \frac{g_{0,\tilde{P}_{m}}(X_{l})}{g_{0,\kappa_{m,l}}(X_{l})}\right)\tilde{P}_{l}'\right]$ depends on the exact values for these numbers.

5. Simulation illustration for RSMD and D-RSMD

Here are one simulation example for illustration (Model (4)).

$$y_{i} = W_{i,1} + W_{i,1} * X_{i} + X_{i} + u_{i}$$
$$W_{i,1} = 2Z_{i} + v_{i,1}$$
$$W_{i,1} * X_{i} = 2Z_{i} * X_{i} + v_{i,1} * X_{i}$$

(a) For RSMD,

$$a'E(\tilde{W}_{i}|Z_{i}) = 2a_{1}Z_{i} - 2a_{1}E(E(Z_{i}|X_{i})|Z_{i})$$

+ $2a_{2}Z_{i}E(X_{i}|Z_{i}) - 2a_{2}E(E(Z_{i}|X_{i})X_{i}|Z_{i})$
= 0

When Z_i and X_i are independent and $E(Z_i)$ is a number (for instance, $E(Z_i) = 1$),

$$a_1 Z_i - a_1 E(Z_i) + a_2 Z_i E(X_i) - a_2 E(X_i) E(Z_i) = 0$$
$$(a_1 + a_2 E(X_i)) Z_i - (a_1 + a_2 E(X_i)) E(Z_i) = 0$$

- i. When $E(Z_i) = 0$, a_1 must be 0 when Z_i is continuous or discrete. a_2 can be any value. Hence, the first parameter is identified.
- ii. When $E(Z_i) = 1$, there are infinite combinations for (a_1, a_2) to make

$$a'E(X_i|Z_i) = 0$$

. Both parameters are not identified.

(b) For D-RSMD, because we have extra terms, it should identify both parameters when RSMD cannot.

All of the simulation results are included in Table 14.

Estimator	Instrument	Med.Bias	MAD	Med.SE	RR	Mean.Bias	Mean.SE	Mean.AHSE
		Example	e 2(a)ii:	Continu	ous ins	trument		
SMD β_1	$(\mathbb{Z}_{i,1},\mathbb{Z}_{i,2})$	0.000	0.005	0.007	0.055	0.000	0.007	0.007
SMD β_2	$(\mathbb{Z}_{i,1},\mathbb{Z}_{i,2})$	0.000	0.006	0.009	0.052	0.000	0.009	0.009
SMD β_1	$Z_{i,1}$	-0.011	0.034	0.065	0.026	-0.021	1.598	37.825
SMD β_2	$Z_{i,1}$	0.042	0.132	0.256	0.041	0.087	6.394	149.586
SMD β_1	$Z_{i,2}$	0.018	0.106	0.209	0.022	-0.050	18.259	783.448
SMD β_2	$Z_{i,2}$	-0.007	0.036	0.070	0.014	0.008	5.521	238.702
TSLS β_1	$(\mathbb{Z}_{i,1},\mathbb{Z}_{i,2})$	0.000	0.004	0.006	0.063	0.000	0.006	0.006
TSLS β_2	$(\mathbb{Z}_{i,1},\mathbb{Z}_{i,2})$	0.000	0.006	0.008	0.055	0.000	0.008	0.008
		Examp	ole 2(a)	iii: Binar	y instr	ument		
SMD β_1	$(Z_{i,1}, Z_{i,2})$	0.000	0.008	0.012	0.052	0.000	0.012	0.012
SMD β_2	$(\mathbb{Z}_{i,1},\mathbb{Z}_{i,2})$	0.000	0.010	0.015	0.047	0.000	0.015	0.015
SMD β_1	$Z_{i,1}$	0.000	0.011	0.017	0.053	0.000	0.017	0.017
SMD β_2	$Z_{i,1}$	-0.001	0.017	0.026	0.049	0.000	0.026	0.026
SMD β_1	$Z_{i,2}$	0.000	0.013	0.020	0.046	0.000	0.020	0.021
SMD β_2	$Z_{i,2}$	0.000	0.014	0.021	0.046	0.000	0.021	0.021
TSLS β_1	$(\mathbb{Z}_{i,1},\mathbb{Z}_{i,2})$	0.000	0.008	0.013	0.054	0.000	0.013	0.013
TSLS β_2	$(\mathbb{Z}_{i,1},\mathbb{Z}_{i,2})$	0.000	0.011	0.017	0.048	0.000	0.017	0.017
		Exam	ple 2(b))i: Binary	/ instru	ument		
SMD β_1	Z_i	0.000	0.013	0.019	0.036	0.000	0.024	0.027
SMD β_2	Z_i	0.000	0.043	0.077	0.039	0.019	0.738	0.596
		Exam	ple 2(b)	ii: Binar	y instr	ument		
SMD β_1	Z_i	0.000	0.046	0.077	0.039	-0.020	0.742	0.602
SMD β_2	Z_i	0.000	0.043	0.077	0.039	0.019	0.738	0.596
		Example	e 5(a)i:	Continuo	ous ins	trument		
RSMD β_1	Z_i	-0.001	0.009	0.013	0.032	0.000	0.171	1.937
RSMD β_2	Z_i	0.000	0.056	0.138	0.006	0.020	4.528	77.709
D-RSMD β_1	Z_i	-0.001	0.009	0.013	0.041	-0.001	0.021	0.114
D-RSMD β_2	Z_i	0.000	0.041	0.066	0.029	0.026	1.093	7.282
		Example	e 5(a)ii:	Continu	ous ins	trument		
RSMD β_1	Z_i	-0.002	0.057	0.137	0.008	-0.039	7.188	120.469
RSMD β_2	Z_i	0.001	0.055	0.137	0.005	0.038	7.331	122.732
D-RSMD β_1	Z_i	-0.001	0.043	0.067	0.031	-0.002	0.970	2.922
D-RSMD β_2	Z_i	0.002	0.041	0.066	0.029	0.001	0.970	2.919

Table 14: Identification Illustration

Notes: Mean.AHSE stands for Mean.Asympt.Heterosk.SE. There are 5000 rounds of simulation and 2000 observations for each simulation study.