# Reduction of electrostatic turbulence in a quasi-helically symmetric stellarator via critical gradient optimization

G. T. Roberg-Clark<sup>1</sup>†, P. Xanthopoulos <sup>1</sup> and G. G. Plunk <sup>1</sup>

<sup>1</sup>Max-Planck-Institut Für Plasmaphysik, D-17491, Greifswald, Germany

(Received xx; revised xx; accepted xx)

We present a stellarator configuration optimized for a large threshold ("critical gradient") for the onset of the ion temperature gradient (ITG) driven mode, which achieves the largest critical gradient we have seen in any stellarator. Above this threshold, gyrokinetic simulations show that the configuration has low turbulence levels over an experimentally relevant range of the drive strength. The applied optimization seeks to maximize the drift curvature, leading to enhanced local-shear stabilization of toroidal ITG modes, and the associated turbulence. These benefits are combined with excellent quasisymmetry, yielding low neoclassical transport and vanishingly small alpha particle losses. Analysis of the resulting configuration suggests a trade-off between magnetohydrodynamic (MHD) and ITG stability.

### 1. Introduction

The excitation of the ion temperature gradient (ITG) mode in magnetically confined fusion devices leads to turbulence that is responsible for energy losses, which reduce the plasma confinement needed for potential fusion performance. For instance, it has been argued that, during operation of the Wendelstein 7-X (W7-X) stellarator in electron heating scenarios, the ITG mode leads to so-called "ion temperature clamping" (Beurskens et al. 2021, 2022), thus preventing the heating of ions in the plasma core above 2 keV. While resolving the location and extent of the fine-scale (ion Larmor radius) fluctuations is difficult in experiments, implementation of diagnostics using phase contrast imaging (Bähner et al. 2021) and Doppler reflectometry (Carralero et al. 2021) has helped to characterize the dependence of W7-X experimental temperature and density profiles on such fluctuations.

In order to lessen the negative effects of the ITG mode, a possible strategy to follow involves the manipulation of the density and electron profiles, as implemented in W7-X using pellet injections (Bozhenkov et al. 2020; Pablant et al. 2020). A stellarator magnetic field may also be shaped to reduce microturbulence losses during steady-state operation. In the case of electron-temperature gradient turbulence, it is argued that multiple field periods are stabilizing via reduction of the parallel connection length (Plunk et al. 2019), while trapped-electron mode turbulence can be ameliorated by requiring that the parallel adiabatic invariant J achieves its maximum on the magnetic axis (Proll et al. 2012), the so-called "Maximum-J" property (Helander et al. 2012; Mackenbach et al. 2022).

While most previous works have exploited magnetic field shaping in order to lower the rate of the ITG transport scaling ("stiffness"), see, e.g., (Mynick *et al.* 2010; Xanthopoulos *et al.* 2014; Hegna *et al.* 2018; Nunami *et al.* 2013; Jorge & Landreman 2021;

Stroteich et al. 2022), the goal of the present work is to increase the linear onset of ITG modes ("critical gradient") (Roberg-Clark et al. 2021), since for configurations with stiff transport, it can determine the radial ion temperature profile (Baumgaertel et al. 2013). Even though the nature of the marginally unstable fluctuations may differ between low-magnetic-shear stellarators (Zocco et al. 2018, 2022; Bhattacharjee et al. 1983) [Floquet-like] and tokamaks (Terry et al. 1982; Romanelli 1989; Biglari et al. 1989; Jenko et al. 2001; Plunk et al. 2014) [Toroidal-like], it seems that the size of the socalled "drift curvature", a factor appearing in the gyrokinetic equation, can be used to predict the critical gradient for most optimized stellarator configurations (Roberg-Clark et al. 2022). Here we leverage this predictive capability to generate a new quasi-helicallysymmetric configuration with low neoclassical transport and large ITG critical gradient. It turns out that turbulent losses from ITG modes above this threshold are suppressed when compared with a well-known quasi-helically symmetric configuration, in particular near the critical gradient. We attribute this enhanced stability, at least in part, to the high onset gradient of localized, toroidal ITG modes, which are further damped by flux expansion and local shear effects.

The paper is structured as follows. In section 2 we define the linear gyrokinetic system used to analyze ITG mode properties. Section 3 describes the optimization procedure used to generate the new configuration, while Section 4 presents results of the optimization, along with a description of the local shear effect. We conclude with an outlook on the apparent competition between ITG and MHD stability in section 5.

# 2. Linear gyrokinetic equation

Following (Plunk et al. 2014), we use the standard gyrokinetic system of equations (Brizard & Hahm 2007) to describe electrostatic fluctuations destabilized along a thin flux tube tracing a magnetic field line. The ballooning transform (Dewar & Glasser 1983) and twisted slicing representation (Roberts & Taylor 1965) are used to separate out the fast perpendicular (to the magnetic field) scale from the slow parallel scale. The magnetic field representation in field following (Clebsch) representation reads,  $\mathbf{B} = \nabla \psi \times \nabla \alpha$ , where  $\psi$  is a toroidal flux surface label and  $\alpha = \vartheta - \iota \phi$  labels the magnetic field line on the surface, with q the safety factor,  $\vartheta$  the poloidal angle, and  $\phi$  the toroidal angle. The perpendicular wave vector is then expressed as  $\mathbf{k}_{\perp} = k_{\alpha} \nabla \alpha + k_{\psi} \nabla \psi$ , where  $k_{\alpha}$  and  $k_{\psi}$  are constants, so the variation of  $\mathbf{k}_{\perp}(l)$  stems from that of the geometric quantities  $\nabla \alpha$  and  $\nabla \psi$ , with  $\ell$  the field-line-following (arc length) coordinate.

We assume Boltzmann-distributed (adiabatic) electrons, thus solving for the perturbed ion distribution  $g_i(v_{\parallel}, v_{\perp}, \ell, t)$ , defined to be the non-adiabatic part of  $\delta f_i$  ( $\delta f_i = f_i - f_{i0}$ ) with  $f_i$  the ion distribution function and  $f_{i0}$  a Maxwellian. The electrostatic potential is  $\phi(\ell)$ , and  $v_{\parallel}$  and  $v_{\perp}$  are the particle velocities parallel and perpendicular to the magnetic field, respectively.

The linear gyrokinetic equation for the ions is written

$$iv_{\parallel} \frac{\partial g}{\partial \ell} + (\omega - \widetilde{\omega}_d)g = \varphi J_0(\omega - \widetilde{\omega}_*) f_0$$
 (2.1)

with the following definitions:  $J_0 = J_0(k_{\perp}v_{\perp}/\Omega) = J_0(k_{\perp}\rho\sqrt{2}v_{\perp}/v_{\rm T})$ ; the ion thermal velocity is  $v_{\rm T} = \sqrt{2T/m}$  and the thermal ion Larmor radius is  $\rho = v_{\rm T}/(\Omega\sqrt{2})$ ; n and T are the background ion density and temperature; q is the ion charge;  $\varphi = q\phi/T$  is the normalized electrostatic potential;  $\Omega = qB/m$  is the cyclotron frequency, with  $B = |\mathbf{B}|$ 

the magnetic field strength. Assuming Boltzmann electrons, the quasineutrality condition is

$$\int d^3 \mathbf{v} J_0 g = n(1+\tau)\varphi, \tag{2.2}$$

where  $\tau = T/(ZT_e)$  with the charge ratio defined as  $Z = q/q_e$ . The equilibrium distribution is the Maxwellian

$$f_0 = \frac{n}{(v_T^2 \pi)^{3/2}} \exp(-v^2/v_T^2),$$
 (2.3)

and we introduce the velocity-dependent diamagnetic frequency

$$\widetilde{\omega}_* = \omega_*^{\mathrm{T}} \left[ \frac{v^2}{v_{\mathrm{T}}^2} - \frac{3}{2} \right] \tag{2.4}$$

where we neglect background density variation and define  $\omega_*^{\mathrm{T}} = (Tk_{\alpha}/q)d\ln T/d\psi$ . The magnetic drift frequency is  $\widetilde{\omega}_d = \mathbf{v}_d \cdot \mathbf{k}_{\perp}$  and the magnetic drift velocity is  $\mathbf{v}_d = \hat{\mathbf{b}} \times ((v_{\perp}^2/2)\nabla \ln B + v_{\parallel}^2 \kappa)/\Omega$ , where  $\kappa = \hat{\mathbf{b}} \cdot \nabla \hat{\mathbf{b}}$ . We take  $\nabla \ln B = \kappa$ , the zero  $\beta$  approximation, for simplicity. We then let

$$\widetilde{\omega}_d = \frac{\mathbf{k}_{\perp} \cdot (\hat{\mathbf{b}} \times \boldsymbol{\kappa}) v_T^2}{\Omega} \left[ \frac{v_{\parallel}^2}{v_{\mathrm{T}}^2} + \frac{v_{\perp}^2}{2v_{\mathrm{T}}^2} \right] = \omega_d(\ell) \left[ \frac{v_{\parallel}^2}{v_{\mathrm{T}}^2} + \frac{v_{\perp}^2}{2v_{\mathrm{T}}^2} \right], \tag{2.5}$$

where the velocity-independent drift frequency  $\omega_d(\ell)$  generally varies along the field line. Positive values of  $\omega_d$  correspond to "bad" curvature, i.e. are destabilizing for ITG modes, assuming  $\omega_*^{\rm T}$  is negative. Assuming  $k_{\psi} = 0$  for simplicity, we then define the drift curvature  $K_d$  by writing

$$\omega_d(\ell) \propto K_d(\ell) \equiv a^2 \nabla \alpha \cdot \hat{\mathbf{b}} \times \kappa,$$
 (2.6)

where  $K_d$  contains the purely geometric variation of the drift frequency and a is the minor radius of the flux surface at the edge. For the purpose of analyzing gyrokinetic simulation results we define the metrics  $g^{yy} = a^2 s_0/(q_0^2)(\nabla \alpha)^2$ , with  $s = \psi/\psi_{edge}$  the toroidal flux normalized to its value at the last closed flux surface  $(q_0)$  is the safety factor on a particular surface  $s = s_0$ , and  $s^{xx} = a^2/(4s_0)(\nabla s)^2$ . Finally, we define the poloidal wavenumber  $s_y = (q_0/\sqrt{s_0})k_\alpha$ .

## 2.1. Averaging effect in the gyrokinetic equation

An integral equation can be derived from Eqns. 2.1-2.2 assuming "outgoing" boundary conditions  $g(v_{\parallel} > 0, \ell = -\infty) = g(v_{\parallel} < 0, \ell = \infty) = 0$ , consistent with ballooning modes that decay as  $|\ell| \to \infty$  (Connor *et al.* 1980; Romanelli 1989). To enforce these conditions we assume the system has non-zero global shear,  $d\epsilon/d\psi \neq 0$ , though it is allowed to be small. One then obtains (Plunk *et al.* 2014)

$$(1+\tau)\varphi(\ell) = \frac{-2i}{v_{\rm T}\sqrt{\pi}} \int_0^\infty \frac{dx_{\parallel}}{x_{\parallel}} \int_0^\infty dx_{\perp} x_{\perp}(\omega - \widetilde{\omega}_*) J_0$$
$$\times \int_{-\infty}^\infty d\ell' J_0' \exp(-x^2 + i \operatorname{sgn}(\ell - \ell') M(\ell', \ell)) \varphi(\ell'), \tag{2.7}$$

where  $x_{\perp} = v_{\perp}/v_{\mathrm{T}}$  and  $x_{\parallel} = v_{\parallel}/v_{\mathrm{T}}$ , sgn gives the sign of its argument,  $J_0 =$ 

 $J_0\left(\sqrt{2b(\ell)}x_\perp\right)$ ,  $J_0'=J_0\left(\sqrt{2b(\ell')}x_\perp\right)$ , and  $b(\ell)=\rho^2k_\perp^2(\ell)$ . The physics of the drift resonance is contained in the factor

$$M(\ell',\ell) = \int_{\ell'}^{\ell} \frac{\omega - \widetilde{\omega}_d(\ell'')}{v_{\rm T} x_{\parallel}} d\ell''. \tag{2.8}$$

The kernel M contains an averaging over  $\ell''$  which suggests the use of a smoothed curvature profile, where small-scale ripples are averaged out and a "coarse-grained" drift curvature can then be substituted into (2.7) (see Roberg-Clark *et al.* (2022) for further discussion).

We neglect particle trapping so  $x_{\parallel}$  and  $x_{\perp}$  do not depend on  $\ell$ . Most evidence indicates that the ITG mode uniformly responds to changes in the parameters  $\tau$  and  $dn/d\psi$  and is stabilized by increasing either of them, except for a relatively small region of parameter space where positive density gradients can destabilize the mode. For simplicity we thus set  $\tau = 1$  and  $\nabla n = 0$ .

#### 2.2. Estimating the ITG mode critical gradient

As found in Roberg-Clark et al. (2022), the geometric dependence of the linear ITG critical gradient can be estimated in a simple way. The drift curvature profile (eqn. 2.6) on a particular magnetic field line (at radial location  $s = s_0$  and field line  $\alpha = \alpha_0$ ) is fitted with quadratic curves in regions of bad curvature, which we refer to as "drift wells". The fitting acts an effective coarse-graining of the geometry, producing a smoothed curvature profile as mentioned in the preceding section. Using the fitted profile, a value of the predicted critical gradient is produced, corresponding to the drift well with the smallest critical gradient. The formula reads:

$$F_{crit} = \frac{a}{L_{T,crit}} = 2.66 \left(\frac{a}{R_{\text{eff}}}\right) \tag{2.9}$$

with  $a/R_{\rm eff}$  the peak value of the drift curvature within the drift well. Here we have ignored the parallel stabilizing term included in Roberg-Clark et al. (2022) as we intend to produce a configuration with a large critical gradient solely by large "drift curvature"; see the later discussion of geometric interpretation in Section 4.1.

# 3. Optimization Method

We use the SIMSOPT software framework (Landreman *et al.* 2021) to generate a vacuum stellarator configuration with quasi-helical (QH) symmetry and a significant linear ITG critical gradient as a result of drift curvature. The stellarator magnetic field is described by a boundary surface given in the Fourier representation

$$R(\vartheta,\phi) = \sum_{m,n} R_{m,n} \cos(m\vartheta - 4n\phi)$$
(3.1)

$$Z(\vartheta,\phi) = \sum_{m,n} Z_{m,n} \sin(m\vartheta - 4n\phi), \tag{3.2}$$

where we have assumed stellarator symmetry and have set the number of field periods to  $n_{fp} = 4$ . Global vacuum solutions are constructed at each iteration by running the VMEC (Hirshman & Whitson 1983) code, which solves the MHD equations using an energy-minimizing principle.

Optimization proceeds by treating the Fourier coefficients in equation (3.1) as parameters and varying them in order to find a least-squares minimization (using a trust region)




Figure 1. The boundary surface of HSK.

of the specified objective function, which reads

$$f = f_{OS} + (A - 4.10)^2 + (F_{crit} - 2.00)^2, \tag{3.3}$$

where

$$f_{QS} = \sum_{s_i} \left\langle \left( \frac{1}{B^3} [(-1 - \iota) \mathbf{B} \times \nabla B \cdot \nabla \psi - G \mathbf{B} \cdot \nabla B] \right)^2 \right\rangle$$
(3.4)

is the quasisymmetry residual taken from (Landreman & Paul 2022), with A = R/athe aspect ratio output by VMEC, and  $a/L_{T,crit}$  as defined in eqn. (2.9) with the field line  $(s_0 = 0.5, \alpha_0 = 0)$  chosen. Here  $G(\psi)$  is  $\mu_0/(2\pi)$  times the poloidal current outside the surface and we have set the toroidal current to zero, while  $\langle \cdot \rangle$  denotes a flux surface average over the individual surfaces  $s = s_i$ . We choose  $B = B(\theta + 4\phi_B)$ , with  $\theta$  and  $\phi_B$ the Boozer poloidal and toroidal angles, corresponding to quasi-helical symmetry with a negative axis helicity. The input equilibrium for the optimization is a "warm start" example file included in SIMSOPT with approximate quasi-helical symmetry,  $n_{fp} = 4$ and aspect ratio A = R/a = 6, with R the major radius and a the minor radius. The target A = 4.10 was chosen to enhance the drift curvature  $K_d$ , while we speculate  $a/L_{T.crit} = 2.00$  to be roughly the maximum achievable linear critical gradient in the absence of shear stabilization (Roberg-Clark et al. 2022). We calculate the residual  $f_{QS}$ on the surfaces s = [0, 0.1, 0.2, 0.3, 0.4, 0.5], thus leaving the region 0.5 < s < 1.0 free. We make this choice knowing that requiring a "precise" degree of quasisymmetry in the entire volume often has the effect of reducing the global magnetic shear in the configuration to vanishingly small values (Landreman & Paul 2022). Similarly to Landreman & Paul (2022), the number of Fourier coefficient parameters are increased in a series of four steps, with the toroidal and poloidal mode numbers in the VMEC calculation chosen to be  $m_{pol} = n_{tor} = [3, 5, 6, 6]$ . The final step is a refinement to see if a better local minimum can be found at the same resolution of six modes.

#### 4. Results

Figure 1 shows a surface plot of the outermost flux surface of the resulting optimized configuration, which we dub "HSK" (Helically-Symmetric Kompakt stellarator). The contours of B are plotted in the Boozer angle plane at s=0.25 in Fig. 2(a) indicating the quasi-helical symmetry. The highly compact configuration has roughly triangular cross




FIGURE 2. Properties of HSK. (a) Contours of B with the trajectory of a field line (black curve near the center) overlaid in Boozer coordinates. (b) Rotational transform as a function of radius. (c) Neoclassical transport coefficient  $\epsilon_{eff}$  as a function of radius.




Figure 3. Surface cuts of HSK taken at constant toroidal angle,  $\phi = \text{const.}$ 

sections (cuts at constant toroidal angle) [Fig. 3] and noticeably lacks a "bean-shaped" cross section at toroidal angle  $\phi=0$ , which most optimized stellarator configurations possess. The spatial separation between the flux surfaces is substantial, indicating low surface compression, while the lack of an indentation on the inboard side at  $\phi=0$  appears to be a result of not optimizing for a vacuum magnetic well. The configuration instead has a sizable magnetic hill, i.e.  $V''(\psi)$ , the second derivative of the surface volume, is positive at all radii. Heuristically speaking, this result is expected from configurations with average bad curvature, i.e. the average value of the drift curvature over the entire surface is positive. HSK has a rotational transform varying from  $\iota=1.3$  to 1.8 (demonstrating significant global magnetic shear) [Fig. 2(b)] and values of the neoclassical transport coefficient  $\epsilon_{eff}$  (Nemov et al. 1999) ranging from 0.4% on axis to 1.6% at the last closed flux surface [Fig. 2(c)], implying excellent neoclassical confinement. Using a symplectic integrator for guiding center trajectories (Albert et al. 2020) in this configuration but




FIGURE 4. Simulation of critical gradients versus model predictions on a log-log scale for flux tubes taken from several stellarator geometries, adapted from figure (1) in (Roberg-Clark *et al.* 2022), with the data point for HSK added. The field line ( $s_0 = 0.5, \alpha_0 = 0$ ) was chosen for all configurations.

rescaled to the same B and minor radius as the ARIES-CS reactor (Mau *et al.* 2008), we find no collisionless alpha particle losses after 10 ms for particles launched from inner radii, suggesting excellent quasisymmetry (figure not shown).

Linear gyrokinetic simulations in local flux tube geometry using the GENE code (Jenko et al. 2000) are performed to determine the critical gradient in the same manner as in Roberg-Clark et al. (2021, 2022), i.e. by reducing the applied temperature gradient until a single, marginally unstable mode remains. An unusually large resolution in  $\mu$  (i.e.  $v_{\perp}$  space) of  $n_{\mu}=32$  was required for numerical convergence of the critical gradient, likely because of the extreme values of curvature near the outboard midplane that imply significant linear phase mixing in  $v_{\perp}$  through the  $\nabla B$  drift term in eqn. (2.5). The linear ITG critical gradient at  $(s_0=0.5,\alpha_0=0)$  [ $k_y\rho=0.3$ ], while not reaching 2 as predicted by the fitting model (eqn. 2.9), nonetheless attains the value of  $a/L_{T,crit}=1.75$ , the highest we have seen in any stellarator (Fig. 4). Thus the simultaneous optimization for quasi-helical symmetry, aspect ratio, and ITG linear critical gradient was successful.

One might reasonably expect this particular ITG optimization strategy (increasing bad curvature) to heavily exacerbate linear growth rates, and thus nonlinear transport, of ITG modes above the linear critical gradient, implying a tradeoff between linear and nonlinear stability. Our results in the next section, however, oppose this intuition, revealing significant ITG stability above the critical gradient.

#### 4.1. Mechanisms of ITG turbulence suppression

One means by which a stellarator configuration can attain large values of the drift curvature  $K_d$  (eqn. 2.6) is by compression of the  $\alpha$  coordinate, i.e. large  $|\nabla \alpha|$ . This is indeed observed for HSK, in the metrics following the magnetic field line at  $(s_0 = 0.5, \alpha_0 = 0)$ , where we plot  $g^{yy} \propto g^{\alpha\alpha} = a^2 |\nabla \alpha|^2$  as a function of arc length  $\ell$ , which attains a value on the outboard midplane  $(\ell = 0)$  larger than 2 [Fig. 5(a)]. Note that this location at the outboard midplane corresponds to  $\phi = 0$  in figure 1(b). For comparison, we also plot the metrics along the same field line in the the HSX stellarator (Talmadge et al. 2008), another QH optimized configuration [Fig. 5(b)].

In a simple limit, when  $\nabla \alpha \cdot \nabla \psi \approx 0$ , and assuming weak variation of the overall magnetic field strength  $B = |\nabla \alpha \times \nabla \psi|$ , the  $\alpha$ -compression is directly linked to reduction of  $g^{xx} \propto g^{\psi\psi}$  (i.e.  $g^{xx} \sim 1/g^{yy}$ ). In such cases, reduced instability growths and turbulent




FIGURE 5. Metrics (defined near the end of section 2) as a function of the field-line-following coordinate  $\ell$  at the flux tube location ( $s_0 = 0.5, \alpha_0 = 0$ ), where HSK was optimized for a high critical gradient. The flux tubes were constructed with two poloidal turns. The vertical axis is in dimensionless units, such that the metrics for each configuration can be compared directly. (a) HSK. (b) HSX.




FIGURE 6. ITG mode linear growth rates as a function of  $k_y \rho_i$  for HSK (a) and HSX (b) for the temperature gradients  $a/L_T = 2, 3$ , and 4, at the flux tube location  $(s_0 = 0.25, \alpha_0 = 0)$ .

transport can be intuitively explained via expansion of flux surfaces, resulting in an effective suppression of the applied temperature gradient, since the physical temperature gradient  $|\nabla T| = dT/d\psi |\nabla \psi|$  scales with  $|\nabla \psi|$  (Plunk & Helander 2022; Helander & Plunk 2021; Stroteich et al. 2022; Angelino et al. 2009).

Typically in a stellarator the vector  $\nabla \alpha$  also develops a substantial component in the direction parallel to  $\nabla \psi$ , attributed to so-called "local shear" of the magnetic field lines (Helander 2014). This effect, distinct from flux expansion, is associated with a stabilization effect (Waltz & Boozer 1993) due to  $k_{\perp}\rho_i$  increasing, *i.e.* due a finite Larmor radius (FLR). Global shear may further amplify this damping through secular increase of  $g^{yy}$  along the field line, although we suspect this is not the dominant effect in the case of HSK. We see that both of these effects are at play with HSK, but note for the case of HSX that the pattern near  $\ell=0$  in the metrics is effectively inverted, i.e.  $|K_{da}|<<1$ ,  $g^{yy}<1$  and  $g^{xx}>1$ , which is typical of most optimized stellarator configurations.

We now carry out linear gyrokinetic simulations comparing HSK with HSX using the GENE code, but this time above marginal stability. We choose the flux tube ( $s_0 =$ 




FIGURE 7. (a) Nonlinear heat fluxes computed by GENE at the flux tube ( $s_0 = 0.25, \alpha_0 = 0$ ) for HSK and HSX, varying the applied temperature gradient. (b) The same plot as (a) but with the vertical axis on a log scale.

 $0.25, \alpha_0 = 0$ ) to check that the benefits of HSK are not confined to the field line optimized for ITG stability at  $(s_0 = 0.5, \alpha_0 = 0)$ . Figure 6(a) shows a strikingly narrow range of unstable wavenumbers for HSK in  $k_y \rho$  at the gradient  $a/L_T = 2$ . Much of the analysis of electrostatic modes in QH stellarators has focused on damped or subdominant eigenmodes (Faber et al. 2018; Terry et al. 2006; Sugama 1999), which can form the basis for nonlinear saturation of ITG turbulence (Pueschel et al. 2016; Hegna et al. 2018; Mckinney et al. 2019). We interpret the narrow linear growth rate spectrum of HSK at  $a/L_T = 2$  to mean that most of the linear eigenmode spectrum is stable at this gradient. It is worth mentioning that HSX sometimes has a favourable scaling of heat flux at these temperature gradients compared to other configurations (Mckinney et al. 2019; Plunk et al. 2017), despite larger linear growth rates. HSX was also not optimized for reduced turbulent transport, so one should not expect it to outperform turbulence-optimized configurations.

A nonlinear gyrokinetic analysis with GENE using the same flux tubes shows that the narrow and small linear growth rates in HSK are complemented by a stark reduction in heat fluxes, a truer figure of merit for confinement. In comparison with HSX, we find heat fluxes are smaller by roughly a factor of 4 at  $a/L_T = 2$  (Fig. 7(b)). While the reduction is not as strong at higher gradients, the curve for HSK remains below that of HSX for the full range of gradients studied. The effect of the optimization is most potent near the critical gradient,  $a/L_T \simeq 2-3$ . This range of gradients is likely the most relevant to experiments, as seen in data for (Beurskens et al. 2021) and in global gyrokinetic simulations of (Bañón Navarro et al. 2022) W7-X. For the same heating power, we can thus expect HSK to have a significantly larger temperature gradient compared to HSX, leading to improved confinement and higher core ion temperatures. Furthermore, HSK has the advantage of a reduced aspect ratio, which implies a favorable confinement time compared to larger aspect ratio devices with the same transport coefficients. Note that the turbulence-optimized WISTELL-C QH configuration presented in Hegna et al. (2022) is likely compared with HSX at the radial location s = 0.5, achieving a factor of 2 reduction in heat fluxes at  $a/L_T = 4$ . The same reduction factor for  $a/L_T = 4$  at  $s_0 = 0.25$  is found when comparing HSK and HSX [Fig. 7(b)].

While bad curvature has markedly increased in HSK relative to HSX, stiffness of ITG transport has not worsened for the range  $2 \leqslant a/L_T \leqslant 4$ . We attribute this surprising result to the fact that the onset of localized, toroidal ITG modes (Zocco *et al.* 2018) appears to have been increased by the combined effect of bad curvature and parallel

stabilization by FLR damping, pushing it beyond the critical gradient  $a/L_T$ , crit = 1.75. It has long been suspected that these modes, rather than extended Floquet or slab-like modes, cause the worst ITG transport (Jenko & Dorland 2002; Zocco *et al.* 2022). Such an increase in the toroidal threshold could be modelled by an additional stabilization term in the expression (2.9) (Jenko *et al.* 2001) and will be explored in future work.

# 5. Balancing MHD and ITG stability

HSK is MHD unstable for reasonably finite values of  $\beta$  so the comparison to a reactor with finite plasma pressure is theoretical. However, it is revealing to see the extent to which ITG turbulent losses can be quenched by abandoning MHD stability in the optimization. This trade-off was hinted at in previous reactor studies carried out for QH stellarators (Bader et al. 2020, 2019). We can gain some intuition for the lack of MHD stability in HSK by comparing to the standard "bean-shaped" cross section in W7-X, which has an indentation on the inboard, and vertical, compressed surfaces on the outboard. The indentation aids the formation of a vacuum magnetic well by strongly reducing the volumetric expansion of the surfaces such that  $V''(\psi) < 0$  (Cooper 1992). The outboard of the bean section has a small drift curvature, a result of large values of  $g^{\psi\psi}$ as well as a small geometric curvature [see discussion of the QIPC stellarator (Subbotin et al. 2006; Beidler et al. 2011), which has a similar bean-shaped section to that of W7-X. Quantitatively, we also know from near-axis theory that the magnitude of the drift curvature contributes to the formation of an unstable magnetic hill (Landreman & Jorge 2020), and that the expression for Mercier stability (governing large-n ballooning modes near rational surfaces) in general contains destabilizing terms proportional to  $1/|\nabla\psi|$ (Landreman & Jorge 2020). Thus the bean cross-section, possessing large values of  $|\nabla \psi|$ , a small geometric curvature, and  $V''(\psi) < 0$ , is generally stable to MHD modes.

In contrast, HSK lacks an indentation and has expanded surfaces on the outboard side, leading to a magnetic hill and significant instability with respect to the Mercier criterion. However, it is plausible that stability of a configuration like HSK could be enhanced through the addition of an indentation and a weakening of the drift curvature on the outboard side. Optimization studies exploring this compromise between MHD and ITG stability are currently underway.

#### 5.1. Discussion

Our optimization strategy for stellarators seems to have simple geometric consequences. The near-ubiquitous "bean-shape" cross section, which is thought to impart significant MHD stability and is often the site of the most detrimental ITG turbulence, can be modified to acquire a more triangular shape with a point at the outboard midplane via increasing the gradient of the binormal coordinate,  $|\nabla \alpha|$ . The resulting increase in "bad" curvature leads to improved linear ITG mode critical gradients, while heat transport at gradients near this threshold is significantly reduced. The feared trade-off between large critical gradients and stiffness of the transport above those thresholds appears not to be a true impediment, as also hinted at by the compact W7-K configuration (Roberg-Clark et al. 2022). Rather than reducing the magnitudes of both drift curvature (eqn. 2.6) and flux surface compression (Mynick et al. 2010; Xanthopoulos et al. 2014), one can increase drift curvature and local shear while improving both the ITG critical gradient and the stiffness of near-marginal transport. The reason appears to be that the critical gradient of localized, toroidal ITG modes is also increased, shielding the configuration from the most detrimental transport losses.

We note that the current work has not taken micro-turbulence effects into account

such as kinetic electron physics of ITG modes (Helander et al. 2015; Proll et al. 2022), non-zero density gradients (Thienpondt et al. 2022), trapped-electron mode turbulence (Faber et al. 2015; Mackenbach et al. 2022), or finite beta effects, which are generally stabilizing (Pueschel et al. 2008; Zocco et al. 2015). However, our goal here is to study a worst-case scenario for ITG modes and how it can be improved from purely geometric considerations, while also ensuring good quasisymmetry. Our optimization strategy and gyrokinetic analysis are local, specific to certain flux tubes, and future work will address ITG stability on the entire surface, taking more locations into account to ensure the optimization succeeds globally.

The most salient compromise to emerge from this work is that between ITG and MHD stability, and future designs will likely have to prioritize which type of stability is most important. We note that a similar optimization trade-off appears to exist between quasisymmetry and MHD stability measures such as the vacuum magnetic well (Landreman & Paul 2022; Landreman 2022). There are promising signs that Mercier stability does not have to be rigidly adhered to, e.g. in global gyrokinetic simulations of kinetic ballooning modes (Mishchenko et al. 2022), in order for heat fluxes at finite beta to match those of traditional MHD-optimized stellarators such as W7-X. We plan to study the stability of HSK to kinetic ballooning modes using this approach to see how detrimental electromagnetic turbulence can be when Mercier stability is strongly violated. Furthermore, there is experimental evidence from the LHD heliotron (Fujiwara et al. 2001) and stellarators such as TJ-II (De Aguilera et al. 2015) and W7-AS (Geiger et al. 2004; Weller et al. 2006) that Mercier-unstable and/or magnetic hill configurations can operate at relatively large  $\beta$  values with no serious loss of confinement, with the caveat that the LHD results may be restricted to low density operational regimes. A wide range of stellarator configurations, straddling the line between rigorous MHD stability and strongly suppressed ITG turbulence, thus appears realizable.

Acknowledgments. The authors thank C. Nührenberg for checking MHD stability of the HSK configuration, R. Jorge for providing the NEAT code to track alpha particle losses, and M. Landreman and B. Medasani for help with the use of SIMSOPT. We thank P. Helander, A. Zocco, and M. Landreman for contributing useful ideas to this work. This research was supported by a grant from the Simons Foundation (No. 560651, G. T. R.-C.). Computing resources at the Cobra cluster at IPP Garching and the Marconi Cluster were used to perform the simulations. This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (Grant Agreement No 101052200 — EUROfusion). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.

#### REFERENCES

Albert, Christopher G., Kasilov, Sergei V. & Kernbichler, Winfried 2020 Symplectic integration with non-canonical quadrature for guiding-center orbits in magnetic confinement devices. *Journal of Computational Physics* 403, 109065, arXiv: 1903.06885.

Angelino, P., Garbet, X., Villard, L., Bottino, A., Jolliet, S., Ghendrih, Ph, Grandgirard, V., McMillan, B. F., Sarazin, Y., Dif-Pradalier, G. & Tran, T. M. 2009 Role of plasma elongation on turbulent transport in magnetically confined plasmas. *Physical Review Letters* **102** (19), 1–4.

Bañón Navarro, A., Di Siena, A., Velasco, J. L., Wilms, F., Merlo, G., Windisch,

- T., LoDestro, L. L., Parker, J. B. & Jenko, F. 2022 First-principles based plasma profile predictions for optimized stellarators pp. 1–6, arXiv: 2210.01667.
- Bader, Aaron, Drevlak, M., Anderson, D. T., Faber, B. J., Hegna, C. C., Likin, K. M., Schmitt, J. C. & Talmadge, J. N. 2019 Stellarator equilibria with reactor relevant energetic particle losses. *Journal of Plasma Physics* 85 (5), 1–18.
- Bader, A., Faber, B. J., Schmitt, J. C., Anderson, D. T., Drevlak, M., Duff, J. M., Frerichs, H., Hegna, C. C., Kruger, T. G., Landreman, M., McKinney, I. J., Singh, L., Schroeder, J. M., Terry, P. W. & Ware, A. S. 2020 Advancing the physics basis for quasi-helically symmetric stellarators. *Journal of Plasma Physics* 1, 1–24.
- Bähner, J.-P., Alcusón, J.A., Hansen, S.K., von Stechow, A., Grulke, O., Windisch, T., Smith, H.M., Huang, Z., Edlund, E.M., Porkolab, M., Beurskens, M.N.A., Bozhenkov, S.A., Ford, O.P., Vanó, L., Langenberg, A., Pablant, N., Plunk, G.G., Bañón Navarro, A. & Jenko, F. 2021 Phase contrast imaging measurements and numerical simulations of turbulent density fluctuations in gas-fuelled ECRH discharges in Wendelstein 7-X. Journal of Plasma Physics 87 (3).
- Baumgaertel, J. A., Hammett, G. W. & Mikkelsen, D. R. 2013 Comparing linear ion-temperature-gradient-driven mode stability of the National Compact Stellarator Experiment and a shaped tokamak. *Physics of Plasmas* **20** (2).
- Beidler, C. D., Allmaier, K., Isaev, M. Yu, Kasilov, S. V., Kernbichler, W., Leitold, G. O., Maassberg, H., Mikkelsen, D. R., Murakami, S., Schmidt, M., Spong, D. A., Tribaldos, V. & Wakasa, A. 2011 Benchmarking of the mono-energetic transport coefficients Results from the International Collaboration on Neoclassical Transport in Stellarators (ICNTS). Nuclear Fusion 51 (7).
- Beurskens, M. N.A., Angioni, C., Bozhenkov, S. A., Ford, O., Kiefer, C., Xanthopoulos, P., Turkin, Y., Alcusón, J. A., Baehner, J. P., Beidler, C., Birkenmeier, G., Fable, E., Fuchert, G., Geiger, B., Grulke, O., Hirsch, M., Jakubowski, M., Laqua, H. P., Langenberg, A., Lazerson, S., Pablant, N., Reisner, M., Schneider, P., Scott, E. R., Stange, T., Von Stechow, A., Stober, J., Stroth, U., Wegner, Th, Weir, G., Zhang, D., Zocco, A., Wolf, R. C. & Zohm, H. 2022 Confinement in electron heated plasmas in Wendelstein 7-X and ASDEX Upgrade; The necessity to control turbulent transport. *Nuclear Fusion* 62 (1).
- Beurskens, M. N.A., Bozhenkov, S. A., Ford, O., Xanthopoulos, P., Zocco, A., Turkin, Y., Alonso, A., Beidler, C., Caivo, I., Carralero, D., Estrada, T., Fuchert, G., Grulke, O., Hirsch, M., Ida, K., Jakubowski, M., Killer, C., Krychowiak, M., Kwak, S., Lazerson, S., Langenberg, A., Lunsford, R., Pablant, N., Pasch, E., Pavone, A., Reimold, F., Romba, Th, Von Stechow, A., Smith, H. M., Windisch, T., Yoshinuma, M., Zhang, D. & Wolf, R. C. 2021 Ion temperature clamping in Wendelstein 7-X electron cyclotron heated plasmas. *Nuclear Fusion* 61 (11), 116072.
- Bhattacharjee, A., Sedlak, J. E., Similon, P. L., Rosenbluth, M. N. & Ross, D. W. 1983 Drift waves in a straight stellarator. *Physics of Fluids* **26** (4), 880–882.
- BIGLARI, H., DIAMOND, P. H. & ROSENBLUTH, M. N. 1989 Toroidal ion-pressure-gradient-driven drift instabilities and transport revisited. *Physics of Fluids B* 1 (1), 109–118.
- Bozhenkov, S. A., Kazakov, Y., Ford, O. P., Beurskens, M. N.A., Alcusón, J., Alonso, J. A., Baldzuhn, J., Brandt, C., Brunner, K. J., Damm, H., Fuchert, G., Geiger, J., Grulke, O., Hirsch, M., Höfel, U., Huang, Z., Knauer, J., Krychowiak, M., Langenberg, A., Laqua, H. P., Lazerson, S., Marushchenko, N. B., Moseev, D., Otte, M., Pablant, N., Pasch, E., Pavone, A., Proll, J. H.E., Rahbarnia, K., Scott, E. R., Smith, H. M., Stange, T., Von Stechow, A., Thomsen, H., Turkin, Yu, Wurden, G., Xanthopoulos, P., Zhang, D. & Wolf, R. C. 2020 High-performance plasmas after pellet injections in Wendelstein 7-X. Nuclear Fusion 60 (6), 066011.
- Brizard, A. J. & Hahm, T. S. 2007 Foundations of nonlinear gyrokinetic theory. *Reviews of Modern Physics* **79** (2), 421–468.
- Carralero, D., Estrada, T., Maragkoudakis, E., Windisch, T., Alonso, J. A., Beurskens, M., Bozhenkov, S., Calvo, I., Damm, H., Ford, O., Fuchert, G., García-Regaña, J. M., Pablant, N., Sánchez, E., Pasch, E. & Velasco, J. L.

- 2021 An experimental characterization of core turbulence regimes in Wendelstein 7-X. *Nuclear Fusion* **61** (9), arXiv: 2105.05107.
- Connor, J. W., Hastie, R. J. & Taylor, J. B. 1980 Stability of general plasma equilibria, III. *Plasma Physics* **22** (7), 757–769.
- Cooper, A. 1992 Variational formulation of the linear MHD stability of 3D plasmas with noninteracting hot electrons. *Plasma Physics and Controlled Fusion* **34** (6), 1011–1036.
- DE AGUILERA, ADRIANA M., CASTEJÓN, FRANCISCO, ASCASÍBAR, ENRIQUE, BLANCO, EMILIO, DE LA CAL, EDUARDO, HIDALGO, CARLOS, LIU, BING, LÓPEZ-FRAGUAS, ANTONIO, MEDINA, FRANCISCO, OCHANDO, MARÍA ANTONIA, PASTOR, IGNACIO, PEDROSA, MARÍA ÁNGELES, VAN MILLIGEN, BOUDEWIJN & VELASCO, JOSÉ LUIS 2015 Magnetic well scan and confinement in the TJ-II stellarator. *Nuclear Fusion* **55** (11).
- Dewar, R. L. & Glasser, A. H. 1983 Ballooning mode spectrum in general toroidal systems. *Physics of Fluids* **26** (10), 3038–3052.
- Faber, B. J., Pueschel, M. J., Terry, P. W., Hegna, C. C. & Roman, J. E. 2018 Stellarator microinstabilities and turbulence at low magnetic shear. *Journal of Plasma Physics* 84 (5), 1–28.
- Faber, B. J., Pueschel, M. J., Proll, J. H.E., Xanthopoulos, P., Terry, P. W., Hegna, C. C., Weir, G. M., Likin, K. M. & Talmadge, J. N. 2015 Gyrokinetic studies of trapped electron mode turbulence in the Helically Symmetric experiment stellarator. *Physics of Plasmas* 22 (7).
- Fujiwara, M, Kawahata, K, Ohyabu, N, Kaneko, O, Komori, A & Yamada, H 2001 Overview of LHD experiments. *Nuclear Fusion* 41, 1355.
- Geiger, Joachim E., Weller, Arthur, Zarnstorff, Michael C., Nührenberg, Carolin, Werner, Andreas Horst Franz & Kolesnichenko, Yaroslav I. 2004 Equilibrium and stability of high- $\beta$  plasmas in Wendelstein 7-AS. Fusion Science and Technology 46 (1), 13–23.
- Hegna, C. C., Anderson, D. T., Bader, A., Bechtel, T. A., Bhattacharjee, A., Cole, M., Drevlak, M., Duff, J. M., Faber, B. J., Hudson, S. R., Kotschenreuther, M., Kruger, T. G., Landreman, M., McKinney, I. J., Paul, E., Pueschel, M. J., Schmitt, J. S., Terry, P. W., Ware, A. S., Zarnstorff, M. & Zhu, C. 2022 Improving the stellarator through advances in plasma theory. *Nuclear Fusion* 62 (4).
- HEGNA, C. C., TERRY, P. W. & FABER, B. J. 2018 Theory of ITG turbulent saturation in stellarators: Identifying mechanisms to reduce turbulent transport. *Physics of Plasmas* **25** (2).
- Helander, Per 2014 Theory of plasma confinement in non-axisymmetric magnetic fields. Reports on Progress in Physics 77, 087001.
- Helander, P., Beidler, C. D., Bird, T. M., Drevlak, M., Feng, Y., Hatzky, R., Jenko, F., Kleiber, R., Proll, J. H.E., Turkin, Y. & Xanthopoulos, P. 2012 Stellarator and tokamak plasmas: A comparison. *Plasma Physics and Controlled Fusion* 54 (12).
- Helander, P., Bird, T., Jenko, F., Kleiber, R., Plunk, G. G., Proll, J. H.E., Riemann, J. & Xanthopoulos, P. 2015 Advances in stellarator gyrokinetics. *Nuclear Fusion* **55** (5).
- Helander, P. & Plunk, G. G. 2021 Upper Bounds on Gyrokinetic Instabilities in Magnetized Plasmas. *Physical Review Letters* **127** (15), 155001.
- HIRSHMAN, S. P. & WHITSON, J. C. 1983 Steepest-descent moment method for three-dimensional magnetohydrodynamic equilibria. *Physics of Fluids* **26** (12), 3553–3568.
- Jenko, F & Dorland, W 2002 Prediction of Significant Tokamak Turbulence at Electron Gyroradius Scales. *Physical Review Lett* **89** (22), 25001–1.
- Jenko, F., Dorland, W. & Hammett, G. W. 2001 Critical gradient formula for toroidal electron temperature gradient modes. *Physics of Plasmas*  $\bf 8$  (9), 4096–4104.
- Jenko, F., Dorland, W., Kotschenreuther, M. & Rogers, B. N. 2000 Electron temperature gradient driven turbulence. *Physics of Plasmas* **7** (5), 1904–1910.
- JORGE, R. & LANDREMAN, M. 2021 Ion-temperature-gradient stability near the magnetic axis of quasisymmetric stellarators, arXiv: 2102.12390.
- Landreman, Matt 2022 Mapping the space of quasisymmetric stellarators using optimized near-axis expansion , arXiv: 2209.11849.

- Landreman, Matt & Jorge, Rogerio 2020 Magnetic well and Mercier stability of stellarators near the magnetic axis. *Journal of Plasma Physics*, arXiv: 2006.14881.
- Landreman, Matt, Medasani, Bharat, Wechsung, Florian, Giuliani, Andrew, Jorge, Rogerio & Zhu, Caoxiang 2021 SIMSOPT: A flexible framework for stellarator optimization. *Journal of Open Source Software* 6 (65), 3525.
- LANDREMAN, MATT & PAUL, ELIZABETH 2022 Magnetic Fields with Precise Quasisymmetry for Plasma Confinement. *Physical Review Letters* 128 (3), 35001.
- MACKENBACH, R. J.J., PROLL, J. H.E. & HELANDER, P. 2022 Available Energy of Trapped Electrons and Its Relation to Turbulent Transport. *Physical Review Letters* **128** (17), 175001, arXiv: 2109.01042.
- Mau, T. K., Kaiser, T. B., Grossman, A. A., Raffray, A. R., Wang, X. R., Lyon, J. F., Maingi, R., Ku, L. P. & Zarnstorff, M. C. 2008 Divertor configuration and heat load studies for the ARIES-CS fusion power plant. *Fusion Science and Technology* **54** (3), 771–786.
- MCKINNEY, I J, PUESCHEL, M J, FABER, B J & HEGNA, C C 2019 A comparison of turbulent transport in a quasi-helical and a quasi-axisymmetric stellarator. *Journal of Plasma Physics* 85, 905850503.
- MISHCHENKO, ALEXEY, BORCHARDT, MATTHIAS, HATZKY, ROMAN, KLEIBER, RALF, KÖNIES, AXEL, NÜHRENBERG, CAROLIN, XANTHOPOULOS, PAVLOS, ROBERG-CLARK, GARETH & PLUNK, GABRIEL 2022 Global gyrokinetic simulations of electromagnetic turbulence in stellarator plasmas. *Journal of Plasma Physics (Submitted)*.
- Mynick, H. E., Pomphrey, N. & Xanthopoulos, P. 2010 Optimizing stellarators for turbulent transport. *Physical Review Letters* **105** (9), 1–4.
- Nemov, V. V., Kasilov, S. V., Kernbichler, W. & Heyn, M. F. 1999 Evaluation of 1/v neoclassical transport in stellarators. *Physics of Plasmas* 6 (12), 4622–4632.
- Nunami, M., Watanabe, T. H. & Sugama, H. 2013 A reduced model for ion temperature gradient turbulent transport in helical plasmas. *Physics of Plasmas* **20** (9).
- Pablant, N., Langenberg, A., Alonso, A., Baldzuhn, J., Beidler, C. D., Bozhenkov, S., Burhenn, R., Brunner, K. J., Dinklage, A., Fuchert, G., Ford, O., Gates, D. A., Geiger, J., Hirsch, M., Höfel, U., Kazakov, Ye O., Knauer, J., Krychowiak, M., Laqua, H., Landreman, M., Lazerson, S., Maassberg, H., Marchuck, O., Mollen, A., Pasch, E., Pavone, A., Satake, S., Schröder, T., Smith, H. M., Svensson, J., Traverso, P., Turkin, Y., Velasco, J. L., Von Stechow, A., Warmer, F., Weir, G., Wolf, R. C. & Zhang, D. 2020 Investigation of the neoclassical ambipolar electric field in ion-root plasmas on W7-X. *Nuclear Fusion* 60 (3).
- Plunk, G. G. & Helander, P. 2022 Energetic bounds on gyrokinetic instabilities. Part 2. Modes of optimal growth. *Journal of Plasma Physics* 88 (3), arXiv: 2201.08707.
- Plunk, G. G., Helander, P., Xanthopoulos, P. & Connor, J. W. 2014 Collisionless microinstabilities in stellarators. III. the ion-temperature-gradient mode. *Physics of Plasmas* 21 (3).
- Plunk, G. G., Xanthopoulos, P. & Helander, P. 2017 Distinct Turbulence Saturation Regimes in Stellarators. *Physical Review Letters* **118** (10), 1–5, arXiv: 1703.03257.
- Plunk, G. G., Xanthopoulos, P., Weir, G. M., Bozhenkov, S. A., Dinklage, A., Fuchert, G., Geiger, J., Hirsch, M., Hoefel, U., Jakubowski, M., Langenberg, A., Pablant, N., Pasch, E., Stange, T. & Zhang, D. 2019 Stellarators Resist Turbulent Transport on the Electron Larmor Scale. *Physical Review Letters* **122** (3), 35002.
- Proll, J.H.E., Plunk, G.G., Faber, B.J., Görler, T., Helander, P., McKinney, I.J., Pueschel, M.J., Smith, H.M. & Xanthopoulos, P. 2022 Turbulence mitigation in maximum-J stellarators with electron-density gradient. *Journal of Plasma Physics* 88 (1).
- Proll, J. H.E., Helander, P., Connor, J. W. & Plunk, G. G. 2012 Resilience of quasiisodynamic stellarators against trapped-particle instabilities. *Physical Review Letters* 108 (24), 7–10.
- PUESCHEL, M. J., FABER, B. J., CITRIN, J., HEGNA, C. C., TERRY, P. W. & HATCH, D. R. 2016 Stellarator Turbulence: Subdominant Eigenmodes and Quasilinear Modeling. Physical Review Letters 116 (8), 1–5.

- Pueschel, M. J., Kammerer, M. & Jenko, F. 2008 Gyrokinetic turbulence simulations at high plasma beta. *Physics of Plasmas* **15**, 102310.
- Roberg-Clark, G. T., Plunk, G. G. & Xanthopoulos, P. 2021 Calculating the linear critical gradient for the ion-temperature-gradient mode in magnetically confined plasmas. *Journal of Plasma Physics* 87, 905870306.
- ROBERG-CLARK, G. T., PLUNK, G. G. & XANTHOPOULOS, P. 2022 Coarse-grained gyrokinetics for the critical ion temperature gradient in stellarators. *Physical Review Research* 4 (3), L032028.
- ROBERTS, K. V. & TAYLOR, J. B. 1965 Gravitational resistive instability of an incompressible plasma in a sheared magnetic field. *Physics of Fluids* 8 (2), 315–322.
- ROMANELLI, F. 1989 Ion temperature-gradient-driven modes and anomalous ion transport in tokamaks. *Physics of Fluids B* **1** (5), 1018–1025.
- Stroteich, Sven, Xanthopoulos, Pavlos, Plunk, Gabriel & Schneider, Ralf 2022 Seeking turbulence optimized configurations for the Wendelstein 7-X stellarator: ion temperature gradient and electron temperature gradient turbulence. *Journal of Plasma Physics* 88, 175880501.
- Subbotin, A. A., Mikhailov, M. I., Shafranov, V. D., Isaev, M. Yu, Nührenberg, C., Nührenberg, J., Zille, R., Nemov, V. V., Kasilov, S. V., Kalyuzhnyj, V. N. & Cooper, W. A. 2006 Integrated physics optimization of a quasi-isodynamic stellarator with poloidally closed contours of the magnetic field strength. *Nuclear Fusion* 46 (11), 921–927.
- Sugama, H. 1999 Damping of toroidal ion temperature gradient modes. *Physics of Plasmas* **6** (9), 3527–3535.
- Talmadge, J. N., Anderson, F. S. B., Anderson, D. T., Deng, C., Guttenfelder, W., Likin, K. M., Lore, J., Schmitt, J. C. & Zhai, K. 2008 Experimental Tests of Quasisymmetry in HSX. *Plasma and Fusion Research* 3, S1002–S1002.
- Terry, P., Anderson, W. & Horton, W. 1982 Kinetic effects on the toroidal ion pressure gradient drift mode. *Nuclear Fusion* 22 (4), 487–497.
- Terry, P. W., Baver, D. A. & Gupta, Sangeeta 2006 Role of stable eigenmodes in saturated local plasma turbulence. *Physics of Plasmas* 13 (2).
- Thienpondt, H., García-Regaña, J. M., Calvo, I., Alonso, J. A., Velasco, J. L., González-Jerez, A., Barnes, M., Brunner, K., Ford, O., Fuchert, G., Knauer, J., Pasch, E., Vanó, L. & Team, the Wendelstein 7-X 2022 Turbulence prevents core particle depletion in stellarators, arXiv: 2209.04194.
- Waltz, R. E. & Boozer, A. H. 1993 Local shear in general magnetic stellarator geometry. *Physics of Fluids B* **5** (7), 2201–2205.
- Weller, A., Sakakibara, S., Watanabe, K. Y., Toi, K., Geiger, J., Zarnstorff, M. C., Hudson, S. R., Reiman, A., Werner, A., Nührenberg, C., Ohdachi, S., Suzuki, Y. & Yamada, H. 2006 Significance of MHD effects in stellarator confinement. Fusion Science and Technology **50** (2), 158–170.
- Xanthopoulos, P., Mynick, H. E., Helander, P., Turkin, Y., Plunk, G. G., Jenko, F., Görler, T., Told, D., Bird, T. & Proll, J. H.E. 2014 Controlling turbulence in present and future stellarators. *Physical Review Letters* 113 (15), 1–4.
- ZOCCO, A., HELANDER, P. & CONNOR, J. W. 2015 Magnetic compressibility and ion-temperature-gradient-driven microinstabilities in magnetically confined plasmas. Plasma Physics and Controlled Fusion 57 (8).
- ZOCCO, ALESSANDRO, PODAVINI, LINDA, GARCÌA-REGAÑA, JOSÉ MANUEL, BARNES, MICHAEL, PARRA, FELIX I., MISHCHENKO, A. & HELANDER, PER 2022 Gyrokinetic electrostatic turbulence close to marginality in the Wendelstein 7-X stellarator. *Physical Review E* 106 (1), 1–6.
- ZOCCO, A., XANTHOPOULOS, P., DOERK, H., CONNOR, J. W. & HELANDER, P. 2018 Threshold for the destabilisation of the ion-temperature-gradient mode in magnetically confined toroidal plasmas. *Journal of Plasma Physics* 84 (1).