

On Modular Gray Map

Manish K Gupta *

Dhirubhai Ambani Institute of Information and Communication Technology,

Post Bag Number 4 , Near Indroda Circle

Gandhinagar, 382007, India

Email:mankg@guptalab.org

October 31, 2022

Abstract

This paper introduces an isometry between the modular rings \mathbb{Z}_{2^s} and $\mathbb{Z}_{2^{s-1}}$ with respect to the homogeneous weights. Certain product of these maps gives Carlet's generalised Gray map and also Vega's Gray map. For $s = 2$ this reduces to popular Gray map. Several interesting properties of these maps are studied. Towards the end we list several interesting problems to work on.

Keywords: Codes over rings, Gray isometry, Gray map.

*A part of this work was presented at the Special Session on Algebraic Coding Theory, Central Section Meeting #995, American Mathematical Society, Athens,OH, 26-27 March 2004.

1 Introduction

In 19th century a cyclic-permuted binary code was invented by a french engineer *Emile Baudot* (now often called *Gray code* as it was patented by *Frank Gray* on 17.03.1953 [13]). This has represented a major advancement in telegraphy. After that it has witnessed various practical applications other than solving puzzles such as Tower of Hanoi and the Brain [29]. It is very recent when it is used as an isometry between \mathbb{Z}_4 and \mathbb{Z}_2^2 with respect to the Lee and Hamming distances to solve a 30 year old coding theory puzzle [17]. This isometry can be extended component-wise between \mathbb{Z}_4^n and \mathbb{Z}_2^{2n} and is usually known as the *Gray map*. Due to an isometry the Gray map has been used extensively to construct good quality binary linear and nonlinear codes [28, 15, 17]. Also the \mathbb{Z}_4 -linearity of various binary codes of even length has been investigated. In [22], Ana *Sălăgean-Mandache* has shown that except for the well-known case $p = s = 2$, it is not possible to construct a weight function on \mathbb{Z}_{p^s} for which \mathbb{Z}_{p^s} is isometric to \mathbb{Z}_p^s with the Hamming metric. A similar result for the Lee metric was established in [10] by S. R. Costa et al. However there exist an isometry between $\mathbb{Z}_{2^s} \rightarrow \mathbb{Z}_2^{2^{s-1}}$ was discovered by Carlet in [8]. Using this *generalized Gray map* he extends the concept of \mathbb{Z}_4 -linear codes to \mathbb{Z}_{2^s} -linear codes. The generalization of these isometries up to finite chain rings was considered by Greferath and Schmidt, Honold and Landjev, San Ling and Blackford and Nechaev in [14, 18, 19, 20]. In [5] Asch and Tilborg has studied the isometry between \mathbb{Z}_{p^2} and \mathbb{Z}_p^p . More recently an isometry between codes over \mathbb{Z}_{2^s} and codes over \mathbb{Z}_4 has been introduced by Vega and Tapia-Recillas in [25, 26]. Further they have studied the condition under which a code over \mathbb{Z}_4 is \mathbb{Z}_{2^s} linear. In this paper we will introduce a Gray isometry between \mathbb{Z}_{2^s} and $\mathbb{Z}_{2^{s-1}}$. It is shown that the generalized Gray map introduced by Carlet is equivalent to the product of $s - 1$ such maps. It also covers the various generalizations introduced earlier. Some basic properties of these maps are studied.

2 Preliminaries And Notations

A *code* \mathcal{C} , of length n , over \mathbb{Z}_{p^s} is simply a subset of $\mathbb{Z}_{p^s}^n$. If in addition it is an additive subgroup of $\mathbb{Z}_{p^s}^n$ we call \mathcal{C} a *linear code* of length n . An element of \mathcal{C} is called a *codeword* of \mathcal{C} and a *generator matrix* of a linear code \mathcal{C} is a matrix whose rows generate \mathcal{C} . The *Hamming weight* $w_H(x)$ of a vector x in $\mathbb{Z}_{p^s}^n$ is the number of non-zero components. The *Lee weight* $w_L(x)$ of a vector $x = (x_1, x_2, \dots, x_n)$ is $\sum_{i=1}^n \min\{|x_i|, |p^s - x_i|\}$. The *Euclidean weight* $w_E(x)$ of a vector x is $\sum_{i=1}^n \min\{x_i^2, (p^s - x_i)^2\}$. The Euclidean weight is useful in connection with lattice constructions. The *Chinese Euclidean weight* $w_{CE}(x)$ of a vector $x \in \mathbb{Z}_m^n$ is $\sum_{i=1}^n \left\{ 2 - 2 \cos \left(\frac{2\pi x_i}{m} \right) \right\}$. This is useful for m -PSK coding. In [9], Constantinescu and Werner introduced the *homogeneous weight* on the residue class rings of integers which was later extended to finite Frobenius rings. The homogeneous weight on \mathbb{Z}_{p^s} is defined as

follows. For $u \neq 0, u \in \mathbb{Z}_{p^s}$

$$w_{HW}(u) = \begin{cases} p^{s-2}(p-1), u \notin p^{s-1}\mathbb{Z}_p & \text{and } w_{HW}(0) = 0. \\ p^{s-1}, u \in p^{s-1}\mathbb{Z}_p \end{cases}$$

The Hamming, Lee, Euclidean and Homogeneous distances $d_H(x, y)$, $d_L(x, y)$, $d_E(x, y)$ and $d_{HW}(x, y)$ between two vectors x and y are $w_H(x-y)$, $w_L(x-y)$, $w_E(x-y)$ and $w_{HW}(x-y)$ respectively. The minimum Hamming, Lee, Euclidean and Homogeneous weights, d_H, d_L, d_E and d_{HW} of \mathcal{C} are the smallest Hamming, Lee, Euclidean and Homogeneous weights among all non-zero codewords of \mathcal{C} , respectively.

Carlet's definition of generalized Gray map can be extended to \mathbb{Z}_{p^s} naturally. The *Generalized Gray map* $\gamma : \mathbb{Z}_{p^s}^n \rightarrow \mathbb{Z}_p^{p^{s-1}n}$ is the coordinate-wise extension of the function γ_g from \mathbb{Z}_{p^s} to $\mathbb{Z}_p^{p^{s-1}}$ defined as follows. Let $u = u_1 + pu_2 + \dots + p^{s-1}u_s \in \mathbb{Z}_{p^s}$ be the finite p -adic expansion of u , where $u_i \in \mathbb{Z}_p$ also let $y_i \in \mathbb{Z}_p$. The map $\gamma_g : (y_1, y_2, \dots, y_{s-1}) \mapsto u_s + \sum_{i=1}^{s-1} u_i y_i \in \mathbb{Z}_p$ defined as a generalized Boolean function evaluated on $GF(p^{s-1})$. It is easy to see that this map is equivalent to the generalized Gray map introduced by Greferath and Schmidt [14] using tensor product. In particular for $s = 2$ this map is also equivalent to a generalized Gray map from \mathbb{Z}_{p^2} to \mathbb{Z}_p^p introduced recently by Asch and Tilborg [5]. The image $\gamma(\mathcal{C})$, of a linear code \mathcal{C} over \mathbb{Z}_{p^s} of length n by the generalized Gray map, is a p -ary code of length $p^{s-1}n$. The *dual code* \mathcal{C}^\perp of \mathcal{C} is defined as $\{x \in \mathbb{Z}_{p^s}^n \mid x \cdot y = 0 \text{ for all } y \in \mathcal{C}\}$ where $x \cdot y$ is the standard inner product of x and y . \mathcal{C} is *self-orthogonal* if $\mathcal{C} \subseteq \mathcal{C}^\perp$ and \mathcal{C} is *self-dual* if $\mathcal{C} = \mathcal{C}^\perp$. Two codes are said to be *equivalent* if one can be obtained from the other by permuting the coordinates and (if necessary) changing the signs of certain coordinates. Codes differing by only a permutation of coordinates are called *permutation-equivalent*.

Any linear code \mathcal{C} over \mathbb{Z}_{p^s} is permutation-equivalent to a code with generator matrix G (the rows of G generate \mathcal{C}) of the form

$$(1) \quad G = \begin{bmatrix} I_{k_0} & A_{01} & A_{02} & \cdots & A_{0s-1} & A_{0s} \\ \mathbf{0} & pI_{k_1} & pA_{12} & \cdots & pA_{1s-1} & pA_{1s} \\ \mathbf{0} & \mathbf{0} & p^2I_{k_2} & \cdots & p^2A_{2s-1} & p^2A_{2s} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & p^{s-1}I_{k_{s-1}} & p^{s-1}A_{s-1s} \end{bmatrix},$$

where A_{ij} are matrices over \mathbb{Z}_{p^s} and the columns are grouped into blocks of sizes $k_0, k_1, \dots, k_{s-1}, k_s$, respectively. Let $k = \sum_{i=0}^{s-1} (s-i)k_i$. Then $|\mathcal{C}| = p^k$ is the number of codewords in the code \mathcal{C} . Note that \mathcal{C} is a free code if and only if $k_i = 0$ for all $i = 1, 2, \dots, s-1$. It is easy to see that a generator matrix for \mathcal{C}^\perp is of the form

$$H = \begin{bmatrix} B_{0s} & B_{0s-1} & \cdots & B_{01} & I_{k_s} \\ pB_{1s} & pB_{1s-1} & \cdots & pI_{k_{s-1}} & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ p^{s-1}B_{s-1s} & p^{s-1}I_{k_1} & \cdots & \mathbf{0} & \mathbf{0} \end{bmatrix},$$

where B_{ij} are matrices over \mathbb{Z}_{p^s} and the columns are grouped into blocks of size k_0, k_1, \dots, k_s . Therefore $|\mathcal{C}^\perp| = p^{sn-k}$. Note that $sn - k = \sum_{i=1}^s ik_i$.

A vector \mathbf{v} is a p -linear combination of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ if $\mathbf{v} = l_1\mathbf{v}_1 + \dots + l_k\mathbf{v}_k$ with $l_i \in \mathbb{Z}_p$ for $1 \leq i \leq k$. A subset $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ of \mathcal{C} is called a p -basis for \mathcal{C} if for each $i = 1, 2, \dots, k-1$, $p\mathbf{v}_i$ is a p -linear combination of $\mathbf{v}_{i+1}, \dots, \mathbf{v}_k$, $2\mathbf{v}_k = 0$, \mathcal{C} is the p -linear span of S and S is p -linearly independent [15]. The number of elements in a p -basis for \mathcal{C} is called the p -dimension of \mathcal{C} . It is easy to verify that the rows of the matrix

$$(2) \quad \mathcal{B} = \begin{bmatrix} I_{k_0} & A_{01} & A_{02} & \cdots & A_{0s-1} & A_{0s} \\ pI_{k_0} & pA_{01} & pA_{02} & \cdots & pA_{0s-1} & pA_{0s} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ p^{s-1}I_{k_0} & p^{s-1}A_{01} & p^{s-1}A_{02} & \cdots & p^{s-1}A_{0s-1} & p^{s-1}A_{0s} \\ \hline \mathbf{0} & pI_{k_1} & pA_{12} & \cdots & pA_{1s-1} & pA_{1s} \\ \mathbf{0} & p^2I_{k_1} & p^2A_{12} & \cdots & p^2A_{1s-1} & p^2A_{1s} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \mathbf{0} & p^{s-1}I_{k_1} & p^{s-1}A_{12} & \cdots & p^{s-1}A_{1s-1} & p^{s-1}A_{1s} \\ \hline \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \hline \mathbf{0} & \mathbf{0} & \mathbf{0} & \cdots & p^{s-1}I_{k_{s-1}} & p^{s-1}A_{s-1s} \end{bmatrix}.$$

form a p -basis for the code \mathcal{C} generated by G given in (1).

A linear code \mathcal{C} over \mathbb{Z}_{p^s} (over \mathbb{Z}_2) of length n , p -dimension k , minimum distance d_H, d_L, d_E, d_{CE} and d_{HW} is called an $[n, k, d_H, d_L, d_E, d_{CE}, d_{HW}]$ ($[n, k, d_H]$) or simply an $[n, k]$ code.

3 Modular Gray Map

Now we define a natural Gray map as follows.

Let

$$\begin{aligned} A_1 &= \{0, 1, 2, 3, \dots, 2^{s-2} - 1\}, \\ A_2 &= \{2^{s-2}, 2^{s-2} + 1, \dots, 2^{s-1} - 1\}, \\ A_3 &= \{2^{s-1}, 2^{s-1} + 1, \dots, 2^{s-1} + 2^{s-2} - 1\} \text{ and} \\ A_4 &= \{2^{s-1} + 2^{s-2}, 2^{s-1} + 2^{s-2} + 1, \dots, 2^{s-1} + 2^{s-1} - 1\} \end{aligned}$$

be the four consecutive disjoint partitions of the integers of \mathbb{Z}_{2^s} . Define the modular generalized Gray map η_g^s on \mathbb{Z}_{2^s} to $\mathbb{Z}_{2^{s-1}}$ as

$$\eta_g^s(i) = \begin{cases} (i, i) & i \in A_1 \\ (i - 2^{s-2}, i) & i \in A_2 \\ (i - 2^{s-2}, i - 2^{s-2}) & i \in A_3 \\ (i - 2^{s-1}, i - 3 \cdot 2^{s-2}) & i \in A_4 \end{cases}$$

Remark 1 Each entry of \mathbb{Z}_{2^s} is mapped onto the first order Reed Muller code of length 2 over \mathbb{Z}_2 [15, 16] generated by the following generator matrix

$$G = \begin{bmatrix} 1 & 1 \\ 0 & 2^{s-2} \end{bmatrix}_{2 \times 2}.$$

Remark 2 For $s = 2$ η_g^s reduces to ordinary Gray map.

Remark 3 Applying η_g^s to a code \mathcal{C} over \mathbb{Z}_{2^s} gives a code $\eta_g^s(\mathcal{C})$ over $\mathbb{Z}_{2^{s-1}}$ of length twice the length of \mathcal{C}

Example 1 For $s = 2, 3, 4$ these maps are given below.

$$\begin{aligned} \mathbb{Z}_4 &\mapsto \mathbb{Z}_2 \\ 0 &\mapsto 00 \\ 1 &\mapsto 01 \\ 2 &\mapsto 11 \\ 3 &\mapsto 10 \end{aligned}$$

$$\begin{aligned} \mathbb{Z}_8 &\mapsto \mathbb{Z}_4 \\ 0 &\mapsto 00 \\ 1 &\mapsto 11 \\ 2 &\mapsto 02 \\ 3 &\mapsto 13 \\ 4 &\mapsto 22 \\ 5 &\mapsto 33 \\ 6 &\mapsto 20 \\ 7 &\mapsto 31 \end{aligned}$$

\mathbb{Z}_{16}	\mapsto	\mathbb{Z}_8
0	\mapsto	00
1	\mapsto	11
2	\mapsto	22
3	\mapsto	33
4	\mapsto	04
5	\mapsto	15
6	\mapsto	26
7	\mapsto	37
8	\mapsto	44
9	\mapsto	55
10	\mapsto	66
11	\mapsto	77
12	\mapsto	40
13	\mapsto	51
14	\mapsto	62
15	\mapsto	73

Some interesting facts about modular Gray maps are collected in the following.

Theorem 1 *The generalized Gray map of Carlet $\Phi_G : \mathbb{Z}_{2^s} \mapsto \mathbb{Z}_2$ can be written as a product of $s - 1$ modular Gray maps i.e., $\Phi_G = \eta_g^s \eta_g^{s-1} \eta_g^{s-2} \dots \eta_g^2$.*

Proof. By remark 1, if we apply the modular gray map η_g^s to the 2-basis generator matrix of the first order Reed Muller code over \mathbb{Z}_{2^s} we get the 2-basis generator matrix of the first order Reed Muller code over $\mathbb{Z}_{2^{s-1}}$ which is the image set of η_g^{s-1} and so on. Thus successive applications yields Carlet's generalized Gray map [15, 16]. □

Remark 4 *It is interesting to note that now we can get modular Gray map between all intermediate rings i.e., between*

$$\mathbb{Z}_{2^s} \mapsto \mathbb{Z}_{2^{s-1}} \mapsto \mathbb{Z}_{2^{s-2}} \mapsto \dots \mapsto \mathbb{Z}_4 \mapsto \mathbb{Z}_2.$$

For example, the Gray map of Tapia-Recillas and Vega [25, 26] between \mathbb{Z}_{2^s} and \mathbb{Z}_4 is equivalent to a map by taking product of $s - 2$ such maps $\eta_g^s \eta_g^{s-1} \dots \eta_g^3$. Clearly our map is much more powerful.

Lemma 1 *The images of the rows of the matrix \mathcal{B} given by (2) under the modular Gray map η_g^s are 2-linearly independent over $\mathbb{Z}_{2^{s-1}}$.*

Proof. After taking appropriate block permutations row-wise to the matrix \mathcal{B} and applying the modular Gray map we see that in the diagonals we have 2-linearly independent vectors over $\mathbb{Z}_{2^{s-1}}$. This proves our claim. \square

Proposition 1 *The modular Gray map η_g^s is an isometry between $\mathbb{Z}_{2^s} \mapsto (\mathbb{Z}_{2^{s-1}})^2$ with respect to the homogeneous weights.*

Proof. Let $u(\neq 0) \in \mathbb{Z}_{2^s}$. Then the homogeneous weight will be given as

$$w_{HW}(u) = \begin{cases} 2^{s-2}, & u \neq 2^{s-1} \\ 2^{s-1}, & u = 2^{s-1}. \end{cases}$$

Thus homogeneous weight of each entry in the sets A_1, A_2, A_3 (except first entry which is 2^{s-1}) and A_4 will be 2^{s-2} . Thus direct computation of homogeneous weight as a rational sum of the entries in the definition yields the desired result. The case of $u = 0$ is trivial. \square

Remark 5 *Note that any product of modular Gray maps η_g^s is also an isometry.*

Remark 6 *The modular Gray map η_g^s can be extended coordinate-wise to a map from $(\mathbb{Z}_{2^s})^n$ to $(\mathbb{Z}_{2^{s-1}})^{2n}$. With abuse of notation we denote this map also by η_g^s .*

3.1 Permuted Modular Gray Map

We can permute the modular Gray map to get some very interesting results like perverseness of Octocode. The modified modular Gray map ξ_g^s is define as

....

Let us modify our examples to get the modify modular Gray map for $s = 2, 3, 4$.

$$\begin{array}{l} \mathbb{Z}_4 \mapsto \mathbb{Z}_2 \\ 0 \mapsto 00 \\ 1 \mapsto 10 \\ 2 \mapsto 11 \\ 3 \mapsto 01 \end{array}$$

$$\begin{array}{l}
\mathbb{Z}_8 \mapsto \mathbb{Z}_4 \\
0 \mapsto 00 \\
1 \mapsto 11 \\
2 \mapsto 20 \\
3 \mapsto 13 \\
4 \mapsto 22 \\
5 \mapsto 31 \\
6 \mapsto 02 \\
7 \mapsto 33
\end{array}$$

$$\begin{array}{l}
\mathbb{Z}_{16} \mapsto \mathbb{Z}_8 \\
0 \mapsto 00 \\
1 \mapsto 11 \\
2 \mapsto 22 \\
3 \mapsto 13 \\
4 \mapsto 40 \\
5 \mapsto 15 \\
6 \mapsto 26 \\
7 \mapsto 17 \\
8 \mapsto 44 \\
9 \mapsto 71 \\
10 \mapsto 62 \\
11 \mapsto 73 \\
12 \mapsto 04 \\
13 \mapsto 75 \\
14 \mapsto 66 \\
15 \mapsto 77
\end{array}$$

Example 2 Consider the generator matrix of the Octocode over \mathbb{Z}_8 .

$$G_8 = \begin{bmatrix} 5 & 7 & 5 & 6 & 1 & 0 & 0 & 0 \\ 5 & 0 & 7 & 5 & 6 & 1 & 0 & 0 \\ 5 & 0 & 0 & 7 & 5 & 6 & 1 & 0 \\ 5 & 0 & 0 & 0 & 7 & 5 & 6 & 1 \end{bmatrix}.$$

Apply the modular Gray map η_g^3 on the rows of G_8 . We get the following generator matrix

$$\eta_g^3(G_8) = \left[\begin{array}{cccccccc|cccccc}
3 & 3 & 3 & 0 & 1 & 0 & 0 & 0 & 1 & 3 & 1 & 2 & 1 & 0 & 0 & 0 \\
3 & 0 & 3 & 3 & 0 & 1 & 0 & 0 & 1 & 0 & 3 & 1 & 2 & 1 & 0 & 0 \\
3 & 0 & 0 & 3 & 3 & 0 & 1 & 0 & 1 & 0 & 0 & 3 & 1 & 2 & 1 & 0 \\
3 & 0 & 0 & 0 & 3 & 3 & 0 & 1 & 1 & 0 & 0 & 0 & 3 & 1 & 2 & 1 \end{array} \right].$$

4 Properties

In this section we list several interesting properties of modular Gray map that one can study.

- General results of Theorems 4, 5, 6, page 305 of [17] these should also generalize the results of Vega et al in [25] and Carlet's result of [8].
- Studying the characterizations of cyclic, quasi-cyclic, consta-cyclic, negacyclic, quasi-twisted codes with respect to modular Gray map in view of the following papers [31, 30, 19, 26, 27].
- Does modular Gray map preserves self orthogonality of codes ? Under what conditions ?
- Applications of both modular and permuted modular Gray map for the constructions of new codes over various modular rings and binary fields. For example to Hamming codes, Simplex codes, Golay codes and other standard codes.
- Is there any connection for the modular Gray map with the Nenchev's Gray map ?
- Other non-trivial applications like by taking the generator matrix of Kerdock codes, Preparata codes, Goethals codes or like codes and applying the modular Gray map.
- Construction of record breaking codes !
- Effect on Torsion and reduction codes !
- New code construction with respect to 2-basis modular Gray map applications !
- Connection with Hensal-uplift !
- Studying the \mathbb{Z}_{2^s} -linearity of standard uplifted codes over $\mathbb{Z}_{2^{s-1}}$.
- Finding other non-trivial properties !
- ...

Note This note is an old contribution and has not been updated with some new results on Gray map. We refer the reader to look into some new literature in the area.

References

- [1] T. Abualrub and R. Oehmke, *On the generators of \mathbb{Z}_4 cyclic codes of length 2^e* . IEEE Trans. Inform. Theory **49** (2003), 2126–2133.
- [2] M. C. Bhandari, M. K. Gupta and A. K. Lal, *On \mathbb{Z}_4 simplex codes and their gray images* Applied Algebra, Algebraic Algorithms and Error-Correcting Codes, AAECC-13, Lecture Notes in Computer Science **1719** (1999), 170–180.
- [3] T. Blackford, *Negacyclic codes over \mathbb{Z}_4 of even length*. IEEE Trans. Inform. Theory **49** (2003), 1417–1424.
- [4] T. Blackford and San Ling, *A transform approach to constacyclic and quasi-twisted codes* preprint(2005)
- [5] B. van Asch and Henk C.A. van Tilborg, *Two "dual" families of nearly-linear codes*. Applicable Algebra in Applied Engineering, Communication and Computing **11** (2001), 313–329.
- [6] A. Bonnetcaze, P. Solé, C. Bachoc and B. Mourrain *Type II codes over \mathbb{Z}_4* . IEEE Trans. Inform. Theory **43** (1997), 969–976.
- [7] J. Borges, C. Fernandez and K. T. Phelps, *Quaternary Reed-Muller codes* IEEE Trans. Inform. Theory **51** (2005), 2686–2691.
- [8] C. Carlet, *\mathbb{Z}_{2^k} -linear codes* IEEE Trans. Inform. Theory **44** (1998), 1543–1547.
- [9] I. Constantinescu and W. Heise, *A new metric for codes over residue class rings of integers* Probl. Pered. Inform. **33**, (1997) pp. 22–28.
- [10] Sueli I. Rodrigues Costa, João Roberto Gerônimo, Reginaldo Palazzo Jr., J. Carmelo Interlando and Marcelo Muniz Silva Alves *The symmetry group of \mathbb{Z}_q^n in the Lee space and \mathbb{Z}_{q^n} -linear code* Applied Algebra, Algebraic Algorithms and Error-Correcting Codes, AAECC-12, Lecture Notes in Computer Science **1255** (1997), 66–77.
- [11] J.H. Conway and N.J.A. Sloane, *Self-dual codes over the integers modulo 4*. J. Combin. Theory Ser. A **62** (1993), 30–45.
- [12] H. Dinh and S. R. Lopez-Permouth, *Cyclic and negacyclic codes over finite chain rings* IEEE Trans. Inform. Theory **50** (2004), 1728–1738.
- [13] F. Gray, *Pulse Code Communication, US Patent*, (March 17, 1953)
- [14] M. Greferath and S. E. Schmidt, *Gray isometries for finite chain rings and a nonlinear ternary $(36, 3^{12}, 15)$ code* IEEE Trans. Inform. Theory **45** (1999), 2522–2524.

- [15] M. K. Gupta, *On Linear Codes over \mathbb{Z}_{2^s}* **PhD Thesis**. Department of Mathematics, IIT Kanpur, India 2001.
- [16] M. K. Gupta, M. C. Bhandari and A. K. Lal, *On linear codes over \mathbb{Z}_{2^s}* Design Codes and Cryptography **36** (2005), 227–244.
- [17] A. R. Hammons, P. V. Kumar, A. R. Calderbank, N. J. A. Sloane, and P. Solé. *The \mathbb{Z}_4 -linearity of kerdock, preparata, goethals, and related codes*. IEEE Trans. Inform. Theory, **40** (1994), 301–319.
- [18] T. Honold and I. Landjev, *Linearly representable codes over chain rings*. Proc. ACCT 6 (Pskov, Russia) (1998), pp. 135–141.
- [19] S. Ling and T. Blackford, *$\mathbb{Z}_{p^{k+1}}$ -linear codes* IEEE Trans. Inform. Theory **48** (2002), 2592–2605.
- [20] A. A. Nechaev and A. S. Kuzmin, *Linearly representable codes* Proc. IEEE Int. Symp. Inform. Theory and Its Applications (1996), pp. 31–34.
- [21] E. M. Rains and N. J. A. Sloane. *Self-Dual Codes* in V. Pless and W.C. Huffman (Eds.) **The Handbook of Coding Theory**. North-Holland, New York, 1998.
- [22] A. Salagean-Mandache, *On the isometries between \mathbb{Z}_p^k and \mathbb{Z}_p^k* IEEE Trans. Inform. Theory **45** (1999), 2146–2148.
- [23] H. Tapia-Recillas, *The Gray map on $GR(p^2, n)$ and Repeated-Root Cyclic Codes*. Lecture Notes in Computer Science, **2948** (2004), 181–196.
- [24] H. Tapia-Recillas, *On the Gray image of linear cyclic codes over Galois rings* Proc. AGCT-10, CIRM, Marseille-Luminy, France, (September 2005).
- [25] H. Tapia-Recillas and G. Vega, *On \mathbb{Z}_{2^k} -linear and quaternary codes*. Siam J. Disc. Math., **17** (2003), 103–113.
- [26] H. Tapia-Recillas and G. Vega, *Some constacyclic codes over \mathbb{Z}_{2^k} -linear and binary quasi-cyclic codes*. Disc. Applied Math., **128** (2003), 305–316.
- [27] G. Vega and J. Wolfmann, *Some families of \mathbb{Z}_4 -cyclic codes*. Finite Fields and Their Applications, **10** (2004), 530–539.
- [28] Z. X. Wan, **Quaternary Codes**. World Scientific, Singapore, 256 pp, 1997.
- [29] web.usna.navy.mil/wdj/gray.htm

- [30] J. Wolfmann, *Binary Images of cyclic codes over \mathbb{Z}_4* IEEE Trans. Inform. Theory **47** (2001), 1773–1779.
- [31] J. Wolfmann, *Negacyclic and cyclic codes over \mathbb{Z}_4* IEEE Trans. Inform. Theory **47** (1999), 2527–2532.