

TFU extensions in LCA groups

Aliakbar Alijani

Abstract

Let ℓ be the category of all locally compact abelian (LCA) groups. Let $G \in \ell$ and $H \subseteq G$. The first Ulm subgroup of G is denoted by $G^{(1)}$ and the closure of H by \overline{H} . A proper short exact sequence $0 \rightarrow A \xrightarrow{\phi} B \xrightarrow{\psi} C \rightarrow 0$ in ℓ is said to be a *TFU* extension if $0 \rightarrow \overline{A^{(1)}} \xrightarrow{\overline{\phi}} \overline{B^{(1)}} \xrightarrow{\overline{\psi}} \overline{C^{(1)}} \rightarrow 0$ is a proper short exact sequence where $\overline{\phi} = \phi|_{\overline{A^{(1)}}}$ and $\overline{\psi} = \psi|_{\overline{B^{(1)}}}$. We introduce some results on *TFU* extensions. Also, we establish conditions under which the *TFU* extensions split.

1 Introduction

Throughout, all groups are Hausdorff topological abelian groups and will be written additively. Let ℓ denote the category of locally compact abelian (LCA) groups with continuous homomorphisms as morphisms. A morphism is called proper if it is open onto its image, and a short exact sequence $0 \rightarrow A \xrightarrow{\phi} B \xrightarrow{\psi} C \rightarrow 0$ in ℓ is said to be proper exact if ϕ and ψ are proper morphisms. In this case the sequence is called an extension of A by C (in ℓ). Following [5], we let $Ext(C, A)$ denote the group of extensions of A by C . The first Ulm subgroup of a group G is denoted by $G^{(1)}$. In this paper, we introduce a new concept on extensions. An extension $0 \rightarrow A \xrightarrow{\phi} B \xrightarrow{\psi} C \rightarrow 0$ in ℓ is called a *TFU* extension if $0 \rightarrow \overline{A^{(1)}} \xrightarrow{\overline{\phi}} \overline{B^{(1)}} \xrightarrow{\overline{\psi}} \overline{C^{(1)}} \rightarrow 0$ be an extension where $\overline{\phi} = \phi|_{\overline{A^{(1)}}}$ and $\overline{\psi} = \psi|_{\overline{B^{(1)}}}$. In Section 1, we show that every extension of a divisible group by an arbitrary LCA group is a *TFU* extension (Corollary 2.4). A subgroup H of a group G is said to be pure if $H \cap nG = nH$ for every positive integer n [3]. An extension $0 \rightarrow A \xrightarrow{\phi} B \xrightarrow{\psi} C \rightarrow 0$ is said to be a pure extension if $\phi(A)$ is pure in B [4]. We show that every pure extension of a group by a compact totally disconnected group is a *TFU* extension (Corollary 2.6). In Section 2, we establish some results on splitting of *TFU* extensions (see Lemma 3.1,3.2,3.3,3.5,3.6.).

The additive topological group of real numbers is denoted by \mathbb{R} , \mathbb{Q} is the group of rationals with the discrete topology and \mathbb{Z} , the group of integers. For any group G , G_0 is the identity component of G and tG , the maximal torsion subgroup of G . For groups G and H , $Hom(G, H)$ is the group of all continuous homomorphisms from G to H , endowed with the compact-open topology. The dual group of G is $\hat{G} = Hom(G, \mathbb{R}/\mathbb{Z})$ and (\hat{G}, S) denoted the annihilator of $S \subseteq G$ in \hat{G} . For more on locally compact abelian groups, see [7].

2 TFU extensions

Let $A, C \in \mathcal{L}$. In this section, we define the concept of a TFU extensions of A by C .

Definition 2.1. An extension $0 \rightarrow A \xrightarrow{\phi} B \xrightarrow{\psi} C \rightarrow 0$ in \mathcal{L} is called a TFU extension if $0 \rightarrow \overline{A^{(1)}} \xrightarrow{\overline{\phi}} \overline{B^{(1)}} \xrightarrow{\overline{\psi}} \overline{C^{(1)}} \rightarrow 0$ is an extension.

Remark 2.1. Let G_1 and G_2 be two groups. An easy calculation shows that $(G_1 \oplus G_2)^{(1)} = G_1^{(1)} \oplus G_2^{(1)}$. So, $\overline{(G_1 \oplus G_2)^{(1)}} = \overline{G_1^{(1)}} \oplus \overline{G_2^{(1)}}$.

Lemma 2.2. For groups $A, C \in \mathcal{L}$, the trivial extension $0 \rightarrow A \rightarrow A \oplus C \rightarrow C \rightarrow 0$ is a TFU extension.

Proof. It is clear by Remark 2.1. □

Lemma 2.3. Let $G \in \mathcal{L}$ and H be a closed, divisible subgroup of G . Then, $\overline{(G/H)^{(1)}} = \overline{G^{(1)}}/H$.

Proof. Let $\pi : G \rightarrow G/H$ be the natural mapping. Then, $\pi(\overline{G^{(1)}}) \subseteq \overline{\pi(G^{(1)})} = \overline{(G/H)^{(1)}}$. But, $\pi(\overline{G^{(1)}}) = \overline{(G^{(1)})}/H$. Hence, $\overline{G^{(1)}}/H \subseteq \overline{(G/H)^{(1)}}$. Now, suppose that $x + H \in \overline{(G/H)^{(1)}}$. We show that $x \in \overline{G^{(1)}}$. Let V be a neighborhood of G containing x and n , an arbitrary positive integer. Then, $y + H \in (V + H)/H \cap (G/H)^{(1)} \neq \phi$ for some $y \in G$. From $y + H \in (V + H)/H$, deduce that $y + H = z + H$ for some $z \in V$. Since H is divisible, so $z = y + nh$ for some $h \in H$. On the other hand, $y + H \in (G/H)^{(1)}$. So, $y + H = ng + H$ for some $g \in G$. Hence, $y = ng + nh_1$ for some $h_1 \in H$. Therefore, $z \in nG$. This shows that $z \in V \cap G^{(1)}$. Hence, $x \in \overline{G^{(1)}}$. □

Corollary 2.4. Every extension of a divisible group by an arbitrary LCA group is a TFU extension.

Proof. It is clear by Lemma 2.3. □

Lemma 2.5. Let $G \in \mathcal{L}$ and H a closed, pure subgroup of G such that $(G/H)^{(1)} = 0$. Then, $G^{(1)} = H^{(1)}$.

Proof. Let $x \in G^{(1)}$ and n an arbitrary positive integer. Then, $x = ng$ for some $g \in G$. So, $x + H \in n(G/H)$. Hence, $x + H \in (G/H)^{(1)} = 0$. This shows that $G^{(1)} \subseteq H$. Since H is pure, $G^{(1)} = H^{(1)}$. □

Corollary 2.6. Every pure extension of a group by a compact totally disconnected group is a TFU extension.

Proof. Let G be a compact totally disconnected group. By Lemma 2.5, it is sufficient to show that $G^{(1)} = 0$. By Theorem 24.26 of [7], \hat{G} is a torsion group. So, $G^{(1)} \subseteq (G, \hat{G}) = 0$ (see Theorem 24.22 of [7]). □

Corollary 2.7. Let $G \in \mathcal{L}$ such that nG is closed in G for all positive integers n . Then, every pure extension of a group by $G/G^{(1)}$ is a TFU extension.

Proof. Let $x + G^{(1)} \in (G/G^{(1)})^{(1)}$ and n be an arbitrary positive integer. Then, $x + G^{(1)} = ng + G^{(1)}$ for some $g \in G$. This shows that $x \in G^{(1)}$. Hence, $(G/G^{(1)})^{(1)} = 0$. \square

The dual of an extension $E : 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is defined by $\hat{E} : 0 \rightarrow \hat{C} \rightarrow \hat{B} \rightarrow \hat{A}$. The following example shows that the dual of a *TFU* extension need not to be a *TFU* extension.

Example 2.8. Consider the extension $E : 0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}_2 \rightarrow 0$. Clearly, E is a *TFU* extension. But, $\hat{E} : 0 \rightarrow \mathbb{Z}_2 \rightarrow \mathbb{R}/\mathbb{Z} \xrightarrow{\times 2} \mathbb{R}/\mathbb{Z} \rightarrow 0$ is not a *TFU* extension.

Recall that two extensions $0 \rightarrow A \xrightarrow{\phi_1} B \xrightarrow{\psi_1} C \rightarrow 0$ and $0 \rightarrow A \xrightarrow{\phi_2} X \xrightarrow{\psi_2} C \rightarrow 0$ are said to be equivalent if there is a topological isomorphism $\beta : B \rightarrow X$ such that the following diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{\phi_1} & B & \xrightarrow{\psi_1} & C & \longrightarrow & 0 \\ & & \downarrow 1_A & & \downarrow \beta & & \downarrow 1_C & & \\ 0 & \longrightarrow & A & \xrightarrow{\phi_2} & X & \xrightarrow{\psi_2} & C & \longrightarrow & 0 \end{array}$$

is commutative.

Lemma 2.9. *An extension equivalent to a TFU extension is a TFU extension.*

Proof. Let

$$\begin{aligned} E_1 : 0 &\rightarrow A \xrightarrow{\phi_1} B \xrightarrow{\psi_1} C \rightarrow 0 \\ E_2 : 0 &\rightarrow A \xrightarrow{\phi_2} X \xrightarrow{\psi_2} C \rightarrow 0 \end{aligned}$$

be two equivalent extensions such that E_1 is a *TFU* extension. Then, there is a topological isomorphism $\beta : B \rightarrow X$ such that $\beta\phi_1 = \phi_2$ and $\psi_2\beta = \psi_1$. Since $\beta(\overline{B^{(1)}}) = \overline{X^{(1)}}$, so

$$\psi_2(\overline{X^{(1)}}) = \psi_2\beta(\overline{B^{(1)}}) = \psi_1(\overline{B^{(1)}}) = \overline{C^{(1)}}$$

Hence, $\overline{\psi_2} : \overline{X^{(1)}} \rightarrow \overline{C^{(1)}}$ is surjective. Since $\phi_2 = \beta\phi_1$ and E_1 is a *TFU* extension, so

$$\psi_2\phi_2(\overline{A^{(1)}}) = \psi_2(\beta\phi_1(\overline{A^{(1)}})) = \psi_1\phi_1(\overline{A^{(1)}}) = 0$$

Hence, $Im\overline{\phi_2} \subseteq Ker\overline{\psi_2}$. Now, we show that $Ker\overline{\psi_2} \subseteq Im\overline{\phi_2}$. Let $x \in \overline{X^{(1)}}$ and $\psi_2(x) = 0$. Then, there exists $b \in \overline{B^{(1)}}$ such that $x = \beta(b)$. Since $\psi_1(b) = \psi_2\beta(b) = \psi_2(x) = 0$ and E_1 is a *TFU* extension, so $b = \phi_1(a)$ for some $a \in \overline{A^{(1)}}$. Hence,

$$\phi_2(a) = \beta\phi_1(a) = \beta(b) = x$$

and $0 \rightarrow \overline{A^{(1)}} \xrightarrow{\overline{\phi_2}} \overline{X^{(1)}} \xrightarrow{\overline{\psi_2}} \overline{C^{(1)}} \rightarrow 0$ is an exact sequence. Since

$$\overline{\psi_2} = \overline{\psi_1}(\overline{\beta})^{-1}, \overline{\phi_2} = \overline{\beta}(\overline{\phi_1})$$

$\overline{\psi_2}$ and $\overline{\phi_2}$ are open. So, E_2 is a *TFU* extension. \square

Lemma 2.10. Let $C \in \ell$ be a torsion-free group, $0 \rightarrow A \xrightarrow{\phi} B \xrightarrow{\psi} C \rightarrow 0$ be a TFU extension and assume

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{\mu} & X & \xrightarrow{\nu} & Y \longrightarrow 0 \\ & & \downarrow 1_A & & \downarrow \theta & & \downarrow f \\ 0 & \longrightarrow & A & \xrightarrow{\phi} & B & \xrightarrow{\psi} & C \longrightarrow 0 \end{array}$$

is the standard pullback diagram in \mathcal{L} (See [5]) such that $\bar{f} : \overline{Y^{(1)}} \rightarrow \overline{C^{(1)}}$ be a proper morphism. Then

$$0 \rightarrow A \xrightarrow{\mu} X \xrightarrow{\nu} Y \rightarrow 0$$

is a TFU extension.

Proof. We have

$$X = \{(y, b) \in Y \oplus B : f(y) = \psi(b)\}.$$

and

$$\mu : a \mapsto (0, \phi(a)), \nu : (y, b) \mapsto y, \theta : (y, b) \mapsto b.$$

Consider the following standard pullback diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \overline{A^{(1)}} & \xrightarrow{\phi'} & N & \xrightarrow{\psi'} & \overline{Y^{(1)}} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow f \\ 0 & \longrightarrow & \overline{A^{(1)}} & \xrightarrow{\bar{\phi}} & \overline{B^{(1)}} & \xrightarrow{\bar{\psi}} & \overline{C^{(1)}} \longrightarrow 0 \end{array}$$

where $N = \{(y, b) \in \overline{Y^{(1)}} \oplus \overline{B^{(1)}} : f(y) = \psi(b)\}$. First, we show that $X^{(1)} = \{(y, b) \in Y^{(1)} \oplus B^{(1)} : f(y) = \psi(b)\}$. Let $(y, b) \in X^{(1)}$ and n an arbitrary positive integer. Then, $y = ny_1$ and $b = nb_1$ for some $y_1 \in Y$, $b_1 \in B$. Also, $f(y_1) = \psi(b_1)$. We have

$$f(y) = f(ny_1) = \psi(nb_1) = \psi(b)$$

Since n be arbitrary, $(y, b) \in \{(y, b) \in Y^{(1)} \oplus B^{(1)} : f(y) = \psi(b)\}$. Conversely, let $(y, b) \in Y^{(1)} \oplus B^{(1)}$, $f(y) = \psi(b)$ and n be an arbitrary positive integer. Then, $y = ny_1$ and $b = nb_1$ for some $y_1 \in Y$, $b_1 \in B$. Since C is torsion-free, $n(f(y_1) - \psi(b_1)) = 0$ deduce that $f(y_1) = \psi(b_1)$. Hence, $(y, b) \in nX$. Since n be arbitrary, $(y, b) \in X^{(1)}$. An easy calculation shows that $\overline{X^{(1)}} = N$. Clearly, $\phi' = \bar{\mu}$ and $\psi' = \bar{\nu}$. Hence, $0 \rightarrow \overline{A^{(1)}} \xrightarrow{\bar{\mu}} \overline{X^{(1)}} \xrightarrow{\bar{\nu}} \overline{Y^{(1)}} \rightarrow 0$ is an extension. \square

Lemma 2.11. Let $A \in \ell$ be a divisible group and $C \in \mathcal{L}$. Then, a pushout of a TFU extension of A by C is a TFU extension.

Proof. Let $E : 0 \rightarrow A \xrightarrow{\phi} B \rightarrow C \rightarrow 0$ be a TFU extension and $f : A \rightarrow G$ a proper morphism. Then, $fE : 0 \rightarrow G \rightarrow X \rightarrow C \rightarrow 0$ is a pushout of E , where $X = (G \oplus B)/H$ and $H = \{(-f(a), \phi(a)); a \in A\}$ (See [5]). Since E is a TFU extension, $E' : 0 \rightarrow A \rightarrow \overline{B^{(1)}} \rightarrow \overline{C^{(1)}} \rightarrow 0$ is an extension. Hence, hE' is an extension where $h : A \rightarrow \overline{G^{(1)}}$ defined by $h(a) = f(a)$ for every $a \in A$. But,

$hE' : 0 \rightarrow \overline{G^{(1)}} \rightarrow Y \rightarrow \overline{C^{(1)}} \rightarrow 0$ where $Y = (\overline{G^{(1)}} \oplus \overline{B^{(1)}})/K$ and $K = \{(-h(a), \phi(a)); a \in A\}$. Clearly, $K = H$. Since H is a closed, divisible subgroup of $G \oplus B$, so by Lemma 2.3 and Remark 2.1, $\overline{X^{(1)}} = (\overline{G^{(1)}} \oplus \overline{B^{(1)}})/H = Y$. Hence, fE is a *TFU* extension. \square

3 Splitting of *TFU* extensions

An extension is called split if it is equivalent to the trivial extension. Let $Ext_{tfu}(C, A)$ be the class of all equivalence classes of *TFU* extensions of A by C . Recall that for groups $A, C \in \ell$, $Ext(C, A) = 0$ (or $Ext_{tfu}(C, A) = 0$) deduce that every extension (or *TFU* extension) of A by C splits. In this section, we establish some conditions on A and C such that $Ext_{tfu}(C, A) = 0$.

Lemma 3.1. *Let A be a discrete group such that $Ext_{tfu}(X, A) = 0$ for all groups $X \in \ell$. Then, A is a reduced group such that A/tA is a divisible group.*

Proof. Let A be a discrete group such that $Ext_{tfu}(X, A) = 0$ for all groups $X \in \ell$. Let D be a divisible subgroup of A and C a connected group. By Corollary 2.10 of [6], we have the following exact sequence

$$Hom(C, A/D) \rightarrow Ext(C, D) \xrightarrow{i_*} Ext(C, A)$$

Since C is a connected group and A/D a discrete group, $Hom(C, A/D) = 0$. Hence, i_* is injective. By Lemma 2.11, $i_*(Ext_{tfu}(C, D)) \subseteq Ext_{tcf}(C, A) = 0$. So, $Ext_{tcf}(C, D) = 0$. By Corollary 2.4, $Ext(C, D) = 0$. Hence, $D = 0$ (see Theorem 3.3 of [6]). So, A is a reduced group. Now, we show that A/tA is a divisible group. By Corollary 2.6, $Pext((\mathbb{Q}/\mathbb{Z}), A) \subseteq Ext_{tfu}((\mathbb{Q}/\mathbb{Z}), A) = 0$. So, $Ext((\mathbb{Q}/\mathbb{Z}), A/tA) = 0$. Hence, A/tA is a divisible group (see the proof of Theorem 2 of [9]). \square

Lemma 3.2. *Let G be a compact group such that $Ext_{tfu}(X, G) = 0$ for all groups $X \in \ell$. Then, $G \cong \bigoplus (\mathbb{R}/\mathbb{Z})^\sigma \bigoplus \mathbb{R}^n \bigoplus M$, where n is a positive integer, σ a cardinal number and M is a direct product of finite cyclic groups.*

Proof. Let G be a compact group and $Ext_{tfu}(X, G) = 0$ for all groups $X \in \ell$. Then, $Ext_{tfu}(C, G) = 0$ for all connected groups $C \in \ell$. Consider the exact sequence $0 \rightarrow G_0 \xrightarrow{i} G \rightarrow G/G_0 \rightarrow 0$. We have the following exact sequence

$$0 = Hom(C, G/G_0) \rightarrow Ext(C, G_0) \xrightarrow{i_*} Ext(C, G)$$

Hence, i_* is injective. By Lemma 2.11, $i_*(Ext_{tfu}(C, G_0)) \subseteq Ext_{tcf}(C, G) = 0$. So, $Ext_{tcf}(C, G_0) = 0$. By Corollary 2.4, $Ext(C, G_0) = 0$. Hence, $G_0 \cong (\mathbb{R}/\mathbb{Z})^\sigma \bigoplus \mathbb{R}^n$ (see Theorem 3.3 of [6]). By Corollary 3.4 of [6], G_0 splits in G . So, $G \cong G_0 \bigoplus G/G_0$. Set $M = G/G_0$. Then, M is a compact totally disconnected group. By Corollary 2.6, $Pext(X, G) \subseteq Ext_{tfu}(X, G) = 0$ for all compact totally disconnected groups X . So, $Pext(X, M) = 0$ for all compact totally disconnected groups X . Let Y be a compact group. By Proposition 2 of [4], we have the following exact sequence

$$0 = Pext(Y/Y_0, M) \rightarrow Pext(Y, M) \rightarrow Pext(Y_0, M)$$

By Theorem 4.2 of [2], $Pext(Y_0, M) = 0$. Hence, $Pext(Y, M) = 0$ for all compact groups $Y \in \ell$. By Proposition 53.4 of [3] and Lemma 2.3 of [8], \hat{M} is a direct sum of finite cyclic groups. Hence, M is a direct product of finite cyclic groups. \square

Lemma 3.3. *Let A be a σ -compact group such that $A^{(1)} = 0$ and C , a divisible group in ℓ . Then, every TFU pure extension of A by C splits.*

Proof. Let $E : 0 \rightarrow A \xrightarrow{\phi} B \xrightarrow{\psi} C \rightarrow 0$ be a TFU pure extension. An easy calculation shows that $B = \overline{B^{(1)}} + \phi(A)$. Since $\phi(A)$ is pure in B , $\overline{B^{(1)}} \cap \phi(A) = 0$. Hence, by Corollary 3.2 of [5], $B = \overline{B^{(1)}} \oplus \phi(A)$. So, E splits. \square

Lemma 3.4. *Let A be a discrete, torsion-free group. Then, $A \cong A^{(1)} \oplus A/A^{(1)}$.*

Proof. Let A be a discrete, torsion-free group. First, we show that $A/A^{(1)}$ is torsion-free. Let $m(a + A^{(1)}) = 0$ for some positive integer m . Then, $ma \in A^{(1)}$. Hence, $ma = mna_1$ for an arbitrary positive integer n . So, $a = na_1$. Since n be arbitrary, $a \in A^{(1)}$. So, $A^{(1)}$ is pure in A . On the other hand, $A^{(1)} \subseteq nA$ for every positive integer n . Hence, $A^{(1)}$ is a divisible group. By Theorem 21.1 of [3], $A \cong A^{(1)} \oplus A/A^{(1)}$. \square

Lemma 3.5. *Let A be a discrete, torsion-free group such that $Ext_{tfu}(A, X) = 0$ for all groups $X \in \ell$. Then, $A \cong (\bigoplus_{\sigma} \mathbb{Z}) \oplus D$ where D is a discrete, torsion-free divisible group.*

Proof. Let A be a discrete, torsion-free group such that $Ext_{tfu}(A, X) = 0$ for all groups $X \in \ell$. By Lemma 3.4, $A \cong A^{(1)} \oplus A/A^{(1)}$. By Lemma 2.10, $Ext_{tfu}(A/A^{(1)}, X) = 0$ for all groups $X \in \ell$. Hence, $Ext(A/A^{(1)}, X) = 0$ for all groups $X \in \ell$ (see Corollary 2.7). By Theorem 3.3 of [?], $A/A^{(1)} \cong \bigoplus_{\sigma} \mathbb{Z}$. \square

Lemma 3.6. *Let G be a compact, torsion-free group. Then, $Ext_{tfu}(G, X) = 0$ for all groups $X \in \ell$ if and only if $G = 0$.*

Proof. Let G be a compact, torsion-free group and $Ext_{tfu}(G, X) = 0$ for all groups $X \in \ell$. Let X be a totally disconnected group in ℓ . Consider the following exact sequence

$$0 = Hom(G_0, X) \rightarrow Ext(G/G_0, X) \xrightarrow{\pi_*} Ext(G, X)$$

By Lemma 2.10, $\pi_*(Ext_{tfu}(G/G_0, X)) \subseteq Ext_{tfu}(G, X) = 0$. So, $Ext_{tfu}(G/G_0, X) = 0$. By Corollary 2.6, $Ext(G/G_0, X) = 0$. Hence, $G = G_0$ (see Theorem 3.5 of [6]). Since \mathbb{Q} is divisible, $Ext(G, \mathbb{Q}) = 0$ which is a contradiction by Lemma 2.9 of [1]. So, $G = 0$. \square

References

[1] A. A. Alijani, On generalized ℓ -cotorsion LCA groups, arXiv:2210.04178[math.GR].

- [2] A. A. Alijani and H. Sahleh, On t-extensions of abelian groups, *Khayyam J. Math*, 1 (5) (2019), 78-86.
- [3] L. Fuchs, *Infinite Abelian Groups*, Vol. 1, Academic Press, (1970).
- [4] R. O. Fulp, Homological study of purity in locally compact groups, *Proc. London Math. Soc*, 21 (1970), 501-512.
- [5] R.O. Fulp and P. Griffith, Extensions of locally compact abelian groups I, *Trans. Amer. Math. Soc*, 154 (1971),341-356.
- [6] R. O. Fulp and P. Griffith, Extensions of locally compact abelian groups II, *Trans. Amer. Math. Soc*, 154 (1971), 357-363.
- [7] E. Hewitt and K. Ross, *Abstract Harmonic Analysis*, vol. 1, 2nd edn, Springer-Verlog, Berlin (1979).
- [8] P. Loth, Pure extensions of locally compact abelian groups, *Rend. Sem. Mat. Univ. Padova*, 116 (2006), 31-40.
- [9] H. Sahleh and A. A. Alijani, The pure injectives and pure projectives in the category of totally disconnected, locally compact abelian groups, *Math. Sci.* 7(49) (2013).