

UMPA, Ecole Normale Supérieure de Lyon
Email: alexei.tsygvintsev@ens-lyon.fr

ON SOME COLLINEAR CONFIGURATIONS IN THE PLANAR THREE-BODY PROBLEM

ALEXEI TSYGVINTSEV

ABSTRACT. In this paper, we investigate collinear configurations in the planar Newtonian three-body problem where the three bodies or their velocities are aligned. These configurations, also known as generalised syzygies, were previously studied by the author in [4] for bounded solutions. Here, we extend our study to negative energy cases and provide new bounds. Among other results, we present an independent proof of Montgomery’s result [3] that, except for Lagrange’s solution, every solution to the zero angular momentum case suffers syzygies. Our approach is algebraic and based on elementary Sturm-Liouville theory.

1. INTRODUCTION

We consider the motion of three points with masses m_1 , m_2 , and m_3 , denoted P_1 , P_2 , and P_3 , respectively, in the plane. These points have positions $(x_i, y_i) \in \mathbb{R}^2$, where $i = 1, 2, 3$.

The Newtonian three-body problem involves finding the motion of three point masses under the influence of their mutual gravitational attraction.

The equations of motion can be written in the complex compact form as follows

$$\ddot{z}_1 = m_2 \frac{z_{21}}{|z_{21}|^3} - m_3 \frac{z_{13}}{|z_{13}|^3}, \quad \ddot{z}_2 = m_3 \frac{z_{32}}{|z_{32}|^3} - m_1 \frac{z_{21}}{|z_{21}|^3}, \quad \ddot{z}_3 = m_1 \frac{z_{13}}{|z_{13}|^3} - m_2 \frac{z_{32}}{|z_{32}|^3}, \quad (1.1)$$

where $z_k = x_k + iy_k \in \mathbb{C}$, $k = 1, 2, 3$ and $z_{kl} = z_k - z_l$. We assume that the total linear momentum is zero:

$$\sum_k m_k \dot{z}_k = \sum_k m_k z_k = 0, \quad (1.2)$$

by placing the centre of mass at the origin of the coordinate system.

The word *syzygy* (from Late Latin *sȳzygia* = “conjunction”) historically been used by astronomers to describe the alignment of celestial bodies, and in this context, it refers to a configuration where all three points lie on a straight line. In general, a solution of the

N -body problem is said to have a syzygy at $t = t_0$ if at that moment all bodies belong to a certain straight line.

Let $t \mapsto z_i(t)$, $i = 1, 2, 3$ be any solution of the equations (1.1) defined for $t \in I = [0, a)$, $a > 0$. In our work [4] we proposed to study the natural generalisation of syzygies by adding the supplementary condition of collinearity of velocities.

Definition 1.1. *The three bodies P_1, P_2, P_3 form a generalised syzygy at the moment $t_0 \in I$ if at least one of the complex triplets $(z_1, z_2, z_3)(t_0)$ (positions) or $(\dot{z}_1, \dot{z}_2, \dot{z}_3)(t_0)$ (velocities) belongs to the same straight line passing through the origin (see Figure 1).*




FIGURE 1. A generalised syzygy: (A) – the positions are collinear (eclipse), (B) – the velocities (dashed arrows) are collinear

Montgomery showed in his work [3] that, except for the Lagrange solution, every solution to the Newtonian three-body problem with zero angular momentum and negative energy suffers encounters a syzygy, which is a collinear configuration of positions. Diacu demonstrated in [2] that the set of initial conditions leading to syzygy solutions in the planar three-body problem is non-empty and open.

In particular, the orbit of one particle crosses the line of the other two and can not be tangent to this line in the transition point. Generalised syzygies may occur more frequently, as they include collinear configurations of velocities as well. In Chapter 2, we present a simple algebraic condition that defines an open set of initial conditions leading to a generalised syzygy, which holds for arbitrary angular momentum.

Non-syzygy solutions with nonzero angular momentum are still relatively unexplored. In Chapter 3, we will demonstrate that in the periodic case, all non-syzygy solutions must satisfy certain special geometric restrictions. Finally, in Chapter 4, we provide a new elementary proof of Montgomery’s result [3] on the existence of syzygies in the zero angular momentum case.

2. SUFFICIENT CONDITION FOR THE EXISTENCE OF GENERALISED SYZYGIES

The goal of this section is to describe a simple geometric condition, imposed on the initial positions and velocities, that guarantees the existence of generalised syzygies.

Definition 2.1. *The configuration of the bodies P_1 , P_2 and P_3 at the moment $t_0 \in I$ is called antisymmetric, if the oriented areas of parallelograms spanned by the vectors $(z_j(t_0), z_k(t_0))$ and $(\dot{z}_j(t_0), \dot{z}_k(t_0))$ are nonzero and have opposite signs for some $j \neq k$ (see Figure 2). This condition is equivalent algebraically to*

$$\begin{vmatrix} x_j & y_j \\ x_k & y_k \end{vmatrix} (t_0) \cdot \begin{vmatrix} \dot{x}_j & \dot{y}_j \\ \dot{x}_k & \dot{y}_k \end{vmatrix} (t_0) = (x_j y_k - y_j x_k)(t_0) \cdot (\dot{x}_j \dot{y}_k - \dot{y}_j \dot{x}_k)(t_0) < 0. \quad (2.1)$$




FIGURE 2. Antisymmetric configuration of three bodies

Proposition 2.1. *The property of being antisymmetric is independent of the choice of the pair of the bodies: once verified for a particular pair of bodies P_j, P_k , $j \neq k$, the condition (2.1) will be also satisfied for all other possible choices of pairs P_n, P_m , $n \neq m$.*

Proof. We observe that the condition (2.1) is invariant under the permutation of j and k . Without loss of generality, we can assume that (2.1) holds for $(j, k) = (1, 2)$ and show that it is true also for the case where $(j, k) = (2, 3)$. The proof for other cases is similar.

The product of two determinants in (2.1) for $j = 2, k = 3$ can be written as

$$\delta = \underbrace{\begin{vmatrix} x_2 & y_2 \\ x_3 & y_3 \end{vmatrix} \cdot \begin{vmatrix} \dot{x}_2 & \dot{y}_2 \\ \dot{x}_3 & \dot{y}_3 \end{vmatrix}}_a = \frac{1}{m_2^2 m_3^2} \begin{vmatrix} m_2 x_2 & m_2 y_2 \\ m_3 x_3 & m_3 y_3 \end{vmatrix} \cdot \begin{vmatrix} m_2 \dot{x}_2 & m_2 \dot{y}_2 \\ m_3 \dot{x}_3 & m_3 \dot{y}_3 \end{vmatrix}. \quad (2.2)$$

As can be seen from the equalities (1.2):

$$\begin{aligned} [m_3 x_3, m_3 y_3] &= -[m_1 x_1, m_1 y_1] - [m_2 x_2, m_2 y_2], \\ [m_3 \dot{x}_3, m_3 \dot{y}_3] &= -[m_1 \dot{x}_1, m_1 \dot{y}_1] - [m_2 \dot{x}_2, m_2 \dot{y}_2]. \end{aligned} \quad (2.3)$$

Therefore, by using (2.2), (2.3), and the elementary properties of determinants, we have:

$$\delta = \frac{1}{m_2^2 m_3^2} \begin{vmatrix} m_2 x_2 & m_2 y_2 \\ -m_1 x_1 & -m_1 y_1 \end{vmatrix} \cdot \begin{vmatrix} m_2 \dot{x}_2 & m_2 \dot{y}_2 \\ -m_1 \dot{x}_1 & -m_1 \dot{y}_1 \end{vmatrix} = \frac{m_1^2}{m_3^2} \underbrace{\begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \end{vmatrix} \cdot \begin{vmatrix} \dot{x}_1 & \dot{y}_1 \\ \dot{x}_2 & \dot{y}_2 \end{vmatrix}}_b \quad (2.4)$$

Obviously, the sign of a in (2.2) and b in (2.4) are the same. That finishes the proof. \square

The author has shown in [4] that the existence of a generalised syzygy can be guaranteed by the antisymmetry condition (2.1), provided that the mutual distances of the three bodies remain bounded. The following theorem generalises this result to the case of negative energy, which includes unbounded (escape) solutions in particular. Moreover, the proof presented here is simpler and more straightforward.

Theorem 2.1. *Let $t \mapsto (z_1(t), z_2(t), z_3(t))$ be a solution of the three-body problem (1.1) with negative energy $H = -\alpha$, $\alpha > 0$. We assume that the initial configuration of the bodies P_1, P_2, P_3 at $t = 0$ is antisymmetric and the solution is collision free for $t \in [0, T]$ where*

$$T(\alpha) = \frac{\pi \Sigma}{\alpha^{3/2}}, \quad \Sigma = \frac{(m_3 m_2)^{3/2}}{m_{32}^{1/2}} + \frac{(m_1 m_3)^{3/2}}{m_{13}^{1/2}} + \frac{(m_2 m_1)^{3/2}}{m_{21}^{1/2}}. \quad (2.5)$$

Then there exists $t \in [0, T]$ such that the three bodies attain a generalised syzygy at time $t = t_0$.

Proof. After introducing the new variables $w_i = m_i z_i$, $i = 1, 2, 3$ the relations (1.2) yield

$$\sum_i w_i = 0. \quad (2.6)$$

Writing

$$w_k = X_k + iY_k, \quad X_k = m_k x_k, \quad Y_k = m_k y_k, \quad k = 1, 2, 3, \quad (2.7)$$

and using (2.6), one derives from the equations (1.1) the following 2×2 matrix equation

$$\ddot{X} = AX, \quad X = \begin{bmatrix} X_1 & Y_1 \\ X_2 & Y_2 \end{bmatrix}, \quad A = \begin{bmatrix} -m_2\rho_3 - m_{13}\rho_2 & m_1\rho_{32} \\ m_2\rho_{31} & -m_1\rho_3 - m_{32}\rho_1 \end{bmatrix}, \quad (2.8)$$

where

$$\rho_1 = 1/|z_{32}|^3, \quad \rho_2 = 1/|z_{13}|^3, \quad \rho_3 = 1/|z_{21}|^3, \quad m_{ij} = m_i + m_j, \quad \rho_{ij} = \rho_i - \rho_j. \quad (2.9)$$

Proposition 2.2. *Both definitions of a generalised syzygy and an antisymmetric configuration, are invariant when replacing (x_i, y_i) by (X_i, Y_i) and (\dot{x}_i, \dot{y}_i) by (\dot{X}_i, \dot{Y}_i) for $i = 1, 2, 3$.*

Proof. It follows immediately from (2.7) and the identity below, verified for any $j \neq k$:

$$\begin{vmatrix} x_j & y_j \\ x_k & y_k \end{vmatrix} \cdot \begin{vmatrix} \dot{x}_j & \dot{y}_j \\ \dot{x}_k & \dot{y}_k \end{vmatrix} = \frac{1}{m_j^2 m_k^2} \begin{vmatrix} X_j & Y_j \\ X_k & Y_k \end{vmatrix} \cdot \begin{vmatrix} \dot{X}_j & \dot{Y}_j \\ \dot{X}_k & \dot{Y}_k \end{vmatrix}. \quad (2.10)$$

□

We consider the two matrices

$$X = \begin{bmatrix} X_1 & Y_1 \\ X_2 & Y_2 \end{bmatrix}, \quad \dot{X} = \begin{bmatrix} \dot{X}_1 & \dot{Y}_1 \\ \dot{X}_2 & \dot{Y}_2 \end{bmatrix}, \quad (2.11)$$

and denote their determinants respectively by $\Delta_1 = \det(X)$ and $\Delta_2 = \det(\dot{X})$.

Let us compute the second-order derivative of the determinant Δ_1 :

$$\dot{\Delta}_1 = \begin{vmatrix} \dot{X}_1 & Y_1 \\ \dot{X}_2 & Y_2 \end{vmatrix} + \begin{vmatrix} X_1 & \dot{Y}_1 \\ X_2 & \dot{Y}_2 \end{vmatrix}, \quad \ddot{\Delta}_1 = \begin{vmatrix} \ddot{X}_1 & Y_1 \\ \ddot{X}_2 & Y_2 \end{vmatrix} + \begin{vmatrix} X_1 & \ddot{Y}_1 \\ X_2 & \ddot{Y}_2 \end{vmatrix} + 2 \begin{vmatrix} \dot{X}_1 & \dot{Y}_1 \\ \dot{X}_2 & \dot{Y}_2 \end{vmatrix}. \quad (2.12)$$

Introducing $x = [X_1, X_2]^T$, $y = [Y_1, Y_2]^T$ and using the equation (2.8), we obtain $\ddot{x} = Ax$, $\ddot{y} = Ay$. Therefore, $\ddot{\Delta}_1$ can be written as follows

$$\ddot{\Delta}_1 = \det([Ax, y]) + \det([x, Ay]) + 2 \det([\dot{x}, \dot{y}]) = \det([Ax, y]) + \det([x, Ay]) + 2\Delta_2. \quad (2.13)$$

The simple algebraic computation shows that $\det([Ax, y]) + \det([x, Ay]) = \text{Tr}(A)\Delta_1$.

Thus, as follows from (2.13),

$$\ddot{\Delta}_1 = \text{Tr}(A)\Delta_1 + 2\Delta_2, \quad (2.14)$$

where

$$\text{Tr}(A) = -(m_{32}\rho_1 + m_{13}\rho_2 + m_{21}\rho_3). \quad (2.15)$$

Lemma 2.1. *Let $t \mapsto z_i(t)$, $i = 1, 2, 3$ be any solution of the three-body problem (1.1) with negative energy $H = -\alpha$, where $\alpha > 0$. Then $\text{Tr}(A) \leq -\alpha^3/\Sigma^2$ along this solution with Σ defined in (2.5).*

Proof. By introducing $r_i = 1/\rho_i^{1/3}$, $i = 1, 2, 3$ the total energy of the three-body problem (1.1) can be written as

$$H = K - U(r), \quad U(r) = m_3m_2r_1 + m_1m_3r_2 + m_2m_1r_3, \quad r = (r_1, r_2, r_3), \quad (2.16)$$

where $K \geq 0$ is the kinetic energy.

Since $K - U = -\alpha$ and $K \geq 0$, one obtains $U \geq \alpha$.

Writing $F(r) = -\text{Tr}(A)(r) = m_{32}r_1^3 + m_{13}r_2^3 + m_{21}r_3^3$, we shall calculate, for any $s > 0$, the minimum of the function $r \mapsto F(r)$ on the compact set

$$K_s = \{r \in \mathbb{R}^3 \mid U(r) = s, r_i \geq 0, i = 1, 2, 3\}. \quad (2.17)$$

Because K_s is convex (triangle) and F is a convex function on K_s (since $d^2F \geq 0$ on K), it is sufficient to determine its local minimum. One computes:

$$\nabla U = (m_3m_2, m_1m_3, m_2m_1), \quad \nabla F = 3(m_{32}r_1^2, m_{13}r_2^2, m_{21}r_3^2). \quad (2.18)$$

The Lagrange multiplier λ found from the equations $\nabla F = \lambda \nabla U$, $U = s$ is given by

$$\lambda = \frac{3s^2}{\Sigma^2}, \quad (2.19)$$

with Σ defined in (2.5).

The corresponding extremum point is

$$r^* = (r_1^*, r_2^*, r_3^*) = \frac{s}{\Sigma} \left(\sqrt{\frac{m_3m_2}{m_{32}}}, \sqrt{\frac{m_1m_3}{m_{13}}}, \sqrt{\frac{m_2m_1}{m_{21}}} \right) \in K_s, \quad (2.20)$$

which is a local minimum of F because $d^2F(r^*)$ is strictly positive and $d^2U = 0$. Thus, by substitution:

$$\min_{r \in K_s} F(r) = F(r^*) = \frac{s^3}{\Sigma^2}. \quad (2.21)$$

Hence, considering $s \geq \alpha$, one shows that $U(r) \geq \alpha$ implies $F(r) = -\text{Tr}(A)(r) \geq \frac{\alpha^3}{\Sigma^2}$. The proof of the Lemma 2.1 is finished. \square

Writing $d = \Delta_2/\Delta_1$, the equation (2.14) can be transformed into the second-order Hill's linear differential equation:

$$\ddot{\Delta}_1 = (\text{Tr}(A) + 2d)\Delta_1. \quad (2.22)$$

We suppose now that the solution of (1.1) $t \mapsto z_i(t)$, $i = 1, 2, 3$ is defined in the interval $[0, T]$, with T given in (2.5). Assuming it starts at $t = 0$ from the antisymmetric configuration and that no generalised syzygy happens for any $0 < t \leq T$ we obtain

$$d(t) = \frac{\Delta_2}{\Delta_1}(t) < 0, \quad \Delta_1(t) \neq 0, \quad \forall t \in [0, T]. \quad (2.23)$$

Therefore, according to (2.22), (2.23) and the Lemma 2.1:

$$\ddot{\Delta}_1 = \phi \Delta_1, \quad \phi = \text{Tr}(A) + 2d \leq -\theta^2, \quad \theta^2 = \alpha^3 / \Sigma^2, \quad t \in [0, T]. \quad (2.24)$$

As follows from the zero comparison theorem of the Sturm-Liouville theory [1], the solution Δ_1 of (2.24) always has a zero between any two consecutive zeros of any solution y of the equation $\ddot{y} = -\theta^2 y$ whose general solution is $y(t) = A \cos(\theta t + \phi_0)$, $A, \phi_0 \in \mathbb{R}$. For a nonzero A , every two consecutive zeros of y are separated by an interval of the length π/θ . Hence, $\Delta_1(t_0) = 0$ for some $0 < t_0 < T = \pi/\theta$ with T given by the formula (2.5). That is in contradiction with (2.23). The proof of Theorem 2.1 is finished. \square

3. EXISTENCE OF SYZYGIES FOR PERIODIC SOLUTIONS

In this section we will show that any periodic solution of the three-body problem will have a syzygy, i.e. exhibits a collinear configuration of the bodies as soon as some geometrical constraints are imposed on the shape of the triangle formed by the bodies. Not every periodic solution has a syzygy; for example, the Lagrange periodic solution has the three bodies situated at the vertices of an equilateral triangle and never aligned.

Definition 3.1. *A periodic solution of the three-body problem (1.1) is called θ -rigid if there exists a non-zero vector $\theta = (\theta_1, \theta_2, \theta_3) \in \mathbb{R}^3$ such that*

$$\theta_1(\rho_3(t) - \rho_2(t)) + \theta_2(\rho_1(t) - \rho_3(t)) + \theta_3(\rho_2(t) - \rho_1(t)) \geq 0, \quad \forall t \in \mathbb{R}, \quad (3.1)$$

where the sum is strictly positive for at least one $t_0 \in \mathbb{R}$.

For instance, let us consider a periodic solution for which

$$|z_{32}(t)| > |z_{13}(t)| \Leftrightarrow \rho_1(t) < \rho_2(t), \quad \forall t \in \mathbb{R}. \quad (3.2)$$

Thus, it is θ -rigid with respect to the vector $(\theta_1, \theta_2, \theta_3) = (0, 0, 1)$.

We note that the Lagrange equilateral solution is not θ -rigid for any choice of θ . Indeed, in this case $\rho_1 = \rho_2 = \rho_3$ and, therefore, the sum in (3.1) is always zero.

One might ask whether it is possible to have a periodic solution without syzygies, in which one side of the triangle is always smaller than the other, and one of the conditions

of the form (3.2) is satisfied. Our next theorem provides a negative response to this question.

Theorem 3.1. *Every θ -rigid periodic solution to the three-body problem (1.1) admits a syzygy.*

Proof. Let $t \mapsto z_i(t)$, $i = 1, 2, 3$ be a periodic solution of the three-body problem (1.1) with period $\tau > 0$.

Let us suppose that it is θ -rigid and has no syzygies i.e. $\Delta_1(t) = \det(X(t)) \neq 0$, $\forall t \in [0, \tau]$. Without loss of generality, we can assume that

$$\Delta_1(t) > 0, \quad \forall t \in [0, \tau]. \quad (3.3)$$

Let $S_i = X_i \dot{Y}_i - \dot{X}_i Y_i$, $i = 1, 2, 3$ be three oriented areas of parallelograms formed by the vectors (X_i, Y_i) and (\dot{X}_i, \dot{Y}_i) . Then, using (2.8) and (2.6), one derives the following equations

$$\dot{S}_1 = m_1 \Delta_1 (\rho_3 - \rho_2), \quad \dot{S}_2 = m_2 \Delta_1 (\rho_1 - \rho_3), \quad \dot{S}_3 = m_3 \Delta_1 (\rho_2 - \rho_1). \quad (3.4)$$

Thus,

$$\frac{d}{dt} \left(\sum_{i=1}^3 \frac{\theta_i \dot{S}_i}{m_i} \right) = \Delta_1 S, \quad S = \theta_1 (\rho_3 - \rho_2) + \theta_2 (\rho_1 - \rho_3) + \theta_3 (\rho_2 - \rho_1), \quad t \in [0, \tau]. \quad (3.5)$$

Integrating (3.5) and using the periodicity of S_i , $i = 1, 2, 3$ we find

$$\int_0^\tau \Delta_1 S dt = 0, \quad (3.6)$$

which obviously contradicts (3.3) and that $\Sigma(t_0) > 0$ for some $t_0 \in [0, \tau]$. The proof is finished. \square

4. ZERO ANGULAR MOMENTUM CASE

Let $\gamma : t \mapsto (z_1(t), z_2(t), z_3(t))$ be a zero angular momentum solution with negative energy of the three-body problem (1.1). In this section we prove that if γ is collision free for a sufficiently long period of time, it will always encounter a syzygy. A similar result was first established by Montgomery in [3] using the shape sphere approach. Our proof is essentially algebraic and is more elementary.

The function $t \mapsto C(t) = \dot{X} X^{-1}$ satisfies, using (2.8), the following matrix Riccati equation

$$\dot{C} + C^2 = A. \quad (4.1)$$

Combined with the Cayley-Hamilton identity

$$C^2 - \text{Tr}(C)C + \det(C)I_2 = 0, \quad (4.2)$$

the equation (4.1) yields

$$\dot{C} + \text{Tr}(C)C = A + \det(C)I_2. \quad (4.3)$$

Applying Liouville's formula to the equation $\dot{X} = CX$, we obtain:

$$\dot{\Delta}_1 = \text{Tr}(C)\Delta_1. \quad (4.4)$$

Multiplying both sides of equation (4.3) by Δ_1 , we get

$$\frac{d}{dt}(\Delta_1 C) = \Delta_1 A + \Delta_1 \det(C)I_2 = \Delta_1 A + \Delta_2 I_2, \quad (4.5)$$

where we have used $\det(C) = \Delta_2/\Delta_1$.

Introducing the adjugate matrix \tilde{X} of X and using $C = \dot{X}X^{-1}$, we compute

$$\Delta_1 C = \dot{X}\tilde{X} = \begin{bmatrix} \dot{X}_1 & \dot{Y}_1 \\ \dot{X}_2 & \dot{Y}_2 \end{bmatrix} \begin{bmatrix} Y_2 & -Y_1 \\ -X_2 & X_1 \end{bmatrix}. \quad (4.6)$$

The matrix $A(\rho_1, \rho_2, \rho_3)$, defined in (2.8) can be written as a linear combination

$$A = \rho_1 A_1 + \rho_2 A_2 + \rho_3 A_3, \quad (4.7)$$

where A_1 , A_2 , and A_3 are 2×2 constant matrices that depend on the masses and are defined by

$$A_1 = \begin{bmatrix} 0 & 0 \\ -m_2 & -m_{32} \end{bmatrix}, \quad A_2 = \begin{bmatrix} -m_{13} & -m_1 \\ 0 & 0 \end{bmatrix}, \quad A_3 = \begin{bmatrix} -m_2 & m_1 \\ m_2 & -m_1 \end{bmatrix}. \quad (4.8)$$

Let $E = M_3(\mathbb{R})$ be the Euclidean space of real 2×2 matrices equipped with the inner product \langle, \rangle defined by $\langle A, B \rangle = \text{Tr}(A^T B)$ for $A, B \in E$.

We check that $A_1, A_2, A_3 \in E$ are linearly independent and span a 3-dimensional vector subspace of E . Furthermore, we have the identity

$$-\frac{A_1 + A_2 + A_3}{M} = I_2, \quad M = \sum m_i. \quad (4.9)$$

Combining equations (4.7), (4.9) and (4.3), we obtain

$$\frac{d}{dt}(\dot{X}\tilde{X}) = \left(\Delta_1 \rho_1 - \frac{\Delta_2}{M}\right) A_1 + \left(\Delta_1 \rho_2 - \frac{\Delta_2}{M}\right) A_2 + \left(\Delta_1 \rho_3 - \frac{\Delta_2}{M}\right) A_3. \quad (4.10)$$

Let us check, with help of the equation (4.10), constancy of the angular momentum. We introduce the matrix

$$L = \begin{bmatrix} -m_3^{-1} & m_1^{-1} + m_3^{-1} \\ -m_2^{-1} - m_3^{-1} & m_3^{-1} \end{bmatrix}. \quad (4.11)$$

It can be checked that $\langle L, A_i \rangle = 0$, $i = 1, 2, 3$.

Taking the inner product of both sides of the equation (4.10) with L , we find that

$$\frac{d}{dt} \langle (\dot{X}\tilde{X}), L \rangle = 0, \quad (4.12)$$

i.e. the function $I = \langle \dot{X}\tilde{X}, L \rangle = \text{const}$ is time independent.

One can easily verify that I , written as a function of variables X_i, Y_j , $1 \leq i, j \leq 2$ (by excluding (X_3, Y_3) with help of (2.6)) is the angular momentum of the three-body problem (1.1) :

$$I = \sum_{i=1}^3 m_i R_i \times \dot{R}_i = \sum_{i=1}^3 \frac{S_i}{m_i}, \quad (4.13)$$

where $R_i = (x_i, y_i)$, $1 \leq i \leq 3$.

The equation (4.5) can be written, by replacing Δ_2 using equation (2.14), in the following form

$$\frac{d}{dt} \left(\dot{X}\tilde{X} - \frac{\dot{\Delta}_1}{2} I_2 \right) = \Delta_1 \left(A - \frac{\text{Tr}(A)}{2} I_2 \right). \quad (4.14)$$

Applying (4.7), we can transform it to

$$\frac{d}{dt} \left(\dot{X}\tilde{X} - \frac{\dot{\Delta}_1}{2} I_2 \right) = \Delta_1 \sum_{i=1}^3 \rho_i \tilde{A}_i, \quad (4.15)$$

where $\tilde{A}_i = A_i - \frac{\text{Tr}(A_i)}{2} I_2$, $\text{Tr}(\tilde{A}_i) = 0$, $1 \leq i \leq 3$.

It can be shown, using (4.8), (4.9), that $\tilde{A}_1 + \tilde{A}_2 + \tilde{A}_3 = 0$. Substituting $\tilde{A}_3 = -\tilde{A}_1 - \tilde{A}_2$ in (4.15) we obtain finally

$$\frac{d}{dt} \left(\dot{X}\tilde{X} - \frac{\dot{\Delta}_1}{2} I_2 \right) = \Delta_1 (\rho_1 - \rho_3) \tilde{A}_1 + \Delta_1 (\rho_2 - \rho_3) \tilde{A}_2, \quad (4.16)$$

where

$$\tilde{A}_1 = \begin{bmatrix} m_{32}/2 & 0 \\ -m_2 & -m_{32}/2 \end{bmatrix}, \quad \tilde{A}_2 = \begin{bmatrix} -m_{13}/2 & -m_1 \\ 0 & m_{13}/2 \end{bmatrix}, \quad m_{ij} = m_i + m_j. \quad (4.17)$$

Moreover, as follows from (4.4), (4.6), one always has

$$\text{Tr}(\dot{X}\tilde{X}) = \dot{\Delta}_1, \quad \det(\dot{X}\tilde{X}) = \Delta_1 \Delta_2. \quad (4.18)$$

Let $I = \langle \dot{X}\tilde{X}, L \rangle = k$ where $k \in \mathbb{R}$ is the constant value of the angular momentum. Computing $\langle \dot{X}\tilde{X}, L \rangle$ using (4.6), it is easy to derive the following equality:

$$\dot{X}\tilde{X} - \frac{\dot{\Delta}_1}{2}I_2 = \frac{b}{m_2}\tilde{A}_1 - \frac{a}{m_1}\tilde{A}_2 - k\frac{m_3}{2}J, \quad (4.19)$$

where

$$a = \begin{vmatrix} X_1 & Y_1 \\ \dot{X}_1 & \dot{Y}_1 \end{vmatrix} = S_1, \quad b = \begin{vmatrix} X_2 & Y_2 \\ \dot{X}_2 & \dot{Y}_2 \end{vmatrix} = S_2, \quad J = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (4.20)$$

Therefore, as follows from the equality (4.19), the equation (4.16) is equivalent to the first two differential equations of the system (3.4).

Theorem 4.1. *Let $t \mapsto (z_1(t), z_2(t), z_3(t))$, $t \in [0, T_1]$ be a zero angular momentum collision-free solution to the three-body problem (1.1) with negative energy $H = -\alpha$, $\alpha > 0$ where*

$$T_1(\alpha) = \frac{\sqrt{2\pi}\Sigma}{\alpha^{3/2}}. \quad (4.21)$$

Then there exists $t_0 \in [0, T_1]$ such that the three bodies form a syzygy at the moment $t = t_0$.

Proof. For $k = 0$, the formula (4.19) is given by

$$\dot{X}\tilde{X} - \frac{\dot{\Delta}_1}{2}I_2 = \frac{b}{m_2}\tilde{A}_1 - \frac{a}{m_1}\tilde{A}_2 = \begin{bmatrix} \beta a + \alpha b & a \\ b & -\beta a - \alpha b \end{bmatrix} = R, \quad (4.22)$$

where

$$\alpha = \frac{1}{2} \left(\frac{m_3}{m_2} + 1 \right), \quad \beta = \frac{1}{2} \left(\frac{m_3}{m_1} + 1 \right), \quad (4.23)$$

and

$$-\det(R) = \beta^2 a^2 + (2\alpha\beta - 1)ab + \alpha^2 b^2. \quad (4.24)$$

As follows from (4.23), one always has $\alpha, \beta > 1/2$. Using these inequalities, it is easy to show that the quadratic form in (4.24), as a function of a and b , is positive and therefore

$$\det(R)(a, b) \leq 0, \quad \forall a, b \in \mathbb{R}. \quad (4.25)$$

Since $\text{Tr}(R) = 0$, the eigenvalues of R are real. Hence, according to (4.22), the eigenvalues λ_1, λ_2 of $\dot{X}\tilde{X}$ are also real. Thus, using (4.18), we have the following inequality holding for arbitrary $a, b \in \mathbb{R}$

$$\text{Tr}(\dot{X}\tilde{X})^2 - 4\det(\dot{X}\tilde{X}) = \dot{\Delta}_1^2 - 4\Delta_1\Delta_2 \geq 0. \quad (4.26)$$

Suppose that the solution $t \mapsto (z_1(t), z_2(t), z_3(t))$ is syzygy free in the interval $[0, T_1]$. In particular, we can suppose

$$\Delta_1(t) > 0, \quad \forall t \in [0, T_1], \quad (4.27)$$

such that the function $\delta : t \mapsto \sqrt{\Delta_1(t)}$ is differentiable in this interval.

Differentiating δ twice using (2.14), we obtain

$$\ddot{\delta} = \eta\delta, \quad \eta = \left(\frac{\text{Tr}(A)}{2} - \frac{\dot{\Delta}_1^2 - 4\Delta_1\Delta_2}{4\Delta_1^2} \right), \quad t \in [0, T_1]. \quad (4.28)$$

According to (4.26) and Lemma 2.1:

$$\eta(t) \leq \frac{\text{Tr}(A)}{2} \leq -\zeta^2, \quad \zeta^2 = \frac{\alpha^3}{2\Sigma^2}, \quad t \in [0, T_1]. \quad (4.29)$$

Comparing (4.28) with $\ddot{y} = -\zeta^2 y$, and using the same Sturm-Liouville argument as in the proof of Theorem 2.1, we conclude that δ admits at least one zero in the interval $[0, \pi/\zeta] = [0, T_1]$. This contradicts our hypothesis (4.27). The proof of Theorem 4.1 is finished. \square

5. CONCLUSION

Montgomery [3] showed the existence of syzygies in the three-body problem for the case of zero angular momentum and negative energy, except for the Lagrange homothetic solutions. Our Theorem 2.1 is free from the restriction on the angular momentum and contains both an easy-to-check sufficient condition and an upper bound on the time instant when the generalised syzygy occurs. Theorem 3.1 deals with the collinear configurations (syzygies) in the periodic case. Under the assumption that the triangle formed by the bodies obeys a geometric restriction called θ -rigidity, we show that the bodies become aligned at some instant, resulting in a syzygy in the corresponding solution. We also hypothesise that a similar result holds for bounded non-periodic solutions. A more in-depth analysis of equations (4.10) and (4.1) may reveal additional interesting properties regarding collinear configurations in the three-body problem.

REFERENCES

- [1] R. Carroll, "Transformation theory and application", North-Holland, 1985
- [2] F.N. Diacu, "On the planar syzygy solutions of the 3-body problem", *Celestial Mech Dyn Astr*, 46, 1989, 119-128
- [3] R. Montgomery, "The zero angular momentum, three-body problem: All but one solution has syzygies", *Ergod. Th. Dynam. Sys.*, 27,2007, 311-340
- [4] A. Tsygvintsev, "On the existence of generalised syzygies in the planar three-body problem", *C. R. Acad. Sci. Paris*, vol 361, 2023, 331-335
- [5] A. Wintner, "The analytical foundations of Celestial Mechanics", Princeton University Press, 1941